

F -MANIFOLD ALGEBRAS AND DEFORMATION QUANTIZATION VIA PRE-LIE ALGEBRAS

JIEFENG LIU, YUNHE SHENG, AND CHENGMING BAI

ABSTRACT. The notion of an F -manifold algebra is the underlying algebraic structure of an F -manifold. We introduce the notion of pre-Lie formal deformations of commutative associative algebras and show that F -manifold algebras are the corresponding semi-classical limits. We study pre-Lie infinitesimal deformations and extension of pre-Lie n -deformation to pre-Lie $(n + 1)$ -deformation of a commutative associative algebra through the cohomology groups of pre-Lie algebras. We introduce the notions of pre- F -manifold algebras and dual pre- F -manifold algebras, and show that a pre- F -manifold algebra gives rise to an F -manifold algebra through the sub-adjacent associative algebra and the sub-adjacent Lie algebra. We use Rota-Baxter operators, more generally O -operators and average operators on F -manifold algebras to construct pre- F -manifold algebras and dual pre- F -manifold algebras.

CONTENTS

1. Introduction	1
2. Preliminaries	3
3. F -manifold algebras and pre-Lie deformations of commutative associative algebras	5
3.1. F -manifold algebras and F -manifold-admissible algebras	5
3.2. Pre-Lie deformation quantization of commutative pre-Lie algebras	9
4. Pre- F -manifold algebras, Rota-Baxter operators and O -operators on F -manifold algebras	11
4.1. Representations of F -manifold algebras	12
4.2. Pre- F -manifold algebras	14
4.3. Rota-Baxter operators and O -operators on F -manifold algebras	15
5. Dual pre- F -manifold algebras and average operators on F -manifold algebras	19
References	21

1. INTRODUCTION

The concept of Frobenius manifolds was introduced by Dubrovin in [14] as a geometrical manifestation of the Witten-Dijkgraaf-Verlinde-Verlinde (WDVV) associativity equations in the 2-dimensional topological field theories. Hertling and Manin weakened the conditions of

⁰MSC: 17A30, 53D55, 17B38

Key words and phrases. F -manifold algebra, pre-Lie deformation quantization, pre- F -manifold algebra, Rota-Baxter operator.

a Frobenius manifold and introduced the notion of an F -manifold in [23]. Any Frobenius manifold is an F -manifold. F -manifolds appear in many fields of mathematics such as singularity theory [22], quantum K-theory [25], integrable systems [10, 11, 30], operad [33] and so on.

Motivated by the study of the operad of the underlying algebraic structure of an F -manifold, Dotsenko introduced the notion of an F -manifold algebra in [13] and showed that the graded object of the filtration of the operad encoding pre-Lie algebras is the operad encoding F -manifold algebras. The slogan for F -manifold algebras promoted by Dotsenko is that “ F -manifold algebras are the same to pre-Lie algebras as Poisson algebras to associative algebras”. It is well-known that Poisson algebras can be understood as semi-classical limits of associative formal deformations of commutative associative algebras. So it is natural to ask whether F -manifold algebras can be understood as certain semi-classical limits.

The first aim of this paper is to answer the above question. We introduce the notion of pre-Lie formal deformations of commutative associative algebras and show that F -manifold algebras are the corresponding semi-classical limits. Note that a commutative pre-Lie algebra is associative and hence the aforementioned formal deformation can put into an extent of “pure” pre-Lie algebras, that is, F -manifold algebras are the semi-classical limits of pre-Lie formal deformations of commutative pre-Lie algebras. This result is parallel to that the semi-classical limit of an associative formal deformation of a commutative associative algebra is a Poisson algebra and illustrates the slogan promoted by Dotsenko. Viewing the commutative associative algebra as a pre-Lie algebra, we show that pre-Lie infinitesimal deformation and extension of pre-Lie n -deformation to pre-Lie $(n + 1)$ -deformation of a commutative associative algebra are classified by the second and the third cohomology groups of the pre-Lie algebra respectively.

The notion of a pre-Poisson algebra was introduced by Aguiar in [1], which combines a Zinbiel algebra and a pre-Lie algebra such that some compatibility conditions are satisfied. Zinbiel algebras (also called dual Leibniz algebras) were introduced by Loday in [28] in his study of the algebraic structure behind the cup product on the cohomology groups of the Leibniz algebra. See [27, 29] for more details on Zinbiel algebras. More importantly, a pre-Poisson algebra gives rise to a Poisson algebra naturally through the sub-adjacent commutative associative algebra of the Zinbiel algebra and the sub-adjacent Lie algebra of the pre-Lie algebra. Conversely, a Rota-Baxter operator action (more generally an \mathcal{O} -operator action) on a Poisson algebra gives rise to a pre-Poisson algebra. In this paper, we introduce the notion of a pre- F -manifold algebra, which also contains a Zinbiel algebra and a pre-Lie algebra, such that some compatibility conditions are satisfied. By the sub-adjacent associative algebra and the sub-adjacent Lie algebra, a pre- F -manifold algebra gives rise to an F -manifold algebra naturally. We further introduce the notion of a Rota-Baxter operator (more generally an \mathcal{O} -operator) on an F -manifold algebra, which is simultaneously a Rota-Baxter operator on the underlying associative algebra and a Rota-Baxter operator on the underlying Lie algebra. See [1, 2, 3, 9, 16, 20, 21] for more details on Rota-Baxter operators and \mathcal{O} -operators. A pre- F -manifold algebra can be obtained through the action of a Rota-Baxter operator (more generally an \mathcal{O} -operator). This coincides with the general theory of splitting of operads [4, 34]. The above relations can be summarized into the

following commutative diagram:

$$\begin{array}{ccc}
 \text{Zinbiel algebra} + \text{pre-Lie algebra} & \longrightarrow & \text{pre-}F\text{-manifold algebra} \\
 \text{sub-adjacent} \downarrow \uparrow \text{Rota-Baxter action} & & \text{sub-adjacent} \downarrow \uparrow \text{Rota-Baxter action} \\
 \text{comm associative algebra} + \text{Lie algebra} & \longrightarrow & F\text{-manifold algebra.}
 \end{array}$$

It is well known that the Koszul duals of the operad PreLie of pre-Lie algebras and the operad Zinb of Zinbiel algebras are the operad Perm of permutative algebras and the operad Leib of Leibniz algebras respectively ([29, 8]). We further introduce the notion of a dual pre- F -manifold algebra, which contains a permutative algebra and a Leibniz algebra such that some compatibility conditions hold. By the action of an average operator on an F -manifold algebra, we can obtain a dual pre- F -manifold algebra.

The paper is organized as follows. In Section 2, we recall pre-Lie algebras, Lie-admissible algebras and the cohomology theory of pre-Lie algebras. In Section 3, first we introduce the notion of F -manifold-admissible algebras, which gives rise to F -manifold algebras. Various examples on F -manifold algebras are given. Then we introduce the notion of pre-Lie formal deformations of commutative associative algebras and show that F -manifold algebras are the corresponding semi-classical limits. Furthermore, we study pre-Lie infinitesimal deformations and extensions of pre-Lie n -deformations to pre-Lie $(n + 1)$ -deformations of a commutative associative algebra. In Section 4, first we study representations of an F -manifold algebra. Then we introduce the notions of pre- F -manifold algebras and Rota-Baxter operators (more generally \mathcal{O} -operators) on an F -manifold algebra. We show that on one hand, an \mathcal{O} -operator on an F -manifold algebra gives a pre- F -manifold algebra, and on the other hand, a pre- F -manifold algebra naturally gives an \mathcal{O} -operator on the sub-adjacent F -manifold algebra. More examples of pre- F -manifold algebras and F -manifold algebras are given. In Section 5, we introduce the notions of dual pre- F -manifold algebras and average operators on an F -manifold algebra and show that an average operator on an F -manifold algebra gives a dual pre- F -manifold algebra naturally.

In this paper, all the vector spaces are over algebraically closed field \mathbb{K} of characteristic 0, and finite dimensional.

Acknowledgements. This research was supported by NSFC (11922110, 11901501, 11931009). C. Bai is also supported by the Fundamental Research Funds for the Central Universities and Nankai ZhiDe Foundation.

2. PRELIMINARIES

In this section, we briefly recall pre-Lie algebras, Lie-admissible algebras and the cohomology theory of pre-Lie algebras.

Pre-Lie algebras are a class of nonassociative algebras appearing in many fields in mathematics and mathematical physics (cf. [5, 6, 8, 12, 26, 31], and the survey [7]). In particular, as pointed out in [13], the operad of pre-Lie algebras is “one of the most famous operads in the literature”.

Definition 2.1. A **pre-Lie algebra** is a pair $(\mathfrak{g}, *_\mathfrak{g})$, where \mathfrak{g} is a vector space and $*_\mathfrak{g} : \mathfrak{g} \otimes \mathfrak{g} \longrightarrow \mathfrak{g}$ is a bilinear multiplication satisfying that for all $x, y, z \in \mathfrak{g}$, the associator $(x, y, z) = (x *_\mathfrak{g} y) *_\mathfrak{g} z - x *_\mathfrak{g} (y *_\mathfrak{g} z) = 0$.

$z - x *_g (y *_g z)$ is symmetric in x, y , i.e.

$$(x, y, z) = (y, x, z), \text{ or equivalently, } (x *_g y) *_g z - x *_g (y *_g z) = (y *_g x) *_g z - y *_g (x *_g z).$$

It is obvious that any associative algebra is a pre-Lie algebra.

Lemma 2.2. ([7]) *A commutative pre-Lie algebra is associative.*

Lemma 2.3. ([7]) *Let $(g, *_g)$ be a pre-Lie algebra. The commutator $[x, y]_g = x *_g y - y *_g x$ defines a Lie algebra structure on g , which is called the **sub-adjacent Lie algebra** of $(g, *_g)$ and denoted by g^c . Furthermore, $L : g \rightarrow \mathfrak{gl}(g)$ defined by*

$$(1) \quad L_x y = x *_g y, \quad \forall x, y \in g$$

gives a representation of g^c on g .

A **Lie-admissible algebra** is a nonassociative algebra $(g, *_g)$ whose commutator algebra is a Lie algebra. More precisely, it is equivalent to the following condition:

$$(2) \quad (x, y, z) - (y, x, z) + (y, z, x) - (z, y, x) + (z, x, y) - (x, z, y) = 0, \quad \forall x, y, z \in g.$$

Obviously, a pre-Lie algebra is a Lie-admissible algebra.

Definition 2.4. *Let $(g, *_g)$ be a pre-Lie algebra and V a vector space. A **representation** of g on V consists of a pair (ρ, μ) , where $\rho : g \rightarrow \mathfrak{gl}(V)$ is a representation of the Lie algebra g^c on V and $\mu : g \rightarrow \mathfrak{gl}(V)$ is a linear map satisfying*

$$(3) \quad \rho(x)\mu(y)u - \mu(y)\rho(x)u = \mu(x *_g y)u - \mu(y)\mu(x)u, \quad \forall x, y \in g, u \in V.$$

Usually, we denote a representation by $(V; \rho, \mu)$. Let $R : g \rightarrow \mathfrak{gl}(g)$ be a linear map with $x \rightarrow R_x$, where the linear map $R_x : g \rightarrow g$ is defined by $R_x(y) = y *_g x$, for all $x, y \in g$. Then $(g; \rho = L, \mu = R)$ is a representation, which we call the **regular representation**. Define two linear maps $L^*, R^* : g \rightarrow \mathfrak{gl}(g^*)$ with $x \rightarrow L_x^*$ and $x \rightarrow R_x^*$ respectively (for all $x \in g$) by

$$(4) \quad \langle L_x^*(\xi), y \rangle = -\langle \xi, x *_g y \rangle, \quad \langle R_x^*(\xi), y \rangle = -\langle \xi, y *_g x \rangle, \quad \forall x, y \in g, \xi \in g^*.$$

Then $(g^*; \rho = \text{ad}^* = L^* - R^*, \mu = -R^*)$ is a representation of (g, \cdot_g) .

The cohomology complex for a pre-Lie algebra $(g, *_g)$ with a representation $(V; \rho, \mu)$ is given as follows ([15]). The set of n -cochains is given by $C^n(g, V) := \text{Hom}(\wedge^{n-1} g \otimes g, V)$, $n \geq 1$. For all $\phi \in C^n(g, V)$, the coboundary operator $d : C^n(g, V) \rightarrow C^{n+1}(g, V)$ is given by

$$(5) \quad \begin{aligned} d\phi(x_1, \dots, x_{n+1}) &= \sum_{i=1}^n (-1)^{i+1} \rho(x_i) \phi(x_1, \dots, \hat{x}_i, \dots, x_{n+1}) \\ &\quad + \sum_{i=1}^n (-1)^{i+1} \mu(x_{n+1}) \phi(x_1, \dots, \hat{x}_i, \dots, x_n, x_i) \\ &\quad - \sum_{i=1}^n (-1)^{i+1} \phi(x_1, \dots, \hat{x}_i, \dots, x_n, x_i *_g x_{n+1}) \\ &\quad + \sum_{1 \leq i < j \leq n} (-1)^{i+j} \phi([x_i, x_j]_g, x_1, \dots, \hat{x}_i, \dots, \hat{x}_j, \dots, x_{n+1}), \end{aligned}$$

for all $x_i \in g$, $i = 1, \dots, n+1$. In particular, we use the symbol d^{reg} to refer the coboundary operator associated to the regular representation. We denote the n -th cohomology group for the coboundary operator d^{reg} by $H_{\text{reg}}^n(g, g)$ and $H_{\text{reg}}(g, g) = \bigoplus_{n \geq 1} H_{\text{reg}}^n(g, g)$.

3. F -MANIFOLD ALGEBRAS AND PRE-LIE DEFORMATIONS OF COMMUTATIVE ASSOCIATIVE ALGEBRAS

3.1. F -manifold algebras and F -manifold-admissible algebras. In this subsection, we introduce the notion of F -manifold-admissible algebras, which give rise to F -manifold algebras. Various examples are given.

Definition 3.1. ([13]) An F -manifold algebra is a triple $(A, \cdot_A, [-, -]_A)$, where (A, \cdot_A) is a commutative associative algebra and $(A, [-, -]_A)$ is a Lie algebra, such that for all $x, y, z, w \in A$, the Hertling-Manin relation holds:

$$(6) \quad P_{x \cdot_A y}(z, w) = x \cdot_A P_y(z, w) + y \cdot_A P_x(z, w),$$

where $P_x(y, z)$ is defined by

$$(7) \quad P_x(y, z) = [x, y \cdot_A z]_A - [x, y]_A \cdot_A z - y \cdot_A [x, z]_A.$$

Remark 3.2. For any elements $z, w \in A$, the Hertling-Manin relation means that the linear map $P_{-}(z, w) : A \rightarrow A$ defined by $P_{-}(z, w)(x) := P_x(z, w)$ is a Hochschild 2-cocycle of the commutative associative algebra (A, \cdot_A) .

Remark 3.3. An F -manifold is a pair (M, \cdot) , where M is a manifold, \cdot is a $C^\infty(M)$ -bilinear, commutative, associative multiplication on the tangent bundle TM , such that the Hertling-Manin relation holds, where the Lie bracket in the Hertling-Manin relation is the usual Lie bracket of vector fields. Thus the underlying algebraic structure of an F -manifold (ignore the geometry of the arguments) is an F -manifold algebra.

Definition 3.4. Let $(A, \cdot_A, [-, -]_A)$ and $(B, \cdot_B, [-, -]_B)$ be two F -manifold algebras. A **homomorphism** between A and B is a linear map $\varphi : A \rightarrow B$ such that

$$(8) \quad \varphi(x \cdot_A y) = \varphi(x) \cdot_B \varphi(y),$$

$$(9) \quad \varphi[x, y]_A = [\varphi(x), \varphi(y)]_B, \quad \forall x, y \in A.$$

Example 3.5. Let $(A, \cdot_A, [-, -]_A)$ and $(B, \cdot_B, [-, -]_B)$ be two F -manifold algebras. Then $(A \oplus B, \cdot_{A \oplus B}, [-, -]_{A \oplus B})$ is an F -manifold algebra, where the product $\cdot_{A \oplus B}$ and bracket $[-, -]_{A \oplus B}$ are given by

$$\begin{aligned} (x_1 + x_2) \cdot_{A \oplus B} (y_1 + y_2) &= x_1 \cdot_A y_1 + x_2 \cdot_B y_2, \\ [(x_1 + x_2), (y_1 + y_2)]_{A \oplus B} &= [x_1, y_1]_A + [x_2, y_2]_B \end{aligned}$$

for all $x_1, y_1 \in A, x_2, y_2 \in B$.

Example 3.6. Let $(A, \cdot_A, [-, -]_A)$ and $(B, \cdot_B, [-, -]_B)$ be two F -manifold algebras. Then $(A \otimes B, \cdot_{A \otimes B}, [-, -]_{A \otimes B})$ is an F -manifold algebra, where the product $\cdot_{A \otimes B}$ and bracket $[-, -]_{A \otimes B}$ are given by

$$\begin{aligned} (x_1 \otimes x_2) \cdot_{A \otimes B} (y_1 \otimes y_2) &= (x_1 \cdot_A y_1) \otimes (x_2 \cdot_B y_2), \\ [x_1 \otimes x_2, y_1 \otimes y_2]_{A \otimes B} &= [x_1, y_1]_A \otimes (x_2 \cdot_B y_2) + (x_1 \cdot_A y_1) \otimes [x_2, y_2]_B \end{aligned}$$

for all $x_1, y_1 \in A, x_2, y_2 \in B$.

Recall that a **Poisson algebra** is a triple $(P, \cdot_P, \{-, -\}_P)$, where (P, \cdot_P) is a commutative associative algebra and $(P, \{-, -\}_P)$ is a Lie algebra, such that the Leibniz rule holds:

$$\{x, y \cdot_P z\}_P = \{x, y\}_P \cdot_P z + y \cdot_P \{x, z\}_P, \quad \forall x, y, z \in P.$$

Example 3.7. Any Poisson algebra is an F -manifold algebra.

The notion of PreLie-Com algebras was given in [18, 32].

Definition 3.8. A **pre-Lie commutative algebra (or PreLie-Com algebra)** is a triple $(A, \cdot_A, *_A)$, where (A, \cdot_A) is a commutative associative algebra and $(A, *_A)$ is a pre-Lie algebra satisfying

$$(10) \quad x *_A (y \cdot_A z) - (x *_A y) \cdot_A z - y \cdot_A (x *_A z) = 0, \quad \forall x, y, z \in A.$$

Corollary 3.9. ([13]) Let $(A, \cdot_A, *_A)$ be a PreLie-Com algebra. Then $(A, \cdot_A, [-, -]_A)$ is an F -manifold algebra, where the bracket $[-, -]_A$ is given by

$$(11) \quad [x, y]_A = x *_A y - y *_A x, \quad \forall x, y \in A.$$

Let $A = \mathbb{K}[x_1, x_2, \dots, x_n]$ be the algebra of polynomials in n variables. Let

$$\mathcal{D}_n = \{\partial_{x_1}, \partial_{x_2}, \dots, \partial_{x_n}\}$$

be the system of derivations over A . For any polynomial $f \in A$, the endomorphisms

$$f\partial_{x_i} : A \longrightarrow A, \quad (f\partial_{x_i})(g) = f\partial_{x_i}(g), \quad \forall g \in A$$

are derivations of A . Denote by $A\mathcal{D}_n = \{\sum_{i=1}^n f_i \partial_{x_i} \mid f_i \in A, \partial_{x_i} \in \mathcal{D}_n\}$ the space of derivations.

Example 3.10. Let A be the algebra of polynomials in n variables. Define $\cdot : A\mathcal{D}_n \times A\mathcal{D}_n \longrightarrow A\mathcal{D}_n$ and $*$: $A\mathcal{D}_n \times A\mathcal{D}_n \longrightarrow A\mathcal{D}_n$ by

$$\begin{aligned} (f\partial_{x_i}) \cdot (g\partial_{x_j}) &= (fg)\delta_{ij}\partial_{x_i}, \\ (f\partial_{x_i}) * (g\partial_{x_j}) &= f\partial_{x_i}(g)\partial_{x_j}, \quad \forall f, g \in A. \end{aligned}$$

Then $(A\mathcal{D}_n, \cdot, *)$ is a PreLie-Com algebra. Furthermore, $(A\mathcal{D}_n, \cdot, [-, -])$ is an F -manifold algebra, where the bracket is given by

$$[f\partial_{x_i}, g\partial_{x_j}] = f\partial_{x_i}(g)\partial_{x_j} - g\partial_{x_j}(f)\partial_{x_i}, \quad \forall f, g \in A.$$

Example 3.11. Let $A = \mathbb{K}[x_1, x_2]$ be the algebra of polynomials in two variables. Besides the PreLie-Com algebra structure given in Example 3.10 on $A\mathcal{D}_2$, there is another PreLie-Com algebra $(A\mathcal{D}_2, \cdot, *)$, where the operations \cdot and $*$ are determined by

$$\begin{aligned} \partial_{x_1} \cdot \partial_{x_1} &= \partial_{x_1}, \quad \partial_{x_1} \cdot \partial_{x_2} = \partial_{x_2} \cdot \partial_{x_1} = \partial_{x_2}, \quad \partial_{x_2} \cdot \partial_{x_2} = 0, \\ (f\partial_{x_i}) * (g\partial_{x_j}) &= f\partial_{x_i}(g)\partial_{x_j}, \quad \forall f, g \in A \ (i, j = 1, 2). \end{aligned}$$

Furthermore, $(A\mathcal{D}_2, \cdot, [-, -])$ is an F -manifold algebra, where the bracket is given by

$$[f\partial_{x_i}, g\partial_{x_j}] = f\partial_{x_i}(g)\partial_{x_j} - g\partial_{x_j}(f)\partial_{x_i}, \quad \forall f, g \in A.$$

In the sequel, we introduce the notion of F -manifold-admissible algebras, which include PreLie-Com algebras as special cases, and can be used to construct F -manifold algebras.

Definition 3.12. An **F -manifold-admissible algebra** is a vector space A equipped with two bilinear maps $\cdot_A : A \times A \rightarrow A$ and $*_A : A \times A \rightarrow A$ such that (A, \cdot_A) is a commutative associative algebra and $(A, *_A)$ is a Lie-admissible algebra satisfying for all $x, y, z \in A$,

$$(12) \quad x *_A (y \cdot_A z) - (x *_A y) \cdot_A z - y \cdot_A (x *_A z) = y *_A (x \cdot_A z) - (y *_A x) \cdot_A z - x \cdot_A (y *_A z).$$

Remark 3.13. Viewing the associative algebra A as a pre-Lie algebra, the condition (12) means that the F -manifold admissible operation $*_A : A \times A \rightarrow A$ is a 2-cocycle of the pre-Lie algebra (A, \cdot_A) with the coefficients in the regular representation, i.e. $d^{\text{reg}}(*_A) = 0$.

Theorem 3.14. *Let $(A, \cdot_A, *_A)$ be an F -manifold-admissible algebra. Then $(A, \cdot_A, [-, -]_A)$ is an F -manifold algebra, where $[-, -]_A$ is given by (11).*

Proof. We only need to verify the Hertling-Manin relation. By (12), we have

$$\begin{aligned}
 P_x(y, z) &= [x, y \cdot_A z]_A - [x, y]_A \cdot_A z - y \cdot_A [x, z]_A \\
 &= x *_A (y \cdot_A z) - (y \cdot_A z) *_A x - (x *_A y) \cdot_A z + (y *_A x) \cdot_A z \\
 &\quad - y \cdot_A (x *_A z) + y \cdot_A (z *_A x) \\
 &= x *_A (y \cdot_A z) - (x *_A y) \cdot_A z - y \cdot_A (x *_A z) - (y \cdot_A z) *_A x \\
 &\quad + (y *_A x) \cdot_A z + y \cdot_A (z *_A x) \\
 &= y *_A (x \cdot_A z) - (y *_A x) \cdot_A z - x \cdot_A (y *_A z) - (y \cdot_A z) *_A x \\
 &\quad + (y *_A x) \cdot_A z + y \cdot_A (z *_A x).
 \end{aligned}$$

By this formula and (12), we have

$$\begin{aligned}
 P_{x \cdot_A y}(z, w) &= z *_A ((x \cdot_A y) \cdot_A w) - (z *_A (x \cdot_A y)) \cdot_A w - (x \cdot_A y) \cdot_A (z *_A w) - (z \cdot_A w) *_A (x \cdot_A y) \\
 &\quad + (z *_A (x \cdot_A y)) \cdot_A w + z \cdot_A (w *_A (x \cdot_A y)) \\
 &= z *_A (x \cdot_A (y \cdot_A w)) - ((z *_A x) \cdot_A y - x \cdot_A (z *_A y) + x *_A (z \cdot_A y) - (x *_A z) \cdot_A y \\
 &\quad - z \cdot_A (x *_A y)) \cdot_A w - (x \cdot_A y) \cdot_A (z *_A w) - (((z \cdot_A w) *_A x) \cdot_A y + x \cdot_A ((z \cdot_A w) *_A y) \\
 &\quad + x *_A ((z \cdot_A w) \cdot_A y) - (x *_A (z \cdot_A w) \cdot_A y) - (z \cdot_A w) \cdot_A (x *_A y)) + ((z *_A x) \cdot_A y \\
 &\quad + x \cdot_A (z *_A y) + x *_A (z \cdot_A y) - (x *_A z) \cdot_A y - z \cdot_A (x *_A y))) \cdot_A w + ((w *_A x) \cdot_A y \\
 &\quad + x \cdot_A (w *_A y) + x *_A (w \cdot_A y) - (x *_A w) \cdot_A y - w \cdot_A (x *_A y)) \cdot_A z \\
 &= y \cdot_A P_x(z, w) + x \cdot_A (-w \cdot_A (z *_A y) - y \cdot_A (z *_A w) - (z \cdot_A w) *_A y + (z *_A y) \cdot_A w \\
 &\quad + (w *_A y) \cdot_A z) + (z *_A (x \cdot_A (y \cdot_A w)) - (z *_A x) \cdot_A (y \cdot_A w) \\
 &\quad - x *_A (z \cdot_A (y \cdot_A w)) + (x *_A z) \cdot_A (y \cdot_A w) + z \cdot_A (x *_A (y \cdot_A w))) \\
 &= y \cdot_A P_x(z, w) + x \cdot_A (-w \cdot_A (z *_A y) - y \cdot_A (z *_A w) - (z \cdot_A w) *_A y + (z *_A y) \cdot_A w \\
 &\quad + (w *_A y) \cdot_A z) + x \cdot_A (z *_A (y \cdot_A w)) \\
 &= y \cdot_A P_x(z, w) + x \cdot_A P_y(z, w).
 \end{aligned}$$

This shows that the Hertling-Manin relation holds. \square

Proposition 3.15. *Let (A, \cdot) be a commutative associative algebra with a derivation D . Then the new product*

$$x *_a y = x \cdot D(y) + a \cdot x \cdot y, \quad \forall x, y \in A$$

*makes $(A, \cdot, *_a)$ being an F -manifold-admissible algebra for any fixed $a \in \mathbb{K}$ or $a \in A$. In particular, for $a = 0$, $(A, \cdot, *_0)$ is a PreLie-Com algebra. Furthermore, $(A, \cdot, [-, -])$ is an F -manifold algebra, where the bracket is given by*

$$[x, y] = x *_a y - y *_a x = x \cdot D(y) - y \cdot D(x), \quad \forall x, y \in A.$$

Proof. It was shown that $(A, *_a)$ is a pre-Lie algebra for $a = 0$ by S.I. Gel'fand [19], for $a \in \mathbb{K}$ by Filippov [17] and for $a \in A$ by Xu [35]. Thus $(A, *_a)$ is a Lie-admissible algebra. Furthermore, by the fact that (A, \cdot) is a commutative associative algebra and D is a derivation on it, we have

$$x *_a (y \cdot z) - (x *_a y) \cdot z - y \cdot (x *_a z)$$

$$\begin{aligned}
&= x \cdot D(y \cdot z) + x \cdot (a \cdot y \cdot z) - (x \cdot D(y) + a \cdot x \cdot y) \cdot z - y \cdot (x \cdot D(z) + a \cdot x \cdot z) \\
&= -a \cdot x \cdot y \cdot z.
\end{aligned}$$

Similarly, we have

$$y *_a (x \cdot z) - (y *_a x) \cdot z - x \cdot (y *_a z) = -a \cdot x \cdot y \cdot z.$$

Thus

$$x *_a (y \cdot z) - (x *_a y) \cdot z - y \cdot (x *_a z) = y *_a (x \cdot z) - (y *_a x) \cdot z - x \cdot (y *_a z),$$

which implies that $(A, \cdot, *_a)$ is an F -manifold-admissible algebra for any fixed $a \in \mathbb{K}$ or $a \in A$. It is obvious that $(A, \cdot, *_0)$ is a PreLie-Com algebra. By Theorem 3.14, $(A, \cdot, [-, -])$ is an F -manifold algebra. \square

Example 3.16. Let A be the algebra of polynomials in n variables. Set $D = \sum_{i=1}^n f_i \partial_{x_i} \in A\mathcal{D}_n$ for $f_i \in A, \partial_{x_i} \in \mathcal{D}_n$. Fixed a polynomial $\kappa \in A$, define $\cdot : A \times A \rightarrow A$ and $*_{\kappa} : A \times A \rightarrow A$ by

$$\begin{aligned}
g \cdot h &= gh, \\
g *_k h &= gD(h) + \kappa gh = \sum_{i=1}^n g f_i \partial_{x_i}(h) + \kappa gh, \quad \forall g, h \in A.
\end{aligned}$$

Then by Proposition 3.15, $(A, \cdot, *_k)$ is an F -manifold-admissible algebra. Furthermore, $(A, \cdot, [-, -])$ is an F -manifold algebra, where the bracket is given by

$$[g, h] = \sum_{i=1}^n f_i (g \partial_{x_i}(h) - h \partial_{x_i}(g)), \quad \forall g, h \in A.$$

Example 3.17. Let A be a 2-dimensional vector space with basis $\{e_1, e_2\}$. Then A with the non-zero multiplication

$$e_1 \cdot e_1 = e_1, \quad e_1 \cdot e_2 = e_2 \cdot e_1 = e_2$$

is a commutative associative algebra. It is straightforward to check that all derivations on (A, \cdot) are determined by

$$D(e_2) = ae_2, \quad \forall a \in \mathbb{K}.$$

Thus by Proposition 3.15, $(A, \cdot, [-, -])$ with the bracket

$$[e_1, e_2] = ae_2$$

is an F -manifold algebra.

Example 3.18. Let A be a 3-dimensional vector space with basis $\{e_1, e_2, e_3\}$. Then A with the non-zero multiplication

$$e_2 \cdot e_3 = e_3 \cdot e_2 = e_1, \quad e_3 \cdot e_3 = e_2$$

is a commutative associative algebra. It is straightforward to check that all derivations on (A, \cdot) are determined by

$$D(e_1) = 3ae_1, \quad D(e_2) = 2be_1 + 2ae_2, \quad D(e_3) = ce_1 + be_2 + ae_3, \quad \forall a, b, c \in \mathbb{K}.$$

Thus by Proposition 3.15, $(A, \cdot, [-, -])$ with the bracket

$$[e_2, e_3] = -ae_1$$

is an F -manifold algebra.

3.2. Pre-Lie deformation quantization of commutative pre-Lie algebras. In this subsection, we introduce the notion of pre-Lie formal deformations of commutative associative algebras (that is, commutative pre-Lie algebras) and show that F -manifold algebras are the corresponding semi-classical limits. This illustrates that F -manifold algebras are the same to pre-Lie algebras as Poisson algebras to associative algebras. Furthermore, we show that pre-Lie infinitesimal deformations and extensions of pre-Lie n -deformations to pre-Lie $(n + 1)$ -deformations of a commutative associative algebra A are classified by the second and the third cohomology group of the pre-Lie algebra A (view the commutative associative algebra A as a pre-Lie algebra).

Let (A, \cdot_A) be a commutative associative algebra. Recall that an **associative formal deformation** of A is a sequence of bilinear maps $\mu_n : A \times A \rightarrow A$ for $n \geq 0$ with μ_0 being the commutative associative algebra product \cdot_A on A , such that the $\mathbb{K}[[\hbar]]$ -bilinear product \cdot_{\hbar} on $A[[\hbar]]$ determined by

$$x \cdot_{\hbar} y = \sum_{n=0}^{\infty} \hbar^n \mu_n(x, y), \quad \forall x, y \in A$$

is associative, where $A[[\hbar]]$ is the set of formal power series of \hbar with coefficients in A . Define

$$\{x, y\} = \mu_1(x, y) - \mu_1(y, x), \quad \forall x, y \in A.$$

It is well known that $(A, \cdot_A, \{-, -\})$ is a Poisson algebra, called the **semi-classical limit** of $(A[[\hbar]], \cdot_{\hbar})$.

Since an associative algebra can be regarded as a pre-Lie algebra, one may look for formal deformations of a commutative associative algebra into pre-Lie algebras, that is, in the aforementioned associative formal deformation, replace the associative product \cdot_{\hbar} by the pre-Lie product, and wonder what additional structure will appear on A . On the other hand, such an approach can be also seen as formal deformations of a commutative pre-Lie algebra into (non-commutative) pre-Lie algebras, which is completely parallel to the associative formal deformations of a commutative associative algebra into (non-commutative) associative algebras. Surprisingly, we find that this is the structure of the F -manifold algebra. Now we give the definition of a pre-Lie formal deformation of a commutative associative algebra.

Definition 3.19. Let (A, \cdot_A) be a commutative associative algebra. A **pre-Lie formal deformation** of A is a sequence of bilinear maps $\mu_k : A \times A \rightarrow A$ for $k \geq 0$ with μ_0 being the commutative associative algebra product \cdot_A on A , such that the $\mathbb{K}[[\hbar]]$ -bilinear product \cdot_{\hbar} on $A[[\hbar]]$ determined by

$$x \cdot_{\hbar} y = \sum_{n=0}^{\infty} \hbar^n \mu_n(x, y), \quad \forall x, y \in A$$

is a pre-Lie algebra product.

Note that the rule of pre-Lie algebra product \cdot_{\hbar} on $A[[\hbar]]$ is equivalent to

$$(13) \quad \sum_{i+j=k} (\mu_i(\mu_j(x, y), z) - \mu_i(x, \mu_j(y, z))) = \sum_{i+j=k} (\mu_i(\mu_j(y, x), z) - \mu_i(y, \mu_j(x, z))), \quad \forall k \geq 0.$$

Theorem 3.20. Let (A, \cdot_A) be a commutative associative algebra and $(A[[\hbar]], \cdot_{\hbar})$ a pre-Lie formal deformation of A . Define

$$[x, y]_A = \mu_1(x, y) - \mu_1(y, x), \quad \forall x, y \in A.$$

Then $(A, \cdot_A, [-, -]_A)$ is an F -manifold algebra. The F -manifold algebra $(A, \cdot_A, [-, -]_A)$ is called the **semi-classical limit** of $(A[[\hbar]], \cdot_{\hbar})$. The pre-Lie algebra $(A[[\hbar]], \cdot_{\hbar})$ is called a **pre-Lie deformation quantization** of (A, \cdot_A) .

Proof. Define the bracket $[-, -]_{\hbar}$ on $A[[\hbar]]$ by

$$[x, y]_{\hbar} = x \cdot_{\hbar} y - y \cdot_{\hbar} x = \hbar[x, y]_A + \hbar^2(\mu_2(x, y) - \mu_2(y, x)) + \cdots, \quad \forall x, y \in A.$$

By the fact that $(A[[\hbar]], \cdot_{\hbar})$ is a pre-Lie algebra, $(A[[\hbar]], [-, -]_{\hbar})$ is a Lie algebra. The \hbar^2 -terms of the Jacobi identity for $[-, -]_{\hbar}$ gives the Jacobi identity for $[-, -]_A$. Thus $(A, [-, -]_A)$ is a Lie algebra.

For $k = 1$ in (13), by the commutativity of μ_0 , we have

$$\mu_0(\mu_1(x, y), z) - \mu_0(x, \mu_1(y, z)) - \mu_1(x, \mu_0(y, z)) = \mu_0(\mu_1(y, x), z) - \mu_0(y, \mu_1(x, z)) - \mu_1(y, \mu_0(x, z)).$$

This is just the equality (12) with $x \cdot_A y = \mu_0(x, y)$ and $x *_A y = \mu_1(x, y)$ for $x, y \in A$. Thus $(A, \cdot_A, *_A)$ is an F -manifold-admissible algebra. By Theorem 3.14, $(A, \cdot_A, [-, -]_A)$ is an F -manifold algebra. \square

In the sequel, we study pre-Lie n -deformations and pre-Lie infinitesimal deformations of commutative associative algebras.

Definition 3.21. Let (A, \cdot_A) be a commutative associative algebra. A **pre-Lie n -deformation** of A is a sequence of bilinear maps $\mu_i : A \times A \rightarrow A$ for $0 \leq i \leq n$ with μ_0 being the commutative associative algebra product \cdot_A on A , such that the $\mathbb{K}[[\hbar]]/(\hbar^{n+1})$ -bilinear product \cdot_{\hbar} on $A[[\hbar]]/(\hbar^{n+1})$ determined by

$$x \cdot_{\hbar} y = \sum_{k=0}^n \hbar^k \mu_k(x, y), \quad \forall x, y \in A$$

is a pre-Lie algebra product.

We call a pre-Lie 1-deformation of a commutative associative algebra (A, \cdot_A) a **pre-Lie infinitesimal deformation** and denote it by (A, μ_1) .

By direct calculations, (A, μ_1) is a pre-Lie infinitesimal deformation of a commutative associative algebra (A, \cdot_A) if and only if for all $x, y, z \in A$

$$(14) \quad \mu_1(x, y) \cdot_A z - x \cdot_A \mu_1(y, z) - \mu_1(x, y \cdot_A z) = \mu_1(y, x) \cdot_A z - y \cdot_A \mu_1(x, z) - \mu_1(y, x \cdot_A z).$$

Equation (14) means that μ_1 is a 2-cocycle for the pre-Lie algebra (A, \cdot_A) , i.e. $d^{\text{reg}}\mu_1 = 0$.

Two pre-Lie infinitesimal deformations $A_{\hbar} = (A, \mu_1)$ and $A'_{\hbar} = (A, \mu'_1)$ of a commutative associative algebra (A, \cdot_A) are said to be **equivalent** if there exists a family of pre-Lie algebra homomorphisms $\text{Id} + \hbar\varphi : A_{\hbar} \rightarrow A'_{\hbar}$ modulo \hbar^2 . A pre-Lie infinitesimal deformation is said to be **trivial** if there exists a family of pre-Lie algebra homomorphisms $\text{Id} + \hbar\varphi : A_{\hbar} \rightarrow (A, \cdot_A)$ modulo \hbar^2 .

By direct calculations, A_{\hbar} and A'_{\hbar} are equivalent pre-Lie infinitesimal deformations if and only if

$$(15) \quad \mu_1(x, y) - \mu'_1(x, y) = x \cdot_A \varphi(y) + \varphi(x) \cdot_A y - \varphi(x \cdot_A y).$$

Equation (15) means that $\mu_1 - \mu'_1 = d^{\text{reg}}\varphi$. Thus we have

Theorem 3.22. There is a one-to-one correspondence between the space of equivalence classes of pre-Lie infinitesimal deformations of A and the second cohomology group $H_{\text{reg}}^2(A, A)$.

It is routine to check that

Proposition 3.23. *Let (A, \cdot_A) be a commutative associative algebra such that $H_{\text{reg}}^2(A, A) = 0$. Then all pre-Lie infinitesimal deformations of A are trivial.*

Definition 3.24. *Let $\{\mu_1, \dots, \mu_n\}$ be a pre-Lie n -deformation of a commutative associative algebra (A, \cdot_A) . A pre-Lie $(n+1)$ -deformation of a commutative associative algebra (A, \cdot_A) given by $\{\mu_1, \dots, \mu_n, \mu_{n+1}\}$ is called an **extension** of the pre-Lie n -deformation given by $\{\mu_1, \dots, \mu_n\}$.*

Theorem 3.25. *For any pre-Lie n -deformation of a commutative associative algebra (A, \cdot_A) , the $\Theta_n \in \text{Hom}(\otimes^3 \mathfrak{g}, \mathfrak{g})$ defined by*

$$(16) \quad \Theta_n(x, y, z) = \sum_{i+j=n+1, i, j \geq 1} (\mu_i(\mu_j(x, y), z) - \mu_i(x, \mu_j(y, z)) - \mu_i(\mu_j(y, x), z) + \mu_i(y, \mu_j(x, z))).$$

is a cocycle, i.e. $d^{\text{reg}}\Theta_n = 0$.

Moreover, the pre-Lie n -deformation $\{\mu_1, \dots, \mu_n\}$ extends into some pre-Lie $(n+1)$ -deformation if and only if $[\Theta_n] = 0$ in $H_{\text{reg}}^3(A, A)$.

Proof. It is obvious that

$$\Theta_n(x, y, z) = -\Theta_n(y, x, z), \quad \forall x, y, z \in A.$$

Thus Θ_n is an element of $C^3(A, A)$. It is straightforward to check that the cochain $\Theta_n \in C^3(A, A)$ is closed.

Assume that the pre-Lie $(n+1)$ -deformation of a commutative associative algebra (A, \cdot_A) given by $\{\mu_1, \dots, \mu_n, \mu_{n+1}\}$ is an extension of the pre-Lie n -deformation given by $\{\mu_1, \dots, \mu_n\}$, then we have

$$\begin{aligned} x \cdot_A \mu_{n+1}(y, z) - y \cdot_A \mu_{n+1}(x, z) + \mu_{n+1}(y, x) \cdot_A z - \mu_{n+1}(x, y) \cdot_A z + \mu_{n+1}(y, x) \cdot_A z \\ - \mu_{n+1}(x, y) \cdot_A z = \sum_{i+j=n+1, i, j \geq 1} (\mu_i(\mu_j(x, y), z) - \mu_i(x, \mu_j(y, z)) - \mu_i(\mu_j(y, x), z) + \mu_i(y, \mu_j(x, z))). \end{aligned}$$

It is obvious that the right-hand side of the above equality is just $\Theta_n(x, y, z)$. We can rewrite the above equality as

$$d^{\text{reg}}\mu_{n+1}(x, y, z) = \Theta_n(x, y, z).$$

We conclude that, if a pre-Lie n -deformation of a commutative associative algebra (A, \cdot_A) extends to a pre-Lie $(n+1)$ -deformation, then Θ_n is coboundary.

Conversely, if Θ_n is coboundary, then there exists an element $\psi \in C^2(A, A)$ such that

$$d^{\text{reg}}\psi(x, y, z) = \Theta_n(x, y, z).$$

It is not hard to check that $\{\mu_1, \dots, \mu_n, \mu_{n+1}\}$ with $\mu_{n+1} = \psi$ generates a pre-Lie $(n+1)$ -deformation of (A, \cdot_A) and thus this pre-Lie $(n+1)$ -deformation is an extension of the pre-Lie n -deformation given by $\{\mu_1, \dots, \mu_n\}$. \square

4. PRE- F -MANIFOLD ALGEBRAS, ROTA-BAXTER OPERATORS AND \mathcal{O} -OPERATORS ON F -MANIFOLD ALGEBRAS

In this section, first we study representations of an F -manifold algebra. Then we introduce the notions of pre- F -manifold algebras and Rota-Baxter operators (more generally \mathcal{O} -operators) on an F -manifold algebra. We show that on one hand, an \mathcal{O} -operator on an F -manifold algebra

gives a pre- F -manifold algebra, and on the other hand, a pre- F -manifold algebra naturally gives an O -operator on the sub-adjacent F -manifold algebra. More examples on pre- F -manifold algebras and F -manifold algebras are given.

4.1. Representations of F -manifold algebras. In this subsection, we introduce the notion of representations of F -manifold algebras.

Let (A, \cdot_A) be a commutative associative algebra. Recall that a **representation** of A on a vector space V is a linear map $\mu : A \rightarrow \mathfrak{gl}(V)$ such that $\mu(x \cdot_A y) = \mu(x) \circ \mu(y)$ for any $x, y \in A$. We will denote a representation of A by $(V; \mu)$. Let $(V; \mu)$ be a representation of a commutative associative algebra (A, \cdot_A) . Define $\mu^* : A \rightarrow \mathfrak{gl}(V^*)$ by

$$\langle \mu^*(x)\alpha, v \rangle = -\langle \alpha, \mu(x)v \rangle, \quad \forall x \in A, \alpha \in V^*, v \in V.$$

Then $(V^*; -\mu^*)$ is a representation of (A, \cdot_A) .

Example 4.1. Let (A, \cdot_A) be a commutative associative algebra. Let \mathcal{L}_x denote the multiplication operator, that is, $\mathcal{L}_x y = x \cdot_A y$, for all $x, y \in A$. Then $(A; \mathcal{L})$ is a representation of (A, \cdot_A) , called the **regular representation**. Furthermore, $(A^*; -\mathcal{L}^*)$ is also a representation of (A, \cdot_A) .

Similarly, let $(\mathfrak{g}, [-, -]_{\mathfrak{g}})$ be a Lie algebra and V a vector space. Let $\rho : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be a linear map. The pair $(V; \rho)$ is called a **representation** of \mathfrak{g} if for all $x, y \in \mathfrak{g}$, we have $\rho([x, y]_{\mathfrak{g}}) = [\rho(x), \rho(y)]$. Let $(V; \rho)$ be a representation of a Lie algebra $(\mathfrak{g}, [-, -]_{\mathfrak{g}})$. Define $\rho^* : \mathfrak{g} \rightarrow \mathfrak{gl}(V^*)$ by

$$\langle \rho^*(x)(\alpha), v \rangle = -\langle \alpha, \rho(x)(v) \rangle, \quad \forall x \in \mathfrak{g}, \alpha \in V^*, v \in V.$$

Then $(V^*; \rho^*)$ is a representation of $(\mathfrak{g}, [-, -]_{\mathfrak{g}})$.

Example 4.2. Define $\text{ad} : \mathfrak{g} \rightarrow \mathfrak{gl}(\mathfrak{g})$ by $\text{ad}_x y = [x, y]_{\mathfrak{g}}$ for all $x, y \in \mathfrak{g}$. Then $(\mathfrak{g}; \text{ad})$ is a representation of $(\mathfrak{g}, [-, -]_{\mathfrak{g}})$, called the **adjoint representation**. Furthermore, $(\mathfrak{g}^*; \text{ad}^*)$ is also a representation of $(\mathfrak{g}, [-, -]_{\mathfrak{g}})$.

Definition 4.3. Let $(A, \cdot_A, [-, -]_A)$ be an F -manifold algebra. A **representation** of A is a triple $(V; \rho, \mu)$ such that $(V; \rho)$ is a representation of the Lie algebra $(A, [-, -]_A)$ and $(V; \mu)$ is a representation of the commutative associative algebra (A, \cdot_A) satisfying

$$(17) \quad R_{\rho, \mu}(x \cdot_A y, z) = \mu(x) \circ R_{\rho, \mu}(y, z) + \mu(y) \circ R_{\rho, \mu}(x, z),$$

$$(18) \quad \mu(P_x(y, z)) = S_{\rho, \mu}(y, z) \circ \mu(x) - \mu(x) \circ S_{\rho, \mu}(y, z),$$

where $R_{\rho, \mu}, S_{\rho, \mu} : A \otimes A \rightarrow \mathfrak{gl}(V)$ are defined by

$$(19) \quad R_{\rho, \mu}(x, y) = \rho(x) \circ \mu(y) - \mu(y) \circ \rho(x) - \mu([x, y]_A),$$

$$(20) \quad S_{\rho, \mu}(x, y) = \mu(x) \circ \rho(y) + \mu(y) \circ \rho(x) - \rho(x \cdot_A y)$$

for all $x, y, z \in A$.

It is straightforward to obtain the following conclusion.

Proposition 4.4. Let $(A, \cdot_A, [-, -]_A)$ be an F -manifold algebra and $(V; \rho, \mu)$ a representation. Then $(A \oplus V, \cdot_{\mu}, [-, -]_{\rho})$ is an F -manifold algebra, where $(A \oplus V, \cdot_{\mu})$ is the semi-direct product commutative associative algebra $A \ltimes_{\mu} V$, i.e.

$$(x_1 + v_1) \cdot_{\mu} (x_2 + v_2) = x_1 \cdot_A x_2 + \mu(x_1)v_2 + \mu(x_2)v_1, \quad \forall x_1, x_2 \in A, v_1, v_2 \in V$$

and $(A \oplus V, [-, -]_\rho)$ is the semi-direct product Lie algebra $A \ltimes_\rho V$, i.e.

$$[x_1 + v_1, x_2 + v_2]_\rho = [x_1, x_2]_g + \rho(x_1)(v_2) - \rho(x_2)(v_1), \quad \forall x_1, x_2 \in A, v_1, v_2 \in V.$$

Example 4.5. Let $(V; \rho, \mu)$ be a representation of a Poisson algebra $(P, \cdot_P, \{-, -\}_P)$, i.e. $(V; \rho)$ is a representation of the Lie algebra $(P, \{-, -\}_P)$ and $(V; \mu)$ is a representation of the commutative associative algebra (P, \cdot_P) satisfying

$$(21) \quad \rho(x) \circ \mu(y) - \mu(y) \circ \rho(x) - \mu([x, y]_A) = 0,$$

$$(22) \quad \mu(x) \circ \rho(y) + \mu(y) \circ \rho(x) - \rho(x \cdot_A y) = 0, \quad \forall x, y, z \in P.$$

Then $(V; \rho, \mu)$ is also a representation of the F -manifold algebra given by this Poisson algebra P .

Let $(V; \rho, \mu)$ be a representation of a Poisson algebra $(P, \cdot_P, \{-, -\}_P)$. Then the triple $(V^*; \rho^*, -\mu^*)$ is also a representation of P . But F -manifold algebras do not have this property. In fact, we have

Proposition 4.6. Let $(A, \cdot_A, [-, -]_A)$ be an F -manifold algebra. If the triple $(V; \rho, \mu)$, where $(V; \rho)$ is a representation of the Lie algebra $(A, [-, -]_A)$ and $(V; \mu)$ is a representation of the commutative associative algebra (A, \cdot_A) , satisfies

$$(23) \quad R_{\rho, \mu}(x \cdot_A y, z) = R_{\rho, \mu}(y, z) \circ \mu(x) + R_{\rho, \mu}(x, z) \circ \mu(y),$$

$$(24) \quad \mu(P_x(y, z)) = T_{\rho, \mu}(y, z) \circ \mu(x) - \mu(x) \circ T_{\rho, \mu}(y, z),$$

where $R_{\rho, \mu}$ is given by (19) and $T_{\rho, \mu} : A \otimes A \rightarrow \mathfrak{gl}(V)$ is defined by

$$(25) \quad T_{\rho, \mu}(x, y) = \rho(y) \circ \mu(x) + \rho(x) \circ \mu(y) - \rho(x \cdot_A y), \quad \forall x, y \in A,$$

then $(V^*; \rho^*, -\mu^*)$ is a representation of A .

Proof. By direct calculations, for all $x, y \in A, v \in V, \alpha \in V^*$, we have

$$\langle R_{\rho^*, -\mu^*}(x, y)(\alpha), v \rangle = \langle \alpha, R_{\rho, \mu}(v) \rangle, \quad \langle S_{\rho^*, -\mu^*}(x, y)(\alpha), v \rangle = \langle \alpha, T_{\rho, \mu}(v) \rangle.$$

Furthermore, we have

$$\begin{aligned} & \langle R_{\rho^*, -\mu^*}(x \cdot_A y, z)(\alpha) + \mu^*(x)R_{\rho^*, -\mu^*}(y, z)(\alpha) + \mu^*(y)R_{\rho^*, -\mu^*}(x, z)(\alpha), v \rangle \\ &= \langle \alpha, R_{\rho, \mu}(x \cdot_A y, z)(v) - R_{\rho, \mu}(y, z)\mu(x)(v) - R_{\rho, \mu}(x, z)\mu(y)(v) \rangle \end{aligned}$$

and

$$\begin{aligned} & \langle -\mu^*(P_x(y, z))(\alpha) + S_{\rho^*, -\mu^*}(y, z)\mu^*(x)(\alpha) + \mu^*(x)S_{\rho^*, -\mu^*}(y, z)(\alpha), v \rangle \\ &= \langle \alpha, \mu(P_x(y, z))(v) - T_{\rho, \mu}(y, z)\mu(x)(v) + \mu(x)T_{\rho, \mu}(y, z)(v) \rangle. \end{aligned}$$

By the hypothesis and the definition of representation, the conclusion follows immediately. \square

Example 4.7. Let $(A, \cdot_A, [-, -]_A)$ be an F -manifold algebra. Then $(A; \text{ad}, \mathcal{L})$ is a representation of A , which is also called the **regular representation**. Furthermore, if the F -manifold algebra also satisfies the following relations:

$$(26) \quad P_{x \cdot_A y}(z, w) = P_y(z, x \cdot_A w) + P_x(z, y \cdot_A w),$$

$$(27) \quad P_x(y, z) \cdot_A w = x \cdot_A Q(y, z, w) - Q(y, z, x \cdot_A w), \quad \forall x, y, z, w \in A,$$

where $Q : \otimes^3 \rightarrow A$ is defined by

$$Q(x, y, z) = [x \cdot_A y, z]_A + [y \cdot_A z, x]_A + [z \cdot_A x, y]_A.$$

Then $(A^*; \text{ad}^*, -\mathcal{L}^*)$ is a representation of A .

Definition 4.8. A coherence F -manifold algebra is an F -manifold algebra such that (26) and (27) hold.

Proposition 4.9. Let $(A, \cdot_A, [-, -]_A)$ be an F -manifold algebra. Suppose that there is a nondegenerate symmetric bilinear form \mathfrak{B} such that \mathfrak{B} is invariant in the following sense

$$(28) \quad \mathfrak{B}(x \cdot_A y, z) = \mathfrak{B}(x, y \cdot_A z), \quad \mathfrak{B}([x, y]_A, z) = \mathfrak{B}(x, [y, z]_A), \quad \forall x, y, z \in A.$$

Then $(A, \cdot_A, [-, -]_A)$ is a coherence F -manifold algebra.

Proof. By the invariance of \mathfrak{B} , we have

$$\mathfrak{B}(P_x(y, z), w) = \mathfrak{B}(z, P_x(y, w)) = \mathfrak{B}(x, Q(y, z, w)), \quad \forall x, y, z, w \in P.$$

By the above relations, for $x, y, z, w_1, w_2 \in P$, we have

$$\begin{aligned} & \mathfrak{B}(P_{x \cdot_A y}(z, w_1) - P_y(z, x \cdot_A w_1) - P_x(z, y \cdot_A w_1), w_2) \\ &= \mathfrak{B}(w_1, P_{x \cdot_A y}(z, w_2) - x \cdot_A P_y(z, w_2) - y \cdot_A P_x(z, w_2)) \end{aligned}$$

and

$$\begin{aligned} & \mathfrak{B}(P_x(y, z) \cdot_A w_1 - x \cdot_A Q(y, z, w_1) + Q(y, z, x \cdot_A w), w_2) \\ &= \mathfrak{B}(w_1, -P_{x \cdot_A y}(z, w_2) + x \cdot_A P_y(z, w_2) + y \cdot_A P_x(z, w_2)). \end{aligned}$$

By the fact that A is an F -manifold algebra and \mathfrak{B} is nondegenerate, we deduce that (26) and (27) hold. Thus $(A, \cdot_A, [-, -]_A)$ is a coherence F -manifold algebra. \square

4.2. Pre- F -manifold algebras. Recall that a **Zinbiel algebra** is a pair (A, \diamond) , where A is a vector space, and $\diamond : A \otimes A \longrightarrow A$ is a bilinear multiplication satisfying that for all $x, y, z \in A$,

$$(29) \quad x \diamond (y \diamond z) = (y \diamond x) \diamond z + (x \diamond y) \diamond z.$$

Lemma 4.10. Let (A, \diamond) be a Zinbiel algebra. Then (A, \cdot) is a commutative associative algebra, where $x \cdot y = x \diamond y + y \diamond x$. Moreover, for $x \in A$, define $\mathfrak{L}_x : A \longrightarrow \text{gl}(A)$ by

$$(30) \quad \mathfrak{L}_x y = x \diamond y, \quad \forall y \in A.$$

Then $(A; \mathfrak{L})$ is a representation of the commutative associative algebra (A, \cdot) .

Now we are ready to give the main notion in this subsection.

Definition 4.11. A pre- F -manifold algebra is a triple $(A, \diamond, *)$, where (A, \diamond) is a Zinbiel algebra and $(A, *)$ is a pre-Lie algebra, such that the following compatibility conditions hold:

$$(31) \quad F_1(x \cdot y, z, w) = x \diamond F_1(y, z, w) + y \diamond F_1(x, z, w),$$

$$(32) \quad (F_1(x, y, z) + F_1(x, z, y) + F_2(y, z, x)) \diamond w = F_2(y, z, x \diamond w) - x \diamond F_2(y, z, w)$$

for all $x, y, z, w \in A$. Here $F_1, F_2 : \otimes^3 A \longrightarrow A$ are defined by

$$(33) \quad F_1(x, y, z) = x * (y \diamond z) - y \diamond (x * z) - [x, y] \diamond z,$$

$$(34) \quad F_2(x, y, z) = x \diamond (y * z) + y \diamond (x * z) - (x \cdot y) * z$$

and the operation \cdot and bracket $[-, -]$ are defined by

$$(35) \quad x \cdot y = x \diamond y + y \diamond x, \quad [x, y] = x * y - y * x.$$

Remark 4.12. If F_1 and F_2 vanish in the definition of a pre- F -manifold algebra $(A, \diamond, *)$, then we obtain Aguiar's notion of a pre-Poisson algebra. See [1] for more details.

Theorem 4.13. Let $(A, \diamond, *)$ be a pre- F -manifold algebra. Then

- (i) $(A, \cdot, [-, -])$ is an F -manifold algebra, where the operation \cdot and bracket $[-, -]$ are given by (35), which is called the **sub-adjacent F -manifold algebra** of $(A, \diamond, *)$ and denoted by A^c .
- (ii) $(A; L, \mathfrak{Q})$ is a representation of the sub-adjacent F -manifold algebras A^c , where L and \mathfrak{Q} are given by (1) and (30), respectively.

Proof. (i) By Lemma 2.3 and Lemma 4.10, we deduce that (A, \cdot) is a commutative associative algebra and $(A, [-, -])$ is a Lie algebra. By direct calculations, we obtain

$$(36) \quad P_x(y, z) = F_1(x, y, z) + F_1(x, z, y) + F_2(y, z, x), \quad \forall x, y, z \in A.$$

By (31), (32) and (36), we have

$$\begin{aligned} & P_{x \cdot y}(z, w) - x \cdot P_y(z, w) - y \cdot P_x(z, w) \\ = & F_1(x \cdot y, z, w) + F_1(x \cdot y, w, z) + F_2(z, w, x \cdot y) - x \cdot (F_1(y, z, w) + F_1(y, w, z) + F_2(z, w, y)) \\ & - y \cdot (F_1(x, z, w) + F_1(x, w, z) + F_2(z, w, x)) \\ = & (F_1(x \cdot y, z, w) - x \diamond F_1(y, z, w) - y \diamond F_1(x, z, w)) \\ & + (F_1(x \cdot y, w, z) - x \diamond F_1(y, w, z) - y \diamond F_1(x, w, z)) \\ & + (F_2(z, w, x \diamond y) - F_1(x, z, w) \diamond y + F_1(x, w, z) \diamond y + F_2(z, w, x) \diamond y + x \diamond F_2(z, w, y)) \\ & + (F_2(z, w, y \diamond x) - F_1(y, z, w) \diamond x + F_1(y, w, z) \diamond x + F_2(z, w, y) \diamond x + y \diamond F_2(z, w, x)) \\ = & 0. \end{aligned}$$

Thus $(A, \cdot, [-, -])$ is an F -manifold algebra.

(ii) By Lemma 2.3, $(A; L)$ is a representation of the sub-adjacent Lie algebra A^c . By Lemma 4.10, $(A; \mathfrak{Q})$ is a representation of the commutative associative algebra (A, \cdot) . Moreover, note that $F_1(x, y, z) = R_{L, \mathfrak{Q}}(x, y)(z)$ and $F_2(x, y, z) = S_{L, \mathfrak{Q}}(x, y)(z)$. Thus (31) implies that (17) holds and, by (36), (32) implies that (18) holds. Thus $(A; L, \mathfrak{Q})$ is a representation of the sub-adjacent F -manifold algebra A^c . \square

Remark 4.14. In fact, the operad PreFMan of pre- F -manifold algebras is the arity splitting of the operad FMan of F -manifold algebras (into two pieces) in the sense of [34] or the disuccessor of FMan in the sense of [4].

4.3. Rota-Baxter operators and \mathcal{O} -operators on F -manifold algebras. The notion of an \mathcal{O} -operator was first given for Lie algebras by Kupershmidt in [24] as a natural generalization of the classical Yang-Baxter equation.

A linear map $T : V \longrightarrow A$ is called an **\mathcal{O} -operator** on a commutative associative algebra (A, \cdot_A) with respect to a representation $(V; \mu)$ if T satisfies

$$(37) \quad T(u) \cdot_A T(v) = T(\mu(T(u))v + \mu(T(v))u), \quad \forall u, v \in V.$$

In particular, an \mathcal{O} -operator on a commutative associative algebra (A, \cdot_A) with respect to the regular representation is called a **Rota-Baxter operator of weight zero** or briefly a **Rota-Baxter operator** on A .

Lemma 4.15. ([3]) *Let (A, \cdot_A) be a commutative associative algebra and $(V; \mu)$ a representation. Let $T : V \rightarrow A$ be an O -operator on (A, \cdot_A) with respect to $(V; \mu)$. Then there exists a Zinbiel algebra structure on V given by*

$$u \diamond v = \mu(T(u))v, \quad \forall u, v \in V.$$

A linear map $T : V \rightarrow \mathfrak{g}$ is called an **O -operator** on a Lie algebra $(\mathfrak{g}, [-, -]_{\mathfrak{g}})$ with respect to a representation $(V; \rho)$ if T satisfies

$$(38) \quad [T(u), T(v)]_{\mathfrak{g}} = T(\rho(T(u))(v) - \rho(T(v))(u)), \quad \forall u, v \in V.$$

In particular, an O -operator on a Lie algebra $(\mathfrak{g}, [-, -]_{\mathfrak{g}})$ with respect to the adjoint representation is called a **Rota-Baxter operator of weight zero** or briefly a **Rota-Baxter operator** on \mathfrak{g} .

Lemma 4.16. ([2]) *Let $T : V \rightarrow \mathfrak{g}$ be an O -operator on a Lie algebra $(\mathfrak{g}, [-, -]_{\mathfrak{g}})$ with respect to a representation $(V; \rho)$. Define a multiplication $*$ on V by*

$$(39) \quad u * v = \rho(T(u))(v), \quad \forall u, v \in V.$$

*Then $(V, *)$ is a pre-Lie algebra.*

Let $(V; \rho, \mu)$ be a representation of an F -manifold algebra $(A, \cdot_A, [-, -]_A)$.

Definition 4.17. (i) *A linear operator $T : V \rightarrow A$ is called an O -operator on A if T is both an O -operator on the commutative associative algebra (A, \cdot_A) and an O -operator on the Lie algebra $(A, [-, -]_A)$;*
(ii) *A linear operator $\mathcal{B} : A \rightarrow A$ is called a **Rota-Baxter operator of weight zero** or briefly a **Rota-Baxter operator** on A , if \mathcal{B} is both a Rota-Baxter operator on the commutative associative algebra (A, \cdot_A) and a Rota-Baxter operator on the Lie algebra $(A, [-, -]_A)$.*

It is obvious that $\mathcal{B} : A \rightarrow A$ is a Rota-Baxter operator on A if and only if \mathcal{B} is an O -operator on A with respect to the representation $(A; \text{ad}, \mathcal{L})$.

Example 4.18. Let $(A, \diamond, *)$ be a pre- F -manifold algebra. Then the identity map Id is an O -operator on A^c with respect to the representation $(A; L, \mathfrak{L})$, where L and \mathfrak{L} are given by (1) and (30) respectively.

An O -operator on an F -manifold algebra gives a pre- F -manifold algebra.

Theorem 4.19. *Let $(A, \cdot_A, [-, -]_A)$ be an F -manifold algebra and $T : V \rightarrow A$ an O -operator on A with respect to the representation $(V; \rho, \mu)$. Define new operations \diamond and $*$ on V by*

$$u \diamond v = \mu(T(u))v, \quad u * v = \rho(T(u))v.$$

*Then $(V, \diamond, *)$ is a pre- F -manifold algebra and T is a homomorphism from V^c to $(A, \cdot_A, [-, -]_A)$.*

Proof. First by the fact that T is an O -operator on the commutative associative algebra (A, \cdot_A) as well as an O -operator on the Lie algebra $(A, [-, -]_A)$ with respect to the representations $(V; \mu)$ and $(V; \rho)$ respectively, we deduce that (V, \diamond) is a Zinbiel algebra and $(V, *)$ is a pre-Lie algebra.

Denote by $[u, v]_T := u * v - v * u$ and $u \cdot_T v := u \diamond v + v \diamond u$, we have $T([u, v]_T) = [T(u), T(v)]_A$ and $T(u \cdot_T v) = T(u) \cdot_A T(v)$. By these facts and (17), for $v_1, v_2, v_3, v_4 \in V$, one has

$$F_1(v_1 \cdot_T v_2, v_3, v_4) - v_1 \diamond F_1(v_2, v_3, v_4) - v_2 \diamond F_1(v_1, v_3, v_4)$$

$$\begin{aligned}
&= R_{\rho,\mu}(T(v_1) \cdot_A T(v_2), T(v_3))(v_4) - \mu(T(v_1))R_{\rho,\mu}(T(v_2), T(v_3))(v_4) \\
&\quad - \mu(T(v_2))R_{\rho,\mu}(T(v_1), T(v_3))(v_4) = 0,
\end{aligned}$$

which implies that (31) holds.

Similarly, by (18), one has

$$\begin{aligned}
&(F_1(v_1, v_2, v_3) + F_1(v_1, v_3, v_2) + F_2(v_2, v_3, v_1)) \diamond v_4 - F_2(v_2, v_3, v_1 \diamond v_4) + v_1 \diamond F_2(v_2, v_3, v_4) \\
&= \mu(P_{T(v_1)}(T(v_2), T(v_3))(v_4) - S_{\rho,\mu}(T(v_2), T(v_3))\mu(T(v_1))(v_4) \\
&\quad + \mu(T(v_1))S_{\rho,\mu}(T(v_2), T(v_3))(v_4) = 0,
\end{aligned}$$

which implies that (32) holds. Thus, $(V, \diamond, *)$ is a pre- F -manifold algebra. It is obvious that T is a homomorphism from V^c to $(A, \cdot_A, [-, -]_A)$. \square

Corollary 4.20. *Let $(A, \cdot_A, [-, -]_A)$ be an F -manifold algebra and $T : V \rightarrow A$ an O -operator on A with respect to the representation $(V; \rho, \mu)$. Then $T(V) = \{T(v) \mid v \in V\} \subset A$ is a subalgebra of A and there is an induced F -manifold algebras structure on $T(V)$ given by*

$$T(u) \diamond T(v) = T(u \diamond v), \quad T(u) * T(v) = T(u * v)$$

for all $u, v \in V$.

Corollary 4.21. *Let $(A, \cdot_A, [-, -]_A)$ be an F -manifold algebra and $\mathcal{B} : A \rightarrow A$ a Rota-Baxter operator. Define new operations on A by*

$$x \diamond y = \mathcal{B}(x) \cdot_A y, \quad x * y = [\mathcal{B}(x), y]_A.$$

*Then $(A, \diamond, *)$ is a pre- F -manifold algebra and \mathcal{B} is a homomorphism from the sub-adjacent F -manifold algebras $(A, \cdot_{\mathcal{B}}, [-, -]_{\mathcal{B}})$ to $(A, \cdot_A, [-, -]_A)$, where $x \cdot_{\mathcal{B}} y = x \diamond y + y \diamond x$ and $[x, y]_{\mathcal{B}} = x * y - y * x$.*

Corollary 4.22. *Consider the F -manifold algebra $(A, \cdot, [-, -])$ given by Proposition 3.15. If $\mathcal{B} : A \rightarrow A$ is a Rota-Baxter operator on the commutative associative algebra (A, \cdot) and satisfies $\mathcal{B} \circ D = D \circ \mathcal{B}$, then $\mathcal{B} : A \rightarrow A$ is a Rota-Baxter operator on the Lie algebra $(A, [-, -])$. Thus $(A, \diamond_{\mathcal{B}}, *_{\mathcal{B}})$ is a pre- F -manifold algebra, where*

$$x \diamond_{\mathcal{B}} y = \mathcal{B}(x) \cdot y, \quad x *_{\mathcal{B}} y = \mathcal{B}(x) \cdot D(y) - y \cdot D(\mathcal{B}(x))$$

and $(A, \cdot_{\mathcal{B}}, [-, -]_{\mathcal{B}})$ is the sub-adjacent F -manifold algebra with

$$x \cdot_{\mathcal{B}} y = x \diamond_{\mathcal{B}} y + y \diamond_{\mathcal{B}} x, \quad [x, y]_{\mathcal{B}} = x *_{\mathcal{B}} y - y *_{\mathcal{B}} x.$$

Example 4.23. We put $A = C^1([0, 1])$. Then $(A, \cdot, [-, -])$ with the following multiplication and bracket

$$\begin{aligned}
f \cdot g &= fg, \\
[f, g] &= fg' - gf', \quad \forall f, g \in A
\end{aligned}$$

is an F -manifold algebra. It is well-known that the integral operator is a Rota-Baxter operator:

$$\mathcal{B} : A \rightarrow A, \quad \mathcal{B}(f)(x) := \int_0^x f(t)dt.$$

It is easy to see that

$$\mathcal{B} \circ \partial_x = \partial_x \circ \mathcal{B} = \text{Id}.$$

Thus $(A, \diamond_{\mathcal{B}}, *_{\mathcal{B}})$ is a pre- F -manifold algebra, where

$$f \diamond_{\mathcal{B}} g = g \int_0^x f(t)dt, \quad f *_{\mathcal{B}} g = g' \int_0^x f(t)dt - f \cdot g$$

and $(A, \cdot_{\mathcal{B}}, [-, -]_{\mathcal{B}})$ is the sub-adjacent F -manifold algebra with

$$f \cdot_{\mathcal{B}} g = f \int_0^x g(t)dt + g \int_0^x f(t)dt, \quad [f, g]_{\mathcal{B}} = f' \int_0^x g(t)dt - g' \int_0^x f(t)dt.$$

Example 4.24. Consider the F -manifold algebra $(A, \cdot, [-, -])$ given by Example 3.18. It is straightforward to check that \mathcal{B} given by

$$\mathcal{B}(e_1) = re_1, \quad \mathcal{B}(e_2) = 2se_1 + \frac{3}{2}re_2, \quad \mathcal{B}(e_3) = te_1 + 3se_2 + 3re_3, \quad r, s, t \in \mathbb{K}$$

is a Rota-Baxter operator on the F -manifold algebra A . Thus $(A, \diamond_{\mathcal{B}}, *_{\mathcal{B}})$ is a pre- F -manifold algebra, where

$$\begin{aligned} e_2 \diamond_{\mathcal{B}} e_3 &= \frac{3}{2}re_1, & e_3 \diamond_{\mathcal{B}} e_2 &= 3re_1, & e_3 \diamond_{\mathcal{B}} e_3 &= 3se_1 + 3re_2; \\ e_2 *_{\mathcal{B}} e_3 &= -\frac{3}{2}are_1, & e_3 *_{\mathcal{B}} e_2 &= 3are_1, & e_3 *_{\mathcal{B}} e_3 &= -3ase_1 \end{aligned}$$

and $(A, \cdot_{\mathcal{B}}, [-, -]_{\mathcal{B}})$ is the sub-adjacent F -manifold algebra with

$$\begin{aligned} e_2 \cdot_{\mathcal{B}} e_3 &= \frac{9}{2}re_1, & e_3 \cdot_{\mathcal{B}} e_3 &= 6se_1 + 6re_2; \\ [e_2, e_3]_{\mathcal{B}} &= -\frac{9}{2}are_1. \end{aligned}$$

At the end of this section, we give a necessary and sufficient condition on an F -manifold algebra admitting a pre- F -manifold algebra structure.

Proposition 4.25. *Let $(A, \cdot_A, [-, -]_A)$ be an F -manifold algebra. There is a pre- F -manifold algebra structure on A such that its sub-adjacent F -manifold algebra is exactly $(A, \cdot_A, [-, -]_A)$ if and only if there exists an invertible \mathcal{O} -operator on $(A, \cdot_A, [-, -]_A)$.*

Proof. If $T : V \rightarrow A$ is an invertible \mathcal{O} -operator on A with respect to the representation $(V; \rho, \mu)$, then the compatible pre- F -manifold algebra structure on A is given by

$$x \diamond y = T(\mu(x)(T^{-1}(y))), \quad x * y = T(\rho(x)(T^{-1}(y)))$$

for all $x, y \in P$.

Conversely, let $(A, \diamond, *)$ be a pre- F -manifold algebra and $(A, \cdot_A, [-, -]_A)$ the sub-adjacent F -manifold algebra. Then the identity map Id is an \mathcal{O} -operator on A with respect to the representation $(A; L, \mathfrak{Q})$. \square

Corollary 4.26. *Let $(A, \cdot_A, [-, -]_A)$ be a coherence F -manifold algebra. Let $\omega \in \wedge^2 A^*$ be a cyclic 2-cocycle in the sense of Connes on the commutative associative algebra (A, \cdot_A) , i.e.*

$$\omega(x \cdot_A y, z) + \omega(y \cdot_A z, x) + \omega(z \cdot_A x, y) = 0, \quad \forall x, y, z \in A,$$

as well as a symplectic structure on the Lie algebra $(A, [-, -]_A)$, i.e.

$$\omega([x, y]_A, z) + \omega([y, z]_A, x) + \omega([z, x]_A, y) = 0, \quad \forall x, y, z \in A.$$

Then there is a compatible pre- F -manifold algebra structure on A given by

$$\omega(x \diamond y, z) = \omega(y, x \cdot_A z), \quad \omega(x * y, z) = \omega(y, [z, x]_A).$$

Proof. Since $(A, \cdot_A, [-, -]_A)$ is a coherence F -manifold algebra, $(A^*; \text{ad}^*, -\mathcal{L}^*)$ is a representation of A . By the fact that ω is a cyclic 2-cocycle, $(\omega^\sharp)^{-1}$ is an \mathcal{O} -operator on the commutative associative algebra (A, \cdot_A) with respect to the representation $(A^*, -\mathcal{L}^*)$, where $\omega^\sharp : A \rightarrow A^*$ is defined by $\langle \omega^\sharp(x), y \rangle = \omega(x, y)$. By the fact that ω is a symplectic structure, $(\omega^\sharp)^{-1}$ is an \mathcal{O} -operator on the Lie algebra $(A, [-, \cdot]_A)$ with respect to the representation (A^*, ad^*) . Thus, $(\omega^\sharp)^{-1}$ is an \mathcal{O} -operator on the coherence F -manifold algebra $(A, \cdot_A, [-, -]_A)$ with respect to the representation $(A^*; \text{ad}^*, -\mathcal{L}^*)$. By Proposition 4.25, there is a compatible pre- F -manifold algebra structure on A given as above. \square

5. DUAL PRE- F -MANIFOLD ALGEBRAS AND AVERAGE OPERATORS ON F -MANIFOLD ALGEBRAS

It is well known that the Koszul dual of the operad PreLie of pre-Lie algebras is the operad Perm of permutative algebras ([29]). Recall that a **permutative algebra** is a pair (A, \bullet) , where A is a vector space and $\bullet : A \otimes A \rightarrow A$ is a bilinear multiplication satisfying that for all $x, y, z \in A$,

$$(40) \quad x \bullet (y \bullet z) = (x \bullet y) \bullet z = (y \bullet x) \bullet z.$$

It is well known that the Koszul dual of the operad Zinb of Zinbiel algebras is the operad Leib of Leibniz algebras ([8]). Recall that a **Leibniz algebra** is a pair $(g, \{-, -\})$, where g is a vector space and $\{-, -\} : g \otimes g \rightarrow g$ is a bilinear bracket satisfying that for all $x, y, z \in g$,

$$(41) \quad \{x, \{y, z\}\} = \{\{x, y\}, z\} + \{y, \{x, z\}\}.$$

Now we give the definition of a dual pre- F -manifold algebra.

Definition 5.1. A **dual pre- F -manifold algebra** is a triple $(A, \bullet, \{-, -\})$, where (A, \bullet) is a permutative algebra and $(A, \{-, -\})$ is a Leibniz algebra, such that for all $x, y, z, w \in A$, the following conditions hold:

$$(42) \quad G_1(x \bullet y, z, w) = x \bullet G_1(y, z, w) + y \bullet G_1(x, z, w),$$

$$(43) \quad G_2(y \bullet x, z, w) = y \bullet G_2(x, z, w) - G_1(y, z, w) \bullet x,$$

$$(44) \quad \{x, y\} \bullet z + \{y, x\} \bullet z = 0,$$

where $G_1, G_2 : \otimes^3 A \rightarrow A$ are defined by

$$(45) \quad G_1(x, y, z) = \{x, y \bullet z\} - \{x, y\} \bullet z - y \bullet \{x, z\},$$

$$(46) \quad G_2(x, y, z) = \{y \bullet z, x\} - z \bullet \{y, x\} - y \bullet \{z, x\}.$$

Recall that an **average operator** on a commutative associative algebra (A, \cdot) is a linear map $\alpha : A \rightarrow A$ such that

$$\alpha(x) \cdot \alpha(y) = \alpha(\alpha(x) \cdot y), \quad \forall x, y \in A.$$

Lemma 5.2. ([1]) Let (A, \cdot) be a commutative associative algebra and $\alpha : A \rightarrow A$ an average operator. Define a multiplication \bullet on A by

$$(47) \quad x \bullet y = \alpha(x) \cdot y, \quad \forall x, y \in A.$$

Then (A, \bullet) is a permutative algebra.

Recall that an **average operator** on a Lie algebra $(\mathfrak{g}, [-, -])$ is a linear map $\alpha : \mathfrak{g} \rightarrow \mathfrak{g}$ such that

$$[\alpha(x), \alpha(y)] = \alpha([\alpha(x), y]), \quad \forall x, y \in \mathfrak{g}.$$

Lemma 5.3. ([1]) *Let $(\mathfrak{g}, [-, -])$ be a Lie algebra and $\alpha : \mathfrak{g} \rightarrow \mathfrak{g}$ an average operator. Define a bracket $\{-, -\}$ on \mathfrak{g} by*

$$(48) \quad \{x, y\} = [\alpha(x), y], \quad \forall x, y \in \mathfrak{g}.$$

Then $(\mathfrak{g}, \{-, -\})$ is a Leibniz algebra.

Definition 5.4. *Let $(A, \cdot_A, [-, -]_A)$ be an F -manifold algebra. A linear operator $\alpha : A \rightarrow A$ is called an **average operator on A** if α is both an average operator on the commutative associative algebra (A, \cdot_A) and an average operator on the Lie algebra $(A, [-, -]_A)$.*

Theorem 5.5. *Let $(A, \cdot_A, [-, -]_A)$ be an F -manifold algebra and $\alpha : A \rightarrow A$ an average operator. Define new operations on A by*

$$(49) \quad x \bullet y = \alpha(x) \cdot_A y \quad \text{and} \quad \{x, y\} = [\alpha(x), y]_A, \quad \forall x, y \in A.$$

Then $(A, \bullet, \{-, -\})$ is a dual pre- F -manifold algebras.

Proof. Let $(A, \cdot_A, [-, -]_A)$ be an F -manifold algebra. Since $\alpha : A \rightarrow A$ is an average operator, we have $[\alpha(x), \alpha(y)]_A = \alpha([\alpha(x), y]_A)$ and $\alpha(x) \cdot_A \alpha(y) = \alpha(\alpha(x) \cdot_A y)$. By Lemma 5.2 and Lemma 5.3, (A, \cdot_A) is a permutative algebra and $(A, [-, -]_A)$ is a Leibniz algebra. By a direct calculation, we have

$$(50) \quad P_{\alpha(x)}(\alpha(y), z) = P_{\alpha(x)}(z, \alpha(y)) = G_1(x, y, z), \quad P_x(\alpha(y), \alpha(z)) = -G_2(x, y, z).$$

By the definition of an average operator and (50), we have

$$\begin{aligned} & P_{\alpha(x) \cdot_A \alpha(y)}(\alpha(z), w) - \alpha(x) \cdot_A P_{\alpha(y)}(\alpha(z), w) - \alpha(y) \cdot_A P_{\alpha(x)}(\alpha(z), w) \\ &= P_{\alpha(x \cdot_A y)}(\alpha(z), w) - x \bullet P_{\alpha(y)}(\alpha(z), w) - y \bullet P_{\alpha(x)}(\alpha(z), w) \\ &= G_1(x \bullet y, z, w) - x \bullet G_1(y, z, w) - y \bullet G_1(x, z, w), \end{aligned}$$

which implies that (42) holds.

Similarly, we have

$$\begin{aligned} & P_{x \cdot_A \alpha(y)}(\alpha(z), \alpha(w)) - x \cdot_A P_{\alpha(y)}(\alpha(z), \alpha(w)) - \alpha(y) \cdot_A P_x(\alpha(z), \alpha(w)) \\ &= P_{y \bullet x}(\alpha(z), w) - x \cdot_A \alpha(G_1(y, z, w)) - y \bullet P_x(\alpha(z), \alpha(w)) \\ &= -G_2(y \bullet x, z, w) - G_1(y, z, w) \bullet x + y \bullet G_2(x, z, w), \end{aligned}$$

which implies that (43) holds.

Since $\alpha([\alpha(x), y]_A) = \alpha([x, \alpha(y)]_A) = [\alpha(x), \alpha(y)]_A$, we have

$$\begin{aligned} \{x, y\} \bullet z + \{y, x\} \bullet z &= \alpha([\alpha(x), y]_A) \cdot_A z + ([\alpha(y), x]_A) \cdot_A z \\ &= ([\alpha(x), \alpha(y)]_A + [\alpha(y), \alpha(x)]_A) \cdot_A z = 0, \end{aligned}$$

which implies that (44) holds. \square

Proposition 5.6. *Consider the F -manifold algebra $(A, \cdot, [-, -])$ given by Proposition 3.15. If $\alpha : A \rightarrow A$ is an average operator on the commutative associative algebra (A, \cdot) and satisfies*

$\alpha \circ D = D \circ \alpha$, then $\alpha : A \longrightarrow A$ is an average operator on the Lie algebra $(A, [-, -])$. Thus $(A, \bullet, \{-, -\})$ is a dual pre- F -manifold algebra, where

$$x \bullet y = \alpha(x) \cdot y \quad \text{and} \quad \{x, y\} = \alpha(x) \cdot D(y) - y \cdot D(\alpha(x)), \quad \forall x, y \in A.$$

Proof. It follows by a direct calculation. \square

Example 5.7. Consider the F -manifold algebra $(A, \cdot, [-, -])$ given by Example 3.18. It is straightforward to check that α given by

$$\alpha(e_1) = re_1, \quad \alpha(e_2) = se_1 + re_2, \quad \alpha(e_3) = te_1 + se_2 + re_3, \quad r, s, t \in \mathbb{K}$$

is an average operator on the F -manifold algebra A . Thus $(A, \bullet, \{-, -\})$ is a dual pre- F -manifold algebra, where

$$\begin{aligned} e_2 \bullet e_3 &= re_1, & e_3 \bullet e_2 &= re_1, & e_3 \bullet e_3 &= se_1 + re_2; \\ \{e_2, e_3\} &= -are_1, & \{e_3, e_2\} &= are_1, & \{e_3, e_3\} &= -ase_1 \quad a, r, s \in \mathbb{K}. \end{aligned}$$

REFERENCES

- [1] M. Aguiar, Pre-Poisson algebras. *Lett. Math. Phys.* 54 (2000), 263-277. [2](#), [15](#), [19](#), [20](#)
- [2] C. Bai, A unified algebraic approach to classical Yang-Baxter equation. *J. Phys. A: Math. Theo.* 40 (2007), 11073-11082. [2](#), [16](#)
- [3] C. Bai, L. Guo and X. Ni, \mathcal{O} -operators on associative algebras and associative Yang-Baxter equations. *Pacific J. Math.* 256 (2012), 257-289. [2](#), [16](#)
- [4] C. Bai, O. Bellier, L. Guo and X. Ni, Splitting of operations, Manin products and Rota-Baxter operators. *Int. Math. Res. Not.* 3 (2013), 485-524. [2](#), [15](#)
- [5] B. Bakalov and V. Kac, Field algebras. *Int. Math. Res. Not.* 3 (2003), 123-159. [3](#)
- [6] R. Bandiera, Formality of Kapranovs brackets in Kähler geometry via pre-Lie deformation theory. *Int. Math. Res. Not.* 21 (2016), 6626-6655. [3](#)
- [7] D. Burde, Left-symmetric algebras and pre-Lie algebras in geometry and physics. *Cent. Eur. J. Math.* 4 (2006), 323-357. [3](#), [4](#)
- [8] F. Chapoton and M. Livernet, Pre-Lie algebras and the rooted trees operad. *Int. Math. Res. Not.* 8 (2001), 395-408. [3](#), [19](#)
- [9] A. Connes and D. Kreimer, Renormalization in quantum field theory and the Riemann-Hilbert problem. I. The Hopf algebra structure of graphs and the main theorem. *Comm. Math. Phys.* 210 (2000), 249-273. [2](#)
- [10] L. David and I. A. B. Strachan, Compatible metrics on a manifold and nonlocal bi-Hamiltonian structures. *Int. Math. Res. Not.* 66 (2004), 3533-3557. [2](#)
- [11] L. David, and I. A. B. Strachan, Dubrovins duality for F -manifolds with eventual identities. *Adv. Math.* 226 (2011), 4031-4060. [2](#)
- [12] V. Dotsenko, S. Shadrin and B. Vallette, Pre-Lie deformation theory. *Mosc. Math. J.* 16 (2016), 505-543. [3](#)
- [13] V. Dotsenko, Algebraic structures of F -manifolds via pre-Lie algebras. *Ann. Mat. Pura Appl.* 198 (2019), 517-527. [2](#), [3](#), [5](#), [6](#)
- [14] B. Dubrovin, Geometry of 2D topological field theories. *Lecture Notes in Math*, 1620 (1995). [1](#)
- [15] A. Dzhumadil'daev, Cohomologies and deformations of right-symmetric algebras. *J. Math. Sci.* 93 (1999), 836-876. [4](#)
- [16] K. Ebrahimi-Fard, L. Guo and D. Kreimer, Integrable renormalization II: the general case. *Ann. Henri Poincare* 6 (2005), 369-395. [2](#)
- [17] V. T. Filippov, A class of simple nonassociative algebras, *Mat. Zametki* 45 (1989), 101-105. [7](#)
- [18] L. Foissy, The Hopf algebra of Fliess operators and its dual pre-Lie algebra. *Commun. Algebra* 43 (2015), 4528-4552. [6](#)
- [19] I. M. Gelfand and I. Ya. Dorfman, Hamiltonian operators and algebraic structures related to them. *Funct. Anal. Appl.* 13 (1979) 248-262. [7](#)

- [20] L. Guo, An introduction to Rota-Baxter algebra. *Surveys of Modern Mathematics*, 4. International Press, Somerville, MA; Higher Education Press, Beijing, 2012. xii+226 pp. [2](#)
- [21] L. Guo and W. Keigher, Baxter algebras and shuffle products. *Adv. Math.* 150 (2000), 117-149. [2](#)
- [22] C. Hertling, Frobenius Manifolds and Moduli Spaces for Singularities. *Cambridge Tracts in Math.* Cambridge University Press, 2002. [2](#)
- [23] C. Hertling and Y. I. Manin, Weak Frobenius manifolds. *Int. Math. Res. Not.* 6 (1999), 277-286. [2](#)
- [24] B. A. Kupersmidt, What a classical r -matrix really is, *J. Nonlinear Math. Phys.* 6 (1999), 448-488. [15](#)
- [25] Y. P. Lee, Quantum K-theory, I: Foundations. *Duke Math. J.* 121 (3) (2004), 389-424. [2](#)
- [26] A. Lichnerowicz and A. Medina, On Lie groups with left-invariant symplectic or Kählerian structures. *Lett. Math. Phys.* 16 (1988), 225-235. [3](#)
- [27] M. Livernet, Rational homotopy of Leibniz algebras. *Manuscripta Math.* 96 (1998), 295-315. [2](#)
- [28] J.-L. Loday, Cup product for Leibniz cohomology and dual Leibniz algebras. *Math. Scand.* 77, Univ. Louis Pasteur, Strasbourg, 1995, pp. 189-196. [2](#)
- [29] J.-L. Loday, Dialgebras. In *Dialgebras and related operads. Lecture Notes in Math.* 1763, Springer, Berlin 2001, 7-66. [2](#), [3](#), [19](#)
- [30] P. Lorenzoni, M. Pedroni and A. Raimondo, F -manifolds and integrable systems of hydrodynamic type. *Arch. Math.* (Brno) 47 (2011), 163-180. [2](#)
- [31] S. Majid and W. Tao, Noncommutative differentials on Poisson-Lie groups and pre-Lie algebras. *Pacific J. Math.* 284 (2016), 213-256. [3](#)
- [32] A. Mansuy, Preordered forests, packed words and contraction algebras. *J. Algebra* 411 (2014), 259-311. [6](#)
- [33] S. A. Merkulov, Operads, deformation theory and F -manifolds. *Asp. Math.* 36 (2004), 213-251. [2](#)
- [34] J. Pei, C. Bai and L. Guo, Splitting of operads and Rota-Baxter operators on operads. *Appl. Cate. Stru.* 25 (2017), 505-538. [2](#), [15](#)
- [35] X. Xu, On simple Novikov algebras and their irreducible modules. *J. Algebra* 185 (1996), 905-934. [7](#)

SCHOOL OF MATHEMATICS AND STATISTICS, NORTHEAST NORMAL UNIVERSITY, CHANGCHUN 130024, JILIN, CHINA
E-mail address: liujf12@126.com

DEPARTMENT OF MATHEMATICS, JILIN UNIVERSITY, CHANGCHUN 130012, JILIN, CHINA
E-mail address: shengyh@jlu.edu.cn

CHERN INSTITUTE OF MATHEMATICS AND LPMC, NANKAI UNIVERSITY, TIANJIN 300071, CHINA
E-mail address: baicm@nankai.edu.cn