

Multiple solutions for superlinear Klein-Gordon-Maxwell equations*

Dong-Lun Wu^{a,b}, Hongxia Lin^c

^aCollege of Science, Southwest Petroleum University,
Chengdu, Sichuan 610500, P.R. China

^bInstitute of Nonlinear Dynamics, Southwest Petroleum University,
Chengdu, Sichuan 610500, P.R. China

^cGeomathematics Key Laboratory of Sichuan Province, Chengdu University of Technology,
Chengdu, Sichuan 610059, P.R. China

Abstract In this paper, we consider the following Klein-Gordon-Maxwell equations

$$\begin{cases} -\Delta u + V(x)u - (2\omega + \phi)\phi u = f(x, u) + h(x) & \text{in } \mathbb{R}^3, \\ -\Delta \phi + \phi u^2 = -\omega u^2 & \text{in } \mathbb{R}^3, \end{cases}$$

where $\omega > 0$ is a constant, $u, \phi : \mathbb{R}^3 \rightarrow \mathbb{R}$, $V : \mathbb{R}^3 \rightarrow \mathbb{R}$ is a potential function. By assuming the coercive condition on V and some new superlinear conditions on f , we obtain two nontrivial solutions when h is nonzero and infinitely many solutions when f is odd in u and $h \equiv 0$ for above equations.

Key words Klein-Gordon-Maxwell equations; Superlinear conditions; Multiple solutions; Variational methods

1 Introduction and main results

In this paper, we considered the following Klein-Gordon-Maxwell equations

$$\begin{cases} -\Delta u + V(x)u - (2\omega + \phi)\phi u = f(x, u) + h(x) & \text{in } \mathbb{R}^3, \\ -\Delta \phi + \phi u^2 = -\omega u^2 & \text{in } \mathbb{R}^3, \end{cases} \quad (1)$$

where $\omega > 0$ is a constant, $u, \phi, V : \mathbb{R}^3 \rightarrow \mathbb{R}$. This type of equation has very interesting physical background which is a model to describe the nonlinear Klein-Gordon field interacting with the electromagnetic field. Along with the development of variational methods, many mathematicians

*D.-L. Wu is partially supported by NSF of China (No.11801472) and the Youth Science and Technology Innovation Team of Southwest Petroleum University for Nonlinear Systems (No.2017CXTD02) and The Science and Technology Innovation Team of Education Department of Sichuan for Dynamical System and its Applications (No.18TD0013). H.X. Lin is partially supported by the NSF of China (No.11701049), the China Postdoctoral Science Foundation (No.2017M622989), and the Scientific fund of the Education Department of Sichuan Province (No.18ZB0069).

E-mail: wudl2008@163.com; linhongxia18@126.com.

used these methods to investigate the existence and multiplicity of solutions for Klein-Gordon-Maxwell equations(see[1–3, 6–21, 24, 25, 27]). In 2001, V. Benci and D. Fortunato [3] considered the following systems

$$\begin{cases} -\Delta u + [m^2 - (\omega + \phi)^2]u = |u|^{q-2}u & \text{in } \mathbb{R}^3, \\ -\Delta \phi + \phi u^2 = -\omega u^2 & \text{in } \mathbb{R}^3. \end{cases} \quad (2)$$

By using the variational methods, they obtained infinitely many solitary wave solutions when $|m| > |\omega|$ and $4 < q < 6$. After this first work, many mathematicians have treated this problem with variant cases with m , ω and q . In 2004, D'Aprile and Mugnai [13] dealt with the case $q \in (2, 4]$ with $\sqrt{\left(\frac{q}{2} - 1\right)}m > \omega > 0$. Some existence and nonexistence results are obtained for problem (1) with variant conditions on m , ω and q (see [1, 2, 13, 14, 21] for more details).

Although we lose the compactness for problem (1) since the problem lies in a unbounded domain, we can also consider this problem in a radial symmetric space $H_r^1(\mathbb{R}^3) := \{u \in H^1(\mathbb{R}^3) : u = u(r), r = |x|\}$. It is known that $H_r^1(\mathbb{R}^3)$ is compactly embedded into $L^s(\mathbb{R}^3)$ for $2 < s < 2^*$. If $V(x)$ is not radial symmetric, there are still some other ways to retrieve the compactness. One classical way is to assume $V(x)$ to be coercive. In this paper, we mainly consider the coercive case. This case has also been studied in many papers(see [5, 9, 12, 15, 17–19]). Many authors(see [12, 18, 22, 23]) considered the following coercive condition.

(V1) Suppose $V \in C(\mathbb{R}^3, \mathbb{R})$, $\inf_{\mathbb{R}^3} V(x) = V_0 > 0$ and there is a constant $r > 0$ such that

$$\lim_{|y| \rightarrow +\infty} \text{meas}(\{x \in \mathbb{R}^3 : |x - y| \leq r, V(x) \leq M\}) = 0, \quad \forall M > 0,$$

where $\text{meas}(\cdot)$ denote the Lebesgue measure. The condition (V1) was introduced by T. Bartsch et al. [[5], Lemma 3.1.] to guarantee the compactness of the embedding. We will use this condition to prove our theorem.

As we know, the growth of the nonlinear terms is important in showing the geometric structure of the corresponding functionals(which is defined in (5)) and the boundedness of the Plais-Smale (PS) sequence. In [10], the author assumed the following condition.

(AR) there exists a constant $\theta > 4$ such that

$$f(x, t)t \geq \theta F(x, t) \quad \text{for } (x, t) \in \mathbb{R}^3 \times \mathbb{R},$$

where $F(x, t) = \int_0^t f(x, s)ds$. This condition is used to show the geometric structure of the corresponding functionals and the boundedness of the (PS) sequences. Hence (AR) has been widely used in many papers to obtain the existence and multiplicity of elliptic problem with variational methods. This condition implies that the growth exponent of the nonlinear terms is more than 4 at infinity. In order to deal with other nonlinearities with 4-superlinear growth, many authors(see[15, 17]) considered the following growth conditions.

(SL1) $f(x, t)/t^3 \rightarrow \infty$ as $|t| \rightarrow +\infty$ uniformly in x ;

(SL2) $\frac{1}{4}f(x, t)t - F(x, t) \geq -Dt^2$ for $D > 0$ and $|t|$ large enough uniformly in x .

(SL1) and (SL2) has been directly or indirectly used to show the existence and multiplicity of solutions for problem (1). However, above conditions on f eliminate many nonlinearities such as $F(x, t) = |t|^{5/2}$.

In order to deal with the nonlinearities with growth exponent between 2 and 4, Chen and Song [9] obtained two solutions for problem (1) under (AR) condition just requiring $\theta > 2$ with coercive potentials when $h(x) \not\equiv 0$. In 2018, Chen and Tang [12] introduced the following superlinear condition which is weaker than (AR) and obtained infinitely many solutions for problem (1).

(WAR) there exist constants $\theta > 2$ and $K > 0$ such that

$$f(x, t)t - \theta F(x, t) + Kt^2 \geq 0 \quad \text{for } (x, t) \in \mathbb{R}^3 \times \mathbb{R}.$$

There are still many functions cannot be included in above conditions. In this paper, we introduce some new superlinear conditions and an example is given to show the difference from previous conditions. Now, we state our main results.

Theorem 1.1. *Suppose that V satisfies (V1) and $f \in C(\mathbb{R}^3 \times \mathbb{R}, \mathbb{R})$ satisfies*

(F1) $f(x, t) = o(|t|)$ as $t \rightarrow 0$ for all $x \in \mathbb{R}^3$;

(F2) $F(x, 0) = 0$ for all $x \in \mathbb{R}^3$ and there exist $\tau \in (2, 6)$ and $d_1 > 0$ such that

$$|f(x, t)| \leq d_1(|t| + |t|^{\tau-1}) \quad \text{for all } (x, t) \in \mathbb{R}^3 \times \mathbb{R};$$

(F3) let $\tilde{F}(x, t) = f(x, t)t - 2F(x, t)$, then there exist $d_2 > 0$ and $r_0 > 0$ such that

$$\tilde{F}(x, t) \geq d_2|t|^\tau \quad \text{for all } |t| \geq r_0 \quad \text{and } x \in \mathbb{R}^3;$$

(F4) $F(x, t)/t^2 \rightarrow +\infty$ as $|t| \rightarrow \infty$ uniformly in x ;

(F5) $f(x, t)t \geq 2F(x, t) \geq 0$ for all $(x, t) \in \mathbb{R}^3 \times \mathbb{R}$.

Then there is a constant $m > 0$ such that, for any $h \not\equiv 0$ satisfying $\|h\|_2 < m$, problem (1) possesses at least two nontrivial solutions.

Theorem 1.2. *Suppose that (V1), (F2)-(F5) hold, $h(x) \equiv 0$ and $f(x, -t) = -f(x, t)$, for all $(x, t) \in \mathbb{R}^3 \times \mathbb{R}$. Then problem (1) possesses infinitely many solutions.*

Remark 1. In 2018, Shi and Chen[19] obtained two solutions for problem (1) with nonzero perturbation by using the combination of a cut-off functional and a Pohozaev type identity. Although $f(x, t)$ was required to satisfy some very weak growth conditions in their paper, they needed some smooth conditions on the gradients of $V(x)$ and $h(x)$, which are not needed in our theorems.

Remark 2. Setting $4 > p > 2$, $0 < \epsilon < p - 2$, consider

$$F(x, t) = |t|^p + a(p - 2)|t|^{p-\epsilon} \sin^2(|t|^\epsilon/\epsilon). \quad (3)$$

For any $\theta > 2$ and $K > 0$, let $\max\left\{0, \frac{p-\theta}{p-2}\right\} < a < 1$ and $t_n = (\epsilon(n\pi + \frac{3\pi}{4}))^{1/\epsilon}$, then

$$\begin{aligned} & f(x, t_n)t_n - \theta F(x, t_n) - Kt_n^2 \\ &= (p - \theta)|t_n|^p + a(p - 2)(p - \theta - \epsilon)|t_n|^{p-\epsilon} \sin^2(|t_n|^\epsilon/\epsilon) + a(p - 2)|t_n|^p \sin(2|t_n|^\epsilon/\epsilon) - Kt_n^2 \\ &= |t_n|^p \left[(p - \theta) - a(p - 2) + \frac{a(p - 2)(p - \theta - \epsilon)}{2|t_n|^\epsilon} \right] - Kt_n^2 \\ &\rightarrow -\infty \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Hence (3) does not satisfy (WAR) or the following condition

(FSL) $F(x, t)/t^4 \rightarrow +\infty$ as $|t| \rightarrow +\infty$ uniformly in x .

However, it is easy to see that, for all $x \in \mathbb{R}^3$,

$$\begin{aligned} f(x, t)t - 2F(x, t) &= (p-2)|t|^p \left[(1 + a \sin(2|t|^\epsilon/\epsilon)) + \frac{a(p-2-\epsilon) \sin^2(|t|^\epsilon/\epsilon)}{|t|^\epsilon} \right] \\ &\geq \frac{1}{2}(1-a)(p-2)|t|^p \end{aligned}$$

for $|t|$ large enough. Then we can check that (3) satisfies conditions (F1)-(F5).

2 Preliminaries

Let

$$E := \left\{ u \in H^1(\mathbb{R}^3) : \int_{\mathbb{R}^3} (|\nabla u|^2 + V(x)u^2) dx < \infty \right\}.$$

Then E is a Hilbert space with the inner product

$$(u, v)_E = \int_{\mathbb{R}^3} (\nabla u \cdot \nabla v + V(x)uv) dx$$

and the norm $\|u\|_E = (u, u)_E^{1/2}$. Define the function space

$$D^{1,2}(\mathbb{R}^3) := \{u \in L^6(\mathbb{R}^3) : |\nabla u| \in L^2(\mathbb{R}^3)\}$$

with the norm

$$\|u\|_{D^{1,2}} := \left(\int_{\mathbb{R}^3} |\nabla u|^2 dx \right)^{\frac{1}{2}}.$$

Under (V1), the embedding $E \hookrightarrow L^s(\mathbb{R}^3)$ are compact for any $s \in [2, 6)$ (see [5]). Hence for each $s \in [2, 6)$, there exists a constant $C_s > 0$ such that

$$\|u\|_{L^s} \leq C_s \|u\|_E, \quad \forall u \in E. \quad (4)$$

Obviously, problem (1) has a variational structure. Consider $J : E \times D^{1,2}(\mathbb{R}^3) \rightarrow \mathbb{R}$ defined by

$$J(u, \phi) = \frac{1}{2} \int_{\mathbb{R}^3} (|\nabla u|^2 - |\nabla \phi|^2 + [V(x) - (2\omega + \phi)\phi]u^2) dx - \int_{\mathbb{R}^3} F(x, u) dx - \int_{\mathbb{R}^3} h(x)u dx.$$

Evidently, J belongs to $C^1(E \times D^{1,2}(\mathbb{R}^3), \mathbb{R})$. Thus, the pair (u, ϕ) is a weak solution of problem (1) if and only if it is a critical point of J in $E \times D^{1,2}(\mathbb{R}^3)$. We can also see that J is strong indefinite. To reduce this functional, we need the following lemmas.

Lemma 2.1. (see[13]) *For every $u \in H^1(\mathbb{R}^3)$ there exists a unique $\phi_u \in D^{1,2}(\mathbb{R}^3)$ which solves the second equation of problem (1). Furthermore,*

- (i) $-\omega \leq \phi_u \leq 0$ in \mathbb{R}^3 ;
- (ii) if u is radially symmetric, ϕ_u is radial too.

We can consider the functional $I : E \rightarrow \mathbb{R}$ defined by $I(u) = J(u, \phi_u)$. Therefore

$$I(u) = \frac{1}{2} \int_{\mathbb{R}^3} (|\nabla u|^2 + V(x)u^2 - \omega \phi_u u^2) dx - \int_{\mathbb{R}^3} F(x, u) dx - \int_{\mathbb{R}^3} h(x)u dx. \quad (5)$$

and we have, for any $u, v \in E$,

$$\langle I'(u), v \rangle = \int_{\mathbb{R}^3} (\nabla u \cdot \nabla v + V(x)uv - (2\omega + \phi_u)\phi_u uv - f(x, u)v - h(x)v) dx. \quad (6)$$

Lemma 2.2. *The following statements are equivalent:*

- (1) $(u, \phi) \in H^1(\mathbb{R}^3) \times D^{1,2}(\mathbb{R}^3)$ is a critical point of J (i.e. (u, ϕ) is a solution of system (1));
- (2) u is a critical point of I and $\phi = \phi_u$.

By Lemmas (2.1) and (2.2), we only need to look for the critical points of I to show the existence and multiplicity of critical points for J . Some details can be found in [18]. Subsequently, we prove our theorems with mountain pass theorem and a abstract critical point theorem introduced by T. Bartsch.

3 Proof of Theorem 1.1

Lemma 3.1. *Suppose that (V1), (F1) and (F2) hold. Let $h \in L^2(\mathbb{R}^3)$, then there exist some constants $\rho, \alpha, m > 0$ such that $I(u)|_{\|u\|_E=\rho} \geq \alpha$ for all h satisfying $\|h\|_{L^2} < m$.*

Proof. The proof is similar to Lemma 3.1 in [9]. □

Lemma 3.2. *Suppose that (V1), (F4), (F5) hold, then there exists $v \in E$ with $\|v\|_E > \rho$ such that $I(v) < 0$, where ρ is given in Lemma 3.1.*

Proof. Set $e \in C_0^\infty(B_1(0), \mathbb{R})$, where $B_1(0) = \{x \in \mathbb{R}^3 : |x| \leq 1\}$, such that $\|e\|_E = 1$ and $A = \frac{2(1+\int_{B_1(0)} \omega^2 e^2 dx)}{\int_{B_1(0)} e^2 dx}$. It follows from (F4) and (F5) that there exists $Q > 0$ such that

$$F(x, t) \geq A(t^2 - Q^2) \quad (7)$$

for all $(x, t) \in B_1(0) \times \mathbb{R}$. By (7) and Lemma 2.1, for every $\eta \in \mathbb{R}^+$, we have

$$\begin{aligned} I(\eta e) &= \frac{\eta^2}{2} \int_{B_1(0)} (|\nabla e|^2 + V(x)e^2 - \omega \phi_{\eta e} e^2) dx - \int_{B_1(0)} F(x, \eta e) dx - \eta \int_{B_1(0)} h(x) e dx \\ &\leq \frac{\eta^2}{2} \left(1 + \int_{B_1(0)} \omega^2 e^2 dx - 2A \int_{B_1(0)} e^2 dx \right) + \frac{4\pi A Q^2}{3} - \eta \int_{B_1(0)} h(x) e dx \\ &\leq -\frac{3\eta^2}{2} \left(1 + \int_{B_1(0)} \omega^2 e^2 dx \right) + \frac{4\pi A Q^2}{3} - \eta \int_{B_1(0)} h(x) e dx. \end{aligned}$$

Then $I(\eta e) \rightarrow -\infty$ as $\eta \rightarrow +\infty$. Therefore, there exists $\eta_0 > 0$ such that $I(\eta_0 e) < 0$ and $\|\eta_0 e\|_E > \rho$. Let $v = \eta_0 e$, we can see $I(v) < 0$, which proves this lemma. □

Lemma 3.3. *Assume that (V1) and (F2) – (F5) hold, then I satisfies the (PS) condition.*

Proof. Suppose that $\{u_n\} \subset E$ is a sequence such that $\{I(u_n)\}$ is bounded and $I'(u_n) \rightarrow 0$ as $n \rightarrow \infty$. Then there exists a constant $\overline{M} > 0$ such that

$$|I(u_n)| \leq \overline{M}, \quad \|I'(u_n)\| \leq \overline{M}. \quad (8)$$

Now we prove that $\{u_n\}$ is bounded in E . Arguing in an indirect way, we assume that $\|u_n\|_E \rightarrow +\infty$ as $n \rightarrow \infty$. Set $w_n = \frac{u_n}{\|u_n\|_E}$. Then $\|w_n\|_E = 1$ and there exists a subsequence of $\{w_n\}$, still denoted by $\{w_n\}$, such that $w_n \rightharpoonup w_0$ in E . Then we have

$$w_n \rightarrow w_0 \text{ in } L^s(\mathbb{R}^3) \text{ for any } s \in [2, 6).$$

Let $\Omega = \{x \in \mathbb{R}^3 \mid |w_0(x)| > 0\}$. If $\text{meas}(\Omega) > 0$, we have $|u_n| \rightarrow +\infty$ as $n \rightarrow \infty$ for a.e. $x \in \Omega$. On one hand, by (F4), (F5) and Fatou's Lemma, we obtain

$$\begin{aligned} \liminf_{n \rightarrow \infty} \int_{\mathbb{R}^3} \frac{F(x, u_n)}{\|u_n\|_E^2} dx &\geq \liminf_{n \rightarrow \infty} \int_{\Omega} \frac{F(x, u_n)}{\|u_n\|_E^2} dx \\ &= \liminf_{n \rightarrow \infty} \int_{\Omega} \frac{F(x, u_n)}{|u_n|^2} |u_n|^2 dx \\ &= +\infty. \end{aligned}$$

On the other hand, by (8) and Lemma 2.1, we get

$$\begin{aligned} \left| \int_{\mathbb{R}^3} \frac{F(x, u_n)}{\|u_n\|_E^2} dx - \frac{1}{2} \right| &= \left| -\frac{I(u_n)}{\|u_n\|_E^2} - \frac{\omega}{2} \int_{\mathbb{R}^3} \frac{\phi_u u_n^2}{\|u_n\|_E^2} dx - \int_{\mathbb{R}^3} \frac{h(x) u_n}{\|u_n\|_E^2} dx \right| \\ &\leq \frac{\overline{M}}{\|u_n\|_E^2} + \frac{\omega^2 C_2}{2} + \frac{C_2 \|h\|_2}{\|u_n\|_E} \rightarrow \frac{\omega^2 C_2}{2} \text{ as } n \rightarrow \infty, \end{aligned}$$

which is a contradiction. Then we have $\text{meas}(\Omega) = 0$, which implies that $w_0 = 0$ a.e. $x \in \mathbb{R}^3$ and $w_n \rightarrow 0$ in $L^s(\mathbb{R}^3)$ ($2 \leq s < 6$). By (5), (6), (F3) and (F5), we obtain

$$\begin{aligned} 2I(u_n) - \langle I'(u_n), u_n \rangle &= \int_{\mathbb{R}^3} (\omega + \phi_{u_n}) \phi_{u_n} u_n^2 dx + \int_{\mathbb{R}^3} \tilde{F}(x, u_n) dx - \int_{\mathbb{R}^3} h(x) u_n dx \\ &\geq \int_{\mathbb{R}^3} (\omega + \phi_{u_n}) \phi_{u_n} u_n^2 dx + d_2 \int_{|u_n| \geq r_0} |u_n|^\tau dx - \int_{\mathbb{R}^3} h(x) u_n dx, \end{aligned}$$

which implies that

$$\lim_{n \rightarrow \infty} \frac{\int_{|u_n| \geq r_0} |u_n|^\tau dx}{\|u_n\|_E^2} = 0. \quad (9)$$

By (6), (9), (F2) and (4), one sees that

$$\begin{aligned} \frac{\overline{M} + 1}{\|u_n\|_E} &\geq \frac{\langle I'(u_n), u_n \rangle}{\|u_n\|_E^2} \\ &\geq 1 - d_1 \int_{\mathbb{R}^3} \left(w_n^2 + \frac{|u_n|^\tau}{\|u_n\|_E^2} \right) dx - \frac{\|h\|_2 \|u_n\|_2}{\|u_n\|_E^2} \\ &\geq 1 - d_1 \int_{\mathbb{R}^3} w_n^2 dx - d_1 \left(\frac{\int_{|u_n| \geq r_0} |u_n|^\tau dx}{\|u_n\|_E^2} + r_0^{\tau-2} \int_{|u_n| \leq r_0} w_n^2 dx \right) - \frac{C_2 \|h\|_2}{\|u_n\|_E} \\ &\rightarrow 1 \text{ as } n \rightarrow \infty, \end{aligned}$$

which is a contradiction, then $\{u_n\}$ is bounded in E . Similar to Lemma 3.3 in [9], we can see that $\{u_n\}$ has a strong convergent subsequence. Then I satisfies the (PS) condition. \square

Proof of Theorem 1.1. In order to obtain two nontrivial solutions for problem (1), we will apply the Ekeland's variational principle and the mountain pass theorem to the functional I . The rest proof of Theorem 1.1 is similar to the proof of Theorem 1.2 in [9].

4 Proof of Theorem 1.2

In order to obtain infinitely many solutions of (1), similar to [18], we shall use the abstract critical point theorem introduced by T. Bartsch in [4]. Let space X be reflexive and separable, then there exist $e_i \in X$ and $f_i \in X^*$ such that $X = \overline{\langle e_i, i \in \mathbb{N} \rangle}$, $X^* = \overline{\langle f_i, i \in \mathbb{N} \rangle}$, $\langle e_i, f_j \rangle = \delta_{i,j}$, where $\delta_{i,j}$ denotes the Kronecker symbol. Subsequently, put

$$X_k = \text{span}\{e_k\}, \quad Y_k = \bigoplus_{i=1}^k X_i, \quad Z_k = \overline{\bigoplus_{i=k}^{\infty} X_i}.$$

Now we state the critical points theorem by T. Bartsch.

Lemma 4.1. *Assume $\phi \in C^1(X, \mathbb{R})$ satisfies the (PS) condition, $\phi(-u) = \phi(u)$. For every $k \in \mathbb{N}$, there exists $\rho_k > r_k > 0$, such that*

- (i) $a_k := \max_{u \in Y_k, \|u\|=\rho_k} \phi(u) \leq 0$;
- (ii) $b_k := \inf_{u \in Z_k, \|u\|=r_k} \phi(u) \rightarrow +\infty$ as $k \rightarrow \infty$.

Then ϕ has a sequence of critical values tending to $+\infty$

Lemma 4.2. *Assume that (V1), (F2), (F4) and (F5) hold. Then for every $k \in \mathbb{N}$, there exists $\rho_k > r_k > 0$, such that (i) $a_k := \inf_{u \in Z_k, \|u\|_E=r_k} I(u) \rightarrow +\infty$ as $k \rightarrow \infty$; (ii) $b_k := \max_{u \in Y_k, \|u\|_E=\rho_k} I(u) \leq 0$.*

Proof. It follows from (F2) that

$$|F(x, t)| \leq d_1 \left(\frac{1}{2} t^2 + \frac{1}{\tau} |t|^\tau \right), \quad \forall (x, t) \in \mathbb{R}^3 \times \mathbb{R}. \quad (10)$$

For any $k \in \mathbb{N}$ and $p \in [2, 6)$, we set

$$\beta_k(p) = \sup_{u \in Z_k, \|u\|_E=1} \|u\|_p.$$

Similar with Lemma 2.8 in [18], we have $\beta_k(p) \rightarrow 0$ as $k \rightarrow \infty$. Letting $r_k = \left(\frac{\tau}{8d_1\beta_k^\tau(\tau)} \right)^{\frac{1}{\tau-2}}$, for any $u \in Z_k$, it follows from (5), Lemma 2.1 and (10) that

$$\begin{aligned} I(u) &= \frac{1}{2} \int_{\mathbb{R}^3} (|\nabla u|^2 + V(x)u^2 - \omega\phi_u u^2) dx - \int_{\mathbb{R}^3} F(x, u) dx \\ &\geq \frac{1}{2} \|u\|_E^2 - d_1 \left(\frac{\beta_k^2(2)}{2} \|u\|_E^2 + \frac{\beta_k^\tau(\tau)}{\tau} \|u\|_E^\tau \right) \\ &\geq \frac{1}{8} \|u\|_E^2 \end{aligned}$$

for k large enough. Thus we obtain $a_k = \inf_{u \in Z_k, \|u\|_E=r_k} I(u) \geq \frac{1}{8} r_k^2 \rightarrow +\infty$ as $k \rightarrow \infty$.

Subsequently, for any $u \in Y_k$ and $\delta > 0$, set

$$\Gamma_\delta(u) = \{x \in \mathbb{R}^3 : |u| \geq \delta \|u\|_E\}.$$

Similar to Lemma 2.6 in [26], there exists $\varepsilon_1 > 0$ such that

$$meas(\Gamma_{\varepsilon_1}(u)) \geq \varepsilon_1.$$

From (F4), there exists $r_\infty > 0$ such that

$$F(x, u) \geq \frac{(1 + \omega^2 C_2^2)}{\varepsilon_1^3} |u|^2 \geq \frac{(1 + \omega^2 C_2^2)}{\varepsilon_1} \|u\|_E^2 \quad (11)$$

for all $u \in Y_k$ and $x \in \Gamma_{\varepsilon_1}(u)$ with $\|u\|_E \geq \frac{r_\infty}{\varepsilon_1}$. We can choose $\rho_k > \max\left\{\frac{r_\infty}{\varepsilon_1}, r_k\right\}$, then for any $u \in Y_k$ with $\|u\|_E = \rho_k$, it follows from (5), (F4), Lemma 2.1, (11) and (F5) that

$$\begin{aligned} I(u) &= \frac{1}{2} \int_{\mathbb{R}^3} (|\nabla u|^2 + V(x)u^2 - \omega \phi_u u^2) dx - \int_{\mathbb{R}^3} F(x, u) dx \\ &\leq \frac{1}{2} \|u\|_E^2 + \frac{\omega^2}{2} \|u\|_2^2 - \int_{\Gamma_{\varepsilon_1}(u)} F(x, u) dx \\ &\leq -\frac{(1 + \omega^2 C_2^2)}{2} \|u\|_E^2, \end{aligned}$$

which means $b_k \leq 0$ for ρ_k large enough. Then we finish the proof of this lemma. \square

Proof of Theorem 1.2. Similar to the proof of Theorem 1.1, we see that I satisfies the (PS) condition. Furthermore, $I(-u) = I(u)$, then we obtain our conclusion by using the Lemma 4.1.

5 Acknowledgments

The authors are grateful to the referees for the helpful comments which improve the writing of the paper. This paper was finished when D.-L. Wu was visiting Utah State University with the support of China Scholarship Council(No.201708515186); he is grateful to the members in the Department of Mathematics and Statistics at Utah State University for their invitation and hospitality.

References

- [1] A. Azzollini, A. Pomponio, Ground state solutions for the nonlinear Klein-Gordon-Maxwell equations, *Topol. Methods Nonlinear Anal.* 35 (1) (2010) 33–42.
- [2] A. Azzollini, L. Pisani, A. Pomponio, Improved estimates and a limit case for the electrostatic Klein-Gordon-Maxwell system, *Proc. Roy. Soc. Edinburgh Sect. A* 141 (3) (2011) 449-463.
- [3] V. Benci, D. Fortunato, The nonlinear Klein-Gordon equation coupled with the Maxwell equations, *Nonlinear Anal.* 47 (2001) 6065-6072.

- [4] T. Bartsch, Infinitely many solutions of a symmetric Dirichlet problem, *Nonlinear Anal.* 20 (10) (1993) 1205-1216.
- [5] T. Bartsch, Z.-Q. Wang, M. Willem, The Dirichlet problem for superlinear elliptic equations, in: *Stationary Partial Differential Equations. Vol. II*, in: *Handb. Differ. Equ.*, Elsevier/North-Holland, Amsterdam, 2005, pp. 1-55.
- [6] D. Cassani, Existence and non-existence of solitary waves for the critical Klein-Gordon equation coupled with Maxwell's equations, *Nonlinear Anal.* 58 (7-8) (2004) 733-747.
- [7] P. C. Carrião, P. L. Cunha, O. H. Miyagaki, Existence results for the Klein-Gordon-Maxwell equations in higher dimensions with critical exponents, *Commun. Pure Appl. Anal.* 10 (2) (2011) 709-718.
- [8] P. C. Carrião, P. L. Cunha, O. H. Miyagaki, Positive ground state solutions for the critical Klein-Gordon-Maxwell system with potentials, *Nonlinear Anal.* 75 (10) (2012) 4068-4078.
- [9] S.J. Chen, S.Z. Song, Multiple solutions for nonhomogeneous Klein-Gordon-Maxwell equations on \mathbb{R}^3 , *Nonlinear Analysis: Real World Applications* 22 (2015) 259-271.
- [10] S.-J. Chen, C.-L. Tang, Multiple solutions for nonhomogeneous Schrödinger-Maxwell and Klein-Gordon-Maxwell equations on \mathbb{R}^3 , *NoDEA Nonlinear Differential Equations Appl.* 17 (5) (2010) 559-574.
- [11] A. M. Candela, A. Salvatore, Multiple solitary waves for non-homogeneous Klein-Gordon-Maxwell equations, in: *More Progresses in Analysis Proceedings of the 5th International ISAAC Congress (Catania, 2005)*, 753-762, 2009.
- [12] S.T. Chen, X.H. Tang, Infinitely many solutions and least energy solutions for Klein-Gordon-Maxwell systems with general superlinear nonlinearity, *Comput. Math. Appl.* 75 (2018) 3358-3366.
- [13] T. D'Aprile, D. Mugnai, Solitary waves for nonlinear Klein-Gordon-Maxwell and Schrödinger-Maxwell equations, *Proc. Roy. Soc. Edinburgh Sect. A* 134 (5) (2004) 893-906.
- [14] T. D'Aprile, D. Mugnai, Non-existence results for the coupled Klein-Gordon-Maxwell equations, *Adv. Nonlinear Stud.* 4 (3) (2004) 307-322.
- [15] L. Ding, L. Li, Infinitely many standing wave solutions for the nonlinear Klein-Gordon-Maxwell system with sign-changing potential, *Comput. Math. Appl.* 68 (2014) 589-595.
- [16] P. d'Avenia, L. Pisani, Nonlinear Klein-Gordon equations coupled with Born-Infeld type equations, *Electron. J. Differential Equations* 26 (2002) 1-13.
- [17] X.M. He, Multiplicity of solutions for a nonlinear Klein-Gordon-Maxwell system, *Acta Appl. Math.* 130 (2014) 237-250.
- [18] L. Li, C.-L. Tang, Infinitely many solutions for a nonlinear Klein-Gordon-Maxwell system, *Nonlinear Anal.* 110 (2014) 157-169.

- [19] H.X. Shi, H.B. Chen, Multiple positive solutions for nonhomogeneous Klein-Gordon-Maxwell equations, *Appl. Math. Comput.* 337 (2018) 504-513.
- [20] K. Teng, K. Zhang, Existence of solitary wave solutions for the nonlinear Klein-Gordon equation coupled with Born-Infeld theory with critical Sobolev exponent, *Nonlinear Anal.* 74 (12) (2011) 4241–4251.
- [21] F.Z. Wang; Ground state solutions for the electrostatic nonlinear Klein-Gordon-Maxwell system, *Nonlinear Anal.* 74 (14) (2011), 4796-4803.
- [22] D.-L. Wu, F. Li, Solutions for fourth-order Kirchhoff type elliptic equations involving concave-convex nonlinearities in \mathbb{R}^N , *Comput. Math. Appl.* 79(2)(2020), 489-499.
- [23] D.-L. Wu, X. Yu, New homoclinic orbits for Hamiltonian systems with asymptotically quadratic growth at infinity, *Qual. Theory Dyn. Syst.* 19 (2020), 22.
- [24] L.X. Wang, S.J. Chen, Two solutions for nonhomogeneous Klein-Gordon-Maxwell system with sign-changing potential; *Elec. Diff. Equat.* 124 (2018) 1-21.
- [25] L.P. Xu, H.B. Chen, Existence and multiplicity of solutions for nonhomogenous Klein-Gordon-Maxwell equations, *Electron. J. Differential Equations* 102 (2015) 1-12.
- [26] Q.Y. Zhang, B. Xu, Multiplicity of solutions for a class of semilinear Schrödinger equations with sign-changing potential, *J. Math. Anal. Appl.* 377 (2011) 834-840.
- [27] J. Zhang, Solutions to the critical Klein-Gordon-Maxwell system with external potential, *J. Math. Anal. Appl.*, 455 (2017) 1152-1177.