

Circumbinary Planets – The Next Steps

David V. Martin^{1,2,3}

¹ *University of Chicago, Department of Astronomy & Astrophysics, 5640 S Ellis Ave, Chicago, IL-60637, USA*

² *The Ohio State University, Department of Astronomy, 100 W 18th Ave, Columbus, OH-43210, USA (E-mail: martin.4096@osu.edu)*

³ *Fellow of the Swiss National Science Foundation*

Abstract. The Kepler mission opened the door to a small but bonafide sample of circumbinary planets. Some initial trends have been identified and used to challenge our theories of planet and binary formation. However, the Kepler sample is not only small but contains biases. I will present a circumbinary plan for the future. Specifically, I will cover the BEBOP radial velocity survey, the latest TESS transit mission and a new technique for digging out small circumbinary planets in archival Kepler photometry.

Key words: exoplanets – circumbinary planets – transits – radial velocities

1. Introduction

Binary stars are common. Exoplanets are common. It is natural to seek planets in binaries. Planets in binary star systems come in two flavours: circumbinary planets on exterior orbits around tight binaries, and circumstellar planets on interior orbits around one of the two components of a wide binary. Here we will only consider circumbinary planets. The discovery of Kepler-16 (Doyle et al., 2011) really kicked off a search which had been anticipated since before the dawn of exoplanet discoveries (Borucki & Summers, 1984; Schneider & Chevreton, 1991). A dozen or so transiting circumbinary planets have been found by this mission (reviews in Welsh & Orosz 2018; Martin 2018), but this paper will look beyond the existing Kepler discoveries.

2. Trends And Open Questions In Circumbinary Planets

The dozen circumbinary planets discovered to date exhibit a few interesting trends and pose a few interesting questions.

1. ***There is a dearth of circumbinary planets orbiting the tightest eclipsing binaries (EBs).*** Most EBs have a very short period ($\sim 2 - 3$ days) but the transiting planets are only around > 7 day binaries (Fig. 1a). Muñoz & Lai (2015); Martin et al. (2015a); Hamers et al. (2016) explained this dearth by invoking a known formation mechanism of tight binaries under the influence of a misaligned third star and Kozai-Lidov cycles. The applicability of this story has been called into question lately by Moe & Kratter




Figure 1.: a) Histogram of Kepler eclipsing binary periods, compared with the periods of binaries known to host circumbinary planets, and the first TESS discovery of TOI-1338/EBLM J0608-59. b) Left: planet and binary periods, Right: planet periapse and binary apoapse. There is a tend for common ratios, which place the planets close to the stability boundary (Holman & Wiegert, 1999). c) Size and period of Kepler and TESS discoveries around both single and multiple stars. Figures reproduced from Kostov et al. (under review) and Martin (2018).

(2018), who deduce that Kozai-Lidov is only responsible for the minority of tight binary formation. More theoretical work is needed. Additionally, Muñoz & Lai (2015); Martin et al. (2015a); Hamers et al. (2016) suggested that any planets found orbiting around very tight binaries would be likely small and/or misaligned, both of which have been difficult to find to date.

2. ***There is an over-abundance of planets orbiting near the dynamical stability limit.*** This is likely the result of migratory formation of the planets, stalling near the edge of an inner disc cavity, which roughly coincides with the dynamical stability limit (Holman & Wiegert, 1999; Kley et al., 2019). Whilst this is not the sole result of an observational bias (Fig. 1b), more detections are needed to determine its statistical significance (Martin & Triaud, 2014; Li et al., 2016). In particular, finding circumbinary planets by radial velocities (Martin et al., 2019) or microlensing (Bennett et al., 2016) would allow for planet detections at longer periods, farther from the stability limit.
3. ***All transiting circumbinary planets are larger than $3R_{\oplus}$.*** This is contrary to the abundant discoveries of small planets around single stars, with a comparison shown in Fig. 1c. If it were a real absence, it would be enlightening, however the lack of small circumbinary planets is a detection bias; the days-amplitude transit timing variations (TTVs, Armstrong et al. 2013) inhibit traditional planet detection techniques based on phase-folding on a fixed period. Only transits of giant planets could be found so far, by eye. Some algorithms have been proposed to find small circumbinary planets, using modified versions of Boxed Least Squares (BLS, Ofir 2008) and the Quasiperiodic Automated Transit Search (QATS, Windemuth et al. 2019b), but no new candidates have been reported yet.

3. A Search For Small Transiting Circumbinary Planets In Kepler

The archival Kepler data remains the best source for finding small circumbinary planets, because of its long four-year baseline, high-precision photometry and well-characterised EB catalog (Prša et al., 2011; Windemuth et al., 2019b). In collaboration with Dan Fabrycky, a new transit search algorithm is being specifically designed for shallow transits of small circumbinary planets. It can successfully recover all known circumbinary planets, and also injected planets slightly smaller than Earth (Fig 2). Planet detection is assisted by a detrending algorithm designed specific to EBs, which accounts for the variable length of circumbinary planet transits as a function of the binary phase. Unique to this transit detection algorithm is building TTVs directly and exactly into the search. For each set of orbital parameters the algorithm produces a quasi-periodic mask of transit times and durations using a rapid N-body algorithm. This mask is

matched to the photometric data similar to the cross correlation technique for high-precision RV fits to spectroscopic data. The N-body-derived mask fully incorporates the three-body geometry and both short and long-term dynamical variations of the planet’s orbit. The search grid has been optimised using principles similar to (Ofir, 2014), but adapted to circumbinary planets.

Roughly two dozen detections are expected if planets have a similar size distribution around one and two stars (preliminary research suggests this is the case for gas giants, Martin & Triaud 2014; Armstrong et al. 2014). Alternatively, it is possible that small circumbinary planets *are rare or non-existent*. This would suggest that super-Earths form in situ rather than with significant migration, helping answer a hotly-debated topic (Ogihara et al., 2015); around single stars such a process is possible but around a binary it would be suppressed (Paardekooper et al., 2012).




Figure 2.: Recovery of an injected 260-day $0.875R_{\oplus}$ circumbinary planet on Kepler-16 (real planet is over $8R_{\oplus}$ with a 228 day period) with the new automated algorithm. Injected transits were created using *BATMAN* (Kreidberg, 2015), with the duration scaled according to the relative planet-star velocity calculated by the *REBOUND* N-body algorithm (Rein & Liu, 2012)

4. The BEBOP radial velocity survey

Between 2013 and 2018 a blind survey for circumbinary planets was run on the Swiss Euler Telescope. It was given the delightful name BEBOP - “Binaries Escorted By Orbiting Planets”. BEBOP uniquely targetted eclipsing, single-lined spectroscopic binaries. The eclipses add preferential biases in both radial velocity amplitude and transit probability (Martin & Triaud, 2015b; Martin, 2017). The single-lined binaries, composed of F/G primaries and M-dwarf secondaries, avoid the difficult problem of spectral contamination, and the need to deconvolve two moving sets of spectral lines. This is different to the SB2 search of TATOOINE (Konacki et al., 2009).

Over 1000 observations taken over more than 60 nights were compiled in Martin et al. (2019). The survey was sensitive down to $0.5M_{\text{Jup}}$, but our lack of detections showed that circumbinary planets are typically sub-Saturn mass (Fig. 3). BEBOP was sensitive to planetary mass companions at periods of several years, much longer than the Kepler discoveries. BEBOP also demonstrated that there was not a large abundance of giant, misaligned planets, which were proposed by Martin & Triaud (2014); Armstrong et al. (2014) as compatible with the Kepler transit results. BEBOP has since been expanded to large programs on HARPS, SOPHIE and ESPRESSO.




Figure 3.: BEBOP detection completeness, detected triple star systems (green circles), known transiting circumbinary planets with roughly-characterised masses (upwards blue triangles) and a known circumbinary brown dwarf (downwards blue triangle). Numbers in white boxes indicate 95% confidence abundance bounds. Figure reproduced from Martin et al. (2019).




Figure 4.: Three primary transits of the circumbinary planet TOI-1338/EBLM J0608-59 and the photodynamical fit with its residuals (observed minus calculated). The variable transit duration, owing to a variable relative velocity between the star and planet, is a smoking-gun signature of a circumbinary planet. Figure reproduced from Kostov et al. (under review).

5. The TESS transit mission

TESS presents different challenges and opportunities when compared with Kepler. TESS is observing most of the sky, in both hemispheres, and hence is targeting many more bright stars so ground-based follow-up is significantly easier. However, two drawbacks are the smaller telescope size (10.5 cm compared with 95 cm) and shorter observing timespans (30 days for most of the TESS field). Only near the ecliptic poles does the TESS timespan increase to almost a year of continuous viewing, owing to the overlap of multiple sectors. Indeed, the single TESS planet found so far is near the continuous viewing zone: TOI-1338/EBLM J0608-59 (Kostov et al. under review, Fig. 4). The planet has very similar properties to the Kepler population of planets (it is highlighted in Fig 1).

A unique aspect of this discovery, compared with the Kepler discoveries, is that the binary was already known and well characterised as a part of the EBLM (Triaud et al., 2017) and BEBOP (Martin et al., 2019) radial velocity surveys, and those measurements were vital to the planet’s characterisation.

TESS is unlikely to significantly break into new parameter spaces of circumbinary planets, due to the shortened observational timespans and inferior photometric precision to Kepler. Most detections will be harder than TOI-1338/EBLM J0608-59. What TESS will hopefully provide though is a significant increase in the statistics of circumbinary planets. The TESS circumbinary planet working group predicts 140 TESS circumbinary planets if we can detect them on a single passing that transits both stars, a “1-2 punch”. This is based on 400,000 eclipsing binaries, a Kepler-like circumbinary planet detection rate of 11/2800, a 30/180 chance of the median circumbinary period transiting during a one month window and a 1/2 chance of the planet transiting both stars (Martin, 2017). Based on Kepler, ~ 50 of these planets are expected to be in the habitable zone.

References

- Armstrong, D., Martin, D. V., Brown, G., et al., Placing limits on the transit timing variations of circumbinary exoplanets. 2013, *Mon. Not. R. Astron. Soc.*, **434**, 3047, DOI: 10.1093/mnras/stt1226
- Armstrong, D. J., Osborn, H. P., Brown, D. J. A., et al., On the abundance of circumbinary planets. 2014, *Mon. Not. R. Astron. Soc.*, **444**, 1873, DOI: 10.1093/mnras/stu1570
- Bennett, D. P., Rhie, S. H., Udalski, A., et al., The First Circumbinary Planet Found by Microlensing: OGLE-2007-BLG-349L(AB)c. 2016, *Astron. J.*, **152**, 125, DOI: 10.3847/0004-6256/152/5/125
- Borucki, W. J. & Summers, A. L., The photometric method of detecting other planetary systems. 1984, *Icarus*, **58**, 121, DOI: 10.1016/0019-1035(84)90102-7
- Doyle, L. R., Carter, J. A., Fabrycky, D. C., et al., Kepler-16: A Transiting Circumbinary Planet. 2011, *Science*, **333**, 1602, DOI: 10.1126/science.1210923
- Hamers, A. S., Perets, H. B., & Portegies Zwart, S. F., A triple origin for the lack of tight coplanar circumbinary planets around short-period binaries. 2016, *Mon. Not. R. Astron. Soc.*, **455**, 3180, DOI: 10.1093/mnras/stv2447
- Holman, M. J. & Wiegert, P. A., Long-Term Stability of Planets in Binary Systems. 1999, *Astron. J.*, **117**, 621, DOI: 10.1086/300695
- Kley, W., Thun, D., & Penzlin, A. B. T., Circumbinary discs with radiative cooling and embedded planets. 2019, *Astron. Astrophys.*, **627**, A91, DOI: 10.1051/0004-6361/201935503

- Konacki, M., Muterspaugh, M. W., Kulkarni, S. R., & Hełminiak, K. G., The Radial Velocity Tatooine Search for Circumbinary Planets: Planet Detection Limits for a Sample of Double-Lined Binary Stars. 2009, *Astrophys. J.*, **704**, 513, DOI: 10.1088/0004-637X/704/1/513
- Kreidberg, L., batman: BASic Transit Model cAlculatioN in Python. 2015, *Publications of the Astronomical Society of the Pacific*, **127**, 11611165, DOI: 10.1086/683602
- Li, G., Holman, M. J., & Tao, M., Uncovering Circumbinary Planetary Architectural Properties from Selection Biases. 2016, *Astrophys. J.*, **831**, 96, DOI: 10.3847/0004-637X/831/1/96
- Martin, D. V., Circumbinary planets - II. When transits come and go. 2017, *Mon. Not. R. Astron. Soc.*, **465**, 3235, DOI: 10.1093/mnras/stw2851
- Martin, D. V. 2018, *Populations of Planets in Multiple Star Systems*, 156
- Martin, D. V., Mazeh, T., & Fabrycky, D. C., No circumbinary planets transiting the tightest Kepler binaries - a possible fingerprint of a third star. 2015a, *Mon. Not. R. Astron. Soc.*, **453**, 3554, DOI: 10.1093/mnras/stv1870
- Martin, D. V. & Triaud, A. H. M. J., Planets transiting non-eclipsing binaries. 2014, *Astron. Astrophys.*, **570**, A91, DOI: 10.1051/0004-6361/201323112
- Martin, D. V. & Triaud, A. H. M. J., Circumbinary planets - why they are so likely to transit. 2015b, *Mon. Not. R. Astron. Soc.*, **449**, 781, DOI: 10.1093/mnras/stv121
- Martin, D. V., Triaud, A. H. M. J., Udry, S., et al., The BEBOP radial-velocity survey for circumbinary planets. I. Eight years of CORALIE observations of 47 single-line eclipsing binaries and abundance constraints on the masses of circumbinary planets. 2019, *Astron. Astrophys.*, **624**, A68, DOI: 10.1051/0004-6361/201833669
- Moe, M. & Kratter, K. M., Dynamical Formation of Close Binaries during the Pre-main-sequence Phase. 2018, *Astrophys. J.*, **854**, 44, DOI: 10.3847/1538-4357/aaa6d2
- Muñoz, D. J. & Lai, D., Survival of planets around shrinking stellar binaries. 2015, *Proceedings of the National Academy of Science*, **112**, 9264, DOI: 10.1073/pnas.1505671112
- Ofir, A., An algorithm for photometric identification of transiting circumbinary planets. 2008, *Mon. Not. R. Astron. Soc.*, **387**, 1597, DOI: 10.1111/j.1365-2966.2008.13336.x
- Ofir, A., Optimizing the search for transiting planets in long time series. 2014, *Astron. Astrophys.*, **561**, A138, DOI: 10.1051/0004-6361/201220860

- Ogihara, M., Morbidelli, A., & Guillot, T., A reassessment of the in situ formation of close-in super-Earths. 2015, *Astron. Astrophys.*, **578**, A36, DOI: 10.1051/0004-6361/201525884
- Paardekooper, S.-J., Leinhardt, Z. M., Thébault, P., & Baruteau, C., How Not to Build Tatooine: The Difficulty of In Situ Formation of Circumbinary Planets Kepler 16b, Kepler 34b, and Kepler 35b. 2012, *Astrophys. J., Lett.*, **754**, L16, DOI: 10.1088/2041-8205/754/1/L16
- Prša, A., Batalha, N., Slawson, R. W., et al., Kepler Eclipsing Binary Stars. I. Catalog and Principal Characterization of 1879 Eclipsing Binaries in the First Data Release. 2011, *Astron. J.*, **141**, 83, DOI: 10.1088/0004-6256/141/3/83
- Rein, H. & Liu, S. F., REBOUND: an open-source multi-purpose N-body code for collisional dynamics. 2012, *Astron. Astrophys.*, **537**, A128, DOI: 10.1051/0004-6361/201118085
- Schneider, J. & Chevreton, M. *A Proposal for the Search for Extra-Solar Planets by Occultation*, ed. J. Heidmann & M. J. Klein, , Vol. **390**, 33–33
- Triaud, A. H. M. J., Martin, D. V., Ségransan, D., et al., The EBLM Project. IV. Spectroscopic orbits of over 100 eclipsing M dwarfs masquerading as transiting hot Jupiters. 2017, *Astron. Astrophys.*, **608**, A129, DOI: 10.1051/0004-6361/201730993
- Welsh, W. F. & Orosz, J. A. 2018, *Two Suns in the Sky: The Kepler Circumbinary Planets*, 34
- Windemuth, D., Agol, E., Carter, J., et al., An automated method to detect transiting circumbinary planets. 2019b, *Mon. Not. R. Astron. Soc.*, **490**, 1313, DOI: 10.1093/mnras/stz2637