

Integer Plane Multiflow Maximisation: Flow-Cut Gap and One-Quarter-Approximation

Naveen Garg, Nikhil Kumar, András Sebő

Indian Institute of Technology, Delhi

naveen@cse.iitd.ac.in

nikhil@cse.iitd.ac.in

CNRS, Laboratoire G-SCOP, Univ. Grenoble Alpes

andras.sebo@grenoble-inp.fr

Abstract. In this paper, we bound the integrality gap and the approximation ratio for maximum plane multiflow problems and deduce bounds on the flow-cut-gap. Planarity means here that the union of the supply and demand graph is planar. We first prove that there exists a multiflow of value at least half of the capacity of a minimum multicut. We then show how to convert any multiflow into a half-integer one of value at least half of the original multiflow. Finally, we round any half-integer multiflow into an integer multiflow, losing again at most half of the value, in polynomial time, achieving a $1/4$ -approximation algorithm for maximum integer multiflows in the plane, and an integer-flow-cut gap of 8.

Keywords: Multicommodity Flow · multiflow · multicut · Network Design · Planar Graphs · flow-cut, integrality gap · approximation algorithm.

1 Introduction

Given an undirected graph $G = (V, E)$ with edge capacities $c : E \rightarrow \mathbb{R}_+$, and some pairs of vertices given as edges of the graph $H = (V, F)$, the *maximum-multiflow problem* with input (G, H, c) , asks for the maximum flow that can be routed in G , simultaneously between the endpoints of edges in F , respecting the capacities c .

This is one of many widely studied variants of the multiflow problem. Other popular variants include demand flows, all or nothing flows, unsplittable flows etc. In this paper, we are mainly interested in the integer version of this problem, and the half-integer or fractional versions also occur as tools. When capacities are 1, the capacity constraint specialises to edge-disjointness, whence the *maximum edge disjoint paths problem* (MEDP) between a given set of pairs is a special case; MEDP is NP-Hard to compute for general graphs, even in very restricted settings like when G is a tree [10].

The edges in F are called *demand edges* (sometimes commodities), those in E are called *supply edges*; accordingly, $H = (V, F)$ is the *demand graph*, and $G = (V, E)$ is the *supply graph*. If $G + H = (V, E \cup F)$ is planar we call the problem a

plane multiflow problem. Plane multiflows have been studied for the past forty years, starting with Seymour [25].

Let $\mathcal{P}_e$ ($e \in E$) be the set of paths in G between the endpoints of e , and $\mathcal{P} := \cup_{e \in E} \mathcal{P}_e$. For $P \in \mathcal{P}_e$, the edge e is said to be the *demand-edge* of P , denoted by e_P . A *multiflow*, or for simplicity a *flow* in this paper is a function $f : \mathcal{P} \rightarrow \mathbb{R}^+$. The flow f is called *feasible*, if $\sum_{\{P \in \mathcal{P} : e \in P\}} f(P) \leq c(e)$ for all $e \in E$. The *value* of a flow f is defined as $|f| := \sum_{P \in \mathcal{P}} f(P)$.

For a path $P \in \mathcal{P}$, we refer to $f(P)$ as the *flow on P* . If the flow on every path is integer or half-integer, we say that the flow is integer or half-integer, respectively. A *circuit* is a connected subgraph with all degrees equal to two.

The value of multiflows (without restrictions on integrality) can be maximised in general, in (strongly) polynomial time [23, 70.6, page 1225] using a linear programming algorithm.

A *multicut* for (G, H) is a set of edges $M \subseteq E$ such that every $P \in \mathcal{P}$ contains at least one edge in M .¹ Multicuts provide the simplest possible and most natural constraints for the dual of the maximum multiflow problem.

It is easy to see that the value of any feasible multiflow is smaller than or equal to the capacity of any multicut. Klein, Mathieu and Zhou [14] prove that computing the minimum multicut is NP-hard if $G + H$ is planar, and they also provide a PTAS.

There is a rich literature on the maximum ratio of a minimum capacity of a multicut over the maximum multiflow. With an abuse of terminology we will call this the (*possibly integer or half-integer*) *flow-cut gap*, sometimes also restricted to subclasses of problems. This is not to be confused with the same term used for demand problems. The integer flow-cut gap is 1 when G is a path and 2 when G is a tree. For arbitrary (G, H) , the flow-cut gap is $\theta(\log |F|)$ [9]. Building on decomposition theorems from Klein, Plotkin and Rao [13], Tardos and Vazirani [26] showed a flow-cut gap of $O(r^3)$ for graphs which do not contain a $K_{r,r}$ minor; note that for $r = 3$ this includes the class of planar graphs. A long line of impressive work, culminated in [24] proving a constant approximation ratio for maximum half-integer flows, which together with [26] implies a constant half-integer flow-cut gap for planar supply graphs. A simple topological obstruction proves that the integer flow-cut gap for planar supply graphs, even when all demand edges are on one face of the graph, also called *Okamura-Seymour instances*, is $\Omega(|F|)$ [10].

These results are often bounded by the *integrality gap* of the multiflow problems for the problem classes, which is the infimum of $\text{MAX}/|f|$ over all instances (G, H, c) of the problem class, and where MAX is the maximum value of an integer multiflow, and f is a multiflow for this same input, and the *half-integrality gap* is defined similarly by replacing “integer” by “half-integer” in the nominator. A ρ -approximation algorithm ($\rho \in \mathbb{R}$) for a maximisation problem is a polynomial algorithm which outputs a solution of value at least ρ times

¹ Given a partition of V so that all demand edges join different classes, the supply edges that join different classes form a multicut, and inclusionwise minimal multicuts are of this form.

the optimum; ρ is also called the *approximation ratio* (or guarantee). All these gaps and ratios are defined analogously for minimisation problems.

Our first result (Section 3, Theorem 1) is an upper bound of 2 for the flow-cut gap (i.e. multicut/multiflow ratio) for plane instances, the missing relation we mentioned. We prove this by relating multicuts to 2-edge-connectivity-augmentation in the plane-dual, and applying a bound of Williamson, Goemans, Mihail and Vazirani [28] for this problem.

We next show (Section 4, Theorem 2) how to obtain a half-integer flow from a given (fractional) flow in plane instances, reducing the problem to a linear program with a particular combinatorial structure, and solving it.

Finally (Section 5, Theorem 4), given any feasible half-integer flow, we show how to extract an integer flow of value at least half of the original, in polynomial time, using an algorithm that 4-colors planar graphs efficiently [21].

These results imply an integrality gap of $1/2$ for maximum half-integer flows, $1/4$ for maximum integer flows, and the same approximation ratios for each. In turn, the flow-cut gap of 2 implies a half-integer flow-cut gap of 4 and an integer flow-cut gap of 8 for plane instances.

Section 6 presents some examples providing lower bounds for multicuts and upper bounds for multiflows. A summary of the results completed by lower bounds and open problems are stated in Section 7. The Appendix is a guide to some of the tools.

These are the first constant approximation ratios and gaps proved for the studied problems.² Approximately at the same time as our submission to IPCO21, a manuscript was published on Arxiv, proving constant ratios for the same problems [12]. The bounds for the gaps and the approximation ratio of [12] are less sharp, but an interesting new rounding method, and a new complexity result appear in it. The latter will occur in the following section.

2 Preliminaries

We detail here some notions, notations, terminology, facts and tools we use, including some preceding results of influence.

Demands, the Cut Condition and Plane Duality

We describe the notion of demand flows, which is closely related to multiflows defined in the previous section. The problem is defined by the quadruple (G, H, c, d) , where G, H, c are as before, *demands* $d : F \rightarrow \mathbb{Z}_+$ are given, and we are looking for a feasible (sometimes in addition integer or half-integer) flow f satisfying $\sum_{P \in \mathcal{P}: e \in P} f(P) = d(e)$ for all $e \in F$.

A *cut* in a graph $G = (V, E)$ is a set of edges of the form $\delta(S) = \delta_E(S) := \{e \in E : e \text{ has exactly one endpoint in } S\}$ for all $S \subseteq V$. Note that $\delta(S) = \delta(V \setminus S)$; $S, V \setminus S$ are called the *shores* of the cut. For a subset $E' \subseteq E$ we use the notation $c(E') := \sum_{e \in E'} c(e)$, and we adopt this usual way of extending a

² A shortened version of this manuscript has been submitted to IPCO21 in November 2019 and will appear in the volume of the conference (LNCS, Springer) in June 2020.

function on single elements to subsets. For instance, $d(F') := \sum_{e \in F'} d(e)$ is the *demand* of the set $F' \subset F$.

A necessary condition for the existence of a feasible multiflow is the so called *Cut Condition*: for every $S \subseteq V$, $c(\delta_E(S)) \geq d(\delta_F(S))$, that is, the capacity of each cut must be at least as large as its demand. The cut condition is not sufficient for a flow in general, but Seymour [25] showed that it is sufficient for an integer flow, provided $G + H$ is planar, the capacities and demands are integer, and their sum is even on the edges incident to any vertex. A half-integer flow follows then for arbitrary integer capacities for such *plane* instances (see Section 1). The same holds for Okamura-Seymour instances [20]. There are more examples, unrelated to planarity, where the cut condition is sufficient to satisfy all demands, and with an integer flow, for instance when all demand edges can be covered by at most two vertices.

Seymour's theorem [25] on the sufficiency of the cut condition and about the existence of integer flows has promoted plane multiflow problems to become one of the targets of investigations. **This paper is devoted to plane multiflow maximisation.** Seymour's proof is based on a nice correspondence to other combinatorial problems through plane duality, that we also adopt.

Middendorf and Pfeiffer [19] showed that the plane multiflow problem (already deciding the existence of edge disjoint paths) is NP-Hard, which immediately implies that *plane multiflow maximisation is also NP-hard* (see Footnote ??). The cut condition can be checked in polynomial time³ so the difficult question to decide is the existence or not of an integer flow when the cut condition is satisfied. As far as multicuts are concerned, we show that their minimisation in plane instances is equivalent to a 2-edge-connectivity augmentation problem in planar graphs.

Following Schrijver [23, p. 27] we denote the dual of the planar graph G by $G^* = (V^*, E^*)$, where V^* is the set of faces of G and each edge $e \in E$ corresponds to an edge $e^* \in E^*$ joining the two faces that share e . For $X \subseteq E$ denote $X^* := \{e^* : e \in X\}$. An important fact we need and use about dualisation: *C is an inclusionwise minimal cut in G if and only if C^* is a circuit in G^* .*

It is tempting to reduce multiflow maximisation to demand flows, using the sufficiency of the cut condition. The plane dual of the question is to maximise the sum of the absolute values of weights (demands) on the negative edges so that there is no negative cycle in the graph. An integer solution to this problem does not imply an integer multiflow, cf. $(C_4, 2K_2)$ ⁴. The complexity of finding such an integer solution was open, has been proved recently in [2, Theorem 2]

³ Seymour's correspondence through dualisation reduces this problem to checking whether F^* is a minimum weight " T_{F^*} -join" (see eg. [23]) in $(G + H)^*$ with weights defined by c and d , where T_{F^*} is the set of odd degree vertices of F^* . This also provides a polynomial separation algorithm for maximising the sum of (not necessarily integer) demands satisfying the cut condition.

⁴ This notation means $G = C_4$ and $H = 2K_2$, where $C_4 = (V, E)$ is the graph consisting of a circuit on V , $|V| = 4$, and $2K_2 = (V, F)$ is its complement, so $G + H = K_4$. The capacities and the demands are defined to be 1. To transform this example to multiflow maximisation with integrality gap $1/2$ we keep the capacities, and wish to

to be NP-hard when $G + H$ is planar after deleting a vertex, [12] proves this when $G + H$ is planar, but it remains open when H consists of one tree and isolated vertices; then, in addition, [15] provides an integer solution whenever the cut condition is satisfied.

Two Lemmas on Laminar Families

This correspondence by plane duality allows to transform any fact on cuts to circuits in the dual and vice versa. For example fractional, half-integer or integer packings of cuts in G^* , where each cut contains exactly one edge of F^* correspond to a fractional, half-integer or integer multiflow in G . We provide now some related definitions, notations and facts. We do this directly on the graph $(G + H)^*$ where they will be used; we denote by V^* the vertices of this graph, that is the faces of $G + H$. So $(G + H)^* = (V^*, E^* \cup F^*)$.

Let $\delta(A) = \delta_{E^* \cup F^*}(A)$, $\delta(B) = \delta_{E^* \cup F^*}(B)$ ($A, B \subseteq V^*$) be two *crossing* cuts, i.e. A and B are neither disjoint nor contain one another, each of which contains exactly one edge of F^* . They can be replaced by $\delta(A^* \cap B^*)$ and $\delta(A^* \cup B^*)$ or $\delta(A^* \setminus B^*)$ and $\delta(B^* \setminus A^*)$ (in the plane dual). It is easy to check that every edge is contained in at most as many cuts after the replacement as before, and if both cuts contain exactly one edge of F^* then this also holds for the replacing cuts. Doing this iteratively and using plane duality, we can convert any feasible flow into another one (without changing the value of the flow) in which no two flow paths cross. Lemma 1 below formalises exactly what we need.

A family of subsets of V^* is called *laminar* if any two of its members are disjoint or one of them contains the other. If for any two members one of them contains the other we say that the family is a *chain*. Given a laminar family $\mathcal{L} \subseteq 2^{V^*}$, a chain $\mathcal{C} \subseteq \mathcal{L}$ is *full* (in $\mathcal{L}$) if $X, Y, Z \in \mathcal{L}$, $X \subseteq Y \subseteq Z$ and $X, Z \in \mathcal{C}$ implies $Y \in \mathcal{C}$. We call a multiflow f *laminar*, if $\{C^* : C = P \cup \{e_P\}, P \in \mathcal{P}, f(P) > 0\} = \{\delta_{E^* \cup F^*}(L) : L \in \mathcal{L}\}$, where $\mathcal{L} \subseteq 2^{V^*}$ is laminar. We state the following well-known Lemma without proof (see the Appendix for references and explanations):

Lemma 1. *For every feasible multiflow f there exists a laminar feasible multiflow f' so that $|f'| = |f|$, and f' can be found in polynomial time.*

The following useful properties are easy to check:

For a family $\mathcal{L}$ of subsets of V^* and $a \neq b \in V^*$, denote $\mathcal{L}(a) := \{L \in \mathcal{L} : a \in L\}$; $\mathcal{L}(a, b) := \{L \in \mathcal{L} : a \in L, b \notin L\}$.

Lemma 2. *Let $\mathcal{L}$ be a laminar family of subsets of V^* . Then*

- a. $|\mathcal{L}| \leq 2(|V^*| - 1) = O(|V|)$.
- b. $\mathcal{L}(a, b) \subseteq \mathcal{L}(a) \subseteq V^*$ both form full chains of subsets of V^* . □

put an upper bound on the demands, otherwise there is an integer flow of value 2. Such an upper bound can be realized by replacing each demand edge by two edges in series, a demand and a supply edge, the latter of capacity 1, getting from $(C_4, 2K_2)$ the MEDP for $(\overline{C_4}, \overline{2K_2})$, with integrality gap $1/2$. This is in general a useful transformation from demand problems to maximisation, for which we will use the same “overline” notation. This is at the same time a reduction of the NP-hardness of plane multiflow feasibility, known to be NP-hard [19] to plane multiflow maximisation.

Integrality in demand flows and stable sets

We first compute a half integer flow of value at least half the fractional flow and then convert this into an integer flow. We now describe an instance which illustrates the difficulty in finding an integer flow. Consider a planar graph with a perfect matching without any nontrivial *tight cut*⁵ to be $(G + H)^*$, and F^* to be any perfect matching in it. Let all capacities be 1. In order to have an upper bound of 1 on each demand, apply the transformation of Footnote 4. Then multiflows can use only the dual edge-sets of stars of vertices in $(G + H)^*$, so an integer multiflow of value k corresponds exactly to a *stable set*, that is, a set of vertices not inducing any edge, of size k in $(G + H)^*$. This indicates that in order to find a large integer flow, we sometimes need to find large stable sets in planar graphs.

Despite the restriction of the paths to those arising from duals of stars of vertices, the gap between the integer and the half-integer max-flow is at least $1/2$ for these bricks: it follows from the 4-color theorem [1] that a stable set of size at least $|V^*|/4$ exists while the maximum half integer flow cannot exceed $|F| = |V^*|/2$. We will be able to reach this ratio in general (see Section 5, Theorem 4), and $(\overline{C_4}, 2\overline{K_2})$, defined as the graph in Footnote 4 but instead of shows that this cannot be improved.

In order to reach this integrality gap of 2 in general, we will actually need to find a stable-set of size $n/4$ in a planar auxiliary graph of n vertices. The maximum stable set problem is NP-hard, but there is a PTAS for it in planar graphs [3], which, combined with the 4-color theorem [1] provides a stable-set of size $n/4$; an alternative is to use the 4-coloring algorithm of Robertson, Sanders, Seymour and Thomas [21] which directly provides a 4-coloring of a planar graph in polynomial time, and the largest color class is clearly of size at least $n/4$. Either of these algorithms can be used as a black-box-tool for rounding half-integer flows, so we state the result:

Lemma 3. *In a planar graph on n vertices, a stable set of size $n/4$ can be found in polynomial time.*

A similar rounding argument appeared in the work of Fiorini, Hardy, Reed and Vetta [6] in the somewhat different context of proving an upper bound of Král and Voss [17] for the ratio between “minimum size of an odd cycle edge transversal” versus the “maximum odd cycle edge packing” using the 4-color theorem [1]. Our procedure in Section 5, occurs to be more general in that it is starting from an *arbitrary, not necessary optimal, half-integer multiflow* for an arbitrary capacity function, and constructs an integer multiflow, with a polynomial algorithm. Similar difficulties approached independently with the 4-color theorem confirm that it may be unavoidable for plane multiflows.

⁵ i.e. a cut with both shores containing more than one vertex, and meeting every perfect matching in exactly one edge. Lovász characterised graphs without nontrivial tight cuts as “bicritical 3-connected graphs” [18] called *bricks*. Such graphs may have arbitrarily many vertices, even under the planarity constraint.

3 Multicuts versus Multiflows via 2-Connectors

We show in this section that the flow-cut gap is at most two for plane instances, via a reduction to the 2-edge-connectivity augmentation problem.

Given $G = (V, E)$, $H = (V, F)$, a 2-connector for H in G is a set of edges $Q \subseteq E$ such that none of the edges $e \in F$ is a cut edge of $(V, Q \cup F)$; equivalently, Q is a 2-connector if and only if each $e \in F$ is contained in a circuit of $Q \cup F$. The 2-edge-connectivity Augmentation Problem (2ECAP) is to find, for given edge costs $c : E \rightarrow \mathbb{Z}_+$ on E , a minimum cost 2-connector.

Let (G, H, c) , $G = (V, E)$, $H = (V, F)$ be the input of a plane multiflow maximisation problem, and $(G + H)^* = (V^*, E^* \cup F^*)$, where V^* is the set of faces of $G + H$. Define $c(e^*) := c(e)$ ($e \in E$).

Lemma 4. *The edge-set $Q \subseteq E$ is a multicut for (G, H) if and only if Q^* is a 2-connector for (V^*, F^*) in (V^*, E^*) .*

Proof. The edge-set $Q \subseteq E$ forms a multicut in G if and only if the endpoints u, v of any edge $e \in F$ are in different components of $(V, E \setminus Q)$, that is, if and only if for all $e \in F$ there exists an inclusionwise minimal cut $C \subseteq Q \cup F$ of $G + H$ such that $e \in C$. But we saw among the preliminaries concerning duality that C is an inclusionwise minimal cut in $G + H$ if and only if C^* is a circuit in $Q^* \cup F^*$. Summarising, $Q \subseteq E$ forms a multicut in G , if and only if for all $e^* \in F^*$ there exists a circuit C^* in $Q^* \cup F^*$ such that $e^* \in C^*$. This means exactly that $Q^* \subseteq E^*$ is a 2-connector for (V^*, F^*) , in (V^*, E^*) , finishing the proof. $\square$

Let (G, H, c) be an instance of multiflow problem with $G + H$ planar. Let $p : 2^{V^*} \rightarrow \{0, 1\}$ with $p(S) = 1$ if and only if $|\delta(S) \cap F^*| = 1$, otherwise $p(S) = 0$ ($S \subseteq V^*$). The following linear program specialises the one investigated in [28]:

$$\begin{aligned} & \text{minimise} && \sum_{e \in E^*} c(e)x(e), \\ & \text{subject to} && \\ & \sum_{e \in \delta(S)} x(e) \geq p(S), && S \subseteq V^*; \\ & x(e) \geq 0 && e \in E^*. \end{aligned} \quad (2ECAP)$$

Since p is $\{0, 1\}$ -valued so are the coordinatewise minimal integer solutions including the integer optima of (2ECAP). *The $\{0, 1\}$ -solutions are exactly the (incidence vectors of) 2-connectors of (V^*, F^*) in (V^*, E^*) .* In the dual linear program of (2ECAP), we have a variable $y(S)$ for all $S \subseteq V^*$, constraints $\sum_{S: e \in \delta(S)} y(S) \leq c_e$ for all $e \in E^*$ and $y(S) \geq 0$ for all $S \subseteq V^*$. The objective is to maximise $\sum_{S \subseteq V^*} p(S)y(S)$. Williamson, Goemans, Mihail and Vazirani [28] developed a primal-dual algorithm finding for given input $(G + H)^*$ and c , an integer primal solution x_{WGMV} to a class of linear programs including (2ECAP), together with a (not necessarily integer) dual solution y_{WGMV} , in polynomial time, proving the following WGMV inequality [28, Lemma 2.1], see also [16, Section 20.4]:

$$\text{LIN} \leq \text{OPT} \leq \sum_{e \in E^*} c(e)x_{\text{WGMV}}(e) \leq 2 \sum_{S \subseteq V^*, p(S)=1} y_{\text{WGMV}}(S) \leq 2 \text{LIN} \leq 2 \text{OPT},$$

where OPT is the minimum cost of a 2-connector, and LIN is the optimum of (2ECAP). We will refer to this algorithm as the *WGMV algorithm*.

Note that the WGMV algorithm works for the class of weakly supermodular functions. A function $h : 2^V \rightarrow \{0, 1\}$ is called *weakly supermodular* if $h(V) = 0$ and for any $A, B \subseteq V$, $h(A) + h(B) \leq \max\{h(A \cap B) + h(A \cup B), h(A \setminus B) + h(B \setminus A)\}$. It can be verified that p defined above is weakly supermodular.

Theorem 1. *Let (G, H, c) be a plane multiflow problem. Then there exists a feasible multiflow f and a multicut Q , such that $c(Q) \leq 2|f|$, where both f and Q can be computed in polynomial time.*

Proof. Recall that the WGMV algorithm finds x_{WGMV} and y_{WGMV} satisfying the WGMV inequality, where x_{WGMV} is the incidence vector of a 2-connector of (V^*, F^*) in (V^*, E^*) , let us denote its plane dual set in $G + H$ by Q_{WGMV} . By Lemma 4 Q_{WGMV} is a multicut.

Consider a set S with $p(S) = 1$, that is, $|\delta(S) \cap F^*| = |\delta(S)^* \cap F| = 1$, all the other sets can be supposed to be absent from the inequalities. Then $\delta(S)^*$ contains a circuit C of $G + H$ containing the unique edge of $|\delta(S)^* \cap F|$. Therefore $C \setminus F$ is a path in G , denote it by P_S . Define a multiflow f in $G + H$ by $f(P_S) = y_{\text{WGMV}}(S)$. The dual feasibility of y_{WGMV} means exactly that the multiflow f is feasible. By our construction and the WGMV inequality we have

$$c(Q_{\text{WGMV}}) = \sum_{e \in E^*} c(e)x_{\text{WGMV}}(e) \leq 2 \sum_{S \subseteq V^*, p(S)=1} y_{\text{WGMV}}(S) = 2|f|.$$

So the multicut $Q := Q_{\text{WGMV}}$ and the multiflow f satisfy the claimed inequality; all operations, including the WGMV algorithm run in polynomial time. $\square$

Note that if y_{WGMV} is half-integer (assuming integer edge-costs), the obtained multiflow is half-integer and a half-integer flow-cut gap of 2 directly follows. There are examples where the WGMV algorithm does not produce a half-integer dual solution, but we do not know of an instance where half-integer flow-cut gap is larger than 2.

4 From Fractional to Half-Integer

We show here how to convert a flow to a half-integer one, loosing at most half of the flow value, provided the capacities are integers.

Theorem 2. *Let (G, H, c) be a plane multiflow problem, where $c : E \rightarrow \mathbb{Z}_+$ is an integer capacity function. Given a feasible multiflow f , there exists a feasible half-integer multiflow f' , such that $|f'| \geq |f|/2$, and it can be computed in polynomial time.*

Proof. By Lemma 1 we can suppose that the given feasible multiflow f is laminar and can be found in polynomial time. Let $\mathcal{L} \subseteq 2^{V^*}$ be the laminar family of cuts in $(G + H)^*$, with $\{C^* : C = P \cup \{e_P\}, f(P) > 0\} = \{\delta_{E^* \cup F^*}(L) : L \in \mathcal{L}\}$

(see Section 2, just above Lemma 1). Denote $f_L := f(\delta(L)^* \setminus F)$, ($L \in \mathcal{L}$). The feasibility of the multiflow f means $\sum_{L \in \mathcal{L}, e \in \delta_{E^*}(L)} f_L \leq c(e)$, that is, $x_L = f_L \in \mathbb{N}_+$ satisfies

$$\sum_{L \in \mathcal{L}} x_L = |f|, \text{ and } \sum_{L \in \mathcal{L}, e \in \delta_{E^*}(L)} x_L \leq c(e), \text{ for all } e \in E, x_L \geq 0, (L \in \mathcal{L}). \quad (1)$$

Clearly, the edge $e = uv$ is contained in exactly those sets $\delta(L)$ ($L \in \mathcal{L}$) for which $L \in \mathcal{L}(u, v) \cup \mathcal{L}(v, u)$, where $\mathcal{L}(a, b) := \{L \in \mathcal{L} : a \in L, b \notin L\}$ ($a, b \in V^*$), and both $\mathcal{L}(u, v)$ and $\mathcal{L}(v, u)$ form full chains (Lemma 2). So the linear program

$$\begin{aligned} & \max \quad \sum_{L \in \mathcal{L}} x_L \\ & \text{subject to} \quad \sum_{L \in \mathcal{L}(u, v)} x_L \leq c(e) \quad \text{for all } (u, v) \in V^* \times V^*, uv = e^* \in E^* \\ & \quad \quad \quad x_L \geq 0 \quad \quad \quad \text{for all } L \in \mathcal{L}, \end{aligned} \quad (2)$$

is a relaxation of (1): for each $u, v \in V^*, uv = e^* \in E^*$ the coefficient vector of (1) associated with e^* is the sum of the coefficient vectors, one for each of (u, v) and (v, u) , of the two inequalities associated with e^* in (2). Both of these ($\mathcal{L}(u, v)$ and $\mathcal{L}(v, u)$) correspond to full chains in the laminar family $\mathcal{L}$, and the right hand side $c(e)$ is repeated for both.

Denote $M = M(f)$ the $2|E| \times |\mathcal{L}|$ coefficient matrix of (2) (without the non-negativity constraints).

According to Edmonds and Giles [5], $\mathcal{L}$ has a rooted tree (arborescence) representation in which the full chains correspond to subpaths of paths from the root, so M is a network matrix. As such, it is totally unimodular by Tutte [27] and (2) has an integer optimum x , computable in polynomial time by [11], [22, Theorem 19.3 (ii), p. 269].

To finish the proof now, note that on the one hand, f_L ($L \in \mathcal{L}$) is a solution to (1), and therefore it is also a feasible solution of the relaxation (2). Since x_L ($L \in \mathcal{L}$) is the maximum of (2), $\sum_{L \in \mathcal{L}} x_L \geq \sum_{L \in \mathcal{L}} f_L = |f|$. According to the two inequalities of (2) associated to e^* , the sum of coefficients of the paths containing any given edge $e \in E$ is at most $c(e) + c(e) = 2c(e)$, so $f' = x/2$ defines a half-integer feasible flow in (G, H, c) (by assigning the flow value $f'(L)$ to the path $\delta(L)^* \setminus F$, so $|f'| \geq |f|/2$, finishing the proof. $\square$

For more explanations and references showing that M is a network matrix, and an alternative direct combinatorial solution of the integer linear program with a simple greedy-type algorithm, see the Appendix.

This proof does not fully exploit the possibilities of totally unimodular matrices: instead of putting $c(e)$ as right hand side for both inequalities of (2) associated with e^* we can put everywhere the smallest integer capacities satisfied by the fractional flow. Since the fractional values on the mentioned two inequalities sum to at most $c(e)$ and not $2c(e)$ we get a sharper result this way. The proof works if we replace the capacities by the rounded up fractional flow, leading to

an error of only 1 compared to the original capacities, due to the round-up. Let us denote by $\underline{1}$ the all 1 function on E , and check this precisely:

Theorem 3. *Let (G, H, c) be a plane multifold problem, where $c : E \rightarrow \mathbb{Z}_+$. Given a feasible multifold f , there exists a feasible integer multifold f' , computable in polynomial time, feasible for the capacity function $c + \underline{1}$, and $|f'| \geq |f|$.*

Proof. Let $f \in \mathbb{R}^{\mathcal{L}}$ be a feasible multifold, and $M = M(f)$ the $2|E| \times |\mathcal{L}|$ coefficient matrix defined in the proof of Theorem 2. Define $d := Mf \in \mathbb{R}^{2|E|}$, and consider the linear program

$$\begin{aligned} \max \quad & \sum_{L \in \mathcal{L}} x_L \\ \text{subject to} \quad & \sum_{L \in \mathcal{L}(u,v)} x_L \leq \lceil d(u,v) \rceil \quad \text{for all } (u,v) \in V^* \times V^*, uv = e^* \in E^* \\ & x_L \geq 0 \quad \text{for all } L \in \mathcal{L}, \end{aligned} \tag{3}$$

where $d(u,v) := \sum_{L \in \mathcal{L}(u,v)} f_{\delta(L)^* \setminus F}$. In words, (3) has exactly the same coefficients as (2), but the right hand sides $d(u,v)$ and $d(v,u)$ of the two inequalities associated with $e \in E$, $e^* = uv$ are defined with the sum of flow values on $\delta(L)^* \setminus F$, for $L \in \mathcal{L}(u,v)$ and $L \in \mathcal{L}(v,u)$ respectively.

Since f is a feasible flow for (G, H, c) , $d(u,v) + d(v,u) \leq c(e)$, so $\lceil d(u,v) \rceil + \lceil d(v,u) \rceil \leq c(e) + 1$, and f is feasible for (3) since the capacities have been defined by rounding up the flow values. On the other hand, the coefficient matrix is totally unimodular, so by [11], [22, Theorem 19.3 (ii), p. 269] the linear program (3) has an integer maximum solution f' , again computable in polynomial time, and $f' \leq c + \underline{1}$, $|f'| \geq |f|$. $\square$

Theorem 2 is an immediate consequence of Theorem 3:

For each $k \in \mathbb{N}$, $k \geq 1$, $\frac{k+1}{2} \leq k$ holds, so (after deleting 0 capacity edges) $(c + \underline{1})/2 \leq c$, and therefore, dividing by 2 the primal optimum of (3), we immediately get a half-integer solution to (2).

This result extends a natural consequence for maximisation of the tight additive integrality gap known for plane demand flow problems with integer capacities and demands satisfying the cut condition: according to a result of Korach and Penn [15], if all demand edges lie in two faces of the supply graph, *there exists an integer multifold satisfying all demands but at most 1*. This readily implies that increasing each capacity by 1, an integer flow of the same value as the maximum flow for the original capacities, can be reached.

From Frank and Szegedi [8] the same can be deduced for demand-edges lying on an arbitrary number k of faces. Indeed, increasing all capacities by 1, *the surplus of the cut condition will be at least k , which is the Frank-Szegedi condition for integer multifolds*. Theorem 3 states that this consequence is actually true in general, without requiring the integrality of the demands, and also for the maximisation problem; the same holds for maximum “packings of T -cuts”.

5 From Half-Integer to Integer

In this section, we show how to round a half-integer flow to an integer one, losing at most one half of the flow value.

Theorem 4. *Let (G, H, c) be a plane multiflow problem, where $G = (V, E)$, $H = (V, F)$ and $c : E \rightarrow \mathbb{Z}_+$. Given a feasible half-integer multiflow f , there exists a feasible integer multiflow f' , computable in polynomial time, and $|f'| \geq |f|/2$.*

Proof. Let (G, H, c) and f be as assumed in the condition, moreover that f is laminar (Lemma 1). We proceed by induction on the integer $2|f|$. We suppose that *all nonzero values $f(P) > 0$ ($P \in \mathcal{P}$) are actually $1/2$* : if $f(P) \geq 1$, we can decrease $f(P)$ by $\lfloor f(P) \rfloor$, as well as all capacities of edges of P , and the statement follows from the induction hypothesis. (Such a step can be repeated only a polynomial number of times, since $|\mathcal{P}| = O(|V|)$ by Lemma 2.)

To choose the values to round we replace $c(e)$, by $c(e)$ parallel edges of capacity 1 each.⁶ Then take the plane dual of the resulting graph, which is $(G + H)^*$ with each edge $e^* \in E^*$ replaced by a path of size $c(e)$. We consider the laminar system $\mathcal{L}$ defining the paths $P \in \mathcal{P}$, $f(P) > 0$ ⁷ in this subdivided graph so that every edge is contained in at most two sets $L \in \mathcal{L}$, and remains laminar (this is clearly possible, since all positive f_P values are $1/2$). For simplicity, we keep the notations of the original graph - as if what we get in this way were the given graph with all capacities equal to 1.

Let $I(\mathcal{L}) = (\mathcal{L}, M)$ be the intersection graph of the cuts defined by $\mathcal{L}$, that is, $M := \{L_1 L_2 : L_1, L_2 \in \mathcal{L}, \delta(L_1) \cap \delta(L_2) \neq \emptyset\}$. We have Claim a. and b. so far, and now we check Claim c.:

- Claim:** a. Each $e \in E^*$ is contained in at most two sets in $\{\delta(L) : L \in \mathcal{L}\}$.
 b. $|f| = |\mathcal{L}|/2$.
 c. $I(\mathcal{L})$ is planar.

To check Claim c., note first its validity if $\mathcal{L}$ consists of disjoint sets. If this does not hold, not even by *complementing* some $L \in \mathcal{L}$ (i.e. replacing it by $V^* \setminus L$), then it is easy to find (possibly after complementation) three sets $L_1 \subset L_0 \subset L_2$ in $\mathcal{L}$. We claim that L_0 is a cut-vertex in $I(\mathcal{L})$. By laminarity, every cut $\delta(L)$ ($L \in \mathcal{L}$) has either a shore A contained in L_0 (like L_1), or a shore B disjoint from L_0 (like $V \setminus L_2$). If there exists an $e \in \delta(A) \cap \delta(B)$, this would mean that e has an endpoint in $A \subseteq L_0$ and the other endpoint in $B \subseteq V \setminus L_0$, so $e \in \delta(L_0)$, contradicting Claim a. So L_0 is a cut vertex, i.e. $\mathcal{L} = \mathcal{L}_1 \cup \mathcal{L}_2$, $\mathcal{L}_1 \cap \mathcal{L}_2 = \{L_0\}$, with no edge between $\mathcal{L}_1$ and $\mathcal{L}_2$ in $I(\mathcal{L})$.

Since the graphs induced by $\mathcal{L}_i$ ($i = 1, 2$) are both defined by flows of smaller value, we can apply the induction hypothesis to them: they are planar, so $I(\mathcal{L})$ is also planar, and the Claim follows.

⁶ This is not an allowed step for a polynomial algorithm, but it will not really be necessary to do it. The choice of the cuts to be rounded down or up will be clear from the proof without actually executing this subdivision. The choices for rounding concern a family of size $O(|V|)$.

⁷ $\{\delta_{E^* \cup F^*}(L)^* : L \in \mathcal{L}\} = \{P \cup e_P : P \in \mathcal{P}, f(P) > 0\}$, as before.

To finish the proof of the theorem using Claim c., find a stable set of size $|\mathcal{L}|/4$ in $\mathcal{L}$, by Lemma 3, and increase the flow on the corresponding paths $\delta(L)^* \setminus F$ to 1, while decreasing the flow on the other paths to 0. This results in a feasible integer flow $|f'| \geq |\mathcal{L}|/4 = |f|/2$, finishing the proof of the bound.

For the computational complexity results, first recall that the support of the half integer laminar flow f obeys Lemma 2a. Then note, that among this linear number of sets, the proof finds in polynomial time at least one fourth of the cuts to round up, while the other cuts are rounded down, so that the capacity constraints are not violated. It is straightforward to mimick the subdivision of edges without doing it, and to compute an input to Lemma 3, in strongly polynomial time. $\square$

6 Examples

In the previous sections we showed upper bounds on the flow-cut gap, lower bounds on the (half)-integrality gap equal to the approximation guarantee of multiflow maximisation. In this Section we present some examples proving lower bounds for multicuts and upper bounds for multiflows, showing the performance of the precedingly proved theorems.

A first simple example occurred already: $(\overline{C_4}, \overline{2K_2})$ (Footnote 4), where the maximisation of demands so that a feasible flow exists can be achieved in integers, but the maximum integer flow is only half of the maximum flow. Another, still simple and small example shows already that a unique max-flow may not even be half-integer (Figure 1).

The gap values of Figure 1 are not best possible. We provide now an infinite class of examples providing asymptotically the best possible values for some of the gaps. For every non-negative integer k we define a graph G_k , where for instance the half-integer flow-cut gap tends to 2 as $k \rightarrow \infty$. (Cheriyān et.al. [4] used these instances to show integrality gap results for the Tree Augmentation Problem.)

Let $G_k = (V_k, E_k)$, $H_k = (V_k, F_k)$, $k \geq 3$ be an instance of the multiflow problem defined as follows: $V_k = \{a_1, a_2, \dots, a_k\} \cup \{b_1, b_2, \dots, b_k\}$, $E_k = \{(a_i, b_i) | i \in [1, k]\} \cup \{(a_i, a_{i+1}) | i \in [1, k-1]\}$ and $F_k = \{(b_i, b_{i+1}) | i \in [1, k-1]\} \cup \{(b_i, a_{i+2}) | i \in [1, k-2]\}$. The capacity of all edges in E_k is 1 (see Fig. 2).

Theorem 5. *The graph $G_k + H_k$ is planar for all $k = 1, 2, \dots$, and the following hold:*

- *The minimum multicut capacity is $k - 1$.*
- *The maximum multiflow value is $2(k - 1)/3$.*
- *The maximum half-integer multiflow value is $k/2$.*
- *The maximum integer multiflow value is $\lfloor k/2 \rfloor$.*

Proof. The minimum multicut capacity is clearly at most $k - 1$, since deleting each edge of the (a_1, a_{k-1}) -path, the endpoints of all demand-edges are separated. We check now $|C| \geq k - 1$ for an arbitrary multicut C , by induction. For $k = 1, 2$ the statement is obvious. Deleting b_1 and a_1 , the remaining (G', H') is




Fig. 1: Black edges are supply while red (thick) ones are demand edges. All supply edges have capacity 1. The maximum flow value is $7/3$, while the value of the maximum half-integer (or integer) flow is 2. The value of minimum multicut is 3. So the (half-) integrality gap of this example is $6/7$, the flow-cut gap is at least $9/7$, the (half) integer flow-cut gap is at least $3/2$.

(isomorphic to) G_{k-1}, H_{k-1} and $C' := C \setminus \{a_1b_1, a_1a_2\}$ is a multicut in it. By the induction hypothesis, the minimum multicut of (G', H') is of size $k - 2$.

Now either both $a_2b_2 \in C'$ and $a_2a_3 \in C'$ in which case C' is not an inclusion-wise minimal multicut for (G', H') , since deleting one of a_1b_1, a_1a_2 we already disconnect the same terminal pairs. So in this case $|C'| > k - 2$, and we are done; or one of a_2b_2, a_2a_3 is not in C' , but then one of a_1b_1, a_1a_2 must be in it, since otherwise b_1 is not disconnected by C from b_2 or from a_3 . So in this case $|C| \geq |C'| + 1 \geq k - 1$, finishing the proof of the first assertion.

For the maximum multiflow value μ , or maximum half-integer and integer multiflow values $\mu_{\text{half}}, \mu_{\text{int}}$, note that the supply edges form a tree, so the set $\mathcal{P}$ of paths between the endpoints of demand edges contains exactly one path P_e for each demand edge e , so $|\mathcal{P}| = 2k - 3$. Defining $f(P_{b_{k-1}b_k}) = 2/3$ and $f(P) = 1/3$ for each other path $P \in \mathcal{P}$, we have $|f| = 2(k - 1)/3$. To prove that this is a maximum flow, note that $e_1 := a_1b_1$ is contained in $P \in \mathcal{P}$ if and only if $e_2 := a_2b_2$ is contained in it, so for any multiflow f ,

$$\alpha := \alpha_f := \sum_{\{P \in \mathcal{P} : e_1 \in P\}} f(P) = \sum_{\{P \in \mathcal{P} : e_2 \in P\}} f(P)$$

Claim: $\mu \leq 2(k - 1)/3$, and if $\alpha \leq 2/3$, then $\mu \leq 2(k - 2)/3 + \alpha$.

Indeed, on the one hand, $\alpha = 2/3 + \varepsilon$ ($0 \leq \varepsilon \leq 1/3$) causes $\alpha' \leq 2/3 - \varepsilon$ in (G', H') isomorphic to (G_{k-1}, H_{k-1}) with max flow value μ' , after the deletion of $\{a_1, b_1\}$, and then by induction, $\mu \leq 2/3 + \varepsilon + \mu' \leq 2/3 + \varepsilon + 2(k - 2)/3 -$




Fig. 2: G_8 : the capacity of supply edges is 1; demand edges are red (thick).

$\varepsilon = 2(k-1)/3$. On the other hand, if $\alpha = 2/3 - \varepsilon$ we have by induction $\mu \leq 2/3 - \varepsilon + \mu' \leq 2/3 - \varepsilon + 2(k-2)/3 = 2(k-1)/3 - \varepsilon$, and the claim is proved, finishing the proof of the second statement.

The proof of the third assertion, concerning μ_{half} works similarly. Defining $f(P_{b_i b_{i+1}}) = 1/2$ ($i = 1, \dots, k-1$), and $f(P_{b_1 a_3}) = 1/2$, otherwise $f(P) = 0$, where f is a feasible half-integer flow, $|f| = k/2$. To prove $\mu_{\text{half}} \leq k/2$ an inductive proof using α works as before: for $\alpha \geq 1/2$ the stronger statement $\mu \leq (k-2)/2 + \alpha$ follows by induction.

Finally, the fourth assertion is immediate from the third one: $\mu_{\text{int}} \leq \mu_{\text{half}} \leq k/2$ is integer, and an integer flow f , $|f| = \lfloor k/2 \rfloor$ is also easy to construct. $\square$

This theorem provides the best known bounds for several of the defined gaps, and in some cases the best possible bounds that are reviewed below.

7 Conclusions

This paper established bounds on the integrality gap of multiflows, developed approximation algorithms for them, and bounded the flow-cut gap. Applying Theorem 1 first, then Theorem 2 to a maximum multifold, and finally Theorem 4 to the result, we arrive at the following summary:

Theorem 6. *There exists a $1/4$ -approximation algorithm for integer plane multifold maximisation, with an integer flow-cut gap of 8; there exists a $1/2$ -approximation algorithm for half-integer plane multiflows with a half-integer flow-cut gap of 4; the flow-cut gap is at most 2; the approximation algorithms provide lower bounds to the integrality or half-integrality gap equal to the approximation ratio.*

a. Multicuts and flow-cut gap: The minimum multicut is NP-hard to find, but has a PTAS by applying [14] after the transformation of Lemma 4. The (“fractional”) flow-cut gap is in the interval $[3/2, 2]$ by Theorem 5 illustrated in Figure 2 and Theorem 6; the half-integer flow-cut gap is at least 2 and at most 4 for the same reason. For the integer flow-cut gap, a lower bound of 2 is shown by

$(\overline{C_4}, \overline{2K_2})$ (Footnote 4). The true value of the integer flow-cut gap is thus wide open in the interval $[2, 8]$.

b. Half-integrality gap: We do not know the complexity of finding a half-integer multiflow of maximum value. The worst case ratio between the maximum value of a half-integer feasible flow and of a fractional flow is in the interval $[1/2, 3/4]$, again by Theorem 2 and Theorem 5. It remains an interesting open problem to pin down the exact half-integrality gap in this interval, and the complexity of finding the maximum value of a half-integer flow.

c. Integrality gap: This lies in the interval $[1/4, 1/2]$, with lower bound given by Theorem 6, and upper bound by $(\overline{C_4}, \overline{2K_2})$ (Footnote 4). Finally, the worst integer flow/half-integer flow ratio is closed: it is exactly $1/2$, as the same example and Theorem 4 show.

SUMMARY	fractional	half-integer	integer
mflow/LP	1	$[1/2, 3/4]^9$	$[1/4, 1/2]^{10}$
mcut/mflow	$(3/2, 2]^{11}$	$(2, 4]^{12}$	$(2, 8]^{11}$
complexity	P	?	NP-hard

Gaps, complexity, and approximation ratios for plane multiflows and multicuts

The approximation ratios for multiflows are the lower bounds of the first row. The second row contains the flow-cut gaps; an upper bound for the approximation ratio for minimum multicuts is provided by the first upper bound in this row (equal to 2), but there is also a PTAS [14] for the minimum multicut problem together with an NP-hardness proof. The complexity results (last row) are known (see Introduction) and the Preliminaries, except for half-integer flows. The Gaps and exact approximation ratios are open in the indicated intervals and the maximum half integer plane multiflow problem is also open.

References

1. K. Appel and W. Haken. A proof of the four color theorem. *Discrete Mathematics*, 16:179–180, 1976.
2. Solal Attias. *Exercices et Problèmes Combinatoires: Connectivité des Réseau et Routing*. Rapport de stage de licence, Grenoble, June-July, 2019.
3. Brenda S. Baker. Approximation algorithms for np-complete problems on planar graphs. *J. ACM*, 41(1):153–180, January 1994.
4. Joseph Cheriyan, Howard Karloff, Rohit Khandekar, and Jochen Könemann. On the integrality ratio for tree augmentation. *Operations Research Letters*, 36(4):399–401, 2008.
5. Jack Edmonds and Giles Rick. A min-max relation for submodular functions in graphs. *Ann. Discrete Math*, 1, pages 185–204, 1977.
6. S. Fiorini, N. Hardy, B. Reed, and A. Vetta. Approximate min-max relations for odd cycles in planar graphs. *Mathematical Programming*, 110(1):71–91, 1995.

⁹ Lower bound: Theorem 2 ; upper bound: Theorem 5 , see Figure 2 .

¹⁰ Lower bound: Theorem 2 , Theorem 4 ; upper bound: $(\overline{C_4}, \overline{2K_2})$.

¹¹ Lower bound: Theorem 5 , see Figure 2; upper bound: Theorem 1

¹² Lower bound: $(\overline{C_4}, \overline{2K_2})$; upper bound: Theorem 6 .

7. András Frank. *Connections in Combinatorial Optimization*. Oxford University Press, 2011.
8. András Frank and Zoltán Szigeti. A note on packing paths in planar graphs. *Mathematical Programming*, 70:201–209, 1995.
9. Naveen Garg, Vijay V Vazirani, and Mihalis Yannakakis. Approximate max-flow min-(multi) cut theorems and their applications. *SIAM Journal on Computing*, 25(2):235–251, 1996.
10. Naveen Garg, Vijay V. Vazirani, and Mihalis Yannakakis. Primal-dual approximation algorithms for integral flow and multicut in trees. *Algorithmica*, 18(1):3–20, 1997.
11. A. J. Hoffman and J.B.Kruskal. Integral boundary points of convex polyhedra. *Linear Inequalities and Related Systems*, pages 223–246, 1956.
12. Chien-Chung Huang, Mathieu Mari, Claire Mathieu, Kevin Schewior, and Jens Vygen. *An Approximation Algorithm for Fully Planar Edge-Disjoint Paths*. <https://arxiv.org/abs/2001.01715>, 2020.
13. Philip Klein, Serge A Plotkin, and Satish Rao. Excluded minors, network decomposition, and multicommodity flow. In *Proceedings of the twenty-fifth annual ACM symposium on Theory of computing*, pages 682–690. ACM, 1993.
14. Philip N Klein, Claire Mathieu, and Hang Zhou. Correlation clustering and two-edge-connected augmentation for planar graphs. In *32nd International Symposium on Theoretical Aspects of Computer Science (STACS 2015)*. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2015.
15. Ephraim Korach and Michal Penn. Tight integral duality gap in the chinese postman problem. *Mathematical Programming*, 55:183–191, 1992.
16. Bernhard Korte and Jens Vygen. *Combinatorial optimization: Theory and Algorithms*, volume 21. Springer-Verlag Berlin and Heidelberg, collection Algorithms and Combinatorics, fifth edition, 2012.
17. D. Král and H. Voss. Edge-disjoint odd cycles in planar graphs. *Journal of Combinatorial Theory, Series B*, 90:107–120, 2004.
18. László Lovász and Michael Plummer. *Matching Theory*. Akadémiai Kiadó and Springer, 1986.
19. Matthias Middendorf and Frank Pfeiffer. On the complexity of the disjoint paths problem. *Combinatorica*, 13(1):97–107, 1993.
20. Haruko Okamura and Paul D Seymour. Multicommodity flows in planar graphs. *Journal of Combinatorial Theory, Series B*, 31(1):75–81, 1981.
21. Neil Robertson, Daniel P Sanders, Paul Seymour, and Robin Thomas. Efficiently four-coloring planar graphs. In *Proceedings of the twenty-eighth annual ACM symposium on Theory of computing*, pages 571–575, 1996.
22. Alexander Schrijver. *Theory of Linear and Integer Programming*. Wiley and Chichester, 1986.
23. Alexander Schrijver. *Combinatorial optimization: polyhedra and efficiency*, volume 24. Springer Science & Business Media, 2003.
24. Loïc Seguin-Charbonneau and F Bruce Shepherd. Maximum edge-disjoint paths in planar graphs with congestion 2. In *2011 IEEE 52nd Annual Symposium on Foundations of Computer Science*, pages 200–209. IEEE, 2011.
25. Paul D Seymour. On odd cuts and plane multicommodity flows. *Proceedings of the London Mathematical Society*, 3(1):178–192, 1981.
26. Éva Tardos and Vijay V. Vazirani. Improved bounds for the max-flow min-multicut ratio for planar and k_r -free graphs. *Inf. Process. Lett.*, 47(2):77–80, 1993.
27. W. T. Tutte. Lectures on matroids. *J. of Research, Nat. Bureau of Standards*, B(69), 1965.

28. David P Williamson, Michel X Goemans, Milena Mihail, and Vijay V Vazirani. A primal-dual approximation algorithm for generalized steiner network problems. *Combinatorica*, 15(3):435–454, 1995.

APPENDIX: Uncrossing, Laminarity and Integrality

Uncrossing, laminar families and the related totally unimodular network matrices and integer solutions to linear programs with constraint-matrices having some of these properties are widely used in combinatorial optimization [23], [16], [7]. We wish to sketch here the main ideas or references about facts or arguments related to proofs, algorithms or complexity results of our work. For some of these we provide an adaptation to the special cases and particular purposes of this paper, with the notation and terminology used here.

The **uncrossing** we use is essentially the same as the one introduced in [18, Section 6.5, p. 235] for odd cuts (more precisely “*T*-cuts”). To restate it recall $G = (V, E)$, $H = (V, F)$, let $G + H$ be planar, and consider cuts in $(G + H)^* = (V^*, E^* \cup F^*)$ of the form $\delta(X) := \delta_{E^* \cup F^*}(X)$. It is easy to check:

Let $\delta(X), \delta(Y) \subseteq E^ \cup F^*$ ($X, Y \subseteq V^*$) be two cuts in $(G + H)^*$, both containing exactly one edge of F^* , which cross. Then every edge of $E^* \cup F^*$ is covered at most as many times by $\delta(X \cap Y)$ and $\delta(X \cup Y)$, and – possibly after complementing X say – both contain exactly one edge of F^* . Equivalently, either $\delta(X \cap Y)$ and $\delta(X \cup Y)$, or $\delta(X \setminus Y)$ and $\delta(X \setminus Y)$ contain exactly one edge of F^* , so they “can replace $\delta(X), \delta(Y)$ ”.*

Another property not to forget about is that *the number (multiplicity) of crossing pairs decreases in this way*.

It is not difficult to see that the series of replacements, called *uncrossing*, is finite even if the cuts are endowed with multiplicity coefficients: then we can uncross X and Y with a multiplicity equal to the minimum of their coefficients. It is also true that uncrossing can be executed in polynomial time in a general setting. (See [23, 60.17, 60.3e] for a proof of finiteness and further references.)

Laminarity is used then throughout the paper. In Section 3 implicitly, through the WGMV algorithm. In Section 4 we can use Edmonds and Giles’s [5] to prove integrality of (2) and (3). The closest to our use is Korte and Vygen’s “one-way cut-incidence matrix” [16, Theorem 28] which directly deduces the total unimodularity of our matrix M : replace every E^* by arcs in both directions and put a capacity of $c(e)$ on both (and infinite capacity on F^*) to have (2), and put $d(u, v), d(v, u)$ on the two arcs to get (3). These can also be deduced from the rooted tree representations of laminar systems and their total unimodularity [7, 1.4.1, Section 4.2.2] and [23, Theorem 13.21, pages 213–216], since the inequalities in (2) and (3) are subpaths of paths from the root, so they define a non-negative network matrix.

Once network matrices have been identified, total unimodularity is known from Tutte [27], included directly in Korte and Vygen’s discussion. Hoffman and Kruskal [11] observed the **integer solvability** of linear programs with totally unimodular coefficient matrices, in polynomial time, and in particular, that of (2) and (3).

After realizing polynomial solvability in multiple ways from known results, one can develop simple direct algorithms for our special case.

In fact a maximum integer solution to $Mx \leq b, x \geq 0$ and a minimum solution to the dual can be realised by a greedy algorithm for arbitrary $b \in \mathbb{N}_+^{2|E|}$ assigning right hand sides independently of one another to each ordered pair (u, v) and (v, u) such that $uv \in E^*$, in the following way:

We restrict ourselves to all 1 costs, as in (2) and (3). Take an inclusionwise minimal set $L \in \mathcal{L}$ and let $x_L := b(u_0, v_0)$ be the minimum of $b(u, v)$ among $u, v \in V^*, u \in L, v \notin L$. Deleting row (u_0, v_0) from M and all variables corresponding to a 1 in this row (involving the deletion of all corresponding columns of M and coordinates of b and c), and updating b with the assignments $b(u, v) := b(u, v) - b(u_0, v_0)$ for all $u \in L, v \notin L$, continue as follows:

By induction, the greedy algorithm determines a primal and dual solution of the same value, a relation kept after the assignment $x_L = b(u_0, v_0)$ and

if so far $y(u, v) = 0$ for each pair $(u, v), u \in L, v \in V^ \setminus L$, assign $y(u_0, v_0) = 1$.*

It is straightforward to check that the equality between the primal and dual solution is kept in this way, we leave this to the reader. The particular structure of our constraints has to be exploited through the following property : $\mathcal{L}(u, v) \supset \mathcal{L}(u_0, v_0)$ for all constraints that contain L and do not disappear after the reduction.