arXiv:2002.11001v1 [math.AP] 25 Feb 2020

An improved Trudinger-Moser inequality involving N–Finsler–Laplacian and L^p norm

Yanjun Liu[a,b]

[a]School of Mathematical Sciences, Nankai University, Tianjin 300071, P. R. China

[b]Albert-Ludwigs-Universität Freiburg, Mathematisches Institut, Eckerstr. 1,

D-79104 Freiburg, Germany

Abstract: Suppose $F:\mathbb{R}^N\to[0,+\infty)$ be a convex function of class $C^2(\mathbb{R}^N\backslash\{0\})$ which is even and positively homogeneous of degree 1. We denote $\gamma_1=\inf\limits_{u\in W_0^{1,N}(\Omega)\backslash\{0\}}\frac{\int_\Omega F^N(\nabla u)dx}{\|u\|_p^N}$. and define the norm $\|u\|_{N,F,\gamma,p}=\left(\int_\Omega F^N(\nabla u)dx-\gamma\|u\|_p^N\right)^{\frac{1}{N}}$. Let $\Omega\subset\mathbb{R}^N(N\geq 2)$ be a smooth bounded domain. Then for $p>1$ and $0\leq\gamma<\gamma_1$, we have

$$\sup_{u\in W_0^{1,N}(\Omega),\|u\|_{N,F,\gamma,p}\leq 1}\int_\Omega e^{\lambda|u|^{\frac{N}{N-1}}}dx<+\infty,$$

where $0<\lambda\leq\lambda_N=N^{\frac{N}{N-1}}\kappa_N^{\frac{1}{N-1}}$ and κ_N is the volume of a unit Wulff ball. Moreover, by using blow-up analysis and capacity technique, we prove that the supremum can be attained for any $0\leq\gamma<\gamma_1$.

Keywords: N–Finsler–Laplacian; Trudinger-Moser inequality; Extremal function; Blow-up analysis; Elliptic regularity theory

MSC2010: 26D10, 35J70, 46E35

1 Introduction and main results

Suppose $\Omega\subset\mathbb{R}^N(N\geq 2)$ be a bounded smooth domain. When $1<p<N$, the Sobolev embedding theorem implies that $W_0^{1,p}(\Omega)\hookrightarrow L^q(\Omega)$ is continuous for $1\leq q\leq\frac{Np}{N-p}$. In particular, $W_0^{1,N}(\Omega)\hookrightarrow L^q(\Omega)$ for $1\leq q<\infty$, but the

Email: liuyj@mail.nankai.edu.cn(Y. Liu).

embedding $W_0^{1,N}(\Omega) \not\hookrightarrow L^\infty(\Omega)$. A counterexample is given by the function $u(x) = (-\ln|\ln|x||)_+$ as Ω is the unit ball. It was proposed independently by Yudovich [55], Pohozaev [38], Peetre [39] and Trudinger [46] that $W_0^{1,N}(\Omega)$ is embedded in the Orlicz space $L_{\varphi_\alpha}(\Omega)$ determined by the Young function $\varphi_\alpha(t) = e^{\alpha|t|^{\frac{N}{N-1}}} - 1$ for some positive number α, it was sharpened by Moser [36] who found the best exponent and proved the following result:

Theorem A There exists a sharp constant $\alpha_N := N^{\frac{N}{N-1}}\omega_N^{\frac{1}{N-1}}$ such that

$$\sup_{u\in W_0^{1,N}(\Omega),\|\nabla u\|_N\le 1}\int_\Omega e^{\alpha|u|^{\frac{N}{N-1}}}dx < +\infty, \forall \alpha \le \alpha_N, \tag{1.1}$$

where ω_N is the volume of unit ball in $\mathbb{R}^N$. Moreover, the supremum in (1.1) is $+\infty$ if $\alpha > \alpha_N$. Related inequalities for unbounded domains were proposed by D. M. Cao [7] in dimension two and J. M. do Ó [15], Adachi-Tanaka [1] in high dimension, however they just considered the subcritical Trudinger-Moser inequality. Ruf [40] (for the case $N = 2$), Li and Ruf [26] (for the general case $N \ge 2$) obtained the Trudinger-Moser inequality in the critical case by replacing the Dirichlet norm with the standard Sobolev norm in $W^{1,N}(\mathbb{R}^N)$. Subsequntly, Masmoudi and Sani [34] derived Trudinger-Moser inequalities with the exact growth condition in $\mathbb{R}^n$, These inequality plays an important role in geometric analysis and partial differential equations, we refer to [10, 12, 4, 13, 23, 16, 34] and references therein. In [24], Lam, Lu and Zhang provide a precise relationship between subcritical and critical Trudinger-Moser inequality. The similar result in Lorentz-Sobolev norms was also proved by Lu and Tang [31]. Trudinger-Moser inequality for first order derivatives was extended to high order derivatives by D. Adams [2] for bounded domains when dimension $N \ge 2$. B. Ruf and F. Sani [41] studied the Adams type inequality with higher derivatives of even orders for unbounded domains in $\mathbb{R}^N$. In [22], Lam and Lu applied a rearrangement-free argument to prove sharp Adams' inequality in general case.

One important problem on Trudinger-Moser inequalities is whether or not extremal functions exist. Existence of extremal functions for the Trudinger-Moser inequality was first obtained by Carleson-Chang [8] when Ω is the unit ball, by M. Struwe [42] when Ω is close to the ball in the sense of measure, then by M. Flucher

[19] and K. Lin [29] when Ω is a general bounded smooth domain. Recently based on the work by Malchiodi and Martinazzi in [33], Mancini and Martinazzi [35] reproved the Carleson and Chang's result by using a new method based on the Dirichlet energy, also allowing for perturbations of the functional. In the entire Euclidean space, existence of extremal functions was proved by Ruf [40] (for the case $N = 2$) and Li and Ruf [26] (for the general case $N \geq 2$). For extremal functions of singular version, Csat*ó* and Roy [11] proved that extremal functions exist in bounded domain of 2 dimension. Li and Yang [27] proved that extremal functions exist in the entire Euclidean space.

Moreover, there are some extensions of the Trudinger-Moser inequality. Let $\alpha_1(\Omega)$ be the first eigenvalue of the Laplacian, Adimurthi and O. Druet [3] proved that

$$\sup_{u\in W_0^{1,2}(\Omega),\int_\Omega |\nabla u|^2 dx\leq 1}\int_\Omega e^{4\pi u^2(1+\alpha\|u\|_2^2)}dx < +\infty$$

for $0 \leq \alpha < \alpha_1(\Omega)$, the supremum is infinity for any $\alpha \geq \alpha_1(\Omega)$. This result was generalized by Yang [51, 52] to the cases of high dimension and a compact Riemannian surface. Lu-Yang [32] and J. Zhu [58] considered the case involving the L^p norm for any $p > 1$. For existence of extremal functions of Adimurthi-Druet type inequalities, they proved in [52, 32] that supremums ($N = 2$) are attained for sufficiently small $\alpha \geq 0$, and that the supremum ($N \geq 3$)[51] is attained for all $\alpha, 0 \leq \alpha < \alpha_1(\Omega)$. Subsequently, J.M. do *Ó* and M. de Souza generalized the similar result in whole Euclidean space [14] and high dimension case [17], and the existence of extremal functions was also obtained. A stronger version was established by Tintarev [45], namely,

$$\sup_{u\in W_0^{1,2}(\Omega),\int_\Omega |\nabla u|^2 dx-\alpha\|u\|_2^2\leq 1}\int_\Omega e^{4\pi u^2}dx < +\infty, \quad 0 \leq \alpha < \alpha_1(\Omega), \tag{1.2}$$

Yang [53] obtained extremal functions for (1.2), which was also extended to singular version (see [54]). In [37], the author extends the result of Tintarev to the higher dimension as the following result:

Theorem B. Let $\Omega \subset \mathbb{R}^N (N \geq 2)$ be a smooth bounded domain and define

$$\alpha(\Omega) = \inf_{u\in W_0^{1,N}(\Omega),u\not\equiv 0}\frac{\int_\Omega |\nabla u|^N dx}{\|u\|_N^N}.$$

Then for any $0 \leq \alpha < \alpha(\Omega)$,

$$\sup_{u\in W_0^{1,N}(\Omega),\int_\Omega |\nabla u|^N dx-\alpha\|u\|_N^N\leq 1}\int_\Omega e^{\alpha_N|u|^{\frac{N}{N-1}}}dx < +\infty, \tag{1.3}$$

where $\alpha_N := N^{\frac{N}{N-1}}\omega_N^{\frac{1}{N-1}}$ and ω_N is the volume of unit ball in $\mathbb{R}^N$.

Another interesting research is that Trudinger-Moser inequality has been generalized to the case of anisotropic norm. In this paper, denote that $F \in C^2(\mathbb{R}^N\backslash 0)$ is a positive, convex and homogeneous function, $F_{\xi_i} = \frac{\partial F}{\partial \xi_i}$ and its polar $F^o(x)$ represents a Finsler metric on $\mathbb{R}^N$. We will replace the isotropic Dirichlet norm $\|u\|_{W_0^{1,N}(\Omega)} = (\int_\Omega |\nabla u|^N dx)^{\frac{1}{N}}$ by the anisotropic Dirichlet norm $(\int_\Omega F^N(\nabla u)dx)^{\frac{1}{N}}$ in $W_0^{1,N}(\Omega)$. In [49], Wang and Xia proved the following result:

Theorem C (Anisotropic Trudinger-Moser Inequality). Let $\Omega \subset \mathbb{R}^N (N \geq 2)$ be a smooth bounded domain. Let $u \in W_0^{1,N}(\Omega)$ and $(\int_\Omega F^N(\nabla u)dx) \leq 1$. Then

$$\sup_{u\in W_0^{1,N}(\Omega),\int_\Omega F^N(\nabla u)dx\leq 1}\int_\Omega e^{\lambda_N u^{\frac{N}{N-1}}}dx < +\infty, \tag{1.4}$$

where $\lambda_N = N^{\frac{N}{N-1}}\kappa_N^{\frac{1}{N-1}}$ and $\kappa_N = \{x \in \mathbb{R}^N : F^o(x) \leq 1\}$. λ_N is sharp in the sense that if $\lambda > \lambda_N$ then there exists a sequence (u_n) such that $\int_\Omega e^{\lambda u^{\frac{N}{N-1}}}dx$ diverges.

The above inequality is related with N-Finsler-Laplacian operator Q_N which is defined by

$$Q_N u := \sum_{i=1}^N \frac{\partial}{\partial x_i}(F^{N-1}(\nabla u)F_{\xi_i}(\nabla u)),$$

when $N = 2$ and $F(\xi) = |\xi|$, Q_2 is just the ordinary Laplacian. This operator is closely related to a smooth, convex hypersurface in $\mathbb{R}^N$, which is called the Wulff shape (or equilibrium crystal shape) of F. This operator Q_N was studied in some literatures, see [5, 6, 18] and the references therein. In [56], they obtained the existence of extremal functions for the sharp geometric inequality (1.4).

Our aim is to establish and find extremal functions for Trudinger-Moser inequality involving N–Finsler–Laplacian and L^p norm. For $p > 1$, we denote

$$\gamma_1 = \inf_{u\in W_0^{1,N}(\Omega),u\not\equiv 0}\frac{\int_\Omega F^N(\nabla u)dx}{\|u\|_p^N},$$

and

$$\|u\|_{N,F,\gamma,p} = \left(\int_\Omega F^N(\nabla u)dx - \gamma\|u\|_p^N\right)^{\frac{1}{N}}.$$

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^N (N \geq 2)$ be a smooth bounded domain. Then for any $0 \leq \gamma < \gamma_1$,*

$$\Lambda_\gamma = \sup_{u \in W_0^{1,N}(\Omega), \|u\|_{N,F,\gamma,p} \leq 1} \int_\Omega e^{\lambda_N |u|^{\frac{N}{N-1}}} dx < +\infty, \tag{1.5}$$

where $\lambda_N = N^{\frac{N}{N-1}} \kappa_N^{\frac{1}{N-1}}$ and κ_N is the volume of a unit Wulff ball.

Remark 1.2. From Theorem 1.1, for $0 \leq \gamma < \gamma_1$, we can derive the following weak version:

$$\sup_{u \in W_0^{1,N}(\Omega), \int_\Omega F^N(\nabla u) dx \leq 1} \int_\Omega e^{\lambda_N (1+\gamma \|u\|_p^N)^{\frac{1}{N-1}} |u|^{\frac{N}{N-1}}} dx < +\infty, \tag{1.6}$$

for the special case $p = N$ in (1.6), we refer reader to [57]. Next, let's show that we have got better result. Indeed, if $0 \leq \gamma < \gamma_1$, $u \in W_0^{1,N}(\Omega)$ and $\int_\Omega F^N(\nabla u) dx \leq 1$. Set $v = (1+\gamma \|u\|_p^N)^{\frac{1}{N}} u \in W_0^{1,N}(\Omega)$, note that F is a positively homogeneous function of degree 1. Thus

$$\begin{aligned} \|v\|_{N,F,\gamma,p}^N &= (1 + \gamma \|u\|_p^N) \int_\Omega F^N(\nabla u) dx - \gamma \|u\|_p^N - \gamma^2 \|u\|_p^{2N} \\ &\leq \int_\Omega F^N(\nabla u) dx \leq 1. \end{aligned}$$

Applying (1.5) to the function v, we obtain (1.6). This implies that (1.5) is a stronger inequality.

Theorem 1.2. *Let $\Omega \subset \mathbb{R}^N (N \geq 2)$ be a smooth bounded domain. Then the supremum*

$$\Lambda_\gamma = \sup_{u \in W_0^{1,N}(\Omega), \|u\|_{N,F,\gamma,p} \leq 1} \int_\Omega e^{\lambda_N |u|^{\frac{N}{N-1}}} dx \tag{1.7}$$

can be attained by $u_0 \in W_0^{1,N}(\Omega) \bigcap C^1(\overline{\Omega})$ with $\|u_0\|_{N,F,\gamma,p} = 1$.

This paper is organized as follows: In Section 2, we give some preliminaries, meanwhile, under anisotropic Dirichlet norm and L^p norm, we establish the Lions type concentration-compactness principle of Trudinger-Moser Inequalities. In Section 3, we give the existence of subcritical maximizers. In Section 4, we analyze the convergence of maximizing sequence and its blow-up behavior, an upper bound is established by capacity estimates. In Section 5, we provide the proof of Theorems 1.1 and 1.2 by contradiction arguments and the construction of test function.

2 Preliminaries

In this section, we will give some preliminaries for our use later. Let $F : \mathbb{R}^N \to [0, +\infty)$ be a convex function of class $C^2(\mathbb{R}^N \backslash \{0\})$ which is even and positively homogeneous of degree 1, so that

$$F(t\xi) = |t|F(\xi) \qquad \text{for any} \quad t \in \mathbb{R}, \quad \xi \in \mathbb{R}^N. \tag{2.1}$$

We also assume that $F(\xi) > 0$ for any $\xi \neq 0$ and $Hess(F^2)$ is positive definite in $\mathbb{R}^N \backslash \{0\}$. Then by Xie and Gong [50], $Hess(F^N)$ is also positive definite in $\mathbb{R}^N \backslash \{0\}$. A typical example is $F(\xi) = (\sum_i |\xi_i|^q)^{\frac{1}{q}}$ for $q \in [1, \infty)$. Let F^o be the support function of $K := \{x \in \mathbb{R}^N : F(x) \leq 1\}$, which is defined by

$$F^o(x) := \sup_{\xi \in K} \langle x, \xi \rangle,$$

so $F^o : \mathbb{R}^N \to [0, +\infty)$ is also a convex, homogeneous function of class $C^2(\mathbb{R}^N \backslash \{0\})$. From [5], F^o is dual to F in the sense that

$$F^o(x) = \sup_{\xi \neq 0} \frac{\langle x, \xi \rangle}{F(\xi)}, \qquad F(x) = \sup_{\xi \neq 0} \frac{\langle x, \xi \rangle}{F^o(\xi)}.$$

Consider the map $\phi : S^{N-1} \to \mathbb{R}^N$, $\phi(\xi) = F_\xi(\xi)$. Its image $\phi(S^{N-1})$ is smooth, convex hypersurface in $\mathbb{R}^N$, which is called the Wulff shape (or equilibrium crystal shape) of F. Then $\phi(S^{N-1}) = \{x \in \mathbb{R}^N | F^o(x) = 1\}$(see [48], Proposition 2.1). Denote $\mathcal{W}_r(x_0) = \{x \in \mathbb{R}^N : F^o(x - x_0) \leq r\}$, we call $\mathcal{W}_r(0)$ is a Wulff ball of radius r with center at 0. We will use the convex symmetrization, which is defined in [5]. The convex symmetrization generalizes the Schwarz symmetrization(see [44]). Let us consider a measured function u on $\Omega \subset \mathbb{R}^N$, one dimensional decreasing rearrangement of u is

$$u^\sharp(t) = \sup\{s \geq 0 : |\{x \in \Omega : u(x) \geq s\}| > t\} \quad \text{for} \quad t \in \mathbb{R}. \tag{2.2}$$

The convex symmetrization of u with respect to F is defined as

$$u^\star(x) = u^\sharp(\kappa_N F^o(x)^N) \quad \text{for} \quad x \in \Omega^\star. \tag{2.3}$$

Here $\kappa_N F^o(x)^N$ is just the Lebesgue measure of a homothetic Wulff ball with radius $F^o(x)$ and $\Omega^\star$ is the homothetic Wulff ball centered at the origin having the same

measure as Ω. In [5], the authors proved a P$\acute{o}$lya-Szeg$\ddot{o}$ principle and a comparison result for solutions of the Dirichlet problem for elliptic equations for the convex symmetrization, which generalizes the classical results for Schwarz symmetrization due to Talenti [44].

Now, we give the definition of anisotropic perimeter of a set with respect to F , a co-area formula and an isoperimetric inequality. Precisely, for a a subset $E \subset \Omega$ and a function of bounded variation $u \in BV(\Omega)$, anisotropic bounded variation of u with respect to F is

$$\int_\Omega |\nabla u|_F = \sup\left\{\int_\Omega u \cdot \mathrm{div}\sigma dx : \sigma \in C_0^1(\Omega;\mathbb{R}^N), F^o(\sigma) \le 1\right\}.$$

Define the anisotropic perimeter of E with respect to F as

$$P_F(E) := \int_\Omega |\nabla \chi_E|_F,$$

where χ_E is the characteristic function of the set E. From the reference [20], we have the co-area formula

$$\int_\Omega |\nabla u|_F = \int_0^\infty P_F(|u| > t)dt \tag{2.4}$$

and the isoperimetric inequality

$$P_F(E) \ge N\kappa_N^{1/N}|E|^{1-\frac{1}{N}}. \tag{2.5}$$

We will establish the Lions type concentration-compactness principle [30] for Trudinger-Moser Inequalities under anisotropic Dirichlet norm and L^p norm, which is the extention of Theorem 1.1 in [9] and Lemma 2.3 in [56].

Lemma 2.1. *Suppose $0 \le \gamma < \gamma_1$. Let $\{u_n\} \subset W_0^{1,N}(\Omega)$ be a sequence such that $\|u\|_{N,F,\gamma,p} = 1$, $u_n \rightharpoonup u \not\equiv 0$ weakly in $W_0^{1,N}(\Omega)$. Then for any*

$$0 < q < q_N(u) := (1 - \|u\|_{N,F,\gamma,p}^N)^{-\frac{1}{N-1}}$$

we have

$$\int_\Omega e^{\lambda_N q|u_n|^{\frac{N}{N-1}}} dx < +\infty \tag{2.6}$$

where $\lambda_N = N^{\frac{N}{N-1}}\kappa_N^{\frac{1}{N-1}}$ and κ_N is the volume of a unit Wulff ball. Moreover, this conclusion fails if $p \ge p_N(u)$.

Proof. Since $\|u\|_{N,F,\gamma,p}=1$, we have

$$\lim_{n\to\infty}\int_\Omega F^N(\nabla u_n)=\lim_{n\to\infty}(1+\gamma\|u_n\|_p^N)=1+\gamma\|u\|_p^N.$$

Let

$$v_n=\frac{u_n}{(\int_\Omega F^N(\nabla u_n)dx)^{\frac{1}{N}}},$$

we have $\int_\Omega F^N(\nabla v_n)dx=1$ and $v_n\rightharpoonup v:=u/(1+\gamma\|u\|_p^N)^{\frac{1}{N}}$ weakly in $W_0^{1,N}(\Omega)$. By Lemma 2.3 of [56], it holds

$$\limsup_{n\to\infty}\int_\Omega e^{\lambda_N r|v_n|^{\frac{N}{N-1}}}dx<+\infty \tag{2.7}$$

for any $0<r<(1-\int_\Omega F^N(\nabla v_n)dx)^{-\frac{1}{N-1}}$. Since $q<q_N(u)=(1-\|u\|_{N,F,\gamma,p}^N)^{-\frac{1}{N-1}}$, we have

$$\begin{aligned}\lim_{n\to\infty}q\int_\Omega F^N(\nabla u_n)dx)^{\frac{1}{N-1}}&=q(1+\gamma\|u\|_p^N)^{\frac{1}{N-1}}\\&<\left(\frac{1+\gamma\|u\|_p^N}{1-(\int_\Omega F^N(\nabla u)dx+\gamma\|u\|_p^N)}\right)^{\frac{1}{N-1}}\\&=(1-\int_\Omega F^N(\nabla v_n)dx)^{-\frac{1}{N-1}}\end{aligned} \tag{2.8}$$

Take $r<(1-\int_\Omega F^N(\nabla v_n)dx)^{-\frac{1}{N-1}}$ such that $q\int_\Omega F^N(\nabla u_n)dx)^{\frac{1}{N-1}}<r$ for n large enough. Thus

$$\int_\Omega e^{\lambda_N q|u_n|^{\frac{N}{N-1}}}dx=\int_\Omega e^{\lambda_N q(\int_\Omega F^N(\nabla u_n)dx)^{\frac{1}{N-1}}|v_n|^{\frac{N}{N-1}}}dx<\int_\Omega e^{\lambda_N r|v_n|^{\frac{N}{N-1}}}dx \tag{2.9}$$

for n large enough. The result is followed from (2.7). □

Now, as the similar process of Theorem 2.2 in [43], we give the following estimate involving N-Finsler-Laplacian and L^p norm.

Lemma 2.2. *Assume that $p>1$ and $0\le\gamma<\gamma_1$. Let $f\in L^1(\Omega)$ and $u\in C^1(\bar\Omega)\bigcap W_0^{1,N}(\Omega)$ satisfies*

$$-Q_Nu=f+\gamma\|u\|_p^{N-p}|u|^{p-2}u\quad in\quad\Omega, \tag{2.10}$$

where $Q_Nu=\mathrm{div}(F^{N-1}(\nabla u)F_{\xi_i}(\nabla u))$. Then for any $1<q<N$, $u\in W_0^{1,q}(\Omega)$ and

$$\|u\|_{W_0^{1,q}(\Omega)}\le C(q,N,\gamma,\gamma_1)\|f\|_{L^1(\Omega)}.$$

Proof. Fix $t>0$. Testing (2.10) by $u^t:=\min\{u,t\}\in W_0^{1,N}(\Omega)$ and integrating by parts, we have

$$\begin{aligned}\int_\Omega F^N(\nabla u^t)dx \leq& \int_\Omega fu^t dx+\gamma\int_\Omega \|u^t\|_p^{N-p}|u^t|^p dx\\ \leq& t\|f\|_{L^1(\Omega)}+\frac{\gamma}{\gamma_1}\int_\Omega F^N(\nabla u^t)dx. \end{aligned}\tag{2.11}$$

Thus

$$\int_\Omega F^N(\nabla u^t)dx\leq \frac{\gamma_1}{\gamma_1-\gamma}t\|f\|_{L^1(\Omega)}.\tag{2.12}$$

Denote $\mathcal{W}_r(0)=\{x\in\mathbb{R}^N: F^o(x)\leq r\}$ be a Wulff ball of the same measure as Ω. Let $v^\star$ be the convex symmetrization of u^t with respect to F and $|\mathcal{W}_\rho(0)|=|x\in\mathcal{W}_r(0): v^\star\geq t\}|$. In [5], the authors proved the P*ó*lya-Szeg*ö* principle

$$\int\limits_{\mathcal{W}_r(0)} F^N(\nabla v^\star)dx\leq\int_\Omega F^N(\nabla u^t)dx.$$

Thus

$$\begin{aligned}\inf_{\phi\in W_0^{1,N}(\mathcal{W}_r(0)),\phi|\mathcal{W}_\rho=t}\int\limits_{\mathcal{W}_r(0)} F^N(\nabla\phi)dx\leq& \int\limits_{\mathcal{W}_r(0)} F^N(\nabla v^\star)dx\\ \leq&\frac{\gamma_1}{\gamma_1-\gamma}t\|f\|_{L^1(\Omega)}.\end{aligned}\tag{2.13}$$

On the other hand, the above infimum can be achieved by

$$\phi_0(x)=\begin{cases} t\log\frac{r}{F^o(x)}/\log\frac{r}{\rho} & in \quad \mathcal{W}_r(0)\backslash\mathcal{W}_\rho(0),\\ t & in\quad \mathcal{W}_\rho(0).\end{cases}\tag{2.14}$$

Since $F(\nabla F^o(x))=1$, through the direct computation, we have

$$\begin{aligned}\int\limits_{\mathcal{W}_r(0)} F^N(\nabla\phi_0)dx=&\int_\rho^r F^N(\frac{t}{\log\frac{r}{\rho}}\frac{\nabla F^o(x)}{-s})\int_{\partial\omega_s}\frac{1}{|\nabla F^o(x)|}d\sigma ds\\ =&\int_\rho^r\frac{t^N}{(\log\frac{r}{\rho})^N}\frac{1}{s^N}\int_{\partial\omega_s}\frac{1}{|\nabla F^o(x)|}d\sigma ds\\ =&\int_\rho^r\frac{t^N}{(\log\frac{r}{\rho})^N}\frac{1}{s^N}N\kappa_N s^{N-1}ds\\ =&\frac{N\kappa_N t^N}{(\log\frac{r}{\rho})^{N-1}}.\end{aligned}\tag{2.15}$$

Hence $\frac{N\kappa_N t^N}{(\log\frac{r}{\rho})^{N-1}} \le \frac{\gamma_1}{\gamma_1-\gamma} t\|f\|_{L^1(\Omega)}$, it holds

$$\begin{aligned}
|\{x\in\Omega : u(x)\ge t\}| = |\mathcal{W}_\rho(0)| =& \kappa_N \rho^N \\
\le& \kappa_N r^N \exp(-N(N\kappa_N)^{\frac{1}{N-1}} t(\frac{\gamma_1-\gamma}{\gamma_1}\|f\|_{L^1}^{-1})^{\frac{1}{N-1}}) \\
\le& |\Omega| \exp(-N(N\kappa_N)^{\frac{1}{N-1}} t(\frac{\gamma_1-\gamma}{\gamma_1}\|f\|_{L^1}^{-1})^{\frac{1}{N-1}}) \\
=& : |\Omega| e^{-NC_1 t}.
\end{aligned} \tag{2.16}$$

For every $0 < b < NC_1$,

$$\begin{aligned}
\int_\Omega e^{bu} dx \le& \int_{\{x:u(x)\le 1\}} e^{bu} dx + \int_{\{x:u(x)\ge 1\}} e^{bu} dx \\
\le& e^b|\Omega| + \sum_{k=1}^{\infty} e^{b(k+1)} |\{x\in\Omega : k\le u\le k+1\}| \\
\le& e^b|\Omega| + e^b|\Omega| \sum_{k=1}^{\infty} e^{(b-NC_1)k} \le C(b)|\Omega|.
\end{aligned} \tag{2.17}$$

From (2.12), we have

$$\int_{\{x:u\le t\}} F^N(\nabla u) dx \le \frac{\gamma_1}{\gamma_1-\gamma} t\|f\|_{L^1(\Omega)}. \tag{2.18}$$

Hence,

$$\begin{aligned}
\int_\Omega \frac{F^N(\nabla u)}{1+u^2} dx =& \int_{\{x:u(x)\le 1\}} \frac{F^N(\nabla u)}{1+u^2} dx + \int_{\{x:u(x)\ge 1\}} \frac{F^N(\nabla u)}{1+u^2} dx \\
\le& \int_{\{x:u(x)\le 1\}} F^N(\nabla u) dx + \sum_{m\ge 0} \int_{\{x:2^m\le u(x)\le 2^{m+1}\}} \frac{F^N(\nabla u)}{u^2} dx \\
\le& \int_{\{x:u(x)\le 1\}} F^N(\nabla u) dx + \sum_{m\ge 0} \frac{1}{2^m} \int_{\{x:2^m\le u(x)\le 2^{m+1}\}} \frac{F^N(\nabla u)}{u} dx \\
\le& \frac{\gamma_1}{\gamma_1-\gamma}\|f\|_{L^1(\Omega)} + \sum_{m\ge 0} \frac{1}{2^m} \frac{2\gamma_1}{\gamma_1-\gamma}\|f\|_{L^1(\Omega)} \\
=& \frac{5\gamma_1}{\gamma_1-\gamma}\|f\|_{L^1(\Omega)},
\end{aligned} \tag{2.19}$$

where we have used the estimate (2.18) for $t = 1$ and $t = 2^{m+1}$ in last inequality. For $1 < q < N$, $\frac{q}{N} + \frac{N-q}{N} = 1$, by Young's inequality, we have

$$\begin{aligned}
\int_\Omega F^q(\nabla u) dx =& \int_\Omega \frac{F^q(\nabla u)}{(1+u^2)^{q/N}} (1+u^2)^{q/N} dx \\
\le& \int_\Omega \frac{F^N(\nabla u)}{1+u^2} dx + \int_\Omega (1+u^2)^{q/(N-q)} dx
\end{aligned} \tag{2.20}$$

The desired bound now follows from (2.17) and (2.19). □

3 Maximizers of the subcritical case

In this section, we will show the existence of the maximizers for Trudinger-Moser in the subcritical case.

Lemma 3.1. *Let $\Omega \subset \mathbb{R}^N (N \geq 2)$ be a smooth bounded domain. Then for any $\epsilon \in (0, \lambda_N)$, the supremum*

$$\Lambda_{\gamma,\epsilon} = \sup_{u \in W_0^{1,N}(\Omega), \|u\|_{N,F,\gamma,p} \leq 1} \int_\Omega e^{(\lambda_N - \epsilon)|u|^{\frac{N}{N-1}}} dx \tag{3.1}$$

can be attained by $u_\epsilon \in W_0^{1,N}(\Omega) \bigcap C^1(\overline{\Omega})$ with $\|u_{\epsilon,n}\|_{N,F,\gamma,p} = 1$. In the distributional sense, u_ϵ satisfies the following equation

$$\begin{cases} -Q_N u_\epsilon - \gamma \|u_\epsilon\|_p^{N-p} u_\epsilon^{p-1} = \frac{1}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N - \epsilon) u_\epsilon^{\frac{N}{N-1}}} & in \quad \Omega, \\ u_\epsilon = 0 \qquad on \quad \partial\Omega, \\ \lambda_\epsilon = \int_\Omega u_\epsilon^{\frac{N}{N-1}} e^{(\lambda_N - \epsilon) u_\epsilon^{\frac{N}{N-1}}} dx. \end{cases} \tag{3.2}$$

Moreover,

$$\liminf_{\epsilon \to 0} \lambda_\epsilon > 0$$

and

$$\liminf_{\epsilon \to 0} \Lambda_{\gamma,\epsilon} = \Lambda_\gamma. \tag{3.3}$$

Proof. Let $u_{\epsilon,n}$ be a maximizing sequence for $\Lambda_{\gamma,\epsilon}$, *i.e.*, $u_{\epsilon,n} \in W_0^{1,N}(\Omega)$, $\|u_{\epsilon,n}\|_{N,F,\gamma,p} \leq 1$ and

$$\int_\Omega e^{(\lambda_N - \epsilon)|u_{\epsilon,n}|^{\frac{N}{N-1}}} dx \to \Lambda_{\gamma,\epsilon} \tag{3.4}$$

as $n \to \infty$. Since $\gamma < \gamma_1$, thus

$$(1 - \frac{\gamma}{\gamma_1}) \int_\Omega F^N(\nabla u_{\epsilon,n}) dx \leq \int_\Omega F^N(\nabla u_{\epsilon,n}) dx - \gamma \|u_{\epsilon,n}\|_p^N \leq 1,$$

which lead to $u_{\epsilon,n}$ is bounded in $W_0^{1,N}(\Omega)$, so there exists some $u_\epsilon \in W_0^{1,N}(\Omega)$ such that up to a subsequence, $u_{\epsilon,n} \rightharpoonup u_\epsilon$ weakly in $W_0^{1,N}(\Omega)$, $u_{\epsilon,n} \to u_\epsilon$ strongly in $L^q(\Omega)$ for any $q \geq 1$, and $u_{\epsilon,n} \to u_\epsilon$ a. e. in Ω. We claim that $u_\epsilon \not\equiv 0$. If otherwise, then

$\limsup_{n\to\infty}\int_\Omega F^N(\nabla u_{\epsilon,n})dx \le 1$. The anisotropic Trudinger-Moser inequality implies $e^{(\lambda_N-\epsilon)|u_{\epsilon,n}|^{\frac{N}{N-1}}}$ is uniformly bounded in $L^s(\Omega)$ for any $1 < s < \frac{\lambda_N}{\lambda_N-\epsilon}$. Thus

$$\Lambda_{\gamma,\epsilon} = \lim_{n\to\infty}\int_\Omega e^{(\lambda_N-\epsilon)|u_{\epsilon,n}|^{\frac{N}{N-1}}}dx = |\Omega|, \tag{3.5}$$

which is impossible. So $u_\epsilon \not\equiv 0$ and Lemma 2.1 implies $\int_\Omega e^{(\lambda_N-\epsilon)|u_{\epsilon,n}|^{\frac{N}{N-1}}}dx$ in $L^s(\Omega)$ for some $s > 1$. Consequently, we have

$$\Lambda_{\gamma,\epsilon} = \lim_{n\to\infty}\int_\Omega e^{(\lambda_N-\epsilon)|u_{\epsilon,n}|^{\frac{N}{N-1}}}dx = \int_\Omega e^{(\lambda_N-\epsilon)|u_\epsilon|^{\frac{N}{N-1}}}dx, \tag{3.6}$$

Then u_ϵ attains the supremum. We claim $\|u_\epsilon\|_{N,F,\gamma,p} = 1$. In fact, if $\|u_\epsilon\|_{N,F,\gamma,p} < 1$, then

$$\Lambda_{\gamma,\epsilon} = \int_\Omega e^{(\lambda_N-\epsilon)|u_\epsilon|^{\frac{N}{N-1}}}dx < \int_\Omega e^{(\lambda_N-\epsilon)|\frac{u_\epsilon}{\|u_\epsilon\|_{N,F,\gamma,p}}|^{\frac{N}{N-1}}}dx \le \Lambda_{N,\epsilon}, \tag{3.7}$$

which is a contradiction. Furthermore, we know that u_ϵ satisfies the Euler–Lagrange equation in distributional sense. By regularity theory obtained in [28], we have $u_\epsilon \in C^1(\overline{\Omega})$.

Since $e^t \le 1 + te^t$ for $t \ge 0$, we have

$$\begin{aligned}
&\int_\Omega e^{(\lambda_N-\epsilon)|u_\epsilon|^{\frac{N}{N-1}}}dx \\
\le&|\Omega| + (\lambda_N-\epsilon)\int_\Omega |u_\epsilon|^{\frac{N}{N-1}} e^{(\lambda_N-\epsilon)|u_\epsilon|^{\frac{N}{N-1}}}dx \\
=&|\Omega| + (\lambda_N-\epsilon)\lambda_\epsilon.
\end{aligned}$$

This leads to $\liminf_{\epsilon\to 0}\lambda_\epsilon > 0$.

Obviously, $\limsup_{\epsilon\to 0}\Lambda_{\gamma,\epsilon} \le \Lambda_\gamma$. On the other hand, for any $u \in W_0^{1,N}(\Omega)$ with $\|u_\epsilon\|_{N,F,\gamma,p} \le 1$, by Fatou's lemma, we have

$$\int_\Omega e^{\lambda_N|u|^{\frac{N}{N-1}}}dx \le \liminf_{\epsilon\to 0}\int_\Omega e^{(\lambda_N-\epsilon)|u_\epsilon|^{\frac{N}{N-1}}}dx \le \liminf_{\epsilon\to 0}\Lambda_{\gamma,\epsilon},$$

which implies that $\liminf_{\epsilon\to 0}\Lambda_{\gamma,\epsilon} \ge \Lambda_\gamma$. Thus $\liminf_{\epsilon\to 0}\Lambda_{\gamma,\epsilon} = \Lambda_\gamma$. □

4 Maximizers of the critical case

In this section, by using blow-up analysis, we analyze the behavior of the maximizers u_ϵ in section 3. Since u_ϵ is bounded in $W_0^{1,N}(\Omega)$, up to a subsequence, we can assume $u_\epsilon \rightharpoonup u_0$ weakly in $W_0^{1,N}(\Omega)$, $u_\epsilon \to u_0$ strongly in $L^q(\Omega)$ for any $q \ge 1$, and $u_\epsilon \to u_0$ a.e. in Ω as $\epsilon \to 0$.

4.1 Blow-up analysis

Let $c_\epsilon = \max_{\overline{\Omega}} u_\epsilon = u_\epsilon(x_\epsilon)$. If c_ϵ is bounded, then for any $u \in W_0^{1,N}(\Omega)$ with $\|u\|_{N,F,\gamma,p} \leq 1$, by Lebesgue dominated convergence theorem, we have

$$\begin{aligned}\int_\Omega e^{\lambda_N |u|^{\frac{N}{N-1}}} dx &= \lim_{\epsilon\to 0} \int_\Omega e^{(\lambda_N-\epsilon)|u|^{\frac{N}{N-1}}} dx \\ &\leq \lim_{\epsilon\to 0} \int_\Omega e^{(\lambda_N-\epsilon)|u_\epsilon|^{\frac{N}{N-1}}} dx \\ &= \int_\Omega e^{\lambda_N |u_0|^{\frac{N}{N-1}}} dx \end{aligned} \tag{4.1}$$

Therefore u_0 is the desired maximizer. Moreover, $u_\epsilon \to u_0$ and $u_0 \in C^1(\overline{\Omega})$ by standard elliptic regularity theory. In the following, we consider another case, we assume $c_\epsilon \to +\infty$ and $x_\epsilon \to x_0$ as $\epsilon \to 0$. We assume $x_0 \in \Omega$, at the end of this section, we shall exclude the case $x_0 \in \partial\Omega$.

Lemma 4.1. *$u_0 \equiv 0$ and $F^N(\nabla u_\epsilon)dx \rightharpoonup \delta_{x_0}$ weakly in the sense of measure as $\epsilon \to 0$, where δ_{x_0} is the Dirac measure at x_0.*

Proof. Suppose $u_0 \not\equiv 0$. Notice that $\liminf_{\epsilon\to 0} \lambda_\epsilon > 0$, by Lemma 2.1 and Hölder inequality, we have $\frac{1}{\lambda_\epsilon} u_\epsilon |u_\epsilon|^{\frac{2-N}{N-1}} e^{(\lambda_N-\epsilon)|u_\epsilon|^{\frac{N}{N-1}}}$ is uniformly bounded in $L^q(\Omega)$ for some $q > 1$. Then by Lemma 2.2 in [56], u_ϵ is uniformly bounded in Ω, which contradicts $c_\epsilon \to +\infty$ as $\epsilon \to 0$. Hence $u_0 \equiv 0$.

Notice that $\int_\Omega F^N(\nabla u_\epsilon)dx = 1 + \gamma\|u_\epsilon\|_p^N \to 1$. If $F^N(\nabla u_\epsilon)dx \rightharpoonup \mu \neq \delta_{x_0}$ in the sense of measure as $\epsilon \to 0$, then there exists $\theta < 1$ and $r > 0$ small enough such that

$$\lim_{\epsilon\to 0} \int\limits_{\mathcal{W}_r(x_0)} F^N(\nabla u_\epsilon)dx < \theta,$$

Consider the cut-off function $\phi \in C_0^\infty(\Omega)$, which is supported in $\mathcal{W}_r(x_0)$ for some $r > 0$, $0 \leq \phi \leq 1$ and $\phi = 1$ in $\mathcal{W}_{\frac{r}{2}}(x_0)$. Since $u_\epsilon \to 0$ in $L^q(\Omega)$ for any $q > 1$, we have

$$\limsup_{\epsilon\to 0} \int_{\mathcal{W}_r(x_0)} F^N(\nabla(\phi u_\epsilon)dx \leq \lim_{\epsilon\to 0} \int_{\mathcal{W}_r(x_0)} F^N(\nabla u_\epsilon dx < \theta.$$

For sufficiently small $\epsilon > 0$, using anisotropic Moser-Trudinger inequality to ϕu_ϵ,we have $e^{\lambda_N u_\epsilon^{\frac{N}{N-1}}} dx$ is uniformly bounded in $L^s(\mathcal{W}_{\frac{r}{2}}(x_0))$ for some $s > 1$. From Lemma 2.2 in [56], u_ϵ is uniformly bounded in $\mathcal{W}_{\frac{r}{2}}(x_0)$, which contradicts to $c_\epsilon \to +\infty$. Hence $F^N(\nabla u_\epsilon)dx \rightharpoonup \delta_0$ as $\epsilon \to 0$. □

Let

$$r_\epsilon^N = \lambda_\epsilon c_\epsilon^{-N/(N-1)} e^{-(\lambda_N-\epsilon)c_\epsilon^{N/(N-1)}}. \tag{4.2}$$

Denote $\Omega_\epsilon = \{x \in \mathbb{R}^N : x_\epsilon + r_\epsilon x \in \Omega\}$. Define

$$\psi_\epsilon(x) = c_\epsilon^{-1} u_\epsilon(x_\epsilon + r_\epsilon x) \tag{4.3}$$

and

$$\varphi_\epsilon(x) = c_\epsilon^{N/(N-1)}(\psi_\epsilon(x) - 1) = c_\epsilon^{1/(N-1)}(u_\epsilon(x_\epsilon + r_\epsilon x) - c_\epsilon). \tag{4.4}$$

We have the following:

Lemma 4.2. *Let $r_\epsilon, \psi_\epsilon(x)$ and φ_ϵ be defined as in (4.2)-(5.3). Then*

(i) there hold $r_\epsilon \to 0$ as $\epsilon \to 0$;

(ii) $\psi_\epsilon(x) \to 1$ in $C^1_{loc}(\mathbb{R}^N)$;

(iii) $\varphi_\epsilon(x) \to \varphi(x)$ in $C^1_{loc}(\mathbb{R}^N)$, where

$$\varphi(x) = -\frac{N-1}{\lambda_N} \log\left(1 + \kappa_N^{\frac{1}{N-1}} F^o(x)^{\frac{N}{N-1}}\right). \tag{4.5}$$

Moreover,

$$\int_{\mathbb{R}^N} e^{\frac{N}{N-1}\lambda_N \varphi} dx = 1. \tag{4.6}$$

Proof. (i) From (4.2), we have

$$\begin{aligned} r_\epsilon^N c_\epsilon^{N/(N-1)} =& \lambda_\epsilon e^{-(\lambda_N-\epsilon)c_\epsilon^{N/(N-1)}} \\ =& \int_\Omega |u_\epsilon|^{\frac{N}{N-1}} e^{(\lambda_N-\epsilon)|u_\epsilon|^{\frac{N}{N-1}}} dx \cdot e^{-(\lambda_N-\epsilon)c_\epsilon^{N/(N-1)}} \\ \le& c_\epsilon^{N/(N-1)} e^{(\frac{\lambda_N}{2}-\epsilon)c_\epsilon^{N/(N-1)}} \int_\Omega e^{\frac{\lambda_N}{2}|u_\epsilon|^{\frac{N}{N-1}}} dx \cdot e^{-(\lambda_N-\epsilon)c_\epsilon^{N/(N-1)}} \\ \le& C c_\epsilon^{N/(N-1)} e^{-\frac{\lambda_N}{2}c_\epsilon^{N/(N-1)}} \to 0, \end{aligned}$$

as $\epsilon \to 0$. Thus $\lim\limits_{\epsilon\to 0} r_\epsilon = 0$.

(ii) By a direction calculation, $0 \le \psi_\epsilon \le 1$ and ψ_ϵ is a weak solution to

$$\begin{aligned} &-\operatorname{div}(F^{N-1}(\nabla\psi_\epsilon)F_\xi(\nabla\psi_\epsilon)) \\ &= c_\epsilon^{-N} e^{(\lambda_N-\epsilon)c_\epsilon^{N/(N-1)}(\psi_\epsilon^{\frac{N}{N-1}}-1)} \psi_\epsilon^{\frac{1}{N-1}} + \gamma c_\epsilon^{p-N} r_\epsilon^N \|u_\epsilon\|_p^{N-p} \psi_\epsilon^{p-1} \qquad in \quad \Omega_\epsilon. \end{aligned}$$

Since $c_\epsilon^{-N} e^{(\lambda_N-\epsilon)c_\epsilon^{N/(N-1)}(\psi_\epsilon^{\frac{N}{N-1}}-1)}\psi_\epsilon^{\frac{1}{N-1}} \le c_\epsilon^{-N} \to 0$ as $\epsilon \to 0$, and

$$\begin{aligned}&\left(\int\limits_{B_{r_\epsilon^{-1}}} (c_\epsilon^{p-N} r_\epsilon^N \|u_\epsilon\|_p^{N-p}\psi_\epsilon^{p-1})^{p/(p-1)}dx\right)^{\frac{p-1}{p}}\\ =&c_\epsilon^{1-N} r_\epsilon^{\frac{N}{p}} \|u_\epsilon\|_p^{N-1} \to 0 \quad as \quad \epsilon \to 0.\end{aligned}$$

Thus we obtain that $\mathrm{div}(F^{N-1}(\nabla\psi_\epsilon)F_\xi(\nabla\psi_\epsilon))$ is bounded in $L^{p/(p-1)}(B_{r_\epsilon^{-1}})$. Using elliptic regularity theory (see[47]), we have $\psi_\epsilon \to \psi$ in $C^1_{loc}(\mathbb{R}^N)$, where ψ is a weak solution to the equation

$$-\mathrm{div}(F^{N-1}(\nabla\psi)F_\xi(\nabla\psi)) = 0 \quad in \quad \mathbb{R}^N.$$

The Liouville theorem (see [21]) implies $\psi \equiv 1$.

(iii) We have

$$\begin{aligned}&-\mathrm{div}(F^{N-1}(\nabla\varphi_\epsilon)F_\xi(\nabla\varphi_\epsilon))\\ &= e^{(\lambda_N-\epsilon)c_\epsilon^{N/(N-1)}(\psi_\epsilon^{\frac{N}{N-1}}-1)}\psi_\epsilon^{\frac{1}{N-1}} + \gamma c_\epsilon^p r_\epsilon^N \|u_\epsilon\|_p^{N-p}\psi_\epsilon^{p-1} \qquad in \quad \Omega_\epsilon.\end{aligned}$$

When $p > N$, for $R > 0$ and sufficiently small ϵ,

$$\begin{aligned}\|u_\epsilon\|_p^{N-p} \le&\left(\int\limits_{B_{Rr_\epsilon}} u_\epsilon^p dx\right)^{N/P-1}\\ =&c_\epsilon^{N-p} r_\epsilon^{N^2/p-N}\left(\int\limits_{B_R}\psi_\epsilon^p dx\right)^{N/P-1}\end{aligned}$$

Since $0 \le \psi_\epsilon \le 1$, $\psi_\epsilon \to 1$ in $C^1_{loc}(\mathbb{R}^N)$, we have $\int\limits_{B_R}\psi^p dx > 0$. Thus

$$c_\epsilon^p r_\epsilon^N \|u_\epsilon\|_p^{N-p}\psi_\epsilon^{p-1} \le 2\left(\int\limits_{B_R}\psi_\epsilon^p dx\right)^{N/P-1} c_\epsilon^N r_\epsilon^{N^2/p} \to 0 \quad as \quad \epsilon \to 0.$$

When $1 < p \le N$, obviously, $c_\epsilon^p r_\epsilon^N \|u_\epsilon\|_p^{N-p}\psi_\epsilon^{p-1} \to 0$ as $\epsilon \to 0$. From regularity theory in [47], up to a subsequence, there exists $\varphi \in C^1(\mathbb{R}^N)$ such that $\varphi_\epsilon \to \varphi$ in

$C^1_{loc}(\mathbb{R}^N)$. From the definition of ψ_ϵ and φ_ϵ, we have

$$\begin{aligned} u_\epsilon(x_\epsilon + r_\epsilon x)^{\frac{N}{N-1}} - c_\epsilon^{\frac{N}{N-1}} &= c_\epsilon^{\frac{N}{N-1}} \left(\psi_\epsilon(x)^{\frac{N}{N-1}} - 1 \right) \\ &= c_\epsilon^{\frac{N}{N-1}} \left((1 + \psi_\epsilon(x) - 1)^{\frac{N}{N-1}} - 1 \right) \\ &= \frac{N}{N-1} \varphi_\epsilon(x) + c_\epsilon^{\frac{N}{N-1}} O((\psi_\epsilon(x) - 1)^2) \\ &= \frac{N}{N-1} \varphi_\epsilon(x) + O(\psi_\epsilon(x) - 1). \end{aligned}$$

By $\epsilon \to 0$, φ satisfies

$$\begin{cases} -\mathrm{div}(F^{N-1}(\nabla\varphi) F_\xi(\nabla\varphi)) = e^{\frac{N}{N-1}\lambda_N \varphi} & in \quad \mathbb{R}^N, \\ \varphi(x) \le \varphi(0) = 0. \end{cases}$$

Combing (4.2) and (4.3), for any $R > 0$, we have

$$\begin{aligned} \int\limits_{\mathcal{W}_R(0)} e^{\frac{N}{N-1}\lambda_N \varphi} dx &= \lim_{\epsilon \to 0} \int\limits_{\mathcal{W}_R(0)} e^{(\lambda_N - \epsilon)(u_\epsilon(x_\epsilon + r_\epsilon x)^{\frac{N}{N-1}} - c_\epsilon^{\frac{N}{N-1}})} dx \\ &= \lim_{\epsilon \to 0} \lambda_\epsilon^{-1} \int\limits_{\mathcal{W}_R r_\epsilon(x_\epsilon)} c_\epsilon^{N/(N-1)} e^{(\lambda_N - \epsilon) u_\epsilon(y)^{\frac{N}{N-1}}} dy \\ &= \lim_{\epsilon \to 0} \lambda_\epsilon^{-1} \int\limits_{\mathcal{W}_R r_\epsilon(x_\epsilon)} u_\epsilon(y)^{N/(N-1)} e^{(\lambda_N - \epsilon) u_\epsilon(y)^{\frac{N}{N-1}}} dy \\ &\le 1. \end{aligned}$$

This leads to

$$\int\limits_{\mathbb{R}^N} e^{\frac{N}{N-1}\lambda_N \varphi} dx \le 1.$$

From co-area formula (2.4) and isoperimetric inequality (2.5), through a simple computation, it follows from Höder inequlity, we have $\int_{\mathbb{R}^N} e^{\frac{N}{N-1}\lambda_N \varphi} dx \ge 1$. Thus

$$\int_{\mathbb{R}^N} e^{\frac{N}{N-1}\lambda_N \varphi} dx = 1, \tag{4.7}$$

which implies φ is symmetric with respect to F^o, i.e., $\varphi(x) = \varphi(F^o(x))$(see [48], Prop. 6.1). Thus we get

$$\varphi(x) = -\frac{N-1}{\lambda_N} \log \left(1 + \kappa_N^{\frac{1}{N-1}} F^o(x)^{\frac{N}{N-1}} \right). \tag{4.8}$$

□

Denote $\mathcal{W}_R(x_\epsilon)$ be a Wulff ball of radius R with center at x_ϵ. For any $0<a<1$, use the notation

$$u_{\epsilon,a}=\min\{u_\epsilon, ac_\epsilon\}.$$

Then we have the following:

Lemma 4.3. *For any $0<a<1$, there holds*

$$\lim_{\epsilon\to 0}\int_\Omega F^N(\nabla u_{\epsilon,a})dx=a.$$

Proof. Testing the equation (3.3), we have

$$\begin{aligned}\int_\Omega F^N(\nabla u_{\epsilon,a})dx=&\int_\Omega F^{N-1}(\nabla u_\epsilon)F_\xi(\nabla u_\epsilon)\cdot u_{\epsilon,a}dx\\ =&-\int_\Omega \mathrm{div}(F^{N-1}(\nabla u_\epsilon)F_\xi(\nabla u_\epsilon))u_{\epsilon,a}dx\\ =&\int_\Omega \frac{1}{\lambda_\epsilon}u_\epsilon^{\frac{1}{N-1}}e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}u_{\epsilon,a}dx+\gamma\int_\Omega \|u_\epsilon\|_p^{N-p}u_\epsilon^{p-1}u_{\epsilon,a}dx.\end{aligned}$$

For any $R>0$, we have $\mathcal{W}_R(x_\epsilon)\subset\{u_\epsilon>ac_\epsilon\}$ for $\epsilon>0$ small enough. Thus

$$\begin{aligned}\int_\Omega F^N(\nabla u_{\epsilon,a})dx=&\int_{\{u_\epsilon\le ac_\epsilon\}}\frac{1}{\lambda_\epsilon}u_\epsilon^{\frac{1}{N-1}}e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}u_{\epsilon,a}dx\\ &+\int_{\{u_\epsilon>ac_\epsilon\}}\frac{1}{\lambda_\epsilon}u_\epsilon^{\frac{1}{N-1}}e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}u_{\epsilon,a}dx+\gamma\int_\Omega \|u_\epsilon\|_p^{N-p}u_\epsilon^{p-1}u_{\epsilon,a}dx\\ >&ac_\epsilon\int\limits_{\mathcal{W}_R(x_\epsilon)}\frac{1}{\lambda_\epsilon}u_\epsilon^{\frac{1}{N-1}}e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}u_{\epsilon,a}dx+o_\epsilon(1).\end{aligned}$$

Set $x=x_\epsilon+r_\epsilon y$, we have

$$\begin{aligned}&ac_\epsilon\int\limits_{\mathcal{W}_R(x_\epsilon)}\frac{1}{\lambda_\epsilon}u_\epsilon^{\frac{1}{N-1}}e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}u_{\epsilon,a}dx\\ =&a\int\limits_{\mathcal{W}_R(0)}\psi_\epsilon(y)^{\frac{1}{N-1}}e^{(\lambda_N-\epsilon)c_\epsilon^{\frac{N}{N-1}}(\psi_\epsilon(y)^{\frac{N}{N-1}}-1)}dy\to\int\limits_{\mathcal{W}_R(0)}e^{\frac{N}{N-1}\lambda_N\varphi}dy.\end{aligned}$$

Letting $R\to\infty$, we derive that

$$\liminf_{\epsilon\to 0}\int_\Omega F^N(\nabla u_{\epsilon,a})dx\ge a.$$

Similarly, we choose $(u_\epsilon-u_{\epsilon,a})$ as a test function of (3.3), we obtain $\liminf\limits_{\epsilon\to 0}\int_\Omega F^N(\nabla(u_\epsilon-u_{\epsilon,a}))\ge 1-a$. Notice that

$$\begin{aligned}\int_\Omega F^N(\nabla u_{\epsilon,a})dx=&\int_\Omega F^N(\nabla u_\epsilon)dx-\int_\Omega F^N(\nabla(u_\epsilon-u_{\epsilon,a}))dx\\ =&1+\gamma\|u_\epsilon\|_p^N-\int_\Omega F^N(\nabla(u_\epsilon-u_{\epsilon,a}))dx.\end{aligned}$$

which implies

$$\limsup_{\epsilon\to 0}\int_\Omega F^N(\nabla u_{\epsilon,a})dx \le a.$$

We have finished the proof of the lemma. □

Lemma 4.4. *There holds*

$$\lim_{\epsilon\to 0}\int_\Omega e^{(\lambda_N-\epsilon)|u_\epsilon|^{\frac{N}{N-1}}}dx \le |\Omega| + \limsup_{\epsilon\to 0}\frac{\lambda_\epsilon}{c_\epsilon^{N/(N-1)}}.$$

Proof. For any $0<a<1$, by Lemma 4.3, we have $\lim_{\epsilon\to 0}\int_\Omega F^N(\nabla u_{\epsilon,a})dx = a < 1$. From anisotropic Trudinger-Moser inequality, $e^{\lambda_N u_{\epsilon,a}^{\frac{N}{N-1}}}$ is bounded in $L^q(\Omega)$ for some $q>1$. Notice that $u_{\epsilon,a}\to 0$ a.e. in Ω, which implies $\lim_{\epsilon\to 0}\int_\Omega e^{\lambda_N u_{\epsilon,a}^{\frac{N}{N-1}}} = |\Omega|$. Hence

$$\begin{aligned}
e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}} &= \int\limits_{\{u_\epsilon\le ac_\epsilon\}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}dx + \int\limits_{\{u_\epsilon> ac_\epsilon\}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}dx\\
&\le \int\limits_{\{u_\epsilon\le ac_\epsilon\}} e^{(\lambda_N-\epsilon)u_{\epsilon,a}^{\frac{N}{N-1}}}dx + \frac{1}{(ac_\epsilon)^{\frac{N}{N-1}}}\int\limits_{\{u_\epsilon> ac_\epsilon\}} u_\epsilon^{\frac{N}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}dx\\
&\le \int_\Omega e^{(\lambda_N-\epsilon)u_{\epsilon,a}^{\frac{N}{N-1}}}dx + \frac{\lambda_\epsilon}{(ac_\epsilon)^{\frac{N}{N-1}}}.
\end{aligned}$$

Letting $\epsilon\to 0$ and $a\to 1$, we conclude our proof. □

As a consequence of Lemma 4.4, for any $\theta < \frac{N}{N-1}$, there holds

$$\lim_{\epsilon\to 0}\frac{\lambda_\epsilon}{c_\epsilon^\theta} = +\infty. \tag{4.9}$$

If otherwise, $\frac{\lambda_\epsilon}{c_\epsilon^{N/(N-1)}}\to 0$ as $\epsilon\to 0$, by Lemma 4.4, we have $\lim_{\epsilon\to 0}\int_\Omega e^{(\lambda_N-\epsilon)|u_\epsilon|^{\frac{N}{N-1}}}dx \le |\Omega|$, which is impossible.

Lemma 4.5. *For any $1<q<N$, $c_\epsilon^{\frac{1}{N-1}}u_\epsilon \rightharpoonup G$ in $W_0^{1,q}(\Omega)$, where G is a distributional solution to*

$$\begin{cases} -\mathrm{div}(F^{N-1}(\nabla G)F_\xi(\nabla G)) = \delta_{x_0} + \gamma\|G\|_p^{N-p}G^{p-1} & \text{in}\quad \Omega,\\ G = 0 \qquad \text{on}\quad \partial\Omega. \end{cases} \tag{4.10}$$

Furthermore, $c_\epsilon^{\frac{1}{N-1}}u_\epsilon\to G$ in $C^1_{loc}(\overline{\Omega}\backslash\{x_0\})$, and G has the form

$$G(x) = -\frac{1}{(N\kappa_N)^{\frac{1}{N-1}}}\log F^o(x-x_0) + A_{x_0} + \xi(x), \tag{4.11}$$

where A_{x_0} is a constant depending only on x_0, $\xi \in C^0(\overline{\Omega}) \bigcap C^1_{loc}(\Omega\backslash\{x_0\})$ and $\xi(x) = O(F^o(x-x_0))$ as $x \to x_0$.

Proof. Firstly, we claim for any function $\phi \in C(\bar{\Omega})$, it holds

$$\lim_{\epsilon\to 0}\int_\Omega \frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}\phi dx = \phi(x_0). \tag{4.12}$$

In fact, for any $0<a<1$ and $R>0$, we have $\mathcal{W}_{Rr_\epsilon}(x_\epsilon) \subset \{u_\epsilon > ac_\epsilon\}$ as $\epsilon>0$ small enough, we denote

$$\begin{aligned}
&\int_\Omega \frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}\phi dx\\
=&\int\limits_{\{u_\epsilon\le ac_\epsilon\}} \frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}\phi dx + \int\limits_{\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} \frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}\phi dx\\
&+\int\limits_{\{u_\epsilon>ac_\epsilon\}\backslash\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} \frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}\phi dx\\
:=&I+II+III.
\end{aligned}$$

By Lemma 4.3 and (4.10), we have

$$\begin{aligned}
I=&\frac{c_\epsilon}{\lambda_\epsilon}\int_{\{u_\epsilon\le ac_\epsilon\}} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}\phi dx\\
\le&\frac{c_\epsilon}{\lambda_\epsilon}\int_\Omega u_{\epsilon,a}^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_{\epsilon,a}^{\frac{N}{N-1}}}\phi dx\\
=&o_\epsilon(1)O(a).
\end{aligned}$$

Making the change in variable $x = x_\epsilon + r_\epsilon y$, since $\phi(x_\epsilon + r_\epsilon\cdot) \to \phi(x_0)$ uniformly in $\mathcal{W}_R(0)$. Note $r_\epsilon^N = \lambda_\epsilon c_\epsilon^{-N/(N-1)} e^{-(\lambda_N-\epsilon)c_\epsilon^{N/(N-1)}}$, together with (4.7), we have

$$\begin{aligned}
II=&\int\limits_{\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} \frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}\phi dx\\
=&\int\limits_{\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} c_\epsilon r_\epsilon^{-N} c_\epsilon^{-N/(N-1)} e^{-(\lambda_N-\epsilon)c_\epsilon^{N/(N-1)}} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}\phi dx\\
=&\int\limits_{\mathcal{W}_R(0)} c_\epsilon r_\epsilon^{-N} c_\epsilon^{-N/(N-1)} c_\epsilon^{1/(N-1)} \psi_\epsilon(y)^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)c_\epsilon^{\frac{N}{N-1}}(\psi_\epsilon(y)^{\frac{N}{N-1}}-1)}\phi(x_\epsilon+r_\epsilon y) r_\epsilon^N dy\\
=&\int\limits_{\mathcal{W}_R(0)} \psi_\epsilon(y)^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)c_\epsilon^{\frac{N}{N-1}}(\psi_\epsilon(y)^{\frac{N}{N-1}}-1)}\phi(x_\epsilon+r_\epsilon y) dy\\
\to&\phi(x_0)
\end{aligned}$$

as $\epsilon \to 0$ and $R \to \infty$.

$$
\begin{aligned}
III = & \int\limits_{\{u_\epsilon > ac_\epsilon\}\backslash \mathcal{W}_{Rr_\epsilon}(x_\epsilon)} \frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}} \phi dx \\
\leq & \|\phi\|_\infty \left(\int\limits_{\{u_\epsilon > ac_\epsilon\}} \frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}} dx - \int\limits_{\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} \frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}} dx \right) \\
\leq & \|\phi\|_\infty \left(\frac{1}{a} \int\limits_{\{u_\epsilon > ac_\epsilon\}} \frac{u_\epsilon^{\frac{N}{N-1}}}{\lambda_\epsilon} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}} dx - \int\limits_{\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} \frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}} dx \right) \\
\leq & \|\phi\|_\infty \left(\frac{1}{a} - \int\limits_{\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} \frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}} dx \right).
\end{aligned}
$$

Thus

$$
\lim_{a\to 1} \lim_{R\to\infty} \lim_{\epsilon\to 0} \int\limits_{\{u_\epsilon > ac_\epsilon\}\backslash \mathcal{W}_{Rr_\epsilon}(x_\epsilon)} \frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}} \phi dx = 0.
$$

Combing the above discussion, we have proved the claim. From (3.3), we have

$$
-Q_N(c_\epsilon^{\frac{1}{N-1}} u_\epsilon) = \frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}} + \gamma \|c_\epsilon^{\frac{1}{N-1}} u_\epsilon\|_p^{N-p} (c_\epsilon^{\frac{1}{N-1}} u_\epsilon)^{p-1}. \tag{4.13}
$$

It follows from (4.12) that $\frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}}$ is bounded in $L^1(\Omega)$. From Lemma 2.2, we know that $c_\epsilon^{\frac{1}{N-1}} u_\epsilon$ is bounded in $W_0^{1,q}(\Omega)$ for any $1 < q < N$. Hence there exists some $G \in W_0^{1,q}(\Omega)$ such that $c_\epsilon^{\frac{1}{N-1}} u_\epsilon \rightharpoonup G$ in $W_0^{1,q}(\Omega)$ for any $1 < q < N$. Multiplying (4.13) with $\phi \in C_0^\infty(\Omega)$, we get

$$
\begin{aligned}
-\int_\Omega \phi Q_N(c_\epsilon^{\frac{1}{N-1}} u_\epsilon) dx = & \int_\Omega \phi \frac{c_\epsilon}{\lambda_\epsilon} u_\epsilon^{\frac{1}{N-1}} e^{(\lambda_N-\epsilon)u_\epsilon^{\frac{N}{N-1}}} dx \\
& + \gamma \|c_\epsilon^{\frac{1}{N-1}} u_\epsilon\|_p^{N-p} \int_\Omega \phi (c_\epsilon^{\frac{1}{N-1}} u_\epsilon)^{p-1} dx.
\end{aligned}
$$

Let $\epsilon \to 0$, using again (4.12), we have

$$
\int_\Omega \nabla\phi F^{N-1}(\nabla G) F_\xi(\nabla G) dx = \phi(x_0) + \gamma \|G\|_p^{N-p} \int_\Omega \phi G^{p-1} dx.
$$

Therefore, in the distributional sense,

$$
-\mathrm{div}(F^{N-1}(\nabla G) F_\xi(\nabla G)) = \delta_{x_0} + \gamma \|G\|_p^{N-p} G^{p-1} \quad \text{in} \quad \Omega.
$$

Applying with the standard elliptic regularity theory in [47], we get $c_\epsilon^{\frac{1}{N-1}} u_\epsilon \to G$ in $C^1_{loc}(\overline{\Omega}\backslash\{x_0\})$. Finally, replacing the right hand term in equation (4.11), we use

the similar discussion as Lemma 4.7 in [57], the asymptotic representation of Green function can immediateiy derived. This complete the proof of Lemma 4.5. □

Next, we will exclude the case $x_0 \in \partial\Omega$. Denote $d_\epsilon = dist(x_\epsilon, \partial\Omega)$ and r_ϵ be defined by (4.2). We obtain

Lemma 4.6. *If $0 \le \gamma < \gamma_1$ and $x_0 \in \partial\Omega$. Then $\frac{r_\epsilon}{d_\epsilon} \to 0$ as $\epsilon \to 0$. Moreover, $c_\epsilon^{\frac{1}{N-1}} u_\epsilon \rightharpoonup 0$ weakly in $W_0^{1,q}(\Omega)(1 < q < N)$ and $c_\epsilon^{\frac{1}{N-1}} u_\epsilon \to 0$ strongly in $C^1(\overline{\Omega}\backslash\{x_0\})$.*

Proof. Firstly, we prove that $\frac{r_\epsilon}{d_\epsilon} \to 0$. If not, there exist some constant δ such that $\frac{r_\epsilon}{d_\epsilon} \ge \delta$ and $y_\epsilon \in \partial\Omega$, $d_\epsilon = |x_\epsilon - y_\epsilon|$. Let

$$\tilde{\psi}_\epsilon = \frac{u_\epsilon(y_\epsilon + r_\epsilon x)}{c_\epsilon}.$$

As the similar procedure of interior case, we have

$$\tilde{\psi}_\epsilon \to 1 \quad \text{in} \quad C^1(B_R^+) \quad \text{for} \quad \|\tilde{\psi}_\epsilon\|_{L^\infty(\overline{B_R^+})} = 1.$$

This is impossible because of $\tilde{\psi}_\epsilon(0) = 0$. Secondly, let $\Omega_\epsilon = \{x \in \mathbb{R}^N : x_\epsilon + r_\epsilon \epsilon \in \Omega\}$, we have know that $\frac{r_\epsilon}{d_\epsilon} \to 0$ by the first step, then $\Omega_\epsilon \to \mathbb{R}^N$. Let φ_ϵ and φ, the same argument as the proof of Lemma 4.2, we get $\varphi_\epsilon \to \varphi$ in $C^1_{loc}(\mathbb{R}^N)$. By the similar process as interior case, we have $c_\epsilon^{\frac{1}{N-1}} u_\epsilon \rightharpoonup \tilde{G}$ in weakly in $W_0^{1,q}(\Omega)(1 < q < N)$ and in $C^1(\overline{\Omega}\backslash\{x_0\})$ with $\tilde{G}$ satisfying $-Q_N\tilde{G} = \gamma\|\tilde{G}\|_p^{N-p}\tilde{G}^{p-1}$ in Ω and $\tilde{G} = 0$ on $\partial\Omega$. By the standard elliptic regularity theory, we have $\tilde{G} \in C^1(\overline{\Omega})$. Since $\gamma < \gamma_1$, test the eqution with function $\tilde{G}$, we get $\tilde{G} \equiv 0$. Thus we have $c_\epsilon^{\frac{1}{N-1}} u_\epsilon \rightharpoonup 0$ weakly in $W_0^{1,q}(\Omega)(1 < q < N)$ and $c_\epsilon^{\frac{1}{N-1}} u_\epsilon \to 0$ strongly in $C^1(\overline{\Omega}\backslash\{x_0\})$. □

Lemma 4.7. *If $0 < \gamma < \gamma_1$, the blow-up point $x_0 \notin \partial\Omega$.*

Proof. Suppose $x_0 \in \partial\Omega$. Then $\|u_\epsilon\|_N^N \to 0$ implies

$$(1 + \gamma\|u_\epsilon\|_p^N)^{-\frac{1}{N-1}} = 1 - \frac{\gamma}{N-1}\|u_\epsilon\|_p^N + O(\|u_\epsilon\|_p^{2N}).$$

Let $w_\epsilon = \frac{u_\epsilon}{\int_\Omega F^N(\nabla u_\epsilon)}$, since $\int_\Omega F^N(\nabla u_\epsilon) = 1 + \gamma\|u_\epsilon\|_p^N$, we have

$$\begin{aligned}
\Lambda_{\gamma,\epsilon} &= \int_\Omega e^{(\lambda_N-\epsilon)|u_\epsilon|^{\frac{N}{N-1}}}dx = \int_\Omega e^{(\lambda_N-\epsilon)(1+\gamma\|u_\epsilon\|_p^N)^{\frac{1}{N-1}}|w_\epsilon|^{\frac{N}{N-1}}}dx \\
&= \int_\Omega e^{[(\lambda_N-\epsilon)(1+\gamma\|u_\epsilon\|_p^N)^{\frac{1}{N-1}}-1]w_\epsilon^{\frac{N}{N-1}}} e^{(\lambda_N-\epsilon)|w_\epsilon|^{\frac{N}{N-1}}}dx \\
&\le \int_\Omega e^{\lambda_N[1-(1+\gamma\|u_\epsilon\|_p^N)^{-\frac{1}{N-1}}]c_\epsilon^{\frac{N}{N-1}}} \int_\Omega e^{\lambda_N|w_\epsilon|^{\frac{N}{N-1}}}dx \\
&\le e^{\lambda_N[\frac{\gamma}{N-1}\|c_\epsilon^{\frac{1}{N-1}}u_\epsilon\|_p^N + c_\epsilon^{-\frac{N}{N-1}}O(\|c_\epsilon^{\frac{1}{N-1}}u_\epsilon\|_p^N)]}\Lambda_0.
\end{aligned}$$

From Lemma 4.6, we have $\|c_\epsilon^{\frac{1}{N-1}}u_\epsilon\|_p^N \to 0$. Thus lettint $\epsilon \to 0$ and using (3.3), we have $\Lambda_\gamma \le \Lambda_0$.

On the other hand, according to the anisotropic Trudinger-Moser inequality, Λ_0 is attained by a function $u \in W_0^{1,N}(\Omega)$ with $\int_\Omega F^N(\nabla u) = 1$. Define $v = u/(1-\gamma\int_\Omega F^N(\nabla u))^{\frac{1}{N}}$. Thus

$$\|v\|_{N,F,\gamma,p} = \left(\int_\Omega F^N(\nabla v)dx - \gamma\|v\|_p^N\right)^{\frac{1}{N}} = 1$$

Since $u \not\equiv 0$ and $\gamma > 0$, we get $|u| \le |v|$ and $|u| \not\equiv |v|$. Thus

$$\Lambda_\gamma \ge \int_\Omega e^{\lambda_N|v|^{\frac{N}{N-1}}}dx > \int_\Omega e^{\lambda_N|u|^{\frac{N}{N-1}}}dx = \Lambda_0,$$

which is a contradiction with $\Lambda_\gamma \le \Lambda_0$. □

4.2 The upper bound estimate

We will use the capacity technique to give an upper bound estimate, which was used by Y. Li [25] and Yang-Zhu [54], our main result of this subsection is an upper bound estimate.

Lemma 4.8. $\Lambda_\gamma \le |\Omega| + \kappa_N e^{\lambda_N A_{x_0}+\sum_{k=1}^{N-1}\frac{1}{k}}$.

Proof. Notice that $x_0 \in \Omega$. Take $\delta > 0$ such that $\mathcal{W}_\delta(x_0) \subset \Omega$. For any $R > 0$, we assume that ϵ is so small that $\delta > Rr_\epsilon$. We denote by $o_\epsilon(1)$ ($o_\delta(1)$; $o_R(1)$) the terms

which tend to 0 as $\epsilon \to 0$ ($\delta \to 0$; $R \to \infty$). From Lemma 4.5, we have

$$\begin{aligned}
\int\limits_{\Omega\setminus\mathcal{W}_\delta(x_\epsilon)} F^N(\nabla u_\epsilon)dx =& \frac{1}{c_\epsilon^{\frac{N}{N-1}}}\Big(\int_{\Omega\setminus\mathcal{W}_\delta(x_\epsilon)} F^N(\nabla G)dx + o_\epsilon(1)\Big)\\
=& \frac{1}{c_\epsilon^{\frac{N}{N-1}}}\Big(\int_{\Omega\setminus\mathcal{W}_\delta(x_\epsilon)} -\mathrm{div}(F^{N-1}(\nabla G)F_\xi(\nabla G))G dx\\
&+ \int_{\partial(\Omega\setminus\mathcal{W}_\delta(x_\epsilon))} GF^{N-1}(\nabla G)\langle F_\xi(\nabla G),\nu\rangle dx + o_\epsilon(1)\Big)\\
=& \frac{1}{c_\epsilon^{\frac{N}{N-1}}}\Big(\gamma\int_{\Omega\setminus\mathcal{W}_\delta(x_\epsilon)} \|G\|_p^{N-p}G^p dx\\
&+ \int_{\partial(\Omega\setminus\mathcal{W}_\delta(x_\epsilon))} GF^{N-1}(\nabla G)\langle F_\xi(\nabla G),\nu\rangle dx + o_\epsilon(1)\Big)\\
=& \frac{1}{c_\epsilon^{\frac{N}{N-1}}}\Big(\gamma\|G\|_p^N - \int_{\partial\mathcal{W}_\delta(x_\epsilon)} GF^{N-1}(\nabla G)\langle F_\xi(\nabla G),\nu\rangle dx + o_\delta(1) + o_\epsilon(1)\Big)\\
=& \frac{1}{c_\epsilon^{\frac{N}{N-1}}}\Big(\gamma\|G\|_p^N - \frac{1}{(N\kappa_N)^{\frac{1}{N-1}}}\log\delta + A_{x_0} + o_\delta(1) + o_\epsilon(1)\Big)\\
=& \frac{1}{c_\epsilon^{\frac{N}{N-1}}}\Big(\gamma\|G\|_p^N - \frac{N}{\lambda_N}\log\delta + A_{x_0} + o_\delta(1) + o_\epsilon(1)\Big).
\end{aligned} \tag{4.14}$$

From (5.3), we have on $\mathcal{W}_{Rr_\epsilon}(x_\epsilon)$ that $u_\epsilon(x) = c_\epsilon^{-\frac{1}{N-1}}\varphi_\epsilon(\frac{x-x_\epsilon}{r_\epsilon}) + c_\epsilon$. Thus

$$\int\limits_{\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} F^N(\nabla u_\epsilon)dx = c_\epsilon^{-\frac{N}{N-1}}\int\limits_{\mathcal{W}_R(0)} F^N(\nabla\varphi_\epsilon)dx = \frac{1}{c_\epsilon^{\frac{N}{N-1}}}\Big(\int\limits_{\mathcal{W}_R(x_0)} F^N(\nabla\varphi)dx + o_\epsilon(1)\Big). \tag{4.15}$$

By a straightforward computation, we have

$$\int\limits_{\mathcal{W}_R(x_0)} F^N(\nabla\varphi)dx = \frac{N}{\lambda_N}\log R + \frac{1}{\lambda_N}\log\kappa_N - \frac{N-1}{\lambda_N}\sum_{k=1}^{N-1}\frac{1}{k} + o_R(1). \tag{4.16}$$

Let $i_\epsilon = \inf_{\partial\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} u_\epsilon$, $s_\epsilon = \sup_{\partial\mathcal{W}_\delta(x_\epsilon)} u_\epsilon$. Since $\psi_\epsilon(x) \to 1$ in $C^1_{loc}(\mathbb{R}^N)$, together with Lemma 4.5, we have $i_\epsilon > s_\epsilon$ for $\epsilon > 0$ small enough. Define a function space

$$\mathbf{S}(i_\epsilon, s_\epsilon) = \{u \in \mathcal{W}_\delta(x_\epsilon)\setminus\overline{\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} : u|_{\partial\mathcal{W}_{Rr_\epsilon}} = i_\epsilon, u|_{\partial\mathcal{W}_\delta(x_\epsilon)} = s_\epsilon\},$$

and $\inf\limits_{u\in\mathbf{S}(i_\epsilon,s_\epsilon)}\int_{B_\delta(x_\epsilon)\setminus B_{Rr_\epsilon}(x_\epsilon)} F^N(\nabla u)dx$ is attained by $h(x)$ satisfying

$$\begin{cases} -Q_N h = 0 & \text{in} \quad \mathcal{W}_\delta(x_\epsilon)\setminus\overline{\mathcal{W}_{Rr_\epsilon}(x_\epsilon)},\\ h|_{\partial\mathcal{W}_{Rr_\epsilon}} = i_\epsilon,\\ h|_{\partial\mathcal{W}_\delta(x_\epsilon)} = s_\epsilon. \end{cases} \tag{4.17}$$

The unique solution is

$$h(x)=\frac{s_\epsilon(\log F^0(x-x_\epsilon)-\log(Rr_\epsilon))+i_\epsilon(-\log\delta-\log F^0(x-x_\epsilon))}{\log\delta-\log(Rr_\epsilon)}, \tag{4.18}$$

and hence

$$\int\limits_{\mathcal{W}_\delta(x_\epsilon)\setminus\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} F^N(\nabla h)dx=N\kappa_N\frac{(i_\epsilon-s_\epsilon)^N}{(\log\delta-\log(Rr_\epsilon))^{N-1}}. \tag{4.19}$$

Define $\tilde{u}_\epsilon=\max\{s_\epsilon,\min\{u_\epsilon,i_\epsilon\}\}$. Then $\tilde{u}_\epsilon\in\mathbf{S}(i_\epsilon,s_\epsilon)$ and $F(\tilde{u}_\epsilon)\le F(u_\epsilon)$ for $\epsilon>0$ small enough. Therefore

$$\begin{aligned}\int\limits_{\mathcal{W}_\delta(x_\epsilon)\setminus\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} F^N(\nabla h)dx=&\int\limits_{\mathcal{W}_\delta(x_\epsilon)\setminus\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} F^N(\nabla\tilde{u}_\epsilon)dx\\ \le&\int\limits_{\mathcal{W}_\delta(x_\epsilon)\setminus\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} F^N(\nabla u_\epsilon)dx\\ =&1+\gamma\|u_\epsilon\|_p^N-\int\limits_{\mathcal{W}_{Rr_\epsilon}(x_\epsilon)} F^N(\nabla u_\epsilon)dx-\int\limits_{\Omega\setminus\mathcal{W}_\delta(x_\epsilon)} F^N(\nabla u_\epsilon)dx.\end{aligned} \tag{4.20}$$

Since $\gamma\|u_\epsilon\|_p^N=\gamma c_\epsilon^{-\frac{N}{N-1}}(\|G\|_p^N+o_\epsilon(1))$, combing (4.14)-(4.16) and (4.19)-(4.20), we obtain

$$\begin{aligned}&\frac{\lambda_N}{N}\frac{(i_\epsilon-s_\epsilon)^{\frac{N}{N-1}}}{\log\frac{\delta}{R}-\log r_\epsilon}\\ \le&\left(1+\frac{\frac{N}{\lambda_N}\log\frac{\delta}{R}-\frac{1}{\lambda_N}\log\kappa_N+\frac{N-1}{\lambda_N}\sum_{k=1}^{N-1}\frac{1}{k}-A_{x_0}+o_\delta(1)+o_\epsilon(1)+o_R(1)}{c_\epsilon^{\frac{N}{N-1}}}\right)^{\frac{1}{N-1}}\\ \le&1+\frac{\frac{N}{\lambda_N}\log\frac{\delta}{R}-\frac{1}{\lambda_N}\log\kappa_N+\frac{N-1}{\lambda_N}\sum_{k=1}^{N-1}\frac{1}{k}-A_{x_0}+o_\delta(1)+o_\epsilon(1)+o_R(1)}{(N-1)c_\epsilon^{\frac{N}{N-1}}}\end{aligned} \tag{4.21}$$

here we use the inequality $(1+t)^{\frac{1}{N-1}}\le 1+\frac{1}{N-1}t$ for any $-1\le t\le 0$ with

$$-1\le\frac{\frac{N}{\lambda_N}\log\frac{\delta}{R}-\frac{1}{\lambda_N}\log\kappa_N+\frac{N-1}{\lambda_N}\sum_{k=1}^{N-1}\frac{1}{k}-A_{x_0}+o_\delta(1)+o_\epsilon(1)+o_R(1)}{c_\epsilon^{\frac{N}{N-1}}}\le 0.$$

On the other hand, using (4.8) and Lemma 4.5, we have

$$\begin{aligned}&(i_\epsilon-s_\epsilon)^{\frac{N}{N-1}}\\ =&c_\epsilon^{\frac{N}{N-1}}\left(1+\frac{\frac{N}{\lambda_N}\log\frac{\delta}{R}-\frac{1}{\lambda_N}\log\kappa_N-A_{x_0}+o_\delta(1)+o_\epsilon(1)+o_R(1)}{c_\epsilon^{\frac{N}{N-1}}}\right)^{\frac{N}{N-1}}\\ \ge&c_\epsilon^{\frac{N}{N-1}}+\frac{N}{N-1}\left(\frac{N}{\lambda_N}\log\frac{\delta}{R}-\frac{1}{\lambda_N}\log\kappa_N-A_{x_0}+o_\delta(1)+o_\epsilon(1))+o_R(1)\right),\end{aligned} \tag{4.22}$$

here we use the inequality $(1+t)^{\frac{N}{N-1}} \geq 1+\frac{N}{N-1}t$ for any $-1 \leq t \leq 0$ with

$$-1 \leq \frac{\frac{N}{\lambda_N}\log\frac{\delta}{R} - \frac{1}{\lambda_N}\log\kappa_N - A_{x_0} + o_\delta(1) + o_\epsilon(1) + o_R(1)}{c_\epsilon^{\frac{N}{N-1}}} \leq 0.$$

Since $\log\frac{\delta}{R} - \log r_\epsilon = \log\frac{\delta}{R} + \frac{\lambda_N-\epsilon}{N}c_\epsilon^{\frac{N}{N-1}} - \frac{1}{N}\log\frac{\lambda_\epsilon}{c_\epsilon^{\frac{N}{N-1}}}$, combing (4.21) and (4.22), we have

$$\begin{aligned}
&\frac{\lambda_N}{N}\left[c_\epsilon^{\frac{N}{N-1}} + \frac{N}{N-1}\left(\frac{N}{\lambda_N}\log\frac{\delta}{R} - \frac{1}{\lambda_N}\log\kappa_N - A_{x_0} + o_\delta(1) + o_\epsilon(1)) + o_R(1)\right)\right] \\
\leq&(\log\frac{\delta}{R} + \frac{\lambda_N-\epsilon}{N}c_\epsilon^{\frac{N}{N-1}} - \frac{1}{N}\log\frac{\lambda_\epsilon}{c_\epsilon^{\frac{N}{N-1}}}) \\
&\times\left[1 + \frac{\frac{N}{\lambda_N}\log\frac{\delta}{R} - \frac{1}{\lambda_N}\log\kappa_N + \frac{N-1}{\lambda_N}\sum_{k=1}^{N-1}\frac{1}{k} - A_{x_0} + o(1)}{(N-1)c_\epsilon^{\frac{N}{N-1}}}\right] \\
\leq&\frac{\lambda_N-\epsilon}{N}c_\epsilon^{\frac{N}{N-1}} + \frac{N}{N-1}\log\frac{\delta}{R} - \frac{1+o(1)}{N}\log\frac{\lambda_\epsilon}{c_\epsilon^{\frac{N}{N-1}}} - \frac{1}{N(N-1)}\log\kappa_N \\
&+\frac{1}{N}\sum_{k=1}^{N-1}\frac{1}{k} - \frac{\lambda_N}{N(N-1)}A_{x_0} + o(1).
\end{aligned} \tag{4.23}$$

Thus

$$\frac{1+o(1)}{N}\log\frac{\lambda_\epsilon}{c_\epsilon^{\frac{N}{N-1}}} \leq \frac{1}{N}\log\kappa_N + \sum_{k=1}^{N-1}\frac{1}{k} + \frac{\lambda_N}{N}A_{x_0} + +o_\delta(1) + o_\epsilon(1)) + o_R(1),$$

which lead to

$$\limsup_{\epsilon\to 0}\frac{\lambda_\epsilon}{c_\epsilon^{\frac{N}{N-1}}} \leq \kappa_N e^{\lambda_N A_{x_0} + \sum_{k=1}^{N-1}\frac{1}{k}}.$$

Recall Lemma 4.4, we have finished the proof. □

5 Proof of main Theorems

Proof of Theorem 1.1. Let $0 \leq \gamma < \gamma_1$. If c_ϵ is bounded, the inequality (4.1) implies that the Theorem holds. If $c_\epsilon \to +\infty$ is bounded, we know that the blow-up point $x_0 \in \Omega$ by Lemma 4.7 and the rusult is followed from Lemma 4.8. □

Proof of Theorem 1.2. Let $0 \leq \gamma < \gamma_1$. We prove that the blow-up phenomena do not occur. In fact, if $c_\epsilon \to +\infty$. In Lemma 4.8, we have got the upper bound of Λ_γ, that is to say

$$\Lambda_\gamma \leq |\Omega| + \kappa_N e^{\lambda_N A_{x_0} + \sum_{k=1}^{N-1}\frac{1}{k}}, \tag{5.1}$$

where x_0 is the blow-up point. If we can construct a sequence $\phi_\epsilon \in W^{1,N}(\Omega)$ with $\|\phi_\epsilon\|^N_{N,F,\gamma,p} = \int_\Omega F^N(\nabla\phi_\epsilon)dx - \gamma\|\phi_\epsilon\|_p^N = 1$, but

$$\int_\Omega e^{\lambda_N|\phi_\epsilon|^{\frac{N}{N-1}}}dx > |\Omega| + \kappa_N e^{\lambda_N A_{x_0}+\sum_{k=1}^{N-1}\frac{1}{k}}. \tag{5.2}$$

This is the contradiction with (5.1), which implies that c_ϵ must be bounded and can be attained by the discussion at the beginning of subsection 4.1. Thus it suffice to construct the sequence ϕ_ϵ such that (5.2) holds when the blow-up phenomena occur. From Lemma 4.5,

$$G(x) = -\frac{1}{(N\kappa_N)^{\frac{1}{N-1}}}\log F^o(x-x_0) + A_{x_0} + \xi(x).$$

Define a sequence of functions

$$\phi_\epsilon(x) = \begin{cases} C + C^{-\frac{1}{N-1}}(-\frac{N-1}{\lambda_N}\log(1+\kappa_N^{\frac{1}{N-1}}(F^o(x-x_0)\epsilon^{-1})^{\frac{N}{N-1}} + B), & x \in \overline{\mathcal{W}_{R\epsilon}(x_0)}, \\ C^{-\frac{1}{N-1}}(G-\eta\xi), & x \in \mathcal{W}_{2R\epsilon}(x_0)\backslash\overline{\mathcal{W}_{R\epsilon}(x_0)}, \\ C^{-\frac{1}{N-1}}G, & \Omega\backslash\mathcal{W}_{2R\epsilon}(x_0), \end{cases} \tag{5.3}$$

where B and C are constants depending only on ϵ, which will be determined later. The cutoff function $\eta \in C_0^1(\mathcal{W}_{2R\epsilon}(x_0))$, $0 \le \eta \le 1$ and $\eta = 1$ in $\mathcal{W}_{R\epsilon}(x_0)$. To ensure $\phi_\epsilon \in W^{1,N}(\Omega)$, we require for all $x \in \partial\mathcal{W}_{R\epsilon}(x_0)$, there holds

$$C + C^{-\frac{1}{N-1}}(-\frac{N-1}{\lambda_N}\log(1+\kappa_N^{\frac{1}{N-1}}R^{\frac{N}{N-1}} + B) = C^{-\frac{1}{N-1}}(-\frac{1}{(N\kappa_N)^{\frac{1}{N-1}}}\log(R\epsilon) + A_{x_0}),$$

which implies

$$B = -C^{\frac{N}{N-1}} + \frac{N-1}{\lambda_N}\log(1+\kappa_N^{\frac{1}{N-1}}R^{\frac{N}{N-1}}) - \frac{N}{\lambda_N}\log(R\epsilon) + A_{x_0}.$$

On one hand, since

$$F^N(\nabla\phi_\epsilon) = C^{-\frac{N}{N-1}}F^N(\nabla\phi_\epsilon)(1+O(R\epsilon))$$

uniformly in $\mathcal{W}_{2R}(x_0)\backslash\mathcal{W}_R(x_0)$ as $\epsilon\to 0$. Thus, by Lemma 4.5, we have

$$\begin{aligned}\int\limits_{\Omega\backslash\mathcal{W}_R(x_0)} F^N(\nabla\phi_\epsilon)dx =& \int\limits_{\Omega\backslash\mathcal{W}_{2R}(x_0)} F^N(\nabla\phi_\epsilon)dx + \int\limits_{\mathcal{W}_{2R}(x_0)\backslash\mathcal{W}_R(x_0)} F^N(\nabla\phi_\epsilon)dx\\ =&C^{-\frac{N}{N-1}}\Big(\int\limits_{\Omega\backslash\mathcal{W}_{2R}(x_0)} F^N(\nabla G)dx + \int\limits_{\mathcal{W}_{2R}(x_0)\backslash\mathcal{W}_R(x_0)} F^N(\nabla G)(1+O(R\epsilon))dx\Big)\\ =&C^{-\frac{N}{N-1}}\Big(\int\limits_{\Omega\backslash\mathcal{W}_R(x_0)} F^N(\nabla G)dx + \int\limits_{\mathcal{W}_{2R}(x_0)\backslash\mathcal{W}_R(x_0)} F^N(\nabla G)O(R\epsilon)dx\Big)\\ =&C^{-\frac{N}{N-1}}\Big(\int_{\Omega\backslash\mathcal{W}_R(x_0)} F^N(\nabla G)dx + O(-R\epsilon\log(R\epsilon))\Big)\\ =&C^{-\frac{N}{N-1}}\left(\gamma\|G\|_p^N - \frac{N}{\lambda_N}\log(R\epsilon) + A_{x_0} + O(-R\epsilon\log(R\epsilon))\right).\end{aligned}$$

On the other hand, through the direct calculation, we have

$$\int\limits_{\mathcal{W}_R(x_0)} F^N(\nabla\phi_\epsilon)dx = C^{-\frac{N}{N-1}}\left(\frac{N-1}{\lambda_N}\log(1+\kappa_N^{\frac{1}{N-1}}R^{\frac{N}{N-1}}) - \frac{N-1}{\lambda_N}\sum_{k=1}^{N-1}\frac{1}{k} + o_R(1)\right).$$

Thus

$$\begin{aligned}\int_\Omega F^N(\nabla\phi_\epsilon)dx =& \int\limits_{\Omega\backslash\mathcal{W}_R(x_0)} + \int\limits_{\mathcal{W}_R(x_0)} F^N(\nabla\phi_\epsilon)dx\\ =&C^{-\frac{N}{N-1}}\left(\gamma\|G\|_p^N - \frac{N}{\lambda_N}\log\epsilon + A_{x_0} + \frac{1}{\lambda_N}\log\kappa_N\right.\\ &\left. - \frac{N-1}{\lambda_N}\sum_{k=1}^{N-1}\frac{1}{k} + O(-R\epsilon\log(R\epsilon))\right),\end{aligned}$$

we also have

$$\|\phi_\epsilon\|_p^N = C^{-\frac{N}{N-1}}\left(\|G\|_p^N + O((R\epsilon)^N(-\log(R\epsilon))^N)\right).$$

Take $R=-\log\epsilon$, we get

$$\begin{aligned}\|\phi_\epsilon\|_{N,F,\gamma,p}^N =& \int_\Omega F^N(\nabla\phi_\epsilon)dx - \gamma\|\phi_\epsilon\|_p^N\\ =&C^{-\frac{N}{N-1}}\left(\frac{N}{\lambda_N}R + A_{x_0} + \frac{1}{\lambda_N}\log\kappa_N - \frac{N-1}{\lambda_N}\sum_{k=1}^{N-1}\frac{1}{k} + O(R^{-\frac{N}{N-1}})\right)\end{aligned} \tag{5.4}$$

Choosing

$$C^{\frac{N}{N-1}} = -\frac{N}{\lambda_N}\log\epsilon + A_{x_0} + \frac{1}{\lambda_N}\log\kappa_N - \frac{N-1}{\lambda_N}\sum_{k=1}^{N-1}\frac{1}{k} + O(R^{-\frac{N}{N-1}}), \tag{5.5}$$

which implies

$$B = \frac{N-1}{\lambda_N}\sum_{k=1}^{N-1}\frac{1}{k} + O(R^{-\frac{N}{N-1}}) + o_R(1). \tag{5.6}$$

Then we can get $\|\phi_\epsilon\|^N_{N,F,\gamma,p} = \int_\Omega F^N(\nabla\phi_\epsilon)dx - \gamma\|\phi_\epsilon\|_p^N = 1$.

Now we estimate $\int_\Omega e^{\lambda_N|\phi_\epsilon|^{\frac{N}{N-1}}}$. Let $F^o(x-x_0) = \epsilon y$, we get

$$\begin{aligned}\int_{\mathcal{W}_{R\epsilon}(x_0)} e^{\lambda_N|\phi_\epsilon|^{\frac{N}{N-1}}} \geq& \kappa_N e^{\lambda_N A_{x_0}+\sum_{k=1}^N\frac{1}{k}+O(R^{-\frac{N}{N-1}})}\int\limits_{\mathcal{W}_R(0)}(1+\kappa_N^{-\frac{1}{N-1}}|y|^{\frac{N}{N-1}})^{-N}dy\\ =&\kappa_N e^{\lambda_N A_{x_0}+\sum_{k=1}^N\frac{1}{k}+O(R^{-\frac{N}{N-1}})}(1+O(R^{-\frac{N}{N-1}}))\\ =&\kappa_N e^{\lambda_N A_{x_0}+\sum_{k=1}^N\frac{1}{k}}+O(R^{-\frac{N}{N-1}}).\end{aligned} \tag{5.7}$$

From the inequality $e^t \geq 1 + \frac{t^{N-1}}{(N-1)!}$, we have

$$\begin{aligned}&\int\limits_{\Omega\setminus\mathcal{W}_{R\epsilon}(x_0)} e^{\lambda_N|\phi_\epsilon|^{\frac{N}{N-1}}}dx\\ \geq&\int\limits_{\Omega\setminus\mathcal{W}_{2R\epsilon}(x_0)}\left(1+\frac{\lambda_N^{N-1}|\phi_\epsilon|^N}{(N-1)!}\right)dx\\ =&|\Omega|-|\mathcal{W}_{2R\epsilon}(x_0)| + C^{-\frac{N}{N-1}}\frac{\lambda_N^{N-1}}{(N-1)!}(\|G\|_N^N + O((R\epsilon)^N(-\log(R\epsilon))^N)\\ =&|\Omega| + C^{-\frac{N}{N-1}}\frac{\lambda_N^{N-1}}{(N-1)!}\|G\|_N^N + O((-\log\epsilon)^{-\frac{N}{N-1}}).\end{aligned} \tag{5.8}$$

Thus

$$\begin{aligned}\int_\Omega e^{\phi_\epsilon}dx =& \int_{\mathcal{W}_{R\epsilon}(x_0)} e^{\lambda_N|\phi_\epsilon|^{\frac{N}{N-1}}}dx + \int\limits_{\Omega\setminus\mathcal{W}_{R\epsilon}(x_0)} e^{\lambda_N|\phi_\epsilon|^{\frac{N}{N-1}}}dx\\ \geq&|\Omega| + \kappa_N e^{\lambda_N A_{x_0}+\sum_{k=1}^N\frac{1}{k}} + C^{-\frac{N}{N-1}}\frac{\lambda_N^{N-1}}{(N-1)!}\|G\|_N^N + O((-\log\epsilon)^{-\frac{N}{N-1}}),\end{aligned} \tag{5.9}$$

By choosing a small $\epsilon > 0$, we conclude $\int_\Omega e^{\phi_\epsilon}dx > |\Omega| + \kappa_N e^{\lambda_N A_{x_0}+\sum_{k=1}^N\frac{1}{k}}$. Hence we finish the proof of Theorem 1.2. □

References

[1] S. Adachi, K. Tanaka, Trudinger type inequalities in $\mathbb{R}^N$ and their best exponents, Proc. Amer. Math. Soc. 128 (2000) 2051-2057.

[2] D. Adams, A sharp inequality of J. Moser for higher order derivatives, Ann. of Math. 128 (1988) 385-398.

[3] Adimurthi, O. Druet, Blow-up analysis in dimension 2 and a sharp form of Moser-Trudinger inequality, Comm. Partial Differential Equations 29 (2004) 295-322.

[4] A. Adimurthi and Y. Yang, An interpolation of Hardy inequality and Trudinger-Moser inequality Int. Math. Res. Notices, 13 (2010), 2394–2426.

[5] A. Alvino, V. Ferone, G. Trombetti, P. Lions, Convex symmetrization and applications, Ann. Inst. H. Poincaré Anal. Non Linéaire 14 (1997) 275-293.

[6] M. Belloni, V. Ferone, B. Kawohl, Isoperimetric inequalities, Wulff shape and related questions for strongly nonlinear elliptic operators, Z. Angew. Math. Phys. 54 (2003) 771-783.

[7] D. M. Cao, Nontrivial solution of semilinear elliptic equation with critical exponent in $\mathbb{R}^2$, Comm. Partial Differential Equations 17 (1992) 407-435.

[8] L. Carleson, A. Chang, On the existence of an extremal function for an inequality of J. Moser, Bull. Sci. Math. 110 (1986) 113-127.

[9] R. Cerny, A. Cianchi, S. Henel, Concentration-compactness principles for Moser-Trudinger inequalities: new results and proofs, Ann. Mat. Pura Appl. 192 (2013) 225-243.

[10] S.Y.A. Chang, P. Yang, The inequality of Moser and Trudinger and applications to conformal geometry, Comm. Pure Appl. Math., 56 (2003) 11351150.

[11] G. Csató, P. Roy, Extremal functions for the singular Moser-Trudinger inequality in 2 dimensions, Calc. Var. Partial Differential Equations 54 (2015) 2341-2366.

[12] D. G. de Figueiredo, O. H. Miyagaki and B. Ruf, Elliptic equations in $\mathbb{R}^2$ with nonlinearities in the critical growth range, Calc. Var. Partial Differ. Equ., 3 (1995), 139–153.

[13] D. G. de Figueiredo, J. M. do Ó, B. Ruf, Elliptic equations and systems with critical Trudinger-Moser nonlinearities, Discrete Contin. Dyn. Syst. 30 (2011) 455-476.

[14] M. de Souza, J.M. do Ó, A sharp Trudinger-Moser type inequality in $\mathbb{R}^2$, Trans. Amer. Math. Soc. 366 (2014) 4513-4549.

[15] J. M. do Ó, N-Laplacian equations in $\mathbb{R}^N$ with critical growth, Abstr. Appl. Anal. 2 (1997) 301-315.

[16] J.M. do Ó, M. de Souza, E. de Medeiros and U. Severo, An improvement for the Trudinger-Moser inequality and applications. J. Differential Equations, 256 (2014) 1317-1349.

[17] J. M. do Ó, M. de Souza, A sharp inequality of Trudinger-Moser type and extremal functions in $H^{1,n}(\mathbb{R}^n)$, J. Differential Equations 258 (2015) 4062-4101.

[18] Vincenzo Ferone, Bernd Kawohl, Remarks on a Finsler-Laplacian, Proc. Amer. Math. Soc. 137 (2009) 247-253.

[19] M. Flucher, Extremal functions for Trudinger-Moser inequality in 2 dimensions, Comment. Math. Helv. 67 (1992) 471-497.

[20] I. Fonseca, S. Müller, A uniqueness proof for the Wulff theorem, Proc. Roy. Soc. Edinburgh Sect. A 119 (1991) 125136.

[21] J. Heinonen, T. Kilpelainen, O. Martio, Nonlinear Potential Theory of Degenerate Elliptic Equations, Oxford University Press, 1993.

[22] N. Lam, G. Lu, A new approach to sharp Moser-Trudinger and Adams type inequalities: A rearrangement-free argument, J. Differential Equations 255 (2013), 298-325.

[23] N. Lam, G. Lu, Elliptic equations and systems with subcritical and critical exponential growth without the Ambrosetti-Rabinowitz condition, J. Geom. Anal., 24 (2014) 118-143.

[24] N. Lam, G. Lu, L. Zhang, Equivalence of critical and subcritical sharp Trudinger-Moser-Adams inequalities, Rev. Mat. Iberoam. 33 (2017), 1219-1246.

[25] Y. Li, Moser-Trudinger inequality on compact Riemannian manifolds of dimension two, J. Partial Differ. Equ. 14 (2001) 163-192.

[26] Y. Li, B. Ruf, A sharp Trudinger-Moser type inequality for unbounded domains in $\mathbb{R}^N$, Indiana Univ. Math. J. 57 (2008) 451-480.

[27] X. Li, Y. Yang, Extremal functions for singular Trudinger-Moser inequalities in the entire Euclidean space, J. Differential Equations 264 (2018) 4901-4943.

[28] G.M. Lieberman, Boundary regularity for solutions of degenerate elliptic equations, Nonlinear Anal. 12 (1988) 1203-1219.

[29] K. Lin, Extremal functions for Moser's inequality, Trans. Amer. Math. Soc. 348 (1996) 2663-2671.

[30] P. L. Lions, The concentration-compactness principle in the calculus of variations, Part I, Rev. Mat. Iberoamericana 1 (1985) 145-201.

[31] G. Lu, H. Tang, Sharp singular Trudinger-Moser inequalities in Lorentz-Sobolev spaces, Adv. Nonlinear Stud. 16 (2016), 581-601.

[32] G. Lu, Y. Yang, The sharp constant and extremal functions for Moser-Trudinger inequalities involving L^p norms, Discrete Contin. Dyn. Syst. 25 (2009) 963-979.

[33] A. Malchiodi, L. Martinazzi, Critical points of the Moser-Trudinger functional on a disk, J. Eur. Math. Soc. (JEMS) 16 (2014) 893-908.

[34] N. Masmoudi, F. Sani, Trudinger-Moser inequalities with the exact growth condition in Rn and application. Commun. Partial Differ. Equ. 40 (2015) 1408-1440.

[35] G. Mancini, L. Martinazzi, The Moser-Trudinger inequality and its extremals on a disk via energy estimates, Calc. Var. Partial Differential Equations 20 (2017) 56-94.

[36] J. Moser, A sharp form of an inequality by N. Trudinger, Indiana Univ. Math. J. 20 (1970) 1077-1092.

[37] V. H. Nguyen, Improved Moser-Trudinger inequality of Tintarev type in dimension n and the existence of its extremal functions, Ann. Global Anal. Geom. 54 (2018) 237-256.

[38] S. Pohozaev, The Sobolev embedding in the special case pl = n, in: Proceedings of the Technical Scientific Conference on Advances of Scientific Research 1964-1965, Mathematics Sections, Moscov. Energet. Inst., Moscow, 1965, pp. 158-170

[39] J. Peetre, Espaces d'interpolation et theoreme de Soboleff, Ann. Inst. Fourier (Grenoble) 16 (1966) 279-317.

[40] B. Ruf, A sharp Trudinger-Moser type inequality for unbounded domains in $\mathbb{R}^2$, J. Funct. Anal. 219 (2005) 340-367.

[41] B. Ruf, F. Sani, Sharp Adams-type inequalities in $\mathbb{R}^n$, Trans. Amer. Math. Soc. 365 (2013) 645-670.

[42] M. Struwe, Critical points of embeddings of $H_0^{1,n}$ into Orlicz spaces, Ann. Inst. H. Poincaré Anal. Non Linéaire 5 (1988) 425-464.

[43] M. Struwe, Positive solution of critical semilinear elliptic equations on non-contractible planar domain, J. Eur. Math. Soc. 2 (2000) 329-388.

[44] G. Talenti, Elliptic equations and rearrangements, Ann. Sc. Norm. Super. Pisa Cl. Sci. 3 (1976) 697-718.

[45] C. Tintarev, Trudinger-Moser inequality with remainder terms, J. Funct. Anal. 266 (2014) 55-66.

[46] N. S. Trudinger, On imbeddings into Orlicz spaces and some applications, J. Math. Mech. 17 (1967) 473-484.

[47] P. Tolksdorf, Regularity for a more general class of quasilinear elliptic equations, J. Differential Equations 51 (1984) 126-150.

[48] G. Wang, C. Xia, A characterization of the Wulff shape by an overdetermined anisotropic PDE. *Arch. Ration. Mech. Anal.* 99 (2011) 99-115.

[49] G. Wang, C. Xia, Blow-up analysis of a Finsler-Liouville equation in two dimensions, J. Differential Equations 252 (2012) 1668-1700.

[50] R. L. Xie, H. J. Gong, A priori estimates and blow-up behavior for solutions of $-Q_N u = Ve^u$ in bounded domain in $\mathbb{R}^N$, Sci. China Math. 59 (2016) 479-492.

[51] Y. Yang, A sharp form of Moser-Trudinger inequality in high dimension, J. Funct. Anal. 239 (2006) 100-126.

[52] Y. Yang, A sharp form of the Moser-Trudinger inequality on a compact Riemannian surface, Trans. Amer. Math. Soc. 359 (2007) 5761-5776.

[53] Y. Yang, Extremal functions for Trudinger-Moser inequalities of Adimurthi-Druet type in dimension two, J. Differential Equations 258 (2015) 3161-3193.

[54] Y. Yang, X. Zhu, Blow-up analysis concerning singular Trudinger-Moser inequalities in dimension two, J. Funct. Anal. 272 (2017) 3347-3374.

[55] V. I. Yudovich, Some estimates connected with integral operators and with solutions of elliptic equations, Sov. Math., Dokl. 2 (1961) 746-749.

[56] C. L. Zhou, C. Q Zhou, Moser–Trudinger inequality involving the anisotropic Dirichlet norm $(\int_\Omega F^N(\nabla u)dx)^{\frac{1}{N}}$ on $W_0^{1,N}(\Omega)$, J. Funct. Anal. 276 (2019) 2901-2935.

[57] C. L. Zhou, Anisotropic Moser-Trudinger inequality involving L^n norm, J. Differential Equations (2019), https://doi.org/10.1016/j.jde.2019.11.066.

[58] J. Zhu, Improved Moser-Trudinger inequality involving L^p norm in n dimensions, Adv. Nonlinear Study 14 (2014) 273-293.