

# Wide-Sense 2-Frameproof Codes\*

Junling Zhou, Wenling Zhou  
 Department of Mathematics  
 Beijing Jiaotong University  
 Beijing 100044, P. R. China  
 jlzhou@bjtu.edu.cn

## Abstract

Various kinds of fingerprinting codes and their related combinatorial structures are extensively studied for protecting copyrighted materials. This paper concentrates on one specialised fingerprinting code named wide-sense frameproof codes in order to prevent innocent users from being framed.

Let  $Q$  be a finite alphabet of size  $q$ . Given a  $t$ -subset  $X = \{x^1, \dots, x^t\} \subseteq Q^n$ , a position  $i$  is called undetectable for  $X$  if the values of the words of  $X$  match in their  $i$ th position:  $x_i^1 = \dots = x_i^t$ . The wide-sense descendant set of  $X$  is defined by  $\text{wdesc}(X) = \{y \in Q^n : y_i = x_i^1, i \in U(X)\}$ , where  $U(X)$  is the set of undetectable positions for  $X$ . A code  $\mathcal{C} \subseteq Q^n$  is called a wide-sense  $t$ -frameproof code if  $\text{wdesc}(X) \cap \mathcal{C} = X$  for all  $X \subseteq \mathcal{C}$  with  $|X| \leq t$ .

The paper improves the upper bounds on the sizes of wide-sense 2-frameproof codes by applying techniques on non 2-covering Sperner families and intersecting families in extremal set theory.

**MSC [2010]:** 05C65, 05D05, 68R05, 94B65

**Keywords:** fingerprinting code, frameproof code, intersecting family, Sperner family

## 1 Introduction

Fingerprinting codes are combinatorial objects that have been studied for more than 20 years due to their applications in digital data copyright protection and their combinatorial interest. Let  $Q$  be a finite alphabet of size  $q$ . In order to protect a copyrighted digital product, a dealer inserts a fingerprint in each copy and then distributes copies to all registered users, where a *fingerprint* is a string  $x = (x_1, \dots, x_n)$  over  $Q$ . The goal of inserting the fingerprint is to personalize the copy given out to the user, and to rule out redistribution. Clearly, an individual user will be deterred from releasing an unauthorized copy. However, a coalition of users may collude in order to produce an unregistered copy. The goal of the coalition is to create a fingerprint of the illegal copy that is unable to identify users from it. We assume that the members of a coalition can only alter those coordinates of the fingerprint in which at least two of their fingerprints differ, and refer to this as the *Marking Assumption*. In this paper we concentrate on  $t$ -frameproof codes, which have the property that no coalition of at most  $t$  users can frame a user not in the coalition.

---

\*Supported by NSFC grants 11571034 and 11971053.

Given a  $t$ -subset  $X = \{x^1, \dots, x^t\} \subseteq Q^n$  we now define the set of descendants of  $X$ . We write  $x_i^j$  for the  $i$ -th component of  $x^j$  for  $1 \leq i \leq n$  and  $1 \leq j \leq t$ . A position  $i$  is called *undetectable* for  $X$  if the values of the words of  $X$  match in their  $i$ -th position:  $x_i^1 = \dots = x_i^t$ . Denote by  $U(X)$  the set of positions undetectable for  $X$ . By the marking assumption, the coalition cannot change the values of undetectable positions. If the position is detectable, then there are several options for the coalition to fill it. We will consider *wide-sense descendant set* defined by

$$\text{wdesc}(X) = \{y \in Q^n : y_i = x_i^1, i \in U(X)\},$$

in contrast to the *narrow-sense descendant set*

$$\text{desc}(X) = \{y \in Q^n : y_i \in \{x_i^1, \dots, x_i^t\}\}.$$

In the literature the wide-sense descendant set could be found under the name of envelope [3], feasible set [9, 27], or Boneh-Shaw descendant [6].

A code  $\mathcal{C} \subseteq Q^n$  of size  $|\mathcal{C}| = m$  is called an  $(n, m, q)$  code. We define  $\mathcal{C}$  to be a *wide-sense  $t$ -frameproof code*, or  $(n, m, q)$   *$t$ -wFP code*, if

$$\text{wdesc}(X) \cap \mathcal{C} = X$$

for all  $X \subseteq \mathcal{C}$  with  $|X| \leq t$ . Replacing  $\text{wdesc}(X)$  with  $\text{desc}(X)$  defines a *narrow-sense  $t$ -frameproof code*.

Boneh and Shaw [9] were the first to define a  $t$ -frameproof code, where they adopted the wide-sense model of descendent sets. In the narrow-sense model, variants of frameproof codes were extensively studied by many researchers. Named after the security properties it guarantees, the following types of fingerprinting codes are well-known: frameproof codes, secure frameproof codes, identifiable parent property codes, traceability codes, and anti-collusion codes, see [1], [3]-[8], [10]-[14], [17, 18], [23]-[29], [30, 31].

It is clear that  $\text{wdesc}(X)$  always strictly contains  $\text{desc}(X)$  if  $2 \leq |X| < q$ . Just as Blackburn [6] said, “This is one reason why the problem of constructing analogues of the codes for Boneh-Shaw descendants is often more difficult than the original problem.” To the best of our knowledge, frameproof codes is the only type of fingerprinting codes that was ever studied in the wide-sense model.

The paper is organized as follows. In Section 2 we introduce Sperner families and intersecting families and we display an upper bound on the size of a Sperner family by its maximum size and minimum size of subsets. In Section 3 we improve the known upper bounds on the sizes of 2-wFP codes, which were previously established by Panoui in her PhD dissertation [22]. Non 2-covering Sperner families generated by all codewords are considered and better upper bounds are established by developing many results on Sperner families and intersecting families. In Section 4 we conclude the paper.

## 2 Sperner families

Stinson and Wei [27] were the first to establish the relationship between Sperner families and  $t$ -wFP codes and then proved that  $m \leq \binom{n}{\lfloor \frac{n}{2} \rfloor} + 1$  for  $(n, m, 2)$  2-wFP codes by applying Sperner’s Theorem ([27, Theorem 5.2]). Panoui [22] developed this idea and presented the equivalence

between a 2-wFP code and the non 2-covering Sperner families generated by all codewords. The upper bounds on the sizes of 2-wFP codes were then improved as follows.

**Lemma 2.1.** [22, Theorem 6.3.8] *Let  $\mathcal{C}$  be an  $(n, m, q)$  2-wFP code.*

- (1) *If  $n$  is even, then  $m \leq \binom{n}{\frac{n}{2}-1} + 1$ .*
- (2) *If  $n$  is odd, then  $m \leq \binom{n-1}{\frac{n-1}{2}} - \frac{n-1}{2}$ .*

The aim of this section is to improve the above upper bounds. We first introduce related definitions in extremal set theory and recall or develop some useful results.

Let  $\mathcal{F}$  be a family of finite sets. If any two distinct sets of  $\mathcal{F}$  are incomparable, that is,  $A \not\subseteq B$  for any  $A, B \in \mathcal{F}$ , then  $\mathcal{F}$  is called an *antichain* or a *Sperner family*. To the other extreme, a *chain* is a set family  $\mathcal{F}$  in which every pair of sets is comparable.

**Theorem 2.2.** (Sperner's Theorem) [2] *Let  $\mathcal{F}$  be a Sperner family over an  $n$ -set. Then*

$$|\mathcal{F}| \leq \binom{n}{\lfloor \frac{n}{2} \rfloor}.$$

The size of a Sperner family which contains a singleton is easily obtained from Sperner's Theorem.

**Proposition 2.3.** [22, Proposition 6.3.4] *Let  $\mathcal{F}$  be a Sperner family over an  $n$ -set. If there exists a set  $F \in \mathcal{F}$  such that  $|F| = 1$ , then*

$$|\mathcal{F}| \leq \binom{n-1}{\lfloor \frac{n-1}{2} \rfloor} + 1.$$

Let  $C = \{A_1, A_2, \dots, A_k\}$  be a chain of subsets of an  $n$ -set, i.e.,  $A_1 \subseteq A_2 \subseteq \dots \subseteq A_k$ . This chain is *symmetric* if  $|A_1| + |A_k| = n$  and  $|A_{i+1}| = |A_i| + 1$  for all  $i = 1, 2, \dots, k-1$ .

**Theorem 2.4.** [19, Theorem 8.3] *The family of all subsets of an  $n$ -set can be partitioned into  $\binom{n}{\lfloor \frac{n}{2} \rfloor}$  mutually disjoint symmetric chains.*

A family  $\mathcal{F}$  of sets is called  *$k$ -intersecting* ( $k \geq 1$ ), if  $|A \cap B| \geq k$  for all  $A, B \in \mathcal{F}$ . An *intersecting family* is a 1-intersecting family. Call the families  $\mathcal{A}$  and  $\mathcal{B}$  *cross- $k$ -intersecting* if  $|A \cap B| \geq k$  holds for all  $A \in \mathcal{A}$  and  $B \in \mathcal{B}$ .  $\mathcal{A}$  and  $\mathcal{B}$  are *cross-intersecting* if they are cross-1-intersecting. Let  $\mathcal{F}$  be a family of subsets of a ground set  $E$ . Then  $\mathcal{F}$  is called *non 2-covering* if for every pair of sets  $A, B \in \mathcal{F}$  we have  $A \cup B \neq E$ .

**Theorem 2.5.** [21] *If  $\mathcal{F}$  is a  $k$ -intersecting Sperner family over an  $n$ -set, then*

$$|\mathcal{F}| \leq \binom{n}{\lfloor \frac{n+k+1}{2} \rfloor}.$$

Let  $\mathcal{C} = \{c^1, c^2, \dots, c^m\}$  be an  $(n, m, q)$  code. For any  $1 \leq i, j \leq m$  and  $i \neq j$ , define  $I(i, j)$  to be the *coincidence set* of  $c^i$  and  $c^j$ , i.e.,

$$I(i, j) = \{k : c_k^i = c_k^j, 1 \leq k \leq n\}.$$

For any  $1 \leq i \leq m$ , define

$$\mathcal{X}_i = \{I(i, j) : i \neq j, 1 \leq j \leq m\}$$

to be the *coincidence family* generated by the codeword  $c^i \in \mathcal{C}$ . Clearly  $|\mathcal{C}| = |\mathcal{X}_i| + 1$  for  $1 \leq i \leq m$ .

**Theorem 2.6.** [22, Lemma 6.3.2, Corollary 6.3.3] Let  $\mathcal{C} = \{c^1, c^2, \dots, c^m\}$  be an  $(n, m, q)$  code. Then,  $\mathcal{C}$  is a 2-wFP code if and only if  $\mathcal{X}_i$  is a non 2-covering Sperner family for any  $1 \leq i \leq m$ .

We have a simple but useful result on the coincidence sets.

**Lemma 2.7.** Let  $\mathcal{C}$  be an  $(n, m, q)$  code. For any three codewords  $c^i, c^j, c^k \in \mathcal{C}$ , we have

$$I(i, j) \cap I(i, k) \subseteq I(j, k) \subseteq (I(i, j) \cap I(i, k)) \cup \overline{I(i, j) \cup I(i, k)}.$$

*Proof.* Firstly let  $p \in I(i, j) \cap I(i, k)$ . Then we have  $c_p^i = c_p^j$  and  $c_p^i = c_p^k$ . Hence  $p \in I(j, k)$  and

$$I(i, j) \cap I(i, k) \subseteq I(j, k).$$

Secondly let  $p \in I(j, k)$  and  $p \notin I(i, j) \cap I(i, k)$ . Clearly we have  $c_p^i \neq c_p^j = c_p^k$ . Hence  $p \notin I(i, j) \cup I(i, k)$  and  $p \in \overline{I(i, j) \cup I(i, k)}$ . It follows that

$$I(j, k) \subseteq (I(i, j) \cap I(i, k)) \cup \overline{I(i, j) \cup I(i, k)}.$$

This completes the proof. □

Let  $\mathcal{F}$  be a family of subsets of an  $n$ -set  $E$ . Let

$$\begin{aligned} l &= \min\{|F| : F \in \mathcal{F}\}, \\ u &= \max\{|F| : F \in \mathcal{F}\} \end{aligned}$$

be the minimum size and maximum size of subsets of  $\mathcal{F}$ . For  $r \geq u$  and  $s \leq l$ , the families

$$\begin{aligned} \nabla_r(\mathcal{F}) &= \{B \subseteq E : |B| = r, \exists F \in \mathcal{F}, F \subseteq B\}, \\ \Delta_s(\mathcal{F}) &= \{B \subseteq E : |B| = s, \exists F \in \mathcal{F}, B \subseteq F\} \end{aligned}$$

are called the  $r$ -shade and  $s$ -shadow of  $\mathcal{F}$ , respectively. When  $\mathcal{F}$  is a family of  $k$ -subsets, the  $(k+1)$ -shade and the  $(k-1)$ -shadow are simply written as  $\nabla(\mathcal{F})$  or  $\Delta(\mathcal{F})$ .

**Lemma 2.8.** [20] If  $\mathcal{A}$  is an intersecting family of  $k$ -subsets of an  $n$ -set, then  $|\Delta\mathcal{A}| \geq |\mathcal{A}|$ .

**Lemma 2.9.** [16, Corollary 2.3.2] Let  $\mathcal{F}$  be a family of  $k$ -subsets of an  $n$ -set where  $k < n$  and  $n \geq 3$ .

(1) If  $k \geq \lceil \frac{n}{2} + 1 \rceil$ , then  $|\Delta\mathcal{F}| - |\mathcal{F}| \geq k - 1 \geq \lceil \frac{n}{2} \rceil$ .

(2) If  $k \leq \lfloor \frac{n}{2} - 1 \rfloor$ , then  $|\nabla\mathcal{F}| - |\mathcal{F}| \geq n - k - 1 \geq \lceil \frac{n}{2} \rceil$ .

**Theorem 2.10.** Let  $\mathcal{F}$  be a Sperner family over an  $n$ -set and let  $l \leq \frac{n}{2} \leq u$ , where  $l$  and  $u$  are the minimum size and the maximum size of subsets in  $\mathcal{F}$ , respectively.

(i) If  $l = \lfloor \frac{n}{2} \rfloor$  and  $u = \lceil \frac{n}{2} \rceil$ , then  $|\mathcal{F}| \leq \binom{n}{\lfloor \frac{n}{2} \rfloor}$ .

(ii) If  $l < \lfloor \frac{n}{2} \rfloor$  and  $u > \lceil \frac{n}{2} \rceil$ , then

$$|\mathcal{F}| \leq \begin{cases} \binom{n}{\frac{n}{2}} - (u-l)\frac{n}{2} - \lfloor \frac{(u-l-1)^2}{4} \rfloor, & \text{if } n \text{ is even,} \\ \binom{\frac{n-1}{2}}{\frac{n-1}{2}} - (u-l-1)\frac{n+1}{2} - \lfloor \frac{(u-l-2)^2}{4} \rfloor, & \text{if } n \text{ is odd.} \end{cases}$$

(iii) If  $l < \lfloor \frac{n}{2} \rfloor$  and  $u = \lceil \frac{n}{2} \rceil$ , then  $|\mathcal{F}| \leq \binom{n}{\lfloor \frac{n}{2} \rfloor} - (\lfloor \frac{n}{2} \rfloor - l)\lceil \frac{n}{2} \rceil - \frac{(\lfloor \frac{n}{2} \rfloor - l)(\lfloor \frac{n}{2} \rfloor - l - 1)}{2}$ .

(iv) If  $l = \lfloor \frac{n}{2} \rfloor$  and  $u > \lceil \frac{n}{2} \rceil$ , then  $|\mathcal{F}| \leq \binom{n}{\lfloor \frac{n}{2} \rfloor} - (u - \lceil \frac{n}{2} \rceil)\lceil \frac{n}{2} \rceil - \frac{(u - \lceil \frac{n}{2} \rceil)(u - \lceil \frac{n}{2} \rceil - 1)}{2}$ .

*Proof.* This is an adaption of [16, Corollary 2.3.3]. The cases  $n = 1, 2$  are trivial, hence we let  $n \geq 3$ . The statement (i) follows immediately from Sperner's Theorem.

In the case that  $l < \lfloor \frac{n}{2} \rfloor$  and  $u > \lceil \frac{n}{2} \rceil$ , we proceed in two steps.

**Step 1:** Replace  $\mathcal{F}$  by  $\mathcal{F}_1 = (\mathcal{F} \setminus \mathcal{G}) \cup \nabla(\mathcal{G})$  where  $\mathcal{G} = \mathcal{F} \cap \binom{[n]}{l}$ . Because  $\mathcal{F}$  is Sperner, we have that  $(\mathcal{F} \setminus \mathcal{G}) \cap \nabla(\mathcal{G}) = \emptyset$  and that  $\mathcal{F}_1$  is a Sperner family for which by Lemma 2.9 we have

$$|\mathcal{F}_1| = |\mathcal{F}| - |\mathcal{G}| + |\nabla(\mathcal{G})| \geq |\mathcal{F}| + n - l - 1.$$

If  $l + 1 \leq \lfloor \frac{n}{2} - 1 \rfloor$  then replace  $\mathcal{F}_1$  by  $\mathcal{F}_2 = (\mathcal{F}_1 \setminus \mathcal{G}_1) \cup \nabla(\mathcal{G}_1)$  where  $\mathcal{G}_1 = \mathcal{F}_1 \cap \binom{[n]}{l+1}$ . After this we obtain a Sperner family  $\mathcal{F}_2$  for which by Lemma 2.9

$$|\mathcal{F}_2| \geq |\mathcal{F}_1| + n - l - 2 \geq |\mathcal{F}| + (n - l - 1) + (n - l - 2).$$

Repeat doing like this until we raise the minimum size of the subsets to  $\lfloor \frac{n}{2} \rfloor$  and we obtain a Sperner family  $\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l}$  satisfying

$$|\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l}| \geq |\mathcal{F}| + (n - l - 1) + (n - l - 2) + \cdots + \lceil \frac{n}{2} \rceil. \quad (1)$$

**Step 2:** We begin to decrease the maximum size of subsets of  $\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l}$ . Replace  $\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l}$  by  $\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l + 1} = (\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l} \setminus \mathcal{H}) \cup \Delta(\mathcal{H})$  where  $\mathcal{H} = \mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l} \cap \binom{[n]}{u}$ . Then we obtain a Sperner family  $\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l + 1}$  for which by Lemma 2.9 we have

$$|\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l + 1}| \geq |\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l}| + u - 1.$$

If  $u - 1 \geq \lceil \frac{n}{2} + 1 \rceil$  then replace  $\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l + 1}$  by  $\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l + 2} = (\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l + 1} \setminus \mathcal{H}_1) \cup \Delta(\mathcal{H}_1)$  where  $\mathcal{H}_1 = \mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l + 1} \cap \binom{[n]}{u-1}$ . Similarly we have

$$|\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l + 2}| \geq |\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l + 1}| + u - 2 \geq |\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l}| + (u - 1) + (u - 2).$$

Repeat this process until we obtain a Sperner family  $\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l + u - \lceil \frac{n}{2} \rceil}$  with maximum size of the subsets being  $\lceil \frac{n}{2} \rceil$  (and all sizes of the subsets being  $\lfloor \frac{n}{2} \rfloor$  or  $\lceil \frac{n}{2} \rceil$ ) and we have

$$\binom{n}{\lfloor \frac{n}{2} \rfloor} \geq |\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l + u - \lceil \frac{n}{2} \rceil}| \geq |\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l}| + (u - 1) + (u - 2) + \cdots + \lceil \frac{n}{2} \rceil. \quad (2)$$

Combining inequality (2) with (1) yields

$$\binom{n}{\lfloor \frac{n}{2} \rfloor} \geq |\mathcal{F}| + (n-l-1) + (n-l-2) + \cdots + \lceil \frac{n}{2} \rceil + (u-1) + (u-2) + \cdots + \lceil \frac{n}{2} \rceil. \quad (3)$$

Whenever  $n$  is even, we further bound (3) by

$$\begin{aligned} \binom{n}{\frac{n}{2}} &\geq |\mathcal{F}| + (u-l)\frac{n}{2} + (1+2+\cdots + (\lfloor \frac{u-l}{2} \rfloor - 1)) + (1+2+\cdots + (\lceil \frac{u-l}{2} \rceil - 1)) \\ &= |\mathcal{F}| + (u-l)\frac{n}{2} + \lfloor \frac{(u-l-1)^2}{4} \rfloor. \end{aligned}$$

Whenever  $n$  is odd, similarly we have

$$\begin{aligned} \binom{n}{\frac{n-1}{2}} &\geq |\mathcal{F}| + (u-l-1)\frac{n+1}{2} + (1+2+\cdots + (\lfloor \frac{u-l-1}{2} \rfloor - 1)) \\ &\quad + (1+2+\cdots + (\lceil \frac{u-l-1}{2} \rceil - 1)) \\ &= |\mathcal{F}| + (u-l-1)\frac{n+1}{2} + \lfloor \frac{(u-l-2)^2}{4} \rfloor. \end{aligned}$$

Then the statement (ii) follows immediately.

Finally we consider (iii) and (iv). If  $l < \lfloor \frac{n}{2} \rfloor$  and  $u = \lceil \frac{n}{2} \rceil$ , then we only need to proceed Step 1 and the Sperner family  $\mathcal{F}_{\lfloor \frac{n}{2} \rfloor - l}$  consists of subsets of sizes  $\lfloor \frac{n}{2} \rfloor$  and  $\lceil \frac{n}{2} \rceil$ . Thus from inequality (1) we have

$$\begin{aligned} \binom{n}{\lfloor \frac{n}{2} \rfloor} &\geq |\mathcal{F}| + (n-l-1) + (n-l-2) + \cdots + \lceil \frac{n}{2} \rceil \\ &= |\mathcal{F}| + (\lfloor \frac{n}{2} \rfloor - l)\lceil \frac{n}{2} \rceil + \frac{(\lceil \frac{n}{2} \rceil - l)(\lceil \frac{n}{2} \rceil - l - 1)}{2}, \end{aligned}$$

proving (iii). Similarly we only proceed Step 2 and prove (iv) for the case that  $l = \lfloor \frac{n}{2} \rfloor$  and  $u > \lceil \frac{n}{2} \rceil$ .  $\square$

**Corollary 2.11.** *Let  $\mathcal{F}$  be a Sperner family over an  $n$ -set. Let  $l$  and  $u$  be the minimum size and the maximum size of subsets in  $\mathcal{F}$ , respectively. For  $l \leq i \leq u$ , define*

$$\begin{aligned} \mathcal{F}_+^i &= \{B : B \in \mathcal{F}, |B| \geq i\}, \\ \mathcal{F}_-^i &= \{B : B \in \mathcal{F}, |B| \leq i\}. \end{aligned}$$

(i) *If  $l \leq \lfloor \frac{n}{2} \rfloor$ , then  $|\nabla_i(\mathcal{F}_-^i)| \geq |\mathcal{F}_-^i| + (i-l)(n-i)$  for any  $l \leq i \leq \lfloor \frac{n}{2} \rfloor$ .*

(ii) *If  $u \geq \lceil \frac{n}{2} \rceil$ , then  $|\Delta_i(\mathcal{F}_+^i)| \geq |\mathcal{F}_+^i| + i(u-i)$  for any  $\lceil \frac{n}{2} \rceil \leq i \leq u$ .*

*Proof.* Obviously the conclusion (i) holds if  $l = \lfloor \frac{n}{2} \rfloor$  and (ii) holds if  $u = \lceil \frac{n}{2} \rceil$ . So we let  $l < \lfloor \frac{n}{2} \rfloor$  in (i) and let  $u > \lceil \frac{n}{2} \rceil$  in (ii).

(i) Analogous to the proof of inequality (1) of Theorem 2.10, we increase the minimum size of the subsets of  $\mathcal{F}_-^i$  to  $i$  step by step and then we have

$$|\nabla_i(\mathcal{F}_-^i)| \geq |\mathcal{F}_-^i| + (n-l-1) + (n-l-2) + \cdots + (n-i) \geq |\mathcal{F}_-^i| + (i-l)(n-i).$$

(ii) Analogous to the proof of inequality (2) of Theorem 2.10, we decrease the maximum size of the subsets of  $\mathcal{F}_+^i$  to  $i$  step by step and then we have

$$|\Delta_i(\mathcal{F}_+^i)| \geq |\mathcal{F}_+^i| + (u-1) + (u-2) + \dots + i \geq |\mathcal{F}_+^i| + i(u-i).$$

□

### 3 Improved upper bounds

Theorem 2.6 establishes the relationship between a 2-wFP code and non 2-covering Sperner families generated by all codewords. Improved upper bounds on the size of a 2-wFP code will be developed in this section.

#### 3.1 Length even

**Lemma 3.1.** *Let  $n \geq 6$  be even and  $\mathcal{F}$  a non 2-covering Sperner family over an  $n$ -set. Denote  $l$  and  $u$  to be the minimum size and the maximum size of subsets in  $\mathcal{F}$ , respectively.*

- (i) *If  $u \geq \frac{n}{2} + 1$ , then  $|\mathcal{F}| \leq \binom{\frac{n}{2}}{\frac{n}{2}-1} - \frac{n}{2}$ .*
- (ii) *If  $l \leq \frac{n}{2} - 2$ , then  $|\mathcal{F}| \leq \binom{\frac{n}{2}}{\frac{n}{2}-1} - \frac{n}{2} - 1$ .*
- (iii) *If  $u = l = \frac{n}{2}$ , then  $|\mathcal{F}| \leq \frac{1}{2} \binom{n}{\frac{n}{2}}$ .*

*Proof.* Let  $\mathcal{F} = \mathcal{A} \cup \mathcal{B}$  be a non 2-covering Sperner family on  $[n]$ , where

$$\begin{aligned}\mathcal{A} &= \{A : A \in \mathcal{F}, |A| \geq \frac{n}{2}\}, \\ \mathcal{B} &= \{B : B \in \mathcal{F}, |B| \leq \frac{n}{2} - 1\}.\end{aligned}$$

- (i) Let  $u \geq \frac{n}{2} + 1$ . By Corollary 2.11, we have

$$|\Delta_{\frac{n}{2}}(\mathcal{A})| \geq |\mathcal{A}| + (u - \frac{n}{2})\frac{n}{2} \geq |\mathcal{A}| + \frac{n}{2}.$$

Denote  $\mathcal{P} = \Delta_{\frac{n}{2}}(\mathcal{A})$ . Because  $\mathcal{F}$  is non 2-covering, we have that  $\mathcal{P}$  is intersecting. Then by Lemma 2.8 we have

$$|\Delta(\mathcal{P})| \geq |\mathcal{P}| \geq |\mathcal{A}| + \frac{n}{2}.$$

By Theorem 2.4, all subsets of  $[n]$  can be partitioned into  $\binom{\frac{n}{2}}{\frac{n}{2}}$  mutually disjoint symmetric chains. If  $\mathcal{B} \neq \emptyset$ , then replace each  $B \in \mathcal{B}$  with  $B'$  in the same symmetric chain, where  $B \subseteq B'$  and  $|B'| = \frac{n}{2} - 1$ . Thus we produce from  $\mathcal{B}$  a new Sperner family  $\mathcal{B}'$  of  $(\frac{n}{2} - 1)$ -subsets. Note also that  $\Delta(\mathcal{P}) \cap \mathcal{B}' = \emptyset$  because  $\mathcal{F} = \mathcal{A} \cup \mathcal{B}$  is Sperner. (Let  $\mathcal{B}' = \emptyset$  if  $\mathcal{B} = \emptyset$ .) As a result,

$$|\mathcal{B}| = |\mathcal{B}'| \leq \binom{\frac{n}{2}}{\frac{n}{2}-1} - |\Delta(\mathcal{P})|.$$

It follows that

$$|\mathcal{F}| = |\mathcal{A}| + |\mathcal{B}| \leq |\mathcal{A}| + \binom{\frac{n}{2}}{\frac{n}{2}-1} - |\Delta(\mathcal{P})| \leq \binom{\frac{n}{2}}{\frac{n}{2}-1} - \frac{n}{2}.$$

(ii) Let  $l \leq \frac{n}{2} - 2$ . Apply Corollary 2.11 (i) to  $\mathcal{B}$ . It follows that

$$|\nabla_{\frac{n}{2}-1}(\mathcal{B})| \geq |\mathcal{B}| + (\frac{n}{2} - 1 - l)(n - (\frac{n}{2} - 1)) \geq |\mathcal{B}| + \frac{n}{2} + 1.$$

In the decomposition of the power set of  $[n]$  into symmetric chains, if  $\mathcal{A} \neq \emptyset$ , then replace each  $A \in \mathcal{A}$  by  $A'$  of the same symmetric chain where  $|A'| = \frac{n}{2}$  to obtain a new family  $\mathcal{A}'$  of  $\frac{n}{2}$ -sets. Since  $\mathcal{F}$  is non 2-covering,  $\mathcal{A}'$  is intersecting and thus  $|\Delta(\mathcal{A}')| \geq |\mathcal{A}'| = |\mathcal{A}|$  by Lemma 2.8. Furthermore, it is easy to see that  $\Delta(\mathcal{A}')$  and  $\nabla_{\frac{n}{2}-1}(\mathcal{B})$  are disjoint because  $\mathcal{F}$  is Sperner. (Let  $\mathcal{A}' = \emptyset$  if  $\mathcal{A} = \emptyset$ .) It follows that

$$|\mathcal{F}| = |\mathcal{A}| + |\mathcal{B}| \leq |\Delta(\mathcal{A}')| + |\nabla_{\frac{n}{2}-1}(\mathcal{B})| - \frac{n}{2} - 1 \leq \binom{n}{\frac{n}{2}-1} - \frac{n}{2} - 1.$$

(iii) If  $u = l = \frac{n}{2}$ , then  $|\mathcal{F}| \leq \frac{1}{2} \binom{n}{\frac{n}{2}}$  because  $\mathcal{F}$  is non 2-covering.  $\square$

For a family  $\mathcal{F}$  of subsets of  $[n]$ , we define its *complement* by  $\overline{\mathcal{F}} = \{\overline{F} : F \in \mathcal{F}\}$ .

**Theorem 3.2.** *Let  $n$  be even and  $n \geq 8$ . Suppose that  $\mathcal{C}$  is an  $(n, m, q)$  2-wFP code. Then*

$$m \leq \binom{n}{\frac{n}{2}-1} - \frac{n}{2} + 1.$$

*Proof.* For  $1 \leq i \leq m$ , let  $\mathcal{X}_i$  be the coincidence family generated by the codeword  $c^i \in \mathcal{C}$ . Then each  $\mathcal{X}_i$  is a non 2-covering Sperner family by Theorem 2.6. Take a fixed  $i \in [m]$  and let  $l$  and  $u$  be the minimum size and the maximum size of subsets in  $\mathcal{X}_i$ , respectively.

By Lemma 3.1, if  $u \geq \frac{n}{2} + 1$ , then  $|\mathcal{X}_i| \leq \binom{n}{\frac{n}{2}-1} - \frac{n}{2}$ ; if  $l \leq \frac{n}{2} - 2$ , then  $|\mathcal{X}_i| \leq \binom{n}{\frac{n}{2}-1} - \frac{n}{2} - 1$ ; if  $u = l = \frac{n}{2}$ , then  $|\mathcal{X}_i| \leq \frac{1}{2} \binom{n}{\frac{n}{2}} = \binom{n-1}{\frac{n}{2}-1}$ . It is easy to show that  $m = |\mathcal{X}_i| + 1 \leq \binom{n}{\frac{n}{2}-1} - \frac{n}{2} + 1$  for these cases. So, to prove the conclusion, we only need to let  $l \geq \frac{n}{2} - 1$ ,  $u \leq \frac{n}{2}$ , and  $(u, l) \neq (\frac{n}{2}, \frac{n}{2})$ . Thus we only need to consider two cases  $u = l = \frac{n}{2} - 1$  and  $(u, l) = (\frac{n}{2}, \frac{n}{2} - 1)$ . Let

$$\begin{aligned} \mathcal{A} &= \{A : A \in \mathcal{X}_i, |A| = \frac{n}{2}\}, \\ \mathcal{B} &= \{A : A \in \mathcal{X}_i, |A| = \frac{n}{2} - 1\}. \end{aligned}$$

Case 1: Let  $u = l = \frac{n}{2} - 1$ . Then  $\mathcal{X}_i = \mathcal{B}$ . We evaluate the upper bound of  $m$  by considering whether  $\mathcal{B}$  is intersecting.

If  $\mathcal{B}$  is intersecting, then consider its complement  $\overline{\mathcal{B}} = \{\overline{B} : B \in \mathcal{B}\}$ . For any  $\overline{A}, \overline{B} \in \overline{\mathcal{B}}$ ,

$$|\overline{A} \cap \overline{B}| = n - |A \cup B| = n - (|A| + |B| - |A \cap B|) = n - (n - 2 - |A \cap B|) \geq 3.$$

Consequently  $\overline{\mathcal{B}}$  is a 3-intersecting Sperner family. By Theorem 2.5, we have

$$m = |\mathcal{X}_i| + 1 = |\mathcal{B}| + 1 \leq \binom{n}{\frac{n}{2}-2} + 1.$$

If  $\mathcal{B}$  is not intersecting, then there exist  $B_1, B_2 \in \mathcal{B}$  such that  $B_1 \cap B_2 = \emptyset$ . Suppose that  $B_1 = I(i, j)$  and  $B_2 = I(i, k)$  where  $i \neq j, k$  and  $1 \leq j, k \leq m$ . By Lemma 2.7,

$$I(j, k) \subseteq (B_1 \cap B_2) \cup \overline{B_1 \cup B_2} = \overline{B_1 \cup B_2}$$



and hence  $|I(j, k)| \leq 2$ . If  $|I(j, k)| = 0$  then  $\mathcal{X}_j$  is not Sperner, contradicting Theorem 2.6. So we have  $|I(j, k)| = 1, 2$ , meaning that  $\mathcal{X}_j$  contains a set of size 1 or 2. Obviously if  $\mathcal{X}_j$  contains a singleton, then by Proposition 2.3 we have  $|\mathcal{X}_j| \leq \binom{n-1}{\frac{n}{2}-1} + 1$  and hence  $m = |\mathcal{X}_j| + 1 \leq \binom{n-1}{\frac{n}{2}-1} + 2$ . If  $|I(j, k)| = 2$ , then the minimum size  $l_j$  of elements of  $\mathcal{X}_j$  satisfies  $l_j \leq 2 \leq \frac{n}{2} - 2$  whenever  $n \geq 8$ . Hence  $|\mathcal{X}_j| \leq \binom{n}{\frac{n}{2}-1} - \frac{n}{2} - 1$  by Lemma 3.1. Thus  $m \leq \binom{n}{\frac{n}{2}-1} - \frac{n}{2}$ . Comparing the upper bounds of  $m$  for  $n \geq 8$  shows  $m \leq \binom{n}{\frac{n}{2}-1} - \frac{n}{2}$  in Case 1.

Case 2: Let  $u = \frac{n}{2}$  and  $l = \frac{n}{2} - 1$ . Then  $\mathcal{X}_i = \mathcal{A} \cup \mathcal{B}$ .

If  $\mathcal{B}$  is not intersecting, then we have  $m \leq \binom{n}{\frac{n}{2}-1} - \frac{n}{2}$  by similar discussions in Case 1.

If  $\mathcal{A}$  and  $\mathcal{B}$  are not cross-intersecting, then there exist  $A \in \mathcal{A}$  and  $B \in \mathcal{B}$  such that  $A \cap B = \emptyset$ . Suppose that  $A = I(i, j)$  and  $B = I(i, k)$  where  $i \neq j, k$  and  $1 \leq j, k \leq m$ . By Lemma 2.7,  $|I(j, k)| \leq |\overline{A \cup B}| = 1$ . Since  $\mathcal{X}_j$  is Sperner, we have  $|I(j, k)| = 1$ . Then by Proposition 2.3 we have  $|\mathcal{X}_j| \leq \binom{n-1}{\frac{n}{2}-1} + 1$  and hence  $m \leq \binom{n-1}{\frac{n}{2}-1} + 2$ .

What remains to bound  $m$  is the subcase that  $\mathcal{B}$  is intersecting and that  $\mathcal{A}$  and  $\mathcal{B}$  are cross-intersecting. By Lemma 2.9,  $|\nabla(\mathcal{B})| \geq |\mathcal{B}| + \frac{n}{2}$ . Note that  $\mathcal{X}_i$  is Sperner and non 2-covering. As a result, we have that  $\mathcal{A} \cap \nabla(\mathcal{B}) = \emptyset$  and that  $\mathcal{F} := \mathcal{A} \cup \nabla(\mathcal{B})$  is a non 2-covering family of  $\frac{n}{2}$ -subsets. Hence

$$\frac{1}{2} \binom{n}{\frac{n}{2}} \geq |\mathcal{F}| = |\mathcal{A} \cup \nabla(\mathcal{B})| = |\mathcal{A}| + |\nabla(\mathcal{B})| \geq |\mathcal{A}| + |\mathcal{B}| + \frac{n}{2},$$

yielding that

$$m = |\mathcal{X}_i| + 1 = |\mathcal{A}| + |\mathcal{B}| + 1 \leq \frac{1}{2} \binom{n}{\frac{n}{2}} - \frac{n}{2} + 1 = \binom{n-1}{\frac{n}{2}-1} - \frac{n}{2} + 1.$$

It follows that  $m \leq \binom{n-1}{\frac{n}{2}-1} + 2$  in Case 2.

To sum up it is immediate that for even  $n \geq 8$  we have

$$m \leq \binom{n}{\frac{n}{2}-1} - \frac{n}{2} + 1.$$

This completes the proof.  $\square$

### 3.2 Length odd

**Lemma 3.3.** *Let  $n$  be odd and  $n \geq 7$ . Suppose that  $\mathcal{F}$  is a non 2-covering Sperner family on  $[n]$ . If  $|F| \geq \frac{n+1}{2}$  for all  $F \in \mathcal{F}$ , then*

$$|\mathcal{F}| \leq \binom{n}{\frac{n+3}{2}}.$$

*Proof.* Since  $\mathcal{F}$  is a non 2-covering Sperner family, for any  $A, B \in \mathcal{F}$  we have

$$n - 1 \geq |A \cup B| = |A| + |B| - |A \cap B| \geq n + 1 - |A \cap B|,$$

implying  $|A \cap B| \geq 2$ . As a result  $\mathcal{F}$  is a 2-intersecting Sperner family. Then applying Theorem 2.5 yields the conclusion.  $\square$

**Lemma 3.4.** *Let  $n$  be odd and  $n \geq 7$ . Suppose that  $\mathcal{C}$  is an  $(n, m, q)$  2-wFP code and  $\mathcal{X}_i$  is the coincidence family generated by the codeword  $c^i \in \mathcal{C}$ . If there is  $i \in [m]$  such that  $|F| \leq \frac{n-1}{2}$  for all  $F \in \mathcal{X}_i$ , then*

$$m \leq \binom{n}{\frac{n+3}{2}} + 1.$$

*Proof.* By Theorem 2.6,  $\mathcal{X}_i$  is a non 2-covering Sperner family. Let  $\mathcal{X}_i = \mathcal{A}_0 \cup \mathcal{A}_1$ , where

$$\begin{aligned}\mathcal{A}_0 &= \{A : A \in \mathcal{X}_i, |A| \leq \frac{n-3}{2}\}, \\ \mathcal{A}_1 &= \{A : A \in \mathcal{X}_i, |A| = \frac{n-1}{2}\}.\end{aligned}$$

If  $\mathcal{A}_1$  is not intersecting, then there exist  $I(i, j), I(i, k) \in \mathcal{A}_1$  ( $1 \leq j, k \leq m, i \neq j, k$ ) such that  $I(i, j) \cap I(i, k) = \emptyset$ . Hence by Lemma 2.7 we have  $I(j, k) \subseteq \overline{I(i, j)} \cup \overline{I(i, k)}$ . Because  $\mathcal{X}_i$  is Sperner, we know that  $|I(j, k)| = 1$ . Apply Proposition 2.3 to have

$$m = |\mathcal{X}_j| + 1 \leq \binom{n-1}{\frac{n-1}{2}} + 2.$$

If  $\mathcal{A}_1$  is intersecting, then let  $\mathcal{F} = \overline{\mathcal{A}_0} \cup \overline{\mathcal{A}_1}$ . Similarly to the proof of Lemma 3.3, we can check that  $\mathcal{F}$  forms a 2-intersecting Sperner family. Hence by Theorem 2.5 we have

$$m = |\mathcal{X}_i| + 1 = |\mathcal{F}| + 1 \leq \binom{n}{\frac{n+3}{2}} + 1.$$

Noting that  $\binom{n-1}{\frac{n-1}{2}} \leq \binom{n}{\frac{n+3}{2}}$  if  $n \geq 7$  yields the conclusion.  $\square$

**Theorem 3.5.** *Let  $\mathcal{C}$  be an  $(n, m, q)$  2-wFP code with  $n$  odd and  $n \geq 7$ . Then*

$$m \leq \begin{cases} \binom{\frac{n-1}{2}}{\frac{n-1}{2}} - \frac{n^2-9}{8} - \lfloor \frac{(n-5)^2}{64} \rfloor, & \text{if } n \equiv 1 \pmod{4}, \\ \binom{\frac{n-1}{2}}{\frac{n-1}{2}} - \frac{(n+1)^2-8}{8} - \lfloor \frac{(n-3)^2}{64} \rfloor, & \text{if } n \equiv 3 \pmod{4}. \end{cases}$$

*Proof.* Let  $\mathcal{C}$  be an  $(n, m, q)$  2-wFP code and  $\mathcal{X}_i$  be the non 2-covering Sperner family generated by the codeword  $c^i$ . Denote

$$\begin{aligned}l_i &= \min\{|A| : A \in \mathcal{X}_i\}, \\ u_i &= \max\{|A| : A \in \mathcal{X}_i\}, \\ d_i &= u_i - l_i.\end{aligned}$$

If there is  $1 \leq i \leq m$  such that  $u_i \leq \frac{n-1}{2}$  or  $l_i \geq \frac{n+1}{2}$ , then by Lemmas 3.3 and 3.4

$$m = |\mathcal{X}_i| + 1 \leq \binom{n}{\frac{n+3}{2}} + 1. \quad (4)$$

Next we let  $l_i \leq \frac{n-1}{2}$  and  $u_i \geq \frac{n+1}{2}$  for all  $1 \leq i \leq m$ . Denote  $d = \min\{d_i : 1 \leq i \leq m\}$  and assume w.l.o.g.  $d_1 = d$ . In the case that  $d \geq \frac{n+1}{2}$ , by Theorem 2.10, it is easy to see that

$$m = |\mathcal{X}_1| + 1 \leq \binom{n}{\frac{n-1}{2}} - \frac{n^2 - 1}{4} - \lfloor \frac{(n-3)^2}{16} \rfloor + 1. \quad (5)$$

In the following proof we let  $d \leq \frac{n-1}{2}$  and then bound  $m$ . Let  $\mathcal{X}_1 = \mathcal{A}_1 \cup \mathcal{A}_2 \cup \mathcal{B}$ , where

$$\begin{aligned} \mathcal{A}_1 &= \{A : A \in \mathcal{X}_1, |A| \leq \frac{n-3}{2}\}, \\ \mathcal{A}_2 &= \{A : A \in \mathcal{X}_1, |A| = \frac{n-1}{2}\}, \\ \mathcal{B} &= \{B : B \in \mathcal{X}_1, |B| \geq \frac{n+1}{2}\}. \end{aligned}$$

We consider three cases as follows.

Case 1: Let  $\mathcal{A}_2 \neq \emptyset$  be not intersecting. Then there exist  $A, B \in \mathcal{A}_2$  such that  $A \cap B = \emptyset$  and  $|A \cup B| = n - 1$ . Suppose that  $A = I(1, j)$  and  $B = I(1, k)$ . Then by Lemma 2.7 we have  $I(j, k) \subseteq (A \cap B) \cup \overline{A \cup B} = \overline{A \cup B}$ , meaning that  $\mathcal{X}_j$  contains a singleton or  $\emptyset$ . Since  $\mathcal{X}_j$  is Sperner, it contains a singleton and hence by Proposition 2.3 we have

$$m = |\mathcal{X}_1| + 1 \leq \binom{n-1}{\frac{n-1}{2}} + 2. \quad (6)$$

Case 2: Let  $\mathcal{A}_1 \neq \emptyset$  and let  $\mathcal{A}_1$  and  $\mathcal{B}$  be not cross-intersecting. Then there exist  $A \in \mathcal{A}_1, B \in \mathcal{B}$  such that  $A \cap B = \emptyset$ . Clearly we have  $|A| \geq \frac{n+1}{2} - d$  ( $2 \leq d \leq \frac{n-1}{2}$ ) and  $|B| \geq \frac{n+1}{2}$ . Suppose that  $A = I(1, j)$  and  $B = I(1, k)$ . Then by Lemma 2.7 we have  $I(j, k) \subseteq \overline{A \cup B}$ . It follows that  $|I(j, k)| \leq n - (\frac{n+1}{2} - d + \frac{n+1}{2}) = d - 1$ , meaning that  $\mathcal{X}_j$  contains an  $r$ -subset where  $1 \leq r \leq d - 1$ . Hence we have  $d_j \geq \frac{n+1}{2} - d + 1$ . Now we apply Theorem 2.10 to bound  $m$ . Let  $d_0$  be an integer with  $1 \leq d_0 \leq \frac{n-3}{2}$ .

Whenever  $d \geq d_0 + 1$ , by Theorem 2.10, noting  $\frac{(d_0-1)(d_0-2)}{2} \geq \lfloor \frac{(d_0-1)^2}{4} \rfloor$ , we have

$$|\mathcal{X}_1| \leq \binom{n}{\frac{n-1}{2}} - d_0 \frac{n+1}{2} - \lfloor \frac{(d_0-1)^2}{4} \rfloor.$$

Whenever  $d \leq d_0$ , we have  $d_j \geq \frac{n+1}{2} - d_0 + 1$  and similarly we have

$$|\mathcal{X}_j| \leq \binom{n}{\frac{n-1}{2}} - (\frac{n+1}{2} - d_0) \frac{n+1}{2} - \lfloor \frac{(\frac{n-1}{2} - d_0)^2}{4} \rfloor.$$

In order to get a better upper bound of  $m = |\mathcal{X}_1| + 1 = |\mathcal{X}_j| + 1$ , we take  $d_0 = \frac{n-1}{4}$  if  $n \equiv 1 \pmod{4}$  and  $d_0 = \frac{n+1}{4}$  if  $n \equiv 3 \pmod{4}$ . By simple reduction we have

$$m \leq \begin{cases} \left( \binom{n}{\frac{n-1}{2}} - \frac{n^2-1}{8} - \lfloor \frac{(n-5)^2}{64} \rfloor \right) + 1, & \text{if } n \equiv 1 \pmod{4}, \\ \left( \binom{n}{\frac{n-1}{2}} - \frac{(n+1)^2}{8} - \lfloor \frac{(n-3)^2}{64} \rfloor \right) + 1, & \text{if } n \equiv 3 \pmod{4}. \end{cases} \quad (7)$$

Case 3: Let  $\mathcal{A}_2$  be intersecting if  $\mathcal{A}_2 \neq \emptyset$  and let  $\mathcal{A}_1$  and  $\mathcal{B}$  be cross-intersecting if  $\mathcal{A}_1 \neq \emptyset$ . Define

$$\mathcal{F} = \overline{\mathcal{A}_1} \cup \overline{\mathcal{A}_2} \cup \mathcal{B},$$

where we let  $\bar{\emptyset} = \emptyset$  if necessary. Then we claim that  $\mathcal{F}$  is a 2-intersecting Sperner family.

Obviously each family of  $\overline{\mathcal{A}}_1, \overline{\mathcal{A}}_2$  and  $\mathcal{B}$  is Sperner. If  $\mathcal{F}$  is not Sperner, then one of the following three possibilities would happen: (a) there is  $A_1 \in \mathcal{A}_1$  and  $A_2 \in \mathcal{A}_2$  such that  $\overline{A_2} \subseteq \overline{A_1}$ , yielding  $A_1 \subseteq A_2$  and contradicting the fact that  $\mathcal{X}_1$  is Sperner; (b) there is  $A \in \mathcal{A}_1$  and  $B \in \mathcal{B}$  such that  $B \subseteq \overline{A}$  or  $\overline{A} \subseteq B$ , yielding  $A \cap B = \emptyset$  or  $A \cup B \supseteq A \cup \overline{A} = [n]$  and contradicting the fact that  $\mathcal{A}_1$  and  $\mathcal{B}$  are cross-intersecting or that  $\mathcal{X}_1$  is non 2-covering; (c) there is  $A \in \mathcal{A}_2$  and  $B \in \mathcal{B}$  such that  $\overline{A} \subseteq B$ , yielding  $A \cup B \supseteq A \cup \overline{A} = [n]$  and contradicting the fact that  $\mathcal{X}_1$  is non 2-covering. It follows that  $\mathcal{F}$  is Sperner.

Next we show that  $\mathcal{F}$  is 2-intersecting. Noting the size of subsets in each family  $\overline{\mathcal{A}}_1, \overline{\mathcal{A}}_2$  and  $\mathcal{B}$ , we readily check that (a)  $\overline{\mathcal{A}}_1$  is 3-intersecting, (b)  $\overline{\mathcal{A}}_2$  is 2-intersecting because  $\mathcal{A}_2$  is intersecting, (c)  $\mathcal{B}$  is 2-intersecting as  $\mathcal{B}$  is non 2-covering, (d)  $\overline{\mathcal{A}}_1$  and  $\overline{\mathcal{A}}_2$  are cross-2-intersecting because  $|A| + |B| \leq n - 2$  for  $A \in \mathcal{A}_1, B \in \mathcal{A}_2$ , and (e)  $\overline{\mathcal{A}}_1 \cup \overline{\mathcal{A}}_2$  and  $\mathcal{B}$  are cross-2-intersecting because for all  $A \in \mathcal{A}_1 \cup \mathcal{A}_2, B \in \mathcal{B}$  we have  $A \cap B \neq A$  ( $\mathcal{X}_i$  Sperner) and

$$|\overline{A} \cap B| = |B| - |A \cap B| \geq |B| - (|A| - 1) \geq \frac{n+1}{2} - \frac{n-3}{2} \geq 2.$$

Now that  $\mathcal{F}$  is a 2-intersecting Sperner family. So by Theorem 2.5 we have

$$m = |\mathcal{X}_1| + 1 = |\mathcal{F}| + 1 \leq \binom{n}{\frac{n+3}{2}} + 1. \quad (8)$$

Comparing the upper bounds in (4)-(8) when  $n \geq 7$  yields the conclusion.  $\square$

## 4 Concluding remarks

In the narrow-sense model, various types of fingerprinting codes were extensively studied by many researchers. However, as far as we know, frameproof codes is the only type of fingerprinting codes that was ever studied in the wide-sense model. We tried the standard probabilistic method (as in [24, 28]) and approaches of hypergraphs (as in [31]) to achieve lower bounds for wide-sense 2-frameproof codes. Unfortunately, we do not achieve a better lower bound than the bound of binary 2-FP codes in [28, Theorem 4.1]. Evaluating a good lower bound for 2-wFP code with general alphabet size is a focus of future work.

The main result of the paper is an improvement on the known upper bounds for 2-wFP codes by applying techniques on non 2-covering Sperner families and intersecting families in extremal set theory. The new bounds (as well as the previous ones by Panoui) do not relate with the alphabet size  $q$ . Not surprisingly, we learn from some small examples that  $q$  usually affects the size of a  $t$ -wFP code. It is worthwhile to examine how to improve the upper bounds by taking the alphabet size into consideration.

**Acknowledgements:** The authors would like to thank the anonymous referees for their helpful comments and valuable suggestions, which have greatly improved the presentation and quality of this paper.

## References

- [1] N. Alon, E. Fischer, and M. Szegedy, Parent-identifying codes, *J. Combinat. Theory A*, 95(2), 349-359, 2001.
- [2] I. Anderson, *Combinatorics of finite sets*, Dover publications, INC. Mineola, New York, 1987.
- [3] A. Barg, G. R. Blakley, and G. A. Kabatiansky, Digital fingerprinting codes: problem statements, constructions, identification of traitors, *IEEE Trans. Inf. Theory*, 49(4), 852-865, 2003.
- [4] A. Barg, G. Cohen, S. Encheva, G. Kabatiansky, and G. Zémor, A hypergraph approach to the identifying parent property: the case of multiple parents, *SIAM Journal on Discrete Mathematics*, 14(3), 423-431, 2001.
- [5] S. R. Blackburn, An upper bound on the size of a code with the  $k$ -identifiable parent property, *J. Combinat. Theory A*, 102(1), 179-185, 2003.
- [6] S. R. Blackburn, Combinatorial schemes for protecting digital content, *Surveys in combinatorics*, In: C.D. Wensley (ed.), London Mathematical Society lecture note series, 307, Cambridge University Press, 43-78, 2003.
- [7] S. R. Blackburn, Frameproof codes, *SIAM Journal on Discrete Mathematics*, 16(3), 499-510, 2003.
- [8] S. R. Blackburn, T. Etzion, and S.-L. Ng. Traceability codes, *J. Combinat. Theory A*, 117(8), 1049-1057, 2010.
- [9] D. Boneh and J. Shaw, Collusion-secure fingerprinting for digital data, *IEEE Trans. Inf. Theory*, 44(5), 1897-1905, 1998.
- [10] Y. Chee, Turán-type problems in group testing, coding theory and cryptography, Department of Computer Science, University of Waterloo, Canada, 1996.
- [11] Y. Chee and X. Zhang, Improved constructions of frameproof codes, *IEEE Trans. Inf. Theory*, 58(8), 5449-5453, 2012.
- [12] M. Cheng and Y. Miao, On anti-collusion codes and detection algorithms for multimedia fingerprinting, *IEEE Trans. Inf. Theory*, 57(7), 4843-4851, 2011.
- [13] B. Chor, A. Fiat, and M. Naor, Tracing traitors, in *Advances in Cryptology-CRYPTO'94*, In: Y.G. Desmedt (ed.), Lecture notes in comput. Sci. 839, Springer, Berlin, 257-270, 1994.
- [14] G. Cohen and S. Encheva, Efficient constructions of frameproof codes, *Electronics Letters*, 36(22), 1840-1842, 2000.
- [15] R. A. Duke, H. Lefmann, and V. Rödl, On uncrowded hypergraphs, *Random Structures and Algorithms*, 6(2-3), 209-212, 1995.
- [16] K. Engel, *Sperner theory (Encyclopedia of Mathematics and its applications book)*, UK: Cambridge University Press, 1997.

- [17] A. Fiat and T. Tassa, Dynamic traitor tracing, *Journal of Cryptology*, 14(3), 211-223, 2001.
- [18] C. Guo, D. R. Stinson, and T. Van Trung, On tight bounds for binary frameproof codes, *Designs, Codes and Cryptography*, 77(2-3), 301-319, 2015.
- [19] S. Jukna, *Extremal Combinatorics: with applications in computer science*, second edition, Springer-Verlag Berlin Heidelberg, 2011.
- [20] G. O. H. Katona, Intersection theorems for systems of finite sets, *Acta Mathematica Academiae Scientiarum Hungarica*, 15(3), 329-337, 1964.
- [21] E. C. Milner. A combinatorial theorem on systems of sets, *Journal of the London Mathematical Society*, s1-43(1), 204-206, 1968.
- [22] A. Panoui, Wide-sense fingerprinting codes and honeycomb arrays, Royal Holloway, University of London, 2012.
- [23] C. Shangguan, J. Ma, G. Ge, New upper bounds for parent-identifying codes and traceability codes, *Designs, Codes and Cryptography*, 86(8), 1727-1737, 2018.
- [24] C. Shangguan, X. Wang, G. Ge, and Y. Miao, New bounds for frameproof codes, *IEEE Trans. Inf. Theory*, 63(11), 7247-7252, 2017.
- [25] J. N. Staddon, D. R. Stinson, and R. Wei, Combinatorial properties of frameproof and traceability codes, *IEEE Trans. Inf. Theory*, 47(3), 1042-1049, 2001.
- [26] D. R. Stinson, T. van Trung, and R. Wei, Secure frameproof codes, key distribution patterns, group testing algorithms and related structures, *Journal of Statistical Planning and Inference*, 86(2), 595-617, 2000.
- [27] D. R. Stinson and R. Wei. Combinatorial properties and constructions of traceability schemes and frameproof codes, *SIAM Journal on Discrete Mathematics*, 11, 41-53, 1998.
- [28] D. R. Stinson, R. Wei, and K. Chen, On generalized separating hash families, *J. Combinat. Theory A*, 115(1), 105-120, 2008.
- [29] T. Van Trung, A tight bound for frameproof codes viewed in terms of separating hash families, *Designs, Codes and Cryptography*, 72(3), 713-718, 2014.
- [30] C. Xing, Asymptotic bounds on frameproof codes, *IEEE Trans. Inf. Theory*, 48(11), 2991-2995, 2002.
- [31] Y. Yang, Y. Zhang, and G. Ge, New lower bounds for secure codes and related hash families: a hypergraph theoretical approach, *IEEE Trans. Inf. Theory*, 63(4), 2446-2453, 2017.