

Quantitative estimate of the continuum approximations of interacting particle systems in one dimension

M. Kimura, P. van Meurs

Abstract

We consider a large class of interacting particle systems in 1D described by an energy whose interaction potential is singular and non-local. This class covers Riesz gases (in particular, log gases) and applications to plasticity and approximation theory of functions. While it is well established that the minimisers of such interaction energies converge to a certain particle density profile as the number of particles tends to infinity, any bound on the rate of this convergence is only known in special cases by means of quantitative estimates. The main result of this paper extends these quantitative estimates to a large class of interaction energies by a different proof. The proof relies on one-dimensional features such as the convexity of the interaction potential and the ordering of the particles. The main novelty of the proof is the treatment of the singularity of the interaction potential by means of a carefully chosen renormalisation.

Keywords: Interacting particle system, calculus of variations, asymptotic analysis
MSC: 82C22, 74Q05, 35A15, 74G10

1 Introduction

We are interested in quantifying the difference between minimisers of interacting particle energies and the minimisers of the related energies for the particle density. The interacting particle energies are given by

$$E_n(\mathbf{x}) = \frac{1}{n^2} \sum_{i=1}^n \sum_{j=0}^{i-1} V(x_i - x_j) + \frac{1}{n} \sum_{i=0}^n U(x_i), \quad (1)$$

where $n + 1$ is the number of particles, and

$$\mathbf{x} := (x_0, x_1, \dots, x_n) \in \Omega := \{\mathbf{y} \in \mathbb{R}^{n+1} : y_0 < y_1 < \dots < y_n\} \quad (2)$$

is the list of ordered particle positions. The energies E_n are the sum of two parts. We interpret the first part as the interaction part, in which V is the interaction potential, and the second part as a confinement term, in which U is the confining potential. Typical examples of V and U are plotted in Figure 1. We aim to keep the assumptions on V and U as weak as possible. These assumptions are as follows.

Assumption 1.1 (V and U). *The interaction potential $V \in L_{loc}^1(\mathbb{R})$ splits as $V = V_a + V_{\text{reg}}$, where*

$$V_a(x) := \begin{cases} -\log|x|, & \text{if } a = 0 \\ |x|^{-a}, & \text{if } 0 < a < 1, \end{cases} \quad (3)$$




Figure 1: Typical examples of V and U .

for a fixed parameter $a \in [0, 1)$, and $V_{\text{reg}} \in C^2(\mathbb{R})$ is such that

$$V \text{ even, } V \text{ convex on } (0, \infty), \quad \lim_{x \rightarrow \infty} \frac{V(x)}{x} = 0. \quad (4)$$

The domain of the confining potential $U : \mathbb{R} \rightarrow [0, \infty]$ is

$$D(U) := \{x \in \mathbb{R} : U(x) < \infty\} = \overline{(z_1, z_2)}$$

for some $-\infty \leq z_1 < z_2 \leq \infty$. It satisfies

$$U \in C^2(D(U)), \quad U \text{ convex on } \mathbb{R}, \quad \min_{\mathbb{R}} U = 0, \quad \lim_{|x| \rightarrow \infty} U(x) = \infty. \quad (5)$$

We interpret Assumption 1.1 as follows. We consider a as a fixed parameter which determines the singularity of V at 0. The part V_{reg} is a regular perturbation which determines the bulk and tails of V . Finite values for z_i correspond to impenetrable barriers for the particle positions.

Given E_n , the related energy for the particle density ρ is given by

$$E : \mathcal{P}(\mathbb{R}) \rightarrow \mathbb{R} \cup \{+\infty\}, \quad E(\rho) := \frac{1}{2} \int_{\mathbb{R}} \int_{\mathbb{R}} V(x-y) d\rho(y) d\rho(x) + \int_{\mathbb{R}} U(x) d\rho(x), \quad (6)$$

where $\mathcal{P}(\mathbb{R})$ is the space of probability measures.

For various choices of V and U , it is known (see, e.g., [ST97, GPPS13, vM18b, vMMP14]) that E_n and E attain their minimal value at some $\mathbf{x}^* \in \Omega$ and $\bar{\rho} \in \mathcal{P}(\mathbb{R})$ respectively, that $\bar{\rho} \in \mathcal{P}(\mathbb{R})$ is unique, and that any sequence of minimisers \mathbf{x}^* (parametrised by n) converges to $\bar{\rho}$ in a suitable topology as $n \rightarrow \infty$. Yet, any *quantitative* estimate between \mathbf{x}^* and $\bar{\rho}$ for finite n is only available for special choices of V and U (see Section 1.3). The aim of this paper is to derive such an estimate for the much larger class of potentials V and U characterised in Assumption 1.1.

In order to give meaning to a quantitative estimate between \mathbf{x}^* and $\bar{\rho}$, we construct from \mathbf{x}^* a probability density function φ^* , and seek to bound $\varphi^* - \bar{\rho}$ in a suitable topology. More generally, for any $\mathbf{x} \in \Omega$, we define a related density function $\varphi \in \mathcal{P}(\mathbb{R})$ given by the piecewise constant function

$$\varphi(y) := \begin{cases} \frac{1/n}{x_i - x_{i-1}} & \text{if } x_{i-1} < y < x_i \text{ for some } i \in \{1, \dots, n\} \\ 0 & \text{otherwise.} \end{cases} \quad (7)$$

We denote by φ^* the density function related to \mathbf{x}^* . The choice of φ is not unique; in Section 1.2 we discuss a different choice based on Voronoi cells. The current choice in (7) is made to ease computations, and has been used in earlier studies; see, e.g., [HCO10].

Figure 2 illustrates typical examples of φ^* and $\bar{\rho}$. Especially in the case where $D(U)$ confines φ^* , the graphs of φ^* and $\bar{\rho}$ are close to each other. This observation is in line with the literature (see, e.g., [GvMPS16, GPPS13, HCO10, HHvM18]). In this paper we wish to finally quantify this observation.




Figure 2: Numerical computations of \mathbf{x}^* for $n = 16$, $V = V_a$ with $a = \frac{1}{2}$ and three different choices of U . The points x_i^* on the horizontal axis are indicated by the vertical edges of each light-gray rectangle (all with area $\frac{1}{n}$). The graph of φ^* is given by the top edges of these rectangles. The black curve is the graph of $\bar{\rho}$. The region where $U = \infty$ is indicated in gray. The values of γ_i are chosen such that $\text{supp } \bar{\rho} = [0, 1]$. The computation of \mathbf{x}^* and $\bar{\rho}$ is explained in Section 7.

1.1 Main result

We estimate $\varphi^* - \bar{\rho}$ in terms of a fractional Sobolev norm. To introduce this norm, we define the fractional Sobolev space on \mathbb{R} by

$$\begin{aligned} H^{-s}(\mathbb{R}) &:= \left\{ \zeta \in \mathcal{S}'(\mathbb{R}) : \int_{\mathbb{R}} (1 + \omega^2)^{-s} |\widehat{\zeta}(\omega)|^2 d\omega < \infty \right\} \\ \|\zeta\|_{H^{-s}(\mathbb{R})}^2 &:= \int_{\mathbb{R}} (1 + \omega^2)^{-s} |\widehat{\zeta}(\omega)|^2 d\omega, \end{aligned} \tag{8}$$

where $s > 0$, $\widehat{\zeta}$ is the Fourier transform of ζ and $\mathcal{S}'(\mathbb{R})$ is the space of tempered distributions, i.e., the dual of the Schwartz space $\mathcal{S}(\mathbb{R})$. The reason for choosing this norm is that it is equivalent to an adjusted energy norm of E for $s = (1 - a)/2$, where we recall that a is the strength of the singularity of V ; see (3). In turn, this adjusted energy norm of $\varphi^* - \bar{\rho}$ can be estimated in terms of energy differences, which are easier to analyse than \mathbf{x}^* itself. The benefit of using the fractional Sobolev norm over the adjusted energy norm is that it is independent of the interaction potential V .

The main theorem of this paper is Theorem 1.2.

Theorem 1.2 (The quantitative estimate). *Let $n \geq 1$. Let E_n and E be as defined in (1) and (6) with potentials V and U satisfying Assumption 1.1 for some $0 \leq a < 1$. Then, the minimal values of E_n and E are attained, and the minimiser $\bar{\rho}$ of E is unique. Moreover,*

there exists $C > 0$ independent of n such that for any minimiser \mathbf{x}^* of E_n ,

$$\|\varphi^* - \bar{\rho}\|_{H^{-(1-a)/2}(\mathbb{R})}^2 \leq C \begin{cases} n^{-1+a} & 0 < a < 1 \\ n^{-1}(\log n)^3 & a = 0, \end{cases}$$

where φ^* is constructed from \mathbf{x}^* by (7).

The available tools for the proof of Theorem 1.2 are the monotonicity and convexity of V and the regularity properties of $\bar{\rho}$ proven in [KvM19] (see Lemma 3.4 below). The difficulty is that no information on \mathbf{x}^* is available, except that \mathbf{x}^* is a minimiser of E_n .

Next we give a sketch of the proof. For simplicity, we assume $a \in (0, 1)$. The proof is divided in 3 steps. Step 1 is a preparatory step. In this step, we show that it is not restrictive to assume that

$$\text{supp } V \text{ is compact,} \quad (9a)$$

$$\text{supp } \bar{\rho} = [0, 1]. \quad (9b)$$

In Step 2, we estimate $\|\varphi^* - \bar{\rho}\|_{H^{-(1-a)/2}(\mathbb{R})}^2$ without using any information on \mathbf{x}^* except that it is the minimiser of E_n . We start by following [KvM19] by rewriting E as the sum of the square of a norm and a linear term, i.e.,

$$E(\rho) = \frac{1}{2} \|\rho\|_V^2 + \int_{\mathbb{R}} U d\rho, \quad \|\rho\|_V^2 := \int_{\mathbb{R}} (V * \rho) d\rho. \quad (10)$$

The norm $\|\cdot\|_V$ is the adjusted energy norm which we mentioned below (8). The ‘adjustment’ refers to the replacement of the interaction potential in Step 1 by a potential V with compact support. We recall from [KvM19, Prop. 3.3] that $\|\cdot\|_V$ is equivalent to the norm on $H^{-(1-a)/2}(\mathbb{R})$. From (10) and the minimality of $\bar{\rho}$ (see Lemma 3.4.(iv)), we obtain

$$\begin{aligned} \|\varphi^* - \bar{\rho}\|_{H^{-(1-a)/2}(\mathbb{R})}^2 &\lesssim \|\varphi^* - \bar{\rho}\|_V^2 \\ &= 2(\bar{\rho}, \bar{\rho} - \varphi^*)_V + \|\varphi^*\|_V^2 - \|\bar{\rho}\|_V^2 \\ &= 2 \int (V * \bar{\rho} + U) d(\bar{\rho} - \varphi^*) + 2E(\varphi^*) - 2E(\bar{\rho}) \\ &\leq 0 + 2(E(\varphi^*) - E(\bar{\rho})). \end{aligned} \quad (11)$$

Then, proving Theorem 1.2 translates into bounding the energy difference in the right-hand side.

To bound this difference, we obtain from the minimality of \mathbf{x}^* that

$$E(\varphi^*) - E(\bar{\rho}) = \underbrace{E(\varphi^*) - E_n(\mathbf{x}^*)}_{T_1} + \underbrace{E_n(\mathbf{x}^*) - E_n(\bar{\mathbf{x}})}_{\leq 0} + \underbrace{E_n(\bar{\mathbf{x}}) - E(\bar{\rho})}_{T_2}, \quad (12)$$

where $\bar{\mathbf{x}} \in \Omega$ can be chosen freely. For an appropriate choice of $\bar{\mathbf{x}}$ (see (34)), we bound T_1 and T_2 from above in Section 4. This is easy for the confinement term of the energy. For the interaction term, we perform a direct computation in which we write $\|\cdot\|_V^2$ explicitly as the integral over the square $(\text{supp } \bar{\rho})^2 = [0, 1]^2$, which we subdivide into the rectangles $(x_{i-1}, x_i) \times (x_{j-1}, x_j)$; see Figure 3. Then, by the monotonicity and convexity of V , we ultimately obtain

$$T_1 \leq \frac{C}{n} + C' E_n^{\text{nn}}(\mathbf{x}^*) \quad \text{and} \quad T_2 \leq C n^{-1+a}, \quad (13)$$

where

$$E_n^{\text{nn}}(\mathbf{x}) := \frac{1}{n^2} \sum_{i=1}^n V(x_i - x_{i-1}) \quad (14)$$

is the part of the interaction term given by all nearest neighbour (superscript ‘nn’) interactions.

Putting the findings of Step 2 together, we obtain

$$\|\varphi^* - \bar{\rho}\|_{H^{-(1-a)/2}(\mathbb{R})}^2 \leq Cn^{-1+a} + C'E_n^{\text{nn}}(\mathbf{x}^*).$$

Hence, it is left to show that

$$E_n^{\text{nn}}(\mathbf{x}^*) \leq Cn^{-1+a}. \quad (15)$$

In Step 3 we prove (15). This is the difficult part of the proof of Theorem 1.2; we consider it as the main mathematical novelty of this paper. Our strategy is to establish the following lower bound on E_n :

$$E_n(\mathbf{x}) - E(\bar{\rho}) \geq E_n^{\text{nn}}(\mathbf{x}) - Cn^{-1+a} \quad \text{for all } \mathbf{x} \in \Omega; \quad (16)$$

see Proposition 5.1. Then, taking $\mathbf{x} = \mathbf{x}^*$, the right-hand side in (16) is bounded from above by T_2 . By the bound on T_2 in (13), we then conclude (15).

The argument in Step 3 is inspired by [PS17, Sec. 2]; we also construct a renormalisation of the norm $\|\cdot\|_V$, but we need to construct a different one to allow for $V_{\text{reg}} \neq 0$ and unbounded $\bar{\rho}$.

We conclude the sketch of the proof by remarking that the case $a = 0$ can be treated analogously; the only difference is that the factor $\log n$ appears at a few places in the estimates.

1.2 Remarks on Theorem 1.2

Here we list several remarks on the statement and the sketch of the proof of Theorem 1.2:

Uniform bound on support of φ^* The proof of Step 1 contains the additional result that $\text{supp } \varphi^*$ is bounded uniformly in n ; see Proposition 3.2. While the proof consists of common arguments in potential theory, we believe that the statement of Proposition 3.2 has merit on its own due to the rather weak assumptions on V and U .

Extension of [KvM19] The statement of Step 1 has further merit; the main results in [KvM19] (on the regularity of $\bar{\rho}$) are stated under the stronger assumption that the tails of V are integrable, in which case $V \in L^1(\mathbb{R})$. Here, in Step 1 we extend these results to the larger class of potentials V specified by Assumption 1.1.

Sharpness of the exponent To test the degree of sharpness of the exponent of n in the estimate in Theorem 1.2, we perform in Section 7 numerical computations to compute $\|\varphi^* - \bar{\rho}\|_{H^{-(1-a)/2}(\mathbb{R})}^2$ for various values of n . Surprisingly, our findings in Section 7 show that in all test cases the numerical values of $\|\varphi^* - \bar{\rho}\|_{H^{-(1-a)/2}(\mathbb{R})}^2$ decrease roughly as a power law n^{-p} with p significantly *larger* than the exponent $1 - a$ in Theorem 1.2. Hence, the exponent $1 - a$ in Theorem 1.2 appears not to be sharp. We comment in Section 1.3 on the loss of sharpness in the proof of Theorem 1.2 in the special case of $V_{\text{reg}} \equiv 0$ (Riesz gases), for which more precise estimates than that in Theorem 1.2 are available.

Choice of distance/norm As mentioned before, our choice of norm circumvents any analysis of fine properties of \mathbf{x}^* , and requires instead in the proof of Theorem 1.2 the lower bound on the energy E_n in (16). Other choices, such as L^p -norms or the Wasserstein distance, appear to require certain properties of \mathbf{x}^* , which makes it more difficult to establish a quantitative estimate in these norms.

Other than mathematical convenience, our choice of norm has a further merit; it provides a quantitative estimate for the particle interaction force on \mathbb{R} induced by the particle densities φ^* and $\bar{\rho}$. These interaction forces are

$$-(V * \varphi^*)' - U' \quad \text{and} \quad -(V * \bar{\rho})' - U'$$

respectively. To obtain a quantitative estimate from Theorem 1.2, we first change V to have property (9a)¹. Then, by [KvM19, Lem. 3.1(iii)] it holds that

$$\exists C > 0 \, \forall \omega \in \mathbb{R} : \widehat{V}(\omega) \leq C(1 + \omega^2)^{-\frac{1-a}{2}}.$$

Hence, writing $\nu := \varphi^* - \bar{\rho}$ and $s := \frac{1-a}{2}$,

$$\|\nu\|_{H^{-s}(\mathbb{R})}^2 = \int_{\mathbb{R}} (1 + \omega^2)^{-s} |\widehat{\nu}(\omega)|^2 d\omega \geq \frac{1}{C^2} \int_{\mathbb{R}} (1 + \omega^2)^s |\widehat{V}(\omega) \widehat{\nu}(\omega)|^2 d\omega = \frac{1}{C^2} \|V * \nu\|_{H^s(\mathbb{R})}^2.$$

Thus,

$$\|(V * \varphi^*)' - (V * \bar{\rho})'\|_{H^{-(1+a)/2}(\mathbb{R})}^2 \leq C' \|V * (\varphi^* - \bar{\rho})\|_{H^{(1-a)/2}(\mathbb{R})}^2 \leq C' \|\varphi^* - \bar{\rho}\|_{H^{-(1-a)/2}(\mathbb{R})}^2,$$

for which Theorem 1.2 gives an upper bound.

Other choice of φ Another commonly used choice for φ than that in (7) is to construct it from a one-dimensional Voronoi tessellation of the points x_i , and to assign to each Voronoi cell a mass of $1/n$. While Voronoi cells easily extend to higher dimensions, even in one dimension they introduce two complications for proving the corresponding estimates in (13). First, an additional choice for the Voronoi cells for x_0 and x_n has to be made. Second, the mass of $\bar{\rho}$ on the Voronoi cells constructed from $\bar{\mathbf{x}}$ may not equal $1/n$, which induces further error terms.

1.3 Position in the literature

Here we put Theorem 1.2 in the context of the literature. In particular, we show how it applies to problems in plasticity and in numerical integration, and how it compares to recent advances on Riesz gases.

Plasticity The paper series started by [GPPS13, Hal11] and continued in [GvMPS16, HHvM18, vM18a, vM18b, vMMP14] studies the connection between models for plasticity of metals and an underlying microscopic model in a one-dimensional setting. This microscopic model is a minimisation problem of a certain E_n of the form (1). In particular, the interaction potential is

$$V(x) = x \coth x - \log(2|\sinh x|),$$

¹Changing V changes the interaction forces, but only on a domain which is a certain distance away from $\text{supp } \bar{\rho}$ and $\text{supp } \varphi^*$.

which fits to Assumption 1.1 with $a = 0$ and $V_{\text{reg}} \not\equiv 0$. While in this paper series the convergence of φ^* to $\bar{\rho}$ as $n \rightarrow \infty$ is established, no quantitative estimates between φ^* and $\bar{\rho}$ were found², which limits the application to plasticity.

The main result of this paper, Theorem 1.2, provides the first quantitative estimate for this microscopic model. The estimate is given by

$$\|\varphi^* - \bar{\rho}\|_{H^{-1/2}}^2 \leq C(U) \frac{(\log n)^3}{n}. \quad (17)$$

For the application, a more detailed dependence of the constant C on U is needed. Since the proof of Theorem 1.2 is constructive, it may be possible to use its steps for constructing an explicit expression for $C(U)$.

Our aim to establish quantitative estimates fits within a recent trend in mathematical research on plasticity. In this trend (see, e.g., [HvMP20]), the starting point is a microscopic particle system which depends on several parameters such as n , the temperature and the lattice spacing between the atoms of the metal. The goal is to identify regions in the space of parameters on which the microscopic system can be approximated by a macroscopic system described in terms of a particle density ρ . To quantify this approximation, quantitative estimates such as that in Theorem 1.2 are required. The main contribution from Theorem 1.2 is that it requires no regularisation of the singularity of V at 0.

Approximation of functions A common problem in the field of approximating a given function is how to choose finitely many sampling points at which to evaluate the function. For analytic functions defined on subsets of the complex plane, it is shown in [TS19, HT19] that the optimal choice of sampling points can be obtained from the minimiser \mathbf{x}^* of E_n for certain potentials V and U which satisfy Assumption 1.1. In particular, the interaction potential is explicitly given by $V(x) = -\log |\tanh x|$, which satisfies Assumption 1.1 with $a = 0$ and $V_{\text{reg}} \not\equiv 0$. Moreover, to bound the error made when replacing the given function by the approximation from the sampling points, it is required to find upper and lower bounds on

$$E_n(\mathbf{x}^*) - E(\bar{\rho}). \quad (18)$$

The currently available bounds on (18) (see [HT19, Thm. 2.3]) are comparable in size to $E_n(\mathbf{x}^*)$ itself. Applying (16) and the estimate on T_2 in (13) yields

$$|E_n(\mathbf{x}^*) - E(\bar{\rho})| \leq C(U) \frac{(\log n)^3}{n},$$

which demonstrates that it may be possible to construct a sharper estimate. In this setting, U depends on n , and thus (similar to the application to plasticity) a more detailed estimate on $C(U)$ is required.

Riesz gases For several other applications in approximation theory, detailed properties of the minimiser \mathbf{x}^* of E_n are desired in the case where $V_{\text{reg}} \equiv 0$ and where the particle positions may be of any dimension. Establishing such properties is the main topic of the paper series by Petrache, Sandier, Serfaty *et al.* ([SS15b, SS15a, PS17] to list a few). In particular, it is found in [PS17, Thm. 4] that

$$E_n(\mathbf{x}^*) - E(\bar{\rho}) = n^{-1+a}(-M + o(1)) \quad \text{as } n \rightarrow \infty \quad (19)$$

²An exception is [vM18a], which establishes a quantitative estimate for a special, n -dependent choice of U .

under appropriate restrictions on the potential U . The constant $M > 0$ is of particular importance. It is explicitly expressed in terms of a maximisation problem on the microscopic configuration of \mathbf{x}^* in the limit $n \rightarrow \infty$.

For the comparison between (19) and Theorem 1.2, it suffices to consider M as a given constant. Even when (19) is restricted to one dimension, it is a more precise result than our estimates in (13) and (16). The reason for obtaining such a precise result is that for $V = V_a$ the extension representation of [CSS08] can be used (see [PS17] for details). For our larger class of potential V , we are not aware of a similar extension representation. Moreover, in this paper, $\bar{\rho}$ need not be bounded, which complicates the estimates (see, e.g., Remark 5.4 below).

Next, we put together the conclusion from (19) that $E_n(\mathbf{x}^*) - E(\bar{\rho})$ decays as n^{-1+a} and the finding from the numerical computations in Section 7 that $\|\varphi^* - \bar{\rho}\|_V^2$ decays faster, namely as n^{-p} with $p \geq -1 + a$. First, we note that our numerical solutions for \mathbf{x}^* satisfy $x_i^* \in \text{supp } \bar{\rho}$ for all $i = 1, \dots, n$ for each tested value of n . This implies that $\text{supp } \varphi^* \subset \text{supp } \bar{\rho}$, and thus by Lemma 3.4(iv) the second inequality in (11) turns into an equality. Substituting this equality in (12), we obtain

$$\underbrace{\|\varphi^* - \bar{\rho}\|_V^2}_{O(n^{-p})} = \underbrace{E(\varphi^*) - E_n(\mathbf{x}^*)}_{T_1} + \underbrace{E_n(\mathbf{x}^*) - E(\bar{\rho})}_{n^{-1+a}(-M+o(1))}. \quad (20)$$

From (20) it is clear that the loss of sharpness in the exponent of n in the estimate in Theorem 1.2 originates from the choice to estimate the energy differences in the right-hand side of (12) without trying to optimise the multiplicative constant in front of n^{-1+a} . Finding this optimal multiplicative constant could result in a sharper estimate than that in Theorem 1.2. However, it seems quite difficult to optimise this constant. At least in the case of Riesz gases in [PS17] where (19) holds, a sharper estimate than that in Theorem 1.2 follows from the right-hand side of (20) if one can show that

$$T_1 = n^{-1+a}(M + o(1)).$$

However, our current estimate $T_1 \leq Cn^{-1+a}$ obtained from the strategy outlined in (13) and (16) is the core of our proof, and we have as of yet no clue on how the corresponding constant C can be characterised as the constant M .

Organisation of the paper In Section 2 we list our notation. In Sections 3, 4 and 5 we establish the required tools to justify the steps in the sketch of the proof of Theorem 1.2. In Section 6 we put these tools together to prove Theorem 1.2. In Section 7 we describe and discuss our numerical findings for the actual dependence of the left-hand side in Theorem 1.2 on n and a .

2 Notation

The following table list the symbols which we use throughout the paper.

\wedge, \vee	$\alpha \wedge \beta := \min\{\alpha, \beta\}$ and $\alpha \vee \beta := \max\{\alpha, \beta\}$	
$(\cdot, \cdot)_V$	inner product constructed from V ; $(f, g)_V = \int (V * g)f$	(31), (32)
$\mathbb{1}_A(x)$	indicator function; $\mathbb{1}_A(x) = 1$ if $x \in A$, and $\mathbb{1}_A(x) = 0$ otherwise	
a	strength of the singularity of V ; $0 \leq a < 1$	(3)

C, C', \dots	some n -independent constants	
E	energy for the particle density	(6)
E_n	interacting particle energy; $E_n : \Omega \rightarrow \mathbb{R}$	(1)
E_n^{nn}	nearest neighbour interactions; part of E_n	(14)
φ	discrete density (piece-wise constant) constructed from $\mathbf{x} \in \Omega$	(7)
Γ	Γ -function; $\Gamma(\alpha) := \int_0^\infty x^{\alpha-1} e^{-x} dx$	
$H^{-s}(\mathbb{R})$	fractional Sobolev space for $s > 0$	(8)
ℓ_i	distance between nearest neighbours in $\mathbf{x} \in \Omega$; $\ell_i := x_i - x_{i-1}$	(35)
m_i	midpoints of nearest neighbours in $\mathbf{x} \in \Omega$; $m_i := \frac{1}{2}(x_i + x_{i-1})$	(35)
n	$n + 1$ is the number of particles; $n \geq 1$	
Ω	space of admissible particle configurations; $\Omega \subset \mathbb{R}^{n+1}$	(2)
$\mathcal{P}(\mathbb{R})$	space of probability measures on \mathbb{R}	
$\bar{\rho}$	the minimiser of E	Lem. 3.4
U	confining potential	Ass. 1.1
V	interaction potential	Ass. 1.1
V_a	singular, homogeneous part of V	(3)
V_{reg}	regular part of V ; $V_{\text{reg}} = V - V_a$	
\mathbf{x}^*	a minimiser of E_n ; $\mathbf{x}^* \in \Omega$	
$\bar{\mathbf{x}}$	particle configuration constructed from $\bar{\rho}$; $\bar{\mathbf{x}} \in \Omega$	(34)

We use the convention that constants denoted by C are independent of n and may change from line to line. In several cases where the estimates are easier to follow when the change in constants is highlighted, we use C', C'', \dots instead.

3 Changing the tails of V

In this section we prove the key Propositions 3.1 and 3.2 which will allow us to justify rigorously Step 1 of the sketch of the proof of Theorem 1.2, i.e., that V may be replaced by a different interaction potential which satisfies (9a). These propositions state that the set of minimisers of E_n and the set of minimisers of E do not depend on the tails of V . Since we consider these propositions of independent interest, we pose them under weaker conditions of V than Assumption 1.1. In addition to proving these propositions, we show that E_n and E attain their minimal values, that the minimiser $\bar{\rho}$ of E is unique, and that $\bar{\rho}$ satisfies several properties.

Minimisers are independent of the tails of V Let $V \in L_{\text{loc}}^1(\mathbb{R})$ satisfy (4) and U satisfy Assumption 1.1. By an affine change of variables, we may assume that $[-1, 1] \subset D(U)$. By (5), there exists a constant $M > 0$ such that

$$U(x) \geq \frac{|x|}{M} - M \quad \text{for all } x \in \mathbb{R}. \quad (21)$$

By (4), we note that, on $(0, \infty)$, V is non-increasing and V' is non-decreasing. Furthermore, $V'(x) \rightarrow 0$ as $x \rightarrow \infty$. Hence, there exists a point of differentiability $R \geq 2$ of V for which

$$V'(R) \geq -\frac{1}{4M} \quad \text{and} \quad \frac{V(R)}{R} \geq -\frac{1}{4M}. \quad (22)$$

Proposition 3.1. *Let $V \in L^1_{loc}(\mathbb{R})$ satisfy (4) and U satisfy Assumption 1.1. Take $M, R > 0$ as in (21) and (22). Then, there exists a constant $S > 0$ independent of $V|_{(R,\infty)}$ such that any minimiser $\bar{\rho}$ of E satisfies $\text{supp } \bar{\rho} \subset [-S, S]$.*

Proof. We start by proving two auxiliary estimates. The first one is given by

$$\inf_{x>0} \left(V(x) + \frac{x}{4M} \right) \geq \min_{0<x\leq R} \left(V(x) + \frac{x}{4M} \right) \wedge 0 =: -N. \quad (23)$$

To prove it, let $x > R$. Since V is convex, the tangent line of V at R is below the graph of V . Then, by (22), we obtain

$$V(x) + \frac{x}{4M} \geq V(R) + (x - R)V'(R) + \frac{x}{4M} \geq -\frac{R}{4M} - \frac{x - R}{4M} + \frac{x}{4M} = 0.$$

This proves (23). We note that the constant $N \geq 0$ does not depend on $V|_{(R,\infty)}$. We set

$$S := 2M(2E(\rho_0) + 2 + 2M + N),$$

where $\rho_0 := \frac{1}{2}\mathbb{1}_{[-1,1]}$ is chosen rather arbitrarily to obtain that the value of $E(\rho_0)$ is finite and independent of $V|_{(R,\infty)}$.

The second auxiliary estimate is given by

$$\frac{V(x - y) + U(x) + U(y)}{2} \geq E(\rho_0) + 1 \quad \text{for all } x, y \text{ such that } |x| \vee |y| \geq S. \quad (24)$$

To prove it, we first note from (21) that

$$\frac{U(x) + U(y)}{2} \geq \frac{|x| + |y|}{2M} - M \geq \frac{S}{2M} - M = 2E(\rho_0) + 2 + M + N.$$

Then, using (23),

$$V(x - y) + \frac{U(x) + U(y)}{2} \geq V(x - y) + \frac{|x - y|}{2M} - M \geq -N - M. \quad (25)$$

Adding the above two estimates, we obtain (24).

Next, we prove Proposition 3.1. Suppose that $\bar{\rho}$ is a minimiser of E such that $\text{supp } \bar{\rho} \not\subset [-S, S]$. Then, $\bar{m} := \bar{\rho}([-S, S]) < 1$. First, we claim that $\bar{m} > 0$. Indeed, if not, then by (24)

$$E(\bar{\rho}) = \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{V(x - y) + U(x) + U(y)}{2} d\bar{\rho}(y) d\bar{\rho}(x) \geq E(\rho_0) + 1,$$

which contradicts with the minimality of $\bar{\rho}$. Using that $\bar{m} > 0$, we set $\rho := \bar{\rho}|_{[-S, S]}/\bar{m} \in \mathcal{P}(\mathbb{R})$, and rely on (24) to estimate

$$\begin{aligned} E(\bar{\rho}) &= \iint_{\mathbb{R}^2 \setminus [-S, S]^2} \frac{V(x - y) + U(x) + U(y)}{2} d\bar{\rho}(y) d\bar{\rho}(x) \\ &\quad + \iint_{[-S, S]^2} \frac{V(x - y) + U(x) + U(y)}{2} \bar{m}^2 d\rho(y) d\rho(x) \\ &\geq (E(\rho_0) + 1)(1 - \bar{m}^2) + \bar{m}^2 E(\rho) \geq (E(\bar{\rho}) + 1)(1 - \bar{m}^2) + \bar{m}^2 E(\rho). \end{aligned}$$

Rearranging terms,

$$E(\bar{\rho}) \geq E(\rho) + \frac{1 - \bar{m}^2}{\bar{m}^2} > E(\rho),$$

which contradicts with the minimality of $\bar{\rho}$. \square

Proposition 3.2. *Let V, U, M, R be as in Proposition 3.1, and let $n \geq 1$. Then, there exists a constant $S > 0$ independent of n and of $V|_{(R, \infty)}$ such that any minimiser $\mathbf{x}^* \in \Omega$ of E_n satisfies $x_i^* \in [-S, S]$ for all $i = 0, \dots, n$.*

Proof. The proof is a discrete version of the proof of Proposition 3.1. We rely again on (23) with the same constant $N \geq 0$. Then, we set

$$S := \left(2N + 3M + \int_{-1}^1 (V + U)(x) dx\right) M. \quad (26)$$

Since \mathbf{x}^* is ordered, it is enough to show that $-S \leq x_0^*$ and $x_n^* \leq S$. Suppose that $x_0^* < -S$ or $x_n^* > S$. We first treat the case in which both $x_0^* < -S$ and $x_n^* > S$ hold, and comment on the remaining case afterwards. We may assume that $U(x_0^*) \leq U(x_n^*)$, because otherwise we can obtain this by applying the variable transformation $x \mapsto -x$.

We will reach a contradiction with the minimality of \mathbf{x}^* by finding a lower energy state in Ω . We construct this state by replacing x_n^* by a more energetically favourable position $y_n \in [-1, 1]$. With this aim, we first compute for any $y \in [-1, 1]$

$$n(E_n(\mathbf{x}^*) - E_n(x_0^*, \dots, x_{n-1}^*, y)) = \underbrace{\frac{1}{n} \sum_{j=0}^{n-1} V(x_n^* - x_j^*) + U(x_n^*)}_{W_n(x_n^*)} - \underbrace{\left(\frac{1}{n} \sum_{j=0}^{n-1} V(y - x_j^*) + U(y) \right)}_{W_n(y)}, \quad (27)$$

where, for ease of notation, we have dropped the convention to have the particles positions ordered in the argument of E_n .

Since W_n is lower semi-continuous and bounded from below on compact sets, it attains its minimum on $[-1, 1]$. We take y_n as a minimiser of W_n over $[-1, 1]$. Then, we estimate

$$2W_n(y_n) \leq \int_{-1}^1 W_n(y) dy = \frac{1}{n} \sum_{j=0}^{n-1} \int_{-1}^1 V(y - x_j^*) dy + \int_{-1}^1 U(y) dy \leq \int_{-1}^1 (V + U)(y) dy.$$

Next we estimate $W_n(x_n^*)$ from below. Using that V is non-increasing on $(0, \infty)$, we obtain

$$W_n(x_n^*) \geq V(x_n^* - x_0^*) + \frac{U(x_n^*) + U(x_0^*)}{4} + \frac{U(x_n^*) - U(x_0^*)}{4} + \frac{1}{2}U(x_n^*). \quad (28)$$

Then, following the estimates in (25) for the first two terms, and applying (21) to the fourth term, we obtain

$$W_n(x_n^*) \geq -(N + M) - 0 + \frac{1}{2} \left(\frac{x_n^*}{M} - M \right) > \frac{S}{2M} - N - \frac{3}{2}M. \quad (29)$$

Collecting our results and substituting them in (27) yields

$$n(E_n(\mathbf{x}^*) - E_n(x_0^*, \dots, x_{n-1}^*, y_n)) > \frac{S}{2M} - N - \frac{3}{2}M - \frac{1}{2} \int_{-1}^1 (V + U)(y) dy,$$

which is non-negative by the choice of S in (26). This contradicts with the minimality of \mathbf{x}^* .

Finally, we treat the case in which either $x_0^* < -S$ or $x_n^* > S$, but not both. Again, by changing variables if needed, we may assume that $x_n^* > S$, and thus $-x_0^* \leq S < x_n^*$. Then,

the same proof can be adopted with a minor modification. This modification is to replace (28) with the following:

$$W_n(x_n^*) \geq V(2x_n^*) + \frac{U(x_n^*) + U(x_n^*)}{4} + \frac{1}{2}U(x_n^*).$$

This results again in (29). \square

Remark 3.3 (E_n attains its minimum). *The existence of minimisers for E_n is included in the proofs of Propositions 3.1 and 3.2. Indeed, since E_n is continuous on Ω and $E_n(\mathbf{x}) \rightarrow \infty$ as $\text{dist}(\mathbf{x}, \partial\Omega) \rightarrow 0$, it remains to be shown that $E_n(\mathbf{x}) \rightarrow \infty$ as $|\mathbf{x}| \rightarrow \infty$. To show this, we write*

$$E_n(\mathbf{x}) = \frac{1}{2n^2} \sum_{i=0}^n \sum_{j \neq i}^n \left(V(x_i - x_j) + \frac{U(x_i) + U(x_j)}{2} \right) + \frac{1}{2n} \sum_{i=0}^n U(x_i)$$

and obtain from (21) and (25) that

$$E_n(\mathbf{x}) \geq -C + \frac{1}{2Mn} \sum_{i=0}^n |x_i| \xrightarrow{|\mathbf{x}| \rightarrow \infty} \infty.$$

Properties of the compactly supported V and $\bar{\rho}$ Let V satisfy Assumption 1.1 and the compact support condition (9a). By Assumption 1.1, there exist constants $b, c > 0$ such that

$$V''(r) \geq cr^{-(2+a)} \quad \text{for all } 0 < r \leq b. \quad (30)$$

Since $\text{supp } V$ is bounded, it is obvious from the convexity of V on $\mathbb{R} \setminus \{0\}$ that $V \geq 0$. Moreover, for any $f, g \in L^2(\mathbb{R})$,

$$\int_{\mathbb{R}} (V * f)g =: (f, g)_V \quad (31)$$

defines an inner product, which induces the Hilbert space

$$\overline{L^2(\mathbb{R})}^{\|\cdot\|_V} \cong H^{-(1-a)/2}(\mathbb{R}). \quad (32)$$

The proof of this is given in [KvM19, Lem. 3.1(iii) and Prop. 3.3], and uses that V satisfies Assumption 1.1 and $V \in L^1(\mathbb{R})$. The following lemma is a simplified version of [KvM19, Thms. 1.4 and 1.5].

Lemma 3.4 (Properties of $\bar{\rho}$). *Let $0 \leq a < 1$. E attains its minimal value on $\mathcal{P}(\mathbb{R})$. Its minimiser is unique, and has a density $\bar{\rho} \in L^1(\mathbb{R})$. Moreover, after applying an appropriate affine change of variables, $\bar{\rho}$ satisfies*

$$(i) \quad \text{supp } \bar{\rho} = [0, 1];$$

$$(ii) \quad \bar{\rho} \in L^1(0, 1) \cap C((0, 1));$$

$$(iii) \quad \exists C > 0 \quad \forall 0 < x < 1 : \bar{\rho}(x) \leq C[x(1-x)]^{-\frac{1-a}{2}};$$

$$(iv) \quad \left\{ \begin{array}{ll} V * \bar{\rho} + U - \bar{C} = 0 & \text{on } [0, 1] \\ V * \bar{\rho} + U - \bar{C} \geq 0 & \text{on } \mathbb{R}. \end{array} \right\}, \text{ where } \bar{C} := \int_{\mathbb{R}} (V * \bar{\rho}) d\bar{\rho} + \int_{\mathbb{R}} U d\bar{\rho} > 0.$$

The purpose of changing variables in Lemma 3.4 is to have $\text{supp } \bar{\rho} = [0, 1]$ instead of some other bounded, closed interval. It is easy to see that Assumption 1.1 and (9a) are invariant under an affine change of variables.

4 Estimates on the terms T_1 and T_2

In this section we fill in the details of Step 2 of the sketch of the proof of Theorem 1.2. We rely on the preparations in Step 1 to assume (9), i.e., that the supports of V and $\bar{\rho}$ are compact.

Most of the estimates in the sketch are already rigorously justified. The only estimates left to prove are the two inequalities in (13), which we recall to be

$$\underbrace{E(\varphi^*) - E_n(\mathbf{x}^*)}_{T_1} \leq \frac{C}{n} + C' E_n^{\text{nn}}(\mathbf{x}^*) \quad \text{and} \quad \underbrace{E_n(\bar{\mathbf{x}}) - E(\bar{\rho})}_{T_2} \leq C n^{-1+a}. \quad (33)$$

We note that $\bar{\mathbf{x}} \in \Omega$ is yet to be constructed, $C, C' > 0$ are independent of n , and E_n^{nn} is defined in (14). Recalling that $\text{supp } \bar{\rho} = [0, 1]$, we define $\bar{\mathbf{x}}$ by

$$\bar{x}_0 = 0, \quad \bar{x}_n = 1, \quad \text{and} \quad \int_{\bar{x}_{i-1}}^{\bar{x}_i} \bar{\rho}(x) dx = \frac{1}{n} \quad \text{for all } i = 1, \dots, n. \quad (34)$$

The structure of the proof of (33) is as follows. First, we split

$$T_2 = \underbrace{E_n(\bar{\mathbf{x}}) - E(\bar{\varphi})}_{T_3} + \underbrace{E(\bar{\varphi}) - E(\bar{\rho})}_{T_4},$$

where $\bar{\varphi}$ is defined from $\bar{\mathbf{x}}$ by (7). We prove that $T_4 \lesssim n^{-1+a}$ in Lemmas 4.2 and 4.3. For T_1 and T_3 , we note that – except for the sign – they are both of the form

$$E(\varphi) - E_n(\mathbf{x}),$$

where \mathbf{x} and φ are related through (7). In Lemma 4.5 we show that

$$-\frac{C}{n} - E_n^{\text{nn}}(\mathbf{x}) \leq E(\varphi) - E_n(\mathbf{x}) \leq \frac{C'}{n} + C'' E_n^{\text{nn}}(\mathbf{x}),$$

which yields

$$T_1 \leq \frac{C}{n} + C' E_n^{\text{nn}}(\mathbf{x}^*) \quad \text{and} \quad T_3 \leq \frac{C}{n} + E_n^{\text{nn}}(\bar{\mathbf{x}}).$$

Finally, in Lemma 4.1(iii) we show that $E_n^{\text{nn}}(\bar{\mathbf{x}}) \leq C n^{-1+a}$, which completes the proof of (33).

Properties of $\bar{\mathbf{x}}$ and the bound on $E_n^{\text{nn}}(\bar{\mathbf{x}})$ We start by introducing some notation. First, for given $\mathbf{x} \in \Omega$, we define

$$m_i := (x_i + x_{i-1})/2, \quad \text{and} \quad \ell_i := x_i - x_{i-1} \quad (35)$$

where $m := (m_1, \dots, m_n) \in \mathbb{R}^n$ lists the midpoints of neighbouring particles, and $\ell := (\ell_1, \dots, \ell_n) \in \mathbb{R}^n$ the distances between them. These quantities are illustrated in Figure 3. For the specific choices $\bar{\mathbf{x}}$ and \mathbf{x}^* , we denote the related midpoints and interparticle distances as \bar{m}_i , $\bar{\ell}_i$ and m_i^* , ℓ_i^* respectively.

Second, we introduce

$$D_n : \Omega \rightarrow [0, \infty), \quad D_n(\mathbf{x}) := \frac{1}{2} \frac{1}{n^2} \sum_{i=1}^n \frac{1}{\ell_i^2} \iint_{(0, \ell_i)^2} V(x - y) dy dx, \quad (36)$$

where ℓ_i depends on \mathbf{x} through (35). We note that

$$E_n^{\text{nn}}(\mathbf{x}) \leq 2D_n(\mathbf{x}). \quad (37)$$




Figure 3: The integration domain.

Lemma 4.1 (Properties of $\bar{\mathbf{x}}$). *There exist constants $C, c > 0$ such that for all $n \geq 1$*

$$(i) \quad \bar{x}_i \geq c \left(\frac{i}{n} \right)^{\frac{2}{1+a}} \quad \text{and} \quad \bar{x}_{n-i} \leq 1 - c \left(\frac{i}{n} \right)^{\frac{2}{1+a}} \quad \text{for all } i = 0, \dots, \left\lfloor \frac{n}{2} \right\rfloor;$$

$$(ii) \quad \bar{\ell}_1 \wedge \bar{\ell}_n \geq c \left(\frac{1}{n} \right)^{\frac{2}{1+a}} \quad \text{and} \quad \bar{\ell}_i \wedge \bar{\ell}_{n+1-i} \geq \frac{c}{n} \left(\frac{i-1}{n} \right)^{\frac{1-a}{1+a}} \quad \text{for all } i = 2, \dots, \left\lfloor \frac{n}{2} \right\rfloor;$$

$$(iii) \quad D_n(\bar{\mathbf{x}}) + E_n^{\text{nn}}(\bar{\mathbf{x}}) \leq Cn^{-1+a}.$$

Proof. For convenience we assume that n is even. Since $\bar{x}_0 = 0$, it is sufficient to consider any $i \geq 1$. Using Lemma 3.4(iii), we find that

$$\frac{i}{n} = \int_0^{\bar{x}_i} \bar{\rho} \leq \int_0^{\bar{x}_i} Cx^{-\frac{1-a}{2}} dx = C' \bar{x}_i^{\frac{1+a}{2}}.$$

This implies the first part of Property (i). The estimate for \bar{x}_{n-i} is found analogously.

Next we bound $\bar{\ell}_i$ from below. For $i = 1$, we find

$$\bar{\ell}_1 = \bar{x}_1 \geq c \left(\frac{1}{n} \right)^{\frac{2}{1+a}}.$$

For $i \geq 2$, we estimate similarly as above

$$\frac{1}{n} = \int_{\bar{x}_{i-1}}^{\bar{x}_i} \bar{\rho} \leq \int_{\bar{x}_{i-1}}^{\bar{x}_i} Cx^{-\frac{1-a}{2}} dx = C' \left(\bar{x}_i^{\frac{1+a}{2}} - \bar{x}_{i-1}^{\frac{1+a}{2}} \right).$$

Inserting $\bar{x}_i = \bar{\ell}_i + \bar{x}_{i-1}$, we obtain

$$\bar{\ell}_i \geq \left(\frac{1}{C'n} + \bar{x}_{i-1}^{\frac{1+a}{2}} \right)^{\frac{2}{1+a}} - \bar{x}_{i-1}. \quad (38)$$

Since $\frac{2}{1+a} > 1$, the function $\psi(t) := t^{2/(1+a)}$ is convex for $t > 0$, and thus $\psi(t+\varepsilon) \geq \psi(t) + \varepsilon\psi'(t)$ for all $t, \varepsilon > 0$. Applying this inequality to (38), and then using Property (i), we obtain

$$\bar{\ell}_i \geq \frac{1}{C'n} \frac{2}{1+a} \left(\bar{x}_{i-1}^{\frac{1+a}{2}} \right)^{\frac{2}{1+a}-1} = \frac{c}{n} \bar{x}_{i-1}^{\frac{1-a}{2}} \geq \frac{c'}{n} \left(\frac{i-1}{n} \right)^{\frac{1-a}{1+a}}.$$

The estimate for $\bar{\ell}_{n+1-i}$ is found analogously.

Finally we prove Property (iii). By (37) it is enough to estimate $D_n(\bar{\mathbf{x}})$. From $V(x) \leq C/|x|^a$ and Property (ii) we obtain

$$\begin{aligned} D_n(\bar{\mathbf{x}}) &= \frac{1}{2} \frac{1}{n^2} \sum_{i=1}^n \iint_{(0,1)^2} V(\bar{\ell}_i(x-y)) dy dx \leq \frac{C}{n^2} \left(\bar{\ell}_1^{-a} + \bar{\ell}_n^{-a} + \sum_{i=2}^{n-1} \bar{\ell}_i^{-a} \right) \\ &\leq C n^{-2+\frac{2a}{1+a}} + C \frac{n^a}{n} \frac{1}{n} \sum_{i=2}^{n/2} \left(\frac{i-1}{n} \right)^{-a \frac{1-a}{1+a}} \leq C n^{-1+a}. \end{aligned}$$

□

The bound on $T_4 = E(\bar{\varphi}) - E(\bar{\rho})$ We recall from (10) that E consists of an interaction part and a confinement part. For $E(\bar{\varphi}) - E(\bar{\rho})$, we bound these terms separately in Lemmas 4.3 and 4.2 respectively.

Lemma 4.2. *For all $n \geq 1$*

$$\int_0^1 U(x) (\bar{\varphi} - \bar{\rho})(x) dx \leq \frac{U(0) + U(1)}{n}.$$

Proof. From (7) and (34) we observe that the densities $\bar{\varphi}$ and $\bar{\rho}$ have mass $1/n$ on $[\bar{x}_{i-1}, \bar{x}_i]$ for each i . We use this to estimate

$$\int_0^1 U(\bar{\varphi} - \bar{\rho}) = \sum_{i=1}^n \int_{\bar{x}_{i-1}}^{\bar{x}_i} U(\bar{\varphi} - \bar{\rho}) \leq \frac{1}{n} \sum_{i=1}^n \left(\max_{[\bar{x}_{i-1}, \bar{x}_i]} U - \min_{[\bar{x}_{i-1}, \bar{x}_i]} U \right). \quad (39)$$

Assume for convenience that \bar{x}_I is a minimiser of U for some $I \in \{1, \dots, n\}$. Then, since U is convex, it is non-increasing on $[0, \bar{x}_I]$, and thus

$$\frac{1}{n} \sum_{i=1}^I \left(\max_{[\bar{x}_{i-1}, \bar{x}_i]} U - \min_{[\bar{x}_{i-1}, \bar{x}_i]} U \right) = \frac{1}{n} \sum_{i=1}^I (U(\bar{x}_{i-1}) - U(\bar{x}_i)) = \frac{1}{n} (U(\bar{x}_0) - U(\bar{x}_I)) = \frac{U(0)}{n}.$$

A similar argument on $[\bar{x}_I, 1]$ yields that the remaining terms of the sum in the right-hand side of (39) equal $U(1)/n$, and the statement of Lemma 4.2 follows. In the case where the interval of minimisers of U is contained in $(\bar{x}_{I-1}, \bar{x}_I)$ for some I , a similar argument applies. □

Lemma 4.3. *There exists a constant $C > 0$ such that for all $n \geq 1$*

$$\|\bar{\varphi}\|_V^2 - \|\bar{\rho}\|_V^2 \leq C n^{-1+a}.$$

Proof. We write

$$\|\bar{\varphi}\|_V^2 - \|\bar{\rho}\|_V^2 = \sum_{i=1}^n \sum_{j=1}^n \int_{\bar{x}_{i-1}}^{\bar{x}_i} \int_{\bar{x}_{j-1}}^{\bar{x}_j} V(x-y) (\bar{\varphi}(x)\bar{\varphi}(y) - \bar{\rho}(x)\bar{\rho}(y)) dydx =: T_5 + T_6 + T_7,$$

where the terms T_5 , T_6 and T_7 correspond to the part of the sum where $i-j=0$, $|i-j|=1$ and $|i-j| \geq 2$ respectively. We bound all these three terms separately. With this aim, we set

$$\bar{\varphi}_i := \bar{\varphi} \mathbf{1}_{(\bar{x}_{i-1}, \bar{x}_i)} \quad \text{for } i = 1, \dots, n,$$

and note that, by (36) and Lemma 4.1(iii),

$$\sum_{i=1}^n \|\bar{\varphi}_i\|_V^2 = 2D_n(\bar{\mathbf{x}}) \leq Cn^{-1+a}.$$

For T_5 we simply estimate

$$\begin{aligned} T_5 &= \sum_{i=1}^n \int_{\bar{x}_{i-1}}^{\bar{x}_i} \int_{\bar{x}_{i-1}}^{\bar{x}_i} V(x-y) (\bar{\varphi}(x)\bar{\varphi}(y) - \bar{\rho}(x)\bar{\rho}(y)) dydx \\ &\leq \sum_{i=1}^n \int_{\bar{x}_{i-1}}^{\bar{x}_i} \int_{\bar{x}_{i-1}}^{\bar{x}_i} V(x-y) \bar{\varphi}(x)\bar{\varphi}(y) dydx = \sum_{i=1}^n \|\bar{\varphi}_i\|_V^2 \leq Cn^{-1+a}. \end{aligned}$$

For T_6 , we similarly obtain

$$\begin{aligned} T_6 &= 2 \sum_{i=1}^{n-1} \int_{\bar{x}_i}^{\bar{x}_{i+1}} \int_{\bar{x}_{i-1}}^{\bar{x}_i} V(x-y) (\bar{\varphi}(x)\bar{\varphi}(y) - \bar{\rho}(x)\bar{\rho}(y)) dydx \\ &\leq 2 \sum_{i=1}^{n-1} (\bar{\varphi}_{i+1}, \bar{\varphi}_i)_V \leq \sum_{i=1}^{n-1} (\|\bar{\varphi}_{i+1}\|_V^2 + \|\bar{\varphi}_i\|_V^2) \leq Cn^{-1+a}. \end{aligned} \tag{40}$$

Finally we estimate T_7 . We note that in the integrals in the terms of T_7 , the singularity of V is avoided. This allows for pointwise evaluation of the integrand. By using that V is even, and non-increasing on the positive axis, we estimate

$$\begin{aligned} T_7 &= 2 \sum_{i=3}^n \sum_{j=1}^{i-2} \left(\int_{\bar{x}_{i-1}}^{\bar{x}_i} \int_{\bar{x}_{j-1}}^{\bar{x}_j} V(x-y) (\bar{\varphi}(x)\bar{\varphi}(y) - \bar{\rho}(x)\bar{\rho}(y)) dydx \right. \\ &\quad \leq 2 \sum_{i=3}^n \sum_{j=1}^{i-2} \left(\int_{\bar{x}_{i-1}}^{\bar{x}_i} \int_{\bar{x}_{j-1}}^{\bar{x}_j} V(\bar{x}_{i-1} - \bar{x}_j) \bar{\varphi}(x)\bar{\varphi}(y) dydx \right. \\ &\quad \quad \left. \left. - \int_{\bar{x}_{i-1}}^{\bar{x}_i} \int_{\bar{x}_{j-1}}^{\bar{x}_j} V(\bar{x}_i - \bar{x}_{j-1}) \bar{\rho}(x)\bar{\rho}(y) dydx \right) \right. \\ &= \frac{2}{n^2} \sum_{i=3}^n \sum_{j=1}^{i-2} (V(\bar{x}_{i-1} - \bar{x}_j) - V(\bar{x}_i - \bar{x}_{j-1})). \end{aligned}$$

We recognise a telescopic series after changing the summation index to $k = i + j - 1$:

$$\begin{aligned}
T_7 &\leq \frac{2}{n^2} \sum_{i=3}^n \sum_{j=1}^{i-2} (V(\bar{x}_{i-1} - \bar{x}_j) - V(\bar{x}_i - \bar{x}_{j-1})) \\
&= \frac{2}{n^2} \sum_{i=3}^n \sum_{k=i}^{2i-3} (V(\bar{x}_{i-1} - \bar{x}_{k-(i-1)}) - V(\bar{x}_i - \bar{x}_{k-i})) \\
&= \frac{2}{n^2} \sum_{k=3}^{2n-3} \sum_{i=\lceil \frac{k+3}{2} \rceil}^{k \wedge n} (V(\bar{x}_{i-1} - \bar{x}_{k-(i-1)}) - V(\bar{x}_i - \bar{x}_{k-i})) \\
&= \frac{2}{n^2} \sum_{k=3}^{2n-3} (V(\bar{x}_{\lceil \frac{k+1}{2} \rceil} - \bar{x}_{\lfloor \frac{k-1}{2} \rfloor}) - V(\bar{x}_{k \wedge n} - \bar{x}_{0 \vee (k-n)})) \\
&\leq \frac{2}{n^2} \sum_{k=3}^{2n-3} V(\bar{x}_{\lceil \frac{k+1}{2} \rceil} - \bar{x}_{\lfloor \frac{k-1}{2} \rfloor}) \leq \frac{4}{n^2} \sum_{i=2}^{n-1} V(\bar{\ell}_i) \leq 4E_n^{\text{nn}}(\bar{\mathbf{x}})
\end{aligned}$$

which, by Lemma 4.1(iii), is bounded by Cn^{-1+a} . \square

The upper and lower bound on $E(\varphi) - E_n(\mathbf{x})$ First, we state and prove the opposite inequality in (37) as an auxiliary result.

Lemma 4.4. *There exists constants $C, C' > 0$ such that for all $n \geq 1$ and all $\mathbf{x} \in \Omega$*

$$D_n(\mathbf{x}) \leq CE_n^{\text{nn}}(\mathbf{x}) + \frac{C'}{n}.$$

Proof. Using $V = V_a + V_{\text{reg}}$, we split $D_n(\mathbf{x})$ in two parts:

$$D_n(\mathbf{x}) = \frac{1}{2} \frac{1}{n^2} \sum_{i=1}^n \frac{1}{\ell_i^2} \iint_{(0, \ell_i)^2} V_a(x - y) dy dx + \frac{1}{2} \frac{1}{n^2} \sum_{i=1}^n \frac{1}{\ell_i^2} \iint_{(0, \ell_i)^2} V_{\text{reg}}(x - y) dy dx.$$

The first part can be computed explicitly. This yields

$$\frac{1}{2} \frac{1}{n^2} \sum_{i=1}^n \frac{1}{\ell_i^2} \iint_{(0, \ell_i)^2} V_a(x - y) dy dx = \frac{C_a}{n^2} \sum_{i=1}^n V_a(\ell_i)$$

for some explicit constant $C_a > 0$. For the second term, we rely on the regularity of V_{reg} to estimate

$$\frac{1}{2} \frac{1}{n^2} \sum_{i=1}^n \frac{1}{\ell_i^2} \iint_{(0, \ell_i)^2} V_{\text{reg}}(x - y) dy dx \leq \frac{1}{n} \left(\frac{1}{2} + C_a \right) \|V_{\text{reg}}\|_{C([-1, 1])} + \frac{C_a}{n^2} \sum_{i=1}^n V_{\text{reg}}(\ell_i).$$

Collecting these findings, we obtain the estimate in Lemma 4.4:

$$D_n(\mathbf{x}) \leq \frac{C}{n^2} \sum_{i=1}^n (V_a + V_{\text{reg}})(\ell_i) + \frac{C'}{n} = CE_n^{\text{nn}}(\mathbf{x}) + \frac{C'}{n}.$$

\square

Lemma 4.5 (Energy bounds on the piecewise constant approximation). *There exists $C \geq 0$ such that for all $n \geq 1$ and all $\mathbf{x} \in \Omega$*

$$-E_n^{\text{nn}}(\mathbf{x}) - \frac{1}{n}(U(0) + U(1)) \leq E(\varphi) - E_n(\mathbf{x}) \leq C \left(E_n^{\text{nn}}(\mathbf{x}) + \frac{1}{n} \right),$$

where φ is the piece-wise constant function constructed from \mathbf{x} by (7).

Proof. We divide the proof in four steps. In Step 1 we bound the confinement part of $E(\varphi) - E_n(\mathbf{x})$, and in Steps 2 – 4 we bound the interaction part. Given \mathbf{x} , we let m_i and ℓ_i be defined by (35) (see Figure 3).

Step 1: bounds on the confinement part. The confinement part of $E(\varphi) - E_n(\mathbf{x})$ is given by

$$F_n(\mathbf{x}) := \int_0^1 U\varphi - \frac{1}{n} \sum_{i=0}^n U(x_i) = \frac{1}{n} \sum_{i=1}^n \frac{1}{\ell_i} \int_{x_{i-1}}^{x_i} U - \frac{1}{n} \sum_{i=0}^n U(x_i).$$

Since $U \geq 0$ is convex, it is easy to see that $-\frac{1}{n}(U(0) + U(1)) \leq F_n(\mathbf{x}) \leq 0$.

In the remainder of the proof, we set $U \equiv 0$ to focus on the interaction part.

Step 2: rewriting $E(\varphi) - E_n(\mathbf{x})$ as a sum of error terms. We show that

$$E(\varphi) - E_n(\mathbf{x}) = D_n(\mathbf{x}) + Q_n(\mathbf{x}) - R_n(\mathbf{x}) - B_n(\mathbf{x}), \quad (41)$$

where the four *non-negative* error terms are given by (36) and

$$\begin{aligned} Q_n(\mathbf{x}) &:= \frac{1}{n^2} \sum_{i=2}^n \sum_{j=1}^{i-1} \left[\frac{1}{\ell_i \ell_j} \int_{x_{i-1}}^{x_i} \int_{x_{j-1}}^{x_j} V(x-y) dy dx - V(m_i - m_j) \right], \\ R_n(\mathbf{x}) &:= \frac{1}{n^2} \sum_{i=2}^n \sum_{j=1}^{i-1} \left(\frac{1}{2} V(x_i - x_j) + \frac{1}{2} V(x_{i-1} - x_{j-1}) - V(m_i - m_j) \right), \\ B_n(\mathbf{x}) &:= \frac{1}{2} \frac{1}{n^2} \sum_{i=1}^n [V(x_i - x_0) + V(x_n - x_{i-1})]. \end{aligned}$$

Indeed, (41) follows from

$$\begin{aligned} E(\varphi) &= \frac{1}{2} \int_0^1 \int_0^1 V(x-y) \varphi(x) \varphi(y) dy dx \\ &= \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \int_{x_{i-1}}^{x_i} \int_{x_{j-1}}^{x_j} V(x-y) \frac{1/n}{x_i - x_{i-1}} \frac{1/n}{x_j - x_{j-1}} dy dx \\ &= \frac{1}{n^2} \sum_{i=2}^n \sum_{j=1}^{i-1} \frac{1}{\ell_i \ell_j} \int_{x_{i-1}}^{x_i} \int_{x_{j-1}}^{x_j} V(x-y) dy dx + D_n(\mathbf{x}) \\ &= \frac{1}{n^2} \sum_{i=2}^n \sum_{j=1}^{i-1} V(m_i - m_j) + (D_n + Q_n)(\mathbf{x}) \\ &= \frac{1}{2n^2} \sum_{i=2}^n \sum_{j=1}^{i-1} (V(x_i - x_j) + V(x_{i-1} - x_{j-1})) + (D_n + Q_n - R_n)(\mathbf{x}) \\ &= E_n(\mathbf{x}) + (D_n + Q_n - R_n - B_n)(\mathbf{x}). \end{aligned} \quad (42)$$

Step 3: the lower bound for $E(\varphi) - E_n(\mathbf{x})$. Since the error terms D_n , Q_n , R_n and B_n are all non-negative, we observe from (41) that it is enough to show that

$$R_n(\mathbf{x}) + B_n(\mathbf{x}) \leq E_n^{\text{nn}}(\mathbf{x}).$$

By using $m_i - m_j \leq x_i - x_{j-1}$, we obtain this estimate from

$$\begin{aligned} R_n(\mathbf{x}) + B_n(\mathbf{x}) &= \frac{1}{n^2} \sum_{i=1}^n \sum_{j=0}^{i-1} V(x_i - x_j) - \frac{1}{n^2} \sum_{i=2}^n \sum_{j=1}^{i-1} V(m_i - m_j) \\ &\leq \frac{1}{n^2} \sum_{i=1}^n \sum_{j=0}^{i-1} V(x_i - x_j) - \frac{1}{n^2} \sum_{i=2}^n \sum_{j=0}^{i-2} V(x_i - x_j) = E_n^{\text{nn}}(\mathbf{x}). \end{aligned}$$

Step 4: the upper bound for $E(\varphi) - E_n(\mathbf{x})$. Since $B_n \geq 0$, it is enough to show that

$$D_n(\mathbf{x}) + Q_n(\mathbf{x}) - R_n(\mathbf{x}) \leq C \left(E_n^{\text{nn}}(\mathbf{x}) + \frac{1}{n} \right).$$

Then, by Lemma 4.4, it suffices to show that $Q_n - R_n \leq 2D_n$. Writing

$$\begin{aligned} Q_n(\mathbf{x}) - R_n(\mathbf{x}) &= \frac{1}{n^2} \sum_{i=2}^n \sum_{j=1}^{i-1} \left[\frac{1}{\ell_i \ell_j} \int_{x_{i-1}}^{x_i} \int_{x_{j-1}}^{x_j} V(x-y) dy dx - \left(\frac{1}{2} V(x_i - x_j) + \frac{1}{2} V(x_{i-1} - x_{j-1}) \right) \right], \end{aligned}$$

we use convexity of V to bound the integral for $i \geq j+2$ by

$$\frac{1}{\ell_i \ell_j} \int_{x_{i-1}}^{x_i} \int_{x_{j-1}}^{x_j} V(x-y) dy dx \leq \frac{1}{2} V(x_{i-1} - x_j) + \frac{1}{2} V(x_i - x_{j-1}).$$

This yields

$$\begin{aligned} Q_n(\mathbf{x}) - R_n(\mathbf{x}) &\leq \frac{1}{n^2} \sum_{i=1}^{n-1} \frac{1}{\ell_{i+1} \ell_i} \int_{x_i}^{x_{i+1}} \int_{x_{i-1}}^{x_i} V(x-y) dy dx \\ &\quad + \frac{1}{2n^2} \left(\sum_{i=3}^n \sum_{j=1}^{i-2} [V(x_{i-1} - x_j) + V(x_i - x_{j-1})] - \sum_{i=2}^n \sum_{j=1}^{i-1} [V(x_i - x_j) + V(x_{i-1} - x_{j-1})] \right). \end{aligned} \tag{43}$$

For the term within parentheses, a change of index readily reveals that the second summation includes all terms of the first summation. We then use $V \geq 0$ to estimate this term from above by 0. The remaining term in (43) can be estimated similarly as in (40). This yields $Q_n - R_n \leq 2D_n$. \square

5 Lower bound on E_n

In this section we prove (16), which is the crucial step in Step 3 of the sketch of the proof of Theorem 1.2. More precisely, we assume that V and $\bar{\rho}$ satisfy (9), and prove the following proposition.

Proposition 5.1 (Lower bound on E_n). *There exists $C > 0$ such that for all $n \geq 1$ and all $\mathbf{x} \in \Omega$*

$$E_n(\mathbf{x}) - E_n^{\text{mn}}(\mathbf{x}) - E(\bar{\rho}) \geq -Cn^{-1+a}.$$

We give the proof of Proposition 5.1 after a preliminary construction of a renormalised norm of $\|\cdot\|_V$. With this aim, we introduce

$$V_n(r) := \begin{cases} V(\frac{1}{n}) + (r - \frac{1}{n})V'(\frac{1}{n}) & \text{if } 0 \leq r \leq \frac{1}{n} \\ V(r) & \text{if } r > \frac{1}{n} \end{cases} \quad (44)$$

with even extension to the negative half-line. Figure 4 illustrates a typical example of V and V_n . Lemma 5.2 lists several basic properties of V_n .




Figure 4: The piecewise-affine regularisation V_n of the interaction potential V .

Lemma 5.2 (Properties of V_n). *There exists a constant $C > 0$ such that for all $n \geq 1$:*

- (i) V_n is non-increasing on $[0, \infty)$;
- (ii) V_n and $V - V_n$ are convex on $(0, \infty)$;
- (iii) $\text{supp}(V - V_n) \subset [-\frac{1}{n}, \frac{1}{n}]$;
- (iv) $V_n(0) \leq Cn^a$;
- (v) For $f \in L^2(\mathbb{R})$, $\|f\|_{V_n} := \sqrt{\int_{\mathbb{R}} (V_n * f)f}$ defines a semi-norm;
- (vi) $V_n \uparrow V$ in $L^p(\mathbb{R})$ as $n \rightarrow \infty$ for any $1 \leq p < \frac{1}{a}$.

Proof. Except for (v), all properties are a direct consequence of the assumptions and properties of V and the definition of V_n in (44). Property (v) can be proven along the lines of [KvM19, Lem. 3.2]; it relies on the Fourier-transform of V_n being non-negative, which easily follows from the other properties of V_n (see [KvM19, Lem. 3.1] for details). \square

Next we establish an auxiliary estimate on $(V - V_n) * \bar{\rho}$.

Lemma 5.3. *There exists $C > 0$ such that for all $n \geq 1$ and all $0 \leq x \leq \frac{1}{2}$*

$$((V - V_n) * \bar{\rho})(x) + ((V - V_n) * \bar{\rho})(1 - x) \leq Cn^{-\frac{1-a}{2}} \min \left\{ 1, [nx - 1]_+^{-\frac{1-a}{2}} \right\}.$$

Proof. Take any $0 \leq x \leq \frac{1}{2}$, and set $\underline{x} := (x - \frac{1}{n}) \vee 0$. By Lemma 5.2(iii))

$$((V - V_n) * \bar{\rho})(x) = \int_{\underline{x}}^{x+\frac{1}{n}} (V - V_n)(x - y) \bar{\rho}(y) dy.$$

Since $V_n \geq 0$, we have that $(V - V_n)(r) \leq V(r) \leq C/|r|^a$ for all $|r| \leq 1$. Together with the upper bound on $\bar{\rho}$ in Lemmas 3.4(iii) and $\bar{x} \geq x - \frac{1}{n}$, we continue the estimate by

$$\int_{\underline{x}}^{x+\frac{1}{n}} (V - V_n)(x - y) \bar{\rho}(y) dy \leq C \int_{x-\frac{1}{n}}^{x+\frac{1}{n}} |x - y|^{-a} |y|^{-\frac{1-a}{2}} dy. \quad (45)$$

Noting that $a > \frac{2}{1+a}$, we apply Hölder's inequality with $p \in (\frac{2}{1+a}, a)$ and conjugate exponent $q = p/(p-1)$. This yields

$$\int_{x-\frac{1}{n}}^{x+\frac{1}{n}} |x - y|^{-a} |y|^{-\frac{1-a}{2}} dy \leq \left(\int_{-\frac{1}{n}}^{\frac{1}{n}} |y|^{-ap} dy \right)^{1/p} \left(\int_{x-\frac{1}{n}}^{x+\frac{1}{n}} |y|^{-\frac{1-a}{2}q} dy \right)^{1/q}.$$

Since $ap < 1$ and $\frac{1-a}{2}q < 1$ by construction, the right-hand side is finite. Noting that the right-hand side is maximal at $x = 0$, we obtain

$$\left(\int_{-\frac{1}{n}}^{\frac{1}{n}} |y|^{-ap} dy \right)^{1/p} \left(\int_{-\frac{1}{n}}^{\frac{1}{n}} |y|^{-\frac{1-a}{2}q} dy \right)^{1/q} = C \left(\frac{1}{n} \right)^{\frac{1}{p}-a} \left(\frac{1}{n} \right)^{\frac{1}{q}-\frac{1-a}{2}} = Cn^{-\frac{1-a}{2}}.$$

In conclusion, the estimates above yield that

$$((V - V_n) * \bar{\rho})(x) \leq Cn^{-\frac{1-a}{2}}$$

for all $0 \leq x \leq \frac{1}{2}$.

To sharpen the bound for $\frac{2}{n} \leq x \leq \frac{1}{2}$, we follow the estimate above until (45), and continue as follows:

$$\begin{aligned} ((V - V_n) * \bar{\rho})(x) &\leq C \int_{x-\frac{1}{n}}^{x+\frac{1}{n}} |x - y|^{-a} |y|^{-\frac{1-a}{2}} dy \\ &\leq C(x - \frac{1}{n})^{-\frac{1-a}{2}} \int_{x-\frac{1}{n}}^{x+\frac{1}{n}} |x - y|^{-a} dy \leq C'n^{-1+a}(x - \frac{1}{n})^{-\frac{1-a}{2}}. \end{aligned}$$

The estimate for $((V - V_n) * \bar{\rho})(1 - x)$ is analogous. \square

Proof of Proposition 5.1. The assertion of Proposition 5.1 is obvious when n is bounded from above by a fixed integer $N \geq 1$. In this case, it suffices to take $C = N^{1-a}E(\bar{\rho})$. Therefore, it is not restrictive to assume that $n \geq 1/b$, where $b > 0$ is as in (30).

Let $\mathbf{x} \in \Omega$ be given, and set

$$\mu_n := \frac{1}{n} \sum_{i=0}^n \delta_{x_i} \quad \text{and} \quad \nu_n := \mu_n - \bar{\rho}.$$

Let

$$\Delta_1 := \{(x_i, x_j) : |i - j| \leq 1\} \subset \mathbb{R}^2$$

be the particle pairs that are left out in the interaction term of $E_n(\mathbf{x}) - E_n^{\text{nn}}(\mathbf{x})$, i.e.

$$E_n(\mathbf{x}) - E_n^{\text{nn}}(\mathbf{x}) = \frac{1}{2} \iint_{\Delta_1^c} V(x - y) d\mu_n(y) d\mu_n(x) + \int U d\mu_n.$$

Then, we use Lemma 3.4(iv) to estimate

$$\begin{aligned} & E_n(\mathbf{x}) - E_n^{\text{nn}}(\mathbf{x}) - E(\bar{\rho}) \\ &= \frac{1}{2} \iint_{\Delta_1^c} V(x - y) d\nu_n(y) d\nu_n(x) + \int (V * \bar{\rho}) d\nu_n + \int U d\nu_n \\ &\geq \frac{1}{2} \iint_{\Delta_1^c} V(x - y) d\nu_n(y) d\nu_n(x) \\ &= \underbrace{\frac{1}{2} \iint_{\Delta_1^c} V_n(x - y) d\nu_n(y) d\nu_n(x)}_{T_8} + \underbrace{\frac{1}{2} \iint_{\Delta_1^c} (V - V_n)(x - y) d\nu_n(y) d\nu_n(x)}_{T_9}, \end{aligned}$$

where V_n is the regularisation introduced in (44). To bound T_8 , we use that $\max_{\mathbb{R}} V_n = V_n(0) \leq Cn^a$ by Lemma 5.2(i),(iv) and that $\|\cdot\|_{V_n}^2$ is a norm (see Lemma 5.2(v)). Then, by the definition of Δ_1 , this yields

$$T_8 = \|\nu_n\|_{V_n}^2 - \frac{n+1}{n^2} V_n(0) - \frac{2}{n^2} \sum_{i=1}^n V_n(x_i - x_{i-1}) \geq -Cn^{-1+a}.$$

It remains to bound T_9 from below by $-Cn^{-1+a}$. We expand $\nu_n = \mu_n - \bar{\rho}$ to rewrite

$$T_9 = \frac{2}{n^2} \sum_{i=2}^n \sum_{j=0}^{n-i} (V - V_n)(x_{j+i} - x_j) - 2 \int ((V - V_n) * \bar{\rho}) d\mu_n + \iint_{\mathbb{R}^2} (V - V_n)(x - y) d\bar{\rho}(y) d\bar{\rho}(x). \quad (46)$$

The third term is non-negative; we bound it from below by 0. For the first two terms, we assume for convenience that n is a multiple of 4, and partition the interval $[0, 1]$ into the closed intervals $I_k := [2\frac{k-1}{n}, 2\frac{k}{n}]$ where $k = 1, \dots, \frac{n}{2}$. Note that these intervals only overlap at their endpoints. Then, we remove the contribution of the interaction between any two particles located in different intervals I_k from the double sum in the right-hand side of (46). Finally, we minimise the right-hand side of (46) over each I_k separately, and relax the constraint that the total number of particles should be $n + 1$. This yields

$$T_9 \geq 2 \sum_{k=1}^{n/2} \min_{N \in \mathbb{N}} \left(\underbrace{\frac{1}{n^2} \min_{2\frac{k-1}{n} \leq y_1 \leq y_2 \leq \dots \leq y_N \leq 2\frac{k}{n}} \sum_{i=2}^N \sum_{j=1}^{N-i} (V - V_n)(y_{j+i} - y_j)}_{T_{10}} - \frac{N}{n} \|(V - V_n) * \bar{\rho}\|_{C(I_k)} \right). \quad (47)$$

We treat both terms within the parentheses separately. For the second term, we apply the bound in Lemma 5.3. Since this bound gives the same estimate for the intervals I_k and

$I_{n/2-k+1}$, we focus on bounding it for $k \leq n/4$. This yields

$$\begin{aligned} \frac{N}{n} \|(V - V_n) * \bar{\rho}\|_{C(I_k)} &\leq C \frac{N}{n} n^{-\frac{1-a}{2}} \min \left\{ 1, [2(k-1) - 1]_+^{-\frac{1-a}{2}} \right\} \\ &\leq C' N n^{-1-\frac{1-a}{2}} \begin{cases} 1 & k = 1 \\ (k-1)^{-\frac{1-a}{2}} & k \geq 2. \end{cases} \end{aligned}$$

In particular, if the minimum over N in (47) is reached below an n -independent value C (i.e., $N \leq C$), then it suffices to bound the first term in parentheses in (47) from below simply by 0. Therefore, we assume next that the minimiser N is sufficiently large; in particular $N \geq 9$. We further assume for simplicity that N is a multiple of 3.

To bound the term T_{10} in (47), we rely on the basic arguments in the theory of i -th neighbour interaction energies with convex interaction potentials. In more detail, first we use that the summands are non-negative to remove the latter terms in the sum over i :

$$T_{10} \geq \frac{1}{n^2} \min_{2^{\frac{k-1}{n}} \leq y_1 \leq y_2 \leq \dots \leq y_N \leq 2^{\frac{k}{n}}} \sum_{i=2}^{N/3} \sum_{j=1}^{N-i} (V - V_n)(y_{j+i} - y_j).$$

Then, we bound the minimum from below by exchanging the sum over i with the minimisation over $(y_j)_j$, i.e.,

$$T_{10} \geq \frac{1}{n^2} \sum_{i=2}^{N/3} \min_{2^{\frac{k-1}{n}} \leq y_1 \leq y_2 \leq \dots \leq y_N \leq 2^{\frac{k}{n}}} \sum_{j=1}^{N-i} (V - V_n)(y_{j+i} - y_j) =: T_{11}.$$

Then, the resulting minimisation problem can be written as a sum over independent minimisation problems. To see this, we change variables in the summation index by $j = (m-1)i + \ell$, and write the sum over j as two sums over ℓ and m . By possibly skipping a few terms for those j that are close to $N-i$, we obtain

$$\begin{aligned} T_{11} &\geq \frac{1}{n^2} \sum_{i=2}^{N/3} \min_{2^{\frac{k-1}{n}} \leq y_1 \leq y_2 \leq \dots \leq y_N \leq 2^{\frac{k}{n}}} \sum_{\ell=1}^i \sum_{m=1}^{\lfloor \frac{N-\ell}{i} \rfloor} (V - V_n)(y_{mi+\ell} - y_{(m-1)i+\ell}) \\ &\geq \frac{1}{n^2} \sum_{i=2}^{N/3} \sum_{\ell=1}^i \min_{2^{\frac{k-1}{n}} \leq y_1 \leq y_2 \leq \dots \leq y_N \leq 2^{\frac{k}{n}}} \sum_{m=1}^{\lfloor \frac{N-\ell}{i} \rfloor} (V - V_n)(y_{mi+\ell} - y_{(m-1)i+\ell}) \\ &= \frac{1}{n^2} \sum_{i=2}^{N/3} \sum_{\ell=1}^i \min_{2^{\frac{k-1}{n}} \leq y_\ell \leq y_{i+\ell} \leq y_{2i+\ell} \leq \dots \leq y_{\lfloor \frac{N-\ell}{i} \rfloor i + \ell} \leq 2^{\frac{k}{n}}} \sum_{m=1}^{\lfloor \frac{N-\ell}{i} \rfloor} (V - V_n)(y_{mi+\ell} - y_{(m-1)i+\ell}) \\ &= \frac{1}{n^2} \sum_{i=2}^{N/3} \sum_{\ell=1}^i \min_{2^{\frac{k-1}{n}} \leq z_0 \leq z_1 \leq \dots \leq z_{\lfloor \frac{N-\ell}{i} \rfloor} \leq 2^{\frac{k}{n}}} \sum_{m=1}^{\lfloor \frac{N-\ell}{i} \rfloor} (V - V_n)(z_m - z_{m-1}), \end{aligned}$$

where in the last equality we change to the variable $z_m := y_{mi+\ell}$. Each such minimisation problem over z_m involves only nearest neighbour interactions with the convex, repelling interaction potential $V - V_n$, which is minimised by the equispaced configuration. Plugging in

the equispaced configuration $z_m = 2^{\frac{k-1}{n}} + 2m/(n \lfloor \frac{N-\ell}{i} \rfloor)$, we get

$$\begin{aligned}
T_{11} &\geq \frac{1}{n^2} \sum_{i=2}^{N/3} \sum_{\ell=1}^i \lfloor \frac{N-\ell}{i} \rfloor (V - V_n) \left(\frac{2}{n \lfloor \frac{N-\ell}{i} \rfloor} \right) \geq \frac{1}{n^2} \sum_{i=2}^{N/3} \sum_{\ell=1}^i \left(\frac{2}{3} \frac{N}{i} - 1 \right) (V - V_n) \left(\frac{3i}{nN} \right) \\
&= \frac{1}{n^2} \sum_{i=2}^{N/3} \left(\frac{2}{3} N - i \right) (V - V_n) \left(\frac{3i}{nN} \right) \geq \frac{N}{3n^2} \sum_{i=2}^{N/3} (V - V_n) \left(\frac{3i}{nN} \right) \\
&\geq \frac{N^2}{9n} \int_{\frac{6}{nN}}^{\frac{1}{n}} (V - V_n)(x) dx = \frac{N^2}{9n^2} \int_{\frac{6}{N}}^1 (V - V_n) \left(\frac{x}{n} \right) dx,
\end{aligned} \tag{48}$$

where in the last inequality we have recognized the sum as a Riemann upper-sum. To estimate the integrand from below, we integrate twice, use that $(V - V_n)(\frac{1}{n}) = 0$ and rely on $V_n'' = 0$ on $(0, \frac{1}{n})$ and the lower bound on V'' on $(0, b) \supset (0, \frac{1}{n})$ in (30) to deduce that

$$(V - V_n) \left(\frac{x}{n} \right) = \int_{\frac{x}{n}}^{\frac{1}{n}} \int_y^{\frac{1}{n}} V''(z) dz dy \geq C \int_{\frac{x}{n}}^{\frac{1}{n}} \int_y^{\frac{1}{n}} z^{-2-a} dz dy = Cn^a \int_x^1 \int_y^1 z^{-2-a} dz dy$$

for all $0 < x < 1$. The double integral in the right-hand side is independent of n , positive, and decreasing as a function of x . Using this and noting that $6/N \leq 2/3$, we continue the estimate in (48) by

$$\frac{N^2}{9n^2} \int_{\frac{6}{N}}^1 (V - V_n) \left(\frac{x}{n} \right) dx \geq CN^2 n^{-2+a} \int_{\frac{2}{3}}^1 \int_x^1 \int_y^1 z^{-2-a} dz dy dx \geq C' N^2 n^{-2+a}.$$

Finally, collecting our estimates in (47), we obtain two constants $C, C' > 0$ such that

$$\begin{aligned}
T_9 &\geq C' \sum_{k=1}^{n/2} \min_{N \in \mathbb{R}} \left(N^2 n^{-2+a} - Ck^{-\frac{1-a}{2}} N n^{-1-\frac{1-a}{2}} \right) \\
&= C' n^{-2+a} \sum_{k=1}^{n/2} \min_{N \in \mathbb{R}} N \left(N - C \left(\frac{k}{n} \right)^{-\frac{1-a}{2}} \right) \\
&= -C' n^{-2+a} \frac{C^2}{4} \sum_{k=1}^{n/2} \left(\frac{k}{n} \right)^{-1+a} \geq -C n^{-1+a}.
\end{aligned}$$

□

Remark 5.4 (The case $\bar{\rho} \leq C$). When $\bar{\rho}$ is bounded, the proof of Proposition 5.1 simplifies significantly. Indeed, if $\bar{\rho}$ is bounded, then instead of Lemma 5.3 the rougher estimate $\|(V - V_n) * \bar{\rho}\|_{C([0,1])} \leq Cn^{-1+a}$ is sufficient, because this estimate gives immediately the desired bound on the second term in (46).

6 Proof of Theorem 1.2

We first treat the case $0 < a < 1$. Given V and U as in Theorem 1.2, we start by constructing a more convenient interaction potential \tilde{V} whose corresponding energies \tilde{E} and \tilde{E}_n have the

same sets of minimisers as E and E_n respectively. Let R and S be as in Proposition 3.1. Let \tilde{V} satisfy Assumption 1.1 such that

$$\tilde{V} = V \text{ on } (0, \max\{2S, R\}] \quad \text{and} \quad \tilde{V} = C \text{ on } [\max\{2S, R\} + 1, \infty)$$

for some constant $C \in \mathbb{R}$. Such a \tilde{V} can be obtained by multiplying $V'(x)$ with a cut-off function, and then integrating from x to ∞ . From the integrability and convexity of \tilde{V} we note that $C = \min_{\mathbb{R}} \tilde{V}$. Then, Proposition 3.1 applies to \tilde{V} with the same constants R and S , and thus any minimiser of

$$\tilde{E}(\rho) := \frac{1}{2} \int_{\mathbb{R}} (\tilde{V} * \rho) d\rho + \int_{\mathbb{R}} U d\rho$$

is also supported in $[-S, S]$. Since by the choice of \tilde{V} it holds that $\tilde{E} = E$ on $\mathcal{P}([-S, S])$, any minimiser of E is a minimiser of \tilde{E} and vice versa. Analogously, we obtain the same conclusion for E_n for a possibly different constant S . Hence, we may replace V in Theorem 1.2 by \tilde{V} . In addition, we may further subtract the constant $C/2 = \min_{\mathbb{R}} \tilde{V}/2$ from \tilde{E} and subtract the constant $(1 + \frac{1}{n})C/2$ from \tilde{E}_n such that the resulting interaction potential has compact support. We denote this potential by \tilde{V} without changing notation.

The existence of minimisers of E_n is shown in Remark 3.3. Since \tilde{V} has compact support, Lemma 3.4 applies. This shows that \tilde{E} has a unique minimiser $\tilde{\rho}$. Moreover, by Lemma 3.4(i) there exists an affine change of variables such that $\text{supp } \tilde{\rho}$ turns into the interval $[0, 1]$. It is not difficult to verify (see [KvM19, Step 1 in the proof of Lem. 6.8]) that under this change of variables, E and E_n (possibly multiplied or shifted by a constant) still satisfy Assumption 1.1, and that the resulting Sobolev norm (8) is equivalent to that before the change of variables. Hence, redefining $V, U, E, E_n, \bar{\rho}, \mathbf{x}^*$ as the potentials, energies and minimisers that appear after the affine change of variables has been applied, we observe that V, U satisfy Assumption 1.1, that the additional property (9) holds, and that $\|\varphi^* - \bar{\rho}\|_{H^{-(1-a)/2}(\mathbb{R})}^2$ has changed by an n -independent multiplicative constant with respect to the original value.

The remaining part of the proof concerns the estimate of $\|\varphi^* - \bar{\rho}\|_{H^{-(1-a)/2}(\mathbb{R})}^2$. This part is already sufficiently detailed in the sketch of the proof given in Section 1.1; see (11)–(16) and the references therein to Sections 4 and 5. This completes the proof of Theorem 1.2 in the case $0 < a < 1$.

Finally, we treat the case $a = 0$. The proof is analogous to the case $0 < a < 1$; the only differences are several minor changes in the computations. All these changes are ramifications of the change in the upper bound on V , which is

$$V(r) \leq C - \log |r|.$$

In Table 2 and in the two items listed below we mention all statements of Sections 3 – 5 which are not literally valid for $a = 0$, and provide the required modification.

1. The constant C' in Lemma 4.4 also contains a contribution from V_a .

Statement	updated estimate
Lemma 4.1(iii)	$D_n(\bar{\mathbf{x}}) + E_n^{\text{nn}}(\bar{\mathbf{x}}) \leq Cn^{-1} \log n$
Lemma 4.3	$\ \bar{\varphi}\ _V^2 - \ \bar{\rho}\ _V^2 \leq Cn^{-1} \log n$
Lemma 5.2(iv)	$V_n(0) \leq \log n + C$
Lemma 5.2(vi)	$V_n \uparrow V$ in $L^p(\mathbb{R})$ as $n \rightarrow \infty$ for any $1 \leq p < \infty$
Lemma 5.3	$\begin{cases} ((V - V_n) * \bar{\rho})(x) + ((V - V_n) * \bar{\rho})(1 - x) \\ \leq Cn^{-1/2}(\log n) \min \left\{ 1, [nx - 1]_+^{-1/2} \right\} \end{cases}$
Proposition 5.1	$E_n(\mathbf{x}) - E_n^{\text{nn}}(\mathbf{x}) - E(\bar{\rho}) \geq -Cn^{-1}(\log n)^3$

Table 2: Changes in the estimates for $a = 0$.

2. In the proof of Proposition 5.1, the estimate in the final display changes as follows:

$$\begin{aligned}
T_9 &\geq C' \sum_{k=1}^{n/2} \min_{N \in \mathbb{R}} \left(N^2 n^{-2} - Ck^{-1/2} N n^{-3/2} \log n \right) \\
&= C' n^{-2} \sum_{k=1}^{n/2} \min_{N \in \mathbb{R}} N \left(N - C(\log n) \left(\frac{k}{n} \right)^{-1/2} \right) \\
&= -\frac{C'}{n} (\log n)^2 \frac{C^2}{4} \frac{1}{n} \sum_{k=1}^{n/2} \left(\frac{k}{n} \right)^{-1} \geq -Cn^{-1} (\log n)^3.
\end{aligned}$$

7 Numerical computations on the rate in Theorem 1.2

The aim of this section is to compare the upper bound of the convergence rate in Theorem 1.2 with the actual convergence rate in concrete examples. These concrete examples are given by specific choices for the potentials V and U for which all quantities except for \mathbf{x}^* can be computed explicitly. With this aim, we take $V_{\text{reg}} = 0$ and U a convex polynomial on $D(U)$. Given the qualitatively different profiles of $\bar{\rho}$ observed in Figure 2, we consider two choices for $D(U)$; a bounded interval (Case 1) and \mathbb{R} (Case 2).

For each of these two cases, the method to test Theorem 1.2 numerically is as follows. First, we compute \mathbf{x}^* by minimizing E_n in (1) numerically with Newton's method for several values of n . One observation we did from this data is that

$$x_i^* \in \text{supp } \bar{\rho} \quad \text{for all } i = 0, 1, \dots, n \quad (49)$$

for each value of n used in our simulations.

Then, instead of using the norm in $H^{-(1-a)/2}$, we use the equivalent norm $\|\cdot\|_V$ to make the computation easier. Indeed, since by (49) and Lemma 3.4(iv) the inequality in (11) becomes an equality, we obtain that

$$e_n := \|\bar{\rho} - \varphi^*\|_V^2 = 2(E(\varphi^*) - E(\bar{\rho})). \quad (50)$$

Now, $E(\bar{\rho})$ can be computed explicitly given that $V_{\text{reg}} = 0$ and U is a polynomial. To compute $E(\varphi^*)$, we set $\mathbf{x} := \mathbf{x}^*$ and $\ell_i := x_i - x_{i-1}$, and obtain from (42) that

$$E(\varphi^*) = \frac{1}{2n^2} \sum_{i=1}^n \sum_{j=1}^n \frac{1}{\ell_i \ell_j} \int_{x_{i-1}}^{x_i} \int_{x_{j-1}}^{x_j} V_a(x - y) dy dx + \frac{1}{n} \sum_{i=1}^n \frac{1}{\ell_i} \int_{x_{i-1}}^{x_i} U(x) dx. \quad (51)$$

Since U is a polynomial, both integrals above can be computed explicitly as a function of \mathbf{x} . Hence, once \mathbf{x}^* is computed numerically, e_n can be computed without any further numerical error (except for machine precision).

Finally, to compare the numerically computed values for e_n with Theorem 1.2, we make the ansatz

$$e_n = Cn^{-p}.$$

Then, $e_n/e_{2n} = 2^p$, and thus

$$p = \frac{\log e_n - \log e_{2n}}{\log 2}. \quad (52)$$

Hence, by taking n as subsequent powers of 2, we can compute p for each pair of subsequent values of e_n , and compare the values of p with the theoretically obtained power $1 - a$.

Case 1: the bounded domain $D(U) = [0, 1]$. We take $D(U) = [0, 1]$ and $U = 0$ on $[0, 1]$. Following the computations in, e.g., [KvM19], we obtain

$$\bar{\rho}(x) = \begin{cases} \frac{1}{\pi} [x(1-x)]^{-\frac{1}{2}} & \text{if } a = 0 \\ \frac{a\Gamma(a)}{\Gamma(\frac{1+a}{2})^2} [x(1-x)]^{-\frac{1-a}{2}} & \text{if } 0 < a < 1 \end{cases}$$

and

$$E(\bar{\rho}) = \begin{cases} \log 2 & \text{if } a = 0 \\ \frac{\pi a \Gamma(a)}{2\Gamma(\frac{1+a}{2})^2 \cos(\frac{a\pi}{2})} & \text{if } 0 < a < 1, \end{cases}$$

where $\Gamma(\alpha) = \int_0^\infty x^{\alpha-1} e^{-x} dx$ is the usual Γ -function.

With $E(\bar{\rho})$ specified, we compute e_n and p in (50) and (52) with the method described above. The results are shown in Table 3 and Figures 5 and 6. We note that $-p$ is the slope of the graphs of e_n in Figure 5. For all four values of a , e_n seems to converge to 0 as $n \rightarrow \infty$. Also, p decreases as a increases. These observations are in line with Theorem 1.2. However, for all four values of a , the computed value of p is significantly larger than the theoretical prediction $1 - a$ from Theorem 1.2.




Figure 5: The numerically computed values for e_n (see (50)) in Cases 1 and 2 for the values $a = 0$ (\bullet), $a = \frac{1}{4}$ (\blacksquare), $a = \frac{1}{2}$ (\blacktriangledown) and $a = \frac{3}{4}$ (\blacktriangle).




Figure 6: Values of p as a function of a in Cases 1 and 2 compared with the theoretical prediction from Theorem 1.2. The n -dependence is removed by taking the average of p over the last four values of n in Table 3.

Case 2: the infinite domain $D(U) = \mathbb{R}$. We take $D(U) = \mathbb{R}$ and

$$U(x) = \gamma_a \left(x - \frac{1}{2}\right)^2, \quad \gamma_a := \begin{cases} 4 & \text{if } a = 0 \\ \frac{2\pi a^2(2+a)\Gamma(a)}{\Gamma(\frac{1+a}{2})^2 \cos(\frac{a\pi}{2})} & \text{if } 0 < a < 1. \end{cases}$$

The constant γ_a is chosen such that $\text{supp } \bar{\rho} = [0, 1]$. Following the computations in, e.g., [KvM19] and [ST97, Chap. IV, Thm. 5.1], we obtain

$$\bar{\rho}(x) = \begin{cases} \frac{8}{\pi} [x(1-x)]^{\frac{1}{2}} & \text{if } a = 0 \\ \frac{4(2+a)a\Gamma(a)}{(1+a)\Gamma(\frac{1+a}{2})^2} [x(1-x)]^{\frac{1+a}{2}} & \text{if } 0 < a < 1 \end{cases}$$

and

$$E(\bar{\rho}) = \begin{cases} \log 2 & \text{if } a = 0 \\ \frac{\pi(2+a)^2 a \Gamma(a)}{2(4+a)\Gamma(\frac{1+a}{2})^2 \cos(\frac{a\pi}{2})} & \text{if } 0 < a < 1. \end{cases}$$

Similar to Case 1, we compute e_n and p . The results are shown in Figure 5 and Table 3. The similarities with Case 1 are that e_n seems to converge to 0 as $n \rightarrow \infty$, that p decreases as a increases, and that the computed value of p is significantly larger than the theoretical prediction $1 - a$ from Theorem 1.2.

We end this section with three quantitative comparisons between Cases 1 and 2:

- For $a = 0$, the values of e_n and p are similar.
- When a increases, the values of e_n are larger in Case 2 than in Case 1 (at least when n is not too large). This is consistent with Figure 2, where the graph of φ^* seems a better match with the graph of $\bar{\rho}$ in Case 1 than in Case 2.
- Yet, the values of p are larger in Case 2, which would imply that for n large enough, the values of e_n in Case 2 are smaller than those in Case 1. A possible reason for this could be the singularities of $\bar{\rho}$ at $x = 0$ and $x = 1$ in Case 1.

Values of p in Case 1					Values of p in Case 2				
n	$a = 0$	$a = \frac{1}{4}$	$a = \frac{1}{2}$	$a = \frac{3}{4}$	n	$a = 0$	$a = \frac{1}{4}$	$a = \frac{1}{2}$	$a = \frac{3}{4}$
2^2	1.60	1.42	1.24	0.23	2^2	1.40	1.26	1.07	0.83
2^3	1.72	1.54	1.17	0.02	2^3	1.52	1.36	1.13	0.82
2^4	1.78	1.59	0.98	0.10	2^4	1.61	1.43	1.16	0.79
2^5	1.82	1.60	0.80	0.24	2^5	1.68	1.48	1.17	0.75
2^6	1.84	1.57	0.72	0.35	2^6	1.73	1.52	1.17	0.70
2^7	1.86	1.52	0.72	0.43	2^7	1.77	1.55	1.15	0.66
2^8	1.87	1.47	0.75	0.48	2^8	1.80	1.56	1.14	0.62
2^9	1.88	1.41	0.79	0.52	2^9	1.82	1.57	1.12	0.59
2^{10}	1.89	1.37	0.83	0.54	2^{10}	1.84	1.58	1.09	0.57
2^{11}	1.90	1.34	0.86	0.55	2^{11}	1.86	1.59	1.07	0.55
2^{12}	1.88	1.33	0.89	0.56	2^{12}	1.87	1.59	1.06	0.54
2^{13}	2.07	1.33	0.91	0.56	2^{13}	1.88	1.58	1.04	0.53
Thm.	≈ 1	0.75	0.50	0.25	Thm.	≈ 1	0.75	0.50	0.25

Table 3: The numerically computed values for p (see (52)) in Cases 1 and 2. The bottom row is the prediction from Theorem 1.2. In Case 1 with $a = 0$ the last two values of p seem off; we expect that this is due to the numerical rounding errors that were made when (51) was computed.

Acknowledgments MK gratefully acknowledge support from JSPS KAKENHI Grant Number 17H02857.

PvM gratefully acknowledges support from the International Research Fellowship of the Japanese Society for the Promotion of Science and the associated JSPS KAKENHI Grant Number 15F15019.

References

- [CSS08] L. A. Caffarelli, S. Salsa, and L. Silvestre. Regularity estimates for the solution and the free boundary of the obstacle problem for the fractional Laplacian. *Inventiones Mathematicae*, 171(2):425–461, 2008.
- [GPPS13] M. G. D. Geers, R. H. J. Peerlings, M. A. Peletier, and L. Scardia. Asymptotic behaviour of a pile-up of infinite walls of edge dislocations. *Archive for Rational Mechanics and Analysis*, 209:495–539, 2013.
- [GR07] I. S. Gradshteyn and I. M. Ryzhik. *Table of Integrals, Series, and Products, Seventh Edition*. Academic Press, Burlington, Massachusetts, 2007.
- [GvMPS16] A. Garroni, P. van Meurs, M. A. Peletier, and L. Scardia. Boundary-layer analysis of a pile-up of walls of edge dislocations at a lock. *Mathematical Models and Methods in Applied Sciences*, 26(14):2735–2768, 2016.
- [Hal11] C. L. Hall. Asymptotic analysis of a pile-up of regular edge dislocation walls. *Materials Science and Engineering: A*, 530:144–148, 2011.

- [HCO10] C. L. Hall, S. J. Chapman, and J. R. Ockendon. Asymptotic analysis of a system of algebraic equations arising in dislocation theory. *SIAM Journal on Applied Mathematics*, 70(7):2729–2749, 2010.
- [HHvM18] C. L. Hall, T. Hudson, and P. van Meurs. Asymptotic analysis of boundary layers in a repulsive particle system. *Acta Applicandae Mathematicae*, 153(1):1–54, 2018.
- [HT19] S. Hayakawa and K. Tanaka. Convergence analysis of approximation formulas for analytic functions via duality for potential energy minimization. *ArXiv: 1906.03133*, 2019.
- [HvMP20] T. Hudson, P. van Meurs, and M. A. Peletier. Atomistic origins of continuum dislocation dynamics. *ArXiv: 2001.06120*, 2020.
- [KvM19] M. Kimura and P. van Meurs. Regularity of the minimiser of one-dimensional interaction energies. *ESAIM: Control, Optimisation and Calculus of Variations*, 2019. to appear.
- [PS17] M. Petrache and S. Serfaty. Next order asymptotics and renormalized energy for Riesz interactions. *Journal of the Institute of Mathematics of Jussieu*, 16(3):501–569, 2017.
- [SS15a] E. Sandier and S. Serfaty. 1D log gases and the renormalized energy: crystallization at vanishing temperature. *Probability Theory and Related Fields*, 162(3-4):795–846, 2015.
- [SS15b] E. Sandier and S. Serfaty. 2D Coulomb gases and the renormalized energy. *The Annals of Probability*, 43(4):2026–2083, 2015.
- [ST97] E. B. Saff and V. Totik. *Logarithmic Potentials with External Fields*. Springer Verlag Berlin Heidelberg, 1997.
- [TS19] K. Tanaka and M. Sugihara. Design of accurate formulas for approximating functions in weighted Hardy spaces by discrete energy minimization. *IMA Journal of Numerical Analysis*, 39(4):1957–1984, 2019.
- [vM18a] P. van Meurs. Convergence rates for discrete-to-continuum limits in 1D particle systems. In *Mathematical Analysis of Continuum Mechanics and Industrial Applications II: Proceedings of the International Conference CoMFoS16*, pages 181–193. Springer, 2018.
- [vM18b] P. van Meurs. Many-particle limits and non-convergence of dislocation wall pile-ups. *Nonlinearity*, 31:165–225, 2018.
- [vMMP14] P. van Meurs, A. Muntean, and M. A. Peletier. Upscaling of dislocation walls in finite domains. *European Journal of Applied Mathematics*, 25(6):749–781, 2014.