

Entanglement transfer from quantum matter to classical geometry in an emergent holographic dual description of a scalar field theory

Ki-Seok Kim^{a,b} and Shinsei Ryu^c

^a*Department of Physics, POSTECH, Pohang, Gyeongbuk 37673, Korea*

^b*Asia Pacific Center for Theoretical Physics (APCTP), Pohang, Gyeongbuk 37673, Korea*

^c*Kadanoff Center for Theoretical Physics, University of Chicago, IL 60637, USA*

ABSTRACT: Applying recursive renormalization group transformations to a scalar field theory, we obtain an effective quantum gravity theory with an emergent extra dimension, described by a dual holographic Einstein-Klein-Gordon action. In the large N limit, where N is the flavor number of the original scalar fields, quantum fluctuations of both dynamical metric and dual scalar fields are suppressed, leading to a classical field theory of the Einstein-scalar type in $(D + 1)$ -spacetime dimensions. We show that this emergent background gravity describes the renormalization group flows of coupling functions in the UV quantum field theory through the extra dimension. More precisely, the IR boundary conditions of the Einstein equations, given by the Gibbons-Hawking-York action, correspond to the renormalization group β -functions of the quantum field theory, where the infinitesimal distance in the extra-dimensional space is identified with an energy scale for the renormalization group transformation. Finally, we also show that this dual holographic formulation describes quantum entanglement in a geometrical way, encoding the transfer of quantum entanglement from quantum matter to classical gravity in the large N limit. We claim that this entanglement transfer serves as a microscopic foundation for the emergent holographic duality description.

Contents

1	Introduction	1
2	Overview	3
2.1	Emergent gravity description for a scalar field theory	3
2.2	Transferring quantum entanglement from matter to geometry	6
3	Derivation of the holographic Landau-Ginzburg effective field theory	7
3.1	The first renormalization group transformation in real space a la Polchinski	7
3.2	Recursive renormalization group transformations a la Sung-Sik Lee	12
3.3	Remarks on the renormalization group transformation for the metric tensor	15
4	Physical interpretation of the emergent dynamical metric tensor: From renormalization group equations of coupling parameters to the evolution equation of the metric tensor	20
5	Evaluation of entanglement entropy based on the heat kernel method	23
5.1	Entanglement entropy	23
5.2	Heat kernel method for entanglement entropy	26
5.3	Invariance of entanglement entropy with respect to renormalization group transformations	28
6	Discussion	29
7	Conclusion	31
A	Emergent geometric description for a scalar lattice field theory in one dimension	32
B	Further developments for the invariance of entanglement entropy with respect to renormalization group transformations	39
B.1	An identity for entanglement entropy at $z = z_f$	39
B.2	Entanglement entropy after the first iteration	41

1 Introduction

Renormalization group transformations lead us to construct an effective field theory of renormalized low-energy fluctuations in terms of renormalized interaction vertices, which result from quantum fluctuations of high-energy modes [1]. This renormalization group

analysis requires that the resulting effective field theory remains invariant in its mathematical form after the renormalization group transformation. As a result, we can extract out how interaction vertices evolve as a function of the renormalization group energy scale, referred to as renormalization group β -functions. In addition, we obtain the so-called Callan-Symanzik equation that a correlation function has to satisfy in the renormalization group transformation [1]. Solving this renormalization group equation for the correlation function, we find a renormalized correlation function in terms of transfer momentum and energy, which encodes anomalous scaling dimensions of renormalized low-energy fluctuations and renormalization group β -functions of interaction vertices.

One may suggest that the renormalization group energy scale is introduced into an effective field theory as an extra dimension, manifesting a renormalization group flow of the effective field theory through the extra dimensional space. Renormalization group flows of interaction vertices are realized along the extra dimension, serving as an effective curved spacetime for renormalized low-energy fluctuations in this effective field theory. Renormalization group flows of correlation functions can be described by introducing external source fields, coupled to conserved currents minimally, into the effective field theory, regarded to be a dual description in this emergent spacetime with an extra dimension. This novel construction for renormalization group transformations reminds us of the holographic duality conjecture [2–8].

In this study, we follow this line of thoughts and suggest a microscopic foundation for the holographic duality conjecture [9–29]. The key idea is the recursive applications of the renormalization group transformations. We follow the real space renormalization group scheme developed by Polchinski (although whether one works in real or momentum space should not matter ultimately). This renormalization group technique was further developed by Sung-Sik Lee, by applying the renormalization group transformations in a recursive way [31–33]. Resorting to this renormalization group analysis, we show that renormalization group flows of interaction vertices appear to form an emergent curved spacetime with an extra dimension. Here, the infinitesimal distance in this extra-dimensional space is identified with the renormalization group energy scale. To be concrete, we apply the Polchinski renormalization group technique a la Sung-Sik Lee to a quantum field theory of bosons with self-interactions in D -spacetime dimensions. We derive an effective dual holographic description of the Einstein-Klein-Gordon theory in $(D + 1)$ -spacetime dimensions, given by a quantum gravity theory of the Einstein-Hilbert action coupled to fluctuating scalar fields dual to the density of the original bosons. It turns out that quantum fluctuations of both dynamical metric and dual scalar fields are suppressed in the large N limit, where N is the number of boson flavors. As a result, we obtain a classical field theory of the Einstein-scalar type in $(D + 1)$ -spacetime dimensions.

Since the quantum field theory is geometrized and described by a classical field theory, it is natural to ask how the quantum mechanical nature of the original theory can be encoded into the classical and geometric description appropriately. We thus study the entanglement entropy [34–39], which measures the quantum correlation between a given spatial region and its complement, by utilizing the classical holographic description. As we will show, it turns out that entanglement is transferred from quantum matter to classical gravity through

recursive renormalization group transformations. We claim that the entanglement transfer serves as a microscopic foundation for the holographic duality conjecture.

2 Overview

2.1 Emergent gravity description for a scalar field theory

Since the present study involves a lot of formulae, discussing our main results briefly with the introduction of the structure of this paper would be helpful for readers to figure out the physics. We start from an effective field theory of self-interacting bosons, where the partition function is

$$Z = \int D\phi_\alpha(x) \exp \left\{ -S_{UV}[\phi_\alpha(x); g_{\mu\nu}^B(x)] \right\} \quad (2.1)$$

and the effective action is

$$\begin{aligned} & S_{UV}[\phi_\alpha(x); g_{\mu\nu}^B(x)] \\ &= \int d^D x \sqrt{g_B} \left\{ g_B^{\mu\nu} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + m^2 \phi_\alpha^2 + \xi R_B \phi_\alpha^2 + \frac{u}{2N} \phi_\alpha^2 \phi_\beta^2 + \frac{\lambda}{2N} T_{\mu\nu} T^{\mu\nu} \right\}. \end{aligned} \quad (2.2)$$

Here, $\phi_\alpha(x)$ is a real scalar field with a flavor index $\alpha = 1, \dots, N$ at a D -dimensional spacetime coordinate x . We point out that the two flavor indices α and β are summed independently. $g_B^{\mu\nu}$ with μ & $\nu = 0, \dots, D-1$ is a background metric tensor to describe a D -dimensional curved spacetime manifold, where the bosonic field lives. g_B is the determinant of the metric tensor, where $d^D x \sqrt{g_B}$ gives an invariant volume factor. For example, we may consider the background metric as follows

$$ds^2 = dr^2 + r^2 d\theta^2 + \delta_{ij} dx_\perp^i dx_\perp^j, \quad g_{rr}^B = 1, \quad g_{\theta\theta}^B = r^2, \quad g_{ij}^{B\perp} = \delta_{ij}, \quad \sqrt{g_B} = r, \quad (2.3)$$

which will be useful for evaluation of the entanglement entropy using the replica technique. Here, i and j run from 2 to $D-1$. m denotes the mass of these scalar bosons and u represents their self-interactions. R_B is the Ricci scalar and ξ is a coupling constant between scalar fields and the background curvature [40]. Finally, the last term describes effective interactions between energy-momentum tensor currents $T^{\mu\nu}$, where λ is the corresponding coupling constant. Although these effective interactions are irrelevant at both Gaussian and Wilson-Fisher fixed points in the perturbative regime [1], we introduce this term explicitly, which plays an important role in uplifting the background geometry into full dynamical degrees of freedom for the emergent bulk geometry [41].

In Section 3, by implementing the Polchinski real-space renormalization group technique [30] to this effective field theory in a recursive way a la Sung-Sik Lee [31–33], we obtain a quantum gravity theory coupled to a dual scalar field in one-dimensional higher spacetime. It is given by the following partition function

$$\begin{aligned} Z = Z_\Lambda \int D\varphi(x, z) Dg_{\mu\nu}(x, z) \exp \left\{ -S_{UV}[g_{\mu\nu}(x, 0), \varphi(x, 0)] - S_{IR}[g_{\mu\nu}(x, z_f), \varphi(x, z_f)] \right. \\ \left. - S_{Bulk}[g_{\mu\nu}(x, z), \varphi(x, z)] \right\}. \end{aligned} \quad (2.4)$$

Here, z is the coordinate of an emergent extra dimension, identified with a renormalization group scale as discussed in the introduction. $g_{\mu\nu}(x, z)$ with μ & $\nu = 0, \dots, D-1$ is an emergent dynamical metric field, where metric tensors involved with this extra dimension are gauge-fixed as $g_{DD}(x, z) = 1$ and $g_{\mu D}(x, z) = 0$. $\varphi(x, z)$ is a dual scalar field, conventionally taken to be $\varphi(x) = \frac{1}{N} \left\langle \sum_{\alpha=1}^N [\phi_{\alpha}(x)]^2 \right\rangle$ in the large N analysis of the quantum field theory [42], where $\langle \hat{O} \rangle$ is the expectation value (ensemble average) of the operator \hat{O} at the Gaussian fixed point.

The dynamics of both emergent metric and dual scalar fields are governed by the following bulk effective action

$$\begin{aligned}
S_{Bulk}[g_{\mu\nu}(x, z), \varphi(x, z)] = N \int_0^{z_f} dz \int d^D x \sqrt{g(x, z)} \{ \\
+ \frac{1}{2u} [\partial_z \varphi(x, z)]^2 + \frac{C_{\varphi}}{2} g^{\mu\nu}(x, z) [\partial_{\mu} \varphi(x, z)] [\partial_{\nu} \varphi(x, z)] + C_{\xi} R(x, z) [\varphi(x, z)]^2 \\
- \frac{1}{2\lambda} \left(\partial_z g^{\mu\nu}(x, z) - \frac{1}{2dz} g^{\mu\nu'}(x, z) (\partial_{\nu'} \partial_{\mu'} G_{xx'}[g_{\mu\nu}(x, z), \varphi(x, z)])_{x' \rightarrow x} g^{\mu'\nu}(x, z) \right)^2 \\
+ \frac{1}{2\kappa} (R(x, z) - 2\Lambda) \}. \tag{2.5}
\end{aligned}$$

Here, the dynamics of $\varphi(x, z)$ is given by rather a conventional form of the bosonic effective action on a curved spacetime manifold [40]. C_{φ} and C_{ξ} are positive constants, which decrease as the mass of the original scalar fields increases. The metric-tensor field is described essentially by the Einstein-Hilbert type action in a gauge-fixed version, which originates from quantum fluctuations of matter fields. This emergent Einstein-Hilbert action is analogous to the notion of induced gravity [43, 44]. We also point out that the dynamics of the metric along the extra dimensional space is modified through $G_{xx'}[g_{\mu\nu}(x, z), \varphi(x, z)]$, which is the Green's function given by quantum fluctuations of heavy-mass scalar fields in the renormalization group transformation,

$$\begin{aligned}
\left\{ - \frac{1}{\sqrt{g(x, z)}} \partial_{\mu} \left(\sqrt{g(x, z)} g^{\mu\nu}(x, z) \partial_{\nu} \right) + \frac{1}{2dz} [m^2 - i\varphi(x, z)] \right\} G_{xx'}[g_{\mu\nu}(x, z), \varphi(x, z)] \\
= \frac{1}{\sqrt{g(x, z)}} \delta^{(D)}(x - x'), \tag{2.6}
\end{aligned}$$

where dz is an infinitesimal parameter of the renormalization group transformation. The evolution dynamics of both metric and dual scalar fields are purely determined by this Green's function in addition to the emergent local symmetry in the dynamical bulk geometry, i.e., $(D+1)$ -dimensional diffeomorphism invariance [31–33], although it is explicitly broken by the gauge fixing. As clearly seen in this expression, quantum fluctuations of both metric and dual scalar fields are suppressed in the large N limit. As a result, both dynamics are described by coupled classical equations of motion in the presence of the emergent extra dimensional space.

Since these coupled equations of motion are second order in derivatives with respect to the extra dimensional space, we need two boundary conditions. The first boundary

condition is described by an effective UV action in Eq. (2.4), given by,

$$S_{UV}[g_{\mu\nu}(x, 0), \varphi(x, 0)] = N \int d^D x \sqrt{g(x, 0)} \times \left\{ \frac{1}{2u} \left(\varphi(x, 0) - i\xi R(x, 0) \right)^2 - \frac{1}{2\lambda} \left(g^{\mu\nu}(x, 0) - g_B^{\mu\nu}(x) \right)^2 \right\}. \quad (2.7)$$

Of course, an actual equation corresponding to this UV boundary condition has to take into account a linear z derivative term at $z = 0$, which originates from the above bulk effective action by the technique of integration-by-parts. For the gravity sector, this boundary term is referred to as a Gibbons-Hawking-York term [45, 46]. The other boundary condition is provided by an effective IR action

$$S_{IR}[g_{\mu\nu}(x, z_f), \varphi(x, z_f)] = N \int d^D x \sqrt{g(x, z_f)} \left\{ \frac{\mathcal{C}_\varphi^f}{2} g^{\mu\nu}(x, z_f) [\partial_\mu \varphi(x, z_f)] [\partial_\nu \varphi(x, z_f)] + \mathcal{C}_\xi^f R(x, z_f) [\varphi(x, z_f)]^2 + \frac{1}{2\kappa_f} \left(R(x, z_f) - 2\Lambda_f \right) \right\}, \quad (2.8)$$

where the original scalar fields have been integrated out, using the gradient expansion, to give this effective action for the dual scalar fields and metric tensors [40, 43, 44] at $z = z_f$. Here, \mathcal{C}_φ^f and \mathcal{C}_ξ^f are positive coefficients, which also decrease as the mass m increases. The effective gravitational constant κ_f and the cosmological parameter Λ_f are similarly determined from the gradient expansion. Also, actual equations to describe the IR boundary condition have to be supported by the linear z -derivative term at $z = z_f$. All these boundary conditions including the Gibbons-Hawking-York term are self-consistently determined by these effective actions. We show all derivations for this emergent holographic construction in section 3.

z_f in Eqs. (2.5) and (2.8) is an IR cutoff, which may be regarded to be inversely proportional to temperature in the renormalization group transformation. This implies that the IR boundary conditions correspond to renormalization group equations. More precisely, as we discuss in Section 4 and appendix A, the IR boundary equations for metric tensors correspond to renormalization group β -functions of interaction vertices. On the other hand, the IR boundary conditions of dual scalar fields does to a dual expression of the Callan-Symanzik equation.

Finally, we point out that the limit $z_f \rightarrow 0$ leads to the conventional large N analysis as follows

$$S_{UV}[g_{\mu\nu}(x, 0), \varphi(x, 0)] + \lim_{z_f \rightarrow 0} S_{IR}[g_{\mu\nu}(x, z_f), \varphi(x, z_f)] = S_{UV}[g_B^{\mu\nu}(x), \varphi(x)], \quad (2.9)$$

where the bulk effective action vanishes in the left-hand-side of the equality and the original scalar fields are integrated out to give an effective action in the large N limit for the right-hand-side of the equality.

2.2 Transferring quantum entanglement from matter to geometry

To figure out how the quantum-entanglement structure can be translated into the emergent classical geometry, we calculate in Section 5, the entanglement entropy [34–39]. Entanglement entropy may be regarded as an entropy of a subsystem, given by the von-Neumann entropy of the reduced density matrix for the subsystem. Here, we perform a field-theory calculation of the entanglement entropy, based on the replica technique in a Riemann surface with a conical singularity. It turns out that the entanglement entropy at the IR cutoff $z = z_f$ can be decomposed into $\mathcal{S}_{EE}^{\phi_\alpha}(z_f)$ of the matter and $\mathcal{S}_{EE}^{GR}(z_f)$ of the emergent gravity as follows

$$\mathcal{S}_{EE}(z_f) = \mathcal{S}_{EE}^{\phi_\alpha}(z_f) + \mathcal{S}_{EE}^{GR}(z_f). \quad (2.10)$$

Resorting to the heat-kernel method, one finds that the entanglement entropy of free bosons with the flavor degeneracy N follows an area law, given by [47]

$$\mathcal{S}_{EE}^{\phi_\alpha}(z_f) = \frac{N}{6(D-2)(4\pi)^{\frac{D}{2}-1}} \frac{\mathcal{A}[\Sigma(z_f)]}{\epsilon^{D-2}}, \quad (2.11)$$

where

$$\mathcal{A}[\Sigma(z_f)] = \int_{\Sigma(z_f)} d^{D-2}x_\perp \sqrt{\gamma(x, z_f)} \quad (2.12)$$

is an area of the subsystem at $z = z_f$. ϵ is a UV cutoff, here corresponding to a microscopic scale for mapping from a discrete version of the holographic Einstein-Klein-Gordon theory to the above continuum field theory Eq. (2.5), as we will discuss in detail in Section 3. On the other hand, the gravity contribution to the entanglement entropy is given by

$$\begin{aligned} \mathcal{S}_{EE}^{GR}(z_f) &= N \int d^{D-2}x_\perp \int_0^{2\pi} d\theta \int_0^\infty dr \int_0^{z_f} dz \sqrt{g(r, x_\perp, z)} T_{\mu\nu}^{Bulk}(r, x_\perp, z) \frac{\partial g_n^{\mu\nu}(r, x_\perp, z)}{\partial n} \Big|_{n=1} \\ &\quad - N \int d^{D-2}x_\perp \int_0^{2\pi} d\theta \int_0^\infty dr \int_0^{z_f} dz \sqrt{g(r, x_\perp, z)} \left\{ \frac{1}{2u} [\partial_z \varphi(r, x_\perp, z)]^2 \right. \\ &\quad + \frac{\mathcal{C}_\varphi}{2} g^{\mu\nu}(r, x_\perp, z) [\partial_\mu \varphi(r, x_\perp, z)] [\partial_\nu \varphi(r, x_\perp, z)] + \mathcal{C}_\xi R(r, x_\perp, z) [\varphi(r, x_\perp, z)]^2 \\ &\quad - \frac{1}{2\lambda} \left(\partial_z g^{\mu\nu}(r, x_\perp, z) - \frac{1}{2dz} g^{\mu\nu'}(r, x_\perp, z) (\partial_{\nu'} \partial_{\mu'} G_{xx'}[g_{\mu\nu}(r, x_\perp, z), \varphi(r, x_\perp, z)])_{x' \rightarrow x} g^{\mu'\nu'}(r, x_\perp, z) \right)^2 \\ &\quad \left. + \frac{1}{2\kappa} (R(r, x_\perp, z) - 2\Lambda) \right\}. \end{aligned} \quad (2.13)$$

Here,

$$T_{\mu\nu}^{Bulk}(r, x_\perp, z) = \frac{1}{\sqrt{g(r, x_\perp, z)}} \frac{\partial \sqrt{g(r, x_\perp, z)} \mathcal{L}_{Bulk}[g_{\mu\nu}(r, x_\perp, z), \varphi(r, x_\perp, z)]}{\partial g^{\mu\nu}(r, x_\perp, z)} \quad (2.14)$$

is an energy-momentum tensor of the $(D+1)$ -dimensional Einstein gravity in Eq. (2.5), given by $S_{Bulk}[g_{\mu\nu}(x, z), \varphi(x, z)] = \int_0^{z_f} dz \int d^Dx \sqrt{g(x, z)} \mathcal{L}_{Bulk}[g_{\mu\nu}(x, z), \varphi(x, z)]$. The quantum average of $\langle T_{\mu\nu}^{Bulk}(r, x_\perp, z) \rangle$ with respect to the quantum gravity action is reduced into its

classical value of $T_{\mu\nu}^g(r, x_\perp, z)$ in the large N limit. n is the replica index to describe the conical singularity of the geometry [47]. The detailed derivation is given in Section 5.

An important observation is that the entanglement entropy does not depend on the IR cutoff z_f at all. In other words, it is a renormalization-group invariant, regardless of z_f . This originates from that fact that the partition function is a renormalization-group invariant. As a result, we find the renormalization-group flow equation for the entanglement entropy, given by

$$\partial_{z_f} \mathcal{S}_{EE}(z_f) = \partial_{z_f} \mathcal{S}_{EE}^{\phi_\alpha}(z_f) + \partial_{z_f} \mathcal{S}_{EE}^{GR}(z_f) = 0. \quad (2.15)$$

This observation leads us to conclude that the entanglement entropy of the matter sector at UV is transferred into that of the classical bulk gravity part at IR. (While we could not verify this relation based on explicit calculations for the general formula, we could check it after the first renormalization group transformation, as discussed in detail in appendix B.) Recalling that an emergent geometry of gapped quantum fields is essentially a cap geometry, where the spacetime does not exist beyond some scale in the extra dimension [33, 48, 49], the entanglement entropy of the matter sector vanishes after the cap geometry in the extra dimensional space. Our conclusion on the entanglement transfer is consistent with a recent study [50].

3 Derivation of the holographic Landau-Ginzburg effective field theory

3.1 The first renormalization group transformation in real space a la Polchinski

We start from an effective scalar field theory on a D -dimensional curved spacetime described by a background metric tensor $g_B^{\mu\nu}$. The partition function is given by [40]

$$Z = \int D\phi_\alpha Dg_{\mu\nu}^{(0)} DT_{\mu\nu}^{(0)} \exp \left[- \int d^D x \sqrt{g^{(0)}} \left\{ g^{\mu\nu(0)} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + m^2 \phi_\alpha^2 + \xi R^{(0)} \phi_\alpha^2 + \frac{u}{2N} \phi_\alpha^2 \phi_\beta^2 - NT_{\mu\nu}^{(0)} (g^{\mu\nu(0)} - g_B^{\mu\nu}) \right\} \right]. \quad (3.1)$$

Here, $T_{\mu\nu}^{(0)}$ is a Lagrange multiplier field to impose the constraint $g^{\mu\nu(0)} = g_B^{\mu\nu}$. For the time being, we do not consider the effective interaction between energy-momentum tensor currents, for simplicity in the presentation of derivations. To deal with self-interactions between bosons, we introduce a dual scalar field $\varphi^{(0)}$ as

$$Z = \int D\phi_\alpha D\varphi^{(0)} Dg_{\mu\nu}^{(0)} DT_{\mu\nu}^{(0)} \exp \left[- \int d^D x \sqrt{g^{(0)}} \left\{ g^{\mu\nu(0)} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + m^2 \phi_\alpha^2 + \xi R^{(0)} \phi_\alpha^2 - i\varphi^{(0)} \phi_\alpha^2 + \frac{N}{2u} \varphi^{(0)2} - NT_{\mu\nu}^{(0)} (g^{\mu\nu(0)} - g_B^{\mu\nu}) \right\} \right] \quad (3.2)$$

by the Hubbard-Stratonovich transformation.

To implement the renormalization group transformation for the dynamics of dual scalar fields, we introduce an auxiliary field $\eta^{(0)}$ in the following way:

$$Z = \int D\phi_\alpha D\varphi^{(0)} D\eta^{(0)} Dg_{\mu\nu}^{(0)} DT_{\mu\nu}^{(0)} \exp \left[- \int d^D x \sqrt{g^{(0)}} \left\{ g^{\mu\nu(0)} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + m^2 \phi_\alpha^2 + \xi R^{(0)} \phi_\alpha^2 - i\varphi^{(0)} \phi_\alpha^2 + \frac{N}{2u} \varphi^{(0)2} + \frac{N}{2u_\eta} \eta^{(0)2} - NT_{\mu\nu}^{(0)} (g^{\mu\nu(0)} - g_B^{\mu\nu}) \right\} \right]. \quad (3.3)$$

We emphasize that the introduction of this auxiliary field does not change any physics except for the normalization constant of the partition function, omitted here for notational simplicity.

Now, we separate low-energy and high-energy dual scalar fields, $\varphi^{(0)}$ and $\chi^{(0)}$, respectively, as follows

$$\varphi^{(0)} \implies \varphi^{(0)} + \chi^{(0)}, \quad \eta^{(0)} \implies c_\varphi^{(0)} \varphi^{(0)} + c_\chi^{(0)} \chi^{(0)}. \quad (3.4)$$

Here, we determine two coefficients of $c_\varphi^{(0)}$ and $c_\chi^{(0)}$ as

$$c_\varphi^{(0)} = \frac{u^{-1}}{\mu^{(0)} u_\eta^{-1/2}}, \quad c_\chi^{(0)} = -\frac{\mu^{(0)}}{u_\eta^{-1/2}}, \quad \mu^{(0)} = \frac{u^{-1/2}}{\sqrt{e^{2\beta^{(0)} dz} - 1}}, \quad (3.5)$$

such that we do not generate mixing terms between such light and heavy dual scalar fields in the mass sector. $\mu^{(0)}$ is an effective mass for the heavy dual scalar field, clarified soon. $\beta^{(0)}$ represents a local speed of coarse graining for the dual scalar field, and dz is an infinitesimal parameter for the renormalization group transformation, also to be clarified below. As a result, we rewrite the mass sector in terms of these light and heavy dual boson fields as follows

$$\frac{N}{2u} \varphi^{(0)2} + \frac{N}{2u_\eta} \eta^{(0)2} \implies \frac{N}{2u} e^{2\beta^{(0)} dz} \varphi^{(0)2} + \frac{N}{2u} \frac{e^{2\beta^{(0)} dz}}{e^{2\beta^{(0)} dz} - 1} \chi^{(0)2}. \quad (3.6)$$

Now, it is clear why we call $\chi^{(0)}$ a heavy dual scalar field, whose mass is given by $\mu^{(0)}$. Rescaling both dual scalar fields in the following way to return the mass sector to its original form,

$$\varphi^{(0)} \implies e^{-\beta^{(0)} dz} \varphi^{(0)}, \quad \chi^{(0)} \implies e^{-\beta^{(0)} dz} \chi^{(0)}, \quad (3.7)$$

we rewrite the partition function in terms of these light and heavy dual scalar bosons as

$$Z = \int D\phi_\alpha D\varphi^{(0)} D\chi^{(0)} Dg_{\mu\nu}^{(0)} DT_{\mu\nu}^{(0)} \exp \left[- \int d^D x \sqrt{g^{(0)}} \left\{ g^{\mu\nu(0)} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + m^2 \phi_\alpha^2 + \xi R^{(0)} \phi_\alpha^2 - ie^{-\beta^{(0)} dz} (\varphi^{(0)} + \chi^{(0)}) \phi_\alpha^2 + \frac{N}{2u} \varphi^{(0)2} + \frac{N}{2u} \frac{1}{e^{2\beta^{(0)} dz} - 1} \chi^{(0)2} - NT_{\mu\nu}^{(0)} (g^{\mu\nu(0)} - g_B^{\mu\nu}) \right\} \right]. \quad (3.8)$$

It is straightforward to perform the Gaussian integration for the heavy dual scalar field, resulting in the partition function

$$Z = \int D\phi_\alpha D\varphi^{(0)} Dg_{\mu\nu}^{(0)} DT_{\mu\nu}^{(0)} \exp \left[-\frac{1}{2} \text{tr}_{xx'} \ln \frac{N}{2u} \frac{1}{e^{2\beta^{(0)} dz} - 1} \right. \\ \left. - \int d^D x \sqrt{g^{(0)}} \left\{ g^{\mu\nu(0)} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + m^2 \phi_\alpha^2 + \xi R^{(0)} \phi_\alpha^2 - ie^{-\beta^{(0)} dz} \varphi^{(0)} \phi_\alpha^2 + \frac{N}{2u} \varphi^{(0)2} \right. \right. \\ \left. \left. + \frac{u}{2N} (1 - e^{-2\beta^{(0)} dz}) \phi_\alpha^2 \phi_\beta^2 - NT_{\mu\nu}^{(0)} (g^{\mu\nu(0)} - g_B^{\mu\nu}) \right\} \right], \quad (3.9)$$

where self-interactions between the original scalar bosons are generated, as expected. To prepare for the second renormalization group transformation, we perform the Hubbard-Stratonovich transformation once again for this newly generated self-interaction term,

$$Z = \int D\phi_\alpha D\varphi^{(0)} D\varphi^{(1)} Dg_{\mu\nu}^{(0)} DT_{\mu\nu}^{(0)} \exp \left[-\frac{1}{2} \text{tr}_{xx'} \ln \frac{N}{2u} \frac{1}{e^{2\beta^{(0)} dz} - 1} \right. \\ \left. - \int d^D x \sqrt{g^{(0)}} \left\{ g^{\mu\nu(0)} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + m^2 \phi_\alpha^2 + \xi R^{(0)} \phi_\alpha^2 - ie^{-\beta^{(0)} dz} \varphi^{(0)} \phi_\alpha^2 + \frac{N}{2u} \varphi^{(0)2} \right. \right. \\ \left. \left. + \frac{N}{2u} \frac{e^{2\beta^{(0)} dz}}{e^{2\beta^{(0)} dz} - 1} \varphi^{(1)2} - i\varphi^{(1)} \phi_\alpha^2 - NT_{\mu\nu}^{(0)} (g^{\mu\nu(0)} - g_B^{\mu\nu}) \right\} \right]. \quad (3.10)$$

Rescaling the dual scalar field as

$$\varphi^{(1)} \implies e^{-\beta^{(0)} dz} \varphi^{(1)} \quad (3.11)$$

and shifting the field as

$$\varphi^{(1)} \implies \varphi^{(1)} - \varphi^{(0)} \quad (3.12)$$

together with $\varphi^{(1)} \implies \varphi^{(1)} - ie^{-\beta^{(0)} dz} \xi R^{(0)}$ and $\varphi^{(0)} \implies \varphi^{(0)} - ie^{-\beta^{(0)} dz} \xi R^{(0)}$, we obtain the following expression of the partition function after the first renormalization group transformation for the dual scalar-field sector

$$Z = \int D\phi_\alpha D\varphi^{(0)} D\varphi^{(1)} Dg_{\mu\nu}^{(0)} DT_{\mu\nu}^{(0)} \exp \left[-\frac{1}{2} \text{tr}_{xx'} \ln \frac{N}{2u} \frac{1}{e^{2\beta^{(0)} dz} - 1} \right. \\ \left. - \int d^D x \sqrt{g^{(0)}} \left\{ g^{\mu\nu(0)} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + m^2 \phi_\alpha^2 - ie^{-\beta^{(0)} dz} \varphi^{(1)} \phi_\alpha^2 \right. \right. \\ \left. \left. + \frac{N}{2u} \left(\varphi^{(0)} - ie^{-\beta^{(0)} dz} \xi R^{(0)} \right)^2 + \frac{N}{2u} \frac{1}{e^{2\beta^{(0)} dz} - 1} (\varphi^{(1)} - \varphi^{(0)})^2 - NT_{\mu\nu}^{(0)} (g^{\mu\nu(0)} - g_B^{\mu\nu}) \right\} \right]. \quad (3.13)$$

Now, we perform the first renormalization group transformation for the original boson sector. As in the renormalization group transformation for the dual scalar-field sector, we introduce an auxiliary field ψ_α into the partition function

$$Z = \int D\phi_\alpha D\psi_\alpha D\varphi^{(0)} D\varphi^{(1)} Dg_{\mu\nu}^{(0)} DT_{\mu\nu}^{(0)} \exp \left[-\frac{1}{2} \text{tr}_{xx'} \ln \frac{N}{2u} \frac{1}{e^{2\beta^{(0)} dz} - 1} \right. \\ \left. - \int d^D x \sqrt{g^{(0)}} \left\{ g^{\mu\nu(0)} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + m^2 \phi_\alpha^2 - ie^{-\beta^{(0)} dz} \varphi^{(1)} \phi_\alpha^2 + M^2 \psi_\alpha^2 \right. \right. \\ \left. \left. + \frac{N}{2u} \left(\varphi^{(0)} - ie^{-\beta^{(0)} dz} \xi R^{(0)} \right)^2 + \frac{N}{2u} \frac{1}{e^{2\beta^{(0)} dz} - 1} (\varphi^{(1)} - \varphi^{(0)})^2 - NT_{\mu\nu}^{(0)} (g^{\mu\nu(0)} - g_B^{\mu\nu}) \right\} \right]. \quad (3.14)$$

We point out again that the introduction of the auxiliary field ψ_α does not change any physics except for the normalization constant of the partition function, omitted here for notational simplicity.

We separate low-energy and high-energy scalar fields, ϕ_α and Φ_α , respectively,

$$\phi_\alpha \implies \phi_\alpha + \Phi_\alpha, \quad \psi_\alpha \implies c_\phi^{(0)} \phi_\alpha + c_\Phi^{(0)} \Phi_\alpha, \quad (3.15)$$

where two coefficients of $c_\phi^{(0)}$ and $c_\Phi^{(0)}$ are given by

$$c_\phi^{(0)} = \frac{(m^2 - ie^{-\beta^{(0)}dz} \varphi^{(1)})}{\mu^{(0)} M}, \quad c_\Phi^{(0)} = -\frac{\mu^{(0)}}{M}, \quad \mu^{(0)} = \frac{(m^2 - ie^{-\beta^{(0)}dz} \varphi^{(1)})^{1/2}}{\sqrt{e^{2\alpha^{(0)}dz} - 1}}. \quad (3.16)$$

Here, as $\beta^{(0)}$, $\alpha^{(0)}$ represents a local speed of coarse graining for the dual scalar field. Then, the mass sector of the original scalar fields reads

$$\begin{aligned} & (m^2 - ie^{-\beta^{(0)}dz} \varphi^{(1)}) \phi_\alpha^2 + M^2 \psi_\alpha^2 \\ & \implies e^{2\alpha^{(0)}dz} (m^2 - ie^{-\beta^{(0)}dz} \varphi^{(1)}) \phi_\alpha^2 + \frac{e^{2\alpha^{(0)}dz}}{e^{2\alpha^{(0)}dz} - 1} (m^2 - ie^{-\beta^{(0)}dz} \varphi^{(1)}) \Phi_\alpha^2. \end{aligned} \quad (3.17)$$

Rescaling both scalar fields as

$$\phi_\alpha \implies e^{-\alpha^{(0)}dz} \phi_\alpha, \quad \Phi_\alpha \implies e^{-\alpha^{(0)}dz} \Phi_\alpha, \quad (3.18)$$

we rewrite the partition function in terms of light and heavy scalar fields as follows

$$\begin{aligned} Z = & \int D\phi_\alpha D\Phi_\alpha D\varphi^{(0)} D\varphi^{(1)} Dg_{\mu\nu}^{(0)} DT_{\mu\nu}^{(0)} \exp \left[-\frac{1}{2} \text{tr}_{xx'} \ln \frac{N}{2u} \frac{1}{e^{2\beta^{(0)}dz} - 1} \right. \\ & - \int d^D x \sqrt{g^{(0)}} \left\{ g^{\mu\nu(0)} [\partial_\mu e^{-\alpha^{(0)}dz} (\phi_\alpha + \Phi_\alpha)] [\partial_\nu e^{-\alpha^{(0)}dz} (\phi_\alpha + \Phi_\alpha)] \right. \\ & + (m^2 - ie^{-\beta^{(0)}dz} \varphi^{(1)}) \phi_\alpha^2 + \frac{1}{e^{2\alpha^{(0)}dz} - 1} (m^2 - ie^{-\beta^{(0)}dz} \varphi^{(1)}) \Phi_\alpha^2 \\ & \left. \left. + \frac{N}{2u} \left(\varphi^{(0)} - ie^{-\beta^{(0)}dz} \xi R^{(0)} \right)^2 + \frac{N}{2u} \frac{1}{e^{2\beta^{(0)}dz} - 1} (\varphi^{(1)} - \varphi^{(0)})^2 - NT_{\mu\nu}^{(0)} (g^{\mu\nu(0)} - g_B^{\mu\nu}) \right\} \right]. \end{aligned} \quad (3.19)$$

Here, effective couplings between light and heavy scalar bosons arise in the kinetic-energy sector.

The next step is to perform the Gaussian integral for the heavy scalar fields. For our simple derivation, we fix a gauge for the local speed of coarse graining as

$$\partial_\mu \alpha^{(0)} = 0. \quad (3.20)$$

The physical meaning of this gauge fixing is that the renormalization group transformation is taken into account in a uniform way, where the coefficient in front of the infinitesimal parameter for the renormalization group transformation does not depend on the spacetime coordinate x . Integrating over the heavy scalar fields with this gauge fixing, we obtain

$$\begin{aligned} Z = & \int D\phi_{\alpha x} D\varphi_x^{(0)} D\varphi_x^{(1)} Dg_{\mu\nu x}^{(0)} DT_{\mu\nu x}^{(0)} \exp \left\{ -S_{\phi\phi}^{(0)} - \Delta S_{\phi\Phi}^{(0)} - S_{\Phi\Phi}^{(0)} - S_{\varphi\varphi}^{(1)} \right. \\ & \left. + N \int d^D x \sqrt{g_x^{(0)}} T_{\mu\nu x}^{(0)} (g_x^{\mu\nu(0)} - g_{Bx}^{\mu\nu}) \right\}. \end{aligned} \quad (3.21)$$

Here, $S_{\phi\phi}^{(0)}$ is an effective action before the first renormalization group transformation, given by

$$S_{\phi\phi}^{(0)} = \int d^D x \sqrt{g_x^{(0)}} \left\{ e^{-2\alpha^{(0)} dz} g_x^{\mu\nu(0)} (\partial_\mu \phi_{\alpha x}) (\partial_\nu \phi_{\alpha x}) + (m^2 - i e^{-\beta^{(0)} dz} \varphi_x^{(1)}) \phi_{\alpha x}^2 \right\}. \quad (3.22)$$

$\Delta S_{\phi\Phi}^{(0)}$ is an effective action to result from the mixing terms between light and heavy scalar fields in the kinetic-energy part, given by

$$\Delta S_{\phi\Phi}^{(0)} = - \int d^D x \sqrt{g_x^{(0)}} \int d^D x' \sqrt{g_{x'}^{(0)}} e^{-2\alpha^{(0)} dz} g_x^{\mu\nu(0)} g_{x'}^{\mu'\nu'(0)} (\partial_\mu \phi_{\alpha x}) (\partial_{\nu'} \partial_{\mu'} G_{xx'}^{(0)}) (\partial_{\nu'} \phi_{\alpha x'}), \quad (3.23)$$

where $G_{xx'}^{(0)}$ is the Green's function of the heavy scalar bosons, and obeys

$$\begin{aligned} & \left\{ - \frac{1}{\sqrt{g_x^{(0)}}} \partial_\mu \left(\sqrt{g_x^{(0)}} g_x^{\mu\nu(0)} \partial_\nu \right) + \frac{e^{(2\alpha^{(0)} - \beta^{(0)}) dz}}{e^{2\alpha^{(0)} dz} - 1} (e^{-2\alpha^{(0)} dz} m^2 - i \varphi_x^{(1)}) \right\} G_{xx'}^{(0)} \\ &= \frac{1}{\sqrt{g_x^{(0)}}} \delta^{(D)}(x - x'). \end{aligned} \quad (3.24)$$

This Green's function plays a central role in the renormalization group transformation, regarded to be the only dynamic information. $S_{\Phi\Phi}^{(0)}$ serves as an effective vacuum action to originate from quantum fluctuations of the heavy scalar fields, given by

$$S_{\Phi\Phi}^{(0)} = \frac{N}{2} \text{tr}_{xx'} \ln e^{-2\alpha^{(0)} dz} \sqrt{g_x^{(0)}} \left\{ - \frac{1}{\sqrt{g_x^{(0)}}} \partial_\mu \left(\sqrt{g_x^{(0)}} g_x^{\mu\nu(0)} \partial_\nu \right) + \frac{e^{(2\alpha^{(0)} - \beta^{(0)}) dz}}{e^{2\alpha^{(0)} dz} - 1} (e^{-2\alpha^{(0)} dz} m^2 - i \varphi_x^{(1)}) \right\}. \quad (3.25)$$

Finally, $S_{\varphi\varphi}^{(1)}$ is an effective action after the first renormalization group transformation for the dual scalar bosons, given by

$$\begin{aligned} S_{\varphi\varphi}^{(1)} &= \frac{1}{2} \text{tr}_{xx'} \ln \frac{N}{2u} \frac{1}{e^{2\beta^{(0)} dz} - 1} \\ &+ \int d^D x \sqrt{g_x^{(0)}} \left\{ \frac{N}{2u} \left(\varphi_x^{(0)} - i e^{-\beta^{(0)} dz} \xi R_x^{(0)} \right)^2 + \frac{N}{2u} \frac{1}{e^{2\beta^{(0)} dz} - 1} (\varphi_x^{(1)} - \varphi_x^{(0)})^2 \right\}. \end{aligned} \quad (3.26)$$

Taking the gradient expansion in the vacuum sector of $S_{\Phi\Phi}^{(0)}$ with respect to the mass

of scalar bosons and keeping all terms up to the linear order in dz , we obtain

$$\begin{aligned}
Z = & \int D\phi_{\alpha x} D\varphi_x^{(0)} D\varphi_x^{(1)} Dg_{\mu\nu x}^{(0)} DT_{\mu\nu x}^{(0)} \exp \left[\right. \\
& \frac{N}{2} \text{tr}_{xx'} \ln(2\alpha^{(0)} dz) + \frac{1}{2} \text{tr}_{xx'} \ln(2\beta^{(0)} dz) - \frac{N}{2} \text{tr}_{xx'} \ln \sqrt{g_x^{(0)}} m^2 - \frac{1}{2} \text{tr}_{xx'} \ln \frac{N}{2u} \\
& - \int d^D x \sqrt{g_x^{(0)}} \left\{ (1 - 2\alpha^{(0)} dz) g_x^{\mu\nu(0)} (\partial_\mu \phi_{\alpha x}) (\partial_\nu \phi_{\alpha x}) + (m^2 - i\varphi_x^{(1)}) \phi_{\alpha x}^2 \right. \\
& - \int d^D x' \sqrt{g_{x'}^{(0)}} g_{x'}^{\mu\nu(0)} g_{x'}^{\mu'\nu'(0)} (\partial_\mu \phi_{\alpha x}) (\partial_\nu \partial_{\mu'} G_{xx'}^{(0)}) (\partial_{\nu'} \phi_{\alpha x'}) \\
& + 2\alpha^{(0)} dz N \left(-\mathcal{C}_\Lambda + \mathcal{C}_R R_x^{(0)} + \frac{\mathcal{C}_\varphi}{2} g_x^{\mu\nu(0)} (\partial_\mu \varphi_x^{(1)}) (\partial_\nu \varphi_x^{(1)}) + \mathcal{C}_\xi R_x^{(0)} \varphi_x^{(1)2} \right) \\
& \left. + \frac{N}{2u} \left(\varphi_x^{(0)} - i\xi R_x^{(0)} \right)^2 + \beta^{(0)} dz \frac{N}{4u} \left(\frac{\varphi_x^{(1)} - \varphi_x^{(0)}}{\beta^{(0)} dz} \right)^2 - NT_{\mu\nu x}^{(0)} (g_x^{\mu\nu(0)} - g_{Bx}^{\mu\nu}) \right\} \left. \right]. \quad (3.27)
\end{aligned}$$

We point out that the Einstein-Hilbert action appears to count vacuum fluctuations of high-energy scalar bosons in the background geometry, known to be the notion of induced gravity [40, 43, 44]. In addition, dual scalar bosons acquire their kinetic energy, where \mathcal{C}_φ and \mathcal{C}_ξ are positive constants. All the coefficients \mathcal{C}_Λ , \mathcal{C}_R , \mathcal{C}_φ , and \mathcal{C}_ξ decrease as the mass of scalar bosons increases. The Green's function $G_{xx'}^{(0)}$ is determined to linear order in dz by the equation

$$\left\{ -\frac{1}{\sqrt{g_x^{(0)}}} \partial_\mu \left(\sqrt{g_x^{(0)}} g_x^{\mu\nu(0)} \partial_\nu \right) + \frac{1}{2\alpha^{(0)} dz} (m^2 - i\varphi_x^{(1)}) \right\} G_{xx'}^{(0)} = \frac{1}{\sqrt{g_x^{(0)}}} \delta^{(D)}(x - x'). \quad (3.28)$$

3.2 Recursive renormalization group transformations a la Sung-Sik Lee

The idea for the second renormalization group transformation is that $S_{\phi\phi}^{(0)} + \Delta S_{\phi\Phi}^{(0)}$ is reformulated as $S_{\phi\phi}^{(1)}$, where the metric tensor is updated to be from $g^{(0)\mu\nu}$ to $g^{(1)\mu\nu}$ appropriately. However, this reformulation turns out not to be straightforward since $\Delta S_{\phi\Phi}^{(0)}$ is a nonlocal action because of $G_{xx'}^{(0)}$, which differs from the local kinetic energy term of $S_{\phi\phi}^{(0)}$. An important observation is that this Green's function is given by an exponential form $\sim \exp \left(- (2\alpha^{(0)} dz)^{-1/2} \sqrt{m^2 - i\varphi_x^{(1)}} |x - x'| \right)$, where the inverse of the decay length is $(2\alpha^{(0)} dz)^{-1/2} \sqrt{m^2 - i\varphi_x^{(1)}}$. As a result, we may only keep the local term in the gradient expansion as follows

$$(\partial_\nu \partial_{\mu'} G_{xx'}^{(0)}) \approx (\partial_\nu \partial_{\mu'} G_{xx'}^{(0)}) \frac{1}{\sqrt{g_{x'}}} \delta^{(D)}(x - x') + \dots, \quad (3.29)$$

where higher gradient terms are in higher orders of dz .

Taking into account this locality approximation justified to linear order in dz we obtain

$$\begin{aligned}
Z = & \int D\phi_\alpha D\varphi^{(0)} D\varphi^{(1)} Dg_{\mu\nu}^{(0)} DT_{\mu\nu}^{(0)} \exp \left[\right. \\
& \frac{N}{2} \text{tr}_{xx'} \ln(2\alpha^{(0)} dz) + \frac{1}{2} \text{tr}_{xx'} \ln(2\beta^{(0)} dz) - \frac{N}{2} \text{tr}_{xx'} \ln \sqrt{g^{(0)} m^2} - \frac{1}{2} \text{tr}_{xx'} \ln \frac{N}{2u} \\
& - \int d^D x \sqrt{g^{(0)}} \left\{ g^{\mu\nu(0)} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + (m^2 - i\varphi^{(1)}) \phi_\alpha^2 \right. \\
& - 2\alpha^{(0)} dz \left(g^{\mu\nu(0)} + \frac{1}{2\alpha^{(0)} dz} g^{\mu\nu'(0)} (\partial_{\nu'} \partial_{\mu'} G_{xx'}^{(0)})_{x' \rightarrow x} g^{\mu'\nu(0)} \right) (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) \\
& + 2\alpha^{(0)} dz N \left(-\mathcal{C}_\Lambda + \mathcal{C}_R R^{(0)} + \frac{\mathcal{C}_\varphi}{2} g^{\mu\nu(0)} (\partial_\mu \varphi^{(1)}) (\partial_\nu \varphi^{(1)}) + \mathcal{C}_\xi R^{(0)} \varphi^{(1)2} \right) \\
& \left. + \frac{N}{2u} \left(\varphi^{(0)} - i\xi R^{(0)} \right)^2 + \beta^{(0)} dz \frac{N}{4u} \left(\frac{\varphi^{(1)} - \varphi^{(0)}}{\beta^{(0)} dz} \right)^2 - NT_{\mu\nu}^{(0)} (g^{\mu\nu(0)} - g_B^{\mu\nu}) \right\} \left. \right]. \quad (3.30)
\end{aligned}$$

Then, it is straightforward to update the metric tensor up to the linear order in dz as follows

$$\begin{aligned}
Z = & \int D\phi_\alpha D\varphi^{(0)} D\varphi^{(1)} Dg_{\mu\nu}^{(0)} DT_{\mu\nu}^{(0)} Dg_{\mu\nu}^{(1)} DT_{\mu\nu}^{(1)} \exp \left[\right. \\
& \frac{N}{2} \text{tr}_{xx'} \ln(2\alpha^{(0)} dz) + \frac{1}{2} \text{tr}_{xx'} \ln(2\beta^{(0)} dz) - \frac{N}{2} \text{tr}_{xx'} \ln \sqrt{g^{(0)} m^2} - \frac{1}{2} \text{tr}_{xx'} \ln \frac{N}{2u} \\
& - \int d^D x \sqrt{g^{(1)}} \left\{ g^{(1)\mu\nu} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + (m^2 - i\varphi^{(1)}) \phi_\alpha^2 \right\} \\
& - N \int d^D x \sqrt{g^{(0)}} \left\{ \frac{1}{2u} \left(\varphi^{(0)} - i\xi R^{(0)} \right)^2 - T_{\mu\nu}^{(0)} (g^{(0)\mu\nu} - g_B^{\mu\nu}) \right\} \\
& - N \int d^D x \sqrt{g^{(0)}} \left\{ 2\alpha^{(0)} dz \left(-\mathcal{C}_\Lambda + \mathcal{C}_R R^{(0)} + \frac{\mathcal{C}_\varphi}{2} g^{\mu\nu(0)} (\partial_\mu \varphi^{(1)}) (\partial_\nu \varphi^{(1)}) + \mathcal{C}_\xi R^{(0)} \varphi^{(1)2} \right) \right. \\
& \left. + \beta^{(0)} dz \frac{N}{4u} \left(\frac{\varphi^{(1)} - \varphi^{(0)}}{\beta^{(0)} dz} \right)^2 - T_{\mu\nu}^{(1)} \left(g^{(1)\mu\nu} - g^{(0)\mu\nu} - g^{(0)\mu\nu'} (\partial_{\nu'} \partial_{\mu'} G_{xx'}^{(0)})_{x' \rightarrow x} g^{(0)\mu'\nu} \right) \right\} \left. \right]. \quad (3.31)
\end{aligned}$$

Here, $T_{\mu\nu}^{(1)}$ is a Lagrange multiplier field to impose the update condition. This completes the first renormalization group transformation.

Based on the above expression, we can estimate the partition function after the $(f-1)$ th renormalization group transformation, given by

$$Z = \int D\phi_\alpha \Pi_{k=0}^f D\varphi^{(k)} Dg_{\mu\nu}^{(k)} DT_{\mu\nu}^{(k)} \exp \left\{ -S_\Lambda - S_{UV} - S_{IR} - S_{Bulk} \right\}. \quad (3.32)$$

Here, S_Λ is an effective action which is UV-divergent, and given by

$$\begin{aligned}
S_\Lambda = & -\frac{N}{2} \sum_{k=1}^f \text{tr}_{xx'} \ln(2dz) - \frac{1}{2} \sum_{k=1}^f \text{tr}_{xx'} \ln(2dz) \\
& + \frac{1}{2} \sum_{k=1}^f \text{tr}_{xx'} \ln \frac{N}{2u} + \frac{N}{2} \sum_{k=1}^f \text{tr}_{xx'} \ln \sqrt{g^{(k-1)} m^2}. \quad (3.33)
\end{aligned}$$

S_{UV} is an effective action imposing UV boundary conditions for both metric and dual scalar fields, given by

$$S_{UV} = N \int d^D x \left\{ -\frac{1}{2u} \sqrt{g^{(0)}} \left(\varphi^{(0)} - \xi R^{(0)} \right)^2 - \sqrt{g^{(0)}} T_{\mu\nu}^{(0)} (g^{(0)\mu\nu} - g_B^{\mu\nu}) \right\}. \quad (3.34)$$

S_{IR} is an effective action implementing IR boundary conditions of both metric and dual scalar fields, given by

$$S_{IR} = \int d^D x \sqrt{g^{(f)}} \left\{ g^{(f)\mu\nu} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + (m^2 + \varphi^{(f)}) \phi_\alpha^2 \right\}. \quad (3.35)$$

S_{Bulk} is an effective action to govern the dynamics of both metric and dual scalar fields, given by

$$\begin{aligned} S_{Bulk} = N(2dz) \sum_{k=1}^f \int d^D x \sqrt{g^{(k-1)}} \left\{ -\frac{1}{2u} \left(\frac{\varphi^{(k)} - \varphi^{(k-1)}}{2dz} \right)^2 \right. \\ \left. - T_{\mu\nu}^{(k)} \left(\frac{g^{(k)\mu\nu} - g^{(k-1)\mu\nu}}{2dz} - \frac{1}{2dz} g^{(k-1)\mu\nu'} (\partial_{\nu'} \partial_{\mu'} G_{xx'}^{(k-1)})_{x' \rightarrow x} g^{(k-1)\mu'\nu} \right) \right. \\ \left. - \mathcal{C}_\Lambda + \mathcal{C}_R R^{(k-1)} - \frac{\mathcal{C}_\varphi}{2} g^{\mu\nu(k-1)} (\partial_\mu \varphi^{(k)}) (\partial_\nu \varphi^{(k)}) - \mathcal{C}_\xi R^{(k-1)} \varphi^{(k)2} \right\}. \quad (3.36) \end{aligned}$$

$G_{xx'}^{(k-1)}$ is the Green's function for the k -th renormalization group transformation, given by

$$\left\{ -\frac{1}{\sqrt{g_x^{(k-1)}}} \partial_\mu \left(\sqrt{g_x^{(k-1)}} g_x^{\mu\nu(k-1)} \partial_\nu \right) + \frac{1}{2dz} (m^2 + \varphi_x^{(k)}) \right\} G_{xx'}^{(k-1)} = \frac{1}{\sqrt{g_x^{(k-1)}}} \delta^{(D)}(x - x'). \quad (3.37)$$

Here, we replaced $\varphi^{(k)}$ with $i\varphi^{(k)}$ and adopt gauge fixing for both uniform speeds of coarse graining as $\alpha^{(k)} = \beta^{(k)} = 1$. More discussions on this gauge fixing will be presented below.

The last step is to rewrite the above partition function in the continuous coordinate representation instead of the discrete variable k . Considering

$$\begin{aligned} (2dz) \sum_{k=1}^f \int d^D x \sqrt{g^{(k-1)}} T_{\mu\nu}^{(k)} \left(\frac{g^{(k)\mu\nu} - g^{(k-1)\mu\nu}}{2dz} \right) \\ \implies \int_0^{z_f} dz \int d^D x \sqrt{g(x, z)} T_{\mu\nu}(x, z) \partial_z g^{\mu\nu}(x, z) \end{aligned} \quad (3.38)$$

with $(2dz) \sum_{k=1}^f \implies \int_0^{z_f} dz$ and $\frac{g^{(k)\mu\nu} - g^{(k-1)\mu\nu}}{2dz} \implies \partial_z g^{\mu\nu}(x, z)$, we obtain

$$\begin{aligned}
Z = Z_\Lambda \int & D\phi_\alpha(x) D\varphi(x, z) Dg_{\mu\nu}(x, z) DT_{\mu\nu}(x, z) \exp \left[\right. \\
& - \int d^D x \sqrt{g(x, z_f)} \left\{ g^{\mu\nu}(x, z_f) [\partial_\mu \phi_\alpha(x)] [\partial_\nu \phi_\alpha(x)] + [m^2 + \varphi(x, z_f)] \phi_\alpha^2(x) \right\} \\
& - N \int d^D x \sqrt{g(x, 0)} \left\{ -\frac{1}{2u} \left(\varphi(x, 0) - \xi R(x, 0) \right)^2 - T_{\mu\nu}(x, 0) \left(g^{\mu\nu}(x, 0) - g_B^{\mu\nu}(x) \right) \right\} \\
& - N \int_0^{z_f} dz \int d^D x \sqrt{g(x, z)} \left\{ \right. \\
& - \frac{1}{2u} [\partial_z \varphi(x, z)]^2 - \frac{C_\varphi}{2} g^{\mu\nu}(x, z) [\partial_\mu \varphi(x, z)] [\partial_\nu \varphi(x, z)] \\
& - C_\xi R(x, z) [\varphi(x, z)]^2 + \frac{1}{2\kappa} \left(R(x, z) - 2\Lambda \right) \\
& \left. - T_{\mu\nu}(x, z) \left(\partial_z g^{\mu\nu}(x, z) - \frac{1}{2dz} g^{\mu\nu'}(x, z) (\partial_{\nu'} \partial_{\mu'} G_{xx'}[g_{\mu\nu}(x, z), \varphi(x, z)])_{x' \rightarrow x} g^{\mu'\nu'}(x, z) \right) \right\} \left. \right], \tag{3.39}
\end{aligned}$$

where all UV divergent terms are absorbed into a normalization constant Z_Λ . For the Einstein-Hilbert action, we took the following replacements

$$C_R \equiv \frac{1}{2\kappa}, \quad \frac{C_\Lambda}{C_R} \equiv 2\Lambda. \tag{3.40}$$

All the dynamical information is encoded into the Green's function, given by

$$\begin{aligned}
& \left\{ -\frac{1}{\sqrt{g(x, z)}} \partial_\mu \left(\sqrt{g(x, z)} g^{\mu\nu}(x, z) \partial_\nu \right) + \frac{1}{2dz} [m^2 + \varphi(x, z)] \right\} G_{xx'}[g_{\mu\nu}(x, z), \varphi(x, z)] \\
& = \frac{1}{\sqrt{g(x, z)}} \delta^{(D)}(x - x'). \tag{3.41}
\end{aligned}$$

As discussed earlier in section 2, the physical description is quite clear in this effective action. The D -dimensional metric tensor evolves through the extra dimension, following the equation of motion

$$\partial_z g^{\mu\nu}(x, z) = \frac{1}{2dz} g^{\mu\nu'}(x, z) (\partial_{\nu'} \partial_{\mu'} G_{xx'}[g_{\mu\nu}(x, z), \varphi(x, z)])_{x' \rightarrow x} g^{\mu'\nu'}(x, z). \tag{3.42}$$

As a result, a fully renormalized metric $g^{\mu\nu}(x, z_f)$ appears in the IR effective action $\mathcal{S}_{IR} = \int d^D x \sqrt{g(x, z_f)} \left\{ g^{\mu\nu}(x, z_f) [\partial_\mu \phi_\alpha(x)] [\partial_\nu \phi_\alpha(x)] + [m^2 + \varphi(x, z_f)] \phi_\alpha^2(x) \right\}$, describing all possible renormalizations such as field renormalization, mass renormalization, and interaction renormalization. This evolution equation plays essentially the same role as renormalization group β -functions.

3.3 Remarks on the renormalization group transformation for the metric tensor

Although the renormalization group flow of the metric tensor results from quantum fluctuations of matter fields, the metric tensor is not fully dynamical in contrast with the

holographic duality conjecture [2–8]. This should be contrasted with the order-parameter field which is promoted to be dynamical in the extra-dimensional space. We recall that the dynamics of the order-parameter field originates from effective self-interactions between matter fields. In this respect we introduce effective tensor-field interactions into the partition function as follows

$$Z = \int D\phi_\alpha \exp \left[- \int d^D x \sqrt{g_B} \left\{ g_B^{\mu\nu} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + m^2 \phi_\alpha^2 + \xi R_B \phi_\alpha^2 + \frac{u}{2N} \phi_\alpha^2 \phi_\beta^2 + \frac{\lambda}{2N} [(\partial^\mu \phi_\alpha) (\partial^\nu \phi_\alpha)] [(\partial_\mu \phi_\beta) (\partial_\nu \phi_\beta)] \right\} \right]. \quad (3.43)$$

It is natural to expect that the effective interaction term $\frac{\lambda}{2N} [(\partial^\mu \phi_\alpha) (\partial^\nu \phi_\alpha)] [(\partial_\mu \phi_\beta) (\partial_\nu \phi_\beta)]$ is irrelevant in the renormalization group sense as long as the coupling constant λ remains to be below a critical value. However, we point out that these tensor-type quantum fluctuations promote the emergent metric tensor to be fully dynamical [41]. Although one can take into account an effective interaction term of the exact energy-momentum tensor in the effective Lagrangian, the above introduction of the effective interaction term is sufficient in discussing which approximation scheme has to be used for the metric renormalization-group transformation.

Performing the Hubbard-Stratonovich transformation for both effective interactions, we obtain

$$Z = \int D\phi_\alpha D\varphi^{(0)} Dg_{\mu\nu}^{(0)} Dt_{\mu\nu}^{(0)} Dg_{\mu\nu} \exp \left[- \int d^D x \sqrt{g^{(0)}} \left\{ (g^{\mu\nu(0)} - ig^{\mu\nu}) (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + (m^2 - i\varphi^{(0)}) \phi_\alpha^2 + \xi R^{(0)} \phi_\alpha^2 + \frac{N}{2u} \varphi^{(0)2} + \frac{N}{2\lambda} g^{\mu\nu} g_{\mu\nu} - N t_{\mu\nu}^{(0)} (g^{\mu\nu(0)} - g_B^{\mu\nu}) \right\} \right], \quad (3.44)$$

where $t_{\mu\nu}^{(0)}$ is a Lagrange multiplier field to impose the initial condition $g^{\mu\nu(0)} = g_B^{\mu\nu}$. Shifting the metric tensor in the following way

$$g^{\mu\nu(0)} \implies g^{\mu\nu(0)} + ig^{\mu\nu}, \quad (3.45)$$

we obtain

$$Z = \int D\phi_\alpha D\varphi^{(0)} Dg_{\mu\nu}^{(0)} Dt_{\mu\nu}^{(0)} Dg_{\mu\nu} \exp \left[- \int d^D x \sqrt{\det[g_{\mu\nu}^{(0)} + ig_{\mu\nu}]} \left\{ g^{\mu\nu(0)} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + (m^2 - i\varphi^{(0)}) \phi_\alpha^2 + \xi R[g_{\mu\nu}^{(0)} + ig_{\mu\nu}] \phi_\alpha^2 + \frac{N}{2u} \varphi^{(0)2} + \frac{N}{2\lambda} g^{\mu\nu} g_{\mu\nu} - N t_{\mu\nu}^{(0)} (g^{\mu\nu(0)} + ig^{\mu\nu} - g_B^{\mu\nu}) \right\} \right]. \quad (3.46)$$

We now perform the path integral $\int Dt_{\mu\nu}^{(0)} Dg_{\mu\nu}$. The approximation that we have to use is to keep quantum fluctuations of the metric tensor up to the linear order. In other words, we neglect the $ig_{\mu\nu}$ contribution in both the determinant and Ricci scalar. As a result, we obtain

$$Z \approx \int D\phi_\alpha D\varphi^{(0)} Dg_{\mu\nu}^{(0)} \exp \left[- \int d^D x \sqrt{g^{(0)}} \left\{ g^{\mu\nu(0)} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + (m^2 - i\varphi^{(0)}) \phi_\alpha^2 + \xi R^{(0)} \phi_\alpha^2 + \frac{N}{2u} \varphi^{(0)2} - \frac{N}{2\lambda} (g^{\mu\nu(0)} - g_B^{\mu\nu}) (g_{\mu\nu}^{(0)} - g_{B\mu\nu}) \right\} \right]. \quad (3.47)$$

To perform the renormalization group transformation for the metric tensor, we introduce an auxiliary field in the following way

$$\begin{aligned}
Z &= \int D\phi_\alpha D\varphi^{(0)} D\varphi^{(1)} Dg_{\mu\nu}^{(0)} DG_{\mu\nu}^{(0)} \\
&\exp \left[- \int d^D x \sqrt{i^{-1} \det[g_{\mu\nu}^{(0)} + g_{B\mu\nu}]} \left\{ i[g^{\mu\nu(0)} + g_B^{\mu\nu}] (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + (m^2 - ie^{-\beta^{(0)} dz} \varphi^{(1)}) \phi_\alpha^2 \right. \right. \\
&+ \frac{N}{2u} \left(\varphi^{(0)} - ie^{-\beta^{(0)} dz} \xi R [i(g_{\mu\nu}^{(0)} + g_{B\mu\nu})] \right)^2 + \frac{N}{2u} \frac{1}{e^{2\beta^{(0)} dz} - 1} (\varphi^{(1)} - \varphi^{(0)})^2 \\
&\left. \left. + \frac{N}{2\lambda} g^{\mu\nu(0)} g_{\mu\nu}^{(0)} + \frac{N}{2\lambda_G} G^{\mu\nu(0)} G_{\mu\nu}^{(0)} \right\} \right], \tag{3.48}
\end{aligned}$$

where the renormalization group transformation for the order-parameter field has been performed and $g^{\mu\nu(0)} \Rightarrow i(g^{\mu\nu(0)} + g_B^{\mu\nu})$ has been taken. Separating slow and fast degrees of freedom in the metric tensor as

$$g^{\mu\nu(0)} \Rightarrow g^{\mu\nu(0)} + \mathcal{G}^{\mu\nu(0)}, \quad G^{\mu\nu(0)} \Rightarrow c_g^{(0)} g^{\mu\nu(0)} + c_G^{(0)} \mathcal{G}^{\mu\nu(0)}, \tag{3.49}$$

where both coefficients are given by

$$c_g^{(0)} = \frac{\lambda^{-1}}{\mu^{(0)} \lambda_G^{-1/2}}, \quad c_G^{(0)} = -\frac{\mu^{(0)}}{\lambda_G^{-1/2}}, \quad \mu^{(0)} = \frac{\lambda^{-1/2}}{\sqrt{e^{2\delta^{(0)} dz} - 1}}, \tag{3.50}$$

and rescaling both degrees of freedom as

$$g^{\mu\nu(0)} \Rightarrow e^{-\delta^{(0)} dz} g^{\mu\nu(0)}, \quad \mathcal{G}^{\mu\nu(0)} \Rightarrow e^{-\delta^{(0)} dz} \mathcal{G}^{\mu\nu(0)}, \tag{3.51}$$

where $\delta^{(0)}$ is the local speed of coarse graining, we have

$$\begin{aligned}
Z &= \int D\phi_\alpha D\varphi^{(0)} D\varphi^{(1)} Dg_{\mu\nu}^{(0)} D\mathcal{G}_{\mu\nu}^{(0)} \exp \left[- \int d^D x \sqrt{i^{-1} \det[e^{\delta^{(0)} dz} (g_{\mu\nu}^{(0)} + e^{-\delta^{(0)} dz} g_{B\mu\nu} + \mathcal{G}_{\mu\nu}^{(0)})]} \right. \\
&\left\{ ie^{-\delta^{(0)} dz} (g^{\mu\nu(0)} + e^{\delta^{(0)} dz} g_B^{\mu\nu} + \mathcal{G}^{\mu\nu(0)}) (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + (m^2 - ie^{-\beta^{(0)} dz} \varphi^{(1)}) \phi_\alpha^2 \right. \\
&+ \frac{N}{2u} \left(\varphi^{(0)} - ie^{-\beta^{(0)} dz} \xi R [ie^{\delta^{(0)} dz} (g_{\mu\nu}^{(0)} + e^{-\delta^{(0)} dz} g_{B\mu\nu} + \mathcal{G}_{\mu\nu}^{(0)})] \right)^2 + \frac{N}{2u} \frac{1}{e^{2\beta^{(0)} dz} - 1} (\varphi^{(1)} - \varphi^{(0)})^2 \\
&\left. \left. + \frac{N}{2\lambda} g^{\mu\nu(0)} g_{\mu\nu}^{(0)} + \frac{N}{2\lambda} \frac{1}{e^{2\delta^{(0)} dz} - 1} \mathcal{G}^{\mu\nu(0)} \mathcal{G}_{\mu\nu}^{(0)} \right\} \right]. \tag{3.52}
\end{aligned}$$

To perform the renormalization group transformation for the metric tensor, we consider the linear approximation again for the $\mathcal{G}_{\mu\nu}^{(0)}$ path integral, where the $\mathcal{G}_{\mu\nu}^{(0)}$ contribution in both the determinant and the Ricci scalar is neglected. As a result, we obtain

$$\begin{aligned}
Z &= \int D\phi_\alpha D\varphi^{(0)} D\varphi^{(1)} Dg_{\mu\nu}^{(0)} \exp \left[- \int d^D x \sqrt{e^{\delta^{(0)} dz} g^{(0)}} \left\{ e^{-\delta^{(0)} dz} g^{\mu\nu(0)} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) \right. \right. \\
&+ (m^2 - ie^{-\beta^{(0)} dz} \varphi^{(1)}) \phi_\alpha^2 + \frac{N}{2u} \left(\varphi^{(0)} - ie^{-\beta^{(0)} dz} \xi R^{(0)} \right)^2 + \frac{N}{2u} \frac{1}{e^{2\beta^{(0)} dz} - 1} (\varphi^{(1)} - \varphi^{(0)})^2 \\
&\left. \left. - \frac{N}{2\lambda} (g^{\mu\nu(0)} - e^{\delta^{(0)} dz} g_B^{\mu\nu}) (g_{\mu\nu}^{(0)} - e^{-\delta^{(0)} dz} g_{B\mu\nu}) + \frac{\lambda}{2N} [1 - e^{-2\delta^{(0)} dz}] (\partial^\mu \phi_\alpha) (\partial^\nu \phi_\alpha) (\partial_\mu \phi_\beta) (\partial_\nu \phi_\beta) \right\} \right], \tag{3.53}
\end{aligned}$$

where an effective interaction term $\frac{\lambda}{2N}[1 - e^{-2\delta^{(0)}dz}](\partial^\mu\phi_\alpha)(\partial^\nu\phi_\alpha)(\partial_\mu\phi_\beta)(\partial_\nu\phi_\beta)$ has been generated. Performing the Hubbard-Stratonovich transformation for the newly generated effective interaction term and updating the metric tensor in the boundary action, we obtain

$$\begin{aligned}
Z = & \int D\phi_\alpha D\varphi^{(0)} D\varphi^{(1)} Dg_{\mu\nu}^{(0)} D\tilde{g}_{\mu\nu}^{(0)} \exp \left[- \int d^D x \sqrt{e^{\delta^{(0)}dz} \tilde{g}^{(0)}} \left\{ e^{-\delta^{(0)}dz} \tilde{g}^{\mu\nu(0)} (\partial_\mu\phi_\alpha)(\partial_\nu\phi_\alpha) \right. \right. \\
& + (m^2 - ie^{-\beta^{(0)}dz} \varphi^{(1)}) \phi_\alpha^2 + \frac{N}{2u} \frac{1}{e^{2\beta^{(0)}dz} - 1} (\varphi^{(1)} - \varphi^{(0)})^2 \\
& - \frac{N}{2\lambda} \frac{1}{e^{2\delta^{(0)}dz} - 1} (\tilde{g}^{\mu\nu(0)} - g^{\mu\nu(0)}) (\tilde{g}_{\mu\nu}^{(0)} - g_{\mu\nu}^{(0)}) \left. \right\} - \int d^D x \sqrt{e^{\delta^{(0)}dz} g^{(0)}} \left\{ \frac{N}{2u} \left(\varphi^{(0)} - ie^{-\beta^{(0)}dz} \xi R^{(0)} \right)^2 \right. \\
& \left. \left. - \frac{N}{2\lambda} (g^{\mu\nu(0)} - e^{\delta^{(0)}dz} g_B^{\mu\nu}) (g_{\mu\nu}^{(0)} - e^{-\delta^{(0)}dz} g_{B\mu\nu}) \right\} \right], \tag{3.54}
\end{aligned}$$

where $\tilde{g}^{\mu\nu(0)}$ is an updated metric tensor.

Finally, we perform the renormalization group transformation for the matter sector as discussed in the previous section. Taking the locality approximation for the Green's function and keeping all terms up to the linear order in dz , we find

$$\begin{aligned}
Z = & \int D\phi_\alpha D\varphi^{(0)} D\varphi^{(1)} Dg_{\mu\nu}^{(0)} D\tilde{g}_{\mu\nu}^{(0)} \\
& \exp \left[- \int d^D x \sqrt{\tilde{g}^{(0)}} \left\{ (\tilde{g}^{\mu\nu(0)} - \tilde{g}^{\mu\delta(0)} (\partial_\delta \partial_{\delta'} G_{xx'}^{(0)})_{x' \rightarrow x} \tilde{g}^{\delta'\nu(0)}) (\partial_\mu\phi_\alpha)(\partial_\nu\phi_\alpha) + (m^2 - i\varphi^{(1)}) \phi_\alpha^2 \right. \right. \\
& + 2\beta^{(0)} dz \frac{N}{2u} \left(\frac{\varphi^{(1)} - \varphi^{(0)}}{2\beta^{(0)} dz} \right)^2 - 2\delta^{(0)} dz \frac{N}{2\lambda} \left(\frac{\tilde{g}^{\mu\nu(0)} - g^{\mu\nu(0)}}{2\delta^{(0)} dz} \right) \left(\frac{\tilde{g}_{\mu\nu}^{(0)} - g_{\mu\nu}^{(0)}}{2\delta^{(0)} dz} \right) \\
& + 2\alpha^{(0)} dz N \left(-\mathcal{C}_\Lambda + \mathcal{C}_R \tilde{R}^{(0)} + \frac{\mathcal{C}_\varphi}{2} \tilde{g}^{\mu\nu(0)} (\partial_\mu\varphi^{(1)}) (\partial_\nu\varphi^{(1)}) + \mathcal{C}_\xi \tilde{R}^{(0)} \varphi^{(1)2} \right) \left. \right\} \\
& - \int d^D x \sqrt{g^{(0)}} \left\{ \frac{N}{2u} \left(\varphi^{(0)} - i\xi R^{(0)} \right)^2 - \frac{N}{2\lambda} (g^{\mu\nu(0)} - g_B^{\mu\nu}) (g_{\mu\nu}^{(0)} - g_{B\mu\nu}) \right\}. \tag{3.55}
\end{aligned}$$

Upgrading the metric tensor $\tilde{g}_{\mu\nu}^{(0)}$ to make the kinetic-energy term be invariant, we rewrite the above expression as follows

$$\begin{aligned}
Z = & \int D\phi_\alpha D\varphi^{(0)} D\varphi^{(1)} Dg_{\mu\nu}^{(0)} Dg_{\mu\nu}^{(1)} \\
& \exp \left[- \int d^D x \sqrt{g^{(1)}} \left\{ g^{\mu\nu(1)} (\partial_\mu\phi_\alpha)(\partial_\nu\phi_\alpha) + (m^2 - i\varphi^{(1)}) \phi_\alpha^2 + 2\beta^{(0)} dz \frac{N}{2u} \left(\frac{\varphi^{(1)} - \varphi^{(0)}}{2\beta^{(0)} dz} \right)^2 \right. \right. \\
& - 2\delta^{(0)} dz \frac{N}{2\lambda} \left(\frac{g^{\mu\nu(1)} - g^{\mu\nu(0)} - g^{\mu\delta(0)} (\partial_\delta \partial_{\delta'} G_{xx'}^{(0)})_{x' \rightarrow x} g^{\delta'\nu(0)}}{2\delta^{(0)} dz} \right) \left(\frac{g_{\mu\nu}^{(1)} - g_{\mu\nu}^{(0)} - g_{\mu\delta}^{(0)} (\partial_\delta \partial_{\delta'} G_{xx'}^{(0)})_{x' \rightarrow x} g_{\delta'\nu}^{(0)}}{2\delta^{(0)} dz} \right) \\
& + 2\alpha^{(0)} dz N \left(-\mathcal{C}_\Lambda + \mathcal{C}_R R^{(1)} + \frac{\mathcal{C}_\varphi}{2} g^{\mu\nu(1)} (\partial_\mu\varphi^{(1)}) (\partial_\nu\varphi^{(1)}) + \mathcal{C}_\xi R^{(1)} \varphi^{(1)2} \right) \left. \right\} \\
& - \int d^D x \sqrt{g^{(0)}} \left\{ \frac{N}{2u} \left(\varphi^{(0)} - i\xi R^{(0)} \right)^2 - \frac{N}{2\lambda} (g^{\mu\nu(0)} - g_B^{\mu\nu}) (g_{\mu\nu}^{(0)} - g_{B\mu\nu}) \right\}. \tag{3.56}
\end{aligned}$$

Repeating all these renormalization group transformations within the recursion frame-

work and rewriting the recursive expression in the continuous variable z , we obtain

$$\begin{aligned}
Z = Z_\Lambda \int & D\phi_\alpha(x) D\varphi(x, z) Dg_{\mu\nu}(x, z) \exp \left[\right. \\
& - \int d^D x \sqrt{g(x, z_f)} \left\{ g^{\mu\nu}(x, z_f) [\partial_\mu \phi_\alpha(x)] [\partial_\nu \phi_\alpha(x)] + [m^2 + \varphi(x, z_f)] \phi_\alpha^2(x) \right\} \\
& - N \int d^D x \sqrt{g(x, 0)} \left\{ - \frac{1}{2u} \left(\varphi(x, 0) - \xi R(x, 0) \right)^2 - \frac{1}{2\lambda} \left(g^{\mu\nu}(x, 0) - g_B^{\mu\nu}(x) \right)^2 \right\} \\
& - N \int_0^{z_f} dz \int d^D x \sqrt{g(x, z)} \left\{ \right. \\
& \quad - \frac{1}{2u} [\partial_z \varphi(x, z)]^2 - \frac{\mathcal{C}_\varphi}{2} g^{\mu\nu}(x, z) [\partial_\mu \varphi(x, z)] [\partial_\nu \varphi(x, z)] - \mathcal{C}_\xi R(x, z) [\varphi(x, z)]^2 \\
& \quad - \frac{1}{2\lambda} \left(\partial_z g^{\mu\nu}(x, z) - \frac{1}{2dz} g^{\mu\nu'}(x, z) (\partial_{\nu'} \partial_{\mu'} G_{xx'} [g_{\mu\nu}(x, z), \varphi(x, z)]) \Big|_{x' \rightarrow x} g^{\mu'\nu'}(x, z) \right)^2 \\
& \quad \left. + \frac{1}{2\kappa} \left(R(x, z) - 2\Lambda \right) \right\} \left. \right]. \tag{3.57}
\end{aligned}$$

Here, the second derivative with respect to z coordinatizing the extra dimension arises in the metric evolution, resulting from essentially the same procedure as that in the order-parameter evolution $[\partial_z \varphi(x, z)]^2$, where the linear approximation for the metric tensor has been used. It is interesting to notice that the linear approximation for quantum fluctuations of the metric tensor gives rise to the highly nonlinear renormalization group flow, shown in Eq. (3.57), where the metric-tensor renormalization is intertwined with the renormalization of the order-parameter field through the Green's function of the matter field. In this case renormalization group β -functions are given by IR boundary conditions, where the linear derivative in z appears from the bulk effective action by the technique of integration-by-parts. In the next section we demonstrate that the evolution equations of the metric tensor along the extra dimensional space are nothing but the renormalization group β -functions.

Before going further, we point out that this holographic dual effective field theory is not covariant but a gauge-fixed version. In other words, the metric-tensor components involved with the extra dimension are gauge-fixed as follows: $g_{DD}(x, z) = 1$ and $g_{\mu D}(x, z) = 0$ with $\mu = 0, \dots, D-1$. It turns out that the gauge choice of $\alpha^{(k)} = 1$ gives rise to $g_{DD}(x, z) = 1$. On the other hand, the gauge freedom in $g_{\mu D}(x, z) = 0$ results from the invariance of the partition function with respect to D -dimensional diffeomorphism after the renormalization group transformation with dz [31–33]. While we do not delve into this issue here, a fully covariant formulation has been constructed in the absence of dual scalar fields, where the D -dimensional Einstein-Hilbert action is uplifted into the $(D+1)$ -dimensional Einstein-Hilbert one via recursive renormalization group transformations [31–33].

We also point out that the order of the renormalization group transformations for the dual and original boson fields is immaterial. We recall that the $dz \rightarrow 0$ limit with $z_f \equiv dz \sum_{k=1}^f = fdz$ controls our recursive renormalization group transformations. The order of the renormalization group transformations does not matter in this controllable limit.

4 Physical interpretation of the emergent dynamical metric tensor: From renormalization group equations of coupling parameters to the evolution equation of the metric tensor

In this section we demonstrate explicitly that renormalization group β -functions in the quantum field theory are nothing but IR boundary conditions in the large N limit of the holographic dual field theory. We do so by comparing our holographic theory and yet another real space renormalization group transformation (the Kadanoff block-spin transformation). For simplicity in the presentation, we switch off the self-interactions and consider a one-dimensional lattice field theory, given by

$$Z = \int D\Phi_i \exp \left[- \int_0^\beta d\tau \sum_{i=1}^M \left\{ (\partial_\tau \Phi_i)^2 - t(\Phi_i \Phi_{i+1} + \Phi_{i+1} \Phi_i) + m^2 \Phi_i^2 \right\} \right]. \quad (4.1)$$

Here, Φ_i is a real scalar field at site i . t is a hopping integral and m is a mass parameter. In the case of one dimensional lattice, it is straightforward to perform the Kadanoff block-spin transformation [51]. Integrating out all dynamical fields at even lattice sites, we obtain an effective field theory for odd-site scalar fields with renormalized hopping and mass parameters. Implementing this renormalization group transformation in a recursive way [48], we obtain the following expression of the partition function

$$\begin{aligned} Z = & \int D\Phi(i, \tau) Dt(z) Dm^2(z) \\ & \times \delta(t(0) - t) \delta(m^2(0) - m^2) \delta\left(\partial_z m^2(z) + \frac{[t(z)]^2}{am^2(z)}\right) \delta\left(\partial_z t(z) + \frac{1}{a}t(z) - \frac{[t(z)]^2}{2am^2(z)}\right) \\ & \times \exp \left[- \int_0^\beta d\tau \sum_{i=1}^M \left\{ (\partial_\tau \Phi(i, \tau))^2 + m^2(z_f) [\Phi(i, \tau)]^2 \right. \right. \\ & \left. \left. - t(z_f) (\Phi(i, \tau) \Phi(i+1, \tau) + \Phi(i+1, \tau) \Phi(i, \tau)) \right\} - \frac{M}{4a} \int_0^{z_f} dz \sum_{i\omega_n} \ln(\omega_n^2 + m^2(z)) \right]. \end{aligned} \quad (4.2)$$

Here, a is a scale for the renormalization group transformation, defined in Eqs. (A.8) and (A.9), which corresponds to dz in the previous section. All details are shown in appendix A, where recursive Kadanoff block-spin transformations have been performed in the presence of self-interactions. We point out that both the hopping and mass parameters are renormalized to appear in the IR boundary action. In this respect two δ -function constraints serve as renormalization group β -functions for the hopping and mass parameters, respectively.

To show the equivalence between the above renormalization group β -functions and the evolution equations for the metric-tensor fields, we recall the gravity formulation, given

by

$$\begin{aligned}
Z &= \int D\phi_\alpha(x) Dg_{\mu\nu}(x, z) \delta\left(g^{\mu\nu}(x, 0) - g_B^{\mu\nu}\right) \\
&\quad \times \delta\left(\partial_z g^{\mu\nu}(x, z) - \frac{1}{2dz} g^{\mu\nu'}(x, z) (\partial_{\nu'} \partial_{\mu'} G_{xx'}[g_{\mu\nu}(x, z)])_{x' \rightarrow x} g^{\mu'\nu'}(x, z)\right) \\
&\quad \times \exp\left[-\int d^D x \sqrt{g(x, z_f)} \left\{g^{\mu\nu}(x, z_f) (\partial_\mu \phi_\alpha(x)) (\partial_\nu \phi_\alpha(x)) + m^2 \phi_\alpha^2(x)\right\}\right. \\
&\quad \left.- \frac{N}{2\kappa} \int_0^{z_f} dz \int d^D x \sqrt{g(x, z)} (R(x, z) - 2\Lambda)\right], \tag{4.3}
\end{aligned}$$

which results from the UV theory of

$$Z = \int D\phi_\alpha(x) \exp\left[-\int d^D x \sqrt{g_B} \left\{g_B^{\mu\nu} (\partial_\mu \phi_\alpha) (\partial_\nu \phi_\alpha) + m^2 \phi_\alpha^2\right\}\right]. \tag{4.4}$$

Comparing each term between Eq. (4.2) and Eq. (4.3), we observe the following correspondences,

$$\sqrt{g(x, z_f)} m^2 = m^2(z_f) - 2t(z_f) \implies \frac{1}{\sqrt{g(x, z_f)}} = \frac{m^2}{m^2(z_f) - 2t(z_f)} \tag{4.5}$$

for the metric determinant,

$$\sqrt{g(x, z_f)} g^{xx}(x, z_f) = t(z_f) \implies g^{xx}(x, z_f) = \frac{m^2}{m^2(z_f) - 2t(z_f)} t(z_f) \tag{4.6}$$

for the $g^{xx}(x, z_f)$ component, and

$$\sqrt{g(x, z_f)} g^{\tau\tau}(x, z_f) = 1 \implies g^{\tau\tau}(x, z_f) = \frac{m^2}{m^2(z_f) - 2t(z_f)} \tag{4.7}$$

for the $g^{\tau\tau}(x, z_f)$ component. We recall the gauge choice

$$g^{zz}(x, z) = 1. \tag{4.8}$$

We also point out the correspondence in the Green's function, given by

$$G[q_0, q_x; t(z), m(z)] = \frac{1}{q_0^2 + t(z)q_x^2 + \frac{m^2(z) - 2t(z)}{2dz}}, \tag{4.9}$$

where q_0 and q_x are temporal and spatial components of the momentum, respectively. Introducing all these correspondences into the evolution equations for the metric tensors, given by

$$\partial_z g^{\tau\tau}(z) = g^{\tau\tau}(z) \left\{ \int dq_0 dq_x \frac{q_0^2}{q_0^2 + t(z)q_x^2 + \frac{m^2(z) - 2t(z)}{2dz}} \right\} g^{\tau\tau}(z) \tag{4.10}$$

and

$$\partial_z g^{xx}(z) = g^{xx}(z) \left\{ \int dq_0 dq_x \frac{q_x^2}{q_0^2 + t(z)q_x^2 + \frac{m^2(z) - 2t(z)}{2dz}} \right\} g^{xx}(z), \tag{4.11}$$

we obtain the following equations for both mass and hopping parameters,

$$\partial_z m^2(z) - 2\partial_z t(z) = -\pi \frac{m^2}{2dz} \left(\int dq_0 \frac{q_0^2}{\sqrt{q_0^2 + 1}} \right) \frac{m^2(z) - 2t(z)}{\sqrt{t(z)}} \quad (4.12)$$

and

$$m^2(z)\partial_z t(z) - t(z)\partial_z m^2(z) = \pi \frac{m^2}{2dz} \left(\int dq_x \frac{q_x^2}{\sqrt{q_x^2 + 1}} \right) \sqrt{t(z)} [m^2(z) - 2t(z)]. \quad (4.13)$$

These two coupled equations are reduced into

$$\partial_z m^2(z) = -\pi \frac{m^2}{2dz} \left(\int dq_0 \frac{q_0^2}{\sqrt{q_0^2 + 1}} \right) \frac{m^2(z) - 2t(z)}{\sqrt{t(z)}} \quad (4.14)$$

and

$$\partial_z t(z) = 0. \quad (4.15)$$

Redefining the mass parameter as

$$M^2(z) \equiv m^2(z) + 2t(z), \quad (4.16)$$

we rewrite Eq. (4.14) as

$$\partial_z M^2(z) = -\pi \frac{m^2}{2dz\sqrt{t}} \left(\int dq_0 \frac{q_0^2}{\sqrt{q_0^2 + 1}} \right) M^2(z). \quad (4.17)$$

To show the equivalence between this equation and the β -function of the mass parameter, we consider the following fixed point, given by

$$\partial_z t(z) = -\frac{1}{a} t(z) + \frac{[t(z)]^2}{2am^2(z)} \implies 0 \quad (4.18)$$

in the β -function of the hopping parameter. Then, the β -function of the mass parameter is

$$\partial_z m^2(z) = -\frac{[t(z)]^2}{am^2(z)} \implies -\frac{4}{a} m^2(z). \quad (4.19)$$

Comparing Eq. (4.17) with Eq. (4.19), we obtain the correspondence between the scale dz of the Polchinski real-space renormalization group transformation and the scale a of the Kadanoff block-spin transformation as follows

$$\frac{2dz}{a} = \pi \frac{m^2}{4\sqrt{t}} \left(\int dq_0 \frac{q_0^2}{\sqrt{q_0^2 + 1}} \right). \quad (4.20)$$

This completes our demonstration that the evolution equations of the metric tensor are nothing but the renormalization-group β -functions of the coupling constants.

5 Evaluation of entanglement entropy based on the heat kernel method

5.1 Entanglement entropy

As we have demonstrated that the quantum field theory is geometrized in the large N limit, we now ask how quantum entanglement in the quantum field theory is encoded into the classical geometry in the holographic dual reformulation. As discussed before, we address this issue by calculating the entanglement entropy.

Entanglement entropy is a subsystem von Neumann entropy $\mathcal{S}_{EE} = -\text{tr}_A \rho_A \ln \rho_A$, given by a reduced density matrix of the subsystem $\rho_A = \text{tr}_B \rho_{A \cup B}$, where $\rho_{A \cup B}$ is the density matrix of a total system $A \cup B$. This von Neumann entropy can be obtained from the Renyi entropy $\mathcal{S}_{RE} = \frac{1}{1-n} \text{tr}_A \rho_A^n$ as follows [34–39]

$$\begin{aligned} \mathcal{S}_{EE}(z_f) &= \lim_{n \rightarrow 1} \frac{1}{1-n} \text{tr}_A \rho_A^n = \lim_{n \rightarrow 1} \frac{\ln Z_n(z_f) - n \ln Z_1(z_f)}{1-n} \\ &= -\frac{[\partial_n Z_n(z_f)]_{n=1}}{Z_1(z_f)} + \ln Z_1(z_f). \end{aligned} \quad (5.1)$$

Here, $Z_1(z_f)$ is the partition function of the emergent holographic dual description with an IR cutoff z_f , defined in the Riemann geometry without a conical singularity and given by the D -dimensional metric of $ds^2 = dr^2 + r^2 d\theta^2 + \delta_{ij} dx_\perp^i dx_\perp^j$ with i & $j = 2, \dots, D-1$. For simplicity, we do not consider the self-interactions in the channel of energy-momentum tensor-currents. Then, the partition function is given by

$$\begin{aligned} Z_1(z_f) &= \int D\phi_\alpha(x) D\varphi(x, z) Dg_{\mu\nu}(x, z) \delta(g^{\mu\nu}(x, 0) - g_B^{\mu\nu}) \\ &\quad \times \delta\left(\partial_z g^{\mu\nu}(x, z) - \frac{1}{2dz} g^{\mu\nu'}(x, z) (\partial_{\nu'} \partial_{\mu'} G_{xx'}[g_{\mu\nu}(x, z), \varphi(x, z)])_{x' \rightarrow x} g^{\mu'\nu}(x, z)\right) \\ &\quad \times \exp\left[-\int d^D x \sqrt{g(x, z_f)} \left\{ g^{\mu\nu}(x, z_f) [\partial_\mu \phi_\alpha(x)] [\partial_\nu \phi_\alpha(x)] + [m^2 - i\varphi(x, z_f)] \phi_\alpha^2(x) \right\}\right. \\ &\quad \left. - N \int d^D x \sqrt{g(x, 0)} \left\{ \frac{1}{2u} (\varphi(x, 0) - i\xi R(x, 0))^2 \right\}\right. \\ &\quad \left. - N \int_0^{z_f} dz \int d^D x \sqrt{g(x, z)} \left\{ \right. \right. \\ &\quad \left. \left. + \frac{1}{2u} [\partial_z \varphi(x, z)]^2 + \frac{\mathcal{C}_\varphi}{2} g^{\mu\nu}(x, z) [\partial_\mu \varphi(x, z)] [\partial_\nu \varphi(x, z)] + \mathcal{C}_\xi R(x, z) [\varphi(x, z)]^2 \right. \right. \\ &\quad \left. \left. + \frac{1}{2\kappa} (R(x, z) - 2\Lambda) \right\} \right]. \end{aligned} \quad (5.2)$$

On the other hand, $Z_n(z_f)$ is the partition function of the emergent holographic dual description with an IR cutoff z_f , defined in the Riemann geometry with a conical singularity and given by the D -dimensional metric of $ds^2 = dr^2 + n^2 r^2 d\theta^2 + \delta_{ij} dx_\perp^i dx_\perp^j$ with i & $j =$

$2, \dots, D-1$ [47]. It is given by

$$\begin{aligned}
Z_n(z_f) &= \int D\phi_\alpha(x) D\varphi_n(x, z) Dg_{\mu\nu}^n(x, z) \delta\left(g_n^{\mu\nu}(x, 0) - g_{Bn}^{\mu\nu}\right) \\
&\times \delta\left(\partial_z g_n^{\mu\nu}(x, z) - \frac{1}{2dz} g_n^{\mu\nu'}(x, z) (\partial_{\nu'} \partial_{\mu'} G_{xx'}^n[g_n^{\mu\nu}(x, z), \varphi_n(x, z)])_{x' \rightarrow x} g_n^{\mu'\nu}(x, z)\right) \\
&\times \exp\left[-\int d^D x \sqrt{g_n(x, z_f)} \left\{ g_n^{\mu\nu}(x, z_f) [\partial_\mu \phi_\alpha(x)] [\partial_\nu \phi_\alpha(x)] + [m^2 - i\varphi_n(x, z_f)] \phi_\alpha^2(x) \right\}\right. \\
&\quad - N \int d^D x \sqrt{g_n(x, 0)} \left\{ \frac{1}{2u} \left(\varphi_n(x, 0) - i\xi R_n(x, 0) \right)^2 \right\} \\
&\quad - N \int_0^{z_f} dz \int d^D x \sqrt{g_n(x, z)} \left\{ \right. \\
&\quad \quad + \frac{1}{2u} [\partial_z \varphi_n(x, z)]^2 + \frac{\mathcal{C}_\varphi}{2} g_n^{\mu\nu}(x, z) [\partial_\mu \varphi_n(x, z)] [\partial_\nu \varphi_n(x, z)] + \mathcal{C}_\xi R_n(x, z) [\varphi_n(x, z)]^2 \\
&\quad \quad \left. + \frac{1}{2\kappa} \left(R_n(x, z) - 2\Lambda \right) \right\}. \tag{5.3}
\end{aligned}$$

Following this prescription, we find that the entanglement entropy is composed of two pieces, given by

$$\mathcal{S}_{EE}(z_f) = \mathcal{S}_{EE}^{\phi_\alpha}(z_f) + \mathcal{S}_{EE}^{GR}(z_f), \tag{5.4}$$

where $\mathcal{S}_{EE}^{\phi_\alpha}(z_f)$ is the entanglement entropy from the matter sector at the IR cutoff $z = z_f$ and $\mathcal{S}_{EE}^{GR}(z_f)$ is that from the emergent geometry also at the IR cutoff $z = z_f$.

It is straightforward to see that the matter contribution is [52]

$$\begin{aligned}
\mathcal{S}_{EE}^{\phi_\alpha}(z_f) &= \int d^{D-2} x_\perp \int_0^{2\pi} d\theta \int_0^\infty dr \sqrt{g(r, x_\perp, z_f)} \left\langle T_{\mu\nu}^{\phi_\alpha}(r, x_\perp, z_f) \right\rangle \frac{\partial g_n^{\mu\nu}(r, x_\perp, z_f)}{\partial n} \Big|_{n=1} \\
&\quad - \frac{N}{2} \text{tr}_{xx'} \ln \sqrt{g(x, z_f)} \left\{ -\frac{1}{\sqrt{g(x, z_f)}} \partial_\mu \left(\sqrt{g(x, z_f)} g^{\mu\nu}(x, z_f) \partial_\nu \right) + m^2 - i\varphi(x, z_f) \right\}. \tag{5.5}
\end{aligned}$$

Here, the energy-momentum tensor is given by

$$\begin{aligned}
T_{\mu\nu}^{\phi_\alpha}(x, z_f) &\equiv \frac{1}{\sqrt{g(x, z_f)}} \frac{\partial \sqrt{g(x, z_f)} \mathcal{L}_{\phi_\alpha}[g_{\mu\nu}(x, z_f), \varphi(x, z_f)]}{\partial g^{\mu\nu}(x, z_f)} \\
&= -2(\partial_\mu \phi_\alpha(x)) (\partial_\nu \phi_\alpha(x)) \\
&\quad + g_{\mu\nu}(x, z_f) \left(g^{\mu'\nu'}(x, z_f) (\partial_{\mu'} \phi_\alpha(x)) (\partial_{\nu'} \phi_\alpha(x)) + [m^2 - i\varphi(x, z_f)] \phi_\alpha^2(x) \right), \tag{5.6}
\end{aligned}$$

where the effective Lagrangian for the matter sector is $\mathcal{L}_{\phi_\alpha}[g_{\mu\nu}(x, z_f), \varphi(x, z_f)] = g^{\mu\nu}(x, z_f) [\partial_\mu \phi_\alpha(x)] [\partial_\nu \phi_\alpha(x)] + [m^2 - i\varphi(x, z_f)] \phi_\alpha^2(x)$. Inserting this energy-momentum tensor into the above expression,

we obtain

$$\begin{aligned}
\mathcal{S}_{EE}^{\phi_\alpha}(z_f) = & N \int d^{D-2}x_\perp \int_0^{2\pi} d\theta \int_0^\infty dr \sqrt{g(r, x_\perp, z_f)} \left\{ -2(\partial_\mu \partial_\nu G[x, x; g_{\mu\nu}(x, z_f), \varphi(x, z_f)]) \right. \\
& + g_{\mu\nu}(r, x_\perp, z_f) \left(g^{\mu'\nu'}(r, x_\perp, z_f) (\partial_{\mu'} \partial_{\nu'} G[x, x; g_{\mu\nu}(x, z_f), \varphi(x, z_f)]) \right. \\
& \left. \left. + [m^2 - i\varphi(x, z_f)] G[x, x; g_{\mu\nu}(x, z_f), \varphi(x, z_f)] \right) \right\} \frac{\partial g_n^{\mu\nu}(r, x_\perp, z_f)}{\partial n} \Big|_{n=1} \\
& - N \int d^{D-2}x_\perp \int_0^{2\pi} d\theta \int_0^\infty dr \sqrt{g(r, x_\perp, z_f)} \left\{ \frac{\mathcal{C}_\varphi^f}{2} g^{\mu\nu}(r, x_\perp, z_f) [\partial_\mu \varphi(r, x_\perp, z_f)] [\partial_\nu \varphi(r, x_\perp, z_f)] \right. \\
& \left. + \mathcal{C}_\xi^f R(r, x_\perp, z_f) [\varphi(r, x_\perp, z_f)]^2 + \frac{1}{2\kappa_f} \left(R(r, x_\perp, z_f) - 2\Lambda_f \right) \right\}, \tag{5.7}
\end{aligned}$$

where $G[x, x'; g_{\mu\nu}(x, z_f), \varphi(x, z_f)] = \left\langle \frac{1}{N} \sum_{\alpha=1}^N \phi_\alpha(x) \phi_\alpha(x') \right\rangle$ is the Green's function of the original scalar fields at $z = z_f$. Both the Klein-Gordon action and the induced Einstein-Hilbert gravity action at $z = z_f$ in the last two lines result from the gradient expansion of the last logarithmic term [43, 44] in Eq. (5.5).

The entanglement entropy from the classical geometry in the presence of dual scalar fields is given by

$$\begin{aligned}
\mathcal{S}_{EE}^{GR}(z_f) = & N \int d^{D-2}x_\perp \int_0^{2\pi} d\theta \int_0^\infty dr \int_0^{z_f} dz \sqrt{g(r, x_\perp, z)} T_{\mu\nu}^{GR}(r, x_\perp, z) \frac{\partial g_n^{\mu\nu}(r, x_\perp, z)}{\partial n} \Big|_{n=1} \\
& - N \int d^{D-2}x_\perp \int_0^{2\pi} d\theta \int_0^\infty dr \int_0^{z_f} dz \sqrt{g(r, x_\perp, z)} \left\{ \frac{1}{2u} [\partial_z \varphi(r, x_\perp, z)]^2 \right. \\
& + \frac{\mathcal{C}_\varphi}{2} g^{\mu\nu}(r, x_\perp, z) [\partial_\mu \varphi(r, x_\perp, z)] [\partial_\nu \varphi(r, x_\perp, z)] + \mathcal{C}_\xi R(r, x_\perp, z) [\varphi(r, x_\perp, z)]^2 \\
& \left. + \frac{1}{2\kappa} \left(R(r, x_\perp, z) - 2\Lambda \right) \right\}. \tag{5.8}
\end{aligned}$$

The energy-momentum tensor of the gravity sector is

$$T_{\mu\nu}^{GR}(x, z) \equiv \frac{1}{\sqrt{g(x, z)}} \frac{\partial \sqrt{g(x, z)} \mathcal{L}_{GR}[g_{\mu\nu}(x, z), \varphi(x, z)]}{\partial g^{\mu\nu}(x, z)} \equiv T_{\mu\nu}^{g\mu\nu}(x, z) + T_{\mu\nu}^\varphi(x, z), \tag{5.9}$$

where the bulk Lagrangian in terms of bulk metric tensor and dual scalar fields is given by

$$\begin{aligned}
\mathcal{L}_{GR}[g_{\mu\nu}(x, z), \varphi(x, z)] = & \frac{1}{2u} [\partial_z \varphi(x, z)]^2 + \frac{\mathcal{C}_\varphi}{2} g^{\mu\nu}(x, z) [\partial_\mu \varphi(x, z)] [\partial_\nu \varphi(x, z)] + \mathcal{C}_\xi R(x, z) [\varphi(x, z)]^2 \\
& + \frac{1}{2\kappa} \left(R(x, z) - 2\Lambda \right). \tag{5.10}
\end{aligned}$$

As a result, the energy-momentum tensor of our classical gravity is [53]

$$\begin{aligned}
2\kappa T_{\mu\nu}^{g\mu\nu}(x, z) = & -2 \left(R_{\mu\nu}(x, z) - \frac{1}{2} g_{\mu\nu}(x, z) R(x, z) + \Lambda g_{\mu\nu}(x, z) \right) \\
& + \frac{1}{g(x, z)} g_{\mu\mu'}(x, z) g_{\nu\nu'}(x, z) \partial_{\mu''} \partial_{\nu''} \left\{ g(x, z) \left(g^{\mu'\nu'}(x, z) g^{\mu''\nu''}(x, z) - g^{\mu'\mu''}(x, z) g^{\nu'\nu''}(x, z) \right) \right\}, \tag{5.11}
\end{aligned}$$

and that of the dual scalar field is

$$\begin{aligned}
T_{\mu\nu}^\varphi(x, z) = & -\frac{2}{u}\delta_{\mu z}\delta_{\nu z}[\partial_z\varphi(x, z)]^2 - 2(1 - \delta_{\mu z}\delta_{\nu z})\mathcal{C}_\varphi[\partial_\mu\varphi(x, z)][\partial_\nu\varphi(x, z)] \\
& + g_{\mu\nu}(x, z)\left(\frac{1}{u}[\partial_z\varphi(x, z)]^2 + \mathcal{C}_\varphi g^{\mu'\nu'}(x, z)[\partial_{\mu'}\varphi(x, z)][\partial_{\nu'}\varphi(x, z)]\right) \\
& - 2\mathcal{C}_\xi\left(R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R\right)[\varphi(x, z)]^2 + 2\mathcal{C}_\xi\left(g_{\mu\nu}\partial_\rho\partial^\rho - \partial_\mu\partial_\nu\right)[\varphi(x, z)]^2. \quad (5.12)
\end{aligned}$$

5.2 Heat kernel method for entanglement entropy

To show the entanglement transfer from matter to geometry through recursive renormalization group transformations, we need to simplify the entanglement-entropy formula of the matter sector. Here, we review the entanglement-entropy formula based on the heat kernel method [47]. An effective free energy on n -sheeted Riemann with a conical singularity can be represented by

$$-\ln Z_n^{\phi_\alpha}(z_f) = -\frac{N}{2} \int_{\epsilon^2}^{\infty} \frac{ds}{s} \text{tr}_{xx'} K(s, x, x'; z_f), \quad (5.13)$$

where $K(s, x, x'; z_f) \equiv \langle xz_f | e^{-s\mathcal{D}} | x'z_f \rangle$ is the corresponding heat kernel with the differential operator $\mathcal{D} \equiv -\frac{1}{\sqrt{g(x, z_f)}}\partial_\mu\left(\sqrt{g(x, z_f)}g^{\mu\nu}(x, z_f)\partial_\nu\right) + m^2$. For the time being, we consider the case of a constant mass, given by $m^2 - i\varphi(x, z_f) \implies m^2$. This heat kernel satisfies an effective heat-diffusion equation $(\partial_s + \mathcal{D})K(s, x, x'; z_f) = 0$ with an initial condition $K(s=0, x, x'; z_f) = \frac{1}{\sqrt{g(x, z_f)}}\delta^{(D)}(x - x')$ – this is the origin of the name of the heat kernel. ϵ is a UV cutoff to cure the UV divergence of the entanglement entropy, which will be clarified below.

To deal with curvature integrals in the presence of a conical singularity, we need to understand the curvature structure of this geometry [47]. Let E_α be an α -fold Riemann sheet, which covers a smooth manifold E along the Killing vector ∂_θ . A codimension two surface Σ is a stationary point ($r = 0$) of this isometry. Then, the space E_α is a direct product near Σ ($r = 0$), given by $\Sigma \times C_\alpha$, where C_α is a two-dimensional cone with angle deficit $\delta = 2\pi(1 - \alpha)$. In other words, the metric of E_α is $ds_{E_\alpha}^2 = ds_{C_\alpha}^2 + ds_\Sigma^2$, where the metric of the two dimensional cone is $ds_{C_\alpha}^2 = g_{rr}^\alpha(r, x_\perp, z)dr^2 + g_{\theta\theta}^\alpha(r, x_\perp, z)d\theta^2$ and the metric of the codimension two surface is $ds_\Sigma^2 = \gamma_{ij}(r, x_\perp, z)dx_\perp^i dx_\perp^j$ at a fixed renormalization group scale z . Outside this singular surface Σ , E_α is reduced to the smooth manifold E , where their curvature tensors coincide. On the other hand, the conical singularity at the surface Σ gives rise to a singular contribution to the curvatures. Here, the positive integer n is analytically continued to a real value α , which results from the abelian isometry generated by the Killing vector ∂_θ .

One may take an expansion for small s in the heat kernel, given by

$$\text{tr}_{xx'} K_{E_\alpha}(s) = \frac{1}{(4\pi s)^{\frac{d}{2}}} \sum_{n=0} a_n s^n. \quad (5.14)$$

Then, we have two types of coefficients in this expansion,

$$a_n = a_n^{reg} + a_n^\Sigma, \quad (5.15)$$

where a_n^{reg} is a regular contribution resulting from the bulk and a_n^Σ is a singular one coming from the surface. Regular bulk coefficients are given by

$$\begin{aligned}
a_0^{reg} &= \int_E d^D x \sqrt{g(x, z_f)}, \\
a_1^{reg} &= \int_E d^D x \sqrt{g(x, z_f)} \left(\frac{1}{6} R(x, z_f) - m^2 \right), \\
a_2^{reg} &= \int_E d^D x \sqrt{g(x, z_f)} \left\{ \frac{1}{180} R_{\mu\nu\alpha\beta}^2(x, z_f) - \frac{1}{180} R_{\mu\nu}^2(x, z_f) \right. \\
&\quad \left. + \frac{1}{6} \frac{1}{\sqrt{g(x, z_f)}} \partial_\mu \left(\sqrt{g(x, z_f)} g^{\mu\nu}(x, z_f) \partial_\nu \right) \left(\frac{1}{5} R(x, z_f) - m^2 \right) + \frac{1}{2} \left(\frac{1}{6} R(x, z_f) - m^2 \right)^2 \right\},
\end{aligned} \tag{5.16}$$

for general curvature tensors [47]. On the other hand, singular surface coefficients are given by [47]

$$\begin{aligned}
a_0^\Sigma &= 0, \\
a_1^\Sigma &= \frac{\pi}{3} \frac{(1-\alpha)(1+\alpha)}{\alpha} \int_{\Sigma(z_f)} d^{D-2} x_\perp \sqrt{\gamma(x, z_f)}, \\
a_2^\Sigma &= \frac{\pi}{3} \frac{(1-\alpha)(1+\alpha)}{\alpha} \int_{\Sigma(z_f)} d^{D-2} x_\perp \sqrt{\gamma(x, z_f)} \left(\frac{1}{6} R(x, z_f) - m^2 \right) \\
&\quad - \frac{\pi}{180} \frac{(1-\alpha)(1+\alpha)(1+\alpha^2)}{\alpha^3} \int_{\Sigma(z_f)} d^{D-2} x_\perp \sqrt{\gamma(x, z_f)} \left(R_{ii}(x, z_f) - 2R_{ijij}(x, z_f) \right).
\end{aligned} \tag{5.17}$$

We recall $ds_\Sigma^2 = \gamma_{ij}(r, x_\perp, z) dx_\perp^i dx_\perp^j$.

Based on this general curvature formula from the heat kernel expansion, we obtain the following expression of the entanglement entropy for the matter sector

$$\begin{aligned}
\mathcal{S}_{EE}^{\phi_\alpha}(z_f) &= (\alpha \partial_\alpha - 1) [-\ln Z_{\phi_\alpha}(\alpha; z_f)] \Big|_{\alpha=1} \\
&\approx -N \frac{\pi}{6} \frac{1}{(4\pi)^{\frac{D}{2}}} \left\{ (\alpha \partial_\alpha - 1) \frac{(1-\alpha)(1+\alpha)}{\alpha} \right\}_{\alpha=1} \left(\int_{\epsilon^2}^\infty \frac{ds}{s} s^{1-\frac{D}{2}} \right) \left\{ \int_{\Sigma(z_f)} d^{D-2} x_\perp \sqrt{\gamma(x, z_f)} \right\}.
\end{aligned} \tag{5.18}$$

This leading contribution results from the a_1^Σ term in Eq. (5.17). Considering that the area of the singular surface $\Sigma(z_f)$ is given by

$$\mathcal{A}[\Sigma(z_f)] = \int_{\Sigma(z_f)} d^{D-2} x_\perp \sqrt{\gamma(x, z_f)} \tag{5.19}$$

at the renormalization group scale $z = z_f$, we obtain the area law for the entanglement entropy of the matter sector

$$\mathcal{S}_{EE}^{\phi_\alpha}(z_f) \approx \frac{N}{6(D-2)(4\pi)^{\frac{D}{2}-1}} \frac{\mathcal{A}[\Sigma(z_f)]}{\epsilon^{D-2}}. \tag{5.20}$$

It is straightforward to see that this formula is reduced to the well known result in two spacetime dimensions, given by

$$\mathcal{S}_{EE}^{\phi_\alpha}(z_f) = \frac{N}{6} \ln \frac{\xi}{\epsilon}. \quad (5.21)$$

Here, ξ is the correlation length proportional to the inverse of the mass parameter m , which appears from the introduction of an IR cutoff in the s -integral, given by $\int_{\epsilon^2}^{\xi^2} \frac{ds}{s}$.

Now, we introduce inhomogeneity of the mass term into the above heat-kernel expansion. Then, spacetime derivatives for $\varphi(x, z_f)$ will appear in both expansion coefficients. As far as a disordered phase is concerned, we speculate that this area law would be preserved. It is an interesting question how $\partial_\mu \varphi(x, z_f)$ modify the entanglement entropy of the matter sector at quantum criticality. In this paper, we continue our discussions for the gapped phase.

Finally, we obtain the total entanglement entropy at $z = z_f$

$$\begin{aligned} \mathcal{S}_{EE}(z_f) &= \mathcal{S}_{EE}^{\phi_\alpha}(z_f) + \mathcal{S}_{EE}^{GR}(z_f) \\ &= \frac{N}{6(D-2)(4\pi)^{\frac{D}{2}-1}} \frac{\mathcal{A}[\Sigma(z_f)]}{\epsilon^{D-2}} \\ &\quad + N \int d^{D-2}x_\perp \int_0^{2\pi} d\theta \int_0^\infty dr \int_0^{z_f} dz \sqrt{g(r, x_\perp, z)} \\ &\quad \times \left(T_{\mu\nu}^\varphi(r, x_\perp, z) + T_{\mu\nu}^{g\mu\nu}(r, x_\perp, z) \right) \frac{\partial g_n^{\mu\nu}(r, x_\perp, z)}{\partial n} \Big|_{n=1} \\ &\quad - N \int d^{D-2}x_\perp \int_0^{2\pi} d\theta \int_0^\infty dr \int_0^{z_f} dz \sqrt{g(r, x_\perp, z)} \left\{ \frac{1}{2u} [\partial_z \varphi(r, x_\perp, z)]^2 \right. \\ &\quad + \frac{\mathcal{C}_\varphi}{2} g^{\mu\nu}(r, x_\perp, z) [\partial_\mu \varphi(r, x_\perp, z)] [\partial_\nu \varphi(r, x_\perp, z)] + \mathcal{C}_\xi R(r, x_\perp, z) [\varphi(r, x_\perp, z)]^2 \\ &\quad \left. + \frac{1}{2\kappa} \left(R(r, x_\perp, z) - 2\Lambda \right) \right\}. \end{aligned} \quad (5.22)$$

In this formula we focus on the fact that the entanglement entropy of the matter sector is given by the area of the subsystem at a fixed renormalization group scale $z = z_f$ while that of the classical gravity part is given by the integral of the extra dimensional space. If the emergent geometry is given by a cap in the extra dimensional space, the area of the subsystem at a given $z = z_f$ vanishes beyond the cap in the extra dimensional space. Instead, the entanglement entropy from the classical geometry is maximized. Actually, the cap geometry has been realized when the corresponding quantum state is gapped [33, 48, 49]. Even if the resulting geometry is given by AdS_{D+1} , the entanglement entropy is transferred from quantum matter to classical gravity.

5.3 Invariance of entanglement entropy with respect to renormalization group transformations

Since the partition function (5.3) has to be invariant with respect to recursive renormalization group transformations, we obtain the following equation

$$\partial_{z_f} \mathcal{S}_{EE}(z_f) = \partial_{z_f} \mathcal{S}_{EE}^{\phi_\alpha}(z_f) + \partial_{z_f} \mathcal{S}_{EE}^{g\mu\nu}(z_f) = 0 \implies \mathcal{S}_{EE}(z_f) = \mathcal{S}_{EE}^{\phi_\alpha}(0). \quad (5.23)$$

In other words, the entanglement entropy is an invariant for renormalization group transformations. Resorting to the full expression Eq. (5.22) of the entanglement entropy, we reach the following identity

$$\begin{aligned}
& \frac{N}{6(D-2)(4\pi)^{\frac{D}{2}-1}} \frac{\partial_{z_f} \mathcal{A}[\Sigma(z_f)]}{\epsilon^{D-2}} \\
& + N \int d^{D-2} x_{\perp} \int_0^{2\pi} d\theta \int_0^{\infty} dr \sqrt{g(r, x_{\perp}, z_f)} \left(T_{\mu\nu}^{\varphi}(r, x_{\perp}, z_f) + T_{\mu\nu}^{g\mu\nu}(r, x_{\perp}, z_f) \right) \frac{\partial g_n^{\mu\nu}(r, x_{\perp}, z_f)}{\partial n} \Big|_{n=1} \\
& - N \int d^{D-2} x_{\perp} \int_0^{2\pi} d\theta \int_0^{\infty} dr \sqrt{g(r, x_{\perp}, z_f)} \left\{ \frac{1}{2u} [\partial_z \varphi(r, x_{\perp}, z_f)]^2 \right. \\
& \quad + \frac{\mathcal{C}_{\varphi}}{2} g^{\mu\nu}(r, x_{\perp}, z_f) [\partial_{\mu} \varphi(r, x_{\perp}, z_f)] [\partial_{\nu} \varphi(r, x_{\perp}, z_f)] + \mathcal{C}_{\xi} R(r, x_{\perp}, z_f) [\varphi(r, x_{\perp}, z_f)]^2 \\
& \quad \left. + \frac{1}{2\kappa} (R(r, x_{\perp}, z_f) - 2\Lambda) \right\} = 0. \tag{5.24}
\end{aligned}$$

We suspect that the bulk gravity contribution may be related with the holographic entanglement entropy of the Ryu-Takayanagi formula [36, 37]. In particular, one may express this identity in terms of fully geometric information such as metric and curvature. In appendix B, we discuss this identity further and verify the first iteration version of this identity.

6 Discussion

One may suggest an effective higher spin gauge theory as a dual holographic description of the $O(N)$ vector model, which has an infinite tower of higher-spin fields without hierarchy in the spectrum [54–57]. Actually, we believe that this speculation would be realized if the $O(N)$ symmetry is taken into account appropriately. An idea how to introduce the role of global symmetry into the present dual holographic description is turning on an external gauge field which couples to the $O(N)$ –symmetry conserved current. Here, the question is how the internal or flavor symmetry can be uplifted into the gauge symmetry with higher spins. Mixing between the internal symmetric space and the curved spacetime coordinate has to occur for the higher spin gauge theory to be realized during the renormalization group transformation. Indeed, the renormalization group transformation results in the mixing between the flavor index and the spacetime coordinate.

We start from an $O(N)$ vector model in a background curved spacetime metric $g_{B\mu\nu}$,

$$\begin{aligned}
Z = \int D\phi_{\alpha} \exp \left[- \int d^D x \sqrt{g_B} \left\{ g_B^{\mu\nu} [\partial_{\mu} \phi_{\alpha} - (q/\sqrt{N}) A_{\mu\alpha\beta}^{ext} \phi_{\beta}] [\partial_{\nu} \phi_{\alpha} - (q/\sqrt{N}) A_{\nu\alpha\gamma}^{ext} \phi_{\gamma}] \right. \right. \\
\left. \left. + m^2 \phi_{\alpha}^2 + \xi R_B \phi_{\alpha}^2 + \frac{u}{2N} \phi_{\alpha}^2 \phi_{\beta}^2 \right\} \right], \tag{6.1}
\end{aligned}$$

where the background $O(N)$ gauge field has been introduced with an $O(N)$ gauge charge q . Under the local $O(N)$ transformation for the matter field given by $\phi_{\alpha} \Rightarrow O_{\alpha\beta} \phi_{\beta}$ with $O_{\alpha\beta} \in O(N)$, the background $O(N)$ gauge field transforms as follows

$$A_{\mu\alpha\beta}^{ext} \Rightarrow O_{\alpha\gamma} A_{\mu\gamma\delta}^{ext} O_{\delta\beta}^T + (q/\sqrt{N})^{-1} (\partial_{\mu} O_{\alpha\gamma}) O_{\gamma\beta}^T, \tag{6.2}$$

where the classical action remains invariant.

Together with the Hubbard-Stratonovich transformation to decompose the self-interaction term, we rewrite the above path integral expression of the partition function in the following way

$$\begin{aligned}
Z &= \int D\phi_\alpha D\varphi^{(0)} Dg_{\mu\nu}^{(0)} Dt_{\mu\nu}^{(0)} DA_{\mu\alpha\beta}^{(0)} DJ_{\mu\alpha\beta}^{(0)} \\
&\exp \left[- \int d^D x \sqrt{g^{(0)}} \left\{ g^{\mu\nu(0)} [\partial_\mu \phi_\alpha - (q/\sqrt{N}) A_{\mu\alpha\beta}^{(0)} \phi_\beta] [\partial_\nu \phi_\alpha - (q/\sqrt{N}) A_{\nu\alpha\gamma}^{(0)} \phi_\gamma] \right. \right. \\
&\left. \left. + (m^2 - i\varphi^{(0)}) \phi_\alpha^2 + \xi R^{(0)} \phi_\alpha^2 + \frac{N}{2u} \varphi^{(0)2} - N t_{\mu\nu}^{(0)} (g^{\mu\nu(0)} - g_B^{\mu\nu}) - \frac{1}{N} J_{\alpha\beta}^{\mu(0)} (A_{\mu\beta\alpha}^{(0)} - A_{\mu\beta\alpha}^{ext}) \right\} \right]. \quad (6.3)
\end{aligned}$$

$t_{\mu\nu}^{(0)}$ and $J_{\alpha\beta}^{\mu(0)}$ are Lagrange multiplier fields to impose two initial conditions of $g^{\mu\nu(0)} = g_B^{\mu\nu}$ and $A_{\mu\beta\alpha}^{(0)} = A_{\mu\beta\alpha}^{ext}$, respectively.

Repeating exactly the same renormalization group transformation as discussed in section 3, we obtain the following expression of the partition function

$$\begin{aligned}
Z &= \int D\phi_\alpha D\varphi^{(0)} D\varphi^{(1)} Dg_{\mu\nu}^{(0)} Dt_{\mu\nu}^{(0)} DA_{\mu\alpha\beta}^{(0)} DJ_{\mu\alpha\beta}^{(0)} \exp \left[- \int d^D x \sqrt{g^{(0)}} \left\{ \right. \right. \\
&\left. \left. [(\partial_\mu \mathbf{I} - (q/\sqrt{N}) \mathbf{A}_\mu^{(0)}) \phi]_\alpha \left(g^{\mu\nu(0)} \mathbf{I}_{\alpha\beta} - g^{\mu\gamma(0)} [(\partial_\gamma \mathbf{I} - (q/\sqrt{N}) \mathbf{A}_{\gamma x}^{(0)})_{\alpha\alpha'} \right. \right. \right. \\
&\quad \left. \left. \times (\partial_{\gamma'} \mathbf{I} - (q/\sqrt{N}) \mathbf{A}_{\gamma' x'}^{(0)})_{\alpha' \beta'} G_{\beta' \beta x x'}^{(0)} \right]_{x' \rightarrow x} g^{\gamma' \nu(0)} \right) [(\partial_\nu \mathbf{I} - (q/\sqrt{N}) \mathbf{A}_\nu^{(0)}) \phi]_\beta \\
&\left. + (m^2 - i\varphi_x^{(1)}) \phi_{\alpha x}^2 + \frac{N}{2u} \left(\varphi^{(0)} - i e^{-\beta(0) dz} \xi R^{(0)} \right)^2 + 2\beta^{(0)} dz \frac{N}{2u} \left(\frac{\varphi^{(1)} - \varphi^{(0)}}{2\beta^{(0)} dz} \right)^2 \right. \\
&\left. + 2\alpha^{(0)} dz N \left(-\mathcal{C}_\Lambda + \mathcal{C}_R R^{(0)} + \frac{\mathcal{C}_\varphi}{2} g^{\mu\nu(0)} (\partial_\mu \varphi^{(1)}) (\partial_\nu \varphi^{(1)}) + \mathcal{C}_\xi R^{(0)} \varphi^{(1)2} + \frac{q^2}{N^2} \mathcal{C}_F \mathbf{F}_{\mu\nu}^{(0)} \mathbf{F}^{\mu\nu(0)} \right. \right. \\
&\quad \left. \left. + \frac{q}{N} \mathcal{C}_{mix} (\mathbf{F}^{\mu\nu(0)})_{\alpha\alpha} (\partial_\mu \varphi^{(1)}) (\partial_\nu \varphi^{(1)}) \right) \right. \\
&\left. - N t_{\mu\nu}^{(0)} (g^{\mu\nu(0)} - g_B^{\mu\nu}) - \frac{1}{N} J_{\alpha\beta}^{\mu(0)} (A_{\mu\alpha\beta}^{(0)} - A_{\mu\beta\alpha}^{ext}) \right\} \right]. \quad (6.4)
\end{aligned}$$

Here,

$$\begin{aligned}
&\int d^D x \sqrt{g^{(0)}} \left\{ 2\alpha^{(0)} dz N \left(-\mathcal{C}_\Lambda + \mathcal{C}_R R^{(0)} + \frac{\mathcal{C}_\varphi}{2} g^{\mu\nu(0)} (\partial_\mu \varphi^{(1)}) (\partial_\nu \varphi^{(1)}) + \mathcal{C}_\xi R^{(0)} \varphi^{(1)2} \right. \right. \\
&\left. \left. + \frac{q^2}{N^2} \mathcal{C}_F \mathbf{F}_{\mu\nu}^{(0)} \mathbf{F}^{\mu\nu(0)} + \frac{q}{N} \mathcal{C}_{mix} (\mathbf{F}^{\mu\nu(0)})_{\alpha\alpha} (\partial_\mu \varphi^{(1)}) (\partial_\nu \varphi^{(1)}) \right) \right\}
\end{aligned}$$

results from the gradient expansion of [40, 43, 44]

$$\begin{aligned}
&-\frac{N}{2} \text{tr}_{xx'} \ln \sqrt{g^{(0)}} \left\{ -\frac{1}{\sqrt{g^{(0)}}} [\partial_\mu \mathbf{I} - (q/\sqrt{N}) \mathbf{A}_\mu^{(0)}] \left(\sqrt{g^{(0)}} g^{\mu\nu(0)} [\partial_\nu \mathbf{I} - (q/\sqrt{N}) \mathbf{A}_\nu^{(0)}] \right) \right. \\
&\left. + \frac{(m^2 - i\varphi_x^{(1)})}{2\alpha^{(0)} dz} \mathbf{I} \right\},
\end{aligned}$$

given by the renormalization group transformation in the matter sector. The bold symbols denote either the vector or matrix of the $O(N)$ symmetric space. $\mathbf{F}^{\mu\nu(0)}$ is the field strength

tensor of the $O(N)$ gauge field. The Green's function is given by

$$\begin{aligned} & \left\{ -\frac{1}{\sqrt{g^{(0)}}} [\partial_\mu \mathbf{I} - (q/\sqrt{N}) \mathbf{A}_\mu^{(0)}] \left(\sqrt{g^{(0)}} g^{\mu\nu(0)} [\partial_\nu \mathbf{I} - (q/\sqrt{N}) \mathbf{A}_\nu^{(0)}] \right) + \frac{(m^2 - i\varphi^{(1)})}{2\alpha^{(0)} dz} \mathbf{I} \right\}_{\alpha\beta} G_{\beta\gamma x x'}^{(0)} \\ &= \frac{\delta_{\alpha\gamma}}{\sqrt{g^{(0)}}} \delta^{(D)}(x - x'). \end{aligned} \quad (6.5)$$

The upgrade rule has to be given by

$$\begin{aligned} & [(\partial_\mu \mathbf{I} - (q/\sqrt{N}) \mathbf{A}_\mu^{(0)}) \phi]_\alpha \left(g^{\mu\nu(0)} \mathbf{I}_{\alpha\beta} - g^{\mu\gamma(0)} [(\partial_\gamma \mathbf{I} - (q/\sqrt{N}) \mathbf{A}_{\gamma x}^{(0)})_{\alpha\alpha'} \right. \\ & \times (\partial_{\gamma'} \mathbf{I} - (q/\sqrt{N}) \mathbf{A}_{\gamma' x'}^{(0)})_{\alpha'\beta'} G_{\beta'\beta x x'}^{(0)}]_{x' \rightarrow x} g^{\gamma'\nu(0)} \left. \right) [(\partial_\nu \mathbf{I} - (q/\sqrt{N}) \mathbf{A}_\nu^{(0)}) \phi]_\beta \\ & \implies [(\partial_\mu \mathbf{I} - (q/\sqrt{N}) \mathbf{A}_\mu^{(1)}) \phi]_\alpha (g^{\mu\nu(1)})_{\alpha\beta} [(\partial_\nu \mathbf{I} - (q/\sqrt{N}) \mathbf{A}_\nu^{(1)}) \phi]_\beta \end{aligned} \quad (6.6)$$

for both $O(N)$ gauge field and metric tensor. An essential feature is that the $O(N)$ symmetric space be intertwined with the spacetime coordinate in order to define the renormalization group transformation appropriately. We believe that this mixing between spacetime and internal symmetric space is the physical origin for the emergence of an effective higher spin gauge theory as a dual holographic description of the $O(N)$ vector model. Since this issue has to be investigated more sincerely, we leave complete descriptions in the future research.

7 Conclusion

In the present study we tried to clarify the underlying physics of the dual holographic formulation for quantum field theories as a non-perturbative description. We uncovered how the emergent geometry takes into account the information of quantum entanglement. Let us recapitulate the essential features of the dual holographic formulation. First, the infinitesimal distance in the emergent extra dimensional space is identified with the renormalization group scale in the recursive renormalization group transformation. Second, the IR boundary condition describes the renormalization group β -functions of all the coupling parameters in the corresponding quantum field theory. In particular, the IR boundary conditions given by metric tensor fields are identified with the renormalization group equations of all interaction vertices and that given by the scalar field is nothing but the Callan-Symanzik equation for an order parameter field in the dual description. Finally, the bulk Einstein-Klein-Gordon equation describes the entanglement transfer from quantum matter to classical geometry in the large N limit, where N is the number of flavors of scalar fields.

The future direction in this line of research would be to solve the resulting coupled equations of motion in the emergent curved spacetime with an extra dimension and to show the appearance of hydrodynamic behaviors at IR, for example, calculating the ratio η/s , where η is shear viscosity and s is thermodynamic entropy [58–61]. Recently, one of the authors has investigated the Kondo effect based on this emergent geometric description and was able to describe the crossover behavior around the Kondo temperature from the UV local-moment fixed point to the IR local Fermi-liquid one in a non-perturbative way [62]. The so-called holographic liquid state may serve as a novel quantum liquid beyond the perturbation theoretical framework.

Acknowledgments

K.-S. Kim was supported by the Ministry of Education, Science, and Technology (No. 2011-0030046) of the National Research Foundation of Korea (NRF) and by TJ Park Science Fellowship of the POSCO TJ Park Foundation. This work was supported by a grant from the Simons Foundation (566116, SR).

A Emergent geometric description for a scalar lattice field theory in one dimension

We start from the following one-dimensional lattice field theory

$$Z = \int D\Phi_i \exp \left[- \int_0^\beta d\tau \sum_{i=1}^M \left\{ (\partial_\tau \Phi_i)^2 - t(\Phi_i \Phi_{i+1} + \Phi_{i+1} \Phi_i) + m^2 \Phi_i^2 + \frac{u}{2} \Phi_i^4 \right\} \right]. \quad (\text{A.1})$$

Here, Φ_i is a real scalar field at site i . t is a hopping integral and m is a mass parameter. u represents the strength of self-interactions between these scalar fields.

To implement recursive Kadanoff block-spin transformations [51] in this lattice field theory, we uplift all coupling parameters into dynamical field variables as follows

$$\begin{aligned} Z = & \int D\Phi_i D\varphi_i^{(0)} D\rho_i^{(0)} Ds_{ii+1}^{(0)} Dt_{ii+1}^{(0)} D\chi_i^{(0)} Dm_i^{2(0)} Dv_i^{(0)} Du_i^{(0)} \\ & \times \exp \left[- \int_0^\beta d\tau \sum_{i=1}^M \left\{ (\partial_\tau \Phi_i)^2 - t_{ii+1}^{(0)} (\Phi_i \Phi_{i+1} + \Phi_{i+1} \Phi_i) + m_i^{2(0)} \Phi_i^2 + \frac{u_i^{(0)}}{2} \rho_i^{(0)2} \right. \right. \\ & \left. \left. + i\varphi_i^{(0)} (\rho_i^{(0)} - \Phi_i^2) + is_{ii+1}^{(0)} (t_{ii+1}^{(0)} - t) + i\chi_i^{(0)} (m_i^{2(0)} - m^2) + iv_i^{(0)} (u_i^{(0)} - u) \right\} \right]. \quad (\text{A.2}) \end{aligned}$$

Here, $s_{ii+1}^{(0)}$, $\chi_i^{(0)}$, and $v_i^{(0)}$ are Lagrange multiplier fields to impose each constraint for hopping, mass, and self-interaction parameters, respectively. $\varphi_i^{(0)}$ is a Lagrange multiplier field to realize the Hubbard-Stratonovich transformation for the self-interaction channel.

Separating all dynamical fields at each site into those at even and odd sites, integrating over all field variables in even sites, and rescaling to return the lattice to the original one, we obtain an effective lattice field theory with renormalized hopping, mass, self-interaction,

and dual scalar fields as follows

$$\begin{aligned}
Z = & \int D\Phi_i D\varphi_i^{(0)} D\rho_i^{(0)} Ds_{ii+1}^{(0)} Dt_{ii+1}^{(0)} D\chi_i^{(0)} Dm_i^{2(0)} Dv_i^{(0)} Du_i^{(0)} \\
& \times \exp \left[-\frac{1}{4} \sum_{i=1}^M \text{tr}_{\tau\tau'} \ln \left(-\partial_\tau^2 + m_i^{2(0)} \right) \right. \\
& - \int_0^\beta d\tau \sum_{i=1}^M \left\{ (\partial_\tau \Phi_i)^2 + \left(m_i^{2(0)} - \frac{2m_i^{2(0)} t_{ii+1}^{(0)2}}{2m_i^{4(0)} + u_i^{(0)}} \right) \Phi_i^2 - i\varphi_i^{(0)} \Phi_i^2 \right. \\
& \left. \left. + \frac{u_i^{(0)}}{2} \rho_i^{(0)2} + i\varphi_i^{(0)} \rho_i^{(0)} + is_{ii+1}^{(0)} (t_{ii+1}^{(0)} - t) + i\chi_i^{(0)} (m_i^{2(0)} - m^2) + iv_i^{(0)} (u_i^{(0)} - u) \right\} \right. \\
& - \int_0^\beta d\tau \sum_{i=1}^M \left\{ -\frac{t_{ii+1}^{(0)2}}{2m_i^{4(0)} + u_i^{(0)}} \left\{ m_i^{2(0)} - \frac{u_i^{(0)} t_{ii+1}^{(0)2}}{m_i^{4(0)}} \left(\Phi_i^2 + \Phi_{i+1}^2 \right) \right\} \left(\Phi_i \Phi_{i+1} + \Phi_{i+1} \Phi_i \right) \right. \\
& \left. \left. + \frac{u_i^{(0)}}{2m_i^{4(0)} + u_i^{(0)}} \left\{ \frac{t_{ii+1}^{(0)4}}{m_i^{4(0)}} \Phi_i^4 + 3 \frac{t_{ii+1}^{(0)4}}{m_i^{4(0)}} \Phi_i^2 \Phi_{i+1}^2 \right\} + \frac{1}{16m_i^{4(0)}} \left(\frac{1}{u_i^{(0)}} + \frac{1}{2m_i^{4(0)}} \right)^{-1} \right\} \right]. \tag{A.3}
\end{aligned}$$

To prepare for the second renormalization group transformation, we rewrite this partition function in terms of updated coupling fields in the following way

$$\begin{aligned}
Z = & \int D\Phi_i D\varphi_i^{(0)} D\rho_i^{(0)} Ds_{ii+1}^{(0)} Dt_{ii+1}^{(0)} D\chi_i^{(0)} Dm_i^{2(0)} Dv_i^{(0)} Du_i^{(0)} \\
& \times D\varphi_i^{(1)} D\rho_i^{(1)} Ds_{ii+1}^{(1)} Dt_{ii+1}^{(1)} D\chi_i^{(1)} Dm_i^{2(1)} Dv_i^{(1)} Du_i^{(1)} \\
& \times \exp \left[-\frac{1}{4} \sum_{i=1}^M \text{tr}_{\tau\tau'} \ln \left(-\partial_\tau^2 + m_i^{2(0)} \right) \right. \\
& - \int_0^\beta d\tau \sum_{i=1}^M \left\{ (\partial_\tau \Phi_i)^2 + m_i^{2(1)} \Phi_i^2 - i\varphi_i^{(0)} \Phi_i^2 - t_{ii+1}^{(1)} \left(\Phi_i \Phi_{i+1} + \Phi_{i+1} \Phi_i \right) \right. \\
& \left. + \frac{u_i^{(0)}}{2} \rho_i^{(0)2} + i\varphi_i^{(0)} \rho_i^{(0)} + is_{ii+1}^{(0)} (t_{ii+1}^{(0)} - t) + i\chi_i^{(0)} (m_i^{2(0)} - m^2) + iv_i^{(0)} (u_i^{(0)} - u) \right. \\
& \left. + \frac{u_i^{(1)}}{2} \left(\rho_i^{(1)2} + 3\rho_i^{(1)} \rho_{i+1}^{(1)} \right) \right. \\
& \left. + i\varphi_i^{(1)} \left(\rho_i^{(1)} - \Phi_i^2 \right) + is_{ii+1}^{(1)} \left[t_{ii+1}^{(1)} - \frac{t_{ii+1}^{(0)2}}{2m_i^{4(0)} + u_i^{(0)}} \left\{ m_i^{2(0)} - \frac{u_i^{(0)} t_{ii+1}^{(0)2}}{m_i^{4(0)}} \left(\rho_i^{(1)} + \rho_{i+1}^{(1)} \right) \right\} \right] \right. \\
& \left. + i\chi_i^{(1)} \left\{ m_i^{2(1)} - \left(m_i^{2(0)} - \frac{2m_i^{2(0)} t_{ii+1}^{(0)2}}{2m_i^{4(0)} + u_i^{(0)}} \right) \right\} \right. \\
& \left. \left. + iv_i^{(1)} \left(u_i^{(1)} - \frac{2u_i^{(0)} t_{ii+1}^{(0)4}}{m_i^{4(0)} (2m_i^{4(0)} + u_i^{(0)})} \right) + \frac{1}{16m_i^{4(0)}} \left(\frac{1}{u_i^{(0)}} + \frac{1}{2m_i^{4(0)}} \right)^{-1} \right\} \right]. \tag{A.4}
\end{aligned}$$

Here, $s_{ii+1}^{(1)}$, $\chi_i^{(1)}$, and $v_i^{(1)}$ are Lagrange multiplier fields to impose the renormalization group flow equations for hopping, mass, and self-interaction fields, respectively. $\varphi_i^{(1)}$ is

a Lagrange multiplier field to realize the Hubbard-Stratonovich transformation for newly generated effective interactions.

Shifting $\varphi_i^{(1)}$ as

$$\varphi_i^{(1)} \implies \varphi_i^{(1)} - \varphi_i^{(0)} \quad (\text{A.5})$$

and keeping only local interactions in the sense of the gradient expansion, we obtain

$$\begin{aligned}
Z = & \int D\Phi_i D\varphi_i^{(0)} D\rho_i^{(0)} Ds_{ii+1}^{(0)} Dt_{ii+1}^{(0)} D\chi_i^{(0)} Dm_i^{2(0)} Dv_i^{(0)} Du_i^{(0)} \\
& \times D\varphi_i^{(1)} D\rho_i^{(1)} Ds_{ii+1}^{(1)} Dt_{ii+1}^{(1)} D\chi_i^{(1)} Dm_i^{2(1)} Dv_i^{(1)} Du_i^{(1)} \\
& \times \exp \left[-\frac{1}{4} \sum_{i=1}^M \text{tr}_{\tau\tau'} \ln \left(-\partial_\tau^2 + m_i^{2(0)} \right) \right. \\
& - \int_0^\beta d\tau \sum_{i=1}^M \left\{ (\partial_\tau \Phi_i)^2 + m_i^{2(1)} \Phi_i^2 - i\varphi_i^{(1)} \Phi_i^2 - t_{ii+1}^{(1)} (\Phi_i \Phi_{i+1} + \Phi_{i+1} \Phi_i) \right. \\
& + \frac{u_i^{(0)}}{2} \rho_i^{(0)2} + i\varphi_i^{(0)} \rho_i^{(0)} + is_{ii+1}^{(0)} (t_{ii+1}^{(0)} - t) + i\chi_i^{(0)} (m_i^{2(0)} - m^2) + iv_i^{(0)} (u_i^{(0)} - u) \\
& + \frac{u_i^{(1)}}{2} \rho_i^{(1)2} + i\rho_i^{(1)} (\varphi_i^{(1)} - \varphi_i^{(0)}) \\
& + is_{ii+1}^{(1)} \left\{ t_{ii+1}^{(1)} - \frac{t_{ii+1}^{(0)2}}{2m_i^{4(0)} + u_i^{(0)}} \left(m_i^{2(0)} - \frac{2u_i^{(0)} t_{ii+1}^{(0)2}}{m_i^{4(0)}} \rho_i^{(1)} \right) \right\} \\
& + i\chi_i^{(1)} \left\{ m_i^{2(1)} - \left(m_i^{2(0)} - \frac{2m_i^{2(0)} t_{ii+1}^{(0)2}}{2m_i^{4(0)} + u_i^{(0)}} \right) \right\} \\
& \left. + iv_i^{(1)} \left(u_i^{(1)} - \frac{u_i^{(0)} t_{ii+1}^{(0)4}}{2m_i^{4(0)} (2m_i^{4(0)} + u_i^{(0)})} \right) + \frac{1}{16m_i^{4(0)}} \left(\frac{1}{u_i^{(0)}} + \frac{1}{2m_i^{4(0)}} \right)^{-1} \right\}. \quad (\text{A.6})
\end{aligned}$$

This completes the first renormalization group transformation.

It is straightforward to extend this expression in a recursive way, given by

$$\begin{aligned}
Z = & \int D\Phi_i \Pi_{l=0}^f D\varphi_i^{(l)} D\rho_i^{(l)} Ds_{ii+1}^{(l)} Dt_{ii+1}^{(l)} D\chi_i^{(l)} Dm_i^{2(l)} Dv_i^{(l)} Du_i^{(l)} \\
& \exp \left[-\frac{1}{4} \sum_{l=1}^f \sum_{i=1}^M \text{tr}_{\tau\tau'} \ln \left(-\partial_\tau^2 + m_i^{2(l-1)} \right) \right. \\
& - \int_0^\beta d\tau \sum_{i=1}^M \left\{ (\partial_\tau \Phi_i)^2 + m_i^{2(f)} \Phi_i^2 - i\varphi_i^{(f)} \Phi_i^2 - t_{ii+1}^{(f)} (\Phi_i \Phi_{i+1} + \Phi_{i+1} \Phi_i) \right. \\
& + \frac{u_i^{(0)}}{2} \rho_i^{(0)2} + i\varphi_i^{(0)} \rho_i^{(0)} + is_{ii+1}^{(0)} (t_{ii+1}^{(0)} - t) + i\chi_i^{(0)} (m_i^{2(0)} - m^2) + iv_i^{(0)} (u_i^{(0)} - u) \\
& + \sum_{l=1}^f \left[\frac{u_i^{(l)}}{2} \rho_i^{(l)2} + i\rho_i^{(l)} (\varphi_i^{(l)} - \varphi_i^{(l-1)}) \right. \\
& + is_{ii+1}^{(l)} \left\{ t_{ii+1}^{(l)} - \frac{t_{ii+1}^{(l-1)2}}{2m_i^{4(l-1)} + u_i^{(l-1)}} \left(m_i^{2(l-1)} - \frac{2u_i^{(l-1)} t_{ii+1}^{(l-1)2}}{m_i^{4(l-1)}} \rho_i^{(l)} \right) \right\} \\
& + i\chi_i^{(l)} \left\{ m_i^{2(l)} - \left(m_i^{2(l-1)} - \frac{2m_i^{2(l-1)} t_{ii+1}^{(l-1)2}}{2m_i^{4(l-1)} + u_i^{(l-1)}} \right) \right\} \\
& \left. \left. + iv_i^{(l)} \left(u_i^{(l)} - \frac{u_i^{(l-1)} t_{ii+1}^{(l-1)4}}{2m_i^{4(l-1)} (2m_i^{4(l-1)} + u_i^{(l-1)})} \right) + \frac{1}{16m_i^{4(l-1)}} \left(\frac{1}{u_i^{(l-1)}} + \frac{1}{2m_i^{4(l-1)}} \right)^{-1} \right] \right\}. \tag{A.7}
\end{aligned}$$

Resorting to

$$a \sum_{l=1}^f \implies \int_0^{z_f} dz \tag{A.8}$$

and

$$\frac{\varphi_i^{(l)} - \varphi_i^{(l-1)}}{a} \implies \partial_z \varphi(i, \tau, z), \tag{A.9}$$

where a is a scale for the renormalization group transformation, we rewrite the above partition function in the following way

$$\begin{aligned}
Z = & \int D\Phi(i, \tau) \int D\varphi(i, \tau, z) D\rho(i, \tau, z) \\
& \times \int Ds(i, i+1, \tau, z) Dt(i, i+1, \tau, z) D\chi(i, \tau, z) Dm^2(i, \tau, z) Dv(i, \tau, z) Du(i, \tau, z) \\
& \times \exp \left\{ -S_{IR} - S_{UV} - S_{Bulk} \right\}. \tag{A.10}
\end{aligned}$$

Here, both IR and UV effective action are given by

$$\begin{aligned}
S_{IR} + S_{UV} = & \int_0^\beta d\tau \sum_{i=1}^M \left\{ \left(\partial_\tau \Phi(i, \tau) \right)^2 + m^2(i, \tau, z_f) [\Phi(i, \tau)]^2 - i\varphi(i, \tau, z_f) [\Phi(i, \tau)]^2 \right. \\
& - t(i, i+1, \tau, z_f) \left(\Phi(i, \tau) \Phi(i+1, \tau) + \Phi(i+1, \tau) \Phi(i, \tau) \right) + \frac{u(i, \tau, 0)}{2} [\rho(i, \tau, 0)]^2 + i\varphi(i, \tau, 0) \rho(i, \tau, 0) \\
& \left. + is(i, i+1, \tau, 0) \left(t(i, i+1, \tau, 0) - t \right) + i\chi(i, \tau, 0) \left(m^2(i, \tau, 0) - m^2 \right) + iv(i, \tau, 0) \left(u(i, \tau, 0) - u \right) \right\}.
\end{aligned} \tag{A.11}$$

The bulk effective action is

$$\begin{aligned}
S_{Bulk} = & \int_0^{z_f} dz \int_0^\beta d\tau \sum_{i=1}^M \left\{ \frac{u(i, \tau, z)}{2a} [\rho(i, \tau, z)]^2 + i\rho(i, \tau, z) \partial_z \varphi(i, \tau, z) \right. \\
& + is(i, i+1, \tau, z) \left\{ \partial_z t(i, i+1, \tau, z) + \frac{1}{a} t(i, i+1, \tau, z) \right. \\
& \left. - \frac{[t(i, i+1, \tau, z)]^2}{a \left(2[m^2(i, \tau, z)]^2 + u(i, \tau, z) \right)} \left(m^2(i, \tau, z) - \frac{2u(i, \tau, z) [t(i, i+1, \tau, z)]^2}{[m^2(i, \tau, z)]^2} \rho(i, \tau, z) \right) \right\} \\
& + i\chi(i, \tau, z) \left\{ \partial_z m^2(i, \tau, z) + \frac{2m^2(i, \tau, z) [t(i, i+1, \tau, z)]^2}{a \left(2[m^2(i, \tau, z)]^2 + u(i, \tau, z) \right)} \right\} \\
& + iv(i, \tau, z) \left(\partial_z u(i, \tau, z) + \frac{1}{a} u(i, \tau, z) - \frac{u(i, \tau, z) [t(i, i+1, \tau, z)]^4}{2a [m^2(i, \tau, z)]^2 \left(2[m^2(i, \tau, z)]^2 + u(i, \tau, z) \right)} \right) \\
& \left. + \frac{u(i, \tau, z)}{8a \left(2[m^2(i, \tau, z)]^2 + u(i, \tau, z) \right)} + \frac{1}{4a} \ln \left(-\partial_\tau^2 + m^2(i, \tau, z) \right) \right\}.
\end{aligned} \tag{A.12}$$

Performing Gaussian integrals for $\rho(i, \tau, z)$, $s(i, i+1, \tau, z)$, $\chi(i, \tau, z)$, and $v(i, \tau, z)$, one can further simplify these expressions as

$$\begin{aligned}
Z = & \int D\Phi(i, \tau) D\varphi(i, \tau, z) Dt(i, i+1, \tau, z) Dm^2(i, \tau, z) Du(i, \tau, z) \\
& \times \delta \left(t(i, i+1, \tau, 0) - t \right) \delta \left(m^2(i, \tau, 0) - m^2 \right) \delta \left(u(i, \tau, 0) - u \right) \\
& \times \delta \left(\partial_z m^2(i, \tau, z) + \frac{2m^2(i, \tau, z) [t(i, i+1, \tau, z)]^2}{a \left(2[m^2(i, \tau, z)]^2 + u(i, \tau, z) \right)} \right) \\
& \times \delta \left(\partial_z u(i, \tau, z) + \frac{1}{a} u(i, \tau, z) - \frac{u(i, \tau, z) [t(i, i+1, \tau, z)]^4}{2a [m^2(i, \tau, z)]^2 \left(2[m^2(i, \tau, z)]^2 + u(i, \tau, z) \right)} \right) \\
& \times \exp \left\{ -S_{IR} - S_{UV} - S_{Bulk} \right\}.
\end{aligned} \tag{A.13}$$

Both IR and UV effective actions are given by

$$\begin{aligned}
S_{IR} + S_{UV} = & \int_0^\beta d\tau \sum_{i=1}^M \left\{ \left(\partial_\tau \Phi(i, \tau) \right)^2 + m^2(i, \tau, z_f) [\Phi(i, \tau)]^2 - i\varphi(i, \tau, z_f) [\Phi(i, \tau)]^2 \right. \\
& \left. - t(i, i+1, \tau, z_f) \left(\Phi(i, \tau) \Phi(i+1, \tau) + \Phi(i+1, \tau) \Phi(i, \tau) \right) + \frac{1}{2u(i, \tau, 0)} [\varphi(i, \tau, 0)]^2 \right\}
\end{aligned} \tag{A.14}$$

and the bulk effective action is

$$\begin{aligned}
S_{Bulk} = & \int_0^{z_f} dz \int_0^\beta d\tau \sum_{i=1}^M \left\{ \frac{3a}{8u(i, \tau, z)} [\partial_z \varphi(i, \tau, z)]^2 \right. \\
& + \frac{a[m^2(i, \tau, z)]^4 \left(2[m^2(i, \tau, z)]^2 + u(i, \tau, z) \right)^2}{8u(i, \tau, z) [t(i, i+1, \tau, z)]^8} \\
& \times \left\{ \partial_z t(i, i+1, \tau, z) + \frac{1}{a} t(i, i+1, \tau, z) - \frac{m^2(i, \tau, z) [t(i, i+1, \tau, z)]^2}{a \left(2[m^2(i, \tau, z)]^2 + u(i, \tau, z) \right)} \right\}^2 \\
& + \frac{a[m^2(i, \tau, z)]^2 \left(2[m^2(i, \tau, z)]^2 + u(i, \tau, z) \right)}{8u(i, \tau, z) [t(i, i+1, \tau, z)]^4} \left(-i\partial_z \varphi(i, \tau, z) \right) \\
& \times \left\{ \partial_z t(i, i+1, \tau, z) + \frac{1}{a} t(i, i+1, \tau, z) - \frac{m^2(i, \tau, z) [t(i, i+1, \tau, z)]^2}{a \left(2[m^2(i, \tau, z)]^2 + u(i, \tau, z) \right)} \right\} \\
& \left. + \frac{u(i, \tau, z)}{8a \left(2[m^2(i, \tau, z)]^2 + u(i, \tau, z) \right)} + \frac{1}{4a} \ln \left(-\partial_\tau^2 + m^2(i, \tau, z) \right) \right\}.
\end{aligned} \tag{A.15}$$

For the presentation and as a natural approximation, we take a uniform ansatz in the following way

$$\begin{aligned}
Z = & \int D\Phi(i, \tau) D\varphi(z) Dt(z) Dm^2(z) Du(z) \delta(t(0) - t) \delta(m^2(0) - m^2) \delta(u(0) - u) \\
& \times \delta \left(\partial_z m^2(z) + \frac{2m^2(z) [t(z)]^2}{a \left(2[m^2(z)]^2 + u(z) \right)} \right) \delta \left(\partial_z u(z) + \frac{1}{a} u(z) - \frac{u(z) [t(z)]^4}{2a [m^2(z)]^2 \left(2[m^2(z)]^2 + u(z) \right)} \right) \\
& \times \exp \left\{ -S_{IR} - S_{UV} - S_{Bulk} \right\},
\end{aligned} \tag{A.16}$$

where the IR and UV effective actions are

$$\begin{aligned}
S_{IR} + S_{UV} = & \int_0^\beta d\tau \sum_{i=1}^M \left\{ \left(\partial_\tau \Phi(i, \tau) \right)^2 + m^2(z_f) [\Phi(i, \tau)]^2 - i\varphi(z_f) [\Phi(i, \tau)]^2 \right. \\
& \left. - t(z_f) \left(\Phi(i, \tau) \Phi(i+1, \tau) + \Phi(i+1, \tau) \Phi(i, \tau) \right) + \frac{1}{2u(0)} [\varphi(0)]^2 \right\}
\end{aligned} \tag{A.17}$$

and the bulk effective action is

$$\begin{aligned}
S_{Bulk} = & \beta M \int_0^{z_f} dz \left\{ \frac{3a}{8u(z)} [\partial_z \varphi(z)]^2 \right. \\
& + \frac{a[m^2(z)]^4 \left(2[m^2(z)]^2 + u(z) \right)^2}{8u(z)[t(z)]^8} \left\{ \partial_z t(z) + \frac{1}{a} t(z) - \frac{m^2(z)[t(z)]^2}{a \left(2[m^2(z)]^2 + u(z) \right)} \right\}^2 \\
& + \frac{a[m^2(z)]^2 \left(2[m^2(z)]^2 + u(z) \right)}{8u(z)[t(z)]^4} \left(-i\partial_z \varphi(z) \right) \left\{ \partial_z t(z) + \frac{1}{a} t(z) - \frac{m^2(z)[t(z)]^2}{a \left(2[m^2(z)]^2 + u(z) \right)} \right\} \\
& \left. + \frac{u(z)}{8a \left(2[m^2(z)]^2 + u(z) \right)} + \frac{1}{4a} \frac{1}{\beta} \sum_{i\omega_n} \ln \left(\omega_n^2 + m^2(z) \right) \right\}. \tag{A.18}
\end{aligned}$$

Finally, we perform the mean-field analysis, where quantum fluctuations of $\varphi(z)$ and $t(z)$ are neglected to be their saddle-point values. Considering

$$\varphi(z) \implies -i\varphi(z) \tag{A.19}$$

for real saddle-point values in $\varphi(z)$, we obtain the mean-field free-energy functional for $\varphi(z)$ and $t(z)$

$$\begin{aligned}
\frac{1}{M} F[\varphi(z), t(z); z_f] = & \frac{1}{2\beta} \sum_{i\omega_n} \frac{1}{M} \sum_q \ln \left\{ \omega_n^2 - 2t(z_f) \cos q + m^2(z_f) - \varphi(z_f) \right\} - \frac{1}{2u(0)} [\varphi(0)]^2 \\
& + \int_0^{z_f} dz \left\{ -\frac{3a}{8u(z)} [\partial_z \varphi(z)]^2 \right. \\
& + \frac{a[m^2(z)]^4 \left(2[m^2(z)]^2 + u(z) \right)^2}{8u(z)[t(z)]^8} \left\{ \partial_z t(z) + \frac{1}{a} t(z) - \frac{m^2(z)[t(z)]^2}{a \left(2[m^2(z)]^2 + u(z) \right)} \right\}^2 \\
& - \frac{a[m^2(z)]^2 \left(2[m^2(z)]^2 + u(z) \right)}{8u(z)[t(z)]^4} \left(\partial_z \varphi(z) \right) \left\{ \partial_z t(z) + \frac{1}{a} t(z) - \frac{m^2(z)[t(z)]^2}{a \left(2[m^2(z)]^2 + u(z) \right)} \right\} \\
& \left. + \frac{u(z)}{8a \left(2[m^2(z)]^2 + u(z) \right)} + \frac{1}{4a} \frac{1}{\beta} \sum_{i\omega_n} \ln \left(\omega_n^2 + m^2(z) \right) \right\}. \tag{A.20}
\end{aligned}$$

The renormalization group equation for the mass parameter is

$$\partial_z m^2(z) = -\frac{2m^2(z)[t(z)]^2}{a \left(2[m^2(z)]^2 + u(z) \right)} \tag{A.21}$$

and that for the interaction vertex is

$$\partial_z u(z) = -\frac{1}{a} u(z) + \frac{u(z)[t(z)]^4}{2a[m^2(z)]^2 \left(2[m^2(z)]^2 + u(z) \right)}. \tag{A.22}$$

Minimizing this free-energy functional with respect to both $\varphi(z)$ and $t(z)$, we obtain their equations of motion, given by the second-order derivatives in the coordinate of the extra dimensional space. On the other hand, their IR boundary conditions are given by the linear derivatives in z , corresponding to the renormalization group equation for the hopping integral and the Callan-Symanzik equation for the order parameter field, respectively.

B Further developments for the invariance of entanglement entropy with respect to renormalization group transformations

B.1 An identity for entanglement entropy at $z = z_f$

Resorting to the evolution equation for the metric tensor along the extra-dimensional space, we rewrite the area law of the matter sector at $z = z_f$ in terms of the Green's function as follows

$$\begin{aligned}\partial_{z_f}\mathcal{A}[\Sigma(z_f)] &= \int_{\Sigma(z_f)} d^{D-2}x_{\perp} \partial_{z_f} \sqrt{\gamma(x, z_f)} \\ &= \frac{1}{2} \int_{\Sigma(z_f)} d^{D-2}x_{\perp} \sqrt{\gamma(x, z_f)} \gamma^{ij}(x, z_f) \partial_{z_f} \gamma_{ij}(x, z_f) \\ &= -\frac{1}{4dz} \int_{\Sigma(z_f)} d^{D-2}x_{\perp} \sqrt{\gamma(x, z_f)} \gamma^{ij}(x, z_f) (\partial_i \partial_j G[x, x; g_{\mu\nu}(x, z_f)]). \quad (\text{B.1})\end{aligned}$$

We recall that the Green's function is given by

$$\begin{aligned}&\left\{ -\frac{1}{\sqrt{g(x, z_f)}} \partial_{\mu} \left(\sqrt{g(x, z_f)} g^{\mu\nu}(x, z_f) \partial_{\nu} \right) + \frac{m^2}{e^{2dz} - 1} \right\} G[x, x'; g_{\mu\nu}(x, z_f)] \\ &= \frac{1}{\sqrt{g(x, z_f)}} \delta^{(D)}(x - x'). \quad (\text{B.2})\end{aligned}$$

One can reformulate this Green's function in terms of the geometric information such as metric and curvature, considering the heat kernel expansion. Introducing the heat kernel $K(s, x, x'; z) \equiv \langle xz | e^{-s\mathcal{D}} | x'z \rangle$ with the differential operator $\mathcal{D} \equiv -\frac{1}{\sqrt{g(x, z)}} \partial_{\mu} \left(\sqrt{g(x, z)} g^{\mu\nu}(x, z) \partial_{\nu} \right) + \frac{m^2}{e^{2dz} - 1}$, which satisfies the diffusion equation $(\partial_s + \mathcal{D})K(s, x, x'; z) = 0$ with an initial condition $K(s = 0, x, x'; z) = \frac{1}{\sqrt{g(x, z)}} \delta^{(D)}(x - x')$, we obtain

$$G(x, x'; z) = \int_{\epsilon^2}^{\infty} ds K(s, x, x'; z). \quad (\text{B.3})$$

Here, ϵ is a UV cutoff to cure the UV divergence of the entanglement entropy, introduced in subsection 5.2. As discussed in the heat-kernel expansion Eq. (5.14), we express the

Green's function in terms of the metric and curvature at $z = z_f$ as

$$\begin{aligned}
G(x, x; z_f) &= \int_{\epsilon^2}^{\infty} ds K(s, x, x; z_f) = \int_{\epsilon^2}^{\infty} ds \sum_{n=0}^{\infty} a_n(x, z_f) s^{\frac{n-D}{2}} \\
&= \int_{\epsilon^2}^{\infty} ds \left\{ a_0(x, z_f) s^{-\frac{D}{2}} + a_2(x, z_f) s^{\frac{2-D}{2}} + a_4(x, z_f) s^{\frac{4-D}{2}} + \dots \right\} \\
&= \int_{\epsilon^2}^{\infty} ds \frac{1}{(4\pi)^{\frac{D}{2}}} \left[s^{-\frac{D}{2}} + \left(\frac{1}{6} R(x, z_f) - \frac{m^2}{e^{2dz} - 1} \right) s^{\frac{2-D}{2}} \right. \\
&\quad + \left\{ \frac{1}{180} R_{\mu\nu\alpha\beta}^2(x, z_f) - \frac{1}{180} R_{\mu\nu}^2(x, z_f) \right. \\
&\quad + \frac{1}{6} \frac{1}{\sqrt{g(x, z_f)}} \partial_\mu \left(\sqrt{g(x, z_f)} g^{\mu\nu}(x, z_f) \partial_\nu \right) \left(\frac{1}{5} R(x, z_f) - \frac{m^2}{e^{2dz} - 1} \right) \\
&\quad \left. \left. + \frac{1}{2} \left(\frac{1}{6} R(x, z_f) - \frac{m^2}{e^{2dz} - 1} \right)^2 \right\} s^{\frac{4-D}{2}} + \dots \right]. \tag{B.4}
\end{aligned}$$

Inserting this expression into Eq. (B.1), we obtain

$$\begin{aligned}
\partial_{z_f} \mathcal{A}[\Sigma(z_f)] &= \\
&- \frac{1}{4dz} \left(\frac{1}{(4\pi)^{\frac{D}{2}}} \int_{\epsilon^2}^{\infty} ds s^{\frac{2-D}{2}} \right) \int_{\Sigma(z_f)} d^{D-2} x_\perp \sqrt{\gamma(x, z_f)} \gamma^{ij}(x, z_f) \partial_i \partial_j \left(\frac{1}{6} R(x, z_f) - \frac{m^2}{e^{2dz} - 1} \right) \\
&- \frac{1}{4dz} \left(\frac{1}{(4\pi)^{\frac{D}{2}}} \int_{\epsilon^2}^{\infty} ds s^{\frac{4-D}{2}} \right) \int_{\Sigma(z_f)} d^{D-2} x_\perp \sqrt{\gamma(x, z_f)} \gamma^{ij}(x, z_f) \\
&\quad \times \partial_i \partial_j \left\{ \frac{1}{180} R_{\mu\nu\alpha\beta}^2(x, z_f) - \frac{1}{180} R_{\mu\nu}^2(x, z_f) \right. \\
&\quad + \frac{1}{6} \frac{1}{\sqrt{g(x, z_f)}} \partial_\mu \left(\sqrt{g(x, z_f)} g^{\mu\nu}(x, z_f) \partial_\nu \right) \left(\frac{1}{5} R(x, z_f) - \frac{m^2}{e^{2dz} - 1} \right) \\
&\quad \left. + \frac{1}{2} \left(\frac{1}{6} R(x, z_f) - \frac{m^2}{e^{2dz} - 1} \right)^2 \right\}. \tag{B.5}
\end{aligned}$$

Combining this expression with that of the gravity sector, we reach the following identity

for the entanglement entropy

$$\begin{aligned}
& \frac{N}{6(D-2)(4\pi)^{\frac{D}{2}-1}} \frac{1}{\epsilon^{D-2}} \frac{1}{4dz} \frac{1}{(4\pi)^{\frac{D}{2}}} \int_{\Sigma(z_f)} d^{D-2}x_{\perp} \sqrt{\gamma(x, z_f)} \gamma^{ij}(x, z_f) \left[\right. \\
& - \left(\int_{\epsilon^2}^{\infty} ds s^{\frac{2-D}{2}} \right) \partial_i \partial_j \left(\frac{1}{6} R(x, z_f) - \frac{m^2}{e^{2dz} - 1} \right) \\
& - \left(\int_{\epsilon^2}^{\infty} ds s^{\frac{4-D}{2}} \right) \partial_i \partial_j \left\{ \frac{1}{180} R_{\mu\nu\alpha\beta}^2(x, z_f) - \frac{1}{180} R_{\mu\nu}^2(x, z_f) \right. \\
& \left. + \frac{1}{6} \frac{1}{\sqrt{g(x, z_f)}} \partial_{\mu} \left(\sqrt{g(x, z_f)} g^{\mu\nu}(x, z_f) \partial_{\nu} \right) \left(\frac{1}{5} R(x, z_f) - \frac{m^2}{e^{2dz} - 1} \right) + \frac{1}{2} \left(\frac{1}{6} R(x, z_f) - \frac{m^2}{e^{2dz} - 1} \right)^2 \right\} \left. \right] \\
& + N \int d^{D-2}x_{\perp} \int_0^{2\pi} d\theta \int_0^{\infty} dr \sqrt{g(r, x_{\perp}, z_f)} \left[\left(T_{\mu\nu}^{\varphi}(r, x_{\perp}, z_f) + T_{\mu\nu}^{g\mu\nu}(r, x_{\perp}, z_f) \right) \frac{\partial g_n^{\mu\nu}(r, x_{\perp}, z_f)}{\partial n} \Big|_{n=1} \right. \\
& - \left\{ \frac{1}{2u} [\partial_z \varphi(r, x_{\perp}, z_f)]^2 + \frac{C_{\varphi}}{2} g^{\mu\nu}(r, x_{\perp}, z_f) [\partial_{\mu} \varphi(r, x_{\perp}, z_f)] [\partial_{\nu} \varphi(r, x_{\perp}, z_f)] + C_{\xi} R(r, x_{\perp}, z_f) [\varphi(r, x_{\perp}, z_f)]^2 \right. \\
& \left. \left. + \frac{1}{2\kappa} \left(R(r, x_{\perp}, z_f) - 2\Lambda \right) \right\} \right] = 0. \tag{B.6}
\end{aligned}$$

We point out that both integrals of $\int_{\epsilon^2}^{\infty} ds s^{(2-D)/2}$ and $\int_{\epsilon^2}^{\infty} ds s^{(4-D)/2}$ diverge for large s . Existence of these IR divergences may be an artifact of the small s -expansion. One may introduce the correlation length into the upper cutoff of these integrals as $\int_{\epsilon^2}^{\xi^2} ds s^{(2-D)/2}$ and $\int_{\epsilon^2}^{\xi^2} ds s^{(4-D)/2}$. We recall that this IR cutoff reproduces the entanglement entropy in $D = 2$, as shown in Eq. (5.21). Now, the identity is expressed in a fully geometric way.

B.2 Entanglement entropy after the first iteration

While it is written in terms of all the geometrical information, it is not clear how to interpret the above lengthy expression. In addition, it is not easy to check out this identity in a brute force way. In this respect we consider the entanglement entropy after the first iteration of the renormalization group transformation and verify the invariance explicitly for the case without all types of interactions.

We recall the partition function after the renormalization group transformation

$$\begin{aligned}
Z^{(1)} &= \int D\phi_{\alpha} Dg_{\mu\nu}^{(1)} \delta \left(g^{(1)\mu\nu} - g_B^{\mu\nu} - g_B^{\mu\nu'} (\partial_{\nu'} \partial_{\mu'} G_{xx'})_{x' \rightarrow x} g_B^{\mu'\nu'} \right) \\
&\times \exp \left[- \int d^D x \sqrt{g^{(1)}} \left\{ g^{(1)\mu\nu} (\partial_{\mu} \phi_{\alpha}) (\partial_{\nu} \phi_{\alpha}) + m^2 \phi_{\alpha}^2 \right\} - \frac{N}{2\kappa} \int d^D x \sqrt{g_B} (R_B - 2\Lambda) \right], \tag{B.7}
\end{aligned}$$

where the coefficients of the gravity action are $2dz\mathcal{C}_R = \frac{1}{2\kappa}$ and $\frac{C_{\Lambda}}{C_R} = 2\Lambda$. Then, the

entanglement entropy is

$$\begin{aligned}
\mathcal{S}_{EE}^{(1)} &= \frac{N}{6(D-2)(4\pi)^{\frac{D}{2}-1}} \frac{1}{\epsilon^{D-2}} \int_{\Sigma^{(1)}} d^{D-2}x_{\perp} \sqrt{g_{\perp}^{(1)}(x)} \\
&+ \frac{N}{2\kappa} \int d^{D-2}x_{\perp} \int_0^{2\pi} d\theta \int_0^{\infty} dr \sqrt{g_B(x)} \left[-2 \left(R_{\mu\nu}^B(x) - \frac{1}{2} g_{\mu\nu}^B(x) R_B(x) + \Lambda g_{\mu\nu}^B(x) \right) \right. \\
&+ \left. \frac{1}{g_B(x)} g_{\mu\mu'}^B(x) g_{\nu\nu'}^B(x) \partial_{\mu''} \partial_{\nu''} \left\{ g_B(x) \left(g_B^{\mu'\nu'}(x) g_B^{\mu''\nu''}(x) - g_B^{\mu'\mu''}(x) g_B^{\nu'\nu''}(x) \right) \right\} \right] \frac{\partial g_{Bn}^{\mu\nu}(r, x_{\perp})}{\partial n} \Big|_{n=1} \\
&- \frac{N}{2\kappa} \int d^{D-2}x_{\perp} \int_0^{2\pi} d\theta \int_0^{\infty} dr \sqrt{g_B(x)} \left(R_B(x) - 2\Lambda \right). \tag{B.8}
\end{aligned}$$

This entanglement entropy has to be identified with

$$\mathcal{S}_{EE} = \frac{N}{6(D-2)(4\pi)^{\frac{D}{2}-1}} \frac{1}{\epsilon^{D-2}} \int_{\Sigma_B} d^{D-2}x_{\perp} \sqrt{g_{\perp}^B(x)}. \tag{B.9}$$

In other words, we have

$$\mathcal{S}_{EE} = \mathcal{S}_{EE}^{(1)}. \tag{B.10}$$

As a result, we have to obtain

$$\begin{aligned}
&\frac{N}{6(D-2)(4\pi)^{\frac{D}{2}-1}} \frac{1}{\epsilon^{D-2}} \left(\int_{\Sigma_B} d^{D-2}x_{\perp} \sqrt{g_{\perp}^B(x)} - \int_{\Sigma^{(1)}} d^{D-2}x_{\perp} \sqrt{g_{\perp}^{(1)}(x)} \right) \\
&= \frac{N}{2\kappa} \int d^{D-2}x_{\perp} \int_0^{2\pi} d\theta \int_0^{\infty} dr \sqrt{g_B(x)} \left[-2 \left(R_{\mu\nu}^B(x) - \frac{1}{2} g_{\mu\nu}^B(x) R_B(x) + \Lambda g_{\mu\nu}^B(x) \right) \right. \\
&+ \left. \frac{1}{g_B(x)} g_{\mu\mu'}^B(x) g_{\nu\nu'}^B(x) \partial_{\mu''} \partial_{\nu''} \left\{ g_B(x) \left(g_B^{\mu'\nu'}(x) g_B^{\mu''\nu''}(x) - g_B^{\mu'\mu''}(x) g_B^{\nu'\nu''}(x) \right) \right\} \right] \frac{\partial g_{Bn}^{\mu\nu}(r, x_{\perp})}{\partial n} \Big|_{n=1} \\
&- \frac{N}{2\kappa} \int d^{D-2}x_{\perp} \int_0^{2\pi} d\theta \int_0^{\infty} dr \sqrt{g_B(x)} \left(R_B(x) - 2\Lambda \right). \tag{B.11}
\end{aligned}$$

Recalling the evolution equation of

$$g^{(1)\mu\nu}(x) = g_B^{\mu\nu}(x) + g_B^{\mu\nu'}(x) (\partial_{\nu'} \partial_{\mu'} G_{xx'})_{x' \rightarrow x} g_B^{\mu'\nu}(x), \tag{B.12}$$

we obtain

$$\sqrt{g^{(1)}(x)} = \sqrt{g_B(x)} \left\{ 1 - \frac{1}{2} g_B^{\mu\nu}(x) (\partial_{\mu} \partial_{\nu} G_{xx'})_{x' \rightarrow x} \right\}. \tag{B.13}$$

As a result, we have

$$\begin{aligned}
&\int_{\Sigma_B} d^{D-2}x_{\perp} \sqrt{g_{\perp}^B(x)} - \int_{\Sigma^{(1)}} d^{D-2}x_{\perp} \sqrt{g_{\perp}^{(1)}(x)} \\
&= \frac{1}{2} \int_{\Sigma_B} d^{D-2}x_{\perp} \sqrt{g_{\perp}^B(x)} g_{B\perp}^{\mu\nu}(x) (\partial_{\mu} \partial_{\nu} G_{xx'})_{x' \rightarrow x}. \tag{B.14}
\end{aligned}$$

The final equation to be verified is

$$\begin{aligned}
& \frac{N}{6(D-2)(4\pi)^{\frac{D}{2}-1}} \frac{1}{2\epsilon^{D-2}} \int_{\Sigma_B} d^{D-2}x_{\perp} \sqrt{g_{\perp}^B(x)} g_{B\perp}^{\mu\nu}(x) (\partial_{\mu} \partial_{\nu} G_{xx'})_{x' \rightarrow x} \\
&= \frac{N}{2\kappa} \int d^{D-2}x_{\perp} \int_0^{2\pi} d\theta \int_0^{\infty} dr \sqrt{g_B(x)} \left[-2 \left(R_{\mu\nu}^B(x) - \frac{1}{2} g_{\mu\nu}^B(x) R_B(x) + \Lambda g_{\mu\nu}^B(x) \right) \right. \\
&\quad \left. + \frac{1}{g_B(x)} g_{\mu\mu'}^B(x) g_{\nu\nu'}^B(x) \partial_{\mu''} \partial_{\nu''} \left\{ g_B(x) \left(g_B^{\mu'\nu'}(x) g_B^{\mu''\nu''}(x) - g_B^{\mu'\mu''}(x) g_B^{\nu'\nu''}(x) \right) \right\} \right] \frac{\partial g_{Bn}^{\mu\nu}(r, x_{\perp})}{\partial n} \Big|_{n=1} \\
&\quad - \frac{N}{2\kappa} \int d^{D-2}x_{\perp} \int_0^{2\pi} d\theta \int_0^{\infty} dr \sqrt{g_B(x)} \left(R_B(x) - 2\Lambda \right). \tag{B.15}
\end{aligned}$$

Here, the background metric is

$$ds^2 = dr^2 + r^2 d\theta^2 + \delta_{ij} dx_{\perp}^i dx_{\perp}^j, \quad g_{rr}^B = 1, \quad g_{\theta\theta}^B = r^2, \quad g_{ij}^{B\perp} = \delta_{ij}, \quad \sqrt{g_B} = r \tag{B.16}$$

with

$$\frac{\partial g_{Bn}^{\mu\nu}(r, x_{\perp})}{\partial n} \Big|_{n=1} = 2r^2 \delta_{\mu\theta} \delta_{\nu\theta} \tag{B.17}$$

and the Green's function is [34, 35]

$$\begin{aligned}
G_{xx'} &\equiv G(r, r', \theta, \theta', x_{\perp} - x'_{\perp}) \\
&= \int \frac{d^{D-2}p_{\perp}}{(2\pi)^{D-2}} e^{ip_{\perp}(x_{\perp} - x'_{\perp})} \frac{1}{2\pi} \sum_{k=0}^{\infty} d_k \int_0^{\infty} d\lambda \lambda \frac{J_k(\lambda r) J_k(\lambda r')}{p_{\perp}^2 + \lambda^2 + m^2/(e^{2dz} - 1)} \cos k(\theta - \theta') \tag{B.18}
\end{aligned}$$

with

$$d_0 = 1, \quad d_{k>0} = 2. \tag{B.19}$$

Inserting this Green's function into Eq. (B.15), the left-hand-side of this equation is written in terms of ϵ and m with Σ_B . On the other hand, the right-hand-side of this equation is given by κ and Λ with a UV cutoff in the r integral. As a result, renormalizing both coefficients of κ and Λ appropriately, one can verify this identity.

References

- [1] M. E. Peskin and D. V. Schroeder, *An Introduction to quantum field theory* (Westview, Boulder, 1995).
- [2] J. M. Maldacena, *The Large N Limit of Superconformal Field Theories and Supergravity*, Int. J. Theor. Phys. **38**, 1113 (1999) [arXiv:hep-th/9711200v3].
- [3] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, *Gauge Theory Correlators from Non-Critical String Theory*, Phys. Lett. B **428**, 105 (1998) [arXiv:hep-th/9802109v2].
- [4] E. Witten, *Anti De Sitter Space And Holography*, Adv. Theor. Math. Phys. **2**, 253 (1998) [arXiv:hep-th/9802150v2].
- [5] M. Bianchi, D. Z. Freedman, and K. Skenderis, *Holographic Renormalization*, Nucl. Phys. B **631**, 159 (2002) [arXiv:hep-th/0112119v2].

- [6] J. de Boer, E. P. Verlinde, and H. L. Verlinde, *On the Holographic Renormalization Group*, J. High Energy Phys. **08** (2000) 003 [arXiv:hep-th/9912012v1].
- [7] E. P. Verlinde and H. L. Verlinde, *RG-Flow, Gravity and the Cosmological Constant*, J. High Energy Phys. **05** (2000) 034 [arXiv:hep-th/9912018v2].
- [8] O. Aharony, S. S. Gubser, J. Maldacena, H. Ooguri, and Y. Oz, *Large N Field Theories, String Theory and Gravity*, Phys. Rep. **323**, 183 (2000) [arXiv:hep-th/9905111v3].
- [9] M. Nozaki, S. Ryu, and T. Takayanagi, *Holographic Geometry of Entanglement Renormalization in Quantum Field Theories*, J. High Energy Phys. **10** (2012) 193 [arXiv:1208.3469v3].
- [10] I. Heemskerk, J. Penedones, J. Polchinski, and J. Sully, *Holography from Conformal Field Theory*, J. High Energy Phys. **10** (2009) 079 [arXiv:0907.0151v3].
- [11] I. Heemskerk and J. Polchinski, *Holographic and Wilsonian Renormalization Groups*, J. High Energy Phys. **06** (2011) 031 [arXiv:1010.1264v2].
- [12] T. Faulkner, H. Liu, and M. Rangamani, *Integrating out geometry: Holographic Wilsonian RG and the membrane paradigm*, J. High Energy Phys. **08** (2011) 051 [arXiv:1010.4036v2].
- [13] R. de Mello Koch, A. Jevicki, K. Jin, and J. P. Rodrigues, *AdS₄/CFT₃ Construction from Collective Fields*, Phys. Rev. D **83**, 025006 (2011) [arXiv:1008.0633v3].
- [14] M. R. Douglas, L. Mazzucato, and S. S. Razamat, *Holographic dual of free field theory*, Phys. Rev. D **83**, 071701 (2011) [arXiv:1011.4926v2].
- [15] R. G. Leigh, O. Parrikar, and A. B. Weiss, *The Holographic Geometry of the Renormalization Group and Higher Spin Symmetries*, Phys. Rev. D **89**, 106012 (2014) [arXiv:1402.1430v2].
- [16] E. Mintun and J. Polchinski, *Higher Spin Holography, RG, and the Light Cone*, arXiv:1411.3151v1.
- [17] F. Pastawski, B. Yoshida, D. Harlow, and J. Preskill, *Holographic quantum error-correcting codes: Toy models for the bulk/boundary correspondence*, J. High Energy Phys. **06** (2015) 149 [arXiv:1503.06237v2].
- [18] X. Wen, G. Y. Cho, P. L. S. Lopes, Y. Gu, X.-L. Qi, and S. Ryu, *Holographic Entanglement Renormalization of Topological Insulators*, Phys. Rev. B **94**, 075124 (2016) [arXiv:1605.07199v1].
- [19] Y. Gu, C. H. Lee, X. Wen, G. Y. Cho, S. Ryu, and X.-L. Qi, *Holographic duality between (2+1)-dimensional quantum anomalous Hall state and (3+1)-dimensional topological insulators*, Phys. Rev. B **94**, 125107 (2016) [arXiv:1605.00570v1].
- [20] T. Faulkner, M. Guica, T. Hartman, R. C. Myers, and M. Van Raamsdonk, *Gravitation from Entanglement in Holographic CFTs*, J. High Energy Phys. **03** (2014) 051 [arXiv:1312.7856v2].
- [21] T. Faulkner, F.M. Haehl, E. Hijano, O. Parrikar, C. Rabideau, and M. Van Raamsdonk, *Nonlinear Gravity from Entanglement in Conformal Field Theories*, J. High Energy Phys. **08** (2017) 057 [arXiv:1705.03026v1].
- [22] B. Czech, *Einstein Equations from Varying Complexity*, Phys. Rev. Lett. **120**, 031601 (2018) [arXiv:1706.00965v1].
- [23] K.-S. Kim and C. Park, *Emergent geometry from field theory: Wilson's renormalization group revisited*, Phys. Rev. D **93**, 121702 (2016) [arXiv:1604.04990v1].

- [24] G. Vidal, *Entanglement renormalization*, Phys. Rev. Lett. **99**, 220405 (2007) [arXiv:cond-mat/0512165v2].
- [25] J. Haegeman, T. J. Osborne, H. Verschelde, and F. Verstraete, *Entanglement Renormalization for Quantum Fields in Real Space*, Phys. Rev. Lett. **110**, 100402 (2013) [arXiv:1102.5524v2].
- [26] G. Vidal, *Understanding Quantum Phase Transitions*, edited by L. D. Carr (Taylor & Francis, Boca Raton, 2010).
- [27] G. Evenbly and G. Vidal, *Strongly Correlated Systems. Numerical Methods*, edited by A. Avella and F. Mancini, Springer Series in Solid-State Sciences Vol. **176** (Springer-Verlag, Berlin, Heidelberg, 2013), Chap. 4.
- [28] G. Evenbly and G. Vidal, *Tensor network states and geometry*, J. Stat. Phys. **145**, 891 (2011) [arXiv:1106.1082v1].
- [29] B. Swingle, *Entanglement renormalization and holography*, Phys. Rev. D **86**, 065007 (2012) [arXiv:0905.1317v1].
- [30] J. Polchinski, *Renormalization and Effective Lagrangians*, Nucl. Phys. B **231** (1984) 269.
- [31] Sung-Sik Lee, *Background independent holographic description : From matrix field theory to quantum gravity*, J. High Energy Phys. **10** (2012) 160 [arXiv:1204.1780v3].
- [32] Sung-Sik Lee, *Quantum Renormalization Group and Holography*, J. High Energy Phys. **01** (2014) 076 [arXiv:1305.3908v2].
- [33] Peter Lunts, Subhro Bhattacharjee, Jonah Miller, Erik Schnetter, Yong Baek Kim, and Sung-Sik Lee, *Ab initio holography*, J. High Energy Phys. **08** (2015) 107 [arXiv:1503.06474v3].
- [34] P. Calabrese and J. Cardy, *Entanglement Entropy and Quantum Field Theory*, J. Stat. Mech. (2004) **P06002** [arXiv:hep-th/0405152v3].
- [35] P. Calabrese and J. Cardy, *Entanglement entropy and conformal field theory*, J. Phys. A **42**, 504005 (2009) [arXiv:0905.4013v2].
- [36] S. Ryu and T. Takayanagi, *Holographic Derivation of Entanglement Entropy from AdS/CFT*, Phys. Rev. Lett. **96**, 181602 (2006) [arXiv:hep-th/0603001v2].
- [37] S. Ryu and T. Takayanagi, *Aspects of Holographic Entanglement Entropy*, J. High Energy Phys. **08** (2006) 045 [arXiv:hep-th/0605073v3].
- [38] J. Eisert, M. Cramer, and M. B. Plenio, *Colloquium: Area laws for the entanglement entropy*, Rev. Mod. Phys. **82**, 277 (2010) [arXiv:0808.3773v4].
- [39] Tatsuma Nishioka, *Entanglement entropy: Holography and renormalization group*, Rev. Mod. Phys. **90**, 035007 (2018) [arXiv:1801.10352v3].
- [40] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, New York, 1982).
- [41] J. Cardy, *The $T\bar{T}$ deformation of quantum field theory as random geometry*, J. High Energy Phys. **10** (2018) 186 [arXiv:1801.06895v6].
- [42] P. M. Chaikin and T. C. Lubensky, *Principles of Condensed Matter Physics* (Cambridge University Press, New York, 1995).

- [43] A. Sakharov, *Vacuum Quantum Fluctuations in Curved Space and the Theory of Gravitation*, Sov. Phys. Dokl. **12**, 1040 (1968), and reprinted in Gen. Rel. Grav. **32**, 365 (2000).
- [44] M. Visser, *Sakharov's induced gravity: a modern perspective*, Mod. Phys. Lett. **A 17**, 977 (2002), and references therein [arXiv:gr-qc/0204062v1].
- [45] James W. York, Jr., *Role of Conformal Three-Geometry in the Dynamics of Gravitation*, Phys. Rev. Lett. **28**, 1082 (1972).
- [46] G. W. Gibbons and S. W. Hawking, *Action integrals and partition functions in quantum gravity*, Phys. Rev. D **15**, 2752 (1977).
- [47] Sergey N. Solodukhin, *Entanglement entropy of black holes*, Living Rev. Relativity. **14** (2011) 8 [arXiv:1104.3712v1].
- [48] K.-S. Kim, M. Park, J. Cho, and C. Park, *Emergent geometric description for a topological phase transition in the Kitaev superconductor model*, Phys. Rev. D **96**, 086015 (2017) [arXiv:1610.07312v9].
- [49] Sung-Sik Lee, *Horizon as Critical Phenomenon*, J. High Energy Phys. **09** (2016) 044 [arXiv:1603.08509v2].
- [50] Sung-Sik Lee, *Emergent gravity from relatively local Hamiltonians and a possible resolution of the black hole information puzzle*, J. High Energy Phys. **10** (2018) 043 [arXiv:1803.00556v4].
- [51] Michael Plischke and Birger Bergersen, *Equilibrium Statistical Physics* (World Scientific Publishing, London, 2006).
- [52] A. Lewkowycz and J. Maldacena, *Generalized gravitational entropy*, J. High Energy Phys. **08** (2013) 090 [arXiv:1304.4926v2].
- [53] A. N. Petrov, *Classical and Quantum Gravity Research*, edited by M.N. Christiansen and T.K. Rasmussen (Nova Science Publishers, New York, 2008), Chap. 2.
- [54] C. Fronsdal, *Massless fields with integer spin*, Phys. Rev. D **18**, 3624 (1978).
- [55] E. S. Fradkin and M. A. Vasiliev, *On the Gravitational Interaction of Massless Higher Spin Fields*, Phys. Lett. B **189**, 89 (1987); E. S. Fradkin and M. A. Vasiliev, *Cubic Interaction in Extended Theories of Massless Higher Spin Fields*, Nucl. Phys. B **291**, 141 (1987).
- [56] I. R. Klebanov and A. M. Polyakov, *AdS dual of the critical $O(N)$ vector model*, Phys. Lett. B **550**, 213 (2002) [arXiv:hep-th/0210114].
- [57] X. Bekaert, N. Boulanger, and P. A. Sundell, *How higher-spin gravity surpasses the spin-two barrier*, Rev. Mod. Phys. **84**, 987 (2012) [arXiv:1007.0435v3].
- [58] G. Policastro, D. T. Son, and A. O. Starinets, *From AdS/CFT correspondence to hydrodynamics*, J. High Energy Phys. **09** (2002) 043 [arXiv:hep-th/0205052v2].
- [59] G. Policastro, D. T. Son, and A. O. Starinets, *From AdS/CFT correspondence to hydrodynamics. II. Sound waves*, J. High Energy Phys. **12** (2002) 054 [arXiv:hep-th/0210220v3].
- [60] Pavel Kovtun, Dam T. Son, and Andrei O. Starinets, *Holography and hydrodynamics: diffusion on stretched horizons*, J. High Energy Phys. **10** (2003) 064 [arXiv:hep-th/0309213v1].
- [61] P. Kovtun, D. T. Son, and A. O. Starinets, *Viscosity in Strongly Interacting Quantum Field Theories from Black Hole Physics*, Phys. Rev. Lett. **94**, 111601 (2005) [arXiv:hep-th/0405231v2].

- [62] Ki-Seok Kim, Suk Bum Chung, Chanyong Park, and Jae-Ho Han, *Emergent holographic description for the Kondo effect: Comparison with Bethe ansatz*, Phys. Rev. D **99**, 105012 (2019) [arXiv:1705.06571v6].