

CHEN'S INEQUALITIES FOR SUBMANIFOLDS IN (κ, μ) -CONTACT SPACE FORM WITH GENERALIZED SEMI-SYMMETRIC NON-METRIC CONNECTIONS

YONG WANG

ABSTRACT. In this paper, we obtain Chen's inequalities for submanifolds in (κ, μ) -contact space form with two kinds of generalized semi-symmetric non-metric connections.

1. INTRODUCTION

H. A. Hayden introduced the notion of a semi-symmetric metric connection on a Riemannian manifold [15]. K. Yano studied a Riemannian manifold endowed with a semi-symmetric metric connection [31]. Some properties of a Riemannian manifold and a hypersurface of a Riemannian manifold with a semi-symmetric metric connection were studied by T. Imai [16, 17]. Z. Nakao [25] studied submanifolds of a Riemannian manifold with semi-symmetric metric connections. N. S. Agashe and M. R. Chafle introduced the notion of a semisymmetric non-metric connection and studied some of its properties and submanifolds of a Riemannian manifold with a semi-symmetric non-metric connection [1, 2]. In [28], we introduced a kind of generalized semi-symmetric non-metric connection as a generalization of a semi-symmetric metric connection and a semi-symmetric non-metric connection and we studied the Einstein warped products and multiply warped products with this generalized semi-symmetric non-metric connection. In [21], Li, He and Zhao introduced another kind of generalized semi-symmetric non-metric connection. They studied submanifolds in a Riemannian manifold with this kind of generalized semi-symmetric non-metric connection.

On the other hand, one of the basic problems in submanifold theory is to find simple relationships between the extrinsic and intrinsic invariants of a submanifold. B. Y. Chen [6, 7, 8, 9, 10] established inequalities in this respect, called Chen inequalities. Afterwards, many geometers studied similar problems for different submanifolds in various ambient spaces; see, for example, [5, 11, 12, 14, 18, 20, 13, 19, 22, 35]. In [23, 26], Mihai and Özgür studied Chen inequalities for submanifolds of real space forms with a semi-symmetric metric connection and a semi-symmetric non-metric connection, respectively. Later, in [35], Zhang, Zhang and Song obtained Chen's inequalities for submanifolds of a Riemannian

2010 *Mathematics Subject Classification*: 53C40; 53C42.

Key words and phrases: Chen's inequalities; (κ, μ) -contact space form; generalized semi-symmetric non-metric connection .

manifold of quasi-constant curvature endowed with a semi-symmetric metric connection. At the meantime, in [33], Zhang *et al.* obtained Chen's inequalities for submanifolds of a Riemannian manifold of nearly quasi-constant curvature endowed with a semi-symmetric non-metric connection. In [24], Mihai and Özgür studied Chen inequalities for submanifolds of complex space forms and Sasakian space forms with a semi-symmetric metric connection. In [3], Ahmad, He, Tang and Zhao obtained Chen's inequalities for submanifolds in (κ, μ) -contact space form with a semi-symmetric metric connection. In [4], Ahmad, Shahzad and Li obtained Chen's inequalities for submanifolds in (κ, μ) -contact space form with a semi-symmetric non-metric connection.

The paper is organized as follows. In Section 2, we give a brief introduction about two kinds of generalized semi-symmetric non-metric connections. In Section 3, we establish Chen's inequalities for submanifolds in (κ, μ) -contact space form with the first generalized semi-symmetric non-metric connection. In Section 4, we establish Chen's inequalities for submanifolds in (κ, μ) -contact space form with the second generalized semi-symmetric non-metric connection.

2. PRELIMINARIES

Let N^{n+p} be an $(n + p)$ -dimensional Riemannian manifold with Riemannian metric g and the Levi-Civita connection $\widehat{\nabla}$. Let $\overline{\nabla}$ be a linear connection defined by

$$(2.1) \quad \overline{\nabla}_X Y = \widehat{\nabla}_X Y + \lambda_1 \pi(Y)X - \lambda_2 g(X, Y)P,$$

where P is a vector field on N and $\pi(X) = g(P, X)$. We call $\overline{\nabla}$ as the first generalized semi-symmetric non-metric connection. When $\lambda_1 = \lambda_2 = 1$, $\overline{\nabla}$ is the semi symmetric metric connection and when $\lambda_1 = 1$, $\lambda_2 = 0$, $\overline{\nabla}$ is the semi symmetric non-metric connection. Let

$$\overline{R}(X, Y)Z := \overline{\nabla}_X \overline{\nabla}_Y Z - \overline{\nabla}_Y \overline{\nabla}_X Z - \overline{\nabla}_{[X, Y]}Z$$

be the curvature tensor of $\overline{\nabla}$. The curvature tensor of $\widehat{\nabla}$, say \widehat{R} , can be defined as the same. Let

$$\alpha(X, Y) = (\widehat{\nabla}_X \pi)(Y) - \lambda_1 \pi(X)\pi(Y) + \frac{\lambda_2}{2}g(X, Y)\pi(P),$$

and

$$\beta(X, Y) = \frac{\pi(P)}{2}g(X, Y) + \pi(X)\pi(Y)$$

be two $(0, 2)$ -tensors, then we have

$$(2.2) \quad \begin{aligned} \overline{R}(X, Y, Z, W) &= \widehat{R}(X, Y, Z, W) + \lambda_1 \alpha(X, Z)g(Y, W) - \lambda_1 \alpha(Y, Z)g(X, W) \\ &+ \lambda_2 g(X, Z)\alpha(Y, W) - \lambda_2 g(Y, Z)\alpha(X, W) + \lambda_2(\lambda_1 - \lambda_2)g(X, Z)\beta(Y, W) \\ &- \lambda_2(\lambda_1 - \lambda_2)g(Y, Z)\beta(X, W). \end{aligned}$$

For simplicity, we denote by $\text{tr}(\alpha) = \lambda$ and $\text{tr}(\beta) = \mu$.

Let M^n be an n -dimensional submanifold of an $(n+p)$ -dimensional Riemannian manifold N^{n+p} . On the submanifold M we consider the induced generalized semi-symmetric non-metric connection denoted by ∇ and the induced Levi-Civita connection denoted by $\widehat{\nabla}$. Let R and \widehat{R} be the curvature tensors of ∇ and $\widehat{\nabla}$. Decomposing the vector field P on N uniquely into its tangent and normal components P^\top and P^\perp , respectively, then we have $P = P^\top + P^\perp$. The Gauss formulas with respect to ∇ and $\widehat{\nabla}$ can be written as:

$$\overline{\nabla}_X Y = \nabla_X Y + h(X, Y), \quad X, Y \in \Gamma(TM),$$

$$\widehat{\nabla}_X Y = \widehat{\nabla}_X Y + \widehat{h}(X, Y), \quad X, Y \in \Gamma(TM),$$

where \widehat{h} is the second fundamental form of M in N and

$$h(X, Y) = \widehat{h}(X, Y) - \lambda_2 g(X, Y) P^\perp.$$

In N^{n+p} we can choose a local orthonormal frame $e_1, \dots, e_n, e_{n+1}, \dots, e_{n+p}$, such that, restricting to M , e_1, \dots, e_n are tangent to M^n . We write $h_{ij}^r = g(h(e_i, e_j), e_r)$. The squared length of h is $\|h\|^2 = \sum_{i,j=1}^n g(h(e_i, e_j), h(e_i, e_j))$ and the mean curvature vector of M associated to ∇ is $H = \frac{1}{n} \sum_{i=1}^n h(e_i, e_i)$. Similarly, the mean curvature vector of M associated to $\widehat{\nabla}$ is $\widehat{H} = \frac{1}{n} \sum_{i=1}^n \widehat{h}(e_i, e_i)$. We have the Gauss equation

$$\begin{aligned} \overline{R}(X, Y, Z, W) &= R(X, Y, Z, W) - g(h(X, W), h(Y, Z)) + g(h(Y, W), h(X, Z)) \\ (2.3) \quad &+ (\lambda_1 - \lambda_2)g(h(Y, Z), P)g(X, W) + (\lambda_2 - \lambda_1)g(h(X, Z), P)g(Y, W). \end{aligned}$$

Let $\Pi \subset T_x M$, $x \in M$, be a 2-plane section and $\Pi = \text{span}\{e_1, e_2\}$ where e_1, e_2 are orthonormal basis of Π . Since $R(X, Y, Z, W) \neq -R(X, Y, W, Z)$, we define $K(\Pi)$ the sectional curvature of M with the induced connection ∇ as follows:

$$(2.4) \quad K(\Pi) = \frac{1}{2}[R(e_1, e_2, e_2, e_1) - R(e_1, e_2, e_1, e_2)],$$

then $K(\Pi)$ is independent of the choice of e_1, e_2 . For any orthonormal basis $\{e_1, \dots, e_n\}$ of the tangent space $T_x M$, the scalar curvature τ at x is defined by

$$(2.5) \quad \tau(x) = \sum_{1 \leq i < j \leq n} K(e_i \wedge e_j) = \frac{1}{2} \sum_{1 \leq i, j \leq n} R(e_i, e_j, e_j, e_i).$$

Let $\widetilde{\nabla}$ be a linear connection called the second generalized semi-symmetric non-metric connection defined by

$$(2.6) \quad \widetilde{\nabla}_X Y = \widehat{\nabla}_X Y + a\pi(X)Y + b\pi(Y)X.$$

Let \widetilde{R} be the curvature tensor of $\widetilde{\nabla}$.

Let $(\widetilde{\nabla}_X \pi)(Y) = \alpha'(X, Y)$ be a $(0, 2)$ -tensor. The following equality states the relationship between \widetilde{R} and \widehat{R} (see [21] (7)):

$$\begin{aligned}
(2.7) \quad \widetilde{\widehat{R}}(X, Y, Z, W) &= \widehat{R}(X, Y, Z, W) - a\alpha'(Y, X)g(Z, W) + a\alpha'(X, Y)g(Z, W) \\
&\quad - b\alpha'(Y, Z)g(X, W) + b\alpha'(X, Z)g(Y, W) \\
&\quad + b^2\pi(Y)\pi(Z)g(X, W) - b^2\pi(X)\pi(Z)g(Y, W).
\end{aligned}$$

We denote by $\text{tr}(\alpha') = \lambda'$. The Gauss formulas with respect to $\widetilde{\nabla}$ can be written as:

$$\widetilde{\nabla}_X Y = \widetilde{\nabla}_X Y + \widetilde{h}(X, Y), \quad X, Y \in \Gamma(TM).$$

Then

$$\widetilde{h}(X, Y) = \widehat{h}(X, Y).$$

By (25) in [21], we have the Gauss equation

$$\begin{aligned}
(2.8) \quad \widetilde{\widehat{R}}(X, Y, Z, W) &= \widetilde{R}(X, Y, Z, W) - g(\widetilde{h}(X, W), \widetilde{h}(Y, Z)) + g(\widetilde{h}(Y, W), \widetilde{h}(X, Z)) \\
&\quad + b\pi(\widetilde{h}(Y, Z))g(X, W) - b\pi(\widetilde{h}(X, Z))g(Y, W).
\end{aligned}$$

3. CHEN INEQUALITIES FOR SUBMANIFOLDS IN (κ, μ) -CONTACT SPACE FORM WITH THE FIRST GENERALIZED SEMI-SYMMETRIC NON-METRIC CONNECTION.

A $(2m+1)$ -dimensional Riemannian manifold (N^{2m+1}, g) has an almost contact metric structure if it admits a $(1, 1)$ -tensor field φ , a vector field ξ and 1-form η satisfying:

$$\varphi^2 X = -X + \eta(X)\xi, \quad \eta(\xi) = 1, \quad \varphi\xi = 0, \quad \eta \circ \varphi = 0$$

$$g(X, \varphi Y) = -g(\varphi X, Y), \quad g(X, \xi) = \eta(X),$$

for any vector fields X, Y on TN . Let Φ denote the fundamental 2-form in N , given by $\Phi(X, Y) = g(X, \varphi Y)$, for all X, Y on TN . If $\Phi = d\eta$, then N is called a *contact metric manifold*. In a contact metric manifold N , the $(1, 1)$ -tensor field h' defined by $2h' = L_\xi \varphi$ is symmetric and satisfies

$$h'\xi = 0, \quad h'\varphi + \varphi h' = 0, \quad \widehat{\nabla}\xi = -\varphi - \varphi h', \quad \text{trace}(h') = \text{trace}(\varphi h') = 0.$$

The (κ, μ) -nullity distribution of a contact metric manifold N is a distribution

$$\begin{aligned}
N(\kappa, \mu) : p \rightarrow N_p(\kappa, \mu) &= \{Z \in T_p N \mid \widehat{R}(X, Y)Z = \kappa[g(Y, Z)X - g(X, Z)Y] \\
&\quad + \mu[g(Y, Z)h'X - g(X, Z)h'Y]\}
\end{aligned}$$

where κ and μ are constants. If $\xi \in N(\kappa, \mu)$, N is called a (κ, μ) -contact metric manifold. A plane section Π in $T_x N$ is called a φ -section if it is spanned by X and φX , where X is a unit tangent vector field orthogonal to ξ . The sectional curvature of a φ -section is called a φ -sectional curvature. If a (κ, μ) -contact metric manifold N has constant φ -section

curvature c , then it is called a (κ, μ) -contact space form and it is denoted by $N(c)$. The curvature tensor \widehat{R} of $N(c)$ with respect to the Levi-civita connection $\widehat{\nabla}$ is expressed by

$$\begin{aligned}
 (3.1) \quad \widehat{R}(X, Y)Z &= \frac{c+3}{4}[g(Y, Z)X - g(X, Z)Y] + \frac{c+3-4\kappa}{4}[\eta(X)\eta(Z)Y - \eta(Y)\eta(Z)X \\
 &\quad + g(X, Z)\eta(Y)\xi - g(Y, Z)\eta(X)\xi] + \frac{c-1}{4}[2g(X, \varphi Y)\varphi Z + g(X, \varphi Z)\varphi Y \\
 &\quad - g(Y, \varphi Z)\varphi X] + \frac{1}{2}[g(h'Y, Z)h'X - g(h'X, Z)h'Y + g(\varphi h'X, Z)\varphi h'Y \\
 &\quad - g(\varphi h'Y, Z)\varphi h'X] - g(X, Z)h'Y + g(Y, Z)h'X + \eta(X)\eta(Z)h'Y - \eta(Y)\eta(Z)h'X \\
 &\quad - g(h'X, Z)Y + g(h'Y, Z)X - g(h'Y, Z)\eta(X)\xi + g(h'X, Z)\eta(Y)\xi \\
 &\quad + \mu[\eta(Y)\eta(Z)h'X - \eta(X)\eta(Z)h'Y + g(h'Y, Z)\eta(X)\xi - g(h'X, Z)\eta(Y)\xi],
 \end{aligned}$$

for vector fields X, Y, Z on $N(c)$. Let M be an n -dimensional submanifold of a $(2m+1)$ -dimensional (κ, μ) -contact space form of constant φ -sectional curvature $N(c)$. For any tangent vector field X to M , we put

$$\varphi X = \hat{P}X + FX,$$

where $\hat{P}X$ and FX are tangential and normal components of φX respectively and we decompose

$$\xi = \xi^\top + \xi^\perp,$$

where ξ^\top and ξ^\perp denote the tangential and normal parts of ξ . Let $(\varphi h')^\top X$ and $(h')^\top X$ denote the tangential parts of $\varphi h'X$ and $h'X$ respectively. We set

$$\vartheta^2 = \sum_{i,j=1}^n g(e_i, \vartheta e_j)^2, \quad \vartheta \in \{\hat{P}, (\varphi h')^\top, (h')^\top\}.$$

We denote by $\beta(\Pi) = g^2(\hat{P}e_1, e_2) = g^2(\varphi e_1, e_2)$, where $\{e_1, e_2\}$ is an orthonormal basis of a 2-plane section Π . Put

$$\gamma(\Pi) = (\eta(e_1))^2 + (\eta(e_2))^2,$$

$$\theta(\Pi) = \eta(e_1)^2 g((h')^\top e_2, e_2) + \eta(e_2)^2 g((h')^\top e_1, e_1) - 2\eta(e_1)\eta(e_2)g((h')^\top e_1, e_2).$$

Then $\gamma(\Pi)$ and $\theta(\Pi)$ do not depend on the choice of orthonormal basis $\{e_1, e_2\}$. We have the following Chen's inequalities:

Theorem 3.1. *Let M^n , $n \geq 3$, be an n -dimensional submanifold of a $(2m+1)$ -dimensional (κ, μ) -contact space form of constant φ -sectional curvature $N(c)$ endowed with a connection $\overline{\nabla}$, then*

$$\begin{aligned}
\tau(x) - K(\Pi) \leq & \frac{(n+1)(n-2)}{2} \frac{c+3}{4} + \frac{c+3-4\kappa}{4} [-(n-1)\|\xi^\top\|^2 + \gamma(\Pi)] \\
& + \frac{c-1}{8} [3\|\widehat{P}\|^2 - 6\beta(\Pi)] + \frac{1}{4} [\|(\varphi h')^\top\|^2 - \|(h')^\top\|^2 \\
& + |\text{trace}((h')^\top)|^2 - |\text{trace}((\varphi h')^\top)|^2] - \frac{1}{2} [\det(h'|_\Pi) - \det((\varphi h')|_\Pi)] \\
& + (n-1)\text{tr}((h')^\top) - \text{tr}(h'|_\Pi) + (1-\mu)[g(\xi^\top, h'\xi^\top) - \text{tr}((h')^\top)\|\xi^\top\|^2 \\
& + \theta(\Pi)] - \frac{\lambda_1 + \lambda_2}{2}(n-1)\lambda - \frac{\lambda_2}{2}(\lambda_1 - \lambda_2)(n-1)\mu \\
& - \frac{\lambda_1 - \lambda_2}{2}(n-1)n\pi(H) + \frac{\lambda_1 + \lambda_2}{2}\text{tr}(\alpha|_\Pi) \\
& + \frac{\lambda_2(\lambda_1 - \lambda_2)}{2}\text{tr}(\beta|_\Pi) + \frac{\lambda_1 - \lambda_2}{2}g(\text{tr}(h|_\Pi), P) \\
& + \frac{n^2(n-2)}{2(n-1)}\|H\|^2.
\end{aligned}$$

Proof. Let $x \in M$ and $\{e_1, \dots, e_n\}$ and $\{e_{n+1}, \dots, e_{n+p}\}$ be orthonormal basis of $T_x M$ and $T_x^\perp M$ respectively. For $X = W = e_i$, $Y = Z = e_j$, $i \neq j$ by (2.2), (2.3) and (3.1), we have

(3.2)

$$\begin{aligned}
R(e_i, e_j, e_j, e_i) = & \frac{c+3}{4} + \frac{c+3-4\kappa}{4} [-\eta(e_i)^2 - \eta(e_j)^2] \\
& + \frac{c-1}{4} [3g(e_i, \varphi e_j)^2] + \frac{1}{2} [g(e_i, \varphi h' e_j)^2 - g(e_i, h' e_j)^2 \\
& + g(e_i, h' e_i)g(e_j, h' e_j) - g(e_i, \varphi h' e_i)g(e_j, \varphi h' e_j)] + g(e_i, h' e_i) \\
& + 2\eta(e_i)\eta(e_j)g(e_i, h' e_j) - g(e_i, h' e_i)\eta(e_j)^2 - g(e_j, h' e_j)\eta(e_i)^2 + g(e_j, h' e_j) \\
& + \mu[g(e_i, h' e_i)\eta(e_j)^2 + g(e_j, h' e_j)\eta(e_i)^2 - 2\eta(e_i)\eta(e_j)g(e_i, h' e_j)] \\
& - \lambda_1\alpha(e_j, e_j) - \lambda_2\alpha(e_i, e_i) - \lambda_2(\lambda_1 - \lambda_2)\beta(e_i, e_i) \\
& - g(h(e_i, e_j), h(e_i, e_j)) + g(h(e_i, e_i), h(e_j, e_j)) - (\lambda_1 - \lambda_2)g(h(e_j, e_j), P).
\end{aligned}$$

Then

$$\begin{aligned}
 (3.3) \quad \tau(x) &= \frac{1}{2} \sum_{1 \leq i \neq j \leq n} R(e_i, e_j, e_j, e_i) \\
 &= \frac{(n-1)n}{2} \frac{c+3}{4} + \frac{3(c-1)}{8} \|\widehat{P}\|^2 - \frac{c+3-4\kappa}{4} (n-1) \|\xi^\top\|^2 \\
 &\quad + \frac{1}{4} [\|(\varphi h')^\top\|^2 - \|(h')^\top\|^2 + |\text{trace}((h')^\top)|^2 - |\text{trace}((\varphi h')^\top)|^2] \\
 &\quad + (n-1) \text{tr}((h')^\top) + (1-\mu) [g(\xi^\top, h' \xi^\top) - \text{tr}((h')^\top) \|\xi^\top\|^2] \\
 &\quad - \frac{\lambda_1 + \lambda_2}{2} (n-1) \lambda - \frac{\lambda_2}{2} (\lambda_1 - \lambda_2) (n-1) \mu \\
 &\quad - \frac{\lambda_1 - \lambda_2}{2} (n-1) n \pi(H) + \sum_{r=n+1}^{n+p} \sum_{1 \leq i < j \leq n} [h_{ii}^r h_{jj}^r - (h_{ij}^r)^2].
 \end{aligned}$$

Let $\Pi = \text{span}\{e_1, e_2\}$ and by (3.2), we have

$$\begin{aligned}
 (3.4) \quad R(e_1, e_2, e_2, e_1) &= \frac{c+3}{4} - \frac{c+3-4\kappa}{4} \gamma(\Pi) \\
 &\quad + \frac{3(c-1)}{4} \beta(\Pi) + \frac{1}{2} [\det(h'|_\Pi) - \det((\varphi h')|_\Pi)] \\
 &\quad + \text{tr}(h'|_\Pi) + (\mu-1) \theta(\Pi) - \lambda_1 \alpha(e_2, e_2) \\
 &\quad - \lambda_2 \alpha(e_1, e_1) - \lambda_2 (\lambda_1 - \lambda_2) \beta(e_1, e_1) \\
 &\quad - (\lambda_1 - \lambda_2) g(h(e_2, e_2), P) + \sum_{r=n+1}^{n+p} [h_{11}^r h_{22}^r - (h_{12}^r)^2].
 \end{aligned}$$

Similarly, we have

$$\begin{aligned}
 (3.5) \quad R(e_1, e_2, e_1, e_2) &= -\left\{ \frac{c+3}{4} - \frac{c+3-4\kappa}{4} \gamma(\Pi) + \frac{3(c-1)}{4} \beta(\Pi) \right. \\
 &\quad \left. + \frac{1}{2} [\det(h'|_\Pi) - \det((\varphi h')|_\Pi)] + \text{tr}(h'|_\Pi) + (\mu-1) \theta(\Pi) \right\} \\
 &\quad + \lambda_1 \alpha(e_1, e_1) + \lambda_2 \alpha(e_2, e_2) + \lambda_2 (\lambda_1 - \lambda_2) \beta(e_2, e_2) \\
 &\quad + (\lambda_1 - \lambda_2) g(h(e_1, e_1), P) - \sum_{r=n+1}^{n+p} [h_{11}^r h_{22}^r - (h_{12}^r)^2].
 \end{aligned}$$

By (2.4),(3.4) and (3.5), we have

$$\begin{aligned}
 (3.6) \quad K(\Pi) &= \frac{c+3}{4} - \frac{c+3-4\kappa}{4}\gamma(\Pi) + \frac{3(c-1)}{4}\beta(\Pi) \\
 &+ \frac{1}{2}[\det(h'|_{\Pi}) - \det((\varphi h')|_{\Pi})] + \text{tr}(h'|_{\Pi}) + (\mu-1)\theta(\Pi) \\
 &- \frac{\lambda_1 + \lambda_2}{2}\text{tr}(\alpha|_{\Pi}) - \frac{\lambda_2(\lambda_1 - \lambda_2)}{2}\text{tr}(\beta|_{\Pi}) \\
 &- \frac{\lambda_1 - \lambda_2}{2}g(\text{tr}(h|_{\Pi}), P) + \sum_{r=n+1}^{n+p} [h_{11}^r h_{22}^r - (h_{12}^r)^2].
 \end{aligned}$$

By Lemma 2.4 in [35], we get

$$(3.7) \quad \sum_{r=n+1}^{n+p} \left[\sum_{1 \leq i < j \leq n} h_{ii}^r h_{jj}^r - h_{11}^r h_{22}^r - \sum_{1 \leq i < j \leq n} (h_{ij}^r)^2 + (h_{12}^r)^2 \right] \leq \frac{n^2(n-2)}{2(n-1)} \|H\|^2.$$

By (3.3),(3.6) and (3.7) we complete the proof. \square

Similar to Corollaries 3.4 in [35], we have

Corollary 3.2. *If P is a tangent vector field on M , then $h = \hat{h}$. In this case, the equality case of Theorem 3.1 holds at a point $x \in M$ if and only if, with respect to a suitable orthonormal basis $\{e_A\}$ at x , the shape operators $A_r = A_{e_r}$ take the following forms:*

$$A_{n+1} = \begin{pmatrix} h_{11}^{n+1} & 0 & 0 & \cdots & 0 \\ 0 & h_{22}^{n+1} & 0 & \cdots & 0 \\ 0 & 0 & h_{11}^{n+1} + h_{22}^{n+1} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & 0 \\ 0 & 0 & 0 & \cdots & h_{11}^{n+1} + h_{22}^{n+1} \end{pmatrix}$$

and

$$A_r = \begin{pmatrix} h_{11}^r & h_{12}^r & 0 & \cdots & 0 \\ h_{12}^r & -h_{11}^r & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & 0 \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix}, r = n+2, \dots, n+p.$$

Theorem 3.3. *Let M^n , $n \geq 3$, be an n -dimensional submanifold of a $(2m+1)$ -dimensional (κ, μ) -contact space form of constant φ -sectional curvature $N(c)$ endowed with a connection $\overline{\nabla}$, then:*

(i) For each unit vector X in $T_x M$ we have

(3.8)

$$\begin{aligned} \text{Ric}(X) \leq & \frac{(n-1)(c+3)}{4} + \frac{3(c-1)}{4} \|\widehat{P}X\|^2 - \frac{c+3-4\kappa}{4} [(n-2)\eta(X)^2 + \|\xi^\top\|^2] \\ & + \frac{1}{2} [\|(\varphi h'X)^\top\|^2 - \|(h'X)^\top\|^2 + g(X, h'X)\text{tr}((h')^\top) - g(X, \varphi h'X)\text{tr}((\varphi h')^\top)] \\ & + (n-2 - \|\xi^\top\|^2 + \mu\|\xi^\top\|^2)g(X, hX) + (1 - \eta(X)^2 + \mu\eta(X)^2)\text{tr}((h')^\top) \\ & + (2 - 2\mu)\eta(X)g(X, h'(\xi^\top)) - \lambda_1\lambda + [\lambda_1 - \lambda_2(n-1)]\alpha(X, X) \\ & - \lambda_2(\lambda_1 - \lambda_2)(n-1)\beta(X, X) - n(\lambda_1 - \lambda_2)\pi(H) + (\lambda_1 - \lambda_2)\pi(h(X, X)) + \frac{n^2}{4}\|H\|^2. \end{aligned}$$

(ii) If $H(x) = 0$, then a unit tangent vector X at x satisfies the equality case of (3.8) if and only if $X \in \mathcal{N}(x) = \{X \in T_x M | h(X, Y) = 0, \forall Y \in T_x M\}$.

(iii) The equality of (3.8) holds for all unit tangent vector at x if and only if $h_{ij}^r = 0, i, j = 1, 2, \dots, n, r = n+1, \dots, n+p$.

Proof. Let $X \in T_x M$ be a unit tangent vector at x . We choose the orthonormal basis $\{e_1, \dots, e_n\}$ such that $e_1 = X$. Then by (3.2), we have

$$\begin{aligned} (3.9) \quad \text{Ric}(X) = & \sum_{j=2}^n R(e_1, e_j, e_j, e_1) = \frac{(n-1)(c+3)}{4} + \frac{3(c-1)}{4} \|\widehat{P}X\|^2 \\ & - \frac{c+3-4\kappa}{4} [(n-2)\eta(X)^2 + \|\xi^\top\|^2] + \frac{1}{2} [\|(\varphi h'X)^\top\|^2 \\ & - \|(h'X)^\top\|^2 + g(X, h'X)\text{tr}((h')^\top) - g(X, \varphi h'X)\text{tr}((\varphi h')^\top)] \\ & + (n-2 - \|\xi^\top\|^2 + \mu\|\xi^\top\|^2)g(X, hX) + (1 - \eta(X)^2 + \mu\eta(X)^2)\text{tr}((h')^\top) \\ & + (2 - 2\mu)\eta(X)g(X, h'(\xi^\top)) - \lambda_1\lambda + [\lambda_1 - \lambda_2(n-1)]\alpha(X, X) \\ & - \lambda_2(\lambda_1 - \lambda_2)(n-1)\beta(X, X) - n(\lambda_1 - \lambda_2)\pi(H) + (\lambda_1 - \lambda_2)\pi(h(X, X)) \\ & + \sum_{r=n+1}^{n+p} \sum_{j=2}^n [h_{11}^r h_{jj}^r - (h_{1j}^r)^2]. \end{aligned}$$

By Lemma 2.5 in [35], we get

$$\sum_{r=n+1}^{n+p} \sum_{j=2}^n h_{11}^r h_{jj}^r \leq \frac{n^2}{4} \|H\|^2,$$

which together with (3.9) gives (3.8). The Proofs of (ii) and (iii) are same as the proofs of (ii) and (iii) in Theorem 5.2 in [35]. \square

Let L be a k -plane section of $T_x M$, $x \in M$ and X a unit vector in L . We choose an orthonormal frame e_1, \dots, e_k of L such that $e_1 = X$. In [7], Chen defined the k -Ricci

curvature of L at X by

$$\text{Ric}_L(X) = K_{12} + K_{13} + \cdots + K_{1k}.$$

For an integer k , $2 \leq k \leq n$, the Riemannian invariant Θ_k on M defined by

$$\Theta_k(x) = \frac{1}{k-1} \inf \{ \text{Ric}_L(X) | L, X \}, \quad x \in M,$$

where L runs over all k -plane sections in $T_x M$ and X runs over all unit vectors in L . We have

$$(3.10) \quad \tau(x) \geq \frac{n(n-1)}{2} \Theta_k(x).$$

Theorem 3.4. *Let M^n , $n \geq 3$, be an n -dimensional submanifold of a $(2m+1)$ -dimensional (κ, μ) -contact space form of constant φ -sectional curvature $N(c)$ endowed with a connection $\bar{\nabla}$, then: for any integer k , $2 \leq k \leq n$, and any point $x \in M$, we have*

$$(3.11) \quad \begin{aligned} n(n-1) \|H\|^2(x) &\geq n(n-1) \Theta_k(x) - \frac{c+3}{4}(n-1)n - \frac{3(c-1)}{4} \|\widehat{P}\|^2 + \frac{c+3-4\kappa}{2}(n-1) \|\xi^\top\|^2 \\ &\quad - \frac{1}{2} [\|(\varphi h')^\top\|^2 - \|(h')^\top\|^2 + |\text{trace}((h')^\top)|^2 - |\text{trace}((\varphi h')^\top)|^2] \\ &\quad - 2(n-1) \text{tr}((h')^\top) - 2(1-\mu) [g(\xi^\top, h' \xi^\top) - \text{tr}((h')^\top) \|\xi^\top\|^2] \\ &\quad + \lambda(n-1)(\lambda_1 + \lambda_2) + \lambda_2(\lambda_1 - \lambda_2)(n-1)\mu + (n-1)n(\lambda_1 - \lambda_2)\pi(H). \end{aligned}$$

Proof. We choose the orthonormal frame $\{e_1, \dots, e_n, e_{n+1}, \dots, e_{n+p}\}$ at x such that the e_{n+1} is in the direction of the mean curvature vector $H(x)$ and $\{e_1, \dots, e_n\}$ diagonalize the shape operator A_{n+1} . Then the shape operators take the following form

$$(3.12) \quad A_{n+1} = \text{diag}(a_1, a_2, \dots, a_n), \quad \text{trace } A_r = 0, \quad r = n+2, \dots, n+p.$$

By (3.3), we have

$$(3.13) \quad \begin{aligned} 2\tau - \frac{c+3}{4}(n-1)n - \frac{3(c-1)}{4} \|\widehat{P}\|^2 + \frac{c+3-4\kappa}{2}(n-1) \|\xi^\top\|^2 \\ - \frac{1}{2} [\|(\varphi h')^\top\|^2 - \|(h')^\top\|^2 + |\text{trace}((h')^\top)|^2 - |\text{trace}((\varphi h')^\top)|^2] \\ - 2(n-1) \text{tr}((h')^\top) - 2(1-\mu) [g(\xi^\top, h' \xi^\top) - \text{tr}((h')^\top) \|\xi^\top\|^2] \\ + \lambda(n-1)(\lambda_1 + \lambda_2) + \lambda_2(\lambda_1 - \lambda_2)(n-1)\mu + (n-1)n(\lambda_1 - \lambda_2)\pi(H) \\ = n^2 \|H\|^2 - \|h\|^2. \end{aligned}$$

By (3.12), we get

$$\|h\|^2 = \sum_{j=1}^n a_j^2 + \sum_{r=n+2}^{n+p} \sum_{1 \leq i, j \leq l} (h_{ij}^r)^2.$$

Since $\sum_{j=1}^n a_j^2 \geq n \|H\|^2$, by (3.10) and (3.13), we completes the proof. \square

Nextly, we prove Casorati inequalities. We denote the Casorati curvature with respect to $\overline{\nabla}$ by

$$\mathcal{C} = \frac{1}{n} \sum_{r=n+1}^{n+p} \sum_{i,j=1}^n (h_{ij}^r)^2.$$

Suppose now that L is an l -dimensional subspace of $T_x M$, $l \geq 2$, and $\{e_1, \dots, e_l\}$ be an orthonormal basis of L . Then the Casorati curvature of the l -plane section L with respect to $\overline{\nabla}$ is given by

$$\mathcal{C}(L) = \frac{1}{l} \sum_{r=n+1}^{n+p} \sum_{i,j=1}^l (h_{ij}^r)^2.$$

Then normalized δ -Casorati curvature $\delta_c(n-1)$ and $\hat{\delta}_c(n-1)$ with respect to $\overline{\nabla}$ are given by

$$[\delta_c(n-1)]_x = \frac{1}{2}\mathcal{C}_x + \frac{n+1}{2n} \inf\{\mathcal{C}(L) | L \text{ a hyperplane of } T_x M\},$$

$$[\hat{\delta}_c(n-1)]_x = 2\mathcal{C}_x - \frac{2n-1}{2n} \sup\{\mathcal{C}(L) | L \text{ a hyperplane of } T_x M\}.$$

Theorem 3.5. *Let M^n , $n \geq 3$, be an n -dimensional submanifold of a $(2m+1)$ -dimensional (κ, μ) -contact space form of constant φ -sectional curvature $N(c)$ endowed with a connection $\overline{\nabla}$, then:*

(i) *The normalized δ -Casorati curvature $\delta_c(n-1)$ satisfies*

(3.14)

$$\begin{aligned} 2\tau \leq & n(n-1)\delta_c(n-1) + \frac{c+3}{4}(n-1)n + \frac{3(c-1)}{4} \|\widehat{P}\|^2 - \frac{c+3-4\kappa}{2}(n-1) \|\xi^\top\|^2 \\ & + \frac{1}{2} [\|(\varphi h')^\top\|^2 - \|(h')^\top\|^2 + |\text{trace}((h')^\top)|^2 - |\text{trace}((\varphi h')^\top)|^2] \\ & + 2(n-1)\text{tr}((h')^\top) + 2(1-\mu)[g(\xi^\top, h'\xi^\top) - \text{tr}((h')^\top) \|\xi^\top\|^2] \\ & - \lambda(n-1)(\lambda_1 + \lambda_2) - \lambda_2(\lambda_1 - \lambda_2)(n-1)\mu - (n-1)n(\lambda_1 - \lambda_2)\pi(H). \end{aligned}$$

Moreover, if P is tangent to M , the equality case of (3.14) holds if and only if M is an invariantly quasi-umbilical submanifold in N , such that with respect to suitable orthonormal tangent frame $\{e_1, \dots, e_n\}$ and normal orthonormal frame $\{e_{n+1}, \dots, e_{n+p}\}$, the shape operator $A_r = A_{e_r}$, $r \in \{n+1, \dots, n+p\}$, taking the following forms:

$$A_{n+1} = \text{diag}(a, a, \dots, a, 2a), \quad A_{n+2} = \dots = A_{n+p} = 0.$$

(ii) The normalized δ -Casorati curvature $\hat{\delta}_c(n-1)$ satisfies

(3.15)

$$\begin{aligned} 2\tau \leq & n(n-1)\hat{\delta}_c(n-1) + \frac{c+3}{4}(n-1)n + \frac{3(c-1)}{4}\|\widehat{P}\|^2 - \frac{c+3-4\kappa}{2}(n-1)\|\xi^\top\|^2 \\ & + \frac{1}{2}[|(\varphi h')^\top|^2 - |(h')^\top|^2 + |\text{trace}((h')^\top)|^2 - |\text{trace}((\varphi h')^\top)|^2] \\ & + 2(n-1)\text{tr}((h')^\top) + 2(1-\mu)[g(\xi^\top, h'\xi^\top) - \text{tr}((h')^\top)\|\xi^\top\|^2] \\ & - \lambda(n-1)(\lambda_1 + \lambda_2) - \lambda_2(\lambda_1 - \lambda_2)(n-1)\mu - (n-1)n(\lambda_1 - \lambda_2)\pi(H). \end{aligned}$$

Moreover, if P is tangent to M , the equality case of (3.14) holds if and only if M is an invariantly quasi-umbilical submanifold in N , such that with respect to suitable orthonormal tangent frame $\{e_1, \dots, e_n\}$ and normal orthonormal frame $\{e_{n+1}, \dots, e_{n+p}\}$, the shape operator $A_r = A_{e_r}$, $r \in \{n+1, \dots, n+p\}$, taking the following forms:

$$A_{n+1} = \text{diag}(2a, 2a, \dots, 2a, a), \quad A_{n+2} = \dots = A_{n+p} = 0.$$

Proof. By (3.3), we have

$$\begin{aligned} (3.16) \quad 2\tau = & \frac{c+3}{4}(n-1)n + \frac{3(c-1)}{4}\|\widehat{P}\|^2 - \frac{c+3-4\kappa}{2}(n-1)\|\xi^\top\|^2 \\ & + \frac{1}{2}[|(\varphi h')^\top|^2 - |(h')^\top|^2 + |\text{trace}((h')^\top)|^2 - |\text{trace}((\varphi h')^\top)|^2] \\ & + 2(n-1)\text{tr}((h')^\top) + 2(1-\mu)[g(\xi^\top, h'\xi^\top) - \text{tr}((h')^\top)\|\xi^\top\|^2] \\ & - \lambda(n-1)(\lambda_1 + \lambda_2) - \lambda_2(\lambda_1 - \lambda_2)(n-1)\mu \\ & - (n-1)n(\lambda_1 - \lambda_2)\pi(H) + n^2\|H\|^2 - n\mathcal{C}. \end{aligned}$$

Let

$$\begin{aligned} (3.17) \quad Q = & \frac{1}{2}n(n-1)\mathcal{C} + \frac{(n-1)(n+1)}{2}\mathcal{C}(L) - 2\tau + \frac{c+3}{4}(n-1)n \\ & + \frac{3(c-1)}{4}\|\widehat{P}\|^2 - \frac{c+3-4\kappa}{2}(n-1)\|\xi^\top\|^2 \\ & + \frac{1}{2}[|(\varphi h')^\top|^2 - |(h')^\top|^2 + |\text{trace}((h')^\top)|^2 - |\text{trace}((\varphi h')^\top)|^2] \\ & + 2(n-1)\text{tr}((h')^\top) + 2(1-\mu)[g(\xi^\top, h'\xi^\top) - \text{tr}((h')^\top)\|\xi^\top\|^2] \\ & - \lambda(n-1)(\lambda_1 + \lambda_2) - \lambda_2(\lambda_1 - \lambda_2)(n-1)\mu \\ & - (n-1)n(\lambda_1 - \lambda_2)\pi(H). \end{aligned}$$

By Lemma 4 in [34], we have $Q \geq 0$, so we get (3.14). Similar to the proof of Theorem 3 in [34], we can prove Theorem 3.5. \square

4. CHEN INEQUALITIES FOR SUBMANIFOLDS IN (κ, μ) -CONTACT SPACE FORM WITH THE SECOND GENERALIZED SEMI-SYMMETRIC NON-METRIC CONNECTION.

Let $\tilde{\tau}$ and $\tilde{K}(\Pi)$ be the scalar curvature and the section curvature with respect to the connection $\tilde{\nabla}$ respectively. Similar to Theorem 3.1, we have

Theorem 4.1. *Let M^n , $n \geq 3$, be an n -dimensional submanifold of a $(2m+1)$ -dimensional (κ, μ) -contact space form of constant φ -sectional curvature $N(c)$ endowed with a connection $\tilde{\nabla}$, then*

$$\begin{aligned} \tilde{\tau}(x) - \tilde{K}(\Pi) &\leq \frac{(n+1)(n-2)}{2} \frac{c+3}{4} + \frac{c+3-4\kappa}{4} [-(n-1)\|\xi^\top\|^2 + \gamma(\Pi)] \\ &\quad + \frac{c-1}{8} [3\|\hat{P}\|^2 - 6\beta(\Pi)] + \frac{1}{4} [\|(\varphi h')^\top\|^2 - \|(h')^\top\|^2] \\ &\quad + |\text{trace}((h')^\top)|^2 - |\text{trace}((\varphi h')^\top)|^2 - \frac{1}{2} [\det(h'|_\Pi) - \det((\varphi h')|_\Pi)] \\ &\quad + (n-1)\text{tr}((h')^\top) - \text{tr}(h'|_\Pi) + (1-\mu)[g(\xi^\top, h'\xi^\top) - \text{tr}((h')^\top)\|\xi^\top\|^2] \\ &\quad + \theta(\Pi)] - \frac{n-1}{2} b\lambda' + \frac{n-1}{2} b^2\|P^\top\|^2 \\ &\quad - \frac{b}{2}(n-1)n\pi(\tilde{H}) + \frac{b}{2}\text{tr}(\alpha'|_\Pi) - \frac{b^2}{2}\|P_\Pi\|^2 \\ &\quad + \frac{b}{2}\pi(\text{tr}(\tilde{h}|_\Pi)) + \frac{n^2(n-2)}{2(n-1)}\|\tilde{H}\|^2. \end{aligned}$$

Proof. Let $x \in M$ and $\{e_1, \dots, e_n\}$ and $\{e_{n+1}, \dots, e_{n+p}\}$ be orthonormal basis of $T_x M$ and $T_x^\perp M$ respectively. For $X = W = e_i$, $Y = Z = e_j$, $i \neq j$ by (2.7), (2.8) and (3.1), we have

(4.1)

$$\begin{aligned} \tilde{R}(e_i, e_j, e_j, e_i) &= \frac{c+3}{4} + \frac{c+3-4\kappa}{4} [-\eta(e_i)^2 - \eta(e_j)^2] \\ &\quad + \frac{c-1}{4} [3g(e_i, \varphi e_j)^2] + \frac{1}{2} [g(e_i, \varphi h' e_j)^2 - g(e_i, h' e_j)^2] \\ &\quad + g(e_i, h' e_i)g(e_j, h' e_j) - g(e_i, \varphi h' e_i)g(e_j, \varphi h' e_j) + g(e_i, h' e_i) \\ &\quad + 2\eta(e_i)\eta(e_j)g(e_i, h' e_j) - g(e_i, h' e_i)\eta(e_j)^2 - g(e_j, h' e_j)\eta(e_i)^2 + g(e_j, h' e_j) \\ &\quad + \mu[g(e_i, h' e_i)\eta(e_j)^2 + g(e_j, h' e_j)\eta(e_i)^2 - 2\eta(e_i)\eta(e_j)g(e_i, h' e_j)] \\ &\quad - b\alpha'(e_j, e_j) + b^2\pi(e_j)^2 - b\pi(\tilde{h}(e_j, e_j)) + \sum_{r=n+1}^{n+p} [\tilde{h}_{ii}^r \tilde{h}_{jj}^r - (\tilde{h}_{ij}^r)^2]. \end{aligned}$$

Then

$$\begin{aligned}
(4.2) \quad \tilde{\tau}(x) = & \frac{(n-1)n}{2} \frac{c+3}{4} + \frac{3(c-1)}{8} \|\widehat{P}\|^2 - \frac{c+3-4\kappa}{4} (n-1) \|\xi^\top\|^2 \\
& + \frac{1}{4} [\|(\varphi h')^\top\|^2 - \|(h')^\top\|^2 + |\text{trace}((h')^\top)|^2 - |\text{trace}((\varphi h')^\top)|^2] \\
& + (n-1) \text{tr}((h')^\top) + (1-\mu) [g(\xi^\top, h' \xi^\top) - \text{tr}((h')^\top) \|\xi^\top\|^2] \\
& - \frac{n-1}{2} b \lambda' + \frac{n-1}{2} b^2 \|P^\top\|^2 - \frac{b}{2} (n-1) n \pi(\tilde{H}) \\
& + \sum_{r=n+1}^{n+p} \sum_{1 \leq i < j \leq n} [\tilde{h}_{ii}^r \tilde{h}_{jj}^r - (\tilde{h}_{ij}^r)^2].
\end{aligned}$$

Let $\Pi = \text{span}\{e_1, e_2\}$ and by (4.1), we have

$$\begin{aligned}
(4.3) \quad \tilde{R}(e_1, e_2, e_2, e_1) = & \frac{c+3}{4} - \frac{c+3-4\kappa}{4} \gamma(\Pi) \\
& + \frac{3(c-1)}{4} \beta(\Pi) + \frac{1}{2} [\det(h'|_\Pi) - \det((\varphi h')|_\Pi)] \\
& + \text{tr}(h'|_\Pi) + (\mu-1) \theta(\Pi) - b \alpha'(e_2, e_2) + b^2 \pi(e_2)^2 \\
& - b \pi(\tilde{h}(e_2, e_2)) + \sum_{r=n+1}^{n+p} [\tilde{h}_{11}^r \tilde{h}_{22}^r - (\tilde{h}_{12}^r)^2]
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
(4.4) \quad \tilde{R}(e_1, e_2, e_1, e_2) = & - \left\{ \frac{c+3}{4} - \frac{c+3-4\kappa}{4} \gamma(\Pi) + \frac{3(c-1)}{4} \beta(\Pi) \right. \\
& + \frac{1}{2} [\det(h'|_\Pi) - \det((\varphi h')|_\Pi)] + \text{tr}(h'|_\Pi) + (\mu-1) \theta(\Pi) \Big\} \\
& + b \alpha'(e_1, e_1) - b^2 \pi(e_1)^2 + b \pi(\tilde{h}(e_1, e_1)) \\
& - \sum_{r=n+1}^{n+p} [\tilde{h}_{11}^r \tilde{h}_{22}^r - (\tilde{h}_{12}^r)^2]
\end{aligned}$$

By (2.4), (4.3) and (4.4), we have

$$\begin{aligned}
(4.5) \quad \tilde{K}(\Pi) = & \frac{c+3}{4} - \frac{c+3-4\kappa}{4} \gamma(\Pi) + \frac{3(c-1)}{4} \beta(\Pi) \\
& + \frac{1}{2} [\det(h'|_\Pi) - \det((\varphi h')|_\Pi)] + \text{tr}(h'|_\Pi) + (\mu-1) \theta(\Pi) \\
& - \frac{b}{2} \text{tr}(\alpha'|_\Pi) + \frac{b^2}{2} \|P_\Pi\|^2 - \frac{b}{2} \pi(\text{tr}(\tilde{h}|_\Pi)) \\
& + \sum_{r=n+1}^{n+p} [\tilde{h}_{11}^r \tilde{h}_{22}^r - (\tilde{h}_{12}^r)^2].
\end{aligned}$$

By (4.2), (4.5) and (3.7) we complete the proof. \square

Similarly to Theorem 3.3, we have

Theorem 4.2. *Let M^n , $n \geq 3$, be an n -dimensional submanifold of a $(2m+1)$ -dimensional (κ, μ) -contact space form of constant φ -sectional curvature $N(c)$ endowed with a connection $\widetilde{\nabla}$, then: for each unit vector X in $T_x M$ we have*

$$\begin{aligned}
 (4.6) \quad \widetilde{\text{Ric}}(X) &\leq \frac{(n-1)(c+3)}{4} + \frac{3(c-1)}{4} \|\widehat{P}X\|^2 - \frac{c+3-4\kappa}{4} [(n-2)\eta(X)^2 + \|\xi^\top\|^2] \\
 &\quad + \frac{1}{2} [\|(\varphi h'X)^\top\|^2 - \|(h'X)^\top\|^2 + g(X, h'X) \text{tr}((h')^\top) - g(X, \varphi h'X) \text{tr}((\varphi h')^\top)] \\
 &\quad + (n-2 - \|\xi^\top\|^2 + \mu \|\xi^\top\|^2) g(X, hX) + (1 - \eta(X)^2 + \mu \eta(X)^2) \text{tr}((h')^\top) \\
 &\quad + (2 - 2\mu) \eta(X) g(X, h'(\xi^\top)) - b\lambda + b\alpha'(X, X) + b^2 \|P^\top\|^2 \\
 &\quad - b^2 \pi(X)^2 - nb\pi(\widetilde{H}) + b\pi(\widetilde{h}(X, X)) + \frac{n^2}{4} \|\widetilde{H}\|^2.
 \end{aligned}$$

Similarly to Theorem 3.4, we have

Theorem 4.3. *Let M^n , $n \geq 3$, be an n -dimensional submanifold of a $(2m+1)$ -dimensional (κ, μ) -contact space form of constant φ -sectional curvature $N(c)$ endowed with a connection $\widetilde{\nabla}$, then: for any integer k , $2 \leq k \leq n$, and any point $x \in M$, we have*

$$\begin{aligned}
 (4.7) \quad n(n-1) \|\widetilde{H}\|^2(x) &\geq n(n-1) \widetilde{\Theta}_k(x) - \frac{c+3}{4} (n-1)n - \frac{3(c-1)}{4} \|\widehat{P}\|^2 + \frac{c+3-4\kappa}{2} (n-1) \|\xi^\top\|^2 \\
 &\quad - \frac{1}{2} [\|(\varphi h')^\top\|^2 - \|(h')^\top\|^2 + |\text{trace}((h')^\top)|^2 - |\text{trace}((\varphi h')^\top)|^2] \\
 &\quad - 2(n-1) \text{tr}((h')^\top) - 2(1-\mu) [g(\xi^\top, h'\xi^\top) - \text{tr}((h')^\top) \|\xi^\top\|^2] \\
 &\quad + (n-1)b\lambda' - (n-1)b^2 \|P^\top\|^2 + b(n-1)n\pi(\widetilde{H}).
 \end{aligned}$$

We denote the Casorati curvature with respect to $\widetilde{\nabla}$ by $\widetilde{\mathcal{C}}$. We denote the normalized δ -Casorati curvature by $\widetilde{\delta}_c(n-1)$ and $\widehat{\delta}_c(n-1)$ with respect to $\widetilde{\nabla}$. Similarly to Theorem 3.5, we have

Theorem 4.4. *Let M^n , $n \geq 3$, be an n -dimensional submanifold of a $(2m+1)$ -dimensional (κ, μ) -contact space form of constant φ -sectional curvature $N(c)$ endowed with a connection $\widetilde{\nabla}$, then:*

(i) *The normalized δ -Casorati curvature $\tilde{\delta}_c(n-1)$ satisfies*
(4.8)

$$\begin{aligned} 2\tilde{\tau} \leq & n(n-1)\tilde{\delta}_c(n-1) + \frac{c+3}{4}(n-1)n + \frac{3(c-1)}{4}\|\widehat{P}\|^2 - \frac{c+3-4\kappa}{2}(n-1)\|\xi^\top\|^2 \\ & + \frac{1}{2}[|(\varphi h')^\top|^2 - |(h')^\top|^2 + |\text{trace}((h')^\top)|^2 - |\text{trace}((\varphi h')^\top)|^2] \\ & + 2(n-1)\text{tr}((h')^\top) + 2(1-\mu)[g(\xi^\top, h'\xi^\top) - \text{tr}((h')^\top)\|\xi^\top\|^2] \\ & - (n-1)b\lambda' + (n-1)b^2\|P^\top\|^2 - b(n-1)n\pi(\tilde{H}). \end{aligned}$$

(ii) *The normalized δ -Casorati curvature $\tilde{\delta}_c(n-1)$ satisfies*
(4.9)

$$\begin{aligned} 2\tilde{\tau} \leq & n(n-1)\tilde{\delta}_c(n-1) + \frac{c+3}{4}(n-1)n + \frac{3(c-1)}{4}\|\widehat{P}\|^2 - \frac{c+3-4\kappa}{2}(n-1)\|\xi^\top\|^2 \\ & + \frac{1}{2}[|(\varphi h')^\top|^2 - |(h')^\top|^2 + |\text{trace}((h')^\top)|^2 - |\text{trace}((\varphi h')^\top)|^2] \\ & + 2(n-1)\text{tr}((h')^\top) + 2(1-\mu)[g(\xi^\top, h'\xi^\top) - \text{tr}((h')^\top)\|\xi^\top\|^2] \\ & - (n-1)b\lambda' + (n-1)b^2\|P^\top\|^2 - b(n-1)n\pi(\tilde{H}). \end{aligned}$$

5. ACKNOWLEDGEMENTS

The author was supported in part by NSFC No.11771070.

REFERENCES

- [1] N.S. Agashe and M.R. Chafle, *A semi-symmetric non-metric connection on a Riemannian manifold*, Indian J. Pure Appl. Math. **23** (1992), 399-409.
- [2] N.S. Agashe and M.R. Chafle, *On submanifolds of a Riemannian manifold with a semi-symmetric non-metric connection*, Tensor **55** (1994), 120-130.
- [3] A. Ahmad, G. He, W. Tang and P. Zhao, *Chen's inequalities for submanifolds in (κ, μ) -contact space form with a semi-symmetric metric connection*, Open Math. **16** (2018), 380-391.
- [4] A. Ahmad, F. Shahzad and J. Li, *Chen's inequalities for submanifolds in (κ, μ) -contact space form with a semi-symmetric non-metric connection*, J. Appl. Math. Phys., bf 6 (2018), 389-404.
- [5] M. Aquib M, J.W. Lee, G.E. Vilcu, D.W. Yoon, *Classification of Casorati ideal Lagrangian submanifolds in complex space forms*, Differ.Geom. Appl. **63** (2019), 30-49.
- [6] B.Y. Chen, *Some pinching and classification theorems for minimal submanifolds*, Arch. Math. (Basel) **60** (1993), 568-578.
- [7] B.Y. Chen, *Relations between Ricci curvature and shape operator for submanifolds with arbitrary codimensions*, Glasg. Math. J. **41** (1999), 33-41.
- [8] B.Y. Chen, *Mean curvature and shape operator of isometric immersions in real space forms*, Glasg. Math. J. **38** (1996), 87-97.
- [9] B.Y. Chen, *A Riemannian invariant and its applications to Einstein manifolds*, Bull. Austral. Math. Soc. **70** (2004), 55-65.

- [10] B.Y. Chen, *A general optimal inequality for warped products in complex projective spaces and its applications*, Proc. Japan Acad. Ser. A Math. Sci. **79** (2002), 96-100.
- [11] P.J. De Smet, F. Dillen, L. Verstraelen and L. Vrancken, *A pointwise inequality in submanifold theory*, Arch. Math. (Brno) **35** (1999), 115-128.
- [12] S. Decu, M. Petrovie-Torgasev, A. Sebekovic and L. Verstraelen, *Ricci and Casorati principal directions of Wintgen ideal submanifolds*, Filomat **28** (2014), 657-661.
- [13] J. Ge and Z. Tang, *A proof of the DDVV conjecture and its equality case*, Pacific J. Math. **237** (2008), 87-95.
- [14] M. Gülbahar, E. Kılıç, S. Keleş and M.M. Tripathi, *Some basic inequalities for submanifolds of nearly quasi-constant curvature manifolds*, Differ. Geom. Dyn. Syst. **16** (2014), 156-167.
- [15] H.A. Hayden, *Subspaces of a space with torsion*, Proc. London Math. Soc. **34** (1932), 27-50.
- [16] T. Imai, *Hypersurfaces of a Riemannian manifold with semi-symmetric metric connection*, Tensor **23** (1972), 300-306.
- [17] T. Imai, *Notes on semi-symmetric metric connections*, Tensor **24** (1972), 293-296.
- [18] C.W. Lee, J.W. Lee, *Inequalities on Sasakian Statistical Manifolds in Terms of Casorati Curvatures*, Mathematics **6** (2018), 259.
- [19] C.W. Lee, J.W. Lee, G.E. Vilcu, *Optimal inequalities for the normalized δ -Casorati curvatures of submanifolds in Kenmotsu space forms*, Advances in Geometry **17** (2017), 355-362.
- [20] J.W. Lee and G.E. Vilcu, *Inequalities for generalized normalized δ -Casorati curvatures of slant submanifolds in quaternionic space forms*, Taiwan. J. Math. **19** (2015), 691-702.
- [21] J. Li, G. He, P. Zhao, *On submanifolds in a Riemannian manifold with a semi-symmetric non-metric connection*, Symmetry **9** (2017), no. 7, Paper No. 112, 10 pp.
- [22] M.A. Lone, M.H. Shahid, G.E. Vilcu, *On Casorati curvatures of submanifolds in pointwise Kenmotsu space forms*, Mathematical Physics, Analysis and Geometry **22** (2019), 2.
- [23] A. Mihai and C. Özgür, *Chen inequalities for submanifolds of real space forms with a semi-symmetric metric connection*, Taiwanese J. Math. **14** (2010), 1465-1477.
- [24] A. Mihai and C. Özgür, *Chen inequalities for submanifolds of complex space forms and Sasakian space forms endowed with semi-symmetric metric connections*, Rocky Mountain J. Math. **41** (2011), no. 5, 1653-1673.
- [25] Z. Nakao, *Submanifolds of a Riemannian manifold with semisymmetric metric connections*. Proc. Amer. Math. Soc. **54** (1976), 261-266.
- [26] C. Özgür, A. Mihai, *Chen inequalities for submanifolds of real space forms with a semi-symmetric non-metric connection*, Canad. Math. Bull. **55** (2012), 611-622.
- [27] K.S. Park, *Inequalities for the Casorati curvatures of real hypersurfaces in some Grassmannians*, Taiwanese J. Math. **22** (2018), 63-77.
- [28] Q. Qu, Y. Wang, *Multiply warped products with a quarter-symmetric connection*, J. Math. Anal. Appl. **431** (2015), 955-987.
- [29] M.M. Tripathi, M. Gülbahar, E. Kılıç, S. Keleş, *Inequalities for scalar curvature of pseudo-Riemannian submanifolds*, J. Geome. Phys. **112** (2017), 74-84.
- [30] G.E. Vilcu, *An optimal inequality for Lagrangian submanifolds in complex space forms involving Casorati curvature*, J. Math. Anal. Appl. **465** (2018), 1209-1222.
- [31] K. Yano, *On semi-symmetric metric connection*. Rev. Roumaine Math. Pures Appl. **15** (1970), 1579-1586.
- [32] H. Yıldırım, L. Vrancken, *$\delta^{\sharp}(2, 2)$ -Ideal centroaffine hypersurfaces of dimension 5*, Taiwanese J. Math. **21** (2017), 283-304.
- [33] P. Zhang, X. Pan, L. Zhang, *Inequalities for submanifolds of a Riemannian manifold of nearly quasi-constant curvature with a semi-symmetric non-metric connection*, Rev. Un. Mat. Argentina **56** (2015), 1-19.

- [34] P. Zhang, L. Zhang, *Casorati inequalities for submanifolds in a Riemannian manifold of quasi-constant curvature with a semi-symmetric metric connection*, Symmetry **8** (2016), no. 4, Art. 19, 9 pp.
- [35] P. Zhang, L. Zhang, W. Song, *Chen's inequalities for submanifolds of a Riemannian manifold of quasi-constant curvature with a semi-symmetric metric connection*, Taiwanese J. Math. **18** (2014), 1841-1862.

Y. Wang

School of Mathematics and Statistics, Northeast Normal University, Changchun 130024, China

Email: wangy581@nenu.edu.cn