

FOUNDATIONS OF FINSLER SPACETIMES FROM THE OBSERVERS' VIEWPOINT

ANTONIO N. BERNAL, MIGUEL ANGEL JAVALOYES, AND MIGUEL SÁNCHEZ

ABSTRACT. Physical foundations for relativistic spacetimes are revisited, in order to check at what extent Finsler spacetimes lie in their framework. Arguments based on inertial observers (as in the foundations of Special Relativity and Classical Mechanics) are shown to correspond with a double linear approximation in the measurement of space and time. While General Relativity appears by dropping the first linearization, Finsler spacetimes appear by dropping the second one. The classical Ehlers-Pirani-Schild approach is carefully discussed and shown to be compatible with the Lorentz-Finsler case. The precise mathematical definition of Finsler spacetime is discussed by using the *space of observers*. Special care is taken in some issues such as: the fact that a Lorentz-Finsler metric would be physically measurable only on the causal directions for a cone structure, the implications for models of spacetimes of some apparently innocuous hypotheses on differentiability, or the possibilities of measurement of a varying speed of light.

MSC: 53C60, 83D05 83A05, 83C05.

Keywords: Finsler spacetime, Ehlers-Pirani-Schild approach, Lorentz symmetry breaking, Very Special Relativity, Signature-changing spacetimes.

CONTENTS

1. Introduction	2
2. The doubly linearized models	5
2.1. Postulates	5
2.2. Groups $O^{(k)}(4, \mathbb{R})$	6
2.3. Linear models of spacetimes	9
2.4. Temporal models and interpretation of $k = -c^2$	11
3. First non-linearization	12
3.1. General case and signature change	13
3.2. Space of observers	14
3.3. Pointwise variation of speed of light	14
3.4. Relativistic vs Leibnizian structures	15
4. Second non-linearization	17
4.1. Background: norms, cones and Lorentz-Finsler metrics	17
4.2. Physical intuitions for Finsler spacetimes	24
5. Comparison with Ehlers-Pirani-Schild approach	26
5.1. Summary of the approach	26
5.2. Keys of compatibility with Finslerian spacetimes	27
5.3. Constructive EPS approach vs observer's approach	32
6. Lorentz symmetry breaking	33

6.1. Modified Special Relativity	34
6.2. Anisotropic speed of light	36
6.3. Matter as anisotropy and Quantum Physics	39
7. Conclusions	40
Acknowledgments	43
References	43

1. INTRODUCTION

A plethora of alternatives to classical General Relativity has been developed since its very beginning. Many of them were motivated by the search of a unified theory which solved disturbing issues of compatibility with Quantum Mechanics (Kaluza-Klein, M-theory, quantum field gravity...) while, since the 90's, unexpected cosmological measurements led to further alternatives (cosmological constant, quintaessence, theories with varying speed of light...). However, the possibility to consider a Finslerian modification of GR has not settled in the mainstream of research and it has been scarcely considered in the literature until recent times (some examples are references [1, 16, 17, 29, 30, 32, 37, 43, 46, 50, 57, 67, 71, 74, 81]). Certainly, the generality of Finsler Geometry in comparison with the Riemannian setup (namely, analogous to the generality of the convex open subsets of an affine space in comparison with the ellipsoids) is a big drawback, as the number of new variables and parameters would seem immeasurable. Nevertheless, this is similar to the generality of General Relativity in comparison with Special one (see Remark 6.1). Anyway, any Finslerian modification of General Relativity would mean to drop the beloved Lorentz invariance not only at global and local levels (as it occurs in General Relativity) but also infinitesimally, i.e. looking such an invariance as a limit symmetry around each event. However, from a fundamental viewpoint, this should not seem too strange: as physical measurements are always approximations, one would not be surprised if the symmetries of the models were only approximations to a more complex reality. Indeed, as we will explain, the existence of some symmetries among observers becomes a natural requirement in order to make direct measurements of space and time. There is no reason to assume that the physical reality will satisfy such requirements in an exact way—even though, certainly, the existence of such approximated symmetries are meaningful and useful for modeling.

In the present article, a physical motivation to consider Finsler spacetimes as models of space and time is developed, and quite a few of related ideas are discussed. We stress the following four guidelines.

1. Approach from the foundations viewpoint. We develop an approach for the foundations of the theories of spacetime starting at the observers viewpoint in Classical Mechanics and Special Relativity (§2–4). Finsler spacetimes are shown to appear by dropping the symmetries of inertial observers in a natural way. Our approach follows the viewpoint in [9] by López and two of the authors in §2 and §3, which includes the celebrated ideas by V. Ignatowski [40, 41, 42] about the foundations of Special Relativity.

Specifically, we argue that the geometric models of spacetime appear from the notion of inertial observers by means of a double linearization of the measuring problem, namely:

- (1) there are inertial frames of reference (IFR) whose changes of coordinates are linear, and
- (2) the symmetries in the change of the timelike coordinate (and, independently, in the three spacelike ones) between two IFR's are encoded in that linear structure.

These assumptions lead to four classic linear 4-dimensional structures, namely Lorentz-Minkowski, Galilei-Newton, dual Galilei-Newton and Euclidean (see §2). Dropping (1) leads from Special to General Relativity (see §3), as well as to other transitions for the other three structures. The latter are also briefly explained here, namely, from Galilean to Leibnizian spacetimes, see §3.4, including signature changing metrics (see §3.1), and the possibility of a pointwise varying speed of light (see §3.3). Dropping (2) leads to Lorentz-Minkowski norms and, then, to Finsler spacetimes, discussed both mathematically and physically in §4.

2. Critical revision of EPS axiomatics. The classical Ehlers, Pirani and Schild (EPS) approach for General Relativity [21] will be revisited (see §5). We show that, certainly, this approach *is compatible* with the existence of Lorentz-Finsler metrics, a possibility already suggested by Tavakol and Van Den Bergh for Berwald spaces [81] (see the discussion in §5.2.5). Such a possibility was ignored in EPS because of a too restrictive development of two steps¹, namely:

- (1) An artificial requirement of smoothability of some combination of radar coordinates, which would forbid null cone structures non-compatible with Lorentzian metrics. This was recently pointed out by Lammërzhal and Perlick's [57] and it is developed here in detail (see §5.2.1).
- (2) A deduction of the existence of a projective structure starting at a general version of the law of inertia. This would exclude the timelike pregeodesics for a Lorentz-Finsler metric (except those of Berwald-type) but, again, the proof crucially relies on an argument of C^2 -differentiability, which is related to non-trivial issues on Finslerian metrics (see §5.2.2).

3. Precise geometric framework. Along the article, a careful mathematical approach is carried out, following [50]. For the convenience of the reader, a brief mathematical summary on (Lorentz) Finsler concepts is also included, §4.1. This allows us to model and to discuss issues on Lorentz-Finsler metrics L which turn out to be important from the physical viewpoint such as:

- (1) *Causal cone domain* (§4). The physically meaningful domain for L is only the causal cone of a cone structure \mathcal{C} .

¹The reason relies on a classical result for any Finsler metric F : its square F^2 is C^2 at the zero section if and only if F comes from a Riemannian metric (see Remark 4.4 (1) and §5.2).

Indeed, even in the classical relativistic case only the future-directed causal directions for a cone \mathcal{C}^+ determined by the metric g contains the elements physically measurable for any (true or *gedanken*) experiment. In Relativity, the Lorentzian scalar product g_p at each event p is determined by its value on the cone \mathcal{C}_p^+ (or on its timelike directions); therefore, a Lorentz metric g can be determined on the whole TM even if, actually, only its value on \mathcal{C}^+ can be measured. However, this is not by any means true for a Lorentz-Finsler metric L , where there is a huge freedom to extend the Lorentz-Finsler metric away from \mathcal{C} .

So, our Lorentz-Finsler metrics will be defined only on a (causal) cone structure².

- (2) *Smoothness*, i.e., differentiability up to some appropriate order. Usually, such a requirement is regarded as a harmless macroscopic approximation to the structure of the spacetime. However, the discussion on EPS above shows that this is not so trivial in the Finslerian case. What is more, other issues appear in the literature:
- The possibility that the cone is smooth and the Lorentz-Finsler metric is smooth only on the timelike directions but cannot be smoothly extended to the cone, which happens in metrics such as Bogoslovski in Very Special Relativity [12] and others [73], see §6.1.
 - The lack of differentiability outside the zero section of Finsler product spacetimes, which may lead to definitions of Finsler static spacetimes which are not smooth in the static direction [16], a fact which can be overcome with our approach to the space of observers, see §4.2 (item 5 (b)).
- (3) *Anisotropic speed of light*. Finsler spacetimes permit different possibilities for a speed of light which may vary not only with the point (an issue already considered even for relativistic spacetimes, §3.3) but also with the direction, §6.2.

4. *Importance of the space of observers*. The relevance of the *space of observers* in Special and General Relativity, its links with the symmetries of the spacetime and the possibility to lift Relativity to this space have been stressed by several authors [34, 36] in the framework of Lorentz violation and Lorentz-Finsler geometry. It is worth emphasizing that the essential role of this space appears explicitly along our development. In the initial discussion of the linearized models, we start with the set \mathbf{S} of *inertial frames of reference* (IFR), which permits even signature changing metrics (§3.1). However, once the symmetries of these models are dropped, only the space of observers \mathbf{O} remains as physically meaningful (Definition 3.1). In a classical relativistic spacetime (M, g) , \mathbf{O} is just the submanifold $\Sigma^g \subset TM$ of all the g -unit vectors in the future timelike cone; thus, each $\Sigma_p^g := \Sigma^g \cap T_pM$ is a hyperbolic space in the tangent space T_pM of each event $p \in M$. Breaking Lorentz symmetry at each p leads to regard Σ^g just as a more general

²This is consistent with our choices in our previous work [46]. There are other reasons for this choice from the purely mathematical viewpoint, as it clarifies the properties of anisotropically conformal metrics, see [50].

pointwise concave hypersurface Σ , which becomes then the indicatrix of a Lorentz-Finsler metric L (see Remark 4.13).

This observers' viewpoint allows one to use geometric methods recently developed in [50] which may have interesting physical applications such as: (a) going from g to L by perturbing the pointwise hyperboloids Σ^g into pointwise concave hypersurfaces Σ (as suggested in §6.3, such a perturbation might be produced by the presence of matter/energy and lead to quantum consequences), (b) to avoid or to smoothen possible singularities in Σ and then in L (showing that known non-smooth physical examples can always be approximated by smooth ones), (c) to construct systematically any Lorentz-Finsler metric from a Riemannian and a Finslerian one or (d) to construct systematically static and stationary metrics (avoiding any problem of smoothability).

In our opinion, the previous ideas support strongly that Finsler spacetimes have become an exciting vast field to explore thoughtfully from both the physical and mathematical viewpoints.

2. THE DOUBLY LINEARIZED MODELS

Next, we develop our approach for the foundations of the theories of spacetime. As a difference with the EPS approach, we will not assume postulates on *the nature of the behavior of the physical objects* which will be measured but on *how we can measure* those physical objects. A posteriori, if we are able to measure by using some sort of symmetry, the spacetime itself will be endowed with the geometric structure which codifies such symmetries.

The first step, to be developed along this section, considers the simplest symmetries for observers, common to both Classical Mechanics and Special Relativity. They will be regarded later as a (linear) idealization.

2.1. Postulates. Let us introduce the approach to the theories of spacetimes following³ [9] (a priori, this is non-quantum, even though quantum links will appear in §6.3).

The physical considerations on the existence of inertial frames of reference are encoded in the following two postulates.

Postulate 1 (Linear approach to spacetime). The physical spacetime is endowed with a structure of affine space Aff on a real vector space V of dimension $n = 4$. Physical observers are able to construct a non-empty set \mathbf{S}_{IFR} of affine frames of reference (each one $R = (O, B)$ composed by a point $O \in \text{Aff}$ and a basis B of V) which are called *inertial frames of reference* (IFR).

Thus, each IFR, R , provides an affine chart, i.e. a bijection $\varphi : \text{Aff} \rightarrow \mathbb{R}^4$, $\varphi(P) = (t(P), x^1(P), x^2(P), x^3(P))$, such that, given another IFR, \bar{R} , the coordinate change $\bar{\varphi} \circ \varphi^{-1} : \mathbb{R}^4 \rightarrow \mathbb{R}^4$ is an affine map. The first coordinate t of each IFR will be called *temporal* and the other three x^i , *spatial*.

³It is worth pointing out that [9] focuses on the viewpoint of General Relativity. So, the first postulate there is different to the one here. Our viewpoint was pointed in the reference [11] (written for a general audience in Spanish) and it is developed further here by introducing concepts such as apparent temporality (Theorem 2.4) or arguments as those on the varying of the speed of light.

The meaning of this first postulate is just that a linear approximation Aff to spacetime is being considered. The postulate also says that physicists will be able to construct *some* of the natural charts of the affine space Aff. The physical process to obtain such charts is not specified, even though the names *temporal* and *spatial* suggest the nature of their measurements.

Our second postulate, based essentially in von Ignatowski's [40], will ensure just that, when making measurements of the temporal coordinate (resp. when making measurements of the spatial coordinates), the viewpoint of two IFR's are interchangeable. This will be reflected by a requirement of symmetry in the corresponding charts. To understand this symmetry easily, let us discuss the bidimensional case $n = 2$. Let R, \bar{R} be two IFR's with coordinates (t, x) and (\bar{t}, \bar{x}) , resp. By Postulate 1:

$$\begin{pmatrix} \bar{t} \\ \bar{x} \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} t \\ x \end{pmatrix} + \begin{pmatrix} e \\ f \end{pmatrix}. \quad (1)$$

The interchangeability of the viewpoints of R and \bar{R} will collect the following physical assertion: the temporal coordinate \bar{t} (resp. the spatial coordinate \bar{x}) of \bar{R} measured by using the physical clock (resp. the rod) of R goes by as the temporal coordinate t (resp. the spatial coordinate x) of R measured by using the physical clock (resp. the rod) of \bar{R} . Mathematically,

$$\partial\bar{t}/\partial t (= a) = \partial t/\partial\bar{t} \quad \text{and} \quad \partial\bar{x}/\partial x (= d) = \partial x/\partial\bar{x}. \quad (2)$$

In dimension $n = 4$, interchangeability between the three spatial coordinates will also be imposed.

Postulate 2 (Time and spatial interchangeability). Let $R, \bar{R} \in \mathbf{S}_{\text{IFR}}$ be two IFR's. Then, their coordinates (t, x_1, x_2, x_3) and $(\bar{t}, \bar{x}_1, \bar{x}_2, \bar{x}_3)$ satisfy:

$$\partial\bar{t}/\partial t = \partial t/\partial\bar{t}, \quad \partial\bar{x}_i/\partial x_j = \partial x_j/\partial\bar{x}_i, \quad \forall i, j = 1, 2, 3. \quad (3)$$

2.2. Groups $O^{(k)}(4, \mathbb{R})$. The linear part of an affine change of coordinates from a first IFR, R , to a second one, \bar{R} , will be called the *transition matrix* A from R to \bar{R} . The second postulate implies that the transition matrices satisfy the condition (4) below, so, in order to obtain all the possibilities, one just needs to solve the following algebraic exercise.

Exercise 1. Let $A \in GL(4, \mathbb{R})$ be a regular 4×4 matrix and A^{-1} its inverse. Write them by using boxes as follows:

$$A = \left(\begin{array}{c|c} a_{00} & \mathbf{a}_h \\ \mathbf{a}_v^t & \hat{A} \end{array} \right), \quad A^{-1} = \left(\begin{array}{c|c} \tilde{a}_{00} & \tilde{\mathbf{a}}_h \\ \tilde{\mathbf{a}}_v^t & \tilde{\hat{A}} \end{array} \right),$$

where $a_{00}, \tilde{a}_{00} \in \mathbb{R}$, $\mathbf{a}_h, \mathbf{a}_v, \tilde{\mathbf{a}}_h, \tilde{\mathbf{a}}_v \in \mathbb{R}^3$, $\hat{A}, \tilde{\hat{A}}$ are 3×3 submatrices, and the superscript t denotes transpose. Then, determine those matrices A satisfying:

$$\tilde{a}_{00} = a_{00} \quad \tilde{\hat{A}} = \hat{A}^t. \quad (4)$$

Such an exercise is solved in [9, §3] in full detail. Next, we will describe the main properties of its solutions⁴.

⁴The reader can consider the simple case $n = 2$ (as in (2)), when $\hat{A} \equiv d$, $\tilde{\hat{A}} \equiv \tilde{d}$ ($d, \tilde{d} \in \mathbb{R}$). The solutions of this case yield all the relevant possibilities. They follow easily

Definition 2.1. Let $S^1 = \mathbb{R} \cup \{\omega\}$ be the circle regarded as the extended real line $\mathbb{R}^* = [-\infty, +\infty]$ with $+\infty$ identified to $-\infty$ as a single point ω . For each $k \in S^1$, consider the matrix

$$I^{(k)} = \left(\begin{array}{c|c} k & 0 \\ \hline 0 & I_3 \end{array} \right) \quad (\text{where } I_3 \text{ is the } 3 \times 3 \text{ identity matrix})$$

and define the group $O^{(k)}(4, \mathbb{R}) \subset GL(4, \mathbb{R})$ as follows:

- if $k \in \mathbb{R}$, $O^{(k)}(4, \mathbb{R}) = \{A \in GL(4, \mathbb{R}) : \det A^2 = 1, A^t I^{(k)} A = I^{(k)}\}$,
- if $k = \omega$, $O^{(\omega)}(4, \mathbb{R}) = \{A \in GL(4, \mathbb{R}) : A^t \in O^{(0)}(4, \mathbb{R})\}$.

We will say that $A \in GL(4, \mathbb{R})$ is *k-congruent* if $A \in O^{(k)}(4, \mathbb{R})$. Accordingly, two IFR's R, R' are *k-congruent* is so is its transition matrix. It is easy to check that any *k-congruent* matrix A is a solution of Exercise 1 as in this case $A^t I^{(k)} = I^{(k)} A^{-1}$. Remarkably, it will turn out that the converse holds except in very exceptional cases (detailed in [9, Prop. 3.1]). Indeed, these exceptional cases will be avoided by using very mild and natural conditions from both the mathematical and physical viewpoints (any of the hypotheses (1)–(4) in the main Theorem 2.4 below).

Remark 2.2. (1) In the case $k \neq 0, \omega$, the equality

$$A^t I^{(k)} A = I^{(k)}$$

implies $\det A^2 = 1$ trivially. What is more, this equality is equivalent to

$$A^{-1} I^{(1/k)} (A^{-1})^t = I^{(1/k)}.$$

Then, the case $k = \omega$ becomes equivalent to taking the limit $k \rightarrow \omega (\equiv \pm\infty)$:

$$O^{(\omega)}(4, \mathbb{R}) = \{A \in GL(4, \mathbb{R}) : \det A^2 = 1, A^{-1} I^{(0)} (A^{-1})^t = I^{(0)}\}.$$

(2) If A is *k-congruent* for two distinct values of k , then so it is for any k . Concretely, let $k_1, k_2 \in S^1$, from [9, Lemma 3.3] (see its part 1 and proof):

$$k_1 \neq k_2 \implies O^{(k_1)}(4, \mathbb{R}) \cap O^{(k_2)}(4, \mathbb{R}) = \bigcap_{k \in S^1} O^{(k)}(4, \mathbb{R}) = \{\pm 1\} \times O(3, \mathbb{R}),$$

where $O(3, \mathbb{R})$ is the usual orthogonal group and

$$\{\pm 1\} \times O(3, \mathbb{R}) := \left(\begin{array}{c|c} \pm 1 & 0 \\ \hline 0 & O(3, \mathbb{R}) \end{array} \right).$$

Now, the relevant solutions to our exercise can be easily described.

Lemma 2.3. *Let $A \in GL(4, \mathbb{R})$ satisfy the hypothesis (4) of Exercise 1.*

- (1) *If the matrix A^2 also satisfies the property (4) then $\det A^2 = 1$.*
- (2) *If $\det A^2 = 1$, then there exists $k \in S^1$ such that A is *k-congruent*, and either k is unique or it can be arbitrarily chosen in S^1 .*
- (3) *Let $A_1, A_2 \in GL(4, \mathbb{R})$ be k_1 - and k_2 -congruent, resp. If k_1 is univocally determined and $A_1 \cdot A_2$ (resp. $A_2 \cdot A_1$) is *k-congruent* for some $k \in S^1$, then $A_1 \cdot A_2$ (resp. $A_2 \cdot A_1$) is k_1 -congruent.*

by noticing that, from the algorithm to compute the inverse matrix:

$$a = \tilde{a} = d / \det A, \quad d = \tilde{d} = a / \det A, \quad \tilde{b} = -b / \det A \quad \text{and} \quad \tilde{c} = -c / \det A.$$

In particular, $d \neq 0 \Leftrightarrow a \neq 0$ and, then, $\det A^2 = 1$. Therefore, this equality would follow by assuming additionally $a > 0$ (i.e., $\partial t / \partial t > 0$ in (3)), which will correspond with the condition of apparent temporality in Theorem 2.4.

Proof. Assertion (1) follows from the sentence above [9, Lemma 3.3] (recall that, as explained at the beginning of the paragraph containing that sentence, *incongruent* means $\det A^2 \neq 1$). For (2), the existence of k follows also from the paragraph above [9, Lemma 3.3] and the uniqueness from part 1 of [9, Lemma 3.3] regarding S_p as a set of two congruent observers and A as the transition matrix between them or from Remark 2.2(2). Assertion (3) follows from part 1 of [9, Lemma 3.3] regarding S_p as a set of three congruent observers with transition matrices A_1 , A_2 and, say, $A_1 \cdot A_2$ (and its inverses). Then, all of them must be k' -congruent for some k' and, as k_1 was univocally determined, $k' = k_1$. \square

Lemma 2.3 implies that, under minimal realistic hypotheses, any set \mathbf{S}_{IFR} of IFR determines (at least) one value of $k \in S^1$. Mathematically, such realistic properties just ensure that $\det A = \pm 1$, which would be related to the conservation of the volume. Such a property might also be postulated directly, nevertheless, there are other physically sound weak hypotheses that imply it.

In order to formulate such hypotheses, recall first that the set \mathbf{S}_{IFR} of IFR obtained from our postulates is rather arbitrary. For example, the unique restriction to its number of elements comes from $\mathbf{S}_{\text{IFR}} \neq \emptyset$; that is, one can remove arbitrarily some elements of \mathbf{S}_{IFR} (but not of all them) and this new set would satisfy the postulates 1 and 2 too. What is more, if there is some $k \in S^1$ such that \mathbf{S}_{IFR} is composed by (a small number of) k -congruent IFR's, one can enlarge \mathbf{S}_{IFR} by acting with the group $O^{(k)}(4, \mathbb{R})$ obtaining a bigger set S of compatible IFR's. Notice that if there were a second $k' \neq k$ such that all IFR's in \mathbf{S}_{IFR} were k' -congruent, a different enlargement S' could also be obtained. These observations suggest the following construction. Given \mathbf{S}_{IFR} , define

$$\mathbf{S}_{\text{IFR}}^* := \bigcap_{\alpha} S_{\alpha}, \quad (5)$$

where each S_{α} is a set of affine reference frames satisfying: (i) S_{α} includes \mathbf{S}_{IFR} , (ii) the change of coordinates between any two elements of S_{α} satisfies the formula (3) in Postulate 2, and (iii) S_{α} is *maximal* (i.e., not included in a bigger set satisfying the previous conditions (i) and (ii)).

Recall: (a) $\mathbf{S}_{\text{IFR}}^* (\supset \mathbf{S}_{\text{IFR}})$ is determined univocally by \mathbf{S}_{IFR} , (b) physically, all the affine reference frames in $\mathbf{S}_{\text{IFR}}^*$ could be regarded as IFR's with the same status as those in \mathbf{S}_{IFR} , and (c) mathematically, one would expect that the transition matrices between all the pairs of elements of $\mathbf{S}_{\text{IFR}}^*$ had a more natural structure than \mathbf{S}_{IFR} .

Theorem 2.4. *Let \mathbf{S}_{IFR} be a set of IFR's (satisfying the Postulates 1 and 2). There exists $k \in S^1$ such that the transition matrix $A \in GL(4, \mathbb{R})$ of each transformation of coordinates between two IFR's, R_1 and R_2 , is k -congruent for all $R_1, R_2 \in \mathbf{S}_{\text{IFR}}$, whenever any of the following hypotheses hold:*

- (1) Conservation of the IFR volume: $\det A = \pm 1$, for any transition matrix A .
- (2) Transitivity: if A is the transition matrix from a first IFR, $R_1 \in \mathbf{S}_{\text{IFR}}$, to a second IFR, $R_2 \in \mathbf{S}_{\text{IFR}}$, then there exists an IFR, R_0 , such that the transition matrix A from R_0 to R_1 is equal to A .

- (3) Action by a group: *the set of transition matrices A between elements of $\mathbf{S}_{\text{IFR}}^*$ (as in (5)) is a subgroup G of $GL(4, \mathbb{R})$.*
- (4) Apparent temporality: *any transition matrix A between elements of \mathbf{S}_{IFR} satisfies $a_{00} > 0$ (with a_{00} as in Exercise 1; recall also the discussion at §2.4).*

Moreover, the existence of such a k implies that the properties (1), (2) and (3) hold, being the group G in (3) either $O^{(k)}(4, \mathbb{R})$ or the intersection of all of them, i.e., $\{\pm 1\} \times O(3, \mathbb{R})$.

Proof. First, let us prove the existence of the required k under the hypothesis (1) and, then, let us check (1) \Leftrightarrow (2) \Leftrightarrow (3) and (1) \Leftrightarrow (4). Under (1), the existence of some k for each A is ensured by part (2) of Lemma 2.3. Then [9, Lemma 3.3(1)] (or part (3) of Lemma 2.3) ensures that one can choose the same k for all the transition matrices A determined by pairs of elements in \mathbf{S}_{IFR} . If the hypothesis (2) holds then A^2 is also a transition matrix between IFR, and part (1) of Lemma 2.3 implies that the hypothesis (1) holds too. Analogously, (3) implies (2) trivially. Finally, (4) implies (1) from [9, Lemma 3.1, item 1(ii)].

For the last assertion, let us check that, when such a k exists, then (3) holds. Indeed, one of the sets S_α in the definition of $\mathbf{S}_{\text{IFR}}^*$, name it S_k , can be chosen such that the group $O^{(k)}(4, \mathbb{R})$ acts transitively on S_k (just choose $R \in \mathbf{S}_{\text{IFR}}$ and take all the affine reference frames R' with transition matrix A in $O^{(k)}(4, \mathbb{R})$). So, when k is univocally determined for one pair of elements $R_1, R_2 \in \mathbf{S}_{\text{IFR}}$, then $\mathbf{S}_{\text{IFR}}^* = S_k$ and the hypothesis (3) holds with the group $G = O^{(k)}(4, \mathbb{R})$. Otherwise, k can be arbitrarily chosen by Lemma 2.3(2), then $\mathbf{S}_{\text{IFR}}^* = \bigcap_{k \in S^1} S_k$ and (3) holds with the group $G = \{\pm 1\} \times O(3, \mathbb{R})$ (see Remark 2.2(2)). \square

2.3. Linear models of spacetimes. Theorem 2.4 implies that, whenever one of its mild hypotheses (1)–(4) holds, the existence of a set \mathbf{S}_{IFR} of IFR's according to Postulates 1 and 2, selects a group $G = O^{(k)}(4, \mathbb{R})$ (or the intersection of all of them). As the spacetime was represented by an affine space Aff on a vector space V by postulate 1, this vector space (and then Aff) will be endowed automatically with the geometric structure invariant by G . Let us study each case.

- (1) Case $k \in (-\infty, 0)$. By the definition of $O^{(k)}(4, \mathbb{R})$, V is naturally endowed with a Lorentzian scalar product $\langle \cdot, \cdot \rangle_1$. Indeed, if $R = (O, B = (e_0, e_1, e_2, e_3))$ is any IFR, then the unique $\langle \cdot, \cdot \rangle_1$ such that B is an orthonormal basis for it, up to the normalization of its first vector

$$\sqrt{|\langle e_0, e_0 \rangle_1|} = \sqrt{-k}, \quad (6)$$

becomes independent of the chosen R . What is more, for $k = -1$, the group $O^{(k)}(4, \mathbb{R})$ is the Lorentz group; otherwise, $O^{(k)}(4, \mathbb{R})$ is conjugate to the Lorentz group. Indeed, putting $k = -c^2$ with $c > 0$, then $I^{(k)} = I^{(c)} \cdot I^{(-1)} \cdot I^{(c)}$, the inverse of $I^{(c)}$ is $I^{(1/c)}$ and

$$O^{(k)}(4, \mathbb{R}) = I^{(1/c)} \cdot O^{(1)}(4, \mathbb{R}) \cdot I^{(c)}. \quad (7)$$

Anyway, the spacetime of Special Relativity is obtained.

- (2) Case $k = \omega$. The group $O^{(k)}(4, \mathbb{R})$ becomes the (non-orthochronous) *Galilean group*

$$O^{(\omega)}(4, \mathbb{R}) := \left\{ \left(\begin{array}{c|c} \pm 1 & 0 \\ \mathbf{a}_v^t & \hat{A} \end{array} \right) : \mathbf{a}_v \in \mathbb{R}^3, \hat{A} \in O(3, \mathbb{R}) \right\}.$$

Thus, the dual basis $B^* = (\phi^0, \phi^1, \phi^2, \phi^3)$ of each IFR contains the same first element ϕ^0 , up to a sign. When a choice in $\{\phi^0, -\phi^0\}$ is carried out, name it $\mathbf{t} : V \rightarrow \mathbb{R}$, then \mathbf{t} is called the *absolute time*. The kernel E of $\pm\phi^0$ is endowed with a scalar product $\langle \cdot, \cdot \rangle_E$ (being the elements (e_1, e_2, e_3) of B an orthonormal basis of $\langle \cdot, \cdot \rangle_E$ for each IFR). Then, E endowed with this scalar product is called the *absolute space*.

Summing up, the spacetime of Galilei-Newton is recovered now.

- (3) Case $k = 0$. The group $O^{(k)}(4, \mathbb{R})$ becomes the *dual Galilean group*⁵

$$O^{(0)}(4, \mathbb{R}) := \left\{ \left(\begin{array}{c|c} \pm 1 & \mathbf{a}_h \\ 0 & \hat{A} \end{array} \right) : \mathbf{a}_h \in \mathbb{R}^3, \hat{A} \in O(3, \mathbb{R}) \right\}.$$

In this case, the basis $B = (e_0, e_1, e_2, e_3)$ of each IFR contains the same first element e_0 , up to a sign. Choosing a sign, this vector defines the *absolute rest observer*. Thus, the kernel (annihilator) of $\pm e_0$ in the dual space V^* (that is, the subspace $E^* := \text{Span}\{\phi^1, \phi^2, \phi^3\}$ of B^* for each IFR) is also independent of the IFR. It is naturally endowed with a scalar product $\langle \cdot, \cdot \rangle_{E^*}$ so that, for each IFR, the set (ϕ^1, ϕ^2, ϕ^3) becomes an orthonormal basis.

Summing up, an a priori aphysical dual of Galilei-Newton spacetime (with a completely analogous geometric structure) is obtained.

- (4) Case $k \in (0, \infty)$. For $k = 1$, the group $O^{(k)}(4, \mathbb{R})$ is the Euclidean orthonormal group⁶; otherwise, $O^{(k)}(4, \mathbb{R})$ is conjugate to this group. Indeed, reasoning as in the case $k < 0$, V is naturally endowed with an Euclidean scalar product $\langle \cdot, \cdot \rangle_0$ and any basis B of an IFR is orthonormal for $\langle \cdot, \cdot \rangle_0$, up to the normalization of its first vector.

Summing up, one obtains the a priori aphysical case when the full spacetime is endowed with a Euclidean scalar product, which is mathematically analogous to the Lorentzian one.

- (5) Case $k \in S^1$ non-unique. In this case, the group is $G = \{\pm 1\} \times O(3, \mathbb{R})$ and, thus, the basis B and its dual B^* for any IFR satisfy all the properties in the previous cases. In particular, choosing a sign, one has an absolute time T , an absolute rest observer e_0 (with $T(e_0) = 1$) and an absolute space $(E, \langle \cdot, \cdot \rangle_E)$ whose dual space can be identified with $(E^*, \langle \cdot, \cdot \rangle_{E^*})$ defined in the case $k = 0$.

⁵This group was studied by Lévi-Leblond [59], who named it *Carrollian group*, after Lewis Carroll. Even though introduced as an academical exercise, recent applications of this group can be found in [24, 20].

⁶It is worth pointing out that, in this case, not only the independent symmetries between time and spatial coordinates in formula (3) hold, but also the crossed symmetries $\partial t / \partial \tilde{x}^i = \partial \tilde{x}^i / \partial t$ and $\partial x^i / \partial \tilde{t} = \partial \tilde{t} / \partial x^i$ appear now.

This case should be regarded as aphysical too⁷ and, being obtained as a “degenerate” case of the previous ones, it will not be taken into account anymore.

2.4. Temporal models and interpretation of $k = -c^2$. Taking into account the previous four models of spacetime which depend on a unique $k \in S^1$, let us revisit the role of the hypothesis of *apparent temporality* in Theorem 2.4.

Recall that apparent temporality was enough to ensure the existence of k in that theorem. However, the Euclidean case $k > 0$ would not be excluded by this hypothesis, because the set \mathbf{S}_{IFR} of all the IFR’s might contain “few” elements (so that only transition matrices A with $a_{00} = \cos \theta$ appeared for values of θ with $\cos \theta > 0$). Moreover, in the other three cases for k , the elements of \mathbf{S}_{IFR} would determine a *time-orientation*⁸ under apparent temporality, but there would still be elements in $\mathbf{S}_{\text{IFR}}^*$ which would not match with the chosen time-orientation. However, when the case $k > 0$ is disregarded a priori (say, regarding it as aphysical), it would be natural to strengthen the hypothesis of apparent temporality into *temporality*, namely: all the transition matrices between pairs of elements of $\mathbf{S}_{\text{IFR}}^*$ in (5) have $a_{00} > 0$. This requirement not only would exclude the group $O^{(k)}(4, \mathbb{R})$ for $k > 0$ but it would also imply a restriction on the group for the other cases. This discussion makes natural the following definition and convention.

Definition 2.5. The linear models of spacetime with $k \in (-\infty, 0) \cup \{\omega, 0\}$ will be called *temporal models*. When only these models are considered, we will assume that *apparent temporality* also holds and, then, the following *convention of temporality* can be assumed with no loss of generality:

- (a) The temporal models are time-oriented.
- (b) All the elements in \mathbf{S}_{IFR} are assumed to lie in the chosen time-orientation.
- (c) \mathbf{S}_{IFR} is assumed to be maximal for the property (b). Thus, depending on the value of k , the *orthochronous* subgroup of the Lorentz (or conjugate to Lorentz), Galilean or dual Galilean group will act freely and transitively on \mathbf{S}_{IFR} .
- (d) When there is no possibility of confusion, $\mathbf{S}_{\text{IFR}}^*$ is regarded as equal to \mathbf{S}_{IFR} in (c).

For temporal models, given a transition matrix A which gives the coordinates (\bar{t}, \bar{x}^j) for \bar{R} from the coordinates (t, x^j) of R , the *velocity* and *speed* of R measured by \bar{R} are, resp.,

$$\mathbf{v} = \mathbf{a}_{\mathbf{v}}/a_{00} \quad |\mathbf{v}| = \sqrt{\sum_{i=1}^3 (v^i)^2} \quad (8)$$

⁷Anyway, it would represent the model of space and time which goes back to Aristoteles. Recall that in that model, one would assume not only the existence of the absolute space and time but also that, for any $P \in \text{Aff}$, there exists a physical observer at P at absolute rest. This would determine the affine line $P + \mathbb{R} \cdot e_0$, which would be regarded as a “space point at any time”.

⁸That is, a choice of one of the two timelike cones when $k < 0$ and one of the two choices of absolute time or absolute rest observer when $k = \omega, 0$, resp.

in the notation of Exercise 1 (see also [9, §5 (2)]).

Proposition 2.6. *For any temporal model, $c := \sqrt{|k|} \in [0, \infty]$ is the supremum of speeds measured between IFR's in \mathbf{S}_{IFR}^* .*

Proof. For $k = 0, \omega$ this follows from (8) taking into account the expression of $O^{(k)}(4, \mathbb{R})$ at each case (see items (2) and (3) at §2.3). For $k \in (-\infty, 0)$, using (7) the first column of A is $(a_{00}, \mathbf{a}_v) = (a_{00}, cb^1, cb^2, cb^3)$ with $\sum_i (b^i)^2 = a_{00}^2 - 1$ (thus, $|\mathbf{v}|^2 = c^2 - c^2/a_{00}^2$) and $a_{00} \geq 1$ unbounded. \square

As one would expect, this supremum is ∞ (i.e. the speeds are unbounded) in the Galilei-Newton case, finite equal to c in the case of Special Relativity and strictly 0 in the dual Galilean case (where all IFR's lie at absolute rest).

Remark 2.7. In principle, it is appealing to call c the *speed of light*. Notice, however, that there is no mention neither to Electromagnetism nor to any other interaction in our approach. Nevertheless, an essential property of electromagnetism can justify that name. Namely, light is described by a wave *which propagates in vacuum*. An obvious natural hypothesis for IFR's is that the vacuum is “equal” for all of them, and, so, any physical scalar quantity measured with respect to the vacuum must yield the same number for all of them. In particular, this would mean that all IFR's must measure the same speed of propagation of the light with respect to the vacuum. As the supremum c is the unique speed equal for all of them, the following definition is justified.

Definition 2.8. For any temporal model, $c = \sqrt{|k|}$ is called the *speed of light*.

Anyway, the following digression about the physical content of this definition may be worthy. If one considered another interaction which also propagated in vacuum (say, gravitation) then the arguments in Remark 2.7 would imply that its speed of propagation c' with respect to vacuum would be the same c as for light. As emphasized by some authors, see [31], there is no logical contradiction assuming that $c \neq c'$ and, thus, this question becomes an experimental issue⁹. In the affirmative, these different interactions might allow one to construct different types of clocks and rods in order to measure the temporal and spatial coordinates. So, the name IFR should include the interactions which allow Postulates 1 and 2 to hold.

3. FIRST NON-LINEARIZATION

General Relativity can be regarded as a first non-linear generalization of Special Relativity. Such nonlinearity comes from the fact that Postulate 1, namely, the global affine character of spacetime, is being dropped and the set of all the events is modeled by a manifold. Nevertheless (as apparent from [9]), Postulate 2 would make still sense if the symmetries stated there are regarded just as infinitesimal ones, at the tangent space of each event.

This idea is well-established in the Lorentz case and it may seem very speculative in the other linear models of spacetimes. However, this will be

⁹However, recent measurements of gravitational waves show that the speed of propagation of light and gravitation are equal with an extraordinary accuracy [60].

developed briefly along this section with a double aim: on the one hand, the role of observers will be emphasized and, on the other, the framework of further issues relevant to the Lorentz-Finsler case will be settled. Only in §4 we will focus on the Lorentz case and will go beyond, in order to reach the Lorentz-Finsler generalization.

3.1. General case and signature change. Assume now that the space-time is described by a (smooth, connected) manifold M and that our postulates are regarded as infinitesimal requirements of symmetry at the tangent space T_pM of each $p \in M$, that is, around each event $p \in M$, one can find a set of coordinate charts such that the relations (3) occur only at p , namely, considering normal coordinates.

Then, we will have a set \mathbf{S}_p of linear bases at each T_pM which will play the role of (linear) IFR's at p . For simplicity, we will assume in what follows:

(i) \mathbf{S}_p determines univocally some $k(p) \in S^1$ (i.e., the degenerate case of non k -congruent solutions of Exercise 1 is skipped),

(ii) \mathbf{S}_p is maximal (i.e., $\mathbf{S}_p = \mathbf{S}_p^*$, consistently with the discussion above Theorem 2.4), and

(iii) Consistently with Definition 2.5, the convention of temporality will be assumed whenever $k(p) \notin (0, \infty)$ (in particular, the notion of future-directed timelike vectors makes sense then).

Moreover, as an extra hypothesis (or third postulate, as in [9]) we assume:

(P3) \mathbf{S}_p varies smoothly in the bundle LM of linear frames¹⁰ of M .

Formally, this means that $\mathbf{S} := \cup_{p \in M} \mathbf{S}_p$ is a smooth bundle embedded in LM (in the sense of a submanifold of LM with the induced topology such that the projection on M is a submersion) so that the function $k : M \rightarrow S^1$ becomes smooth.

In general, one obtains then a signature changing metric g which is Lorentzian (resp. Riemannian) in the set $-\infty < k < 0$ (resp. $0 < k < \infty$). Following the terminology in [9, 10], in the closed subset determined by $k = \omega$, one has a *Leibnizian structure*, that is, a non-vanishing 1-form Ω (*absolute time form*) on M and a Riemannian metric h in the subbundle $\ker(\Omega)$ of TM , being then $(\ker(\Omega), h)$ the *absolute space*¹¹. Analogously, the region $k = 0$ is endowed with an anti-Leibnizian structure, consisting in a non-vanishing vector field W (*absolute rest field*) on M and a Riemannian metric h^* on the subbundle $\ker(W)$ of the cotangent bundle TM^* .

Let g be the semi-Riemannian (Lorentzian or Riemannian) metric in the region $k \neq 0, \omega$ and g^* the (physically equivalent) metric induced in the cotangent space. It is worth emphasizing that, in the region $k = 0$, g can be extended as a degenerate metric and g^* cannot; however, g^* matches smoothly with h^* on $\ker(W)$. Analogously, in the region $k = \omega$, g^* can be extended as a degenerate metric, while g matches smoothly with h on $\ker(\Omega)$.

¹⁰ LM contains all the (ordered) linear bases of T_pM for all $p \in M$.

¹¹Such a structure is equivalent to having the 1-form Ω and a positive semidefinite 2-contravariant tensor T of rank 3 with $i_\Omega T := T(\Omega, \cdot) \equiv 0$, studied in [55]. Indeed, such a T induces a Riemannian metric in the dual of $\ker(\Omega)$ and, then, in $\ker(\Omega)$. Conversely, the Leibnizian structure yields a Riemannian metric on $\ker(\Omega)$ and then in its dual; this yields the tensor T by imposing that its radical is $\text{Span}\{\Omega\}$.

Summing up, this first non-linear generalization of the IFR setting yields as a general model of spacetime a geometry governed by the smooth function k . Whenever $k \neq 0, \omega$, a semi-Riemannian metric g and its equivalent dual metric g^* are obtained; in the regions $k = 0$ or $k = \omega$ either the metric g or g^* are extended as a degenerate metric and additional geometric structures appear¹². The transition among these elements is smooth, as so is \mathbf{S} .

3.2. Space of observers. For convenience, let us introduce a new element by taking the most important information from \mathbf{S} .

Definition 3.1. The *space of observers* is the subset \mathbf{O} of TM containing the first vector of each basis at \mathbf{S} , and the *space of observers at p* is $\mathbf{O}_p := \mathbf{O} \cap T_pM$. That is:

- (i) In the region $k > 0$, \mathbf{O} contains all the unit vectors for g , and each \mathbf{O}_p is a sphere.
- (ii) In the region $k < 0$, \mathbf{O} contains all the future-directed timelike unit vectors for g , and each \mathbf{O}_p is a hyperboloid.
- (iii) In the region $k = \infty$, \mathbf{O} is equal to $\Omega^{-1}(1)$, and each \mathbf{O}_p is an affine hyperplane not containing 0.
- (iv) In the region $k = 0$, \mathbf{O} is equal to the absolute rest vector field W (so, each \mathbf{O}_p is a subset containing a single non-zero tangent vector).

Remark 3.2. If $n = \dim M$, then $\dim LM = n(n + 1)$ and $\mathbf{S} \subset LM$ is always a submanifold with $\dim \mathbf{S} = (n + 1)n/2$. Nevertheless, \mathbf{O} must be regarded as a subset of TM . Then, it becomes a smooth manifold of dimension $2n - 1$ in the region $k \neq 0$ but it collapses to a submanifold of dimension n when $k = 0$.

The transition from Lorentzian to Riemannian through a region with $k = \omega$ can be easily understood by looking at \mathbf{O} (see Fig. 1). We will not be interested in the transition through a region with $k = 0$. However, this could be described in a completely analogous way by defining a dual space of observers (constructed by picking at each point p the first element of the elements in the bases which are dual to those in \mathbf{S}_p).

3.3. Pointwise variation of speed of light. In the region $-\infty < k < 0$ the function $c(p) = \sqrt{|k(p)|}$ might be understood as a variation of the speed of light with the point at M . Such a possibility has been speculated since the beginning of General Relativity, and was put forward in the 90's in relation to cosmological inflation and the horizon problem (see for example [3, 6, 69, 72], as well as some criticism in [22, 84]). In order to avoid the circularity of using the light to define the units to measure its speed, Barrow and Magueijo [7] argue that only the variation of adimensional constants would have a true physical meaning; so, the variation of c should be regarded as a variation of the (adimensional) fine structure constant $\alpha = e^2/\hbar c 4\pi\epsilon_0$.

Without deepening into these questions, some comments about varying c in our framework are in order. Recall first that, in the affine case obtained

¹²Recall that models of signature changing metrics have been studied at least since the influential “no boundary” proposal by Hartle & Hawking [35], see for example [19, 87]. Moreover, the existence of an “absolute time” in the transition region has also been pointed out by several authors [52, §2] (see also [85]).




FIGURE 1. Signature changing spacetime on $M = \mathbb{R}^2$. The natural coordinate basis $B = (e_0, e_1) \equiv (\partial_t, \partial_x)$ is assumed to lie on \mathbf{S} at each point. The matrices of the metric g and g^* are $\begin{pmatrix} k & 0 \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} 1/k & 0 \\ 0 & 1 \end{pmatrix}$, respectively, with $k(t, x) = 1/x \in S^1 \setminus \{0\}$. The space of observers changes from a hyperbola to a line and to a circumference.

by assuming Postulate 1, to assume also Postulate 2 would imply that all the IFR's would be using the same units for measurement and, then, the specific value of c would depend of the chosen units. Indeed, the natural interpretation of the group $O^{(k)}(4, \mathbb{R})$ for $k = -c^2 \in (-\infty, 0)$ is just the Lorentz group in some appropriate coordinates. Thus, if one regarded the affine space Aff as a manifold and took different units at each point, then this could not be interpreted as a variable speed of light.

To measure a varying speed of light would rely on the possibility to compare the units of measurement at different points. In the affine case such a comparison would be possible if the interactions were invariant by translations (an unlikely possibility). In general, one would need measurements involving magnitudes which are dimensionally independent (in particular, this would be achieved by measuring adimensional constants, as commented above). In principle, this might be achieved by measuring essentially different interactions, as in the case of light and gravity propagation (see below Definition 2.8)¹³. Anyway, as we will see, the Lorentz-Finsler viewpoint will open other possibilities by using infinitesimal anisotropy.

3.4. Relativistic vs Leibnizian structures. A priori, the cases $-\infty < k < 0$ and $k = \omega$ (with constant k , and then c) are the physically interesting ones, either as a model of spacetime or as an approximation to this model. Next, they will be briefly compared.

¹³In the more speculative case of a signature changing metric, the speed of light would change necessarily in the regions $k = 0, \omega$. So, the possibility to measure a varying speed of light when $-\infty < k < 0$ would imply that the collapse of the lightcones (to a line or a hyperplane) could be measured gradually when approaching to those regions.

In the first case, we will assume $k = -1$ on all M , after our discussion in the previous subsection. So, one has a time-oriented Lorentzian metric g and \mathbf{O} is a fibered space on M with fiber the hyperbolic space. Moreover, the Levi-Civita connection ∇^g is canonically associated with g , and any other affine connection ∇ parallelizing g (i.e., satisfying $\nabla_\mu g_{\nu\rho} = 0$) must be non-symmetric. This selects ∇^g and provides a sense of free fall and lightlike geodesics (compare with EPS later).

In the case $k = \omega$, the Leibnizian structure consisting in the absolute time form Ω and the absolute space $(\ker(\Omega), h)$ on M described in subsection 3.1 is obtained. These structures were studied systematically in [55] and [10]. In this setting, one considers *Galilean connections*, that is, affine connections ∇ which parallelize both, Ω and h . It is worth pointing out that the set of all the Galilean connections has the same degrees of freedom as the set of all affine connections (symmetric or not) parallelizing a Lorentzian metric g . However, a symmetric Galilean connection will exist if and only if Ω is closed ($d\Omega = 0$), that is, locally $\Omega = d\mathbf{t}$ for some function \mathbf{t} . Nevertheless, in this case, there is no univocally determined symmetric connection. Moreover, this happens even if Ω is exact, that is, $\Omega = d\mathbf{t}$ for some function \mathbf{t} defined globally on M , which will be called the *absolute time* (\mathbf{t} is unique up to an additive constant). Indeed, an explicit Koszul-type formula reconstructs all the symmetric Galilean connections in terms of two data¹⁴ [10, Cor. 28], namely: the *gravitational field* (a vector field in the absolute space, that is, a section of the bundle $\ker(\Omega) \rightarrow M$) and the *vorticity* (a skew symmetric 2-form on the vector bundle $\ker(\Omega)$).

In conclusion, relativistic spacetimes are preferred to Leibnizian ones from the viewpoint of foundations, because of two celebrated properties: (a) they permit to model a finite speed of propagation in vacuum (recall that observers appear now at each event as infinitesimal approximations to IFR's and, so, the arguments in Remark 2.7 apply), and (b) they select a unique affine connection in the set of all the connections parallelizing the geometric structure, while Leibnizian ones require the gravitational and vorticity fields as an extra input.

In the next section, we will focus only on relativistic spacetimes and the Finslerian extensions. Nevertheless, some previous elements serve as a background for the Lorentz and Lorentz-Finsler cases and they can be compared a posteriori with them (see Table 1). We point out a pair of them so that the interested reader might come back here later:

(1) Leibnizian structure (Ω, h) vs cone triple (Ω, T, F) (which is useful to define and to handle any cone structure \mathcal{C} , Lorentz or Lorentz-Finsler, see Definition 4.6, Remark 4.7(2)). Notice that when F comes from a Riemannian metric h , then the Leibnizian structure can be regarded as a sort of limit when $\lambda \rightarrow \infty$ of the triples $(\lambda\Omega, T/\lambda, F)$, which “open” the cone \mathcal{C} .

(2) Chronometric vs EPS approach to spacetime (§5). The Leibnizian structure (Ω, h) (eventually, with $\Omega = d\mathbf{t}$) gives a chronometric approach to spacetime, in a similar way as the Lorentz metric does in Relativity. However, the former requires an additional input (an affine connection) in

¹⁴Such a formula can be extended to include non-symmetric connections by adding as a third datum a suitable component of the torsion, see [10, Th. 27].

order to define free fall. So, the EPS approach (at least the axioms which do not consider light propagation) might also be interesting in the Leibnizian case. In contrast, the Lorentz-Finsler metric L will provide timelike and lightlike geodesics in a very similar way as the Lorentz metric g , in spite of the differences between the Levi-Civita ∇^g and the anisotropic connection (see footnote 21) determined by L .

Note 3.3. *Newton-Leibniz controversy.* To end this section it is worth pointing out that the notion of Leibnizian structure provides a precise mathematical description of a historical controversy between Leibniz and Newton. Roughly speaking, Leibniz criticized Newton's arguments about IFR's by pointing out that the Euclidean space perceived by an observer is equal even after a rotation of the observer's coordinates. So, he claimed that one could not detect whether these axes are being rotated at different times. Newton replied that spinning water in a bucket would detect whether the observer is rotating or not. From the mathematical viewpoint, Newton was using the structure of a Galilei-Newton spacetime, as described in §2.3 (that is, the linear quadratic classical space + time approximation in Table 1). So, the overall affine structure of the (four dimensional) spacetime yields a natural affine connection, which can be used to detect rotation. Leibniz, however, is considering physical spacetime only as a manifold endowed with a Leibnizian structure (that is, he drops the spacetime affine structure and considers only the pointwise quadratic first nonlinear space + time in Table 1). So, with these elements, no affine connection is determined, and rotation cannot be measured. Summing up, Leibniz was right pointing out that, only with the Leibnizian structure on M at hand, no Galilean connection is selected¹⁵. However, Newton did select such a connection by guessing the further affine structure of M .

4. SECOND NON-LINEARIZATION

4.1. Background: norms, cones and Lorentz-Finsler metrics. In order to show rigorously the emergence of the notion of Finsler spacetime, some purely geometric elements are stressed first. Even though some of them are elementary, they will be necessary to make precise discussions. So, the experimented reader can skip some parts and come back when necessary.

The first ones come from classical norms on a (finite-dimensional, real) n -vector space V and Finsler Geometry; they are carefully explained in [48].

Definition 4.1. A *Minkowski norm* on V is a map $F_0 : V \rightarrow \mathbb{R}$ satisfying

- (i) positiveness: $F_0(v) \geq 0$, with equality if and only if $v = 0$,
- (ii) positive homogeneity: $F_0(\lambda v) = \lambda F_0(v)$ for all $\lambda > 0$,
- (iii) strongly convex indicatrix: F_0 is smooth away from 0 and the *fundamental tensor field* g defined as the Hessian of $\frac{1}{2}F_0^2$ is positive definite on $V \setminus \{0\}$.

Remark 4.2. Notice about this definition:

¹⁵In our opinion, this justifies the name *Leibnizian* used here (following [10]), compare with [55, 76].

MODEL	Linear space: affine Aff with vector V (translation-invariant elements) Geodesics \equiv straight lines		Smooth connected manifold M (pointwise dependent elements)	
	Quadratic forms (doubly linear)	No quadratic restriction	First nonlinearizat.: pointwise quadratic	Second nonlinearizat.: no quadratic restriction
SPACE	Euclidean scalar product g_E on V Symmetry $O(n)$	Minkowski norm $\ \cdot\$ (drop parallelogram identity + reversibility)	Riemannian metric g_0 Unit sphere bundle = pointwise ellipsoid. Levi-Civita natural mathematical choice.	Finsler metric $F(L = F^2)$ Indicatrix = pointwise strongly convex hypers. Cartan connection
Space + time Classic	Galilei-Newton (\mathbf{t}, g_E) - \mathbf{t} absolute time (on V): Non-zero linear form - $(\text{Ker } \mathbf{t}, g_E)$ absolute space: g_E Eucl. scalar product on $\text{Ker } \mathbf{t}$. Symmetry: orthochr. Galilean group	Non-quadratic Galilei-Newton $(\mathbf{t}, \ \cdot\)$ -Replace g_E in Galilei-Newton by a norm. Not developed (as far as we know)	Leibnizian structure: Non-vanishing 1-form Ω (eventually, $\Omega = dt$) with a Riemannian metric on the bundle $\text{Ker}(\Omega)$. Required to choose a linear connection parallelizing Ω and g_R)	Leibniz-Finsler str. -Replace the Riemannian metric on $\text{Ker}(\Omega)$ by a Finslerian one. Not developed (as far as we know)
Space-time Relat.	Special Relat. (g_L, C) - g_L Lorentzian scalar product $(+, \dots, +, -)$ - C time-orientation (choice of one between 2 cones) Symmetry : orthochr. Lorentz group $O_1^\uparrow(n)$	Modified Special Relat. (L_0, C_0) - C_0 cone - L_0 Lorentz-norm on C_0 causal vectors (eventually C_0 determined from L_0) No symmetry but includes the case VSR (with proper a subgroup of $O_1^\uparrow(n)$)	General Relat. g_1: Pointwise smooth Lorentzian scalar product g_1 continuously time-oriented Levi-Civita connection: free fall, lighthlike pregeodesics, gravitational force	Finsler space-time L (with a cone str. C) Pointwise smooth Lorentz-Minkowski norm. Geodesics determined by Cartan (and Chern etc.) connection. C pregeodesics independent of L Anisotropy with causal directions (possibly due to matter/energy)

TABLE 1. Classical models of non-quantum space and time (linear models and their non-linearizations)

- (1) *Positive homogeneity.* This requirement only for $\lambda > 0$ enhances the applications of Finsler Geometry¹⁶, and it will be enough for our purposes. Positive homogeneity implies that F_0 is univocally determined by its *indicatrix* (unit sphere) $\Sigma_0 := F^{-1}(1)$. In particular, the full homogeneity of F_0 becomes equivalent to the symmetry of Σ_0 with respect to the origin.
- (2) *Smoothness.* The standard definition of norm implies that they are only continuous. We assume smoothness (say, C^∞ , pointing out the cases when lower regularity becomes relevant) away from 0. Using (ii), this is clearly equivalent to the smoothness of Σ_0 .
- (3) *Role of triangle inequality.* It is not imposed directly, however:
 - (a) Triangle inequality becomes equivalent (for any 1-homogeneous function smooth away from 0) to the *convexity of Σ_0* (i.e., its inner-pointing second fundamental form σ is positive semidefinite). Moreover, it is also equivalent to the *convexity of the open unit ball $B_0 := F_0^{-1}([0, 1])$* (all the segments connecting points $u, v \in B_0$ are included in B_0).
 - (b) The strict triangle inequality becomes equivalent to the *strict convexity of Σ_0* (the hyperplane tangent to Σ_0 at each point p only intersects Σ_0 at p). Moreover, it is also equivalent to the *strict convexity of the closed unit ball \bar{B}_0* (segments connecting points $u, v \in \bar{B}_0$ are included in the open ball B_0 up to the endpoints u, v).
 - (c) Assuming (i) and (ii), the hypothesis (iii) becomes equivalent to the *strong convexity of Σ* (σ is positive definite), which is more restrictive than its strict convexity.
- (4) *Conic Minkowski norms.* These norms are as in Definition 4.1 just by allowing the map F_0 to be defined only on a cone domain (see Definition 4.5 below) A_0 of V . All the previous considerations on the triangle inequality extend trivially to such conic Minkowski norms.
- (5) *Scalar products.* Norms coming from (Euclidean) scalar products are Minkowski. Conversely, a Minkowski norm comes from a scalar product under one (and, then, both) of the following properties:
 - (a) The classical parallelogram identity holds.
 - (b) F_0^2 is C^2 -smooth at zero [86, Proposition 4.1].
 Recall also that, clearly, any norm coming from a (Euclidean or Lorentzian) scalar product is determined by its value on a cone domain.

Definition 4.3. A *Finsler metric* F on a manifold M is a function $F : TM \rightarrow \mathbb{R}$ satisfying: (i) F is smooth away from the zero section $\mathbf{0} \subset TM$ and (ii) the restriction F_p of F to each tangent space T_pM , $p \in M$, is a Minkowski norm.

Remark 4.4. Notice about this definition:

- (1) *2-homogeneity.* Taking F^2 instead of F , Finsler metrics can be defined alternatively as positive 2-homogeneous functions (this will be convenient for their Lorentzian extensions). What is more, then

¹⁶Including those for relativistic stationary spacetimes, see [26] and references therein.

the C^2 -smoothability of F^2 at 0 would imply that it comes from a Riemannian metric (recall Remark 4.2 (5)).

- (2) *Role of the indicatrix.* As F is determined by its indicatrix $F^{-1}(1)$, a Finsler metric can be defined alternatively as a smooth hypersurface Σ embedded in TM satisfying appropriate conditions, namely: (a) Σ intersects transversely¹⁷ each T_pM and (b) this intersection $\Sigma_p := \Sigma \cap T_pM$ is a strongly convex compact connected embedded hypersurface whose inner domain B_p (such that $\Sigma_p = \partial B_p$, where ∂ denotes the boundary in V)¹⁸ contains the zero vector 0_p .
- (3) *Fundamental tensor on a vector bundle.* Each F_p defines a fundamental tensor field on $T_pM \setminus \{0\}$ and, so, a 2-covariant tensor on each fiber of the (slit) tangent bundle $\pi : TM \setminus \mathbf{0} \rightarrow M$. We will use the letter g to denote such a tensor field, so that, for each $v \in TM \setminus \mathbf{0}$, g_v will be a tensor on T_pM , being $p = \pi(v)$. Clearly, the definition of Finsler metric and fundamental tensor can be extended to any vector bundle, not necessarily the tangent one.

The rest of elements involves the Lorentz-Finsler case, and we follow [46]. We start with the definition of cone. For our purposes, the next one is enough. A more intrinsic definition can be seen in [50, Def. 2.1] (the equivalence and related optimal assumptions are analyzed in [50, §2.1]).

Definition 4.5. A (strong) *cone* \mathcal{C}_0 in V is any embedded hypersurface which can be constructed as follows: choose a hyperplane $\Pi \subset V$ which does not contain 0 and a strongly convex compact connected embedded $(n-2)$ -hypersurface $S_0 \subset \Pi$, take all the half-lines from 0 to the points of S_0 and define \mathcal{C}_0 as the union of all these half-lines except 0.

Then, the *cone domain* is the open subset A_0 obtained analogously by taking the (open) half-lines from 0 to each point of the inner domain B_0 of S_0 in Π (so that the boundary of A_0 in $V \setminus \{0\}$ is \mathcal{C}_0).

Definition 4.6. A *cone structure* on a manifold M is a smooth embedded hypersurface $\mathcal{C} \subset TM$ such that: (a) \mathcal{C} intersects transversely each T_pM and (b) this intersection $\mathcal{C}_p := \mathcal{C} \cap T_pM$ is a cone on T_pM . Then, the *cone (structure) domain* is $A = \cup_{p \in M} A_p$, where each A_p is the cone domain of \mathcal{C}_p . A vector $v \in A$ (resp. $v \in \mathcal{C}$; $v \in A \cup \mathcal{C}$) is called *timelike* (resp. *lightlike*; *causal*).

Remark 4.7. Notice about this definition:

- (1) *Smoothness and transversality.* Intuitively, a cone structure is just to smoothly put a cone at each T_pM , $p \in M$. From the formal viewpoint, however, this cannot be deduced only from the smoothness of \mathcal{C} , making necessary assumption (a) (recall Rem. 4.4 (2) and footnote 17, or the discussion around [15, Fig. 2]).
- (2) *Cone triples.* Any cone structure \mathcal{C} can be determined (in a highly non-unique way) by means of a cone triple (Ω, T, F) , where Ω is any *timelike* 1-form on M (i.e. $\Omega(v) > 0$ for all causal v), T is

¹⁷For the role of the condition of transversality see [15, Prop. 12] and [50, Def. 2.7, Rem. 2.8].

¹⁸Such a B_p exists by Jordan-Brower theorem.

any timelike vector field with $\Omega(T) \equiv 1$ and F is the unique Finsler metric on $\ker(\Omega)$ such that $F(w)T + w \in \mathcal{C}$ for any $w \in \ker(\Omega) \setminus \mathbf{0}$ [50, §2.4]. Conversely, any (Ω, T, F) with $\Omega(T) \equiv 1$ and F Finsler on $\ker(\Omega)$ is the cone triple of some cone structure \mathcal{C} .

- (3) *Extended classical Causality.* \mathcal{C} allows one to extend basic elements of Causality of spacetimes such as the chronological \ll , strict causal $<$, causal \leq and horismotic \rightarrow relations ($p \rightarrow q$ when $p < q$ and $p \not\ll q$) and, thus, the chronological/causal futures and pasts of a point, $I^+(p), I^-(p) / J^+(p), J^-(p)$. In particular, *cone geodesics* are defined as locally horismotic curves, and they generalize the future-directed lightlike pregeodesics associated with the conformal structure of any Lorentz metric.

In the following, we will say that a function is smooth in a manifold with boundary (contained in a regular manifold M) if it can be (locally) extended to a smooth function on an open subset of M .

Definition 4.8. Let \mathcal{C}_0 be a cone on V and $\bar{A}_0 = \mathcal{C}_0 \cup A_0$. A (*properly*) *Lorentz-Minkowski norm* with cone \mathcal{C}_0 is a smooth map $L_0 : \bar{A}_0 \rightarrow \mathbb{R}$ satisfying:

- (i) $L_0(v) \geq 0$ with equality if and only if $v \in \mathcal{C}_0$.
- (ii) L_0 is positive 2-homogeneous: $L_0(\lambda v) = \lambda^2 v$ for all $v \in \bar{A}_0$ and $\lambda > 0$.
- (iii) The fundamental tensor g obtained as the Hessian of $\frac{1}{2}L_0$ has Lorentzian signature $(+, -, \dots, -)$ on A_0 .

Remark 4.9. Consistently with the positive definite case, let us observe the following:

- (1) A less redundant definition for L_0 (as well as for the Lorentz-Finsler metric L below) can be carried out without prescribing the cone \mathcal{C}_0 , see [50, Def. 3.1, 3.5].
- (2) Two homogeneity for L_0 is preferred to 1-homogeneity because of the general equality $L_0(v) = g_v(v, v)$. Notice also that the Lorentzian signature is changed with respect to previous sections and, consistently, if L_0 is smoothly extended around any $v \in \mathcal{C}_0$, then L_0 must become negative away from \bar{A}_0 .
- (3) L_0 is determined by its indicatrix $\Sigma_0 = L_0^{-1}(1)$, which is now *strongly concave* and *asymptotic to \mathcal{C}_0* . Indeed, a Lorentz-Finsler metric could be defined alternatively as a strongly concave hypersurface Σ_0 in A_0 which is asymptotic to some cone structure \mathcal{C}_0 under the mild technical condition that the map $A_0 \ni v \mapsto L_0(v)$ such that $v/L_0(v) \in \Sigma_0$, extend smoothly to \mathcal{C}_0 with non-degenerate¹⁹ g .
- (4) All the properties related to the triangle inequality in the positive definite case (which were associated with the convexity of the indicatrix and held for conic Minkowski norms, Remark 4.2 (3)) are automatically translated now as reverse triangle inequalities in the Lorentz-Finsler case (associated with the concaveness of Σ_0).

¹⁹These conditions would be satisfied by hypersurfaces suitably C^2 -close to the space of observers \mathbf{O} of any relativistic spacetime (notice that some issues appear involving the extendability of L to the cone and whether the cone is prescribed or not), and they can be constructed for any cone (recall Rem. 4.11(4) below).

- (5) Even though $\bar{A}_0 \subset V \setminus \{0\}$, L_0 can be continuously extended to 0 ($L_0(0) = 0$). However, the smoothness of this extension²⁰ depends on whether L_0 comes from a Lorentzian scalar product, as in the positive definite case.
- (6) It is possible to smoothly extend L (preserving the 2-homogeneity) to an open conic subset A_0^* which contains \bar{A}_0 (recall that $0 \notin \bar{A}_0$). This extension is far from unique, but the fundamental tensor in the boundary is well-determined.

Definition 4.10. Let \mathcal{C} be a cone structure on M and $\bar{A} = \mathcal{C} \cup A$. A (*properly*) *Lorentz-Finsler metric* with cone \mathcal{C} is a smooth map $L : \bar{A} \rightarrow \mathbb{R}$ satisfying that the restriction L_p of L to each $T_p M \cap \bar{A}$ is a Lorentz-Minkowski norm. Then, (M, L) is a (*properly*) *Finsler spacetime*.

Remark 4.11. The following results on Finsler spacetimes will be relevant:

- (1) Any Lorentz-Finsler metric can be extended to $TM \setminus \mathbf{0}$ as a smooth 2-homogeneous function with fundamental tensor g of Lorentzian signature, see [67]. However, such an extension is highly non-unique and, as we will see, it is not justified by direct measures of observers.
- (2) Given L , timelike and lightlike geodesics are naturally defined and they satisfy local maximizing properties which extend those of relativistic spacetimes (recall Rem. 4.9(4) and [50, Prop. 6.5]). In particular, the lightlike pregeodesics of L coincide with the cone geodesics of \mathcal{C} [50, §6.2].
- (3) Thus, all the Lorentz-Finsler metrics with the same cone structure have the same lightlike pregeodesics. Two such metrics L_1, L_2 are called *anisotropically equivalent* and they satisfy $L_2 = \mu L_1$ for some 0-homogeneous function $\mu > 0$ on \bar{A} [50, §3.3].
- (4) Any cone structure \mathcal{C} is associated with a Lorentz-Finsler metric L (and, then, with its anisotropically equivalent class). Indeed, if \mathcal{C} is determined by a cone triple (Ω, T, F) , one can construct such an L starting at the map

$$G(v) := \Omega(v)^2 - F(\pi_2(v))^2, \quad \forall v \in \bar{A}, \quad (9)$$

where $\pi_2 : TM = \text{Span}(T) \oplus \ker(\Omega) \rightarrow \ker(\Omega)$ is the natural projection. G satisfies all the required properties of L except the differentiability on $\text{Span}(T)$, the latter because of the lack of differentiability of F^2 at 0 when it is not Riemannian. Indeed, the indicatrix $G^{-1}(1) \subset A$ is not smooth precisely on T , that is, only at the point T_p on each p . However, standard techniques of smoothability for convex functions allow one to smoothen G around T obtaining the required L [50, §5.2].

- (5) The lack of differentiability of G above is analogous to the well-known lack of differentiability of any product of (non-Riemannian) Finsler manifolds. Indeed, if (M_1, F_1) is a Finsler manifold then $dt^2 \oplus (\pm F_1^2)$ are not smooth as Finsler or Lorentz-Finsler metrics on

²⁰ Recall that, for any function L_0 on $\bar{A}_0 \cup \{0\} \subset V$ (with A_0 a cone domain), the elementary definition of existence of a differential map at 0 makes sense because 0 is an accumulation point of the domain of L_0 and its uniqueness is guaranteed because A_0 contains n independent directions converging to 0.

$\mathbb{R} \times M_1$ along the direction ∂_t . This problem prevents the extension to the Lorentz-Finsler case of the trivial procedure to construct a relativistic product spacetime starting at a Riemannian manifold.

- (6) Given a Lorentz-Finsler metric, there exists a univocally determined A -anisotropic connection which is torsion-free and parallel. Moreover, when we consider a properly Lorentz-Finsler metric, this A -anisotropic connection can be extended to an open subset A^* which contains $\bar{A} \setminus \mathbf{0}$. As the extension away from \bar{A} is highly non-unique, we will speak about \bar{A} -anisotropic connections. When $A = TM \setminus \mathbf{0}$, we will just say *anisotropic connection*²¹.

Due to this last item, the definitions of some classes of Finsler spacetimes such as the static ones have included the possibility to have some non-smooth directions [16, 17, 58]. However, the smoothing procedure mentioned in part (4) is also applicable to these cases. This shows that, from the foundations viewpoint, the motivation for non-smooth metrics is not stronger for the Lorentz-Finsler case than for classical relativistic spacetimes [50, §4.2, §4.4].

Note 4.12. Nevertheless, there are some physical considerations (see §6.1) which lead to examples where L is not smoothly extendible to the cone structure \mathcal{C} , even if:

- (i) its cone \mathcal{C} is smooth (so, the cone geodesics are well defined), and
- (ii) the A -anisotropic connection can be smoothly extended to \mathcal{C} (so, the Finslerian curvature tensors are well-defined on \mathcal{C}).

Such examples could also be included in our definition of Lorentz-Finsler metrics and spacetimes, as all the relevant geometric properties remain. However, we will consider for simplicity that L is smooth at \mathcal{C} and, when this property does not hold, we refer to them as *improper* and we will discuss whether (i) and (ii) hold then. Accordingly (and consistently with [50, Def. 3.1]), an *improper* Lorentz-Minkowski norm L_0 satisfies all the properties in Def. 4.8 but the differentiability at $L_0^{-1}(0)$.

Remarkably, a large class of spacetimes satisfying both conditions (i) and (ii) can be found following [73]. Namely, they hold for any two-homogeneous function L defined on the set of causal vectors \bar{A} determined by a cone structure \mathcal{C} such that: (a) L is zero on \mathcal{C} and determines a Lorentz-Finsler metric in the interior A of \bar{A} and (b) there is a power of L which is smooth on \mathcal{C} with non-degenerate Hessian (notice that, in [73], L is assumed to be defined on the whole TM).

Anyway, there are some examples of Finsler spacetimes in the literature that do not even satisfy our weaker definition of improper Lorentz-Finsler spacetime, such as Randers spacetimes or those introduced by Kostelecky [53, 54], which are the effective model of some particles with no GR background (see the discussion in [50, Appendix A, B]).

²¹ Essentially, this is a connection where, formally, the Christoffel symbols of a chart (U, φ) depend also on the direction and, so, they are functions on $TU \cap A \subset TM \setminus \mathbf{0}$, which are positive homogeneous of degree zero. The name and a thorough study of A -anisotropic connections were given in [44, 45]; see also [63, 64] for a study of connections on fiber bundles from a more general viewpoint.

4.2. Physical intuitions for Finsler spacetimes. Next, our aim is to justify physically our definition of Finsler spacetime (Def. 4.10), supported by some mathematical properties pointed out above. The first consideration is that Postulate 2 should be regarded now as an approximate symmetry at each point, in a similar way as the affine structure of Postulate 1 has been regarded as an approximate symmetry to the structure of a relativistic spacetime²². This means that, now, one cannot find a set of coordinate charts such that the relations (3) occur at each p ; however, one would expect that we will not be far from this situation (at least in regions of spacetime free of extremely exotic or violent situations). Consistently, we will not have the sets \mathbf{S}_p of linear bases at each T_pM playing the role of (linear) IFR at p . However, one would expect that the set of observers \mathbf{O} introduced in Def. 3.1 will still make sense and will be “close” to the space of observers for a relativistic spacetime. As the latter is a hyperboloid (asymptotic to a quadratic cone) at each point p , now, \mathbf{O}_p should be a strongly concave hypersurface asymptotic to some cone structure defining a Lorentz-Minkowski norm at p (see Rem. 4.9 (3)) and, moreover, \mathbf{O} should be identified as the indicatrix Σ of a Lorentz-Finsler metric L .

Remark 4.13. The previous discussion leads us to a Lorentz-Finsler metric L with indicatrix Σ equal to \mathbf{O} which lies exactly under our Def. 4.10 (including also the improper case explained in Note 4.12). The way to arrive at this definition from the viewpoint of symmetries can be summarized as follows.

(1) Following [34], consider the connected parts of the identity $ISO_1(4)$, $SO_1(4)$, $ISO(3)$, $SO(3)$ of the Poincaré, Lorentz, Euclidean and orthogonal groups, resp. In Special Relativity, the homogeneous spaces obtained as the quotients $ISO_1(4)/SO_1(4)$, $ISO_1(4)/ISO(3)$, $ISO_1(4)/SO(3)$ are, respectively, the spacetime, the space of all the (rest) spaces (i.e., the space of all the spacelike hyperplanes, being the standard rest space $ISO(3)/SO(3)$) and the space of observers \mathbf{O} (being the space of velocities $SO_1(4)/SO(3)$). Here, \mathbf{O} is metrically identifiable with $\mathbb{R}^4 \times H_+^3$.

(2) In General Relativity, \mathbf{O} is identified with the set Σ^g of all the future-directed unit vectors. Σ^g is a subbundle of TM whose fibers are affine hyperboloids at each tangent space. Such hyperboloids characterize g univocally so that the information of g is codified in \mathbf{O} .

(3) For the space of observers \mathbf{O} in the Lorentz-Finsler setting, Σ^g is replaced with a hypersurface Σ satisfying formal properties analogous to Σ^g (but dropping its pointwise symmetries) so that it characterizes a Lorentz-Finsler metric.

Next, let us discuss more carefully the physical grounds of Def. 4.3:

(1) The fact that L is defined only on a cone domain A and it is extended continuously to \mathcal{C} comes from the nature of the space of observers.

Recall that, then, one has timelike geodesics (Rem. 4.11 (2)) and, thus, freely falling observers. At least from a trivial mathematical

²² Even though we focus on the relativistic case, (disregarding the Leibnizian case and the other possibilities), one could also consider a *Leibniz-Finsler structure* (Ω, h) on a manifold M , where h would be now a Finsler metric on $\text{Ker}(\Omega)$ instead of a Riemannian one, according to Table 1.

viewpoint, this is enough to determine L and, then, the fundamental tensor g on the cone domain A .

Notice that, given an observer $v \in \Sigma_p$, the tensor g_v is then also obtained *on the directions* of $T_v\Sigma_p$. In principle, g_v (which can be obtained just from Σ) could be measured, as it comprises properties of neighboring observers.

- (2) The smooth extensibility of both L and the fundamental tensor g (as a non-degenerate one) to the cone structure \mathcal{C} appears as a natural approximation (in principle, one would expect to remain close to the situation in a relativistic spacetime) which mathematically ensures that \mathcal{C} is truly a cone (with S_0 in Def. 4.5 satisfying strong convexity). Moreover, then L also determines lightlike geodesics which, up to reparametrization, are inherent to the cone structure. The improper case of Finsler spacetimes satisfying the properties (i) and (ii) in Note 4.12 would also satisfy all these properties about geodesics and cones.

Then, as a consequence, the behaviour of lightlike geodesics becomes completely analogous to the classical relativistic case. Indeed, Lorentz-Finsler metrics with the same cone structure are also related by an “anisotropic conformal factor μ ” (Rem. 4.11(3)) and the cone structure \mathcal{C} also allows one to mimic the relativistic behaviour of Causality (Rem 4.7(3)).

- (3) The physical considerations in the two previous items are also assumed in standard Relativity. Namely, observers are always expected to measure only massive or massless particles, that is, elements with velocities in a causal cone. In General Relativity, this is apparent from the EPS formulation, where radar coordinates are systematically used with this aim (see the next section). Certainly, the metric tensor g is assumed to be defined on all the directions in the relativistic case but the underlying reason is that g is *fully determined by its value along the causal vectors* (Rem. 4.2(5)). This is not by any means true in the Lorentz-Finsler case, even if L can be extended to the whole TM (recall Rem. 4.11(1)).
- (4) When a spacelike separation in a direction l is going to be measured by an observer v , it seems natural to consider $g_v(l, l)$; so, it would be irrelevant whether L is not defined outside the cone.

Indeed, from a purely geometric viewpoint, $T_v\Sigma_p$ would be naturally regarded as the rest space of the observer v at p , and g_v would be the unique metric available there, even though the physical process to measure it might not be obvious. It is worth pointing out here Ishikawa’s claim in [43] that $g_v(l, l)$ can be measured assuming that the physical light rays are those of g_v . Indeed, this author criticizes Beem’s definition of light rays, who constructed them by using the lightlike vectors on the cone \mathcal{C} . Anyway, in our opinion, Ishikawa’s claim needs further physical support.

- (5) It is worth emphasizing that no issue on smoothability occurs with Σ , which can be assumed smooth (as in Rem. 4.2 2) in most interesting cases. Indeed:

(a) The Lorentz-Finsler metric L cannot be C^2 -extended to 0, in agreement with the behavior of norms in both the positive definite case and the Lorentz-Finsler one (Rem. 4.2(5)). However, no physical nor mathematical reason seems to require the smoothability of L at 0 (compare with the EPS approach in §5.2.1 below).

(b) Product metrics $-dt^2 + F^2$ or, with more generality, the rough Lorentz-Finsler version of static spacetimes $-\Lambda(x)dt^2 + F^2(x, y)$, with natural coordinates (x, y) at TM , are never smooth at ∂_t whenever F is Finsler but not Riemannian. Consequently, some authors have included the possible existence of non-smooth directions as a fundamental ingredient of Lorentz-Finsler metrics (see for example [16, 17, 58]). Nevertheless, as explained in Rem. 4.11, parts (4) and (5), general smoothing procedures can be applied. What is more, a natural definition of (smooth) static spacetimes as well as an explicit procedure to construct locally all of them are available at [50, §4.2].

(c) Other issues of non-smoothness appear when modelling some specific physical situations (Very Special Relativity, birefringence) and will be considered in §6.1.

5. COMPARISON WITH EHLERS-PIRANI-SCHILD APPROACH

5.1. Summary of the approach. EPS approach [21] constructs step by step each geometric structure of physical spacetime (until reaching the metric) by means of physically motivated axioms:

- (1) Spacetime becomes a *differential manifold* M endowed with a *cone structure* \mathcal{C} . Essentially, this is obtained by means of axioms on light propagation which involve messages and echoes between particles.

Indeed, these axioms allow one to find *radar coordinates* with respect to (*freely falling, massive*) *particles*, the latter represented by a class of unparametrized curves, which provide the structure of differentiable manifold, see EPS axioms D_1 — D_4 . Then, the cone structure \mathcal{C} is obtained by using two axioms, L_1, L_2 , on the local character of light propagation around each event e . Indeed, L_1 states that given any particle P with some parameter t which passes through e , it follows that any event p ($p \notin P$) can be connected with the particle by exactly two light rays²³, while L_2 distinguishes two connected components for light rays. Moreover, L_1 also states that, if these two rays cross the curve at the events e_1, e_2 , then $g(p) := -t(e_1)t(e_2)$ is required to be smooth in a small neighborhood of e . EPS claims that, then, \mathcal{C} will come from the *conformal structure* of some *Lorentz metric* (a particular case of our Def. 4.6) and, so, we can speak about \mathcal{C} -*timelike* directions.

- (2) Spacetime is endowed with a *projective structure* \mathcal{P} . This is achieved by means of two axioms, P_1, P_2 , which model the free fall of particles.

²³Along the events $\tilde{e} \in P$, all the light rays from \tilde{e} would trivially cross P at \tilde{e} ; so, the function g below would be trivially extended as $g(\tilde{e}) = -t(\tilde{e})^2$. However, the points on P would be excluded in order to define the differentiable structure of the manifold by using radar coordinates (recall the example in footnote 24 below).

The first one states only the existence of a unique particle, represented by means of an (unparametrized) curve, for each event e and \mathcal{C} -timelike direction at e . The second axiom states that, around each event e , one can find coordinates \bar{x}^i such that *any* particle through e admits a parametrization $\bar{x}(\bar{u})$ satisfying:

$$\left. \frac{d^2 \bar{x}^i}{d\bar{u}^2} \right|_e = 0. \quad (10)$$

This equality is regarded as an *infinitesimal law of inertia* (consistently with Trautman [82]). By using (10), EPS argues that a *projective structure*, which is claimed to be compatible with some affine connection \mathcal{A} , must appear. As a consequence, not only the original particles would be recovered as pregeodesics of \mathcal{A} but one would also obtain *pregeodesics at any direction*, timelike or not.

- (3) Spacetime is a *Weyl space* $(M, \mathcal{C}, \mathcal{A})$, where \mathcal{A} is an affine connection *compatible* with the cone structure \mathcal{C} , in the sense that the lightlike \mathcal{C} -pregeodesics are also \mathcal{A} -pregeodesics. This is carried out by means of their axiom C , which matches particles and light rays.

Specifically, this axiom assumes that, around each event e , any point in the \mathcal{C} -chronological future of e lies on a particle through e . This will imply that the lightlike \mathcal{C} -pregeodesics of the conformal structure (namely, the \mathcal{C} -cone geodesics, see Remark 4.11 2) are also pregeodesics for the projective structure \mathcal{P} in the step (2). Then, EPS claims that such a compatibility selects a unique affine connection \mathcal{A} compatible with the projective structure.

- (4) Spacetime is endowed with a (*time-oriented*) *Lorentzian metric* g , up to an overall (constant) scalar factor. This is obtained by means of a *Riemannian axiom*, which takes into account that \mathcal{A} has its own parallel transport and its curvature tensor; the axiom imposes the compatibility of (one of) these two elements with g .

Indeed, they state that the Riemannian compatibility of $(M, \mathcal{C}, \mathcal{A})$ is equivalent to any of the following conditions: (a) the vectors obtained by \mathcal{A} -parallel transport of a single one v at $p \in M$ along two curves with the same endpoint q have the same norm at q (computed with any of the homothetic scalar products compatible with \mathcal{C}_q), or (b) using Jacobi fields to construct arbitrarily close particles, the proper times of two of such particles are linearly related at first order, that is, the regular ticking of a clock for the first particle implies the regular ticking for the second one.

About these axioms and proofs, EPS admits: “a fully rigorous formalization has not yet been achieved”. Next, we will focus just on the relation of EPS approach with Lorentz-Finsler metrics. For progress on EPS approach, see for example [77].

5.2. Keys of compatibility with Finslerian spacetimes. The fact that a Finslerian spacetime can fulfill the EPS axioms was already pointed out by Tavakol & Van den Berg [81], who considered the case of Berwald spaces. Now, our aim is to revisit precisely the compatibility of the four EPS steps with Finslerian elements, as well as [81].

5.2.1. *EPS step (1)*. Recently, Lammärzhal and Perlick [57] have argued against the role of smoothness of the function $g(p)$ at e in the step (1). This differentiability becomes essential, because the equalities $g(e) = 0, g_{,a}(e) = 0$ allow EPS to find a metric tensor $g_{,ab}(e)$ compatible with \mathcal{C} .

Indeed, there are subtle differences at this point in comparison with the introduction of radar coordinates, which are used to settle the smooth (C^3) manifold structure of the spacetime. Certainly, EPS were aware of the existence of non-trivial subtleties, as one can read at the beginning of their subsection *Differential Topology*: “The reason that we do not take this structure [smooth manifold] for granted is that differentiability plays a crucial role in our introduction of null cones (...) and in the infinitesimal version of the law of free fall”. The following three items must be taken into account in the EPS development:

(i) The axioms D_1 — D_4 , which allow one to define radar coordinates, should apply to particles P, Q which do not intersect. Otherwise, spurious differential issues might appear even in the case of Lorentz-Minkowski spacetime²⁴.

(ii) Axiom L_1 , however, considers the functions $p \mapsto t(e_1), p \mapsto t(e_2)$ (which would be radar coordinates for some particle P through e) defined even on P. Moreover, this axiom ensures that the particular combination $g(p) = -t(e_1)t(e_2)$ is C^2 -differentiable on P too.

(iii) In the discussion above [21, Lemma 1], they explain that $t(e_1) = t(e_2) = 0$ occurs if and only if $p = e = e_1 = e_2$ (thus, $p \in P$) and they focus on this case. Then, EPS argues first that the differential $g_{,a}(e)$ must be 0 by applying L_2 and, using C^2 differentiability, they show that the light directions must lie in the quadratic cone of the lightlike vectors of $g_{,ab}(e)$.

Recall, however, that there is *no physical justification* about why g must be differentiable or C^2 . Notice that g is constructed from the functions $e \mapsto t(e_1)$ and $e \mapsto t(e_2)$, which are not smooth even in the Lorentz-Minkowski spacetime (see Example 5.1 below). This assumption on the product $t(e_1)t(e_2)$ yields a posteriori the quadratic character of the cone, forbidding more general cone structures.

From a purely mathematical viewpoint, the smoothness issue on the radar coordinates above would be similar to the differentiability of the radial coordinate r of a normed vector space at 0: r is never smooth at 0 and r^2 is smooth if and only if the norm comes from a Euclidean scalar product (Rem. 4.2(5)). So, such an a priori assumption would be completely *unjustified from a mathematical viewpoint* too, indeed:

- (a) There are norms with an analytic indicatrix (thus, analytic away from 0) which do not come from a scalar product. For example, on \mathbb{R}^2 , when the indicatrix is equal to the curve in polar coordinates $\rho(\theta) = 1 + \epsilon \sin \theta$ for small $\epsilon > 0$ (so that it is strongly convex).
- (b) Euclidean scalar products are very particular cases of analytic norms.

²⁴ For example, let P be the t -axis and $Q = \{(t, x = t/2, y = 0, z = 0) : t \in \mathbb{R}\}$. A message from Q to P would yield the map $t \mapsto t/2$ if $t \leq 0$ and $t \mapsto 3t/2$ if $t \geq 0$ (see the Example 5.1 below) which is not smooth at 0, in contradiction with D_2 (recall also footnote 23).

That is, *the apparently mild EPS requirement of differentiability at 0 becomes even stronger than analyticity for a norm.*

Example 5.1. Let us see the role of smoothability for the EPS function g obtained by using a pair of radar coordinates with respect to a particle (according to EPS, one should take two pairs of radar coordinates by choosing two particles). We will work on $M = \mathbb{R} \times \mathbb{R}^3$. Let $t : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{R}$ be the natural projection, consider any Minkowski norm F_0 on \mathbb{R}^3 and take spherical-type coordinates (r, θ, φ) on \mathbb{R}^3 (up to suitable points) with θ, φ , the usual spherical angles and $r \equiv F_0$; then, extend the functions r, θ, φ to $\mathbb{R} \times \mathbb{R}^3$ in a t -independent way. Let \mathcal{C} be the natural (constant) cone structure given by $t(p) = r(p)$ and regard the t -axis as a particle P. The corresponding radar coordinates are $t \pm r$ and thus, the EPS g is $g(p) = -t^2(p) + r^2(p)$. This function is smooth at 0 if and only if F_0 comes from a Euclidean scalar product²⁵. Anyway, the cone structure is smooth, because it is determined by the cone triple (dt, ∂_t, F_0) and, so, it is compatible with a smooth Lorentz-Finsler metric L (indeed, a Lorentz-Minkowski norm), see Rem. 4.11(4). As stressed in the item (5) below Rem. 4.13, the fact that $-dt^2 + F_0^2$ is not smooth at ∂_t neither contradicts the existence of a smooth L nor introduces any issue of smoothability.

5.2.2. *EPS step (2).* The way how EPS deduces the existence of the projective structure \mathcal{P} from the infinitesimal law of inertia (10) consists in rewriting this last formula in arbitrary coordinates to obtain [21, formula (7)]

$$\ddot{x}^a + \Pi_{bc}^a \dot{x}^b \dot{x}^c = \lambda \dot{x}^a \quad (11)$$

where λ depends on the parameterization $x^a(u)$ of the curve and Π_{bc}^a depend on x^a . These functions are called the *projective coefficients*, as they would determine a projective structure \mathcal{P} compatible with some affine connection.

However, if one allowed the functions Π_{bc}^a to depend on the direction of the velocities \dot{x}^j , then $\Pi_{bc}^a(x^i, \dot{x}^j)$ could represent the formal Christoffel symbols for a Lorentz-Finsler metric L (indeed, for its A -anisotropic connection, see Remark 4.11, item 6). Thus, the solutions of (11) would be pregeodesics for L which satisfy the law of inertia (10), up to the following issue of C^2 -differentiability of the chart coordinates at the origin.

The existence of *normal coordinates in \mathcal{C} -timelike directions* (which is ensured for any A -anisotropic connection²⁶) would be the natural mathematical translation for the law of inertia. However, the Christoffel symbols of a Lorentz-Finsler metric might not be even continuous at the origin by the trivial reason that these symbols may depend on the direction but they cannot vary along each direction (they are homogeneous of degree 0). Thus, its exponential map is not guaranteed to be C^2 at the origin unless the anisotropic connection is affine (i.e., it does not depend on the direction). It

²⁵Of course, one could introduce a spurious differential structure on \mathbb{R}^4 so that r^2 becomes smooth for a non-Euclidean F_0 , but this would not be natural by any means.

²⁶In principle, the normal coordinates can be defined when the anisotropic connection is defined for all the vectors in $TM \setminus 0$, but it is always possible to extend the \bar{A} -anisotropic connection to all directions locally (see [50, Remark 6.3], where the Lorentz-Finsler case is considered in detail). These coordinates are obtained using the exponential map in a neighborhood as in [50, Lemma 6.2].

is known that, for a positive definite Finsler metric, this happens if and only if the metric is of Berwald type²⁷ [2] (see also [4, Ex. 5.3.5]) and this can be extended to the Lorentz-Finsler case. Indeed, this type of metrics provides the Lorentz-Finsler examples beyond EPS suggested in the literature, see §5.2.5.

Summing up, we emphasize: (a) the coordinates provided by the exponential map of a Lorentz-Finsler metric at any event e are smooth along the half-lines starting at e and they satisfy (10), and (b) to exclude anisotropic connections because of their lack of smoothness at 0 is a subtle mathematical issue and (as in the discussion of the Step 1 in §5.2.1) this is not justified in EPS neither physically nor mathematically. Thus, the law of inertia should be regarded as compatible with Lorentz-Finsler metrics according to our definition (where the directions outside the causal cone are not taken into account), including even the improper case in Note 4.12).

5.2.3. *EPS step (3)*. The compatibility of $(\mathcal{C}, \mathcal{P})$ as a Weyl space with a (unique) affine connection \mathcal{A} obtained by using EPS axiom C becomes a subtle question. On the one hand, Trautman [83] claimed the necessity of a detailed proofs in his review on the reprinted EPS article and, shortly after, this author and V. Matveev [66] characterized when a pair $(\mathcal{C}, \mathcal{P})$ is compatible. On the other hand, the notion of Weyl space as the triple $(M, \mathcal{C}, \mathcal{A})$ given by EPS does not coincide with the standard one of Weyl geometry²⁸. Some authors questioned whether such an EPS structure permits to define a standard Weyl one as well as EPS development at this step. However, very recently, this question has been positively answered by Matveev and Scholtz [65], vindicating the EPS approach.

We emphasize that the EPS compatibility axiom C can be stated with no modification in the case that \mathcal{C} is any cone structure and \mathcal{P} is the projective class of pregeodesics of any \bar{A} -anisotropic connection defined on all the \mathcal{C} -causal directions (as already commented, \mathcal{C} determines intrinsically cone geodesics extending those in EPS conformal cones, Remark 4.7(3)). So, the possibility to extend previous results to this setting should be explored.

5.2.4. *EPS step (4)*. In the EPS spirit, the *Riemann axiom* would be any (minimum, physically well-motivated) assumption making a compatible triple $(\mathcal{C}, \mathcal{P}, \mathcal{A})$ also compatible with a Lorentzian metric, as the conditions labelled (a) and (b) at step (4). However, in order to state now a *Finslerian axiom*, one should notice that these conditions involve \mathcal{A} and, so, they might depend on the way how the previous step is solved.

Anyway, it is worth pointing out some reasons which would support the convenience of such a Finslerian axiom. On the mathematical side, the results collected in Rem. 4.11 (parts (3) and (4)) show a natural consistency: (i) any \mathcal{C} can be associated with a Lorentz-Finsler metric L , (ii) any other associated L' is anisotropically related to L , and (iii) the lightlike pregeodesics

²⁷This means that its Chern-Rund connection defines an affine connection on the underlying manifold, see [80] for quite a few of characterizations.

²⁸In modern language, a Weyl geometry on M is a conformal structure \mathcal{C} endowed with a connection on the \mathbb{R}^+ -principle bundle $P \rightarrow \mathcal{C}$, where the fiber of P at each \mathcal{C}_p is the class of homothetic Lorentzian scalar products compatible with \mathcal{C}_p (see for example [27]); such a notion was considered in references on EPS as [23].

of all the associated Lorentz-Finsler metrics agree with the cone geodesics of \mathcal{C} . On the physical side, the standard chronometric approach is reduced to the determination of the indicatrix of the observers at each event and this would depend only on the behaviour of clocks and measurements of proper time²⁹. Notice that, in the Finslerian case, this behaviour would not be restricted by any condition of quadratic compatibility (but only by a mild overall concaveness and asymptoticity to \mathcal{C}).

5.2.5. *Finslerian examples strictly compatible with EPS.* As we have explained, the requirement of C^2 smoothability at 0 for cones and geodesics is the main gap in the EPS approach. However, Tavakol and van der Berg [81] showed Finslerian examples which are even compatible with this requirement. Next, let us analyze these and other possible examples of Finsler EPS compatible (FEPS) spacetimes.

A very simple FEPS example would be the following. Consider an affine space endowed with any Lorentz norm L_0 with the same cone as a Lorentzian scalar product $\langle \cdot, \cdot \rangle$ (L_0 can be obtained by perturbing the indicatrix of $\langle \cdot, \cdot \rangle$, as explained in Remark 4.11, item (4)). Then, the cone and geodesics of L_0 would satisfy all the EPS axioms, including those of C^2 smoothness at 0. Here, the key is that the affine parallel transport preserves both, the indicatrix of L_0 and $\langle \cdot, \cdot \rangle$.

Remark 5.2. Tavakol & van der Berg examples also obey this pattern, even though they are more refined and interesting. Indeed, they are Berwald type spacetimes constructed by using an auxiliary Lorentz metric g . The fact that they are FEPS examples becomes apparent, because they have the same cone and geodesics as g .

However, we emphasize that these FEPS examples are *not* in contradiction with the EPS conclusions. Indeed, the above examples only show that the physical elements \mathcal{C} , \mathcal{A} , under the EPS restrictions, may be compatible with two different geometric structures, the Lorentz g and Lorentz-Finsler L metrics. To decide which of them would be physically more appropriate would depend on further physical input. In absence of such input, the use of g would be mathematically simpler. Nevertheless, this input might appear from the measurements of proper time, as suggested at the end of §5.2.4.

In order to obtain a true Finslerian contradiction with EPS conclusions, one should construct a Lorentz-Finsler metric L with associated cone \mathcal{C} and anisotropic connection \mathcal{A} satisfying:

- (i) the EPS C^2 requirements,
- (ii) the cone geodesics of \mathcal{C} are pregeodesics of \mathcal{A} ,
- (iii) \mathcal{C} is invariant under the \mathcal{A} -parallel transport, and
- (iv) \mathcal{A} is not compatible with any Lorentz metric.

However, the following known results on linear algebra and Finsler metrics suggest the difficulty to find such a contradiction. Notice that the Finslerian results have been obtained in the positive definite case (the last one after the original EPS paper) and their suitable extensions to the Lorentz-Finsler case is not always clear:

²⁹Compare with EPS claim (1) in §5.3 below.

(a) *The square of a norm is C^2 at 0 if and only if it comes from an Euclidean scalar product* ([86], §4.2, item 5 (b)). As a consequence, the C^2 requirement (i) implies the Lorentzian character of the cones, §5.2.1.

(b) *A linear map between two Lorentzian vector spaces is homothetic if and only if it preserves the lightcones*³⁰. As a consequence, if \mathcal{C} is compatible with a Lorentzian metric g (as established in (a)), the preservation of \mathcal{C} under \mathcal{A} -transport in (iii) implies that this transport must be a g -homothety; in particular, the Riemann axiom (its version (a) in §5.1, item 4) is satisfied.

(c) *The exponential of a Finsler metric is smooth at 0 if and only if it is Berwald* [2]. As a consequence, the law of the inertia (with the C^2 requirement (i)) would imply that only Berwald-type Lorentz-Finsler metrics could be admitted, §5.2.2).

(d) *All Finsler metrics of Berwald type metric are affinely equivalent to a Riemann space*, that is, their affine connections are Levi-Civita for Riemannian metrics (Szabó, [79]).

Notice that, in the case that a suitable Lorentz-Finsler version of this last result existed (taking into account, eventually, the requirement (ii)), this would imply that FEPS is also compatible with a Lorentz metric, that is, the requirement (iv) could not be fulfilled if (i), (ii), (iii) held.

Remark 5.3. Recently, Fuster et al [28] have shown that there are Berwald-type Finsler spacetimes which are not affinely equivalent to a Lorentz metric. However, they contain non smooth directions; this must be taken into account for the comparison with Szabó's result or the possible contradiction with EPS. Anyway, they show a minimal violation of smoothness. Indeed, their examples include improper Lorentz-Finsler metrics L , satisfying both (i) and (ii) in Note 4.12 and, moreover, they satisfy that some power L^r (with $r > 1$ and integer) is smooth even at the lightlike directions of their cone, see Remark 6.3.

Remark 5.4. Recently, Hohmann et al [39] have classified the Berwald spacetimes which are spatially homogeneous and isotropic. Among them, they have found a genuinely Finslerian class (with cones equal to classical FLWR spacetimes). As a proper Finslerian extension of relativistic cosmological spacetimes, the interest of this FEPS class is remarkable (even if it is not clear that they yield a true contradiction with EPS or not).

5.3. Constructive EPS approach vs observer's approach. In order to compare EPS approach and ours, notice first that EPS distinguishes between a *chronometric* approach à la Synge [78] and their *constructive* approach. The former one regards the concepts of particle and standard clock as basic, and introduces the metric g as fundamental. So, it regards as primitive an easily measurable physical quantity (proper time) and a single geometric structure (the metric), the latter encoding all the other geometric elements in a simple way. As a consequence of these advantages, the chronometrical approach is very economical. However, EPS also pointed out drawbacks such as:

- (1) the impossibility to construct the metric from the behavior of the clocks alone,

³⁰See for example [8, §2.3].

- (2) the inclusion by hand of the hypothesis that metric geodesics will correspond with free motion and, then,
- (3) the expectation that the clocks constructed by means of freely falling particles and light rays will agree with the metric clocks.

This motivated their constructive approach starting at basic elements (events, particles, light rays) and axioms close to the physical experience. Certainly, EPS aimed to deduce the metric structure from their axioms. However, the difficulties found in some points (as explained in §5.2.3, the step (3) would have been solved only very recently) as well as the necessity to introduce a Riemannian axiom at the end, makes the procedure somewhat awkward.

In contrast, our approach is neither chronometric nor constructive; instead, it only appeals to the way how we measure. As such a procedure is complex, one starts at the ideal situation when some symmetries among measurements are assumed (our two postulates). Under our viewpoint if such symmetries did not hold at all, it would not be clear even the meaning of the verb “to measure”. However, in the case that the symmetries can be invoked as an approximation, the meaning of measurements can be recovered. Then, the emergence of some geometric structures resembles a sort of experimental Klein’s Erlangen program.

Notice that only hypotheses on the way of taking coordinates of space and time (inertial reference frames, observers) were assumed. It is noteworthy that only some few possibilities emerged for the geometry of spacetime when these symmetries hold in a strict way. From the standard physical viewpoint (close to philosophical realism), the fact that the space, time and matter allow us to measure in some specific way should be interpreted as an evidence about the power of the emerged geometric structures in order to describe the physical spacetime.

Anyway, it is also worth noticing that our final geometric model of spacetime (a manifold endowed with a Lorentz-Finsler metric defined only on the set \bar{A} of causal vectors for a cone structure) is compatible with EPS approach. Indeed, as shown in the previous subsection, EPS excluded the properly Finslerian case only due to two mathematical subtleties about unjustified restrictions of smoothness in radar coordinates (step (1)) and the law of inertia (step (2)). As pointed out in our discussion at §5.2.4, in the case that \mathcal{C} (or the Weyl pair $(\mathcal{C}, \mathcal{P})$ in the step (4)) were not assumed to be compatible with a Lorentz metric, the Riemannian axiom might be replaced by a Finslerian one which would involve only the behaviour of clocks.

Finally, we emphasize that EPS approach also gives a strong support to our hypothesis that, in principle, the Lorentz-Finsler metric must be defined only at the causal directions in \bar{A} : no basic element in the EPS approach (particle, light rays, radar coordinates, echoes) involves non-causal directions.

6. LORENTZ SYMMETRY BREAKING

The implications of the introduction of Finslerian geometry may be more transparent if we focus on the Lorentz symmetry breaking which occurs when Lorentz-Finsler norms are used to extend Special Relativity (i.e., when one considers only the second non-linearization in Table 1). We will center

around this breaking from our theoretical viewpoint; for a more experimental one, a review on tests of Lorentz invariance (which includes Lorentz-Finsler possibilities and discussions on von Ignatowski approach) was updated in 2013 by Liberati [61].

6.1. Modified Special Relativity. Assume that the spacetime has a structure of affine n -space Aff and it is endowed with a Lorentz-Minkowski norm L_0 rather than a Lorentz scalar product $\langle \cdot, \cdot \rangle_1$. Roughly speaking, this is a generalization of Special Relativity where, instead of dropping Postulate 1 (as in General Relativity), we are dropping Postulate 2. Thus, one has affine reference frames but no IFR's; however, one can still assume that any physically relevant vector basis B will be composed of a timelike vector with respect to the cone \mathcal{C}_0 associated with L_0 and three non-causal ones spanning a spacelike hyperplane Π ($\Pi \cap \mathcal{C}_0 = \emptyset$).

Remark 6.1. There is a mathematical analogy between the transition from $\langle \cdot, \cdot \rangle_1$ to L_0 and the one from Special to General Relativity. The latter goes from the point-independent $\langle \cdot, \cdot \rangle_1$ to a Lorentz metric g_p which depends on the point p in an n -manifold M . In the former transition the vector space V associated with Aff is endowed with a Lorentzian metric g_v which depends on the direction of $v \in \bar{A}_0$ for some cone structure \mathcal{C}_0 . What is more, the independence of g_v with the radial direction ($g_v = g_{\lambda v}$ for $\lambda > 0$) makes relevant only the variation of v on a topological $(n - 1)$ -spherical cap.

6.1.1. VSR and GVSR. The transition from $\langle \cdot, \cdot \rangle_1$ to L_0 appears naturally in the so-called *Very Special Relativity* (VSR). This was introduced by Cohen and Glashow [18] who realized that most physical theories (including those satisfying the charge-parity symmetry) which are invariant under certain proper subgroups of the Poincaré group have the symmetries of Special Relativity. Thus, the cases when VSR does not imply Special Relativity appear as a convenient arena to test violations of Lorentz invariance. Remarkably, Bogoslovsky [12] had already studied the most general transformations which preserve the massless wave equation and he found the invariant metric:

$$L_{\text{Bog}} = \langle \cdot, \cdot \rangle_1^{(1-b)} (\beta \otimes \beta)^b, \quad (12)$$

where β is a $\langle \cdot, \cdot \rangle_1$ -lightlike dual vector and $0 \leq b < 1$, a constant³¹.

Remark 6.2. (1) When L_{Bog} is restricted to the future causal cone \mathcal{C}_0 of $\langle \cdot, \cdot \rangle_1$, then it becomes a Lorentz-Minkowski norm, up to the requirement of differentiability at the lightlike vectors, that is, L_{Bog} is an *improper* Lorentz-Minkowski norm according to Note 4.12. Indeed, L_{Bog} is not smooth at \mathcal{C}_0 , but it trivially satisfies the properties (i) and (ii) of that note as, in this case, the A -anisotropic Chern connection of L_{Bog} is the affine connection of the Euclidean space.

(2) Recall that the restriction of L_{Bog} to the causal \mathcal{C}_0 -vectors is natural not only because of the physical reasons discussed in the previous sections, but also because the vectors where L_{Bog} vanishes include the $\langle \cdot, \cdot \rangle_1$ -spacelike

³¹ β would correspond with the direction of propagation of the wave, b with a parameter for a conformal transformation of $\langle \cdot, \cdot \rangle_1$ which preserves the wave equation and L_{Bog} with a Finsler metric invariant by this transformation, see also [13] for further information.

ones in the kernel of β , and these vectors do not seem to admit any natural interpretation as directions of light rays.

As a generalization of VSR for curved spaces, *General Very Special Relativity (GVSR)* drops the invariance of VSR by translations. This was introduced by Gibbons et al. [32], who pointed out the Finslerian character of GVSR. Relevant examples of Lorentz-Finsler metrics in VSR and GVSR have been recently found, see [29, 30] and references therein.

Remark 6.3. A natural generalization of Bogoslovski metric to GVSR is obtained by regarding $\langle \cdot, \cdot \rangle_1$ and β as a Lorentz metric and arbitrary 1-form on a manifold M . Fuster et al. [28] even consider the generalization obtained by multiplying the latter by a homogeneous factor type $(c + m\beta^2 / \langle \cdot, \cdot \rangle_1)^p$, where $c, m, p \in \mathbb{R}$. Among this type of metrics, they found the Berwald spacetimes non-affinely equivalent to a Lorentz one cited in Remark 5.3.

6.1.2. *Smoothability at the cone and birefringence.* By starting at our previous study of Bogoslovsky metric, we can go further in the issue of the differentiability of the Lorentz-Finsler metrics at the cone, by comparing our approach with the one introduced by Pfeifer and Wohlfart (PW) [73, §A], which has been modified sometimes [37, 39].

These authors considered a definition of Lorentz-Finsler spacetime and metric which permits degenerate directions. This definition becomes consistent with our notion of improper Lorentz-Finsler metric in Note 4.12 and the conditions (i) and (ii) therein. Essentially, PW considers, instead of a Lorentz-Finsler metric L as above, a function L_r which is r -homogeneous for some $r \geq 2$, and they relax the non-degeneracy of the fundamental tensor g allowing a set of zero-measure where it degenerates. Remarkably, the smoothness of L_r does not imply the smoothness of the two-homogeneous function $L = L_r^{2/r}$ along the cone \mathcal{C} . Nevertheless, the A -anisotropic connection (which is well defined on a dense set of timelike vectors) can be then extended to the lightlike ones (see [73, Th. 2]). In this case, $L = L_r^{2/r}$ lies under our definition of improper Lorentz-Finsler metric with a connection extendible to \mathcal{C} .

However, for most choices of b , Bogoslovsky metric (12) is an example which does not lie under PW definition, in spite of having a regular cone and a connection extendible to it (indeed, both of them the same as Lorentz-Minkowski spacetime). Nevertheless, they remain under the definition in the variants [37, 39] and they are always improper Lorentz-Finsler in the sense of Note 4.12, which seems to provide a suitable geometric framework for these cases. Indeed, let us analyze a generalization of Bogoslovsky metrics from norms to arbitrary manifolds considered in [30]. Let $L_{\text{Bog}} = g(\cdot, \cdot)^{(1-b)} (\beta \otimes \beta)^b$, where g is a (time-oriented) Lorentzian metric and β a 1-form in a manifold M ; notice that, whenever β remains g -causal, the future cone \mathcal{C} of g agrees with the lightlike vectors for L_{Bog} and this metric is well-defined on all the g -causal vectors. Let $r = 1/(1-b)$ and $L_{\text{Bog}}^r = g(\cdot, \cdot) (\beta \otimes \beta)^m$, with

$m = b/(1 - b)$. Then,

$$\begin{aligned} g_v^{L_{\text{Bog}}^r}(u, w) &= \beta(v)^m g(u, w) + m\beta(v)^{m-1}(g(v, u)\beta(w) + g(v, w)\beta(u)) \\ &\quad + \frac{1}{2}m(m-1)g(v, v)\beta(v)^{m-2}\beta(u)\beta(w). \end{aligned}$$

It is not difficult to see that $g^{L_{\text{Bog}}^r}$ has the same signature as g when $\beta(v) > 0$ (use for example the criterion in [50, Prop. 4.10]), but it is trivially equal to zero, when $\beta(v) = g(v, v) = 0$ and $1/2 < b < 1$ (observe that in such a case, $m > 1$). As a consequence, if β is always g -timelike, the generalized Bogoslovsky metric is always a Finsler spacetime according to PW definition. When β is g -lightlike, there will be lightlike directions of L_{Bog} which do not satisfy the conditions of PW, no matter if the connection is extendible to the (regular, Lorentzian) cone \mathcal{C} or not; however, they will be improper Lorentz-Finsler metrics and satisfy also the definitions in [37, 39].

An issue beyond the lack of smoothness is birefringence. This phenomenon occurs in some crystals and it is described by using two cones, each one with a Lorentz or Lorentz-Finsler metric. It is related with the dispersion of the light with different wavelengths in the crystal. Some authors have pointed out the possibility that these dispersions occur also as a constitutive element of the spacetime [56, 74].

One way to describe the lightrays when there is birefringence is by using the product of two Lorentz metrics $L = \sqrt{L_1 L_2}$. Essentially, the lightrays are described then by the lightlike geodesics of this product; indeed, when one of the metrics L_1 vanishes and the other does not, then a metric anisotropically conformal to L_1 is obtained. However, some additional subtleties appear. For example, when the lightcones $\mathcal{C}_1, \mathcal{C}_2$ of the metrics are one inside the other, say $\mathcal{C}_1 < \mathcal{C}_2$, this product is an improper Lorentz-Finsler spacetime on the domain A_1 determined by the interior cone \mathcal{C}_1 (see [50, Appendix A.5]). Notice, however, that the situation would be more complex when the position of the cones is arbitrary. Assuming that the intersection $A_1 \cap A_2$ is non-empty at every $p \in M$, then each $(A_1)_p \cap (A_2)_p$ is convex. However, its boundary may have non-smooth directions and L would become an improper Lorentz-Finsler metric.

Under our viewpoint, the existence of different light cones may be a worthy possibility (see the discussions around Def. 2.8). However, in principle, our mathematical framework would consider separately the cones. Indeed, a possible way to describe phenomena related to the dispersion of light would be to introduce a space $\bar{M} = M \times \mathbb{R}^+$ with an extra dimension representing the refractive index n . Then, a Lorentz-Finsler metric L_n would appear for each n and the different cone structures \mathcal{C}_n on $TM \times \{n\}$ would project on TM . The birefringent model would correspond with an effective description of polarization by using two refractive index, that is, the projection on M of a limit case on \bar{M} where only two values of n would become relevant.

6.2. Anisotropic speed of light. In subsection 3.3, the possibility of a pointwise variation of c was discussed for Lorentz metrics. As explained there, an additional element to the metric structure (such as a pointwise




FIGURE 2. The directions $u_1, u_2 \in S_v$ may have $g_v(u_1, u_1) \neq g_v(u_2, u_2)$ and, so, the observer v could conclude $c_v(u_1) \neq c_v(u_2)$, i.e., the speed of light depends on the direction.

measurement of the fine structure constant α) was germane. Next we will consider some different possibilities for the measurement of a varying speed of light (VSL) proper of the Lorentz-Finsler case.

The underlying reason of the difficulty to measure a VSL in the Lorentzian case relied on the fact that the Levi-Civita parallel transport is a conformal transformation (indeed, an isometry), thus, mapping always affinely lightlike cones into lightlike cones. A first possibility in the Finslerian case is:

(VSL1) Lightlike cones at different points may be non-affinely equivalent³². Clearly, this should be an indicator of the existence of different speeds of light at different points. Anyway, at the end such a possibility would be possible because a Lorentz-Finsler metric L provides a breaking of Lorentz symmetry at each point. This would turn out in the existence of anisotropies of the speed of light emitted from a single event p in different directions. So, let us focus on this possibility, which includes Lorentz-Minkowski norms in affine spaces.

(VSL2) At an event $p \in M$, a single observer $v \in \Sigma_p$ finds distinct speeds of light at different directions at its *rest space* ($T_v \Sigma_p$ endowed with g_v).

At least from a purely geometric viewpoint, this could happen as follows. The cone C_p will intersect the rest space $T_v \Sigma_p$ at some strongly convex $(n-2)$ -hypersurface S_v , say, *the sky observed by*³³ v , see Fig. 2. Then, for $u \in S_v$ the value of $c_v(u) := \sqrt{g_v(u, u)}$ can be regarded as a u -dependent speed of light measured by v (namely, the spacelike length covered by the light in the direction of u in a unit of time).

³² From a mathematical viewpoint, the property that lightcones are affinely diffeomorphic is a Berwald-type property. Recall that one of the characterizations of Berwald manifolds in the class of the Finsler ones is the existence of a torsion free derivative operator such that the parallel translations with respect to it preserve the Finsler norms of tangent vectors [80, Prop. 6]; in particular, the norms at different points are isometric.

³³ Equally, the rest space and the sky could be regarded as the hyperplane $T_v^0 \Sigma$ parallel to $T_v \Sigma$ through the origin $0 \in T_p M$ and the projection S_v^0 of S_v along the direction v_p into $T_v^0 \Sigma$, respectively. This is a usual identification in General Relativity, [75].




FIGURE 3. In the plane π spanned by the observers v, v' , the tangent lines to Σ_p in π , l and l' differ. Then if $u \in S_v$ and $u' \in S_{v'}$, possibly, $g_v(u, u) \neq g_{v'}(u', u')$. So the observers v, v' measure different speeds of light in their common plane π .

Remark 6.4. (1) This u -dependent speed of light appears because of the anisotropies of Σ_p . Thus (in contrast to (VSL1)) it might happen even for a Lorentz-Finsler metric compatible with the cone structure of Lorentz-Minkowski spacetime (or any other Lorentzian manifold).

Indeed, at each $p \in M$, the metric g_v depends on the space of observers Σ_p close to v . So, if Σ_p were the space of observers for the Lorentz-Minkowski metric L , we could perturb it around some $v \in \Sigma_p$ in order to obtain the space of observers Σ'_p of an anisotropically equivalent Lorentz-Finsler metric L' satisfying:

$$v \in \Sigma_p \cap \Sigma'_p, \quad \text{and} \quad T_v \Sigma_p = T_v \Sigma'_p.$$

Then, the skies of v for L and L' are equal but, in general, $g_v \neq g'_v$ and $c_v(u) \neq c'_v(u)$.

(2) It is also worth pointing out that two different observers $v, v' \in \Sigma_p$ will span a single plane $\Pi \subset T_p M$ which can be regarded as a timelike one for both g_v and $g_{v'}$. The intersections of Π with the rest spaces $T_v \Sigma_p, T_{v'} \Sigma_p$ will give two lines l and l' (which are spacelike for g_v and $g_{v'}$, respectively). Even though l and l' are different they would represent the “spacelike direction where the other observers lies”. However, the speed of light in the (consistently oriented) directions of l and l' may differ, that is, $c_v(u) \neq c_{v'}(u')$ for $u \in l$ and $u' \in l'$ (see Fig. 3).

The possibility to measure (VSL2) might be somewhat naïve because: (a) experimental difficulties for the measurements of the involved geometric elements g_v, S_v (or the relation between g_v and $g_{v'}$), might appear, and (b) in the case that \mathcal{C} were compatible with a Lorentzian metric, then one should speak on anisotropies of the space for massive particles (or, eventually, for measurement instruments) rather than for the propagation of light.

Anyway, there is an *anisotropic propagation of the light* in the case of a breaking of the conformal Lorentz symmetry, namely:

(VSL3) At an event $p \in M$, the cone \mathcal{C}_p is not compatible with any Lorentz scalar product.

In principle, this could be measured by using the trajectories of lightrays even in the case of Lorentz-Minkowski norms on an affine space (so that the lightrays are straight lines). Indeed, when L comes from a Lorentz scalar product g , then g_v depends only on p ($g_v \equiv g_p$), S_v becomes a sphere in $T_v\Sigma$ centered at 0 of radius $r = 1$, the second fundamental form σ (with respect to the inner direction) of S_v can be identified with the restriction of g_p/r^2 to S_v and the speed of light is regarded naturally as isotropic. However, in the case of a Lorentz-Minkowski norm L , the second fundamental form σ_u at some $u \in S_v$ may satisfy, for example, $\sigma_u > g_v/g_v(u, u)$ (as quadratic forms on T_uS_v). Then, the vectors of S_v close to u can be regarded as “shorter” than those in the Lorentz metric case, that is:

the speed of light measured by $v \in \Sigma_p$ at the direction $u \in S_v := T_v\Sigma_p \cap C_p$ is bigger (resp. smaller) than the speed of light in the directions close to u when $\sigma_u > g_v/g_v(u, u)$ (resp. $\sigma_u < g_v/g_v(u, u)$).

More precisely, if $\lambda (> 0)$ is the eigenvalue of σ_u in the direction $w \in T_uS_v$ then $1/\lambda$ would rate the increasing of the speed along the direction w .

We emphasize that the previous procedure would allow the observer v to realize that an anisotropy holds either in Σ_p or in C_v . The fact that g_v only depends on the behavior of Σ_p around v prevents to disregard the first case. However:

C_v is compatible with a Lorentz scalar product if and only if S_v is an ellipsoid,

and the latter property can be checked in purely affine terms on $T_v\Sigma$ (namely, it holds when it vanishes the cubic form $C(X, Y, Z) = \nabla_X \sigma^\xi(Y, Z)$ ³⁴ constructed from the second fundamental form σ^ξ and the induced connection ∇ , both for the Blaschke normal ξ , see [70, Theorem II.4.5].

Remark 6.5. The property $\nabla \sigma^\xi \neq 0$ implies the intrinsic anisotropy of the speed of light, but it does not assign an “absolute” speed of light $c_v(u)$ (which would depend on the Lorentz metric L as in (VSL2)). However, one has the possibility to measure variations on the speed of light around each u . The qualitative behavior of such variations rely on the cone structure instead of the metric (compare with Remark 6.4).

6.3. Matter as anisotropy and Quantum Physics. Clearly, a Lorentz-Minkowski norm or properly Lorentz-Finsler metric would appear if some type of anisotropy were detected in physical spacetime (see for example [71] and references therein). However, we emphasize:

The existence of an anisotropy does not mean necessarily a “pre-existing spacelike anisotropy of empty space”. Indeed, the existence of matter induces anisotropies in causal directions, and this might be reflected in the indicatrix of L .

³⁴Observe that the cubic form coincides with the Matsumoto tensor of the pseudo-Minkowski norm having the affine hypersurface as indicatrix up to multiplication by a function (see for example [68] or [51]). The Matsumoto tensor is zero when the pseudo-Finsler metric comes from a scalar product.

This possibility is stressed in our formalism, as L is defined only on causal directions. Even though this idea is quite speculative, let us explain it briefly.

Consider first that an event $p \in M$ is crossed by a particle γ , $\gamma(0) = p$, with mass $m > 0$. In this case, $\gamma'(0)$ selects a privileged direction at p , and this would introduce an anisotropy in the space of observers \mathbf{O}_p (with respect to a background Lorentz metric). This perturbation might be made quantitative in some ways; for example, by introducing a perturbation in the curvature of \mathbf{O}_p around p proportional to m . In the case of having a stress energy tensor T in an initial background Lorentzian metric g , algebraic properties of T (as the energy density or pressure for perfect fluids) might induce the perturbation of \mathbf{O}_p .

These perturbations, even if tiny, might have interest at Planck scale. Indeed, it is commonplace to assume that nonlinear modifications of linear Schrödinger equation might lead to an effective collapse which resolves the measurement problem (see for example [25, §7]). So, the nonlinear framework of Finsler spacetimes opens possibilities in this direction which are worth to be studied further.

Remark 6.6. Recent examples of Finslerian spacetimes, as the model of relativistic kinetic gases in [38], can be understood also under the above viewpoint. Their authors explain that an ensemble of a large number of P individual interacting and gravitating point particles can be described at three levels:

- (1) individual particles,
- (2) description as a kinetic gas, by using a 1-particle distribution function (1PDF), which retains information about velocities, and
- (3) description as a fluid, where velocities at each point are also averaged.

That reference develops the second viewpoint, where a Lorentz-Finsler model emerges. However, one should take into account that, certainly, the individual particle description is the extreme idealization of the gas, as these particles should be quantum objects. So, the Lorentz-Finsler metric might be directly the most natural description as a semi-classical limit.

7. CONCLUSIONS

Along this article, we have obtained goals in the following three directions:

- (1) A revision of the foundations of the theories of non-quantum spacetime from the viewpoint of how space and time are being measured, carried out in three parts.

1a. In the first one (doubly linearized models, §2) the previous approaches in this direction [40, 9] have been sharpened and simplified, and the four compatible models of spacetime have been concisely described. In particular, we have introduced the hypothesis of *apparent temporality*. This hypothesis is enough to obtain the models with no additional hypotheses on, for example, group actions, Theorem 2.4. Moreover, it will yield time-orientability in three of the models (the temporal ones) and it will underlie our definition of Finsler spacetime, where the Lorentz-Finsler metric is defined only on the causal vectors of a single cone structure. The other two parts consider their natural non-linear generalizations.

1b. The first non-linearization §3 is carried out in the spirit of the generalization from Special to General Relativity. In a natural way, the previous four models lead to a signature-changing metric, with Leibnizian structures (and their dual) in the degenerate part and to pointwise variations $c(p)$ of the speed of light which are briefly discussed. It is worth pointing out that, consistently with the discussion at the end of §2, here $c(p)$ appears as the supremum of velocities between observers at each event p ; however, it becomes identifiable with the speed of propagation of the light because it propagates in vacuum (and $c(p)$ is the unique common speed different to 0 measurable by all the observers at p).

1c. Focusing in the Relativistic case, the second non-linearization §4 is obtained just by removing the relativistic quadratic restriction (intrinsic to Lorentzian metrics) on the space of observers. This leads directly to our definition of Finsler spacetime. Its mathematical background and subtleties (including issues on differentiability specific to the Finsler case which will be relevant later) are also introduced concisely.

(2) A critical revision of EPS approach §5 with a triple aim.

2a. The first aim was to examine which EPS assumptions forbid non-relativistic Lorentz-Finsler metrics to emerge, taking into account previous studies [81, 57]. We have found that these assumptions appear neatly at two steps (§5.2.1, 5.2.2) and they have the same origin: they impose certain conditions of C^2 -differentiability at 0 (in each tangent space T_pM) of some geometric quantities which, by its very nature, forbids any anisotropy and, mathematically, leads to the quadratic restriction on the metric (the latter, essentially, by an elementary computation in [86, Proposition 4.1]). Intuitively, this condition can be understood as follows: if one has any element in a vector space depending only on the direction (as the fundamental tensor of a non-Riemannian Finslerian metric or the Christoffel symbols of a non-affine anisotropic connection) then this element cannot be even continuous at 0, as this vector can be regarded as the limit of vectors coming from different directions. Of course, such a condition would not be reasonable from a mathematical viewpoint (it would exclude as non-smooth even all the analytic Finsler metrics) but also from a physical one. Indeed, it would be even preferable to assume directly the isotropy in different directions as a physical assumption, as such an isotropy might be natural in some cases. In contrast, the assumption on C^2 -differentiability at 0 a priori may be misleading and it interferes with the assumption on radar coordinates (which is regarded as involved by many authors, see for example recent [65, footnote 7]). For the sake of completeness, we have also studied the Finslerian examples which are compatible with the EPS axioms (including C^2 differentiability at 0, as in [81]) and discussed at what extent they contradict EPS conclusions, §5.2.5.

2b. The second aim was to compare EPS, as well as the standard chronometric approach, with ours. As an important difference between the philosophies of the previous approaches and ours, our postulates do not involve *the physical objects which will be measured* but *the way how we can measure* physical objects. Indeed, the possibility to make meaningful measurements of the physical spacetime relies on the existence of some mild symmetries

among the observers, so that different measurements (carried out at different events and by different observers at each event) can be compared. As stressed here, such symmetries become then apparent in the observers space \mathbf{O} and, then, allow one to determine some geometries for the physical spacetime. The fact that the exact symmetries of \mathbf{O} in the initial linearized model may be only approximate, leads to General Relativity, modified Special Relativity and the general model of Finsler spacetimes.

2c. As an extra bonus of the previous two aims, EPS approach can be also used to obtain Lorentz-Finsler metrics for the geometry of spacetime. Indeed, removing the criticized hypotheses of C^2 smoothability, any Lorentz-Finsler metric L will be compatible with the two first steps of EPS. The other two steps should justify the uniqueness of L up to an overall factor. These steps would be involved mathematically (indeed, the third one would have been justified for the original EPS approach one only recently [65]). However, as suggested in §5.2.4, only the behavior of clocks would be enough to construct \mathbf{O} and, then, to characterize L . Even though this behavior becomes natural in the chronometric approach rather than in EPS, the main objection of these authors to chronometrics (part (1) in §5.3) would be solved. It is also worth emphasizing that, in this way, our procedure becomes simple and rigorous at all the stages.

(3) A summary of some issues related to Lorentz symmetry breaking discussed from the introduced viewpoint. This includes:

3a. Very Special Relativity and Pfeifer & Wohlfart (PW) definition of Finsler spacetimes §6.1. They are particular cases of Finsler spacetimes with non-smooth lightlike directions (and, so, they do not satisfy properly our definition of Lorentz-Finsler metric. However, they are endowed with a regular cone structure \mathcal{C} and an isotropic connection extendible to \mathcal{C} and, so, most of their relevant properties hold (see Note 4.12). The case of Bogoslovsky metric and its generalization to arbitrary manifolds is studied specifically. Moreover, the way to fit the phenomenon of birefringence in our setting is also discussed.

3b. Three ways to detect the possibility that the speed of light varied with the direction §6.2. The first one would be a pointwise variation which would go beyond the one discussed in General Relativity, which relies on the possibility that a cone structure has cones at different points non-affinely isomorphic. The other two ways focus on the Lorentz symmetry breaking at each point $p \in M$. The first one is a geometric analysis which would detect the anisotropies of the Lorentz-Finsler metric L (and, then, of the measured speed of light) in different situations, namely: when a single observer looks at different spacelike directions (Fig. 2) and when two observers at p compare their spacelike measurements (Fig. 3). Because of these anisotropies of L , the measured speeds of the light might be different even for a cone structure compatible with a quadratic (relativistic) cone. Thus, the other procedure focuses on the specific properties of the cone and would detect its lack of quadraticity.

3c. A justification of Lorentz-Finsler anisotropy. Typically, Finslerian anisotropy is considered as a spacelike anisotropy. Notice, however, that our Lorentz-Finsler metrics are not even defined on spacelike directions. As

extensively argued along the article, Lorentz-Finsler anisotropies appear on the space of observers. So, it is natural to think that they might be associated with the distribution of mass and energy. These might be anisotropic even if one thought that a “background isotropic vacuum” existed. In this vein, a possible link with Quantum Mechanics is suggested and further developments on this issue might be worthy.

Summing up, this paper tries to provide physical grounds and precise mathematical formulations for the development of Lorentz-Finsler geometry and its relativistic applications. It is worth emphasizing that the applications, however, go beyond the relativistic setting. For example, an extra bonus has its roots in analogue gravity [5]. Indeed, the classical non-relativistic problem of Zermelo navigation is better understood by using Lorentz-Finsler metrics and the corresponding Fermat principle [15, 50]. Then, on the one hand, the classical Finslerian/Zermelo viewpoint has applications to spacetimes [14, 47] and, on the other, the Lorentz-Finsler viewpoint has applications for issues such as the propagation of fire spreading, quantum navigation and classical Finsler Geometry [62, 33, 49]. So, Lorentz-Finsler geometry and its applications appears as a fascinating area to be developed further.

ACKNOWLEDGMENTS

The authors warmly acknowledge discussions on this topic with Prof. Volker Perlick (Univ. of Bremen), Christian Pfeifer (Univ. of Tartu) and Nicoletta Voicu (Univ. of Brasov).

MAJ was partially supported by MICINN/FEDER project with reference PGC2018-097046-B-I00 and Fundación Séneca project with reference 19901/GERM/15, Spain, and MS by Spanish MINECO/FEDER project reference MTM2016-78807-C2-1-P. This work is included in the framework of the Programme of Excellence Groups of the Region of Murcia, Spain funded by Fundación Séneca, Science and Technology Agency of the Region of Murcia.

REFERENCES

- [1] A. B. AAZAMI, AND M. A. JAVALOYES. Penrose’s singularity theorem in a Finsler spacetime. *Classical Quantum Gravity*, 33(2):025003, 22, (2016).
- [2] H. AKBAR-ZADEH. Sur les espaces de Finsler à courbures sectionnelles constantes. *Acad. Roy. Belg. Bull. Cl. Sci. (5)*, 74(10):281–322, (1988).
- [3] A. ALBRECHT, AND J. MAGUEIJO. A time varying speed of light as a solution to cosmological puzzles. *Phys. Rev. D* 59(4):043516, (1999).
- [4] D. BAO, S.-S. CHERN, Z. SHEN. *An introduction to Riemann-Finsler geometry*. Graduate Texts in Mathematics, 200. Springer-Verlag, New York, 2000.
- [5] C. BARCELÓ, S. LIBERATI, AND M. VISSER. Analogue Gravity. *Living Rev. Relativ.* 14, 3, (2011). <https://doi.org/10.12942/lrr-2011-3>.
- [6] J.D. BARROW. Cosmologies with varying light speed. *Phys. Rev. D* 59(4):043515, (1999).
- [7] J.D. BARROW, AND J. MAGUEIJO. Varying- α theories and solutions to the Cosmological Problems. *Phys. Lett. B* 443(1-4): 104–110, (1998).
- [8] J. K. BEEM, P. E. EHRLICH, AND K. L. EASLEY. *Global Lorentzian geometry*, volume 202 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker, Inc., New York, Second Edition, 1996.
- [9] A. N. BERNAL, M. P. LÓPEZ, AND M. SÁNCHEZ. Fundamental Units of Length and Time. *Found. Phys.* 32(1):77–108, (2002).

- [10] A. N. BERNAL, AND M. SÁNCHEZ. Leibnizian, Galilean and Newtonian structures of spacetime. *J. Math. Phys.* 44:1129–1149, (2003).
- [11] A. N. BERNAL, AND M. SÁNCHEZ. Un paseo por las geometrías del espaciotiempo en el centenario de la Relatividad General, *Gaceta RSME*, 18(3):521–542, (2015).
- [12] G. BOGOSLOVSKY. A special-relativistic theory of the locally anisotropic space-time. *Il Nuovo Cimento B Series* 40:99–115, (1977).
- [13] G. BOGOSLOVSKY. The rest momentum as an additional property of a massive particle in Finsler space-time. *J. Phys.: Conf. Ser.* 1051:012007, 11, (2018).
- [14] E. CAPONIO, M. A. JAVALOYES, AND M. SÁNCHEZ. On the interplay between Lorentzian causality and Finsler metrics of Randers type. *Rev. Mat. Iberoam.* 27(3):919-952, (2011).
- [15] E. CAPONIO, M. A. JAVALOYES, AND M. SÁNCHEZ. Wind Finslerian structures: from Zermelo’s navigation to the causality of spacetimes. *Arxiv e-prints*, 1407.5494.
- [16] E. CAPONIO, AND G. STANCARONE. Standard static Finsler spacetimes. *Int. J. Geom. Methods Mod. Phys.*, 13(4):1650040, 25, (2016).
- [17] E. CAPONIO, AND G. STANCARONE. On Finsler spacetimes with a timelike Killing vector field. *Classical Quantum Gravity*, 35(8):085007, 28, (2018).
- [18] A.G. COHEN, AND S. L. GLASHOW. Very special relativity. *Phys. Rev. Lett.* 97(2):021601 (2006).
- [19] T. DRAY, G. ELLIS, C. HELLABY, AND C. A. MANOGUE. Gravity and signature change. *Gen. Relativity Gravitation*, 29 (5):591–597, (1997).
- [20] C. DUVAL, G. W. GIBBONS, P. A. HORVATHY, AND P. M. ZHANG. Carroll versus Newton and Galilei: Two dual non-Einsteinian concepts of time. *Classical Quantum Gravity* 31(8):085016, 24, (2014).
- [21] J. EHLERS, F. A. E. PIRANI, AND A. SCHILD. Republication of: The geometry of free fall and light propagation. *Gen. Relativity Gravitation* 44 (6):1587–1609, (2012). (Original article: J. Ehlers, F. A. E. Pirani and A. Schild, in: General Relativity, papers in honour of J. L. Synge. Edited by L. O’Reifeartaigh. Oxford, Clarendon Press 1972, pp. 63–84.)
- [22] G. F. R. ELLIS. Note on Varying Speed of Light Cosmologies. *Gen. Relativity Gravitation* 39 (4):511–520, (2007).
- [23] L. FATIBENE AND M. FRANCAVIGLIA. Weyl Geometries and Timelike Geodesics. *Int. J. Geom. Meth. Mod. Phys.* 9(5):1220006, 9, (2012).
- [24] J. FIGUEROA-O-FARRILL, R. GRASSIE, AND S. PROHAZKA. Geometry and BMS Lie algebras of spatially isotropic homogeneous spacetimes. *J. High Energy Physics* 8:119, 92, (2019).
- [25] F. FINSTER, AND J. KLEINER. Causal Fermion Systems as a Candidate for a Unified Physical Theory. *J. Phys.: Conf. Ser.* 626:012020, (2015).
- [26] J. L. FLORES, J. HERRERA, AND M. SÁNCHEZ. Gromov, Cauchy and Causal Boundaries for Riemannian, Finslerian and Lorentzian Manifolds. *Mem. Amer. Math. Soc.* 226(1064): 76, (2013).
- [27] G. FOLLAND. Weyl structures. *J. Differential Geometry*, 4:145–153, (1970).
- [28] A. FUSTER, S. HEEFER, C. PFEIFER, N. VOICU, On the non metrizable of Berwald Finsler spacetimes. *Arxiv e-prints*: 2003.02300v1.
- [29] A. FUSTER, AND C. PABST. Finsler pp-waves. *Phys. Rev. D* 94(10): 104072, (2016).
- [30] A. FUSTER, C. PABST, AND C. PFEIFER. Berwald spacetimes and very special relativity. *Phys. Rev. D* 98(8): 084062, (2018).
- [31] R.P. GEROCH. Faster than light? In: *Advances in Lorentzian geometry*. (Edited by Matthias Plaue, Alan Rendall and Mike Scherfner.) AMS/IP Studies in Advanced Mathematics, 49. International Press, Somerville, MA, (2011).
- [32] G. W. GIBBONS, J. GOMIS, AND C. N. POPE. General very special relativity is Finsler geometry. *Phys. Rev. D*, 76(8):081701, 5, (2007).
- [33] G. W. GIBBONS. A Spacetime Geometry picture of Forest Fire Spreading and of Quantum Navigation. *ArXiv e-prints*, arXiv:1708.02777 [gr-qc], (2017).
- [34] S. GIELEN, AND D. K. WISE. Lifting general relativity to observer space. *J. Math. Phys.* 54(5):052501, (2013).

- [35] J.B. HARTLE, AND S.W. HAWKING. Wave function of the Universe. *Phys. Rev. D* (3) 28(12):2960–2975, (1983).
- [36] M. HOHMANN. Spacetime and observer space symmetries in the language of Cartan geometry. *J. Math. Phys.* 57(8):082502, 28, (2016).
- [37] M. HOHMANN, C. PFEIFER, AND N. VOICU. Finsler gravity action from variational completion. *Phys. Rev. D* 100: 064035, (2019).
- [38] M. HOHMANN, C. PFEIFER, AND N. VOICU. Relativistic kinetic gases as direct sources of gravity. *Phys. Rev. D* 101: 024062, (2020).
- [39] M. HOHMANN, C. PFEIFER, AND N. VOICU. Cosmological Berwald Spacetimes. Arxiv e-prints: 2003.02299v1.
- [40] W. V. IGNATOWSKY. Einige allgemeine Bemerkungen über das Relativitätsprinzip. *Phys. Zeits.* 11:972–976, (1910). Available at https://de.wikisource.org/wiki/Einige_allgemeine_Bemerkungen_%C3%BCber_das_Relativit%C3%A4tsprinzip
- [41] W. V. IGNATOWSKY. Das Relativitätsprinzip. *Arch. Math. Phys.* 17:1–24, (1911) .
- [42] W. V. IGNATOWSKY. Das Relativitätsprinzip. *Arch. Math. Phys.* 18: 17–40, (1911).
- [43] H. ISHIKAWA. Note on Finslerian relativity. *J. Math. Phys.*, 22(5):995–1004, (1981).
- [44] M. A. JAVALOYES. Anisotropic tensor calculus. *Int. J. Geom. Methods Mod. Phys.*, 16(2):1941001, 26, (2019).
- [45] M. A. JAVALOYES. Curvature computations in Finsler Geometry using a distinguished class of anisotropic connections. *Mediterr. J. Math.* (to be published), arxiv: 1904.07178, (2020).
- [46] M. A. JAVALOYES, AND M. SÁNCHEZ. Finsler metrics and relativistic spacetimes. *Int. J. Geom. Methods Mod. Phys.*, 11(9):1460032, 15, (2014).
- [47] M. A. JAVALOYES, AND M. SÁNCHEZ. Some criteria for wind Riemannian completeness and existence of Cauchy hypersurfaces. In: *Lorentzian geometry and related topics*, 117–151, Springer Proc. Math. Stat., 211, Springer, Cham, (2017).
- [48] M. A. JAVALOYES, AND M. SÁNCHEZ. On the definition and examples of Finsler metrics. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 13(3):813–858, (2014).
- [49] M. A. JAVALOYES, AND M. SÁNCHEZ. Wind Riemannian spaceforms and Randers-Kropina metrics of constant flag curvature. *Eur. J. Math.* 3(4):1225–1244, (2017).
- [50] M. A. JAVALOYES, AND M. SÁNCHEZ. On the definition and examples of cones and Finsler spacetimes. *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM*, 114(1):30, 46, (2020).
- [51] M. A. JAVALOYES AND H. VITÓRIO. Some properties of Zermelo navigation in pseudo-Finsler metrics under an arbitrary wind. *Houston Journal of Mathematics*, 44(4):1147–1179, (2018).
- [52] M. KOSSOWSKI AND M. KRIELE. Signature type change and absolute time in general relativity. *Classical Quantum Gravity* 10, 1157, (1993).
- [53] V. A. KOSTELECKÝ. Riemann-Finsler geometry and Lorentz-violating kinematics. *Phys. Lett. B*, 701(1):137–143, (2011).
- [54] V. A. KOSTELECKÝ, N. RUSSELL, AND R. TSO. Bipartite Riemann-Finsler geometry and Lorentz violation. *Phys. Lett. B*, 716(3-5):470–474, (2012).
- [55] H.P. KÜNZLE. Galilei and Lorentz structures on space-time: comparison of the corresponding geometry and physics. *Ann. Inst. H. Poincaré Sect. A (N.S.)*, 17:337–362, (1972).
- [56] C. LAMMÉRZAHN, AND F. W. HEHL. Riemannian light cone from vanishing birefringence in premetric vacuum electrodynamics. *Phys. Rev.D* 70:105022, (2004).
- [57] C. LAMMÉRZAHN, AND V. PERLICK. Finsler geometry as a model for relativistic gravity. *Int. J. Geom. Methods Mod. Phys.* 15, Supp. 1: 1850166, 20, (2018).
- [58] C. LAEMMÉRZAHN, V. PERLICK, AND W. HASSE. Observable effects in a class of spherically symmetric static Finsler spacetimes. *Phys. Rev. D*, 86:104042, (2012).
- [59] J-M. LÉVY LEBLOND. Une nouvelle limite non-relativiste du groupe de Poincaré, *Ann. Inst. H. Poincaré Sect. A (N. S.)*, tome 3 (1):1–12, (1965).
- [60] LIGO Scientific Collaboration and Virgo Collaboration, Fermi Gamma-ray Burst Monitor, and INTEGRAL. Gravitational Waves and Gamma-Rays from a Binary Neutron Star Merger: GW170817 and GRB 170817A. *Astrophysical J. Lett.* 848:L13, 27, (2017).

- [61] S. LIBERATI. Tests of Lorentz invariance: a 2013 update. *Classical Quantum Gravity* 30(13):133001, 51, (2013).
- [62] S. MARKVORSEN. A Finsler geodesic spray paradigm for wildfire spread modelling. *Nonlinear Anal. Real World Appl.*, 28:208–228, (2016).
- [63] E. MARTÍNEZ, J.F. CARIÑENA, AND W. SARLET. Derivations of differential forms along the tangent bundle projection. *Diff. Geom. Appl.*, 2:17–43, (1992).
- [64] E. MARTÍNEZ, J.F. CARIÑENA, AND W. SARLET. Derivations of differential forms along the tangent bundle projection. Part II. *Diff. Geom. Appl.*, 3:1–29, (1993).
- [65] V. MATVEEV, AND A. SCHOLZ. Light cone and Weyl compatibility of conformal and projective structures. *ArXiv e-prints, arXiv:2001.0149 [math.DG]*, (2020).
- [66] V. MATVEEV, AND A. TRAUTMAN. A criterion for compatibility of conformal and projective structures. *Commun. Math. Phys.* 329:821–825, (2014).
- [67] E. MINGUZZI. An equivalence of Finslerian relativistic theories. *Rep. Math. Phys.*, 77(1):45–55, (2016).
- [68] X. MO, AND L. HUANG. On characterizations of Randers norms in a Minkowski space. *Internat. J. Math.*, 21:523–535, (2010).
- [69] J. MOFFAT. Superluminary Universe: A Possible Solution to the Initial Value Problem in Cosmology. *Int. J. Mod. Phys. D* 2(3):351–366, (1993).
- [70] K. NOMIZU, AND T. SASAKI. *Affine differential geometry*, vol. 111 of Cambridge Tracts in Mathematics, Cambridge University Press, Cambridge, 1994.
- [71] V. PERLICK. Fermat Principle in Finsler Spacetimes. *Gen. Relativity Gravitation*, 38(2): 365–380, (2006).
- [72] J.P. PETIT. An interpretation of cosmological model with variable light velocity *Mod. Phys. Lett. A* 3(16):1527-1532, (1998).
- [73] C. PFEIFER, AND M. WOHLFARTH. Causal structure and electrodynamics on Finsler space- times. *Phys. Rev. D*, 84:044039, (2011).
- [74] C. PFEIFER. Finsler spacetime geometry in Physics. *Int. J. Geom. Methods Mod. Phys.* 16, Supp. 2: 1941004, 18, (2019).
- [75] R. K. SACHS AND H. H. WU. *General relativity for mathematicians*, Springer-Verlag, New York, 1977.
- [76] J. B. SANCHO DE SALAS. Characterization of Levi-Civita and Newton-Cartan connections in dimension 2. *Diff. Geom. Appl.* 68:101583, 15, (2020).
- [77] J. STACHEL. Conformal and projective structures in general relativity. *Gen. Relativity Gravitation* 43:3399–3409, (2011).
- [78] J.L. SYNGE. *Relativity: the special theory*. North Holland, Amsterdam (1960). *Relativity: the general theory*. North Holland, Amsterdam (1964).
- [79] A. SZABÓ. Positive definite Berwald spaces. *Tensor, N.S.* 35:25–39, (1981).
- [80] J SZILASI, RL LOVAS, AND DC KERTÉSZ. Ten ways to Berwald manifolds —and some steps beyond. *ArXiv preprints, arXiv:1106.2223*, (2011).
- [81] R. TAVAKOL, AND N. VAN DEN BERGH. Finsler spaces and the underlying geometry of space-time. *Phys. Lett. A* 112(1-2):23–25, (1985).
- [82] A. TRAUTMAN. The general theory of relativity. *Usp. fiz. Nauk* 89:3–37, (1966). [English translation: *Soviet Phys. Usp.* 9, 319-39 (1966).]
- [83] A. TRAUTMAN. Editorial note to: J. Ehlers, F. A. E. Pirani and A. Schild, The geometry of free fall and light propagation. *Gen. Relativity Gravitation* 44:1581–1586, (2012).
- [84] J.-P. UZAN. Fundamental constants and their variation: Observational status and theoretical motivations. *Reviews of Modern Physics* 75(2):403–455, (2003).
- [85] B. VAKILIA, AND S. JALALZADEHB. Signature transition in Einstein-Cartan cosmology. *Physics Letters B* 726:28–32, (2013).
- [86] F. W. WARNER. The conjugate locus of a Riemannian manifold. *Amer. J. Math.*, 87:575–604, (1965).
- [87] A. WHITE, S. WEINFURTNER, AND M. VISSER. Signature change events: a challenge for quantum gravity? *Classical Quantum Gravity* 27(4):045007, (2010).

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA, FACULTAD DE CIENCIAS,
UNIVERSIDAD DE GRANADA,
CAMPUS FUENTENUEVA S/N,
18071 GRANADA, SPAIN
E-mail address: anbernal@no-gravity.eu

DEPARTAMENTO DE MATEMÁTICAS,
UNIVERSIDAD DE MURCIA,
CAMPUS DE ESPINARDO,
30100 ESPINARDO, MURCIA, SPAIN
E-mail address: majava@um.es

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA, FACULTAD DE CIENCIAS,
UNIVERSIDAD DE GRANADA,
CAMPUS FUENTENUEVA S/N,
18071 GRANADA, SPAIN
E-mail address: sanchezm@ugr.es