

SU(2|1) supersymmetric spinning models of chiral superfields

Stepan Sidorov

*Bogoliubov Laboratory of Theoretical Physics, JINR,
141980 Dubna, Moscow region, Russia*
sidorovstepan88@gmail.com

Abstract

We construct SU(2|1), $d = 1$ supersymmetric models based on the coupling of dynamical and semi-dynamical (spin) multiplets, where the interaction term of both multiplets is defined on the generalized chiral superspace. The dynamical multiplet is defined as a chiral multiplet $(\mathbf{2}, \mathbf{4}, \mathbf{2})$, while the semi-dynamical multiplet is associated with a multiplet $(\mathbf{4}, \mathbf{4}, \mathbf{0})$ of the mirror type.

PACS: 11.30.Pb, 12.60.Jv

Keywords: supersymmetric mechanics, deformation, spin variables

1 Introduction

A new class of systems of $\mathcal{N} = 4$, $d = 1$ supersymmetric quantum mechanics called “Kähler oscillator” was introduced by S. Bellucci and A. Nersessian in [1]. They studied supersymmetric oscillator models on Kähler manifolds with Wess-Zumino (WZ) type terms responsible for the presence of a constant magnetic field. It turned out, the presence of oscillator and WZ terms deforms the standard $\mathcal{N} = 4$ Poincaré supersymmetry to the so called “weak supersymmetry” [2]. In our works [3, 4], we showed that the deformed superalgebra of weak supersymmetry corresponds to $SU(2|1)$ supersymmetry. In these papers we initiated a study of deformed supersymmetric quantum mechanics by employing superfield approach based on the worldline supersymmetry $SU(2|1)$ with a mass dimension deformation parameter m . In the limit $m = 0$, models of the standard $\mathcal{N} = 4$ supersymmetric mechanics are restored. Indeed, within this framework we reproduced the models studied earlier at the component level [1, 2, 5] and obtained the new ones, including those constructed via harmonic superspace [6] (see also [7]).

During the study of $SU(2|1)$ supersymmetric mechanics we revealed few peculiar features. One of them is the generalization of a chiral condition given by [4]

$$(\cos \lambda \bar{\mathcal{D}}_i - \sin \lambda \mathcal{D}_i) \varphi = 0, \quad (1.1)$$

where \mathcal{D}_i and $\bar{\mathcal{D}}_i$ are $SU(2|1)$ covariant derivatives. It describes a new type of the chiral multiplet $(\mathbf{2}, \mathbf{4}, \mathbf{2})$ defined on the generalized chiral superspace and depending on two deformation parameters: λ and m . Exactly this multiplet is a basis for the construction of supersymmetric Kähler oscillator models [1], where the frequency of oscillator and the strength of external magnetic field are identified with

$$\omega = \frac{m \sin 2\lambda}{2}, \quad B = m \cos 2\lambda. \quad (1.2)$$

However, both parameters disappear in the limit $m = 0$, because the rotation parameter λ becomes just an external automorphism parameter of the standard $\mathcal{N} = 4$ Poincaré supersymmetry. One needs to point out that the corresponding Hamiltonian does not commute with supercharges and is identified with internal $U(1)$ generator of $SU(2|1)$. It can be treated as a central charge only when $\lambda = 0$. Recently, particular superintegrable Kähler oscillator models were considered [8].

With the lapse of time we found out other distinguished features of deformed supersymmetric quantum mechanics. An important one concerns $\mathcal{N} = 4$ “mirror multiplets” [9]. The standard $\mathcal{N} = 4$ multiplets have their mirror counterparts characterized by the interchange of two $SU(2)$ groups which form $SO(4)$ automorphism group of the standard $\mathcal{N} = 4$ Poincaré supersymmetry. Since this interchange has no essential impact on Poincaré supersymmetry, $\mathcal{N} = 4$ multiplets and their mirror counterparts are mutually equivalent when dealing with only one multiplet from such a pair. Deformation to $SU(2|1)$ supersymmetry breaks this equivalence, because the first $SU(2)$ group becomes subgroup of $SU(2|1)$ and the second one is broken. It means that $SU(2|1)$ multiplets differ from their mirror counterparts. We showed this difference in details on the example of $(\mathbf{4}, \mathbf{4}, \mathbf{0})$ multiplets [6].

The main goal of the present paper is to employ $SU(2|1)$ superfield approach to spinning models of chiral superfields instead of harmonic ones. The $SU(2|1)$ supersymmetric spinning models of harmonic superfields [7, 10] followed the construction elaborated in

[11, 12], with the dynamical multiplet $(\mathbf{1}, \mathbf{4}, \mathbf{3})$ and the semi-dynamical multiplet $(\mathbf{4}, \mathbf{4}, \mathbf{0})$. The most important property is that both dynamical and semi-dynamical multiplets admit a description in the analytic harmonic superspace [13], like as the $\mathcal{N} = 4$, $d = 1$ gauge multiplet [14]. The latter one allows to introduce the U(1) gauge symmetry transformations. The gauge invariant coupling term is constructed as an integral over the analytic harmonic superspace and provides the interaction of two multiplets. This term is identified with WZ term and involves semi-dynamical spin variables ¹. Extra SU(2) symmetries are defined in terms of these variables, with respect to which physical states of quantum models carry additional spin quantum numbers. Generalization of the construction to matrix superfields [11] give rise to $\mathcal{N} = 4$ supersymmetric extensions of the widely known Calogero system [16] (see also the review [17]).

In this paper we consider the coupling of the dynamical multiplet $(\mathbf{2}, \mathbf{4}, \mathbf{2})$ and the semi-dynamical multiplet $(\mathbf{4}, \mathbf{4}, \mathbf{0})$ of the mirror type. Both multiplets satisfy the generalized chiral condition (1.1) that allows to introduce the interaction term as a superpotential term. We construct SU(2|1) supersymmetric models based on this coupling, undeformed versions of which were studied in [18] ². To compare the results with [18], we consider an example of a model on the pseudo-sphere SU(1, 1)/U(1) (the Lobachevsky space).

2 SU(2|1) chiral superfields

In [4], we considered a special type of SU(2|1) superspace defined for the standard form of the superalgebra $su(2|1)$:

$$\begin{aligned} \{Q^i, \bar{Q}_j\} &= 2mI_j^i + 2\delta_j^i \mathcal{H}, & [I_j^i, I_l^k] &= \delta_j^k I_l^i - \delta_l^i I_j^k, \\ [I_j^i, \bar{Q}_l] &= \frac{1}{2} \delta_j^i \bar{Q}_l - \delta_l^i \bar{Q}_j, & [I_j^i, Q^k] &= \delta_j^k Q^i - \frac{1}{2} \delta_j^i Q^k, \\ [\mathcal{H}, \bar{Q}_l] &= \frac{m}{2} \bar{Q}_l, & [\mathcal{H}, Q^k] &= -\frac{m}{2} Q^k. \end{aligned} \quad (2.1)$$

The bosonic subgroup includes of SU(2) subgroup composed of I_j^i ($i = 1, 2$) and U(1) generator \mathcal{H} associated with Hamiltonian. Factorizing the SU(2) subgroup, the superspace is defined as

$$\frac{\text{SU}(2|1)}{\text{SU}(2)} \sim \frac{\{\mathcal{H}, Q^i, \bar{Q}_j, I_j^i\}}{\{I_j^i\}} = \{t, \theta_i, \bar{\theta}^j\}. \quad (2.2)$$

The odd ϵ^i transformations are given by

$$\begin{aligned} \delta t &= i \left(\epsilon_k \bar{\theta}^k + \bar{\epsilon}^k \theta_k \right), & \delta \theta_i &= \epsilon_i + 2m \bar{\epsilon}^k \theta_k \theta_i, & \delta \bar{\theta}^j &= \bar{\epsilon}^j - 2m \epsilon_k \bar{\theta}^k \bar{\theta}^j, \\ (\theta_i) &= \bar{\theta}^i, & (\epsilon_i) &= \bar{\epsilon}^i. \end{aligned} \quad (2.3)$$

In the limit $m = 0$, the standard “flat” supersymmetry transformations are restored. One can easily pick out the chiral subspace $\{t_L, \theta_i\}$, where

$$t_L = t + i \bar{\theta}^k \theta_k - im \left(\bar{\theta}^k \theta_k \right)^2, \quad \delta t_L = 2i \bar{\epsilon}^k \theta_k, \quad \delta \theta_i = \epsilon_i + 2m \bar{\epsilon}^k \theta_k \theta_i. \quad (2.4)$$

¹Some recent $\mathcal{N} = 4$ deformed and undeformed supersymmetric models with spin variables were studied at the component level in [15].

²In this paper, superfield description for a semi-dynamical multiplet proceeds from the fermionic multiplet $(\mathbf{0}, \mathbf{4}, \mathbf{4})$. One can make the same comment as in the footnote 7 of [19].

We also define the generalized chiral coordinates

$$\hat{t}_L = t + i \bar{\theta}^k \hat{\theta}_k, \quad \hat{\theta}_i = \left(\cos \lambda \theta_i e^{\frac{i}{2} m t} + \sin \lambda \bar{\theta}_i e^{-\frac{i}{2} m t} \right) \left(1 - \frac{m}{2} \bar{\theta}^k \theta_k \right). \quad (2.5)$$

They are closed under the following supersymmetry transformations:

$$\begin{aligned} \delta \hat{\theta}_i &= \cos \lambda \left(\epsilon_i e^{\frac{i}{2} m \hat{t}_L} + m \bar{\epsilon}^k \hat{\theta}_k \hat{\theta}_i e^{-\frac{i}{2} m \hat{t}_L} \right) + \sin \lambda \left(\bar{\epsilon}_i e^{-\frac{i}{2} m \hat{t}_L} + m \epsilon^k \hat{\theta}_k \hat{\theta}_i e^{\frac{i}{2} m \hat{t}_L} \right), \\ \delta \hat{t}_L &= 2i \left(\cos \lambda \bar{\epsilon}^k \hat{\theta}_k e^{-\frac{i}{2} m \hat{t}_L} - \sin \lambda \epsilon^k \hat{\theta}_k e^{\frac{i}{2} m \hat{t}_L} \right). \end{aligned} \quad (2.6)$$

Chiral superfields, defined on the left chiral subspace $\{\hat{t}_L, \hat{\theta}_i\}$, satisfy the chiral condition

$$\bar{\mathcal{D}}_j \varphi(\hat{t}_L, \hat{\theta}) = 0, \quad (2.7)$$

where the covariant derivatives $\bar{\mathcal{D}}_i$ and $\tilde{\mathcal{D}}^i$ are modified as

$$\bar{\mathcal{D}}_i = \cos \lambda \bar{\mathcal{D}}_i - \sin \lambda \mathcal{D}_i, \quad \tilde{\mathcal{D}}^i = \cos \lambda \mathcal{D}^i + \sin \lambda \bar{\mathcal{D}}^i, \quad \lambda \in [0, \pi/2). \quad (2.8)$$

In the basis $\{\hat{t}_L, \hat{\theta}_j, \bar{\theta}^i\}$, the covariant derivative $\bar{\mathcal{D}}_i$ is written as

$$\bar{\mathcal{D}}_j = - \left[1 + \frac{m \cos 2\lambda}{2} \bar{\theta}^k \hat{\theta}_k - \frac{m \sin 2\lambda}{4} \left(\bar{\theta}^k \bar{\theta}_k + \hat{\theta}_k \hat{\theta}^k \right) - \frac{m^2}{16} \hat{\theta}_i \hat{\theta}^i \bar{\theta}^j \bar{\theta}_j \right] \frac{\partial}{\partial \bar{\theta}^j}. \quad (2.9)$$

Here, we ignore the matrix $SU(2)$ generators \tilde{I}_j^i , because the generalized chiral superfields $\bar{\mathcal{D}}_i$ cannot carry any external $SU(2)$ indices³.

The $SU(2|1)$ invariant measure is written as

$$d\zeta = dt d^2\theta d^2\bar{\theta} \left(1 + 2m \bar{\theta}^k \theta_k \right). \quad (2.10)$$

One can write this measure in the basis $\{t, \hat{\theta}_i, \bar{\theta}^j\}$ as

$$d\zeta = dt d^2\hat{\theta} d^2\bar{\theta} \left[1 + m \cos 2\lambda \bar{\theta}^k \hat{\theta}_k - \frac{m \sin 2\lambda}{2} \left(\hat{\theta}_k \hat{\theta}^k + \bar{\theta}^k \bar{\theta}_k \right) \right]. \quad (2.11)$$

The chiral measure $dt_L d^2\hat{\theta}$ is also invariant under the transformations (2.6):

$$d\zeta_L = dt_L d^2\hat{\theta}, \quad \delta(d\zeta_L) = 0. \quad (2.12)$$

2.1 Dynamical multiplet (2,4,2)

The multiplet $(2, 4, 2)$ is described by a complex superfield Φ subjected to the chirality condition [4]

$$\bar{\mathcal{D}}_j \Phi = 0. \quad (2.13)$$

Its general solution reads

$$\Phi(\hat{t}_L, \hat{\theta}) = z + \sqrt{2} \hat{\theta}_k \xi^k + \hat{\theta}_k \hat{\theta}^k B. \quad (2.14)$$

³The explanation is given in Appendix B of [21].

The superfield is not deformed, but transformation properties of its components are still deformed:

$$\begin{aligned}
\delta z &= -\sqrt{2} \cos \lambda \epsilon_k \xi^k e^{\frac{i}{2}mt} - \sqrt{2} \sin \lambda \bar{\epsilon}_k \xi^k e^{-\frac{i}{2}mt}, \\
\delta \xi^k &= \sqrt{2} \bar{\epsilon}^k (i \cos \lambda \dot{z} - \sin \lambda B) e^{-\frac{i}{2}mt} - \sqrt{2} \epsilon^k (i \sin \lambda \dot{z} + \cos \lambda B) e^{\frac{i}{2}mt}, \\
\delta B &= -\sqrt{2} \cos \lambda \bar{\epsilon}_k \left(i \dot{\xi}^k + \frac{m}{2} \xi^k \right) e^{-\frac{i}{2}mt} + \sqrt{2} \sin \lambda \epsilon_k \left(i \dot{\xi}^k - \frac{m}{2} \xi^k \right) e^{\frac{i}{2}mt}. \quad (2.15)
\end{aligned}$$

The deformation parameters m and λ also appear in (2.11).

Superfield kinetic action for the chiral superfield Φ is given by

$$S_{\text{kin.}} = \frac{1}{4} \int d\zeta K(\Phi, \bar{\Phi}). \quad (2.16)$$

where $K(\Phi, \bar{\Phi})$ is a Kähler potential. The component Lagrangian is written as

$$\begin{aligned}
\mathcal{L}_{\text{kin.}} &= g \dot{z} \dot{\bar{z}} + \frac{i}{2} g \left(\xi^i \dot{\xi}_i - \bar{\xi}^i \dot{\bar{\xi}}_i \right) + g B \bar{B} - \frac{B}{2} \partial_z g \bar{\xi}^k \bar{\xi}_k - \frac{\bar{B}}{2} \partial_z g \xi_k \xi^k \\
&+ \frac{i}{2} (\dot{z} \partial_z g - z \partial_z g) \xi^k \bar{\xi}_k + \frac{1}{4} \partial_z \partial_z g \xi_i \xi^i \bar{\xi}^j \bar{\xi}_j - \frac{i}{2} m \cos 2\lambda (\dot{z} \partial_z K - z \partial_z K) \\
&- \frac{m \cos 2\lambda}{2} g \xi^k \bar{\xi}_k - \frac{m \sin 2\lambda}{2} (\bar{B} \partial_z K + B \partial_z K) \\
&+ \frac{m \sin 2\lambda}{4} (\partial_z \partial_z K \xi_i \xi^i + \partial_z \partial_z K \bar{\xi}^j \bar{\xi}_j). \quad (2.17)
\end{aligned}$$

with the metric $g(z, \bar{z}) := \partial_z \partial_{\bar{z}} K(z, \bar{z})$.

2.2 Semi-dynamical mirror multiplet (4,4,0)

In [6], we studied the deformed mirror multiplet $(\mathbf{4}, \mathbf{4}, \mathbf{0})$ in the framework of harmonic superspace. Here we consider its generalization to the chiral superspace (2.5). The generalized mirror multiplet $(\mathbf{4}, \mathbf{4}, \mathbf{0})$ is described by a pair of chiral superfields Y^A ($A = 1, 2$) satisfying the constraints

$$\bar{\mathcal{D}}_i Y^A = 0, \quad \tilde{\mathcal{D}}^i \bar{Y}^A = 0, \quad \tilde{\mathcal{D}}_i Y^A = \bar{\mathcal{D}}_i \bar{Y}^A, \quad \overline{(Y^A)} = \bar{Y}_A. \quad (2.18)$$

Their solution reads

$$Y^A \left(\hat{t}_L, \hat{\theta} \right) = y^A + \sqrt{2} \hat{\theta}_i \psi^{iA} + i \hat{\theta}_k \hat{\theta}^k \dot{y}^A, \quad (2.19)$$

where

$$\overline{(y^A)} = \bar{y}_A, \quad \overline{(\psi^{iA})} = \psi_{iA}. \quad (2.20)$$

The corresponding component field transformations are

$$\begin{aligned}
\delta y^A &= -\sqrt{2} \cos \lambda \epsilon_k \psi^{kA} e^{\frac{i}{2}mt} - \sqrt{2} \sin \lambda \bar{\epsilon}_k \psi^{kA} e^{-\frac{i}{2}mt}, \\
\delta \bar{y}_A &= \sqrt{2} \cos \lambda \bar{\epsilon}^k \psi_{kA} e^{-\frac{i}{2}mt} - \sqrt{2} \sin \lambda \epsilon^k \psi_{kA} e^{\frac{i}{2}mt}, \\
\delta \psi^{kA} &= \sqrt{2} i \bar{\epsilon}^k (\cos \lambda \dot{y}^A - \sin \lambda \dot{\bar{y}}^A) e^{-\frac{i}{2}mt} - \sqrt{2} i \epsilon^k (\cos \lambda \dot{y}^A + \sin \lambda \dot{\bar{y}}^A) e^{\frac{i}{2}mt}. \quad (2.21)
\end{aligned}$$

The superpotential term can be written as

$$S_{\text{pot.}} = \frac{\mu}{2} \int d\zeta_{\text{L}} h(Y^A) + \frac{\mu}{2} \int d\zeta_{\text{R}} h(\bar{Y}_A). \quad (2.22)$$

It yields Wess-Zumino type Lagrangian:

$$S_{\text{pot.}} = \int dt \mathcal{L}_{\text{WZ}}, \quad \mathcal{L}_{\text{WZ}} = \mu \left[i \dot{y}^A \partial_A h(y^A) + \frac{1}{2} \psi^{iA} \psi_i^B \partial_A \partial_B h(y^A) + \text{c.c.} \right]. \quad (2.23)$$

One can consider the function $h(y^A)$ in (2.23) as a holomorphic subset of the solutions found in [6], where the general construction of Wess-Zumino type Lagrangians was given.

The superpotential term (2.23) contains no deformation parameters. So, these terms are invariant as well under the standard ($m = 0$) supersymmetry transformations. Indeed, the superpotential term must be invariant under the transformations closed on the centerless $\mathcal{N} = 4$ super Virasoro algebra [20], which is an infinite dimensional superalgebra possessing both deformed and undeformed superalgebras.

2.3 Gauged mirror multiplet (4,4,0)

By analogy with [7], we can also introduce that the left chiral superfield Y^A is subjected to the local $U(1)$ transformations

$$\begin{aligned} (Y^A)' &= e^{(\Lambda - \bar{\Lambda})\mathcal{I}} Y^A, & \mathcal{I} Y^A &= \frac{1}{2} (\sigma_3)_B^A Y^B, \\ (\bar{Y}^A)' &= e^{(\Lambda - \bar{\Lambda})\mathcal{I}} \bar{Y}^A, & \mathcal{I} \bar{Y}^A &= \frac{1}{2} (\sigma_3)_B^A \bar{Y}^B, \end{aligned} \quad (2.24)$$

where \mathcal{I} is a $U(1)$ generator and $\Lambda = \Lambda(\hat{t}_{\text{L}}, \hat{\theta})$, $\bar{\Lambda} = \bar{\Lambda}(\hat{t}_{\text{R}}, \hat{\bar{\theta}})$. Gauged superfields satisfy the new gauge invariant constraints

$$\left(\tilde{\mathcal{D}}^i + [\tilde{\mathcal{D}}^i, X] \mathcal{I} \right) Y^A = 0, \quad \left(\tilde{\mathcal{D}}^i - [\tilde{\mathcal{D}}^i, X] \mathcal{I} \right) \bar{Y}^A = 0, \quad (2.25)$$

$$\left(\tilde{\mathcal{D}}^i - [\tilde{\mathcal{D}}^i, X] \mathcal{I} \right) Y^A = \left(\tilde{\mathcal{D}}^i + [\tilde{\mathcal{D}}^i, X] \mathcal{I} \right) \bar{Y}^A, \quad (2.26)$$

where the real superfield X is a gauge superfield transforming as

$$X' = X + \Lambda + \bar{\Lambda}. \quad (2.27)$$

From the constraints (2.25) and (2.26), one can obtain that the gauge superfield X is subjected to the additional constraint

$$\tilde{\mathcal{D}}_{(i} \tilde{\mathcal{D}}_{j)} X = 0. \quad (2.28)$$

Note that this condition is invariant with respect to the gauge transformations (2.27). Hence, it is solved by

$$\begin{aligned} X(t, \hat{\theta}, \hat{\bar{\theta}}) &= x + \sqrt{2} \left(\hat{\theta}_k \bar{\chi}^k + \hat{\bar{\theta}}^k \chi_k \right) + 2 \hat{\bar{\theta}}_k \hat{\theta}^k \mathcal{A} + \hat{\theta}_k \hat{\theta}^k D + \hat{\bar{\theta}}^k \hat{\bar{\theta}}_k \bar{D} \\ &+ \sqrt{2} i \hat{\bar{\theta}}^k \hat{\theta}_k \left(\hat{\theta}_i \dot{\chi}^i - \hat{\bar{\theta}}^i \dot{\chi}_i \right) - \frac{1}{4} \hat{\theta}_i \hat{\theta}^i \hat{\bar{\theta}}^j \hat{\bar{\theta}}_j \ddot{x}, \end{aligned} \quad (2.29)$$

where

$$\overline{(x)} = x, \quad \overline{(\mathcal{A})} = \mathcal{A}, \quad \overline{(D)} = \bar{D}, \quad \overline{(\chi^i)} = \bar{\chi}_i. \quad (2.30)$$

This superfield describes the mirror multiplet $(\mathbf{1}, \mathbf{4}, \mathbf{3})$ that differs from the ordinary multiplet $(\mathbf{1}, \mathbf{4}, \mathbf{3})$ [3] because of the deformation. Using the U(1) gauge freedom (2.27), we can choose the WZ gauge:

$$X_{\text{WZ}} = 2\bar{\theta}^k \hat{\theta}_k \mathcal{A}, \quad \mathcal{A}' = \mathcal{A} - \dot{\alpha}. \quad (2.31)$$

Thus, it can be interpreted as a mirror counterpart of the ‘‘topological’’ gauge multiplet [14] described by the harmonic superfield V^{++} in the WZ gauge. One can introduce accompanying chiral superfields

$$\mathcal{V}_{\text{WZ}}(\hat{t}_{\text{L}}, \hat{\theta}) = \hat{\theta}_k \hat{\theta}^k \mathcal{A}, \quad \bar{\mathcal{V}}_{\text{WZ}}(\hat{t}_{\text{R}}, \bar{\theta}) = \bar{\theta}^k \bar{\theta}_k \mathcal{A}, \quad (2.32)$$

satisfying

$$\tilde{D}_i X_{\text{WZ}} = \tilde{\bar{D}}_i \bar{\mathcal{V}}_{\text{WZ}}, \quad \tilde{\bar{D}}_i X_{\text{WZ}} = -\tilde{D}_i \mathcal{V}_{\text{WZ}}. \quad (2.33)$$

The triplet of superfields, given by (2.31) and (2.32), has a harmonic superfield description in the flat superspace $m = 0$ with respect to the second SU'(2) subgroup [9]. One could ascribe SU'(2) indices to this triplet as $\mathcal{V}^{(i'j')}$ and harmonize them. In the deformed supersymmetric mechanics we have no such description, because the second SU'(2) symmetry is broken.

According to the chiral condition (2.25), the superfield solution (2.19) is modified as

$$\begin{aligned} Y^A(t, \hat{\theta}, \bar{\theta}) &= e^{-X\mathcal{I}} Y_{\text{L}}^A(\hat{t}_{\text{L}}, \hat{\theta}), & (Y_{\text{L}}^A)' &= e^{2\Lambda\mathcal{I}} Y_{\text{L}}^A, \\ \bar{Y}^A(t, \hat{\theta}, \bar{\theta}) &= e^{X\mathcal{I}} \bar{Y}_{\text{R}}^A(\hat{t}_{\text{R}}, \bar{\theta}), & (\bar{Y}_{\text{R}}^A)' &= e^{-2\bar{\Lambda}\mathcal{I}} \bar{Y}_{\text{R}}^A. \end{aligned} \quad (2.34)$$

Solving (2.26), the left chiral superfield Y_{L}^A has the θ -expansion

$$Y_{\text{L}}^A(\hat{t}_{\text{L}}, \hat{\theta}) = y^A + \sqrt{2} \hat{\theta}_i \psi^{iA} + i \hat{\theta}_k \hat{\theta}^k \nabla_t \bar{y}^A, \quad (2.35)$$

where

$$\nabla_t = \partial_t + 2i\mathcal{A}\mathcal{I}. \quad (2.36)$$

It is necessary to replace time derivatives by ∇_t in the transformations (2.21). The residual local $\alpha = \alpha(t)$ transformations of component fields must be written as

$$(y^A)' = e^{2i\alpha\mathcal{I}} y^A, \quad (\psi^{iA})' = e^{2i\alpha\mathcal{I}} \psi^{iA}, \quad (\nabla_t \bar{y}^A)' = e^{2i\alpha\mathcal{I}} \nabla_t \bar{y}^A, \quad \mathcal{A}' = \mathcal{A} - \dot{\alpha}. \quad (2.37)$$

Taking into account these transformations and the supersymmetry transformations (2.6), one can check that the real superfield (2.31) transforms as (2.27) with the parameter

$$\Lambda(\hat{t}_{\text{L}}, \hat{\theta}) = i\alpha(\hat{t}_{\text{L}}) + 2\mathcal{A}(\hat{t}_{\text{L}}) \left(\cos \lambda \epsilon^k \hat{\theta}_k e^{-\frac{i}{2} m \hat{t}_{\text{L}}} - \sin \lambda \epsilon^k \hat{\theta}_k e^{\frac{i}{2} m \hat{t}_{\text{L}}} \right). \quad (2.38)$$

The superfield Y^A transforms with the same parameter according to (2.24).

Finally, the superpotential (2.22) must be written in terms of $Y^1 Y^2 \equiv Y_L^1 Y_L^2$, since it is the only gauge invariant object defined on the left chiral subspace. One can define also the invariant F-term

$$S_{\text{FI}} = -\frac{c}{4} \left[\int d\zeta_L \mathcal{V}_{\text{WZ}} + \int d\zeta_R \bar{\mathcal{V}}_{\text{WZ}} \right] \Rightarrow \mathcal{L}_{\text{FI}} = -c \mathcal{A}, \quad c = \text{const}, \quad (2.39)$$

which is a counterpart of Fayet-Iliopoulos term defined on the analytic harmonic superspace [14].

3 Coupling of dynamical and semi-dynamical multiplets

In this section we consider the coupling of dynamical and semi-dynamical multiplets identified with the chiral multiplet $(\mathbf{2}, \mathbf{4}, \mathbf{2})$ and the gauged mirror multiplet $(\mathbf{4}, \mathbf{4}, \mathbf{0})$, respectively.

3.1 Interacting superpotential term

According to (2.24), the simplest interacting superpotential term reads

$$S_{\text{int.}} = \frac{\mu}{2} \int d\zeta_L Y^1 Y^2 f(\Phi) + \frac{\mu}{2} \int d\zeta_R \bar{Y}_1 \bar{Y}_2 \bar{f}(\bar{\Phi}), \quad (3.1)$$

where f is an arbitrary holomorphic function of Φ . The Lagrangian is then given by

$$\begin{aligned} \mathcal{L}_{\text{int.}} = & \mu \left[i (y^1 \nabla \bar{y}_1 - y^2 \nabla \bar{y}_2) f + \psi^{i1} \psi_i^2 f + B y^1 y^2 \partial_z f \right. \\ & \left. + \xi^i (\psi_{i1} y^1 - \psi_{i2} y^2) \partial_z f - \frac{\xi_i \bar{\xi}^i}{2} y^1 y^2 \partial_z \partial_z f + \text{c.c.} \right]. \end{aligned} \quad (3.2)$$

It is straightforward to check that the Lagrangian is invariant under the local U(1) transformations (2.37).

3.2 Total Lagrangian

The total Lagrangian of the interacting model is given by the sum of the kinetic Lagrangian (2.17), the interacting superpotential term (3.2) and Fayet-Iliopoulos term (2.39):

$$\mathcal{L} = \mathcal{L}_{\text{kin.}} + \mathcal{L}_{\text{int.}} + \mathcal{L}_{\text{FI}}. \quad (3.3)$$

Eliminating the auxiliary fields B and ψ^{iA} by their equations of motion and performing the following redefinition

$$\begin{aligned} y^1 &= v (f + \bar{f})^{-\frac{1}{2}}, & \bar{y}_1 &= \bar{v} (f + \bar{f})^{-\frac{1}{2}}, \\ y^2 &= \bar{w} (f + \bar{f})^{-\frac{1}{2}}, & \bar{y}_2 &= w (f + \bar{f})^{-\frac{1}{2}}, \\ \xi^i &= g^{-\frac{1}{2}} \eta^i, & \bar{\xi}_j &= g^{-\frac{1}{2}} \bar{\eta}_j, \end{aligned} \quad (3.4)$$

we obtain the total on-shell Lagrangian (up to full time derivatives) as

$$\begin{aligned}
\mathcal{L} = & g \dot{z}\dot{z} + \frac{i}{2} (\eta^i \dot{\eta}_i - \dot{\eta}^i \bar{\eta}_i) + \frac{i}{2} \mu (v\dot{v} + w\dot{w} - \dot{v}\bar{v} - \dot{w}\bar{w}) \\
& - \frac{i}{2} m \cos 2\lambda (\dot{z} \partial_{\bar{z}} K - \dot{z} \partial_z K) + \frac{i\mu (\dot{z} \partial_{\bar{z}} \bar{f} - \dot{z} \partial_z f)}{2(f + \bar{f})} (v\bar{v} - w\bar{w}) \\
& + \frac{i}{2} (\dot{z} \partial_z g - \dot{z} \partial_{\bar{z}} g) g^{-1} \eta^k \bar{\eta}_k - \frac{m \cos 2\lambda}{2} \eta^k \bar{\eta}_k - \frac{\mu \partial_z f \partial_{\bar{z}} \bar{f}}{(f + \bar{f})^2} (v\bar{v} - w\bar{w}) g^{-1} \eta^k \bar{\eta}_k \\
& - \frac{\mu v\bar{w}}{(f + \bar{f})} \left[\frac{\partial_z \partial_z f}{2} - \frac{\partial_z f \partial_z f}{(f + \bar{f})} - \frac{\partial_z f}{2} g^{-1} \partial_z g \right] g^{-1} \eta_i \eta^i \\
& - \frac{\mu w\bar{v}}{(f + \bar{f})} \left[\frac{\partial_{\bar{z}} \partial_{\bar{z}} \bar{f}}{2} - \frac{\partial_{\bar{z}} \bar{f} \partial_{\bar{z}} \bar{f}}{(f + \bar{f})} - \frac{\partial_{\bar{z}} \bar{f}}{2} g^{-1} \partial_{\bar{z}} g \right] g^{-1} \bar{\eta}^j \bar{\eta}_j \\
& + \frac{m \sin 2\lambda}{4} [(\partial_z \partial_z K - g^{-1} \partial_z K \partial_z g) g^{-1} \eta_i \eta^i + (\partial_{\bar{z}} \partial_{\bar{z}} K - g^{-1} \partial_{\bar{z}} K \partial_{\bar{z}} g) g^{-1} \bar{\eta}^j \bar{\eta}_j] \\
& - g^{-1} \left(\frac{\mu v\bar{w} \partial_z f}{(f + \bar{f})} - \frac{m \sin 2\lambda}{2} \partial_z K \right) \left(\frac{\mu w\bar{v} \partial_{\bar{z}} \bar{f}}{(f + \bar{f})} - \frac{m \sin 2\lambda}{2} \partial_{\bar{z}} K \right) \\
& + \frac{1}{4} (\partial_z \partial_{\bar{z}} g - g^{-1} \partial_z g \partial_{\bar{z}} g) g^{-2} \eta_i \eta^i \bar{\eta}^j \bar{\eta}_j + [\mu (v\bar{v} + w\bar{w}) - c] \mathcal{A}. \tag{3.5}
\end{aligned}$$

Looking at the last term, one concludes that the U(1) gauge field \mathcal{A} plays the role of a Lagrange multiplier enforcing the constraint

$$\mu (v\bar{v} + w\bar{w}) - c = 0. \tag{3.6}$$

3.3 Hamiltonian and Noether charges

Our next step is a formulation of a classical Hamiltonian mechanics. We perform Legendre transformation with the exclusion of the Lagrange multiplier \mathcal{A} . As a result, we obtain the classical Hamiltonian

$$\begin{aligned}
\mathcal{H} = & g^{-1} \left[p_z - \frac{i}{2} m \cos 2\lambda \partial_z K + \frac{i\mu \partial_z f (v\bar{v} - w\bar{w})}{2(f + \bar{f})} + \frac{i}{2} g^{-1} \partial_z g \eta^i \bar{\eta}_i \right] \\
& \times \left[p_{\bar{z}} + \frac{i}{2} m \cos 2\lambda \partial_{\bar{z}} K - \frac{i\mu \partial_{\bar{z}} \bar{f} (v\bar{v} - w\bar{w})}{2(f + \bar{f})} - \frac{i}{2} g^{-1} \partial_{\bar{z}} g \bar{\eta}^j \eta_j \right] \\
& + \frac{m \cos 2\lambda}{2} \eta^k \bar{\eta}_k + \frac{\mu \partial_z f \partial_{\bar{z}} \bar{f}}{(f + \bar{f})^2} (v\bar{v} - w\bar{w}) g^{-1} \eta^k \bar{\eta}_k \\
& + \frac{\mu v\bar{w}}{(f + \bar{f})} \left[\frac{\partial_z \partial_z f}{2} - \frac{\partial_z f \partial_z f}{(f + \bar{f})} - \frac{\partial_z f}{2} g^{-1} \partial_z g \right] g^{-1} \eta_k \eta^k \\
& + \frac{\mu w\bar{v}}{(f + \bar{f})} \left[\frac{\partial_{\bar{z}} \partial_{\bar{z}} \bar{f}}{2} - \frac{\partial_{\bar{z}} \bar{f} \partial_{\bar{z}} \bar{f}}{(f + \bar{f})} - \frac{\partial_{\bar{z}} \bar{f}}{2} g^{-1} \partial_{\bar{z}} g \right] g^{-1} \bar{\eta}^k \eta_k \\
& - \frac{m \sin 2\lambda}{4} [(\partial_z \partial_z K - g^{-1} \partial_z K \partial_z g) g^{-1} \eta_i \eta^i + (\partial_{\bar{z}} \partial_{\bar{z}} K - g^{-1} \partial_{\bar{z}} K \partial_{\bar{z}} g) g^{-1} \bar{\eta}^j \eta_j] \\
& + g^{-1} \left(\frac{\mu v\bar{w} \partial_z f}{(f + \bar{f})} - \frac{m \sin 2\lambda}{2} \partial_z K \right) \left(\frac{\mu w\bar{v} \partial_{\bar{z}} \bar{f}}{(f + \bar{f})} - \frac{m \sin 2\lambda}{2} \partial_{\bar{z}} K \right)
\end{aligned}$$

$$-\frac{1}{4} (\partial_z \partial_{\bar{z}} g - g^{-1} \partial_z g \partial_{\bar{z}} g) g^{-2} \eta_i \eta^i \bar{\eta}^j \bar{\eta}_j, \quad (3.7)$$

and the secondary constraint (3.6). Poisson (Dirac) brackets are imposed as

$$\begin{aligned} \{p_z, z\}_{\text{P.B.}} &= -1, & \{p_{\bar{z}}, \bar{z}\}_{\text{P.B.}} &= -1, & \{\eta^i, \bar{\eta}_j\}_{\text{P.B.}} &= -i\delta_j^i, \\ \{v, \bar{v}\}_{\text{P.B.}} &= i\mu^{-1}, & \{w, \bar{w}\}_{\text{P.B.}} &= i\mu^{-1}. \end{aligned} \quad (3.8)$$

SU(2|1) supercharges are

$$\begin{aligned} Q^i &= \sqrt{2} e^{\frac{i}{2}mt} g^{-\frac{1}{2}} \left\{ \cos \lambda \eta^i \left[p_z - \frac{i}{2} m \partial_z K + \frac{i\mu \partial_z f (v\bar{v} - w\bar{w})}{2(f + \bar{f})} + \frac{i}{2} g^{-1} \partial_z g \eta^k \bar{\eta}_k \right] \right. \\ &\quad \left. - \sin \lambda \bar{\eta}^i \left[p_{\bar{z}} - \frac{i}{2} m \partial_{\bar{z}} K - \frac{i\mu \partial_{\bar{z}} \bar{f} (v\bar{v} - w\bar{w})}{2(f + \bar{f})} - \frac{i}{2} g^{-1} \partial_{\bar{z}} g \eta^j \bar{\eta}_j \right] \right\} \\ &\quad - \frac{\sqrt{2} i \mu e^{\frac{i}{2}mt} g^{-\frac{1}{2}}}{(f + \bar{f})} (\cos \lambda w \bar{v} \partial_z \bar{f} \bar{\eta}^i - \sin \lambda v \bar{w} \partial_z f \eta^i), \\ \bar{Q}_j &= \sqrt{2} e^{-\frac{i}{2}mt} g^{-\frac{1}{2}} \left\{ \cos \lambda \bar{\eta}_j \left[p_{\bar{z}} + \frac{i}{2} m \partial_{\bar{z}} K - \frac{i\mu \partial_{\bar{z}} \bar{f} (v\bar{v} - w\bar{w})}{2(f + \bar{f})} - \frac{i}{2} g^{-1} \partial_{\bar{z}} g \eta^k \bar{\eta}_k \right] \right. \\ &\quad \left. + \sin \lambda \eta_j \left[p_z + \frac{i}{2} m \partial_z K + \frac{i\mu \partial_z f (v\bar{v} - w\bar{w})}{2(f + \bar{f})} + \frac{i}{2} g^{-1} \partial_z g \eta^k \bar{\eta}_k \right] \right\} \\ &\quad - \frac{\sqrt{2} i \mu e^{-\frac{i}{2}mt} g^{-\frac{1}{2}}}{(f + \bar{f})} (\cos \lambda v \bar{w} \partial_z f \eta_j + \sin \lambda w \bar{v} \partial_z \bar{f} \bar{\eta}_j). \end{aligned} \quad (3.9)$$

With respect to the brackets (3.8), they close on the superalgebra $su(2|1)$:

$$\begin{aligned} \{Q^i, \bar{Q}_j\}_{\text{P.B.}} &= -i(2mI_j^i + 2\delta_j^i \mathcal{H}), & \{I_j^i, I_l^k\}_{\text{P.B.}} &= -i(\delta_j^k I_l^i - \delta_l^i I_j^k), \\ \{I_j^i, \bar{Q}_l\}_{\text{P.B.}} &= -i\left(\frac{1}{2}\delta_j^i \bar{Q}_l - \delta_l^i \bar{Q}_j\right), & \{I_j^i, Q^k\}_{\text{P.B.}} &= -i\left(\delta_j^k Q^i - \frac{1}{2}\delta_j^i Q^k\right), \\ \{\mathcal{H}, Q^i\}_{\text{P.B.}} &= \frac{i}{2} m Q^i, & \{\mathcal{H}, \bar{Q}_j\}_{\text{P.B.}} &= -\frac{i}{2} m \bar{Q}_j, \end{aligned} \quad (3.10)$$

where the SU(2) generators are

$$I_j^i = \eta^i \bar{\eta}_j - \frac{\delta_j^i}{2} \eta^k \bar{\eta}_k. \quad (3.11)$$

It should be noted that the Hamiltonian (3.7) can always be redefined as a central charge when $\lambda = 0$ (see the example in the next Section).

There is another SU(2) subgroup that is generated by spin variables as

$$S_3 = \frac{\mu}{2} (v\bar{v} - w\bar{w}), \quad S_+ = \mu v\bar{w}, \quad S_- = \mu w\bar{v}. \quad (3.12)$$

Indeed, one can check these generators form the $su(2)$ algebra:

$$\{S_3, S_{\pm}\}_{\text{P.B.}} = \mp i S_{\pm}, \quad \{S_+, S_-\}_{\text{P.B.}} = -2i S_3. \quad (3.13)$$

Since the Hamiltonian (3.7) contains these SU(2) generators, it commutes only with the quadratic Casimir operator

$$C_{\text{SU}(2)} = S_+ S_- + (S_3)^2. \quad (3.14)$$

Taking into account (3.6), the Casimir operator is determined by the constant

$$C_{\text{SU}(2)} = \frac{c^2}{4}. \quad (3.15)$$

Its quantum counterpart (up to the ordering ambiguity) is given by ($c \approx 2s$)

$$C_{\text{SU}(2)} = s(s+1), \quad (3.16)$$

where s is a spin of the quantum system.

4 Model on SU(1,1)/U(1)

Let us consider the model on the coset space SU(1,1)/U(1), that is in fact a deformation of the model studied in [18]. It is given by the Kähler potential

$$K = -\frac{1}{\gamma} \log(1 - \gamma z \bar{z}), \quad \gamma > 0, \quad (4.1)$$

that leads to the metric

$$g = \frac{1}{(1 - \gamma z \bar{z})^2}. \quad (4.2)$$

To eliminate the terms $\sim \mu v \bar{w} \eta_k \bar{\eta}^k$ in (3.7), we choose the holomorphic function $f(z)$ as ⁴

$$f = \frac{1 + \sqrt{\gamma} z}{1 - \sqrt{\gamma} z}. \quad (4.3)$$

To simplify the notations, we also perform a redefinition in the phase space that preserves the brackets (3.8):

$$\begin{aligned} v &\rightarrow (1 - \sqrt{\gamma} z)^{1/2} (1 - \sqrt{\gamma} \bar{z})^{-1/2} v, & \bar{v} &\rightarrow (1 - \sqrt{\gamma} z)^{-1/2} (1 - \sqrt{\gamma} \bar{z})^{1/2} \bar{v}, \\ \bar{w} &\rightarrow (1 - \sqrt{\gamma} z)^{1/2} (1 - \sqrt{\gamma} \bar{z})^{-1/2} \bar{w}, & w &\rightarrow (1 - \sqrt{\gamma} z)^{-1/2} (1 - \sqrt{\gamma} \bar{z})^{1/2} w, \\ p_z &\rightarrow p_z - \frac{i\sqrt{\gamma}\mu(v\bar{v} - w\bar{w})}{2(1 - \sqrt{\gamma}z)}, & p_{\bar{z}} &\rightarrow p_{\bar{z}} + \frac{i\sqrt{\gamma}\mu(v\bar{v} - w\bar{w})}{2(1 - \sqrt{\gamma}\bar{z})}. \end{aligned} \quad (4.4)$$

Thereby, the Hamiltonian reads

$$\begin{aligned} \mathcal{H} &= (1 - \gamma z \bar{z})^2 \left[\bar{\pi} - \frac{i(1 - \cos 2\lambda) m z}{2(1 - \gamma z \bar{z})} \right] \left[\pi + \frac{i(1 - \cos 2\lambda) m \bar{z}}{2(1 - \gamma z \bar{z})} \right] + \frac{m \cos 2\lambda}{2} \eta^k \bar{\eta}_k \\ &+ \gamma \mu (v\bar{v} - w\bar{w}) \eta^k \bar{\eta}_k + \frac{\gamma m \sin 2\lambda}{2} \left(\bar{z}^2 \eta_k \eta^k + z^2 \bar{\eta}^k \bar{\eta}_k \right) - \frac{\gamma}{2} \eta_i \eta^i \bar{\eta}^j \bar{\eta}_j \\ &+ \left(\sqrt{\gamma} \mu v \bar{w} - \frac{m \sin 2\lambda}{2} \bar{z} \right) \left(\sqrt{\gamma} \mu w \bar{v} - \frac{m \sin 2\lambda}{2} z \right), \end{aligned} \quad (4.5)$$

⁴In the limit $\gamma = 0$ to the flat space, the coupling with the spin multiplet disappears since the holomorphic function (4.3) becomes trivial: $f = 1$.

where

$$\begin{aligned}\pi &= p_z - \frac{i\bar{z}}{1-\gamma z\bar{z}} \left[\frac{m}{2} + \frac{\gamma\mu}{2} (v\bar{v} - w\bar{w}) - \gamma\eta^i\bar{\eta}_i \right], \\ \bar{\pi} &= p_{\bar{z}} + \frac{iz}{1-\gamma z\bar{z}} \left[\frac{m}{2} + \frac{\gamma\mu}{2} (v\bar{v} - w\bar{w}) - \gamma\eta^i\bar{\eta}_i \right].\end{aligned}\quad (4.6)$$

Supercharges are written in the form

$$\begin{aligned}Q^i &= \sqrt{2} e^{\frac{i}{2}mt} (1-\gamma z\bar{z}) \left[\cos\lambda \eta^i \pi - \sin\lambda \bar{\eta}^i \left(\bar{\pi} - \frac{imz}{1-\gamma z\bar{z}} \right) \right] \\ &\quad - \sqrt{2\gamma} e^{\frac{i}{2}mt} i\mu (\cos\lambda w\bar{v} \bar{\eta}^i - \sin\lambda v\bar{w} \eta^i), \\ \bar{Q}_j &= \sqrt{2} e^{-\frac{i}{2}mt} (1-\gamma z\bar{z}) \left[\cos\lambda \bar{\eta}_j \bar{\pi} + \sin\lambda \eta_j \left(\pi + \frac{im\bar{z}}{1-\gamma z\bar{z}} \right) \right] \\ &\quad - \sqrt{2\gamma} e^{-\frac{i}{2}mt} i\mu (\cos\lambda v\bar{w} \eta_j + \sin\lambda w\bar{v} \bar{\eta}_j).\end{aligned}\quad (4.7)$$

SU(2) generators are given by the same expression (3.11).

Limit $\lambda = 0$. In distinct from [18], the Hamiltonian (4.5) contains terms ($\sim m \sin 2\lambda$) breaking the target space symmetry SU(1,1). For instance, the corresponding bosonic Hamiltonian is characterized by a term such as the oscillator term $m^2 \sin^2 2\lambda z\bar{z}/4$. We can define only the remaining U(1) generator J_3 ,

$$J_3 = i(\bar{z}p_{\bar{z}} - zp_z) + \frac{\mu}{2}(v\bar{v} - w\bar{w}) - \eta^k \bar{\eta}_k, \quad (4.8)$$

that commutes with all SU(2|1) generators. Thus, to restore the symmetry SU(1,1) we take the limit $\lambda = 0$. In this limit, we can introduce an external U(1) generator F given by the expression

$$F = \frac{1}{2} \left[\eta^k \bar{\eta}_k - \mu(v\bar{v} - w\bar{w}) \right], \quad (4.9)$$

that rotates the relevant supercharges for $\lambda = 0$ as

$$\{F, Q^i|_{\lambda=0}\}_{\text{P.B.}} = -\frac{i}{2} Q^i|_{\lambda=0}, \quad \{F, \bar{Q}^j|_{\lambda=0}\}_{\text{P.B.}} = \frac{i}{2} \bar{Q}^j|_{\lambda=0}. \quad (4.10)$$

Hence, we can define a central charge generator of the extended superalgebra $su(2|1) \oplus u(1)$ as

$$H = \mathcal{H}|_{\lambda=0} + mF. \quad (4.11)$$

and take it as a new Hamiltonian of the system, while the generator F becomes an internal U(1) generator of the extended superalgebra.

Deformed SU(1,1) symmetry generators are written as

$$\begin{aligned}J_3 &= i(\bar{z}p_{\bar{z}} - zp_z) + \frac{\mu}{2}(v\bar{v} - w\bar{w}) - \eta^k \bar{\eta}_k, \\ J_+ &= p_{\bar{z}} - \gamma z^2 p_z - iz \left[\frac{m}{2} + \frac{\gamma\mu}{2} (v\bar{v} - w\bar{w}) - \gamma\eta^k \bar{\eta}_k \right], \\ J_- &= p_z - \gamma \bar{z}^2 p_{\bar{z}} + i\bar{z} \left[\frac{m}{2} + \frac{\gamma\mu}{2} (v\bar{v} - w\bar{w}) - \gamma\eta^k \bar{\eta}_k \right].\end{aligned}\quad (4.12)$$

Indeed, they commute with all $SU(2|1)$ generators including F , and form the central extension of the $su(1,1)$ algebra:

$$\{J_3, J_{\pm}\}_{\text{P.B.}} = \pm iJ_{\pm}, \quad \{J_+, J_-\}_{\text{P.B.}} = -2i\gamma J_3 - im, \quad (4.13)$$

where the deformation parameter m is a central charge. Defining its Casimir operator

$$C_{\text{SU}(1,1)} = J_+ J_- - \gamma \left(J_3 + \frac{m}{2\gamma} \right)^2, \quad (4.14)$$

we obtain the following expression for the Hamiltonian (4.11):

$$H = \gamma C_{\text{SU}(2)} + C_{\text{SU}(1,1)} + \frac{m^2}{4\gamma}. \quad (4.15)$$

In the limit $m = 0$, we reproduce the model studied in [18].

In the limit $\gamma = 0$, the generators J_{\pm} become magnetic translation generators forming the Heisenberg algebra with the external $U(1)$ automorphism group generator J_3 :

$$\{J, J_{\pm}\}_{\text{P.B.}} = \pm iJ_{\pm}, \quad \{J_+, J_-\}_{\text{P.B.}} = -im, \quad \{J_3, J_{\pm}\}_{\text{P.B.}} = \pm iJ_{\pm}. \quad (4.16)$$

Spin variables drop out of the model leaving only the dynamical multiplet $(\mathbf{2}, \mathbf{4}, \mathbf{2})$ [3].

5 Conclusions

In this paper, we proposed new models of single-particle $SU(2|1)$ supersymmetric mechanics with the use of dynamical, semi-dynamical and gauge multiplets. As an alternative of harmonic superspace, we exploited the generalized chiral superspace (2.5). The generalized chiral multiplet $(\mathbf{2}, \mathbf{4}, \mathbf{2})$ was introduced in [4]. For the mirror multiplet $(\mathbf{4}, \mathbf{4}, \mathbf{0})$, we defined the gauge invariant constraints (2.25) and (2.26). As turned out, the gauge superfield X , bonded with these constraints, satisfies (2.28) and describes the mirror multiplet $(\mathbf{1}, \mathbf{4}, \mathbf{3})$. As an example, we considered in details the deformed model on the pseudo-sphere $SU(1,1)/U(1)$.

The most natural question is whether it is possible to consider superconformal models. During our study of superconformal models of the trigonometric type [21], we suggested a simple criteria for construction of superconformal models. With this experience we tried to obtain superconformal symmetry for considered models, but it is impossible at least for a single-particle model. For example, if we take two chiral dynamical multiplets then suitable superconformal models can possibly be obtained. This may be a good challenge for future research.

It would be interesting to study the mirror counterpart of the multiplet $(\mathbf{1}, \mathbf{4}, \mathbf{3})$ [3]. Another interesting problem is the application of the gauging procedure [7] to the deformed multiplets $(\mathbf{4}, \mathbf{4}, \mathbf{0})$ [6]. It can give rise to general actions of the ordinary and mirror multiplets $(\mathbf{3}, \mathbf{4}, \mathbf{1})$ ⁵.

Acknowledgments

The author is indebted to Evgeny Ivanov for interest and valuable comments. The author also thanks Sergey Fedoruk and Armen Nersessian for useful discussions. This research was supported by the Russian Science Foundation Grant No 16-12-10306.

⁵ $SU(2|1)$ constraints for the ordinary one were solved in Appendix C of [22].

References

- [1] S. Bellucci, A. Nersessian, *(Super)oscillator on $CP^{**}N$ and constant magnetic field*, Phys. Rev. **D67** (2003) 065013, [Erratum-ibid. **D71** (2005) 089901], arXiv:hep-th/0211070;
S. Bellucci, A. Nersessian, *Supersymmetric Kähler oscillator in a constant magnetic field*, Proc. of the Workshop “Supersymmetries and Quantum Symmetries (SQS’03)”, p. 379-384, Dubna, Russia, July 24-29, 2003, arXiv:hep-th/0401232.
- [2] A.V. Smilga, *Weak supersymmetry*, Phys. Lett. **B585** (2004) 173, arXiv:hep-th/0311023.
- [3] E. Ivanov, S. Sidorov, *Deformed supersymmetric mechanics*, Class. Quant. Grav. **31** (2014) 075013, arXiv:1307.7690 [hep-th].
- [4] E. Ivanov, S. Sidorov, *Super Kähler oscillator from $SU(2|1)$ superspace*, J. Phys. **A47** (2014) 292002, arXiv:1312.6821 [hep-th].
- [5] C. Romelsberger, *Counting chiral primaries in $N=1$, $d=4$ superconformal field theories*, Nucl. Phys. **B747** (2006) 329, arXiv:hep-th/0510060;
C. Romelsberger, *Calculating the superconformal index and Seiberg duality*, arXiv:0707.3702 [hep-th].
- [6] E. Ivanov, S. Sidorov, *$SU(2|1)$ mechanics and harmonic superspace*, Class. Quant. Grav. **33** (2016) 055001, arXiv:1507.00987 [hep-th].
- [7] S. Fedoruk, E. Ivanov, *Gauged spinning models with deformed supersymmetry*, JHEP **1611** (2016) 103, arXiv:1610.04202 [hep-th].
- [8] H. Shmavonyan, *\mathbb{C}^N -Smorodinsky-Winternitz system in a constant magnetic field*, Phys. Lett. **A383** (2019) 1223-1228, arXiv:1804.03721 [hep-th];
E. Ivanov, A. Nersessian, H. Shmavonyan, *$\mathbb{C}\mathbb{P}^N$ -Rosochatius system, superintegrability, supersymmetry*, Phys. Rev. **D99** (2019) 085007, arXiv:1812.00930 [hep-th];
E. Ivanov, A. Nersessian, S. Sidorov, H. Shmavonyan, *Symmetries of deformed supersymmetric mechanics on Kähler manifolds*, Phys. Rev. **D101** (2020) 025003, arXiv:1911.06290 [hep-th].
- [9] E. Ivanov, J. Niederle, *Bi-harmonic superspace for $N=4$ mechanics*, Phys. Rev. **D80** (2009) 065027, arXiv:0905.3770 [hep-th].
- [10] S. Fedoruk, E. Ivanov, S. Sidorov, *Deformed supersymmetric quantum mechanics with spin variables*, JHEP **1801** (2018) 132, arXiv:1710.02130;
S. Fedoruk, E. Ivanov, O. Lechtenfeld, S. Sidorov, *Quantum $SU(2|1)$ supersymmetric Calogero-Moser spinning systems*, JHEP **1804** (2018) 043, arXiv:1801.00206 [hep-th].
- [11] S. Fedoruk, E. Ivanov, O. Lechtenfeld, *Supersymmetric Calogero models by gauging*, Phys. Rev. **D79** (2009) 105015, arXiv:0812.4276 [hep-th].
- [12] S. Fedoruk, E. Ivanov, O. Lechtenfeld, *$OSp(4|2)$ Superconformal Mechanics*, JHEP **0908** (2009) 081, arXiv:0905.4951 [hep-th];

- S. Fedoruk, E. Ivanov, O. Lechtenfeld, *New $D(2, 1, \alpha)$ Mechanics with Spin Variables*, JHEP **1004** (2010) 129, [arXiv:0912.3508 \[hep-th\]](#).
- [13] E. Ivanov, O. Lechtenfeld, *$N=4$ supersymmetric mechanics in harmonic superspace*, JHEP **0309** (2003) 073, [arXiv:hep-th/0307111](#).
- [14] F. Delduc, E. Ivanov, *Gauging $N=4$ supersymmetric mechanics*, Nucl. Phys. **B753** (2006) 211-241, [arXiv:hep-th/0605211](#);
F. Delduc, E. Ivanov, *Gauging $N=4$ supersymmetric mechanics II: $(1, 4, 3)$ models from the $(4, 4, 0)$ ones*, Nucl. Phys. **B770** (2007) 179-205, [arXiv:hep-th/0611247](#).
- [15] N. Kozyrev, S. Krivonos, O. Lechtenfeld, A. Sutulin, *$SU(2|1)$ supersymmetric mechanics on curved spaces*, JHEP **1805** (2018) 175, [arXiv:1712.09898 \[hep-th\]](#);
S. Krivonos, O. Lechtenfeld, A. Sutulin, *N -extended supersymmetric Calogero models*, Phys. Lett. **B784** (2018) 137-141, [arXiv:1804.10825 \[hep-th\]](#);
A. Galajinsky, O. Lechtenfeld, *Spinning extensions of $D(2, 1, \alpha)$ superconformal mechanics*, JHEP **1903** (2019) 069, [arXiv:1902.06851 \[hep-th\]](#).
- [16] A. Polychronakos, *Integrable systems from gauged matrix models*, Phys. Lett. **B266** (1991) 29-34.
- [17] A. Polychronakos, *Physics and Mathematics of Calogero particles*, J. Phys. **A39** (2006) 12793-12846, [arXiv:hep-th/0607033](#).
- [18] S. Bellucci, N. Kozyrev, S. Krivonos, A. Sutulin, *$N=4$ chiral supermultiplet interacting with a magnetic field*, Phys. Rev. **D85** (2012) 065024, [arXiv:1112.0763 \[hep-th\]](#).
- [19] E. Ivanov, *Harmonic Superfields in $\mathcal{N} = 4$ Supersymmetric Quantum Mechanics*, SIGMA **7** (2011) 015, [arXiv:1102.2288 \[hep-th\]](#).
- [20] N.L. Holanda, F. Toppan, *Four types of (super)conformal mechanics: D -module reps and invariant actions*, J. Math. Phys. **55** (2014) 061703, [arXiv:1402.7298 \[hep-th\]](#).
- [21] E. Ivanov, S. Sidorov, F. Toppan, *Superconformal mechanics in $SU(2|1)$ superspace*, Phys. Rev. **D91** (2015) 085032, [arXiv:1501.05622 \[hep-th\]](#).
- [22] E. Ivanov, O. Lechtenfeld, S. Sidorov, *$SU(2|2)$ supersymmetric mechanics*, JHEP **1611** (2016) 031, [arXiv:1609.00490 \[hep-th\]](#).