

A REMARK ON A CONJECTURE OF ERDŐS AND STRAUS

YOUSSEF LAZAR

ABSTRACT. The aim of this note is to show that given a positive integer $n \geq 5$, the positive integral solutions of the diophantine equation $4/n = 1/x + 1/y + 1/z$ cannot have solution such that x and y are coprime with $xy < \sqrt{z/2}$. The proof uses the continued fraction expansion of $4/n$.

1. THE RESULT

Given a positive integer n we are interested in the following diophantine equation

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}. \quad (1)$$

where x, y, z are three positive integers to be found. The question of finding positive integer solutions (x, y, z) for this equation was raised by P. Erdős and E. Straus (see e.g. [1]). This problem has attracted a lot of attention and despite many efforts it is still widely open. For an account of the main contributions on this subject we refer the reader to [2]. Usual reductions allow us to assume that n is an odd prime number, thus one can assume that n is an odd prime number p . We are looking for integral primitive solutions $(x, y) \in \mathbb{N}$ of the diophantine problem, when z is a fixed positive number

$$\left| \frac{4}{p} - \frac{x+y}{xy} \right| = \frac{1}{z}. \quad (2)$$

Our main result shows that a solution (x, y, z) of (2) cannot have its largest coordinate, say z , too far away from the two other coordinates x and y , provided $\gcd(x, y) = 1$. This gives a certain piece of information regarding the localization of the lattice points which are solution of the problem. Let us denote by $f(p)$ the number of triples of solutions to (1) with $n = p$ or equivalent to (2). Recently Elsholtz and Tao gave in [2] precise bounds for averages of the form $\sum_{p \leq N} f(p)$, the study of the counting function $f(p)$ reduces to count solutions triples of two different types

- *Type I* solutions of (1) where p divides x and $\gcd(p, yz) = 1$.
- *Type II* solutions of (1) where p divides y, z and $\gcd(p, x) = 1$.

We denote by $f_I(p)$ (resp. $f_{II}(p)$) the number of solutions of (1) of type I (resp. of type II). It was stressed in the same work that for any odd prime number one has the relation

$$f(p) = 3f_I(p) + 3f_{II}(p).$$

We introduce a new type which we call the type III which are the solutions of (1) where $xy < \sqrt{z/2}$ and $\gcd(x, y) = 1$. Analogously we denote by $f_{III}(p)$ the number of solutions of (1) of type III. Our main result is that for any prime number greater than 3, we have $f_{III}(p) = 0$, more precisely

Theorem 1.1. *Given an arbitrary prime number $p \geq 5$, there are no triple of positive integers (x, y, z) which is solution of (2) in the range $xy < \sqrt{z/2}$ and with $\gcd(x, y) = 1$.*

Proof of the Theorem. Suppose we fix an arbitrarily large integer $z_0 > 0$ in the range $xy < \sqrt{z/2}$ and let us try to solve the following diophantine equation with $(x, y) \in \mathbb{N}^2$,

$$\left| \frac{4}{p} - \frac{x+y}{xy} \right| = \frac{1}{z_0}. \quad (3)$$

In particular the equation in (3) gives rise to the following inequality

$$\left| \frac{4}{p} - \frac{x+y}{xy} \right| < \frac{1}{2(xy)^2}. \quad (4)$$

The conclusion of the theorem will follow from the fact that such x and y would never exist. Since we assume that $\gcd(x, y) = 1$ then we have $\gcd(x+y, xy) = 1$. We need the following classical result of the theory of continued fraction which can be found for instance in [3] (Theorem 19).

Lemma 1.2. *Let m, n be two positive integers and suppose $\gcd(r, s) = 1$. If*

$$\left| \frac{m}{n} - \frac{r}{s} \right| < \frac{1}{2s^2}.$$

Then $\frac{r}{s}$ is one of the convergent of $\frac{m}{n}$.

By Lemma 1.2 we infer from (4) that the rational number $\frac{x+y}{xy}$ must be one of the convergents of $4/p$. If we write the continued fraction expansion of $4/p = [0; a_1, \dots, a_l]$, we can say that there exists some $1 \leq k \leq l$ such that

$$\frac{x+y}{xy} = c_k\left(\frac{4}{p}\right) = [0; a_1, \dots, a_k] = \frac{p_k}{q_k}.$$

Since x, y play a symmetric role, we can assume that $x \leq y$. Our fractions are reduced thus we deduce that we might have $x+y = p_k$ and $xy = q_k$. The fact that such x and y might exist relies on the solvability in \mathbb{N} of the following quadratic equation

$$X^2 - p_k X + q_k = 0.$$

The discriminant $D_k = p_k^2 - 4q_k$ cannot vanish otherwise p_k and q_k will fail to be coprime. If $D_k = p_k^2 - 4q_k > 0$, then a couple (x, y) of rational solutions of the equation

$$\frac{4}{p} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z_0} \quad (5)$$

is given by

$$x_k = \frac{p_k - \sqrt{D_k}}{2} \quad \text{and} \quad y_k = \frac{p_k + \sqrt{D_k}}{2}.$$

Necessarily $z_0 = \frac{p}{4 - pc_k}$. Indeed,

$$\frac{4}{p} - \frac{1}{x_k} - \frac{1}{y_k} - \frac{1}{z_0} = \frac{4}{p} - c_k - \frac{4 - pc_k}{p} = 0.$$

Regarding x_k and y_k , both they are in the quadratic field $\mathbb{Q}[\sqrt{D_k}]$, so these are not necessarily integers. In order to obtain integral solutions we are forced to assume that D_k has a square root which is an odd integer. In others words, $D_k = a^2$ where a is an odd integer. In this case, we obtain that a triple of solutions (x_k, y_k, z_0) which is given by

$$x_k = \frac{p_k - a}{2} \quad y_k = \frac{p_k + a}{2}.$$

Note that necessarily

$$z_0 = \left| \frac{1}{4/p - c_k} \right| = \frac{1}{r_k}$$

where $r_k = |4/p - c_k|$ is the k -th error term in the continued fraction approximation. It is well known (see e.g.) that

$$r_k = \frac{1}{q_k(x_{k+1} + q_{k-1})}$$

where

$$x_{k+1} = [a_{k+1}; a_{k+2}, \dots, a_l].$$

Hence, the only possible triples of solutions of (3) in the range given above with a fixed z_0 must take the following form

$$x_k = \frac{p_k - a}{2} \quad y_k = \frac{p_k + a}{2} \quad \text{and} \quad z_0 = q_k(x_{k+1}q_k + q_{k-1})$$

where $\frac{p_k}{q_k}$ is one of the convergents of the continued fraction expansion $4/p = [a_1; a_2, \dots, a_l]$ and provided $p_k^2 - 4q_k = a^2$ with a being an odd integer. We will show that the latter condition can never be fulfilled.

To proceed we take advantage from the fact that the convergents of $4/p$ can only assume specific values which are given in the following lemma,

Lemma 1.3. *For any prime number $p \geq 5$ set*

$$a_1 = \begin{cases} \frac{p-1}{4} & \text{if } p \equiv 1 \pmod{4} \\ \frac{p-3}{4} & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

We have two cases,

- (a) *if $p \equiv 1 \pmod{4}$, then $\frac{4}{p} = [0; a_1, 4]$ and the convergents are $\{0, \frac{4}{p-1}, \frac{4}{p}\}$.*
- (b) *If $p \equiv 3 \pmod{4}$, then $\frac{4}{p} = [0; a_1, 1, 3]$ and the convergents are $\{0, \frac{4}{p-3}, \frac{4}{p+1}, \frac{4}{p}\}$.*

Proof. The continued fraction of a rational number is entirely determine by the eucliden algorithm between 4 and p .

(a) Suppose $p \equiv 1 \pmod{4}$, we perform to the division algorithm

$$\begin{aligned} 4 &= p(0) + 4 \\ p &= 4(a_1) + 1 \\ 4 &= 1(4) + 0. \end{aligned}$$

Here $a_1 = \lfloor \frac{p}{4} \rfloor = \frac{p-1}{4}$, thus $\frac{4}{p} = [0; a_1, 4]$. The successive convergents are given by $c_0 = 0$, $c_1 = \frac{1}{a_1} = \frac{4}{p-1}$ and $c_2 = \frac{1}{a_1 + \frac{1}{4}} = \frac{4}{p}$.

(b) Suppose $p \equiv 3 \pmod{4}$, the division algorithm again shows that

$$\begin{aligned} 4 &= p(0) + 4 \\ p &= 4(a_1) + 3 \\ 4 &= 3(1) + 1 \\ 3 &= 1(3) + 0. \end{aligned}$$

Therefore $\frac{4}{p} = [0; a_1, 1, 3]$. The corresponding convergents are $c_0 = 0$, $c_1 = \frac{4}{p-3}$,
 $c_2 = \frac{1}{a_1 + 1} = \frac{1}{\frac{p-3}{4} + 1} = \frac{4}{p+1}$ and $c_3 = \frac{1}{a_1 + \frac{1}{1+\frac{1}{3}}} = \frac{1}{a_1 + \frac{3}{4}} = \frac{4}{p}$. This proves the Lemma.

We are ready to conclude. In both case, Lemma 1.3 shows that all the non-trivial convergents of $4/p$ (i.e. other than 0 and $4/p$) are egyptian fractions, in particular $p_1 = 1$ (case (a)) and $p_1 = p_2 = 1$ (case (b)). It follows that the $X^2 - p_k X + q_k = 0$ is not solvable in \mathbb{N} since $D_k = 1 - 4q_k < 0$ in all the cases. Hence in the range $xy < \sqrt{z/2}$ there are no solution of (2) with $\gcd(x, y) = 1$. This finishes the proof of Theorem 1.1.

2. CONCLUDING REMARKS

Let p be a prime number, and let us introduce the subset

$$E_p := \{(x, y, z) \in \mathbb{N}^3 \mid 4/p = 1/x + 1/y + 1/z\}.$$

Then p is solution of (2) if and only if $E_p \neq \emptyset$. So the validity of the Erdos-Straus conjecture amounts to prove that $E_p \neq \emptyset$ for every prime p . The Theorem 1.1 tells us that for any prime $p \geq 5$ the set

$$R_p := E_p \cap \{(x, y, z) \in \mathbb{N}^3 \mid \gcd(x, y) = 1, xy < \sqrt{z/2}\}$$

is empty, in other words $f_{III}(p) = 0$. Now suppose we are looking for solutions of (1) with $0 < x, y, z \leq N$, so we have a total number of possibilities equal to $(N-1)^3$ from which we have to remove the elements of $\{(x, y, z) \in [1, N]^3 \mid \gcd(x, y) = 1, xy < \sqrt{z/2}\}$. Let us denote by A_N this subset and put $a_N = |A_N|$. An estimate of a_N can be obtained. Indeed by slicing we get,

$$a_N = \sum_{1 \leq z \leq N} \left| \left\{ 1 \leq x, y \leq \lfloor \sqrt{z/2} \rfloor \mid \gcd(x, y) = 1, xy < \sqrt{z/2} \right\} \right|.$$

The inner sum counts the number of primitive lattice points under the hyperbola of equation $y = \sqrt{z/2} x^{-1}$. Thus we can write

$$a_N = \sum_{1 \leq z \leq N} \sum_{x \leq \lfloor \sqrt{z/2} \rfloor} \sum_{xy \leq \lfloor \sqrt{z/2} \rfloor} \varphi(y).$$

We need the following estimate which can be found in [5] (Thm 3.4) as $X \rightarrow \infty$

$$\sum_{n \leq X} \varphi(n) = \frac{6}{\pi^2} X^2 + O(X \ln X).$$

This asymptotics leads us to

$$a_N = \frac{3}{\pi^2} \sum_{1 \leq z \leq N} \sum_{x \leq \lfloor \sqrt{z/2} \rfloor} \frac{z}{x^2} + O\left(\frac{\sqrt{z}}{2x} \left(\ln \frac{z^{1/2}}{x}\right)\right).$$

Using crude bounds we get

$$\frac{3}{\pi^2} N(1 + o(1)) \ll a_N \ll \frac{1}{2\sqrt{2}\pi^2} N^{5/2} + O(N^{3/2} \ln N).$$

As a conclusion, the previous discussion shows that the number of lattices points to be discarded from $E_p \cap [1, N]^3$ is at an order of magnitude of at least N and at most $N^{5/2}$ lattices points inside the cube $[1, N]^3$.

Further questions. It would be interesting to find an analog of the main result for $5/p$ instead of $4/p$, which is also a conjecture due to Sierpinski. More generally, one can also try to check what is happening for fractions of the form a/p where a is a integer which does not divide p . The continued fraction expansion will depend on the class of $a \pmod p$ and in this case Lemma 1.3 might be less trivial than the $4/p$ -case.

REFERENCES

- [1] P. Erdős, *Az $1/x_1 + 1/x_2 + \dots + 1/x_n = a/b$ egyenlet egész szanu megoldasairól*, Mat. Lapok **1** (1950), 192-210h.
- [2] C. Elsholtz and T. Tao, *Counting the number of solutions of the Erdos-Straus equation on unit fractions*, J. Aust. Math. Soc. **94** (2013), 50-105.
- [3] A. Ya. Khinchin, *Continued Fractions*, Dover, **100** (1997).
- [4] D. Li, *On the equation $4/n = 1/x + 1/y + 1/z$* , J. Number Theory, **13** (1981), 485-494.
- [5] G. Tenenbaum, *Introduction to Analytic and Probabilistic Number Theory*, Graduate Texts in Mathematics, 3rd Ed. Amer. Math. Soc. **163** (2015).

E-mail address: ylazar77@gmail.com