## A surprise in the updated list of stellar perturbers of long period comets motion.

Rita Wysoczańska\* and Piotr A. Dybczyński\*\* and Magdalena Polińska\*\*\*

Astronomical Observatory Institute, Faculty of Physics, A. Mickiewicz University, Słoneczna 36, Poznań, Poland

November 6, 2021

#### ABSTRACT

Context. The second Gaia data release (Gaia DR2) provided us with the precise five-parameter astrometry for 1.3 billion of sources. As stars passing close to the Solar System are thought to be responsible for influencing the dynamical history of long period comets, we update and extend the list of stars that could potentially perturb motion of these comets.

Aims. We aim to announce a publicly available database containing an up to date list of stars and stellar systems potentially perturbing long period comets motion. We add new objects and revise previously published lists. Special emphasis was placed on stellar systems.

Methods. Using the astrometry, preferably from Gaia DR2, augmented with data from other sources, we calculate spatial positions and velocities for each star. To filter studied objects on the basis of their minimal heliocentric distances we numerically integrate motion of stars under the Galactic potential and their mutual interactions.

Results. We announce the updated list of stellar perturbers of cometary motion, including masses of perturbers along with the publicly available database interface. These data are ready to be used with the observed long period comets orbits to study an individual influence of a whole sample of perturbers, or specific stars, on a dynamical past or future of a specific comet. 147 new perturbers were

Conclusions. We demonstrate that a new set of potential perturbers constitute an important tool in studies of cometary dynamics. The usage of our data significantly changes results of the past and future comet motion analysis. We point out a puzzling object in our list, star ALS 9243. The Gaia DR2 astrometry suggests a very close encounter of this star with the Sun however, its astrophysical parameters result in a completely different current distance of ALS 9243 and its large mass.

Movember 6, 2021 **DODUCED Context**. The second *Gaia* data release (*Gaia* DR2) provided us we As stars passing close to the Solar System are thought to be response wighted and extend the list of stars that could potentially perturb *Aims*. We aim to announce a publicly available database containing long period comets motion. We add new objects and revise previou Discussion on masses estimation is included. **Methods**. Using the astrometry, preferably from Gaia DR2, augm and velocities for each star. To filter studied objects on the basis motion of stars under the Galactic potential and their mutual intera *Results*. We announce the updated list of stellar perturbers of cover available database interface. These data are ready to be used with the usage of our data significantly changes results of the past and units, star ALS 9243. The *Gaia* DR2 astrometry suggests a very parameters result in a completely different current distance of ALS. **Key words**. Astronomical databases – Stars: general objects appeared and they significantly change results of dynamical history studies of many LPCs. This research is highly advanced and a detailed paper on LPCs past and future dynamics under the influence of nearby stars and the overall Galactic potential is in preparation.

In this paper we describe in detail the updated version of the list of potential stellar perturbers introduced in the abovementioned paper. We also announce a simple publicly available database interface to access this list.

It should be stressed that our aim is not to determine and study minimal heliocentric distances of passing stars. We use these nominal minimal distances only as a filtering tool while composing the list. For that reason we do not perform any error budget analysis for these values, as well as for other parameters of the encounters. The uncertainty estimations should be performed individually for each particular star - comet interaction, for example in a similar way as in Wysoczańska et al. (2020). All data necessary to perform such an analysis are included in the announced database.

In Sect. 2 we describe sources used while selecting potentially interesting stars. In Sect. 3 methods applied to single stars and problems concerning estimations of their masses are discussed as perturber mass is crucial in examining mutual interactions between a star and a comet. An in-depth analysis of properties of a single star ALS 9243, a new, puzzling but potentially strong stellar perturber is presented in Sect. 4. Sect. 5 and 6 focus on multiple systems, the most interesting and troublesome systems are described. In Sect. 7 a brief description of a public database where our results are presented is given. In Sect. 8 we include several examples on how the new list of stellar perturbers changes the results of cometary dynamics studies. We also present the dependence of these results on the assumed mass of ALS 9243. We conclude in Sect. 9 with a short discussion on main results, issues encountered and prospect for the future.

e-mail: rita.wysoczanska@amu.edu.pl

e-mail: dybol@amu.edu.pl

<sup>\*\*\*</sup> e-mail: polinska@amu.edu.pl

#### 2. Compiling the list of perturbers

Using modern data on stellar positions and kinematics we decided to check again all stars mentioned in several published papers on stellar encounters with the Solar System for their minimal distances from the Sun. Our initial list of stars consists of (these sources partially overlap):

- 156 stars (with the proximity threshold (PT) of 5 pc) from García-Sánchez et al. (2001), based on HIPPARCOS catalogue (ESA 1997),
- 46 stars (PT=2.5 pc) listed in Dybczyński (2006), based on ARIHIP catalogue (Wielen et al. 2001),
- 142 stars (PT=5 pc) analyzed in Jiménez-Torres et al. (2011), based on HIPPARCOS catalogue (ESA 1997)
- 90 stars or stellar systems (PT=3.5 pc) Dybczyński & Królikowska (2015), based on XHIP catalogue (Anderson & Francis 2012),
- 40 stars (PT=2 pc) Dybczyński & Berski (2015), based on HIP2 catalogue (van Leeuwen 2007),
- 42 stars (PT=2 pc) found by Bailer-Jones (2015a), based on HIPPARCOS (ESA 1997), HIP2 (van Leeuwen 2007) and Tycho-2 (Høg et al. 2000) catalogues,
- 166 stars (PT=10 pc) listed by Bailer-Jones (2018), based on *Gaia* DR1 (Gaia Collaboration et al. 2016) and TGAS (Lindegren et al. 2016) catalogues,
- 3379 stars for PT=10 pc listed in Bailer-Jones et al. (2018a), based on *Gaia* DR2 catalogue (Gaia Collaboration et al. 2018),
- 3865 potential stellar perturbers (PT=10 pc) found by us in a subset of *Gaia* DR2 containing over 7 million stars with measured radial velocities,
- 3441 objects (PT=10 pc) selected from all stars with known radial velocity and parallax found in the SIMBAD database in October 2018 (over 2.2 million stars were checked).

Stars from the last two sources were pre-selected according to the linear motion approximation. Then we concatenate the last three sources, numerically integrate each star under the Galactic potential, obtain its minimal heliocentric distance and exclude all stars passing farther than 4 pc from the Sun. This refinement (i.e. applying PT=4 pc) left, in the combined list of the last three sources mentioned above, only 487 stars selected from *Gaia* DR2 and 522 stars selected from the SIMBAD database (these two sets partially overlap). Stars mentioned earlier in at least one of the listed papers, even if new data were adopted and their minimal heliocentric distances increased drastically, were kept for the record. Finally we obtained a list of 820 unique perturbers with 147 new objects which for the first time are identified as potential perturbers.

#### 2.1. Models and methods

While we use the minimal Sun – star distance only as a filtering tool for completing our list of potential LPCs motion perturbers, it might be important for the reader to know how we calculate this value. For all stars and stellar systems the final, nominal smallest heliocentric distance is obtained by a numerical integration of equations of motion formulated in a rectangular Galactocentric frame. We use a reference frame transformation parameters and the Galactic position and velocity of the Sun described in Dybczyński & Berski (2015).

The main difference between our approach and the one presented in the mentioned paper is that, where possible, while calculating rectangular coordinates of stars, we adopted the distance estimates presented in Bailer-Jones et al. (2018b). This way estimated distances were collected for 742 single stars and 85 components of multiple systems. For the stars not included in *Gaia* DR2 we relied on the formula 3 proposed in Bailer-Jones (2015b).

As it concerns a model of the Galactic potential, we used the one proposed by Irrgang et al. (2013).

At first, after completing the preliminary list of 820 objects, we numerically integrated pairs: the Sun with a star or stellar system under the Galactic potential to the past or future depending on the radial velocity sign and recorded the minimal Sun – star distance. After rejecting all perturbers with this distance greater than 4 pc (this leaves 643 objects) we repeated the calculation as a single numerical integration now having 644 interacting bodies (all perturbers plus the Sun).

It should be stressed here, that finding the minimal Sun – star distance serves here only as an approximate tool for filtering potential perturbers. These perturbers are intended to be used in dedicated studies of cometary dynamics, where a detailed analysis of the uncertainties should be carefully done taking into account both stellar and cometary data errors. An example of such an analysis one can found in Wysoczańska et al. (2020).

Readers interested just in Sun – star encounters and their uncertainties should consult other papers, for example Dybczyński & Berski (2015) and Bailer-Jones et al. (2018a).

#### 3. Single stars

Single stars were proceeded in a standard way, described in Section 2.1. In a great majority of cases we use the astrometry from *Gaia* DR2 together with the radial velocity from the same catalogue, if available. When a star is absent in this source we use the SIMBAD and VIZIER databases to find the most appropriate data. Having positions, proper motions, distance estimates, and radial velocities, we calculate rectangular components of the spatial position and velocity. Since we are collecting potential perturbers of the long period comets motion, we need one more parameter - a mass of the perturber.

#### 3.1. Masses

To complete the list of stellar perturbers it was necessary to obtain stellar masses. As *Gaia* DR2 do not provide us with masses of stars, we had to search for them in other sources.

We were unable to find a catalogue or literature source of masses which would cover all stars in question, therefore, our choice was to gather as many different sources of stellar mass estimates as possible, even if, for particular stars, these sources overlap. This approach facilitated a verification of mass estimates and showed whether there is a compliance between different sources and methods.

Below, we describe specific sources and methods which allowed us to obtain stellar mass estimates with clear indication how many masses were collected with each of these methods.

405 stars from our list have their mass estimates presented in Bailer-Jones et al. (2018a).

572 masses of single stars can be found in Anders et al. (2019). These two above-mentioned sources contain only masses of stars included in *Gaia* DR2.

We also performed our own estimations. For M dwarfs we used a formula from Benedict et al. (2016) which allows us to

estimate masses M as a function of the absolute brightness  $M_K$ 

$$M = C_0 + C_1(M_K - x_0) + C_2(M_K - x_0)^2 + C_3(M_K - x_0)^3 + C_4(M_K - x_0)^4,$$
(1)

were polynomial coefficients are  $C_0 = 0.2311$ ,  $C_1 = -0.1352$ ,  $C_2 = 0.0400$ ,  $C_3 = 0.0038$ ,  $C_4 = -0.0032$  and the magnitude offset equals  $x_0 = 7.5$ . We use the K-band because it better agrees with the model, as stated in Benedict et al. (2016). Using this method we were able to obtain masses for 74 single stars from our list of perturbers.

Masses of main sequence dwarfs with known effective temperatures were estimated by us utilising formulas from Eker et al. (2018):

- for ultra low masses in range  $0.179 < M/M_{\odot} \le 0.45$  we use the following formula

$$\log(L) = 2.028(135)\log(M) - 0.976(070)$$
(2)

– for very low masses in range  $0.45 < M/M_{\odot} \le 0.72$ 

 $\log(L) = 4.572(319)\log(M) - 0.102(076)$ (3)

- for low masses in range  $0.72 < M/M_{\odot} \le 1.05$ 

 $\log(L) = 5.743(413)\log(M) - 0.007(026) \tag{4}$ 

- for intermediate masses in range  $1.05 < M/M_{\odot} \le 2.40$ 

 $\log(L) = 4.329(087)\log(M) + 0.010(019)$ <sup>(5)</sup>

- for high masses in range  $2.40 < M/M_{\odot} \le 7$ 

$$\log(L) = 3.967(143)\log(M) + 0.093(083) \tag{6}$$

- for very high masses in range  $7 < M/M_{\odot} \le 31$ 

$$\log(L) = 2.865(155)\log(M) + 1.105(176). \tag{7}$$

For each star its mass was calculated with all the abovementioned formula and then checked for which formula the estimated mass falls within its range of validity. Thanks to these formulas we managed to obtain 402 stellar masses which were further verified whether they meet the conditions stated in Table 5 from Eker et al. (2018). After the verification we ended up with 310 masses obtained with this method.

Formulas from Eker et al. (2018) have to be used in conjunction with formulas from (Andrae et al. 2018) which allow to calculate a radius and a luminosity of the star when only an effective temperature is given.

Additionally, utilising the luminosity in J band, 473 masses were obtained from the tables created by Pecaut & Mamajek (2013). Using the same tables 445 masses were gathered basing on the luminosity in  $K_s$  band and 495 masses when luminosity in V band was used. Each time, when possible, it was checked whether the effective temperature matches the calculated mass.

Because *Gaia* DR2 provides us with luminosities in *G* band it was necessary to convert them into other bands. The following formulas from *Gaia* DR2 documentation Gaia Collaboration et al. (2018) were used:

- to convert into V band

$$G - V = -0.01760 - 0.006860(G_{BP} - G_{RP}) - 0.1732(G_{BP} - G_{RP})^2$$
(8)

- to convert into J band

$$G - J = -0.01883 + 1.394(G_{BP} - G_{RP}) - 0.07893(G_{BP} - G_{RP})^2$$
(9)

- to convert into  $K_s$  band

$$G - K_s = -0.01885 + 2.092(G_{BP} - G_{RP}) - 0.1345(G_{BP} - G_{RP})^2.$$
(10)

For 702 single stars their mass estimates were directly obtained from Pecaut & Mamajek (2013) tables using only an effective temperature given.

The TESS catalogue (Stassun et al. 2018; Muirhead et al. 2018) was also used as a source of star mass estimates. From TESS1 325 masses were gathered, while in TESS2 there were no masses of stars in question.

Using all the above-mentioned sources, for most of the stars, we have obtained several, sometimes different, mass estimates. For a great majority of them (572 from among 783 single stars) we finally decided to use the mass estimates from Anders et al. (2019). For objects missing in this source we decided to take a mean of the gathered values after the most extreme ones and these most flawed have been excluded.

Masses of components of multiple systems were obtained in a similar way depending on the data availability. 76 stellar masses of components of multiple systems recognised by *Gaia* DR2 were gathered from Anders et al. (2019). We also used tens of other sources found through the SIMBAD and VIZIER databases. In some cases mass estimates were taken from papers that describe the specific multiple system.

### 4. New, puzzling but potentially strong stellar perturber: ALS 9243

Using the astrometry from the Gaia DR2 catalogue and the radial velocity from the SIMBAD database we found that a star ALS 9243, never mentioned in earlier papers in a context of being a cometary motion perturber, 2.5 Myr ago passed as close as 0.25 pc from the Sun. But the main reason for being surprised was the estimated mass of this star: according to the spectral type O9 – B0 and the luminosity class IV repeated in the literature we should assume its mass to be over 15 solar masses! Such a close passage of such a massive star that took place only 2.5 Myr ago would have made a strong influence on the observed long period comet orbits and probably on the Solar System as a whole. At first, we classified this object as a multiple star due to the information from the SIMBAD database but later it appeared that its multiplicity is rather not confirmed. Although this object can be found in the WDS catalogue (Mason et al. 2001a) basing on observations done by Aldoretta et al. (2015), there is no indication of a name of the alleged second component and data necessary to calculate its position and velocity.

#### 4.1. What we know about the star ALS 9243

The star in question was probably first mentioned and named in 1965 during the completion of the 'Luminous Stars in the Northern Milky Way' catalogue (Nassau et al. 1965; Hardorp et al. 1965). The star was designated as LS VI -04 19, which reads: Luminous Stars, volume six, declination zone -04, star number 19. This was an objective prism survey aimed at young stars. They quote OB as the 'estimated spectral type'. Almost forty years later an 'all sky' database of OB stars was collected by Reed

(2003, 2005) and he assigned a new name to this star: ALS 9243. This name will be used throughout the present work.

Later on, this star has been also included in large modern catalogues: Tycho-2 (as TYC 4809-2410-1, Høg et al. 2000), 2MASS (as J06593022-0448438, Cutri et al. 2003) and finally *Gaia* DR2 (as DR2 3101630187797866112, Gaia Collaboration et al. 2018).

In an elegant paper by Graham (1971) the photometric distance to this star was first estimated, as well as its radial velocity. In the last row of his Table I one can find a distance modulus equal  $11.9^m$  (which is equivalent to the distance of 2.4 kpc, other distance estimates are presented in Table 1) and  $v_r = 49.5$  km s<sup>-1</sup>. In the same year (Crampton 1971) the star was for the first time associated with the H II region and its spectral classification was narrowed down to B0 IV. A year later Crampton (1972) published radial velocity measurements of the star in question, again using the objective prism, ranging from 26 to 55 km s<sup>-1</sup> during a ten day interval.

Recently a paper by Anders et al. (2019) appeared, where distances and astrophysical parameters of large number of *Gaia* DR2 stars are recalculated. For ALS 9243 they obtained a distance of about 93 pc and a mass of only 0.65 M<sub> $\odot$ </sub>. These results strongly depend on *Gaia* DR2 results for this star. However, it should be noted, that the most preferred *Gaia* DR2 (Gaia Collaboration et al. 2018) astrometry quality indicator, a re-normalised unit weight error (RUWE)<sup>1</sup> for ALS 9243 equals 6.97 while the upper RUWE limit for 'good' astrometric solutions is  $1.4^2$ .

#### 4.2. Atmospheric parameters of ALS 9243

We tried to solve this puzzling inconsistency in ALS 9243 parameters. In January 2020 on our kind request three spectra with the fiber fed echelle spectrograph ESPERO connected to the 2-m telescope in Rozhen National Astronomical Observatory (Bonev et al. (2017)) with resolving power  $R \sim 40\,000$  and in range from 410 to 950 nm were obtained. In our (still simplified and approximate) analysis presented below we used only one spectrum observed on 9th January due to its better quality, where measured signal-to-noise ratio was between 30-40.

The atmospheric parameters: an effective temperature  $T_{\rm eff}$ , a surface gravity log g and a projected rotational velocity  $v \sin i$  were calculated using the ISPEC code (Blanco-Cuaresma 2019; Blanco-Cuaresma et al. 2014). The observed spectrum was compared with a grid of fluxes BSTAR2006 (Lanz & Hubeny 2007) created with TLUSTY model atmospheres and SYNSPEC spectra. We used stellar atmosphere models which are metal line-blanketed, non-LTE, plane-parallel, and we examine hydrostatic atmospheres.

At first, the effective temperature and the surface gravity were estimated using the Balmer lines H $\alpha$  and H $\beta$ . For hot stars ( $T_{\rm eff} > 8000$  K) Balmer lines are sensitive to the log g parameter thus both  $T_{\rm eff}$  and log g parameters were derived simultaneously. Additionally, we assumed a microturbulence velocity of 2 km s<sup>-1</sup> and a macroturbulence of 0 km s<sup>-1</sup>. The metallicity [M/H] value was fixed to 0.0 dex. In our calculation we also used the six neutral and ionized helium lines (He I, He II), which are well visible in the ALS 9243 spectrum, such as He II (468.6, 471.4 nm) and He I (501.6, 587.5, 667.8, 706.7 nm). The ob-



**Fig. 1.** Comparison of the observed spectrum (black color) and synthetic ones (with different colors) of the H $\beta$  region and He I lines. For He I lines different colors correspond to synthetic spectra calculated for various  $T_{\rm eff}$ , log g and vsini within the error limits (respectively aquamarine:  $T_{\rm eff} = 26\,000$  K, logg = 3.7 dex , vsini = 10 km s<sup>-1</sup>; blue:  $T_{\rm eff} = 28\,000$  K, log g = 3.9 dex, vsini = 15 km s<sup>-1</sup>; red:  $T_{\rm eff} = 30\,000$  K, log g = 4.2 dex, vsini = 20 km s<sup>-1</sup>).

tained effective temperature is  $28\,000 \pm 2\,000$  K,  $\log g = 3.9 \pm 0.3$  and  $v\sin i = 15 \pm 5$  km s<sup>-1</sup>. The comparison of the observed and synthetic spectra within the error limits of H $\alpha$  and two He lines is shown in Fig. 1.

To evaluate the uncertainties of all determined parameters we took into account the difference in values calculated separately from the lines. The obtained uncertainties are mainly caused by low signal-to-noise ratio and continuum normalization process of the echelle spectra during which it is difficult to recover the original line profiles. The estimated atmospheric parameters for ALS 9243 object should be verified in the future from a spectrum of a better quality, with the signal-to-noise ratio of at least 100.

Our temperature measurement is in good agreement with most of the previous spectral type determinations (Table 2). The obtained temperature and log g correlate with B0 spectral type and subgiant luminosity class IV. According to Straižys (1992) tables this corresponds to the mass of ~ 22  $M_{\odot}$ .

#### 4.3. Call for observations

The trigonometric distance obtained by *Gaia* DR2 seems to be highly inconsistent with the luminosity and the visual magnitude of such a massive star. The star should be much farther. On the other hand, the J-H color index (fluxes in J and H band were taken from Cutri et al. (2003), values of 9.692 and 9.553 respectively, were adopted) of the star suggests a lower temperature of about 6800 K. All this contradictory results could be explained for example with the extremely high extinction. In the line of sight we have a molecular cloud SH2-287. However, the cloud might be in the star background or closer to us, and the distance of the SH2-287 is estimated to be 2.1 kpc (Neckel & Staude 1992). The striking discrepancy between the trigonometric and photometric results should be explained with future observations.

We would like to encourage observers to make an effort to clarify this situation by taking high quality positional and spectro-photometric observations of ALS 9243. Our preliminary attempt revealed that it is a difficult object as it is not so bright

<sup>1</sup> Lindegren, 2018, see: L. Considerations for the Tech. Use of DR2 Astrometry, available rep., at http://www.rssd.esa.int/doc\_fetch.php?id=3757412

<sup>&</sup>lt;sup>2</sup> see *Gaia* Data Release 2 Documentation (release 1.2), Section 14.1.2

Table 1. ALS 9243 distance estimates and measurements

distance [pc]	method	ref
2400	photometric	Graham (1971)
$3300 \pm 800$	photometric	Avedisova & Kondratenko (1984)
31 ( −21 +∞)	trig. parallax	Tycho-1 (ESA 1997)
86 (-35 + 83)	photometric	Ammons et al. (2006)
256	photometric	Pickles & Depagne (2010)
2900	photometric	Garmany et al. (2015)
3200	photometric	Aldoretta et al. (2015)
443	photometric	Stassun et al. (2018)
94.7 (-3.5 + 3.7)	trig. parallax	Gaia DR2 (Gaia Collaboration et al. 2018)
94.5 (-3.3 + 4.0)	trig. parallax	Bailer-Jones et al. (2018b, based on <i>Gaia</i> DR2)
93 (-4 + 6.0)	'photo-astrometric'	Anders et al. (2019, based on Gaia DR2)

Table 2. ALS 9243 spectral classifications

Teff [K]	Spectrum	Luminosity class	$\mathbf{M}_V$	ref
_	OB	-	-	Nassau et al. (1965)
-	_	-	$-3.4^{m}$	Graham (1971)
_	B0	IV	-	Crampton (1971)
_	09.5	V	$-2.29^{m}$	Georgelin et al. (1973)
_	B0.5	V	-	Mayer & Macák (1973)
_	09.5	V	-	Moffat et al. (1979)
6608	_	_	_	Ammons et al. (2006)
_	F5	V	-	Pickles & Depagne (2010)
_	_	-	$-3.6^{m}$	Garmany et al. (2015)
_	O9.7	IV	-	Aldoretta et al. (2015)
7773	_	V	-	Stassun et al. (2018)
6185	_	V	_	Gaia Collaboration et al. (2018)
5461	_	—	-	Anders et al. (2019)

and it is visible on the background (or surrounded by) a H II nebula. The most striking controversy is the distance: 0.1 or over 2 kpc. Also its mass estimates vary from 0.5 up to over 20  $M_{\odot}$ .

For now, in the absence of a clear explanation of the discrepancies found in the literature, we decided to keep this star in our list, use the *Gaia* DR2 astrometry and a 'compromise' mass of  $2 M_{\odot}$ .

#### 5. Multiples

Almost all previously published lists of stars passing through the close solar neighbourhood (García-Sánchez et al. (2001), Dybczyński (2006), Jiménez-Torres et al. (2011), Dybczyński & Berski (2015), Bailer-Jones (2015a), Bailer-Jones et al. (2018a), and Bailer-Jones (2018)) contain only objects considered as single stars even if they are, in reality, parts of multiple systems. Treating components of multiple systems as single stars often leads to misleading conclusions. While a particular component of multiple seems to encounter the Sun at a very small distance, a center of mass of this multiple can even move in another direction.

Because stellar systems are statistically more massive than single stars, and therefore can act as more significant perturbers, in our work we tried to analyse as many cases of multiplicity as possible. Each star from our list was checked whether it is a component of a system. An identification of multiple systems was done utilizing the SIMBAD database and this explains why our search for multiples was generally limited to stars mentioned in earlier papers. New potential perturbers found only thanks to the *Gaia* cannot be checked for multiplicity – to the best of our knowledge no studies were published identifying all cases (or even the significant part of cases) of multiple systems in *Gaia* DR2.

While the SIMBAD database facilitated our work on this subject, we had to check each case carefully by an extended research. Some stars identified in SIMBAD as components of multiple systems are, in fact, single stars, having for example completely different parallaxes. Their alleged multiplicity can, for example, remain from the time where they were observed close to other stars and thought to be dynamically bound to them. On the other hand, although some stars are definitely parts of multiple systems, we sometimes had to treat the whole system as a single star due to the data incompleteness.

It is important to mention that the most reliable center of mass kinematic parameters can be obtained only when the fiveparameter astrometry and the radial velocity are given for all components for the same epoch which is nearly never the case.

For most of the multiple systems we calculated their center of mass parameters with data available in *Gaia* DR2 or with data available in the SIMBAD database. Only in four cases specific systems were described in dedicated papers and we relied on the data found therein.

In Sect. 6 several interesting cases of stellar system investigated by us are described. These systems were either thoroughly examined by us and for the first time classified as multiple stellar perturbers (despite the fact that some individual components of these systems were previously suggested as stellar perturbers) or, by considering their multiplicity, we ruled them out from the list of potential stellar perturbers. Where possible, a comparison with the results found in papers listed in Sect. 2 is given.

#### 6. Multiples - special cases

#### 6.1. Algol

Algol, know also as  $\beta$  Persei, is a very bright hierarchical system. It consists of a close binary stellar system with a more distant tertiary component. Algol's components were not separated in the SIMBAD database. In the past it was treated as a single star and was classified as stellar perturber by many authors: García-Sánchez et al. (2001), Dybczyński (2006), Jiménez-Torres et al. (2011), and Dybczyński & Królikowska (2015). Despite a rather distant passage near the Sun (over 3 pc) this perturber is rather important in near-parabolic comet motion studies due to its large systemic mass and a very small systemic velocity relative to the Sun.

In *Gaia* DR2 catalogue one of the components of this system has its right ascension and declination measured but there are no parallax, proper motion, and radial velocity data and, more importantly, there are no clues which component was observed.

Taking this into account, we decided to rely on data found in the another source. Based on observations focused on Algol and UX Arietis, Peterson et al. (2011) published a parallax, a declination, a right ascension, both proper motion components and a radial velocity of the center of mass of this triple system. We adopted these values which are listed in Table 3.

In view of these values, the Algol system, with its total mass equal to 6.0  $M_{\odot}$ , is the most massive perturber in our list. In fact we use the proximity threshold of 4 pc when constructing our list of perturbers to keep the Algol system included. Using the input values presented in Table 3 we have obtained a minimal distance of 3.78 pc during the closest approach to the Sun which took place 13.06 Myr ago with the relative velocity of only 2.17 km s<sup>-1</sup>. Two last values mentioned makes Algol's encounter the oldest and slowest one from among all perturbers in our list. It is worth to mention that the data from Peterson et al. (2011) significantly changed these parameters. The previously used values, derived on the basis of Lestrade et al. (1999) or Hipparcos catalogue read: the minimal distance of ~ 3 pc, the closest approach at 6 – 7 Myr ago with the relative velocity of 4 km s<sup>-1</sup>.

#### 6.2. $\rho$ Orionis

 $\rho$  Orionis is a spectroscopic binary classified as stellar perturber by Dybczyński & Królikowska (2015). It was then treated as a single star, as its components are not listed separately in XHIP Anderson & Francis (2012) which was the only source of the 6D stellar data used by the authors.

Thanks to *Gaia* DR2 we were able to update data concerning this system. Both of its components were identified by us in *Gaia* DR2 catalogue as *Gaia* DR2 3235349837026718976 and *Gaia* DR2 3235349940105933568 objects. The second one do not have the radial velocity measured in *Gaia* DR2, so we adopted the value from Malaroda et al. (2006). We used masses from Tokovinin (2018) where it is suggested that  $\rho$  Orionis is a triple system, but, due to lack of any positional information about the third component, we decided to assume that it is a binary system.

New data allowed us to calculate the center of mass parameters and the new parameters of the approach. While in Dybczyński & Królikowska (2015) the minimal distance from the Sun was equal to 3.23 pc, now it is over 17 pc. The encounter happened 2.60 Myr ago at the relative velocity of 46.14 km s<sup>-1</sup>.

As it can be seen, an improvement of the quality of the data and the confirmation of the binary character of the object in question cancelled the importance of  $\rho$  Orionis as a stellar perturber of cometary motion.

#### 6.3. Ross 614

Ross 614 was at first discovered as a single star by Ross (1927). Then, Reuyl (1936) detected the second component of this very low mass system which consists of red dwarfs. Later it was a subject of extensive studies, the most recent ones were conducted by Ségransan et al. (2000), Gatewood et al. (2003), and Kervella et al. (2019).

Only one component of Ross 614 can be found in the *Gaia* DR2 catalogue, where it is identified as *Gaia* DR2 3117120863523946368, but it does not have its radial velocity measured. For the second component, even in the SIMBAD database, data are incomplete.

Although above-mentioned papers in-depth examine nature of this stellar system, data found therein are not sufficient enough for our purpose.

For this reason, our decision was to take masses of both components from Anders et al. (2019) but use astrometry done only for Ross 614A. Incomplete data from *Gaia* DR2 were augmented with radial velocity from Gontcharov (2006).

With these values adopted our calculations show that this system encountered the Sun vicinity 0.09 Myr ago at a distance of 3.25 pc. Relative velocity at the time of approach was  $27.18 \text{ km s}^{-1}$ .

In comparison with results obtained in the past, minimal heliocentric distance of Ross 614 increased. In XHIP catalogue Anderson & Francis (2012), so also in Dybczyński & Królikowska (2015), it was equal to 3.03 pc. In order to obtain more reliable parameters of the approach new astrometry for the second component (Ross 614B) is needed.

#### 6.4. a Canis Majoris

 $\alpha$  Canis Majoris, known also as Sirius is a visual binary containing Sirius A which is the brightest star in the sky and Sirius B, the nearest white dwarf. There was a long-lasting discussion whether there is a third body in that system because of irregularities in orbits of Sirius A and B. This possibility was probably ruled out by extensive studies, see for example Bond et al. (2017).

Sirius was earlier classified as stellar perturber by García-Sánchez et al. (2001), Dybczyński (2006), Jiménez-Torres et al. (2011), and Dybczyński & Królikowska (2015) but in none of these papers the multiplicity was considered.

In *Gaia* DR2 only one component is included as *Gaia* DR2 2947050466531873024, but it does not have radial velocity measured and there is also no radial velocity in SIMBAD database. For that reason, because we were unable to calculate center of mass parameters, we decided to use values found in Gatewood & Gatewood (1978) augmented with new measurements of masses from Bond et al. (2017). The values adopted here are listed in Table 4 were positions and proper motions are given in relation to 1950.0 epoch in FK4 frame. They were further recalculated to be consistent with other data.

From these data we obtained parameters of the encounter with the Sun which will happen at 2.41 pc in 0.06 Myr. This result is generally in agreement with minimal heliocentric distances obtained earlier. The relative velocity at the time of the approach will be equal to  $18.49 \text{ km s}^{-1}$  which makes it a rela-

#### Table 3. Algol system center of mass parameters

parameter	value	unit
parallax	$34.7 \pm 0.6$	mas
primary mass	3.70	$M_{\odot}$
secondary mas	0.79	$M_{\odot}$
tertiary mass	$1.51 \pm 0.02$	$M_{\odot}$
right ascension proper motion	$2.70 \pm 0.07$	mas yr <sup>-1</sup>
declination proper motion	$-0.80 \pm 0.09$	mas yr <sup>-1</sup>
radial velocity	2.1	$\rm km~s^{-1}$
right ascension	$3^{h}08^{m}10.13241 \pm 0.7$	mas
declination	$40^{\circ}57'20''3353 \pm 0.6$	mas

Table 4.  $\alpha$  Canis Majoris center of mass parameters (position and proper motions for 1950.0 epoch in the FK4 frame).

parameter	value	unit
parallax	$0.3777 \pm 0.0031$	mas
primary mass	$2.063 \pm 0.023$	$M_{\odot}$
secondary mas	$1.018 \pm 0.011$	$M_{\odot}$
right ascension proper motion	-0.0379	s yr <sup>-1</sup>
declination proper motion	-1.211	" $yr^{-1}$
radial velocity	-7.6	$km s^{-1}$
right ascension	$6^{h}42^{m}56^{s}.73$	
declination	-16°38′46″4	

tively slow passage.  $\alpha$  Canis Majoris system is one of the more massive objects on our perturber list.

#### 6.5. y Leonis

The WDS catalogue (Mason et al. (2001a)) identifies four components of  $\gamma$  Leonis system. We conducted in-depth investigation to verify whether these components belong to the system.

While two of them (WDS J10200+1950A and WDS J10200+1950B) have exactly the same parallax of 25.96 mas, for the third one (WDS J10200+1950Ca,Cb) the parallax equals 201.3683 mas, and for the forth one (WDS J10200+1950D) it is measured to be 1.4566 mas. All these values were taken form the SIMBAD database. As it can be seen, only the first two components actually create a binary system. In the light of available data, two later components were treated by us as single stars that happen to be visually close to the system.

We decided to calculate the center of mass parameters for  $\gamma^1$  Leonis and  $\gamma^2$  Leonis. None of the components is in *Gaia* DR2 catalogue, so all data were taken from the SIM-BAD database.  $\gamma^1$  Leonis has its mass (1.41  $M_{\odot}$ ) estimated in Niedzielski et al. (2016). For  $\gamma^2$  Leonis we have to adopt a crude mass estimate of 1.50  $M_{\odot}$  basing on a spectral type. For this system we calculated the minimal heliocentric distance and it came to be 33.32 pc and will occur in 0.26 Myr at the relative velocity equal to 73.06 km s<sup>-1</sup>. These values ruled  $\gamma$  Leonis out from our final list of stellar perturbers.

WDS J10200+1950Ca,Cb is treated as a single star and as such it is included in the RECONS<sup>3</sup> list of the one hundred nearest star systems. It is also included in *Gaia* DR2 catalogue as *Gaia* DR2 625453654702751872 were it does not have the radial velocity measured. We augmented data from *Gaia* DR2 with a radial velocity from SIMBAD database and a mass of 0.467  $M_{\odot}$  from TESS1 catalogue. Our results show that this star encountered the Sun 0.21 Myr ago at the minimal distance of 3.41 pc and the relative velocity of 17.14 km s<sup>-1</sup>.

For WDS J10200+1950D, identified also as *Gaia* DR2 625453856566097024, data from *Gaia* DR2 were used and 1.04  $M_{\odot}$  mass from Deka-Szymankiewicz et al. (2018) was adopted. The minimal distance of 638.01 pc was obtained and we can state that this star is definitely not a stellar perturber of long period comets motion.

These examples show that, when it comes to multiple systems, we can not even rely on data found in databases concerning multiple systems and a careful investigation of each alleged component is necessary.

#### 6.6. a Centauri

 $\alpha$  Centauri system with its three components is the nearest stellar system to the Sun. It comprise of  $\alpha$  Cen A, a solar-like star,  $\alpha$  Cen B which is a cooler dwarf, and Proxima, a cool red dwarf, recently recognized to be a host of the nearest exoplanet – Proxima Centauri b.

All of its components were classified as stellar perturbers in García-Sánchez et al. (2001), Dybczyński (2006), Jiménez-Torres et al. (2011), Dybczyński & Królikowska (2015), and Bailer-Jones (2015a), but only in Dybczyński & Królikowska (2015)  $\alpha$  Centauri was treated as a multiple system and center of mass parameters were calculated to obtain the minimal heliocentric distance.

Since  $\alpha$  Cen A and  $\alpha$  Cen B are not included in *Gaia* DR2 catalogue and Proxima does not have its radial velocity measured in *Gaia* DR2 we based our calculation on the heliocentric coordinates given in the Galactic frame found in Kervella et al. (2017), summarized in Table 5.

From these values we obtained the center of mass parameters for the system and the parameters of the closest approach to the Sun which will happen in 0.03 Myr at the distance of 0.97 pc, and with the relative velocity of 32.35 km s<sup>-1</sup>. This result is generally in agreement with the values previously published for components of  $\alpha$  Centauri.

<sup>&</sup>lt;sup>3</sup> http://www.recons.org/

Table 5. Heliocentric coordinates and space velocity components of  $\alpha$  Centauri AB and Proxima in the Galactic frame

<b>parameter</b> (unit)	$\alpha$ Cen	Proxima
X (pc)	$0.95845 \pm 0.00078$	$0.90223 \pm 0.00043$
Y (pc)	$-0.93402 \pm 0.00076$	$0.93599 \pm 0.00045$
Z (pc)	$-0.01601 \pm 0.00001$	$-0.04386 \pm 0.00002$
XV (kms <sup>-1</sup> )	$-29.291 \pm 0.026$	$-29.390 \pm 0.027$
YV (kms <sup>-1</sup> )	$1.710 \pm 0.020$	$1.883 \pm 0.018$
ZV (kms <sup>-1</sup> )	$13.589 \pm 0.013$	$13.777 \pm 0.009$
mass $(M_{\odot})$	$2.0429 \pm 0.0072$	$0.1221 \pm 0.0022$

#### 6.7. HD 239960

HD 239960, known also as Kruger 60, is a visual binary comprising of two M spectral type stars and it is supposed to be a host of a planetary system (see for example Bonavita et al. (2016)). This stellar system was earlier identified as stellar perturber but its multiplicity has never been taken into account.

Recently both components of Kruger 60 were observed by *Gaia* and a new astrometry is available in the *Gaia* DR2 catalogue. Components of the system are designated as *Gaia* DR2 2007876324466455424 and *Gaia* DR2 2007876324472098432. For both of them the radial velocity is missing in this source.

Values from *Gaia* DR2 catalogue can be augmented with radial velocities found in the SIMBAD database and other sources. SIMBAD database contain values from the General Catalogue of Stellar Radial Velocities (Wilson (1953)), -24.0 km s<sup>-1</sup> for HD 239960A and -28.0 km s<sup>-1</sup> for HD 239960B, error of radial velocity is in both cases estimated to be 5 km s<sup>-1</sup>. There is no indication on epochs of observation when these values were measured. In the literature it is possible to find values of radial velocities of these stars ranging from -36.0 to -16 km s<sup>-1</sup> but often without any information on which component was observed.

Because the orbital period of Kurger 60 is estimated to be only 44.6 years (Bonavita et al. 2016) we aimed to use positions, proper motions, and radial velocities referred to the same epoch. As it was at first impossible, we decided to use available data and calculate position and velocity of the center of mass. Positions, proper motions, parallaxes from *Gaia* DR2 were used in conjunction with radial velocities from Wilson (1953) and masses found in Bonavita et al. (2016).

While working on the list of stellar perturbers described herein a second interstellar comet 2I/Brisov was discovered. We were involved in a study on its origin and Kruger 60 seemed to be a good candidate (for more details see Dybczyński et al. (2019)). An in-depth investigation on data available for this stellar system showed us that they are insufficient to obtain reliable results. Thanks to Fabo Feng (via private communication) we have been given an access to the new, unpublished right ascension, declination, parallax, proper motions and the radial velocity of the center of mass of Kruger 60 which were calculated using PEXO package (Feng et al. 2019) basing on data from HIPPAR-COS (van Leeuwen 2007), WDS catalogue (Mason et al. 2001b) and the recent LCES HIRES/Keck survey (Butler et al. 2017).

These data were further used to obtain parameters of the approach to the Sun. Kruger 60 is a future perturber. It will reach its minimal heliocentric distance of 1.81 pc in 0.09 Myr with the relative velocity equal to  $38.03 \text{ km s}^{-1}$ . This is a slightly smaller distance than presented in the recent paper by Bailer-Jones (2015a) were Kruger 60 was classified as a close approaching star but its multiplicity was not considered.



**Fig. 2.** Histogram of minimal heliocentric distances. Distances were divided into two subsets. Green bars correspond to the encounters from the past, blue ones depict the minimal distances of approaches which will occur in the future. 643 stars with the minimal distances smaller than 4.0 pc were considered.

#### 7. On-line database for stellar perturbers

After collecting all necessary data we prepared a simple database containing all the data with their uncertainties and sources. We also included heliocentric rectangular position and velocity components in the Galactic frame together with the adopted mass estimates of all 820 considered perturbers. The whole set of these objects was numerically integrated back and forth in time taking into account all their mutual interactions and including the Galactic overall potential, as described in Dybczyński & Berski (2015). Our results reveal that 714 of our perturbers encountered or will encounter the Sun within a distance smaller than 10 pc and 643 were or will be closer than 4.0 pc. We finally accepted this later proximity threshold as the one defining the potential perturber, just to keep the Algol system in our list. This system is important because of its large mass of  $6 M_{\odot}$  and extremely small relative velocity of 2 km s<sup>-1</sup>. Finally, we keep data for all 820 objects in our database but we name only 643 of them as 'potential perturbers' of the near-parabolic comet motion. It is worth to mention that the final list includes 147 new objects for the first time qualified as potential stellar perturbers of long period comets motion.

The distribution of the minimal distances between these stars and the Sun is presented in Fig. 2. The number of the past perturbers is slightly greater than the number of the future ones due to a disproportion in data from the SIMBAD database (there are many more objects with positive radial velocity there). Such a disproportion can also be noticed in data from the *Gaia* mis-



Fig. 3. Nominal relative velocities, minimal heliocentric distances, and masses of objects included in our list. 613 stars were plotted. See text for a explanation.

sion, see for example results in Bailer-Jones et al. (2018a) when the sample is limited to stars with minimal heliocentric distance smaller than 4 pc.

The obtained minimal distances from the Sun are also included in the database for all objects. Objects with the small minimal heliocentric distance will be used by us in the long period comets dynamical studies. All the remaining objects are placed in our database just for the record since they were mentioned in earlier papers. Some of them might return to the list of potential perturbers when new data will be gathered. As a consequence, one can trace how an importance of a specific object changes due to the improvement of data quality.

In Fig. 3 we also present a three-parameter statistic of stars close passages near the Sun which includes the most important nominal parameters from the point of the star – comet interaction: minimal distances, relative velocities and masses of the perturbers. To increase the readability of this plot we additionally restrict ourselves to stars passing the Sun with a relative velocity smaller than 200 km s<sup>-1</sup>. The purpose of this figure is only to graphically illustrate the content of our database, i.e. nominal parameters of potential perturbers. Readers interested in a detailed distribution of stars approaching the Sun and in the completeness of our current knowledge in this field should consult an extensive study by Bailer-Jones (2018).

Our database is publicly available at the address: https://pad2.astro.amu.edu.pl/pub with a simple interface to access the data and their uncertainties, and the sources.

Various lists and statistics are also available and crucial results are made available for the download.

# 8. Usefulness and importance of the new list of stellar perturbers

As it was already stated, the aim of collecting the list of stars and stellar systems announced in this paper is to provide a tool for studying past and future motion of long period comets outside the planetary zone. We carefully selected all potential stellar perturbers of the LPCs motion basing on the contemporary sources of stellar data, mainly the *Gaia* DR2 catalogue (Gaia Collaboration et al. 2018). The *Gaia* mission results are revolutionary in this respect – for example, before this mission we had no more than 130 thousand stellar parallaxes at our disposal, most of them from the Hipparcos catalogue (van Leeuwen 2007). Now we can use over 1.3 billion parallaxes (see for example Bailer-Jones et al. 2018b) in our search for potential LPCs motion perturbers.

The last paper that described the past and future motion of LPCs under simultaneous Galactic and stellar perturbations (Królikowska & Dybczyński 2017) used a list of 90 potential stellar perturbers selected on the basis of data from the XHIP catalogue (Anderson & Francis 2012) which was a compilation of the Hipparcos mission results augmented with radial velocities available at that time. In that paper Królikowska & Dybczyński

**Table 6.** Examples of the LPCs dynamical history results obtained with different models. For four representative comets we present the perihelion distance values: of the original orbit (the first row), at a previous perihelion but calculated without stellar perturbations (the second row), at a previous perihelion as obtained in Królikowska & Dybczyński (2017) (the third row) and in the last row also previous perihelion but obtained with the new list of perturbers. In the last three rows a previous perihelion distance is presented in the form of three deciles, 10, 50, and 90 percent.

	C/1993 F1		C/1997 BA6		C/1999 N4		C/2006 E1					
q <sub>original</sub>	5.899	50 +/- 0.	00007	3.4403	71 +/- 0.	000006	5.5047	39 +/- 0.	000006	6.036	08 +/- 0.	00001
q <sub>prev</sub> (Galaxy only)	7.33	8.16	9.76	15.90	19.54	24.74	6.36	6.44	6.54	22.78	31.80	48.86
q <sub>prev</sub> old model	7.12	7.92	9.44	16.89	20.67	26.08	6.33	6.41	6.50	21.55	29.96	46.00
q <sub>prev</sub> new model	46.27	58.31	68.13	3.10	3.92	7.36	72.07	74.29	75.87	21.91	25.69	52.01

(2017) studied a hundred of LPCs with large osculating perihelion distances and found no strong stellar perturbation effects.

Now we offer the updated list of 643 potential perturbers. Some of them can perturb cometary motion in a spectacular way, see Wysoczańska et al. (2020). But the effect of the newly obtained list of perturbers on motion of many more comets is also noticeable and very important. A paper describing this in great detail is in preparation, here we only show selected results for four comets to serve as an example. In Table 6 we present previous perihelion distances (one orbital period to the past) for C/1993 F1, C/1997 BA6, C/1999 N4, and C/2006 E1, obtained using three different dynamical models. The starting perihelion distances of the original orbits are presented in the first row. Next three rows consists of previous perihelion distance values obtained without stellar perturbations, with an old dynamical model used in Królikowska & Dybczyński (2017) and using the list of stellar perturbers announced in this paper, respectively. These values are presented here as three deciles (10-50-90 percent) since the distribution of clones used for the uncertainty estimation is far from being Gaussian.

In Królikowska & Dybczyński (2017) a comet is classified as dynamically old if its previous perihelion distance is below 10 au. On the other hand, it is called dynamically new if the previous perihelion distance appears to be greater than 20 au. As it is easy to note, the usage of the new list of stellar perturbers reversed this classification for three of the presented comets. Such a change is noticed in a large percentage of almost 300 studied LPCs and will be described in a future paper (Dybczyński & Królikowska, in preparation).

But there exist one important problem with the list of stellar perturbers announced in this paper – its name is ALS 9243. As it was described in detail in Sect. 4 there exist a fundamental inconsistency in data available for this star: its distance obtained with the *Gaia* mission strongly disagrees with its previously published spectral type and luminosity class. This controversy may lead to two solutions:

- The distance based on *Gaia* DR2 is completely wrong, the star is much farther (say, over 2 kpc) and this completely rules out this star from a list of potential perturbes.
- The astrometric results presented in *Gaia* DR2 are approximately correct, which makes this star an important perturber since its nominal minimal distance from the Sun is as small as 0.25 pc and the approach occurred over 2 Myr ago, so almost all the observed LPCs can be affected it. But, to be able to calculate this effect, we need the mass of this star.

As it was described in detail in Sect. 4, the most often quoted spectral type and luminosity class of ALS 9243 is O9 IV (see

**Table 7.** The influence of the assumed ALS 9243 mass on the previous perihelion distance of the same comets as presented in Table 6. The first row, labelled 'O', presents the starting perihelion distance of the original orbit. The last row, labelled 'G', presents the previous perihelion distance when all stellar perturbations were omitted. In the remaining rows the first column shows the assumed mass of ALS 9243 expressed in solar masses.

	C/1993 F1	C/1997 BA6	C/1999 N4	C/2006 E1
O:	5.8995	$\begin{array}{r} 3.4404\\ 1486.4459\\ 361.4909\\ 60.0877\\ 3.7164\\ 5.5040\\ 17.6174\\ 19.6341\end{array}$	5.5139	6.0361
18.5	58.0720		73.7724	2513.8460
10.0	58.0730		73.7736	656.4899
5.0	58.0735		73.7743	137.7216
2.0	58.0738		73.7747	22.0914
1.0	58.0740		73.7748	15.6289
0.0	58.0741		73.7749	25.1433
G:	8.1556		6.4411	31.8632

Table 2) which results in the mass of 18.5 solar masses. Since these astrophysical parameters are in contradiction with the *Gaia* DR2 astrometry and such a close passage of a hot and massive star 2 Myr ago seems improbable, we have to make a trade-off. For now, waiting for new data, we decided to use the *Gaia* DR2 astrometry but we assume the mass of ALS 9243 to be equal to 2.0 solar masses.

The correct mass value of ALS 9243 might be crucial for the LPCs dynamical history studies as we show in Table 7. Here we used nominal orbits of the same comets as shown in Table 6 however, we calculate their previous perihelion distance assuming different masses for ALS 9243. In the first row, labelled 'O', the starting value of the perihelion distance is shown. In the next six rows the previous perihelion distance of these four comets are presented, calculated using the assumed mass of ALS 9243 shown in the first column. For the sake of additional comparison, in the last row, labelled 'G', we present the previous perihelion distance obtained without stellar perturbations at all, only as a result of the galactic perturbations.

One can easily observe that comets C/1993 F1 and C/1999 N4 are practically not affected by ALS 9243 at all. But the remaining two comets are affected significantly by this star and their previous perihelion distance depends strongly on its assumed mass.

#### 9. Conclusions and prospects

Due to a great increase of our knowledge on the Galactic neighbourhood of the Sun we were able to significantly correct and update the list of potential stellar perturbers of long period comets motion. The full list of analysed objects includes 751 single stars and 69 stellar systems. Among them 643 objects appeared to have their minimal heliocentric distance smaller than 4 pc and therefore are classified by us as potential perturbers of LPCs motion

Our updated list consists both of new stars or stellar systems found by us in a manner described in Sect. 2 and of stars that were previously classified as stellar perturbers in the earlier papers mentioned in Sect. 2. Objects from the later group were thoroughly examined whether new or more accurate data are available. Thanks to the improvement in the quality of data we were able to verify the importance of each perturber, how it changes due to new measurements, and, more importantly, check whether the star in question is a component of a multiple system. In some cases taking the multiplicity into account resulted in removing the perturber from the list, in other cases, this approach just changed the expected value of the minimal heliocentric distance and therefore the potential influence of this particular perturber on LPCs motion.

The examples presented in Sect. 6 show the importance of taking the multiplicity into account and many drawbacks of still incomplete data which sometimes limited and hindered us from calculating center of mass parameters based on data concerning all known components of the considered system. Main issues are: lack of radial velocities measurements and masses (which is also applicable to single stars), different epochs of measurements of the positions and proper motions of components of the systems which can lead to unrealistic results, and incomplete data on multiplicity of systems, especially these with stars for the first time observed by Gaia mission. These issues may be addressed with the future *Gaia* data releases.

During gathering of data necessary to calculate minimal heliocentric distances of the stars, we came across a star designated as ALS 9243 which, basing on available measurements (from Gaia DR2 catalogue augmented with radial velocity from SIM-BAD database), 2.5 Myr ago visited the vicinity of the Sun at the distance equal to 0.25 pc. This object has never been mentioned as a stellar perturber before and before the Gaia mission its distance was often estimated to be more than 1 kpc. Because of the discrepancies in mass estimates, we decided to check the atmospheric parameters of the star in question. Our results indicates that this star seems to be very massive, even up to 22  $M_{\odot}$ which is inconsistent with its distance presented in Gaia DR2. We hope that, in the nearest future, we will be given an opportunity to obtain consistent data and verify the significance of this perturber. Before the clarification of this puzzling case, we decided to use Gaia DR2 astrometry and adopt 2.0  $M_{\odot}$  as the mass of ALS 9243 in our database.

All data used by us (with sources) and our results are gathered in a small publicly available database of potential perturbers. This might be a useful tool in future dynamical studies of near-parabolic comets dynamics.

Acknowledgements. We would like to thank the anonymous referee for constructive suggestions that greatly improved the quality of this manuscript. This research was partially supported by the project 2015/17/B/ST9/01790 founded by the National Science Center in Poland. MP wants to thank Ewa Niemczura from the Astronomical Institute of the University of Wrocław, for a valuable advice. PAD would like to thank Marcin Gawroński from the Astronomical Institute of the Nicolaus Copernicus University in Toruń for his valuable comments on the nature of ALS 9243. We would like to thank Kiril Stoyanov and Dragomir Marchev who provided us with spectra of ALS 9243 and Wojciech Dimitrow for insightful comments on the interpretation of these spectra. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. This research has also made use of the VizieR catalogue access tool, CDS, Strasbourg, France (DOI: 10.26093/cds/vizier). This work has made use of data from the European Space Agency (ESA) mission Gaia (https://www.cosmos.esa.int/gaia), processed by the Gaia Data Processing and Analysis Consortium (DPAC, https://www.cosmos.esa.int/web/gaia/dpac/consortium). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the Gaia Multilateral Agreement.

#### References

- Aldoretta, E. J., Caballero-Nieves, S. M., Gies, D. R., et al. 2015, AJ, 149, 26
- Ammons, S. M., Robinson, S. E., Strader, J., et al. 2006, ApJ, 638, 1004
- Anders, F., Khalatyan, A., Chiappini, C., et al. 2019, A&A, 628, A94
- Anderson, E. & Francis, C. 2012, Astronomy Letters, 38, 331
- Andrae, R., Fouesneau, M., Creevey, O., et al. 2018, A&A, 616, A8
- Avedisova, V. S. & Kondratenko, G. I. 1984, Nauchnye Informatsii, 56, 59
- Bailer-Jones, C. A. L. 2015a, A&A, 575, A35
- Bailer-Jones, C. A. L. 2015b, PASP, 127, 994
- Bailer-Jones, C. A. L. 2018, A&A, 609, A8
- Bailer-Jones, C. A. L., Rybizki, J., Andrae, R., & Fouesneau, M. 2018a, A&A, 616, A37
- Bailer-Jones, C. A. L., Rybizki, J., Fouesneau, M., Mantelet, G., & Andrae, R. 2018b, AJ, 156, 58
- Benedict, G., Henry, T., Franz, O., et al. 2016, AJ, 152, 141
- Blanco-Cuaresma, S. 2019, MNRAS, 486, 2075
- Blanco-Cuaresma, S., Soubiran, C., Heiter, U., & Jofré, P. 2014, A&A, 569, A111
- Bonavita, M., Desidera, S., Thalmann, C., et al. 2016, A&A, 593, A38
- Bond, H. E., Schaefer, G. H., Gilliland, R. L., et al. 2017, ApJ, 840, 70
- Bonev, T., Markov, H., Tomov, T., et al. 2017, Bulgarian Astronomical Journal, 26.67
- Butler, R. P., Vogt, S. S., Laughlin, G., et al. 2017, AJ, 153, 208
- Crampton, D. 1971, AJ, 76, 260
- Crampton, D. 1972, MNRAS, 158, 85
- Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, VizieR Online Data Catalog, II/246
- Deka-Szymankiewicz, B., Niedzielski, A., Adamczyk, M., et al. 2018, A&A, 615. A31
- Dybczyński, P. A. 2006, A&A, 449, 1233
- Dybczyński, P. A. & Berski, F. 2015, MNRAS, 449, 2459
- Dybczyński, P. A. & Królikowska, M. 2015, MNRAS, 448, 588
- Dybczyński, P. A., Królikowska, M., & Wysoczańska, R. 2019, arXiv e-prints, arXiv:1909.10952
- Eker, Z., Bakış, V., Bilir, S., et al. 2018, MNRAS, 479, 5491
- ESA. 1997, The HIPPARCOS and TYCHO catalogues, ESA HIPPARCOS Space Astrometry Mission (ESA Publications Division, ESA SP Series vol no: 1200, Noordwijk, Netherlands)
- Feng, F., Lisogorskyi, M., Jones, H. R. A., et al. 2019, ApJS, 244, 39
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, A&A, 616, A1
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., Prusti, T., et al. 2016, A&A, 595. A2
- García-Sánchez, J., Weissman, P. R., Preston, R. A., et al. 2001, A&A, 379, 634
- Garmany, C. D., Glaspey, J. W., Bragança, G. A., et al. 2015, AJ, 150, 41

- Gatewood, G. D. & Gatewood, C. V. 1978, ApJ, 225, 1530 Gatewood, G. D. & Gatewood, C. V. 1978, ApJ, 225, 1510 Georgelin, Y. M., Georgelin, Y. P., & Roux, S. 1973, A&A, 25, 337
- Gontcharov, G. A. 2006, Astronomy Letters, 32, 759
- Graham, J. A. 1971, AJ, 76, 1079
- Hardorp, J., Theile, I., & Voigt, H. H. 1965, Hamburger Sternw. Warner & Swasey Obs., C05, 0
- Høg, E., Fabricius, C., Makarov, V. V., et al. 2000, A&A, 355,
- Irrgang, A., Wilcox, B., Tucker, E., & Schiefelbein, L. 2013, Astronomy and Astrophysics, 549, A137
- Jiménez-Torres, J. J., Pichardo, B., Lake, G., & Throop, H. 2011, MNRAS, 418, 1272
- Kervella, P., Arenou, F., Mignard, F., & Thévenin, F. 2019, A&A, 623, A72

Kervella, P., Thévenin, F., & Lovis, C. 2017, A&A, 598, L7

- Królikowska, M. & Dybczyński, P. A. 2017, MNRAS, 472, 4634
- Lanz, T. & Hubeny, I. 2007, ApJS, 169, 83
- Lestrade, J., Preston, R. A., Jones, D. L., et al. 1999, A&A, 344, 1014
- Lindegren, L., Lammers, U., Bastian, U., et al. 2016, A&A, 595, A4
- Malaroda, S., Levato, H., & Galliani, S. 2006, yCat, III
- Mason, B. D., Wycoff, G. L., Hartkopf, W. I., Douglass, G. G., & Worley, C. E. 2001a, AJ, 122, 3466

Article number, page 11 of 12

- Mason, B. D., Wycoff, G. L., Hartkopf, W. I., Douglass, G. G., & Worley, C. E. 2001b, AJ, 122, 3466
- Mayer, P. & Macák, P. 1973, Bulletin of the Astronomical Institutes of Czechoslovakia, 24, 50
- Moffat, A. F. J., Fitzgerald, M. P., & Jackson, P. D. 1979, A&AS, 38, 197
- Muirhead, P. S., Dressing, C. D., Mann, A. W., et al. 2018, AJ, 155, 180 Nassau, J. J., Stephenson, C. B., & McConnell, D. J. 1965, Hamburger Sternw. Warner & Swasey Obs., C06, 0
- Neckel, T. & Staude, H. J. 1992, A&A, 254, 339
- Niedzielski, A., Deka-Szymankiewicz, B., Adamczyk, M., et al. 2016, A&A, 585, A73
- Pecaut, M. J. & Mamajek, E. E. 2013, ApJS, 208, 9
- Peterson, W. M., Mutel, R. L., Lestrade, J.-F., Güdel, M., & Goss, W. M. 2011, ApJ, 737, 104
- Pickles, A. & Depagne, É. 2010, PASP, 122, 1437
- Reed, B. C. 2003, AJ, 125, 2531
- Reed, C. 2005, VizieR Online Data Catalog, V/125
- Reuyl, D. 1936, AJ, 45, 133
- Ross, F. E. 1927, AJ, 37, 193
- Ségransan, D., Delfosse, X., Forveille, T., et al. 2000, A&A, 364, 665
- Stassun, K. G., Oelkers, R. J., Pepper, J., et al. 2018, ApJ, 156, 102
- Straižys, V. 1992, Multicolor stellar photometry (Pachart Pub. House, Tucson)
- Tokovinin, A. 2018, VizieR Online Data Catalog, J/ApJS/235/6
- van Leeuwen, F., ed. 2007, Astrophysics and Space Science Library, Vol. 350, Hipparcos, the New Reduction of the Raw Data
- Wielen, R., Schwan, H., Dettbarn, C., et al. 2001, Veroeffentlichungen des Astronomischen Rechen-Instituts Heidelberg, 40, 1
- Wilson, R. E. 1953, General Catalogue of Stellar Radial Velocities (Carnegie Institute, Washington D.C. Publication 601)
- Wysoczańska, R., Dybczyński, P. A., & Królikowska, M. 2020, MNRAS, 491, 2119