

STAFFANS-WEISS PERTURBATIONS FOR MAXIMAL L^p -REGULARITY IN BANACH SPACES

AHMED AMANSAG, HAMID BOUNTI, ABDERRAHIM DRIOUICH, AND SAID HADD

ABSTRACT. In this paper we show that the concept of maximal L^p -regularity is stable under a large class of unbounded perturbations, namely Staffans-Weiss perturbations. To that purpose, we first prove that the analyticity of semigroups is preserved under this class of perturbations, which is a necessary condition for the maximal regularity. In UMD spaces, \mathcal{R} -boundedness conditions are exploited to give conditions guaranteeing the maximal regularity. For non-reflexive Banach space, a condition is imposed to the Dirichlet operator associated to the boundary value problem to prove the maximal regularity. A Pde example illustrating the theory and an application to a class of non-autonomous perturbed boundary value problems are presented.

1. INTRODUCTION

In this paper we investigate the maximal L^p -regularity of evolution equations of the type

$$(1) \quad \begin{cases} \dot{z}(t) = A_m z(t) + Pz(t) + f(t), & t \geq 0, \\ z(0) = 0, \\ Gz(t) = Kz(t), & t \geq 0, \end{cases}$$

where $A_m : Z \subset X \rightarrow X$ is a linear closed operator in a Banach space X with domain $D(A_m) = Z$, $P : Z \rightarrow X$ is an additive linear perturbation of A_m , $G, K : Z \rightarrow U$ are linear boundary operators (U is another Banach space) and $f \in L^p(\mathbb{R}^+, X)$ with $p \geq 1$ is a real number. Actually, we assume that $A := A_m$ with domain $D(A) = \ker(G)$ is a generator of a strongly continuous semigroup $\mathbb{T} := (\mathbb{T}(t))_{t \geq 0}$ on X .

The concept of maximal regularity has been the subject of several works for many years, e.g. [11, 16, 12, 13], and the monograph [15]. The main purpose of these works is to give sufficient conditions on the operator A so as the problem (1) with $P \equiv 0$ and $K \equiv 0$, which can be written as

$$(2) \quad \dot{z} = Az + f, \quad z(0) = 0,$$

has a maximal L^p -regularity. A necessary condition for the maximal regularity of the evolution equation (2) is that A is a generator of an analytic semigroup. This condition is also sufficient only if X is a Hilbert space, see [13]. In the case of UMD Banach space X , Weis [43] introduced necessary and sufficient conditions in terms of \mathcal{R} -boundedness of the operator A for the maximal L^p -regularity of (2).

1991 *Mathematics Subject Classification.* 35K90, 47D06, 93C05, 93C73.

Key words and phrases. Maximal L^p -regularity, Unbounded perturbations, Boundary value problems, Operator semigroups, UMD spaces, Nonreflexive space.

Return now to our initial boundary problem (1). This later can be reformulated as

$$(3) \quad \dot{z} = (\mathcal{A} + P)z + f, \quad z(0) = 0,$$

where $\mathcal{A} : D(\mathcal{A}) \subset X \rightarrow X$ is the linear operator defined by

$$\mathcal{A} := A_m, \quad D(\mathcal{A}) = \{x \in Z : Gx = Kx\}.$$

To study maximal regularity of the problem (3), we first study maximal regularity of the problem

$$(4) \quad \dot{z} = \mathcal{A}z + f, \quad z(0) = 0.$$

In addition to our assumption at the beginning of this section, we also suppose that $G : Z \rightarrow U$ is surjective. Let then $\mathbb{D}_\lambda \in \mathcal{L}(U, Z)$ ($\lambda \in \rho(A)$) be the Dirichlet operator associated with A_m and G , see Section 3. In order to state our main results on well-posedness and maximal L^p -regularity of the problem (3), we select $B := (\lambda I - A_{-1})\mathbb{D}_\lambda \in \mathcal{L}(U, X_{-1})$ for $\lambda \in \rho(A)$, where X_{-1} is the extrapolation space associated to A and X , and $A_{-1} : X \rightarrow X_{-1}$ is the extension of A to X , which is a generator of a strongly continuous semigroup on X_{-1} . We assume that B is a p -admissible control operator for A , see the next section for the definition and notation. The case where K is bounded, i.e. $K \in \mathcal{L}(X, U)$, is studied in the recent paper [5]. In the present paper, we assume that the boundary operator K is unbounded, that is $K : Z \subsetneq X \rightarrow U$. This situation is quite difficult which needs additional assumption to treat the well-posedness and maximal L^p -regularity. According to [21], if we assume that $(A, B, K|_{D(A)})$ is regular on X, U, U with $I_U : U \rightarrow U$ as an admissible feedback, then the problem (4) is well-posed on the Banach space X . Moreover, if the problem (2) has a maximal L^p -regularity and $\|\lambda \mathbb{D}_\lambda\| \leq \kappa$ for any $\lambda > \lambda_0$, where $\lambda_0 \in \mathbb{R}$ and $\kappa > 0$ are constants, then the problem (4) has also the maximal L^p -regularity on a non reflexive Banach space X , see Theorem 4.8. On the other hand, we assume that $(A, B, P|_{D(A)})$ generates a regular linear system on X, U, X . Then, in Theorem 3.3, we prove that the problem (3) is well-posed. Corollary 2 shows that the problem (3) has the maximal L^p -regularity. If X is a UMD space then we use \mathcal{R} -boundedness to prove the maximal L^p -regularity for the evolution equation (3), see Theorem 4.6 and Corollary 2. We mention that in [22], the authors proved perturbation theorems for sectoriality and \mathcal{R} -sectoriality in general Banach spaces. They gives conditions on intermediate spaces Z and W such that, for an operator $S : Z \rightarrow W$ of small norm, the operator $A + S$ is sectorial (resp. \mathcal{R} -sectorial) provided A is sectorial (resp. \mathcal{R} -sectorial). Their results are obtained by factorizing $S = BC$. As \mathcal{R} -sectoriality implies maximal regularity in UMD spaces, these theorems yield to maximal regularity perturbation only in UMD spaces.

In the next section, we first recall the necessary material about feedback theory of infinite dimensional linear systems. We then use this theory to prove the well-posedness of the evolution equation (1) in Section 3. Our main results on maximal L^p -regularity for the problem (1) are gathered in Section 4.3 and Section 4.4. The last section is devoted to apply the obtained results to a class of non-autonomous boundary problems.

Notation. Hereafter $p, q \in [1, \infty]$ and $T > 0$ are real numbers such that $\frac{1}{p} + \frac{1}{q} = 1$. If X is a Banach space, we denote by $L^p([0, T]; X)$ the space of all X -valued

Bochner integrable functions. For any $\theta \in (0, \pi)$, Σ_θ is the following sector:

$$\Sigma_\theta := \{z \in \mathbb{C}^*; |\arg z| < \theta\}.$$

For any $\alpha \in \mathbb{R}$, the right half-plane is defined by

$$\mathbb{C}_\alpha =: \{z \in \mathbb{C}; \operatorname{Re} z > \alpha\}.$$

Given a semigroup $\mathbb{T} := (\mathbb{T}(t))_{t \geq 0}$ generated by an operator $A : D(A) \subset X \rightarrow X$, we will always denote by $\omega_0(\mathbb{T})$ (or $\omega_0(A)$) the growth bound of this semigroup. The resolvent set of A is denoted by $\rho(A)$. Preferably, we denote the resolvent operator of A by $R(\lambda, A) := (\lambda - A)^{-1}$ for any $\lambda \in \rho(A)$, where the notation $\lambda - A$ means $\lambda I - A$.

2. FEEDBACK THEORY OF INFINITE DIMENSIONAL LINEAR SYSTEMS

In this section, we gather definitions and results from feedback theory of infinite dimensional linear systems mainly developed in the references [39, 40, 41, 46]. We also give some new development of this theory. Hereafter, X and U are Banach spaces and $p \in [1, \infty[$.

It is known (see e.g. [39, 40]) that partial differential equations with boundary control and point observation can be reformulated as the following distributed linear system

$$(5) \quad \begin{cases} \dot{x}(t) = Ax(t) + Bu(t), & t \geq 0 \\ y(t) = Kx(t), \\ x(0) = x_0, \end{cases}$$

where $A : D(A) \subset Z \subset X \rightarrow X$ is the generator of a strongly continuous semigroup $\mathbb{T} := (\mathbb{T}(t))_{t \geq 0}$ on X with Z is a Banach space continuously and densely embedded in X , $B \in \mathcal{L}(U, X_{-1})$ is a control operator such that

$$R(\lambda, A_{-1})B \in \mathcal{L}(U, Z), \quad \lambda \in \rho(A),$$

and $K \in \mathcal{L}(Z, U)$ is an observation operator. Here X_{-1} is the completion of X with respect to the norm $\|R(\lambda, A) \cdot\|$. We recall that we can extend \mathbb{T} to another strongly continuous semigroup $\mathbb{T}_{-1} := (\mathbb{T}_{-1}(t))_{t \geq 0}$ on X_{-1} with generator $A_{-1} : X \rightarrow X_{-1}$, the extension of A to X (see e.g. [17, chap.2]). The mild solution of the system (5) is given by:

$$(6) \quad x(t) = \mathbb{T}(t)x_0 + \int_0^t \mathbb{T}_{-1}(t-s)Bu(s)ds \quad x_0 \in X,$$

where the integral is taken in X_{-1} . Formally, the well-posedness of the system (5) means that the state satisfies $x(t) \in X$ for any $t \geq 0$, the observation function y is extended to a locally p -integrable function $y \in L^p_{loc}([0, \infty), U)$ satisfying the following property: for any $\tau > 0$, there exists a constant $c_\tau > 0$ such that

$$\|y\|_{L^p([0, \tau], U)} \leq c_\tau (\|x_0\| + \|u\|_{L^p([0, \tau], U)}),$$

for any initial state $x_0 \in X$ and any control function $u \in L^p_{loc}([0, \infty), U)$. In order to mathematically explain this concept, let us define

$$C := Ki \in \mathcal{L}(D(A), U).$$

where i is the canonical injection from $D(A)$ to Z .

We also need the following definition.

Definition 2.1. (i) $B \in \mathcal{L}(U, X_{-1})$ is called p -admissible control operator for A , if there exists $t_0 > 0$ such that :

$$\Phi_{t_0} u := \int_0^{t_0} \mathbb{T}_{-1}(t-s)Bu(s)ds \in X$$

for any $u \in L_{loc}^p([0, \infty), U)$. We also say that (A, B) is p -admissible.

(ii) $C \in \mathcal{L}(D(A), Y)$ is called p -admissible observation operator for A , if there exist $\alpha > 0$ and $\kappa := \kappa_\alpha > 0$ such that:

$$(7) \quad \int_0^\alpha \|C\mathbb{T}(t)x\|_Y^p dt \leq \kappa^p \|x\|^p,$$

for all $x \in D(A)$. We also say that (C, A) is p -admissible.

Let us now describe some consequences of this definition. If B is p -admissible control operator for A , then by the closed graph theorem one can see that for any $t \geq 0$,

$$\Phi_t \in \mathcal{L}(L^p([0, t], U), X).$$

This implies that the state of the system (5) satisfies $x(t) = \mathbb{T}(t)x_0 + \Phi_t u \in X$ for any $t \geq 0$, $x_0 \in X$ and $u \in L_{loc}^p([0, \infty), U)$. According to [45], for all $0 < \tau_1 \leq \tau_2$,

$$(8) \quad \|\Phi_{\tau_1}\| \leq \|\Phi_{\tau_2}\|.$$

Now if C is p -admissible observation operator for A , then due to (7), the map $\Psi_\infty : D(A) \rightarrow L_{loc}^p([0, \infty), U)$ defined by $\Psi_\infty x := C\mathbb{T}(\cdot)x$, can be extended to a bounded operator $\Psi_\infty : X \rightarrow L_{loc}^p([0, \infty), U)$. For any $x \in X$ and $t \geq 0$, we define the family $\Psi_t x := \Psi_\infty x$ on $[0, t]$. Then for all $t \geq 0$,

$$\Psi_t \in \mathcal{L}(X, L_{loc}^p([0, \infty), U)).$$

On the other hand, let us consider the linear operator

$$D(C_\Lambda) := \left\{ x \in X : \lim_{s \rightarrow +\infty} sCR(s, A)x \text{ exists in } U \right\}$$

$$C_\Lambda x := \lim_{s \rightarrow +\infty} sCR(s, A)x.$$

Clearly, $D(A) \subset D(C_\Lambda)$ and $C_\Lambda = C$ on $D(A)$. This shows that C_Λ is in fact an extension of C , called the *Yosida* extension of C w.r.t. A . We note that if C is p -admissible for A , then $\text{Range}(\mathbb{T}(t)) \subset D(C_\Lambda)$ and $(\Psi_\infty x)(t) = C_\Lambda \mathbb{T}(t)x$ for any $x \in X$ and a.e. $t \geq 0$.

In the sequel, we assume that B and C are p -admissible for A and set

$$W_{0,loc}^{2,p}([0, \infty), U) := \left\{ u \in W_{loc}^{2,p}([0, \infty), U) : u(0) = 0 \right\}.$$

This space is dense in $L_{loc}^p([0, \infty), U)$. Remark that for any $u \in W_{0,loc}^{2,p}([0, \infty), U)$, $t \geq 0$ and by assuming $0 \in \rho(A)$ (without loss of generality) and using an integration by parts, we have

$$\Phi_t u = R(0, A_{-1})Bu(t) - R(0, A)\Phi_t \dot{u} \in Z.$$

On the other hand, using the fact that $KR(0, A_{-1})B \in \mathcal{L}(U)$, $CR(0, A) \in \mathcal{L}(X, U)$ and (8), the application $(t \mapsto K\Phi_t u) \in L_{loc}^p([0, \infty), U)$ for any $u \in W_{0,loc}^{2,p}([0, \infty), U)$. Thus we have defined an application

$$(9) \quad \mathbb{F}_\infty : W_{0,loc}^{2,p}([0, \infty), U) \rightarrow L_{loc}^p([0, \infty), U), \quad u \mapsto \mathbb{F}_\infty u = K\Phi.u.$$

Definition 2.2. [44] *Let B and C be p -admissible control and observation operators for A , respectively. We say that the triple (A, B, C) generates a well-posed linear system on X, U, U , if the operator \mathbb{F}_∞ defined by (9) satisfies the following property: For any $\alpha > 0$ there exists a constant $\vartheta_\alpha > 0$ such that for all $u \in W_{0,loc}^{2,p}([0, \infty), U)$,*

$$(10) \quad \|\mathbb{F}_\infty u\|_{L^p([0,\alpha],U)} \leq \vartheta_\alpha \|u\|_{L^p([0,\alpha],U)}.$$

The operator \mathbb{F}_∞ is called the extended input-output operator associated to (A, B, C) .

If the triple (A, B, C) generates a well-posed linear system on X, U, U , then we have two folds: first the state of (5) satisfies $x(t) \in X$ for all $t \geq 0$, and second \mathbb{F}_∞ has an extension $\mathbb{F}_\infty \in \mathcal{L}(L_{loc}^p([0, \infty), U))$, due to (10). Observe that the observation function y verifies

$$(11) \quad y(\cdot) := y(\cdot; x_0, u) = CT(\cdot)x_0 + \mathbb{F}_\infty u = (\Psi_\infty \mathbb{F}_\infty) \begin{pmatrix} x_0 \\ u \end{pmatrix},$$

for all $x_0 \in D(A)$ and $u \in W_{0,loc}^{2,p}([0, \infty), U)$. By density of $D(A) \times W_{0,loc}^{2,p}([0, \infty), U)$ in $X \times L_{loc}^p([0, \infty), U)$, the function y is extended to a function $y \in L_{loc}^p([0, \infty), U)$ such that

$$y = \Psi_\infty x_0 + \mathbb{F}_\infty u, \quad \forall (x_0, u) \in X \times L_{loc}^p([0, \infty), U).$$

We now turn out to give a representation of the observation function y in terms of the observation operator C and the state $x(\cdot)$. To that purpose Weiss [46, 47] introduced the following subclass of well-posed linear systems.

Definition 2.3. *Let the triple (A, B, C) generates a well-posed linear system on X, U, U with extended input-output operator \mathbb{F}_∞ . This system is called regular (with feedthrough $D = 0$) if :*

$$\lim_{\tau \rightarrow 0} \frac{1}{\tau} \int_0^\tau (\mathbb{F}_\infty u_{z_0})(s) ds = 0,$$

with $u_{z_0}(s) = z_0$ for all $s \geq 0$, is a constant control function.

According to Weiss [46, 47], if (A, B, C) generates a regular system on X, U, U , then the state and the observation function of the linear system (5) satisfy

$$(12) \quad x(t) \in D(C_\Lambda) \quad \text{and} \quad y(t) = C_\Lambda x(t),$$

for any initial state $x(0) = x_0 \in X$, any control function $u \in L^p([0, \infty), U)$ and a.e. $t \geq 0$.

Definition 2.4. *Let a triple (A, B, C) generates a well-posed system on X, U, U with extended input-output operator \mathbb{F}_∞ . Define*

$$\mathbb{F}_\tau u := \mathbb{F}_\infty u \quad \text{on} \quad [0, \tau].$$

The identity operator $I_U : U \rightarrow U$ is called an admissible feedback for Σ if the operator $I - \mathbb{F}_{t_0} : L^p([0, t_0], U) \rightarrow L^p([0, t_0], U)$ admits a (uniformly) bounded inverse for some $t_0 > 0$ (hence all $t_0 > 0$).

A consequence of Definition 2.4 is that the feedback law $u = y(\cdot; x_0, u)$ has a sense. In fact, due to (11) this is equivalent to $(I - \mathbb{F}_\tau)u = \Psi_\tau x_0$ on $[0, \tau]$. As $I - \mathbb{F}_\tau$ is invertible in $L^p([0, \tau], U)$, then the equation $u = y(\cdot; x_0, u)$ has a unique solution and this solution $u \in L^p([0, \tau], U)$ is given also by

$$u(t) = C_\Lambda x(t), \quad \text{a.e. } t \geq 0,$$

due to (12). Using (6), the state $x(\cdot)$ satisfies the following variation of constants formula

$$x(t) = \mathbb{T}(t)x_0 + \int_0^t \mathbb{T}_{-1}(t-s)BC_\Lambda x(s)ds,$$

for any $x_0 \in X$ and any $t \geq 0$. Now put set

$$\mathbb{T}^{cl}(t)x_0 := x(t), \quad t \geq 0.$$

Then by using the definition of C_0 -semigroups one can see that $(\mathbb{T}^{cl}(t))_{t \geq 0}$ is a C_0 -semigroup on X . More precisely, we have the following perturbation theorem due to Weiss [46] in Hilbert spaces and to Staffans [40, Chap.7] in Banach spaces.

Theorem 2.5. *Let (A, B, C) generates a regular linear system with the identity operator $I_U : U \rightarrow U$ an admissible feedback operator. Then the operator*

$$(13) \quad \begin{aligned} A^{cl} &:= A_{-1} + BC_\Lambda \\ D(A^{cl}) &:= \{x \in D(C_\Lambda); (A_{-1} + BC_\Lambda)x \in X\} \end{aligned}$$

generates a C_0 -semigroup $(\mathbb{T}^{cl}(t))_{t \geq 0}$ on X such that $\text{Range}(\mathbb{T}^{cl}(t)) \subset D(C_\Lambda)$ for a.e. $t > 0$, and for any $\alpha > 0$, there exists $c_\alpha > 0$ such that for all $x_0 \in X$,

$$(14) \quad \|C_\Lambda \mathbb{T}^{cl}(\cdot)x_0\|_{L^p([0, \alpha], U)} \leq c_\alpha \|x_0\|.$$

Moreover, this semigroup satisfies

$$(15) \quad \mathbb{T}^{cl}(t)x_0 = \mathbb{T}(t)x_0 + \int_0^t \mathbb{T}_{-1}(t-s)BC_\Lambda \mathbb{T}^{cl}(s)x_0 ds \quad x_0 \in X, \quad t \geq 0.$$

In addition (A^{cl}, B, C_Λ) generates a regular system.

Definition 2.6. *Let (A, B, C) generates a regular linear system on X, U, U with the identity operator $I_U : U \rightarrow U$ as an admissible feedback. The operator*

$$\mathbb{P}^{sw} := BC : D(C_\Lambda) \subset X \rightarrow X_{-1}$$

is called the **Staffans-Weiss perturbation** of A .

It is not difficult to see that if one of the operators B or C is bounded (i.e. $B \in \mathcal{L}(U, X)$ or $C \in \mathcal{L}(X, U)$) and the other is p -admissible then the triple (A, B, C) generates a regular linear system on X, U, U with the identity operator $I_U : U \rightarrow U$ as an admissible feedback. As application of the Staffans-Weiss theorem (Theorem 2.5), we distinct two subclasses of perturbations as follows:

Remark 1. (i) *We take $B \in \mathcal{L}(X, U)$ and $C \in \mathcal{L}(D(A), U)$ a p -admissible observation operator for A . According to Theorem 2.5, the operator $A^{cl} := A + BC$ with domain $D(A^{cl}) = D(A)$ is a generator of a strongly continuous semigroup $\mathbb{T}^{cl} := (\mathbb{T}^{cl}(t))_{t \geq 0}$ on X such that $\mathbb{T}^{cl}(t)X \subset D(C_\Lambda)$ for a.e. $t > 0$, the estimate (14) holds, and*

$$(16) \quad \mathbb{T}^{cl}(t)x = \mathbb{T}(t)x + \int_0^t \mathbb{T}(t-s)BC_\Lambda \mathbb{T}^{cl}(s)x ds, \quad t \geq 0, \quad x \in X.$$

On the other hand, it is shown in [19], that the semigroup \mathbb{T}^{cl} satisfy also the following formula

$$(17) \quad \mathbb{T}^{cl}(t)x = \mathbb{T}(t)x + \int_0^t \mathbb{T}^{cl}(t-s)BC_\Lambda \mathbb{T}(s)x ds, \quad t \geq 0, \quad x \in X.$$

Using Hölder inequality one can see that there exist $\alpha_0 > 0$ and $\gamma \in (0, 1)$ such that

$$\int_0^{\alpha_0} \|BCT(t)x\| dt \leq \gamma \|x\|,$$

for all $x \in D(A)$.

The following operator

$$\mathbb{P}^{mv} := BC : D(A) \rightarrow X$$

is called a **Miyadera-Voigt perturbation** for A ; (see e.g. [17, p.195]).

- (ii) We take $C \in \mathcal{L}(X, U)$ and $B \in \mathcal{L}(U, X_{-1})$ a p -admissible control operator for A . Then the part of the operator $A_{-1} + BC$ in X generates a strongly continuous semigroup on X satisfying all properties of Theorem 2.5. In this case the operator

$$\mathbb{P}^{ds} := BC : X \rightarrow X_{-1}$$

is called **Desch-Schappacher perturbation** for A (see e.g. [17, p.182]).

3. WELL-POSEDNESS OF PERTURBED BOUNDARY VALUE PROBLEMS

The object of this section is to investigate the well-posedness of the perturbed boundary value problem defined by (1). We first rewrite (1) as non-homogeneous perturbed Cauchy problem of the form (3). Then the well-posedness of (1) can be obtained if for example the operator

$$(18) \quad \mathcal{A} := A_m, \quad D(\mathcal{A}) = \{x \in Z : Gx = Kx\},$$

generates a strongly continuous semigroup on X and that P is a p -admissible observation operator for A (see Remark 1 (i)). Recently, the authors of [21] introduced conditions on A_m, G and K for which \mathcal{A} is a generator. To be more precise, assume that

(H1) $G : Z \rightarrow U$ is onto,

(H2) The operator defined by $A := A_m$ and $D(A) := \ker(G)$, generates a C_0 -semigroup $(\mathbb{T}(t))_{t \geq 0}$ on X .

According to Greiner [18], these conditions imply that for any $\lambda \in \rho(A)$ the restriction of G to $\ker(\lambda - A_m)$ is invertible. We then define

$$\mathbb{D}_\lambda := (G|_{\ker(\lambda - A_m)})^{-1} \in \mathcal{L}(U, X), \quad \lambda \in \rho(A).$$

This operator is called the *Dirichlet operator*. Define the operators :

$$(19) \quad \begin{aligned} B &:= (\lambda - A_{-1})\mathbb{D}_\lambda \in \mathcal{L}(U, X_{-1}), \\ C &:= Ki \in \mathcal{L}(D(A), U), \end{aligned}$$

where i is the canonical injection from $D(A)$ to Z .

Using the resolvent equation, and the fact that for all $\lambda, \mu \in \rho(A)$ (see [18, Lemma 1.3])

$$(20) \quad \mathbb{D}_\lambda = (I - (\lambda - \mu)R(\mu, A))\mathbb{D}_\mu,$$

it is easy to see that the operator B does not depend on λ .

In the rest of this paper, C_Λ denotes the Yosida extension of C with respect to A .

It is shown in [21, Lemma.3.6] that if A, B, C as above and if (A, B, C) generates a regular linear system on X, U, U , then we have

$$(21) \quad Z \subset D(C_\Lambda) \quad \text{and} \quad (C_\Lambda)|_Z = K.$$

If H is the transfer function of this system and $\alpha > \omega_0(A)$ then

$$(22) \quad H(\lambda) = C_\Lambda R(\lambda, A_{-1})B = C_\Lambda \mathbb{D}_\lambda = K \mathbb{D}_\lambda,$$

for any $\lambda \in \mathbb{C}$ with $\text{Re}\lambda > \alpha$. Moreover, we have

$$(23) \quad \lim_{s \rightarrow +\infty} H(s) = 0.$$

We also assume the following

(H3) the triple (A, B, C) generates a regular linear system on X, U, U with $I_U : U \rightarrow U$ as an admissible feedback operator.

We have the following perturbation theorem (see [21] for the proof).

Theorem 3.1. *Let assumptions (H1) to (H3) be satisfied. Then, the following assertions hold:*

(i) *The operator $(\mathcal{A}, D(\mathcal{A}))$ defined by (18) coincides with the following operator*

$$A^{cl} := A_{-1} + BC_\Lambda, \quad D(A^{cl}) = \{x \in D(C_\Lambda) : (A_{-1} + BC_\Lambda)x \in X\}.$$

(ii) *The operator $(\mathcal{A}, D(\mathcal{A}))$ generates a strongly continuous semigroup $(\mathbb{T}^{cl}(t))_{t \geq 0}$ on X as in Theorem 2.5.*

(iii) *For any $\lambda \in \rho(A)$ we have*

$$\lambda \in \rho(\mathcal{A}) \Leftrightarrow 1 \in \rho(\mathbb{D}_\lambda K) \Leftrightarrow 1 \in \rho(K \mathbb{D}_\lambda).$$

(iv) *Finally for $\lambda \in \rho(A) \cap \rho(\mathcal{A})$:*

$$R(\lambda, \mathcal{A}) = (I - \mathbb{D}_\lambda K)^{-1} R(\lambda, A).$$

Under the assumptions of Theorem 3.1, the mild solution of the problem (4) is given by

$$(24) \quad z(t) = \mathbb{T}^{cl}(t)x + \int_0^t \mathbb{T}^{cl}(t-s)f(s)ds,$$

for any $t \geq 0$, $x \in X$ and $f \in L^p(\mathbb{R}^+, X)$.

Before giving another useful expression of z in term of the initial semigroup \mathbb{T} , we need the following very useful result proved in [19, prop.3.3].

Lemma 3.2. *Let $(\mathbb{S}(t))_{t \geq 0}$ be a strongly continuous semigroup on X with generator $(\mathbb{G}, D(\mathbb{G}))$. Let $\Upsilon \in \mathcal{L}(D(\mathbb{G}), X)$ be a p -admissible observation operator for \mathbb{G} . Denote by Υ_Λ the Yosida extension of Υ with respect to \mathbb{G} . Then*

$$(\mathbb{S} * f)(t) := \int_0^t \mathbb{S}(t-s)f(s)ds \in D(\Upsilon_\Lambda), \quad \text{a.e. } t \geq 0,$$

$$\|\Upsilon_\Lambda(\mathbb{S} * f)\|_{L^p([0, \alpha], X)} \leq c(\alpha)\|f\|_{L^p([0, \alpha], X)},$$

for $\alpha > 0$, $f \in L^p_{loc}([0, \infty), X)$ and a constant $c(\alpha)$ independent of f such that $c(\alpha) \rightarrow 0$ as $\alpha \rightarrow 0$.

Proposition 1. *Let assumptions of Theorem 3.1 be satisfied. Let $\alpha > 0$ and $f \in L^p(\mathbb{R}^+, X)$. The non-homogenous Cauchy problem (4) is well-posed and its mild solution satisfies for any initial condition $x \in X$,*

$$(25) \quad \begin{aligned} z(t) &\in D(C_\Lambda) \quad \text{a.e. } t > 0, \\ \|C_\Lambda z(\cdot)\|_{L^p([0, \alpha], X)} &\leq c(\alpha) (\|x\| + \|f\|_{L^p([0, \alpha], X)}), \\ z(t) &= \mathbb{T}(t)x + \int_0^t \mathbb{T}_{-1}(t-s)BC_\Lambda z(s)ds + \int_0^t \mathbb{T}(t-s)f(s)ds, \quad t \geq 0, \end{aligned}$$

where $c(\alpha) > 0$ is a constant independent of f .

Proof. By virtue of Theorem 3.1, \mathbb{T}^{cl} the semigroup generated by \mathcal{A} and let $z : [0, +\infty) \rightarrow X$ be the mild solution of the problem (4) given by (24). According to Theorem 2.5, we know that C_Λ is an admissible observation operator for \mathcal{A} . We denote by $C_{\Lambda, \mathcal{A}}$ the Yosida extension of C_Λ with respect to \mathcal{A} . Then $D(C_{\Lambda, \mathcal{A}}) \subset D(C_\Lambda)$ and $C_{\Lambda, \mathcal{A}} = C_\Lambda$ on $D(C_{\Lambda, \mathcal{A}})$. In fact, let $x \in D(C_{\Lambda, \mathcal{A}})$ and $s > 0$ sufficiently large. Then by first taking Laplace transform on both sides of (15) and second applying sC_Λ , we obtain

$$(26) \quad sC_\Lambda R(s, \mathcal{A})x = sCR(s, A) + H(s)C_\Lambda sR(s, \mathcal{A})x,$$

where we have used (22). Remark that

$$\|H(s)C_\Lambda sR(s, \mathcal{A})x\| \leq \|H(s)\| (\|C_\Lambda sR(s, \mathcal{A})x - C_{\Lambda, \mathcal{A}}x\| + \|C_{\Lambda, \mathcal{A}}x\|).$$

Hence, by (23) and the fact that $x \in D(C_{\Lambda, \mathcal{A}})$, we obtain

$$\lim_{s \rightarrow +\infty} H(s)C_\Lambda sR(s, \mathcal{A})x = 0.$$

Now from (26), we deduce that $x \in D(C_\Lambda)$ and $C_{\Lambda, \mathcal{A}}x = C_\Lambda x$. Let $x \in X$, $\alpha > 0$ and $f \in L^p([0, \alpha], X)$. The fact that C_Λ is p -admissible for \mathcal{A} , then by using (24) and Lemma 3.2, we obtain $z(t) \in D(C_{\Lambda, \mathcal{A}})$ for a.e. $t > 0$. This shows that $z(t) \in D(C_\Lambda)$ and $C_\Lambda z(t) = C_{\Lambda, \mathcal{A}}z(t)$ for a.e. $t > 0$. The estimation in (25) follows immediately from (14) and Lemma 3.2. Let us prove the last property in (25). By density there exists $(f_n)_n \subset \mathcal{C}([0, \alpha], D(\mathcal{A}))$ such that $f_n \rightarrow f$ in $L^p([0, \alpha], X)$ as $n \rightarrow \infty$. We put

$$(27) \quad z_n(t) = \mathbb{T}^{cl}(t)x + \int_0^t \mathbb{T}^{cl}(t-s)f_n(s)ds, \quad t \geq 0.$$

Using Hölder inequality, it is clear that $\|z_n(t) - z(t)\| \rightarrow 0$ as $n \rightarrow \infty$. Now let us prove that z_n satisfies the third assertion in (25). In fact, the estimation in (25) implies that

$$\|C_\Lambda z_n(\cdot) - C_\Lambda z(\cdot)\|_{L^p([0, \alpha], X)} \leq c_\alpha \|f_n - f\|_{L^p([0, \alpha], X)} \xrightarrow{n \rightarrow \infty} 0.$$

On the other hand, using the expression of the semigroup \mathbb{T}^{cl} given in (15), change of variable and Fubini theorem we obtain

$$(28) \quad \begin{aligned} z_n(t) &= \mathbb{T}^{cl}(t)x + \int_0^t \mathbb{T}(t-s)f_n(s)ds + \int_0^t \mathbb{T}_{-1}(t-\tau)B \int_0^\tau C_\Lambda \mathbb{T}^{cl}(\tau-s)f_n(s)dsd\tau \\ &= \mathbb{T}(t)x + \int_0^t \mathbb{T}(t-s)f_n(s)ds \\ &\quad + \int_0^t \mathbb{T}_{-1}(t-\tau)B \left(C_\Lambda \mathbb{T}^{cl}(\tau)x + \int_0^\tau C_\Lambda \mathbb{T}^{cl}(\tau-s)f_n(s)ds \right) d\tau. \end{aligned}$$

For simplicity we assume that $0 \in \rho(\mathcal{A})$. Then we have

$$\begin{aligned} C_\Lambda \int_0^\tau \mathbb{T}^{\text{cl}}(\tau - s) f_n(s) ds &= C_\Lambda (-\mathcal{A})^{-1} (-\mathcal{A}) \int_0^\tau \mathbb{T}^{\text{cl}}(\tau - s) f_n(s) ds \\ &= C_\Lambda (-\mathcal{A})^{-1} \int_0^\tau \mathbb{T}^{\text{cl}}(\tau - s) (-\mathcal{A}) f_n(s) ds \\ &= \int_0^\tau C_\Lambda (-\mathcal{A})^{-1} \mathbb{T}^{\text{cl}}(\tau - s) (-\mathcal{A}) f_n(s) ds \\ &= \int_0^\tau C_\Lambda \mathbb{T}^{\text{cl}}(\tau - s) f_n(s) ds. \end{aligned}$$

Now replacing this in (28), and using (27), we obtain

$$z_n(t) = \mathbb{T}(t)x + \int_0^t \mathbb{T}_{-1}(t-s) BC_\Lambda z(s) ds + \int_0^t \mathbb{T}(t-s) f_n(s) ds, \quad t \geq 0.$$

Put

$$\varphi(t) = \mathbb{T}(t)x + \int_0^t \mathbb{T}_{-1}(t-s) BC_\Lambda z(s) ds + \int_0^t \mathbb{T}(t-s) f(s) ds, \quad t \geq 0.$$

Then for any $t \in [0, \alpha]$, we have

$$\|z_n(t) - \varphi(t)\| \leq \gamma_\alpha (\|C_\Lambda z_n(\cdot) - C_\Lambda z(\cdot)\|_{L^p([0, \alpha], X)} + \|f_n - f\|_{L^p([0, \alpha], X)}),$$

due to the admissibility of B for A and Hölder inequality. This shows that $\|z_n(t) - \varphi(t)\| \rightarrow 0$ as $n \rightarrow \infty$, and hence $z = \varphi$. \square

Now we can state the main result of this section.

Theorem 3.3. *Let assumptions of Theorem 3.1 be satisfied. In addition, let $P : Z \rightarrow X$ such that (A, B, \mathbb{P}) generates a regular linear system on X, U, X , where $\mathbb{P} = Pi$. The following assertions hold:*

- (i) *The operator $P \in \mathcal{L}(D(\mathcal{A}), X)$ is a p -admissible observation operator for \mathcal{A} , hence the operator $(\mathcal{A} + P, D(\mathcal{A}))$ generates a strongly continuous semigroup on X .*
- (ii) *The boundary problem (1) is well-posed and its mild solution $z : [0, +\infty) \rightarrow X$ satisfies:*

$$\begin{aligned} z(t) &\in D(\mathbb{P}_\Lambda) \quad \text{a.e. } t \geq 0, \\ z(t) &= \mathbb{T}^{\text{cl}}(t)x + \int_0^t \mathbb{T}^{\text{cl}}(t-s) (\mathbb{P}_\Lambda z(s) + f(s)) ds, \end{aligned}$$

for any $t \geq 0$, initial condition $x \in X$ and $f \in L^p([0, \infty), X)$, where \mathbb{P}_Λ is the Yosida extension of P w.r.t \mathcal{A} .

Proof. (i) We first remark from (21) that $Z \subset D(\mathbb{P}_{0, \Lambda})$ and $P = \mathbb{P}_{0, \Lambda}$ on Z , where $\mathbb{P}_{0, \Lambda}$ denotes the Yosida extension of \mathbb{P} w.r.t. \mathcal{A} . Let $x \in D(\mathcal{A})$ and $\alpha > 0$. The facts that (A, B, \mathbb{P}) is regular and using (14), we obtain

$$(29) \quad \begin{aligned} &\int_0^t \mathbb{T}_{-1}(t-s) BC_\Lambda \mathbb{T}^{\text{cl}}(s)x \in D(\mathbb{P}_{0, \Lambda}) \quad \text{a.e. } t \geq 0, \text{ and} \\ &\left\| \mathbb{P}_{0, \Lambda} \int_0^\cdot \mathbb{T}_{-1}(t-s) BC_\Lambda \mathbb{T}^{\text{cl}}(s)x \right\|_{L^p([0, \alpha], X)} \leq \beta(\alpha) \|x\|, \end{aligned}$$

for some constant $\beta(\alpha) > 0$ depending only on α .
On the other hand, appealing to (15), we obtain

$$\begin{aligned} P\mathbb{T}^{cl}(t)x &= \mathbb{P}_{0,\Lambda}\mathbb{T}^{cl}(t)x \\ &= \mathbb{P}_{0,\Lambda}\mathbb{T}(t)x + \mathbb{P}_{0,\Lambda}\int_0^t \mathbb{T}_{-1}(t-s)BC_\Lambda\mathbb{T}^{cl}(s)x. \end{aligned}$$

Combining (29) and the p -admissibility of \mathbb{P} for A yield the p -admissibility of P for \mathcal{A} . According to Remark 1 (i), the operator $(\mathcal{A} + P, D(\mathcal{A}))$ generates a strongly continuous semigroup on X . The assertion (ii) follows from [19, thm.5.1] \square

4. PERTURBATION THEOREMS FOR MAXIMAL REGULARITY

4.1. Maximal regularity. Let $\mathbb{G} : D(\mathbb{G}) \subset X \rightarrow X$ be the generator of a strongly continuous semigroup $\mathbb{S} := (\mathbb{S}(t))_{t \geq 0}$ on a Banach space X . Consider the following non-homogeneous abstract Cauchy problem:

$$(30) \quad \begin{cases} \dot{z}(t) = \mathbb{G}z(t) + f(t), & 0 < t \leq T \\ z(0) = 0, \end{cases}$$

where $f : [0, T] \rightarrow X$ a measurable function.

Definition 4.1. We say that the operator \mathbb{G} (or the problem (30)) has the maximal L^p -regularity on the interval $[0, T]$, and we write $\mathbb{G} \in \mathcal{MR}_p(0, T; X)$, if for all $f \in L^p([0, T], X)$, there exists a unique $z \in W^{1,p}([0, T], X) \cap L^p([0, T], D(\mathbb{G}))$ which verifies (30).

By "maximal" we mean that the applications f , $\mathbb{G}z$ and z have the same regularity. Due to the closed graph theorem, if $\mathbb{G} \in \mathcal{MR}_p(0, T; X)$ then

$$(31) \quad \|\dot{z}\|_{L^p([0, T], X)} + \|z\|_{L^p([0, T], X)} + \|\mathbb{G}z\|_{L^p([0, T], X)} \leq C\|f\|_{L^p([0, T], X)},$$

for a constant $C > 0$ independent of f .

It is well-known that a necessary condition for the maximal L^p -regularity is that \mathbb{G} generates an analytic semigroup. According to De Simon [13] this condition is also sufficient if X is a Hilbert space. On the other hand, it is shown in [16] that if $\mathbb{G} \in \mathcal{MR}_p(0, T; X)$ for one $p \in [1, \infty]$ then $\mathbb{G} \in \mathcal{MR}_q(0, T; X)$ for all $q \in]1, \infty[$. Moreover if $\mathbb{G} \in \mathcal{MR}_p(0, T; X)$ for one $T > 0$, then $\mathbb{G} \in \mathcal{MR}_p(0, T'; X)$ for all $T' > 0$. Hence we simply write $\mathbb{G} \in \mathcal{MR}(0, T; X)$.

Remark 2. (i) Let $\mathcal{C}([0, T]; D(\mathbb{G}))$ be the space of all continuous functions from $[0, T]$ to $D(\mathbb{G})$, which is a dense space of $L^p([0, T]; X)$. It is well-known (see [42] (2.a) or [29] 1.5)) that \mathbb{G} has maximal L^p -regularity on $[0, T]$ if and only if $(\mathbb{S}(t))_{t \geq 0}$ is analytic and the operator \mathcal{R} defined by

$$(32) \quad (\mathcal{R}f)(t) := \mathbb{G} \int_0^t \mathbb{S}(t-s)f(s)ds \quad f \in \mathcal{C}([0, T], D(\mathbb{G})),$$

extends to a bounded operator on $L^p([0, T]; X)$. As we will see in our main results, this characterization is very useful if one works in general Banach spaces.

(ii) It is known (see [16]) that if $\mathbb{G} \in \mathcal{MR}(0, T; X)$ then for every $\lambda \in \mathbb{C}$, $\mathbb{G} + \lambda \in \mathcal{MR}(0, T; X)$, hence without loss of generality, we will assume through this paper that our generators satisfy $\omega_0(\mathbb{G}) < 0$.

In order to recall another characterization of maximal regularity, we need some preliminary definitions.

Definition 4.2. We say that a Banach space X is a UMD-space if for some (hence all) $p \in (1, \infty)$, $\mathcal{H} \in \mathcal{L}(L^p(\mathbb{R}, X))$ where

$$(\mathcal{H}f)(t) = \frac{1}{\pi} \lim_{\epsilon \rightarrow 0} \int_{|s| > \epsilon} \frac{f(t-s)}{s} ds, \quad t \in \mathbb{R}, \quad f \in \mathcal{S}(\mathbb{R}, X),$$

where $\mathcal{S}(\mathbb{R}, X)$ is the Schwartz space.

Classical UMD-spaces are Hilbert spaces and L^p -spaces, where $p \in (1, \infty)$. It is to be noted that every UMD-space is a reflexive space (see [2]).

Definition 4.3. A set $\tau \subset \mathcal{L}(X, Y)$ is called \mathcal{R} -bounded if there is a constant $C > 0$ such that for all $n \in \mathbb{N}$, $T_1, \dots, T_n \in \tau$, $x_1, \dots, x_n \in X$,

$$\int_0^1 \left\| \sum_{j=1}^n r_j(s) T_j x_j \right\|_Y ds \leq C \int_0^1 \left\| \sum_{j=1}^n r_j(s) x_j \right\|_X ds,$$

where $(r_j)_{j \geq 1}$ is a sequence of independent $\{-1; 1\}$ -valued random variables on $[0, 1]$ (e.g. Rademacher variables).

Remark 3. Here we give examples of \mathcal{R} -bounded sets. We let A be the generator of a bounded analytic semigroup on a Banach space X , $B : U \rightarrow X_{-1}$ and $C : D(A) \subset X \rightarrow U$ are linear bounded operators, where U is another (boundary) Banach space. We assume that (A, B, C) generates a regular linear system on X, U, U with transfer function

$$H(\lambda) := C_\Lambda R(\lambda, A_{-1})B \in \mathcal{L}(U), \quad \lambda \in \rho(A),$$

where C_Λ is the Yosida extension of C with respect to A , see Section 2. It has been observed in [23, p.513] that the set $\{H(is) : s \neq 0\}$ is \mathcal{R} -bounded.

The following remarkable result is due to Weis [43]

Theorem 4.4. Let \mathbb{G} be the generator of a bounded analytic semigroup in a UMD-space X . Then \mathbb{G} has maximal L^p -regularity for some (hence all) $p \in (1, \infty)$ if and only if the set $\{sR(is, \mathbb{G}); s \neq 0\}$ is \mathcal{R} -bounded.

4.2. Robustness of analyticity under Staffans-Weiss perturbation. In this part, we study the analyticity of the perturbed semigroup $(\mathbb{T}^{cl}(t))_{t \geq 0}$ defined in (3.1). We then assume, as in Section 3, that **(H1)**, **(H2)** and **(H3)** are satisfied.

Theorem 4.5. Let assumptions **(H1)** to **(H3)** be satisfied. Then the operator $(\mathcal{A}, D(\mathcal{A}))$ defined by (18) generates a strongly continuous semigroup which is analytic whenever the semigroup generated by A is.

Proof. According to Theorem 3.1 (i) \mathcal{A} is a generator of a strongly continuous semigroup $\mathbb{T}^{cl} := (\mathbb{T}^{cl}(t))_{t \geq 0}$ on X . Now assume that A generates an analytic semigroup \mathbb{T} on X . Then by [38, Theorem 12.31], there exist constants $\beta \in \mathbb{R}$ and $M_1 > 0$ such that $\mathbb{C}_\beta \subset \rho(A)$ and

$$(33) \quad \|R(\lambda, A)\| \leq \frac{M_1}{|\lambda - \beta|}, \quad \lambda \in \mathbb{C}_\beta.$$

In [31, Lemma 2.3], it is proved that the admissibility of C (and B by the same method) for A , imply that there exists $\gamma \in]\frac{\pi}{2}, \pi[$ such that

$$(34) \quad \sup_{z \in \Sigma_\gamma} |z|^{\frac{1}{q}} \|CR(z, A)\| < +\infty \quad \text{and} \quad \sup_{z \in \Sigma_\gamma} |z|^{\frac{1}{p}} \|R(z, A_{-1})B\| < +\infty,$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Put

(35)

$$M_2 := \sup_{z \in \mathbb{C}_0} |z|^{\frac{1}{q}} \|CR(z, A)\| < +\infty \quad \text{and} \quad M_3 := \sup_{z \in \mathbb{C}_0} |z|^{\frac{1}{p}} \|R(z, A_{-1})B\| < +\infty,$$

Let $\omega_1 := \max\{\omega_0(A), \omega_0(\mathcal{A})\}$. Due to Theorem 3.1 (ii), for any $\lambda \in \mathbb{C}_{\omega_1}$, we have

$$\begin{aligned} R(\lambda, \mathcal{A}) &= (I - \mathbb{D}_\lambda K)^{-1} R(\lambda, A) \\ (36) \quad &= R(\lambda, A) + R(\lambda, A_{-1})B(I_U - C_\Lambda \mathbb{D}_\lambda)^{-1} CR(\lambda, A). \end{aligned}$$

On the other hand, $(I_U - C_\Lambda \mathbb{D}_\lambda)^{-1} = I + \mathcal{H}^{cl}(\lambda)$, where \mathcal{H}^{cl} is the transfer function of the (closed-loop) regular linear system generated by $(\mathcal{A}, B, C_\Lambda)$. Hence there exists $\alpha > \omega_0(\mathcal{A})$ such that

$$(37) \quad \nu := \sup_{\mathbb{C}_\alpha} \|(I_U - C_\Lambda \mathbb{D}_\lambda)^{-1}\| < \infty.$$

Now let $\omega_2 := \max\{0, \alpha, \beta, \omega_1\}$. Combining (33), (35), (36) and (37), we obtain

$$\|R(\lambda, \mathcal{A})\| \leq \frac{\tilde{M}}{|\lambda - \omega_2|},$$

for all $\lambda \in \mathbb{C}_{\omega_2}$, where $\tilde{M} = M_1 + \nu M_2 M_3$. Finally, by virtue of [38, Theorem 12.31] the semigroup generated by \mathcal{A} is analytic. \square

4.3. Staffans-Weiss perturbation for maximal regularity in UMD spaces.

We are still working under the setting of Section 3 and we will study the maximal regularity of the operator \mathcal{A} in the case of UMD spaces.

Theorem 4.6. *Let conditions (H1) to (H3) be satisfied with X, U be UMD-spaces and A generates a bounded analytic semigroup. Assume that there exists $\omega > \omega_1$ such that the sets $\{s^{\frac{1}{p}} R(\omega + is, A_{-1})B; s \neq 0\}$ and $\{s^{\frac{1}{q}} CR(\omega + is, A); s \neq 0\}$ are \mathcal{R} -bounded. If $A \in \mathcal{MR}(0, T; X)$ then $\mathcal{A} \in \mathcal{MR}(0, T; X)$.*

Proof. Assume that $A \in \mathcal{MR}(0, T; X)$ and let $\omega > \omega_1$ such that the sets $\{s^{\frac{1}{p}} R(\omega + is, A_{-1})B; s \neq 0\}$ and $\{s^{\frac{1}{q}} CR(\omega + is, A); s \neq 0\}$ are \mathcal{R} -bounded. Denote $\mathcal{A}^\omega := -\omega + \mathcal{A}$ and $A^\omega := -\omega + A$ with domains $D(\mathcal{A}^\omega) = D(\mathcal{A})$ and $D(A^\omega) = D(A)$, respectively. These operators are generators of analytic semigroups on X . We first observe that $A^\omega \in \mathcal{MR}(0, T; X)$. To prove our theorem it suffice to show that $\mathcal{A}^\omega \in \mathcal{MR}(0, T; X)$. Clearly, $\omega_0(A^\omega) = \omega_0(A) - \omega < 0$ and $\omega_0(\mathcal{A}^\omega) = \omega_0(\mathcal{A}) - \omega < 0$, so that $i\mathbb{R} \setminus \{0\} \subset \rho(A^\omega) \cap \rho(\mathcal{A}^\omega)$. It is not difficult to show that (A^ω, B, C) is also a regular linear system on X, U, U with the identity operator $I_U : U \rightarrow U$ as an admissible feedback. Now according to Theorem 2.5, the following operator

$$A^{cl, \omega} := A_{-1}^\omega + BC_\Lambda, \quad D(A^{cl, \omega}) = \{x \in D(C_\Lambda) : (A_{-1}^\omega + BC_\Lambda)x \in X\}.$$

As $A_{-1}^\omega = A_{-1} - \omega$, then $D(A^{cl, \omega}) = D(\mathcal{A})$, and $A^{cl, \omega} = \mathcal{A}^\omega$ due to Theorem 3.1 (i). As in (36) we have

(38)

$$\begin{aligned} sR(is, \mathcal{A}^\omega) &= sR(is, A^\omega) + s^{\frac{1}{p}} R(is, A_{-1}^\omega)B(I - H^\omega(is))^{-1} s^{\frac{1}{q}} CR(is, A^\omega) \\ &= sR(is, A^\omega) + s^{\frac{1}{p}} R(\omega + is, A_{-1})B(I - H^\omega(is))^{-1} s^{\frac{1}{q}} CR(\omega + is, A), \end{aligned}$$

where $H^\omega(\lambda) = C_\Lambda R(\lambda, A_{-1}^\omega)B$, $\lambda \in \rho(A)$, is the transfer function of the regular linear system generated by (A^ω, B, C) . Using the assumptions, the equation (38)

and Theorem 4.4, it suffice only to show that the set $\{(I - H^\omega(is))^{-1} : s \neq 0\}$ is \mathcal{R} -bounded. In fact, by Theorem 3.1 (i) and the condition **(H3)** the triple operator $(\mathcal{A}^\omega, B, C_\Lambda)$ generates a regular linear system and its associated transfer function is

$$H^{cl,\omega}(is) = (I - H^\omega(is))^{-1} H^\omega(is), \quad s \neq 0,$$

which implies that

$$(I - H^\omega(is))^{-1} = I_U + H^{cl,\omega}(is), \quad s \neq 0.$$

According to Remark 3, the set $\{H^{cl,\omega}(is) : s \neq 0\}$ is \mathcal{R} -bounded. Hence $\{(I - H^\omega(is))^{-1} : s \neq 0\}$ is \mathcal{R} -bounded. This ends the proof. \square

4.3.1. *Example: PDE with a boundary conditions.* Let $p, q \in (1, \infty)$ such that $\frac{1}{p} + \frac{1}{q} = 1$ and $1 < p < 3$. In this section, we deal with the following problem:

$$(39) \quad \begin{cases} \frac{\partial w(t, s)}{\partial t} = \frac{\partial^2 w(t, s)}{\partial s^2} + f(t) & t \in [0, T] \quad s \in (0, 1), \\ \frac{\partial w(t, 1)}{\partial s} = w(t, 1) - w(t, 0), \\ \frac{\partial w(t, 0)}{\partial s} = 0, \\ w(0, s) = 0. \end{cases}$$

Let $X = L^p(0, 1)$ and $A_m g := g''$ with domain $Z = \{g \in W^{2,p}(0, 1), g'(0) = 0\}$. We define on Z the operator:

$$\begin{aligned} G : Z &\rightarrow \mathbb{R} \\ g &\mapsto g'(0). \end{aligned}$$

We also define the unbounded operator $K : Z \rightarrow \mathbb{R}$ by $Kg := g(1) - g(0)$. One can see that under this setting, the problem (39) can be transformed to the following :

$$\begin{cases} \dot{x}(t) = A_m x(t), & 0 \leq t \leq T, \quad x(0) = 0 \\ Gx(t) = Kx(t), & 0 \leq t \leq T. \end{cases}$$

Define the operator:

$$A = A_m, \quad D(A) = KerG = \{g \in W^{2,p}(0, 1), g'(0) = 0 \text{ and } g'(1) = 0\}.$$

It is well-known that the operator A has maximal L^p - regularity.

We set:

$$\mathbb{D}_\lambda := (G|_{ker(\lambda - A_m)})^{-1}, \quad B := (\lambda - A_{-1})\mathbb{D}_\lambda, \quad C := Ki$$

Now the main result is the following

Theorem 4.7. *The operator A^{cl} defined by :*

$$A^{cl}g = g'', \quad D(A^{cl}) := \{g \in W^{2,p}(0, 1), g'(0) = 0 \text{ and } g'(1) = g(1) - g(0)\}$$

has maximal L^r -regularity for every $r > 1$ and the problem (39) has a unique solution $w \in W^{1,r}([0, T]; X) \cap L^r([0, T]; D(A^{cl}))$ such that:

$$\|\dot{w}\|_{L^r([0, T]; X)} + \|w\|_{L^r([0, T]; X)} + \|w''\|_{L^r([0, T]; X)} \leq C\|f\|_{L^r([0, T]; X)},$$

for some constant $C > 0$ anf $f \in L^r([0, T]; X)$.

Proof. We will show that all conditions of Theorem 4.6 are satisfied. According to [1, Theorem 2.4], to show that the triple (A, B, C) generates a regular linear system, all we have to show is that there exist $\beta \in [0, 1]$ and $\gamma \in (0, 1]$ such that :

- (i) $\text{Range}(\mathbb{D}_\lambda) \subset F_{1-\beta}^A$ (where F_α^A is the Favard space of A of order α , see [17])
- (ii) $D[(\lambda - A)^\gamma] \subset D(C_\lambda)$
- (iii) $\beta + \gamma < 1$.

Let us prove the above three propositions:

- (i) By simple calculation, one can see that there exists $\lambda_0 > 0$ such that

$$\sup_{\lambda > \lambda_0} \|\lambda^{\frac{p+1}{2p}} \mathbb{D}_\lambda\| < +\infty.$$

This fact implies (see [1, Lemma A.1]) that $\text{Range}(\mathbb{D}_\lambda) \subset F_{\frac{p+1}{2p}}^A$ for some $\lambda \in \rho(A)$.

We take $\beta = 1 - \frac{p+1}{2p} = \frac{p-1}{2p}$.

- (ii) According to [1, Lemma A.2], we have to show that for $\alpha \in (0, 1)$ and every $\rho \geq \rho_0 > 0$ we have

$$|Cg| \leq M(\rho^\alpha \|g\|_p + \rho^{\alpha-1} \|g''\|_p) \quad g \in D(A),$$

for some constant $M > 0$.

For $g \in D(A)$ we know that

$$g(1) = g(0) + \int_0^1 g'(s) ds$$

then we can easily show that :

$$|g(1) - g(0)| \leq \|g'\|_p.$$

By [17, Example III.2.2], we have for $\epsilon > 0$

$$\|g'\|_p \leq \frac{\rho}{\epsilon} \|g\|_p + \epsilon \|g''\|_p.$$

By taking $\rho = \epsilon^{-3}$, $\alpha = \frac{1}{3}$ and $\gamma \in (\frac{1}{3}, \frac{1}{p})$, we show the assertion.

- (iii) Clearly we have $\beta + \gamma < 1$

Hence, [1, Theorem 2.4] implies that the operators B and C are r -admissible for $\frac{2p}{p+1} < r < \frac{1}{\gamma}$ and the triple (A, B, C) generates a regular system and the operator identity is an admissible feedback. For instance we can take $r = 2$.

To show that the sets $\{s^{\frac{1}{2}}CR(is, A); s \neq 0\}$ and $\{s^{\frac{1}{2}}R(is, A_{-1})B; s \neq 0\}$ are \mathcal{R} -bounded, it is sufficient to show that C and B are l -admissible (l -admissibility is more general than admissibility, see [23] for definitions), namely, it is sufficient to show that the sets $\{\sqrt{s}CR(s, A); s > 0\}$ and $\{\sqrt{s}R(s, A_{-1})B; s > 0\}$ are \mathcal{R} -bounded (see [23], page 514), which is equivalent to show that the sets $\{\sqrt{s}CR(s+1, A); s > 0\}$ and $\{\sqrt{s}R(s+1, A_{-1})B; s > 0\}$ are \mathcal{R} -bounded

In order to prove that $\{\sqrt{s}CR(s+1, A); s > 0\}$ is \mathcal{R} -bounded, we follow the example in [23] (page 528) and the technique used there to show the \mathcal{R} -boundedness of the above set, namely it suffices to find a space \tilde{Z} such that $D(A) \subset \tilde{Z} \subset X$ and C is bounded in $\|\cdot\|_{\tilde{Z} \rightarrow \mathbb{R}}$ and the set $\{\sqrt{s}R(s+1, A); s > 0\}$ is \mathcal{R} -bounded in $\mathcal{L}(\tilde{Z}, X)$, this holds, by the same example, for $\tilde{Z} = W^{1,p}(0, 1)$ since C is bounded in $\mathcal{L}(\tilde{Z}, \mathbb{R})$.

Now to show the \mathcal{R} -boundedness of $\{\sqrt{s}R(s+1, A_{-1})B; s > 0\}$ we follow the same example, we show that the set $\{\sqrt{s}B^*R(s+1, A^*); s > 0\}$ is \mathcal{R} -bounded. Let

first determine B^* , we proceed again as in [23] and multiplying the first equation in (39) with a fixed $v \in C^\infty([0, 1])$ and integrating by parts we obtain:

$$\langle w'(t, \cdot), v \rangle_{(0,1)} = \langle w(t, \cdot), v'' \rangle_{(0,1)} + Gw(t, \cdot) \delta_1 v,$$

this means that $B^* = \delta_1$. By the same argument used to show that $\{\sqrt{s}CR(s+1, A); s > 0\}$ is \mathcal{R} -bounded, we show that $\{\sqrt{s}B^*R(s+1, A^*); s > 0\}$ is \mathcal{R} -bounded.

All hypothesis of the theorem are satisfied, then the operator A^{cl} defined by :

$$A^{cl}g = g'', \quad D(A^{cl}) := \{g \in W^{2,p}(0, 1), g'(0) = 0 \quad \text{and} \quad g'(1) = g(1) - g(0)\}$$

has maximal L^2 -regularity, thus maximal L^r -regularity for every $r > 1$ and then problem (39) has unique solution $u \in W^{1,r}([0, T]; X) \cap L^r([0, T]; D(A^{cl}))$ such that:

$$\begin{aligned} & \left[\int_0^T \left(\int_0^1 \left\| \frac{\partial w(t, s)}{\partial t} \right\|^p ds \right)^{r/p} dt \right]^{1/r} + \left[\int_0^T \left(\int_0^1 \|w(t, s)\|^p ds \right)^{r/p} dt \right]^{1/r} \\ & + \left[\int_0^T \left(\int_0^1 \left\| \frac{\partial^2 w(t, s)}{\partial s^2} \right\|^p ds \right)^{r/p} dt \right]^{1/r} \\ & \leq C \left[\int_0^T \left(\int_0^1 \|f(t, s)\|^p ds \right)^{r/p} dt \right]^{1/r} \end{aligned}$$

for some constant $C > 0$. □

Remark 4. Besides Hilbert spaces there is another interesting situation, where bounded sets of operators are automatically \mathcal{R} -bounded. For example if F is a Banach space of type 2 and E is a Banach space of cotype 2, then every bounded set $\tau \subset \mathcal{L}(E, F)$ is \mathcal{R} -bounded (see. [14]). Thanks to (34), for all $\omega > \omega_0(A)$ the sets $\{s^{\frac{1}{2}}CR(\omega + is, A); s \neq 0\}$ and $\{s^{\frac{1}{2}}R(\omega + is, A_{-1})B; s \neq 0\}$ are bounded. So, to prove their \mathcal{R} -boundedness for the previous example, we distinguish two cases (case $p = 2$ is trivial):

- (i) If $2 < p < 3$, then $X = L^p(0, 1)$ is of cotype 2, hence the set $\{s^{\frac{1}{2}}CR(is, A); s \neq 0\}$ is \mathcal{R} -bounded since $U = \mathbb{R}$ is of type 2 and it suffices only to show \mathcal{R} -boundedness of $\{s^{\frac{1}{2}}R(\omega + is, A_{-1})B; s \neq 0\}$
- (ii) If $1 < p < 2$, then $X = L^p(0, 1)$ is of type 2, hence the set $\{s^{\frac{1}{2}}R(\omega + is, A_{-1})B; s \neq 0\}$ is \mathcal{R} -bounded since $U = \mathbb{R}$ is of cotype 2 and it suffices that only to show that $\{s^{\frac{1}{2}}CR(is, A); s \neq 0\}$ is \mathcal{R} -bounded.

4.4. Staffans-Weiss perturbation for maximal regularity in non-reflexive spaces. In this section, we assume that X is a non-reflexive Banach space. The following result shows the maximal regularity of the perturbed boundary value problem (1) in the case of $P \equiv 0$.

Theorem 4.8. Let assumptions (H1) to (H3) be satisfied. Assume additionally that there exists $\lambda_0 \in \mathbb{R}$ such that

$$(40) \quad \kappa_0 := \sup_{\lambda > \lambda_0} \|\lambda \mathbb{D}_\lambda\| < +\infty.$$

If $A \in \mathcal{MR}(0, T; X)$ then $\mathcal{A} \in \mathcal{MR}(0, T; X)$.

Proof. As $A \in \mathcal{MR}(0, T; X)$, there exists $\mathcal{R} \in \mathcal{L}(L^p([0, T], X))$ such that

$$(\mathcal{R}f)(t) = A \int_0^t \mathbb{T}(t-s)f(s)ds,$$

for all $f \in L^p([0, T], X)$ and a.e $t \geq 0$. By Theorem 4.5, \mathcal{A} generates an analytic semigroup \mathbb{T}^{cl} on X . We then can define the following operator

$$(\mathcal{R}^{cl}f)(t) := \mathcal{A} \int_0^t \mathbb{T}^{cl}(t-s)f(s)ds \quad f \in C([0, T]; D(\mathcal{A})).$$

Our objective is to show that the operator \mathcal{R}^{cl} admits a bounded extension on $L^p([0, T]; X)$. Let us consider the Yosida approximation operators of \mathcal{A} by

$$\mathcal{A}_n := nAR(n, \mathcal{A}) = n^2R(n, \mathcal{A}) - nI,$$

for any $n \in \mathbb{N}$ such that $n > \omega_0(\mathcal{A})$.

From (36) and for any sufficiently large integer n , one can write

$$(41) \quad \mathcal{A}_n = nAR(n, A) + n^2\mathbb{D}_n(I - C_\Lambda\mathbb{D}_n)^{-1}CR(n, A) \quad .$$

Consider the following operators

$$(\mathcal{R}_n^{cl}f)(t) := \mathcal{A}_n \int_0^t \mathbb{T}^{cl}(t-s)f(s)ds,$$

for $f \in C([0, T]; D(\mathcal{A}))$ and $n \in \mathbb{N}$ such that $n > \omega_0(\mathcal{A})$.

We have (see [17]),

$$(\mathcal{R}^{cl}f)(t) = \lim_{n \rightarrow \infty} (\mathcal{R}_n^{cl}f)(t),$$

for every $t \in [0, T]$.

Using Proposition 1, we obtain

$$(\mathcal{R}_n^{cl}f)(t) = \mathcal{A}_n \left(\int_0^t \mathbb{T}(t-s)f(s)ds + \int_0^t \mathbb{T}_{-1}(t-s)C_\Lambda z(s)ds \right),$$

where

$$z(t) = \int_0^t \mathbb{T}^{cl}(t-s)f(s)ds, \quad t \geq 0.$$

Using (41), for large n ,

$$\begin{aligned} (\mathcal{R}_n^{cl}f)(t) &= nAR(n, A) \int_0^t \mathbb{T}(t-s)f(s)ds \\ &\quad + n^2\mathbb{D}_n(I - C_\Lambda\mathbb{D}_n)^{-1}CR(n, A) \int_0^t \mathbb{T}(t-s)f(s)ds \\ &\quad + nAR(n, A) \int_0^t \mathbb{T}_{-1}(t-s)BC_\Lambda z(s)ds \\ &\quad + n^2\mathbb{D}_n(I - C_\Lambda\mathbb{D}_n)^{-1}CR(n, A) \int_0^t \mathbb{T}_{-1}(t-s)BC_\Lambda z(s)ds \\ &:= I_n^1(t) + I_n^2(t) + I_n^3(t) + I_n^4(t). \end{aligned}$$

Let us estimate separately the four terms on the left of the above equality. Maximal regularity of A yields

$$\begin{aligned} \int_0^T \|I_n^1(t)\|^p dt &= \|\mathcal{R}(nR(n, A)f)\|_{L^p([0, T], X)}^p \\ &\leq c_T \|f\|_{L^p([0, T], X)}^p, \end{aligned}$$

for a constant $c_T > 0$ independent of f .

On the other hand, combining (37), (40) and Lemma 3.2 we obtain

$$\begin{aligned} \int_0^T \|I_n^2(t)\|^p dt &= \int_0^T \left\| n\mathbb{D}_n(I - C_\wedge \mathbb{D}_n)^{-1} C nR(n, A) \int_0^t \mathbb{T}(t-s)f(s)ds \right\|^p dt \\ &\leq (\kappa\nu)^p \int_0^T \left\| C \int_0^t \mathbb{T}(t-s)nR(n, A)f(s)ds \right\|^p dt \\ &\leq (\kappa\nu\gamma_T)^p \|f\|_{L^p}^p := c_1(T, p) \|f\|_{L^p}^p, \end{aligned}$$

where $c_1(T, p)$ is a constant depending on T and p .

Now we estimate $I_n^3(t)$

$$\begin{aligned} \int_0^T \|I_n^3(t)\|^p dt &= \int_0^T \left\| A \int_0^t \mathbb{T}(t-s)n\mathbb{D}_n C_\Lambda z(s)ds \right\|^p dt \\ &= \|\mathcal{R}(n\mathbb{D}_n C_\Lambda z(\cdot))\|_{L^p([0, T], X)}^p \\ &\leq c_T \kappa_0^p \|C_\Lambda z(\cdot)\|_{L^p([0, T], U)}^p \\ &\leq c_2(T, p) \|f\|_{L^p}^p, \end{aligned}$$

due to (40) and Proposition 1, where $c_2(T, p) > 0$ is a constant depending only on T and p .

Similarly, we have

$$\begin{aligned} \int_0^T \|I_n^4(t)\|^p dt &= \int_0^T \left\| n\mathbb{D}_n(I - C_\Lambda \mathbb{D}_n)^{-1} C \int_0^t \mathbb{T}(t-s)nR(n, A)BC_\Lambda z(s)ds \right\|^p dt \\ &\leq (\nu\kappa_0)^p \int_0^T \left\| C \int_0^t \mathbb{T}(t-s)n\mathbb{D}_n C_\Lambda z(s)ds \right\|^p dt \\ &\leq c_3(T, p) \|f\|_{L^p}^p, \end{aligned}$$

by (37), (40), Lemma 3.2 and Proposition 1, where $c_3(T, p) > 0$ is a constant depending only on T and p . Finally one can conclude that

$$\int_0^T \|(\mathcal{R}_n^{cl} f)(t)\|^p dt \leq \tilde{C}_p \|f\|_{L^p}^p,$$

for some constant $C(\tilde{T}, p) > 0$ depending only on T and p .

The fact that $\|(\mathcal{R}_n^{cl} f)(t)\|^p \rightarrow \|(\mathcal{R}^{cl} f)(t)\|^p$ for all $t \in [0, T]$, Fatou's Lemma implies

$$\begin{aligned} \int_0^T \|(\mathcal{R}^{cl} f)(t)\|^p dt &\leq \liminf_{n \rightarrow \infty} \int_0^T \|(\mathcal{R}_n^{cl} f)(t)\|^p dt \\ &\leq \tilde{C}_p \|f\|_{L^p}^p. \end{aligned}$$

Thus \mathcal{R}^{cl} can be extended to a bounded operator on $L^p([0, T]; X)$. \square

Remark 5. *In the case of Desch-Schappacher perturbation (bounded perturbation at the boundary) we have obtained in [5, Theorem] the maximal regularity without appealing to the condition (40). However for Staffans-Weiss perturbation (unbounded perturbation at the boundary), our proof of maximal regularity is essentially based on this condition. We mention that (40) cannot take place if we work in a reflexive Banach space. In fact, this estimation can be expressed as $\sup_{\lambda > \lambda_0} \|\lambda R(\lambda, A_{-1})B\| < +\infty$*

which is equivalent to $\text{Range}(B) \subset F_1^{A-1}$ (see [32, Remark 10], [25]). It is important to recall that the control operator B is strictly unbounded, i.e. $\text{Range}(B) \cap X = \{0\}$,

since it comes from the boundary (see [39]). So, if X is reflexive then $F_1^{A-1} = X$ and consequently $\text{Range}(B) \subset X$ which is contradictory since B is not identically nul. Thus all these facts force us to work with a non-reflexive Banach space

Now we state the previous theorem in the case of X being a GT -space (e.g. if $X = L^1$ or $X = C(K)$), which is a non-reflexive space. Since we are not concerned in this paper with GT -spaces and H^∞ -calculus, we omit definitions: all details can be found in [27].

Corollary 1. *Let assumptions of Theorem 4.8 be satisfied. Let X or X^* be a GT -space. If A is an H^∞ -sectorial operator on X with $\omega_H(A) < \frac{\pi}{2}$ then $\mathcal{A} \in \mathcal{MR}(0, T; X)$.*

Proof. According the Theorem 7.5 in [27], if A is an H^∞ -sectorial operator on X with $\omega_H(A) < \frac{\pi}{2}$ then A has maximal L^p -regularity for all $1 < p < \infty$, which implies by Theorem 4.8 that $A^{cl} \in \mathcal{MR}(0, T; X)$. \square

We end this section with the following result given the maximal L^p -regularity for the evolution equation (1) (or equivalently (3)).

Corollary 2. *Let conditions (H1) to (H3) be satisfied on Banach spaces X, U and $A \in \mathcal{MR}(0, T; X)$. Moreover, we assume that (A, B, \mathbb{P}) generates a regular linear system on X, U, X with $\mathbb{P} \in \mathcal{L}(D(A), X)$ is the restriction of $P : Z \rightarrow X$ on $D(A)$. Then the operator $\mathcal{A} + P \in \mathcal{MR}(0, T; X)$ (or equivalently the evolution equation (1) has maximal L^p -regularity) if one of the following conditions hold:*

- (i) X is a non reflexive Banach space and there exists $\lambda_0 \in \mathbb{R}$ such that

$$\sup_{\lambda > \lambda_0} \|\lambda \mathbb{D}_\lambda\| < +\infty.$$

- (ii) X and U are UMD spaces and the sets $\{s^{\frac{1}{p}} R(\omega + is, A_{-1})B; s \neq 0\}$ and $\{s^{\frac{1}{q}} CR(\omega + is, A); s \neq 0\}$ are \mathcal{R} -bounded for $\omega > \omega_1$.

Proof. First of all the operator P is a p -admissible observation operator for \mathcal{A} , due to Theorem 3.3. Thanks to Theorem 4.8 (resp. Theorem 4.6), the assertion (i) (resp. (ii)) implies that \mathcal{A} has maximal regularity. Finally, appealing to [5, Theorem 3] we obtain $\mathcal{A} + P \in \mathcal{MR}(0, T; X)$ which completes the proof. \square

5. APPLICATIONS TO NON-AUTONOMOUS PROBLEM

The object of this section is to apply our obtained results to prove maximal regularity of some classes of non-autonomous problem. First, we present the definition of maximal L^p -regularity in the non-autonomous case.

Let X be a Banach space and $\{A(t), t \in [0, T]\}$ a family of closed linear operators on X . For a measurable function $f : [0, T] \rightarrow X$, we consider the non-autonomous Cauchy problem:

$$(NCP) \quad \begin{cases} \dot{x}(t) = A(t)x(t) + f(t), & t \in [0, T] \\ x(0) = 0. \end{cases}$$

Definition 5.1. *For $p \in (1, \infty)$, we say that $\{A(t), t \in [0, T]\}$ or (NCP) has L^p -maximal regularity if for all $f \in L^p([0, T], X)$, there exists a unique solution of (NCP) $x \in W^{1,p}([0, T], X)$ such that $t \mapsto A(t)x(t) \in L^p([0, T], X)$.*

Maximal L^p -regularity in the non-autonomous case is quite different and less understood. However, several results have been established. We review some of them separating between the case where $D(A(t))$ does not depend on t and the case where $D(A(t))$ is varying with respect to t . In the first case, that of $D(A(t)) = D(A(0))$, Prüss and Schnaubelt [37] and Amann [3] proved that (NCP) has L^p -maximal regularity under the conditions that $t \mapsto A(t)$ is continuous and $A(t)$ has L^p -maximal regularity for every $t \in [0, T]$. Arendt et al. [6] generalized this result to relative continuous function $t \mapsto A(t)$. When the operators $A(t)$ have time-dependent domains and X is Hilbert space, Hieber and Monniaux [24] showed that (NCP) has L^p -maximal regularity whenever the family $\{A(t), t \in [0, T]\}$ satisfies the so-called Acquistapace Terreni conditions and every $A(t)$ has L^p -maximal regularity. This result was extended by Portal and Štrkalj [34] to a UMD spaces by using the concept of \mathcal{R} -boundedness.

The following theorem, due to Prüss and Schnaubelt [37], provides a characterization of maximal regularity by means of continuity.

Theorem 5.2. *Let $A(t)$, $t \in [0, T]$, generates an analytic semigroup of negative type on X , $D(A(t)) = D(A(0)) =: X_1$, and $A(\cdot) \in C([0, T], \mathcal{L}(X_1, X))$, where X_1 is endowed with the graph norm of $A(0)$. Assume additionally that for every $t \in [0, T]$, $A(t)$ has maximal L^p -regularity. Then $\{A(t), t \in [0, T]\}$ has maximal L^p -regularity on X .*

In the following, let X , Z and U be Banach spaces such that $Z \subset X$ with dense and continuous embedding. We consider the non-autonomous problem

$$\begin{cases} \dot{z}(t) = A_m(t)z(t) + f(t), & t \in [0, T], \\ z(0) = 0, \\ Gz(t) = Kz(t), & t \in [0, T], \end{cases}$$

where $A_m(t) : Z \subset X \rightarrow X$, $t \in [0, T]$, are linear closed operators in a Banach space X with domain $D(A_m(t)) := Z$, $G, K : Z \rightarrow U$ are linear boundary operators and $f \in L^p(\mathbb{R}^+, X)$ with $p \geq 1$ is a real number. Define the operator

$$\mathcal{A}(t) := A_m(t) \text{ with } D(\mathcal{A}(t)) := \{x \in Z; Gx = Kx\} =: \mathcal{D}.$$

We define the non-autonomous *Dirichlet map*

$$\mathbb{D}_\lambda(t) := (G|_{\text{Ker}(\lambda - A_m(t))})^{-1} \in \mathcal{L}(U, X), \quad \lambda \in \rho(A(t)),$$

and the λ -independent operators

$$(42) \quad B(t) := (\lambda - A_{-1}(t))\mathbb{D}_\lambda(t) \in \mathcal{L}(U, X_{-1}),$$

and

$$C := Ki,$$

where $i : D \rightarrow Z$ is the canonical injection from D to Z .

The following proposition asserts that continuity is invariant to cross boundary perturbation.

Proposition 2. *Let assumptions (H1) to (H3) be satisfied for every $t \in [0, T]$ by the operators $A(t), B(t)$ and C . Assume additionally (H4): There is $\mu_0 > \omega_0(A(0))$ such that $\text{Ker}(\mu_0 - A_m(t))$ does not depend on t . If $A(\cdot) \in C([0, T], \mathcal{L}(D, X))$ then $\mathcal{A}(\cdot) \in C([0, T], \mathcal{L}(D, X))$.*

Remark 6. (i) Assumption **(H4)** implies that $\mathbb{D}_{\mu_0}(t)$ does not depend on t and we set $\mathbb{D}_{\mu_0} = \mathbb{D}_{\mu_0}(t)$ for all $t \in [0, T]$. Hence, from the discussion above (ii) As for autonomous case, the definition of $B(t)$ does not depend on λ , so we can consider $B(t) = (\mu_0 - A_{-1}(t))\mathbb{D}_{\mu_0}$ for all $t \in [0, T]$. Thanks to Theorem 3.1, we have

$$\mathcal{A}(t) = A_{-1}(t) + B(t)C_\wedge \quad \text{for all } t \in [0, T].$$

By injecting $B(t)$ into above, we obtain

$$\mathcal{A}(t) - \mu_0 = A_{-1}(t) - \mu_0 + (\mu_0 - A_{-1}(t))\mathbb{D}_{\mu_0}C_\wedge = (A_{-1}(t) - \mu_0)(I - \mathbb{D}_{\mu_0}C_\wedge).$$

A simple calculation leads to $x \in D(\mathcal{A}(t)) \Leftrightarrow (I - \mathbb{D}_{\mu_0}C_\wedge)x \in D(A(t))$. Hence

$$(43) \quad \mathcal{A}(t) - \mu_0 = (A(t) - \mu_0)(I - \mathbb{D}_{\mu_0}C_\wedge).$$

Proof. (of Proposition 2) As consequences of the continuity of $A(\cdot)$ on $[0, T]$ (see [37] page 409): the type of the semigroup generated by $A(t)$ does not depend on t and the graph norms of the operators $A(t)$ are uniformly equivalent (and so we equip D with the norm $\|\cdot\|_D = \|(\mu_0 - A(t))\cdot\|$). Thus, the completions of X with respect to the norms $\|\cdot\|_{-1}^{A(t)} := \|R(\lambda, A(t))\cdot\|$ with $t \in [0, T]$ and $\lambda \in \rho(A(t))$ are constant and denoted simply X_{-1} . Denote by $A_{-1}(t)$ the extension of $A(t)$ to X_{-1} . Assumptions **(H1)**, **(H2)** and **(H3)** and Theorem 3.1 imply that for all $t \in [0, T]$ the operator $\mathcal{A}(t)$ generates a strongly continuous semigroup on X . Then we equip \mathcal{D} with the norm $\|\cdot\|_{\mathcal{D}} := \|(\mu_0 - \mathcal{A}(0))\cdot\|$. Put $\mathcal{P}_{\mu_0} := I - \mathbb{D}_{\mu_0}C_\wedge$ and define on D the norm $\|\cdot\|_D := \|(\mu_0 - A(0))\cdot\|$. By virtue of (43) for all $x \in \mathcal{D}$ we have

$$\begin{aligned} \|x\|_{\mathcal{D}} &= \|(\mu_0 - \mathcal{A}(0))x\| \\ &= \|(\mu_0 - A(0))\mathcal{P}_{\mu_0}x\| \\ &= \|\mathcal{P}_{\mu_0}x\|_D. \end{aligned}$$

On the other hand, for every $t, s \in [0, T]$ we have

$$\begin{aligned} \|\mathcal{A}(t) - \mathcal{A}(s)\|_{\mathcal{L}(\mathcal{D}, X)} &= \sup_{\|x\|_{\mathcal{D}}=1} \|\mathcal{A}(t)x - \mathcal{A}(s)x\| \\ &= \sup_{\|x\|_{\mathcal{D}}=1} \|(A(t) - A(s))\mathcal{P}_{\mu_0}x\| \\ &= \sup_{\|\mathcal{P}_{\mu_0}x\|_D=1} \|(A(t) - A(s))\mathcal{P}_{\mu_0}x\| \\ &\leq \sup_{\|z\|_D=1} \|(A(t) - A(s))z\| \\ &= \|A(t) - A(s)\|_{\mathcal{L}(D, X)}. \end{aligned}$$

Thus, the continuity of $\mathcal{A}(\cdot)$ can be deduced from the one of $A(\cdot)$. \square

Similarly, the continuity of $\mathcal{A}(\cdot)$ on $[0, T]$ (see [37] page 409) yields the graph norms of the operators $\mathcal{A}(t)$ are uniformly equivalent. In particular the type of the semigroup generated by $\mathcal{A}(t)$ does not depend on t .

Now, we present our perturbation result for the non-autonomous maximal regularity.

Theorem 5.3. *Let assumptions of Proposition 2 be satisfied. If for every $t \in [0, T]$, $A(t)$ has maximal L^p -regularity and $A(\cdot) \in C([0, T], \mathcal{L}(D, X))$. Then the family $\{\mathcal{A}(t), t \in [0, T]\}$ has maximal L^p -regularity if one of the following conditions holds:*

- (i) X is a non reflexive Banach space and for every $t \in [0, T]$, there exists $\lambda_0 \in \mathbb{R}$ such that

$$\sup_{\lambda > \lambda_0} \|\lambda \mathbb{D}_\lambda(t)\| < +\infty.$$

- (ii) X and U are UMD spaces, the sets

$$\{s^{\frac{1}{p}}R(\omega + is, A_{-1}(t))B(t); s \neq 0\} \text{ and } \{s^{\frac{1}{q}}CR(\omega + is, A(t)); s \neq 0\}, t \in [0, T],$$

are \mathcal{R} -bounded for some $\omega > \max\{\omega_0(A(0)), \omega_0(\mathcal{A}(0))\}$.

Proof. Thanks to Proposition 2, we have $\mathcal{A}(\cdot) \in C([0, T], \mathcal{L}(D, X))$. If condition (i) (resp. (ii)) is satisfied then by virtue of Theorem 4.8 (resp. Theorem 4.6) for the operator $\mathcal{A}(t)$ has maximal L^p -regularity every $t \in [0, T]$. Applying Theorem 5.2, we obtain the desired result. \square

Remark 7. *Let A_m, A, \mathcal{A}, B and C be as in Section 4 and let assumptions of Theorem 4.6 be satisfied. We define the operators $A_m(t) = a(t)A_m$, $A(t) = a(t)A$ and $\mathcal{A}(t) = a(t)\mathcal{A}$ where $a \in C([0, T])$ with $a(t) > \alpha > 0$ for all $t \in [0, T]$. Then clearly, all assumptions of Theorem 5.3 are verified. Moreover, condition (ii) is also satisfied and then the family $\{\mathcal{A}(t), t \in [0, T]\}$ has maximal L^p -regularity. So, Theorem 5.3 asserts the maximal regularity of the non-autonomous version associated to our example considered in Section 4.3.1.*

REFERENCES

- [1] M. Adler, M. Bombieri, and K.-J. Engel, *Perturbation of analytic semigroups and applications to partial differential equations*, J. Evol. Equ. (2016).
- [2] H. Amann, *Linear and Quasilinear Parabolic Problems*. Vol. I. Birkhuser, 1995.
- [3] H. Amann, Maximal regularity for nonautonomous evolution equations, *Adv. Nonlinear Stud.* 4 (2004) 417–430.
- [4] H. Amann, M. Hieber, G. Simonett : *Bounded H^∞ -calculus for elliptic operators*. Differential Integral Equations 7 (1994), no. 3-4, 613-653.
- [5] A. Amansag, H. Bounit, A. Driouich, S. Hadd. On the maximal regularity for perturbed autonomous and non-autonomous evolution equations, J. Evol. Equ. (2019). doi.org/10.1007/s00028-019-00514-8
- [6] W. Arendt, R. Chill, S. Fornaro and C. Poupaud. L^p -maximal regularity for nonautonomous evolution equations. *J. Differ. Equations*, 237:1–26, 2007.
- [7] T. Bárta, *Analytic solutions of Volterra equations via semigroups*, Semigroup Forum 76, no. 1, 2008, 142 - 148.
- [8] T. Bárta, *On R -sectorial derivatives on Bergman spaces*, Bull. Austral. Math. Soc. 77, 2008, 305 - 313.
- [9] T. Bárta, (2008) *Smooth solutions of Volterra equations via semigroups*. Bulletin of the Australian Mathematical Society, 78(2), 249-260. doi:10.1017/S0004972708000683
- [10] H. Bounit, A. Driouich, O. El-Mennaoui, *A direct approach to the weighted admissibility of observation operators for bounded analytic semigroups*. Semigroup Forum (2011) 83: 52. https://doi.org/10.1007/s00233-011-9315-9
- [11] P. Cannarsa and V. Vespri, *On maximal L^p regularity for the abstract Cauchy problem*, Boll. Un. Mat. Ital. B (6) 5 (1986), 165 - 175.
- [12] T. Coulhon and D. Lamberton, *Régularité L^p pour les équations d'évolution*, Séminaire d'Analyse Fonctionnelle 1984/1985, Publ. Math. Univ. Paris VII 26 (1986), 155-165.
- [13] L. De Simon, *Un'applicazione della teoria degli integrali singolari allo studio delle equazioni differenziali lineari astratte del primo ordine*, Rend. Sem. Mat., Univ. Padova 99 (1964), 205-223.

- [14] J. Lindenstrauss, L. Tzafriri *Classical Banach spaces II* L., Classical Banach spaces II, Springer(1979).
- [15] R. Denk, M. Hieber, and J. Pruss, R-Boundedness, *Fourier Multipliers and Problems of Elliptic and Parabolic Type*, Memoirs Amer. Math. Soc., vol. 166, Amer. Math. Soc., Providence, R.I., 2003.
- [16] G. Dore, *Maximal regularity in L^p spaces for an abstract Cauchy problem*. Adv. Differ. Equat. 5(1-3), 293-322 (2000).
- [17] K.-J. Engel and R. Nagel. *One-Parameter Semigroups for Linear Evolution Equations*, Springer-Verlag, New York, Berlin, Heidelberg, 2000.
- [18] G. Greiner. *Perturbing the boundary conditions of a generator*. Houston J. Math., 18:405-425, 2001.
- [19] S. Hadd. *Unbounded perturbations of C_0 -semigroups on Banach spaces and applications*. Semigroup Forum, 70:451-465, 2005.
- [20] S. Hadd and A. Idrissi, On the admissibility of observation for perturbed C_0 -semigroups on Banach spaces, *Systems Control Letters*. vol. 55, no. 1, 1-7, 2006.
- [21] S. Hadd, R. Manzo, A. Rhandi. *Unbounded perturbations of the generator domain*. Discrete & Continuous Dynamical Systems - A, 2015, 35 (2) : 703-723. doi: 10.3934/dcds.2015.35.703
- [22] B.H. Haak, M. Haase, P.C. Kunstmann *Perturbation, interpolation and maximal regularity*. Adv. Differential Equations 11 (2006), no. 2, 201-240. <http://projecteuclid.org/euclid.ade/1355867717>
- [23] B.H. Haak, P.C. Kunstmann *Admissibility of Unbounded Operators and Wellposedness of Linear Systems in Banach Spaces*, Integr. equ. oper. theory (2006) 55: 497. <https://doi.org/10.1007/s00020-005-1416-y>
- [24] M. Hieber and S. Monniaux. Pseudo-differential operators and maximal regularity results for non-autonomous parabolic equations. *Proc. Amer. Math. Soc.* 128:1047-1053, 2000.
- [25] H. Bounit, F. Maragh and H. Hammouri. Pseudo-differential operators and maximal regularity results for non-autonomous parabolic equations. *Eur. Jou. Cont. Soc.* 128:1047-1053, 2014.
- [26] B. Jacob, J.R. Partington, *Admissibility of control and observation operators for semigroups: a survey*, in: J.A. Ball, J.W. Helton, M.Klaus, L. Rodman (Eds.), Current Trends in Operator Theory and its Applications, Proceedings of IWOTA 2002, Operator Theory: Advances and Applications, vol.149, Birkhuser, Basel, pp.199-221, 2004.
- [27] N. Kalton, L. Weis: The H^∞ -calculus and sums of closed operators. Math. Ann. 321(2), 319-345 (2001)
- [28] T. Kato, *Perturbation Theory for Linear Operators*, Springer-Verlag, 1980.
- [29] P. C. Kunstmann and L. Weis, *Maximal L^p regularity for parabolic equations, Fourier multiplier theorems and H^∞ functional calculus*, Levico Lectures, Proceedings of the Autumn School on Evolution Equations and Semigroups (M. Iannelli, R. Nagel, S. Piazzera eds.), vol. 69, Springer Verlag, Heidelberg, Berlin, 2004, pp. 65-320
- [30] P. C. Kunstmann and L. Weis, *Perturbation theorems for maximal L^p -regularity*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 30 (2001), 415-435.
- [31] C. Le Merdy: The Weiss conjecture for bounded analytic semigroups. J. Lond. Math. Soc. (2) 67(3), 715-738 (2003)
- [32] F. Maragh, H. Bounit, A. Fadili, H. Hammouri; *On the admissible control operators for linear and bilinear systems and the Favard spaces*. Bulletin of the Belgian Mathematical Society - Simon Stevin, 21:711-732., (4) 2014.
- [33] A. Pazy: *Semigroups of Linear Operators and Applications to Partial Differential Equations*. Springer-Verlag, Berlin (1983).
- [34] P. Portal, Ž. Štrkalj, Pseudodifferential operators on Bochner spaces and an application, Math. Z. 253 (2006) 805-819.
- [35] J. Prüss, Sohr, Hermann. *Imaginary powers of elliptic second order differential operators in L^p -spaces*. Hiroshima Math. J. 23 (1993), no. 1, 161-192.
- [36] J. Prüss: *Evolutionary Integral Equations and Applications*. Monographs in Mathematics, vol. 87. Birkhäuser, Basel (1993).
- [37] J. Prüss and R. Schnaubelt. Solvability and maximal regularity of parabolic evolution equations with coefficients continuous in time. *J. Math. Anal. Appl.* 256:405-430, 2001.
- [38] M. Renardy and R. C. Rogers, *An introduction to partial differential equations*, 2nd ed., Texts in Applied Mathematics, vol. 13, Springer-Verlag, New York, 2004.

- [39] D. Salamon. *Infinite-dimensional linear system with unbounded control and observation: a functional analytic approach*. Trans.Amer.Math. Soc., 300:383-431, 1987.
- [40] O.J. Staffans. *Well-Posed Linear Systems*.Cambridge Univ. Press,Cambridge, 2005.
- [41] M. Tucsnak and G. Weiss, *Observation and Control for Operator Semigroups*, Birkhuser, 2009.
- [42] L. Weis, *A new approach to maximal L^p -regularity*, Proc. 6th International Conference on Evolution Equations, G. Lumer and L. Weis, eds, Dekker, New York (2000), 195-214.
- [43] L. Weis, *Operator-valued Fourier multiplier theorems and maximal L^p -regularity*, Math. Ann.,319 (2001), 735-758.
- [44] G. Weiss, *Admissible observation operators for linear semigroups*,Israel J. Math. 65 (1989) 17-43.
- [45] G. Weiss, *Admissibility of unbounded control operators*, SIAM J. Control Optim. 27 (1989) 527-545.
- [46] G. Weiss. *Regular linear systems with feedback. Mathematics of Control, Signals, and Systems*, 7:23-57, 1994.
- [47] G. Weiss. *Transfer functions of regular linear systems. Part I: Characterization of regularity*. Trans. Amer. Math. Soc., 342:827-854, 1994.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, IBN ZOHR UNIVERSITY, HAY DAKHLA, BP8106, 80000-AGADIR, MOROCCO