

# POSITIVE ALTERNATING SUMS OF INTEGER PARTITIONS

MOHAMED EL BACHRAOUI

ABSTRACT. We combine an extended version of Bailey's transform with an identity of Bressoud and with some identities of Berkovich and Warnaar to prove a variety of positivity results for alternating sums involving partition functions.

## 1. INTRODUCTION

Throughout the paper  $n$  is a nonnegative integer. The  $q$ -shifted factorials are given by

$$(a; q)_0 = (a)_0 = 1, \quad (a; q)_n = (a)_n = \prod_{j=0}^{n-1} (1 - aq^j), \quad (a; q)_\infty = (a)_\infty = \prod_{j=0}^{\infty} (1 - aq^j),$$

and the  $q$ -binomial coefficient is given for any nonnegative integers  $M$  and  $N$  by

$$\begin{bmatrix} N \\ M \end{bmatrix}_q = \begin{bmatrix} N \\ M \end{bmatrix} = \begin{cases} \frac{(q)_N}{(q)_M (q)_{N-M}} & \text{if } N \geq M, \\ 0 & \text{otherwise.} \end{cases}$$

It is well-known that for integers  $N \geq M \geq 0$ , the  $q$ -binomial coefficient  $\begin{bmatrix} N \\ M \end{bmatrix}$  is a polynomial of degree  $N - M$  which has nonnegative integer coefficients. As usual we let  $p(n)$  denote the number of partitions of  $n$ . Information about  $q$ -binomial coefficients and integer partition can be found in the book by Andrews [3]. Following [12] an overpartition of  $n$  is a partition with the extra condition that the first occurrence of any part may be overlined or not and the number of such overpartitions is written  $\bar{p}(n)$ . As to generating functions, we have

$$(1) \quad \sum_{n=0}^{\infty} p(n)q^n = \frac{1}{(q; q)_\infty} \text{ and } \sum_{n=0}^{\infty} \bar{p}(n)q^n = \frac{(-q; q)_\infty}{(q; q)_\infty}, \quad (p(0) = \bar{p}(0) = 1).$$

Alternating sums of integer partitions have received much attention by mathematicians in recent years. For paper on this subject which are related to our work, see for instance [6, 7, 14, 15]. Merca [15] proved that for all  $n > 0$

$$p(n-5) - p(n-2) - p(n-1) + p(n) \geq 0$$

which Andrews and Merca [6] generalized by showing that for  $n, k > 0$ ,

$$(2) \quad (-1)^{k-1} \sum_{j=0}^{k-1} (-1)^j \left( p(n - j(3j+1)/2) - p(n - j(3j+5)/2 - 1) \right) \geq 0.$$

---

*Date:* April 9, 2024.

*2000 Mathematics Subject Classification.* 33A30; 05A10;

*Key words and phrases.* Partitions; overpartitions; Bailey transform; positivity.

Their proof is partition-theoretic but it is based on the following truncated sum

$$(3) \quad \frac{1}{(q; q)_\infty} \sum_{j=0}^{k-1} (-1)^j q^{j(3j+1)/2} (1 - q^{2j+1}) = 1 + (-1)^{k-1} \sum_{n=1}^{\infty} \frac{q^{\binom{k}{2} + (k+1)n}}{(q; q)_n} \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}.$$

To derive the foregoing identity, Andrews and Merca used induction along with the  $q$ -binomial theorem and the basic properties of the  $q$ -binomial coefficients. Guo and Zeng [14] established the following truncated sum

$$(4) \quad \frac{(-q)_\infty}{(q)_\infty} \left( 1 + 2 \sum_{j=1}^k (-1)^j q^{j^2} \right) = 1 + (-1)^k \sum_{n=k+1}^{\infty} \frac{(-q)_k (-1)_{n-k} q^{(k+1)n}}{(q)_n} \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}$$

and as a consequence they obtained

$$(5) \quad (-1)^k \left( \bar{p}(n) + 2 \sum_{j=0}^k (-1)^j \bar{p}(n - j^2) \right) \geq 0.$$

The authors' proof for (4) relies on the following formula of Andrews [2]

$$\sum_{j=0}^n \frac{(b)_j (1 - bq^{2j}) (b/a)_j a^j q^{j^2}}{(1-b)(q)_j (aq)_j} = \frac{(bq)_n}{(aq)_n} \sum_{j=0}^n \frac{(b/a)_j a^j q^{(n+1)j}}{(q)_j}$$

which is a limiting case of Watson  $q$ -Whipple transformation, see Gasper and Rahman [13, III. 18] and Andrews [2, Lemma 2]. Besides, Guo and Zeng [14, p. 702] stated that it is an open problem to give partition-theoretic interpretation for (5). Recently, Andrews and Merca [7] provided such interpretations by first rewriting (4) in the following form

$$(6) \quad \begin{aligned} & \frac{(-q)_\infty}{(q)_\infty} \left( 1 + 2 \sum_{j=1}^k (-1)^j q^{j^2} \right) \\ &= 1 + 2(-1)^k \frac{(-q)_k}{(q)_k} \sum_{j=0}^{\infty} \frac{q^{(k+1)(k+j+1)} (-q^{k+j+2})_\infty}{(1 - q^{j+k+1}) (q^{j+k+2})_\infty} \end{aligned}$$

and then explaining the right-hand side of this formula in terms integer partitions. The main argument in the authors' proofs for (6) is the following formula which is due to Rogers and Fine (see [16, p. 15])

$$(7) \quad \sum_{j=0}^{\infty} \frac{(a)_j c^j}{(b)_j} = \sum_{j=0}^{\infty} \frac{(a)_j (acq/b)_j (bc)^j q^{j^2-j} (1 - acq^{2j})}{(b)_j (c)_{j+1}}.$$

Furthermore, Andrews and Merca [7] showed that both (3) and (4) can be established as instances of (7). In this paper we shall establish positivity results for alternating sums of the partition functions  $p(n)$  and  $\bar{p}(n)$ . We do that by showing that their corresponding generating functions have nonnegative coefficients. Specifically, we will prove the following main results.

**Theorem 1.** *There holds*

$$\frac{1}{(q; q)_\infty} \sum_{n=0}^{\infty} (-1)^n q^{\frac{13n^2+11n}{2}} (1 - q^{2n+1}) = \sum_{n=0}^{\infty} \frac{q^{n^2+2n}}{(q^2; q)_{2n+1}} \sum_{k=0}^n q^{k^2+k} \begin{bmatrix} n \\ k \end{bmatrix}_q.$$

**Corollary 1.**

$$\sum_{j=0}^n (-1)^j \left( p(n - j(13j + 11)/2) - p(n - j(13j + 15)/2 - 1) \right) \geq 0.$$

**Theorem 2.** *We have*

$$\begin{aligned} & \frac{1}{(q; q)_\infty} \sum_{n=0}^{\infty} (-1)^n q^{4n(10n-1)+4} (1 - q^{16n+8}) \\ &= \sum_{n=0}^{\infty} \frac{(-q^{-1}; q^2)_n}{(q; q)_{2n+3}} \sum_{k=0}^n \frac{q^{(n-2k)^2+5n+2k} (-q^4; q^4)_{k+1}}{(1 - q^{4k+2}) (-q^{3-2n}; q^2)_{2k}} \left[ \begin{matrix} 2n \\ 2k \end{matrix} \right]_{q^2}. \end{aligned}$$

**Corollary 2.**

$$\sum_{j=0}^n (-1)^j \left( \bar{p}(n - 4j(10j + 1) - 4) - \bar{p}(n - 4j(10j + 3) - 12) \right) \geq 0.$$

**Theorem 3.** *We have*

$$\begin{aligned} & \frac{1}{(q; q)_\infty} \sum_{n=0}^{\infty} (-1)^n q^{8n(3n+1)} (1 - q^{32n+16}) \\ &= \sum_{n=0}^{\infty} \frac{(-q^{-1}; q^2)_n}{(q^2; q^2)_n (q; q^2)_{n+2}} \sum_{k=0}^n \frac{q^{(n-2k)^2+5n+6k} (1 + q^{4k+4})}{1 - q^{4k+2}} \left[ \begin{matrix} 2n \\ 2k \end{matrix} \right]_{q^2}. \end{aligned}$$

**Corollary 3.**

$$\sum_{j=0}^n (-1)^j \left( p(n - 8j(3j + 1)) - p(n - 8j(3j + 5) - 16) \right) \geq 0.$$

**Theorem 4.** *There holds*

$$\begin{aligned} & \frac{1}{(q; q)_\infty} \sum_{n=0}^{\infty} (-1)^n q^{6n(9n+7)} (1 - q^{24n+12}) \\ &= \sum_{n=0}^{\infty} \frac{(-q^{-1}; q^2)_n}{(q^2; q^2)_n (q; q^2)_{n+3}} \sum_{k=0}^n \frac{q^{(n-2k)^2+7n+2k} (1 + q^4 + q^8) (1 + q^{6+4k})}{(1 - q^{2k+2}) (1 - q^{2k+4}) (-q^{3-2n}; q^2)_{2k}} \left[ \begin{matrix} 2n \\ 2k \end{matrix} \right]_{q^2}. \end{aligned}$$

**Corollary 4.**

$$\sum_{j=0}^n (-1)^j \left( \bar{p}(n - 6j(9j + 7)) - \bar{p}(n - 6j(9j + 11) - 12) \right) \geq 0.$$

The key argument of our proofs is the following extension of the Bailey's transform.

**Lemma 1.** [8, Lemma 1] *Let  $d, e, m$  and  $n$  be positive integers such that  $d \mid m$ . Assuming convergence of the series, if*

$$B_n = \sum_{j=0}^{\lfloor dn/m \rfloor} A_{ej} U_{dn-mj} V_{dn+mj} \quad \text{and} \quad C_n = \sum_{j=mn/d}^{\infty} D_{ej} U_{dj-mn} V_{dj+mn},$$

*then*

$$\sum_{n=0}^{\infty} A_{en} C_n = \sum_{n=0}^{\infty} B_n D_{en}.$$

Note that the case  $d = e = m = 1$  in Lemma 1 gives the classical Bailey's transform [9]. For a survey on Bailey's transform along with applications we refer for instance to [4, 5, 20, 22, 10]. We need the following well-known result

$$(8) \quad \sum_{k=n}^{\infty} \frac{a^k q^{k^2}}{(q)_{k-n} (aq)_{k+n}} = \frac{a^n q^{n^2}}{(aq)_{\infty}},$$

see for instance Bressoud [11, p. 215], Schilling and Warnaar [17, (1.3)], and Warnaar [21, (1.5)]. Note that according to Schilling and Warnaar [17], the pair of sequences  $(\delta_k)_{k \geq 0}$  and  $(\gamma_k)_{k \geq 0}$  where

$$\delta_k = a^k q^{k^2} \quad \text{and} \quad \gamma_k = \frac{a^k q^{k^2}}{(aq)_{\infty}}$$

is a *conjugate Bailey pair* as they are connected through the relation (8). We refer the reader to [17, 21] for an survey on Bailey pairs, conjugate Bailey pairs, and applications.

## 2. PROOF OF THEOREM 1 AND COROLLARY 1

We need the following identity of Bressoud [11, (3.4)]

$$(9) \quad \sum_{k=0}^n \frac{a^k q^{k^2} \begin{bmatrix} n \\ k \end{bmatrix}_q (a\gamma q; q)_n}{(a\gamma q; q)_k (a\gamma q; q)_{n-k}} \\ = \frac{(a^2 q; q)_{2n} (q; q)_n}{(aq; q)_n} \sum_{k=0}^{\lfloor n/2 \rfloor} \frac{a^{2k} \gamma^k q^{2k^2} (1 - aq^{2k}) (a; q)_k (\gamma^{-1}; q)_k}{(1-a)(q; q)_k (a\gamma q; q)_k (q; q)_{n-2k} (a^2 q; q)_{n+2k}}.$$

We want to apply Lemma 1 to (9) as follows. Let  $m = 2$ ,  $d = e = 1$  and let

$$A_n = \frac{a^{2n} \gamma^n q^{2n^2} (1 - aq^{2n}) (a; q)_n (\gamma^{-1}; q)_n}{(1-a)(q; q)_n (a\gamma q; q)_n}, \quad D_n = a^{2n} q^{n^2}, \\ U_n = \frac{1}{(q; q)_n}, \quad V_n = \frac{1}{(a^2 q; q)_n}.$$

Then

$$B_n = \sum_{k=0}^{\lfloor n/2 \rfloor} A_k U_{n-2k} V_{n+2k} \\ = \sum_{k=0}^{\lfloor n/2 \rfloor} \frac{a^{2k} \gamma^k q^{2k^2} (1 - aq^{2k}) (a; q)_k (\gamma^{-1}; q)_k}{(1-a)(q; q)_k (a\gamma q; q)_k (q; q)_{n-2k} (a^2 q; q)_{n+2k}} \\ = \frac{(aq; q)_n}{(a^2 q; q)_{2n} (q; q)_n} \sum_{k=0}^n \frac{a^k q^{k^2} \begin{bmatrix} n \\ k \end{bmatrix}_q (a\gamma q; q)_n}{(a\gamma q; q)_k (a\gamma q; q)_{n-k}}$$

where the last identity follows by (9). Moreover, by virtue of identity (8) we have

$$C_n = \sum_{k=2n}^{\infty} D_k U_{k-2n} V_{k+2n} = \sum_{k=2n}^{\infty} \frac{a^{2k} q^{k^2}}{(q; q)_{k-2n} (a^2 q; q)_{k+2n}} = \frac{a^{4n} q^{4n^2}}{(a^2 q; q)_{\infty}}.$$

Then by Lemma 1, we get

$$(10) \quad \sum_{n=0}^{\infty} B_n D_n = \sum_{n=0}^{\infty} \frac{a^{2n} q^{n^2} (aq; q)_n (a\gamma q; q)_n}{(a^2 q; q)_{2n} (q; q)_n} \sum_{k=0}^n \frac{a^k q^{k^2} \begin{bmatrix} n \\ k \end{bmatrix}_q}{(a\gamma q; q)_k (a\gamma q; q)_{n-k}}$$

$$= \sum_{n=0}^{\infty} A_n C_n = \frac{1}{(a^2 q; q)_{\infty}} \sum_{n=0}^{\infty} \frac{a^{6n} \gamma^n q^{6n^2} (1 - q^{2n}) (a; q)_n (\gamma^{-1}; q)_n}{(1 - a)(q; q)_n (a \gamma q; q)_n}.$$

Letting  $\gamma \rightarrow 0$  in (10) yields

$$(11) \quad \frac{1}{(a^2 q; q)_{\infty}} \sum_{n=0}^{\infty} \frac{(-1)^n a^{6n} q^{\frac{13n^2-n}{2}} (1 - a q^{2n}) (a; q)_n}{(1 - a)(q; q)_n} \\ = \sum_{n=0}^{\infty} \frac{a^{2n} q^{n^2} (a q; q)_n}{(a^2 q; q)_{2n} (q; q)_n} \sum_{k=0}^n a^k q^{k^2} \begin{bmatrix} n \\ k \end{bmatrix}_q.$$

Now let  $a = q$  in (11) and simplify to obtain

$$\frac{1}{(q; q)_{\infty}} \sum_{n=0}^{\infty} (-1)^n (1 - q^{2n+1}) q^{\frac{13n^2+11n}{2}} = (1 - q) \sum_{n=0}^{\infty} \frac{q^{n^2+2n}}{(q; q)_{2n+2}} \sum_{k=0}^n q^{k^2+k} \begin{bmatrix} n \\ k \end{bmatrix}_q.$$

That is

$$\frac{1}{(q; q)_{\infty}} \sum_{n=0}^{\infty} (-1)^n (1 - q^{2n+1}) q^{\frac{13n^2+11n}{2}} = \sum_{n=0}^{\infty} \frac{q^{n^2+2n}}{(q^2; q)_{2n+1}} \sum_{k=0}^n q^{k^2+k} \begin{bmatrix} n \\ k \end{bmatrix}_q,$$

which completes the proof of the theorem. As to the corollary, simply note that the left-hand side of previous formula is the generating function of

$$\sum_{j=0}^n (-1)^j \left( p(n - j(13j + 11)/2) - p(n - j(13j + 15)/2 - 1) \right)$$

and that its right-hand side has nonnegative coefficients.

### 3. PROOF OF THEOREM 2 AND COROLLARY 2

We need the following formula of Berkovich and Warnaar [5, p. 36]

$$\sum_{k=0}^{\lfloor n/4 \rfloor} \frac{(1 - a^2 q^{16k}) (a^2, a^2 q^4/b; q^8)_k (q^{-n}; q)_{4k}}{(1 - a^2) (q^8, b q^4; q^8)_k (a q^{n+1}; q)_{4k}} (b q^{4n-2})^k \\ = \frac{q^n (-q^{-1}; q^2)_n (a q; q)_n}{(-q; q)_n (a q; q^2)_n} \sum_{k \geq 0} \frac{(b^{1/2}, -b^{1/2}, -a q^4, q^{-2n}, q^{2-2n}; q^4)_k q^{4k}}{(q^4, b, a q^2, -q^{3-2n}, -q^{5-2n}; q^4)_k}$$

which is equivalent with

$$(12) \quad \sum_{k=0}^{\lfloor n/4 \rfloor} \frac{(1 - a^2 q^{16k}) (a^2, a^2 q^4/b; q^8)_k (q; q)_n (a q; q)_n q^{\binom{4k}{2} - 2k} b^k}{(1 - a^2) (q^8, b q^4; q^8)_k (q; q)_{n-4k} (a q; q)_{n+4k}} \\ = \frac{q^n (-q^{-1}; q^2)_n (a q; q)_n}{(-q; q)_n (a q; q^2)_n} \sum_{k \geq 0} \frac{(b^{1/2}, -b^{1/2}, -a q^4, q^{-2n}, q^{2-2n}; q^4)_k q^{4k}}{(q^4, b, a q^2, -q^{3-2n}, -q^{5-2n}; q^4)_k}.$$

We apply Lemma 1 to (12) as follows. Let  $m = 4$ ,  $d = e = 1$  and let

$$A_n = \frac{b^n q^{4n(2n-1)} (1 - a^2 q^{16n}) (a^2, a^2 q^4/b; q^8)_n}{(1 - a^2) (q^8, b q^4; q^8)_n}, \quad D_n = a^n q^{n^2}, \\ U_n = \frac{1}{(q; q)_n}, \quad V_n = \frac{1}{(a q; q)_n}.$$

Then on the one hand by (12),

$$\begin{aligned} B_n &= \sum_{j=0}^{\lfloor n/4 \rfloor} A_j U_{n-4j} V_{n+4j} \\ &= \frac{q^n(-q^{-1}; q^2)_n}{(-q; q)_n (q; q)_n (aq; q^2)_n} \sum_{k \geq 0} \frac{(b^{1/2}, -b^{1/2}, -aq^4, q^{-2n}, q^{2-2n}, q^4)_k q^{4k}}{(q^4, b, aq^2, -q^{3-2n}, -q^{5-2n}, q^4)_k}. \end{aligned}$$

On the other hand by (8),

$$C_n = \sum_{k=4n}^{\infty} \frac{a^k q^{k^2}}{(q; q)_{k-4n} (aq; q)_{k+4n}} = \frac{a^{4n} q^{16n^2}}{(aq; q)_{\infty}}.$$

Now apply Lemma 1 to these sequences to obtain from

$$\sum_{n=0}^{\infty} B_n D_n = \sum_{n=0}^{\infty} A_n C_n$$

that

$$\begin{aligned} (13) \quad & \sum_{n=0}^{\infty} \frac{a^n q^{n^2+n} (-q^{-1}; q^2)_n}{(q^2; q^2)_n (aq; q^2)_n} \sum_{k \geq 0} \frac{q^{4k} (b^{1/2}, -b^{1/2}, -aq^4, q^{-2n}, q^{2-2n}, q^4)_k}{(q^4, b, aq^2, -q^{3-2n}, -q^{5-2n}, q^4)_k} \\ &= \frac{1}{(aq; q)_{\infty}} \sum_{n=0}^{\infty} a^{4n} b^n q^{24n^2-4n} \frac{(1 - a^2 q^{16n})(a^2, a^2 q^4/b; q^8)_n}{(1 - a^2)(q^8, bq^4; q^8)_n} \end{aligned}$$

Letting in (13)  $b \rightarrow 0$  and then  $a = q^4$  we obtain

$$\begin{aligned} & \frac{1}{(q^5; q)_{\infty}} \sum_{n=0}^{\infty} (-1)^n q^{40n^2-4n+4} \frac{1 - q^{16n+8}}{1 - q^8} \\ &= \sum_{n=0}^{\infty} \frac{q^{n^2+5n} (-q^{-1}; q^2)_n}{(q^2; q^2)_n (q^5; q^2)_n} \sum_{k \geq 0} \frac{(-q^8; q^4)_k (q^{-2n}; q^2)_{2k}}{(q^4; q^4)_k (q^6; q^4)_k (-q^{3-2n}; q^2)_{2k} q^{4k}}, \end{aligned}$$

which after some straightforward simplification yields

$$\begin{aligned} & \frac{1}{(q)_{\infty}} \sum_{n=0}^{\infty} (-1)^n q^{40n^2-4n+4} (1 - q^{16n+8}) \\ &= \sum_{n=0}^{\infty} \frac{q^{n^2+5n} (-q^{-1}; q^2)_n}{(q; q)_{2n+3}} \sum_{k=0}^n \frac{q^{4k^2+2k-4nk} (-q^4; q^4)_{k+1} (q^2; q^2)_{2n}}{(q^2; q^2)_{2k+1} (q^2; q^2)_{2n-2k} (-q^{3-2n}; q^2)_{2k}}, \end{aligned}$$

or equivalently,

$$\begin{aligned} (14) \quad & \frac{1}{(q)_{\infty}} \sum_{n=0}^{\infty} (-1)^n q^{40n^2-4n+4} (1 - q^{16n+8}) \\ &= \sum_{n=0}^{\infty} \frac{(-q^{-1}; q^2)_n}{(q; q)_{2n+3}} \sum_{k=0}^n \frac{q^{(n-2k)^2+5n+2k} (-q^4; q^4)_{k+1}}{(1 - q^{4k+2})(-q^{3-2n}; q^2)_{2k}} \left[ \begin{matrix} 2n \\ 2k \end{matrix} \right]_{q^2}. \end{aligned}$$

This proves the theorem. Concerning the corollary, multiply both sides of (14) by  $(-q; q)_{\infty}$ . Then by (1) the left-hand side of the obtained formula is the generating function of

$$\sum_{j=0}^n (-1)^j \left( \bar{p}(n - 4j(10j + 1) - 4) - \bar{p}(n - 4j(10j + 3) - 12) \right).$$

Moreover, it is easily seen that the right-hand side of the obtained formula has nonnegative coefficients. This completes the proof.

#### 4. PROOF OF THEOREM 3 AND COROLLARY 3

As to this result we shall appeal to the following formula of Berkovich and Warnaar [10, p. 36]

$$\begin{aligned} & \sum_{k=0}^{\lfloor n/4 \rfloor} \frac{(1 - a^4 q^{32k})(a^4; q^{16})_k (q^{-n}; q)_{4k}}{(1 - a^4)(q^{16}; q^{16})_k (aq^{n+1}; q)_{4k}} (-q^{4n-6})^k \\ &= q^n \frac{(aq^{-1}; q^2)_n (aq; q)_n}{(-q; q)_n (aq; q^2)_n} \sum_{k \geq 0} \frac{(iq^{-2}, -iq^{-2}, -aq^4, q^{-2n}, q^{2-2n}; q^4)_k}{(q^4, -q^4, aq^2, q^{3-2n}, q^{5-2n}; q^4)_k} q^{8k} \end{aligned}$$

which after simplification yields

$$\begin{aligned} (15) \quad & \sum_{k=0}^{\lfloor n/4 \rfloor} (-1)^k q^{8k^2-8k} \frac{(1 - a^4 q^{32k})(a^4; q^{16})_k}{(1 - a^4)(q^{16}; q^{16})_k (q; q)_{n-4k} (aq; q)_{n+4k}} \\ &= \frac{q^n (-q^{-1}; q^2)_n}{(q^2; q^2)_n (aq; q^2)_n} \sum_{k \geq 0} \frac{q^{8k} (-q^4; q^8)_k (q^{-2n}; q^2)_{2k} (-aq^4; q^4)_k}{(-q^8; q^8)_k (aq^2; q^4)_k (q^{3-2n}; q^2)_{2k}}. \end{aligned}$$

Let  $m = 4$  and  $d = e = 1$  and let

$$\begin{aligned} A_n &= \frac{(-1)^n q^{2n(4n-4)} (1 - a^4 q^{32n})(a^4; q^{16})_n}{(1 - a^4)(q^{16}; q^{16})_n}, \quad D_n = a^n q^{n^2}, \\ U_n &= \frac{1}{(q; q)_n}, \quad V_n = \frac{1}{(aq; q)_n}. \end{aligned}$$

Then by (15),

$$\begin{aligned} B_n &= \sum_{j=0}^{\lfloor n/4 \rfloor} A_j U_{n-4j} V_{n+4j} \\ &= \frac{q^n (-q^{-1}; q^2)_n}{(q^2; q^2)_n (aq; q^2)_n} \sum_{k \geq 0} \frac{q^{8k} (-q^4; q^8)_k (q^{-2n}; q^2)_{2k} (-aq^4; q^4)_k}{(-q^8; q^8)_k (aq^2; q^4)_k (q^{3-2n}; q^2)_{2k}} \end{aligned}$$

and by (8),

$$C_n = \sum_{k=4n}^{\infty} \frac{a^k q^{k^2}}{(q; q)_{k-4n} (aq; q)_{k+4n}} = \frac{a^{4n} q^{16n^2}}{(aq; q)_{\infty}}.$$

Thus by virtue of Lemma 1

$$\sum_{n=0}^{\infty} B_n D_n = \sum_{n=0}^{\infty} A_n C_n$$

and therefore

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{a^n q^{n^2+n} (-q^{-1}; q^2)_n}{(q^2; q^2)_n (aq; q^2)_n} \sum_{k \geq 0} \frac{q^{8k} (-q^4; q^8)_k (q^{-2n}; q^2)_{2k} (-aq^4; q^4)_k}{(-q^8; q^8)_k (q^{3-2n}; q^2)_{2k} (aq^2; q^4)_k} \\ &= \frac{1}{(aq; q)_{\infty}} \sum_{n=0}^{\infty} (-1)^n a^{4n} q^{24n^2-8n} \frac{(1 - a^4 q^{32n})(a^4; q^{16})_n}{(1 - a^4)(q^{16}; q^{16})_n}. \end{aligned}$$

Letting in the foregoing identity  $a = q^4$  and simplifying give

$$\begin{aligned} & \frac{1}{(q^5; q)_\infty} \sum_{n=0}^{\infty} (-1)^n q^{24n^2+8n} \frac{1 - q^{32n+16}}{1 - q^8} \\ &= \sum_{n=0}^{\infty} \frac{q^{n^2+5n} (-q^{-1}; q^2)_n}{(q^2; q^2)_n (q^5; q^2)_n} \sum_{k \geq 0} \frac{1 + q^{4k+4}}{1 + q^4} \frac{q^{4k^2+6k-4nk} (-q^4; q^8)_k (q^2; q^2)_{2n}}{(q^4; q^4)_k (q^6; q^4)_k (q^2; q^2)_{2n-2k} (q^{3-2n}; q^2)_{2k}} \end{aligned}$$

which by rearranging becomes

$$\begin{aligned} & \frac{1}{(q; q)_\infty} \sum_{n=0}^{\infty} (-1)^n q^{8n(3n+1)} (1 - q^{32n+16}) \\ &= \sum_{n=0}^{\infty} \frac{(-q^{-1}; q^2)_n}{(q^2; q^2)_n (q; q^2)_{n+2}} \sum_{k=0}^n \frac{q^{(n-2k)^2+5n+6k} (1 + q^{4k+4})}{1 - q^{4k+2}} \begin{bmatrix} 2n \\ 2k \end{bmatrix}_{q^2}. \end{aligned}$$

which proves the theorem. The corollary follows immediately as the left-hand side of the foregoing formula is the generating function of

$$\sum_{j=0}^n (-1)^j \left( p(n - 8j(3j + 1)) - p(n - 8j(3j + 5) - 16) \right)$$

and its right-hand side has nonnegative coefficients.

## 5. PROOF OF THEOREM 4 AND COROLLARY 4

The proof relies on the following relation due to Berkovich and Warnaar [10, p. 36]

$$\begin{aligned} & \sum_{k=0}^{\lfloor n/6 \rfloor} \frac{(1 - a^2 q^{24k})(a^2; q^{12})_k (q^{-n}; q)_{6k}}{(q^{12}; q^{12})_k (aq^{n+1}; q)_{6k}} (-a^2 q^{6n-3})^k \\ &= \frac{q^n (-q^{-1}; q^2)_n (aq; q)_n}{(-q; q)_n (aq; q^2)_n} \sum_{k \geq 0} \frac{q^{4k} (a^{2/3}, \omega a^{2/3}, \omega^2 a^{2/3}, -aq^4, q^{-2n}, q^{2-2n}; q^4)_k}{(q^4, a, -a, aq^2, -q^{3-2n}, -q^{5-2n}; q^4)_k} \end{aligned}$$

where  $\omega$  is a third root of unity. Then the previous formula becomes

$$\begin{aligned} (16) \quad & \sum_{k=0}^{\lfloor n/6 \rfloor} (-1)^k a^{2k} q^{18k^2-6k} \frac{(1 - a^2 q^{24k})(a^2; q^{12})_k}{(1 - a^2)(q^{12}; q^{12})_k (q; q)_{n-6k} (aq; q)_{n+6k}} \\ &= \frac{q^n (-q^{-1}; q^2)_n}{(q^2; q^2)_n (aq; q^2)_n} \sum_{k \geq 0} \frac{q^{4k} (a^{2/3}, \omega a^{2/3}, \omega^2 a^{2/3}; q^4)_k (q^{-2n}; q^2)_{2k} (-aq^4; q^4)_k}{(q^4, a, -a, aq^2; q^4)_k (-q^{3-2n}; q^2)_{2k}}. \end{aligned}$$

Let  $m = 6$ ,  $d = e = 1$  and let

$$\begin{aligned} A_n &= \frac{(-1)^n a^{2n} q^{3n(6n-2)} (1 - a^2 q^{24n})(a^2; q^{12})_n}{(1 - a^2)(q^{12}; q^{12})_n}, \quad D_n = a^n q^{n^2}, \\ U_n &= \frac{1}{(q; q)_n}, \quad V_n = \frac{1}{(aq; q)_n}. \end{aligned}$$

Thus by Lemma 1 applied to these items we find

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{a^n q^{n^2+n} (-q^{-1}; q^2)_n}{(q^2; q^2)_n (aq; q^2)_n} \sum_{k \geq 0} \frac{q^{4k} (a^{2/3}, \omega a^{2/3}, \omega^2 a^{2/3}, -aq^4; q^4)_k (q^{-2n}; q^2)_{2k}}{(q^4, a, -a, aq^2; q^4)_k (-q^{3-2n}; q^2)_{2k}} \\ &= \frac{1}{(aq; q)_\infty} \sum_{n=0}^{\infty} (-1)^n a^{8n} q^{54n^2-6n} \frac{(1 - a^2 q^{24n})(a^2; q^{12})_n}{(1 - a^2)(q^{12}; q^{12})_n}. \end{aligned}$$



Letting  $a = q^6$  in the forgoing formula we find after easy simplification, including basic facts on  $\omega$ ,

$$\begin{aligned} & \frac{1}{(q^7; q)_\infty} \sum_{n=0}^{\infty} (-1)^n q^{54n^2+42n} \frac{1 - q^{24n+12}}{1 - q^{12}} \\ &= \sum_{n=0}^{\infty} \frac{q^{n^2+7n}(-q^{-1}; q^2)_n}{(q^2; q^2)_n (q^7; q^2)_n} \sum_{k \geq 0} \frac{q^{4k}(1 + q^4 + q^8)(-q^{10}; q^4)_k (q^{-2n}; q^2)_{2k}}{(q^{12}; q^8)_k (q^8; q^4)_k (-q^{3-2n}; q^2)_{2k}}, \end{aligned}$$

or equivalently,

$$\begin{aligned} & \frac{1}{(q^6; q)_\infty} \sum_{n=0}^{\infty} (-1)^n q^{54n^2+42n} (1 - q^{24n+12}) \\ &= \sum_{n=0}^{\infty} \frac{q^{n^2+7n}(-q^{-1}; q^2)_n}{(q^2; q^2)_n (q^7; q^2)_n} \sum_{k \geq 0} \frac{q^{4k}(1 + q^4 + q^8)(1 + q^{6+4k})(q^{-2n}; q^2)_{2k}}{(q^6; q^4)_k (q^8; q^4)_k (-q^{3-2n}; q^2)_{2k}}, \end{aligned}$$

which yields,

$$\begin{aligned} & \frac{1}{(q; q)_\infty} \sum_{n=0}^{\infty} (-1)^n q^{54n^2+42n} (1 - q^{24n+12}) \\ &= \sum_{n=0}^{\infty} \frac{q^{n^2+7n}(-q^{-1}; q^2)_n}{(q^2; q^2)_n (q; q^2)_{n+3}} \sum_{k \geq 0} \frac{q^{4k^2+2k-4nk}(1 + q^4 + q^8)(1 + q^{6+4k})(q^2; q^2)_{2n}}{(q^2; q^4)_{k+1} (q^4; q^4)_{k+1} (q^2; q^2)_{2n-2k} (-q^{3-2n}; q^2)_{2k}}. \end{aligned}$$

That is,

$$\begin{aligned} & \frac{1}{(q; q)_\infty} \sum_{n=0}^{\infty} (-1)^n q^{54n^2+42n} (1 - q^{24n+12}) \\ &= \sum_{n=0}^{\infty} \frac{(-q^{-1}; q^2)_n}{(q^2; q^2)_n (q; q^2)_{n+3}} \sum_{k=0}^n \frac{q^{(n-2k)^2+7n+2k}(1 + q^4 + q^8)(1 + q^{6+4k})}{(1 - q^{2k+2})(1 - q^{2k+4})(-q^{3-2n}; q^2)_{2k}} \begin{bmatrix} 2n \\ 2k \end{bmatrix}_{q^2} \end{aligned}$$

which confirms the theorem. To prove the corollary, first multiply both sides of the previous identity by  $(-q; q)_\infty$ . Next observe that the left-hand side of the obtained formula is the generating function of

$$\sum_{j=0}^n (-1)^j \left( \bar{p}(n - 6j(9j + 7)) - \bar{p}(n - 6j(9j + 11) - 12) \right)$$

and the right-hand side of the obtained formula has nonnegative coefficients. This completes the proof.

## 6. CONCLUDING REMARKS AND OPEN QUESTIONS

Each one of our proofs for Theorems 1-4 relies on substituting specific values for the parameter  $a$  which lead to the appropriate formulas. By choosing different values for  $a$  one might derive other important results. For instance, upon letting in the relation (11)  $a \rightarrow 1$  instead of  $a = q$ , we get after simplification

$$(17) \quad \frac{1}{(q; q)_\infty} \sum_{n=0}^{\infty} (-1)^n (1 + q^n) q^{\frac{13n^2-n}{2}} = \sum_{n=0}^{\infty} \sum_{k=0}^n \frac{q^{n^2+k^2}}{(q; q)_{2n}} \begin{bmatrix} n \\ k \end{bmatrix}_q$$

or equivalently

$$(18) \quad \frac{1}{(q; q)_\infty} \sum_{n=-\infty}^{\infty} (-1)^n q^{\frac{13n^2+n}{2}} = \sum_{n=0}^{\infty} \sum_{k=0}^n \frac{q^{n^2+k^2}}{(q; q)_{2n}} \begin{bmatrix} n \\ k \end{bmatrix}_q.$$

In addition, Andrews [1, formula (5.8) for  $k = 1$ ] and Warnaar [20, Theorem 1.2 for  $k = 4$ ] evaluated the right-hand side of (18) as follows

$$(19) \quad \sum_{n=0}^{\infty} \sum_{k=0}^n \frac{q^{n^2+k^2}}{(q; q)_{2n}} \begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{(q^6, q^7, q^{13}; q^{13})_{\infty}}{(q; q)_{\infty}}.$$

Then by a combination of (18) and (19) we deduce

$$(20) \quad \frac{1}{(q; q)_{\infty}} \sum_{n=-\infty}^{\infty} (-1)^n q^{\frac{13n^2+n}{2}} = \frac{(q^6, q^7, q^{13}; q^{13})_{\infty}}{(q; q)_{\infty}}$$

which is also easily obtained by Jacobi's triple product identity [13, p. 15]

$$(21) \quad (zq, z^{-1}q, q^2; q^2)_{\infty} = \sum_{n=-\infty}^{\infty} (-1)^n q^{n^2} z^n$$

applied to  $z = q^{1/2}$  and  $q$  replaced by  $q^{13/2}$ . Therefore both (18) and (20) imply that the power series

$$\frac{1}{(q; q)_{\infty}} \sum_{n=-\infty}^{\infty} (-1)^n q^{\frac{13n^2+n}{2}}$$

has nonnegative coefficients. Besides, it turns out that for any nonnegative odd integer  $M$  we find by (21) that

$$\frac{1}{(q; q)_{\infty}} \sum_{n=-\infty}^{\infty} (-1)^n q^{\frac{Mn^2+n}{2}} = \frac{(q^{(M-1)/2}, q^{(M+1)/2}, q^M; q^M)_{\infty}}{(q; q)_{\infty}}$$

showing that the left-hand side of this formula has nonnegative integer coefficients. However, we do not how to arrive at this fact using Lemma 1. This would suggest a general pattern for our proofs in Theorems 1-4. Obviously, the first challenging part is how to obtain a terminating identity of the form

$$\sum_{k=0}^{\lfloor n/m \rfloor} \frac{A_k(a, q)}{(q; q)_{n-mk} (aq; q)_{n+mk}} = B_n(a, q),$$

where  $A_k(a, q)$  and  $B_n(a, q)$  are rational functions of  $a$  and  $q$ . Furthermore, as each one of the alternating sums in Corollaries 1-4 are nonnegative, it is natural to ask the question whether combinatorial interpretations exist for such sums.

**Acknowledgment.** The author thanks Wadim Zudilin for bringing his attention to the papers by Warnaar [20, 22] and the paper by Berokich and Warnaar [10] along with other papers. The author is grateful to him and to Ole Warnaar for reading carefully a previous version of the paper and for fruitful comments and interesting suggestions which manifestly improved the presentation and quality of this paper.

## REFERENCES

- [1] G.E. Andrews, *Multiple series Rogers–Ramanujan type identities*, Pacific J. Math. 114 (1984), 267–283.
- [2] G. E. Andrews, *The fifth and seventh order mock theta functions*, Trans. Amer. Math. Soc. 293 (1986), no. 1, 113–134.
- [3] G.E. Andrews, *The Theory of Partitions*, Cambridge University Press, Cambridge, 1998.
- [4] G. E. Andrews, *Bailey's transform, lemma, chains and tree*, Proceedings of Special Functions Conference, Tempe, 2000.

- [5] G. E. Andrews and A. Berkovich, *The WP-Bailey tree and its implications*, J. London Math. Soc. (2) 66 (2002), no. 3, 529–549.
- [6] G.E. Andrews and M. Merca, *The truncated pentagonal number theorems*, J. Combin. Theory. Ser. A 119 (2012) 1639–1643.
- [7] G.E. Andrews and M. Merca, *Truncated theta series and a problem of Guo and Zeng*, J. Combin. Theory. Ser. A 154 (2018) 610–619.
- [8] M. El Bachraoui, *Further extended Bailey’s transforms and applications to some classical  $q$ -series identities*, Submitted.
- [9] W.N. Bailey, *Some identities in combinatory analysis*, Proc. London Math. Soc. (2) 49 (1947) 421–435.
- [10] A. Berkovich and S.O. Warnaar *Positivity preserving transformations for  $q$ -binomial coefficients*, Trans. Amer. Math. Soc. 357 (2005) 2291–2351.
- [11] D.M. Bressoud, *Some identities for terminating  $q$ -series*, Math. Proc. Cambridge Philos. Soc. 81 (1981) 211–223.
- [12] S. Corteel and J. Lovejoy, *Overpartitions*, Trans. Amer. Math. Soc. 356 (2004) 1623–1635.
- [13] G. Gasper and M. Rahman, *Basic Hypergeometric series*, Cambridge University Press, 2004.
- [14] V.J.W Guo and J. Zeng, *Two truncated identities of Gauss*, J. Combin. Theory. Ser. A 120 (2013) 700–707.
- [15] M. Merca, *Fast algorithm for generating ascending compositions*, J. Math. Model. Algorithms 11 (1) (2012) 89–104.
- [16] L.J. Rogers, *On two theorems of combinatory analysis and some allied identities*, Proc. Lond. Math. Soc., Ser. 2 16 (1916) 315–336.
- [17] A. Schilling and S.O. Warnaar, *A higher level Bailey lemma: proof and application*, Ramanujan J. 2 (1998) 327–349.
- [18] D. Shanks, *A short proof of an identity of Euler*, Proc. Amer. Math. Soc. 2 (1951) 747–749.
- [19] L. J. Slater, *Generalized Hypergeometric Functions*, Cambridge University, Cambridge 1966.
- [20] S.O. Warnaar *The generalized Borwein conjecture. I. The Burge transform*, Contemporary Mathematics 291 (2001) 243–267.
- [21] S.O. Warnaar *50 years of Baileys lemma*, in Alg. Combinatorics and Appl., A. Betten et al eds., Springer, Berlin, 2001.
- [22] S.O. Warnaar *The generalized Borwein conjecture. II. Refined  $q$ -trinomial coefficients*, Discrete Mathematics 272 (2003) 215–258.

DEPT. MATH. SCI, UNITED ARAB EMIRATES UNIVERSITY, PO Box 15551, AL-AIN, UAE  
*E-mail address:* melbachraoui@uaeu.ac.ae