

ON THE BEHAVIOR OF UNBOUNDED COLLATZ SEQUENCES

JORGE SALAZAR

ABSTRACT. The aim of this paper is to show a peculiar behavior of a (hypothetical) Collatz sequence going to infinity. We study the associated Syracuse sequence (the odd elements of the former) and show that the limit set of a conveniently normalized sequence is the whole unit interval. In particular, for any positive integer there is a subsequence whose elements' expansions in base 3 begin (from the left) with the expansion of the given number.

1. INTRODUCTION

The Collatz operation is $Col(a) = 3a + 1$, if a is odd and $Col(a) = a/2$ if a is even. In 1937 Lothar Collatz introduced this operation and conjectured that any sequence of its iterations eventually becomes the loop $1, 4, 2, 1$. The conjecture is also known as the “ $3x+1$ conjecture” or the “Syracuse problem.” Terence Tao [2] gave an enthusiastic slide presentation of this conjecture and he himself have important contributions [1].

Since $3a + 1$ is even, we can divide by 2 in the same step and consider instead the iterations of

$$Col_2(a) = \begin{cases} \frac{1}{2}(3a + 1), & \text{if } a \text{ is odd,} \\ \frac{1}{2}a, & \text{if } a \text{ is even.} \end{cases}$$

Given $a_0 \geq 1$, denote $a_\alpha = Col_2(a_{\alpha-1})$, $\alpha \geq 1$, the sequence of iterations. Let's denote q_α the biggest power in the development of a_α in base 3. i.e.

$$(1.1) \quad 1 \leq 3^{-q_\alpha} a_\alpha < 3$$

Theorem 1.1. *Assume that for some integer $a_0 \geq 1$, the Collatz sequence verifies $\lim_{\alpha \rightarrow \infty} a_\alpha = \infty$. Let ψ be any positive integer. Then there are infinitely many indexes α such that $[3^{p-q_\alpha} a_\alpha] = \psi$, where $p = \left\lfloor \frac{\ln \psi}{\ln 3} \right\rfloor$.*

($[\cdot]$ is the integer part of a given number.)

The equality $[3^{p-q_\alpha} a_\alpha] = \psi$ means that the first p digits (from the left) of a_α in base 3 match those of ψ .

2020 *Mathematics Subject Classification.* Number Theory.

Key words and phrases. Collatz sequence, Syracuse sequence, Collatz conjecture.

In the whole paper, $(a_\alpha)_{\alpha \in \mathbb{N}}$ is a **fixed** orbit of Col_2 going to infinity. i.e.

$$(1.2) \quad \forall \alpha \geq 1, a_\alpha = Col_2(a_{\alpha-1}) \text{ and } \lim_{\alpha \rightarrow \infty} a_\alpha = \infty.$$

2. BOUNDED VERSION OF THE COLLATZ PROCESS

For all $\alpha \geq 1$, put

$$(2.1) \quad c_\alpha = 3^{-q_\alpha} \cdot a_\alpha.$$

The following definitions are intended to obtain c_α as the iterations of a single operation starting from c_1 . i.e. We need to understand how the Collatz operation works on c_α , for $\alpha \geq 1$.

Definition 2.1. Denote

$$(2.2) \quad \mathbf{N} = \bigcup_{r=0}^{\infty} 3^{-r} \mathbb{N}.$$

the set of all the rational numbers with a finite expansion in base 3

For $b \in \mathbf{N}$, define

$$(2.3) \quad Col_3(b) = \begin{cases} \frac{b}{2}, & \text{if } \frac{b}{2} \in \mathbf{N}, \\ \frac{b}{2} + \frac{3^{-q-1}}{2}, & \text{if } \frac{b}{2} \notin \mathbf{N}, \end{cases}$$

where

$$(2.4) \quad q = \min \{r \geq 0; 3^r b \in \mathbb{N}\}.$$

Note: If $\frac{b}{2} \notin \mathbf{N}$, the base 3 expansion of $\frac{b}{2}$ ends in an infinite string of 1's, starting at the position $-q-1$. In that case, we “round up” to $\frac{b+3^{-q-1}}{2}$ and $Col_3(b) \in \mathbf{N}$. When the rounding occurs, the last position is $-q-1$ and its value is 2. Working with Col_3 , we take advantage of the long division procedure (in base 3) by starting with division by 2, without caring about the parity of the number until we get to the last digit.

For future reference, observe that for all $b \in \mathbf{N}$ and $n \in \mathbb{N}$,

$$(2.5) \quad 2^{-n}b \leq Col_3^n(b) \leq 2^{-n}b + 3^{-q},$$

where q is defined by (2.4).

Definition 2.2. For all $b \in \mathbf{N} \cap [1, 3)$, define

$$(2.6) \quad Col_4(b) = \begin{cases} Col_3(b), & \text{if } 2 \leq b < 3, \\ 3 \cdot Col_3(b), & \text{if } 1 \leq b < 2. \end{cases}$$

Clearly, for all $\alpha \geq 1$,

$$(2.7) \quad c_{\alpha+1} = Col_4(c_\alpha).$$

(If a_0 is not divisible by 3, (2.7) is also true for $\alpha = 0$.)

The usefulness of this version of the Collatz iteration (Col_4) is that we can multiply by the appropriate power of 3 after any number of iterations of Col_3 , as we can see in the next Lemma.

Lemma 2.3. *For all $n \in \mathbb{N}$, for all $b \in \mathbb{N} \cap [1, 3)$,*

$$Col_4^n(b) = 3^k \cdot Col_3^n(b),$$

where

$$(2.8) \quad k = - \left\lceil \frac{\ln Col_3^n(b)}{\ln 3} \right\rceil.$$

Indeed, for all α , the coefficients of a_α , c_α and

$$b_{\alpha+1} = Col_3(b_\alpha), \quad \alpha \geq 1, \quad b_1 = c_1,$$

in base 3, are the same, just shifted.

The main ingredient in the proof of Theorem 1.1 is the following result (interesting on their own), which will be proved in Section 4.

Proposition 2.4.

$$(2.9) \quad \liminf_{\alpha \rightarrow \infty} c_\alpha = 1.$$

3. PRELIMINARY RESULTS

Lemma 3.1. *Let $\phi, \theta \in \mathbb{N} \cap [1, 3)$. Suppose there is an integer k such that*

$$(3.1) \quad \left\lceil 3^k \phi \right\rceil = \left\lceil 3^k \theta \right\rceil.$$

Assume that $\left\lceil 3^k \phi \right\rceil \neq 3^k \phi$ and $\left\lceil 3^k \theta \right\rceil \neq 3^k \theta$ (irrelevant if $\phi = \theta$). Then, for all $m \in \mathbb{N}$,

$$(3.2) \quad \left\lceil \frac{3^k}{2^n} \phi \right\rceil = \left\lceil 3^k Col_3^m(\phi) \right\rceil = \left\lceil 3^k Col_3^m(\theta) \right\rceil = \left\lceil \frac{3^k}{2^n} \theta \right\rceil.$$

Proof. Write ϕ and θ in base 3, and divide them by 2 using the long division procedure.

(The assumption $\left\lceil 3^k \phi \right\rceil \neq 3^k \phi$ and $\left\lceil 3^k \theta \right\rceil \neq 3^k \theta$ ensures that the roundings on all the iterations occur outside of the first k digits (in base 3).

Observation: Since $Col_3^m(b) \rightarrow 0$, when $m \rightarrow \infty$, the Lemma 3.1 becomes irrelevant for m big enough. To keep it meaningful, we restrict it to $m \leq n$, where n is defined by

$$(3.3) \quad \left\lceil 3^k Col_3^n(\phi) \right\rceil = \left\lceil 3^k Col_3^n(\theta) \right\rceil = 1.$$

After the next iteration, all the meaningful digits of $Col_3^{n+1}(\phi)$ and $Col_3^{n+1}(\theta)$ (in base 3) disappear from the “window” we are observing (the first k positions) and there is nothing more we can say from (3.1).

$$\left[3^k Col_3^{n+1}(\phi)\right] = \left[3^k Col_3^{n+1}(\theta)\right] = 0.$$

Definition 3.2. Given an integer $\xi \geq 1$, put $q_0 = \left\lceil \frac{\ln \xi}{\ln 3} \right\rceil$ and denote

$$(3.4) \quad \Phi_\xi = \left\{ b \in \mathbb{N} \setminus \bigcup_{r=0}^{q_0} 3^{-r} \mathbb{N}; [3^{q_0} b] = \xi \right\}$$

the set of all the numbers $b \in [1, 3)$, whose digital expansion in base 3 matches the digits in the expansion of ξ , in the same order, but in a shifted position. In particular, $\Phi_\xi \subset [1, 3)$.

Corollary 3.3. For $b \in \Phi_\xi$ and $n \in \mathbb{N}$, define

$$(3.5) \quad k_1 = - \left\lceil \frac{\ln Col_3^n(b)}{\ln 3} \right\rceil \text{ and } k_2 = - \left\lceil \frac{\ln b}{\ln 3} - \frac{n \ln 2}{\ln 3} \right\rceil.$$

Then, as long as $k_1 \leq q_0$ or $k_2 \leq q_0$, we have $k_1 = k_2$ and their common value, k , does not depend on $b \in \Phi_\xi$, only on n .

In particular, for all pairs (n, k) , $k \leq q_0$ verifying (3.5),

$$(3.6) \quad Col_4^n(b) = 3^k \cdot Col_3^n(b)$$

for all $b \in \Phi_\xi$. (See Lemma 2.3.)

For further reference, let's write this conclusion in other form.

Corollary 3.4. The pairs (n, k) in Corollary 3.3 are exactly the solutions of

$$(3.7) \quad 1 \leq 3^k \cdot Col_3^n(b) < 3 \text{ and/or } 1 \leq 3^k \cdot b \cdot 2^{-n} < 3.$$

for which $k \leq q_0$.

Note that given n , k is unique, while for each k , there might be two values of n verifying (3.7), but not more than two.

Lemma 3.5. There is a sequence (n_j, k_j) (increasing in both components), such that for all real number x , $1 < x < 3$, there is $j_0 \in \mathbb{N}$ such that for all $j \geq j_0$,

$$(3.8) \quad k_j \frac{\ln 3}{\ln 2} < n_j \leq k_j \frac{\ln 3}{\ln 2} + \frac{\ln x}{\ln 2}.$$

(In particular, (n_j, k_j) verifies $1 \leq 3^k \cdot x \cdot 2^{-n} < 3$, see (3.7).)

Proof. Since $\frac{\ln 3}{\ln 2}$ is irrational,

$$\limsup_{k \rightarrow \infty, k \in \mathbb{N}} \left(k \cdot \frac{\ln 3}{\ln 2} - \left\lceil k \cdot \frac{\ln 3}{\ln 2} \right\rceil \right) = 1.$$

So, there is an increasing sequence $k_j \in \mathbb{N}$ such that

$$\lim_{j \rightarrow \infty} k_j \cdot \frac{\ln 3}{\ln 2} - \left[k_j \cdot \frac{\ln 3}{\ln 2} \right] = 1.$$

Put

$$n_j = \left[k_j \cdot \frac{\ln 3}{\ln 2} \right] + 1.$$

So,

$$(3.9) \quad k_j \cdot \frac{\ln 3}{\ln 2} < n_j < k_j \cdot \frac{\ln 3}{\ln 2} + 1.$$

Since $\ln x > 0$ and

$$\lim_{j \rightarrow \infty} n_j - k_j \cdot \frac{\ln 3}{\ln 2} = 0,$$

there exist j_0 such that for all $j \geq j_0$,

$$(3.10) \quad n_j < k_j \cdot \frac{\ln 3}{\ln 2} + \frac{\ln x}{\ln 2}.$$

By (3.9) and (3.10), we have (3.8).

4. PROOFS

4.1. Proof of Proposition 2.4. Assume that $l = \liminf_{\alpha \rightarrow \infty} c_\alpha > 1$.

Let (n_j, k_j) be a sequence given by Lemma 3.5 and fix $0 < \epsilon < \frac{\ln l}{\ln 2}$. By (3.8), there is j_0 such that for all $j \geq j_0$,

$$(4.1) \quad k_j \cdot \frac{\ln 3}{\ln 2} < n_j < k_j \cdot \frac{\ln 3}{\ln 2} + \epsilon.$$

For later reference, note that

$$(4.2) \quad n_j \cdot \frac{\ln 2}{\ln 3} - \epsilon \cdot \frac{\ln 2}{\ln 3} < k_j < n_j \cdot \frac{\ln 2}{\ln 3}.$$

Since $\epsilon \cdot \frac{\ln 2}{\ln 3} < \frac{\ln l}{\ln 3} < 1$,

$$(4.3) \quad k_j = \left[n_j \cdot \frac{\ln 2}{\ln 3} \right].$$

Since $\lim_{j \rightarrow \infty} 3^{-k_j} [3^{k_j} l - 1] = l$, we choose $j_1 \geq j_0$ such that

$$(4.4) \quad \ln 3^{-k_{j_1}} [3^{k_{j_1}} l - 1] > \epsilon \ln 2.$$

Now, choose a sequence $(c_{\alpha_i})_{i \in \mathbb{N}}$, converging to l , such that for all i ,

$$(4.5) \quad \left[3^{k_{j_1}} c_{\alpha_i} \right] = \left[3^{k_{j_1}} l \right].$$

If no such sequence exist, necessarily $l \in \mathbf{N}$ and any sequence c_{α_i} going to l approach l by below. In that case, we can choose c_{α_i} such that for all i ,

$$(4.6) \quad \left[3^{k_{j_1}} c_{\alpha_i} \right] = \left[3^{k_{j_1}} l - 1 \right].$$

The key point is we have chosen $c_{\alpha_i} \rightarrow l$ such that $\left[3^{k_{j_1}} c_{\alpha_i} \right]$ does not depend on i . In other words, taking $\xi = \left[3^{k_{j_1}} c_{\alpha_i} \right]$ (for some i),

$$(4.7) \quad (c_{\alpha_i})_{i \in \mathbf{N}} \subseteq \Phi_\xi.$$

Now, observe that

$$(4.8) \quad c_{\alpha_i} \geq 3^{-k_{j_1}} \left[3^{k_{j_1}} c_{\alpha_i} \right] \geq 3^{-k_{j_1}} \left[3^{k_{j_1}} l - 1 \right].$$

Then, by (4.4), for all i ,

$$(4.9) \quad \ln c_{\alpha_i} > \epsilon \ln 2$$

and, by (4.2),

$$(4.10) \quad -k_{j_1} < \frac{\epsilon \ln 2}{\ln 3} - \frac{n_{j_1} \ln 2}{\ln 3} < \frac{\ln c_{\alpha_i}}{\ln 3} - \frac{n_{j_1} \ln 2}{\ln 3}.$$

Therefore, by Corollary 3.3,

$$(4.11) \quad k = - \left[\frac{\ln Col_3^{n_{j_1}}(c_{\alpha_i})}{\ln 3} \right] = - \left[\frac{\ln c_{\alpha_i}}{\ln 3} - \frac{n_{j_1} \ln 2}{\ln 3} \right] \leq k_{j_1}$$

is constant and, by (3.6),

$$(4.12) \quad Col_4^{n_{j_1}}(c_{\alpha_i}) = 3^k \cdot Col_3^{n_{j_1}}(c_{\alpha_i}) \leq 3^{k_{j_1}} \cdot Col_3^{n_{j_1}}(c_{\alpha_i}).$$

By (2.5),

$$(4.13) \quad 2^{-n_{j_1}} c_{\alpha_i} \leq Col_3^{n_{j_1}}(c_{\alpha_i}) \leq 2^{-n_{j_1}} c_{\alpha_i} + 3^{-q_{\alpha_i}},$$

where q_{α_i} is given by (2.1) (or equivalently by (2.4)). Note that $q_{\alpha_i} \rightarrow \infty$, since $a_{\alpha} \rightarrow \infty$.

Putting (4.12) and (4.13) together,

$$(4.14) \quad Col_4^{n_{j_1}}(c_{\alpha_i}) \leq \frac{3^{k_{j_1}}}{2^{n_{j_1}}} c_{\alpha_i} + 3^{k_{j_1} - q_{\alpha_i}}.$$

Finally, since

$$(4.15) \quad Col_4^{n_j}(c_{\alpha_i}) = c_{\alpha_i + n_j}$$

then

$$(4.16) \quad l \leq \liminf_{i \rightarrow \infty} c_{\alpha_i + n_j} \leq \frac{3^{k_j}}{2^{n_j}} l.$$

Which is a contradiction, since $\frac{3^{k_j}}{2^{n_j}} < 1$.

4.2. Proof of Theorem 1.1. Given an integer $\psi \geq 1$, set

$$(4.17) \quad p = \left\lceil \frac{\ln \psi}{\ln 3} \right\rceil \quad \text{and} \quad \psi_0 = 3^{-p} \psi.$$

Put

$$(4.18) \quad \tau = \frac{\ln \psi_0}{\ln 2} - \left\lfloor \frac{\ln \psi_0}{\ln 2} \right\rfloor \left(\frac{\ln 3}{\ln 2} - 1 \right).$$

In other words,

$$(4.19) \quad \tau = \begin{cases} \frac{\ln \psi_0}{\ln 2}, & \text{if } \left\lfloor \frac{\ln \psi_0}{\ln 2} \right\rfloor = 0, \\ \frac{\ln \psi_0}{\ln 2} + 1 - \frac{\ln 3}{\ln 2}, & \text{if } \left\lfloor \frac{\ln \psi_0}{\ln 2} \right\rfloor = 1. \end{cases}$$

Since $1 \leq \psi_0 < 3$, τ is completely defined and $0 \leq \tau < 1$.

By the irrationality of $\frac{\ln 3}{\ln 2}$, there is a sequence k_j such that

$$(4.20) \quad \lim_{j \rightarrow \infty} k_j \cdot \frac{\ln 3}{\ln 2} - \left\lfloor k_j \cdot \frac{\ln 3}{\ln 2} \right\rfloor = \tau.$$

For all $\epsilon > 0$, there is j_0 such that for all $j \geq j_0$,

$$(4.21) \quad 2^{-\epsilon} \cdot 2^\tau < \frac{3^{k_j}}{2^{n_j}} < 2^\epsilon \cdot 2^\tau,$$

where

$$(4.22) \quad n_j = \left\lfloor k_j \cdot \frac{\ln 3}{\ln 2} \right\rfloor.$$

Now, fix

$$(4.23) \quad 0 < \epsilon < \frac{\ln(1 + 3^{-p-2})}{\ln 2}.$$

Since

$$(4.24) \quad 2^\tau = \begin{cases} \psi_0 & \text{if } 1 \leq \psi_0 < 2, \\ \frac{2}{3}\psi_0 & \text{if } 2 \leq \psi_0 < 3. \end{cases},$$

multiplying all members of (4.21) by 3^p , for all j big enough, we obtain

$$(4.25) \quad \psi - \frac{1}{3} < 2^{-\epsilon} \psi < \frac{3^{k_j+p}}{2^{n_j}} < 2^\epsilon \psi < \psi + \frac{1}{3}$$

if $1 \leq \psi_0 < 2$, or

$$(4.26) \quad \psi - \frac{1}{3} < 2^{-\epsilon} \psi < \frac{3^{k_j+1+p}}{2^{n_j+1}} < 2^\epsilon \psi < \psi + \frac{1}{3}$$

otherwise.

Now, fix j verifying (4.21) and take $\xi = 3^{k_j+q}$ in Definition 3.2, $q \in \mathbb{N}$ will be determined later. By Proposition 2.4, there is a subsequence

$$(4.27) \quad (c_{\alpha_i})_{i \in \mathbb{N}} \subset \Phi_{3^{k_j+q}}.$$

By Corolary 3.3, for all i ,

$$(4.28) \quad Col_4^{n_j}(c_{\alpha_i}) = 3^{k_j} Col_3^{n_j}(c_{\alpha_i}).$$

Since $1 < c_{\alpha_i} < 1 + 3^{-k_j-q}$,

$$(4.29) \quad 2^{-n_j} < Col_3^{n_{j1}}(c_{\alpha_i}) < 2^{-n_{j1}} + 2 \cdot 3^{-k_j-q}.$$

Then,

$$(4.30) \quad \frac{3^{k_j}}{2^{n_j}} < Col_4^{n_j}(c_{\alpha_i}) < \frac{3^{k_j}}{2^{n_j}} + \frac{2}{3^q}.$$

If $1 \leq \psi_0 < 2$, put $q = p+1$. By (4.25) and (4.30), since $Col_4^{n_j}(c_{\alpha_i}) = c_{\alpha_i+n_j}$, for all i ,

$$(4.31) \quad \psi - \frac{1}{3} < 3^p c_{\alpha_i+n_j} < \psi + 1.$$

Then, for all i ,

$$(4.32) \quad \left[3^{p-q\alpha_i+n_j} a_{\alpha_i+n_j} \right] = \left[3^p c_{\alpha_i+n_j} \right] = \psi.$$

If $2 \leq \psi_0 < 3$, note that

$$(4.33) \quad 1 < \frac{3^{k_j}}{2^{n_j}} < 2$$

Choose $q \geq p+2$, such that

$$(4.34) \quad 1 < \frac{3^{k_j}}{2^{n_j}} < c_{\alpha_i+n_j} < \frac{3^{k_j}}{2^{n_j}} + \frac{2}{3^q} < 2.$$

Since $[c_{\alpha_i+n_j}] = 1$, then $[Col_3(c_{\alpha_i+n_j})] = 0$ and

$$(4.35) \quad c_{\alpha_i+n_j+1} = Col_4(c_{\alpha_i+n_j}) = 3 \cdot Col_3(c_{\alpha_i+n_j})$$

$$(4.36) \quad \frac{3}{2} \cdot c_{\alpha_i+n_j} \leq c_{\alpha_i+n_j+1} \leq \frac{3}{2} \cdot c_{\alpha_i+n_j} + \frac{3^{-q\alpha_i+n_j}}{2}.$$

Then,

$$(4.37) \quad \frac{3^{k_j+1}}{2^{n_j+1}} < c_{\alpha_i+n_j+1} < \frac{3^{k_j+1}}{2^{n_j+1}} + \frac{1}{3^{q-1}} + \frac{3^{-q\alpha_i+n_j}}{2}.$$

By (4.26),

$$(4.38) \quad \psi - \frac{1}{3} < 3^p \cdot c_{\alpha_i+n_j+1} < \psi + 1,$$

(since $q_\alpha \rightarrow \infty$, we can assume that $3^{p+1-q\alpha_i+n_j} < 1$).

Then, for all i ,

$$(4.39) \quad \left[3^{p-q_{\alpha_i+n_j+1}} a_{\alpha_i+n_j+1} \right] = \left[3^p c_{\alpha_i+n_j+1} \right] = \psi.$$

An immediate consequence of Theorem 1.1 is the following corollary.

Corollary 4.1.

$$(4.40) \quad \overline{\{c_\alpha\}_{\alpha \in \mathbb{N}}} = [1, 3].$$

Let's define an order relation for integers $\phi \geq 1$ and $\theta \geq 1$ in the following way

$$(4.41) \quad \phi \preceq \theta \iff \phi = \left\lceil 3^{p-q} \theta \right\rceil,$$

where $p = \left\lceil \frac{\ln \phi}{\ln 3} \right\rceil$ and $q = \left\lceil \frac{\ln \theta}{\ln 3} \right\rceil$. We finish by noticing the following corollary.

Corollary 4.2. *For any integer $\phi \geq 1$, there is a subsequence $(a_{\alpha_i})_{i \in \mathbb{N}}$ increasing with respect to the order $\preceq$ defined above, minored by ϕ .*

REFERENCES

1. Terence Tao, *Almost all orbits of the collatz map attain almost bounded values*, <https://arxiv.org/abs/1909.03562v2>, 2019.
2. ———, *The notorious collatz conjecture*, <https://terrytao.files.wordpress.com/2020/02/collatz.pdf>, 2020.

DMAT, UNIVERSIDADE DE ÉVORA, ÉVORA - PORTUGAL
Email address: salazar@uevora.pt