arXiv:2003.05860v1 [math.SG] 12 Mar 2020

Real Topological Recursions and WDVV Relations

Aleksey Zinger*

September 19, 2021

Abstract

Following a question of K. Hori at K. Fukaya's 60th birthday conference, we relate the recently established WDVV-type relations for real Gromov-Witten invariants to topological recursion relations in a real setting. We also describe precisely the connections between the relations themselves previously observed by A. Alcolado.

0 Introduction

Gromov-Witten invariants of a compact symplectic manifold (X,ω) are counts of J-holomorphic curves in X for an ω-tame almost complex structure J. They are governed by Kontsevich-Manin's axioms [17]. By these axioms, relations between homology classes on the Deligne-Mumford moduli space $\overline{\mathcal{M}}_{g,l}$ of stable nodal complex genus g curves with l marked points lift directly to relations between Gromov-Witten invariants. As an example, the equality of the points in the 0-th homology of $\overline{\mathcal{M}}_{0,4}\approx\mathbb{P}^1$ represented by the curves on the top line of Figure 1 implies that the counts of the nodal J-holomorphic curves in X represented by the two diagrams on this line are also the same. Along with the Splitting Axiom 2.2.6 in [17], this equality leads to quadratic relations between counts of genus 0 J-holomorphic curves in different degrees (homology classes). These relations can be expressed in terms of partial differential equations (10), known as WDVV equations, on the generating function Φ_ω of (9) for the genus 0 Gromov-Witten invariants of (X,ω). The WDVV equations, first obtained in [17] and mathematically confirmed for semi-positive symplectic manifolds in [20], completely determine counts of rational curves of a specified degree and passing through specified constraints in many smooth algebraic varieties from elementary input; see Sections 5.2 in [17], 10 in [20], and 3 in [15]. The trivial relation in $H_0(\overline{\mathcal{M}}_{0,4})$ represented by the top line in Figure 1 is a basic example of a tautological recursion relation of Theorem 1.

A real symplectic manifold is a triple (X,ω,ϕ) consisting of a symplectic manifold (X,ω) and an anti-symplectic involution ϕ on (X,ω), i.e. a diffeomorphism ϕ of X with itself so that $\phi^2=\mathrm{id}_X$ and $\phi^*\omega=-\omega$. The fixed locus X^ϕ of ϕ is then a Lagrangian submanifold of (X,ω). The prototypical example is the complex projective space $\mathbb{P}^n$ with the Fubini-Study symplectic form $\omega=\omega_{\mathrm{FS}}$ and the standard complex conjugation

$$\phi=\tau_n\colon\mathbb{P}^n\longrightarrow\mathbb{P}^n,\qquad \tau_n\big([Z_0,\ldots,Z_n]\big)=\big[\overline{Z_0},\ldots,\overline{Z_n}\big].$$

A J-holomorphic curve C in (X,ω,ϕ) is called real if $\phi(C)=C$. Counts of real rational J-holomorphic curves through generic k-tuples of points in a topological component Y of X^ϕ and

*Partially supported by NSF grants DMS 1500875 and 1901979




Figure 1: Bordisms relations between real codimension 2 co-oriented cycles of nodal curves on $\overline{\mathcal{M}}_{0,4}$, $\mathbb{R}\overline{\mathcal{M}}_{0,1,2}$, and $\mathbb{R}\overline{\mathcal{M}}_{0,0,3}$, respectively.

l pairs of ϕ-conjugate points in $X - X^\phi$ dependent only on the degree of the curves, k, and l were first defined in [25] for compact real symplectic fourfolds and in [26] for compact real symplectic sixfolds with Y orientable. Building on earlier work of Fukaya-Oh-Ohta-Ono [11] and C.-C. Liu [18] on the structure and orientability of moduli spaces of stable J-holomorphic maps from bordered Riemann surfaces, Solomon [21] and Georgieva [12] interpret Welschinger's invariants from moduli-theoretic perspectives. The (genus 0) real Gromov-Witten invariants of (X, ω, ϕ) are the counts of real J-holomorphic curves arising from these perspectives. The two moduli-theoretic interpretations of Welschinger's invariants have led to extensions of these counts to counts of real rational J-holomorphic curves with two-dimensional conjugate constraints in sixfolds and to counts with even-dimensional conjugate constraints only in higher-dimensional real symplectic manifolds satisfying certain topological conditions.

The two moduli-theoretic interpretations of Welschinger's invariants have also opened the door to at least potentially obtaining relations between Welschinger's invariants by lifting relations between homology classes on the Deligne-Mumford moduli space $\mathbb{R}\overline{\mathcal{M}}_{0,k,l}$ of stable nodal *real* genus 0 curves with k real marked points and l pairs (z_i^+, z_i^-) of conjugate marked points. Building on [21], Solomon's note [22] predicts a pair of WDVV-type relations for Welschinger's invariants of real symplectic fourfolds and the existence of similar relations for Welschinger's invariants of real symplectic sixfolds (whenever they are defined). Alcolado's thesis [1] predicts an explicit pair of WDVV-type relations for analogues of Welschinger's invariants in much wider settings, including those in which such invariants are yet to be defined mathematically. A WDVV-type relation for the real Gromov-Witten invariants defined in [12] is obtained in [13] by establishing the relation between the one-dimensional homology classes on $\mathbb{R}\overline{\mathcal{M}}_{0,0,3}$ represented by the bottom line in Figure 1, lifting it to a relation between counts of real nodal J-holomorphic curves in (X, ω, ϕ), and

expressing these counts in terms of counts of real and complex irreducible curves. The WDVV-type relations for Welschinger's invariants of real symplectic fourfolds predicted in [22] are established in [4] by lifting relations between cycles in $\mathbb{R}\overline{\mathcal{M}}_{0,1,2}$ and $\mathbb{R}\overline{\mathcal{M}}_{0,0,3}$, but over non-orientable forgetful morphisms (3) from moduli spaces of real maps. The same general approach is used in [7] to obtain WDVV-type relations for Welschinger's invariants of real symplectic sixfolds with some symmetry. As shown in [6], these relations completely determine Welschinger's invariants of some important real symplectic fourfolds and sixfolds from elementary input and lead to lower bounds in the real enumerative geometry of rational curves. In the case of $(\mathbb{P}^3, \omega_{\mathrm{FS}}, \tau_3)$, these lower bounds fit nicely with Kollár's vanishing results [16]; see Section 2.6 in [7].

We recall the standard tautological recursion relation of Theorem 1 and its connection with WDVV relations between the standard (complex) Gromov-Witten invariants in Section 1. We formulate a pair of real topological recursion relations in Theorem 2 and describe their connection with the WDVV-type relations between real Gromov-Witten invariants established in [13, 4, 7]. This in particular affirmatively answers a question of K. Hori on whether these real WDVV-type relations arise from some real topological recursion relations. In contrast to the relations of Theorem 1, the relations of Theorem 2 for the generally unorientable moduli spaces $\mathbb{R}\overline{\mathcal{M}}_{0,k,l}$ involve *co-oriented* cycles, i.e. cycles with orientations on the normal bundles. We recall the standard WDVV PDEs and the two sets of real WDDV PDEs in Section 3. In Section 4, we show that the standard WDVV PDEs and one of the sets of real WDDV PDEs implies the other set when there are nonzero real genus 0 Gromov-Witten invariants with at least 2 real marked point constraints. The statement of Theorem 3 in this section and its proof formalize the observation and reasoning originally due to A. Alcolado (as far as the author is aware); they are included in this note for dissemination purposes. Theorem 2 is established in Section 6, after a standard proof of Theorem 1 is recalled in Section 5.

The author would like to thank Xujia Chen for enlightening discussions on real WDVV relations, the organizers of K. Fukaya's 60th birthday conference for their invitation, and K. Hori for raising the question answered by Theorem 2 in the present note.

1 The $\mathbb{C}$ case

For $l \in \mathbb{Z}^{\geq 0}$, we define

$$[l] = \{1, 2, \ldots, l\}.$$

For $l \geq 3$ and $i \in [l]$, let $\psi_i \in H^2(\overline{\mathcal{M}}_{0,l})$ be the first Chern class of the universal cotangent line bundle L_i^* for the first marked point as usual. For a partition $I \sqcup J$ of $[l]$, we denote by $D_{I,J} \subset \overline{\mathcal{M}}_{0,l}$ the closure of the subspace of two-component curves so that one of the components carries the marked points indexed by I and the other component carries the marked points indexed by J. In particular, $D_{I,J}$ is a complex divisor in the compact complex manifold $\overline{\mathcal{M}}_{0,l}$; it is empty if either $|I| < 2$ or $|J| < 2$.

Theorem 1 ($\mathbb{C}$ Topological Recursion Relations)**.** *If $i, j \in [l] - \{1\}$ are distinct, then*

$$\psi_1 = \sum_{I \sqcup J = [l] - \{1,i,j\}} \mathrm{PD}_{\overline{\mathcal{M}}_{0,l}}\big(D_{1I,ijJ}\big) \in H^2\big(\overline{\mathcal{M}}_{0,l}; \mathbb{Z}\big). \tag{1}$$

Since $\overline{\mathcal{M}}_{0,l}$ and $D_{1I,ijJ}$ are canonically oriented (by their complex orientations), it is customary to drop $\mathrm{PD}_{\overline{\mathcal{M}}_{0,l}}$. For $l=4$, (1) gives

$$D_{12,34} = \psi_1 = D_{13,24} \in H_0\big(\overline{\mathcal{M}}_{0,4};\mathbb{Z}\big); \tag{2}$$

see the top diagram in Figure 1.

Let (X,ω) be a compact symplectic manifold, $B\in H_2(X;\mathbb{Z})$, and J be an ω-tame almost complex structure on X. We then denote by $\overline{\mathfrak{M}}_{0,l}(B;J)$ the moduli space of stable degree B J-holomorphic maps from genus 0 connected nodal Riemann surfaces with l marked points. For $l'\in[l]-[2]$, there is a natural forgetful morphism

$$\mathfrak{f}\colon \overline{\mathfrak{M}}_{0,l}(B;J) \longrightarrow \overline{\mathcal{M}}_{0,l'} \tag{3}$$

dropping the map component and the last $l-l'$ marked points of each stable map. The genus 0 degree B Gromov-Witten invariants of (X,ω) are obtained by intersecting the l natural evaluation morphisms from the moduli space $\overline{\mathfrak{M}}_{0,l}(B;J)$ with l cycles in X in general position.

For $l\ge 4$, the homology relation (2) lifts via the $l'=4$ case of the morphism (3) to a homology relation between cycles on $\overline{\mathfrak{M}}_{0,l}(B;J)$ generically consisting of maps from two-component curves as on the top line in Figure 1. The latter relation in turn translates into a relation between counts of two-component genus 0 degree B J-holomorphic curves passing through l cycles. Replacing the node condition by the splitting of the diagonal as in (8), one then obtains a quadratic relation between the genus 0 Gromov-Witten invariants of (X,ω). The collection of these quadratic relations taken over all B and possible choices of the constraints is equivalent to the PDEs (10) and to the associativity of the quantum product. However, the original proof of the associativity in [20] instead involved defining genus 0 Gromov-Witten invariants for each element of $\overline{\mathcal{M}}_{0,4}$ and showing that they are in fact independent of the chosen element (this is more rigid than taking a path between $D_{12,34}$ and $D_{13,24}$ and lifting it to paths in the moduli spaces of maps).

Let $\mathfrak{o}^c_{D_{1I,ijJ}}$ denote the orientation of the normal bundle $\mathcal{N}D_{1I,ijJ}$ induced by the complex orientations of $\overline{\mathcal{M}}_{0,l}$ and $D_{1I,ijJ}$. The relation (2) can then be written as

$$\mathrm{PD}_{\overline{\mathcal{M}}_{0,4}}\big(D_{12,34},\mathfrak{o}^c_{D_{12,34}}\big) = \psi_1 = \mathrm{PD}_{\overline{\mathcal{M}}_{0,4}}\big(D_{13,24},\mathfrak{o}^c_{D_{13,24}}\big) \in H^2\big(\overline{\mathcal{M}}_{0,4};\mathbb{Z}\big).$$

The equality of the left- and right-hand sides above depends only on choosing the co-orientations $\mathfrak{o}^c_{D_{12,34}}$ and $\mathfrak{o}^c_{D_{12,34}}$ consistently, not on the orientations of $\overline{\mathcal{M}}_{0,4}$ and $D_{1I,ijJ}$. This form of (2) is more convenient for lifting by maps between non-orientable spaces.

2 The $\mathbb{R}$ case

For $k,l\in\mathbb{Z}^{\ge 0}$ with $k+2l\ge 2$ and $i\in[l]$, let $\psi_i\in H^2(\mathbb{R}\overline{\mathcal{M}}_{0,k,l})$ be the first Chern class of the universal cotangent line bundle L_i^* for the first marked point z_i^+ in the i-th conjugate pair (z_i^+,z_i^-) of marked points. For a partition $I\sqcup J$ of $[l]$, we denote by $\mathbb{R}D_{I,J}\subset\mathbb{R}\overline{\mathcal{M}}_{0,k,l}$ the closure of the subspace of three-component curves, with one real component (preserved by the involution) and a conjugate pair of components (interchanged by the involution), so that the conjugate components carry the conjugate pairs of marked points indexed by I and the real component carries the conjugate pairs of marked points indexed by J (along with all real marked points). In particular, $\mathbb{R}D_{I,J}$ is a compact submanifold of real codimension 2; it is empty unless $|I|\ge 2$ and either $J\neq\emptyset$ or $k>0$. Each

non-empty $\mathbb{R}D_{I,J}$ has precisely $2^{|I|-1}$ topological components indexed by the unordered partitions of the marked points z_i^+ with $i\in I$ between the two conjugate components of each domain curve.

The normal bundle $\mathcal{N}\mathbb{R}D_{I,J}$ of $\mathbb{R}D_{I,J}$ in $\mathbb{R}\overline{\mathcal{M}}_{0,k,l}$ consists of conjugate pairs of smoothings of the two nodes. We denote by $\mathfrak{o}^c_{\mathbb{R}D_{I,J}}$ the co-orientation of $\mathbb{R}D_{I,J}$ in $\mathbb{R}\overline{\mathcal{M}}_{0,k,l}$ obtained from the complex orientation of the smoothings of the node separating off the conjugate component carrying the marked point z_i^+ with the smallest value of i in I.

Theorem 2 ($\mathbb{R}$ Topological Recursion Relations)**.** *If $k>0$, then*

$$\psi_1 = \sum_{I\sqcup J=[l]-\{1\}} \mathrm{PD}_{\mathbb{R}\overline{\mathcal{M}}_{0,k,l}}\big(\mathbb{R}D_{1I,J}, \mathfrak{o}^c_{\mathbb{R}D_{1I,J}}\big) \in H^2\big(\mathbb{R}\overline{\mathcal{M}}_{0,k,l};\mathbb{Z}\big).$$

If $i\in[l]-\{1\}$, then

$$\psi_1 = \sum_{I\sqcup J=[l]-\{1,i\}} \mathrm{PD}_{\mathbb{R}\overline{\mathcal{M}}_{0,k,l}}\big(\mathbb{R}D_{1I,iJ}, \mathfrak{o}^c_{\mathbb{R}D_{1I,iJ}}\big) \in H^2\big(\mathbb{R}\overline{\mathcal{M}}_{0,k,l};\mathbb{Z}\big).$$

For $(k,l)=(1,2)$ and $(k,l)=(0,3)$, Theorem 2 gives

$$\mathrm{PD}_{\mathbb{R}\overline{\mathcal{M}}_{0,1,2}}\big(\mathbb{R}D_{12,\emptyset}, \mathfrak{o}^c_{\mathbb{R}D_{12,\emptyset}}\big) = \psi_1 = 0 \in H^2\big(\mathbb{R}\overline{\mathcal{M}}_{0,1,2};\mathbb{Z}\big), \tag{4}$$

$$\mathrm{PD}_{\mathbb{R}\overline{\mathcal{M}}_{0,0,3}}\big(\mathbb{R}D_{12,3}, \mathfrak{o}^c_{\mathbb{R}D_{12,3}}\big) = \psi_1 = \mathrm{PD}_{\mathbb{R}\overline{\mathcal{M}}_{0,0,3}}\big(\mathbb{R}D_{13,2}, \mathfrak{o}^c_{\mathbb{R}D_{13,2}}\big) \in H^2\big(\mathbb{R}\overline{\mathcal{M}}_{0,0,3};\mathbb{Z}\big). \tag{5}$$

The equality of the left- and right-hand sides of (4), depicted by the middle row in Figure 1, is immediate because each of the two sides of (4) represents a point in $\mathbb{R}\overline{\mathcal{M}}_{0,1,2}\approx\mathbb{RP}^2$. The equality of the left- and right-hand sides of (5) in H_1 of the orientable space $\mathbb{R}\overline{\mathcal{M}}_{0,0,3}$, depicted by the bottom row in Figure 1, was established in [13] based on a direct geometric study of $\mathbb{R}\overline{\mathcal{M}}_{0,0,3}$.

Let (X,ω,ϕ) be a compact real symplectic manifold. Define

$$H_2^\phi(X;\mathbb{Z}) \equiv \big\{B'\in H_2(X;\mathbb{Z})\colon \phi_*(B')=-B'\big\}, \quad H^{2*}(X;\mathbb{Q})^\phi_\pm \equiv \big\{\gamma\in H^{2*}(X;\mathbb{Q})\colon \phi^*\gamma=\pm\gamma\big\}.$$

Let $B\in H_2(X;\mathbb{Z})$ and J be an ω-tame almost complex structure on X such that $\phi^*J=-J$. We then denote by $\overline{\mathfrak{M}}^\phi_{0,k,l}(B;J)$ the moduli space of stable degree B J-holomorphic maps u from genus 0 connected nodal symmetric Riemann surfaces (Σ,σ) with k real marked points $x_i\in\Sigma^\sigma$ and l pairs of conjugate marked points $z_i^\pm\in\Sigma-\Sigma^\sigma$ such that $\phi\circ u=u\circ\sigma$. This space is empty unless $B\in H_2^\phi(X;\mathbb{Z})$. For $k'\in[k]$ and $l'\in[l]$ with $k'+2l'\ge3$, there is a natural forgetful morphism

$$\mathfrak{f}\colon \overline{\mathfrak{M}}^\phi_{0,k,l}(B;J) \longrightarrow \mathbb{R}\overline{\mathcal{M}}_{0,k',l'} \tag{6}$$

dropping the map component, the last k–k' real marked points of each stable map, and the last l–l' pairs of conjugate marked points. The genus 0 real degree B Gromov-Witten invariants of (X,ω,ϕ), whenever defined, are obtained by intersecting the k evaluation morphisms from $\overline{\mathfrak{M}}^\phi_{0,k,l}(B;J)$ to X^ϕ and the l evaluation morphisms to X (taken at z_i^+) with k points in X^ϕ and l cycles in X in general position. In contrast to the $\mathbb{C}$ case of Section 1, the total evaluation morphism

$$\mathrm{ev}\equiv\big(\mathrm{ev}_1^{\mathbb{R}},\ldots,\mathrm{ev}_k^{\mathbb{R}},\mathrm{ev}_1^+,\ldots,\mathrm{ev}_l^+\big)\colon \overline{\mathfrak{M}}^\phi_{0,k,l}(B;J) \longrightarrow (X^\phi)^k\times X^l \tag{7}$$




Figure 2: The left-hand sides of the two identities represent codimension 2 cycles with co-orientations in $\mathbb{R}\overline{\mathcal{M}}_{0,1,2}$ and $\mathbb{R}\overline{\mathcal{M}}_{0,0,3}$ that bound co-oriented hypersurfaces $\Upsilon_{1,2}$ and $\Upsilon_{0,3}$, respectively, and their lifts to $\overline{\mathfrak{M}}^{\phi}_{0,k,l}(B;J)$. The right-hand sides represent the corrections to the associated lifted relations from the crossings of the lifts of $\Upsilon_{1,2}$ and $\Upsilon_{0,3}$ with the codimension 1 strata $\mathcal{S}$ that obstruct the relative orientability of (7). The involutions in all cases are the reflections around the horizontal center line.

is generally not even relatively orientable in the present setting.

If $k=0$, $l\ge 3$, and (X,ϕ) satisfies certain topological conditions specified in [12, 8], the homology relation (5) lifts via the $(k',l')=(0,3)$ case of the morphism (6) to a homology relation between cycles on $\overline{\mathfrak{M}}^{\phi}_{0,0,l}(B;J)$ generically consisting of maps from three-component curves, with one real component and a conjugate pair of components as on the bottom line in Figure 1. The latter relation in turn translates into a relation between counts of real three-component genus 0 degree B J-holomorphic curves passing through l cycles. Replacing the node conditions by the splitting of the diagonal as in (8) yields the bilinear relations between the real genus 0 Gromov-Witten invariants of (X,ω,ϕ) with conjugate pairs of insertions and the standard genus 0 Gromov-Witten invariants of (X,ω) obtained in [13]. The collection of these bilinear relations taken over all B and possible choices of the constraints is equivalent to a subset of the PDEs (13); see Section 3 for more details.

If $k\ge 1$, the $(k',l')=(1,2),(0,3)$ cases of the morphisms (6) are generally not relatively orientable. The equality of the left- and right-hand sides of (4) as co-oriented cycles in $\mathbb{R}\overline{\mathcal{M}}_{0,1,2}$ and the equality of the left- and right-hand sides of (5) as co-oriented cycles $\mathbb{R}\overline{\mathcal{M}}_{0,0,3}$ are lifted in [4, 7] over these morphisms *along with* co-oriented bordisms $\Upsilon_{1,2}\subset\mathbb{R}\overline{\mathcal{M}}_{0,1,2}$ and $\Upsilon_{0,3}\subset\mathbb{R}\overline{\mathcal{M}}_{0,0,3}$ between the two sides. The intersections of the lifts of $\Upsilon_{1,2}$ and $\Upsilon_{0,3}$ with the codimension 1 strata $\mathcal{S}$ that obstruct the relative orientability of (7) determine corrections to lifting the relations from $\mathbb{R}\overline{\mathcal{M}}_{0,1,2}$ and $\mathbb{R}\overline{\mathcal{M}}_{0,0,3}$ directly. All such codimension 1 strata $\mathcal{S}$ consist of J-holomorphic maps from a pair of real components joined at a pair of real points which satisfy a certain numerical condition $\epsilon(\mathcal{S})\cong_4 2$ as on the right-hand sides of the two equalities in Figure 2. The corrected lifted relations in turn translate to relations between counts of real three- and two-component genus 0 degree B J-holomorphic curves as in Figure 2 passing through k points in X^{ϕ} and l cycles in X. Splitting

the nodal counts into counts of irreducible curves yields relations between the real genus 0 Gromov-Witten invariants of (X,ω,ϕ) and the standard genus 0 Gromov-Witten invariants of (X,ω). The collection of these relations taken over all B and possible choices of the constraints is equivalent to the PDEs (12) and (13).

The open/real WDVV-type relations in dimension 2 obtained in [4] have the same structure as predicted in [22], but differ by crucial signs. The former lead to the recursions for Welshinger's invariants predicted in [22] due to the comparison between the invariants of [25] and [21] predicted in [22] also being off by signs. The approach to the proof of real WDVV-type relations proposed in [22] is analogous in spirit to [20] and involves defining a count of real curves for each element of $\mathbb{R}\overline{\mathcal{M}}_{0,1,2}$ and $\mathbb{R}\overline{\mathcal{M}}_{0,0,3}$, showing that these counts depend only on the homology classes of the insertions, and are independent of the choice of the element in each of the two moduli spaces. The desired invariance in fact does not hold in the case of $\mathbb{R}\overline{\mathcal{M}}_{0,1,2}$. The proposed approach itself is more rigid than the lifting-of-cobordisms approach introduced in [4] and does not connect as readily with the topological recursion relations.

Let (X,ω,ϕ) be a compact real symplectic sixfold with a Spin-structure on X^ϕ. The real genus 0 Gromov-Witten invariants (11) of (X,ω,ϕ) appearing in [7] differ from those in [21] by uniform multiples (dependent only on the number l of conjugate pairs of marked points) and vanish on insertions from $H^2(X;\mathbb{Q})^\phi_+$ and $H^4(X;\mathbb{Q})^\phi_-$. The $k=0$ invariants (11) defined in [12] agree with the $k=0$ invariants appearing in [7] when the insertions are taken in $H^0(X;\mathbb{Q})$, $H^2(X;\mathbb{Q})$, and $H^6(X;\mathbb{Q})$, but vanish on $H^4(X;\mathbb{Q})^\phi_+$. In particular, the invariants of [12] in the case of real symplectic sixfolds with insertions in $H^4(X;\mathbb{Q})$ are not the same as the $k=0$ cases of the invariants arising from [21, 7].

3 Complex and real WDVV PDEs

Given a compact oriented even-dimensional manifold X, we choose a basis $\mu_1^\bigstar,\ldots,\mu_N^\bigstar$ for $H^{2*}(X;\mathbb{Q})$ consisting of homogeneous elements. Let $(g_{ij})_{i,j}$ be the $N\times N$-matrix given by

$$g_{ij}=\big\langle \mu_i^\bigstar\mu_j^\bigstar,[X]\big\rangle$$

and $(g^{ij})_{ij}$ be its inverse. Thus, the Poincare dual of the diagonal $\Delta_X\subset X^2$ is given by

$$\mathrm{PD}_{X^2}\big(\Delta_X\big)=\sum_{i,j\in[N]} g^{ij}\mu_i^\bigstar\times\mu_j^\bigstar\in H^2(X^2;\mathbb{Q})\big/\big(H^{\mathrm{odd}}(X;\mathbb{Q})\times H^{\mathrm{odd}}(X;\mathbb{Q})\big); \tag{8}$$

see Theorem 11.11 in [19]. For a tuple $\mathbf{t}\equiv(t_1,\ldots,t_N)$ of formal variables, define

$$\mu_{\mathbf{t}}^\bigstar=\mu_1^\bigstar t_1+\ldots+\mu_N^\bigstar t_N\,.$$

Suppose in addition ω is a symplectic form on X. Let

$$\Lambda_\omega=\big\{(\Psi\colon H_2(X;\mathbb{Z})\longrightarrow\mathbb{Q})\colon \big|\{B\in H_2(X;\mathbb{Z})\colon \Psi(B)\neq 0,\ \omega(B)<E\}\big|<\infty\ \forall\, E\in\mathbb{R}\big\}.$$

We write an element Ψ of Λ_ω as

$$\Psi=\sum_{B\in H_2(X;\mathbb{Z})}\Psi(B)q^B$$

and multiply two such elements as powers series in q with the exponents in $H_2(X;\mathbb{Z})$. For $B\in H_2(X;\mathbb{Z})$, we denote by

$$\langle\ldots\rangle_B^\omega\colon \bigoplus_{l=0}^\infty H^{2*}(X;\mathbb{Q})^{\oplus l}\longrightarrow\mathbb{Q}$$

the genus 0 degree B Gromov-Witten invariants of (X,ω) and extend them linearly over the formal variables t_i above. Define $\Phi_\omega\in\Lambda_\omega[[t_1,\ldots,t_N]]$ by

$$\Phi_\omega(t_1,\ldots,t_N)=\sum_{\substack{B\in H_2(X;\mathbb{Z})\\ l\in\mathbb{Z}^{\ge0}}}\left\langle\underbrace{\mu_{\mathbf{t}}^\bigstar,\ldots,\mu_{\mathbf{t}}^\bigstar}_{l}\right\rangle_B^\omega\frac{q^B}{l!}\,. \tag{9}$$

By Gromov's Compactness Theorem, the coefficients of the powers of $t_1,\ldots,t_N$ in Φ_ω lie in Λ_ω. The relations on the genus 0 Gromov-Witten invariants of (X,ω) obtained by lifting the homology relation on $\overline{\mathcal{M}}_{0,4}$ represented by the top line in Figure 1 is equivalent to the set of WDVV PDEs

$$\sum_{1\le i,j\le N}\!\!\big(\partial_{t_a}\partial_{t_b}\partial_{t_i}\Phi_\omega\big)g^{ij}\big(\partial_{t_j}\partial_{t_c}\partial_{t_d}\Phi_\omega\big)=\sum_{1\le i,j\le N}\!\!\big(\partial_{t_a}\partial_{t_c}\partial_{t_i}\Phi_\omega\big)g^{ij}\big(\partial_{t_j}\partial_{t_b}\partial_{t_d}\Phi_\omega\big) \tag{10}$$

with $a,b,c,d=1,\ldots,N$.

Let ϕ be an anti-symplectic involution on (X,ω). Define

$$\mathfrak{d}\colon H_2(X)\longrightarrow H_2(X)^\phi_-,\qquad \mathfrak{d}(B)=B-\phi_*(B)\,.$$

For a (relatively) Spin or Pin-structure $\mathfrak{s}$ on (the tangent bundle of) X^ϕ, $B\in H_2(X;\mathbb{Z})$, and $k\in\mathbb{Z}^{\ge0}$, we denote by

$$\langle\ldots\rangle_{\mathfrak{s};B;k}^{\omega;\phi}\colon\bigoplus_{l=0}^\infty H^{2*}(X;\mathbb{Q})^{\oplus l}\longrightarrow\mathbb{Q} \tag{11}$$

the real genus 0 degree B Gromov-Witten invariants of (X,ω,ϕ) with k real marked points associated with $\mathfrak{s}$, whenever they are defined. We extend them linearly over the formal variables t_i above. Define $\Phi_\omega^\phi\in\Lambda_\omega[[t_1,\ldots,t_N]]$ and $\Omega_{\omega;\mathfrak{s}}^\phi\in\Lambda_\omega[[t_1,\ldots,t_N,u]]$ by

$$\Phi_\omega^\phi(t_1,\ldots,t_N)=\sum_{\substack{B\in H_2(X;\mathbb{Z})\\ l\in\mathbb{Z}^{\ge0}}}\left(\sum_{\substack{B'\in H_2(X;\mathbb{Z})\\ \mathfrak{d}(B')=B}}\left\langle\underbrace{\mu_{\mathbf{t}}^\bigstar,\ldots,\mu_{\mathbf{t}}^\bigstar}_{l}\right\rangle_{B'}^\omega\right)\frac{q^B}{l!}\,,$$

$$\Omega_{\omega;\mathfrak{s}}^\phi(t_1,\ldots,t_N,u)=\sum_{\substack{B\in H_2(X;\mathbb{Z})\\ k,l\in\mathbb{Z}^{\ge0}}}\left\langle\underbrace{\mu_{\mathbf{t}}^\bigstar,\ldots,\mu_{\mathbf{t}}^\bigstar}_{l}\right\rangle_{\mathfrak{s};B,k}^{\omega;\phi}\frac{2^{1-l}q^Bu^k}{k!l!}\,.$$

By Gromov's Compactness Theorem and the assumption that $\phi^*\omega=-\omega$, the inner sum in the definition of Φ_ω^ϕ has finitely nonzero terms. For the same reason, the coefficients of the powers of $t_1,\ldots,t_N,u$ in Φ_ω^ϕ and $\Omega_{\omega;\mathfrak{s}}^\phi$ lie in Λ_ω.

Alcolado's thesis [1] predicts that the real genus 0 Gromov-Witten invariants of (X,ω,ϕ) satisfy the set of extended WDVV PDEs

$$\sum_{1\le i,j\le N}\big(\partial_{t_a}\partial_{t_b}\partial_{t_i}\Phi^\phi_\omega\big)g^{ij}\big(\partial_{t_j}\partial_u\Omega^\phi_{\omega;\mathfrak{s}}\big)+\big(\partial_{t_a}\partial_{t_b}\Omega^\phi_{\omega;\mathfrak{s}}\big)\big(\partial_u^2\Omega^\phi_{\omega;\mathfrak{s}}\big)=\big(\partial_{t_a}\partial_u\Omega^\phi_{\omega;\mathfrak{s}}\big)\big(\partial_{t_b}\partial_u\Omega^\phi_{\omega;\mathfrak{s}}\big), \tag{12}$$

$$\begin{aligned}\sum_{1\le i,j\le N}&\big(\partial_{t_a}\partial_{t_b}\partial_{t_i}\Phi^\phi_\omega\big)g^{ij}\big(\partial_{t_j}\partial_{t_c}\Omega^\phi_{\omega;\mathfrak{s}}\big)+\big(\partial_{t_a}\partial_{t_b}\Omega^\phi_{\omega;\mathfrak{s}}\big)\big(\partial_{t_c}\partial_u\Omega^\phi_{\omega;\mathfrak{s}}\big)\\ &=\sum_{1\le i,j\le N}\big(\partial_{t_a}\partial_{t_c}\partial_{t_i}\Phi^\phi_\omega\big)g^{ij}\big(\partial_{t_j}\partial_{t_b}\Omega^\phi_{\omega;\mathfrak{s}}\big)+\big(\partial_{t_a}\partial_{t_c}\Omega^\phi_{\omega;\mathfrak{s}}\big)\big(\partial_{t_b}\partial_u\Omega^\phi_{\omega;\mathfrak{s}}\big)\end{aligned} \tag{13}$$

with $a,b,c=1,\ldots,N$, with appropriate definitions of the real invariants. Alcolado's predictions have been confirmed

(C1) in [4] if (X,ω,ϕ) is a real symplectic fourfold, $\mathfrak{s}$ is a Pin$^-$-structure on X^ϕ, and the real invariants are defined as in [4];

(C2) in [7] if (X,ω,ϕ) is a real symplectic sixfold with a finite-order automorphism which restricts to an orientation-reversing diffeomorphism of X^ϕ and acts trivially on the $(-\phi)_*$-invariant part of $H_2(X,Y;\mathbb{Z})$, $\mathfrak{s}$ is a Spin-structure on X^ϕ, and the real invariants are defined as in [7].

A specialization of (13) is obtained in [13] for a real symplectic $2n$-fold (X,ω,ϕ) with n odd which admits a complex vector E with a conjugation $\widetilde{\phi}$ lifting ϕ so that

$$\begin{aligned}w_1(E^{\widetilde{\phi}})^2&=w_2(TX^\phi)\qquad\text{and}\\ \big\langle 2\mu(E,E^{\widetilde{\phi}})+\mu(X,X^\phi),\beta\big\rangle&\in 4\mathbb{Z}\quad\forall\,\beta\in H_2(X,X^\phi;\mathbb{Z})\text{ with }\phi_*\beta=-\beta,\end{aligned} \tag{14}$$

where $E^{\widetilde{\phi}}\longrightarrow X^\phi$ is the fixed locus of the $\widetilde{\phi}$-action on E and $\mu(X,X^\phi)$ and $\mu(E,E^{\widetilde{\phi}})$ are the Maslov indices of the pairs (TX,TX^ϕ) and $(E,E^{\widetilde{\phi}})$, respectively. A Spin-structure $\mathfrak{s}$ on $2E^{\widetilde{\phi}}\oplus TX^\phi$ (along with some additional data if the equality on the second line of (14) does not hold for all $\beta\in H_2(X,X^\phi;\mathbb{Z})$) determines real genus 0 Gromov-Witten invariants (11) with $k=0$; see Theorem 1.6 in [12]. We formally define the functionals (11) with $k\ge 1$ to be zero. Suppose $\mu_1^\bigstar,\ldots,\mu_{N_-}^\bigstar$ and $\mu_{N_-+1}^\bigstar,\ldots,\mu_N^\bigstar$ are bases for $H^{2*}(X;\mathbb{Q})^\phi_-$ and $H^{2*}(X;\mathbb{Q})^\phi_+$, respectively. Theorem 2.1 in [13] is then equivalent to the PDEs (13) with $a\in[N]-[N_-]$ and $b,c\in[N_-]$. By Theorem 2.2 in [13], $\partial_{t_a}\partial_{t_b}\Omega^\phi_{\omega;\mathfrak{s}}$ and $\partial_{t_a}\partial_{t_c}\Omega^\phi_{\omega;\mathfrak{s}}$ in (13) vanish with the definitions in [12] because $\mu_{t_a}^\bigstar\in H^{2*}(X;\mathbb{Q})^\phi_+$.

Motivated by [9, 10], Solomon and Tukachinsky [23] use bounding chains in Fukaya's A_∞-algebras to define counts of J-holomorphic disks in a compact symplectic manifold (X,ω) with boundary in a compact Lagrangian Y with a relative Spin-structure $\mathfrak{s}$. According to Theorem 3 in [24], these counts satisfy the real WDVV PDEs (12) and (13) if the generating function $\Omega^\phi_{\omega;\mathfrak{s}}$ is defined without the factors of 2^{1-l}. As noted in Remark 2.7 of [13] and Theorem 2.10 of [5], the use of a relative Spin structure $\mathfrak{s}$ to orient the moduli spaces of disks leads to the necessity to twist the generating function Φ^ϕ_ω by $(-1)^{\langle w_2(\mathfrak{s}),B'\rangle}$ for the purposes of (12) and (13) (the authors of [23, 24] are currently re-adjusting their signs to account for this). This twist can be avoided if there exists a complex vector bundle $E\longrightarrow X$ and a totally real subbundle $E^{\mathbb{R}}\longrightarrow Y$ of $E|_Y$ of maximal rank so that $w_1(E^{\mathbb{R}})^2=w_2(TY)$. A Spin-structure $\mathfrak{s}$ on $2E^{\mathbb{R}}\oplus TY$ then determines orientations on the

moduli spaces of disk maps to (X,Y) as in Section 7 in [12]. If the construction of [23] were carried out with these orientations, instead of those induced by relative Spin structures as in Section 8.1 of [11], then the PDEs (12) and (13) would hold in [24] without the above sign twist. The degree m holomorphic line $E\equiv\mathcal{O}_{\mathbb{P}^{2m-1}}(m)$ over $\mathbb{P}^{2m-1}$ with the standard conjugation $\widetilde{\phi}$ satisfies (14) and can thus be used to orient the moduli space of stable disk and real maps as in [12]. Theorem 10 in [24], stating the PDEs (12) and (13) for $(\mathbb{P}^{2m-1},\tau_{2m-1})$, requires the use of the resulting orientations, rather than those induced by the associated relative Spin-structures. By Theorem 1.4 in [14], these orientations on the main stratum $\mathfrak{M}_{0,k,l}^{\tau_{2m-1}}(d;J)$ of the moduli space of stable real degree d holomorphic maps to $(\mathbb{P}^{2m-1},\tau_{2m-1})$ are different if and only if $m\notin 2\mathbb{Z}$ and $md\cong 1,2 \bmod 4$.

4 Relation between the PDEs (12) and (13)

We now show that the PDEs (12) often imply the PDEs (13).

Theorem 3. *If the functional (11) is nonzero for some $k\ge 2$, then the relations (12) imply the relations (13).*

Proof. The WDVV PDE (10) for the Gromov-Witten invariants of (X,ω) gives

$$\sum_{1\le l,m\le N}\big(\partial_{t_a}\partial_{t_b}\partial_{t_l}\Phi_\omega^\phi\big)g^{lm}\big(\partial_{t_m}\partial_{t_c}\partial_{t_i}\Phi_\omega^\phi\big)=\sum_{1\le l,m\le N}\big(\partial_{t_a}\partial_{t_c}\partial_{t_l}\Phi_\omega^\phi\big)g^{lm}\big(\partial_{t_m}\partial_{t_b}\partial_{t_i}\Phi_\omega^\phi\big) \tag{15}$$

for all $a,b,c,i=1,\ldots,N$. Multiplying (12) by $\partial_{t_c}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi$ gives

$$\begin{aligned}\big(\partial_u^2\Omega_{\omega;\mathfrak{s}}^\phi\big)\big(\partial_{t_a}\partial_{t_b}\Omega_{\omega;\mathfrak{s}}^\phi\big)\big(\partial_{t_c}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi\big)&=\big(\partial_{t_a}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi\big)\big(\partial_{t_b}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi\big)\big(\partial_{t_c}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi\big)\\&\quad-\big(\partial_{t_c}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi\big)\sum_{1\le i,j\le N}\big(\partial_{t_a}\partial_{t_b}\partial_{t_i}\Phi_\omega^\phi\big)g^{ij}\big(\partial_{t_j}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi\big)\,.\end{aligned} \tag{16}$$

Replacing (a,b) in (12) with (c,m) and multiplying the resulting equation by $\partial_{t_a}\partial_{t_b}\partial_{t_l}\Phi_\omega^\phi g^{lm}$ with $c,l,m=1,\ldots,N$, we obtain

$$\begin{aligned}\big(\partial_u^2\Omega_{\omega;\mathfrak{s}}^\phi\big)\big(\partial_{t_a}\partial_{t_b}\partial_{t_l}\Phi_\omega^\phi\big)g^{lm}\big(\partial_{t_m}\partial_{t_c}\Omega_{\omega;\mathfrak{s}}^\phi\big)&=\big(\partial_{t_c}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi\big)\big(\partial_{t_a}\partial_{t_b}\partial_{t_l}\Phi_\omega^\phi\big)g^{lm}\big(\partial_{t_m}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi\big)\\&\quad-\big(\partial_{t_a}\partial_{t_b}\partial_{t_l}\Phi_\omega^\phi\big)g^{lm}\sum_{1\le i,j\le N}\big(\partial_{t_c}\partial_{t_m}\partial_{t_i}\Phi_\omega^\phi\big)g^{ij}\big(\partial_{t_j}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi\big)\,.\end{aligned} \tag{17}$$

Summing up (17) over all $l,m=1,\ldots,N$ and adding (16) to the result, we find that

$$\begin{aligned}\big(\partial_u^2\Omega_{\omega;\mathfrak{s}}^\phi\big)\big(\text{LHS of (13)}\big)&=\big(\partial_{t_a}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi\big)\big(\partial_{t_b}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi\big)\big(\partial_{t_c}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi\big)\\&\quad-\sum_{1\le i,j\le N}\Bigg(\sum_{1\le l,m\le N}\big(\partial_{t_a}\partial_{t_b}\partial_{t_l}\Phi_\omega^\phi\big)g^{lm}\big(\partial_{t_m}\partial_{t_c}\partial_{t_i}\Phi_\omega^\phi\big)\Bigg)g^{ij}\big(\partial_{t_j}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi\big)\,.\end{aligned} \tag{18}$$

The right-hand side of (13) times $\partial_u^2\Omega_{\omega;\mathfrak{s}}^\phi$ is given by the right-hand side of (18) with the roles of b and c interchanged. Thus,

$$\begin{aligned}&\big(\partial_u^2\Omega_{\omega;\mathfrak{s}}^\phi\big)\Big(\big(\text{LHS of (13)}\big)-\big(\text{RHS of (13)}\big)\Big)\\&=\sum_{1\le i,j\le N}\Bigg(\sum_{1\le l,m\le N}\Big(\big(\partial_{t_a}\partial_{t_c}\partial_{t_l}\Phi_\omega^\phi\big)g^{lm}\big(\partial_{t_m}\partial_{t_b}\partial_{t_i}\Phi_\omega^\phi\big)-\big(\partial_{t_a}\partial_{t_b}\partial_{t_l}\Phi_\omega^\phi\big)g^{lm}\big(\partial_{t_m}\partial_{t_c}\partial_{t_i}\Phi_\omega^\phi\big)\Big)\Bigg)g^{ij}\big(\partial_{t_j}\partial_u\Omega_{\omega;\mathfrak{s}}^\phi\big).\end{aligned}$$

Combing this statement with (15), we conclude that

$$\big(\partial_u^2\Omega^\phi_{\omega;\mathfrak{s}}\big)\Big(\big(\text{LHS of (13)}\big)-\big(\text{RHS of (13)}\big)\Big)=0\in\Lambda_\omega[[t_1,\ldots,t_N,u]]. \tag{19}$$

This establishes the claim. □

5 Proof of Theorem 1

By symmetry, it is sufficient to establish this proposition for $i=2$ and $j=3$. Since $\overline{\mathcal{M}}_{0,3}$ is a single point, Theorem 1 holds for $l=3$.

Let $l\ge3$ and

$$\mathfrak{f}\colon\overline{\mathcal{M}}_{0,l+1}\longrightarrow\overline{\mathcal{M}}_{0,l}$$

be the forgetful morphism dropping the last marked point. For each partition $I\cup J$ of $[l]$ with $|I|,|J|\ge2$, the restriction of $\mathfrak{f}$ to the complement of $D_{I(l+1),J}\cap D_{I,J(l+1)}$ in $\overline{\mathcal{M}}_{0,l+1}$ is transverse to $D_{I,J}$. Since

$$\mathfrak{f}^{-1}\big(D_{I,J}\big)=D_{I(l+1),J}\cup D_{I,J(l+1)},$$
$$\big\{\mathrm{d}\mathfrak{f}|_{D_{I(l+1),J}-D_{I,J(l+1)}}\big\}^*\mathfrak{o}^c_{D_{I,J}}=\mathfrak{o}^c_{D_{I(l+1),J}},\quad \big\{\mathrm{d}\mathfrak{f}|_{D_{I,J(l+1)}-D_{I(l+1),J}}\big\}^*\mathfrak{o}^c_{D_{I,J}}=\mathfrak{o}^c_{D_{I,J(l+1)}},$$

and $D_{I(l+1),J}\cap D_{I,J(l+1)}$ is of real codimension 2 in $D_{I(l+1),J}$ and $D_{I,J(l+1)}$,

$$\mathfrak{f}^*\big(\mathrm{PD}_{\overline{\mathcal{M}}_{0,l}}\big(D_{I,J}\big)\big)=\mathrm{PD}_{\overline{\mathcal{M}}_{0,l+1}}\big(D_{I(l+1),J}\big)+\mathrm{PD}_{\overline{\mathcal{M}}_{0,l+1}}\big(D_{I,J(l+1)}\big)\in H^2\big(\overline{\mathcal{M}}_{0,l+1};\mathbb{Z}\big). \tag{20}$$

On the other hand, the section

$$s\in\Gamma\big(\overline{\mathcal{M}}_{0,l+1};L_1^*\otimes\mathfrak{f}^*L_1\big),\qquad s(w)=\mathrm{d}\mathfrak{f}(w),$$

vanishes only along the divisor $D_{1(l+1),[l]-\{1\}}$. Since $\mathrm{d}s$ induces an orientation-preserving isomorphism from $(\mathcal{N}D_{1(l+1),[l]-\{1\}},\mathfrak{o}^c_{D_{1(l+1),[l]-\{1\}}})$ to $L_1^*\otimes\mathfrak{f}^*L_1|_{D_{1(l+1),[l]-\{1\}}}$, we conclude that

$$\psi_1=\mathfrak{f}^*\psi_1+\mathrm{PD}_{\overline{\mathcal{M}}_{0,l+1}}\big(D_{1(l+1),[l]-\{1\}}\big)\in H^2\big(\overline{\mathcal{M}}_{0,l+1};\mathbb{Z}\big). \tag{21}$$

Combining (20) with (21), we obtain the identity of Theorem 1 by induction.

6 Proof of Theorem 2

By symmetry, it is sufficient to assume that $i=2$ in the second statement. Since $\mathbb{R}\overline{\mathcal{M}}_{0,1,1}$ is a single point and $\mathbb{R}\overline{\mathcal{M}}_{0,0,2}$ is a circle, the first statement of this proposition holds for $(k,l)=(1,1)$ and the second for $(k,l)=(0,2)$.

If $k+2l\ge3$, let

$$\mathfrak{f}\colon\mathbb{R}\overline{\mathcal{M}}_{0,k,l+1}\longrightarrow\mathbb{R}\overline{\mathcal{M}}_{0,k,l}\qquad\text{and}\qquad\mathfrak{f}_{\mathbb{R}}\colon\mathbb{R}\overline{\mathcal{M}}_{0,k+1,l}\longrightarrow\mathbb{R}\overline{\mathcal{M}}_{0,k,l}$$

be the forgetful morphisms dropping the last conjugate pair of marked points and the last real marked point, respectively. By the same reasoning as for (20),

$$\begin{aligned}\mathfrak{f}^*\big(\mathrm{PD}_{\mathbb{R}\overline{\mathcal{M}}_{0,k,l}}\big(\mathbb{R}D_{I,J},\mathfrak{o}^c_{\mathbb{R}D_{I,J}}\big)\big)=&\,\mathrm{PD}_{\mathbb{R}\overline{\mathcal{M}}_{0,k,l+1}}\big(\mathbb{R}D_{I(l+1),J},\mathfrak{o}^c_{\mathbb{R}D_{I(l+1),J}}\big)\\&+\mathrm{PD}_{\mathbb{R}\overline{\mathcal{M}}_{0,k,l+1}}\big(\mathbb{R}D_{I,J(l+1)},\mathfrak{o}^c_{\mathbb{R}D_{I,J(l+1)}}\big)\in H^2\big(\mathbb{R}\overline{\mathcal{M}}_{0,k,l+1};\mathbb{Z}\big)\end{aligned} \tag{22}$$

for each partition $I\cup J$ of $[l]$ with $|I|\ge 2$ and either $J\neq\emptyset$ or $k>0$. We also have

$$\mathfrak{f}_{\mathbb{R}}^*\big(\mathrm{PD}_{\mathbb{R}\overline{\mathcal{M}}_{0,k,l}}\big(\mathbb{R}D_{I,J},\mathfrak{o}^c_{\mathbb{R}D_{I,J}}\big)\big)=\mathrm{PD}_{\mathbb{R}\overline{\mathcal{M}}_{0,k+1,l}}\big(\mathbb{R}D_{I,J},\mathfrak{o}^c_{\mathbb{R}D_{I,J}}\big)\in H^2\big(\mathbb{R}\overline{\mathcal{M}}_{0,k+1,l};\mathbb{Z}\big) \tag{23}$$

for each partition $I\cup J$ of $[l]$ with $|I|\ge 2$ and either $J\neq\emptyset$ or $k>0$. By the same reasoning as for (21),

$$\begin{gathered}\psi_1=\mathfrak{f}^*\psi_1+\mathrm{PD}_{\mathbb{R}\overline{\mathcal{M}}_{0,k,l+1}}\big(\mathbb{R}D_{1(l+1),[l]-\{1\}},\mathfrak{o}^c_{\mathbb{R}D_{1(l+1),[l]-\{1\}}}\big)\in H^2\big(\mathbb{R}\overline{\mathcal{M}}_{0,k,l+1};\mathbb{Z}\big),\\ \psi_1=\mathfrak{f}_{\mathbb{R}}^*\psi_1\in H^2\big(\mathbb{R}\overline{\mathcal{M}}_{0,k+1,l};\mathbb{Z}\big).\end{gathered} \tag{24}$$

Combining (22) and (23) with (24), we obtain the two identities of Theorem 2 by induction.

Department of Mathematics, Stony Brook University, Stony Brook, NY 11794
azinger@math.stonybrook.edu

References

[1] A. Alcolado, *Extended Frobenius manifolds and the open WDVV equations*, PhD thesis, McGill, 2017

[2] E. Brugallé and P. Georgieva, *Pencils of quadrics and Gromov-Witten-Welschinger invariants of* $\mathbb{CP}^3$, Math. Ann. 365 (2016), no. 1–2, 363-380

[3] E. Brugallé and G. Mikhalkin, *Enumeration of curves via floor diagrams*, C. R. Math. Acad. Sci. Paris 345 (2007), no. 6, 329-334

[4] X. Chen, *Steenrod pseudocycles, lifted cobordisms, and Solomon's relations for Welschinger's invariants*, math/1809.08919v2

[5] X. Chen, *A geometric depiction of Solomon-Tukachinsky's construction of open GW-invariants*, math/1912.04119

[6] X. Chen and A. Zinger, *WDVV-type relations for Welschinger's invariants: applications*, math/1809.08938v2

[7] X. Chen and A. Zinger, *WDVV-type relations for disk Gromov-Witten invariants in dimension 6*, math/1904.04254

[8] M. Farajzadeh Tehrani, *Counting genus zero real curves in symplectic manifolds*, Geom. Topol. 20 (2016), no. 2, 629-695

[9] K. Fukaya, *Cyclic symmetry and adic convergence in Lagrangian Floer theory*, Kyoto J. Math. 50 (2010), no. 3, 521–590

[10] K. Fukaya, *Counting pseudo-holomorphic discs in Calabi-Yau 3-folds*, Tohoku Math. J. 63 (2011), no. 4, 697–727

[11] K. Fukaya, Y.-G. Oh, H. Ohta, and K. Ono, *Lagrangian Intersection Theory: Anomaly and Obstruction*, AMS Studies in Advanced Mathematics 46, 2009 (early versions, late 1990s)

[12] P. Georgieva, *Open Gromov-Witten invariants in the presence of an anti-symplectic involution*, Adv. Math. 301 (2016), 116-160

[13] P. Georgieva and A. Zinger, *Enumeration of real curves in $\mathbb{CP}^{2n-1}$ and a WDVV relation for real Gromov-Witten invariants*, Duke Math. J. 166 (2017), no. 17, 3291-3347

[14] P. Georgieva and A. Zinger, *Real Gromov-Witten theory in all genera and real enumerative geometry: properties*, J. Symplectic Geom. 17 (2019), no. 4, 1083-1158

[15] L. Göttsche and R. Pandharipande, *The quantum cohomology of blow-ups of $\mathbb{P}^2$ and enumerative geometry*, J. Differential Geom. 48 (1998), no. 1, 61-90

[16] J. Kollár, *Example of vanishing Gromov-Witten-Welschinger invariants*, J. Math. Sci. Univ. Tokyo 22 (2015), no. 1, 261-278

[17] M. Kontsevich and Y. Manin, *Gromov-Witten classes, quantum cohomology, and enumerative geometry*, Comm. Math. Phys. 164 (1994), no. 3, 525–562

[18] C.-C. Liu, *Moduli of J-holomorphic curves with Lagrangian boundary condition and open Gromov-Witten invariants for an S^1-pair*, math/0210257

[19] J. Milnor and J. Stasheff, *Characteristic Classes*, Annals of Mathematics Studies 76, Princeton University Press, 1974

[20] Y. Ruan and G. Tian, *A mathematical theory of quantum cohomology*, J. Differential Geom. 42 (1995), no. 2, 259–367

[21] J. Solomon, *Intersection theory on the moduli space of holomorphic curves with Lagrangian boundary conditions*, math/0606429

[22] J. Solomon, *A differential equation for the open Gromov-Witten potential*, note 2007

[23] J. Solomon and S. Tukachinsky, *Differential forms, Fukaya A_∞ algebras, and Gromov-Witten axioms*, math/1608.01304; *Point-like bounding chains in open Gromov-Witten theory*, math/1608.02495

[24] J. Solomon and S. Tukachinsky, *Relative quantum cohomology*, math/1906.04795v2

[25] J.-Y. Welschinger, *Invariants of real symplectic 4-manifolds and lower bounds in real enumerative geometry*, Invent. Math. 162 (2005), no. 1, 195-234

[26] J.-Y. Welschinger, *Spinor states of real rational curves in real algebraic convex 3-manifolds and enumerative invariants*, Duke Math. J. 127 (2005), no. 1, 89-121; *Enumerative invariants of strongly semipositive real symplectic six-manifolds*, math/0509121v2