

EMPIRICAL MEASUREMENT OF COSMIC LUMINOSITY-ANGULAR DISTANCE RELATION

A PREPRINT

✉ **Juan De Vicente**

Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas (CIEMAT),
Avda. Complutense 40, E-28040, Madrid, Spain
juan.vicente@ciemat.es

May 4, 2023

ABSTRACT

One of the keys to understanding the universe is the proper measurement of the relation between the luminosity distance d_L and the angular diameter distance d_A . In 1933 Etherington deduced from general relativity the reciprocity equation $d_L = d_A(1+z)^\gamma$, with $\gamma = 1$ for a local (non-expanding) universe. This relation has been adapted to an expanding universe with the value $\gamma = 2$ by the introduction of the concept of comoving distance (d_M). In this work, we developed the method *Cosmic Redshift Inference* to measure experimentally the value of γ defined above independently of any possible galaxy evolution. The method has been applied to 1.2 million galaxy of SDSS DR15 obtaining the value $\gamma \simeq 1$.

Keywords Cosmology: theory · Galaxies: distances and redshifts · cosmological parameters · cosmic background radiation · dark matter · dark energy

1 Introduction

During the 20th century were established the foundations of modern cosmology. The field equations of general relativity were formulated by Einstein [1915]. The definition of new metrics based on the cosmological principle with the properties of homogeneity and isotropy allowed the physicists the application of Einstein's field equations to the universe. While Einstein defined a static metric, Friedmann [1922] deduces mathematically a non-stationary model with a time-dependent factor $a(t)$. The solution was independently derived by Lemaître [1927] interpreting $a(t)$ as a scale factor of an expanding universe. The work was completed by Robertson [1933] and Walker [1937] in what is known as the *Friedmann-Lemaître-Robertson-Walker* (FLRW) metric.

Contemporaneously to these achievements, a correlation between redshifts and distances for extragalactic sources was found by Hubble [1929]. The origin of this correlation was subject of intense debate between proponents of non-expanding and expanding universes on the 1930s (Kragh [2017]). The fault of the Einstein's static universe to explain the redshift of galaxies leans the balance to the FLRW metric, whose time dependent factor $a(t)$ can directly explain the cosmological redshift (Tolman [1934]). The FLRW model describes a solutions to the Einstein's field equations for a homogeneous and isotropic universe. The evolution and fate of the Universe depends on the nature of different density components, i.e., radiation, matter, curvature and dark energy. But as shown below, the FLRW metric also support a non-expanding universe accounting for the observed cosmological redshift.

Different cosmological tests were proposed to probe whether the Universe is expanding or remains static. Tolman [1930] predicted that in an expanding universe, the surface brightness of a receding source with redshift z will be dimmed by $\sim (1+z)^{-4}$. Consequently to Tolman's prediction, the equation $d_L = d_A(1+z)^\gamma$, with $\gamma = 2$ was established between *luminosity distance* d_L and *angular diameter distance* d_A for a expanding universe. In spite of some attempts (Lubin and Sandage [2001], Sandage [2010]), these relations have not found a conclusive experimental support from cosmological surveys. On the other hand, the time dilation of Type Ia supernovae light curves suggested by Wilson

[1939] and confirmed by Goldhaber et al. [2001] are assumed in favor of cosmological expansion though the same phenomenon can be described in a non-expanding universe as shown below.

In this work, we have developed the method *Cosmic Redshift Inference* (CRI) to determine experimentally the value of γ . The method has been applied to 1.2 million galaxies from SDSS DR15 sample (Aguado et al. [2019]), finding $\gamma \simeq 1$ which support a non-expanding universe rather than an expanding one. As Tolman [1934] wrote "it is observations rather than hypothesis that must dictate the final nature of our cosmological theory". Therefore, we have to find a non-expanding solution within the general relativity –different to the unsuccessful Einstein's static universe– to conciliate the theory and the experimental results. The most conservative approach is to look into the accepted FLRW metric. Note that this metric was first mathematically developed by Friedmann [1922] and admits a different interpretation to the expanding one given by Lemaître [1927].

The rest of the paper is organized as follows: In Section 2 some basic distance definitions of the *standard model* are reviewed. Section 3 describes a new method to measure the relation between the luminosity distance and the angular diameter distance. Weakness of the current expanding FLRW model are shown in Section 4. Section 5 shows the non-expanding interpretation of the FLRW metric. The conclusions are presented in Section 6. In Appendix A, a possible explanation of the redshift within a non-expanding universe is discussed.

2 Standard model of cosmology

According to the standard model, the *Friedmann-Lemaître-Robertson-Walker* (FLRW) metric along with the Einstein's field equation of *general relativity* describe a homogeneous and isotropic expanding universe. The FLRW metric is given by

$$-c^2 d\tau^2 = -c^2 dt^2 + a(t)^2 \left[\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right] \quad (1)$$

being

$$d\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2 \quad (2)$$

where k describes the curvature while $a(t)$ is the scale factor accounting for the universe expansion. There are different distance ladders relating theory and observations. Let us to provide a brief summary of some distance definitions and their relations with normalized densities ($\Omega_M, \Omega_r, \Omega_\Lambda, \Omega_k$), corresponding to matter, radiation, cosmological constant and curvature (Hogg [1999]). The first Friedmann equation can be expressed from the Hubble parameter H at any time, and the Hubble constant H_0 today as

$$\frac{\dot{a}(t)^2}{a(t)^2} = H^2 = H_0^2 E(z)^2 \quad (3)$$

where

$$E(z) = \sqrt{\Omega_K(1+z)^2 + \Omega_\Lambda + \Omega_M(1+z)^3 + \Omega_r(1+z)^4} \quad (4)$$

By integrating Eq. 1 along with Eq. 3 one can obtain the line of sight *comoving distance* d_C as

$$d_C = d_H \int_0^z \frac{dz'}{E(z')} \quad (5)$$

where $d_H = c/H_0 = 3000h^{-1} Mpc$ is the *Hubble distance*. From the same equations one can get the *transverse comoving distance* d_M as

$$d_M = \begin{cases} d_H \frac{1}{\sqrt{\Omega_k}} \sinh[\sqrt{\Omega_k} d_C / d_H] & \text{for } \Omega_k > 0 \\ d_C & \text{for } \Omega_k = 0 \\ d_H \frac{1}{\sqrt{|\Omega_k|}} \sin[\sqrt{|\Omega_k|} d_C / d_H] & \text{for } \Omega_k < 0 \end{cases} \quad (6)$$

With respect to observable quantities, the *angular diameter distance* d_A is defined as the ratio between the object physical size S and its angular size θ

$$d_A = \frac{S}{\theta} \quad (7)$$

The *angular diameter distance* is related to the *transverse comoving distance* by

$$d_M = d_A(1 + z) \quad (8)$$

where z is the redshift. On the other hand, the *luminosity distance* defines the relation between the bolometric flux energy f received at earth from an object, to its bolometric luminosity L by means of

$$f = \frac{L}{4\pi d_L^2} \quad (9)$$

or finding d_L

$$d_L = \sqrt{\frac{L}{4\pi f}} \quad (10)$$

The relation between d_L and d_M is given by

$$d_L = d_M(1 + z) \quad (11)$$

and taking into account Eq. 8

$$d_L^2 = d_A^2(1 + z)^4 \quad (12)$$




Figure 1: Standard Model luminosity-angular distances relation: Angular diameter distance (d_A), comoving distance (d_C) and luminosity distance (d_L) for a flat universe. d_A is the distance at emission, d_C is the distance at reception and d_L account for the distance elongation due to universe expansion ($\sim (1 + z)$), time dilation and wavelength redshifting ($\sim (1 + z)$). The relation $d_L = d_A(1 + z)^2$ can be deduced from the figure.

There are four $(1+z)$ factors affecting to flux energy diminution (Fig. 1). Two come from the elongation of the initial distance d_A by a factor of $(1+z)$ due to universe expansion according to the inverse square law. Another factor comes from the time dilation due to universe expansion that reduces the photon emission/reception rate by $(1+z)^{-1}$. The last factor comes from the cosmological wavelength redshift that decrease the energy of photons by $(1+z)^{-1}$. Therefore, a relevant relation is established between the *angular diameter distance* and the *luminosity distance* in the *expanding universe* as

$$d_L = d_A(1+z)^2 \quad (\text{expanding universe}) \quad (13)$$

Eq. 13 is commonly known as Etherington distance-duality relation.

3 Cosmic Redshift Inference: Empirical determination of the luminosity-angular distances relation

In this section we derive a photometric redshift method based on the cosmological luminosity-angular distances relation. The luminosity distance can be expressed as

$$d_L = \sqrt{\frac{L}{4\pi f_L}} \quad (14)$$

and since the angular distance corresponds to the distance at emission for both non-expanding and expanding universes, it can be expressed also as

$$d_A = \sqrt{\frac{L}{4\pi f_A}} \quad (15)$$

where f_A corresponds to the hypothetical flux that would be measured at a distance d_A without neither expansion nor redshift. Dividing both expressions we have

$$\frac{d_L}{d_A} = \sqrt{\frac{f_A}{f_L}} \quad (16)$$

On the other hand, the luminosity-angular distances relation is given by

$$(1+z)^\gamma = \frac{d_L}{d_A} \quad (17)$$

with $\gamma = 2$ for an expanding universe and $\gamma = 1$ for a non-expanding one. Substituting Eq. 16 in Eq. 17 we have

$$(1+z)^\gamma = \left[\frac{f_A}{f_L} \right]^{1/2} \quad (18)$$

Taking base10 logarithm and multiplying by 2.5 in both sides of the equation we have

$$5\gamma \log(1+z) = -2.5 \log f_L - (-2.5 \log f_A) \quad (19)$$

and defining




Figure 2: Cosmic Redshift Inference: A linear regression is performed to measure the value of γ . The redshift vector should meet $\|\mathbf{v}_\gamma\| = 1$. It occurs for $\gamma \simeq 1$

$m_L = -2.5 \log f_L$ as the luminosity magnitude
 $m_A = -2.5 \log f_A$ as the angular magnitude
 $m_\gamma = 5\gamma \log(1+z)$ as the redshift magnitude

we have

$$m_L = m_A + m_\gamma \quad (20)$$

The equation can also be expressed for common multi-band surveys in a vectorial form as

$$\mathbf{m}_L = \mathbf{m}_A + \mathbf{m}_\gamma \quad (21)$$

Note that the luminosity magnitude has two independent components: \mathbf{m}_A that depends on the luminosity of the source and the distance at emission, and \mathbf{m}_γ that depends exclusively on redshift. Multiplying Eq. 21 by an unitary vector \mathbf{v}_γ in the direction of \mathbf{m}_γ , we obtain

$$\mathbf{m}_L \cdot \mathbf{v}_\gamma = \mathbf{m}_A \cdot \mathbf{v}_\gamma + \mathbf{m}_\gamma \cdot \mathbf{v}_\gamma \quad (22)$$

As $\mathbf{m}_\gamma \cdot \mathbf{v}_\gamma = m_\gamma$ and $\mathbf{m}_A \cdot \mathbf{v}_\gamma = 0$ since both vectors are orthogonal, the expression becomes

$$m_\gamma = \mathbf{m}_L \cdot \mathbf{v}_\gamma \quad (23)$$

To assess the validity of this expression, we need a galaxy sample with spectroscopic redshift (m_γ) and properly measured photometric magnitudes for an arbitrary number of bands (\mathbf{m}_L). A regression can be applied to this sample to determine \mathbf{v}_γ . The success of this approximation requires a high correlation between both sides of Eq. 23. The proper value of γ is the one whose corresponding computed \mathbf{v}_γ is unitary as assumed above ($\|\mathbf{v}_\gamma\| = 1$).

We have resorted to SDSS DR15 that provides simultaneously spectra and photometric measurements for about 1.2 million galaxies. To ensure a uniform treatment for all galaxies, we select *De Vaucouleurs* magnitudes ($deVMag_{ugriz}$) which achieves accurate measurement of the flux of the bulge, the most luminous part of the galaxies. Thus, $\mathbf{m} = (deVMag_u, deVMag_g, deVMag_r, deVMag_i, deVMag_z, 1)$, where the last component will account in the regression for the unmatched zero definition between redshift and magnitudes.

We have applied the regression (Eq. 23) to this sample obtaining a parameter vector \mathbf{v}_γ that provides high correlation ($c = 0.91$) between both sides of the equation for any value of γ . Nevertheless, only one value of γ meets the second




Figure 3: Cosmic Redshift Inference: Redshift reconstruction after applying $\|\mathbf{v}_\gamma\| = 1$: (a) bias of $\delta(z - z_\gamma)$ distribution. (b) σ_{68} of $\delta(z - z_\gamma)/(1+z)$ distribution




Figure 4: Scatter plot from the redshift reconstruction after applying $\|\mathbf{v}_\gamma\| = 1$: (a) Non-expanding universe (b) Expanding universe

condition: $\|\mathbf{v}_\gamma\| = 1$ (Fig. 2). We can see that this value corresponds to $\gamma \simeq 1$ which is not expected for an expanding universe but for a non-expanding one.

Another way to determine the proper value of γ is to inversely reconstruct the value of the redshift z_γ after normalizing the measured value of \mathbf{v}_γ . Finding z from the definition of m_γ and taking into account Eq. 23 we have

$$z_\gamma = 10^{\frac{m_\gamma \cdot \mathbf{v}_\gamma}{s_\gamma}} - 1 \quad (24)$$

In such a way that we can compare z_γ with the spectroscopic value z . Fig. 3 shows the results of redshift reconstruction along a range of γ values. The Fig. 3a corresponds to the bias for $\delta z = (z - z_\gamma)$ distribution which shows a minimum at $\gamma \simeq 1$ which corresponds to a non-expanding universe. In Fig. 3b the ordinate axis represents σ_{68} that is a measurement of the dispersion of the distribution $\delta z = (z - z_\gamma)/(1+z)$. It corresponds to the width of the distribution measured with respect to the median, in which 68% of the galaxies are enclosed.

$$\sigma_{68} = \frac{1}{2}(P_{84} - P_{16}) \quad (25)$$

where P_{16} and P_{84} are the 84th and the 16th percentile of the cumulative distribution respectively. The abscissa axis corresponds to γ . The minimum deviation is also at $\gamma \simeq 1$.

Fig. 4 shows the scatter plot of redshift reconstruction for $\gamma = 1$ (*non-expanding universe*) and $\gamma = 2$ (*expanding universe*). The comparison support a non-expanding universe.

4 Weakness of expansion interpretation of FLRW metric

At this point we can rise the question:

What are the equations that support the expanding universe? The expanding universe rest on the metric tensor $g_{\mu\nu}$ for a homogeneous and isotropic universe given by the FLRW metric and on the Einstein's field equation.

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4}T_{\mu\nu} \quad (26)$$

But that is not all, the expanding universe requires the additional support from the equation

$$d_L = d_M(1 + z) \quad (27)$$

which relates the luminosity distance d_L with the transverse comoving distance d_M , equation not derived from Einstein's Field equations in spite of both the energy-momentum tensor $T_{\mu\nu}$ and the luminosity distance d_L are related to energy fluxes. Note that it is not $a(t)$ but the definition of this comoving distance d_M and the consideration of r in FLRW metric as a comoving coordinate that force the FLRW model to be an expanding one.

A similar question arise with respect to Etherington relation. Fig. 5 shows the relation between the luminosity distance Δ and the angular diameter distance Δ' derived by Etherington. Thus, the Etherington equation deduced from general relativity was the reciprocity theorem for a non-expanding universe:

$$d_L = d_A(1 + z) \quad (\text{non - expanding universe}) \quad (28)$$

Note that this equation can be adapted for an expanding universe by the introduction of an additional intermediate variable d_M (comoving distance) –not considered by Etherington–, and taking into account Eq. 27 that combined with Eq. 29

$$d_M = d_A(1 + z) \quad (29)$$

gives

$$d_L = d_A(1 + z)^2 \quad (\text{expanding universe}) \quad (30)$$

equation known as Etherington duality relation.

Therefore, it is by the definition of these two external equations (Eq. 27 and Eq. 29) –non derived from general relativity– that the expanding paradigm is supported.

$$\Delta = \left(1 + \frac{\delta\lambda}{\lambda}\right) \Delta'. \quad . \quad . \quad . \quad . \quad . \quad (23)$$

Two conclusions follow from the last equation :—

(i.) The distance computed from estimates of brightness will agree exactly with the distance computed from estimates of size only when the star has no apparent radial velocity.

(ii.) For spherical objects of the same absolute size and brightness, moving variously at varying distances from the observer, the ratio of the apparent diameter to the square root of the apparent brightness is proportional to $(1 + \delta\lambda/\lambda)$,

Figure 5: Etherington equation (snapshot from his paper).

5 Non-expanding interpretation of FLRW metric

Let us to consider the FLRW metric given by

$$-c^2 d\tau^2 = -c^2 dt^2 + a(t)^2 \left[\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right] \quad (31)$$

Let us to analyze the behavior of light rays in FLRW metric considering a non-expanding universe (Fig. 6). Let the origin of coordinates be at the observer O, and considers an extended cosmological object (galaxy) initially located at a distance d_A from O at time of emission t_e . According to general relativity, light rays follow null geodesics where $d\tau^2 = 0$. Substituting this value in Eq. 31, light rays follow the equation

$$c^2 dt^2 = a(t)^2 \left[\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right] \quad (32)$$

Note from Fig. 6 that in a non-expanding universe the light rays that will arrive to the observer O in the future are those pointing initially towards the observer at time of emission t_e (leaving apart non pure cosmological effects as gravitational lensing or astrophysical events). These rays will maintain the same direction up to arriving to the observer, i.e., $\theta = cte$, $d\theta = 0$, $\phi = cte$, $d\phi = 0$ in Eq. 2, and hence $d\Omega = 0$. Substituting $d\Omega = 0$ in Eq. 32, the light rays that will arrive to the observer meet

$$c^2 dt^2 = a(t)^2 \frac{dr^2}{1 - kr^2} \quad (33)$$

Integrating Eq. 33 from $t_e = 0$ we have

$$\int_0^t \frac{cdt}{a(t)} = \int_0^{d_M} \frac{dr}{\sqrt{1 - kr^2}} \quad (34)$$

Note that in a non-expanding universe, the comoving coordinate r of the FLRW model loses its meaning and r can be interpreted as luminosity (distance) coordinate. In this way, the comoving distance d_M can be substituted by the luminosity distance d_L in all equation. Therefore

$$d_L = d_M \quad (\text{non - expanding universe}) \quad (35)$$

and

$$\int_0^t \frac{cdt}{a(t)} = \int_0^{d_L} \frac{dr}{\sqrt{1 - kr^2}} \quad (36)$$

Thus, in the case of a flat ($k = 0$) non-expanding universe, we would have




Figure 6: Geodesics for light rays in a flat non-expanding FLRW universe.

$$d_L = \int_0^t \frac{cdt}{a(t)} \quad (37)$$

which would be the factor responsible of time dilation and cosmological redshift in a non-expanding universe. For the case where c is constant, we can define

$$\Delta t' = \int_0^t \frac{dt}{a(t)} \quad (38)$$

as time dilation in such a way that $d_A = c\Delta t$ and $d_L = c\Delta t'$.

On the other hand, Eq. 6 transforms to Eq. 39

$$d_L = \begin{cases} d_H \frac{1}{\sqrt{\Omega_k}} \sinh[\sqrt{\Omega_k} d_C / d_H] & \text{for } \Omega_k > 0 \\ d_c & \text{for } \Omega_k = 0 \\ d_H \frac{1}{\sqrt{|\Omega_k|}} \sin[\sqrt{|\Omega_k|} d_C / d_H] & \text{for } \Omega_k < 0 \end{cases} \quad (39)$$

that directly relates the first Friedmann equation with the observable d_L derived from FLRW metric (Eq. 37) without the need to define additional equations detached from general relativity. Let us to express the FLRW metric in another form. Dividing Eq. 31 by $a(t)^2$ we have Eq. 40. Note that in this form it is more clear the time dilation nature of $a(t)$ rather than the role of a scale factor for an expanding universe.

$$-\left(\frac{cd\tau}{a(t)}\right)^2 = -\left(\frac{cdt}{a(t)}\right)^2 + \left[\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2\right] \quad (40)$$

In a expanding universe, the first Friedmann equation constraints the form of the scale factor $a(t)$ with the different species of the universe as radiation, matter, curvature and cosmological constant. According to the present experimental results and the non-expanding interpretation of the FLRW model, $a(t)$ corresponds to another unidentified property of vacuum –different from expansion– which also depends on the relative content of these species along cosmic time.

The stability of the non-expanding universe resides on the acceptance of the cosmological principle since $a(t)$ is no more related to the size of the universe. The standard model demonstrates that the geodesics for a free particle (galaxy) corresponds to fixed FLRW comoving coordinates. The same demonstration applies to non-expanding-FLRW by interpreting comoving coordinates as luminosity (distance) ones. Thus, geodesics for a free particle in a non-expanding universe corresponds to fixed FLRW luminosity coordinates, which are spatially fixed but affected by a time dilation in the reception of signals. Therefore, it is the own FLRW metric that ensure the stability of the non-expanding universe.

6 Conclusions

In the 1930s, early after the discovery of the redshift-distance relation, a debate emerged among physicist regarding the feasibility of a non-expanding or an expanding universe. Tolman proposed a surface brightness test as a mean to differentiate an expanding from a non-expanding universe. The test predicts the relation $\mu \sim (1+z)^{-4}$ for an expanding universe. It corresponds to $d_L = d_A(1+z)^\gamma$, with $\gamma = 2$ for an expanding universe and $\gamma = 1$ for a non-expanding one. Up to now, non-conclusive test have been performed on galaxy survey in this respect due to the unknown possible galaxies' evolution.

In this work we have developed Cosmic Redshift Inference, a photometric redshift method that it is able to isolate the dimming due to the redshift from the measured magnitude, allowing us to measure the value of γ . The method has been applied to 1.2 million galaxy of the public SDSS DR15 catalog. The result obtained $\gamma \simeq 1$ supports a non-expanding universe.

The expanding universe is supported by the Einstein's Field Equations and the FLRW metric. In this work, we note that the expanding interpretation of the FLRW metric also requires some external equation based on the definition of the comoving concept. Stripping away this supplement from the FLRW model, uncover the non-expanding essence

of the metric while still preserves $a(t)$, the time dependent function responsible of the redshift. Thus, the non-expanding universe determined experimentally can still be accommodated within the FLRW model. It only requires the interpretation of the ad hoc comoving coordinates as the observed luminous ones.

The research is not over and the physical cause of the cosmological redshift have to be determined. In Appendix A, we explore the magnetic permeability variation with cosmic time as a possible cause of redshift. The growth of the magnetic permeability would drive to a decrease of speed of light while preserving the observed gross atomic structure of distant galaxies.

Acknowledgements

Funding support for this work was provided by the Autonomous Community of Madrid through the project TEC2SPACE-CM (S2018/NMT-4291).

This paper uses data from public SDSS DR-15. Funding for the Sloan Digital Sky Survey IV has been provided by the Alfred P. Sloan Foundation, the U.S. Department of Energy Office of Science, and the Participating Institutions. SDSS-IV acknowledges support and resources from the Center for High-Performance Computing at the University of Utah. The SDSS web site is www.sdss.org.

SDSS-IV is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS Collaboration including the Brazilian Participation Group, the Carnegie Institution for Science, Carnegie Mellon University, the Chilean Participation Group, the French Participation Group, Harvard-Smithsonian Center for Astrophysics, Instituto de Astrofísica de Canarias, The Johns Hopkins University, Kavli Institute for the Physics and Mathematics of the Universe (IPMU) / University of Tokyo, the Korean Participation Group, Lawrence Berkeley National Laboratory, Leibniz Institut für Astrophysik Potsdam (AIP), Max-Planck-Institut für Astronomie (MPIA Heidelberg), Max-Planck-Institut für Astrophysik (MPA Garching), Max-Planck-Institut für Extraterrestrische Physik (MPE), National Astronomical Observatories of China, New Mexico State University, New York University, University of Notre Dame, Observatório Nacional / MCTI, The Ohio State University, Pennsylvania State University, Shanghai Astronomical Observatory, United Kingdom Participation Group, Universidad Nacional Autónoma de México, University of Arizona, University of Colorado Boulder, University of Oxford, University of Portsmouth, University of Utah, University of Virginia, University of Washington, University of Wisconsin, Vanderbilt University, and Yale University.

References

- Albert Einstein. Die feldgleichungen der gravitation. *Sitzung der physikalische-mathematischen Klasse*, 25:844–847, 1915.
- Alexander Friedmann. Über die krümmung des raumes. *Zeitschrift für Physik*, 10:377–386, 1922.
- G Lemaître. Lemaître, g., 1927, ann. soc. sci. bruxelles, ser. 1 47, 49. *Ann. Soc. Sci. Bruxelles, Ser. 1*, 47:49, 1927.
- Howard Percy Robertson. Relativistic cosmology. *Reviews of modern Physics*, 5(1):62, 1933.
- Arthur Geoffrey Walker. On milne’s theory of world-structure. *Proceedings of the London Mathematical Society*, 2(1): 90–127, 1937.
- Edwin Hubble. A relation between distance and radial velocity among extra-galactic nebulae. *Proceedings of the National Academy of Sciences*, 15(3):168–173, 1929.
- Helge Kragh. Is the universe expanding? fritz zwicky and early tired-light hypotheses. *Journal of Astronomical History and Heritage*, 20(1):2–12, 2017.
- RC Tolman. Relativity, thermodynamics, and cosmology. clarendon, 1934.
- Richard C Tolman. On the estimation of distances in a curved universe with a non-static line element. *Proceedings of the National Academy of Sciences*, 16(7):511–520, 1930.
- Lori M Lubin and Allan Sandage. The tolman surface brightness test for the reality of the expansion. iv. a measurement of the tolman signal and the luminosity evolution of early-type galaxies. *The Astronomical Journal*, 122(3):1084, 2001.
- Allan Sandage. The tolman surface brightness test for the reality of the expansion. v. provenance of the test and a new representation of the data for three remote hubble space telescope galaxy clusters. *The Astronomical Journal*, 139(2): 728, 2010.
- O.C Wilson. supernovae clock. *The Astrophysical Journal*, 90:634, 1939.

- Gerson Goldhaber, DE Groom, A Kim, G Aldering, P Astier, A Conley, SE Deustua, R Ellis, S Fabbro, AS Fruchter, et al. Timescale stretch parameterization of type ia supernova b-band light curves. *The Astrophysical Journal*, 558 (1):359, 2001.
- David Sánchez Aguado, Romina Ahumada, Andres Almeida, Scott F Anderson, Brett H Andrews, Borja Anguiano, Erik Aquino Ortíz, Alfonso Aragón-Salamanca, Maria Argudo-Fernández, Marie Aubert, et al. The fifteenth data release of the sloan digital sky surveys: first release of manga-derived quantities, data visualization tools, and stellar library. *The Astrophysical Journal Supplement Series*, 240(2):23, 2019.
- David W Hogg. Distance measures in cosmology. *arXiv preprint astro-ph/9905116*, 1999.
- Fritz Zwicky. On the possibilities of a gravitational drag of light. *Physical Review*, 34(12):1623, 1929.
- John W Moffat. Superluminary universe: A possible solution to the initial value problem in cosmology. *International Journal of Modern Physics D*, 2(03):351–365, 1993.
- Andreas Albrecht and Joao Magueijo. Time varying speed of light as a solution to cosmological puzzles. *Physical Review D*, 59(4):043516, 1999.
- Robert H Dicke. Gravitation without a principle of equivalence. *Reviews of Modern Physics*, 29(3):363, 1957.
- Jef Van Royen. General relativity: Varying speed of light from the friedmann equation. *Journal of Cosmology*, 27: 15389–15406, 2021.
- Giuseppe Pipino. Variable speed of light with time and general relativity. *Journal of High Energy Physics, Gravitation and Cosmology*, 7(2):742–760, 2021.
- Peter I Wold. On the redward shift of spectral lines of nebulae. *Physical Review*, 47(3):217, 1935.

A Exploring variable magnetic permeability with cosmic time as cause of cosmological redshift

The physical causes of the redshift have to be identified among the different proposals. The most popular alternative approach to expansion as the cause of the redshift is the tired light (Zwicky [1929]), where photons lose energy while traveling to earth. Another possibility to be explored is the variable speed of light (VSL) with cosmic time. Let us to extend about this possibility in this section. The cosmological principle assumes a universe spatially homogeneous and spatially isotropic. It does not state that the universe is the same over time. Thus, according to the cosmological principle we can allow a space property to change overtime. That is the case of the scale factor in the expanding universe or the speed of light in the non-expanding one. There are different VSL theories as those addressing the horizon problem (Moffat [1993], Albrecht and Magueijo [1999]) or the ones allowing the variation of speed of light between free-falling observers (Dicke [1957]). Other depart from FLRW metric as the one that assumes both expansion and VLS (Van Royen [2021])(which would requires a value of $\gamma \geq 3$, far from our measurement) or those assuming photons emitted at higher speed of light at earlier times, but maintaining such high velocities up to earth (Pipino [2021]), events not observed experimentally.

Though less known, there is a plausible alternative explanation to redshift based on variable speed of light (VSL) with cosmic time (Wold [1935]). Such approach would still require a spatially constant speed of light among all free falling observes as the general relativity demands. In this case, from Eq. 37 we can write

$$d_L = \int_0^t c(t)dt = c_0 \int_0^t \frac{dt}{a(t)} \quad (41)$$

being c_0 the current value of the speed of light.

The process of photon redshift based on a speed of light decreasing with cosmic time can be described as follow: A galaxy emits a photon at speed c_z due to an electron transition between atomic levels at its corresponding energy $h\nu_0$, being lambda stretched out at emission due to the equation $\lambda_z = c_z/\nu_0$. In the travel of the photon to earth, λ_z remains constant, while the frequency decrease up to ν_z due to speed of light drop $\nu_z = c_0/\lambda_z$.

Note that Eq. 41 implies $d_L = d_A$, and the assumed Eq. 17 changes to

$$(1+z)^\gamma = \frac{d_L}{d_0} \quad (42)$$

being d_0 the angular distance d_A that we would have if the speed of light were constant along cosmic time.

Given that the speed of light is

$$c = \frac{1}{\sqrt{\epsilon_0 \mu_0}} \quad (43)$$

some of the vacuum properties either dielectric permittivity ϵ_0 or magnetic permeability μ_0 have to change with cosmic time. Since the gross atomic structure of redshifted galaxies and its corresponding energy levels depend on ϵ_0 , we assume that it is $\mu_0 = \mu_0(t)$ that depends on cosmic time in the form

$$a(t) = \sqrt{\mu_0(t)} \quad (44)$$

in such a way that only magnetic fields and the fine structure constant would be affected along cosmic time. Thus, the time dependent function $a(t)$ defined in *FLRW* metric would not correspond to an expansion but to a known property of vacuum, the square root of the magnetic permeability. Consequently, the speed of light would decrease with cosmic time $c(t) = 1/\sqrt{\epsilon_0 \mu_0(t)}$ while the electric permittivity ϵ_0 remains constant allowing the observed atomic structures. The model can be denominated $\mu_0 - FLRW$ to differentiate it from other possible alternatives.

Note that $\mu_0 - FLRW$ universe does not change the *FLRW* equations but reinterpreted them. Thus, $\mu_0 - FLRW$ may assume the main ideas of the standard model as that the early universe was hotter and dominated by radiation, but the drop in universe temperature would not be due to expansion but to the drop in the speed of light with cosmic time. Thus, the cosmic microwave background would have been emitted at the same energy as in the standard model with values $\nu_z = \nu_{cmb}(1 + z_{cmb})$, $c_z = c_0(1 + z_{cmb})$ and $\lambda_{cmb} = c_z/\nu_z$, and is received as c_0 , ν_{cmb} and λ_{cmb} .