arXiv:2003.06164v4 [hep-th] 8 May 2020

Classification of six-dimensional Leibniz algebras $\mathcal{E}_3$

Ladislav Hlavatý*
Faculty of Nuclear Sciences and Physical Engineering,
Czech Technical University in Prague,
Czech Republic

February 6, 2022

Abstract

Leibniz algebras $\mathcal{E}_n$ were introduced as algebraic structure underlying U-duality. Algebras $\mathcal{E}_3$ derived from Bianchi three-dimensional Lie algebras are classified here. Two types of algebras are obtained: Six-dimensional Lie algebras that can be considered extension of semi-Abelian four-dimensional Drinfel'd double and unique extensions of non-Abelian Bianchi algebras. For all of the algebras explicit forms of generalized frame fields are given.

1 Introduction

Extensions of Poisson–Lie T-dualities to non-perturbative symmetries of string theories are so called U-dualities (for review see e.g.[1]). Algebraic structures underlying U-duality were suggested in [2] and [3] as Leibniz algebras $\mathcal{E}_n$ obtained as extensions of n-dimensional Lie algebra defining non-symmetric product $\circ$ in $[n+n(n-1)/2]$-dimensional vector space[1] that satisfies Leibniz identity

$$X \circ (Y \circ Z)) = (X \circ Y) \circ Z + Y \circ (X \circ Z). \tag{1}$$

*hlavaty@fjfi.cvut.cz
[1]for $n \leq 4$

In those papers examples of these Leibniz algebras derived from two-dimensional and four-dimensional Lie algebras are given. Goal of the present note is to write down all algebras that can be derived from three dimensional Lie algebras whose classification given by Bianchi is well known.

Namely, let $(T_a, T^{a_1a_2})$, $a, a_1, a_2 \in 1, \ldots, n$, $T^{a_2a_1} = -T^{a_1a_2}$ is a basis of $[n + n(n-1)/2]$-dimensional vector space. The algebra product given in [2] is

$$\begin{aligned} T_a \circ T_b &= f_{ab}{}^c \, T_c \,, \\ T_a \circ T^{b_1b_2} &= f_a{}^{b_1b_2c} \, T_c + 2 \, f_{ac}{}^{[b_1} \, T^{b_2]c} \,, \\ T^{a_1a_2} \circ T_b &= -f_b{}^{a_1a_2c} \, T_c + 3 \, f_{[c_1c_2}{}^{[a_1} \, \delta^{a_2]}_{b]} \, T^{c_1c_2} \,, \\ T^{a_1a_2} \circ T^{b_1b_2} &= -2 \, f_d{}^{a_1a_2[b_1} \, T^{b_2]d} \,, \end{aligned} \tag{2}$$

where $f_{ab}{}^c$ are structure coefficients of n-dimensional Lie algebra and $f_a{}^{b_1b_2b_3} = f_a{}^{[b_1b_2b_3]}$. Moreover, bilinear forms on $\mathcal{E}_n$ are defined

$$\langle T_a, \, T^{b_1b_2} \rangle_c = 2! \, \delta^{[b_1}_a \, \delta^{b_2]}_c \,, \qquad \langle T^{a_1a_2}, \, T^{b_1b_2} \rangle_{c_1 \cdots c_4} = 4! \, \delta^{a_1}_{[c_1} \, \delta^{a_2}_{c_2} \, \delta^{b_1}_{c_3} \, \delta^{b_2}_{c_4]} \,. \tag{3}$$

2 Bianchi-Leibniz algebras

We are going to classify Leibniz algebras $\mathcal{E}_3$ derived from three-dimensional Lie algebras. In this case $f_a{}^{b_1b_2b_3} = f_a \, \varepsilon^{b_1b_2b_3}$ where ε is totally antisymmetric Levi-Civita symbol. Non-vanishing bilinear forms are

$$\langle T_1, \, T^{12} \rangle_2 = \langle T_1, \, T^{13} \rangle_3 = \langle T_2, \, T^{23} \rangle_3 = 1,$$

$$\langle T_2, \, T^{12} \rangle_1 = \langle T_3, \, T^{13} \rangle_1 = \langle T_3, \, T^{23} \rangle_2 = -1.$$

First of all we shall show that for dimension three the Leibniz identities are satisfied only for unimodular Lie Algebras[2], i.e. $f_{ab}{}^b = 0$. Indeed, Leibniz identity

$$T^{23} \circ (T_1 \circ T_1) = (T^{23} \circ T_1) \circ T_1 + T_1 \circ (T^{23} \circ T_1)$$

and definitions (2) give

$$0 = 2 \, (f_{12}{}^2 + f_{13}{}^3)^2 \, T^{23} = 2 \, (f_{1b}{}^b)^2 \, T^{23}$$

[2]This is not true in general as can be shown explicitely for dimension four or by two-dimensional example in [2].

Class	a	n_1	n_2	n_3
B**1**	0	0	0	0
B**2**	0	1	0	0
B**3**	1	0	1	-1
B**4**	1	0	0	1
B**5**	1	0	0	0
B$\mathbf{6}_0$	0	1	-1	0
B$\mathbf{6}_a\,(a>0, a\neq 1)$	a	0	1	-1
B$\mathbf{7}_0$	0	1	1	0
B$\mathbf{7}_a\,(a>0)$	a	0	1	1
B**8**	0	1	1	-1
B**9**	0	1	1	1

Table 1: Bianchi algebras

and similarly for cyclic permutation of $(1, 2, 3)$.

Next point in our computations is the well known classification of 3–dimensional real Lie algebras. Non–isomorphic Lie algebras can be divided into eleven classes, traditionally known as Bianchi algebras. Their Lie algebra products are (see e.g. [4])

$$[X_1, X_2] = -aX_2 + n_3X_3,\ [X_2, X_3] = n_1X_1,\ [X_3, X_1] = n_2X_2 + aX_3, \quad (4)$$

where the parameters a, n_1, n_2, n_3 have the values given in the Table 1. Unimodular Bianchi algebras are those with $a = 0$, i.e. B**1** (Abelian), B**2** (Heisenberg), B$\mathbf{6}_0$ (Euclidean), B$\mathbf{7}_0$ (Poincare), B**8** (so(2,1)), and B**9** (so(3)).

Inserting (4) and (2) into Leibniz identities (1) we get

$$n_j f_k = 0, \quad j, k = 1, 2, 3. \quad (5)$$

This can be shown inspecting e.g. identities (1) for

$$X = T_1,\ Y = T^{23},\ Z = T^{12},$$

and

$$X = T_1,\ Y = T^{23},\ Z = T^{13}.$$

We get

$$n_2 f_1\, T^{13} + n_2 f_2\, T^{23} = 0,$$

$$n_3 f_1\, T^{12} + n_3 f_3\, T^{23} = 0,$$

so that

$$n_2 f_1 = 0, \quad n_2 f_2 = 0, \quad n_3 f_1 = 0, \quad n_3 f_3 = 0. \tag{6}$$

By cyclic permutation of $(1,2,3)$ we get (5) and it is easy to check that these conditions are sufficient for satisfaction of all Leibniz identities (1). Solution of conditions (5) is either $n_j = 0,\ j = 1,2,3$ or $f_k = 0,\ k = 1,2,3$. It means that we get two types of Bianchi-Leibniz algebras.

The first type are algebras depending only on f_k with products

$$\begin{aligned} T_a \circ T_b &= 0\,, \\ T_a \circ T^{b_1 b_2} &= f_a\, \varepsilon^{b_1 b_2 c}\, T_c\,, \\ T^{a_1 a_2} \circ T_b &= -f_b\, \varepsilon^{a_1 a_2 c}\, T_c\,, \\ T^{a_1 a_2} \circ T^{b_1 b_2} &= -2\, f_d\, \varepsilon^{a_1 a_2 [b_1}\, T^{b_2] d}\,. \end{aligned} \tag{7}$$

It is rather easy to check that this product is antisymmetric so that they are six-dimensional Lie algebras. The simplest one is Abelian where all $n_j = 0$ and $f_k = 0$. If at least one of f_k is not zero then by linear transformation from $E_3 = SL(3) \times SL(2)$ we can achieve $f_1 = 1,\ f_2 = f_3 = 0$ so that

$$\begin{aligned} [T_a, T_b] &= 0\,, \\ [T_1 \circ T^{b_1 b_2}] &= 2\, \varepsilon^{b_1 b_2 c}\, T_c\,, \\ [T^{23}, T^{12}] = 2\, T^{12}, &\quad [T^{23}, T^{13}] = 2\, T^{13}\,. \end{aligned} \tag{8}$$

The Bianchi-Leibniz algebras of the second type depend only on n_j whose values are given in the Table 1. It means that they are in one to one correspondence with the unimodular Bianchi algebras. Their products are

$$\begin{aligned} T_a \circ T_b &= [T_a, T_b]\,, \\ T_a \circ T^{b_1 b_2} &= \delta_a^{b_1} \varepsilon_{a b_2 c}\, n_{b_2}\, T^{b_1 c} - \delta_a^{b_2} \varepsilon_{a b_1 c}\, n_{b_1}\, T^{b, c}\,, \\ T^{a_1 a_2} \circ T_b &= 0\,, \\ T^{a_1 a_2} \circ T^{b_1 b_2} &= 0\,. \end{aligned} \tag{9}$$

Explicit forms of products $T_a \circ T^{b_1 b_2}$ are

$$T_1 \circ T^{12} = -T_3 \circ T^{23} = n_2\, T^{13},$$

$$T_1 \circ T^{13} = T_2 \circ T^{23} = -n_3\, T^{12},$$

$$T_2 \circ T^{12} = T_3 \circ T^{13} = n_1\, T^{23}.$$

Maximal isotropic algebras in both types of algebras are generated by $\{T_1, T_2, T_3\}$, $\{T^{12}, T^{13}, T^{23}\}$ and $\{T_1, T^{23}\}$, $\{T_2, T^{13}\}$, $\{T_3, T^{12}\}$.

As mentioned in [2], under some conditions we can choose a subalgebra of dimension $2(n-1)$ of the Leibnitz algebra $\mathcal{E}_n$ that is Lie algebra of Drinfel'd double. Leibniz algebra then can be considered as an extension of Drinfel'd double of dimension $2(n-1)$. Namely, if we can decompose the generators $\{T_a\}$ as $\{T_{\dot a}, T_z\}$ and $\{T^{ab}\}$ as $\{T^{\dot a\dot b},\, T^{\dot a z}\}$ $(\dot a = 1, \ldots, n-1)$ so that

$$f_{ab}{}^{z} = 0\,, \quad f_{az}{}^{b} = 0\,, \quad f_{z}{}^{b_1 b_2 b_3} = 0\,, \quad f_{\dot a}{}^{\dot b_1 \dot b_2 \dot b_3} = 0\,, \tag{10}$$

then the subalgebra spanned by

$$(T_{\dot A}) \equiv (T_{\dot a},\, T^{\dot a}) \qquad (T^{\dot a} \equiv T^{\dot a z}) \tag{11}$$

becomes Lie algebra of Drinfel'd double with the bilinear form

$$\langle T_{\dot a},\, T^{\dot b}\rangle := \langle T_{\dot a},\, T^{\dot b}\rangle_z = \delta_{\dot a}^{\dot b}\,. \tag{12}$$

However for $n = 3$, the conditions (10) and unimodularity are satisfied only for Abelian Bianchi algebra B**1**. It means that Leibniz algebras (7) can be considered extensions of Lie algebras of four-dimensional Drinfel'd doubles [5] generated by T_1, T_2, T^{13}, T^{23} and

$$[T_1, T_2] = 0, \quad [T^{13}, T^{23}] = -f_1\, T^{13} - f_2\, T^{23},$$

$$\langle T_1,\, T^{13}\rangle = \langle T_2,\, T^{23}\rangle = 1.$$

2.1 Generalized frame fields

Now we will present explicit forms of the so called generalized frame fields $E_A{}^I$ required to satisfy

$$\hat{\pounds}_{E_A} E_B{}^I = -\mathcal{F}_{AB}{}^C\, E_C{}^I, \tag{13}$$

where $\mathcal{F}_{AB}{}^C$ are structure constants of the Bianchi-Leibnitz algebras,

$$A, B, C, I \in (1, 2, 3, \{1,2\}, \{1,3\}, \{2,3\}).$$

The generalized Lie derivative can be expressed by

$$\left(\hat{\pounds}_V W^I\right) = \begin{pmatrix} \pounds_v w^i \\ \frac{(\pounds_v w_2 - \iota_w \mathrm{d}v_2)_{i_1 i_2}}{\sqrt{2!}} \end{pmatrix}, \tag{14}$$

generalized vectors V and W are parameterized as

$$V^I = \begin{pmatrix} v^i \\ \frac{v_{i_1 i_2}}{\sqrt{2!}} \end{pmatrix}, \tag{15}$$

$v_2 = \frac{1}{2} v_{ij} \mathrm{d}x^i \wedge \mathrm{d}x^j$ and similarly for W and $\widetilde{w}$.

Generalized frame fields ${E_A}^I$ have block triangular form

$${E_A}^I = \begin{pmatrix} {e_a}^i & 0 \\ E^{a_1 a_2 i} & E^{a_1 a_2}{}_{i1,i_2} \end{pmatrix} = \begin{pmatrix} {e_a}^i & 0 \\ -\frac{\Pi^{a_1 a_2 b} e^i_b}{\sqrt{2!}} & {r_{[i_1}}^{[a_1} {r_{i_2]}}^{a_2]} \end{pmatrix}, \tag{16}$$

where ${e_a}^i$ are components of right-invariant vector fields with respect to Bianchi groups, ${r_i}^a$ are components of right-invariant 1-forms and $\Pi^{a_1 a_2 b}$ is the so called Nambu-Poisson tensor that in the dimension three is $\pi(x_1, x_2, x_3)\, \varepsilon^{a_1 a_2 b}$ where ε is the Levi-Civita symbol.

From the formula (16) follows explicit form of matrices ${E_A}^I$ for the first type algebras (7), namely

$${E_A}^I = \begin{pmatrix} \mathbf{1}_3 & 0 \\ \Pi_3 & \frac{1}{2}\mathbf{1}_3 \end{pmatrix} \tag{17}$$

where $\mathbf{1}_3$ is three-dimensional unit matrix,

$$\Pi_3 = \begin{pmatrix} 0 & 0 & -\pi(x_1, x_2, x_3) \\ 0 & \pi(x_1, x_2, x_3) & 0 \\ -\pi(x_1, x_2, x_3) & 0 & 0 \end{pmatrix}, \tag{18}$$

and $\pi(x_1, x_2, x_3) = f_1 x_1 + f_2 x_2 + f_3 x_3 + const$ where f_a are constants appearing in (7).

Explicit form of matrices ${E_A}^I$ for the second type algebras (9) is

$${E_A}^I = \begin{pmatrix} M_1 & 0 \\ 0 & \frac{1}{2} M_2 \end{pmatrix}$$

where

$$M_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -n_1 x_2 & x_1 & 1 \end{pmatrix},$$

$$M_2 = \begin{pmatrix} 1 & -x_1 & -n_1 x_2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

for Bianchi algebras B**2**, B$\mathbf{6}_0$, B$\mathbf{7}_0$.

$$M_1 = \begin{pmatrix} 1 & 0 & 0 \\ -\sinh x_1 \tanh x_2 & \cosh x_1 & \sinh x_1 \operatorname{sech}\ x_2 \\ -\cosh x_1 \tanh x_2 & \sinh x_1 & \cosh x_1 \operatorname{sech}\ x_2 \end{pmatrix},$$

$$M_2 = \begin{pmatrix} \cosh x_1 & -\sinh x_1 \cosh x_2 & -\sinh x_2 \cosh x_1 \\ -\sinh x_1 & \cosh x_1 \cosh x_2 & \sinh x_1 \sinh x_2 \\ 0 & 0 & \cosh x_2 \end{pmatrix}$$

for Bianchi B**8**.

$$M_1 = \begin{pmatrix} 1 & 0 & 0 \\ \sin x_1 \tan x_2 & \cos x_1 & -\sin x_1 \sec x_2 \\ -\cos x_1 \tan x_2 & \sin x_1 & \cos x_1 \sec x_2 \end{pmatrix},$$

$$M_2 = \begin{pmatrix} \cos x_1 & -\sin x_1 \cos x_2 & -\sin x_2 \cos x_1 \\ \sin x_1 & \cos x_1 \cos x_2 & -\sin x_1 \sin x_2 \\ 0 & 0 & \cos x_2 \end{pmatrix}$$

for Bianchi B**9**.

3 Malek-Thompson modification of frame fields

In the paper [3] another algebra underlying U-duality was presented starting from a more general form of the frame field

$$E_A{}^I = \begin{pmatrix} e_a{}^i & 0 \\ -\frac{\Pi^{a_1 a_2 b} e^i_b}{\sqrt{2!}} & \alpha\, r_{[i_1}{}^{[a_1}\, r_{i_2]}{}^{a_2]} \end{pmatrix} \tag{19}$$

that for $\alpha = 1$ coincides with (16). This ansatz for $\alpha = e^{Z_a x^a}$ leads to modification of the algebra (2) to the form [2]

$$\begin{aligned} T_a \circ T_b &= f_{ab}{}^c\, T_c\,, \\ T_a \circ T^{b_1 b_2} &= f_a{}^{b_1 b_2 c}\, T_c + 2\, f_{ac}{}^{[b_1}\, T^{b_2]c} - Z_a\, T^{b_1 b_2}\,, \\ T^{a_1 a_2} \circ T_b &= -f_b{}^{a_1 a_2 c}\, T_c + 3\, f_{[c_1 c_2}{}^{[a_1}\, \delta^{a_2]}_{b]}\, T^{c_1 c_2} + 3\, Z_{[b}\, \delta^{a_1 a_2}_{c_1 c_2]}\, T^{c_1 c_2}\,, \\ T^{a_1 a_2} \circ T^{b_1 b_2} &= -2\, f_d{}^{a_1 a_2 [b_1}\, T^{b_2]d} \end{aligned} \tag{20}$$

This modification enables to define the algebras $\mathcal{E}_3$ also for the non-unimodular Bianchi algebras.

Leibniz identities for this generalized algebra in case $n = 3$ require $Z_a = -f_{ab}{}^b$ so that they admit also the non-unimodular Bianchi algebras B**3**, B**4**, B**5**, B$\mathbf{6}_a$, B$\mathbf{7}_a$ beside those given in the preceding Section. In these cases $n_1 = 0$ and $f_a{}^{bcd} = 0$, $Z_1 = -f_{1b}{}^b = 2a$, $Z_2 = Z_3 = 0$. In other words, non-vanishing products of these algebras are

$$\begin{aligned} [T_1, T_2] = -aT_2 + n_3 T_3,\ [T_2, T_3] = 0,\ [T_3, T_1] = n_2 T_2 + aT_3\,, \\ T_1 \circ T^{12} = -a\, T^{12} + n_2\, T^{13}\,, \\ T_1 \circ T^{13} = -a\, T^{13} - n_3\, T^{12}\,, \\ T_2 \circ T^{23} = -a\, T^{13} - n_3\, T^{12}\,, \\ T_3 \circ T^{23} = a\, T^{12} - n_2\, T^{13}, \end{aligned} \tag{21}$$

where the values of parameters a, n_2, n_3 are given in the Table 1.

3.1 Twisted generalized frame fields

It is understandable that for structure constants $\mathcal{F}'_{AB}{}^C$ of the modified algebra (21) the generalized frame fields $E_A{}^I$ of the form (16) do not satisfy relations (13). For example, the formula (16) for Bianchi algebra B**5** gives

$$E_A{}^I = diag(1, e^{x_1}, e^{x_1}, \frac{1}{2}, e^{-x_1}, \frac{1}{2}, e^{-x_1}, \frac{1}{2}, e^{-2x_1})$$

and

$$\hat{\pounds}_{E_1} E_6{}^I = (0, 0, 0, 0, 0, e^{-2x_1}).$$

But, as follows from (21),

$$\mathcal{F}'_{1,6}{}^C E_C{}^I = (0, 0, 0, 0, 0, 0).$$

To satisfy relations

$$\hat{\pounds}_{E'_A} E'_B{}^I = -\mathcal{F}'_{AB}{}^C \, E'_C{}^I , \tag{22}$$

where $\mathcal{F}'_{AB}{}^C$ are structure coeficients of the algebra (21) the generalized frame fields must be modified by a twist matrix T as $E'_A{}^I = E_A{}^J T_J{}^I$ [2, 3]. It concerns the non-unimodular Bianchi algebras B**3**, B**4**, B**5**, B$\mathbf{6}_a$, B$\mathbf{7}_a$ and the twist matrix is

$$T_J{}^I = \begin{pmatrix} \mathbf{1}_3 & 0 \\ 0 & e^{2a\,x_1}\mathbf{1}_3 \end{pmatrix}. \tag{23}$$

where a is given in the Table 1.

Corresponding twisted generalized frame fields are

$$E'_A{}^I = e^{x_1} \begin{pmatrix} e^{-x_1} & 0 & 0 & 0 & 0 & 0 \\ 0 & \cosh x_1 & \sinh x_1 & 0 & 0 & 0 \\ 0 & \sinh x_1 & \cosh x_1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{2}\cosh x_1 & -\frac{1}{2}\sinh x_1 & 0 \\ 0 & 0 & 0 & -\frac{1}{2}\sinh x_1 & \frac{1}{2}\cosh x_1 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{2}e^{-x_1} \end{pmatrix}$$

for B**3**,

$$E'_A{}^I = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{x_1} & -e^{x_1}x_1 & 0 & 0 & 0 \\ 0 & 0 & e^{x_1} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{e^{x_1}}{2} & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{2}e^{x_1}x_1 & \frac{e^{x_1}}{2} & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{2} \end{pmatrix}$$

for B**4**,

$$E'_A{}^I = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{x_1} & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{x_1} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{e^{x_1}}{2} & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{e^{x_1}}{2} & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{2} \end{pmatrix}$$

for B**5**,

$$E'_A{}^I = e^{a\,x_1}\begin{pmatrix} e^{-a\,x_1} & 0 & 0 & 0 & 0 & 0 \\ 0 & \cosh x_1 & \sinh x_1 & 0 & 0 & 0 \\ 0 & \sinh x_1 & \cosh x_1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{2}\cosh x_1 & -\frac{1}{2}\sinh x_1 & 0 \\ 0 & 0 & 0 & -\frac{1}{2}\sinh x_1 & \frac{1}{2}\cosh x_1 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{2}e^{-a\,x_1} \end{pmatrix}$$

for B$\mathbf{6}_a$,

$$E'_A{}^I = e^{a\,x_1}\begin{pmatrix} e^{-a\,x_1} & 0 & 0 & 0 & 0 & 0 \\ 0 & \cos x_1 & -\sin x_1 & 0 & 0 & 0 \\ 0 & \sin x_1 & \cos x_1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{2}\cos x_1 & -\frac{1}{2}\sin x_1 & 0 \\ 0 & 0 & 0 & \frac{1}{2}\sin x_1 & \frac{1}{2}\cos x_1 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{2}e^{-a\,x_1} \end{pmatrix}$$

for B$\mathbf{7}_a$.

These twisted generalized frame fields then satisfy relations (22).

4 Conclusions

We have classified six-dimensional Leibniz algebras (2) and (20) starting from Bianchi classification of three-dimensional Lie algebras.

Up to linear transformations from $E_3 = SL(3) \times SL(2)$ we have obtained seven inequivalent algebras (2). Two of them, obtained from Abelian Lie algebra B$\mathbf{1}$, are six-dimensional Lie algebras (7) that can be considered extensions of semi-Abelian four-dimensional Drinfel'd double. The other five are unique Leibniz extensions (9) of unimodular Bianchi algebras B$\mathbf{2}$, B$\mathbf{6}_0$, B$\mathbf{7}_0$, B$\mathbf{8}$, B$\mathbf{9}$ (see Table 1). For all of these algebras we have calculated explicit forms of generalized frame fields and checked that they satisfy relations (13).

Beside that we have obtained five inequivalent generalized algebras (20) corresponding to the Bianchi algebras B$\mathbf{3}$, B$\mathbf{4}$, B$\mathbf{5}$, B$\mathbf{6}_a$, B$\mathbf{7}_a$. Their products are given by relations (21). In this case generalized frame fields given by (16) must be twisted by matrix (23) to satisfy relations (22) for the generalized algebra.

As we have complete classification of algebras $\mathcal{E}_3$ they represent analogs of Manin triples obtained for $n = 3$ in [6]. In case we want to study non-Abelian U-dualities or pluralities we have to look for their mutual relation given by linear transformations, more precisely, representations of the group $E_3 = SL(3) \times SL(2)$. Unfortunately, it seems that no such relations exist. Certainly there is no linear transformation between algebras (7) and (9) or (21) as the former ones are Lie and the latter not. Attempts to find a linear transformations inside these two classes by brute force failed as well. It seems that the only possibility to find non-Abelian U-dualities or pluralities for six-dimensional Leibniz algebras is a further generalization of the form (20).

References

[1] N.A. Obers, B. Piolin *U-duality and M-theory*, Phys. Rept. 318 (1999) 113, [hep-th/9809039]

[2] Yuho Sakatani, *U-duality extension of Drinfel'd double*, Prog. Theor. Exp. Phys. Volume 2020, Issue 2, February 2020, 023B08, [arXiv:1911.06320]

[3] E. Malek and D. C. Thompson, *Poisson-Lie U-duality in Exceptional Field Theory*, JHEP 0420 (2020) 058, [arXiv:1911.07833].

[4] L.D. Landau and E.M. Lifshitz, *The Classical Theory of Fields*, Pergamon Press, 1987

[5] L. Hlavatý and L. Šnobl, *Poisson–Lie T–dual models with two–dimensional targets*, Mod. Phys. Lett. A17 (2002) 429, [hep-th/0110139].

[6] L. Hlavatý and L. Šnobl, *Classification of 6-dimensional Manin triples,* [math.QA/0202209]