The existence of *p*-convex tensor products of $L_p(X)$ -spaces for the case of an arbitrary measure

A. Ya. Helemskii

Abstract

We obtain a far-reaching generalization (in several directions) of the theorem of A. Lambert on the existence of the projective tensor product of operator sequence spaces. This result is obtained in the context of spaces, generalizing p-multinormed spaces of Dales et al. which are based on an arbitrary, perhaps non-discrete measure.

Keywords: L-space, L-boundedness, p-convex tensor product, convenient measure space, inflation.

Mathematics Subject Classification (2010): 46L07, 46M05.

1. Introduction

The main result of the present paper is the following theorem. All notions that participate in it, will be gradually explained. Below X denotes an arbitrary measure space which is not atomic with finite set of atoms and which is supposed, for simplicity, to be separable.

The main Theorem. An arbitrary pair of near- $L_p(X)$ -spaces, where $1 \le p < \infty$, has a p-convex tensor product.

A far-away predecessor of this result is a theorem of A. Lambert [15, §3.1.1] on projective tensor products of his "operator sequence spaces"; the latter are situated, in a sense, between normed spaces and abstract operator spaces. Afterwards, a group of mathematicians (Dales, Polyakov, Daws, Pham, Ramsden, Laustsen, Oikhberg, Troitsky; see [6] and also [3, 4, 5]) introduced more general structures than Lambert spaces, called *p*-multinormed spaces; $p \in [1, \infty]$. After their papers, in the frame-work of the so-called non-coordinate approach, the

^{*}This paper was written with the support of the Russian Foundation for Basic Research (grant no. 19-01-00447).

L-spaces were introduced in [13]. The latter, provided $L := L_p(X)$, can be considered as "multinormed spaces based on arbitrary measures". Indeed, in the case of $X := \mathbb{N}$ with the counting measure they transform to *p*-multinormed spaces of [6] (and if, in addition, p := 2, to Lambert spaces).

"The main theorem" was proved in [13] under the additional condition that the set of atoms in X is either empty or infinite. For a time there was a suspicion that for an arbitrary X the theorem is false. (It was based on the known bad properties of $L_p(X)$ as a tensor factor, see [8, §12.1]). However, recently T. Oikhberg kindly sent to the author a preprint [17], where he has constructed an isomorphism between the categories of ℓ_p -spaces and $L_p(X)$ -spaces for arbitrary X. Apart from the independent value of this result, its proof was based on a construction that, as it happened, allowed us to dispense of the afore-mentioned additional condition on X. As a matter of fact, one can even consider tensor products of the so-called "stratified spaces", more general than $L_p(X)$ -spaces. But this is outside of the scope of this paper.

The author is indebted to T. Oikhberg and N. T. Nemesh for valuable discussions.

2. *L*-spaces and *L*-boundedness.

As usual, we denote by $\mathcal{B}(E)$ the space of all bounded operators on a normed space E, endowed with the operator norm. Two projections P and Q on E are called *transversal*, if PQ = QP = 0. The symbol \otimes is used for the algebraic tensor product of linear spaces and linear operators, and also for elementary tensors.

Choose and fix (so far arbitrary) normed space L, which we shall call the *bse* space.

Our principal example of a base space is $L_p(X)$, where $1 \le p < \infty$, and X is a measure space, which is not reduced to a finite set of atoms, or, equivalently, $L_p(X)$ is infinite-dimensional. To make our text shorter, we shall always assume that all our measures have a countable basis.

The *amplification* of a given linear space E is the tensor product $L \otimes E$. Usually we shortly denote it by LE, and an elementary tensor, say $\xi \otimes x; \xi \in L, x \in E$, by ξx . Note that LE is a left module over the algebra \mathcal{B} with the outer multiplication " \cdot ", well defined by $a \cdot (\xi x) := a(\xi)x$.

Definition 2.1. A norm on LE is called L-norm on E, if the left $\mathcal{B}(L)$ -

module LE is contractive, that is if we always have the estimate $||a \cdot u|| \leq ||a|| ||u||$. This estimate, as well as an equivalent estimate $||a \otimes \mathbf{1}_E|| \leq ||a||$, will be called *contractibility property*. The space E, endowed by an L-norm, is called \mathbf{L} -space. If we only know that we have the indicated estimate for operators of rank 1, we speak about near-L-norms and, accordingly, about near-L-spaces.

Remark 2.2. The class of near-*L*-spaces that we shall need in the proof of the main theorem, is bigger than the class of *L*-spaces. Let $L := L_p(X)$, *E* be a normed space, and the norm on *LE* is given by the identification of that space with the corresponding dense subspace in $L_p(X, E)$. Then we obtain a near-*L*space but, generally speaking, not an *L*-space: the indicated estimate fails already in the case $p = 2, E = \ell_1$ (see [8, p.147]).

As to the papers, cited above, they consider, after the translation into the "index-free" language, the case $X := \mathbb{N}$ with the counting measure. In particular, the notion of an $L_p(X)$ -space for such an X is equivalent to that of a p-multinormed space in [6].

Proposition 2.3. Suppose that E is a normed space, and a cross-norm $\|\cdot\|$ is given on LE. Then $\|\cdot\|$ is a near-L-norm on E iff for all $f \in L^*$ we have $\|f \otimes \mathbf{1}_E\| \leq \|f\|$.

 $\exists \text{ Every } a \in \mathcal{B}(L) \text{ of rank 1 acts as } \eta \mapsto f(\zeta)\xi \text{ for some } \xi \text{ and } f, \text{ so } ||a|| = ||f|| ||\xi||. \text{ It is easy to verify that } a \cdot u = \xi \Big((f \otimes \mathbf{1}_E) u \Big), \text{ for all } u \in LE. \text{ Therefore the estimates } ||a \cdot u|| \leq ||a|| ||u|| \text{ and } ||(f \otimes \mathbf{1}_E) u|| \leq ||f|| ||u|| \text{ are equivalent. } \triangleright$

A near-*L*-space *E* becomes a normed space in the "classical" sense, if for $x \in E$, we set $||x|| := ||\xi x||$, where $\xi \in L$ is an arbitrary vector with $||\xi|| = 1$. Clearly, the result does not depend on the choice of ξ . The obtained normed space is called the *underlying space* of a given *L*-space, and the latter is called an *L*-quantization of a former. We use such a term by analogy with quantizations in operator space theory; see, e.g., [9], [10] or [12].

It is easy to verify that the complex plane \mathbb{C} has the only *L*-quantization, given by the identification of $L\mathbb{C}$ with *L*. However, as a rule, general normed spaces have a lot of *L*-quantizations. In particular, by endowing *LE* with the norm of (non-completed) projective, respectively injective tensor product of normed spaces, we obtain two, generally speaking, different *L*-quantization, called maximal, respectively minimal. (See [13] for details.)

Suppose we are given an operator $\varphi: E \to F$ between linear spaces. Denote,

for brevity, the operator $\mathbf{1}_L \otimes \varphi : LE \to LF$ (taking ξx to $\xi \varphi(x)$) by φ_∞ and call it *amplification* of φ . Obviously, φ_∞ is a morphism of left $\mathcal{B}(L)$ -modules.

Definition 2.4. An operator $\varphi : E \to F$ between *L*-spaces is called *L*bounded or *L*-contractive, if the operator φ_{∞} is bounded or contractive, respectively.

As to numerous examples and counterexamples see, e.g., [13], and also [6].

To define amplifications of bilinear operators, we need a certain additional structure, called in what follows \diamondsuit -operation or "diamond operation" on L. This is a bilinear operator $\diamondsuit: L \times L \to L$ of norm one. We shall write $\xi \diamondsuit \eta$ instead of $\diamondsuit(\xi, \eta)$.

For "most" X, $L_p(X)$ has a natural, in a sense, diamond operation (see [13, §3], and also, in the case of a discrete measure, [15, §1.2.2]). But we emphasize that our main theorem is valid for arbitrary \diamond .

Now let $\mathcal{R} : E \times F \to G$ be a bilinear operator between linear spaces. Its *amplification* is the bilinear operator $\mathcal{R}_{\infty} : LE \times LF \to LG$, well-defined (because of the bilinearity) on elementary tensors by $\mathcal{R}_{\infty}(\xi x, \eta y) = (\xi \Diamond \eta) \mathcal{R}(x, y)$.

Definition 2.5. A bilinear operator \mathcal{R} between *L*-spaces is called *L*-bounded or *L*-contractive, if its amplification is (just) bounded, or contractive, respectively.

In the case $L = \ell_2$ and a particular \diamondsuit , taking sequences $\{\xi_n\}$ and $\{\eta_n\}$ into (arbitrarily renumerated) double sequence $\{\xi_n\eta_m\}; m, n \in \mathbb{N}$, we obtain, in equivalent terms, the definition of an *L*-bounded bilinear operator, given by Lambert.

Again, see [13] for numerous examples.

3. *p*-convex tensor product and preliminaries of its existence

From now on, and up to the end of the paper, we assume that $L := L_p(X); p \in [1, \infty)$ (i.e. we are within the context of our main example of L), and that we fix an arbitrary \diamond -operation on our base space.

Let Y be a measurable subset in X. Consider the projection $P_Y \in \mathcal{B}(L)$, acting as $f \mapsto f\chi$, where χ is a characteristic function of Y. A projection of that kind will be called *proper*. Clearly, two proper projections are transversal iff the intersection of the respective measurable subsets has measure 0. Let E be a linear space. We call a projection $P \in \mathcal{B}(L)$ a support of an element $u \in LE$, if $P \cdot u = u$.

Definition 3.1. A near-*L*-space *E* is called *p*-convex, if for any $u, v \in LE$, with transversal proper supports, we have $||u + v|| \leq (||u||^p + ||v||^p)^{\frac{1}{p}}$.

The introduced class of *L*-spaces, being a generalization for arbitrary p of column operator spaces, is, in our opinion, the most interesting. For the special case $L := \ell_p$, the given definition is equivalent to the definition of a p-convex p-multinormed space, given in [6]. Also it worth mentioning, in this connection, the theory of p-operator spaces of Daws [7]; see also earlier papers of Pisier [19] and Le Merdy [16].

As an example, one can easily show that every L-space with the minimal quantization is p-convex. Another example is provided by the near-L-space from Remark 2.2.

Now let E and F be two *arbitrary chosen* near-L-spaces.

Definition 3.2. A pair (Θ, θ) , consisting of a *p*-convex *L*-space Θ and an *L*contractive bilinear operator $\theta : E \times F \to \Theta$, is called (non-completed) *p*-convex tensor product of *E* and *F* if, for every *p*-convex *L*-space *G* and every *L*-bounded bilinear operator $\mathcal{R} : E \times F \to G$, there exists a unique *L*-bounded operator $R : \Theta \to G$ such that the diagram

$$\begin{array}{c} E \times F \\ \downarrow \theta \\ \Theta \\ \hline R \\ \end{array} \xrightarrow{\mathcal{R}} G \end{array}$$

is commutative, and moreover we have $||R_{\infty}|| = ||\mathcal{R}_{\infty}||$.

In what follows, the property of the pair in question will be called *the universal* property.

We emphasize that Θ and G are supposed (in comparison to E and F) to be L-spaces, and not just near-L-spaces.

Remark 3.3. We see that the *L*-spaces Θ and *G* are assumed to be *p*-convex. Other assumptions lead to other types of tensor products. For instance, if we shall take the class of all *L*-spaces, we shall come to an essentially different concept, the so-called general tensor product of our *E* and *F*. This variety has its own existence theorem; this is Theorem 4.6 in [13]. Nevertheless *p*-convex tensor products, being in the case p = 2 intimately connected with the projective tensor products of operator spaces, discovered by Blecher/Paulsen [1] and Effros/Ruan [11], seem to be most interesting.

Thus, all notions that participate in the formulation of our main theorem, are explained, and we can proceed to its proof.

As it was mentioned in Introduction, this theorem earlier was proved under the additional assumption that X either has no atoms or has an infinite set of atoms. Such a measure space we shall call *convenient*.

We recall the construction of our desired tensor product in the case of a convenient X. Take, as underlying linear space of Θ , just $E \otimes F$, and as θ the canonical bilinear operator $\vartheta : (x, y) \mapsto x \otimes y$. So, our task is to introduce a suitable norm on $L(E \otimes F)$.

We first need an "extended" version of our fixed diamond operation, this time between elements of amplifications of linear spaces. Namely, for $u \in LE, v \in LF$ we consider the element $u \diamondsuit v := \vartheta_{\infty}(u, v) \in L(E \otimes F)$. In other words, this "diamond operation" is well defined by $\xi x \diamondsuit \eta y := (\xi \diamondsuit \eta)(x \otimes y)$, with $\xi, \eta \in L, x \in E, y \in F$.

An isometry on L will be called *proper*, if its image is the image of a proper projection. Two isometries will be called *disjoint*, if the intersection of their images is $\{0\}$.

As is well known (in equivalent terms), if X is convenient, then $L_p(X)$ possesses an infinite family of mutually disjoint proper isometries. See, e.g., [2, Cor. 9.12.18] and also [14, §14] or [21, III.A].

The following preparatory statement is crucial in our construction.

Proposition 3.4 ([13, Prop. 5.6]). Let X be convenient. Then every $U \in L(E \otimes F)$ can be represented as

$$a \cdot \sum_{k=1}^{n} I_k \cdot (u_k \diamondsuit v_k),$$

where $a \in \mathcal{B}(L), u_k \in LE, v_k \in LF$ and I_k are pairwise disjoint proper isometries on L.

Now we have the right to take $U \in L(E \otimes F)$ and assign to it the number

$$||U||_{pL} := \inf \left\{ ||a|| \left(\sum_{k=1}^{n} ||u||^{p} ||v||^{p} \right)^{\frac{1}{p}} \right\},\$$

where the infimum is taken over all possible representations of U in the indicated form. It turns out that it is just what we need:

Theorem 3.5. ([13, Theorem 5.18]). The function $U \mapsto ||U||_{pL}$ is a L-norm on $E \otimes F$, and the pair $(E \otimes_{pL} F, \vartheta)$, where $E \otimes_{pL} F$ denotes $E \otimes F$, endowed with the indicated L-norm, is a p-convex tensor product of E and F.

Remark 3.6. It was assumed in the cited theorem that E and F are L-spaces, and the \diamond -operation has the property $\|\xi \diamond \eta\| = \|\xi\| \|\eta\|$. However one can easily notice that its proof uses the estimate $\|a \otimes \mathbf{1}_E\| \leq \|a\|$; $a \in \mathcal{B}(L)$ from Definition 2.1 only for operators of rank 1, and only the property of \diamond to have norm 1.

We proceed to the main contents of the present paper. How can one behave, if X is not convenient, that is the set of its atoms is not empty and finite? It turns out that it is possible to reduce the "unconvenient" case to the "convenient" one.

For an arbitrary linear space, say H, let us consider the algebraic direct sum of a countable family of its copies. So, it consists of eventually zero sequences $\bar{\xi} = (\xi_1, \xi_2, ...); \xi_k \in H$. If H has a norm, we set $\|\bar{\xi}\| := (\sum_k \|\xi_k\|^p)^{\frac{1}{p}}$ and call the resulting normed space standard extension of H.

Now, for our fixed X, we denote by $\mathbb{N}X$ the measure space which is the disjoint union of a countable family of copies of X: $\mathbb{N}X := X_1 \sqcup X_2 \sqcup \cdots$. Clearly, $\mathbb{N}X$ is convenient. Therefore the space $L_p(\mathbb{N}X)$ satisfies, with $\mathbb{N}X$ in the role of X, the conditions of Theorem 3.5.

Let \mathbb{L} be the algebraic direct sum of a countable family of copies of L. Then we have the right to consider on the spaces $\mathbb{L}E$ and $\mathbb{L}F$ the norm of the standard extension of LE and LF, respectively.

We do not know, whether an arbitrary L-space is also a \mathbb{L} -space with respect to the norm of the standard extension of the given L-norm; may be not. Nevertheless, the following fact is valid.

Proposition 3.7. If E is a near-L-space, then it is also a near-L-space.

⊲ It is easy to verify that the norm on LE, as well as the norm on LE, is a cross–norm with respect to the norm of the underlying space of the given near-L-space. Therefore, by virtue of Proposition 2.3, it suffices to show that for every $\mathbf{f} \in \mathbb{L}^*$ we have $\|\mathbf{f} \otimes \mathbf{1}_E\| \leq \|\mathbf{f}\|$. In what follows, we omit the easy case p = 1.

For $\xi \in L$ and $n \in \mathbb{N}$ we denote by $\overline{\xi}^n \in \mathbb{L}$ the sequence with the *n*-th term ξ and all others zeroes. Introduce the functionals $f_n : L \to \mathbb{C} : \xi \mapsto \mathbf{f}(\overline{\xi}^n)$. Fix, for a moment, *n* and consider an element $\overline{u} = (u_1, u_2, ...) \in \mathbb{L}E$ with $u_n := \xi x$ for some $\xi \in L, x \in E$ and $u_m = 0$ for $m \neq n$. We see that $\mathbf{f} \otimes \mathbf{1}_E(\bar{u}) = \sum_m (f_m \otimes \mathbf{1}_E)(u_m)$. Since sums of such elements give the whole $\mathbb{L}E$, the same equality is valid for all $\bar{u} \in \mathbb{L}E$.

But since we know what is E, the same Proposition 2.3 gives $||f_m \otimes \mathbf{1}_E|| \leq ||f_m||$, for all m. Further, it is known (and easy to verify) that $||\mathbf{f}|| = (\sum_m ||f_m||^q)^{\frac{1}{q}}$, q is the number, conjugate to p. Therefore for every $\bar{v} = (v_1, ...) \in \mathbb{L}E$ we obtain

$$\|(\mathbf{f} \otimes \mathbf{1}_E)(\bar{v})\| \le \sum_m \|f_m\| \|v_m\| \le (\sum_m \|f_m\|^q)^{\frac{1}{q}}) (\sum_m \|v_m\|^p)^{\frac{1}{p}}) = \|\mathbf{f}\| \|\bar{v}\|.$$

The spaces L and \mathbb{L} are connected by the isometry $J : L \to \mathbb{L} : \xi \mapsto (\xi, 0, 0, ...)$ and the coisometry $Q : \mathbb{L} \to L : (\xi, \xi_2, ..., \xi_n, ...) \mapsto \xi$; of course, $QJ = \mathbf{1}_L$. For every linear space G we shall write J_G instead of $J \otimes \mathbf{1}_G : LG \to \mathbb{L}G$ and Q_G instead of $Q \otimes \mathbf{1}_G : \mathbb{L}G \to LG$.

Our task is to construct a pair $(\Theta, \theta : E \times F \to \Theta)$, satisfying the conditions of Definition 3.1. We shall show that, similarly to Theorem 3.5, we can take $E \otimes F$ as the underlying linear space of Θ , and the canonical bilinear operator as θ .

Where to look for the desired norm on $L(E \otimes F)$?

Using the recipe of Proposition 3.7, we transform E and F into near- \mathbb{L} -spaces. Then we introduce the bilinear operator $\bar{\Diamond} : \mathbb{L} \times \mathbb{L} \to \mathbb{L}$ by

$$\bar{\xi}\bar{\Diamond}\bar{\eta} := J(Q_E\bar{\xi}\Diamond Q_F\bar{\eta}).$$

Clearly, it is a diamond operation on \mathbb{L} .

But since $\mathbb{N}X$ (though, perhaps, not X) is convenient, there exists a p-convex tensor product of E and F as that of near- \mathbb{L} -spaces with respect to any diamond operation on \mathbb{L} ; in particular, we choose $\overline{\Diamond}$. Moreover, as the \mathbb{L} -space Θ we can take $E \otimes F$ with the respective norm on $\mathbb{L}(E \otimes F)$) that we shall denote by $\|\cdot\|_{p\mathbb{L}}$.

Finally, we introduce a norm on $L(E \otimes F)$, induced by the injection $J_{E \otimes F}$. In other words, for $U \in L(E \otimes F)$ we set $||U||_{pL} := ||J_{E \otimes F}(U)||_{p\mathbb{L}}$.

This norm will turn out to be our desired L-norm on the desired tensor product. If there is no danger of confusion, we shall omit indices in the notation of the respective norms.

We must verify the needed requirements.

Take $a \in \mathcal{B}(L), U \in L(E \otimes F)$ and set $\bar{a} := JaQ \in \mathcal{B}(\mathbb{L}), \bar{U} := J_{E \otimes F}(U) \in \mathbb{L}(E \otimes F)$. Then

$$||a \cdot U|| = ||(Ja \otimes \mathbf{1}_{E \otimes F})(U)|| = ||(JaG) \cdot (J_{E \otimes F}(U))|| = ||\bar{a} \cdot \bar{U}|| \le ||\bar{a}|| ||\bar{U}|| = ||a|| ||U||,$$

so that we have the contractibility property.

If $U_k \in L(E \otimes F)$; k = 1, 2 have transversal proper projections P_k in L, then $\mathbb{P}_k := JP_kQ$ are transversal proper projections in \mathbb{L} , that are supports of $J_{E \otimes F}U_k$. Therefore the *p*-convexity of $E \otimes F$ as an \mathbb{L} -space implies the *p*-convexity of $E \otimes F$ as an L-space.

Finally, for $u \in LE, v \in LF$ the equality $J_{E\otimes F}(u \diamond v) = J_E u \diamond J_F v$ and \mathbb{L} -contractibility of ϑ imply that $||u \diamond v|| \leq ||u|| ||v||$, that is the desired Lcontractibility of ϑ .

Now the main thing remains: the universal property. In this connection, the following notion will be useful.

Definition 3.8. Let G be a p-convex L-space and simultaneously a p-convex L-space. Then the latter space is called an *inflation* of the former space, if J_G is an isometry, and Q_G is a coisometry.

For example, if G is the minimal L-space, then it is easy to show that G as the minimal L-space is an inflation of the former. As another example, suppose that G belongs to the class SQ_p , i.e. it is a subspace of a quotient space of some $L_p(Y)$. We make it an L-space and an L-space by the identification of LG and $\mathbb{L}G$ with the corresponding subspaces in $L_p(X, G)$ and $L_p(\mathbb{N}X, G)$. Then the second space is an inflation of the first one. The required properties follow from Theorems 1.35 and 1.41 in [6].

In the following three propositions we suppose that G is a given L-space that has an inflation, and we fix the latter.

Let $\mathcal{R}: E \times F \to G$ be a bilinear operator which is *L*-bounded as an operator between near-*L*-spaces with respect to our initial \diamond -operation. We denote by $\overline{\mathcal{R}}$ the same bilinear operator as an operator between near-*L*-spaces. Speaking about its *L*-boundedness, we mean the diamond operation $\overline{\diamond}$, defined above.

Proposition 3.9. Our $\overline{\mathcal{R}}$ is also \mathbb{L} -bounded, and we have $\|\overline{\mathcal{R}}_{\infty}\| = \|\mathcal{R}_{\infty}\|$. \triangleleft The estimate $\|\overline{\mathcal{R}}_{\infty}\| \leq \|\mathcal{R}_{\infty}\|$ follows from the formula

$$\bar{\mathcal{R}}_{\infty}(\bar{u},\bar{v}) = J_G\Big(\mathcal{R}_{\infty}(Q_E\bar{u},Q_F\bar{v})\Big); \bar{u} \in \mathbb{L}E, \bar{v} \in \mathbb{L}F,$$

which is an easy corollary of the definition of \Diamond . The inverse estimate follows from the formula

$$\mathcal{R}_{\infty}(u,v) = Q_G\Big(\bar{\mathcal{R}}_{\infty}(J_E u, J_F v)\Big),$$

an easy corollary of the obvious equality $\xi \Diamond \eta = Q(J\xi \bar{\Diamond} J\eta)$. \triangleright

Consider the operators $R_{\infty} : L(E \otimes F) \to LG$ and $\bar{R}_{\infty} : \mathbb{L}(E \otimes F) \to \mathbb{L}G$. These are the amplifications of the operator $R : E \otimes F \to G$ that is associated with \mathcal{R} and $\bar{\mathcal{R}}$, respectively.

Proposition 3.10. We have $||R_{\infty}|| \leq ||\bar{R}_{\infty}||$.

 \lhd This estimate follows from the formula

$$R_{\infty}(U) = Q_G\Big(\bar{R}_{\infty}(J_{E\otimes F}(U))\Big)$$

Obviously, one should only verify the latter equality on U of the form $\xi(x \otimes y); \xi \in L, x \in E, y \in F$. Then

$$R_{\infty}(U) = (QJ\xi)\bar{R}(x\otimes y) = Q_G\Big(\bar{R}_{\infty}(J\xi(x\otimes y)\Big) = Q_G\Big(\bar{R}_{\infty}(J_{E\otimes F}(U)\Big).$$

Proposition 3.11. If G is as above, then for an arbitrary L-bounded bilinear operator $\mathcal{R} : E \times F \to G$ and the associated linear operator $R : E \otimes F \to G$ we have $||R_{\infty}|| = ||\mathcal{R}_{\infty}||$.

 \triangleleft Consider G with the L-norm of the given inflation. Because of the universal property of the tensor product of our E and F as near-L-spaces, we have $\|\bar{\mathcal{R}}_{\infty}\| = \|\bar{\mathcal{R}}_{\infty}\|$. Combining this with two previous propositions, we obtain the estimate $\|\mathcal{R}_{\infty}\| \leq \|\mathcal{R}_{\infty}\|$.

Further, it follows from the formula $u \diamondsuit v = Q_{E \otimes F}(J_E u \diamondsuit J_F v)$, which can be easily verified on elementary tensors, that for all $u \in LE, v \in LF$ we have

$$\|\mathcal{R}_{\infty}(u,v)\| = \|R_{\infty}(u\Diamond v)\| \le \|R_{\infty}\| \|Q_{E\otimes F}(J_E u\bar{\Diamond}J_F v)\| \le \|R_{\infty}\| \|J_E u\bar{\Diamond}J_F v\|.$$

But $\vartheta : (x, y) \mapsto x \otimes y$ is \mathbb{L} -contractive with respect to the corresponding near- \mathbb{L} -norms and the operation $\bar{\Diamond}$. Therefore $\|J_E u \bar{\Diamond} J_F v\| \leq \|J_E u\| \|J_F v\|$. Consequently we have $\|\mathcal{R}_{\infty}(u, v)\| \leq \|R_{\infty}\| \|u\| \|v\|$, that is $\|\mathcal{R}_{\infty}\| \leq \|R_{\infty}\|$. \triangleright

4. Existence of inflations and completion of the proof of the main theorem

Thus, we see that for concluding the proof of the main theorem it suffices to know that every L-space has at least one inflation. This for some time we did not know. It is natural to begin with the testing of the standard extension of the given L-norm. However the existence of near-L-spaces that are not L-spaces (see Remark 2.2) makes one to have doubts; it seems to us that it does not fit. Nevertheless, by virtue of a recent result of T. Oikhberg, mentioned in Introduction, one can show that inflations do always exist. Indeed it is easily seen that, for arbitrary measure spaces X and Y with infinite-dimensional separable $L_p(X)$ and $L_p(Y)$, his argument actually allows us to construct a certain $L_p(Y)$ -norm on some G, embarking from a given $L_p(X)$ -norm on the same G. We use his method for a proof of the following fact.

Theorem 4.1. Let X be as in the formulation of the main theorem. Then for $L := L_p(X); p \in [1, \infty)$ every L-space has an inflation.

Before the proof, we note that we shall construct an inflation, which essentially differs from the standard extension of the given L-space; see the discussion above.

In what follows, if Z is a measurable subset of some measure space, say Y, we shall denote its normalized in $L_p(Y)$ characteristic function by $\widehat{\chi}(Z)$.

 $\triangleleft \triangleleft$ We need the following preparatory statement.

Lemma. (Here we strictly follow the argument of Oikhberg). Let Y be an arbitrary measure space, \tilde{L} a subspace in $L_p(Y)$, which is the linear span of several characteristic functions of measurable sets. Then there exists a projection on \tilde{L} in $\mathcal{B}((L_p(Y)))$ of norm 1.

⊲ There exist disjoint subsets of non-zero measure Z_k in Y, such that $\tilde{L} = span\{\hat{\chi}(Z_k)\}$. Take in $L_q(Y) = L_p(Y)^*$ (here q is the conjugate number to p) norm 1 functions $\tilde{\xi}_k$, such that $\tilde{\xi}_k = 0$ outside Z_k , and $\langle \tilde{\xi}_k, \hat{\chi}(Z_k) \rangle = 1$. Consider the operator $P: L_p(Y) \to L_p(Y) : \eta \mapsto \sum_{k=1}^n \langle \tilde{\xi}_k, \eta \rangle \hat{\chi}_k$. Clearly, P is identical on \tilde{L} . Further, since for $\eta \in L_p(Y)$ we have $P(\eta) = \sum_{k=1}^n \langle \tilde{\xi}(Z_k), \eta_k \rangle \hat{\chi}(Z_k)$, where $\eta_k = \eta$ on Z_k and $\eta_k = 0$ outside Z_k , we easily see that $\|P(\eta)\| \leq \|\eta\|$. ⊳

So, we are given, for $L := L_p(X)$, an *L*-space *G*. At first we want to introduce a certain norm on $\mathbb{L}^0 G$, where \mathbb{L}^0 is a dense subspace in \mathbb{L} , consisting of simple functions.

Every $u \in \mathbb{L}^0 G$ can be represented, for some family $Y_k; r = 1, ..., n$ of pairwise disjoint subsets of non-zero measure in $\mathbb{N}X$, as $\sum_{k=1}^n \widehat{\chi}(Y_k)x_k, x_k \in G$. Take in X an arbitrary family Z_k of pairwise disjoint subsets of non-zero measure and set $v := \sum_{k=1}^n \widehat{\chi}(Z_k)x_k \in LG$. We put ||u|| := ||v||.

The subsequent argument consists of several natural stages.

1. The number ||u|| does not depend on the choice of the subsets Z_k .

 \triangleleft Let Z'_k be another family, and $v' := \sum_{k=1}^n \widehat{\chi}(Z'_k) x_k \in LG$. Consider the operator $J : span\{\widehat{\chi}(Z_k)\} \to span\{\widehat{\chi}(Z'_k)\} : \widehat{\chi}(Z_k) \mapsto \widehat{\chi}(Z'_k)$; clearly, it is an

isometric isomorphism. By the preceding lemma, there exists a projection P: $L \to span\{\widehat{\chi}(Z_k)\}$ of norm 1. Therefore ||JP|| = 1, so that the contractibility property for L-spaces implies $||v'|| = ||JPv|| \le ||v||$. A similar argument provides the inverse estimate. \triangleright

2. The number ||u|| does not depend on the representation of u as a sum of elementary tensors of the indicated form.

If we have another representation of our u, then, breaking the subsets, corresponding to both families, into the same disjoint unions and using the linear independence of the respective characteristic functions, we see that both representations lead to the same representation. To show that the resulting representation gives the same number as the initial one, it suffices, in its turn, to show that the number does not change after breaking one of the initial subsets into two disjoint subsets of non-zero measure, say, after breaking Y_1 into Y' and Y''. Thus, the new representation has the form $\widehat{\chi}(Y')z_1 + \widehat{\chi}(Y'')z_2 + \sum_{k=2}^n \widehat{\chi}(Y_k)x_k$ for some $z_1, z_2 \in G$. Since the tensor factors $\widehat{\chi}(Y')$ and $\widehat{\chi}(Y'')$ are linearly independent, it follows that $z_l = \lambda_l x_1; l = 1, 2$ for some $\lambda_1, \lambda_2 \in \mathbb{C}$.

Recall the subsets $Z_k \subset X$. By stage 1, we can assume that we can break Z_1 into two disjoint sets of non-zero measure, say, Z' and Z''. Therefore, if we consider the indicated new representation of our u, then the mentioned recipe gives the number ||v''||, where $v'' := \lambda_1 \hat{\chi}(Z') x_1 + \lambda_2 \hat{\chi}(Z'') x_1 + \sum_{k=2}^n \hat{\chi}(Z_k) x_k$.

Obviously we have $\|\lambda_1 \hat{\chi}(Z') + \lambda_2 \hat{\chi}(Z'')\|^p = |\lambda_1|^p + |\lambda_2|^p = 1$. This easily implies that there exist operators J_1, J_2 of norm 1, acting on the space $span\{\hat{\chi}(Z'), \hat{\chi}(Z''), \hat{\chi}(Z_k); k = 2, ..., n\}$ and such that

$$J_1(\widehat{\chi}(Z_1) = \lambda_1 \widehat{\chi}(Z') + \lambda_2 \widehat{\chi}(Z'') \quad \text{and} \quad J_2(\lambda_1 \widehat{\chi}(Z') + \lambda_2 \widehat{\chi}(Z'')) = \widehat{\chi}(Z_1).$$

Further, the lemma provides a projection $P: L \to span\{\widehat{\chi}(Z'), \widehat{\chi}(Z''), \widehat{\chi}(Z_k); k = 2, ..., n\}$ of norm 1. Therefore, the contractibility property for G as for an L-space, being applied to the operators J_1P and J_2P , gives ||v|| = ||v''||. \triangleright

So, stages 1 and 2 together show that the number ||u||; $u \in \mathbb{L}^0 G$ is well defined.

3. The function $u \mapsto ||u||$ is an \mathbb{L}^0 -norm on G.

 \triangleleft Obviously, this function is a norm on $\mathbb{L}^0 G$. So, it remains to show that for $a \in \mathcal{B}(\mathbb{L}^0)$ and $u \in \mathbb{L}^0 G$ we have $||a \cdot u|| \leq ||a|| ||u||$.

Let u has its initial representation, and $a \cdot u$ is represented as $\sum_{l=1}^{m} \widehat{\chi}(Y_l^1) y_l$, where Y_l^1 are some pairwise disjoint subsets in $\mathbb{N}X$ of non-zero measure. Take Z_k as before and choose arbitrarily one more family $Z_l^1; l = 1, ..., m$ of pairwise disjoint subsets of non-zero measure in X.

Consider the operators $I : span\{\widehat{\chi}(Y_k)\} \to span\{\widehat{\chi}(Z_k)\} : \widehat{\chi}(Y_k) \mapsto \widehat{\chi}(Z_k)$ and $J : span\{\widehat{\chi}(Y_l^1)\} \to span\{\widehat{\chi}(Z_l^1)\} : \widehat{\chi}(Y_l^1) \mapsto \widehat{\chi}(Z_l^1)$. Clearly, both are isometric isomorphisms. Further, the construction of the norm on $\mathbb{L}^0 G$ exactly means that $\|u\| = \|(I \otimes \mathbf{1}_G)u\|$ and $\|a \cdot u\| = \|(J \otimes \mathbf{1}_G)(a \cdot u)\|$.

Our lemma provides projections $P : L \to span\{\widehat{\chi}(Z_k)\}$ and $Q : \mathbb{L} \to span\{\widehat{\chi}(Z_l^1)\}$ of norm 1. We easily see that $((P \otimes \mathbf{1}_G)(I \otimes \mathbf{1}_G))(u) = (I \otimes \mathbf{1}_G)(u)$ and $(Q \otimes \mathbf{1}_G)(a \cdot u) = a \cdot u$. Now set $b := JQaI^{-1}P : L \to L$. Then we have $b \cdot ((I \otimes \mathbf{1}_G)u) = (JQa \otimes \mathbf{1}_G)u = (J \otimes \mathbf{1}_G)(a \cdot u)$. Therefore the contractibility property of an L-space G gives $||a \cdot u|| = ||b \cdot ((I \otimes \mathbf{1}_G)u)|| \le ||b|| ||(I \otimes \mathbf{1}_G)u|| \le ||a|| ||u||$.

4. There exists an \mathbb{L} -norm on G which is an extension of the constructed \mathbb{L}^{0} -norm.

⊲ Take some $u = \sum_{k=1}^{K} \xi_k x_k \in \mathbb{L}G$; $\xi_k \in \mathbb{L}, x_k \in G$. Consider a sequence $u_n := \sum_k \xi_{k,n} x_k \in \mathbb{L}^0 G$, where $\xi_{k,n}$ converges to ξ_k in \mathbb{L} . Clearly, the sequence $||u_n||$ converges. Denote its limit by ||u||; obviously, it does not depend on the choice of $\xi_{k,n}$.

Take another representation of u, say $u = \sum_{l=1}^{M} \eta_l y_l$, and denote by ||u||'the number, corresponding to this representation. Take linearly independent $z_r \in G; r = 1, ..., N$, such that $x_k = \sum_{r=1}^{N} \lambda_{kr} z_r$ and $y_l = \sum_{r=1}^{N} \mu_{lr} z_r$ for some $\lambda_{k,r}, \mu_{l,r} \in \mathbb{C}$. We come, in particular, to the representation $u = \sum_{r=1}^{N} (\sum_k \lambda_{k,r} \xi_k) z_k$; this leads to some number, denoted by ||u||''.

Recalling $\xi_{k,n}$, we see that ||u||'' is the limit of the sequence $||\sum_{r=1}^{N} (\sum_{k} \lambda_{k,r} \xi_{k,n}) z_r||$, that clearly coincides with $||u_n||$ above. Therefore ||u||'' = ||u||.

Similarly, ||u||' is the number, corresponding to the representation u as $\sum_{r=1}^{N} (\sum_{l} \mu_{l,r} \eta_{l}) z_{k}$. But, since z_{r} are linearly independent, $\sum_{k} \lambda_{k,r} \xi_{k} = \sum_{l} \mu_{l,r} \eta_{l}$. Consequently, ||u||' coincides with ||u||'', and hence with ||u||.

Thus, we have a well defined function $u \mapsto ||u||$ on $\mathbb{L}G$, which is obviously a seminorm. Let us show that for all $a \in \mathcal{B}(\mathbb{L})$ and $u \in \mathbb{L}G$ we have $||a \cdot u|| \le ||a|| ||u||$.

At first suppose that $u \in \mathbb{L}^0 G$. Let $u = \sum_{k=1}^N \xi_k x_k$, where $\xi_k \in \mathbb{L}^0$, $x_k \in G$. By the lemma, there is a projection of norm 1, say P, of \mathbb{L} on a linear span of several characteristic functions of measurable sets, such that $u = P \cdot u$.

Take $\epsilon > 0$. Since all $a(\xi_k)$ can be approximated by simple functions, the same lemma gives a projection of norm 1, say Q, on a linear span of several

characteristic functions of measurable sets, such that $||a \cdot \xi_k - Qa \cdot \xi_k|| < \epsilon$. Consider an operator on \mathbb{L} , acting as QaP. Then stage 3 gives $||(QaP) \cdot u|| \le ||QaP|| ||u|| \le$ ||a|| ||u||. Hence $||a \cdot u|| \le ||a \cdot u - (QaP) \cdot u|| + ||(QaP) \cdot u|| < \epsilon + ||a|| ||u||$. This, of course, implies $||a \cdot u|| \le ||a|| ||u||$.

Finally, let u be an arbitrary element in $\mathbb{L}G$, and $u = \sum_{k=1}^{N} \xi_k x_k$, where $\xi_k \in \mathbb{L}, x_k \in G$. Since a is bounded, we see that for some sequence $u_n \in \mathbb{L}^0 G$ we simultaneously have $u_n \to u$ and $a \cdot u_n \to a \cdot u$; $n \to \infty$. Therefore taking limit in the already obtained estimate $||a \cdot u_n|| \leq ||a|| ||u_n||$ we have our desired estimate.

Now let us show that our seminorm is actually a norm.

Take $u \neq 0$ and represent it as $\sum_{k=1}^{N} \xi_k x_k$, with linearly independent $\xi_k \in \mathbb{L}$ and $x_1 \neq 0$. There exist $a \in \mathcal{B}(\mathbb{L})$ such that $a(\xi_1) = \xi_1$ and $a(\xi_k) = 0; k > 1$, and also, for every k = 1, ..., N a sequence $\xi_{k,n} \in \mathbb{L}^0; n \in \mathbb{N}$, converging to ξ_k . Set $u_n := \sum_{k=1}^{N} \xi_{k,n} x_k$; we have $||a \cdot u_n|| \geq ||a(\xi_{1,n}) x_1|| - \sum_{k=2}^{N} ||a(\xi_{k,n}) x_k||$. But clearly $a(\xi_{1,n})$ converges to ξ_1 , and $a(\xi_{k,n})$ converges to 0 for other k. Hence for sufficiently big n we have $||a \cdot u_n|| \geq \epsilon$ for some $\epsilon > 0$. Combining this with the estimation above, we obtain that $||u_n|| \geq \epsilon/||a||$. Therefore ||u|| > 0. \triangleright

5. The p-convexity is preserved by passing from L-spaces to \mathbb{L} -spaces.

 \triangleleft Suppose, at first, that u_1, u_2 with transversal supports lie in $\mathbb{L}^0 G$. By definition of the norm in $\mathbb{L}^0 G$, there exist a family $Z_1^1, ..., Z_N^1, Z_1^2, ..., Z_M^2$ of pairwise disjoin subsets of non-zero measure in X such that, for some $v_1, v_2 \in LG$ of the form $v_1 := \sum_{k=1}^N \widehat{\chi}(Z_k^1) x_k, v_2 := \sum_{k=1}^M \widehat{\chi}(Z_l^2) y_l$, respectively, we have $||u_1|| = ||v_1||$, $||u_2|| = ||v_2||$ and $||u_1 + u_2|| = ||v_1 + v_2||$. Since v_1, v_2 obviously have transversal supports, and our L-space is p-convex, we have $||u_1 + u_2||^p \leq ||u_1||^p + ||u_2||^p$.

Now take arbitrary $u_1, u_2 \in \mathbb{L}G$ with transversal supports. Clearly for k = 1, 2there exist a sequence $u_k^n \in \mathbb{L}^0 G$ with the same support as u_k , converging to u_k . Then, passing to limits, we obtain the desired estimate. \triangleright

It is clear that all statements and arguments in stages 1-5 are valid, if we replace $\mathbb{L} := L_p(\mathbb{N}X)$ by $L_p(Y)$ for an arbitrary measure space Y. Now we concentrate on our concrete situation.

End of the proof. It remains to show that J_G is an isometry, and Q_G is a coisometry.

 \triangleleft Take at first $u \in L^0G$ and represent it as $\sum_{k=1}^N \widehat{\chi}(Y_k) x_k$ with pairwise disjoint $Y_k \subset X$. Then $J_G(u)$, as an element of the subspace \mathbb{L}^0G of $\mathbb{L}G$ has the same representation, only now we must consider Y_k as subsets in the first summand X_1

in $\mathbb{N}X = X_1 \sqcup X_2 \sqcup \ldots$ Therefore, calculating $||J_G(u)||$ by the prescribed recipe, we can take as Z_k the initial Y_k , and the same x_k . But then the respective v is just u, therefore $||J_G(u)|| = ||u||$. Thus, the restriction of $J_G : LG \to \mathbb{L}G$ on a dense subspace in LG is an isometry, so the same is true for J_G .

Turn to Q_G . Since we have $Q_G J_G u = u$ for all $u \in LG$, and J_G is an isometry, we only have to show that Q_G is contractive.

Take $\bar{u} \in \mathbb{L}$ and observe that $J_G Q_G \bar{u} = (JQ) \cdot \bar{u}$. Therefore $||Q_G \bar{u}|| = ||J_G Q_G \bar{u}|| \le ||JQ|| ||\bar{u}|| = ||\bar{u}||$. \triangleright

The proof of Theorem 4.1 is concluded. $\triangleright \triangleright$

Combining this theorem with Proposition 3.11, we obtain our main theorem.

5. *p*-convex tensor product as a functor

Now let us do some final observation. Recall an important notion in geometry of normed spaces. Suppose that we assign to every pair, say E, F, of normed spaces the space $E \otimes F$ endowed with some norm. The most interesting are assignments, satisfying the so-called *metric mapping property* [8, §12] (see also "uniform crossnorms" in [20, §6.1]): for every bounded operators $\varphi : E_1 \to F_1, \psi : E_2 \to F_2$ we have $\|\varphi \otimes \psi\| \leq \|\varphi\| \|\psi\|$. (In other terms, such an assignment, extended to bounded operators, is a bifunctor on the category of normed spaces and contractive operators.) We shall show that the *p*-convex tensor product has a natural analogue of that "functorial property" for near-*L*-spaces. As usual, $L := L_p(X)$, and the only condition on *X* is that *L* is infinite-dimensional and separable.

Proposition 5.1. Let $\varphi : E_1 \to E_2$, $\psi : F_1 \to F_2$ be L-bounded operators between near-L-spaces. Then the operator $\varphi \otimes \psi : E_1 \otimes_{pL} E_2 \to F_1 \otimes_{pL} F_2$ is Lbounded, and we have $\|(\varphi \otimes \psi)_{\infty}\| \leq \|\varphi_{\infty}\| \|\psi_{\infty}\|$.

⊲ We first assume that X is convenient. Then, taking $U \in L(E_1 \otimes E_2)$, we have the right to present it as $a \cdot \sum_k I_k \cdot u_k \diamondsuit v_k$; $a \in \mathcal{B}(L), u_k \in LE_1, v_k \in LF_1$ with mutually disjoint proper isometries I_k (see Proposition 3.4). Then, since amplifications of our operators are morphisms of left $\mathcal{B}(L)$ -modules, we have $(\varphi \otimes \psi)_{\infty}(U) = a \cdot \sum_k I_k \cdot (\varphi \otimes \psi)_{\infty}(u_k \diamondsuit v_k)$. By the formula $(\varphi \otimes \psi)_{\infty}(u \diamondsuit v) =$ $\varphi_{\infty}(u) \diamondsuit \psi_{\infty}(v)$, easily verified on elementary tensors, the latter expression is $a \cdot \sum_k I_k \cdot (\varphi_{\infty}(u_k) \diamondsuit \psi_{\infty}(v_k)$. Therefore, by definition of the norm on $F_1 \otimes_{pL} F_2$ for convenient X, we have

$$\|(\varphi \otimes \psi)_{\infty}(U)\| \le \|a\| \Big(\sum_{k} \|\varphi_{\infty}(u_{k})\|^{p} \|\psi_{\infty}(v_{k})\|^{p}\Big)^{\frac{1}{p}} \le \|a\| \Big(\sum_{k} (\|\varphi_{\infty}\|\|u_{k}\|)^{p} \|(\|\psi_{\infty}\|\|v_{k}\|)^{p}\Big)^{\frac{1}{p}} \le \|a\| \|\varphi_{\infty}\| \|\psi_{\infty}\| \Big(\sum_{k} \|u_{k}\|)^{p} \|v_{k}\|^{p}\Big)^{\frac{1}{p}}.$$

It remains to take the respective infimum over all representations of U in the prescribed form.

Turn to an arbitrary X. Using the standard extension of given L-norms, consider our four spaces as near-L-spaces (see Proposition 3.7). Denote our given operators as acting between near-L-spaces as $\tilde{\varphi}$ and $\tilde{\psi}$, respectively. Thus, since $\mathbb{L} = L_p(\mathbb{N}X)$, and $\mathbb{N}X$ is convenient, we have $\|(\tilde{\varphi} \otimes \tilde{\psi})_{\infty}\| \leq \|\tilde{\varphi}_{\infty}\| \|\tilde{\psi}_{\infty}\|$. But, using the definition of near-L-norms on our four spaces, we easily obtain that $\|\tilde{\varphi}_{\infty}\| = \|\varphi_{\infty}\|$ and $\|\tilde{\psi}_{\infty}\| = \|\psi_{\infty}\|$. At the same time for $U \in L(E_1 \otimes_{pl} E_2)$ we have

$$\|(\varphi \otimes \psi)_{\infty}(U)\| = \|J_{F_1 \otimes F_2} \Big(\varphi \otimes \psi)_{\infty}(U)\Big)\| = \|\Big((\varphi \otimes \psi)_{\infty}(U), 0, 0, ...\Big)\| = \|(\widetilde{\varphi} \otimes \widetilde{\psi})_{\infty}(U, 0, 0, ...)\| \le \|(\widetilde{\varphi} \otimes \widetilde{\psi})_{\infty}\|\|(U, 0, 0, ...)\| = \|(\widetilde{\varphi} \otimes \widetilde{\psi})_{\infty}\|\|U\|,$$

and consequently $\|(\varphi \otimes \psi)_{\infty}\| \leq \|(\widetilde{\varphi} \otimes \widetilde{\psi})_{\infty}\|$. The desired estimate immediately follows. \triangleright

References

- D. P. Blecher, V. I. Paulsen, "Tensor products of operator spaces", J. Funct. Anal., No. 2, 99 (1991), 262-292.
- [2] V. Bogachev, *Measure Theory*, v.1,2 (Springer, New York, 2007).
- [3] H. G. Dales, M. E. Polyakov, "Multi-normed spaces", Dissertationes Math. (Rozprawy Mat.), 488 (2012), 165 pp.
- [4] H. G. Dales, M. Daws, H. L. Pham, P. Ramsden, "Multi-norms and the injectivity of $L^p(G)$ ", J. London Math. Soc., (2), 86 (2012), 779-809.
- [5] H. G. Dales, M. Daws, H. L. Pham, P. Ramsden, "Equivalence of multinorms", Dissertationes Math. (Rozprawy Mat.), 498 (2014), 53 pp.

- [6] H. G. Dales, N. J. Laustsen, T. Oikhberg, V. G. Troitsky, Multi-norms and Banach lattices. Dissertationes Math. (Rozprawy Mat.), 524 (2017), 115 pp.
- [7] M. Daws, "p-operator spaces and Figá-Talamanca-Herz algebras", J. Oper. Theory, 63 (2010), 47-83.
- [8] A. Defant, K. Floret, Tensor norms and Operator Ideals, North-Holland, Amsterdam, 1993.
- [9] E. G. Effros, "Advances in quantized functional analysis", Proc. ICM Berkeley, 1986.
- [10] E. G. Effros, Z.-J. Ruan, Operator Spaces, Clarendon Press, Oxford, 2000.
- [11] E. G. Effros, Z.-J. Ruan, "A new approach to operator spaces", Canad. Math. Bull., 34 (1991), 329-337.
- [12] A. Ya. Helemskii, *Quantum Functional Analysis*, American Mathematical Society, Providence, R.I., 2010).
- [13] A. Ya. Helemskii, "Multi-normed spaces, based on non-discrete measures, and their tensor products", Izvestiya RAN: Ser. Mat, 82:2, (2018), 23–25.
- [14] H. E. Lacey, The Isometric Theory of Classical Banach Spaces, Springer-Verlag, Berlin, 1974.
- [15] A. Lambert, *Operatorfolgenräume*, Dissertation, Saarbrüken, 2002.
- [16] C. Le Merdy, "Factorization of p-completely bounded multilinear maps", Pacific J. Math., 172 (1966), 187-213.
- [17] N. Oikhberg, "Discretization in L_p -normed spaces", Preprint, 2019.
- [18] V. I. Paulsen, Completely Bounded Maps and Operator Algebras, Cam. Univ. Press, Cambridge, 2002. 2004.
- [19] J. Pisier, "Completely bounded maps between sets of Banach space operators", Indiana Univ. Math. J. 39 (1990), 249-277.
- [20] R. A. Ryan, Introduction to Tensor Products of Banach Spaces, Springer-Verlag, Berlin, 2002.

[21] P. Wojtaszczyk, Banach spaces for analysts, Cam. Univ. Press, Cambridge, 1991.

A. Ya. Helemskii Faculty of Mechanics and Mathematics Moscow State (Lomonosov) University Moscow 119991 Leninskie Gory E-mail: helemskii@rambler.ru