

Comments on a new ‘full colour’ parton shower

Jack Holguin,^{a,b} Jeffrey R. Forshaw,^{a,b} Simon Plätzer.^{b,c}

^a*Consortium for Fundamental Physics, School of Physics & Astronomy,
University of Manchester, Manchester M13 9PL, United Kingdom*

^b*Erwin Schrödinger Int. Institute for Mathematics and Physics,
University of Vienna, 1090 Wien, Austria*

^c*Particle Physics, Faculty of Physics,
University of Vienna, 1090 Wien, Austria*

E-mail: `jack.holguin@manchester.ac.uk`,
`jeffrey.forshaw@manchester.ac.uk`, `simon.plaetzer@univie.ac.at`

ABSTRACT: A new parton shower algorithm has been presented with the claim of providing soft-gluon resummation at ‘full colour’ [1]. In this paper we show that the algorithm does not succeed in this goal. We show that full colour accuracy requires the Sudakov factors to be defined at amplitude level and that the simple parton-shower unitarity argument employed in [1] is not sufficient.

1 Introduction

Over recent years much attention has been devoted to the development of parton showers with ‘full colour’ evolution [2–7]. The study of these has multiple motivations: most importantly, reducing theoretical uncertainties in parton showers will be crucial for precision phenomenology at future colliders. Currently, parton showers provide some of the largest sources of uncertainty in experimental analyses, e.g. [8]. There has also been a growth in interest towards developing tools for the formal resummation of observables sensitive to the complexity of the non-abelian structure of the strong interaction, specifically observables with non-global or super-leading logarithms [9–14]. These will play an important role in advancing parton shower algorithms. In this context, a widely available ‘full colour’ parton shower would be a powerful tool.

In this letter we comment on the formalism for resumming complex colour structures employed recently in [1]. A similar approach was previously put forward by one of the present authors and collaborators [3, 5]. The authors of [1] describe their formalism as being capable of producing “numerical resummation at full color in the strongly ordered soft gluon limit.” We will examine this claim in what follows.

Let us be clear on what we mean by leading and sub-leading colour. A general observable can be written

$$\Sigma(L) = \sum_{n=0}^{\infty} (N_c \alpha_s)^n \sum_{m=0}^{n+1} C_{n,m}(L) , \quad (1.1)$$

where L is some large logarithm. The coefficients $C_{n,m}$ can be expanded:

$$C_{n,m} = \underbrace{C_{n,m}^{(0)}}_{\text{LC}_{\Sigma}} + \underbrace{\frac{1}{N_c} C_{n,m}^{(1)}}_{\text{NLC}_{\Sigma}} + \underbrace{\frac{1}{N_c^2} C_{n,m}^{(2)}}_{\text{NNLC}_{\Sigma}} + \dots \quad (1.2)$$

and a ‘full colour’ shower should be able to compute all of the $C_{n,m}^{(i)}$ at a stated logarithmic accuracy.¹ We will show that the formalism of [1] generally fails to compute the NNLC_Σ terms, even in the strongly-ordered soft gluon approximation. Note also that, for many observables, the NLC_Σ term vanishes, so that the dominant sub-leading colour corrections occur at NNLC_Σ. It is also important to appreciate that the colour expansion defined in Eq. (1.2) is very weak in its ambition. Just as in the case of logarithmic resummation, more ambitious would be to perform a resummation of towers of enhanced corrections. In which case an expansion of the form of Eq. (1.2) would be exponentiated.

2 Summary of the new ‘full colour’ parton shower

We will briefly summarize the algorithm advocated in [1] and we largely follow their notation. The amplitude for an n -parton hard process is $|M_n\rangle$ and $|m_{n+k}\rangle$ is the amplitude

¹Or in a specified kinematic limit, e.g. the strongly-ordered soft gluon limit.

after dressing with k soft gluons. Real emissions are accounted for recursively according to

$$\langle m_{n+k} | m_{n+k} \rangle = \langle m_{n+k-1} | \mathbf{\Gamma}_{n+k-1}(\mathbf{1}) | m_{n+k-1} \rangle = \langle M_n | \mathbf{\Gamma}_n(\dots \mathbf{\Gamma}_{n+k-2}(\mathbf{\Gamma}_{n+k-1}(\mathbf{1})) \dots) | M_n \rangle, \quad (2.1)$$

where

$$\mathbf{\Gamma}_n(\mathbf{\Gamma}) = - \sum_{\substack{i,j=1 \\ i \neq j}}^n \mathbf{T}_i \mathbf{\Gamma} \mathbf{T}_j \omega_{ij}, \quad \omega_{ij} = \frac{s_{ij}}{s_{iq}s_{qj}} \quad (2.2)$$

and $s_{ij} = 2p_i \cdot p_j$ in terms of the momenta of the partons i and j . The radiation pattern for a single emission, q , is then determined by

$$\frac{d\sigma_{n+k+1}}{\sigma_{n+k}} = d\Phi_{+1} 8\pi\alpha_s \frac{\langle m_{n+k} | \mathbf{\Gamma}_{n+k}(\mathbf{1}) | m_{n+k} \rangle}{\langle m_{n+k} | m_{n+k} \rangle}, \quad (2.3)$$

where $d\Phi_{+1}$ is a phase-space measure and parametrises the momentum map from a state of $n+k$ partons to a state of $n+k+1$ partons. Its details are not needed for our discussion. Virtual corrections are encoded via a no-emission probability, i.e. via a typical parton-shower cross-section-level Sudakov factor, defined though unitarity as

$$\int_{t'}^t \frac{d\kappa_{ij}^2}{\sigma_{n+k}} \int \frac{d\sigma_{n+k+1}}{d\kappa_{ij}^2} \Pi(\kappa_{ij}^2, t) = 1 - \Pi(t', t), \quad (2.4)$$

where $\kappa_{ij}^2 = \omega_{ij}^{-1}$ plays the role of the ordering variable. This equation has the solution

$$\Pi^{(k)}(t', t) = \prod_{\substack{i,j=1 \\ i \neq j}}^{n+k} \Pi_{ij}(t', t), \quad (2.5)$$

where

$$\Pi_{ij}(t', t) = \exp \left(- \int_{t'}^t \frac{d\kappa_{ij}^2}{\kappa_{ij}^2} \int \frac{8\pi d\Phi_{+1}}{d\kappa_{ij}^2} \alpha_s \frac{\langle m_{n+k} | \mathbf{T}_i \mathbf{T}_j | m_{n+k} \rangle}{\langle m_{n+k} | m_{n+k} \rangle} \right), \quad (2.6)$$

is the no-emission probability for a single dipole (i, j) . The overall no-emission probability dresses the real emission matrix elements defined in Eq. (2.1) according to

$$\langle m_{n+k}; t | m_{n+k}; t \rangle = \Pi^{(k)}(t, t_k) \dots \Pi^{(1)}(t_2, t_1) \Pi^{(0)}(t_1, Q^2) \langle m_{n+k} | m_{n+k} \rangle, \quad (2.7)$$

where t_i is the ordering variable associated with the i th emission and Q^2 is the hard scale.

3 The problem with Sudakovs

In this section we show that defining Sudakov factors through cross-section-level unitarity gives rise to two compounding errors in colour. The first error is in the computation of loops, the second is in the computation of the interplay between loops and real emissions. These errors make the inclusion of Coulomb terms impossible, since they always appear as a pure

(abelian) phase in the amplitude. Firstly, we address the computation of loops (resummed into Sudakov factors). The role of Sudakov factors in full-colour evolution of amplitudes has been extensively studied [4, 6, 7, 13–17]. Ignoring Coulomb terms (including them only makes matters more complicated), Sudakov factors² should dress a general amplitude as

$$\begin{aligned}
& \langle m_{n+k}; t' | m_{n+k}; t' \rangle \\
&= \langle m_{n+k}; t | e^{-\int_{t'}^t d\kappa^2 \int \frac{4\pi d\Phi+1}{d\kappa^2} \alpha_s \Gamma_{n+k}(\mathbf{1})} e^{-\int_{t'}^t d\kappa^2 \int \frac{4\pi d\Phi+1}{d\kappa^2} \alpha_s \Gamma_{n+k}^\dagger(\mathbf{1})} | m_{n+k}; t \rangle, \\
&= \frac{\langle m_{n+k}; t | e^{-\int_{t'}^t d\kappa^2 \int \frac{8\pi d\Phi+1}{d\kappa^2} \alpha_s \Gamma_{n+k}(\mathbf{1})} | m_{n+k}; t \rangle}{\langle m_{n+k}; t | m_{n+k}; t \rangle} \langle m_{n+k}; t | m_{n+k}; t \rangle, \\
&\neq \Pi^{(k)}(t', t) \langle m_{n+k}; t | m_{n+k}; t \rangle.
\end{aligned} \tag{3.1}$$

The not equals to sign represents the first error in [1].

We will now attempt to explicate this error and its consequences by giving it two different interpretations. Firstly, we will show how this error can be thought of as a straightforward linear algebra error. Secondly, we will present some fixed-order calculations that show this error corresponds to miscalculating NNLC_Σ diagrams with two or more loops. To begin the linear algebra interpretation, let us rewrite the pertinent term from Eq. (3.1) as

$$\begin{aligned}
& \frac{\langle m_{n+k}; t | e^{-\int_{t'}^t d\kappa^2 \int \frac{8\pi d\Phi+1}{d\kappa^2} \alpha_s \Gamma_{n+k}(\mathbf{1})} | m_{n+k}; t \rangle}{\langle m_{n+k}; t | m_{n+k}; t \rangle} \\
&= \frac{\text{Tr}(|m_{n+k}; t\rangle \langle m_{n+k}; t| e^{\mathbf{V}})}{\text{Tr}(|m_{n+k}; t\rangle \langle m_{n+k}; t|)} \equiv \text{Tr}_{\text{norm}}(e^{\mathbf{V}}),
\end{aligned} \tag{3.2}$$

where Tr_{norm} is a normalised trace, such that $\text{Tr}_{\text{norm}} \mathbf{1} = 1 \neq N$ where N is the dimension of the matrix. In this notation we can write

$$\Pi^{(k)}(t', t) = e^{\text{Tr}_{\text{norm}}(\mathbf{V})}. \tag{3.3}$$

This definition is the source of the error. Motivated by cross-section-level arguments of unitarity, it is implicitly assumed that

$$\text{Tr}_{\text{norm}}(e^{\mathbf{V}}) = e^{\text{Tr}_{\text{norm}}(\mathbf{V})}, \tag{3.4}$$

which is wrong.

As a trivial example of how this sort of error could give problems, consider $\text{Tr} e^{\mathbf{1}_N} = Ne$ whereas $e^{\text{Tr} \mathbf{1}_N} = e^N$. However, the error from using a normalised trace is more subtle, since $\text{Tr}_{\text{norm}} e^{\mathbf{1}_N} = e^{\text{Tr}_{\text{norm}} \mathbf{1}_N} = e$. To see where the actual problem arises, consider a toy model where $\mathbf{V} = \alpha_s N_c (\mathbf{1} + N_c^{-1} \delta \mathbf{V})$ and $\delta \mathbf{V}$ is not diagonal. In this case, the $\alpha_s N_c \mathbf{1}$ piece

²The argument of the Sudakov exponent is the real part of the one-loop cusp anomalous dimension [13, 14]. Depending on the choice of ordering variable, path ordering should be implied. See Section 2 of [6] for more details.

plays the role of the leading colour part of the Sudakov and $\alpha_s \delta \mathbf{V}$ the sub-leading colour part. The result is that

$$\text{Tr}_{\text{norm}}(e^{\mathbf{V}}) = e^{\text{Tr}_{\text{norm}}(\mathbf{V})} + \sum_{n \geq 2} \mathcal{O}(\alpha_s^n N_c^{n-2} (\text{Tr}_{\text{norm}} \delta \mathbf{V}^2 - (\text{Tr}_{\text{norm}} \delta \mathbf{V})^2)). \quad (3.5)$$

The important difference arises because $(\text{Tr}_{\text{norm}} \delta \mathbf{V})^n \neq \text{Tr}_{\text{norm}}(\delta \mathbf{V}^n)$ for $n \geq 2$. From this argument it is clear that errors will occur, starting with the computation of NNLC_Σ .

Now let us now give a physical interpretation of the error by expanding Eq. (3.2) to $\mathcal{O}(\alpha_s^2)$. The $\mathcal{O}(\alpha_s^2)$ term corresponds to dressing a general hard process at fixed order with two strongly ordered soft loops. The correct amplitude is

$$\begin{aligned} & \sum_{\substack{i,j=1 \\ i \neq j}}^n \int_{t'}^t d\kappa_{ij}^2 \int \frac{8\pi d\Phi_{+1}}{d\kappa_{ij}^2} \alpha_s \sum_{\substack{k,l=1 \\ k \neq l}}^n \int_{\kappa_{ij}^2}^t d\kappa_{kl}^2 \int \frac{8\pi d\Phi_{+1}}{d\kappa_{kl}^2} \alpha_s \\ & \times \text{Tr}_{\text{norm}}(\mathbf{T}_i \cdot \mathbf{T}_j \mathbf{T}_k \cdot \mathbf{T}_l) \langle m_{n+k}; t | m_{n+k}; t \rangle. \end{aligned} \quad (3.6)$$

Now, we can expand $\Pi^{(k)}(t', t) \langle m_{n+k}; t | m_{n+k}; t \rangle$ to the same order. We find

$$\begin{aligned} & \frac{1}{2} \sum_{\substack{i,j=1 \\ i \neq j}}^n \int_{t'}^t d\kappa_{ij}^2 \int \frac{8\pi d\Phi_{+1}}{d\kappa_{ij}^2} \alpha_s \text{Tr}_{\text{norm}}(\mathbf{T}_i \cdot \mathbf{T}_j) \sum_{\substack{k,l=1 \\ k \neq l}}^n \int_{t'}^t d\kappa_{kl}^2 \int \frac{8\pi d\Phi_{+1}}{d\kappa_{kl}^2} \alpha_s \\ & \times \text{Tr}_{\text{norm}}(\mathbf{T}_k \cdot \mathbf{T}_l) \langle m_{n+k}; t | m_{n+k}; t \rangle. \end{aligned} \quad (3.7)$$

These two expressions are only equal when $n+k \leq 3$ because the colour matrices are then proportional to identity matrices. However, for multiplicities of coloured partons greater than 3 they differ by NNLC_Σ pieces. This error occurs because writing a matrix element in the form of Eq. (3.7) implicitly assumes that $[\mathbf{T}_i \cdot \mathbf{T}_j, \mathbf{T}_i \cdot \mathbf{T}_k] \approx 0$, which is only correct up to NLC_Σ terms. For example, consider the case of $e^+e^- \rightarrow q\bar{q}g_1g_2$ (for which the NLC_Σ term is zero). To illustrate the point consider the limit that both gluons were emitted from the quark. In this limit a NNLC_Σ error emerges due to the non-vanishing of

$$\begin{aligned} & \alpha_s^2 \text{Tr}_{\text{norm}}(\mathbf{T}_q \cdot \mathbf{T}_{g_1} \mathbf{T}_{g_1} \cdot \mathbf{T}_{g_2}) - \alpha_s^2 \text{Tr}_{\text{norm}}(\mathbf{T}_q \cdot \mathbf{T}_{g_1}) \text{Tr}_{\text{norm}}(\mathbf{T}_{g_1} \cdot \mathbf{T}_{g_2}) \\ & = \alpha_s^2 \frac{N_c^6 + 3N_c^4 - 14N_c^2 + 2}{4N_c^2(N_c^2 - 1)^2} = \frac{(N_c \alpha_s)^2}{4} \left(\frac{1}{N_c^2} + \frac{5}{N_c^4} + \dots \right). \end{aligned} \quad (3.8)$$

Similar errors arise from other emission topologies. The non-vanishing commutator is also the reason why Coulomb terms do not cancel and, as a result, underpins the origin of super-leading logarithms [9].

The second error compounds the first. Let us now consider the evolution of an amplitude to a new scale whilst emitting a single gluon:

$$\begin{aligned} & \langle m_{n+k+1}; t'' | m_{n+k+1}; t'' \rangle \\ & = \int_{t''}^{t'} d\kappa^2 \int \frac{8\pi d\Phi_{+1}}{d\kappa^2} \alpha_s \langle m_{n+k}; t | e^{-\int_{t'}^t d\kappa^2 \int \frac{4\pi d\Phi_{+1}}{d\kappa^2}} \alpha_s \Gamma_{n+k}(\mathbf{1}) \Gamma_{n+k}(\mathbf{1}) \\ & \times e^{-\int_{t'}^t d\kappa^2 \int \frac{4\pi d\Phi_{+1}}{d\kappa^2}} \alpha_s \Gamma_{n+k}^\dagger(\mathbf{1}) | m_{n+k}; t \rangle. \end{aligned} \quad (3.9)$$

In order to recombine the two exponentials into a single Sudakov that builds $\Pi^{(k)}(t', t)$ one must assume $[\Gamma_{n+k}(\mathbf{1}), e^{\mathbf{V}}] \approx 0$. For the same reasons as those described above, this is again a NNLC $_{\Sigma}$ error. Where the previous error was in the higher order colour of loop diagrams (≥ 2 loops), this error is in the higher order colour from the interplay between (≥ 1) loops and emissions. Consequently, the algorithm does correctly generate real emissions in the absence of any loop corrections. It also correctly generates one-loop contributions that dress the softest real emission but fails thereafter.

4 Conclusions

QCD colour dynamics beyond leading colour is highly non-trivial and its correct inclusion generally requires an amplitude-level approach that goes beyond the simple treatment of virtual corrections presented in [1].

Acknowledgements

The authors want to thank the Erwin Schrödinger Institute for support during the period when this and related work has been carried out. This work has received funding from the UK Science and Technology Facilities Council (grant no. ST/P000800/1), the European Union’s Horizon 2020 research and innovation programme as part of the Marie Skłodowska-Curie Innovative Training Network MCnetITN3 (grant agreement no. 722104), and in part by the by the COST actions CA16201 “PARTICLEFACE” and CA16108 “VBSCAN”. JH thanks the UK Science and Technology Facilities Council for the award of a studentship.

References

- [1] S. Hoeche and D. Reichelt, *Numerical resummation at full color in the strongly ordered soft gluon limit*, 2001.11492.
- [2] Z. Nagy and D. E. Soper, *Parton shower evolution with subleading color*, *JHEP* **06** (2012) 044, [1202.4496].
- [3] S. Plätzer and M. Sjödahl, *Subleading N_c improved Parton Showers*, *JHEP* **07** (2012) 042, [1201.0260].
- [4] R. Ángeles Martínez, M. De Angelis, J. R. Forshaw, S. Plätzer and M. H. Seymour, *Soft gluon evolution and non-global logarithms*, *JHEP* **05** (2018) 044, [1802.08531].
- [5] S. Plätzer, M. Sjödahl and J. Thorén, *Color matrix element corrections for parton showers*, *JHEP* **11** (2018) 009, [1808.00332].
- [6] J. R. Forshaw, J. Holguin and S. Plätzer, *Parton branching at amplitude level*, *JHEP* **08** (2019) 145, [1905.08686].
- [7] Z. Nagy and D. E. Soper, *Parton showers with more exact color evolution*, *Phys.Rev.D* **99** (2019) 054009, [1902.02105].
- [8] P. Azzi et al., *Report from Working Group 1, CERN Yellow Rep. Monogr.* **7** (2019) 1–220, [1902.04070].

- [9] J. R. Forshaw, A. Kyrieleis and M. H. Seymour, *Super-leading logarithms in non-global observables in QCD: Colour basis independent calculation*, *JHEP* **09** (2008) 128, [[0808.1269](#)].
- [10] A. Banfi, G. Corcella and M. Dasgupta, *Angular ordering and parton showers for non-global QCD observables*, *JHEP* **03** (2007) 050, [[hep-ph/0612282](#)].
- [11] J. R. Forshaw, M. H. Seymour and A. Siódmok, *On the Breaking of Collinear Factorization in QCD*, *JHEP* **11** (2012) 066, [[1206.6363](#)].
- [12] Y. Hagiwara, Y. Hatta and T. Ueda, *Hemisphere jet mass distribution at finite N_c* , *Phys. Lett.* **B756** (2016) 254–258, [[1507.07641](#)].
- [13] S. Caron-Huot, *Resummation of non-global logarithms and the BFKL equation*, *JHEP* **03** (2018) 036, [[1501.03754](#)].
- [14] T. Becher, M. Neubert, L. Rothen and D. Y. Shao, *Factorization and Resummation for Jet Processes*, *JHEP* **11** (2016) 019, [[1605.02737](#)].
- [15] Z. Nagy and D. E. Soper, *Parton showers with quantum interference: Leading color, with spin*, *JHEP* **07** (2008) 025, [[0805.0216](#)].
- [16] S. Plätzer, *Summing Large- N Towers in Colour Flow Evolution*, *Eur. Phys. J.* **C74** (2014) 2907, [[1312.2448](#)].
- [17] Z. Nagy and D. E. Soper, *Effects of subleading color in a parton shower*, *JHEP* **07** (2015) 119, [[1501.00778](#)].