

UNIFORM ELLIPTICITY AND p - q GROWTH

CRISTIANA DE FILIPPIS AND FRANCESCO LEONETTI

ABSTRACT. Fix any two numbers p and q , with $1 < p < q$; we give an example of an integral functional enjoying uniform ellipticity and p - q growth.

CONTENTS

1. Introduction	1
2. Example	3
3. Preliminary results	4
4. Proof of Theorem 2.1	5
5. Another example	8
References	9

1. INTRODUCTION

We consider integral functionals

$$(1.1) \quad \int_{\Omega} f(Du(x))dx,$$

where $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}^N$, Ω is bounded and open and f is continuous and nonnegative. About f we assume p - q growth

$$(1.2) \quad c_1|z|^p - c_2 \leq f(z) \leq c_3|z|^q + c_4,$$

where c_1, c_2, c_3, c_4, p, q are constants with $c_1, c_3 \in (0, +\infty)$, $c_2, c_4 \in [0, +\infty)$ and $1 < p < q$. In this framework it is usual to assume that

$$(1.3) \quad c_5(\mu + |z|)^{p-2} \leq \langle DDf(z) \frac{\lambda}{|\lambda|}, \frac{\lambda}{|\lambda|} \rangle,$$

and

$$(1.4) \quad |DDf(z)| \leq c_6(\mu + |z|)^{q-2},$$

where c_5, c_6, μ are constants with $c_5, c_6 \in (0, +\infty)$ and $\mu \in [0, 1]$. Retaining only the informations about the growth in the large of the second derivative, as prescribed by (1.3)-(1.4), leads to the following bound on the ratio between the highest and the lower eigenvalue of DDf :

$$(1.5) \quad \mathcal{R}(z) := \frac{\text{highest eigenvalue of } DDf(z)}{\text{lowest eigenvalue of } DDf(z)} \leq c_7(\mu + |z|)^{q-p},$$

2010 *Mathematics Subject Classification.* 35J60, 35J70.

Key words and phrases. Regularity, uniform ellipticity, p - q -growth.

Acknowledgements. C. De Filippis is supported by the Engineering and Physical Sciences Research Council (EPSRC): CDT Grant Ref. EP/L015811/1. F. Leonetti is supported by MIUR, GNAMPA, INdAM, UNIVAQ. .

for some positive constant c_7 . The right hand side of (1.5), evidently blows up as $|z| \rightarrow \infty$, given that, in general, $q > p$. On the other hand, if by any chance the integrand f features certain structural properties which make $\mathcal{R}(z)$ bounded from above by a constant non-depending, in particular, from z , then we have uniform ellipticity. We are concerned with regularity of minimizers $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}^N$ of (1.1); in this framework of p - q growth, the following bound sometimes appears

$$(1.6) \quad q < p + c(n, p),$$

where $c(n, p)$ is positive and tends to 0 when the dimension n tends to $+\infty$; see [2, 6, 9–11, 13, 16, 18] and [17, Section 6]; see also [7, Section 6.2] where a simple argument is given. Now we assume the following structure condition

$$(1.7) \quad f(z) = g(|z|),$$

with $g : [0, +\infty) \rightarrow [0, +\infty)$. Some papers require $g(0) = 0$, $g \in C^2((0, +\infty)) \cap C^1([0, +\infty))$ with $g'(t) > 0$ for $t > 0$; moreover, [3, 5, 8, 15] ask for

$$(1.8) \quad 0 < m \leq \frac{g''(t)t}{g'(t)} \leq M < +\infty \quad \forall t > 0.$$

Note that [1] requires (1.8) with $1 \leq m$; on the other hand, [14] asks for $M \leq 1$. In [4] they consider splitting densities $f(Du) = a(|(D_1u, \dots, D_{n-1}u)|) + b(|D_nu|)$ and they require (1.8) for both a and b . We remark that $g' > 0$ and (1.8) forces $g'' > 0$, so g must be strictly convex; on the other hand, (1.8) allows p - q growth whatever p and q are: in this paper we fix p and q with $1 < p < q$, no matter how far they are, and we show a convex function g verifying (1.8), with p - q growth. In [3] we find Theorem 1.15 that says

Theorem 1.1. *We assume that $g(0) = 0$ and $g \in C^2((0, +\infty)) \cap C^1([0, +\infty))$; moreover, $g'(t) > 0$ for $t > 0$ and (1.8) holds true. If $u \in W_{loc}^{1,1}(\Omega, \mathbb{R}^N)$ is a local minimizer of (1.1) under the structure condition (1.7) with g as before, then u is locally Lipschitz continuous in Ω .*

We are going to show an example for the previous Theorem 1.1: fix p and q with $1 < p < q$, then we give g satisfying all the assumptions of Theorem 1.1 with the chosen p and q : the restriction (1.6) does not apply! Moreover, such a g gives an f for which we have uniform ellipticity; indeed, let g be any function in $C^2((0, +\infty))$ with $g'(t) > 0$ for $t > 0$, satisfying assumption (1.8); then, for the corresponding f given by (1.7), we have

$$\frac{\partial f}{\partial z_i^\alpha}(z) = g'(|z|) \frac{z_i^\alpha}{|z|}$$

and

$$\frac{\partial^2 f}{\partial z_i^\alpha \partial z_j^\beta}(z) = \left[g''(|z|) - \frac{g'(|z|)}{|z|} \right] \frac{z_i^\alpha z_j^\beta}{|z|^2} + \frac{g'(|z|)}{|z|} \delta^{\alpha\beta} \delta_{ij},$$

so that

$$(1.9) \quad \left\langle DDf(z) \frac{\lambda}{|\lambda|}, \frac{\lambda}{|\lambda|} \right\rangle = \left[g''(|z|) - \frac{g'(|z|)}{|z|} \right] \left\langle \frac{z}{|z|}, \frac{\lambda}{|\lambda|} \right\rangle^2 + \frac{g'(|z|)}{|z|};$$

if we consider first the case $[\dots] \geq 0$ and then the other case $[\dots] < 0$, using (1.8), we get

$$(1.10) \quad \frac{\text{highest eigenvalue of } DDf(z)}{\text{lowest eigenvalue of } DDf(z)} \leq \max \left\{ M; \frac{1}{m} \right\};$$

so, we are in the uniform ellipticity regime. So, after fixing p and q at will in $(1, +\infty)$, we are going to write an example of functional with p - q growth and uniform ellipticity. For $1 < p < q$, set $a = \frac{p+q}{2}$ and $b = \frac{q-p}{2}$. Then, we have $a, b > 0$, $1 < p = a - b < a + b = q$ and we can use the function g defined in the next section 2.

2. EXAMPLE

We fix $a, b \in (0, +\infty)$ with

$$(2.1) \quad 1 < a - b.$$

We consider $g : [0, +\infty) \rightarrow [0, +\infty)$ such that

$$(2.2) \quad g(t) = t^{a+b \sin(\varphi(t))},$$

where $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is given by

$$(2.3) \quad \varphi(t) = \begin{cases} \frac{3}{2}\pi & \text{if } t \in (-\infty, 1], \\ \frac{3}{2}\pi + \varepsilon \ln \ln(e + (t-1)^4) & \text{if } t \in (1, +\infty), \end{cases}$$

with $\varepsilon > 0$. Note that

$$(2.4) \quad \varphi'(t) = \begin{cases} 0 & \text{if } t \in (-\infty, 1], \\ \frac{\varepsilon}{\ln(e+(t-1)^4)} \frac{4(t-1)^3}{e+(t-1)^4} & \text{if } t \in (1, +\infty) \end{cases}$$

and

$$(2.5) \quad \varphi''(t) = \begin{cases} 0 & \text{if } t \in (-\infty, 1], \\ \varepsilon \left\{ \frac{-1}{[\ln(e+(t-1)^4)]^2} \left[\frac{4(t-1)^3}{e+(t-1)^4} \right]^2 + \frac{1}{\ln(e+(t-1)^4)} \frac{12(t-1)^2 e - 4(t-1)^6}{[e+(t-1)^4]^2} \right\} & \text{if } t \in (1, +\infty) \end{cases}$$

so that $\varphi \in C^2(\mathbb{R})$. Note that $\varphi'(t) > 0$ when $t > 1$; moreover, $\lim_{t \rightarrow +\infty} \varphi(t) = +\infty$. Then $\varphi(t)$ increases and takes all the values of the interval $[\frac{3}{2}\pi, +\infty)$. This means that, in (2.2), the exponent $a + b \sin(\varphi(t))$ oscillates between $a - b$ and $a + b$ infinitely many times as t goes from 0 to $+\infty$; then $g(t)$ has $a - b$ growth from below and $a + b$ growth from above. As far as ε is concerned, we require that

$$(2.6) \quad 0 < \varepsilon < \min \left\{ 1; \frac{a-1-b}{224b} \right\}.$$

We are going to prove the next

Theorem 2.1. *Let us consider $a, b \in (0, +\infty)$ verifying (2.1); we take $g(t)$ given by (2.2) where φ is defined in (2.3) and ε satisfies (2.6). Then $g : [0, +\infty) \rightarrow [0, +\infty)$, $g(0) = 0$, $g(t) > 0$ for $t > 0$, $g \in C^1([0, +\infty)) \cap C^2((0, +\infty))$, $\lim_{t \rightarrow 0^+} \frac{g(t)}{t} = 0$, $\lim_{t \rightarrow +\infty} \frac{g(t)}{t} = +\infty$, $g'(0) = 0$ and*

$$(2.7) \quad 0 < \{-b8\varepsilon + a - b\} \frac{g(t)}{t} \leq g'(t) \leq \{b8\varepsilon + a + b\} \frac{g(t)}{t}$$

for every $t > 0$; moreover,

$$(2.8) \quad 0 < \{-b8\varepsilon + a - b\} t^{a-b-1} \leq g'(t) \leq \{b8\varepsilon + a + b\} [t^{a-b-1} + t^{a+b-1}]$$

for every $t > 0$. As far as g'' is concerned, we get

$$(2.9) \quad 0 < \{-224b\varepsilon + a - 1 - b\} \frac{g'(t)}{t} \leq g''(t) \leq \{224b\varepsilon + a - 1 + b\} \frac{g'(t)}{t}$$

for every $t > 0$, thus g is strictly convex in $[0, +\infty)$.

The present example is a modification of the one given in [12, 19]; in the present example the small new parameter ε appears and it makes possible to get convexity and p - q growth with any p and q .

3. PRELIMINARY RESULTS

We need some preliminary estimates.

Lemma 3.1. *For all $t \in (1, \infty)$ there holds that:*

$$(3.1) \quad 0 < \frac{(t-1)^3}{e + (t-1)^4} < 1.$$

Proof. If $1 < t \leq 2$, then $0 < t-1 \leq 1$ so that

$$0 < \frac{(t-1)^3}{e + (t-1)^4} \leq \frac{1}{e + (t-1)^4} \leq \frac{1}{e} < 1.$$

If $2 < t$, then $1 < t-1$ so that

$$0 < \frac{(t-1)^3}{e + (t-1)^4} < \frac{(t-1)^4}{e + (t-1)^4} < 1.$$

The two cases give (3.1). □

Lemma 3.2. *For all $t \in (1, \infty)$ there holds that:*

$$(3.2) \quad 0 < \frac{(t-1)^3 t}{e + (t-1)^4} < 2.$$

Proof. We write $t = (t-1) + 1$ and we get

$$0 < \frac{(t-1)^3 t}{e + (t-1)^4} = \frac{(t-1)^3(t-1)}{e + (t-1)^4} + \frac{(t-1)^3}{e + (t-1)^4} < \frac{(t-1)^4}{e + (t-1)^4} + 1 < 1 + 1,$$

where we used (3.1). □

Lemma 3.3. *For all $t \in (1, \infty)$ there holds that:*

$$(3.3) \quad 0 < \frac{(t-1)^2 t^2}{e + (t-1)^4} < 4.$$

Proof. If $1 < t < 2$, then $0 < t-1 < 1$ so that

$$0 < \frac{(t-1)^2 t^2}{e + (t-1)^4} < \frac{4}{e + (t-1)^4} < \frac{4}{e} < \frac{4}{2} = 2.$$

If $2 \leq t$, then $t \leq 2(t-1)$ so that

$$0 < \frac{(t-1)^2 t^2}{e + (t-1)^4} \leq \frac{(t-1)^2 4(t-1)^2}{e + (t-1)^4} = \frac{4(t-1)^4}{e + (t-1)^4} < 4.$$

The two cases give (3.3). □

Lemma 3.4. *For all $t \in (1, \infty)$ there holds that:*

$$(3.4) \quad 0 < \frac{\ln t}{\ln(e + (t-1)^4)} < 1.$$

Proof. If $1 < t \leq e$, then

$$0 < \frac{\ln t}{\ln(e + (t-1)^4)} \leq \frac{\ln e}{\ln(e + (t-1)^4)} < \frac{\ln e}{\ln e} = 1.$$

If $e < t$, then $t < (t-1)^2$: indeed, this last inequality is equivalent to $0 < t^2 - 3t + 1$; the two solutions of the equation $t^2 - 3t + 1 = 0$ are $\frac{3-\sqrt{5}}{2}$ and $\frac{3+\sqrt{5}}{2}$; note that $5 < 5, 29 = (2, 3)^2$, so that $\frac{3+\sqrt{5}}{2} < \frac{3+2,3}{2} = 2,65 < e$; then $e < t$ implies $0 < t^2 - 3t + 1$ and $t < (t-1)^2$. This last inequality allows us to write

$$0 < \frac{\ln t}{\ln(e + (t-1)^4)} < \frac{\ln((t-1)^2)}{\ln(e + (t-1)^4)} < \frac{\ln((t-1)^2)}{\ln((t-1)^4)} = \frac{2}{4}.$$

The two cases give (3.4). □

In this section 3, φ is given by (2.3) with any $\varepsilon > 0$: in the forthcoming lemmas, no restriction from above on ε is required.

Lemma 3.5. *Let $\varepsilon > 0$ be any number and φ be the function in (2.3). Then,*

$$(3.5) \quad 0 < \varphi'(t)t \ln t \leq 8\varepsilon \quad \forall t \in (1, +\infty).$$

Proof. We take into account formula (2.4) and estimates (3.4), (3.2):

$$\begin{aligned} 0 < \varphi'(t)t \ln t &= \frac{\varepsilon}{\ln(e + (t-1)^4)} \frac{4(t-1)^3}{e + (t-1)^4} t \ln t = \\ &= \varepsilon \frac{\ln t}{\ln(e + (t-1)^4)} \frac{4(t-1)^3 t}{e + (t-1)^4} \leq 8\varepsilon. \end{aligned}$$

□

Lemma 3.6. *Let $\varepsilon > 0$ be any number and φ be the function in (2.3). Then,*

$$(3.6) \quad 0 < \varphi'(t)t \leq 8\varepsilon \quad \forall t \in (1, +\infty).$$

Proof. We take into account formula (2.4) and estimate (3.2):

$$\begin{aligned} 0 < \varphi'(t)t &= \frac{\varepsilon}{\ln(e + (t-1)^4)} \frac{4(t-1)^3}{e + (t-1)^4} t = \\ &= \varepsilon \frac{1}{\ln(e + (t-1)^4)} \frac{4(t-1)^3 t}{e + (t-1)^4} \leq 8\varepsilon. \end{aligned}$$

□

Lemma 3.7. *Let $\varepsilon > 0$ be any number and φ be the function in (2.3). Then,*

$$(3.7) \quad |\varphi''(t)|t^2 \ln t \leq 128\varepsilon \quad \forall t \in (1, +\infty).$$

Proof. We take into account formula (2.5) and estimates (3.2), (3.3), (3.4):

$$\begin{aligned} |\varphi''(t)|t^2 \ln t &\leq \frac{\varepsilon \ln t}{[\ln(e + (t-1)^4)]^2} \left[\frac{4(t-1)^3 t}{e + (t-1)^4} \right]^2 + \\ &= \frac{\varepsilon \ln t}{\ln(e + (t-1)^4)} \frac{12e(t-1)^2 t^2 + 4(t-1)^6 t^2}{[e + (t-1)^4]^2} \leq \varepsilon(4^3 + 48 + 16) = 128\varepsilon. \end{aligned}$$

□

4. PROOF OF THEOREM 2.1

Definitions (2.2) and (2.3) say that, when $t \in [0, 1]$, $\varphi(t) = \frac{3}{2}\pi$ and $g(t) = t^{a-b}$; condition (2.1) guarantees that $1 < a - b$ so that

$$(4.1) \quad g(0) = 0,$$

$$(4.2) \quad \lim_{t \rightarrow 0^+} \frac{g(t)}{t} = 0,$$

$$(4.3) \quad g'(0) = 0;$$

moreover, $g(t) > 0$ for $t > 0$. We recall that, for $t > 1$, $t^{a-b} \leq g(t)$; again, condition (2.1) guarantees that $1 < a - b$ so that

$$(4.4) \quad \lim_{t \rightarrow +\infty} \frac{g(t)}{t} = +\infty.$$

Up to now, $g \in C^0([0, +\infty))$. For $t > 0$ we have

$$(4.5) \quad g(t) = t^{a+b \sin(\varphi(t))} = e^{[a+b \sin(\varphi(t))] \ln t},$$

so that

$$(4.6) \quad g'(t) = e^{[a+b \sin(\varphi(t))] \ln t} \left\{ [b \cos(\varphi(t))] \varphi'(t) \ln t + [a + b \sin(\varphi(t))] \frac{1}{t} \right\} = \frac{g(t)}{t} \{ [b \cos(\varphi(t))] \varphi'(t) t \ln t + [a + b \sin(\varphi(t))] \}.$$

If $t \in (0, 1]$, then $\varphi(t) = \frac{3}{2}\pi$ and $\varphi'(t) = 0$, so that

$$(4.7) \quad g'(t) = \frac{g(t)}{t} [a - b] = [a - b] t^{a-b-1};$$

again, condition (2.1) guarantees that $1 < a - b$ so that

$$(4.8) \quad \lim_{t \rightarrow 0^+} g'(t) = 0.$$

Then, $g \in C^1([0, +\infty))$. Using formula (4.6), when $t > 0$, we have

$$(4.9) \quad g''(t) = \frac{g'(t)t - g(t)}{t^2} \{ [b \cos(\varphi(t))] \varphi'(t) t \ln t + [a + b \sin(\varphi(t))] \} + \frac{g(t)}{t} \{ [-b \sin(\varphi(t))] \varphi'(t) \varphi''(t) t \ln t + [b \cos(\varphi(t))] [\varphi''(t) t \ln t + \varphi'(t) (\ln t + 1)] + [b \cos(\varphi(t))] \varphi'(t) \}.$$

Then $g \in C^2((0, +\infty))$. Now we are going to estimate $g'(t)$ by means of $\frac{g(t)}{t}$. First of all, we consider the case $t \in (0, 1]$: we can use formula (4.7) and we get $g'(t) = (a - b) \frac{g(t)}{t}$. After that, we deal with $t > 1$; we use formula (4.6) and estimate (3.5):

$$(4.10) \quad \begin{aligned} \frac{g(t)}{t} \{ -b8\varepsilon + a - b \} &\leq \\ \underbrace{\frac{g(t)}{t} \{ [b \cos(\varphi(t))] \varphi'(t) t \ln t + [a + b \sin(\varphi(t))] \}}_{g'(t)} &\leq \\ \frac{g(t)}{t} \{ b8\varepsilon + a + b \}. \end{aligned}$$

Note that $-b8\varepsilon + a - b < a - b < a + b < b8\varepsilon + a + b$; then

$$(4.11) \quad \frac{g(t)}{t} \{ -b8\varepsilon + a - b \} \leq g'(t) \leq \frac{g(t)}{t} \{ b8\varepsilon + a + b \} \quad \forall t > 0.$$

Up to now, we only used $a, b > 0$, $1 < a - b$ and $\varepsilon > 0$. Assumption (2.6) guarantees that $\varepsilon < \frac{a-1-b}{224b}$; then $8b\varepsilon < 224b\varepsilon < a - 1 - b$, so that $1 < -b8\varepsilon + a - b$; this and positivity of g give $g'(t) > 0$ when $t > 0$. Moreover, (4.11) can be written as follows

$$(4.12) \quad \frac{1}{b8\varepsilon + a + b} g'(t) \leq \frac{g(t)}{t} \leq \frac{1}{-b8\varepsilon + a - b} g'(t) \quad \forall t > 0.$$

We note that

$$(4.13) \quad t^{a-b} \leq \underbrace{t^{a+b \sin(\varphi(t))}}_{g(t)} \leq t^{a+b} \quad \forall t > 1;$$

then we use estimates (4.11), (4.13) and positivity of $-b8\varepsilon + a - b$:

$$(4.14) \quad t^{a-b-1} \{-b8\varepsilon + a - b\} \leq g'(t) \leq t^{a+b-1} \{b8\varepsilon + a + b\} \quad \forall t > 1.$$

We keep in mind that $g'(t) = (a-b) \frac{g(t)}{t} = (a-b)t^{a-b-1}$ for $t \in (0, 1]$; moreover, $-b8\varepsilon + a - b < a - b < a + b < b8\varepsilon + a + b$; then

$$(4.15) \quad t^{a-b-1} \{-b8\varepsilon + a - b\} \leq g'(t) \leq [t^{a-b-1} + t^{a+b-1}] \{b8\varepsilon + a + b\} \quad \forall t > 0.$$

We divide by t and we get

$$(4.16) \quad t^{a-b-2} \{-b8\varepsilon + a - b\} \leq \frac{g'(t)}{t} \leq [t^{a-b-2} + t^{a+b-2}] \{b8\varepsilon + a + b\} \quad \forall t > 0.$$

We need to estimate $g''(t)t$; to this aim, we use (4.9):

$$(4.17) \quad \begin{aligned} g''(t)t &= \left[g'(t) - \frac{g(t)}{t} \right] \{ [b \cos(\varphi(t))] \varphi'(t)t \ln t + a + b \sin(\varphi(t)) \} + \\ &\quad \frac{g(t)}{t} \{ [-b \sin(\varphi(t))] \varphi'(t)t \varphi'(t)t \ln t + \\ &\quad [b \cos(\varphi(t))] [\varphi''(t)t^2 \ln t + \varphi'(t)t(\ln t + 2)] \}. \end{aligned}$$

We keep in mind (4.6) and we can write as follows

$$(4.18) \quad \begin{aligned} g''(t)t &= g'(t) \{ [b \cos(\varphi(t))] \varphi'(t)t \ln t + a - 1 + b \sin(\varphi(t)) \} + \\ &\quad \frac{g(t)}{t} \{ [-b \sin(\varphi(t))] \varphi'(t)t \varphi'(t)t \ln t + \\ &\quad [b \cos(\varphi(t))] [\varphi''(t)t^2 \ln t + \varphi'(t)t(\ln t + 2)] \}. \end{aligned}$$

For simplicity, define

$$\begin{aligned} \Phi_1(t) &:= [b \cos(\varphi(t))] \varphi'(t)t \ln t + a - 1 + b \sin(\varphi(t)) \\ \Phi_2(t) &:= -b \sin(\varphi(t)) \varphi'(t)t \varphi'(t)t \ln t + [b \cos(\varphi(t))] [\varphi''(t)t^2 \ln t + \varphi'(t)t(\ln t + 2)], \end{aligned}$$

in such a way that (4.18) reads as

$$(4.19) \quad g''(t)t = g'(t)\Phi_1(t) + \frac{g(t)}{t}\Phi_2(t).$$

By (3.5), (3.6) and (3.7) we estimate for $t > 1$

$$\begin{aligned} -8\varepsilon b + a - 1 - b &\leq \Phi_1(t) \leq 8\varepsilon b + a - 1 + b; \\ |\Phi_2(t)| &\leq b8\varepsilon 8\varepsilon + b[128\varepsilon + 8\varepsilon + 16\varepsilon] = b\varepsilon[64\varepsilon + 152]. \end{aligned}$$

Now we estimate $g''(t)t$ from below; when $t > 1$ we keep in mind positivity of g' , g and estimates for Φ_1 , Φ_2 : we have

$$g''(t)t = g'(t)\Phi_1(t) + \frac{g(t)}{t}\Phi_2(t) \geq g'(t)\{-8\varepsilon b + a - 1 - b\} + \frac{g(t)}{t}(-b)\varepsilon[64\varepsilon + 152] =: (I);$$

now we use the right hand side of (4.12) and we get

$$(I) \geq g'(t) \left\{ -8\varepsilon b + a - 1 - b + \frac{-b\varepsilon[64\varepsilon + 152]}{-8\varepsilon b + a - b} \right\} =: (II);$$

now we use (2.6): $\varepsilon < 1$ gives $64\varepsilon + 152 < 216$ and $\varepsilon < \frac{a-1-b}{224b}$ gives $1 < -b8\varepsilon + a - b$, so that

$$(4.20) \quad \frac{b\varepsilon\{64\varepsilon + 152\}}{-b8\varepsilon + a - b} < 216b\varepsilon;$$

then

$$(II) \geq g'(t) \{-8\varepsilon b + a - 1 - b - 216\varepsilon b\};$$

this means that, for $t > 1$ we have

$$g''(t)t \geq g'(t) \{-224\varepsilon b + a - 1 - b\}.$$

Note that we required $-224b\varepsilon + a - 1 - b > 0$ in our assumption (2.6).

When $t \in (0, 1]$, we have $\varphi(t) = \frac{3}{2}\pi$, $\varphi'(t) = 0 = \varphi''(t)$; then $g''(t)t = g'(t)(a - 1 - b)$. Moreover, g' is positive and

$$(4.21) \quad a - 1 - b > a - 1 - b - 224b\varepsilon > 0,$$

then,

$$(4.22) \quad g''(t)t \geq g'(t) \{-224\varepsilon b + a - 1 - b\} \quad \forall t > 0.$$

Since $g'(t) > 0$ when $t > 0$, this last inequality guarantees that $g''(t) > 0$ for all $t > 0$; then g' strictly increases in $(0, +\infty)$; since g' is continuous in $[0, +\infty)$, then g' strictly increases in $[0, +\infty)$: this guarantees that g is strictly convex in $[0, +\infty)$.

Now we estimate $g''(t)t$ from above; when $t > 1$ we keep in mind positivity of g' , g and estimates for Φ_1, Φ_2 : we have

$$g''(t)t = g'(t)\Phi_1(t) + \frac{g(t)}{t}\Phi_2(t) \leq g'(t)\{8\varepsilon b + a - 1 + b\} + \frac{g(t)}{t}b\varepsilon[64\varepsilon + 152] =: (III);$$

now we use the right hand side of (4.12) and we get

$$(III) \leq g'(t) \left\{ 8\varepsilon b + a - 1 + b + \frac{b\varepsilon[64\varepsilon + 152]}{-8\varepsilon b + a - b} \right\} =: (IV);$$

we use (4.20) and we get

$$(IV) \leq g'(t) \{8\varepsilon b + a - 1 + b + 216\varepsilon b\};$$

this means that, for $t > 1$ we have

$$g''(t)t \leq g'(t) \{224\varepsilon b + a - 1 + b\}.$$

When $t \in (0, 1]$, we have $\varphi(t) = \frac{3}{2}\pi$, $\varphi'(t) = 0 = \varphi''(t)$; then $g''(t)t = g'(t)(a - 1 - b)$. Moreover, g' is positive so that

$$(4.23) \quad g''(t)t \leq g'(t) \{224\varepsilon b + a - 1 + b\} \quad \forall t > 0.$$

This ends the proof of Theorem 2.1. □

5. ANOTHER EXAMPLE

Now we give an example in the subquadratic case by modifying a little bit the previous example of section 2: we introduce an additional restriction on a, b and we select a smaller ε . More precisely, We fix $a, b \in (0, +\infty)$ with (2.1) as in section 2; moreover, we require, in addition,

$$(5.1) \quad a + b < 2.$$

We consider $g : [0, +\infty) \rightarrow [0, +\infty)$ given by (2.2) with φ as in (2.3) with $\varepsilon > 0$ satisfying (2.6) as in section 2; moreover, we require, in addition,

$$(5.2) \quad \varepsilon < \frac{2-a-b}{224b}.$$

Please, note that (5.1) gives $0 < 2 - a - b$, so the requirement (5.2) is in accordance with $0 < \varepsilon$ and it implies

$$224b\varepsilon + a - 2 + b < 0.$$

This and the right hand side of (2.9) in Theorem 2.1 give

Theorem 5.1. *Let us consider $a, b \in (0, +\infty)$ verifying (2.1), (5.1); we consider $g(t)$ given by (2.2) where φ is defined in (2.3) and ε satisfies (2.6), (5.2). Then*

$$g''(t) - \frac{g'(t)}{t} \leq \{224b\varepsilon + a - 2 + b\} \frac{g'(t)}{t} < 0 \quad \forall t > 0$$

and we get $M = 1$ in the right hand side of (1.8). Since

$$\left(\frac{g'(t)}{t}\right)' = \frac{g''(t)t - g'(t)}{t^2} = \left(g''(t) - \frac{g'(t)}{t}\right) \frac{1}{t} < 0,$$

we get

$$t \rightarrow \frac{g'(t)}{t} \text{ strictly decreases in } (0, +\infty).$$

REFERENCES

- [1] Baroni P. - *Riesz potential estimates for a general class of quasilinear equations* - Calc. Var. Partial Differential Equations **53** (2015), 803-846.
- [2] Baroni P., Colombo M., Mingione G. - *Harnack inequalities for double phase functionals* - Nonlinear Anal., **121** (2015), 206-222.
- [3] Beck, L., Mingione, G. - *Lipschitz bounds and non-uniform ellipticity* - Comm. Pure Appl. Math. (2020). <https://doi.org/10.1002/cpa.21880>
- [4] Bildhauer M., Fuchs M. - *Variational integrals of splitting-type: higher integrability under general growth conditions* - Ann. Mat. Pura Appl., **188** (2009), 467-496.
- [5] Cianchi A., Maz'ya V. - *Global boundedness of the gradient for a class of nonlinear elliptic systems* - Arch. Ration. Mech. Anal. **212** (2014), 129-177.
- [6] Cupini G., Marcellini P., Mascolo E. - *Regularity of minimizers under limit growth conditions* - Nonlinear Anal., **153** (2017), 294-310.
- [7] De Filippis C., Mingione G. - *On the Regularity of Minima of Non-autonomous Functionals* - J. Geom. Anal. (2019) <https://doi.org/10.1007/s12220-019-00225-z>
- [8] Diening L., Ettwein F. - *Fractional estimates for non-differentiable elliptic systems with general growth* - Forum Math. **20** (2008), 523-556.
- [9] Eleuteri M., Marcellini P., Mascolo E. - *Lipschitz estimates for systems with ellipticity conditions at infinity* - Ann. Mat. Pura Appl., **195** (2016), 1575-1603.
- [10] Esposito L., Leonetti F., Mingione G. - *Higher integrability for minimizers of integral functionals with (p, q) growth* - J. Differential Equations, **157** (1999), 414-438.
- [11] Esposito L., Leonetti F., Mingione G. - *Sharp regularity for functionals with (p, q) growth* - J. Differential Equations, **204** (2004), 5-55.
- [12] Fusco N., Sbordone C. - *Higher integrability of the gradient of minimizers of functionals with non standard growth conditions* - Comm. Pure Appl. Math., **43** (1990), 673-683.
- [13] Fusco N., Sbordone C. - *Some remarks on the regularity of minima of anisotropic integrals* - Comm. P. D. E., **18** (1993), 153-167.
- [14] Leonetti F., Mascolo E., Siepe F. - *Everywhere regularity for a class of vectorial functionals under sub-quadratic general growth conditions* - J. Math. Anal. Appl., **287** (2003), 593-608.
- [15] Lieberman G. - *The natural generalization of the natural conditions of Ladyzhenskaya and Ural'tseva* Miniconference on Operators in Analysis, 151-158, Centre for Mathematics and its Applications, Mathematical Sciences Institute, The Australian National University, Canberra AUS, 1990. <https://projecteuclid.org/euclid.pcma/1416335069>
- [16] Marcellini P. - *Regularity of minimizers of integrals of the calculus of variations with nonstandard growth conditions* - Arch. Rational Mech. Anal., **105** (1989), 267-284.
- [17] Mingione G. - *Regularity of minima: an invitation to the dark side of the calculus of variations* - Appl. Math., **51** (2006), 355-425.
- [18] Moscarriello G., Nania L. - *Hölder continuity of minimizers of functionals with nonstandard growth conditions* - Ricerche Mat., **40** (1991), 259-273.

- [19] Talenti G. - *Boundedness of minimizers* - Hokkaido Math. J., **19** (1990), 259-279.

CRISTIANA DE FILIPPIS, MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, ANDREW WILES BUILDING,
RADCLIFFE OBSERVATORY QUARTER, WOODSTOCK ROAD, OXFORD, OX26GG, OXFORD, UNITED KINGDOM
E-mail address: `Cristiana.DeFilippis@maths.ox.ac.uk`

FRANCESCO LEONETTI, UNIVERSITÀ DI L'AQUILA, DIPARTIMENTO DI INGEGNERIA E SCIENZE DELL'INFORMAZIONE
E MATEMATICA, VIA VETOIO SNC, 67100 L'AQUILA, ITALY
E-mail address: `leonetti@univaq.it`