

Control of chaos with minimal information transfer

Christoph Kawan*

Abstract

This paper studies set-invariance and stabilization of hyperbolic sets over rate-limited channels for discrete-time control systems. We first investigate structural and control-theoretic properties of hyperbolic sets, in particular such that arise by adding small control terms to uncontrolled systems admitting (classical) hyperbolic sets. Then we derive a lower bound on the invariance entropy of a hyperbolic set in terms of the difference between the unstable volume growth rate and the measure-theoretic fiber entropy of associated random dynamical systems. We also prove that our lower bound is tight in two extreme cases. Furthermore, we apply our techniques to the problem of local uniform stabilization to a hyperbolic set. Finally, we discuss an example built on the Hénon horseshoe.

Keywords: Control under data-rate constraints; stabilization; discrete-time nonlinear systems; uniform hyperbolicity; invariance entropy; escape rates; SRB measures

1 Introduction

1.1 General introduction

Hyperbolicity is one of the most important paradigms in the modern theory of dynamical systems as it provides a way of understanding the mechanisms leading to erratic behavior of trajectories in chaotic systems. The first traces of the hyperbolic theory are usually located in Poincaré’s prize memoir on the three-body problem in celestial mechanics [56]. It took, however, almost 80 years after Poincaré’s work until a general axiomatic definition of hyperbolicity was presented by Stephen Smale [61]. This definition arose from the desire to explain chaotic phenomena observed in the study of differential equations modeling real-world engineering systems [11, 47], and it built on the concept of Anosov diffeomorphisms studied before by the Russian school. Smale’s notion of a uniformly hyperbolic set was soon generalized in different directions to cover a great variety of systems. We do

*Institute of Informatics, LMU Munich, Oettingenstraße 67, 80538 München, Germany; e-mail: christoph.kawan@lmu.de

not attempt to give an account of all these research threads. The reader may consult Hasselblatt [33], Katok & Hasselblatt [39] and Hasselblatt & Pesin [34] to obtain a comprehensive overview of the still ongoing research in hyperbolic dynamics.

For control engineers, a particularly interesting research direction rooted in hyperbolic dynamics was initiated by Ott, Grebogi & Yorke [55] and is known under the term *control of chaos*. While chaoticity in most cases is considered an unpleasant behavior in an engineering system, in the control of chaos its features are exploited to stabilize a system with low energy use. Here one uses the abundance of unstable periodic orbits on a hyperbolic set to pick one of these orbits and keep the system on a nearby orbit via small “kicks” (control actions), applied at the right time to drive the state closer to the stable manifold of the periodic orbit. One of various applications of this method can be found in space exploration, where it is used to stabilize space probes at unstable Lagrangian points of the solar system, see e.g. [60].

A relatively new and vibrant subfield of control, in which hyperbolic dynamics is likely to play a key role, is the *control under communication constraints*. Motivated by real-world applications suffering from informational bottlenecks in the communication links between sensors and controllers or controllers and actuators, many researchers have studied the problem of characterizing the minimal requirements on a communication network necessary for achieving a desired control goal by a proper coder-controller design. Two of the most often cited examples, in which data-rate constraints constitute the bottleneck, are the control of large-scale networked systems, where the communication resources have to be distributed among many agents (see e.g. [36, 54, 49]), and the coordinated control of unmanned underwater vehicles, where the medium water makes high-rate communication difficult.




Figure 1: Control of a system over a finite-capacity channel

A conceptually simple though highly non-trivial scenario allowing to study some of the essential aspects of the problem is depicted in Fig. 1. Here a controller receives state information, collected by sensors, through a finite-capacity communication channel and the goal is to stabilize the system. In the figure, $x(t)$ denotes the state of the system at time t , $s(t_k)$ is a symbol sent through the (noiseless) channel at the

sampling time t_k , and $u(\cdot)$ is the control input generated by the controller, based on the knowledge of the transmitted symbols. A now classical result focusing on linear system models is known as the *data-rate theorem*. Proven under a great variety of different assumptions on the system model, communication protocol and stabilization objective, it yields the unambiguous answer that there is a minimal channel capacity, given by the log-sum of the open-loop unstable eigenvalues, above which the stabilization objective can be achieved. The fact that this number appears in the theory of dynamical systems as the topological or measure-theoretic entropy of a linear system [8] has motivated researchers to look for further and deeper connections between the data-rate-constrained stabilization problem and ergodic theory, when the dynamics is nonlinear.

These investigations led to the introduction of various notions of “control entropy” which are quantities defined in terms of the open-loop system, resembling topological or measure-theoretic entropy in dynamical systems. Such entropy notions are particularly successful for the description of the minimal channel capacity when the stabilization objective can be achieved in a repetitive way, i.e., via a coding and control protocol that repeats precisely the same tasks periodically in time. An example for such an objective is set-invariance. Indeed, if a coding and control scheme achieves invariance of a certain set in a time interval $[0, \tau]$, then the same scheme can be applied again after time τ to achieve invariance on $[\tau, 2\tau]$, etc. The notion of *topological feedback entropy*, introduced in Nair et al. [53], captures the smallest average data rate above which a compact controlled invariant set can be made invariant by a coding and control scheme that operates over a noiseless discrete channel transmitting state information from the coder to the controller.

A related, in fact equivalent [16], notion of entropy was introduced in Colonius & Kawan [15] under the name *invariance entropy*. While topological feedback entropy is defined in an open-cover fashion similar to the definition of topological entropy by Adler, Konheim & McAndrew [1], invariance entropy is defined via so-called spanning sets of control inputs. The idea is simple: if the controller receives n bits of information, it can distinguish at most 2^n different states, hence generate at most 2^n different control inputs, and consequently, the number of necessary control inputs to achieve invariance (on a finite time interval) is a measure for the required information. Hence, the invariance entropy of a compact controlled invariant set Q is defined as

$$h_{\text{inv}}(Q) := \lim_{\tau \rightarrow \infty} \frac{1}{\tau} \log_2 r_{\text{inv}}(\tau, Q),$$

where $r_{\text{inv}}(\tau, Q)$ denotes the minimal number of control inputs necessary to achieve invariance of Q on a time interval of length τ .

A theory aimed at the description of invariance entropy in terms of dynamical characteristics of the system (such as Lyapunov exponents), has been developed to a certain extent in [22, 25, 42, 43], but mainly for continuous-time systems. In particular, the papers [22, 25] demonstrate that uniform hyperbolicity and con-

trollability assumptions together allow for the derivation of a closed-form expression for $h_{\text{inv}}(Q)$ in terms of instability characteristics on Q such as the sum of the unstable Lyapunov exponents or relative/conditional entropy of the associated skew-product system relative to the left-shift on the space of admissible control inputs. This theory has been successfully applied to right-invariant systems on flag manifolds of semisimple Lie groups [24, 26]. Other aspects of invariance entropy and generalizations thereof have been studied in a number of papers, including [12, 13, 20, 21, 37, 44, 64, 65].

In this paper, we study a discrete-time setting in which we introduce the notion of a uniformly hyperbolic set for a control system, with the ultimate goal to provide a closed-form expression for the invariance entropy of such sets. Although we are not able to achieve this goal in the fully general case, we provide a lower bound together with proofs for two special cases that this bound is tight under additional controllability assumptions. Moreover, via the introduced techniques we provide a necessity result for the local uniform stabilization of a control system to a hyperbolic set (of the autonomous system associated to a fixed control value). This result can be seen as an extension of the local stabilization result presented in [53] for the asymptotic stabilization to an equilibrium point. At the same time, it closes a gap in the proof presented in [53] and provides a new interpretation of a classical escape-rate formula in the theory of dynamical systems [9, 67]. Technical details of our proof program are presented in the next subsection.

Some general references for the theory of control under communication constraints are the books [50, 29, 69] and the survey papers [2, 28, 54].

1.2 Structure and contents of the paper

In this paper, we study discrete-time, time-invertible control systems of the form

$$x_{t+1} = f(x_t, u_t), \quad t \in \mathbb{Z} \quad (1)$$

with states in a Riemannian manifold M and controls in a compact and connected metric space U . Under appropriate regularity assumptions, the system (1) induces a continuous skew-product system $\Phi = (\Phi_t)_{t \in \mathbb{Z}}$ (called *control flow*) on the extended state space $\mathcal{U} \times M$ with $\mathcal{U} := U^{\mathbb{Z}}$ (equipped with the product topology), with the left shift operator θ acting on \mathcal{U} as the driving system. The transition map of (1) is denoted by φ so that

$$\Phi_t(u, x) = (\theta^t u, \varphi(t, x, u)), \quad \Phi_t : \mathcal{U} \times M \rightarrow \mathcal{U} \times M.$$

Frequently, we also write $\varphi_{t,u} = \varphi(t, \cdot, u)$.

A *uniformly hyperbolic set* of (1) is a compact all-time controlled invariant subset $Q \subset M$ that admits a splitting of its extended tangent bundle into a stable and an unstable subbundle which are continuous and allow for uniform estimates of

contraction and expansion rates. The difference to the classical autonomous case is that the stable and unstable subspaces, in general, depend on $(u, x) \in \mathcal{U} \times M$ and not only on x . Similar notions of uniformly hyperbolic sets are studied in the theory of random dynamical systems (RDS), where the driving system models the influence of the noise on the dynamics [32, 46, 48]. Several classical tools from the theory of hyperbolic dynamical systems are available to study uniformly hyperbolic sets of control systems, in particular the *stable manifold theorem*, the *shadowing lemma* and the *Bowen-Ruelle volume lemma*, cf. Subsection 3.2. A uniformly hyperbolic set Q of (1) can be lifted to the extended state space $\mathcal{U} \times M$ by putting

$$L(Q) := \{(u, x) \in \mathcal{U} \times M : \varphi(\mathbb{Z}, x, u) \subset Q\},$$

which is a compact invariant set of the control flow Φ . Then $E^-(u, x)$ and $E^+(u, x)$ denote the stable and unstable subspace at $(u, x) \in L(Q)$, respectively.

Section 3 is devoted to the study of structural and control-theoretic properties of uniformly hyperbolic sets. While the first two subsections introduce the necessary definitions and tools, the third one contains the actual analysis.

Subsection 3.3: Our analysis starts with the study of the *u-fibers* $Q(u) = \{x \in M : \varphi(\mathbb{Z}, x, u) \subset Q\}$, $u \in \mathcal{U}$. Assuming that $L(Q)$ is isolated invariant, the shadowing lemma can be used to prove that all fibers $Q(u)$ are nonempty and homeomorphic to each other. The set-valued mapping $u \mapsto Q(u)$ from \mathcal{U} into the space of closed subsets of Q is, in general, upper semicontinuous (even without the assumption of uniform hyperbolicity). To derive a lower bound on $h_{\text{inv}}(Q)$, we require $u \mapsto Q(u)$ to be lower semicontinuous as well. This assumption can be verified in a “small-perturbation” setting, where we fix a constant control $u^0 \in U$, assume that the diffeomorphism $f(\cdot, u^0) : M \rightarrow M$ admits an isolated invariant uniformly hyperbolic set Λ , and then restrict the control range to a small neighborhood of u^0 in U . By standard perturbation results (see e.g. [48]), one shows that the so-defined control system admits a uniformly hyperbolic set Q whose u^0 -fiber coincides with Λ . We also study the controllability properties on a set Q that arises in this way. Assuming that Λ is topologically transitive and combining classical results from discrete-time control with shadowing arguments, we obtain under analyticity and accessibility assumptions that Q has nonempty interior and complete controllability holds on an open and dense subset of Q . The proof uses the theory of accessibility and universally regular controls developed in Albertini & Sontag [3] and Sontag & Wirth [63]. It remains an open question if the fiber map $u \mapsto Q(u)$ is lower semicontinuous for a more general class of uniformly hyperbolic sets.

In Section 4, we derive a lower bound on the invariance entropy of a uniformly hyperbolic set in terms of dynamical characteristics of associated random dynamical systems. We also discuss the tightness of the bound under additional controllability assumptions.

Subsection 4.1: If the fiber map $u \mapsto Q(u)$ of a compact all-time controlled invariant set Q is lower semicontinuous, we can derive a lower bound on $h_{\text{inv}}(Q)$ in terms of a uniform rate of escape from the ε -neighborhoods of the u -fibers. This lower bound is based on the observation that the sets

$$Q^\pm(u, \tau) := \{x \in M : \varphi(t, x, u) \in Q \text{ for } -\tau < t < \tau\}$$

shrink down to the u -fiber $Q(u)$ as τ tends to infinity, and this shrinking process is uniform with respect to u if the fiber $Q(u)$ depends continuously on u in the Hausdorff metric (which is equivalent to simultaneous upper and lower semicontinuity). If $\mathcal{S} \subset \mathcal{U}$ is a (τ, Q) -spanning set, i.e., a set of control inputs guaranteeing invariance on the time interval between 0 and $\tau - 1$, then Q is covered by the sets

$$Q(u, \tau) := \{x \in M : \varphi(t, x, u) \in Q \text{ for } 0 \leq t < \tau\}, \quad u \in \mathcal{S},$$

which are related by a time shift to the sets $Q^\pm(u, \tau)$. Finally, introducing the sets

$$Q(u, \tau, \varepsilon) := \{x \in M : \text{dist}(\varphi(t, x, u), Q(\theta^t u)) \leq \varepsilon, 0 \leq t < \tau\},$$

a careful analysis of these relations leads to the estimate

$$h_{\text{inv}}(Q) \geq -\liminf_{\tau \rightarrow \infty} \sup_{u \in \mathcal{U}} \frac{1}{\tau} \log \text{vol}(Q(u, \tau, \varepsilon)) \quad (2)$$

which holds true for every $\varepsilon > 0$ provided that Q has positive volume.

Subsection 4.2: Using a classical idea from the study of escape rates [9, 67], one can estimate the volume in (2) in the following way:

$$\text{vol}(Q(u, \tau, \varepsilon)) \leq \text{const} \cdot \sum_{x \in F_{u, \tau, \delta}} J^+ \varphi_{\tau, u}(x)^{-1}. \quad (3)$$

Here, $F_{u, \tau, \delta} \subset Q(u)$ is a (u, τ, δ) -separated set¹ for a small $\delta > 0$, and

$$J^+ \varphi_{\tau, u}(x) = |\det D\varphi_{\tau, u}(x)|_{E^+(u, x)} : E^+(u, x) \rightarrow E^+(\Phi_\tau(u, x))|$$

denotes the *unstable determinant* of the linearization. The main idea behind this estimate is to cover the set $Q(u, \tau, \varepsilon)$ with Bowen-balls of order τ and radius δ , and estimate the volumes of these balls via the Bowen-Ruelle volume lemma. A shadowing argument allows to move the centers of these balls to $Q(u)$. Here, the required uniform hyperbolicity on Q is fully exploited via the use of shadowing and hyperbolic volume estimates.

¹That is, for any two $x, y \in F_{u, \tau, \delta}$ with $x \neq y$ one has $d(\varphi(t, x, u), \varphi(t, y, u)) > \delta$ for some $0 \leq t < \tau$.

Subsections 4.3 and 4.4: To make use of the estimate (3), two intermediate steps are taken, the first of which consists in interchanging the order of limit inferior and supremum in (2). This would be unproblematic if the functions $v_\tau^\varepsilon(u) := \log \text{vol}(Q(u, \tau, \varepsilon))$, $\tau > 0$, would define a continuous subadditive cocycle over the shift (\mathcal{U}, θ) for some $\varepsilon > 0$. Since we are not able to prove this, we introduce families of functions $w_\tau^\delta : \mathcal{U} \rightarrow \mathbb{R}$ that are indeed subadditive cocycles over the shift and approximate v_τ^ε in a certain sense (see Proposition 4.2 for details). Together with the continuity of $u \mapsto v_\tau^\varepsilon(u)$ (Lemma 4.3) this allows to prove that the order of limit and supremum in (2) can be interchanged under the limit for $\varepsilon \downarrow 0$. As a consequence,

$$h_{\text{inv}}(Q) \geq - \lim_{\varepsilon \downarrow 0} \sup_{u \in \mathcal{U}} \liminf_{\tau \rightarrow \infty} \frac{1}{\tau} \log \text{vol}(Q(u, \tau, \varepsilon)). \quad (4)$$

Subsection 4.5: The second step consists in rewriting (4) via ergodic growth rates with respect to shift-invariant probability measures on \mathcal{U} . This leads to

$$h_{\text{inv}}(Q) \geq - \lim_{\varepsilon \downarrow 0} \sup_{P \in \mathcal{M}(\theta)} \liminf_{\tau \rightarrow \infty} \frac{1}{\tau} \int \log \text{vol}(Q(u, \tau, \varepsilon)) \, dP(u), \quad (5)$$

where $\mathcal{M}(\theta)$ denotes the set of all θ -invariant Borel probability measures. In the proof of (5), we use again the approximate subadditivity of $(v_\tau^\varepsilon)_{\tau \in \mathbb{Z}_+}$ established in Proposition 4.2 together with standard arguments used in the context of subadditive cocycles [52]. Here, it is important to point out that each $P \in \mathcal{M}(\theta)$ together with the transition map φ formally induces an RDS over $(\mathcal{U}, \mathcal{B}(\mathcal{U}), P, \theta)^2$ that we denote by (φ, P) . In this context, growth rates of the form

$$\lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int \log \text{vol}(Q(u, \tau, \varepsilon)) \, dP(u)$$

are known as *random escape rates*, see [48].

Subsection 4.6: A further lower bound on $h_{\text{inv}}(Q)$ is derived from (5) and (3) via arguments taken from the standard proof of the variational principle for pressure of RDS, cf. [7]. Essentially, the growth rates of $\#F_{u, \tau, \delta}$ and $J^+ \varphi_{\tau, u}(x)$ are separated and we end up with the estimate

$$h_{\text{inv}}(Q) \geq \inf_{\mu \in \mathcal{M}(\Phi|_{L(Q)})} \left[\int \log J^+ \varphi_{1, u}(x) \, d\mu(u, x) - h_\mu(\varphi, (\pi_{\mathcal{U}})_* \mu) \right], \quad (6)$$

where the infimum is taken over all Φ -invariant Borel probability measures μ , supported on $L(Q)$, and $(\pi_{\mathcal{U}})_* \mu$ denotes the marginal of μ on \mathcal{U} . Moreover,

²Here, $\mathcal{B}(\mathcal{U})$ denotes the Borel σ -algebra on \mathcal{U} .

$h_\mu(\varphi, (\pi_{\mathcal{U}})_*\mu)$ is the measure-theoretic entropy of the RDS $(\varphi, (\pi_{\mathcal{U}})_*\mu)$ with respect to its invariant measure μ (which is the term that captures the growth rate of $\#F_{u,\tau,\delta}$). The well-known Margulis-Ruelle inequality [4] guarantees that

$$h_\mu(\varphi, (\pi_{\mathcal{U}})_*\mu) \leq \int \log J^+ \varphi_{1,u}(x) d\mu(u, x)$$

so that the lower bound (6) is always nonnegative. A natural interpretation of the involved terms is that $\int \log J^+ \varphi_{1,u}(x) d\mu(u, x)$ measures the total instability of the dynamics on Q (seen by the measure μ), while $h_\mu(\varphi, (\pi_{\mathcal{U}})_*\mu)$ measures the part of the instability not leading to exit from Q . This makes perfect sense, since $h_{\text{inv}}(Q)$ measures the control complexity necessary for preventing exit from Q . The fact that we are taking the infimum over all measures might be related to the characterization of invariance entropy as the minimal data rate amongst all coding and control strategies which lead to invariance of Q .

Subsection 4.7: A natural question arising from (6) is whether the infimum on the right-hand side is attained as a minimum. Using the property of expansivity which holds on every uniformly hyperbolic set, this can be verified, and hence

$$h_{\text{inv}}(Q) \geq \int \log J^+ \varphi_{1,u}(x) d\hat{\mu}(u, x) - h_{\hat{\mu}}(\varphi, (\pi_{\mathcal{U}})_*\hat{\mu}) \quad (7)$$

for a (not necessarily unique) measure $\hat{\mu} \in \mathcal{M}(\Phi|_{L(Q)})$. This inequality has interesting consequences, since it allows us to obtain a better understanding of the case when $h_{\text{inv}}(Q) = 0$. Indeed, if $h_{\text{inv}}(Q) = 0$, then

$$h_{\hat{\mu}}(\varphi, (\pi_{\mathcal{U}})_*\hat{\mu}) = \int \log J^+ \varphi_{1,u}(x) d\hat{\mu}(u, x)$$

which exhibits $\hat{\mu}$ as an SRB measure of the RDS $(\varphi, (\pi_{\mathcal{U}})_*\hat{\mu})$. It seems plausible that conversely the existence of an SRB measure implies the existence of some sort of attractor inside Q which, under additional controllability assumptions, would force $h_{\text{inv}}(Q)$ to be zero. For the case of a hyperbolic set as constructed in the small-perturbation setting, this is proved in Theorem 4.17 (in Subsection 4.9).

Subsection 4.8: The lower bound (6) can also be expressed in purely topological terms. Via the variational principle for the pressure of RDS, we can first write it in the form

$$h_{\text{inv}}(Q) \geq - \sup_{P \in \mathcal{M}(\theta)} \pi_{\text{top}}(\varphi^Q, P; -\log J^+ \varphi),$$

where $\pi_{\text{top}}(\varphi^Q, P; -\log J^+ \varphi)$ is the topological pressure with respect to the potential $-\log J^+ \varphi$ of the bundle RDS defined by fixing the measure P on \mathcal{U} and restricting Φ to the invariant set $L(Q)$. Letting

$$\pi_\alpha(u, \tau, \varepsilon) := \sup \left\{ \sum_{x \in F} 2^{\sum_{s=0}^{\tau-1} \alpha(\Phi_s(u, x))} : F \subset Q(u) \text{ is } (u, \tau, \varepsilon)\text{-separated} \right\},$$

we can derive the identity

$$\sup_{P \in \mathcal{M}(\theta)} \pi_{\text{top}}(\varphi^Q, P; -\log J^+ \varphi) = \sup_{u \in \mathcal{U}} \lim_{\varepsilon \downarrow 0} \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log \pi_{-\log J^+ \varphi}(u, \tau, \varepsilon),$$

where we use again that the involved quantities can be approximated by subadditive cocycles. This purely topological expression can possibly serve as a hint how to prove an achievability result, i.e., a sufficiency result for the required data rate to make Q invariant.

Subsection 4.9: We discuss the tightness of the obtained lower bound for $h_{\text{inv}}(Q)$, which in two extreme cases can be made very plausible. The first case occurs when the fibers $Q(u)$ are finite. Then, the measure-theoretic entropy term in the lower bound vanishes and the tightness has been proved in [25] for the continuous-time case under accessibility and controllability assumptions. It is more or less obvious that the same proof works in discrete time. For the small-perturbation setting, this is demonstrated in Theorem 4.14. The other case is the one in which $L(Q)$ supports an SRB measure for one of the RDS (φ, P) , implying that the lower bound vanishes. In this case, it should be possible to find an attractor inside Q so that controllability on Q would make it possible to steer from every initial state into the associated basin of attraction, where no further control actions are necessary, leading to $h_{\text{inv}}(Q) = 0$. Again, in the small-perturbation setting we can provide a proof, see Theorem 4.17.

In Section 5, we prove a result on the necessary average data rate for local uniform stabilization to a uniformly hyperbolic set Λ of the diffeomorphism $f_0 = f(\cdot, u^0)$ with $u^0 \in U$. From the analysis of the preceding section, it almost immediately follows that (under mild regularity assumptions) a lower bound on the data rate is given by the negative topological pressure of f_0 with respect to the negative unstable log-determinant on Λ . This quantity is well-studied in the theory of hyperbolic dynamical systems and, in particular, appears as the rate at which volume escapes from a small neighborhood of an Axiom A basic set [9, 67]. For the case when Λ is a periodic orbit, we prove that our lower bound is tight. For the case when Λ is topologically transitive and supports an SRB measure, it is trivially tight, because this implies that Λ is an attractor.

Section 6 presents an example built on the so-called *Hénon horseshoe*, a non-attracting uniformly hyperbolic set of a map from the Hénon family. Our small-perturbation results allow to study uniformly hyperbolic sets that arise by adding small control terms to the given Hénon map. In particular, numerical studies are available which provide estimates for the escape rate from a small neighborhood of the Hénon horseshoe that in turn yield estimates for the invariance entropy of its small perturbations as well as for the smallest average data rate necessary for stabilization to the horseshoe.

Section 7 presents some open questions and the Appendix (Sections A and B)

contains auxiliary results and supplementary material.

1.3 Remarks, interpretation and further directions

The results presented in this paper. Our results should not be seen first and foremost from a practical point of view (of applicability to engineering problems), but from the viewpoint of a theoretical understanding of stabilization over rate-limited channels. They relate the control-theoretic quantity h_{inv} to quantities that are well studied and of utmost importance in the theory of dynamical systems. Moreover, they give these dynamical quantities a new, control-theoretic interpretation. This should be an inspiration for the search for further relations of similar nature. In particular, in the context of stochastic control systems and stochastic stabilization objectives, it is very likely that weaker and by nature probabilistic/ergodic forms of hyperbolicity such as *non-uniform hyperbolicity* [33, Ch. 5] are helpful to derive similar and even more interesting results.

The role of hyperbolicity (advantages and disadvantages). The assumption of uniform hyperbolicity provides us with tools and techniques that allow to derive very clean and precise results. Additionally, uniform hyperbolicity guarantees the robustness that is necessary for a control strategy to work properly with regard to parameter uncertainties and external noise, cf. [23]. On the other hand, uniform hyperbolicity is a property that is hard to check for a concrete model, although some numerical approaches to this problem exist, see e.g. [6]. Moreover, most systems are not uniformly hyperbolic but exhibit some weaker form of hyperbolicity. Hence, the uniformly hyperbolic case should be seen only as a first step towards a more general theory.

Extension to noisy systems. For noisy systems of the form

$$x_{t+1} = f(x_t, u_t, w_t)$$

with reasonably small bounded noise w_t , it is conceivable that a finite-time analysis leads to comparable results on the minimal data rate for stabilization to a hyperbolic set of the unperturbed system $x_{t+1} = f(x_t, u_t, 0)$. In this case, a time horizon T needs to be chosen small enough so that the noise does not dominate over the control within a time interval of length T . The expected result would then characterize a trade-off between the noise amplitude and the time horizon, respectively, the achievable data rate.

History and new contributions. Many of the ideas and results in this paper have appeared before in other publications:

- The idea of estimating invariance entropy from below by an escape rate has first appeared in [40].
- For continuous-time systems, uniformly hyperbolic sets in the sense of this paper turn out to be quite simple, namely, their u -fibers are finite [41]. In [25], a closed-form expression for the invariance entropy of uniformly hyperbolic control sets of continuous-time systems has been derived. In particular, the derivation of the lower bound already contains some of the ideas involved in the paper at hand, and Theorem 4.14 (the achievability result) mainly uses ideas developed in [25].
- In [22], a special class of partially hyperbolic controlled invariant sets has been introduced and a lower bound for their invariance entropy has been derived. Most of the ideas leading to the estimate (6) are already contained in [22].

The genuinely new contributions of the paper at hand are the following:

- The notion of an isolated (controlled) invariant set used in [22] has been weakened. Instead of assuming an isolatedness condition on the state space M , we now assume that the lift $L(Q)$ in $\mathcal{U} \times M$ is an isolated invariant set of the control flow, which is a weaker and more natural condition.
- The “small-perturbation” construction of a uniformly hyperbolic set, presented in Subsection 3.3 (although well-known in another context) has not been presented before. This construction sheds some light on the assumption of lower semicontinuity of the fiber map which was already used in [22]. Moreover, the analysis of the controllability properties on such a set is new and, to the best of my knowledge, this has not been studied before although somewhat related ideas can be found in Colonius & Du [14].
- The results in Subsection 4.7 are new. In particular, the relation between vanishing invariance entropy and the existence of SRB measures (Corollary 4.10 and Theorem 4.17) is a new contribution of this paper.
- The topological characterization of the lower bound in Subsection 4.8 is another novel contribution.
- The achievability results (Theorem 4.14 and Theorem 4.17) and the local stabilization results (Theorem 5.1 and Theorem 5.4) have not appeared before.
- The example built on the Hénon horseshoe presented in Section 6 is another new contribution.

2 Preliminaries

2.1 Notation

By $|A|$ we denote the cardinality of a set A . Logarithms are by default taken to the base 2. We write \mathbb{Z} , \mathbb{Z}_+ and $\mathbb{Z}_{>0}$ for the sets of integers, nonnegative integers and positive integers, respectively. By $[a; b]$, $(a; b)$, $(a; b]$ and $[a; b)$ we denote the closed, open and half-open intervals in \mathbb{Z} , respectively. The notation $\mathbb{1}_A$ stands for the indicator function of a subset A of some space X , i.e., $\mathbb{1}_A(x) = 1$ if $x \in A$ and $\mathbb{1}_A(x) = 0$ otherwise. If X and Y are two spaces, we write $\pi_X : X \times Y \rightarrow X$ and $\pi_Y : X \times Y \rightarrow Y$ for the corresponding canonical projections $\pi_X(x, y) = x$ and $\pi_Y(x, y) = y$, respectively.

All manifolds in this paper are assumed to be connected and smooth, i.e., equipped with a C^∞ differentiable structure. If M is a manifold, we write $T_x M$ for its tangent space at x . Also, Riemannian metrics are always assumed to be smooth. Given a manifold equipped with a Riemannian metric, we write $|\cdot|$ for the induced norm on each tangent space. Moreover, $d(\cdot, \cdot)$ denotes the induced distance function and $\text{vol}(\cdot)$ the associated volume measure. Finally, we write \exp_x for the Riemannian exponential map at x .

In any metric space (X, d) , we write $B_\varepsilon(x)$ for the open ε -ball centered at x , $\text{dist}(x, A) = \inf_{y \in A} d(x, y)$ for the distance of a point x to a set A , and $N_\varepsilon(A) = \{x \in X : \text{dist}(x, A) \leq \varepsilon\}$ for the closed ε -neighborhood of a set A . The open ε -neighborhood, in contrast, is denoted by $N_\varepsilon^\circ(A)$. Moreover, we use the notation $d_H(A, B)$ for the Hausdorff distance of two sets A, B :

$$d_H(A, B) = \max\{\text{Dist}(A, B), \text{Dist}(B, A)\}, \quad \text{Dist}(A, B) = \sup_{a \in A} \text{dist}(a, B).$$

For any set $A \subset X$, we write $\text{cl } A$, $\text{int } A$ and ∂A for the closure, interior and boundary of A , respectively. Moreover, we write $\text{diam}(A) = \sup_{x, y \in A} d(x, y)$ for the diameter of A . If X and Y are two metric spaces, $C^0(X, Y)$ stands for the space of all continuous mappings $f : X \rightarrow Y$.

If $T : X \rightarrow Y$ is a measurable map between measurable spaces (X, \mathcal{F}_X) and (Y, \mathcal{F}_Y) , respectively, we write T_* for the operator induced by T on the set of measures on (X, \mathcal{F}_X) , i.e., $(T_*\mu)(B) = \mu(T^{-1}(B))$ for every measure μ on (X, \mathcal{F}_X) and all $B \in \mathcal{F}_Y$. We write $\mathcal{M}(T)$ for the set of all T -invariant Borel probability measures of a continuous map $T : X \rightarrow X$ on a compact metric space X . By $\mathcal{B}(X)$ we denote the Borel σ -algebra of a metric space X and by $\text{supp}(\mu)$ the support of a Borel probability measure μ . The notation δ_x stands for the Dirac measure at a point x . If (Ω, \mathcal{F}, P) is a probability space, the *Shannon entropy* of a finite or countably infinite measurable partition \mathcal{A} of Ω is defined as

$$H_P(\mathcal{A}) := - \sum_{A \in \mathcal{A}} P(A) \log P(A).$$

For two partitions \mathcal{A} and \mathcal{B} , the conditional entropy of \mathcal{A} given \mathcal{B} is defined by

$$H_\mu(\mathcal{A}|\mathcal{B}) := \sum_{B \in \mathcal{B}} \mu(B) H_{\mu_B}(\mathcal{A}),$$

where $\mu_B(\cdot) := \mu(\cdot \cap B)/\mu(B)$.

Let A be a real $n \times m$ matrix. Then $\text{rk } A$ denotes the rank of A . If A is a square matrix, we let $\text{spec}(A)$ denote the spectrum of A . If A is a continuous linear operator between normed vector spaces, we write $\|A\|$ for its operator norm.

2.2 Some concepts from dynamical systems

We recall some concepts from the theory of dynamical systems.

For a homeomorphism $T : X \rightarrow X$ on a compact metric space (X, d) , we use the following notions:

- A point $x \in X$ is called *periodic* if there exists $n \in \mathbb{Z}_{>0}$ such that $T^n(x) = x$. Any n with this property is called a *period* of x . The smallest such n is called the *minimal period*.
- T is called *topologically transitive* if for every pair of nonempty open sets $U, V \subset X$ there exists an $n \in \mathbb{Z}_{>0}$ such that $T^{-n}(U) \cap V \neq \emptyset$. If X has no isolated points, this is equivalent to the existence of a point $x_0 \in X$ whose forward orbit $\{x_0, T(x_0), T^2(x_0), \dots\}$ is dense in X .
- For $\varepsilon > 0$, an ε -chain for T is a finite sequence of points x_0, x_1, \dots, x_n in X with $n \in \mathbb{Z}_{>0}$, satisfying $d(T(x_i), x_{i+1}) \leq \varepsilon$ for $i = 0, 1, \dots, n-1$.
- A set $A \subset X$ is *chain transitive* if for all $\varepsilon > 0$ and $x, y \in A$ there exists an ε -chain of the form $x = x_0, x_1, \dots, x_n = y$, where the intermediate points x_1, \dots, x_{n-1} are not necessarily elements of A . If we can always choose the intermediate points in A , we call A *internally chain transitive*. We say that T is *chain transitive* if X is a chain transitive set. The maximal chain transitive sets of T are called the *chain components*.
- A point $x \in X$ is called *chain recurrent* if for every $\varepsilon > 0$ there exists an ε -chain from x to x . The *chain recurrent set* of T is the set of all chain recurrent points.
- A subset $A \subset X$ is called *invariant* if $T(A) = A$. A closed invariant set A is called *isolated invariant* if there is a neighborhood N of A (called an *isolating neighborhood*) such that $T^n(x) \in N$ for all $n \in \mathbb{Z}$ implies $x \in A$.
- An *additive cocycle* over (X, T) is a mapping $\alpha : \mathbb{Z}_+ \times X \rightarrow \mathbb{R}$, $(n, x) \mapsto \alpha_n(x)$, satisfying $\alpha_{n+m}(x) = \alpha_n(x) + \alpha_m(T^n(x))$ for all $n, m \in \mathbb{Z}_+$. If only the inequality $\alpha_{n+m}(x) \leq \alpha_n(x) + \alpha_m(T^n(x))$ holds, we call α a *subadditive cocycle* over (X, T) .

- The *nonwandering set* of T is defined as the set of all $x \in X$ such that for every neighborhood N of x there is an $n \in \mathbb{Z}_{>0}$ with $T^n(N) \cap N \neq \emptyset$.

Next, we recall the concept of a random dynamical system. Let (Ω, \mathcal{F}, P) be a complete probability space and $\theta : \Omega \rightarrow \Omega$, $\omega \mapsto \theta\omega$, a P -preserving invertible map. Further, let (X, \mathcal{B}) be a Polish space and $\mathcal{E} \subset \Omega \times X$ a measurable subset. A *bundle random dynamical system (bundle RDS)* over $(\Omega, \mathcal{F}, P, \theta)$ is generated by mappings $f_\omega : \mathcal{E}_\omega \rightarrow \mathcal{E}_{\theta\omega}$ so that the map $(\omega, x) \mapsto f_\omega(x)$ is measurable, where $\mathcal{E}_\omega := \{x \in X : (\omega, x) \in \mathcal{E}\}$ (the ω -fiber of \mathcal{E}). The map $\Phi : \mathcal{E} \rightarrow \mathcal{E}$ defined by $\Phi(\omega, x) := (\theta\omega, f_\omega(x))$ is called the *skew-product transformation* of the bundle RDS. If $\mathcal{E} = \Omega \times X$, we simply speak of a *random dynamical system (RDS)*. An *invariant measure* μ of the bundle RDS is a probability measure on \mathcal{E} with marginal P on Ω , invariant under Φ . Any such μ disintegrates as $d\mu(\omega, x) = d\mu_\omega(x)dP(\omega)$ with P -almost everywhere defined sample measures μ_ω on \mathcal{E}_ω . The invariance of μ can also be expressed by the identities $(f_\omega)_*\mu_\omega = \mu_{\theta\omega}$ for P -almost all $\omega \in \Omega$. We write $\mathcal{M}_P(\Phi; \mathcal{E})$ for the set of all invariant probability measures of a given bundle RDS. For the entropy theory of bundle RDS, we refer the reader to [46, Sec. 1.1].

3 Hyperbolic sets of control systems

3.1 Setup

We study a discrete-time control system

$$\Sigma : \quad x_{t+1} = f(x_t, u_t) \tag{8}$$

with a right-hand side $f : M \times U \rightarrow M$ satisfying the following assumptions:

- M is a smooth d -dimensional manifold for some $d \in \mathbb{Z}_{>0}$.
- U is a compact and connected metrizable space.
- The map $f_u : M \rightarrow M$, defined by $f_u(x) := f(x, u)$, is a C^1 -diffeomorphism for every $u \in U$, and its derivative $Df_u(x)$ depends (jointly) continuously on (u, x) .
- Both f and $(x, u) \mapsto f_u^{-1}(x)$ are continuous maps on $M \times U$.

The space of *admissible control sequences* for Σ is defined by

$$\mathcal{U} := U^{\mathbb{Z}} = \{u = (u_t)_{t \in \mathbb{Z}} : u_t \in U, \forall t \in \mathbb{Z}\}. \tag{9}$$

The following facts are well-known and can be found in standard textbooks on set-theoretic topology.

3.1 Facts: Equipped with the product topology induced by the topology of U , the space \mathcal{U} is compact, connected and metrizable. If d_U is a metric on U , an induced product metric on \mathcal{U} is given by

$$d_{\mathcal{U}}(u, v) := \sum_{t \in \mathbb{Z}} \frac{1}{2^{|t|}} d_U(u_t, v_t).$$

Sometimes, we will use the notation $d_{\mathcal{U} \times M}$ for a product metric on $\mathcal{U} \times M$, e.g.

$$d_{\mathcal{U} \times M}((u, x), (v, y)) = d_{\mathcal{U}}(u, v) + d(x, y),$$

where d is a given metric on M .

The *left shift operator* $\theta : \mathcal{U} \rightarrow \mathcal{U}$ is defined by

$$(\theta u)_t := u_{t+1} \quad \text{for all } u = (u_t)_{t \in \mathbb{Z}} \in \mathcal{U}.$$

The *transition map* $\varphi : \mathbb{Z} \times M \times \mathcal{U} \rightarrow M$ associated with Σ is given by

$$\varphi(t, x, u) := \begin{cases} f_{u_{t-1}} \circ \cdots \circ f_{u_1} \circ f_{u_0}(x) & \text{if } t > 0, \\ x & \text{if } t = 0, \\ f_{u_t}^{-1} \circ \cdots \circ f_{u_{-2}}^{-1} \circ f_{u_{-1}}^{-1}(x) & \text{if } t < 0. \end{cases}$$

Together, θ and φ constitute a *skew-product system* called the *control flow* of Σ :³

$$\Phi : \mathbb{Z} \times \mathcal{U} \times M \rightarrow \mathcal{U} \times M, \quad (t, u, x) \mapsto \Phi_t(u, x) := (\theta^t u, \varphi(t, x, u)).$$

We also introduce the notation $\varphi_{t,u}(x) := \varphi(t, x, u)$, $\varphi_{t,u} : M \rightarrow M$, for each pair $(t, u) \in \mathbb{Z} \times \mathcal{U}$. Obviously, $\varphi_{t,u}$ is a C^1 -diffeomorphism.

3.2 Proposition: The maps θ , φ and Φ satisfy the following properties:

- (a) $\theta : \mathcal{U} \rightarrow \mathcal{U}$ is a homeomorphism.
- (b) $\varphi(t, \cdot, \cdot) : M \times \mathcal{U} \rightarrow M$, $(x, u) \mapsto \varphi(t, x, u)$, is continuous for every $t \in \mathbb{Z}$.
- (c) φ is a cocycle over the base (\mathcal{U}, θ) , i.e., it satisfies
 - (i) $\varphi(0, x, u) = x$ for all $(u, x) \in \mathcal{U} \times M$,
 - (ii) $\varphi(t + s, x, u) = \varphi(s, \varphi(t, x, u), \theta^t u)$ for all $t, s \in \mathbb{Z}$, $(u, x) \in \mathcal{U} \times M$.
- (d) Φ is a dynamical system on $\mathcal{U} \times M$, i.e., $\Phi_0(u, x) = (u, x)$ and $\Phi_{t+s}(u, x) = \Phi_s(\Phi_t(u, x))$ for all $t, s \in \mathbb{Z}$ and $(u, x) \in \mathcal{U} \times M$.
- (e) For each $(t, u) \in \mathbb{Z} \times \mathcal{U}$, the derivative of $\varphi_{t,u}$ depends continuously on $(u, x) \in \mathcal{U} \times M$.

³Although the word “flow” is typically used for continuous-time systems, we also use it here for lack of a better name.

(f) The periodic points of θ are dense in \mathcal{U} and θ is chain transitive.

All statements except for the very last one follow easily from the assumptions. Hence, we only remark that the chain transitivity of θ follows from the fact that the periodic points are dense combined with the connectedness of \mathcal{U} . Indeed, every periodic point is trivially chain recurrent. Since the chain recurrent set is closed, it thus equals \mathcal{U} . By [18, Prop. 3.3.5(iii)], a closed set which is chain recurrent and connected is chain transitive.

Observe that the cocycle property (item (c) above) implies that the inverse of $\varphi_{t,u}$ is given by $\varphi_{t,u}^{-1} = \varphi_{-t, \theta^t u}$.

Since $(\Phi_t)_{t \in \mathbb{Z}}$ is a dynamical system on $\mathcal{U} \times M$, we have $\Phi_t = (\Phi_1)^t$ for all $t \in \mathbb{Z}$, i.e., the system is completely determined by its time-1 map. This justifies to write Φ not only for the sequence $(\Phi_t)_{t \in \mathbb{Z}}$ but also for the time-1 map Φ_1 .

Sometimes, we also need to require a higher regularity of the system with respect to x . We say that the system Σ is of *regularity class C^2* if for each $u \in U$ the map f_u is a C^2 -diffeomorphism with first and second derivatives depending continuously on $(u, x) \in U \times M$.

We call a set $Q \subset M$ *all-time controlled invariant* if for every $x \in Q$ there is a $u \in \mathcal{U}$ such that $\varphi(\mathbb{Z}, x, u) \subset Q$. To such Q , we associate its *all-time lift*

$$L(Q) := \{(u, x) \in \mathcal{U} \times M : \varphi(\mathbb{Z}, x, u) \subset Q\}.$$

It is easy to see that $L(Q)$ is an invariant set of the control flow Φ which is compact if and only if Q is compact. We define the *u-fibers* of Q by

$$Q(u) := \{x \in M : \varphi(\mathbb{Z}, x, u) \subset Q\}, \quad u \in \mathcal{U}.$$

The following properties of the u -fibers are easy to derive:

- Each u -fiber $Q(u)$ is compact (but not necessarily nonempty).
- For all $t \in \mathbb{Z}$ and $u \in \mathcal{U}$, the following relation holds:

$$\varphi_{t,u}(Q(u)) = Q(\theta^t u).$$

- The set $\mathcal{U}_Q := \{u \in \mathcal{U} : Q(u) \neq \emptyset\}$ is compact and θ -invariant.
- The set-valued map $u \mapsto Q(u)$, defined on \mathcal{U}_Q , is upper semicontinuous (but not necessarily lower semicontinuous).

The map $u \mapsto Q(u)$ as defined above will be called the *fiber map* of Q .

Now we introduce the notion of uniform hyperbolicity which requires an additional structure on the smooth manifold M , namely a Riemannian metric. However, choosing a different metric only results in the change of the constant c in condition (H2) below, and hence the notion of uniform hyperbolicity is metric-independent (see also Proposition B.1 in the Appendix).

3.3 Definition: A nonempty compact all-time controlled invariant set Q is called uniformly hyperbolic (or simply hyperbolic) if for every $(u, x) \in L(Q)$ there is a decomposition

$$T_x M = E^-(u, x) \oplus E^+(u, x)$$

as a direct sum, satisfying the following properties:

(H1) The decomposition is invariant in the sense that

$$D\varphi_{t,u}(x)E^\pm(u, x) = E^\pm(\Phi_t(u, x)) \quad \text{for all } (u, x) \in L(Q), t \in \mathbb{Z}.$$

(H2) There are constants $c \geq 1$ and $\lambda \in (0, 1)$ such that for all $(u, x) \in L(Q)$ and $t \in \mathbb{Z}_+$ the following inequalities hold:

$$\begin{aligned} |D\varphi_{t,u}(x)v| &\leq c\lambda^t|v| \quad \text{for all } v \in E^-(u, x), \\ |D\varphi_{-t,u}(x)v| &\leq c\lambda^t|v| \quad \text{for all } v \in E^+(u, x). \end{aligned}$$

(H3) The dimensions of the subspaces $E^-(u, x)$ and $E^+(u, x)$ are constant over $(u, x) \in L(Q)$.

An easy consequence of the hypotheses (H1) and (H2) is that the subspaces $E^\pm(u, x)$ vary continuously with (u, x) . This continuity statement can be expressed, e.g., in terms of the projections $\pi_{u,x}^\pm : T_x M \rightarrow E^\pm(u, x)$ along the respective complementary subspace, whose components in each coordinate chart are continuous functions of (u, x) . An implication of the continuity is that, also without hypothesis (H3), the dimensions of $E^\pm(u, x)$ are locally constant. Actually, our only reason to require (H3) is that we can avoid to include this as an extra assumption in many results that follow. For obvious reasons, we call $E^-(u, x)$ the *stable subspace* and $E^+(u, x)$ the *unstable subspace* at (u, x) , respectively. The case that one of the subspaces $E^\pm(u, x)$ is zero-dimensional is possible and we do not exclude it from the definition. Some elementary properties of hyperbolic sets are proved in Section B of the Appendix.

If the space U of control values is a singleton $\{u\}$, Definition 3.3 reduces to the classical definition of a uniformly hyperbolic set for the diffeomorphism f_u . In this case, we speak of a *classical hyperbolic set*.

A fundamental quantity used to describe the minimal required data rate above which a set can be rendered invariant by an appropriately designed coder-controller pair is known by the name *invariance entropy*. We now recall its definition. A pair (K, Q) of sets $K \subset Q \subset M$ is called an *admissible pair* (for Σ) if for each $x \in K$ there is a $u \in \mathcal{U}$ with $\varphi(\mathbb{Z}_+, x, u) \subset Q$. A set $\mathcal{S} \subset \mathcal{U}$ is called (τ, K, Q) -spanning for some $\tau \in \mathbb{Z}_{>0}$ if for every $x \in K$ there is a $u \in \mathcal{S}$ with $\varphi(t, x, u) \in Q$ for all $t \in [0; \tau)$. The *invariance entropy* of (K, Q) is defined by

$$h_{\text{inv}}(K, Q) := \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log r_{\text{inv}}(\tau, K, Q), \quad (10)$$

where $r_{\text{inv}}(\tau, K, Q)$ denotes the minimal cardinality of a (τ, K, Q) -spanning set. Associated data-rate theorems that characterize the smallest average data rate required to make Q invariant in terms of $h_{\text{inv}}(K, Q)$ can be found in [43, Thm. 2.4] and [23, Thm. 8]. If $K = Q$, then the lim sup in (10) is a limit and we also write $h_{\text{inv}}(Q)$ instead of $h_{\text{inv}}(Q, Q)$.

3.2 Tools from the hyperbolic theory

In this subsection, we present the main results from the hyperbolic theory that we use in our proofs. Throughout, we assume that a Riemannian metric on M is fixed.

First, we introduce the concepts of pseudo-orbits and shadowing. Consider the control system Σ . A two-sided sequence $(u^t, x_t)_{t \in \mathbb{Z}}$ in $\mathcal{U} \times M$ is called an α -pseudo-orbit for some $\alpha > 0$ if⁴

$$u^{t+1} = \theta u^t \quad \text{and} \quad d(\varphi_{1, u^t}(x_t), x_{t+1}) \leq \alpha \quad \text{for all } t \in \mathbb{Z}.$$

Hence, any pseudo-orbit is a real orbit in the u -component, but not necessarily in the x -component where we allow jumps of size at most α in each step of time. We say that a Φ -orbit $(\theta^t u, \varphi_{t, u}(x))_{t \in \mathbb{Z}}$ β -shadows a pseudo-orbit $(u^t, x_t)_{t \in \mathbb{Z}}$ if

$$u = u^0 \quad \text{and} \quad d(\varphi_{t, u}(x), x_t) \leq \beta \quad \text{for all } t \in \mathbb{Z}.$$

The shadowing lemma roughly says that in a small neighborhood of a hyperbolic set, every α -pseudo-orbit is β -shadowed by a real orbit if $\alpha = \alpha(\beta)$ is chosen small enough. The complete and precise statement is as follows. A proof can be found in Meyer & Zhang [51].

3.4 Theorem: *Let Q be a hyperbolic set of Σ . Then there is a neighborhood $\mathcal{N} \subset \mathcal{U} \times M$ of $L(Q)$ such that the following holds:*

- (a) *For every $\beta > 0$, there is an $\alpha > 0$ such that every α -pseudo-orbit in \mathcal{N} is β -shadowed by an orbit.*
- (b) *There is $\beta_0 > 0$ such that for every $\beta \in (0, \beta_0)$ the β -shadowing orbit in (a) is unique.*
- (c) *If $L(Q)$ is an isolated invariant set of the control flow, then the unique β -shadowing orbit in (b) is completely contained in $L(Q)$.*

Another extremely useful property of hyperbolic sets is called *expansivity*. It is an easy consequence of the stable manifold theorem, see [51, Thm. 2.1].

⁴To avoid abuse of notation, we use a superscript for the u -component, because u_t already denotes the t -th component of the sequence u .

3.5 Theorem: Let Q be a hyperbolic set of Σ . Then there exists $\delta > 0$ such that for all $(u, x) \in L(Q)$ and $y \in M$ the following implication holds: If $d(\varphi(t, x, u), \varphi(t, y, u)) \leq \delta$ for all $t \in \mathbb{Z}$, then $x = y$. Any constant δ with this property is called an *expansivity constant*.

For all $u \in \mathcal{U}$, $x \in M$, $\varepsilon > 0$ and $\tau \in \mathbb{Z}_{>0}$, we introduce the *Bowen-ball*

$$B_\varepsilon^{u, \tau}(x) := \{y \in M : d(\varphi(t, x, u), \varphi(t, y, u)) \leq \varepsilon \text{ for } t = 0, 1, \dots, \tau - 1\}.$$

We call $B_\varepsilon^{u, \tau}(x)$ the *Bowen-ball of order τ and radius ε , centered at x and associated with the control u* . Observe that this is the usual closed ε -ball in the metric

$$d^{u, \tau}(x, y) := \max_{t \in [0; \tau)} d(\varphi(t, x, u), \varphi(t, y, u)),$$

which is compatible with the topology of M .

The (*Bowen-Ruelle*) *volume lemma* provides asymptotically precise estimates for the volumes of Bowen-balls centered in hyperbolic sets. It requires a little more regularity in the state variable. A detailed proof can be found in [25].

3.6 Theorem: Assume that Σ is of regularity class C^2 and let Q be a hyperbolic set of Σ . Then, for every sufficiently small $\varepsilon > 0$, the following estimates hold for all $(u, x) \in L(Q)$ and $\tau \in \mathbb{Z}_{>0}$ with some constant $C_\varepsilon \geq 1$:

$$C_\varepsilon^{-1} |\det D\varphi_{\tau, u}(x)|_{E^+(u, x)}|^{-1} \leq \text{vol}(B_\varepsilon^{u, \tau}(x)) \leq C_\varepsilon |\det D\varphi_{\tau, u}(x)|_{E^+(u, x)}|^{-1}.$$

Since the determinant that appears in the above estimates will be used frequently, we introduce an abbreviation for it:

$$J^+ \varphi_{\tau, u}(x) := |\det D\varphi_{\tau, u}(x)|_{E^+(u, x)} : E^+(u, x) \rightarrow E^+(\Phi_\tau(u, x))|.$$

We also call this function the *unstable determinant*. It is easy to see that

- $(u, x) \mapsto J^+ \varphi_{\tau, u}(x)$ is continuous for every $\tau \in \mathbb{Z}$ and
- $J^+ \varphi_{\tau_1 + \tau_2, u}(x) = J^+ \varphi_{\tau_1, u}(x) \cdot J^+ \varphi_{\tau_2, \theta^{\tau_1} u}(\varphi_{\tau_1, u}(x))$ for all $\tau_1, \tau_2 \in \mathbb{Z}$ and $(u, x) \in L(Q)$. That is, $\log J^+ \varphi$ is an additive cocycle over $(\Phi|_{L(Q)}, L(Q))$.

3.3 Properties of hyperbolic sets

In this subsection, we first prove the following theorem on the structure of hyperbolic sets. Then, we provide a way of constructing hyperbolic sets via small “control-perturbations” of diffeomorphisms. Finally, we study controllability properties of these sets.

3.7 Theorem: *Let Q be a hyperbolic set of the control system Σ and assume that its all-time lift $L(Q)$ is an isolated invariant set of the control flow. Then all fibers $Q(u)$, $u \in \mathcal{U}$, are nonempty and homeomorphic to each other.*

Proof: The shadowing lemma yields a neighborhood $\mathcal{N} \subset \mathcal{U} \times M$ of $L(Q)$, a $\beta > 0$ and an $\alpha = \alpha(\beta) > 0$ so that every α -pseudo-orbit in \mathcal{N} is β -shadowed by a unique orbit in $L(Q)$. Let $\varepsilon = \varepsilon(\alpha) > 0$ be small enough so that $N_{3\varepsilon}(L(Q)) \subset \mathcal{N}$ and so that for all $u, v \in U$ and $x \in Q$ we have

$$d_U(u, v) \leq \varepsilon \quad \Rightarrow \quad d(f_u(x), f_v(x)) \leq \alpha.$$

This is possible by compactness of $L(Q)$ and uniform continuity of f on the compact set $Q \times U$, respectively.

In the following, we will use the metric

$$d_\infty(u, v) := \sup_{t \in \mathbb{Z}} d_U(u_t, v_t)$$

on \mathcal{U} , which in general is not compatible with the product topology. We claim that $d_\infty(u, v) \leq \varepsilon$ implies that $Q(u)$ and $Q(v)$ are homeomorphic. If $Q(u)$ and $Q(v)$ are both empty, there is nothing to show. Hence, let us assume that $Q(u) \neq \emptyset$. Then choose $x \in Q(u)$ arbitrarily and consider the two-sided sequence $x_t := \varphi(t, x, u)$, $t \in \mathbb{Z}$, which lies in Q and, by the choice of ε , satisfies

$$d(\varphi(1, x_t, \theta^t v), x_{t+1}) = d(f_{v_t}(x_t), f_{u_t}(x_t)) \leq \alpha \quad \text{for all } t \in \mathbb{Z}.$$

Hence, the sequence $(\theta^t v, x_t)_{t \in \mathbb{Z}}$ is an α -pseudo-orbit which is 3ε -close to the orbit $(\theta^t u, x_t)_{t \in \mathbb{Z}}$ that is completely contained in $L(Q)$. (A simple computation shows that $d_\infty(u, v) \leq \varepsilon$ implies $d_{\mathcal{U}}(\theta^t u, \theta^t v) \leq 3\varepsilon$ for all $t \in \mathbb{Z}$.) By the choice of ε , there exists a unique orbit $(\theta^t v, \varphi(t, y, v))_{t \in \mathbb{Z}}$ in $L(Q)$ which β -shadows $(\theta^t v, x_t)_{t \in \mathbb{Z}}$, i.e., $y \in Q(v)$ and

$$d(\varphi(t, y, v), \varphi(t, x, u)) \leq \beta \quad \text{for all } t \in \mathbb{Z}.$$

We can thus define the mapping

$$h_{uv} : Q(u) \rightarrow Q(v), \quad x \mapsto y$$

that sends a point $x \in Q(u)$ to the unique point $y \in Q(v)$ given by the shadowing lemma. Since the roles of u and v can be interchanged, we also have a mapping $h_{vu} : Q(v) \rightarrow Q(u)$, defined analogously, which must be the inverse of h_{uv} by the uniqueness of shadowing orbits. It remains to prove the continuity of h_{uv} . To this end, consider a sequence $x_k \rightarrow x$ in $Q(u)$ and let $y_k := h_{uv}(x_k)$, $y := h_{uv}(x)$. Then $\varphi_{t,u}(x_k) \rightarrow \varphi_{t,u}(x)$ for each $t \in \mathbb{Z}$. Let $\delta > 0$ be an expansivity constant according to Theorem 3.5. We prove the continuity of h_{uv} under the assumption that $\beta \leq \delta/3$. For every $t \in \mathbb{Z}$, let $k_0(t)$ be large enough so that

$$d(\varphi_{t,u}(x_k), \varphi_{t,u}(x)) \leq \frac{\delta}{3} \quad \text{for all } k \geq k_0(t).$$

Then for every $t \in \mathbb{Z}$ and $k \geq k_0(t)$ we obtain

$$\begin{aligned} d(\varphi_{t,v}(y_k), \varphi_{t,v}(y)) &\leq d(\varphi_{t,v}(y_k), \varphi_{t,u}(x_k)) + d(\varphi_{t,u}(x_k), \varphi_{t,u}(x)) \\ &\quad + d(\varphi_{t,u}(x), \varphi_{t,v}(y)) \leq 2\beta + \frac{\delta}{3} \leq \delta. \end{aligned}$$

Hence, for any limit point $y_* \in Q(v)$ of the sequence (y_k) , it follows that

$$d(\varphi_{t,v}(y_*), \varphi_{t,v}(y)) \leq \delta \quad \text{for all } t \in \mathbb{Z},$$

implying $y_* = y$ by the choice of δ . We thus obtain $y_k \rightarrow y$ which proves the continuity of h_{uv} .

We have shown that up to homeomorphisms $Q(u)$ is locally constant on the metric space $(U^{\mathbb{Z}}, d_{\infty})$. Since U is connected, Lemma A.1 implies that $(U^{\mathbb{Z}}, d_{\infty})$ is connected as well. It thus follows that all u -fibers are homeomorphic to each other. In particular, this implies that none of them is empty. \square

3.8 Remark: For a constant control $u \in \mathcal{U}$, the fiber $Q(u)$ is a compact hyperbolic set of the diffeomorphism f_u . Assuming a little more regularity, namely that f_u is a $C^{1+\alpha}$ -diffeomorphism for some $\alpha > 0$, there are only two alternatives for the fiber $Q(u)$: either it coincides with the whole state space M (in which case M is compact and $Q = M$) or it has Lebesgue measure zero (see [10, Cor. 5.7]). It is unclear if the same is true for every u -fiber. The homeomorphisms h_{uv} constructed in the above proof can only be expected to be Hölder continuous which does not allow for a statement on the Lebesgue measure.

In addition to the above result, we would like to prove that the fiber map $u \mapsto Q(u)$ is continuous with respect to the Hausdorff metric on the space of nonempty closed subsets of Q . However, in the general case, it is completely unclear how to do this or whether it is true. Instead, we only prove it for hyperbolic sets that are sufficiently “small”. At the same time, we provide a method to construct examples of hyperbolic sets.

Consider the control system Σ , fix a control value $u^0 \in U$ and assume that the following holds:

- The diffeomorphism $f_{u^0} : M \rightarrow M$ has a compact isolated invariant set $\Lambda \subset M$ which is hyperbolic (in the classical sense).
- The metric space U is locally connected at u^0 . That is, every neighborhood of u^0 contains a connected neighborhood.

The following lemma, which is standard in the hyperbolic theory, will turn out to be useful (see also [48, Lem. 1.2]).

3.9 Lemma: *Under the given assumptions, there exist a neighborhood $U_0 \subset U$ of u^0 , a neighborhood $N \subset M$ of Λ , and numbers $\rho_0 > 0$, $C_0 > 0$ and $\alpha_0 \in (0, 1)$ such that the following holds for every $T \in \mathbb{Z}_+$: If $u \in U_0^{\mathbb{Z}}$, $x, y \in N$ and $\varphi(t, x, u), \varphi(t, y, u) \in N$ with $d(\varphi(t, x, u), \varphi(t, y, u)) \leq \rho_0$ for all t with $|t| \leq T$, then $d(x, y) \leq C_0 \alpha_0^T$.*

Proof: Let $T_\Lambda M = E^- \oplus E^+$ be the hyperbolic splitting on Λ . We choose the neighborhoods U_0 and N such that the following holds:

- The hyperbolic splitting on Λ can be extended continuously⁵ to the neighborhood N and there is $C > 0$ such that

$$\|v\|_0 := \max\{|v^-|, |v^+|\} \leq C|v|$$

whenever $v \in T_x M$, $x \in N$ and $v^\pm \in E_x^\pm$ with $v = v^- + v^+$. Let $\lambda_0 \in (0, 1)$ be the hyperbolic constant on Λ and assume that the constant c equals 1 (i.e., contraction is seen in one step of time), which can always be achieved by using an adapted Riemannian metric (see, e.g., [9, Lem. 3.1]).

- There are $\rho_0, a_0, \varepsilon_0 > 0$ satisfying $\lambda_0 < a_0 < 1$ and $0 < \varepsilon_0 < \min\{\frac{1}{2}(1 - a_0), \frac{1}{2}(a_0^{-1} - 1)\}$ such that the following holds: If $x \in N$ and $u \in U_0$ with $\varphi_{1,u}(x) \in N$, then

$$\tilde{\varphi}_{u,x} := \exp_{\varphi_{1,u}(x)}^{-1} \circ \varphi_{1,u} \circ \exp_x : \{v \in T_x M : |v| \leq \rho_0\} \rightarrow T_{\varphi_{1,u}(x)} M$$

is well-defined and, writing

$$D\tilde{\varphi}_{u,x}(0) = \begin{pmatrix} A_{u,x}^{--} & A_{u,x}^{-+} \\ A_{u,x}^{+-} & A_{u,x}^{++} \end{pmatrix} : E_x^- \oplus E_x^+ \rightarrow E_{\varphi_{1,u}(x)}^- \oplus E_{\varphi_{1,u}(x)}^+,$$

we can express $\tilde{\varphi}_{u,x}$ as

$$\tilde{\varphi}_{u,x}(\cdot) = \begin{pmatrix} A_{u,x}^{--} & 0 \\ 0 & A_{u,x}^{++} \end{pmatrix} + R_{u,x}(\cdot),$$

where $\|A_{u,x}^{--}\| \leq a_0$, $\|(A_{u,x}^{++})^{-1}\| \leq a_0$ and $R_{u,x}$ is a Lipschitz map whose Lipschitz constant with respect to $\|\cdot\|_0$ is not bigger than ε_0 . Indeed, this follows from the fact that the derivative of $R_{u,x}(\cdot)$ at the origin is determined by $A_{u,x}^{-+}$ and $A_{u,x}^{+-}$ which have arbitrarily small norms if we choose ρ_0 , N and U_0 small enough. Moreover, we make our choices so that the map

$$\tilde{\varphi}_{u,x}^- := \exp_x^{-1} \circ \varphi_{-1,\theta u} \circ \exp_{\varphi_{1,u}(x)} : \{v \in T_{\varphi_{1,u}(x)} M : |v| \leq \rho_0\} \rightarrow T_x M$$

has analogous properties.

⁵It is a standard fact in the theory of hyperbolic systems that such an extension always exists. However, the extended splitting might no longer be invariant.

Now, if $u \in U_0^{\mathbb{Z}}$ and $x, y \in N$ are as in the formulation of the lemma, let us write $v := \exp_x^{-1}(y) = v^- + v^+ \in E_x^- \oplus E_x^+$ and assume without loss of generality that $|v^+| \geq |v^-|$. Then we can show that

$$(a_0^{-1} - \varepsilon_0)^T \|v\|_0 \leq \|\exp_{\varphi_{T,u}(x)}^{-1}(\varphi_{T,u}(y))\|_0 \leq C\rho_0.$$

The second inequality follows from

$$\begin{aligned} \|\exp_{\varphi_{T,u}(x)}^{-1}(\varphi_{T,u}(y))\|_0 &\leq C|\exp_{\varphi_{T,u}(x)}^{-1}(\varphi_{T,u}(y))| \\ &= Cd(\varphi_{T,u}(x), \varphi_{T,u}(y)) \leq C\rho_0. \end{aligned}$$

The first one can be shown as follows. Using that $|v^+| \geq |v^-|$, we obtain

$$\begin{aligned} |\tilde{\varphi}_{u,x}(v)^+| &= |A_{u,x}^{++}v^+ + R_{u,x}(v)^+| \geq |A_{u,x}^{++}v^+| - |R_{u,x}(v)^+| \\ &\geq a_0^{-1}|v^+| - \|R_{u,x}(v)\|_0 \geq a_0^{-1}|v^+| - \varepsilon_0\|v\|_0 = (a_0^{-1} - \varepsilon_0)|v^+|. \end{aligned}$$

Similarly,

$$\begin{aligned} |\tilde{\varphi}_{u,x}(v)^-| &= |A_{u,x}^{--}v^- + R_{u,x}(v)^-| \leq |A_{u,x}^{--}v^-| + |R_{u,x}(v)^-| \\ &\leq a_0|v^-| + \|R_{u,x}(v)\|_0 \leq a_0|v^-| + \varepsilon_0|v^+| \leq (a_0 + \varepsilon_0)|v^+|. \end{aligned}$$

This implies

$$\frac{|\tilde{\varphi}_{u,x}(v)^+|}{|\tilde{\varphi}_{u,x}(v)^-|} \geq \frac{a_0^{-1} - \varepsilon_0}{a_0 + \varepsilon_0} > 1.$$

Hence, we can repeat these arguments and obtain the claimed inequality inductively.⁶ These inequalities imply

$$\begin{aligned} d(x, y) = |v| &= |v^- + v^+| \leq 2 \max\{|v^-|, |v^+|\} \\ &= 2\|v\|_0 \leq 2C\rho_0(a_0^{-1} - \varepsilon_0)^{-T}. \end{aligned}$$

Hence, the statement of the lemma holds with $C_0 := 2C\rho_0$ and $\alpha_0 := (a_0^{-1} - \varepsilon_0)^{-1}$, where we observe that $a_0^{-1} - \varepsilon_0 > a_0^{-1} - (a_0^{-1} - 1)/2 = (a_0^{-1} + 1)/2 > 1$. \square

The next proposition describes the dynamics of the control system that we obtain by restricting the control values to a small neighborhood of u^0 , when we also consider a small neighborhood of Λ . Essentially, this is [48, Thm. 1.1] (the corresponding result for RDS).

3.10 Proposition: *Consider the control system Σ under the given assumptions. Then there are $\beta_0 > 0$ and a compact, connected neighborhood $U_0 \subset U$ of u^0 such that the following holds:*

⁶In the case when $|v^-| \geq |v^+|$, the maps $\tilde{\varphi}_{u,x}^-$ come into play.

(a) For each $u \in \mathcal{U}_0 := U_0^{\mathbb{Z}}$ and each $x \in \Lambda$, there is a unique $x_u \in M$ with

$$d(\varphi(t, x, u^0), \varphi(t, x_u, u)) \leq \beta_0 \quad \text{for all } t \in \mathbb{Z}.$$

(b) For any $\beta \in (0, \beta_0)$, one can shrink U_0 so that (a) holds with β in place of β_0 .

(c) For every $u \in \mathcal{U}_0$, define $\Lambda_u := \{x_u : x \in \Lambda\}$ and $h_u : \Lambda \rightarrow \Lambda_u, x \mapsto x_u$. Then Λ_u is compact and h_u is a homeomorphism.

(d) The family of maps $\{h_u\}_{u \in \mathcal{U}_0}$ has the following properties:

- (i) $\varphi_{1,u}(\Lambda_u) = \Lambda_{\theta u}$ and $h_{\theta u} \circ \varphi_{1,u^0} = \varphi_{1,u} \circ h_u$ for all $u \in \mathcal{U}_0$.
- (ii) The family $\{h_u\}_{u \in \mathcal{U}_0}$ is equicontinuous. That is, for any $\varepsilon > 0$ there is $\delta > 0$ so that $d(x, y) < \delta$ implies $d(h_u(x), h_u(y)) < \varepsilon$ for all $x, y \in \Lambda$ and $u \in \mathcal{U}_0$. The analogous property holds for the family $\{h_u^{-1}\}_{u \in \mathcal{U}_0}$.
- (iii) The map $H : \mathcal{U}_0 \rightarrow C^0(\Lambda, M), u \mapsto h_u$, is continuous, when $C^0(\Lambda, M)$ is equipped with the topology of uniform convergence.

Proof: We put $\tilde{\Lambda} := \{u^0\} \times \Lambda \subset \mathcal{U} \times M$, where we regard u^0 as the constant sequence $(\dots, u^0, u^0, u^0, \dots) \in \mathcal{U}$, and observe that $\tilde{\Lambda}$ is a hyperbolic set of the control system Σ . Now we choose a neighborhood $\mathcal{N} \subset \mathcal{U} \times M$ of $\tilde{\Lambda}$ and a constant $\beta_0 > 0$ satisfying the following properties:

- $N_{3\beta_0}(\Lambda)$ is an isolating neighborhood of Λ for f_{u^0} .
- If $d(\varphi(t, x, u^0), \varphi(t, y, u^0)) \leq 2\beta_0$ for some $x, y \in \Lambda$ and all $t \in \mathbb{Z}$, then $x = y$, which is possible by expansivity on hyperbolic sets.
- There is $\alpha = \alpha(\beta_0)$ so that every α -pseudo-orbit of Σ , contained in \mathcal{N} , is uniquely β_0 -shadowed by an orbit.

Subsequently, we choose a compact, connected neighborhood U_0 of u^0 (where we use the assumption that U is locally connected at u^0) small enough so that

$$(u, x) \in U_0^{\mathbb{Z}} \times N_{2\beta_0}(\Lambda) \Rightarrow d(\varphi_{1,u}(x), \varphi_{1,u^0}(x)) \leq \alpha \text{ and } (u, x) \in \mathcal{N}. \quad (11)$$

This is possible by the uniform continuity of $\varphi(1, \cdot, \cdot)$ on the compact set $N_{2\beta_0}(\Lambda) \times U$. Now fix $u \in \mathcal{U}_0$ and $x \in \Lambda$. Defining $x_t := \varphi(t, x, u^0) = f_{u^0}^t(x)$, $t \in \mathbb{Z}$, we find that $(\theta^t u, x_t)_{t \in \mathbb{Z}}$ is an α -pseudo-orbit in \mathcal{N} , since

$$d(\varphi(1, x_t, \theta^t u), x_{t+1}) = d(\varphi_{1,\theta^t u}(x_t), \varphi_{1,u^0}(x_t)) \leq \alpha \quad \text{for all } t \in \mathbb{Z}.$$

Hence, there exists a unique point $x_u \in M$ such that

$$d(\varphi(t, x, u^0), \varphi(t, x_u, u)) \leq \beta_0 \quad \text{for all } t \in \mathbb{Z}.$$

This proves (a).

Statement (b) follows from item (b) of the shadowing lemma (Theorem 3.4).

Statement (c) is seen as follows. First, the compactness of Λ_u follows from the continuity of h_u established in (d)(ii). The invertibility of h_u follows from the choice of β_0 , since $h_u(x) = h_u(y)$ implies $d(\varphi(t, x, u^0), \varphi(t, y, u^0)) \leq 2\beta_0$ for all $t \in \mathbb{Z}$. Since any invertible and continuous map between compact metric spaces is a homeomorphism, (c) is proved.

It remains to prove (d). To prove (d)(i), pick $x_u = h_u(x) \in \Lambda_u$. Then $d(\varphi(t, x_u, u), \varphi(t, x, u^0)) \leq \beta_0$ for all $t \in \mathbb{Z}$. By the cocycle property of φ , this is equivalent to $d(\varphi(t, \varphi_{1,u}(x_u), \theta u), \varphi(t, \varphi_{1,u^0}(x), u^0)) \leq \beta_0$ for all $t \in \mathbb{Z}$. Hence, (a) implies that $\varphi_{1,u}(x_u) = h_{\theta u}(\varphi_{1,u^0}(x)) \in \Lambda_{\theta u}$.

To prove (d)(ii), we assume that U_0 is chosen small enough such that the statement of Lemma 3.9 holds with a neighborhood N of Λ and constants ρ_0, C_0, α_0 . Moreover, we choose β_0 small enough such that $\beta_0 \leq \rho_0/3$ and $N_{\beta_0}(\Lambda) \subset N$. Now, for a given $\varepsilon > 0$, we choose $T \in \mathbb{Z}_{>0}$ satisfying $C_0\alpha_0^T < \varepsilon$. We let further $\delta > 0$ be small enough so that $x, y \in \Lambda, d(x, y) < \delta$ implies

$$d(\varphi(t, x, u^0), \varphi(t, y, u^0)) \leq \beta_0 \quad \text{for all } |t| \leq T.$$

Then $d(x, y) < \delta$ and $-T \leq t \leq T$ implies

$$\begin{aligned} d(\varphi_{t,u}(h_u(x)), \varphi_{t,u}(h_u(y))) &\leq d(\varphi_{t,u}(h_u(x)), \varphi_{t,u^0}(x)) \\ &\quad + d(\varphi_{t,u^0}(x), \varphi_{t,u^0}(y)) + d(\varphi_{t,u^0}(y), \varphi_{t,u}(h_u(y))) \leq 3\beta_0 \leq \rho_0. \end{aligned}$$

Hence, Lemma 3.9 yields $d(h_u(x), h_u(y)) \leq C_0\alpha_0^T < \varepsilon$. The proof of equicontinuity of $\{h_u^{-1}\}$ follows the same lines. Here we need to assume that δ is chosen small enough so that $d(\varphi(t, x, u), \varphi(t, y, u)) \leq \beta_0$ for $|t| \leq T$ whenever $u \in \mathcal{U}_0, x, y \in \Lambda_u$ and $d(x, y) < \delta$. This is possible by the uniform continuity of $\varphi(t, \cdot, \cdot)$ on the compact set $N_{\beta_0}(\Lambda) \times \mathcal{U}_0$.

To prove (d)(iii), consider a sequence $u_k \rightarrow u$ in \mathcal{U}_0 . By (ii) and the Arzelà-Ascoli Theorem, every subsequence of $(h_{u_k})_{k \in \mathbb{Z}_{>0}}$ has a limit point. That is, there exists a homeomorphism $h : \Lambda \rightarrow h(\Lambda)$ so that the subsequence converges uniformly to h . If $h_{u_{k_n}} \rightarrow h$ as $n \rightarrow \infty$, then for every $t \in \mathbb{Z}$ and $x \in \Lambda$ we obtain

$$d(\varphi(t, h(x), u), \varphi(t, x, u^0)) = \lim_{n \rightarrow \infty} d(\varphi(t, h_{u_{k_n}}(x), u_{k_n}), \varphi(t, x, u^0)) \leq \beta_0.$$

By statement (a), this implies $h = h_u$. Hence, $h_{u_{k_n}}$ converges to h_u , proving that H is continuous. \square

Using the above proposition, we can show the existence of a hyperbolic set with isolated invariant lift for the given control system with restricted control range U_0 .

3.11 Theorem: *Given the conclusions of the Proposition 3.10, the control system*

$$\Sigma^0 : \quad x_{t+1} = f(x_t, u_t), \quad u \in \mathcal{U}_0 \quad (12)$$

has a hyperbolic set $Q \subset M$ with the following properties:

- (a) $Q(u) = \Lambda_u$ for all $u \in \mathcal{U}_0$.
- (b) The all-time lift $L(Q)$ is an isolated invariant set for the control flow of Σ^0 .
- (c) The fiber map $u \mapsto Q(u)$ is continuous when \mathcal{U}_0 is equipped with the product topology.

Proof: We define

$$Q := \{h_u(x) : x \in \Lambda, u \in \mathcal{U}_0\} = \bigcup_{u \in \mathcal{U}_0} \Lambda_u.$$

We first prove that Q is compact and all-time controlled invariant. Consider the map $\alpha : \mathcal{U}_0 \times \Lambda \rightarrow M$, $(u, x) \mapsto h_u(x)$. From the continuity of $u \mapsto h_u$, it easily follows that α is continuous implying that $Q = \alpha(\mathcal{U}_0 \times \Lambda)$ is compact. All-time controlled invariance follows from Proposition 3.10(d)(i), which implies $\varphi(t, h_u(x), u) = h_{\theta^t u}(\varphi(t, x, u^0))$ for all $t \in \mathbb{Z}$. With standard arguments from the hyperbolic theory of dynamical systems (see, e.g., [39, Prop. 6.4.6]), one can show that for each $(u, h_u(x)) \in Q$ there is a splitting

$$T_{h_u(x)}M = E_{u, h_u(x)}^- \oplus E_{u, h_u(x)}^+$$

which is invariant and uniformly hyperbolic (provided that all relevant constants and neighborhoods are chosen small enough).

Now we prove (a). It is clear that $\Lambda_u \subset Q(u)$ for all $u \in \mathcal{U}_0$. To prove the converse, take $x \in Q(u)$ and recall the notation introduced in the proof of Proposition 3.10(a). The sequence $x_t := \varphi(t, x, u)$, $t \in \mathbb{Z}$, then yields the α -pseudo-orbit $(u^0, x_t)_{t \in \mathbb{Z}}$ in \mathcal{N} by (11). Hence, there exists a unique $y \in M$ such that

$$d(\varphi(t, y, u^0), x_t) \leq \beta_0 \quad \text{for all } t \in \mathbb{Z}.$$

Since $N_{3\beta_0}(\Lambda)$ is an isolating neighborhood of Λ for f_{u^0} , this implies $y \in \Lambda$. Then, by the uniqueness of shadowing orbits, it follows that $x = h_u(y) \in \Lambda_u$.

To prove (b), we show that the open set $\mathcal{U}_0 \times N_{2\beta_0}^\circ(\Lambda)$ is an isolating neighborhood of $L(Q)$. Since every $x \in Q$ satisfies $\text{dist}(x, \Lambda) \leq \beta_0$ by definition, this is a neighborhood of $L(Q)$. If $(\theta^t u, \varphi(t, x, u)) \in \mathcal{U}_0 \times N_{2\beta_0}^\circ(\Lambda)$ for all $t \in \mathbb{Z}$, then (11) implies that $x_t := \varphi(t, x, u)$ satisfies $d(\varphi_{1, u^0}(x_t), x_{t+1}) \leq \alpha$ and $(\theta^t u, x_t) \in \mathcal{N}$ for all $t \in \mathbb{Z}$. Hence, there exists a unique $y \in M$ with $d(\varphi(t, y, u^0), \varphi(t, x, u)) \leq \beta_0$ for all $t \in \mathbb{Z}$. We thus have

$$\text{dist}(\varphi(t, y, u^0), \Lambda) \leq d(\varphi(t, y, u^0), \varphi(t, x, u)) + \text{dist}(\varphi(t, x, u), \Lambda) \leq 3\beta_0$$

for all $t \in \mathbb{Z}$. Since $N_{3\beta_0}(\Lambda)$ is an isolating neighborhood of Λ , it follows that $y \in \Lambda$, implying $x = h_u(y)$ and $(u, x) \in L(Q)$ as desired.

Finally, we prove (c). Since the fiber map is always upper semicontinuous, it remains to prove its lower semicontinuity. Let $u \in \mathcal{U}_0$ and $x \in Q(u)$. Consider

a sequence $u_k \rightarrow u$ in \mathcal{U}_0 . Since $Q(u) = \Lambda_u$, we have $x = h_u(x')$ for some $x' \in \Lambda$. By Proposition 3.10(d)(iii), we know that $h_{u_k}(x') \rightarrow h_u(x')$. Since $h_{u_k}(x') \in \Lambda_{u_k} = Q(u_k)$, we have proved the lower semicontinuity at u . \square

In the following, we study controllability properties on the set Q as constructed above. Moreover, we are interested in finding out under which conditions Q has nonempty interior.

We start with some definitions and a technical lemma. A *controlled ε -chain* from a point $x \in M$ to a point $y \in M$ consists of a finite sequence of points $x = x_0, x_1, \dots, x_r = y$, $r \geq 1$, and controls $u_0, u_1, \dots, u_{r-1} \in \mathcal{U}$ so that $d(\varphi(1, x_i, u_i), x_{i+1}) \leq \varepsilon$ for $i = 0, 1, \dots, r-1$. We say that *chain controllability* holds on a set $E \subset M$ if any two points $x, y \in E$ can be joined by a controlled ε -chain for every $\varepsilon > 0$. For more details on this concept, see [17, 66].

3.12 Lemma: *Assume that the restriction of f_{u^0} to the hyperbolic set Λ is topologically transitive. Then chain controllability holds on the set Q as constructed above. In particular, for any $x, y \in Q$ and $\varepsilon > 0$, a controlled ε -chain (x_i, u^i) from x to y can be chosen so that $(u^i, x_i) \in L(Q)$ for all i . Moreover, the all-time lift $L(Q)$ is internally chain transitive.*

Proof: We use that topological transitivity implies chain transitivity (easy to see).⁷ To prove the assertion, pick two points $x, y \in Q$ and write them as $x = h_u(x')$, $y = h_v(y')$ with $x', y' \in \Lambda$ and $u, v \in \mathcal{U}_0$. We now choose δ -chains of equal length r from u to v in \mathcal{U}_0 and from x' to y' in Λ , respectively. That is, we pick $x' = z'_0, z'_1, \dots, z'_r = y'$ in Λ and $u = w^0, w^1, \dots, w^r = v$ in \mathcal{U}_0 such that

$$d(f_{u^0}(z'_i), z'_{i+1}) \leq \delta \quad \text{and} \quad d_{\mathcal{U}}(\theta w^i, w^{i+1}) \leq \delta$$

for all $i \in [0; r)$. This is possible by chain transitivity of f_{u^0} on Λ and of θ on \mathcal{U}_0 , respectively (see Proposition 3.2(f) for the latter). The reason why we can choose the length r identical for both chains is that \mathcal{U}_0 contains fixed points. Indeed, we can let every δ -chain in \mathcal{U}_0 run through a fixed point and at this fixed point we can stop as long as we want to (introducing an arbitrary number of trivial jumps).

Now, we define

$$z_i := h_{w^i}(z'_i) \in Q(w^i), \quad i = 0, 1, \dots, r.$$

Observe that $z_0 = h_{w^0}(z'_0) = h_u(x') = x$ and $z_r = h_{w^r}(z'_r) = h_v(y') = y$. We claim that if $\delta = \delta(\varepsilon)$ is chosen small enough, then (z_i, w^i) , $i = 0, 1, \dots, r$, is a controlled ε -chain from x to y , i.e.,

$$d(\varphi(1, z_i, w^i), z_{i+1}) \leq \varepsilon, \quad i = 0, 1, \dots, r-1. \quad (13)$$

⁷In fact, from the shadowing lemma it follows that topological transitivity is equivalent to chain transitivity on an isolated invariant hyperbolic set of a diffeomorphism.

We can check this as follows:

$$\begin{aligned}
d(\varphi(1, z_i, w^i), z_{i+1}) &= d(\varphi_{1,w^i}(h_{w^i}(z'_i)), h_{w^{i+1}}(z'_{i+1})) \\
&= d(h_{\theta w^i}(\varphi_{1,u^0}(z'_i)), h_{w^{i+1}}(z'_{i+1})) \\
&\leq d(h_{\theta w^i}(\varphi_{1,u^0}(z'_i)), h_{w^{i+1}}(\varphi_{1,u^0}(z'_i))) \\
&\quad + d(h_{w^{i+1}}(\varphi_{1,u^0}(z'_i)), h_{w^{i+1}}(z'_{i+1})).
\end{aligned}$$

We know that h_u depends continuously on u . By compactness of \mathcal{U}_0 , we even have uniform continuity. This implies that we can choose δ small enough so that the first term becomes smaller than $\varepsilon/2$ for all i . By equicontinuity of the maps h_u , we can choose δ also small enough so that the second term becomes smaller than $\varepsilon/2$ for all i . Altogether, we have proved (13).

To show the last statement, observe that by choosing $(u, x), (v, y) \in L(Q)$, the same construction as above yields the $(\varepsilon + \delta)$ -chain (w^i, x_i) from (u, x) to (v, y) , which is completely contained in $L(Q)$. \square

To make use of the chain controllability and also for later purposes, it is important to know when Q has nonempty interior.⁸ To provide a quite general and checkable sufficient condition, we need to recall some concepts and a result from Sontag & Wirth [63].

The system Σ is called *analytic* if the state space M is a real-analytic manifold, U is a compact subset of some \mathbb{R}^m , satisfying $U = \text{cl int } U$, and the restriction of f to $M \times \text{int } U$ is a real-analytic map.

For a fixed $t \in \mathbb{Z}_{>0}$, a pair $(x, u) \in M \times (\text{int } U)^t$ is called *regular* if

$$\text{rk } \frac{\partial \varphi(t, \cdot, \cdot)}{\partial u}(x, u) = d = \dim M,$$

where $\varphi(t, \cdot, \cdot)$ is regarded as a map from $M \times (\text{int } U)^t$ to M so that $\frac{\partial \varphi(t, \cdot, \cdot)}{\partial u}(x, u)$ is a $d \times tm$ matrix.

A control sequence u of length $t > 0$ is called *universally regular* if (x, u) is regular for every $x \in M$. We write $S(t)$ for the set of all universally regular control sequences $u \in (\text{int } U)^t$.

We write $\mathcal{O}_t^+(x) = \{\varphi(t, x, u) : u \in \mathcal{U}\}$ for $t \geq 0$, and $\mathcal{O}^+(x) = \bigcup_{t \geq 0} \mathcal{O}_t^+(x)$ for the *forward orbit* of a point $x \in M$. If we only allow control sequences taking values in a subset $\tilde{U} \subset U$, we also write $\mathcal{O}_t^+(x; \tilde{U})$ and $\mathcal{O}^+(x; \tilde{U})$, respectively. The *negative orbit* of x is the set $\mathcal{O}^-(x) = \{\varphi(t, x, u) : t \leq 0, u \in \mathcal{U}\}$.

The system Σ is called *forward accessible from x* if $\text{int } \mathcal{O}^+(x) \neq \emptyset$. It is called *forward accessible* if it is forward accessible from every point.

Note that by Sard's theorem $\text{int } \mathcal{O}^+(x; \text{int } U) \neq \emptyset$ is equivalent to the existence of $t \in \mathbb{Z}_{>0}$ and $u \in (\text{int } U)^t$ such that $\text{rk } \frac{\partial \varphi(t, \cdot, \cdot)}{\partial u}(x, u) = d$.

⁸For our main result on invariance entropy, we need to assume that Q has positive volume.

We then have the following result from [63, Prop. 1].

3.13 Theorem: *Let the following assumptions hold:*

- (i) *The system Σ is analytic.*
- (ii) *Σ is uniformly forward accessible with control range $\text{int } U$, i.e., there exists $t_0 \in \mathbb{Z}_{>0}$ such that $\text{int } \mathcal{O}_{t_0}^+(x; \text{int } U) \neq \emptyset$ for all $x \in M$.*

*Then the set $S(t)$ is dense in U^t for all t large enough.*⁹

Under the assumptions of this theorem, imposed on the system Σ^0 , we obtain that the hyperbolic set Q as constructed above has nonempty interior.

3.14 Proposition: *Consider the set Q from Theorem 3.11 and assume that Σ is an analytic system and that Σ^0 is uniformly forward accessible with control range $\text{int } U_0$. Then Q has nonempty interior.*

Proof: Choose t_* large enough such that $S(t)$ (defined with respect to Σ^0) is nonempty for all $t \geq t_*$. Since Λ is hyperbolic and isolated invariant, there exists a periodic orbit in Λ (this is an implication of the Anosov Closing Lemma [39, Thm. 6.4.15]), say $\{f_{u_0}^t(x_0)\}$. Let $\tau \in \mathbb{Z}_{>0}$ denote its period and assume w.l.o.g. that $\tau \geq t_*$. Now pick a universally regular $u^* \in (\text{int } U_0)^\tau$. By periodic continuation, we can extend u^* to a τ -periodic sequence in $(\text{int } U_0)^\mathbb{Z}$ that we also denote by u^* .

Now consider the point $x^* := h_{u^*}(x_0) \in Q(u^*)$. By Proposition 3.10(d), we have

$$\varphi_{\tau, u^*}(x^*) = \varphi_{\tau, u^*}(h_{u^*}(x_0)) = h_{\theta^\tau u^*}(f_{u_0}^\tau(x_0)) = h_{u^*}(x_0) = x^*.$$

Hence, the trajectory $\varphi(\cdot, x^*, u^*)$ is τ -periodic. Using the regularity, we can find $\delta = \delta(\varepsilon) > 0$ so that every $y \in B_\delta(x^*)$ can be steered to every $z \in B_\delta(x^*)$ in time τ via some control sequence $u = u(y, z)$ of length τ , so that the controlled trajectory $(u_t, \varphi(t, y, u))_{t=0}^\tau$ is never further away from $(u_t^*, \varphi(t, x^*, u^*))$ than ε .¹⁰ By choosing $y = z$ and using periodic continuation again, we obtain a τ -periodic trajectory on the full time axis that completely evolves in the ε -neighborhood of Q and, by choosing ε small enough, we can also achieve that $u_t \in U_0$ for all $t \in \mathbb{Z}$. Since $L(Q)$ is isolated invariant, this implies that the trajectory evolves in Q , hence $y \in Q(u)$. This, in turn, implies $B_\delta(x^*) \subset Q$, which completes the proof. \square

Now we study the controllability properties of Σ^0 on the set Q . To formulate the next proposition, we introduce the *core* of a subset $Y \subset M$ as

$$\text{core}(Y) := \{y \in \text{int } Y : \text{int}(\mathcal{O}^-(y) \cap Y) \neq \emptyset \text{ and } \text{int}(\mathcal{O}^+(y) \cap Y) \neq \emptyset\}.$$

⁹The result in [63] actually makes a much stronger statement, which we do not use in our paper.

¹⁰This is a consequence of the implicit function theorem, cf. [62, Thm. 7].

3.15 Proposition: Consider the hyperbolic set Q from Theorem 3.11 for the control system Σ^0 . Additionally, let the following assumptions hold:

- (a) Λ is a topologically transitive set of f_u^0 .
- (b) Q has nonempty interior.

Then complete controllability holds on $\text{core}(Q)$.

Proof: By using the construction in the proof of Lemma 3.12, we can produce bi-infinite controlled ε -chains passing through any two given points in Q . Let $(x_t, w^t)_{t \in \mathbb{Z}}$ be such a controlled chain, that is

$$(w^t, x_t) \in L(Q) \quad \text{and} \quad d(\varphi(1, x_t, w^t), x_{t+1}) \leq \varepsilon \quad \text{for all } t \in \mathbb{Z}.$$

We define another control sequence $w^* \in \mathcal{U}_0$ by putting

$$w_t^* := w_0^t \quad \text{for all } t \in \mathbb{Z}.$$

In this way, $(\theta^t w^*, x_t)_{t \in \mathbb{Z}}$ becomes an ε -pseudo-orbit, since

$$\begin{aligned} d(\varphi(1, x_t, \theta^t w^*), x_{t+1}) &= d(f_{w_t^*}(x_t), x_{t+1}) \\ &= d(f_{w_0^t}(x_t), x_{t+1}) = d(\varphi(1, x_t, w^t), x_{t+1}) \leq \varepsilon. \end{aligned}$$

We want to apply the shadowing lemma to shadow such chains, but we need to make sure that they are close enough to $L(Q)$. Recalling that we constructed the chains with $d_{\mathcal{U}}(\theta w^t, w^{t+1}) \leq \delta$ (where δ only depends on ε), we find that

$$\begin{aligned} d_{\mathcal{U} \times M}((\theta^t w^*, x_t), (w^t, x_t)) &= d_{\mathcal{U}}(\theta^t w^*, w^t) \\ &= \sum_{s \in \mathbb{Z}} \frac{1}{2^{|s|}} d_U(w_{t+s}^*, w_s^t) = \sum_{s \in \mathbb{Z}} \frac{1}{2^{|s|}} d_U(w_0^{t+s}, w_s^t) \\ &= \sum_{s \in \mathbb{Z}} \frac{1}{2^{|s|}} d_U(w_0^{t+s}, (\theta^s w^t)_0) \leq \sum_{s \in \mathbb{Z}} \frac{1}{2^{|s|}} d_{\mathcal{U}}(w^{t+s}, \theta^s w^t). \end{aligned}$$

Now we can split the sum into a finite and an infinite part, the latter being small because of the factor $2^{-|s|}$, and the first being small due to the choice of δ . To be more precise, to achieve that the sum becomes smaller than a given $\gamma > 0$, first pick $s_0 > 0$ large enough so that

$$\text{diam } \mathcal{U} \sum_{|s| > s_0} \frac{1}{2^{|s|}} \leq \frac{\gamma}{2}.$$

Then choose $\delta > 0$ small enough so that for all $|s| \leq s_0$ we have the following:

- If $s > 0$, then

$$\frac{1}{2^s} d_{\mathcal{U}}(w^{t+s}, \theta^s w^t) \leq \frac{1}{2^s} \sum_{i=0}^{s-1} d_{\mathcal{U}}(\theta^i w^{t+s-i}, \theta^i \theta w^{t+s-i-1}) \leq \frac{\gamma}{2(2s_0 + 1)},$$

which is possible, since $\{\theta^i\}_{i=0}^{s_0-1}$ is a uniformly equicontinuous family and $d_{\mathcal{U}}(w^{t+s-i}, \theta w^{t+s-i-1}) \leq \delta$.

- If $s < 0$, then

$$\frac{1}{2^{-s}} d_{\mathcal{U}}(w^{t+s}, \theta^s w^t) \leq \frac{1}{2^{-s}} \sum_{i=s}^{-1} d_{\mathcal{U}}(\theta^i w^{t+s-i}, \theta^i \theta w^{t+s-i-1}) \leq \frac{\gamma}{2(2s_0 + 1)},$$

which is possible by similar reasons as used in the former case.

Altogether, $d_{\mathcal{U} \times M}((\theta^t w^*, x_t), (w^t, x_t)) \leq \gamma$. Hence, it follows that the ε -pseudo-orbit $(\theta^t w^*, x_t)_{t \in \mathbb{Z}}$, for δ sufficiently small, can be β -shadowed by a real orbit in $L(Q)$ of the form $(\theta^t w^*, \varphi(t, z, w^*))_{t \in \mathbb{Z}}$:

$$(w^*, z) \in L(Q) \quad \text{and} \quad d(\varphi(t, z, w^*), x_t) \leq \beta \quad \text{for all } t \in \mathbb{Z}.$$

This implies that for any given points $x, y \in Q$ we find a trajectory in Q starting in an arbitrarily small neighborhood of x and ending (after a finite time) in an arbitrarily small neighborhood of y . Now assume that $x, y \in \text{core}(Q)$. Pick points $x' \in \text{int}(\mathcal{O}^+(x) \cap Q)$ and $y' \in \text{int}(\mathcal{O}^-(y) \cap Q)$ and a trajectory starting at some $x'' \in \text{int}(\mathcal{O}^+(x) \cap Q)$ and ending in $y'' \in \text{int}(\mathcal{O}^-(y) \cap Q)$ (obtained by shadowing a chain from x' to y'). Then one can steer from x to x'' , from x'' to y'' and from y'' to y . This proves the controllability statement. \square

It is important to understand how large $\text{core}(Q)$ is. From [3], we know that $\text{core}(Q)$ is always an open set under mild assumptions on the system.

3.16 Lemma: Assume that $U \subset \mathbb{R}^m$ for some $m \in \mathbb{Z}_{>0}$ and $U_0 = \text{cl int } U_0$. Furthermore, let $f : M \times U \rightarrow M$ be of class C^1 . Then $\text{core}(Q) \neq \emptyset$ implies that $\text{core}(Q)$ is open in M and dense in Q .

Proof: Consider the sets

$$\begin{aligned} \mathcal{O}^-(\text{core}(Q)) &= \{x \in M : \exists y \in \text{core}(Q), u \in \mathcal{U}_0, t \geq 0 \text{ s.t. } \varphi(t, x, u) = y\}, \\ \mathcal{O}^+(\text{core}(Q)) &= \{y \in M : \exists x \in \text{core}(Q), u \in \mathcal{U}_0, t \geq 0 \text{ s.t. } \varphi(t, x, u) = y\}. \end{aligned}$$

Since $\text{core}(Q)$ is nonempty by assumption and open by [3, Lem. 7.8], the preceding proof shows that $\mathcal{O}^-(\text{core}(Q))$ is open and dense in Q . Moreover, every $x \in \mathcal{O}^-(\text{core}(Q))$ satisfies $\text{int}(Q \cap \mathcal{O}^+(x)) \neq \emptyset$. The set $\mathcal{O}^+(\text{core}(Q))$ is also open and dense in Q by the preceding proof and every point $x \in \mathcal{O}^+(\text{core}(Q))$ satisfies $\text{int}(Q \cap \mathcal{O}^-(x)) \neq \emptyset$. Hence, $\mathcal{O}^-(\text{core}(Q)) \cap \mathcal{O}^+(\text{core}(Q)) = \text{core}(Q)$ and the assertion follows. \square

We can thus formulate the following corollary.

3.17 Corollary: Consider the hyperbolic set Q from Theorem 3.11 for the control system Σ^0 . Additionally, let the following assumptions hold:

- (a) $U \subset \mathbb{R}^m$ for some $m \in \mathbb{Z}_{>0}$ and $U_0 = \text{cl int } U_0$.
- (b) $f : M \times U \rightarrow M$ is of class C^1 .
- (c) Λ is a topologically transitive set of f_{u^0} .
- (d) $\text{core}(Q)$ is nonempty.

Then complete controllability holds on an open and dense subset of Q .

The following proposition provides a sufficient condition for $\text{core}(Q) \neq \emptyset$.

3.18 Proposition: Assume that the given system is analytic and forward accessible.¹¹ Then $\text{int } Q \neq \emptyset$ implies $\text{core}(Q) \neq \emptyset$.

Proof: By [3, Lem. 5.1], on an open and dense subset of M the Lie algebra rank condition (introduced in [3, p. 5]) is satisfied. Let W denote the intersection of this set with $\text{int } Q$. Now we pick a point $z \in W$ and a $\gamma > 0$ so that $B_\gamma(z) \subset W$. Consider a bi-infinite ε -pseudo-orbit whose x -component passes through $B_{\gamma/3}(z)$ infinitely many times. By shadowing this pseudo-orbit (choosing ε sufficiently small), we can find an orbit starting in some $x \in \text{int } Q$ that passes through $B_{\gamma/2}(z)$ infinitely many times. Then there exists a sequence of points $x_k \in \mathcal{O}_{n_k}^+(x) \cap B_{\gamma/2}(z)$, where $n_k \rightarrow \infty$. We may assume that x_k converges to some point $y \in \text{cl } B_{\gamma/2}(z)$. Since $y \in W$, the Lie algebra rank condition holds at y . By [3, Lem. 4.1] and the subsequent remarks, one can reach from x an open set in every neighborhood of y . This implies $\text{int}(Q \cap \mathcal{O}^+(x)) \neq \emptyset$. Since the same construction works in backward time, we conclude that also $\text{int}(Q \cap \mathcal{O}^-(x)) \neq \emptyset$. Hence, $x \in \text{core}(Q)$. \square

3.19 Corollary: Let the following assumptions hold for the control system Σ^0 and the hyperbolic set Q from Theorem 3.11:

- (a) Σ^0 is analytic and uniformly forward accessible.
- (b) Λ is a topologically transitive set of f_{u^0} .

Then complete controllability holds on $\text{core}(Q)$, which is an open and dense subset of Q .

In Section 6, we will show by an example how uniform forward accessibility can be checked for a concrete system with a finite number of computations.

¹¹It is actually enough to assume that the system is forward accessible from one point $x \in M$. Then, by analyticity it is forward accessible from all x in an open and dense set, which is enough for the conclusion of the proposition.

4 Invariance entropy of hyperbolic sets

In this section, we derive a lower bound on the invariance entropy of a hyperbolic set in terms of dynamical quantities.

4.1 A first lower estimate on invariance entropy

Let Q be a compact all-time controlled invariant set of Σ . For $u \in \mathcal{U}_Q$, $\tau \in \mathbb{Z}_{>0}$ and $\varepsilon > 0$, we define

$$Q(u, \tau, \varepsilon) := \{x \in M : \text{dist}(\varphi_{t,u}(x), Q(\theta^t u)) \leq \varepsilon, \forall 0 \leq t < \tau\}.$$

Hence, $Q(u, \tau, \varepsilon)$ is the set of all initial states so that the trajectory under u stays ε -close to the corresponding fiber in the time interval $[0; \tau)$.

The following lemma provides a first lower estimate on invariance entropy under the assumption that the fiber map is lower semicontinuous.

4.1 Lemma: *Let Q be a compact all-time controlled invariant set of Σ and assume that the fiber map $u \mapsto Q(u)$, defined on \mathcal{U}_Q , is lower semicontinuous. Then, for every compact set $K \subset Q$ with positive volume and every $\varepsilon > 0$, we have*

$$h_{\text{inv}}(K, Q) \geq -\liminf_{\tau \rightarrow \infty} \sup_{u \in \mathcal{U}_Q} \frac{1}{\tau} \log \text{vol}(Q(u, \tau, \varepsilon)).$$

Proof: For all $\tau \in \mathbb{Z}_{>0}$ and $u \in \mathcal{U}$, we define the sets

$$\begin{aligned} Q(u, \tau) &:= \{x \in M : \varphi(t, x, u) \in Q, \forall 0 \leq t < \tau\}, \\ Q^\pm(u, \tau) &:= \{x \in M : \varphi(t, x, u) \in Q, \forall -\tau < t < \tau\}, \\ V(u, \tau) &:= \{v \in \mathcal{U} : u_t = v_t, \forall -\tau < t < \tau - 1\}. \end{aligned}$$

The set $Q^\pm(u, \tau)$ can be characterized as

$$Q^\pm(u, \tau) = \bigcup_{v \in V(u, \tau)} Q(v).$$

Indeed, if $x \in Q^\pm(u, \tau)$, then by all-time controlled invariance, the control sequence u can be modified outside of the interval $(-\tau; \tau - 1)$ so that $\varphi(\mathbb{Z}, x, u^*) \subset Q$, where u^* denotes the modified sequence. Hence, $x \in Q(u^*)$. Conversely, if $x \in Q(u^*)$ for some u^* which coincides with u on $(-\tau; \tau - 1)$, then clearly $x \in Q^\pm(u, \tau)$.

Now let $\varepsilon > 0$. Since the fiber map $u \mapsto Q(u)$ is always upper semicontinuous, the assumption of lower semicontinuity implies its continuity with respect to the Hausdorff metric. Since \mathcal{U}_Q is compact, we even have uniform continuity. Hence, there exists $\delta > 0$ so that $d_{\mathcal{U}}(u, v) < \delta$ (for any $u, v \in \mathcal{U}$) implies

$$Q(v) \subset N_\varepsilon(Q(u)).$$

We choose $\tau_0 \in \mathbb{Z}_{>0}$ large enough so that $V(u, \tau_0) \subset B_\delta(u)$ for all $u \in \mathcal{U}$, which is possible by definition of the product topology. This implies

$$Q^\pm(u, \tau_0) = \bigcup_{v \in V(u, \tau_0)} Q(v) \subset N_\varepsilon(Q(u)) \quad \text{for all } u \in \mathcal{U}.$$

Now let $\mathcal{S} \subset \mathcal{U}$ be a minimal $(2\tau_0 + t, K, Q)$ -spanning set for some $t \in \mathbb{Z}_+$. We may assume without loss of generality that \mathcal{S} is finite and contained in \mathcal{U}_Q . Then

$$K \subset \bigcup_{u \in \mathcal{S}} Q(u, 2\tau_0 + t). \quad (14)$$

We claim that

$$\varphi_{s, \theta^{\tau_0} u}(\varphi_{\tau_0, u}(Q(u, 2\tau_0 + t))) \subset Q^\pm(\theta^{s+\tau_0} u, \tau_0) \quad \text{for all } s \in [0; t].$$

Indeed, let x be an element of the left-hand side. Then we can write $x = \varphi(s + \tau_0, y, u)$ for some $y \in Q(u, 2\tau_0 + t)$. Hence,

$$\varphi(r, x, \theta^{s+\tau_0} u) = \varphi(r + s + \tau_0, y, u) \in Q \quad \text{for all } r \in [-\tau_0 - s; \tau_0 + t - s]$$

and $(-\tau_0; \tau_0) \subset [-\tau_0 - s; \tau_0 + t - s]$ for all $s \in [0; t]$. We thus have

$$\begin{aligned} \varphi_{\tau_0, u}(Q(u, 2\tau_0 + t)) &\subset \bigcap_{s=0}^{t-1} \varphi_{s, \theta^{\tau_0} u}^{-1} [Q^\pm(\theta^{s+\tau_0} u, \tau_0)] \\ &\subset \bigcap_{s=0}^{t-1} \varphi_{s, \theta^{\tau_0} u}^{-1} [N_\varepsilon(Q(\theta^{s+\tau_0} u))] = Q(\theta^{\tau_0} u, t, \varepsilon). \end{aligned}$$

Together with (14), this yields

$$0 < \text{vol}(K) \leq |\mathcal{S}| \cdot \max_{u \in \mathcal{S}} \text{vol}(\varphi_{\tau_0, u}^{-1}(Q(\theta^{\tau_0} u, t, \varepsilon))).$$

Observing that the volume change of a set affected by the application of $\varphi_{\tau_0, u}^{-1}$ (within some compact domain) does not change the exponential volume growth rate, this estimate implies

$$0 \leq h_{\text{inv}}(K, Q) + \liminf_{t \rightarrow \infty} \sup_{u \in \mathcal{U}_Q} \frac{1}{t} \log \text{vol}(Q(u, t, \varepsilon)),$$

which is equivalent to the desired inequality. \square

4.2 Bowen-balls, measure-theoretic entropy and pressure

In this subsection, we assume throughout that Q is a hyperbolic set for Σ so that $L(Q)$ is an isolated invariant set of the control flow. Moreover, we assume that Σ is of regularity class C^2 .

For $u \in \mathcal{U}$, $\tau \in \mathbb{Z}_{>0}$ and $\varepsilon > 0$, we say that a set $E \subset M$ (u, τ, ε) -spans another set $K \subset M$ if for each $x \in K$ there is $y \in E$ with $d^{u, \tau}(x, y) \leq \varepsilon$. In other words, the Bowen-balls of order τ and radius ε centered at the points in E cover the set K . A set $F \subset M$ is called (u, τ, ε) -separated if $d^{u, \tau}(x, y) > \varepsilon$ for all $x, y \in F$ with $x \neq y$.

We will use Bowen-balls in order to estimate $\text{vol}(Q(u, \tau, \varepsilon))$ as follows. For a small number $\delta > 0$, we let $F_{u, \tau, \delta}$ be a maximal (u, τ, δ) -separated subset of the u -fiber $Q(u)$. By compactness of $Q(u)$, $F_{u, \tau, \delta}$ is finite. Moreover, it is easy to see that a maximal (u, τ, δ) -separated subset of some set also (u, τ, δ) -spans this set.¹²

Now, for an arbitrary $x \in Q(u, \tau, \varepsilon)$, pick $x_* \in Q(u)$ and $x^* \in Q(\theta^{\tau-1}u)$ so that $d(x, x_*) \leq \varepsilon$ and $d(\varphi(\tau-1, x, u), x^*) \leq \varepsilon$. Then we consider the sequence $(x_t)_{t \in \mathbb{Z}}$ defined by

$$x_t := \begin{cases} \varphi(t, x_*, u) & \text{if } t < 0, \\ \varphi(t, x, u) & \text{if } 0 \leq t \leq \tau - 2, \\ \varphi(t - (\tau - 1), x^*, \theta^{\tau-1}u) & \text{if } t \geq \tau - 1. \end{cases}$$

The joint sequence $(\theta^t u, x_t)_{t \in \mathbb{Z}}$ is an ε -pseudo-orbit. Since $(\theta^t u, x_t) \in L(Q)$ for all $t < 0$ and $t \geq \tau - 1$ and $(\theta^t u, x_t)$ is ε -close to some point in $L(Q)$ for all $t \in [0; \tau - 2]$, for ε small enough the shadowing lemma yields a point $z \in Q(u)$ so that¹³

$$d(\varphi(t, x, u), \varphi(t, z, u)) \leq \beta \quad \text{for all } t \in [0; \tau].$$

This implies $x \in B_\beta^{u, \tau}(z)$. Now pick some $y \in F_{u, \tau, \delta}$ so that $d^{u, \tau}(y, z) \leq \delta$. Then $x \in B_{\beta+\delta}^{u, \tau}(y)$. We conclude that

$$Q(u, \tau, \varepsilon) \subset \bigcup_{y \in F_{u, \tau, \delta}} B_{\beta+\delta}^{u, \tau}(y).$$

If β and δ are chosen small enough, we can thus apply the volume lemma in order to estimate

$$\text{vol}(Q(u, \tau, \varepsilon)) \leq C_{\beta+\delta} \sum_{y \in F_{u, \tau, \delta}} J^+ \varphi_{\tau, u}(y)^{-1}. \quad (15)$$

To turn this into a meaningful estimate for $h_{\text{inv}}(K, Q)$, a significant amount of additional work is necessary.

First, we need to pay attention to the fact that the control flow can be regarded as a *random dynamical system*, once we equip the space \mathcal{U} with a Borel probability measure P , invariant under θ . We denote such a random dynamical system briefly by (φ, P) .¹⁴ An *invariant measure* of (φ, P) is a Borel probability measure μ on $\mathcal{U} \times M$ satisfying the following two properties:

¹²This can easily be proved by contradiction.

¹³We have to be a little bit careful when we consider $x_{\tau-1} = x^*$. Note that $d(\varphi(\tau-1, x, u), \varphi(\tau-1, z, u)) \leq d(\varphi(\tau-1, x, u), x^*) + d(x^*, \varphi(\tau-1, z, u)) \leq \varepsilon + \beta$. Hence, we should replace β with $\beta - \varepsilon$.

¹⁴Be aware that (φ, P) is an RDS on a purely formal level. We actually do not consider any randomness here.

- Φ preserves the measure μ , i.e., $\Phi_*\mu = \mu$.
- The marginal of μ on \mathcal{U} coincides with P , i.e., $(\pi_{\mathcal{U}})_*\mu = P$.

By the disintegration theorem, each invariant measure μ admits a disintegration into sample measures μ_u on $(M, \mathcal{B}(M))$, defined for P -almost all $u \in \mathcal{U}$. That is,

$$d\mu(u, x) = d\mu_u(x)dP(u).$$

To each invariant measure μ , we can associate the *measure-theoretic entropy* $h_\mu(\varphi, P)$. Let \mathcal{A} be a finite Borel partition of M . An induced dynamically defined sequence of (finite Borel) partitions of M is given by

$$\mathcal{A}(u, \tau) := \bigvee_{t=0}^{\tau-1} \varphi_{t,u}^{-1} \mathcal{A} = \left\{ A_0 \cap \varphi_{1,u}^{-1}(A_1) \cap \dots \cap \varphi_{\tau-1,u}^{-1}(A_{\tau-1}) : A_s \in \mathcal{A}, \forall s \right\}.$$

The entropy associated with the partition \mathcal{A} is defined as

$$h_\mu(\varphi, P; \mathcal{A}) := \lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int_{\mathcal{U}} H_{\mu_u}(\mathcal{A}(u, \tau)) dP(u),$$

where $H_{\mu_u}(\cdot)$ denotes the Shannon entropy of a partition and the limit exists because of subadditivity, see [7].

The measure-theoretic entropy of (φ, P) with respect to μ is then defined as

$$h_\mu(\varphi, P) := \sup_{\mathcal{A}} h_\mu(\varphi, P; \mathcal{A}) \in [0, \infty],$$

the supremum taken over all finite Borel partitions of M . A related quantity is the *measure-theoretic pressure* of (φ, P) with respect to μ and a μ -integrable “potential” $\alpha : \mathcal{U} \times M \rightarrow \mathbb{R}$, defined as

$$\pi_\mu(\varphi, P; \alpha) := h_\mu(\varphi, P) + \int \alpha d\mu.$$

Our aim is to prove the following lower bound for the invariance entropy:

$$h_{\text{inv}}(K, Q) \geq \inf_{P \in \mathcal{M}(\theta)} \inf_{\mu \in \mathcal{M}_P(\Phi; L(Q))} -\pi_\mu(\varphi, P; -\log J^+ \varphi), \quad (16)$$

where $J^+ \varphi$ denotes the function $(u, x) \mapsto J^+ \varphi_{1,u}(x)$ and $\mathcal{M}_P(\Phi; L(Q))$ the set of all invariant probability measures of the bundle RDS that is defined by the restriction of Φ to $L(Q)$ together with the measure P on \mathcal{U} .¹⁵ By the definition of pressure, this estimate is equivalent to

$$h_{\text{inv}}(K, Q) \geq \inf_{P \in \mathcal{M}(\theta)} \inf_{\mu \in \mathcal{M}_P(\Phi; L(Q))} \left[\int \log J^+ \varphi_{1,u}(x) d\mu(u, x) - h_\mu(\varphi, P) \right].$$

¹⁵Observe that the sets \mathcal{E}_ω in the definition of a bundle RDS in Subsection 2.2 here are precisely the u -fibers $Q(u)$.

Obviously, the double infimum can be written as a single infimum as follows:

$$h_{\text{inv}}(K, Q) \geq \inf_{\mu \in \mathcal{M}(\Phi|_{L(Q)})} \left[\int \log J^+ \varphi_{1,u}(x) d\mu(u, x) - h_\mu(\varphi, (\pi_{\mathcal{U}})_* \mu) \right].$$

By the *Margulis-Ruelle inequality* [4], this lower bound is always nonnegative.

We propose the following interpretation of the terms involved in the right-hand side of the above estimate:

- $\int \log J^+ \varphi_{1,u}(x) d\mu(u, x)$: the total instability of the dynamics on $L(Q)$ seen by the measure μ .
- $h_\mu(\varphi, (\pi_{\mathcal{U}})_* \mu)$: the part of the instability not leading to exit from Q .
- $\inf_{\mu \in \mathcal{M}(\Phi|_{L(Q)})}$: the infimum over all possible control strategies to make Q invariant.

The first item is obvious. The second one can be justified by observing that the entropy with respect to a measure supported on $L(Q)$ captures the complexity of the fiber dynamics which is constituted by the trajectories that completely evolve within Q (here the definition of entropy for a bundle RDS as discussed in [46, Sec. 1.1] is helpful for a precise understanding). Finally, the third item hopefully will be justified by future work on achievability results (upper bounds for invariance entropy) which are still missing for the general case.

From now on, we will frequently use the following three assumptions on the compact all-time controlled invariant set Q :

- (A1) Q is uniformly hyperbolic.
- (A2) $L(Q)$ is an isolated invariant set of the control flow.
- (A3) The fiber map $u \mapsto Q(u)$ is lower semicontinuous.

4.3 Construction of approximating subadditive cocycles

Let the assumptions (A1)–(A3) be satisfied for the compact all-time controlled invariant set Q . For every $\varepsilon > 0$, we define the function

$$v^\varepsilon : (\tau, u) \mapsto v_\tau^\varepsilon(u) := \log \text{vol}(Q(u, \tau, \varepsilon)), \quad v^\varepsilon : \mathbb{Z}_{>0} \times \mathcal{U} \rightarrow \mathbb{R}.$$

It would be useful if v^ε was a subadditive cocycle over the system (\mathcal{U}, θ) . This cannot be expected, however. Instead, we approximate v^ε by subadditive cocycles.

For a fixed $u \in \mathcal{U}$, let $\mathcal{A} = (\mathcal{A}_t)_{t=0}^\infty$ be a sequence so that \mathcal{A}_t is an open cover of the compact set $Q(\theta^t u)$, i.e., a collection of subsets of $Q(\theta^t u)$, open relative to

$Q(\theta^t u)$, whose union equals $Q(\theta^t u)$. We write

$$\mathcal{A}^\tau := \bigvee_{t=0}^{\tau-1} \varphi_{t,u}^{-1}(\mathcal{A}_t), \quad \tau \in \mathbb{Z}_{>0}.$$

This is the collection of all sets of the form

$$A_0 \cap \varphi_{1,u}^{-1}(A_1) \cap \dots \cap \varphi_{\tau-1,u}^{-1}(A_{\tau-1}), \quad A_t \in \mathcal{A}_t.$$

Observe that \mathcal{A}^τ is an open cover of $Q(u)$. We define

$$w_\tau^{\mathcal{A}}(u) := \log \inf \left\{ \sum_{A \in \alpha} \sup_{x \in A} J^+ \varphi_{\tau,u}(x)^{-1} : \alpha \text{ is a finite subcover of } \mathcal{A}^\tau \text{ for } Q(u) \right\},$$

which is well-defined, because $J^+ \varphi_{\tau,u}(x)$ is only evaluated at points $(u, x) \in L(Q)$. We write $\mathcal{A}(\tau)$ for the shifted sequence $(\mathcal{A}_\tau, \mathcal{A}_{\tau+1}, \mathcal{A}_{\tau+2}, \dots)$.

Now let α be a finite subcover of \mathcal{A}^{τ_1} for $Q(u)$ and β a finite subcover of $\mathcal{A}(\tau_1)^{\tau_2}$ for $Q(\theta^{\tau_1} u)$. Then

$$\begin{aligned} & \sum_{C \in \alpha \vee \varphi_{\tau_1,u}^{-1}(\beta)} \sup_{z \in C} J^+ \varphi_{\tau_1+\tau_2,u}(z)^{-1} \\ &= \sum_{C \in \alpha \vee \varphi_{\tau_1,u}^{-1}(\beta)} \sup_{z \in C} \left[J^+ \varphi_{\tau_1,u}(z)^{-1} \cdot J^+ \varphi_{\tau_2, \theta^{\tau_1} u}(\varphi_{\tau_1,u}(z))^{-1} \right] \\ &\leq \sum_{(A,B) \in \alpha \times \beta} \left[\sup_{x \in A} J^+ \varphi_{\tau_1,u}(x)^{-1} \right] \cdot \left[\sup_{y \in B} J^+ \varphi_{\tau_2, \theta^{\tau_1} u}(y)^{-1} \right] \\ &= \sum_{A \in \alpha} \left[\sup_{x \in A} J^+ \varphi_{\tau_1,u}(x)^{-1} \right] \cdot \sum_{B \in \beta} \left[\sup_{y \in B} J^+ \varphi_{\tau_2, \theta^{\tau_1} u}(y)^{-1} \right]. \end{aligned}$$

Hence, if we choose α and β so that the corresponding sums are close to their infima, we see that

$$w_{\tau_1+\tau_2}^{\mathcal{A}}(u) \leq w_{\tau_1}^{\mathcal{A}}(u) + w_{\tau_2}^{\mathcal{A}(\tau_1)}(\theta^{\tau_1} u), \quad (17)$$

where we use that $\alpha \vee \varphi_{\tau_1,u}^{-1}(\beta)$ is a subcover of $\mathcal{A}^{\tau_1+\tau_2}$ for $Q(u)$.

For a fixed (small) $\delta > 0$, let $\mathcal{A}(u) = (\mathcal{A}_t(u))_{t=0}^\infty$ be the unique sequence so that $\mathcal{A}_t(u)$ consists of all open δ -balls in $Q(\theta^t u)$ and put

$$w_\tau^\delta(u) := w_\tau^{\mathcal{A}(u)}(u).$$

Then we have the following result which shows that the family of functions $w^\delta : \mathbb{Z}_{>0} \times \mathcal{U} \rightarrow \mathbb{R}$, $(\tau, u) \mapsto w_\tau^\delta(u)$, consists of subadditive cocycles that can be used to approximate v^ε .

4.2 Proposition: *The functions w^δ have the following properties, where the constants in (b) and (c) come from the volume lemma (Theorem 3.6):*

(a) *For every $\delta > 0$, the function $(\tau, u) \mapsto w_\tau^\delta(u)$ is a subadditive cocycle over (\mathcal{U}, θ) .*

(b) *For every $\delta > 0$, there exists $\varepsilon > 0$ so that for all $u \in \mathcal{U}$ and $\tau \in \mathbb{Z}_{>0}$:*

$$v_\tau^\varepsilon(u) \leq \log C_{3\delta} + w_\tau^\delta(u)$$

(c) *For every $\gamma > 0$, there exists $\delta > 0$ so that for all $u \in \mathcal{U}$ and $\tau \in \mathbb{Z}_{>0}$:*

$$w_\tau^\delta(u) - \log C_{\delta/2} \leq \tau\gamma + v_\tau^{\delta/2}(u)$$

(d) *For all $\varepsilon > 0$ small enough and $\delta \in (0, \varepsilon)$, there exist a constant $\tilde{C}_\delta > 0$ and $T \in \mathbb{Z}_{>0}$ so that for all $u \in \mathcal{U}$ and $\tau > 2T$:*

$$v_\tau^\varepsilon(u) \leq \tilde{C}_\delta + v_{\tau-2T}^\delta(\theta^T u)$$

Proof: (a) This follows directly from (17).

(b) Choose $\varepsilon > 0$ small enough so that every ε -pseudo-orbit contained in an ε -neighborhood of $L(Q)$ is δ -shadowed by an orbit in $L(Q)$. For an arbitrary $u \in \mathcal{U}$, let $\mathcal{A} = \mathcal{A}(u)$ and let $F \subset Q(u)$ be a maximal $(u, \tau, 2\delta)$ -separated set. Then each member of \mathcal{A}^τ contains at most one element of F . Indeed, if there were two such elements x_1 and x_2 , then

$$d(\varphi(t, x_1, u), \varphi(t, x_2, u)) < 2\delta \quad \text{for } t = 0, 1, \dots, \tau - 1$$

in contradiction to the separation property. Hence, for every finite subcover α of \mathcal{A}^τ we have

$$\sum_{x \in F} J^+ \varphi_{\tau, u}(x)^{-1} \leq \sum_{A \in \alpha} \sup_{x \in A} J^+ \varphi_{\tau, u}(x)^{-1}.$$

By (15), we can estimate

$$v_\tau^\varepsilon(u) \leq \log C_{3\delta} + \log \sum_{A \in \alpha} \sup_{x \in A} J^+ \varphi_{\tau, u}(x)^{-1},$$

which implies the assertion.

(c) For the given $\gamma > 0$ choose $\delta > 0$ small enough so that

$$\frac{J^+ \varphi_{1, u}(x_1)}{J^+ \varphi_{1, u}(x_2)} \leq 2^\gamma \tag{18}$$

for all x_1, x_2 in Q satisfying $d(x_1, x_2) \leq 2\delta$ and all $u \in \mathcal{U}$, which is possible by uniform continuity of $(x, u) \mapsto \log J^+ \varphi_{1, u}(x)$ on the compact set $Q \times \mathcal{U}$.

Let $\mathcal{A} := \mathcal{A}(u)$ and consider a finite (u, τ, δ) -spanning set E for $Q(u)$, contained in $Q(u)$. For each $z \in E$, consider $A_t(z) \in \mathcal{A}_t$ so that $B_\delta(\varphi(t, z, u)) = A_t(z)$ for $t = 0, 1, \dots, \tau - 1$. Let

$$C(z) := \bigcap_{t=0}^{\tau-1} \varphi_{t,u}^{-1}(A_t(z)) \in \mathcal{A}^\tau,$$

which is an open Bowen-ball centered at z and intersected with $Q(u)$. The definition of $C(z)$ together with (18) implies

$$\sup_{x \in C(z)} J^+ \varphi_{\tau,u}(x)^{-1} \leq 2^{\tau\gamma} \cdot J^+ \varphi_{\tau,u}(z)^{-1}.$$

Since the sets $C(z)$, $z \in E$, form a finite subcover of \mathcal{A}^τ for $Q(u)$,

$$w_\tau^\delta(u) \leq \tau\gamma + \log \sum_{z \in E} J^+ \varphi_{\tau,u}(z)^{-1}.$$

Since a maximal (u, τ, δ) -separated set is also (u, τ, δ) -spanning and the corresponding Bowen-balls of radius $\delta/2$ are disjoint and contained in $Q(u, \tau, \delta/2)$, the volume lemma implies

$$w_\tau^\delta(u) \leq \tau\gamma + \log C_{\delta/2} + v_\tau^{\delta/2}(u).$$

(d) Fix ε and δ as in the statement. We claim that there exists $T \in \mathbb{Z}_{>0}$ so that for all $u \in \mathcal{U}$ and $x \in M$ the following implication holds:

$$\max_{-T < t < T} \text{dist}(\varphi(t, x, u), Q(\theta^t u)) \leq \varepsilon \quad \Rightarrow \quad \text{dist}(x, Q(u)) < \delta. \quad (19)$$

Suppose to the contrary that for every $T \in \mathbb{Z}_{>0}$ there are $u_T \in \mathcal{U}$ and $x_T \in M$ with

$$\text{dist}(\varphi(t, x_T, u_T), Q(\theta^t u_T)) \leq \varepsilon \text{ for } |t| < T \text{ and } \text{dist}(x_T, Q(u_T)) \geq \delta.$$

By compactness of \mathcal{U} , we may assume that $u_T \rightarrow u \in \mathcal{U}$ and by compactness of small closed neighborhoods of Q , we may assume that $x_T \rightarrow x \in M$. For arbitrary $t \in \mathbb{Z}$, we have $\text{dist}(\varphi(t, x_T, u_T), Q(\theta^t u_T)) \leq \varepsilon$ whenever $T > |t|$. Since $\varphi(t, \cdot, \cdot)$, $Q(\cdot)$ and $\text{dist}(\cdot, \cdot)$ are continuous functions, this implies $\text{dist}(\varphi(t, x, u), Q(\theta^t u)) \leq \varepsilon$ for all $t \in \mathbb{Z}$ and $\text{dist}(x, Q(u)) \geq \delta$. For each $t \in \mathbb{Z}$, pick $y_t \in Q(\theta^t u)$ so that $d(\varphi(t, x, u), y_t) \leq \varepsilon$. Then $\Phi_t(u, x) = (\theta^t u, \varphi(t, x, u))$ is ε -close to $(\theta^t u, y_t) \in L(Q)$. Hence, if $\varepsilon > 0$ is small enough so that $N_\varepsilon(L(Q))$ is an isolating neighborhood of $L(Q)$, then $(u, x) \in L(Q)$, which contradicts $\text{dist}(x, Q(u)) \geq \delta$.

Now choose T according to (19) and let $x \in Q(u, \tau, \varepsilon)$ for some $\tau > 2T$. We want to show that $\varphi_{T,u}(x) \in Q(\theta^T u, \tau - 2T, \delta)$. To show this, let $x_s := \varphi(s, \varphi(T, x, u), \theta^T u) = \varphi(T + s, x, u)$ for $0 \leq s < \tau - 2T$ and observe that

$$\text{dist}(\varphi(r, x_s, \theta^{T+s} u), Q(\theta^r \theta^{T+s} u))$$

$$= \text{dist}(\varphi(T+r+s, x, u), Q(\theta^{T+s+r}u)) \leq \varepsilon$$

whenever $|r| < T$, since $0 < T+r+s < T+(T-1)+(\tau-2T) = \tau-1$. By (19), this implies $\text{dist}(x_s, Q(\theta^s u)) < \delta$ for $0 \leq s < \tau-2T$, hence $x \in Q(\theta^T u, \tau-2T, \delta)$. It follows that $\varphi_{T,u}(Q(u, \tau, \varepsilon)) \subset Q(\theta^T u, \tau-2T, \delta)$, and therefore

$$\begin{aligned} v_\tau^\varepsilon(u) &= \log \text{vol}(Q(u, \tau, \varepsilon)) \leq \log \text{vol}(\varphi_{T,u}^{-1}(Q(\theta^T u, \tau-2T, \delta))) \\ &\leq \log \max_{(u,x) \in \mathcal{U} \times N_\delta(Q)} |\det D\varphi_{T,u}^{-1}(x)| + v_{\tau-2T}^\delta(\theta^T u), \end{aligned}$$

which completes the proof of (d). \square

We do not know if the functions w_τ^δ are continuous, which would be desirable to carry out the proofs in the following subsections. This can be compensated, however, by the following two lemmas.

4.3 Lemma: *The function $u \mapsto \text{vol}(Q(u, \tau, \varepsilon))$ is continuous for all $\tau \in \mathbb{Z}_{>0}$ and $\varepsilon > 0$.*

Proof: Putting $Q^\varepsilon := N_\varepsilon(Q)$ and $A_t(u) := N_\varepsilon(Q(\theta^t u))$, we write the volume of $Q(u, \tau, \varepsilon)$ as

$$\text{vol}(Q(u, \tau, \varepsilon)) = \int_{Q^\varepsilon} \mathbb{1}_{A_0(u)}(x) \mathbb{1}_{A_1(u)}(\varphi_{1,u}(x)) \cdots \mathbb{1}_{A_{\tau-1}(u)}(\varphi_{\tau-1,u}(x)) \, dx.$$

For brevity, we write $g_t(u, x) := \mathbb{1}_{A_t(u)}(\varphi_{t,u}(x))$. We fix $u \in \mathcal{U}$ and prove the continuity of $\text{vol}(Q(\cdot, \tau, \varepsilon))$ at u . To this end, first observe that for arbitrary $\tilde{u} \in \mathcal{U}$ we have

$$\begin{aligned} &|\text{vol}(Q(u, \tau, \varepsilon)) - \text{vol}(Q(\tilde{u}, \tau, \varepsilon))| \\ &\leq \left| \int_{Q^\varepsilon} (g_0(u, x) g_1(u, x) \cdots g_{\tau-1}(u, x) - g_0(\tilde{u}, x) g_1(u, x) \cdots g_{\tau-1}(u, x)) \, dx \right| \\ &+ \left| \int_{Q^\varepsilon} (g_0(\tilde{u}, x) g_1(u, x) \cdots g_{\tau-1}(u, x) \right. \\ &\quad \left. - g_0(\tilde{u}, x) g_1(\tilde{u}, x) g_2(u, x) \cdots g_{\tau-1}(u, x)) \, dx \right| \\ &+ \cdots \\ &+ \left| \int_{Q^\varepsilon} (g_0(\tilde{u}, x) \cdots g_{\tau-1}(\tilde{u}, x) g_{\tau-1}(u, x) - g_0(\tilde{u}, x) \cdots g_{\tau-1}(\tilde{u}, x)) \, dx \right| \\ &\leq \sum_{t=0}^{\tau-1} \int_{Q^\varepsilon} |g_t(u, x) - g_t(\tilde{u}, x)| \, dx. \end{aligned}$$

For a fixed $t \in [0; \tau)$, the integral

$$\int_{Q^\varepsilon} |g_t(u, x) - g_t(\tilde{u}, x)| \, dx = \int_{Q^\varepsilon} |\mathbb{1}_{A_t(u)}(\varphi_{t,u}(x)) - \mathbb{1}_{A_t(\tilde{u})}(\varphi_{t,\tilde{u}}(x))| \, dx$$

is not larger than the volume of the symmetric set difference

$$\left[\varphi_{t,u}^{-1}(A_t(u)) \setminus \varphi_{t,\tilde{u}}^{-1}(A_t(\tilde{u})) \right] \cup \left[\varphi_{t,\tilde{u}}^{-1}(A_t(\tilde{u})) \setminus \varphi_{t,u}^{-1}(A_t(u)) \right]. \quad (20)$$

We show that the volumes of these two sets become arbitrarily small as $\tilde{u} \rightarrow u$:

(i) We write the first term in (20) as

$$\varphi_{t,u}^{-1}(A_t(u)) \setminus \varphi_{t,\tilde{u}}^{-1}(A_t(\tilde{u})) = \varphi_{t,u}^{-1} \left(A_t(u) \setminus \varphi_{t,u}(\varphi_{t,\tilde{u}}^{-1}(A_t(\tilde{u}))) \right).$$

Since u is fixed, it suffices to show that the volume of $A_t(u) \setminus \varphi_{t,u}(\varphi_{t,\tilde{u}}^{-1}(A_t(\tilde{u})))$ tends to zero as $\tilde{u} \rightarrow u$. Using the notation $I_\rho(B) := \{x \in \text{int } B : \text{dist}(x, \partial B) \geq \rho\}$ for any subset $B \subset M$, it is enough to show that

$$A_t(u) \setminus \varphi_{t,u}(\varphi_{t,\tilde{u}}^{-1}(A_t(\tilde{u}))) \subset A_t(u) \setminus I_\rho(A_t(u)) \quad (21)$$

for an arbitrarily small $\rho > 0$ as $\tilde{u} \rightarrow u$, by continuity of the measure and $\text{vol}(\partial A_t(u)) = 0$ (see Lemma A.2). The inclusion (21) is implied by

$$\varphi_{t,\tilde{u}} \circ \varphi_{t,u}^{-1}(I_\rho(A_t(u))) \subset A_t(\tilde{u}) = N_\varepsilon(Q(\theta^t \tilde{u})).$$

Take $x \in I_\rho(A_t(u))$ and let $y \in Q(\theta^t u)$ be a point that minimizes the distance $d(x, y)$, i.e., $d(x, y) = \text{dist}(x, Q(\theta^t u))$. Let $\tilde{y} \in Q(\theta^t \tilde{u})$ be chosen so that $d(y, \tilde{y}) \leq d_H(Q(\theta^t u), Q(\theta^t \tilde{u}))$. Then

$$\begin{aligned} d(\varphi_{t,\tilde{u}} \circ \varphi_{t,u}^{-1}(x), \tilde{y}) &\leq d(\varphi_{t,\tilde{u}} \circ \varphi_{t,u}^{-1}(x), x) \\ &\quad + d(x, y) + d_H(Q(\theta^t \tilde{u}), Q(\theta^t u)). \end{aligned}$$

If we can show that this sum becomes smaller than ε (independently of the choice of x) as $d_{\mathcal{U}}(u, \tilde{u})$ becomes sufficiently small, we are done. The third term becomes small by continuity of $Q(\cdot)$ and θ . The first term becomes small by the continuity properties of φ . Indeed, $\varphi(t, \cdot, \cdot)$ is uniformly continuous on an appropriately chosen compact set, showing that $d(\varphi_{t,\tilde{u}}(\varphi_{t,u}^{-1}(x)), \varphi_{t,u}(\varphi_{t,\tilde{u}}^{-1}(x))) \rightarrow 0$ as $\tilde{u} \rightarrow u$, uniformly with respect to x . Now $x \in I_\rho(A_t(u))$ implies that the second term is smaller than and uniformly bounded away from ε . This implies the assertion.

(ii) Consider now the second term in (20). Writing

$$\varphi_{t,\tilde{u}}^{-1}(A_t(\tilde{u})) \setminus \varphi_{t,u}^{-1}(A_t(u)) = \varphi_{t,u}^{-1}(\varphi_{t,u} \circ \varphi_{t,\tilde{u}}^{-1}(A_t(\tilde{u})) \setminus A_t(u)),$$

we see that it suffices to prove that the volume of $\varphi_{t,u} \circ \varphi_{t,\tilde{u}}^{-1}(A_t(\tilde{u})) \setminus A_t(u)$ tends to zero as $\tilde{u} \rightarrow u$. From the continuity of φ it follows that

$$\varphi_{t,u} \circ \varphi_{t,\tilde{u}}^{-1}(A_t(\tilde{u})) \subset N_\rho(A_t(\tilde{u}))$$

for any given $\rho > 0$ if $d_{\mathcal{U}}(\tilde{u}, u)$ is sufficiently small. Hence,

$$\varphi_{t,u} \circ \varphi_{t,\tilde{u}}^{-1}(A_t(\tilde{u})) \setminus A_t(u) \subset N_{\rho+\varepsilon}(Q(\theta^t \tilde{u})) \setminus N_{\varepsilon}(Q(\theta^t u)).$$

From the Hausdorff convergence $Q(\theta^t \tilde{u}) \rightarrow Q(\theta^t u)$, it follows that $N_{\rho+\varepsilon}(Q(\theta^t \tilde{u})) \subset N_{2\rho+\varepsilon}(Q(\theta^t u))$ if $d_{\mathcal{U}}(u, \tilde{u})$ is small enough, implying

$$\varphi_{t,u} \circ \varphi_{t,\tilde{u}}^{-1}(A_t(\tilde{u})) \setminus A_t(u) \subset N_{2\rho+\varepsilon}(Q(\theta^t u)) \setminus N_{\varepsilon}(Q(\theta^t u)).$$

By continuity of the measure and Lemma A.2, the volume of the right-hand side certainly tends to zero as $\rho \rightarrow 0$.

The proof is complete. \square

4.4 Lemma: For every $\delta > 0$ small enough, there exist constants $-\infty < \underline{w} < 0 < \overline{w} < \infty$ such that

$$\underline{w} \leq \frac{1}{\tau} w_{\tau}^{\delta}(u) \leq \overline{w} \quad \text{for all } (\tau, u) \in \mathbb{Z}_{>0} \times \mathcal{U}.$$

Proof: By item (c) of Proposition 4.2, we can choose $\delta > 0$ small enough so that

$$\begin{aligned} \frac{1}{\tau} w_{\tau}^{\delta}(u) &\leq \frac{1}{\tau} \log C_{\delta/2} + 1 + \frac{1}{\tau} v_{\tau}^{\delta/2}(u) \\ &\leq \log C_{\delta/2} + 1 + \max\{0, \log \text{vol}(N_{\delta/2}(Q))\} =: \overline{w} < \infty. \end{aligned}$$

On the other hand, the definition of w^{δ} implies

$$\begin{aligned} \frac{1}{\tau} w_{\tau}^{\delta}(u) &\geq \frac{1}{\tau} \inf \left\{ \log |\alpha| + \log \min_{(u,x) \in L(Q)} J^+ \varphi_{\tau,u}(x)^{-1} : \alpha \dots \right\} \\ &\geq \frac{1}{\tau} \log \min_{(u,x) \in L(Q)} J^+ \varphi_{\tau,u}(x)^{-1} \geq \min_{(u,x) \in L(Q)} \log J^+ \varphi_{1,u}(x)^{-1} \\ &=: \underline{w} > -\infty. \end{aligned}$$

This completes the proof. \square

4.4 Interchanging limit inferior and supremum

Recall that for all compact sets $K \subset Q$ of positive volume, in Lemma 4.1 we have proved the estimate

$$h_{\text{inv}}(K, Q) \geq - \liminf_{\tau \rightarrow \infty} \sup_{u \in \mathcal{U}} \frac{1}{\tau} v_{\tau}^{\varepsilon}(u).$$

Our next aim is to prove that the limit inferior and the supremum on the right-hand side can be interchanged. First observe that the estimate

$$\liminf_{\tau \rightarrow \infty} \sup_{u \in \mathcal{U}} \frac{1}{\tau} v_{\tau}^{\varepsilon}(u) \geq \sup_{u \in \mathcal{U}} \liminf_{\tau \rightarrow \infty} \frac{1}{\tau} v_{\tau}^{\varepsilon}(u)$$

is trivial on the one hand and useless on the other, since for obtaining a lower estimate of $h_{\text{inv}}(K, Q)$ only the converse inequality can be used. The following proposition shows that under the limit for $\varepsilon \downarrow 0$, the converse inequality holds.

4.5 Proposition: *Under the assumptions (A1)–(A3), for any compact set $K \subset Q$ of positive volume*

$$h_{\text{inv}}(K, Q) \geq - \limsup_{\varepsilon \downarrow 0} \liminf_{u \in \mathcal{U}} \liminf_{\tau \rightarrow \infty} \frac{1}{\tau} \log \text{vol}(Q(u, \tau, \varepsilon)). \quad (22)$$

Proof: Fix $\gamma > 0$ and choose $\delta = \delta(\gamma) > 0$ according to Proposition 4.2(c). Then choose $\varepsilon = \varepsilon(\delta) \in (0, \delta/2)$ according to Proposition 4.2(b). In particular, this implies

$$v_\tau^\varepsilon(u) - \log C_{3\delta} \leq w_\tau^\delta(u) \leq \tau\gamma + \log C_{\delta/2} + v_\tau^{\delta/2}(u) \quad (23)$$

for all $u \in \mathcal{U}$ and $\tau \in \mathbb{Z}_{>0}$. We define

$$S := \sup \left\{ \lambda \in \mathbb{R} : \exists u_k \in \mathcal{U}, t_k \rightarrow \infty \text{ with } \lambda = \lim_{k \rightarrow \infty} \frac{1}{t_k} v_{t_k}^\varepsilon(u_k) \right\}.$$

This number is finite, since (23) together with Lemma 4.4 implies

$$\frac{1}{t} v_t^\varepsilon(u) \leq \log C_{3\delta} + \bar{w} \quad \text{for all } t \geq 1.$$

Moreover, S is independent of ε (as long as ε is small enough), which follows from Proposition 4.2(d). Now consider a sequence $\rho_k \downarrow 0$ and sequences of $u_k \in \mathcal{U}$ and $t_k \rightarrow \infty$ such that

$$\frac{1}{t_k} w_{t_k}^\delta(u_k) > S - \rho_k \quad \text{for all } k \geq 0,$$

which is possible by (23). We put $\tilde{\rho}_k := 1/\sqrt{t_k}$. By Lemma A.3, we find times $t_k^* < t_k$ so that

$$\frac{1}{l} w_l^\delta(\theta^{t_k^*} u_k) > S - \rho_k - \tilde{\rho}_k \quad \text{for } 0 < l \leq t_k - t_k^*,$$

where $t_k - t_k^* \geq \sqrt{t_k}/(2\omega)$ and $\omega = \max\{-\underline{w}, \bar{w}\}$ (see Lemma 4.4). Using (23) again, this leads to

$$\frac{1}{l} v_l^{\delta/2}(\theta^{t_k^*} u_k) > S - \rho_k - \tilde{\rho}_k - \frac{1}{l} \log C_{\delta/2} - \gamma \quad \text{for } 0 < l \leq t_k - t_k^*.$$

We put $\tilde{u}_k := \theta^{t_k^*} u_k$, $\tilde{t}_k := t_k - t_k^* \rightarrow \infty$. By compactness, we may assume that $\tilde{u}_k \rightarrow \tilde{u}$ for some $\tilde{u} \in \mathcal{U}$. Fix $t \in \mathbb{Z}_{>0}$ and $\rho > 0$. Then, for k large enough, $t \leq \tilde{t}_k$ and, by continuity of $v_t^{\delta/2}(\cdot)$ (see Lemma 4.3),

$$|v_t^{\delta/2}(\tilde{u}) - v_t^{\delta/2}(\tilde{u}_k)| < \rho.$$

We thus obtain

$$\begin{aligned}\frac{1}{t}v_t^{\delta/2}(\tilde{u}) &= \frac{1}{t}v_t^{\delta/2}(\tilde{u}_k) + \left(\frac{1}{t}v_t^{\delta/2}(\tilde{u}) - \frac{1}{t}v_t^{\delta/2}(\tilde{u}_k)\right) \\ &> S - \rho_k - \tilde{\rho}_k - \frac{1}{t}\log C_{\delta/2} - \gamma - \frac{\rho}{t}.\end{aligned}$$

Letting $t \rightarrow \infty$, this yields

$$\liminf_{t \rightarrow \infty} \frac{1}{t}v_t^{\delta/2}(\tilde{u}) \geq S - \gamma.$$

Now choose for each $t \in \mathbb{Z}_{>0}$ some $u_t^* \in \mathcal{U}$ with $\sup_{u \in \mathcal{U}} v_t^\varepsilon(u)/t = v_t^\varepsilon(u_t^*)/t$, which is possible by continuity of $v_t^\varepsilon(\cdot)$. Then, using Proposition 4.2(d),

$$\begin{aligned}\liminf_{t \rightarrow \infty} \sup_{u \in \mathcal{U}} \frac{1}{t}v_t^\varepsilon(u) &= \liminf_{t \rightarrow \infty} \frac{1}{t}v_t^\varepsilon(u_t^*) \\ &\leq S \leq \gamma + \liminf_{t \rightarrow \infty} \frac{1}{t}v_t^{\delta/2}(\tilde{u}) \\ &\leq \gamma + \sup_{u \in \mathcal{U}} \liminf_{t \rightarrow \infty} \frac{1}{t}v_t^{\delta/2}(u) \\ &\leq \gamma + \sup_{u \in \mathcal{U}} \liminf_{t \rightarrow \infty} \frac{1}{t}(\tilde{C}_\varepsilon + v_{t-2T}^\varepsilon(u)) \\ &= \gamma + \sup_{u \in \mathcal{U}} \liminf_{t \rightarrow \infty} \frac{1}{t}v_t^\varepsilon(u).\end{aligned}$$

Together with the estimate of Lemma 4.1, this yields

$$h_{\text{inv}}(K, Q) \geq -\liminf_{t \rightarrow \infty} \sup_{u \in \mathcal{U}} \frac{1}{t}v_t^\varepsilon(u) \geq -\gamma - \sup_{u \in \mathcal{U}} \liminf_{t \rightarrow \infty} \frac{1}{t}v_t^\varepsilon(u).$$

We can choose γ arbitrarily small, which also enforces ε to become arbitrarily small. Hence, the desired inequality follows. \square

4.5 An estimate in terms of random escape rates

Our next goal is to replace the supremum over $u \in \mathcal{U}$ in the right-hand side of (22) by a supremum over all θ -invariant probability measures to obtain the estimate

$$h_{\text{inv}}(K, Q) \geq -\lim_{\varepsilon \downarrow 0} \sup_{P \in \mathcal{M}(\theta)} \liminf_{\tau \rightarrow \infty} \frac{1}{\tau} \int \log \text{vol}(Q(u, \tau, \varepsilon)) dP(u).$$

Once this is accomplished, we can prove the desired lower bound (16) in terms of pressure by standard methods from thermodynamic formalism.

Before we prove the desired inequality, we note that any limit of the form

$$\lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int \log \text{vol}(Q(u, \tau, \varepsilon)) dP(u),$$

if it exists, is called a *random escape rate* for the RDS (φ, P) (see [48]).

The main ideas of the proof of the following proposition are taken from [52, Lem. A.6] (a result on abstract subadditive cocycles).

4.6 Proposition: *Under the assumptions (A1)–(A3), for any compact set $K \subset Q$ of positive volume*

$$h_{\text{inv}}(K, Q) \geq -\lim_{\varepsilon \downarrow 0} \sup_{P \in \mathcal{M}(\theta)} \liminf_{\tau \rightarrow \infty} \frac{1}{\tau} \int \log \text{vol}(Q(u, \tau, \varepsilon)) \, dP(u). \quad (24)$$

Proof: We fix $\gamma > 0$, choose $\delta = \delta(\gamma) > 0$ according to Proposition 4.2(c) and $\varepsilon = \varepsilon(\delta) > 0$ according to Proposition 4.2(b). Then we pick an arbitrary $u \in \mathcal{U}$ and let

$$\beta := \liminf_{t \rightarrow \infty} \frac{1}{t} v_t^\varepsilon(u).$$

Now we consider the sequence of Borel probability measures on the measurable space $(\mathcal{U}, \mathcal{B}(\mathcal{U}))$ defined by

$$P_t := \frac{1}{t} \sum_{s=0}^{t-1} \delta_{\theta^s u}, \quad t \in \mathbb{Z}_{>0}.$$

Since \mathcal{U} is compact, there exists a weak* limit point P of $(P_t)_{t>0}$. With standard arguments, one shows that P is θ -invariant. Then the following chain of inequalities holds for any fixed $r \in \mathbb{Z}_{>0}$:

$$\begin{aligned} \beta &\leq \liminf_{t \rightarrow \infty} \frac{1}{t} w_t^\delta(u) \\ &\leq \liminf_{t \rightarrow \infty} \frac{1}{tr} \sum_{s=0}^{t-r} w_r^\delta(\theta^s u) \\ &= \liminf_{t \rightarrow \infty} \frac{1}{tr} \sum_{s=0}^{t-1} w_r^\delta(\theta^s u) \\ &\leq \frac{1}{r} \log C_{\delta/2} + \gamma + \liminf_{t \rightarrow \infty} \frac{1}{tr} \sum_{s=0}^{t-1} v_r^{\delta/2}(\theta^s u) \\ &= \frac{1}{r} \log C_{\delta/2} + \gamma + \liminf_{t \rightarrow \infty} \frac{1}{r} \int v_r^{\delta/2} \, dP_t. \end{aligned}$$

The first line follows from Proposition 4.2(b) and the fourth from item (c) of the same proposition. The third line uses that w_r^δ is bounded on \mathcal{U} and the last line simply uses the definition of P_t . It remains to prove the second inequality. To this end, for each s in the range $0 \leq s < r$ let us choose integers q_s, r_s such that $t = s + q_s r + r_s$ with $q_s \geq 0$ and $0 \leq r_s < r$. By Lemma A.4,

$$\sum_{s=0}^{r-1} \sum_{j=0}^{q_s-1} w_r^\delta(\theta^{s+jr} u) = \sum_{s=0}^{t-r} w_r^\delta(\theta^s u).$$

Hence, using subadditivity, we find that

$$\begin{aligned} rw_t^\delta(u) &\leq \sum_{s=0}^{r-1} \left(w_s^\delta(u) + \sum_{j=0}^{q_s-1} w_r^\delta(\theta^{s+jr}u) + w_{r_s}^\delta(\theta^{s+q_sr}u) \right) \\ &= \sum_{s=0}^{r-1} w_s^\delta(u) + \sum_{s=0}^{t-r} w_r^\delta(\theta^s u) + \sum_{s=0}^{r-1} w_{r_s}^\delta(\theta^{t-rs} u). \end{aligned}$$

Dividing both sides by tr and letting $t \rightarrow \infty$ completes the proof of the second inequality above. We have thus proved the estimate

$$\liminf_{t \rightarrow \infty} \frac{1}{r} \int v_r^{\delta/2} dP_t \geq \beta - \gamma - \frac{1}{r} \log C_{\delta/2}$$

for all $r \in \mathbb{Z}_{>0}$. By continuity of $v_r^{\delta/2}(\cdot)$, this implies

$$\frac{1}{r} \int v_r^{\delta/2} dP \geq \beta - \gamma - \frac{1}{r} \log C_{\delta/2}.$$

According to Proposition 4.2(d), choose $T \in \mathbb{Z}_{>0}$ such that $v_r^{\delta/2}(u) \leq \tilde{C} + v_{r-2T}^\varepsilon(\theta^T u)$, which yields

$$\frac{\tilde{C}}{r} + \frac{1}{r} \int v_{r-2T}^\varepsilon dP \geq \beta - \gamma - \frac{1}{r} \log C_{\delta/2},$$

where we use that P is θ -invariant. Letting $r \rightarrow \infty$, we arrive at

$$\beta \leq \gamma + \liminf_{r \rightarrow \infty} \frac{1}{r} \int v_r^\varepsilon dP,$$

which implies

$$\sup_{u \in \mathcal{U}} \liminf_{t \rightarrow \infty} \frac{1}{t} v_t^\varepsilon(u) \leq \gamma + \sup_{P \in \mathcal{M}(\theta)} \liminf_{t \rightarrow \infty} \frac{1}{t} \int v_t^\varepsilon dP.$$

Since γ can be chosen arbitrarily small, this together with Proposition 4.5 leads to the desired estimate. \square

4.6 An estimate in terms of pressure

To complete the proof of the lower bound, we need to relate the random escape rate bound from Proposition 4.6 to the pressure of the associated random dynamical systems. This is accomplished by the following theorem whose proof follows the proof of the variational principle for the pressure of random dynamical systems [7]. The idea to use these arguments to compute escape rates can be found in many works, including [9, 48, 67].

4.7 Theorem: Assume that the control system Σ is of regularity class C^2 and let Q be a compact all-time controlled invariant set of Σ satisfying the following assumptions:

- (A1) Q is uniformly hyperbolic.
- (A2) $L(Q)$ is an isolated invariant set of the control flow.
- (A3) The fiber map $u \mapsto Q(u)$ is lower semicontinuous.

Then for every compact set $K \subset Q$ of positive volume, the invariance entropy satisfies

$$h_{\text{inv}}(K, Q) \geq \inf_{\mu \in \mathcal{M}(\Phi|_{L(Q)})} \left[\int \log J^+ \varphi_{1,u}(x) d\mu(u, x) - h_\mu(\varphi, (\pi_{\mathcal{U}})_* \mu) \right]. \quad (25)$$

Proof: To simplify some arguments, we assume without loss of generality that the manifold M is compact. Fix some $P \in \mathcal{M}(\theta)$ and sufficiently small $\varepsilon, \delta > 0$. Let $F_{u,t,\delta} \subset Q(u)$ be a maximal (u, t, δ) -separated set for each $u \in \mathcal{U}$ and $t \in \mathbb{Z}_{>0}$. By (15), this implies

$$v_t^\varepsilon(u) \leq \log C_{\beta+\delta} + \log \sum_{x \in F_{u,t,\delta}} J^+ \varphi_{t,u}(x)^{-1}. \quad (26)$$

We define sequences of probability measures on $(M, \mathcal{B}(M))$ by

$$\eta_t^u := \frac{\sum_{x \in F_{u,t,\delta}} 2^{-\log J^+ \varphi_{t,u}(x)} \delta_x}{\sum_{x \in F_{u,t,\delta}} 2^{-\log J^+ \varphi_{t,u}(x)}}, \quad t \in \mathbb{Z}_{>0}, u \in \mathcal{U}$$

and

$$\nu_t^u := \frac{1}{t} \sum_{s=0}^{t-1} \varphi(-s, \cdot, u)_*^{-1} \eta_t^{\theta^{-s}u}, \quad t \in \mathbb{Z}_{>0}.$$

We can choose the sets $F_{u,t,\delta}$ such that η_t^u depends measurably on u (see [7, Proof of Thm. 6.1]), implying that we can define probability measures σ_t on $\mathcal{U} \times M$ by $d\sigma_t(u, x) := d\eta_t^u(x) dP(u)$.

Observe that for any $A \in \mathcal{B}(\mathcal{U} \times M)$ we have

$$\begin{aligned} \frac{1}{t} \sum_{s=0}^{t-1} (\Phi_s)_* \sigma_t(A) &= \frac{1}{t} \sum_{s=0}^{t-1} \sigma_t(\Phi_s^{-1}(A)) \\ &= \frac{1}{t} \sum_{s=0}^{t-1} \int_{\mathcal{U}} \int_M \mathbb{1}_{\Phi_s^{-1}(A)}(u, x) d\eta_t^u(x) dP(u) \\ &= \frac{1}{t} \sum_{s=0}^{t-1} \int_{\mathcal{U}} \int_M \mathbb{1}_A(\theta^s u, \varphi(s, x, u)) d\eta_t^u(x) dP(u) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{t} \sum_{s=0}^{t-1} \int_{\mathcal{U}} \int_M \mathbb{1}_A(v, \varphi(s, x, \theta^{-s}v)) \, d\eta_t^{\theta^{-s}v}(x) dP(v) \\
&= \frac{1}{t} \sum_{s=0}^{t-1} \int_{\mathcal{U}} \int_M \mathbb{1}_A(v, y) \, d[\varphi(s, \cdot, \theta^{-s}v)_* \eta_t^{\theta^{-s}v}](y) dP(v) \\
&= \int_{\mathcal{U}} \int_M \mathbb{1}_A(v, y) \, d\nu_t^v(y) dP(v).
\end{aligned}$$

Hence, the measures ν_t^u are the sample measures of $\mu_t := \frac{1}{t} \sum_{s=0}^{t-1} (\Phi_s)_* \sigma_t$. By weak* compactness, there exists a limit point μ of the sequence $(\mu_t)_{t>0}$. Then μ is a Φ -invariant measure with marginal P on \mathcal{U} , i.e., an invariant measure of the random dynamical system (φ, P) . Indeed, for any $g \in C^0(\mathcal{U} \times M, \mathbb{R})$ and an appropriate subsequence $(t_k)_{k>0}$, we have

$$\begin{aligned}
(\Phi_* \mu - \mu)(g) &= \lim_{k \rightarrow \infty} \left| \int_{\mathcal{U} \times M} g(\Phi(u, x)) \, d\mu_{t_k}(u, x) - \int_{\mathcal{U} \times M} g(u, x) \, d\mu_{t_k}(u, x) \right| \\
&= \lim_{k \rightarrow \infty} \left| \frac{1}{t_k} \sum_{s=0}^{t_k-1} \int_{\mathcal{U}} \int_M (g(\Phi(u, x)) - g(u, x)) \, d[\varphi(-s, \cdot, u)_*^{-1} \eta_{t_k}^{\theta^{-s}u}](x) dP(u) \right| \\
&= \lim_{k \rightarrow \infty} \left| \frac{1}{t_k} \sum_{s=0}^{t_k-1} \int_{\mathcal{U}} \int_M (g(\Phi(\theta^s u, x)) - g(\theta^s u, x)) \, d[(\varphi_{s,u})_* \eta_{t_k}^u](x) dP(u) \right| \\
&= \lim_{k \rightarrow \infty} \left| \frac{1}{t_k} \sum_{s=0}^{t_k-1} \int_{\mathcal{U}} \int_M (g(\Phi_{s+1}(u, x)) - g(\Phi_s(u, x))) \, d\eta_{t_k}^u(x) dP(u) \right| \\
&= \lim_{k \rightarrow \infty} \frac{1}{t_k} \left| \int_{\mathcal{U}} \left[\int_M g(\Phi_{t_k}(u, x)) \, d\eta_{t_k}^u(x) - \int_M g(u, x) \, d\eta_{t_k}^u(x) \right] dP(u) \right| \\
&\leq \lim_{k \rightarrow \infty} \frac{1}{t_k} \int_{\mathcal{U}} \int_M |g(\Phi_{t_k}(u, x)) - g(u, x)| \, d\eta_{t_k}^u(x) dP(u) \\
&\leq \lim_{k \rightarrow \infty} \frac{2}{t_k} \max_{(u,x) \in \mathcal{U} \times M} |g(u, x)| = 0,
\end{aligned}$$

showing that μ is Φ -invariant. The fact that $(\pi_{\mathcal{U}})_* \mu = P$ follows from the continuity of the operator $(\pi_{\mathcal{U}})_*$.

By Lemma A.5, we can choose a finite Borel partition $\mathcal{P} = \{P_1, \dots, P_k\}$ of M with $\text{diam}(P_i) < \delta$ and $(\pi_M)_* \mu(\partial P_i) = 0$ for $i = 1, \dots, k$. Since $(\pi_M)_* \mu(\partial P_i) = \int \mu^u(\partial P_i) \, dP(u)$, where μ^u are the sample measures of μ , we have $\mu^u(\partial P_i) = 0$ for P -almost all $u \in \mathcal{U}$.

Put $\gamma_t(u, x) := -\log J^+ \varphi_{t,u}(x)$ and $S_t(u) := \sum_{x \in F_{u,t,\delta}} 2^{\gamma_t(u,x)}$. Since each element of $\bigvee_{s=0}^{t-1} \varphi(s, \cdot, u)^{-1} \mathcal{P}$ contains at most one element of $F_{u,t,\delta}$, we obtain for P -almost all $u \in \mathcal{U}$ that

$$H_{\eta_t^u} \left(\bigvee_{s=0}^{t-1} \varphi_{s,u}^{-1} \mathcal{P} \right) - \int (-\gamma_t(u, x)) \, d\eta_t^u(x)$$

$$\begin{aligned}
&= - \sum_{x \in F_{u,t,\delta}} \frac{2^{\gamma_t(u,x)}}{S_t(u)} \log \frac{2^{\gamma_t(u,x)}}{S_t(u)} + \sum_{x \in F_{u,t,\delta}} \frac{2^{\gamma_t(u,x)}}{S_t(u)} \log 2^{\gamma_t(u,x)} \\
&= \sum_{x \in F_{u,t,\delta}} \frac{2^{\gamma_t(u,x)}}{S_t(u)} \log S_t(u) = \log S_t(u).
\end{aligned}$$

Now consider $q, t \in \mathbb{Z}_{>0}$ with $1 < q < t$ and let $a(r)$ denote the integer part of $(t-r)/q$ for $0 \leq r \leq q-1$. Then

$$\bigvee_{s=0}^{t-1} \varphi_{s,u}^{-1} \mathcal{P} = \bigvee_{i=0}^{a(r)-1} \varphi_{r+iq,u}^{-1} \bigvee_{j=0}^{q-1} \varphi_{j,\theta^{r+iq}u}^{-1} \mathcal{P} \vee \bigvee_{s \in R} \varphi_{s,u}^{-1} \mathcal{P},$$

where the set R satisfies $|R| \leq 2q$. Hence, using elementary properties of Shannon entropy, we conclude that

$$H_{\eta_t^u} \left(\bigvee_{s=0}^{t-1} \varphi_{s,u}^{-1} \mathcal{P} \right) \leq \sum_{i=0}^{a(r)-1} H_{(\varphi_{r+iq,u})_* \eta_t^u} \left(\bigvee_{j=0}^{q-1} \varphi_{j,\theta^{r+iq}u}^{-1} \mathcal{P} \right) + 2q \log k.$$

Summing over $r = 0, 1, \dots, q-1$, we obtain

$$\begin{aligned}
q \log S_t(u) &\leq \sum_{s=0}^{t-1} H_{(\varphi_{s,u})_* \eta_t^u} \left(\bigvee_{j=0}^{q-1} \varphi_{j,\theta^s u}^{-1} \mathcal{P} \right) \\
&\quad + 2q^2 \log k - q \int (-\gamma_t(u, x)) \, d\eta_t^u(x).
\end{aligned} \tag{27}$$

Using the notation

$$h_{s,q}^t(u) := H_{(\varphi_{-s,u}^{-1})_* \eta_t^{\theta^{-s}u}} \left(\bigvee_{j=0}^{q-1} \varphi_{j,u}^{-1} \mathcal{P} \right),$$

we find that¹⁶

$$\frac{1}{t} \sum_{s=0}^{t-1} h_{s,q}^t(u) \leq H_{\nu_t^u} \left(\bigvee_{j=0}^{q-1} \varphi_{j,u}^{-1} \mathcal{P} \right)$$

and

$$\frac{1}{t} \sum_{s=0}^{t-1} H_{(\varphi_{s,u})_* \eta_t^u} \left(\bigvee_{j=0}^{q-1} \varphi_{j,\theta^s u}^{-1} \mathcal{P} \right) = \frac{1}{t} \sum_{s=0}^{t-1} h_{s,q}^t(\theta^s u).$$

Integrating both sides over u and using θ -invariance of P leads to

$$\frac{1}{t} \sum_{s=0}^{t-1} \int H_{(\varphi_{s,u})_* \eta_t^u} \left(\bigvee_{j=0}^{q-1} \varphi_{j,\theta^s u}^{-1} \mathcal{P} \right) \, dP(u) \leq \int H_{\nu_t^u} \left(\bigvee_{j=0}^{q-1} \varphi_{j,u}^{-1} \mathcal{P} \right) \, dP(u).$$

¹⁶Here we use the elementary property $\sum_i \alpha_i H_{\mu_i}(\mathcal{P}) \leq H_{\sum_i \alpha_i \mu_i}(\mathcal{P})$ of Shannon entropy for convex combinations of measures.

Dividing (27) by t and integrating, we thus obtain

$$\begin{aligned} \frac{q}{t} \int \log S_t(u) \, dP(u) &\leq \int H_{\nu_t^u} \left(\bigvee_{j=0}^{q-1} \varphi_{j,u}^{-1} \mathcal{P} \right) dP(u) \\ &\quad + 2 \frac{q^2}{t} \log k - q \int (-\gamma_1(u, x)) \, d\mu_t(u, x), \end{aligned} \quad (28)$$

where we use that

$$\begin{aligned} \frac{1}{t} \int \int \gamma_t(u, x) \, d\eta_t^u(x) dP(u) &= \frac{1}{t} \int \int \sum_{s=0}^{t-1} \gamma_1(\Phi_s(u, x)) \, d\eta_t^u(x) dP(u) \\ &= \frac{1}{t} \sum_{s=0}^{t-1} \int \int \gamma_1(\theta^s u, x) \, d[\varphi(s, \cdot, u)_* \eta_t^u](x) dP(u) \\ &= \frac{1}{t} \sum_{s=0}^{t-1} \int \int \gamma_1(\theta^s u, x) \, d[\varphi(-s, \cdot, \theta^s u)_*^{-1} \eta_t^u](x) dP(u) \\ &= \int \int \gamma_1(\theta^s u, x) \, d\nu_t^{\theta^s u}(x) dP(u) \\ &= \int \int \gamma_1(u, x) \, d\nu_t^u(x) dP(u) = \int \gamma_1(u, x) \, d\mu_t(u, x). \end{aligned}$$

Letting $t \rightarrow \infty$ (respectively, an appropriate subsequence) in (28), we thus obtain

$$\begin{aligned} q \liminf_{t \rightarrow \infty} \frac{1}{t} \int \log S_t(u) \, dP(u) &\leq \int H_{\mu^u} \left(\bigvee_{j=0}^{q-1} \varphi_{j,u}^{-1} \mathcal{P} \right) dP(u) \\ &\quad - q \int (-\gamma_1(u, x)) \, d\mu(u, x), \end{aligned}$$

where we use that γ_1 is continuous and $\mu^u(\partial P_i) = 0$ for P -almost all u and all $P_i \in \mathcal{P}$. Using (26), we arrive at

$$\begin{aligned} &\liminf_{t \rightarrow \infty} \frac{1}{t} \int v_t^\varepsilon(u) \, dP(u) \\ &\leq \liminf_{t \rightarrow \infty} \frac{1}{t} \int \log S_t(u) \, dP(u) \\ &\leq \frac{1}{q} \int H_{\mu^u} \left(\bigvee_{j=0}^{q-1} \varphi_{j,u}^{-1} \mathcal{P} \right) dP(u) - \int \log J^+ \varphi_{1,u}(x) \, d\mu(u, x). \end{aligned}$$

Since this estimate holds for all $q \in \mathbb{Z}_{>0}$, we can let $q \rightarrow \infty$ and obtain

$$\begin{aligned} \liminf_{t \rightarrow \infty} \frac{1}{t} \int v_t^\varepsilon(u) \, dP(u) &\leq h_\mu(\varphi, P; \mathcal{P}) - \int \log J^+ \varphi_{1,u}(x) \, d\mu(u, x) \\ &\leq h_\mu(\varphi, P) - \int \log J^+ \varphi_{1,u}(x) \, d\mu(u, x). \end{aligned}$$

It remains to show that μ is supported on $L(Q)$. By construction, $\text{supp}(\sigma_t) \subset L(Q)$ for every $t \in \mathbb{Z}_{>0}$, which implies $\text{supp}(\mu_t) \subset L(Q)$ by Φ -invariance of $L(Q)$, and consequently $\text{supp}(\mu) \subset L(Q)$. Together with Proposition 4.6, this yields the desired estimate. \square

4.7 Optimal measures

A natural question that arises from the estimate (25) is whether the infimum is attained as a minimum. Indeed, we will show that this always holds, which leads to an interesting conclusion.

A sufficient condition for the existence of the minimum is the upper semicontinuity of the functional¹⁷

$$\mu \mapsto h_\mu(\varphi, (\pi_{\mathcal{U}})_*\mu) - \int \log J^+ \varphi \, d\mu,$$

where the space of Φ -invariant measures is equipped with the standard weak*-topology. Since the integrand in the last term is a continuous function, it follows that this term is continuous in μ . Hence, it suffices to prove the upper semicontinuity of the measure-theoretic entropy.

4.8 Lemma: *The functional $\mu \mapsto h_\mu(\varphi, (\pi_{\mathcal{U}})_*\mu)$, defined on $\mathcal{M}(\Phi|_{L(Q)})$, is upper semicontinuous.*

Proof: Throughout the proof, we say that a partition has *zero μ -boundary* if the μ -measure of the boundary of each member of the partition vanishes. The proof proceeds in two steps.

Step 1: Writing $\mathcal{F} := \pi_{\mathcal{U}}^{-1}(\mathcal{B}(\mathcal{U}))$ (which is a Φ -invariant σ -algebra on $\mathcal{U} \times M$), we will use the following alternative characterization of the measure-theoretic entropy (with respect to a partition \mathcal{P}) due to [7, Thm. 3.1], using conditional entropy:

$$h_\mu(\varphi, (\pi_{\mathcal{U}})_*\mu; \mathcal{P}) = h_\mu(\Phi; \mathcal{P}|\mathcal{F}) := \lim_{\tau \rightarrow \infty} \frac{1}{\tau} H_\mu \left(\bigvee_{s=0}^{\tau-1} \Phi_{-s}(\{\mathcal{U}\} \times \mathcal{P}) | \mathcal{F} \right). \quad (29)$$

We fix $\mu_0 \in \mathcal{M}(\Phi|_{L(Q)})$ and prove that $h_\mu(\varphi, (\pi_{\mathcal{U}})_*\mu; \mathcal{P})$ depends upper semicontinuously on μ at μ_0 if \mathcal{P} has zero $(\pi_M)_*\mu_0$ -boundary.

To this end, first note that due to subadditivity the limit in (29) can be written as the infimum over $\tau \in \mathbb{Z}_{>0}$. Since the infimum over upper semicontinuous functions is upper semicontinuous, it suffices to prove the upper semicontinuity of the function

$$\mu \mapsto H_\mu \left(\bigvee_{s=0}^{\tau-1} \Phi_{-s}(\{\mathcal{U}\} \times \mathcal{P}) | \mathcal{F} \right)$$

¹⁷Here we use that an upper semicontinuous function defined on a compact space attains its maximum.

at μ_0 for each fixed τ . By the definition of conditional entropy (see [27, Def. 1.4.5]), we have

$$H_\mu\left(\bigvee_{s=0}^{\tau-1} \Phi_{-s}(\{\mathcal{U}\} \times \mathcal{P}) \middle| \mathcal{F}\right) = \inf\left\{H_\mu\left(\bigvee_{s=0}^{\tau-1} \Phi_{-s}(\{\mathcal{U}\} \times \mathcal{P}) \middle| \mathcal{R}\right) : \mathcal{R} \preceq \mathcal{F}\right\}, \quad (30)$$

where the infimum is taken over all countable partitions \mathcal{R} whose elements belong to \mathcal{F} . Hence, it is sufficient to prove that

$$\mu \mapsto H_\mu\left(\bigvee_{s=0}^{\tau-1} \Phi_{-s}(\{\mathcal{U}\} \times \mathcal{P}) \middle| \mathcal{R}\right)$$

is upper semicontinuous for each partition \mathcal{R} as above. Recall that for any partitions \mathcal{A} and \mathcal{B} , the conditional entropy is defined by

$$H_\mu(\mathcal{A}|\mathcal{B}) = \sum_{B \in \mathcal{B}} \mu(B) H_{\mu_B}(\mathcal{A}),$$

where $\mu_B(\cdot) = \mu(\cdot \cap B)/\mu(B)$. As long as both partitions \mathcal{A} and \mathcal{B} have zero μ_0 -boundaries, the Portmanteau-Theorem tells us that $\mu \mapsto H_\mu(\mathcal{A}|\mathcal{B})$ is continuous at μ_0 . Applying this fact to our problem, we see that we are fine if we can restrict ourselves to partitions \mathcal{R} with zero μ_0 -boundaries (observing that $\{\mathcal{U}\} \times \mathcal{P}$ has zero μ_0 -boundary and thus also the joint partitions $\bigvee_{s=0}^{\tau-1} \Phi_{-s}(\{\mathcal{U}\} \times \mathcal{P})$). By a general fact, see [27, Fact 6.6.6], we can find a so-called refining sequence of partitions \mathcal{R}_k , $k \in \mathbb{Z}_{>0}$, with zero μ_0 -boundaries so that the infimum in (30) is approached along this sequence for every μ (see [27, Lem. 1.7.11]). Hence, we have proved that $h_\mu(\varphi, P; \mathcal{P})$ is upper semicontinuous at μ_0 if \mathcal{P} has zero $(\pi_M)_*\mu_0$ -boundary.

Step 2: To complete the proof, it suffices to show that for every $\mu_0 \in \mathcal{M}(\Phi|_{L(Q)})$ there exists a finite measurable partition \mathcal{P} of M with zero $(\pi_M)_*\mu_0$ -boundary so that

$$h_\mu(\varphi, (\pi_{\mathcal{U}})_*\mu) = h_\mu(\varphi, (\pi_{\mathcal{U}})_*\mu; \mathcal{P}) \quad \text{for all } \mu \in \mathcal{M}(\Phi|_{L(Q)}).$$

This follows from expansivity. Indeed, to understand this, we need to regard the restriction of Φ to $L(Q)$ as a bundle random dynamical system over the base (\mathcal{U}, θ) . Then we can write the entropy above as

$$h_\mu(\varphi, (\pi_{\mathcal{U}})_*\mu; \mathcal{P}) = \lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int H_{\mu_u}\left(\bigvee_{s=0}^{\tau-1} \varphi_{s,u}^{-1} \hat{\mathcal{P}}(\theta^s u)\right) dP(u),$$

where $\hat{\mathcal{P}}(u) = \{Q(u) \cap P : P \in \mathcal{P}\}$, see [46, Formula (1.1.5)]. We call the partition \mathcal{P} a *generator* if the sequence of partitions $\varphi_{t,u}^{-1} \hat{\mathcal{P}}(\theta^t u)$, $t \in \mathbb{Z}$, generates the Borel σ -algebra of $Q(u)$ for all $u \in \mathcal{U}$. Assume that $\text{diam}(P) < \delta$ for all $P \in \mathcal{P}$, where $\delta > 0$ is an expansivity constant for the hyperbolic set Q . Then $\hat{\mathcal{P}}^\infty(u)$ contains arbitrarily fine partitions of $Q(u)$, and thus it generates the Borel σ -algebra. By [46, Thm. 1.1.2], it follows that the entropy is attained on the partition \mathcal{P} for

every $\mu \in \mathcal{M}(\Phi|_{L(Q)})$, and from Lemma A.5 it follows that for every fixed μ_0 we can find a partition \mathcal{P} with zero $(\pi_M)_*\mu_0$ -boundary and diameter smaller than δ . \square

Hence, we have the following corollary of Theorem 4.7.

4.9 Corollary: *Let Q be a compact all-time controlled invariant set of Σ satisfying (A1)–(A3). Then there exists $\hat{\mu} \in \mathcal{M}(\Phi|_{L(Q)})$ so that for every compact set $K \subset Q$ with positive volume*

$$h_{\text{inv}}(K, Q) \geq \int \log J^+ \varphi_{1,u}(x) d\hat{\mu}(u, x) - h_{\hat{\mu}}(\varphi, (\pi_{\mathcal{U}})_*\hat{\mu}).$$

An SRB measure of an RDS is an invariant probability measure whose conditional probabilities on the unstable manifolds are absolutely continuous with respect to the Lebesgue measure on these manifolds, see [68] or [46, Def. 3.2.2] for a precise definition. SRB measures μ can also be characterized by the equality $h_{\mu} = \lambda^+(\mu)$, where $\lambda^+(\mu)$ is a short-cut for the integral over the sum of the positive Lyapunov exponents, see [5] or [46, Thm. 3.2.4]. In our case, this equality can be written as

$$h_{\mu}(\varphi, P) = \int \log J^+ \varphi d\mu.$$

This easily implies the following corollary.

4.10 Corollary: *Assume that the control system Σ is of regularity class C^2 . Let Q be a compact all-time controlled invariant set of Σ satisfying (A1)–(A3). Then $h_{\text{inv}}(K, Q) = 0$ for some compact set $K \subset Q$ of positive volume implies the existence of $P \in \mathcal{M}(\theta)$ so that the associated random dynamical system (φ, P) admits an SRB measure supported on $L(Q)$.*

Proof: By Corollary 4.9 and the Margulis-Ruelle inequality [4], $h_{\text{inv}}(K, Q) = 0$ implies the identity

$$h_{\hat{\mu}}(\varphi, (\pi_{\mathcal{U}})_*\hat{\mu}) = \int \log J^+ \varphi d\hat{\mu}$$

which is equivalent to $\hat{\mu}$ being an SRB measure for the random dynamical system $(\varphi, (\pi_{\mathcal{U}})_*\hat{\mu})$ (see [5, Thm. 2.6]).¹⁸ \square

4.8 A purely topological characterization

For certain purposes, it may be useful to have a purely topological characterization of the lower bound of Theorem 4.7. To obtain such a characterization, we first

¹⁸The obligatory integrability condition $\int (\log^+ \|\varphi_{1,u}\|_{C^2} + \log^+ \|\varphi_{-1,u}\|_{C^2}) dP(u) < \infty$ is trivially satisfied by compactness of \mathcal{U} and continuous dependence of the derivatives on u . Here we assume again without loss of generality that M is compact.

recall the definition of *topological pressure* of the bundle random dynamical system that is obtained by restricting Φ to $L(Q)$ and fixing a measure $P \in \mathcal{M}(\theta)$. Let $\alpha : L(Q) \rightarrow \mathbb{R}$ be a continuous function. For $u \in \mathcal{U}$, $\tau \in \mathbb{Z}_{>0}$ and $\varepsilon > 0$, we put

$$\pi_\alpha(u, \tau, \varepsilon) := \sup \left\{ \sum_{x \in F} 2^{\sum_{s=0}^{\tau-1} \alpha(\Phi_s(u, x))} : F \subset Q(u) \text{ is } (u, \tau, \varepsilon)\text{-separated} \right\}.$$

It can be shown that $\pi_\alpha(\cdot, \tau, \varepsilon)$ is measurable for each $\varepsilon > 0$ and $\tau \in \mathbb{Z}_{>0}$ with respect to the completed Borel σ -algebra on \mathcal{U} (see [7, Lem. 5.3]). We then put

$$\begin{aligned} \pi_{\text{top}}(\varphi^Q, P, \varepsilon; \alpha) &:= \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \int \log \pi_\alpha(u, \tau, \varepsilon) dP(u), \\ \pi_{\text{top}}(\varphi^Q, P; \alpha) &:= \lim_{\varepsilon \downarrow 0} \pi_{\text{top}}(\varphi^Q, P, \varepsilon; \alpha), \end{aligned} \quad (31)$$

where φ^Q denotes the bundle RDS given by the restriction of Φ to $L(Q)$. The variational principle (see [7, Thm. 6.1] or [46, Ch. 5, Thm. 1.2.13]) then implies

$$\pi_{\text{top}}(\varphi^Q, P; \alpha) = \sup_{\mu \in \mathcal{M}_P(\Phi; L(Q))} \pi_\mu(\varphi, P; \alpha).$$

Hence, we can write our lower bound as

$$h_{\text{inv}}(K, Q) \geq - \sup_{P \in \mathcal{M}(\theta)} \pi_{\text{top}}(\varphi^Q, P; -\log J^+ \varphi). \quad (32)$$

Now we prove the main result of this subsection, which replaces the supremum over the measures P with a supremum over control sequences.

4.11 Proposition: *It holds that*

$$\sup_{P \in \mathcal{M}(\theta)} \pi_{\text{top}}(\varphi^Q, P; -\log J^+ \varphi) = \sup_{u \in \mathcal{U}} \lim_{\varepsilon \downarrow 0} \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log \pi_{-\log J^+ \varphi}(u, \tau, \varepsilon).$$

Proof: Let us write $\alpha := -\log J^+ \varphi$. By the derivation of our lower bound, we know that

$$\limsup_{\varepsilon \downarrow 0} \liminf_{u \in \mathcal{U}} \lim_{\tau \rightarrow \infty} \frac{1}{\tau} \log \text{vol}(Q(u, \tau, \varepsilon)) \leq \sup_{P \in \mathcal{M}(\theta)} \pi_{\text{top}}(\varphi^Q, P; \alpha). \quad (33)$$

Now let $F_{u, \tau, \varepsilon} \subset Q(u)$ be an arbitrary (u, τ, ε) -separated subset. If $y \in B_\varepsilon^{u, \tau}(x)$ for some $x \in F_{u, \tau, \varepsilon}$, then $d(\varphi(t, y, u), \varphi(t, x, u)) \leq \varepsilon$ implying $\text{dist}(\varphi(t, y, u), Q(\theta^t u)) \leq \varepsilon$ for $0 \leq t < \tau$. Since the Bowen-balls $B_{\varepsilon/2}^{u, \tau}(x)$, $x \in F_{u, \tau, \varepsilon}$, are disjoint, it follows by the volume lemma that

$$\text{vol}(Q(u, \tau, \varepsilon)) \geq \sum_{x \in F_{u, \tau, \varepsilon}} \text{vol}(B_{\varepsilon/2}^{u, \tau}(x)) \geq C_{\varepsilon/2}^{-1} \sum_{x \in F_{u, \tau, \varepsilon}} J^+ \varphi_{\tau, u}(x)^{-1}$$

$$= C_{\varepsilon/2}^{-1} \sum_{x \in F_{u,\tau,\varepsilon}} 2^{\sum_{s=0}^{\tau-1} \alpha(\Phi_s(u,x))}.$$

Hence,

$$\log \text{vol}(Q(u, \tau, \varepsilon)) \geq \log C_{\varepsilon/2}^{-1} + \log \sum_{x \in F_{u,\tau,\varepsilon}} 2^{\sum_{s=0}^{\tau-1} \alpha(\Phi_s(u,x))}. \quad (34)$$

Since this holds true for every (u, τ, ε) -separated subset of $Q(u)$, we obtain

$$\liminf_{\tau \rightarrow \infty} \frac{1}{\tau} \log \text{vol}(Q(u, \tau, \varepsilon)) \geq \liminf_{\tau \rightarrow \infty} \frac{1}{\tau} \log \pi_\alpha(u, \tau, \varepsilon).$$

By [46, Ch. 5, Prop. 1.2.6], it does not matter if we replace \liminf with \limsup in the definition of topological pressure, and hence, in combination with (33) it follows that

$$\sup_{P \in \mathcal{M}(\theta)} \pi_{\text{top}}(\varphi^Q, P; \alpha) \geq \sup_{u \in \mathcal{U}} \lim_{\varepsilon \downarrow 0} \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log \pi_\alpha(u, \tau, \varepsilon).$$

Here we also use that the limit for $\varepsilon \downarrow 0$ can be written as the supremum over $\varepsilon > 0$, and two suprema can be interchanged.

To prove the converse inequality, it suffices to show that for every $P \in \mathcal{M}(\theta)$,

$$\pi_{\text{top}}(\varphi^Q, P; \alpha) \leq \sup_{u \in \mathcal{U}} \lim_{\varepsilon \downarrow 0} \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log \pi_\alpha(u, \tau, \varepsilon).$$

Using the definitions and (34), for the left-hand side we obtain

$$\begin{aligned} \pi_{\text{top}}(\varphi^Q, P; \alpha) &= \lim_{\varepsilon \downarrow 0} \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \int \log \pi_\alpha(u, \tau, \varepsilon) dP(u) \\ &\leq \lim_{\varepsilon \downarrow 0} \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \int \log \text{vol}(Q(u, \tau, \varepsilon)) dP(u) \\ &\leq \lim_{\varepsilon \downarrow 0} \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \sup_{u \in \mathcal{U}} \log \text{vol}(Q(u, \tau, \varepsilon)). \end{aligned}$$

Exactly as in the proof of Proposition 4.5, we can interchange the limit superior and the supremum, hence

$$\pi_{\text{top}}(\varphi^Q, P; \alpha) \leq \limsup_{\varepsilon \downarrow 0} \limsup_{u \in \mathcal{U}} \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log \text{vol}(Q(u, \tau, \varepsilon)).$$

As already shown in (15), for a maximal (u, τ, ε) -separated set $F_{u,\tau,\varepsilon} \subset Q(u)$ we have the inequality

$$\text{vol}(Q(u, \tau, \varepsilon)) \leq C_{\beta+\varepsilon} \sum_{x \in F_{u,\tau,\varepsilon}} 2^{\sum_{s=0}^{\tau-1} \alpha(\Phi_s(u,x))} \leq C_{\beta+\varepsilon} \pi_\alpha(u, \tau, \varepsilon),$$

implying

$$\pi_{\text{top}}(\varphi^Q, P; \alpha) \leq \limsup_{\varepsilon \downarrow 0} \limsup_{u \in \mathcal{U}} \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log \pi_\alpha(u, \tau, \varepsilon).$$

Since the limit in ε is a supremum and two suprema can be interchanged, the result is proved. \square

We close this subsection with a related result that is interesting for the evaluation of the lower bound in the case when Q is a (very) small perturbation of a hyperbolic set of a diffeomorphism.

4.12 Proposition: *Let Q be the hyperbolic set constructed in the small-perturbation setting of Theorem 3.11. Then the function*

$$u \mapsto \lim_{\varepsilon \downarrow 0} \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log \pi_{-\log J^+ \varphi}(u, \tau, \varepsilon)$$

is continuous at u^0 in the sup-metric d_∞ on \mathcal{U}_0 .

Proof: Let $u \in \mathcal{U}_0$ and let $F \subset \Lambda$ be a (u^0, τ, ε) -separated set for some $\tau \in \mathbb{Z}_{>0}$ and $\varepsilon > 0$. Consider the set $\tilde{F} := h_u(F) \subset Q(u)$. Since the family $\{h_u^{-1}\}_{u \in \mathcal{U}_0}$ is equicontinuous and $h_{\theta u} \circ f_{u^0} \equiv \varphi_{1,u} \circ h_u$ (see Proposition 3.10), we can choose $\delta = \delta(\varepsilon) > 0$ (independent of u) so that \tilde{F} is (u, τ, δ) -separated.

Moreover, since $(u, x) \mapsto \log J^+ \varphi_{1,u}(x)$ is uniformly continuous on $L(Q)$, by choosing β in Proposition 3.10(b) small enough, we obtain for all $x \in \Lambda$, $u \in \mathcal{U}_0$ sufficiently close to u^0 in the d_∞ -distance and $\tau \in \mathbb{Z}_{>0}$ that

$$\begin{aligned} \log \frac{J^+ \varphi_{\tau,u}(h_u(x))}{J^+ \varphi_{\tau,u^0}(x)} &= \sum_{s=0}^{\tau-1} (\log J^+ \varphi_{1,\theta^s u}(\varphi_{s,u}(h_u(x))) - \log J^+ \varphi_{1,u^0}(f_{u^0}^s(x))) \\ &\leq \sum_{s=0}^{\tau-1} \tilde{\beta} = \tau \tilde{\beta} \end{aligned}$$

for some $\tilde{\beta} > 0$ that becomes arbitrarily small as β and $d_\infty(u, u^0)$ do. Hence, we can estimate

$$\sum_{x \in F} J^+ \varphi_{\tau,u^0}(x)^{-1} \leq 2^{\tau \tilde{\beta}} \sum_{y \in \tilde{F}} J^+ \varphi_{\tau,u}(y)^{-1},$$

which implies

$$\pi_{-\log J^+ \varphi}(u^0, \tau, \varepsilon) \leq 2^{\tau \tilde{\beta}} \pi_{-\log J^+ \varphi}(u, \tau, \delta).$$

Since this holds for all $\tau > 0$, we have

$$\limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log \pi_{-\log J^+ \varphi}(u^0, \tau, \varepsilon) \leq \tilde{\beta} + \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log \pi_{-\log J^+ \varphi}(u, \tau, \delta).$$

In fact, $\tilde{\beta}$ was chosen independently of ε so that this inequality still holds if we send ε and δ to zero. Interchanging the roles of u and u^0 , we see that also the converse inequality holds. This completes the proof. \square

As a consequence of the above proposition, the lower bound obtained for $h_{\text{inv}}(K, Q)$ converges to the topological pressure on Λ (with respect to $-\log J^+ f_{u^0}$) as the size of the neighborhood U_0 shrinks to zero.

4.9 Achievability

There are good reasons to expect that our lower bound for invariance entropy also becomes an upper bound under additional controllability assumptions, i.e., that average data rates arbitrarily close to the lower bound are achievable by proper coder-controller designs. In the following two extreme cases, this can be made very plausible:

- Assume that the u -fibers of Q are finite. As the main result of [41] shows, this is always the case for hyperbolic sets of continuous-time systems.¹⁹ In this framework, we have derived a formula for $h_{\text{inv}}(K, Q)$ in [25] which is analogous to our lower bound. To obtain this result, we needed to assume that the hyperbolic set is the closure of a maximal set of approximate controllability and that the Lie algebra rank condition (guaranteeing local accessibility) is satisfied on Q . Observe that in the case of finite u -fibers, the entropy term $h_\mu(\varphi, (\pi_U)_*\mu)$ in our lower bound vanishes, because finite fibers cannot support positive entropy. Hence, in this case

$$h_{\text{inv}}(K, Q) \geq \inf_{\mu} \int \log J^+ \varphi \, d\mu.$$

The theory of subadditive cocycles (see, e.g., [52, App. A]) shows that this is equivalent to

$$h_{\text{inv}}(K, Q) \geq \inf_{(u,x) \in L(Q)} \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log J^+ \varphi_{\tau,u}(x).$$

In the continuous-time case, an analogous upper bound is obtained by stabilizing the system around regular periodic trajectories in $\text{int } Q$. Via arguments originating from [53], this leads to upper bounds which approximate all growth rates of the form

$$\limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log J^+ \varphi_{\tau,u}(x), \quad (u, x) \in L(Q).$$

It is more or less obvious that the same proof techniques also work in discrete time. However, since the genericity of universally regular control sequences is needed to carry out some details of the proof, similar assumptions

¹⁹This may seem strange, but follows from the definition of uniform hyperbolicity without a one-dimensional center bundle.

as in Theorem 3.13 (analyticity and uniform forward accessibility, in particular) are necessary. The special case when Q is constructed as in the small-perturbation setting of Theorem 3.11 is handled by Theorem 4.14 below.

- The opposite extreme case is that the set $L(Q)$ supports an SRB measure for one of the random dynamical systems (φ, P) . In this case, as we have seen in Corollary 4.10, the lower bound vanishes. On the other hand, the existence of an SRB measure should imply the existence of some sort of attractor in Q . But if such an attractor exists, then appropriate controllability assumptions will guarantee that one can steer the system from any initial state in K into the basin of attraction, by using only finitely many different control sequences. Once the system has entered the basin of attraction, no further control actions are necessary, which leads to $h_{\text{inv}}(K, Q) = 0$. In the small-perturbation setting, this is shown by Theorem 4.17 below.

For the general case, a concrete idea how to prove an achievability result is missing although it is clear that one has to consider coding and control strategies that stabilize the system at the u -fibers (possibly periodic u 's will do as in the case of finite fibers). However, stabilization around particular trajectories would lead to data rates that are too large to match the lower bound. Hence, appropriate coding and control strategies should keep the state x_t close to $Q(\theta^t u)$ without following the same trajectory for every initial state (due to shadowing it cannot be avoided to follow *some* trajectory, however).

For completeness, we provide proofs for the two simplest cases of the achievability result. First, we handle the case when Q is constructed by a small control-perturbation of a hyperbolic periodic orbit.

The following lemma, taken from [25, Prop. 9], will be used.

4.13 Lemma: *Consider the dynamical system $(U^{\mathbb{Z}}, \theta)$. For every $\varepsilon > 0$, there exists $\delta > 0$ such that every δ -chain of $(U^{\mathbb{Z}}, \theta)$ is ε -shadowed by a real orbit. Moreover, if the δ -chain is periodic, a periodic shadowing orbit with the same period exists.*

4.14 Theorem: *Consider the control system Σ and let $Q \subset M$ be a hyperbolic set constructed as in the small-perturbation setting of Theorem 3.11 for the control system Σ^0 given by (12). Additionally, assume that the following conditions are satisfied:*

- (B1) *The topologically transitive and hyperbolic set Λ of f_{u^0} is a periodic orbit.*
- (B2) *The system Σ^0 is real-analytic and uniformly forward accessible.*
- (B3) *$L(Q)$ is a chain component of the control flow of Σ^0 .*²⁰

²⁰Recall that by Lemma 3.12 $L(Q)$ is an internally chain transitive set. Here we are only adding

Then for any compact set $K \subset \text{core}(Q)$ of positive volume, it holds that

$$h_{\text{inv}}(K, Q) = \inf_{(u,x) \in L(Q)} \limsup_{t \rightarrow \infty} \frac{1}{t} \log J^+ \varphi_{t,u}(x). \quad (35)$$

Proof: Throughout the proof, we denote the right-hand side of (35) by γ . We also recall that

$$\alpha_t(u, x) := \log J^+ \varphi_{t,u}(x), \quad \alpha_t : L(Q) \rightarrow \mathbb{R}$$

is a continuous additive cocycle over $\Phi|_{L(Q)}$. For notational reasons, we write u^* instead of u^0 for the fixed constant control that leads to the hyperbolic periodic orbit Λ .

Step 1: We prove that the right-hand side of the inequality (25) (our lower bound) equals γ under Assumption (B1). In fact, for this conclusion we only need that the u -fibers $Q(u)$ are finite. First, we prove that the measure-theoretic entropy $h_\mu(\varphi, P)$ vanishes for every $P \in \mathcal{M}(\theta)$ and every invariant measure μ of the RDS (φ, P) with $\text{supp}(\mu) \subset L(Q)$. The variational principle for bundle RDS (see [46, Thm. 1.2.13]) implies the inequality $h_\mu(\varphi, P) \leq h_{\text{top}}(\varphi, P)$, where the right-hand side is the topological entropy of the bundle RDS. Since $h_{\text{top}}(\varphi, P)$ is defined via the growth rates of maximal (u, τ, ε) -separated subsets of the u -fibers and these are finite, it vanishes. Hence, to complete the first step it remains to show that

$$\inf_{\mu \in \mathcal{M}(\Phi|_{L(Q)})} \int \log J^+ \varphi \, d\mu = \gamma. \quad (36)$$

This follows immediately from the theory of subadditive cocycles, applied to α , see [52, App. A]. As a consequence, Theorem 4.7 immediately yields

$$h_{\text{inv}}(K, Q) \geq \gamma. \quad (37)$$

Step 2: We introduce the set

$$L_{\text{per}}(Q) := \{(u, x) \in L(Q) : \Phi_\tau(u, x) = (u, x) \text{ for some } \tau \in \mathbb{Z}_{>0}\}$$

of periodic elements of $L(Q)$ and prove that

$$\gamma = \inf_{(u,x) \in L_{\text{per}}(Q)} \lim_{t \rightarrow \infty} \frac{1}{t} \log J^+ \varphi_{t,u}(x). \quad (38)$$

The proof of this identity uses the concept of the *Morse spectrum* of an additive cocycle. Let us therefore first recall some definitions. Consider an ε -chain ζ given by points $(u^0, x_0), \dots, (u^\tau, x_\tau)$ in $L(Q)$.²¹ The *finite-time Morse exponent* of the chain ζ is defined as

$$\lambda(\zeta) := \frac{1}{\tau} \sum_{t=0}^{\tau-1} \alpha_1(u^t, x_t).$$

the requirement that $L(Q)$ is maximal with this property. This is probably satisfied if Λ is an Axiom A basic set.

²¹Recall that we use upper indexes for the u -components to avoid abuse of notation.

The Morse spectrum of the cocycle α is the set

$$S_{\text{Mo}}(\alpha) := \bigcap_{\varepsilon > 0} \text{cl} \{ \lambda(\zeta) : \zeta \text{ is an } \varepsilon\text{-chain in } L(Q) \}.$$

From Assumption (B3) and [58, Thm. 3.2], we know that $S_{\text{Mo}}(\alpha)$ is a compact interval which equals

$$S_{\text{Mo}}(\alpha) = \left\{ \int \alpha_1 d\mu : \mu \in \mathcal{M}(\Phi|_{L(Q)}) \right\}.$$

In particular, by (36) this shows that

$$\gamma = \inf S_{\text{Mo}}(\alpha). \quad (39)$$

By [45, Lem. 8], it suffices to consider periodic ε -chains in the definition of the Morse spectrum, i.e., such with $(u^0, x_0) = (u^\tau, x_\tau)$. Let

$$S_{\text{Mo,Per}}(\alpha) := \bigcap_{\varepsilon > 0} \text{cl} \{ \lambda(\zeta) : \zeta \text{ is a periodic } \varepsilon\text{-chain in } L(Q) \}.$$

Then (39) together with [45, Lem. 8] yields

$$\gamma = \inf S_{\text{Mo,Per}}(\alpha).$$

Now consider a periodic ε -chain ζ in $L(Q)$, given by $(u^0, x_0), (u^1, x_1), \dots, (u^\tau, x_\tau) = (u^0, x_0)$. We want to prove the existence of a τ -periodic point $(u, x) \in L(Q)$ which shadows the ε -chain ζ in the sense that

$$d_{\mathcal{U} \times M}(\Phi_t(u, x), (u^t, x_t)) \leq \beta, \quad t = 0, 1, \dots, \tau - 1, \quad (40)$$

where $\beta > 0$ is given and $\varepsilon = \varepsilon(\beta)$ must be chosen sufficiently small. If we have found such a point (u, x) , it follows that

$$\begin{aligned} \left| \lambda(\zeta) - \lim_{t \rightarrow \infty} \frac{1}{t} \alpha_t(u, x) \right| &= \left| \frac{1}{\tau} \sum_{t=0}^{\tau-1} \alpha_1(u^t, x_t) - \frac{1}{\tau} \alpha_\tau(u, x) \right| \\ &= \frac{1}{\tau} \left| \sum_{t=0}^{\tau-1} (\alpha_1(u^t, x_t) - \alpha_1(\Phi_t(u, x))) \right| \\ &\leq \max_{0 \leq t < \tau} |\alpha_1(u^t, x_t) - \alpha_1(\Phi_t(u, x))|, \end{aligned} \quad (41)$$

where we use that

$$\lim_{t \rightarrow \infty} \frac{1}{t} \alpha_t(u, x) = \frac{1}{\tau} \alpha_\tau(u, x)$$

by the τ -periodicity of (u, x) . It is clear that the last expression in (41) can be made arbitrarily small if β is chosen small enough due to (40) and the uniform continuity of α_1 on the compact set $L(Q)$. Hence, we obtain (38) as desired. To

find the periodic point (u, x) , we proceed in two steps. First, we use the periodic shadowing property of the shift operator on \mathcal{U}_0 described in Lemma 4.13. This property yields a τ -periodic control sequence $u \in \mathcal{U}_0$ such that

$$d_{\mathcal{U}}(\theta^t u, u^t) \leq \delta \quad \text{for } t = 0, 1, \dots, \tau - 1 \quad (42)$$

for any fixed $\delta > 0$ if $\varepsilon = \varepsilon(\delta)$ is chosen small enough. Now observe that the bi-infinite sequence $(\theta^t u, x_t)_{t \in \mathbb{Z}}$, where the finite sequence of x_t 's is continued τ -periodically in both directions, is a pseudo-orbit, since

$$\begin{aligned} d(\varphi(1, x_t, \theta^t u), x_{t+1}) &\leq d(\varphi(1, x_t, \theta^t u), \varphi(1, x_t, u^t)) + d(\varphi(1, x_t, u^t), x_{t+1}) \\ &\leq d(\varphi(1, x_t, \theta^t u), \varphi(1, x_t, u^t)) + d_{\mathcal{U} \times M}(\Phi(u^t, x_t), (u^{t+1}, x_{t+1})). \end{aligned}$$

The first term can be made arbitrarily small by uniform continuity of $\varphi(1, \cdot, \cdot)$ on the compact set $Q \times U_0$ and (42). The second term is smaller than ε by assumption. Moreover, the pseudo-orbit $(\theta^t u, x_t)$ is close to $L(Q)$, because $\theta^t u$ is close to u^t and $(u^t, x_t) \in L(Q)$. Hence, the shadowing lemma yields a true orbit of the form $(\theta^t u, \varphi(t, y, u))_{t \in \mathbb{Z}}$ with $(u, y) \in L(Q)$ which shadows $(\theta^t u, x_t)$, and thus (u^t, x_t) . By uniqueness of shadowing orbits and a shifting argument, it easily follows that this orbit is τ -periodic. The proof of Step 2 is complete.

Step 3: We introduce the set

$$L_{\text{per,reg}}(Q) := \{(u, x) \in L_{\text{per}}(Q) : u_t \in \text{int } U_0 \ \forall t \in \mathbb{Z}, x \in \text{int } Q, (x, u) \text{ is regular}\},$$

where regularity is understood as controllability of the linearization on the time interval $[0; \tau]$ with $\tau > 0$ denoting the minimal period of (u, x) . We prove that

$$\gamma = \inf_{(u, x) \in L_{\text{per,reg}}(Q)} \lim_{t \rightarrow \infty} \frac{1}{t} \log J^+ \varphi_{t, u}(x).$$

To prove this identity, we exploit the genericity of universally regular control sequences as guaranteed by Theorem 3.13 and Assumption (B2). Pick an arbitrary point $(u, x) \in L_{\text{per}}(Q)$ of period τ . We claim that the point (u, x) can be approximated by a sequence $(u^n, x_n) \in L_{\text{per,reg}}(Q)$. We may assume that τ is large enough so that $S(\tau)$, the set of universally regular control sequences in $(\text{int } U_0)^\tau$, is dense in U_0^τ . Hence, we find a sequence $(u^n)_{n \in \mathbb{Z}_{>0}}$ in $S(\tau)$ such that $u^n \rightarrow u$. Here we also think of u^n being extended τ -periodically in both directions so that $u^n \in \mathcal{U}_0$. Using the homeomorphisms h_u introduced in Proposition 3.10, let $x = h_u(z)$ for some z on the periodic orbit Λ . Let $|\Lambda| = \tau^*$ so that τ^* is the minimal period of z . Then

$$h_u(z) = x = \varphi_{\tau, u}(x) = \varphi_{\tau, u}(h_u(z)) = h_{\theta^\tau u}(f_{u^*}^\tau(z)) = h_u(f_{u^*}^\tau(z)).$$

Since h_u is injective, it follows that $z = f_{u^*}^\tau(z)$. Hence, τ must be an integer multiple of τ^* . Now put $x_n := h_{u^n}(z) \in Q(u^n)$. Then

$$\varphi_{\tau, u^n}(x_n) = \varphi_{\tau, u^n}(h_{u^n}(z)) = h_{\theta^\tau u^n}(f_{u^n^*}^\tau(z)) = h_{u^n}(z) = x_n.$$

As $u^n \rightarrow u$ and $v \mapsto h_v$ is continuous, we have $x_n = h_{u^n}(z) \rightarrow h_u(z) = x$. It remains to prove that $x_n \in \text{int } Q$ for each n . Fix n and note that by universal regularity of u^n the linearization along the controlled periodic orbit associated with (u^n, x_n) is controllable on the time interval $[0; \tau]$. Assume to the contrary that $x_n \in \partial Q$. Then local controllability (implied by the controllability of the linearization) leads to periodic trajectories starting in Q , leaving Q and then returning to Q , which stay arbitrarily close to the trajectory $\varphi(\cdot, x^n, u^n)$ at all times. For instance, one can first steer in time τ from x^n to a point $y \notin Q$ close to x^n . Then one steers from y back to x^n in time τ , which leads to a 2τ -periodic trajectory starting in x^n which is not completely contained in Q . Moreover, this trajectory is controlled by a control sequence arbitrarily close to u^n . This contradicts the fact that $L(Q)$ is isolated invariant. Hence, the claim is proved. As $(u^n, x_n) \rightarrow (u, x)$, we also have $\frac{1}{\tau}\alpha_\tau(u^n, x_n) \rightarrow \frac{1}{\tau}\alpha_\tau(u, x)$, which completes the proof of Step 3.

Step 4: To complete the proof of the theorem, it suffices to show that

$$h_{\text{inv}}(K, Q) \leq \lim_{t \rightarrow \infty} \frac{1}{t} \alpha_t(u, x) \quad \text{for all } (u, x) \in L_{\text{per,reg}}(Q).$$

This can be shown by the arguments in [53, Thm. 3] adapted to the case of a periodic orbit (instead of an equilibrium point). Also note that for the continuous-time case these arguments have already been adapted in [42, Thm. 4.3]. Here, it is important that for a fixed $(u, x) \in L_{\text{per,reg}}(Q)$, one can steer from any initial state $x_0 \in K$ to an arbitrarily small neighborhood of x in finite time without leaving Q . To prove this, we use Assumption (B3) again. By Corollary 3.19, we know that we can steer from any $x_0 \in K$ arbitrarily close to x via some trajectory $\varphi(\cdot, x_0, u)$, $u \in \mathcal{U}_0$. For some $\tau > 0$, we have $\varphi(\tau, x_0, u) \in Q$. Assume to the contrary that $\varphi(t, x_0, u) \notin Q$ for some $0 < t < \tau$. We prove that this contradicts that $L(Q)$ is a maximal chain transitive set. It is well-known that Q , as the projection of $L(Q)$ to M , is a maximal set of all-time controlled invariance and chain controllability (see [17, Thm. 4.1.4] for the continuous-time case). However, the set $Q \cup \{\varphi(s, x_0, u) : s \in [0; \tau]\}$ also has these two properties as one can easily check. This contradicts maximality, and hence concludes the proof. \square

In the following, we show how to handle the case when Q is constructed as in the small-perturbation setting and one of the RDS (φ, P) admits an SRB measure on $L(Q)$. To prove the corresponding result, we need some additional concepts and results from the hyperbolic theory.

Consider the control system Σ . For any $(u, x) \in \mathcal{U} \times M$, the *local unstable manifold of size $\varepsilon > 0$* is given by

$$W_{u,\varepsilon}^+(x) = \{y \in M : d(\varphi(-t, x, u), \varphi(-t, y, u)) \leq \varepsilon, \forall t \geq 0\}.$$

If Q is a hyperbolic set of Σ , the *stable manifold theorem* tells us that for all $(u, x) \in L(Q)$, $W_{u,\varepsilon}^+(x)$ is an embedded submanifold of M with

$$T_x W_{u,\varepsilon}^+(x) = E^+(u, x).$$

In particular, all unstable manifolds have the same dimension d^+ .

We will further use the following notation:

$$B(u, \tau, \delta) := \bigcup_{z \in Q(u)} B_\delta^{u, \tau}(z).$$

That is, $B(u, \tau, \delta)$ is the union of all Bowen-balls centered in $Q(u)$ of order τ and radius δ .

By [48, Eq. (3.1)], we have the following lemma on the topological pressure of associated random dynamical systems, which is actually a simple consequence of the volume lemma in combination with the shadowing lemma.

4.15 Lemma: *Let $P \in \mathcal{M}(\theta)$. Then, for all sufficiently small $\delta > 0$, we have*

$$\pi_{\text{top}}(\varphi^Q, P; -\log J^+ \varphi) = \lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int \log \text{vol}(B(u, \tau, \delta)) \, dP(u).$$

We also need the so-called *second volume lemma* [48, Lem. A.1], which reads as follows.

4.16 Lemma: *Let Σ be of regularity class C^2 and assume that Q is a hyperbolic set of Σ . Then, for $\varepsilon, \delta > 0$ small enough, there is a constant $C_{\varepsilon, \delta} > 0$ such that*

$$C_{\varepsilon, \delta}^{-1} \leq \frac{\text{vol}(B_\delta^{u, \tau}(y))}{\text{vol}(B_\varepsilon^{u, \tau}(x))} \leq C_{\varepsilon, \delta}$$

whenever $(u, x) \in L(Q)$, $\tau \geq 0$ and $y \in B_\varepsilon^{u, \tau}(x)$.

Now, we can formulate and prove our main result.

4.17 Theorem: *Consider the control system Σ and assume that it is of regularity class C^2 . Let $Q \subset M$ be a hyperbolic set constructed as in the small-perturbation setting of Theorem 3.11 for the control system Σ^0 given by (12). Additionally, assume that the following conditions are satisfied:*

- (C1) *The isolated invariant and hyperbolic set Λ of f_{u^0} is topologically transitive.*
- (C2) *For some $P \in \mathcal{M}(\theta)$, the RDS (φ, P) admits an invariant probability measure μ , supported on $L(Q)$, which satisfies*

$$h_\mu(\varphi, P) = \int \log J^+ \varphi \, d\mu. \quad (43)$$

- (C3) *Λ is contained in $\text{int } Q$.²²*

²²A sufficient condition is given in Theorem 5.1.

(C4) $L(Q)$ is a chain component of the control flow of Σ^0 .

(C5) $U \subset \mathbb{R}^m$ and f is of class C^1 .

Then, for any compact set $K \subset \text{core}(Q)$, we have

$$h_{\text{inv}}(K, Q) = 0.$$

Proof: The proof proceeds in five steps. The first four of them will show that Λ is an attractor. The last step then uses this fact to show that the invariance entropy vanishes.

Step 1: We prove the following auxiliary statement:

If $W_{u,\varepsilon}^+(x) \subset Q(u)$ for some $(u, x) \in L(Q)$, then Λ is an attractor, i.e., there exists an arbitrarily small neighborhood V of Λ with $f_{u^0}(V) \subset V$.

To this end, consider the set

$$A := h_u^{-1}(W_{u,\varepsilon}^+(x)) = \{h_u^{-1}(y) : d(\varphi(-t, x, u), \varphi(-t, y, u)) \leq \varepsilon, \forall t \geq 0\}$$

which is a well-defined subset of Λ by assumption. Using the uniform equicontinuity of the family $\{h_u^{-1}\}$ (see Proposition 3.10), we can see that $A \subset W_{u^0,\delta}^+(h_u^{-1}(x))$ if $\varepsilon = \varepsilon(\delta) > 0$ is small enough (taking a smaller ε does not hurt). Indeed, this follows from

$$d(f_{u^0}^{-t}(h_u^{-1}(y)), f_{u^0}^{-t}(h_u^{-1}(x))) = d(h_{\theta^{-t}u}^{-1}(\varphi(-t, y, u)), h_{\theta^{-t}u}^{-1}(\varphi(-t, x, u))).$$

Since h_u^{-1} is a homeomorphism, A is a topological submanifold of $W_{u^0,\delta}^+(h_u^{-1}(x))$ of dimension d^+ . By the invariance-of-domain theorem, then A must be open in $W_{u^0,\delta}^+(h_u^{-1}(x))$.²³ Consequently, we can find some $\eta > 0$ small enough such that $W_{u^0,\eta}^+(h_u^{-1}(x)) \subset A \subset \Lambda$. Now, we invoke [9, Lem. 4.9], which shows that this implies that Λ is an attractor (under Assumption (C1)).

Step 2: We prove another auxiliary result:

²³We need to show that every $z \in A$ has a neighborhood in A which is open in $W_{u^0,\delta}^+(x)$. To this end, let $V \subset A$ be a neighborhood of z which is homeomorphic to \mathbb{R}^{d^+} via a homeomorphism $\phi : V \rightarrow \mathbb{R}^{d^+}$. But z also has a neighborhood $\tilde{V} \subset W_{u^0,\delta}^+(x)$, open relative to $W_{u^0,\delta}^+(x)$, which is Euclidean. Let $\tilde{\phi} : \tilde{V} \rightarrow \mathbb{R}^{d^+}$ be the associated homeomorphism. Without loss of generality, we can assume that $V \subset \tilde{V}$. Then we consider the map $\psi := \tilde{\phi}|_V \circ \phi^{-1} : \mathbb{R}^{d^+} \rightarrow \mathbb{R}^{d^+}$. This map is continuous and injective. Now the invariance-of-domain-theorem tells us that the image $U := \psi(\mathbb{R}^{d^+})$ is open in \mathbb{R}^{d^+} . Then we know that $\tilde{\phi}^{-1}(U)$ is open in \tilde{V} , implying that $\tilde{\phi}^{-1}(U)$ is open in $W_{u^0,\delta}^+(x)$. At the same time, $\tilde{\phi}^{-1}(U) = \phi^{-1}(\mathbb{R}^{d^+}) = V$. Hence, V is the desired neighborhood.

If Λ is not an attractor, then there exists a constant $\gamma > 0$ such that for every $(u, x) \in L(Q)$ there is $y \in W_{u,\varepsilon}^+(x)$ with $\text{dist}(y, Q(u)) \geq \gamma$.

To this end, fix $u \in \mathcal{U}$ and consider for every $\beta > 0$ the set

$$V_\beta(u) := \{x \in Q(u) : \text{dist}(y, Q(u)) > \beta \text{ for some } y \in W_{u,\varepsilon}^+(x)\}.$$

This set is open in $Q(u)$, because $W_{u,\varepsilon}^+(x)$ depends continuously on x by the stable manifold theorem. When β decreases, $V_\beta(u)$ increases. By Step 1, we have $W_{u,\varepsilon}^+(x) \not\subset Q(u)$ for all $x \in Q(u)$. Hence, for every $x \in Q(u)$ there is $\beta(x) > 0$ with $x \in V_{\beta(x)}(u)$, and thus $Q(u) = \bigcup_{\beta>0} V_\beta(u)$. By compactness, $V_\beta(u) = Q(u)$ for some $\beta > 0$. We choose $\beta(u)$ as the supremum over all such β .

Now assume to the contrary that there is a sequence $(u^n)_{n \in \mathbb{N}}$ in \mathcal{U} such that $\beta(u^n) \rightarrow 0$ as $n \rightarrow \infty$. By compactness of \mathcal{U} , we may assume that $u^n \rightarrow u^*$ for some $u^* \in \mathcal{U}$. Then there are $x_n \in Q(u^n)$ with $W_{u^n,\varepsilon}^+(x_n) \subset N_{(1/n)+\beta(u^n)}(Q(u^n))$. We can also assume that x_n converges to some $x_* \in Q(u^*)$. As $u \mapsto Q(u)$ and $(u, x) \mapsto W_{u,\varepsilon}^+(x)$ are continuous (the latter holds by the stable manifold theorem, see [51]) and $\beta(u^n) \rightarrow 0$, it follows that $W_{u^*,\varepsilon}^+(x_*) \subset Q(u^*)$, a contradiction. Hence, we can put $\gamma := \inf_{u \in \mathcal{U}} \beta(u)$.

Step 3: We prove that Assumption (C2) implies

$$\pi_{\text{top}}(\varphi^Q, P; -\log J^+ \varphi) = 0. \quad (44)$$

The variational principle for the pressure of bundle RDS [46] tells us that

$$\pi_{\text{top}}(\varphi^Q, P; -\log J^+ \varphi) = \sup_{\nu} \left[h_{\nu}(\varphi, P) - \int \log J^+ \varphi \, d\nu \right],$$

where the supremum is taken over all invariant probability measures ν of the RDS (φ, P) which are supported on $L(Q)$. The Margulis-Ruelle inequality [4] says that

$$h_{\nu}(\varphi, P) \leq \int \log J^+ \varphi \, d\nu$$

for every ν , and hence $\pi_{\text{top}}(\varphi^Q, P; -\log J^+ \varphi) \leq 0$. Thus, from (43) the equality (44) immediately follows.

Step 4: We now prove by contradiction that Λ is an attractor. We thus assume that Λ is not an attractor and show that this leads to $\pi_{\text{top}}(\varphi^Q, P; -\log J^+ \varphi) < 0$ in contradiction to (44).

Given a small $\varepsilon > 0$, choose $\gamma > 0$ as in Step 2. Pick $T \in \mathbb{N}$ such that

$$\varphi_{T,u}(W_{u,\gamma/4}^+(x)) \supset W_{\theta^T u, \varepsilon}^+(\varphi_{T,u}(x)) \quad \text{for all } (u, x) \in L(Q). \quad (45)$$

This is possible due to uniform contraction rates on unstable manifolds. Let $E \subset Q(u)$ be (u, τ, γ) -separated for some $u \in \mathcal{U}$. For $x \in E$, there is $y(x, \tau) \in B_{u, \gamma/4}^\tau(x)$ with

$$\text{dist}(\varphi_{\tau+T, u}(y(x, \tau)), Q(\theta^{\tau+T}u)) > \gamma,$$

since $\varphi_{\tau, u}(B_{\gamma/4}^{u, \tau}(x)) \supset W_{\theta^{\tau}u, \gamma/4}^+(\varphi_{\tau, u}(x))$ (easy to see) and (by (45))

$$\varphi_{T, \theta^{\tau}u}W_{\theta^{\tau}u, \gamma/4}^+(\varphi_{\tau, u}(x)) \supset W_{\Theta^{T+\tau}u, \varepsilon}^+(\varphi_{T+\tau, u}(x)).$$

Choose $\delta \in (0, \gamma/4)$ such that $d(\varphi_{T, u}(y), \varphi_{T, u}(z)) < \gamma/2$ whenever $u \in \mathcal{U}$ and $d(y, z) < \delta$. Then

$$\begin{aligned} B_\delta^{u, \tau}(y(x, \tau)) &\subset B_{\gamma/2}^{u, \tau}(x), \\ \varphi_{\tau+T, u}(B_\delta^{u, \tau}(y(x, \tau))) &\cap N_{\gamma/2}(Q(\theta^{\tau+T}u)) = \emptyset. \end{aligned}$$

Hence,

$$B_\delta^{u, \tau}(y(x, \tau)) \cap B(u, \tau + T, \gamma/2) = \emptyset.$$

Using Lemma 4.16, this leads to

$$\begin{aligned} \text{vol}(B(u, \tau, \gamma/2)) - \text{vol}(B(u, \tau + T, \gamma/2)) &\geq \sum_{x \in E} \text{vol}(B_\delta^{u, \tau}(y(x, \tau))) \\ &\geq C_{3\gamma/2, \delta} \sum_{x \in E} \text{vol}(B_{3\gamma/2}^{u, \tau}(x)) \geq C_{3\gamma/2, \delta} \text{vol}(B(u, \tau, \gamma/2)). \end{aligned}$$

Therefore, setting $C := C_{3\gamma/2, \delta}$, we obtain

$$\text{vol}(B(u, \tau + T, \gamma/2)) \leq (1 - C) \cdot \text{vol}(B(u, \tau, \gamma/2)),$$

where we observe that $C \in (0, 1)$ by our choice of δ . By Lemma 4.15, this implies

$$\begin{aligned} \pi_{\text{top}}(\varphi^Q, P; -\log J^+ \varphi) &= \lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int \log \text{vol}(B(u, \tau, \gamma/2)) \, dP(u) \\ &\leq \frac{1}{T} \log(1 - C) < 0 \end{aligned}$$

in contradiction to (44). We have thus proven that Λ is an attractor under (C1) and (C2).

Step 5: We prove that $h_{\text{inv}}(K, Q) = 0$ for every $K \subset \text{core}(Q)$. We know that Λ is an attractor and by Assumption (C3) we have $\Lambda \subset \text{int } Q$. Hence, there exists an open neighborhood V of Λ with $f_{u^0}(V) \subset V \subset Q$. By complete controllability on $\text{core}(Q)$ (guaranteed by Proposition 3.15) and since $\text{core}(Q)$ is dense in Q (guaranteed by Assumption (C5) and Lemma 3.16), for every $x \in K$ we find $u^x \in \mathcal{U}_0$ and $\tau_x \in \mathbb{Z}_+$ so that $\varphi(\tau_x, x, u^x) \in V$. By continuity, we can choose a neighborhood V_x of x so that $\varphi(\tau_x, V_x \cap K, u^x) \subset V$. Moreover, by Assumption (C4), the involved trajectories do not leave Q (using the same arguments as in the

proof of Theorem 4.14). Since K is compact, we can choose a finite subcover of the cover $\{V_x\}_{x \in K}$, say $\{V_{x_1}, \dots, V_{x_r}\}$. Let $\tau \geq \max\{\tau_{x_i} : i = 1, \dots, r\}$. Then the set $\mathcal{S} \subset \mathcal{U}_0$ consisting of the control sequences

$$u_t^i := \begin{cases} u_t^{x_i} & \text{for } 0 \leq t \leq \tau_{x_i}, \\ u^0 & \text{for } \tau_{x_i} < t \leq \tau \end{cases}, \quad i = 1, \dots, r$$

is a (τ, K, Q) -spanning set by construction. Hence, $r_{\text{inv}}(\tau, K, Q) \leq r$ for all τ large enough, implying $h_{\text{inv}}(K, Q) = 0$. \square

5 Stabilization to a hyperbolic set

Consider again the control system Σ and assume that $U \subset \mathbb{R}^m$ with $U = \text{cl int } U$. Further assume that the right-hand side $f : M \times U \rightarrow M$ is continuously differentiable.

We fix a control value $u^0 \in \text{int } U$. As in Subsection 3.3, we assume that the diffeomorphism $f_0 := f_{u^0} : M \rightarrow M$ has an isolated invariant hyperbolic set Λ . Instead of “blowing up” this set to a hyperbolic set Q of Σ and asking for invariance of Q , we now consider the related control objective of locally stabilizing Σ to Λ .

Given a discrete noiseless channel, we say that Σ is *locally uniformly stabilizable* to Λ if for every $\varepsilon > 0$ there is a $\delta > 0$ and a coder-controller operating over the given channel and achieving that

$$\sup_{t \geq 0, x_0 \in N_\delta(\Lambda)} \text{dist}(x_t, \Lambda) \leq \varepsilon \quad \text{and} \quad \sup_{x_0 \in N_\delta(\Lambda)} d_\infty(u_t, u^0) \leq \varepsilon.$$

That is, whenever the initial state x_0 is close enough to Λ , the controller keeps x_t within a distance of ε to Λ for all times via a control sequence that is ε -close to u^0 at all times.

We borrow here the channel model considered in Nair et al. [53] of a *discrete noiseless channel* with a time-varying coding alphabet \mathcal{A}_t – at each time instant $t \in \mathbb{Z}_+$, one symbol from \mathcal{A}_t is transmitted without error or delay. The associated *average data rate* is

$$R_{\text{av}} := \liminf_{\tau \rightarrow \infty} \frac{1}{\tau} \sum_{t=0}^{\tau-1} \log |\mathcal{A}_t|.$$

Both coder and decoder/controller may use past knowledge, but a detailed description of these components is not necessary for the proof of the following theorem. The only thing important is that at time t the controller cannot use any other information than what has been sent through the channel until time t .

5.1 Theorem: *Let the following assumptions be satisfied for the system Σ :*

- (i) *There is a $\tau \in \mathbb{Z}_{>0}$ so that for every $x \in \Lambda$ the pair (x, u_0^τ) is regular, where $u_0^\tau = (u_0, u_0, \dots, u_0) \in U^\tau$.*

(ii) Σ is locally uniformly stabilizable to Λ over a discrete noiseless channel.

Then the channel must support an average data rate satisfying

$$R_{\text{av}} \geq -P_{\text{top}}((f_0)|_{\Lambda}, -\log J^+ f_0).$$

Proof: We restrict the control range to the closed ε -ball around u^0 in U , where $\varepsilon > 0$ is small enough so that a hyperbolic set Q^ε as in Theorem 3.11 can be constructed for the associated control system Σ^ε . In particular, we know that Λ is the u^0 -fiber of Q^ε . We prove that assumption (i) implies $\Lambda \subset \text{int } Q^\varepsilon$. To this end, pick an arbitrary $x \in \Lambda$. From the assumption, it follows that there exist $\tau > 0$ and a neighborhood $B_{\delta_x}(x)$ so that every $y \in B_{\delta_x}(x)$ can be steered to $f_0^\tau(x)$ in τ steps of time via a controlled trajectory that is never further away from $L(Q^\varepsilon)$ than a given $\rho > 0$. On the other hand, we can choose δ_x small enough so that every $y \in B_{\delta_x}(x)$ can be reached from $x' := f_0^{-\tau}(x)$ via a controlled trajectory with the same property. Hence, for every $y_0 \in B_{\delta_x}(x)$ we can construct a controlled trajectory $(u_t, y_t)_{t \in \mathbb{Z}}$ so that $d_U(u_t, u^0) \leq \rho$ and $d(y_t, f_0^t(x)) \leq \rho$ for all $t \in \mathbb{Z}$. Here the assumption that $u^0 \in \text{int } U$ guarantees that $u_t \in \text{cl } B_\varepsilon(u^0) \cap U$ for $\rho \leq \varepsilon$. Since $L(Q^\varepsilon)$ is isolated invariant, this implies $y_0 \in Q^\varepsilon$. Then $\bigcup_{x \in \Lambda} B_{\delta_x}(x)$ is an open neighborhood of Λ contained in Q^ε .

Since $\Lambda \subset \text{int } Q^\varepsilon$, we can choose $\varepsilon' > 0$ such that $N_{\varepsilon'}(\Lambda) \subset Q^\varepsilon$. If a coder-controller achieves that

$$\sup_{t \geq 0, x_0 \in N_\delta(\Lambda)} \text{dist}(x_t, \Lambda) \leq \varepsilon'$$

for some $\delta \in (0, \varepsilon')$ with controls taking values in $\text{cl } B_\varepsilon(u^0)$, then the set of different control sequences \mathcal{S}_τ generated by the controller in the time interval $[0; \tau)$ is a $(\tau, N_\delta(\Lambda), Q^\varepsilon)$ -spanning set for Σ^ε . Since the number of control sequences the controller can generate is bounded by the amount of information it receives through the channel, the cardinality of \mathcal{S}_τ satisfies

$$|\mathcal{S}_\tau| \leq \prod_{t=0}^{\tau-1} |\mathcal{A}_t|$$

so that the analysis of Section 4 shows that

$$R_{\text{av}} \geq - \sup_{u \in (\text{cl } B_\varepsilon(u^0))^\mathbb{Z}} \lim_{\rho \downarrow 0} \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log \pi_{-\log J^+ \varphi}(u, \tau, \rho).$$

Proposition 4.12 implies that the right-hand side of this inequality converges to $-P_{\text{top}}((f_0)|_{\Lambda}, -\log J^+ f_0)$ as ε tends to zero, which completes the proof. \square

5.2 Remark: The preceding theorem contains as a special case the stabilization to a hyperbolic equilibrium point x_0 of f_0 . In this case, assumption (i) reduces to the

controllability of the linearization at (x_0, u^0) , and the lower bound reduces to

$$\log J^+ f_0(x_0) = \log |\det Df_0(x_0)|_{E_{x_0}^+}| = \sum_{\lambda \in \text{spec}(Df_0(x_0))} \max\{0, d_\lambda \log |\lambda|\},$$

where the sum is taken over the eigenvalues λ of $Df_0(x_0)$ with associated multiplicities d_λ . This lower bound was claimed in [53, Thm. 3] to hold (without the hyperbolicity assumption), but the proof presented there contains a gap.²⁴

5.3 Remark: The topological pressure $P_{\text{top}}((f_0)|_\Lambda, -\log J^+ f_0)$ is a well-studied quantity in the theory of dynamical systems. In particular, it is equal to the escape rate from a small neighborhood N of Λ , see [9, 67]:

$$P_{\text{top}}((f_0)|_\Lambda, -\log J^+ f_0) = \lim_{\tau \rightarrow \infty} \frac{1}{\tau} \log \text{vol}(\{x : f_0^t(x) \in N, 0 \leq t < \tau\}).$$

In the case when Λ reduces to a hyperbolic periodic orbit, the techniques applied to prove [53, Thm. 3] and [42, Thm. 4.3] together with Theorem 5.1 almost immediately yield the following data-rate theorem.

5.4 Theorem: *Consider the control system Σ and let the following assumptions be satisfied:*

- $U \subset \mathbb{R}^m$ with $U = \text{cl int } U$ and $f : M \times U \rightarrow M$ is continuously differentiable.
- For some $u^0 \in \text{int } U$ and $f_0 := f_{u^0}$, there exists a hyperbolic periodic orbit $\Lambda = \{x_0, f_0(x_0), \dots, f^{\tau-1}(x_0)\}$.²⁵
- The linearization of Σ along the orbit Λ is controllable on the time interval $[0; \tau]$.

Then the smallest average data rate R_0 above which Σ is locally uniformly stabilizable to Λ is given by

$$R_0 = \frac{1}{\tau} \sum_{\lambda \in \text{spec}(Df_0^\tau(x_0))} \max\{0, d_\lambda \log |\lambda|\}, \quad (46)$$

where we sum over the different eigenvalues λ of $Df_0^\tau(x_0)$ with associated multiplicities d_λ .

²⁴Actually, [53] studies asymptotic stabilization, but in the analysis of the minimal data rate there is no essential difference between stabilization and asymptotic stabilization.

²⁵Observe that a hyperbolic periodic orbit is always an isolated invariant set.

Proof: By the variational principle for pressure, we have

$$P_{\text{top}}((f_0)|_{\Lambda}, -\log J^+ f_0) = \sup_{\mu \in \mathcal{M}((f_0)|_{\Lambda})} \left[h_{\mu}(f_0) - \int \log J^+ f_0 \, d\mu \right].$$

Note that the measure-theoretic entropy $h_{\mu}(f_0)$ vanishes, because Λ is finite, and the only invariant probability measure on the periodic orbit is the one which puts equal mass to all points of Λ . Hence,

$$\begin{aligned} P_{\text{top}}((f_0)|_{\Lambda}, -\log J^+ f_0) &= -\frac{1}{\tau} \sum_{t=0}^{\tau-1} \log J^+ f_0(f_0^t(x_0)) \\ &= \frac{1}{\tau} \log J^+ f_0^{\tau}(x_0) = \frac{1}{\tau} \sum_{\lambda \in \text{spec}(Df_0^{\tau}(x_0))} \max\{0, d_{\lambda} \log |\lambda|\}, \end{aligned}$$

where we use that the unstable subspace at (u^0, x_0) is the sum of the generalized eigenspaces of the linear operator $Df_0^{\tau}(x_0) : T_{x_0}M \rightarrow T_{x_0}M$ corresponding to unstable eigenvalues. Consequently, Theorem 5.1 yields

$$R_0 \geq \frac{1}{\tau} \sum_{\lambda \in \text{spec}(Df_0^{\tau}(x_0))} \max\{0, d_{\lambda} \log |\lambda|\}.$$

For the converse inequality, we observe that the δ -neighborhood of Λ reduces to the union of the δ -balls around the points $x_0, f_0(x_0), \dots, f_0^{\tau-1}(x_0)$. Then the techniques used in the proof of [42, Thm. 4.3] (adapted to the discrete-time setting) show how one can use the regularity assumption to keep the state within an ε -neighborhood of the periodic orbit for all times with an average data rate arbitrarily close to the right-hand side of (46) if ε is chosen small enough (which actually works also without the hyperbolicity assumption). \square

6 An example built on the Hénon horseshoe

Consider the map

$$f(x, y) := (5 - 0.3y - x^2, x), \quad f : \mathbb{R}^2 \rightarrow \mathbb{R}^2,$$

which is a member of the Hénon family [35], one of the most-studied classes of dynamical systems that exhibit chaotic behavior.

Obviously, f is a polynomial, hence real-analytic diffeomorphism of \mathbb{R}^2 . We extend f to a control system with additive control:

$$\Sigma : \begin{pmatrix} x_{t+1} \\ y_{t+1} \end{pmatrix} = \begin{pmatrix} 5 - 0.3y_t - x_t^2 + u_t \\ x_t + v_t \end{pmatrix}, \quad u_t^1 + v_t^2 \leq 1.$$

According to [57, Thm. 4.2], the nonwandering set of f is a topologically transitive hyperbolic set Λ , contained in the square centered at the origin with side length

$$R := 1.3 + \sqrt{(1.3)^2 + 20}.$$

It is also known that in this case there exists an isolating neighborhood of Λ (cf. [9, Thm. 3.9]). The construction in Subsection 3.3 yields an $\varepsilon > 0$ so that the system

$$\Sigma^\varepsilon : \begin{pmatrix} x_{t+1} \\ y_{t+1} \end{pmatrix} = \begin{pmatrix} 5 - 0.3y_t - x_t^2 + u_t \\ x_t + v_t \end{pmatrix}, \quad u_t^2 + v_t^2 \leq \varepsilon^2$$

admits a hyperbolic set Q^ε which contains Λ as its 0-fiber such that $L(Q^\varepsilon)$ is isolated invariant and the fiber map is continuous.

It can also be shown that Q^ε has nonempty interior by applying Theorem 3.13, since one can clearly reach a set of nonempty interior from every $(x, y) \in Q^\varepsilon$ in only one step of time. Hence, uniform forward accessibility holds. It thus follows that $\text{int } Q^\varepsilon \neq \emptyset$ by Proposition 3.14 and also that complete controllability holds on an open and dense subset of Q^ε by Corollary 3.19. Figure 2 shows a numerical approximation of the set Q^ε for $\varepsilon = 0.08$ computed with the software tool SCOTS [59].




Figure 2: The set Q^ε for $\varepsilon = 0.08$.

Hence, Theorem 4.7 is applicable to Q^ε and we know that

$$h_{\text{inv}}(Q^\varepsilon) \geq \inf_{\mu \in \mathcal{M}(\Phi|_{L(Q^\varepsilon)})} \left[\int \log J^+ \varphi \, d\mu - h_\mu(\varphi, (\pi_{\mathcal{U}})_* \mu) \right].$$

As we have seen before, for $\varepsilon \rightarrow 0$, the right-hand side converges to $-P_{\text{top}}(f|_{\Lambda}; -\log J^+ f)$. Numerical studies from Froyland [30, Table 2], based on Ulam's method, suggest that

$$P_{\text{top}}(f|_{\Lambda}; -\log J^+ f) \approx -0.696.$$

Hence, according to our considerations in Subsection 4.9, we expect that $h_{\text{inv}}(Q^\varepsilon) \approx 0.696$ for all sufficiently small ε (and the same estimate should hold for the minimal average data rate for local stabilization to Λ).

It is also possible to work with a scalar control and consider the system

$$\Sigma' : \begin{pmatrix} x_{t+1} \\ y_{t+1} \end{pmatrix} = \begin{pmatrix} 5 - 0.3y_t - x_t^2 + u_t \\ x_t \end{pmatrix}, \quad |u_t| \leq 1.$$

In this case, some work is needed to check uniform forward accessibility. According to [38, Thm. 3], we need to check that $\dim L^+(x, y) = 2$ for all $(x, y) \in \mathbb{R}^2$, where

$$L^+ = \text{Lie}\{\text{Ad}_0^k X_u^+ : k \geq 0, u \in (-1, 1)\},$$

the Lie algebra generated by the vector fields $\text{Ad}_0^k X_u^+$, defined by

$$\text{Ad}_0^k X_u^+(x, y) = \left. \frac{\partial}{\partial v} \right|_{v=0} f_0^{-k} \circ f_u^{-1} \circ f_{u+v} \circ f_0^k(x, y).$$

A simple computation yields

$$f_u^{-1}(x, y) = (y, \frac{1}{0.3}(5 - y^2 - x + u)).$$

Hence, we can compute

$$\begin{aligned} \text{Ad}_0^0 X_u^+(x, y) &= \left. \frac{\partial}{\partial v} \right|_{v=0} f_u^{-1} \circ f_{u+v}(x, y) \\ &= \left. \frac{\partial}{\partial v} \right|_{v=0} f_u^{-1}(5 - 0.3y - x^2 + u + v, x) \\ &= \left. \frac{\partial}{\partial v} \right|_{v=0} (x, \frac{1}{0.3}(0.3y - v)) = (0, -\frac{1}{0.3}). \end{aligned}$$

In particular,

$$f_u^{-1}(f_{u+v}(x, y)) = (x, y - \frac{1}{0.3}v).$$

This can be used to compute

$$\begin{aligned} \text{Ad}_0^1 X_u^+(x, y) &= \left. \frac{\partial}{\partial v} \right|_{v=0} f_0^{-1} \circ f_u^{-1} \circ f_{u+v} \circ f_0(x, y) \\ &= \left. \frac{\partial}{\partial v} \right|_{v=0} f_0^{-1} \circ f_u^{-1} \circ f_{u+v}(5 - 0.3y - x^2, x) \\ &= \left. \frac{\partial}{\partial v} \right|_{v=0} f_0^{-1}(5 - 0.3y - x^2, x - \frac{1}{0.3}v) \end{aligned}$$

$$\begin{aligned}
&= \frac{\partial}{\partial v} \Big|_{v=0} \left(x - \frac{1}{0.3}v, \frac{1}{0.3} \left(5 - \left(x - \frac{1}{0.3}v \right)^2 - 5 + 0.3y + x^2 \right) \right) \\
&= \left(-\frac{1}{0.3}, \frac{2}{(0.3)^2}x \right).
\end{aligned}$$

Since L_+ contains all linear combinations of the vector fields $\text{Ad}_0^0 X_u^+$ and $\text{Ad}_0^1 X_u^+$, we see that $L^+(x, y) = \mathbb{R}^2$ for all $(x, y) \in \mathbb{R}^2$. Hence, forward accessibility holds (which can easily be seen to be uniform) and our statements about Σ also hold for Σ' .

7 Open questions

The results presented in this paper leave many questions open. In the author's opinion, the most important ones are the following:

- Are there non-trivial hyperbolic sets of control systems which do not arise by the small-perturbation construction?
- Is the fiber map $u \mapsto Q(u)$ lower semicontinuous for a general hyperbolic set with isolated invariant lift? Is it at least lower semicontinuous in the special case when the fibers are finite (but not singletons)?
- Can the results of Subsection 3.3 about the controllability properties on the set Q obtained by the small-perturbation construction be generalized to C^∞ (instead of analytic) systems?²⁶
- Are the hyperbolic sets constructed from small perturbations *control sets* under the assumptions of Corollary 3.19? That is, are they maximal with the property of complete approximate controllability? (This is most probably equivalent to $L(Q)$ being a maximal chain transitive set.)
- How can we prove a general achievability result?
- Can our results about local stabilization be generalized to the continuous-time case, when Λ is a hyperbolic set (with one-dimensional center bundle) of a system given by an ordinary differential equation?

A Some auxiliary results

The proof of the following lemma was provided by Niels J. Diepeveen.²⁷

²⁶In the continuous-time case, results on genericity of universally regular controls exist for C^∞ -systems [19] and have been used in the continuous-time analysis of invariance entropy, see [25].

²⁷See <https://mathoverflow.net/questions/332191/>

A.1 Lemma: Let (X, d) be a compact metric space and let $X^{\mathbb{Z}}$ be equipped with the sup-metric

$$d_{\infty}(x, y) := \sup_{n \in \mathbb{Z}} d(x_n, y_n) \quad \text{for all } x = (x_n), y = (y_n).$$

Then $(X^{\mathbb{Z}}, d_{\infty})$ is connected if and only if (X, d) is connected.

Proof: The projection $(x_n)_{n \in \mathbb{Z}} \mapsto x_0$ from $X^{\mathbb{Z}}$ to X is continuous and surjective. Hence, connectedness of $X^{\mathbb{Z}}$ implies connectedness of X . To see that the converse holds, assume that X is connected and let $F \subset X^{\mathbb{Z}}$ be the subset of all sequences that assume only finitely many values. Since (X, d) is totally bounded, F is dense in $(X^{\mathbb{Z}}, d_{\infty})$. Hence, it suffices to prove that F is connected. To this end, we fix arbitrary $a, b \in F$ and construct a connected subset of F that contains a and b . Let P be a finite partition of \mathbb{Z} into subsets on which both a and b are constant and consider the map $i : X^P \rightarrow F$ given by $i(x)_n := x([n]_P)$, where $[n]_P$ denotes the unique element of P containing n . The map i is an isometric embedding of X^P (equipped with the sup-metric) into F and both a and b are contained in its image. Since X^P is a finite product of copies of X , connectedness of X implies connectedness of X^P (using the fact that the product topology coincides with the uniform topology on finite products). Hence, $i(X^P)$ is the desired subset. \square

A.2 Lemma: Let M be a Riemannian manifold and $K \subset M$ a nonempty compact subset. Then for every $\varepsilon > 0$, the boundary of $N_{\varepsilon}(K)$ has volume zero.

Proof: We give the proof for $M = \mathbb{R}^n$ with the Euclidean metric induced by the Euclidean norm $\|\cdot\|$. The general case can be proved by replacing straight lines with geodesics. Hence, let $K \subset \mathbb{R}^n$ be a nonempty compact set and $\varepsilon > 0$. Take $x \in \partial N_{\varepsilon}(K)$ and fix a point $y \in K$ such that $\text{dist}(x, K) = \|x - y\| = \varepsilon$. We claim that the open ball $B_{\varepsilon}(y)$ is contained in $N_{\varepsilon}(K)$ and does not contain any point from $\partial N_{\varepsilon}(K)$. Indeed, if $z \in B_{\varepsilon}(y)$, then $\text{dist}(z, K) \leq \|z - y\| < \varepsilon$ and all points $w \in \partial N_{\varepsilon}(K)$ satisfy $\text{dist}(w, K) = \varepsilon$ implying $\|w - y\| \geq \varepsilon$. Let $r \in (0, \varepsilon)$. Then the intersection $B_r(x) \cap B_{\varepsilon}(y)$ contains the ball $B_{r/2}(tx + (1-t)y)$ with $t := 1 - r/(2\varepsilon)$. Indeed, if $w \in B_{r/2}(tx + (1-t)y)$, then

$$\begin{aligned} \|w - x\| &\leq \|w - tx - (1-t)y\| + \|tx + (1-t)y - x\| \\ &< \frac{r}{2} + (1-t)\|x - y\| = \frac{r}{2} + \frac{r}{2\varepsilon}\varepsilon = r, \\ \|w - y\| &\leq \|w - tx - (1-t)y\| + \|tx + (1-t)y - y\| \\ &< \frac{r}{2} + t\|x - y\| = \frac{r}{2} + \left(\varepsilon - \frac{r}{2}\right) = \varepsilon. \end{aligned}$$

Hence, for all $r \in (0, \varepsilon)$ we have

$$\frac{\text{vol}(B_r(x) \cap \partial N_{\varepsilon}(K))}{\text{vol}(B_r(x))} \leq \frac{cr^n - c(r/2)^n}{cr^n} = 1 - 2^{-n} < 1.$$

This proves that the density $d(x) = \lim_{r \downarrow 0} \text{vol}(B_r(x) \cap \partial N_\varepsilon(K)) / \text{vol}(B_r(x))$ is less than one wherever it exists on $\partial N_\varepsilon(K)$. Lebesgue's density theorem asserts that $d(x) = 1$ at almost every point of $\partial N_\varepsilon(K)$. This can only be the case if $\text{vol}(\partial N_\varepsilon(K)) = 0$. \square

The next lemma is essentially taken from [31, Lem. 2.4].

A.3 Lemma: *Let $f : X \rightarrow X$ be a map on some set X and $v : \mathbb{Z}_+ \times X \rightarrow \mathbb{R}$ a subadditive cocycle over f , i.e.,*

$$v_{n+m}(x) \leq v_n(x) + v_m(f^n(x)) \quad \text{for all } x \in X, n, m \in \mathbb{Z}_+.$$

Additionally suppose that

$$\omega := \sup_{(n,x) \in \mathbb{Z}_{>0} \times X} \frac{1}{n} |v_n(x)| < \infty. \quad (47)$$

Then for every $x \in X$, $n \in \mathbb{Z}_{>0}$ and $\varepsilon \in (0, 2\omega)$ there is a time $0 \leq n_1 < n$ with

$$\frac{1}{k} v_k(f^{n_1}(x)) > \frac{1}{n} v_n(x) - \varepsilon \quad \text{for all } 0 < k \leq n - n_1.$$

Moreover, $n - n_1 \geq (\varepsilon n) / (2\omega) \rightarrow \infty$ for $n \rightarrow \infty$.

Proof: We write $\sigma := v_n(x)/n$ and define

$$\gamma := \min_{0 < k \leq n} \frac{1}{k} v_k(x).$$

If $\gamma \geq \sigma - \varepsilon$, the assertion follows with $n_1 = 0$. For $\gamma < \sigma - \varepsilon$, observing that the minimum cannot be attained at $k = n$, let

$$n_1 := \max \left\{ k \in (0, n) \cap \mathbb{Z} : \frac{1}{k} v_k(x) \leq \sigma - \varepsilon \right\},$$

implying $v_{n_1}(x)/n_1 \leq \sigma - \varepsilon$. We obtain

$$\begin{aligned} \varepsilon &\leq \frac{1}{n} v_n(x) - \frac{1}{n_1} v_{n_1}(x) = \frac{1}{n} v_{n_1+(n-n_1)}(x) - \frac{1}{n_1} v_{n_1}(x) \\ &\leq \frac{1}{n} (v_{n_1}(x) + v_{n-n_1}(f^{n_1}(x))) - \frac{1}{n_1} v_{n_1}(x) \\ &= \frac{1}{n} \left(-\frac{n-n_1}{n_1} v_{n_1}(x) + \frac{n-n_1}{n-n_1} v_{n-n_1}(f^{n_1}(x)) \right) \\ &= \frac{n-n_1}{n} \left(\frac{1}{n-n_1} v_{n-n_1}(f^{n_1}(x)) - \frac{1}{n_1} v_{n_1}(x) \right) \leq 2\omega \frac{n-n_1}{n}. \end{aligned}$$

This implies

$$n - n_1 \geq \frac{\varepsilon n}{2\omega} \rightarrow \infty \quad \text{for } n \rightarrow \infty.$$

For $0 < k \leq n - n_1$ we have $v_{k+n_1}(x)/(k + n_1) > \sigma - \varepsilon$ and this yields

$$\begin{aligned} \frac{1}{k} v_k(f^{n_1}(x)) &\geq \frac{1}{k} (v_{k+n_1}(x) - v_{n_1}(x)) \\ &> \frac{1}{k} ((k + n_1)(\sigma - \varepsilon) - n_1(\sigma - \varepsilon)) = \sigma - \varepsilon, \end{aligned}$$

completing the proof. \square

A.4 Lemma: Let $n > m$ be positive integers. For each i in the range $0 \leq i < m$ choose integers q_i, r_i such that $n = i + q_i m + r_i$ with $q_i \geq 0$ and $0 \leq r_i < m$. Then

$$\{0, 1, \dots, n - m\} = \{i + jm : 0 \leq i < m, 0 \leq j < q_i\},$$

and all integers in the set on the right-hand side are uniquely parametrized by i and j .

Proof: It is clear that $(i_1, j_1) \neq (i_2, j_2)$ implies $i_1 + j_1 m \neq i_2 + j_2 m$, since $0 \leq i_1, i_2 < m$. Hence, it suffices to show that the two sets are equal. To this end, we first show that $i + jm \leq n - m$, whenever $0 \leq i < m$ and $0 \leq j < q_i$. Since $j < q_i$, we have $(j + 1)m \leq q_i m + r_i$. Adding i on both sides yields $(i + jm) + m \leq n$, or equivalently $i + jm \leq n - m$.

Conversely, let us show that every number l between 0 and $n - m$ can be written as $i + jm$ with $0 \leq i < m$ and $0 \leq j < q_i$. To this end, let i, j be the unique nonnegative integers so that $l = i + jm$ with $0 \leq i < m$. We need to show that $j < q_i$. This is equivalent to

$$l = i + jm < i + q_i m = n - r_i.$$

This inequality holds, because $l < n - (m - 1) \leq n - r_i$ using that $0 \leq r_i < m$. \square

A.5 Lemma: Let (X, d) be a compact metric space and let μ be a Borel probability measure on X . Then, for any $\delta > 0$ there exists a finite measurable partition $\xi = \{C_1, \dots, C_k\}$ of X with $\text{diam}(C_i) < \delta$ and $\mu(\partial C_i) = 0$ for $i = 1, \dots, k$.

Proof: For each $x \in X$, let us consider the disjoint uncountable union $\bigcup_{\varepsilon \in (0, \delta)} \partial B_\varepsilon(x)$, which has finite measure. We assume to the contrary that $\mu(\partial B_\varepsilon(x))$ is positive for every $\varepsilon \in (0, \delta)$. Then $(0, \delta)$ is the (countable) union of the sets $I_n := \{\varepsilon \in (0, \delta) : \mu(\partial B_\varepsilon(x)) > 1/n\}$, $n \in \mathbb{Z}_{>0}$. Hence, one of these sets must be uncountable, which is a contradiction. Thus, for each $x \in X$ there is $\varepsilon = \varepsilon(x) \in (0, \delta)$ with $\mu(\partial B_\varepsilon(x)) = 0$. By compactness, there exists a cover of X consisting of finitely many of such balls, say B_1, \dots, B_k . From this cover we can construct the desired partition by $C_1 := \text{cl } B_1$, $C_i := \text{cl } B_i \setminus \bigcup_{j=1}^{i-1} \text{cl } B_j$ for $i > 1$. Then $\xi := \{C_1, \dots, C_k\}$ satisfies $\bigcup_{i=1}^k \partial C_i \subset \bigcup_{i=1}^k \partial B_i$, and hence $\mu(\bigcup_{i=1}^k \partial C_i) = 0$. \square

B Elementary properties of hyperbolic sets

The following proposition answers some questions that immediately arise from the definition of a hyperbolic set (Definition 3.3).

B.1 Proposition: *The following statements hold:*

- (i) *The definition of a hyperbolic set is independent of the choice of the Riemannian metric on M . In fact, only the constant c depends on the choice of the metric.*
- (ii) *The inequality for tangent vectors $v \in E^+(u, x)$ expressed in (H2) is equivalent to:*

$$|D\varphi_{t,u}(x)v| \geq c^{-1}\lambda^{-t}|v| \quad \text{for all } (u, x) \in L(Q), v \in E^+(u, x), t \in \mathbb{Z}_+.$$

- (iii) *The subspaces $E^\pm(u, x)$ depend continuously on $(u, x) \in L(Q)$, meaning that the projections*

$$\pi_{u,x}^\pm : T_x M \rightarrow E^\pm(u, x)$$

along the respective complementary subspace depend continuously on (u, x) .

Proof: (i) This follows from the fact that any two Riemannian metrics are equivalent on the compact set Q which is shown as follows. Let g and h be two Riemannian metrics on M . Let $S_h Q$ denote the unit tangent bundle over Q with respect to h , i.e., the closed subspace of the tangent bundle that consists of all tangent vectors $v \in T_x M$ satisfying $x \in Q$ and $h(v, v) = 1$. Observe that $S_h Q$ is compact. Since g is continuous, there are constants $0 < \alpha \leq \beta < \infty$ with $\alpha \leq g(v, v) \leq \beta$ for all $v \in S_h Q$. Then, for any $0 \neq v \in T_x M, x \in Q$, we have

$$g(v, v) = h(v, v) \cdot g\left(\frac{v}{\sqrt{h(v, v)}}, \frac{v}{\sqrt{h(v, v)}}\right),$$

which in turn implies

$$\alpha h(v, v) \leq g(v, v) \leq \beta h(v, v).$$

Hence, writing $|\cdot|_h$ and $|\cdot|_g$ for the norms associated with g and h , respectively, the inequality $|D\varphi_{t,u}(x)v|_g \leq c\lambda^t|v|_g$ implies

$$\begin{aligned} |D\varphi_{t,u}(x)v|_h &= h(D\varphi_{t,u}(x)v, D\varphi_{t,u}(x)v)^{1/2} \\ &\leq \frac{1}{\sqrt{\alpha}} |D\varphi_{t,u}(x)v|_g \leq \frac{1}{\sqrt{\alpha}} c\lambda^t |v|_g \leq \sqrt{\frac{\beta}{\alpha}} \lambda^t |v|_h. \end{aligned}$$

This implies the statement.

(ii) Given $v \in E^+(u, x)$, by (H1), we have $D\varphi_{t,u}(x)v \in E^+(\Phi_t(u, x))$. Hence, (H2) implies

$$|D\varphi_{-t,\theta^t u}D\varphi_{t,u}(x)v| \leq c\lambda^t |D\varphi_{t,u}(x)v|$$

for every $t \in \mathbb{Z}_+$. From the cocycle property of φ it follows that

$$D\varphi_{-t,\theta^t u}(\varphi_{t,u}(x))D\varphi_{t,u}(x)v = D(\varphi_{-t,\theta^t u} \circ \varphi_{t,u})(x)v = D(\text{id})v = v,$$

implying $|D\varphi_{t,u}(x)v| \geq c^{-1}\lambda^{-t}|v|$. Going backwards through these inequalities, the other direction of the equivalence follows.

(iii) Let d^- denote the common dimension of the stable subspaces and let $(u_k, x_k)_{k \in \mathbb{Z}_{\geq 0}}$ be a sequence in $L(Q)$, converging to some $(u, x) \in L(Q)$. We choose an orthonormal basis $(v_k^{(1)}, \dots, v_k^{(d^-)})$ of each $E^-(u_k, x_k)$. By compactness of the unit tangent bundle over Q , passing to a subsequence if necessary²⁸ yields the convergence $v_k^{(i)} \rightarrow v^{(i)}$ for some $v^{(i)} \in T_x M$, $i = 1, \dots, d^-$. For each i and n , the inequality $|D\varphi_{t,u_k}(x_k)v_k^{(i)}| \leq c\lambda^t |v_k^{(i)}|$ carries over to the limit for $k \rightarrow \infty$, since $(u, x, v) \mapsto D\varphi_{t,u}(x)v$ is a continuous map by our assumptions on the system. Hence, $|D\varphi_{t,u}(x)v^{(i)}| \leq c\lambda^t |v^{(i)}|$ holds for all $t \geq 0$. Since the subspace $E^-(u, x)$ is characterized uniquely by these inequalities, it follows that $v^{(i)} \in E^-(u, x)$. Hence, $(v^{(1)}, \dots, v^{(d^-)})$ is an orthonormal basis of $E^-(u, x)$, which implies the assertion (similarly for $E^+(u, x)$). \square

B.2 Remark: Item (ii) in the above proposition shows that the “contraction in backward time” property of E^+ can equivalently be expressed as “expansion in forward time”. Hence, one might ask why we should not use this expansion property to define a hyperbolic set (as it is more intuitive and we are mainly interested in the behavior of the system in forward time). The answer to this question is that the expansion property does not uniquely characterize the unstable subspaces. Expansion also happens outside of the unstable subspaces, while contraction in backward time does not, as we have used in the proof of item (iii).

Acknowledgements

Particular thanks go to Serdar Yüksel and Nicolás Garcia for their interest in my work and their feedback on the manuscript. Moreover, I want to thank Adriano Da Silva with whom I have developed, in several collaborations, many of the ideas that went into this paper. Special thanks also go to Niels J. Diepeveen who contributed the proof of Lemma A.1, to Thomas Barthelmé who helped me to confirm an argument in the proof of Theorem 4.17, and to Mahendra Singh Tomar who created Figure 2. Finally, I want to mention Fritz Colonius and Matthias Gundlach for comments leading to an improvement of the paper.

²⁸Observe: proving the convergence for a subsequence is enough. Indeed, if continuity at (u, x) would not hold, then there would exist a sequence $(u_k, x_k) \rightarrow (u, x)$ so that for *no* subsequence the convergence $E^-(u_k, x_k) \rightarrow E^-(u, x)$ would be true.

References

- [1] R. L. Adler, A. G. Konheim, M. H. McAndrew. *Topological entropy*. Trans. Am. Math. Soc. 114 (1965), 309–319.
- [2] B. R. Andrievskii, A. S. Matveev, A. L. Fradkov. *Control and estimation under information constraints: toward a unified theory of control, computation, and communications*. (Russian) Avtomat. i Telemekh. 2010, no. 4, 34–99; translation in Autom. Remote Control 71 (2010), no. 4, 572–633.
- [3] F. Albertini, E. D. Sontag. *Discrete-time transitivity and accessibility: analytic systems*. SIAM J. Control Optim. 31 (1993), no. 6, 1599–1622.
- [4] J. Bahnmüller, T. Bogenschütz. *A Margulis-Ruelle inequality for random dynamical systems*. Arch. Math. (Basel) 64 (1995), no. 3, 246–253.
- [5] J. Bahnmüller, P.–D. Liu. *Characterization of measures satisfying the Pesin entropy formula for random dynamical systems*. J. Dynam. Differential Equations 10 (1998), no. 3, 425–448.
- [6] G. O. Berger, R. M. Jungers. *Formal methods for computing hyperbolic invariant sets for nonlinear systems*. IEEE Control Syst. Lett. 4 (2019), no. 1, 235–240.
- [7] T. Bogenschütz. *Entropy, pressure, and a variational principle for random dynamical systems*. Random Comput. Dynam. 1 (1992/93), no. 1, 99–116.
- [8] R. Bowen. *Entropy for group endomorphisms and homogeneous spaces*. Trans. Am. Math. Soc. 153 (1971), 401–414.
- [9] R. Bowen. *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms*. Second revised edition. Lecture Notes in Math. 470, Springer, Berlin, 2008.
- [10] R. Bowen, D. Ruelle. *The ergodic theory of Axiom A flows*. Invent. Math. 29 (1975), no. 3, 181–202.
- [11] M. L. Cartwright, J. E. Littlewood. *On non-linear differential equations of the second order: I*. J. Lond. Math. Soc. 20 (1945), 180–189.
- [12] F. Colonius. *Metric invariance entropy and conditionally invariant measures*. Ergodic Theory Dynam. Systems 38 (2018), no. 3, 921–939.
- [13] F. Colonius, A. J. Santana, J. Cossich. *Invariance pressure for control systems*. J. Dynam. Differential Equations 31 (2019), no. 1, 1–23.
- [14] F. Colonius, W. Du. *Hyperbolic control sets and chain control sets*. J. Dynam. Control Systems 7 (2001), no. 1, 49–59.

- [15] F. Colonius, C. Kawan. *Invariance entropy for control systems*. SIAM J. Control Optim. 48 (2009), no. 3, 1701–1721.
- [16] F. Colonius, C. Kawan, G. N. Nair. *A note on topological feedback entropy and invariance entropy*. Systems Control Lett. 62 (2013), no. 5, 377–381.
- [17] F. Colonius, W. Kliemann. *The Dynamics of Control*. Birkhäuser, Boston, 2000.
- [18] F. Colonius, W. Kliemann. *Dynamical Systems and Linear Algebra*. Graduate Studies in Mathematics 158, AMS, Providence, RI, 2014.
- [19] J.-M. Coron. *Linearized control systems and applications to smooth stabilization*. SIAM J. Control Optim. 32 (1994), no. 2, 358–386.
- [20] A. Da Silva. *Invariance entropy for random control systems*. Math. Control Signals Systems 25 (2013), no. 4, 491–516.
- [21] A. Da Silva. *Outer invariance entropy for linear systems on Lie groups*. SIAM J. Control Optim. 52 (2014), no. 6, 3917–3934.
- [22] A. Da Silva, C. Kawan. *Invariance entropy for a class of partially hyperbolic sets*. Math. Control Signals Systems 30 (2018), no. 4, 40 pp.
- [23] A. Da Silva, C. Kawan. *Robustness of critical bit rates for practical stabilization of networked control systems*. Automatica J. IFAC 93 (2018), 397–406.
- [24] A. Da Silva, C. Kawan. *Hyperbolic chain control sets on flag manifolds*. J. Dyn. Control Syst. 22 (2016), no. 4, 725–745.
- [25] A. Da Silva, C. Kawan. *Invariance entropy of hyperbolic control sets*. Discrete Contin. Dyn. Syst. 36 (2016), no. 1, 97–136.
- [26] A. Da Silva, C. Kawan. *Lyapunov exponents and partial hyperbolicity of chain control sets on flag manifolds*. Israel J. Math. 232 (2019), no. 2, 947–1000.
- [27] T. Downarowicz. *Entropy in Dynamical Systems*. New Mathematical Monographs 18, Cambridge University Press, Cambridge, 2011.
- [28] M. Franceschetti, P. Minero. *Elements of information theory for networked control systems*. In Information and Control in Networks (pp. 3–37). Springer International Publishing, 2014.
- [29] S. Fang, J. Chen, H. Ishii. *Towards Integrating Control and Information Theories: from Information-Theoretic Measures to Control Performance Limitations*. Vol. 465, Springer, 2016.
- [30] G. Froyland. *Using Ulam’s method to calculate entropy and other dynamical invariants*. Nonlinearity 12 (1999), no. 1, 79–101.

- [31] L. Grüne. *A uniform exponential spectrum for linear flows on vector bundles*. J. Dynam. Differential Equations 12 (2000), no. 2, 435–448.
- [32] V. M. Gundlach, Y. Kifer. *Random hyperbolic systems*. Stochastic dynamics (Bremen, 1997), 117–145, Springer, New York, 1999.
- [33] B. Hasselblatt. *Hyperbolic dynamical systems*. Handbook of dynamical systems, Vol. 1A, 239–319, North-Holland, Amsterdam, 2002.
- [34] B. Hasselblatt, Y. Pesin. *Partially hyperbolic dynamical systems*. Handbook of dynamical systems. Vol. 1B, 1–55, Elsevier, Amsterdam, 2006.
- [35] M. Hénon. *A two-dimensional mapping with a strange attractor*. In: The Theory of Chaotic Attractors, 94–102. Springer, 1976.
- [36] J. Hespanha, P. Naghshtabrizi, Y. Xu. *A survey of recent results in networked control systems*. IEEE Proc. 95 (2007), 138–162.
- [37] Y. Huang, X. Zhong. *Carathéodory–Pesin structures associated with control systems*. Systems Control Lett. 112 (2018), 36–41.
- [38] B. Jakubczyk, E. D. Sontag. *Controllability of nonlinear discrete-time systems: A Lie-algebraic approach*. SIAM J. Control Optim. 28 (1990), no. 1, 1–33.
- [39] A. Katok, B. Hasselblatt. *Introduction to the Modern Theory of Dynamical Systems*. Encyclopedia of Mathematics and its Applications, 54. Cambridge University Press, Cambridge, 1995.
- [40] C. Kawan. *Lower bounds for the strict invariance entropy*. Nonlinearity 24 (2011), no. 7, 1909–1935.
- [41] C. Kawan. *On the structure of uniformly hyperbolic chain control sets*. Systems Control Lett. 90 (2016), 71–75.
- [42] C. Kawan. *Invariance entropy of control sets*. SIAM J. Control Optim. 49 (2011), no. 2, 732–751.
- [43] C. Kawan. *Invariance Entropy for Deterministic Control Systems. An Introduction*. Lecture Notes in Math. 2089. Springer, Cham, 2013.
- [44] C. Kawan, J.–Ch. Delvenne. *Network entropy and data rates required for networked control*. IEEE Trans. Control Netw. Syst. 3 (2015), no. 1, 57–66.
- [45] C. Kawan, T. Stender. *Growth rates for semiflows on Hausdorff spaces*. J. Dynam. Differential Equations 24 (2012), 369–390.
- [46] Y. Kifer, P.–D. Liu. *Random dynamics*. Handbook of dynamical systems, Vol. 1B, 379–499, Elsevier B. V., Amsterdam, 2006.

- [47] N. Levinson. *A second order differential equation with singular solutions*. Ann. of Math. (2) 50 (1949), 127–153.
- [48] P.-D. Liu. *Random perturbations of Axiom A basic sets*. J. Statist. Phys. 90 (1998), no. 1-2, 467–490.
- [49] J. Lunze (Ed.) *Control Theory of Digitally Networked Dynamic Systems*. Vol. 1. Springer, 2014.
- [50] A. S. Matveev, A. V. Savkin. *Estimation and Control over Communication Networks*. Birkhäuser, Boston, 2008.
- [51] K. R. Meyer, X. Zhang. *Stability of skew dynamical systems*. J. Differential Equations 132 (1996), no. 1, 66–86.
- [52] I. Morris. *Mather sets for sequences of matrices and applications to the study of joint spectral radii*. Proc. Lond. Math. Soc. (3) 107 (2013), no. 1, 121–150.
- [53] G. N. Nair, R. J. Evans, I. M. Y. Mareels, and W. Moran. *Topological feedback entropy and nonlinear stabilization*. IEEE Trans. Automat. Control 49 (2004), no. 9, 1585–1597.
- [54] G. N. Nair, F. Fagnani, S. Zampieri, S., R. J. Evans. *Feedback control under data rate constraints: An overview*. Proceedings of the IEEE 95 (2007), no. 1, 108–137.
- [55] E. Ott, C. Grebogi, J. A. Yorke. *Controlling chaos*. Phys. Rev. Lett. 64 (1990), no. 11, 1196–1199.
- [56] H. Poincaré. *Sur le problème des trois corps et les équations de la dynamique*. Acta Math. 13 (1890), no. 1, A3–A270.
- [57] C. Robinson. *Dynamical Systems. Stability, Symbolic Dynamics, and Chaos*. CRC Press, Boca Raton, FL, 1995.
- [58] L. A. B. San Martin, L. Seco. *Morse and Lyapunov spectra and dynamics on flag bundles*. Ergodic Theory Dynam. Systems 30 (2010), no. 3, 893–922.
- [59] M. Rungger, M. Zamani. *SCOTS: A tool for the synthesis of symbolic controllers*. In Proceedings of the 19th Int. Conf. on Hybrid Systems: Computation and control, 2016, 99–104 [Online]. Available: <https://www.hyconsys.com/software/scots/>
- [60] J. Sliz, A. Süli, T. Kovacs. *Control of chaos in the vicinity of the Earth-Moon L5 Lagrangian point to keep a spacecraft in orbit*. Astron. Nachrichten 336 (2015), no. 1, 23–31.
- [61] S. Smale. *Differentiable dynamical systems*. Bull. Amer. Math. Soc. 73 (1967), no. 6, 747–817.

- [62] E. D. Sontag. *Mathematical Control Theory. Deterministic Finite-Dimensional Systems*. 2nd Edition. Texts in Applied Mathematics 6. Springer, New York, 1998.
- [63] E. D. Sontag, F. R. Wirth. *Remarks on universal nonsingular controls for discrete-time systems*. Systems Control Lett. 33 (1998), no. 2, 81–88.
- [64] M. S. Tomar, M. Zamani. *Compositional quantification of invariance feedback entropy for networks of uncertain control systems*. IEEE Control Syst. Lett. 4 (2020), no. 4, 827–832.
- [65] T. Wang, Y. Huang, H.-W. Sun. *Measure-theoretic invariance entropy for control systems*. SIAM J. Control Optim. 57 (2019), no. 1, 310–333.
- [66] F. R. Wirth. *Dynamics and controllability of nonlinear discrete-time control systems*. IFAC Proceedings Volumes 31 (1998), no. 17, 267–272.
- [67] L.-S. Young. *Large deviations in dynamical systems*. Trans. Amer. Math. Soc. 318 (1990), no. 2, 525–543.
- [68] L.-S. Young. *What are SRB measures, and which dynamical systems have them?*. J. Statistical Physics 108 (2002), nos. 5–6, 733–754.
- [69] S. Yüksel, T. Başar. *Stochastic Networked Control Systems: Stabilization and Optimization under Information Constraints*. New York, NY: Springer-Birkhäuser, 2013.