

Weak value vector potential with gauge invariance

Sunkyu Yu, Xianji Piao and Namkyoo Park*

Photonic Systems Laboratory, Department of Electrical and Computer Engineering, Seoul National University, Seoul 08826, Korea

E-mail: nkpark@snu.ac.kr

Received xxxxxx

Accepted for publication xxxxxx

Published xxxxxx

Abstract

The conservation of quantities or dynamics under coordinate transformations, known as gauge invariance, has been the foundation of theoretical frameworks in physics. The finding of gauge-invariant quantities has provided a new way of understanding physical problems, as shown in geometric and topological interpretations of quantum phenomena with the Berry phase, or in the separation of quark and gluon contributions in quantum chromodynamics. Here, focusing on the Berry phase, we extract a new gauge-invariant quantity. By employing different pre- and post-selections in the Berry phase in the context of weak values, we derive the gauge-invariant vector potential from the Berry connection that is originally gauge-dependent, and show that the obtained vector potential corresponds to the weak value of the projected momentum operator in the parameter space. The local nature of this quantity is demonstrated with the Aharonov-Bohm effect, proving that this gauge-invariant vector potential can be interpreted as the only source of the Berry curvature in the magnetic field. This weak value decomposition approach could lead to the identification of new measurable quantities from traditionally unobservable quantities.

Keywords: Berry phase, weak value, Aharonov-Bohm effect

1. Introduction

Gauge invariance, having deep roots in the foundation of electromagnetism [1] since H. Weyl's seminal paper [2,3], constitutes one of the pillars of modern physics. The finding of invariant quantities or dynamics under gauge transformation allows the composition of an observable system with measurable quantities to be determined [4]. The most introductory and historical example is found in classical electromagnetics [5], as shown in the comparison between the gauge-invariant, measurable quantities of the fields (\mathbf{E} , \mathbf{H}) and the gauge-dependent, non-unique quantities of the potentials (\mathbf{A} , V). Other examples include the separation of quark and gluon contributions to the nucleon spin, using gauge-invariant decomposition in quantum chromodynamics [6].

The importance of gauge invariance can also be found in the Berry phase [7,8], which describes geometrically defined phase accumulation based on the adiabatic change of a system in a certain parameter space. In his original paper [7] that provided a framework for the "geometrization" of physical problems [8,9] also in relation to topological theory [10,11], M. V. Berry focused on the vector-potential-like term known as the Berry connection, $\mathbf{A}_n = i\langle u_n | \nabla_{\mathbf{R}} | u_n \rangle$, where $|u_n\rangle$ is the n^{th} system eigenstate and \mathbf{R} is the parameter space. Similar to the electromagnetic vector potential \mathbf{A} , Berry connection \mathbf{A}_n is gauge-dependent for the transformation of the wavefunction $\mathbf{A}_n \rightarrow \mathbf{A}_n - \nabla_{\mathbf{R}} \zeta_n(\mathbf{R})$ with $|u_n\rangle \rightarrow \exp[i\zeta_n(\mathbf{R})]|u_n\rangle$ and had been ignored for a long time prior to the identification of the Berry phase [7,8] – the *cyclic integral* of \mathbf{A}_n in the \mathbf{R} -space associated with gauge invariance. Although the concept of the Berry phase has been extended to non-adiabatic (Aharonov-

Anandan phase) [12] and non-cyclic (Pancharatnam phase) [13-15] evolution in the ray space, the Berry connection \mathbf{A}_n or other geometric connections [12-14], which are the sources of quantum geometric quantities including Berry phase and curvature, have remained gauge-dependent, similar to \mathbf{A} in electromagnetics.

Considering the representation of $\mathbf{A}_n = \langle u_n | i \nabla_{\mathbf{R}} | u_n \rangle$, \mathbf{A}_n can be interpreted as the expectation value of $|u_n\rangle$ for the operator $i \nabla_{\mathbf{R}}$, when the operator is observable. This relation naturally inspires the *extended* representation of \mathbf{A}_n with the concept of a “weak value” [16-18], which describes the information of a physical quantity through a weak measurement. Because such a weak measurement with a weakly coupled system and probe permits an almost undisturbed initial state [17], we can envision the measurement of a certain expectation value through a set of weak values from multiple weak measurements. In terms of the measurements related to gauge invariance, the question then arises: Does each “partial” weak value have the same gauge condition as that of the original expectation value? The answer is directly connected to the finding of new measurable quantities from an unobservable one in terms of the concept of a weak value. Notably, although the weak value of the gauge-invariant geometric *phase* has been studied [19-21], the application of weak values to the gauge-dependent, unobservable Berry *connection* has not been considered.

In this paper, we extract the gauge-invariant vector potential from the weak value representation of the Berry connection. By employing different pre- and post-selected states [16-18], we revisit the derivation of the Berry phase to achieve the weak value expression of the Berry connection \mathbf{A}_n , which yields the decomposition of \mathbf{A}_n into gauge-dependent and gauge-invariant weak-value vector potentials. This decomposition provides a measurable condition for the mutually projected momentum weak value in the parameter space \mathbf{R} , for an arbitrary definition of the post selection (or, an arbitrary weak measurement condition). As a representative example, we apply this approach to the Aharonov-Bohm effect to illustrate the local nature of the newly found gauge-invariant weak value. Finally, we discuss the topological nature of the gauge-invariant weak value, showing that this gauge-invariant weak value becomes the only source of the gauge-invariant Berry curvature in the Aharonov-Bohm effect with a *nonzero* magnetic field, or more generally, in the post-selection with parameter state $\langle \mathbf{R} |$. By revealing a new measurable geometric quantity in the form of weak value vector potentials, which is well-defined for an arbitrary weak measurement, our results provide novel insight into the measurements of geometric and topological properties in quantum mechanics [8,10] and optics [9,11,22].

2. Weak-value decomposition of Berry connection

Following the derivation of the Berry phase [7,8,23], consider a time-dependent Hamiltonian $H = H(t)$. We focus on the Hamiltonians which can be described by a set of parameters $\mathbf{R} = (R_1, R_2, \dots)$, where \mathbf{R} represents a set of conserved quantities such as position or momentum vectors that allow for the state vector representation $|\mathbf{R}\rangle$ defined in a complex Hilbert space. The Hamiltonian for a spinless particle under a uniform magnetic field is a representative example of such Hamiltonians: the spatially varying form of the Hamiltonian due to the Landau gauge [23,24]. The Hamiltonians for a broad class of topological phenomena, including the quantum Hall effect [25] or quantum spin Hall effect [26], and the momentum-dependent Hamiltonians [27] are also described by the state-vector-type parameter \mathbf{R} .

Using this set of parameters \mathbf{R} , the time-dependent evolution of the Hamiltonian can then be expressed as $H = H(\mathbf{R}(t))$ through the time-dependent parameters $\mathbf{R}(t) = (R_1(t), R_2(t), \dots)$. For adiabaticity, we consider non-degenerate eigenstates $|u_n(\mathbf{R}(t))\rangle$ of the Hamiltonian $H(t)$. Also, for simplicity, we assume the normalized eigenstate in time $\langle u_n(t) | u_n(t) \rangle = 1$. When a state $|\psi(\mathbf{R}(t))\rangle$ is initially in the n^{th} eigenstate $|u_n(\mathbf{R}(t))\rangle$ satisfying $H(\mathbf{R})|u_n(\mathbf{R})\rangle = E_n(\mathbf{R})|u_n(\mathbf{R})\rangle$, the exact time-varying solution of $|\psi(\mathbf{R})\rangle$ becomes [23]

$$|\psi(\mathbf{R})\rangle = e^{i\theta_n(t)} e^{i\gamma_n(t)} |u_n(\mathbf{R})\rangle + \varepsilon \sum_{m \neq n} c_m(t) |u_m(\mathbf{R})\rangle, \quad (1)$$

where $\theta_n(t) = -(1/\hbar) \int_0^t E_n(\mathbf{R}(t')) dt'$ is the dynamical phase, $\gamma_n(t)$ is the Berry phase, ε denotes the deviation from the adiabatic limit, and c_m is the transition coefficient for the m^{th} eigenstate. With the time-dependent Schrödinger equation $i\hbar \partial_t |\psi(\mathbf{R})\rangle = H(\mathbf{R}) |\psi(\mathbf{R})\rangle$, it follows that

$$\left(\frac{\partial}{\partial t} + i \frac{\partial \gamma_n(t)}{\partial t} \right) |u_n(\mathbf{R})\rangle = -e^{-i(\theta_n + \gamma_n)} \varepsilon \sum_{m \neq n} \left[\left(\frac{i}{\hbar} c_m E_m + \frac{\partial c_m}{\partial t} \right) |u_m(\mathbf{R})\rangle + c_m \frac{\partial}{\partial t} |u_m(\mathbf{R})\rangle \right]. \quad (2)$$

The second-order term of $c_m \partial_t |u_m\rangle$ can be ignored with the adiabatic condition [23]. In the derivation of the Berry phase, the multiplication by the initial bra eigenstate $\langle u_n(\mathbf{R}) |$ in Eq. (2), which corresponds to the identical pre-selection and post-selection in the weak value definition [16-18,28], leads to $\partial_t \gamma_n(t) = i \langle u_n | \partial_t | u_n \rangle$. Note that the time-varying state $|\psi(\mathbf{R})\rangle$ in Eq. (1) does not have to be normalized because all of the coefficients c_m ($m \neq n$) vanishes when the bra eigenstate $\langle u_n(\mathbf{R}) |$ is applied to Eq. (2). Applying the chain rule $\partial_t = (\partial \mathbf{R} / \partial t) \nabla_{\mathbf{R}}$ and integrating the result $\partial_t \gamma_n(t) = i \langle u_n | \partial_t | u_n \rangle$ with respect to time then provides the Berry phase of $\gamma_n(\mathbf{C}) = \oint \mathbf{A}_n d\mathbf{R}$, which is gauge-invariant modulo 2π for the cyclic integral along the closed loop \mathbf{C} , where $\mathbf{A}_n = i \langle u_n | \nabla_{\mathbf{R}} | u_n \rangle$ is the Berry connection [7,8,23].

To interpret the Berry connection \mathbf{A}_n as the expectation value of the transient state $|u_n\rangle$, we develop the notion of the canonical momentum operator in the general “parameter space \mathbf{R} ” from the abstract definition of \mathbf{R} . For the wavefunction $|\psi(\mathbf{R})\rangle$ defined in the N -dimensional parameter space $\mathbf{R} = (R_1, R_2, \dots, R_N)$, we introduce the \mathbf{R} -translation operator $\mathbf{T}_R(\Delta\mathbf{R})$, which leads to the transformation of a wavefunction for the changes in parameters $\Delta\mathbf{R} = (\Delta R_1, \Delta R_2, \dots, \Delta R_N)$, as $\mathbf{T}_R(\Delta\mathbf{R})|\psi(\mathbf{R})\rangle = |\psi(\mathbf{R} - \Delta\mathbf{R})\rangle$, such as spatial translation $\mathbf{T}_x(\Delta\mathbf{x})|\psi(\mathbf{x})\rangle = |\psi(\mathbf{x} - \Delta\mathbf{x})\rangle$ for $\mathbf{R} = \mathbf{x}$, time translation $\mathbf{T}_t(\Delta t)|\psi(t)\rangle = |\psi(t - \Delta t)\rangle$ for scalar $R = t$, and momentum translation $\mathbf{T}_k(\Delta\mathbf{k})|\psi(\mathbf{k})\rangle = |\psi(\mathbf{k} - \Delta\mathbf{k})\rangle$. Following the definition of the canonical momentum operator as the “generator of spatial translation”, we define the canonical momentum operator in the \mathbf{R} -space \mathbf{p}_R as the “generator of parameter translation”, where p_R^m is the momentum operator for the “ m^{th} parameter” R_m :

$$p_R^m = i\hbar \lim_{\Delta R_m \rightarrow 0} \frac{\mathbf{T}_R(\Delta R_m) - \mathbf{I}}{\Delta R_m}, \quad (3)$$

with the identity operator \mathbf{I} . The application of the m^{th} momentum operator to the wavefunction then gives $p_R^m|\psi(\mathbf{R})\rangle = -i\hbar(\partial/\partial R_m)|\psi(\mathbf{R})\rangle$, and thus, the canonical momentum operator in the \mathbf{R} -space is defined as $\mathbf{p}_R = -i\hbar\nabla_{\mathbf{R}}$. Because the unitary \mathbf{R} -translation operator $\mathbf{T}_R(\Delta\mathbf{R})$ guarantees the Hermitian operator \mathbf{p}_R (see Appendix A for the proof), the application of the weak value theory based on the Hermitian operator [16-18,28] is thus allowed for the unitary translation of the wavefunction in the \mathbf{R} -space, covering a wide range of fundamental parameter translations, such as the spatial translation ($\mathbf{R} = \mathbf{x}$) for the real-space momentum and the time translation ($\mathbf{R} = t$) for the Hamiltonian.

For the weak-value formulation using the Hermitian operator \mathbf{p}_R , we introduce a new degree of freedom: a *generalized* and *arbitrary* post-selection with $\langle\varphi(\mathbf{R})|$ to the Berry connection, instead of the conventional post-selection with $\langle u_n|$. The time derivative of the Berry phase $\partial_t\gamma_n(t)$ can be re-expressed as $\partial_t\gamma_n = i\langle\varphi|u_n\rangle\langle u_n|\partial_t|u_n\rangle/\langle\varphi|u_n\rangle$ with the condition of $\langle\varphi|u_n\rangle \neq 0$, which derives the following form from the identity operator $\mathbf{I} = \sum|u_m\rangle\langle u_m|$, as (see Appendix B for the direct application of the post-selection to the state of the time-dependent Schrödinger equation):

$$\frac{\partial\gamma_n(t)}{\partial t} = \frac{i\langle\varphi|\frac{\partial}{\partial t}|u_n\rangle - i\sum_{m \neq n} \langle\varphi|u_m\rangle\langle u_m|\frac{\partial}{\partial t}|u_n\rangle}{\langle\varphi|u_n\rangle}. \quad (4)$$

By applying the chain rule of $\partial_t = (\partial\mathbf{R}/\partial t)\nabla_{\mathbf{R}}$ to Eq. (4) and integrating the result with respect to time, we arrive at the “weak value picture” of the Berry connection \mathbf{A}_n for $\gamma_n(C) = \oint \mathbf{A}_n d\mathbf{R}$, as the sum of two weak-value vector potentials $\mathbf{A}_n = \mathbf{A}_n^S + \mathbf{A}_n^{\text{MP}}$, which is the first main result of this paper:

$$\mathbf{A}_n^S = i \frac{\langle\varphi|\nabla_{\mathbf{R}}|u_n\rangle}{\langle\varphi|u_n\rangle}, \quad (5)$$

$$\mathbf{A}_n^{\text{MP}} = -i \sum_{m \neq n} \frac{\langle\varphi|u_m\rangle\langle u_m|\nabla_{\mathbf{R}}|u_n\rangle}{\langle\varphi|u_n\rangle} = \sum_{m \neq n} \mathbf{A}_n^{\text{MP}-m}, \quad (6)$$

where each weak value representation is obtained with the operator $\nabla_{\mathbf{R}}$ and $|u_m\rangle\langle u_m|\nabla_{\mathbf{R}}$. Although \mathbf{A}_n^S , \mathbf{A}_n^{MP} , and $\partial_t\gamma_n$ in Eqs. (4-6) are defined for every time t , we focus on the closed loops in the parameter space to examine the Berry phase, which is gauge-dependent when the loop is not closed. Using the relationship $\mathbf{A}_n = \mathbf{A}_n^S + \mathbf{A}_n^{\text{MP}}$, the Berry phase defined along the closed loop C can be separated into two parts in the weak value expansion: $\gamma_n(C) = \oint \mathbf{A}_n^S \cdot d\mathbf{R} + \oint \mathbf{A}_n^{\text{MP}} \cdot d\mathbf{R}$. We emphasize that this decomposition itself is not arbitrary but determined by the pre-defined post-selection state $\langle\varphi(\mathbf{R})|$, while the state $\langle\varphi(\mathbf{R})|$ can be arbitrary selected. For example, the post-selection $\langle\varphi| = \langle u_n|$ vanishes \mathbf{A}_n^{MP} , leading to $\mathbf{A}_n = \mathbf{A}_n^S$. The quantity \mathbf{A}_n^{MP} is thus the inherent result of the weak value picture, originating from $\langle\varphi| \neq \langle u_n|$.

With the \mathbf{R} -space momentum operator \mathbf{p}_R , the vector potential $\mathbf{A}_n^S = -(1/\hbar)[\langle\varphi|\mathbf{p}_R|u_n\rangle/\langle\varphi|u_n\rangle]$ corresponds to the “momentum weak value” of the initial eigenstate in the \mathbf{R} -space, weakly measuring the canonical momentum of $|u_n\rangle$ with an arbitrarily post-selected state $\langle\varphi(\mathbf{R})|$. On the other hand, with the projection operator to the m^{th} eigenstate $\mathbf{P}_m = |u_m\rangle\langle u_m|$, \mathbf{A}_n^{MP} is the sum of $\mathbf{A}_n^{\text{MP}-m} = (1/\hbar)\langle\varphi|\mathbf{P}_m\mathbf{p}_R|u_n\rangle/\langle\varphi|u_n\rangle$ ($m \neq n$), which corresponds to the “mutually-projected momentum weak values” of the m^{th} eigenstate, weakly measuring the \mathbf{R} -space momentum lying along $|u_m\rangle$ with a postselected state $\langle\varphi(\mathbf{R})|$. The reformulation $\mathbf{A}_n = \mathbf{A}_n^S + \mathbf{A}_n^{\text{MP}} = -(1/\hbar)\langle u_n|\mathbf{p}_R|u_n\rangle$ then shows that the Berry connection \mathbf{A}_n , which is the expectation value of the \mathbf{R} -space momentum operator, can be decomposed into the sum of the self (\mathbf{A}_n^S) and mutually projected (\mathbf{A}_n^{MP}) weak value momenta in the \mathbf{R} -space. Furthermore, the weak value vector potential \mathbf{A}_n^{MP} is an intrinsic quantity originating from the weak value representation because $\mathbf{A}_n^{\text{MP}} \neq \mathbf{0}$ only when pre- and post-selected states are different, as $|\varphi\rangle \neq |u_n\rangle$.

With the condition of $\mathbf{A}_n = \mathbf{A}_n^S + \mathbf{A}_n^{\text{MP}}$, we further investigate the gauge condition of Eqs. (5) and (6) to answer the question, “Does each “partial” weak value have the same gauge condition as that of the original expectation value?” First, it is well known [7,8,23] that \mathbf{A}_n is a gauge-dependent quantity, as shown in the change of $\mathbf{A}_n \rightarrow \mathbf{A}_n - \nabla_{\mathbf{R}}\zeta_n(\mathbf{R})$ with the gauge transformation of $|u_n\rangle \rightarrow \exp[i\zeta_n(\mathbf{R})]|u_n\rangle$. For the weak value form $\mathbf{A}_n = \mathbf{A}_n^S + \mathbf{A}_n^{\text{MP}}$, if we apply the gauge transformation as $|\varphi\rangle \rightarrow \exp[i\zeta_\varphi(\mathbf{R})]|\varphi\rangle$ and $|u_m\rangle \rightarrow \exp[i\zeta_m(\mathbf{R})]|u_m\rangle$, a clear distinction is observed between each weak value vector potential \mathbf{A}_n^S and \mathbf{A}_n^{MP} . \mathbf{A}_n^S reflects the apparent gauge dependency as $\mathbf{A}_n^S \rightarrow \mathbf{A}_n^S - \nabla_{\mathbf{R}}\zeta_n(\mathbf{R})$, which is the same as that in the original Berry connection $\mathbf{A}_n \rightarrow \mathbf{A}_n - \nabla_{\mathbf{R}}\zeta_n(\mathbf{R})$. In contrast, \mathbf{A}_n^{MP} and its elements $\mathbf{A}_n^{\text{MP}-m}$ represent the full gauge invariance without any ambiguity, as $\mathbf{A}_n^{\text{MP}} \rightarrow$

\mathbf{A}_n^{MP} and $\mathbf{A}_n^{\text{MP}-m} \rightarrow \mathbf{A}_n^{\text{MP}-m}$ (see Appendix C for the detailed proof).

This gauge transformation shows that the gauge dependence of the Berry connection \mathbf{A}_n originates exclusively from \mathbf{A}_n^{S} . The gauge invariance *only* for well-defined modulo 2π of the Berry phase $\oint \mathbf{A}_n d\mathbf{R}$ then originates from the gauge condition of the closed integral of \mathbf{A}_n^{S} as $\oint \mathbf{A}_n^{\text{S}} \cdot d\mathbf{R}$. In contrast, \mathbf{A}_n^{MP} , and more generally, *any* combined operation involving elemental weak values $\mathbf{A}_n^{\text{MP}-m}$, constitute locally measurable quantities due to its full gauge invariance. This finding demonstrates that when the nonzero condition of \mathbf{A}_n^{MP} ($|\varphi\rangle \neq |u_n\rangle$) is satisfied, we can extract new measurable (or gauge-invariant) quantities from the originally gauge-dependent one, by exploiting the nonconservation of the gauge condition during the weak-value decomposition.

3. Gauge-dependent weak value for Berry phase

To illustrate the impact of separating gauge-dependent and gauge-invariant parts in the Berry connection \mathbf{A}_n , we restrict our discussion to the post-selection by the ‘‘parameter state’’ $\langle\varphi(\mathbf{R})| = \langle\mathbf{R}|$. For such a particular type of the post-selection, the eigenstate $|u_n(\mathbf{R})\rangle$ (or its linear superposition $|\psi(\mathbf{R})\rangle$) and the parameter state $|\mathbf{R}\rangle$ have to be included in the same inner product (or Hilbert) space, satisfying the relation of $\langle\mathbf{R}|u_n(\mathbf{R})\rangle = u_n(\mathbf{R})$.

The most illustrative example of the parameter state for the post-selection can be found in the Hilbert space of a spinless particle, where the inner product between a position basis $|\mathbf{r}\rangle$ and a wavefunction $|\psi\rangle$ is defined. Because the Hamiltonian H of a spinless particle is described by the position representation $H = H(\mathbf{r})$, we can set the parameter state as $|\mathbf{R}\rangle = |\mathbf{r}\rangle$, and a scalar wavefunction is defined by the inner product $\langle\mathbf{r}|\psi\rangle = \psi(\mathbf{r})$. The time-dependent variation of the parameter state $|\mathbf{R}(t)\rangle = |\mathbf{r}(t)\rangle$ then represents the time-dependent coordinate transformation, which describes the moving frame or moving potential, which will be shown in the example of Fig. 1.

The application of a position basis as the post-selection $\langle\varphi(\mathbf{R})| = \langle\mathbf{R}| = \langle\mathbf{r}|$ has been widely studied in the subfield of weak value theory, such as superoscillation [29-32] that is described well in Berry’s weak value definition of local momentum [33]. Furthermore, as shown in the definition of a wavefunction in reciprocal space $\psi(\mathbf{p}) = \langle\mathbf{p}|\psi\rangle$, the parameter state for a lattice momentum $\langle\mathbf{R}| = \langle\mathbf{p}|$ can also be defined in the momentum representation. To describe this generality, we maintain the notation of $|\mathbf{R}\rangle$ as the parameter state for the post-selection.

Equations (5) and (6) with the post-selection of $\langle\varphi(\mathbf{R})| = \langle\mathbf{R}|$ result in the forms of

$$\mathbf{A}_n^{\text{S}} = i \frac{\nabla_{\mathbf{R}} u_n(\mathbf{R})}{u_n(\mathbf{R})}, \quad (7)$$

$$\mathbf{A}_n^{\text{MP}} = -i \sum_{m \neq n} \frac{u_m(\mathbf{R}) \langle u_m | \nabla_{\mathbf{R}} | u_n \rangle}{u_n(\mathbf{R})}, \quad (8)$$

From Eqs. (7,8), we now demonstrate that each weak value of \mathbf{A}_n^{S} and \mathbf{A}_n^{MP} plays a distinct role in the emergence of quantum geometric quantities: Berry phase and curvature. Although Berry connection and Berry curvature are interrelated through $\mathbf{B}_n = \nabla_{\mathbf{R}} \times \mathbf{A}_n$, it does not mean that the nonzero \mathbf{B}_n should be applied to a particle directly for the emergence of Berry phase obtained from Berry connection as $\gamma_n(C) = \oint \mathbf{A}_n d\mathbf{R}$, as demonstrated in the singular existence of the magnetic field in the Aharonov-Bohm effect [34] and recent work in the momentum space [35]. This condition implies the possibility of controlling each geometric quantity (Berry phase and curvature) independently in a given regime, by separating the source of each geometric quantity in terms of the decomposed parts of the Berry connection \mathbf{A}_n^{S} and \mathbf{A}_n^{MP} .

For this purpose, we revisit a representative example of the Berry phase: the Aharonov-Bohm effect [34] that connects electromagnetic and geometric vector potentials. Consider a solenoid with a uniform magnetic field \mathbf{B}_0 (Fig. 1) and $\mathbf{B} = 0$ outside the solenoid, having the singular spatial distribution. For the evolution of a charged particle, the arbitrary scalar potential $V(\mathbf{r} - \mathbf{R}(t))$ that confines the particle [7,23] is assumed for the coordinate origin $\mathbf{R}(t)$. The Hamiltonian H , including the nonzero *electromagnetic* vector potential \mathbf{A} , is then

$$H = \frac{1}{2m} \left[\frac{\hbar}{i} \nabla_{\mathbf{r}} - q\mathbf{A}(\mathbf{r}) \right]^2 + V(\mathbf{r} - \mathbf{R}) \quad (9)$$

for the eigenvalue equation $H|u_n(\mathbf{r}, \mathbf{R})\rangle = E_n|u_n(\mathbf{r}, \mathbf{R})\rangle$ with $\mathbf{R}(t)$, which is the parameter space of the Berry phase in this example defined by the position of a particle. The post-selection of $\langle\varphi(\mathbf{R})| = \langle\mathbf{R}|$ in this example then corresponds to the weak measurement of a certain quantum state along the time-varying position of the charged particle $\mathbf{R}(t)$.




Figure 1. Illustration of the Aharonov-Bohm effect to describe the gauge invariance of the vector potential \mathbf{A}_n^{MP} . \mathbf{r} is the position vector, and $\mathbf{R}(t)$ is the time-dependent evolution

of the scalar potential center for $V(\mathbf{r} - \mathbf{R}(t))$ along the contour C_R . C_r is the contour for the integral of \mathbf{A}_n^{MP} in real space. The results of the contour integrals along C_R and C_r are shown in Appendix E.

With $\mathbf{B} = 0$, Eq. (9) can be solved by Peierls substitution [23,34] $|u_n(\mathbf{r}, \mathbf{R})\rangle = \exp[ig(\mathbf{r}, \mathbf{R})]|v_n\rangle$, where $g(\mathbf{r}, \mathbf{R}) = (q/\hbar)\int_{\mathbf{R}}^{\mathbf{r}} \mathbf{A}(\mathbf{r}')d\mathbf{r}'$, leading to the time-independent Schrödinger equation $\{-[\hbar^2/(2m)]\nabla_{\mathbf{r}}^2 + V(\mathbf{r} - \mathbf{R})\}|v_n\rangle = E_n|v_n\rangle$. It is worth noting that the eigenstate $|v_n\rangle$ is a function of $(\mathbf{r} - \mathbf{R})$, and its complex phase with respect to \mathbf{R} is identical up to constant. With the post-selection of $\langle\varphi(\mathbf{R})| = \langle\mathbf{R}|$, the gauge-invariant vector potential \mathbf{A}_n^{MP} then becomes (see Appendix D for the detailed derivation)

$$\mathbf{A}_n^{\text{MP}} = -i\frac{\nabla_{\mathbf{R}}v_n(\mathbf{r} - \mathbf{R})}{v_n(\mathbf{r} - \mathbf{R})} = -i\nabla_{\mathbf{R}}(\log v_n(\mathbf{r} - \mathbf{R})), \quad (10)$$

while $\mathbf{A}_n^{\text{S}} = (q/\hbar)\mathbf{A} - \mathbf{A}_n^{\text{MP}}$, confirming the well-known relation between the geometric ($\mathbf{A}_n^{\text{S}} + \mathbf{A}_n^{\text{MP}}$) and electromagnetic (\mathbf{A}) vector potentials.

Because the parameter space $\mathbf{R}(t)$ is the *particle position* for the coordinate origin of $V(\mathbf{r} - \mathbf{R}(t))$ with $\nabla_{\mathbf{R}}|v_n(\mathbf{r} - \mathbf{R})\rangle = -\nabla_{\mathbf{r}}|v_n(\mathbf{r} - \mathbf{R})\rangle$, \mathbf{A}_n^{MP} corresponds to the measurable complex-valued local momentum of a stationary wavefunction $|v_n\rangle$ following Berry's definition [33], as $\mathbf{A}_n^{\text{MP}} = (1/\hbar)\mathbf{p}_{\mathbf{R}}v_n/v_n = -(1/\hbar)\mathbf{p}_{\mathbf{r}}v_n/v_n$, where $\mathbf{p}_{\mathbf{R}} = -i\hbar\nabla_{\mathbf{R}}$ and $\mathbf{p}_{\mathbf{r}} = -i\hbar\nabla_{\mathbf{r}}$. Figure 2 represents examples of complex-valued \mathbf{A}_n^{MP} , which are invariant under gauge transformation. The system is composed of an ideal square potential well defined by $V(\mathbf{r} - \mathbf{R})$ around the confined magnetic field \mathbf{B}_0 (Fig. 2a). For the bound states $|v_n\rangle$ (Fig. 2b,c), the transverse (x -axis) component of \mathbf{A}_n^{MP} becomes purely imaginary (Fig. 2d,e), while it has real solutions for unbound states. Each node of the bound state results in the divergence of \mathbf{A}_n^{MP} , which is the imaginary-valued counterpart of the superoscillation [32]: superamplification or superdecay along the transverse plane. The longitudinal (y -axis) component of \mathbf{A}_n^{MP} could be real or imaginary, dependent on the eigenvalue E_n .

It is also noted that \mathbf{A}_n^{MP} does not affect the accumulation of the Berry phase due to its local nature (see Appendix E); any line integrals of \mathbf{A}_n^{MP} are independent of the speed or trajectory of system evolution (C_R in Fig. 1) or the integral path in real space (C_r in Fig. 1), and thus, the closed loop leads to zero for these integrals independent from the presence of the electromagnetic vector potential \mathbf{A} . Therefore, the Berry phase, which is purely geometric along the closed loop, is solely determined by the unobservable quantity \mathbf{A}_n^{S} , the gauge-dependent part in the Berry connection \mathbf{A}_n , as $\gamma_n(C) = \oint \mathbf{A}_n \cdot d\mathbf{R} = \oint \mathbf{A}_n^{\text{S}} \cdot d\mathbf{R}$.




Figure 2. Examples of the gauge-invariant vector potential \mathbf{A}_n^{MP} . (a) A schematic for the Aharonov-Bohm effect with an ideal square potential well defined by $V(\mathbf{r} - \mathbf{R})$. (b,c) Bound states $|v_n\rangle$ and (d,e) the corresponding x -axis component of \mathbf{A}_n^{MP} which is purely imaginary: (b,d) 0th mode and (c,e) 2nd mode. Blue dashed (or orange solid) lines denote real (or imaginary) parts in (d,e). Black solid lines in (b-e) present the shape of the potential. The x -axis (or y -axis) denotes the transverse (or longitudinal) direction to the evolution of $\mathbf{R}(t)$. The spatial coordinate is normalized as $X = (2m/\hbar)x$. The y -axis component of \mathbf{A}_n^{MP} is determined by the eigenvalue.

4. Gauge-invariant weak value for Berry curvature

In contrast to the dominant role of \mathbf{A}_n^{S} in the Berry phase, the role of \mathbf{A}_n^{MP} from the post-selection by $\langle\mathbf{R}|$ is confirmed with the Berry curvature. For the nonzero Berry curvature $\mathbf{B}_n = \nabla_{\mathbf{R}} \times \mathbf{A}_n$ (e.g., nonzero \mathbf{B} outside the solenoid in Fig. 1), we cannot utilize Peierls substitution $|u_n(\mathbf{r}, \mathbf{R})\rangle = \exp[ig(\mathbf{r}, \mathbf{R})]|v_n\rangle$ for the zero magnetic field because the line integral of $g(\mathbf{r}, \mathbf{R}) = (q/\hbar)\int_{\mathbf{R}}^{\mathbf{r}} \mathbf{A}(\mathbf{r}')d\mathbf{r}'$ depends on the path [23]. Instead, from Eqs. (7) and (8), the weak value representation of the Berry curvature $\mathbf{B}_n = \nabla_{\mathbf{R}} \times \mathbf{A}_n$ is decomposed into two parts as $\mathbf{B}_n = \mathbf{B}_n^{\text{S}} + \mathbf{B}_n^{\text{MP}}$, where $\mathbf{B}_n^{\text{S}} = \nabla_{\mathbf{R}} \times \mathbf{A}_n^{\text{S}}$ and $\mathbf{B}_n^{\text{MP}} = \nabla_{\mathbf{R}} \times \mathbf{A}_n^{\text{MP}}$. From the relation of $\mathbf{B}_n^{\text{S}} = \mathbf{O}$, it is noted that the Berry curvature exclusively originates from the gauge-invariant vector potential as $\mathbf{B}_n = \mathbf{B}_n^{\text{MP}} = \nabla_{\mathbf{R}} \times \mathbf{A}_n^{\text{MP}}$ (see Appendix F for the

condition of the zero curvature of \mathbf{A}_n^S for the general post-selection), where the partial weak value for the m^{th} eigenstate ($m \neq n$) leads to the partial Berry curvature $\mathbf{B}_n^{\text{MP}-m}$:

$$\mathbf{B}_n^{\text{MP}-m} = -i \frac{u_m(\mathbf{R})}{u_n(\mathbf{R})} \times \left[\langle \nabla_{\mathbf{R}} u_m | \times | \nabla_{\mathbf{R}} u_n \rangle - \nabla_{\mathbf{R}} \left(\log \frac{u_m(\mathbf{R})}{u_n(\mathbf{R})} \right) \times \langle u_m | \nabla_{\mathbf{R}} | u_n \rangle \right], \quad (11)$$

and \mathbf{B}_n is the sum of $\mathbf{B}_n^{\text{MP}-m}$. This is also an alternative expression of the Berry curvature [8] originating from the weak value formulation using the post-selection of $\langle \varphi(\mathbf{R}) | = \langle \mathbf{R} |$. We emphasize that the exclusive role of the measurable vector potential \mathbf{A}_n^{MP} in determining the Berry curvature \mathbf{B}_n , and the exclusive role of the unobservable vector potential \mathbf{A}_n^S in determining the Berry phase γ_n do not contradict with the relationship $\mathbf{B}_n = \nabla_{\mathbf{R}} \times \mathbf{A}_n$ or its Stokes integral form. Because the nonzero Berry curvature is not a necessary condition for the Berry phase [35] as shown in an Aharonov-Bohm [34] or molecular Aharnovo-Bohm [36] effects, for example, zero Berry curvature (with $\mathbf{A}_n^{\text{MP}} = \mathbf{O}$) can also provide the Berry phase (with $\mathbf{A}_n^S \neq \mathbf{O}$).

5. Conclusion

In conclusion, we have shown that the definition of the gauge-invariant vector potential \mathbf{A}_n^{MP} can be extracted from the weak value decomposition of the gauge-dependent Berry connection \mathbf{A}_n in terms of the momentum representation in the parameter space. With the post-selection of the parameter space state $\langle \mathbf{R} |$, we demonstrated the local nature of \mathbf{A}_n^{MP} independent of the ‘‘speed’’ or ‘‘trajectory’’ of system evolution in the Aharonov-Bohm effect, which exhibits the exclusive vector potential origin \mathbf{A}_n^S of the Berry phase. We also proved that \mathbf{A}_n^{MP} plays an exclusive role in the emergence of the Berry curvature, as $\mathbf{B}_n = \nabla \times \mathbf{A}_n^{\text{MP}}$.

In the context of finding a measurable quantity from an unobservable one, the suggested decomposition process using the concept of weak values is distinct from closed-loop integrals for geometric phases [7,12,14] or differential operators applied to scalar and vector potentials in electromagnetics [1,2]. The separation of the gauge-dependent and gauge-invariant parts of the Berry connection suggests that this process will enable the decomposition of a general gauge-dependent quantity into partial weak values, which may include gauge-invariant, measurable quantities. Considering the significant roles of the Berry connection, phase and curvature in topological phenomena, a new viewpoint and representation of the vector potential origins of the Berry phase and curvature will provide a new framework for the topological understanding of quantum mechanics [8,10] and optics [9,11] in terms of the independent control of quantum geometric quantities. Due to the emergence of the momentum weak value, which has played a critical role in the de Broglie-

Bohm representation [37,38] and Madelung’s hydrodynamic form of quantum mechanics [39], this result will also stimulate a geometric view on Bohmian mechanics and hydrodynamic representation. Furthermore, applying the weak-value decomposition to an arbitrary quantum-mechanical operator can also be envisaged, not only to reveal a new gauge-invariant quantity but also to subdivide existing quantities with new physical pictures, as shown in \mathbf{A}_n^{MP} and its role in the decomposition of quantum geometric quantities.

Acknowledgements

This work was supported by the National Research Foundation of Korea (NRF) through the Global Frontier Program (GFP, 2014M3A6B3063708) funded by the Korean government. S. Yu was supported by the Basic Science Research Program (2016R1A6A3A04009723), and X. Piao and N. Park were supported by the Korea Research Fellowship Program (KRF, 2016H1D3A1938069) through the NRF, all funded by the Korean government.

Appendix A. Hermitian condition of R-space momentum operator

As the generalization of the relation between the spatial translation operator and canonical momentum operator [24], the \mathbf{R} -translation operator $\mathbf{T}_{\mathbf{R}}(\Delta R_m)$ for the m^{th} parameter R_m can be expressed as

$$\mathbf{T}(\Delta R_m) = 1 - \frac{i}{\hbar} p_{\mathbf{R}}^m \Delta R_m + \dots \quad (S1)$$

Equation (S1) becomes Eq. (3) in the main text, or $\mathbf{T}_{\mathbf{R}}(\Delta R_m) = 1 - (i/\hbar)p_{\mathbf{R}}^m \Delta R_m$, for the infinitesimal translation ΔR_m . This relation leads to $\mathbf{T}_{\mathbf{R}}(\Delta R_m)^\dagger = 1 + (i/\hbar)p_{\mathbf{R}}^{m\dagger} \Delta R_m$, while $\mathbf{T}_{\mathbf{R}}(-\Delta R_m) = \mathbf{T}_{\mathbf{R}}(\Delta R_m)^{-1} = 1 + (i/\hbar)p_{\mathbf{R}}^m \Delta R_m$. Therefore, the Hermitian condition of the \mathbf{R} -space momentum operator $p_{\mathbf{R}}^{m\dagger} = p_{\mathbf{R}}^m$ is achieved with the unitary operator $\mathbf{T}_{\mathbf{R}}(\Delta R_m)$, as $\mathbf{T}_{\mathbf{R}}(\Delta R_m)^\dagger = \mathbf{T}_{\mathbf{R}}(\Delta R_m)^{-1} = \mathbf{T}_{\mathbf{R}}(-\Delta R_m)$. The application of the weak value theory [16-18,28], which requires the Hermitian condition of the operator, is then allowed with the unitary translation of the wavefunction on the parameter space \mathbf{R} .

Appendix B. Direct application of $\langle \varphi(\mathbf{R}) |$ to the time-dependent Schrödinger equation

The result of Eq. (4) in the main text can also be obtained by applying the post-selection with $\langle \varphi(\mathbf{R}) |$ in place of $\langle u_n(\mathbf{R}) |$ to the state of the time-dependent Schrödinger equation. From Eq. (2) in the main text, we obtain

$$\begin{aligned} & \langle \varphi | \frac{\partial}{\partial t} | u_n \rangle + i \frac{\partial \gamma_n}{\partial t} \langle \varphi | u_n \rangle \\ & = -e^{-i(\theta_n + \gamma_n)} \varepsilon \sum_{m \neq n} \left(\frac{i}{\hbar} c_m E_m + \frac{\partial c_m}{\partial t} \right) \langle \varphi | u_m \rangle \end{aligned} \quad (S2)$$

The coefficients on the right side of Eq. (S2) are obtained by multiplying Eq. (2) in the main text by the bra eigenstate $\langle u_p(\mathbf{R})|$ ($p \neq n$) as follows:

$$\langle u_p | \frac{\partial}{\partial t} | u_n \rangle = -e^{-i(\theta_n + \gamma_n)} \mathcal{E} \left(\frac{i}{\hbar} c_p E_p + \frac{\partial c_p}{\partial t} \right). \quad (\text{S3})$$

Using Eq. (S3), we can re-express Eq. (S2) as follows:

$$\begin{aligned} \langle \varphi | \frac{\partial}{\partial t} | u_n \rangle + i \frac{\partial \gamma_n}{\partial t} \langle \varphi | u_n \rangle \\ = \sum_{m \neq n} \langle u_m | \frac{\partial}{\partial t} | u_n \rangle \langle \varphi | u_m \rangle = \sum_{m \neq n} \langle \varphi | u_m \rangle \langle u_m | \frac{\partial}{\partial t} | u_n \rangle \end{aligned} \quad (\text{S4})$$

The time derivative of the Berry phase $\partial_i \gamma_n$ then takes the form of Eq. (4) in the main text with the condition of $\langle \varphi | u_n \rangle \neq 0$.

Appendix C. Gauge conditions of the weak value vector potentials

To investigate the gauge condition of each weak value vector potential, we apply the gauge transformation to the weak value form $\mathbf{A}_n = \mathbf{A}_n^S + \mathbf{A}_n^{\text{MP}}$, as $|\varphi\rangle \rightarrow \exp[i\zeta_\varphi(\mathbf{R})]|\varphi\rangle$, $|u_m\rangle \rightarrow \exp[i\zeta_m(\mathbf{R})]|u_m\rangle$, and $|u_n\rangle \rightarrow \exp[i\zeta_n(\mathbf{R})]|u_n\rangle$. The momentum weak value \mathbf{A}_n^S then becomes

$$\begin{aligned} \mathbf{A}_n^{S\text{-gauge}} &= i \frac{\langle \varphi | e^{-i\zeta_\varphi(\mathbf{R})} \nabla_{\mathbf{R}} e^{i\zeta_n(\mathbf{R})} | u_n \rangle}{\langle \varphi | e^{-i\zeta_\varphi(\mathbf{R})} e^{i\zeta_n(\mathbf{R})} | u_n \rangle} \\ &= i \frac{e^{i[\zeta_n(\mathbf{R}) - \zeta_\varphi(\mathbf{R})]} \langle \varphi | (\nabla_{\mathbf{R}} | u_n \rangle + i | u_n \rangle \nabla_{\mathbf{R}} \zeta_n(\mathbf{R}))}{e^{i[\zeta_n(\mathbf{R}) - \zeta_\varphi(\mathbf{R})]} \langle \varphi | u_n \rangle} \quad (\text{S5}) \\ &= \mathbf{A}_n^S - \nabla_{\mathbf{R}} \zeta_n(\mathbf{R}) \end{aligned}$$

$$\begin{aligned} \mathbf{A}_n^{\text{MP}} &= -i \sum_{m \neq n} \frac{\langle \varphi | e^{ig(\mathbf{r}, \mathbf{R})} | v_m \rangle \langle v_m | e^{-ig(\mathbf{r}, \mathbf{R})} (\nabla_{\mathbf{R}} | v_n \rangle + i | v_n \rangle \nabla_{\mathbf{R}} g(\mathbf{r}, \mathbf{R})) e^{ig(\mathbf{r}, \mathbf{R})}}{\langle \varphi | e^{ig(\mathbf{r}, \mathbf{R})} | v_n \rangle} \\ &= -i \sum_{m \neq n} \frac{\langle \varphi | v_m \rangle \langle v_m | (\nabla_{\mathbf{R}} | v_n \rangle + i | v_n \rangle \nabla_{\mathbf{R}} g(\mathbf{r}, \mathbf{R}))}{\langle \varphi | v_n \rangle} \\ &= -i \sum_{m \neq n} \frac{\langle \varphi | v_m \rangle \langle v_m | \nabla_{\mathbf{R}} | v_n \rangle}{\langle \varphi | v_n \rangle} \\ &= -i \sum_m \frac{\langle \varphi | v_m \rangle \langle v_m | \nabla_{\mathbf{R}} | v_n \rangle}{\langle \varphi | v_n \rangle} + i \langle v_n | \nabla_{\mathbf{R}} | v_n \rangle \\ &= -i \frac{\langle \varphi | \nabla_{\mathbf{R}} | v_n \rangle}{\langle \varphi | v_n \rangle} + i \langle v_n | \nabla_{\mathbf{R}} | v_n \rangle \end{aligned} \quad (\text{S7})$$

Because $\nabla_{\mathbf{R}} | v_n(\mathbf{r} - \mathbf{R}) \rangle = -\nabla_{\mathbf{r}} | v_n(\mathbf{r} - \mathbf{R}) \rangle$, we can change the second term in Eq. (S7) to the expectation value of the momentum $i \langle v_n | \nabla_{\mathbf{R}} | v_n \rangle = -i \langle v_n | \nabla_{\mathbf{r}} | v_n \rangle = \langle v_n | \mathbf{p}_{\mathbf{r}} | v_n \rangle / \hbar$, which becomes zero [23] in the stationary state $|v_n\rangle$. Therefore, by applying the post-selected state $\langle \varphi(\mathbf{R}) | = \langle \mathbf{R} |$, Eq. (S7) becomes

$$\mathbf{A}_n^{\text{MP}} = -i \frac{\langle \mathbf{R} | \nabla_{\mathbf{R}} | v_n \rangle}{\langle \mathbf{R} | v_n \rangle} = -i \frac{\nabla_{\mathbf{R}} v_n(\mathbf{r} - \mathbf{R})}{v_n(\mathbf{r} - \mathbf{R})}, \quad (\text{S8})$$

leading to Eq. (10) in the main text.

which illustrates the identical gauge dependency $-\nabla_{\mathbf{R}} \zeta_n(\mathbf{R})$ to that of the original Berry connection \mathbf{A}_n . In contrast, the projected momentum weak value of the m^{th} eigenstate $\mathbf{A}_n^{\text{MP}-m}$ becomes

$$\begin{aligned} \mathbf{A}_n^{\text{MP}-m\text{-gauge}} &= -i \frac{\langle \varphi | e^{-i\zeta_\varphi(\mathbf{R})} e^{i\zeta_m(\mathbf{R})} | u_m \rangle \langle u_m | e^{-i\zeta_m(\mathbf{R})} \nabla_{\mathbf{R}} e^{i\zeta_n(\mathbf{R})} | u_n \rangle}{\langle \varphi | e^{-i\zeta_\varphi(\mathbf{R})} e^{i\zeta_n(\mathbf{R})} | u_n \rangle} \\ &= -i \frac{e^{i[\zeta_n(\mathbf{R}) - \zeta_\varphi(\mathbf{R})]} \langle \varphi | u_m \rangle \langle u_m | (\nabla_{\mathbf{R}} | u_n \rangle + i | u_n \rangle \nabla_{\mathbf{R}} \zeta_n(\mathbf{R}))}{e^{i[\zeta_n(\mathbf{R}) - \zeta_\varphi(\mathbf{R})]} \langle \varphi | u_n \rangle} \\ &= -i \frac{\langle \varphi | u_m \rangle \langle u_m | \nabla_{\mathbf{R}} | u_n \rangle + i \langle u_m | u_n \rangle \nabla_{\mathbf{R}} \zeta_n(\mathbf{R})}{\langle \varphi | u_n \rangle} = \mathbf{A}_n^{\text{MP}-m} \end{aligned} \quad (\text{S6})$$

Because the $\mathbf{A}_n^{\text{MP}-m}$ values for all $m \neq n$ are gauge-invariant, their sum \mathbf{A}_n^{MP} is also gauge-invariant.

Appendix D. Derivation of \mathbf{A}_n^{MP} for the Aharonov-Bohm effect

We start from $|u_{n,m}(\mathbf{r}, \mathbf{R})\rangle = \exp[ig(\mathbf{r}, \mathbf{R})]|v_{n,m}\rangle$, where $g(\mathbf{r}, \mathbf{R}) = (q/\hbar) \int_{\mathbf{R}}^{\mathbf{r}} \mathbf{A}(\mathbf{r}') d\mathbf{r}'$ is defined by the electromagnetic vector potential \mathbf{A} and $\nabla_{\mathbf{R}} g(\mathbf{r}, \mathbf{R}) = -(q/\hbar) \mathbf{A}(\mathbf{R})$. $\mathbf{A}_n^{\text{MP}-m}$ then becomes

Appendix E. Contour integral of \mathbf{A}_n^{MP}

To examine the effects of \mathbf{A}_n^{MP} for $\gamma_n^{\text{MP}} = \int \mathbf{A}_n^{\text{MP}} d\mathbf{R}$, we define two gauge-invariant phase evolutions $\gamma_n^{\text{MP}-\mathbf{r}}$ and $\gamma_n^{\text{MP}-\mathbf{R}}$ driven by \mathbf{A}_n^{MP} , for arbitrary (including non-cyclic) contours, as follows:

$$\begin{aligned} \gamma_n^{\text{MP}-\mathbf{r}}(\mathbf{r}) \Big|_{C_{\mathbf{r}}} &= -i \int_{C_{\mathbf{r}}} \nabla_{\mathbf{R}} (\log v_n(\mathbf{r} - \mathbf{R})) d\mathbf{R} \\ &= -i \log \frac{v_n(\mathbf{r} - \mathbf{R}_{\mathbf{f}})}{v_n(\mathbf{r} - \mathbf{R}_{\mathbf{s}})} \end{aligned} \quad (\text{S9})$$

$$\begin{aligned}\gamma_n^{\text{MP-R}}(\mathbf{R})\Big|_{C_r} &= i \int_{C_r} \nabla_{\mathbf{r}} (\log v_n(\mathbf{r}-\mathbf{R})) d\mathbf{r} \\ &= -i \log \frac{v_n(\mathbf{r}_s-\mathbf{R})}{v_n(\mathbf{r}_f-\mathbf{R})}\end{aligned}\quad (\text{S10})$$

These two complex-valued phase functions represent distinct observation conditions for \mathbf{A}_n^{MP} in terms of the measurement positions (\mathbf{r} or \mathbf{R}) and the evolutions (C_R or C_r): $\gamma_n^{\text{MP-r}}(\mathbf{r})$ for the phase accumulation at *real-space* position \mathbf{r} with the *system* evolution along C_R , and $\gamma_n^{\text{MP-R}}(\mathbf{R})$ for the phase accumulation measured along the *real-space* contour C_r with the *system* position \mathbf{R} . By virtue of the gauge invariance of \mathbf{A}_n^{MP} , both phase accumulations are automatically gauge-invariant regardless of whether the contours defined for the potential (C_R) or position (C_r) are cyclic. Furthermore, because the line integral is defined only with the initial and final state of $v_n(\mathbf{r}-\mathbf{R})$, both $\gamma_n^{\text{MP-r}}$ and $\gamma_n^{\text{MP-R}}$ are independent of the speed or trajectory of system evolution. \mathbf{A}_n^{MP} therefore does not contribute to the accumulation of the Berry phase, because $\gamma_n^{\text{MP-r}}(\mathbf{r}) = \gamma_n^{\text{MP-R}}(\mathbf{R}) = 0$ for the closed loops C_R and C_r . We also note that the phase induced by the gauge-invariant \mathbf{A}_n^{MP} in a non-cyclic evolution process is apparently distinct from the case of a Pancharatnam phase [14], which is dependent on the trajectory composing the closed curve with the geodesic curve in the ray space.

Appendix F. Zero curvature condition of \mathbf{A}_n^S

To separate the source of the Berry curvature in terms of weak-value vector potentials, it is necessary to assign $\mathbf{B}_n^S = \nabla_{\mathbf{R}} \times \mathbf{A}_n^S = \mathbf{O}$ by introducing suitable post-selections. From Eq. (5) in the main text, the general expression of \mathbf{B}_n^S is

$$\begin{aligned}\mathbf{B}_n^S &= \nabla_{\mathbf{R}} \times \mathbf{A}_n^S \\ &= i \frac{\langle \nabla_{\mathbf{R}} \phi | \times | \nabla_{\mathbf{R}} u_n \rangle - \langle \nabla_{\mathbf{R}} \phi | u_n \rangle \times \langle \phi | \nabla_{\mathbf{R}} u_n \rangle}{\langle \phi | u_n \rangle},\end{aligned}\quad (\text{S11})$$

which results in the zero curvature condition of $\langle \nabla_{\mathbf{R}} \phi | \times | \nabla_{\mathbf{R}} u_n \rangle = \langle \nabla_{\mathbf{R}} \phi | u_n \rangle \times \langle \phi | \nabla_{\mathbf{R}} u_n \rangle$. It is noted that the post-selection by the parameter state $\langle \phi(\mathbf{R}) | = \langle \mathbf{R} |$ satisfies this condition.

References

- [1] J. D. Jackson and L. B. Okun, Historical roots of gauge invariance, *Rev. Mod. Phys.* **73**, 663 (2001).
- [2] H. Weyl, Elektron und gravitation. I, *Z. Phys.* **56**, 330 (1929).
- [3] L. O’Raifeartaigh and N. Straumann, Gauge theory: Historical origins and some modern developments, *Rev. Mod. Phys.* **72**, 1 (2000).
- [4] P. A. M. Dirac, *Lectures on quantum mechanics* (Courier Corporation, 2001), Vol. 2.
- [5] J. D. Jackson, *Classical electrodynamics* (John Wiley & Sons, 2012).
- [6] X. Ji, Gauge-invariant decomposition of nucleon spin, *Phys. Rev. Lett.* **78**, 610 (1997).
- [7] M. V. Berry, Quantal phase factors accompanying adiabatic changes, *Proc. R. Soc. Lond. A* **392**, 45 (1984).
- [8] D. Xiao, M.-C. Chang, and Q. Niu, Berry phase effects on electronic properties, *Rev. Mod. Phys.* **82**, 1959 (2010).
- [9] S. Slussarenko, A. Alberucci, C. P. Jisha, B. Piccirillo, E. Santamato, G. Assanto, and L. Marrucci, Guiding light via geometric phases, *Nat. Photon.* **10**, 571 (2016).
- [10] M. Z. Hasan and C. L. Kane, Colloquium: topological insulators, *Rev. Mod. Phys.* **82**, 3045 (2010).
- [11] L. Lu, J. D. Joannopoulos, and M. Soljačić, Topological photonics, *Nat. Photon.* **8**, 821 (2014).
- [12] Y. Aharonov and J. Anandan, Phase change during a cyclic quantum evolution, *Phys. Rev. Lett.* **58**, 1593 (1987).
- [13] S. Pancharatnam, Generalized theory of interference and its applications. Part. I. Coherent pencils, *Proc. Indiana Acad. Sci.* **44**, 398 (1956).
- [14] J. Samuel and R. Bhandari, General setting for Berry’s phase, *Phys. Rev. Lett.* **60**, 2339 (1988).
- [15] H. M. Wong, K. M. Cheng, and M.-C. Chu, Quantum geometric phase between orthogonal states, *Phys. Rev. Lett.* **94**, 070406 (2005).
- [16] Y. Aharonov, D. Z. Albert, and L. Vaidman, How the result of a measurement of a component of the spin of a spin-1/2 particle can turn out to be 100, *Phys. Rev. Lett.* **60**, 1351 (1988).
- [17] J. Dressel, M. Malik, F. M. Miatto, A. N. Jordan, and R. W. Boyd, Colloquium: Understanding quantum weak values: Basics and applications, *Rev. Mod. Phys.* **86**, 307 (2014).
- [18] B. Tamir and E. Cohen, Introduction to weak measurements and weak values, *Quanta* **2**, 7 (2013).
- [19] E. Sjöqvist, Geometric phase in weak measurements, *Phys. Lett. A* **359**, 187 (2006).
- [20] S. Tamate, H. Kobayashi, T. Nakanishi, K. Sugiyama, and M. Kitano, Geometrical aspects of weak measurements and quantum erasers, *New J. Phys.* **11**, 093025 (2009).
- [21] H. Kobayashi, S. Tamate, T. Nakanishi, K. Sugiyama, and M. Kitano, Direct observation of geometric phases using a three-pinhole interferometer, *Phys. Rev. A* **81**, 012104 (2010).
- [22] K. Y. Bliokh, F. Rodríguez-Fortuño, F. Nori, and A. V. Zayats, Spin-orbit interactions of light, *Nat. Photon.* **9**, 796 (2015).
- [23] D. J. Griffiths, *Introduction to quantum mechanics* (Prentice-Hall, Englewood Cliffs, NJ, 1995).
- [24] J. J. Sakurai, S.-F. Tuan, and E. D. Commins, *Modern quantum mechanics* (AAPT, 1995).
- [25] M. E. Cage, K. Klitzing, A. Chang, F. Duncan, M. Haldane, R. Laughlin, A. Pruisken, and D. Thouless, *The quantum Hall effect* (Springer Science & Business Media, 2012).
- [26] C. L. Kane and E. J. Mele, Quantum spin Hall effect in graphene, *Phys. Rev. Lett.* **95**, 226801 (2005).
- [27] Z.-Q. Feng, Momentum dependence of the symmetry potential and its influence on nuclear reactions, *Physical Review C* **84**, 024610 (2011).

- [28] B. E. Svensson, Pedagogical review of quantum measurement theory with an emphasis on weak measurements, *Quanta* **2**, 18 (2013).
- [29] Y. Aharonov, J. Anandan, S. Popescu, and L. Vaidman, Superpositions of time evolutions of a quantum system and a quantum time-translation machine, *Phys. Rev. Lett.* **64**, 2965 (1990).
- [30] M. V. Berry, Evanescent and real waves in quantum billiards and Gaussian beams, *J. Phys. A* **27**, L391 (1994).
- [31] A. Kempf, Black holes, bandwidths and Beethoven, *J. Math. Phys.* **41**, 2360 (2000).
- [32] M. Berry and S. Popescu, Evolution of quantum superoscillations and optical superresolution without evanescent waves, *J. Phys. A* **39**, 6965 (2006).
- [33] M. Berry, Five momenta, *Eur. J. Phys.* **34**, 1337 (2013).
- [34] Y. Aharonov and D. Bohm, Significance of electromagnetic potentials in the quantum theory, *Phys. Rev.* **115**, 485 (1959).
- [35] F. Liu and K. Wakabayashi, Novel topological phase with a zero berry curvature, *Phys. Rev. Lett.* **118**, 076803 (2017).
- [36] C. A. Mead, The molecular Aharonov—Bohm effect in bound states, *Chem. Phys.* **49**, 23 (1980).
- [37] P. R. Holland, *The quantum theory of motion: an account of the de Broglie-Bohm causal interpretation of quantum mechanics* (Cambridge university press, 1995).
- [38] S. Yu, X. Piao, and N. Park, Bohmian Photonics for Independent Control of the Phase and Amplitude of Waves, *Phys. Rev. Lett.* **120**, 193902 (2018).
- [39] E. Madelung, Quantentheorie in hydrodynamischer Form, *Z. Phys.* **40**, 322 (1927).