

From Lie algebra crossed modules to tensor hierarchies

Sylvain Lavau^{*a,b} and Jim Stasheff^c

^a*Euler Institute (EIMI) & Steklov Institute, Fontanka River Embankment 27, 191023, St-Petersburg, Russia.*

^b*Aristotle University of Thessaloniki, Department of Mathematics, 541 24, Thessaloniki, Greece.*

^c*109 Holly Dr, Lansdale, PA 19446, USA.*

Abstract

The present paper, though inspired by the use of tensor hierarchies in theoretical physics, establishes their mathematical credentials, especially as genetically related to Lie algebra crossed modules. Gauging procedures in supergravity rely on a pairing – the *embedding tensor* – between a Leibniz algebra and a Lie algebra. Two such algebras, together with their embedding tensor, form a triple called a *Lie-Leibniz triple*, of which Lie algebra crossed modules are particular cases. This paper is devoted to showing that any Lie-Leibniz triple induces a differential graded Lie algebra – its associated *tensor hierarchy* – whose restriction to the category of Lie algebra crossed modules is the canonical assignment associating to any Lie algebra crossed module its corresponding unique 2-term differential graded Lie algebra. This shows that Lie-Leibniz triples form natural generalizations of Lie algebra crossed modules and that their associated tensor hierarchies can be considered as some kind of ‘lie-ization’ of the former. We deem the present construction of such tensor hierarchies clearer and more straightforward than previous derivations. We stress that such a construction suggests the existence of further well-defined Leibniz gauge theories.

Keywords: Differential graded Lie (super)-algebra, Leibniz algebra, Lie algebra crossed module.

2020 Mathematics Subject Classification: 17B70, 17A32, 18G45.

Contents

| | |
|--|-----------|
| Introduction | 2 |
| 2 Lie-Leibniz triples and Lie algebra crossed modules | 4 |
| 2.1 Leibniz algebras and Lie-Leibniz triples | 4 |
| 2.2 A relationship between Lie-Leibniz triples and Lie algebra crossed modules | 10 |
| 3 The tensor hierarchy associated to a Lie-Leibniz triple | 16 |
| 3.1 Associating a graded Lie algebra to a Lie-Leibniz triple | 16 |
| 3.2 Existence and uniqueness of the differential | 23 |
| 3.3 The differential graded Lie algebra structure on \mathbb{T} | 30 |
| 3.4 Proof of Theorem 1 and open questions | 39 |
| A Corrigendum (in collaboration with Jakob Palmkvist) | 45 |

*Corresponding author: lavau@math.univ-lyon1.fr

1 Introduction

Gauged models of supergravity involve a Lie algebra of symmetries \mathfrak{g} , of which only a sub-algebra \mathfrak{h} is promoted to become a gauge algebra. To preserve the symmetries manifest when performing their computations, physicists stick to a formulation of the theory that involves the bigger Lie algebra \mathfrak{g} . One of the consequences is that the gauge theory needs the addition of several fields of higher form degree—and their associated field strengths, leading to what physicists call a *tensor hierarchy* [16, 17, 18, 43]. Mathematically, it is a (possibly infinite) tower of \mathfrak{g} -modules – that by convention physicists take to be positively graded but that, in the present paper, we take to be negatively graded – such that the space at degree $k > 1$ depends exclusively on the spaces of lower degree. These spaces contain the redundant information carried by the fields of higher degree; they are necessary to preserve the covariance of the theory. The need for a unified perspective on gauging procedures in supergravity, as well as in Double and Exceptional field theories, has been salient in theoretical physics for some years now [4, 8, 12, 23].

On the mathematical side, another approach to tensor hierarchies has been recently explored: that of *Leibniz algebras* [3, 30, 31]. A Leibniz algebra is a vector space V equipped with a bilinear product \circ which is a derivation of itself, i.e. it satisfies the Leibniz rule, hence the denomination [2, 35]:

$$x \circ (y \circ z) = (x \circ y) \circ z + y \circ (x \circ z) \quad (1)$$

The construction of tensor hierarchies relies on the existence of *embedding tensors*. If \mathfrak{g} is a Lie algebra and if a Leibniz algebra V is a \mathfrak{g} -module, an embedding tensor is a map $\Theta : V \rightarrow \mathfrak{g}$ satisfying some consistency conditions – called the linear and quadratic constraints, respectively. The quadratic constraint implies that $\mathfrak{h} := \text{Im}(\Theta)$ is a sub-Lie algebra of \mathfrak{g} , whereas what we call the linear constraint requires that the Leibniz product on V is generated by the action of \mathfrak{h} on V :

$$x \circ y = \Theta(x) \cdot y \quad (2)$$

In other words, the embedding tensor lifts the adjoint action to \mathfrak{g} :

$$\begin{array}{ccc} & & \mathfrak{g} \\ & \nearrow \Theta & \downarrow \rho \\ V & \xrightarrow{\text{ad}} & \text{End}(V) \end{array}$$

where ρ represents the action of \mathfrak{g} on V . We call such compatible data $(\mathfrak{g}, V, \Theta)$ *Lie-Leibniz triples*. Equation (2) is an important aspect of these algebraic structures, because it draws a parallel with the notion of *Lie algebra crossed module*. The latter, also called *differential crossed module* [1], appear to be precisely those Lie-Leibniz triples such that V is a Lie algebra and Θ is \mathfrak{g} -equivariant. The introduction of the notion of Lie algebra crossed module was historically preceded by the study of crossed modules of discrete groups, see [24] for an historical account.

It has been proved that any Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ gives rise canonically to a non-positively graded differential graded Lie algebra [31], which might coincide with the one defined from Borchers algebras in [38]. Unfortunately, the construction presented in [31] is not very satisfying, mathematically speaking, because the role of \mathfrak{g} – although central in the Lie-Leibniz triple – is not salient in the resulting differential graded Lie algebra. In the present paper we provide an alternative, simpler construction, that generalizes the results in [31] and that additionally reinstates the prominent role of \mathfrak{g} . The result is a differential graded Lie algebra $\mathbb{T} = \bigoplus_{i=-1}^{\infty} T_{-i}$ concentrated in degrees lesser than or equal to $+1$:

$$\dots \longrightarrow T_{-3} \xrightarrow{\partial_{-2}} T_{-2} \xrightarrow{\partial_{-1}} V[1] \xrightarrow{\partial_0=\Theta} \mathfrak{g} \xrightarrow{\partial_{+1}} R_{\Theta}[-1]$$

where $R_\Theta[-1]$ is the cyclic \mathfrak{g} -submodule of $\text{Hom}(V, \mathfrak{g})$ generated by Θ , the degree having been shifted by $+1$. We call this differential graded Lie algebra \mathbb{T} the *tensor hierarchy associated to* $(\mathfrak{g}, V, \Theta)$. The negatively graded part $T_\bullet = \bigoplus_{i=1}^\infty T_{-i}$ is obtained as a quotient of the free graded Lie algebra of $V[1]$ by a particular graded ideal, as was proposed by earlier work in theoretical physics [10, 20]. A priori, it is not an exact sequence, although we raise the question that under some circumstances it might be (see Conjecture 1).

Although it is injective on objects, the function that assigns to each Lie-Leibniz triple its associated tensor hierarchy is not functorial on the category **Lie-Leib** of Lie-Leibniz triples. However, we note that its restriction to the full subcategory **Lie** \times **Mod** of Lie algebra crossed modules has a very particular image. More precisely, any Lie algebra crossed module $\mathfrak{c} \xrightarrow{\Theta} \mathfrak{g}$ is sent by this function to its corresponding canonical 2-term differential graded Lie algebra [1] or, more precisely, to the following 3-term differential graded Lie algebra:

$$\mathfrak{c} \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{0} \mathbb{R}[-1] \quad (3)$$

Here, $\mathbb{R}[-1]$ is the irreducible trivial representation of \mathfrak{g} , to which the \mathfrak{g} -submodule $R_\Theta \subset \text{Hom}(\mathfrak{c}, \mathfrak{g})$ is isomorphic, since Θ is \mathfrak{g} -equivariant by definition of Lie algebra crossed modules. We observe that this assignment defines a faithful functor on **Lie** \times **Mod**. It is thus reasonable to look for the biggest subcategory of **Lie-Leib** on which the restriction of the function assigning to each Lie-Leibniz triple its associated tensor hierarchy is functorial.

By construction, the tensor hierarchy associated to a Lie-Leibniz triple encodes the symmetric part of the Leibniz product:

$$\{x, y\} = \frac{1}{2}(x \circ y + y \circ x) \quad (4)$$

This bracket vanishes when V is a Lie algebra, so in particular when $(\mathfrak{g}, V, \Theta)$ is a Lie algebra crossed module, resulting in the cochain complex (3). The prominence of the symmetric bracket (4) and the \mathfrak{g} -action in the construction of the tensor hierarchy associated to a given Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ can be seen more clearly in the fact that the first space of the hierarchy, namely T_{-2} , is the quotient of $S^2(V)$ by the biggest \mathfrak{g} -submodule of $\text{Ker}(\{.,.\}) \subset S^2(V)$, denoted K . We say that a morphism of Lie-Leibniz triples $(\varphi, \chi) : (\mathfrak{g}, V, \Theta) \rightarrow (\mathfrak{g}', V', \Theta')$ is *compatible* if it sends $K \subset \text{Ker}(\{.,.\}) \subset S^2(V)$ to $K' \subset \text{Ker}(\{.,.\}') \subset S^2(V')$. Then, Lie-Leibniz triples together with compatible morphisms form a wide subcategory **compLie-Leib** of the category **Lie-Leib**, and **Lie** \times **Mod** is a full subcategory of both: **Lie** \times **Mod** \subset **compLie-Leib** \subset **Lie-Leib**. We now state the main result of this paper¹:

Theorem 1. *The functor*

$$\begin{aligned} \overline{G} : \mathbf{Lie} \times \mathbf{Mod} &\longrightarrow \mathbf{DGLie}_{\leq 1} \\ (\mathfrak{c} \xrightarrow{\Theta} \mathfrak{g}) &\longmapsto (\mathfrak{c} \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{0} \mathbb{R}[-1]) \end{aligned}$$

*can be canonically extended to an injective-on-objects function $G : \mathbf{Lie-Leib} \rightarrow \mathbf{DGLie}_{\leq 1}$ such that the restriction of this function to the wide sub-category **compLie-Leib** is a faithful functor. Moreover, **compLie-Leib** is the biggest wide subcategory of **Lie-Leib** (with respect to inclusion) such that this functorial property holds.*

This theorem shows that, while Lie-Leibniz triples form a natural generalization of Lie algebra crossed modules, tensor hierarchies materialize as a generalization of the differential graded Lie algebra formulation of such Lie algebra crossed modules, as given in the cochain complex (3). The paper is divided into two sections: Section 2 presents the main mathematical concepts and ideas, while Section 3 is mostly technical

¹After several conversations with Jakob Palmkvist at MITP (Mainz) in January 2022, it appeared that Theorem 1 was erroneous, as $\mathbf{DGLie}_{\leq 1}$ should be \mathbf{DGLie} . We corrected this mistake and provided more details about the reasons why this is so in a Corrigendum [32], which can be found as Appendix A of this preprint.

as it contains the details of the proof of Theorem 1. To begin with, Lie-Leibniz triples are introduced and studied in subsection 2.1. Then, the relationship between Lie algebra crossed modules and Lie-Leibniz triples is established in subsection 2.2, which concludes with an explanation of Theorem 1. Section 3 is subdivided into four parts: building the graded vector space T_\bullet and equipping it with a graded Lie bracket in subsection 3.1, showing that it moreover admits a non-trivial differential in subsection 3.2, proving that the differential and the graded Lie bracket are compatible in subsection 3.3, and finally proving Theorem 1 in the final subsection 3.4. Though the construction(s) and proofs are technical, they proceed by a sequence of steps, each of which is meaningful and fairly transparent. In order for the reader to fully appreciate the first steps of the construction of the tensor hierarchy, we have introduced a detailed example at the end of subsection 3.3 (see Example 7). The paper concludes on a discussion of the algebraic structures involved and open questions. Clarifying the relationship between Lie algebra crossed modules and Lie-Leibniz triples opens the possibility to better understand higher gauge theories. Indeed, the content of this paper suggests that a unified perspective on standard gauge theories and Leibniz gauge theories [3, 23, 42] can emerge through the use of Lie-Leibniz triples and tensor hierarchies.

2 Lie-Leibniz triples and Lie algebra crossed modules

2.1 Leibniz algebras and Lie-Leibniz triples

The beginning of the section involves well-known facts about Leibniz algebras [2, 35] but then we introduce novel notions that will be thoroughly used in the paper. The field of scalars will be \mathbb{R} throughout the paper.

Definition 2. *A (left) Leibniz algebra is a vector space V together with a bilinear operation $\circ : V \otimes V \rightarrow V$ satisfying the relation:*

$$x \circ (y \circ z) = (x \circ y) \circ z + y \circ (x \circ z) \quad (5)$$

A Leibniz subalgebra of V is a subspace U which is stable under the Leibniz product.

A morphism of Leibniz algebras, or Leibniz algebra morphism, between (V, \circ) and (V', \circ') is a linear map $\chi : V \rightarrow V'$ that commutes with the respective Leibniz products:

$$\chi(x \circ y) = \chi(x) \circ' \chi(y) \quad (6)$$

*We call **Leib** the category of Leibniz algebras, together with Leibniz algebra morphisms.*

Let (V, \circ) be a Leibniz algebra. The Leibniz product \circ can be split into its skew-symmetric part, denoted $[\cdot, \cdot] : \wedge^2 V \rightarrow V$, and its symmetric part, denoted $\{\cdot, \cdot\} : S^2(V) \rightarrow V$, defined by:

$$[x, y] := \frac{1}{2}(x \circ y - y \circ x) \quad \text{and} \quad \{x, y\} := \frac{1}{2}(x \circ y + y \circ x)$$

for any $x, y \in V$, so that:

$$x \circ y = [x, y] + \{x, y\} \quad (7)$$

Example 1. *A Lie algebra is a Leibniz algebra whose product is skew-symmetric, i.e. such that $\{\cdot, \cdot\} = 0$. The category **Lie** of Lie algebras with Lie algebra morphisms is a full subcategory of **Leib**.*

Definition 3. *A left ideal of V (or ideal for short) is a subspace \mathcal{K} of V that satisfies the following condition:*

$$V \circ \mathcal{K} \subset \mathcal{K}$$

A left ideal \mathcal{K} of V whose left \circ -product with V is trivial, i.e. such that $\mathcal{K} \circ V = 0$, is called central.

Example 2. The sub-space \mathcal{I} of V generated by elements of the form $\{x, x\}$ is an ideal called the ideal of squares of V . A direct application of Equation (5) shows that it is central. The set \mathcal{Z} of all central elements of V , defined by:

$$\mathcal{Z} = \left\{ x \in V \mid x \circ y = 0 \text{ for all } y \in V \right\}$$

is a central ideal of V , and is the biggest such. It is called the centre of V .

In a general Leibniz algebra, the symmetric part of the bracket is not associative nor does the skew-symmetric part of the bracket satisfy the Jacobi identity (otherwise it would be a Lie bracket), but using Equation (5), we have:

$$[x, [y, z]] + [y, [z, x]] + [z, [x, y]] = -\frac{1}{3} \left(\{x, [y, z]\} + \{y, [z, x]\} + \{z, [x, y]\} \right) \quad (8)$$

for every $x, y, z \in V$. Since the right hand side lies in the ideal of squares \mathcal{I} , we deduce that for any central ideal \mathcal{K} satisfying the following inclusions:

$$\mathcal{I} \subset \mathcal{K} \subset \mathcal{Z},$$

the Leibniz product canonically induces a Lie algebra structure on the quotient V/\mathcal{K} . The Lie algebra V/\mathcal{I} is the biggest such, whereas V/\mathcal{Z} is the smallest. The quotient map $V \rightarrow V/\mathcal{I}$ is universal with respect to morphisms of Leibniz algebras from V to any Lie algebra [35].

A (symmetric) representation of a (left) Leibniz algebra is the natural generalization of the notion of Lie algebra representation:

Definition 4. A (symmetric) representation of a Leibniz algebra V is the data of a vector space E and a morphism of Leibniz algebras $\rho : V \rightarrow \text{End}(E)$.

In particular, since $\text{End}(E)$ is a Lie algebra, the linear map ρ satisfies the following identity:

$$\rho(x \circ y) = [\rho(x), \rho(y)] \quad (9)$$

for every $x, y \in V$. Since the right-hand side is fully skew-symmetric, so should be the left-hand side. Then, the ideal of squares \mathcal{I} is necessarily in the kernel of the map ρ . This is true for every symmetric representation of V . In particular, this implies that such representations correspond to representations of the Lie algebra V/\mathcal{I} [35].

As an example, one may consider the adjoint action of V on itself. Indeed, the Leibniz algebra V is a representation of V through the *adjoint map* – denoted ad – and defined by:

$$\begin{aligned} \text{ad} : V &\longrightarrow \text{End}(V) \\ x &\longmapsto \text{ad}_x : y \longmapsto x \circ y \end{aligned}$$

One can rewrite the Leibniz identity (5) as:

$$\text{ad}_{\text{ad}_x(y)}(z) = \text{ad}_x(\text{ad}_y(z)) - \text{ad}_y(\text{ad}_x(z)) \quad (10)$$

which, in turn, can be compactly summarized as:

$$\text{ad}_{\text{ad}_x(y)} = [\text{ad}_x, \text{ad}_y] \quad (11)$$

where the bracket on the right hand side is the Lie bracket on $\text{End}(V)$. Since $\text{ad}_x(y) = x \circ y$ on the left hand side, this equation proves that the adjoint action indeed induces a representation of V on itself.

Moreover, the Leibniz identity (5) shows that the map ad lands in the derivations of V . A *derivation* of a Leibniz algebra (V, \circ) is a linear map $\delta : V \rightarrow V$ such that:

$$\delta(x \circ y) = \delta(x) \circ y + x \circ \delta(y) \quad (12)$$

The space of derivations of V , denoted $\text{Der}(V)$, is a Lie subalgebra of $\text{End}(V)$. If δ is a derivation of V , then it is a derivation of the symmetric and the skew-symmetric parts of the Leibniz product. These results imply that both the ideal of squares \mathcal{I} and the centre \mathcal{Z} are stable by every derivation of V . The image of the adjoint action $\text{ad} : V \rightarrow \text{Der}(V)$ is a Lie subalgebra of $\text{Der}(V)$ called the *inner derivations of V* , and denoted $\text{inn}(V)$. The kernel of this map is the centre \mathcal{Z} of the Leibniz algebra. Hence, the adjoint map defines a Lie algebra isomorphism between V/\mathcal{Z} and $\text{inn}(V)$.

We are now interested in the relationship between the adjoint action of a Leibniz algebra and the embedding tensor of gauging procedures in supergravity. In these theories, the (super)-symmetries are controlled by a Lie algebra \mathfrak{g} and the field content of the physical model lies in various \mathfrak{g} -modules [16, 18]. It turns out that the gauge fields A_μ do not necessarily take values in the adjoint representation of \mathfrak{g} . This is rather unusual, compared to classical Yang-Mills gauge theories where such fields take values in the adjoint representation of \mathfrak{g} . Since gauge transformations are local symmetries generated by the action of gauge fields on the Lagrangian, physicists have developed a natural set-up to couple the representation in which such fields take values with the Lie algebra \mathfrak{g} of global symmetries, see e.g. [23].

More precisely, the basic data consist of a Lie algebra \mathfrak{g} and a \mathfrak{g} -module V , in which gauge fields take values. Then physicists observe that under some conditions the Lie algebra structure on \mathfrak{g} can be lifted to a Leibniz algebra structure on V [23]. Although the elements of V would play the role of gauge fields, their associated field strength – as given by the usual definition $F = dA + A \wedge A$ – are not covariant under gauge transformations, due precisely to the absence of skew-symmetry of the Leibniz algebra structure. This observation forces physicists to add higher fields to the model, taking values in additional \mathfrak{g} -modules, hence ending up with a tower of spaces that physicists call *tensor hierarchy* [16, 18]. This is a new manifestation of higher gauge theories in theoretical physics.

Physicists lift the Lie algebra structure of \mathfrak{g} to a Leibniz algebra structure on V by using what they call an *embedding tensor*. It is a linear mapping $\Theta : V \rightarrow \mathfrak{g}$ satisfying some conditions which, if satisfied, ensure the compatibility between the Lie algebra structure on \mathfrak{g} and the Leibniz algebra structure on V in a sense that resembles what occurs in a Lie algebra crossed module. Due to their importance in supergravity theories and in the present paper, triples of elements $(\mathfrak{g}, V, \Theta)$ satisfying these conditions deserve their own name:

Definition 5. A Lie-Leibniz triple is a triple $(\mathfrak{g}, V, \Theta)$ where:

- \mathfrak{g} is a Lie algebra,
- V is a \mathfrak{g} -module equipped with a Leibniz algebra structure $\circ : V \otimes V \rightarrow V$,
- $\Theta : V \rightarrow \mathfrak{g}$ is a linear mapping called the embedding tensor, that satisfies two compatibility conditions.

The first one is the linear constraint:

$$x \circ y = \Theta(x) \cdot y \quad (13)$$

The second one is called the quadratic constraint:

$$\Theta(x \circ y) = [\Theta(x), \Theta(y)]_{\mathfrak{g}} \quad (14)$$

where $[\cdot, \cdot]_{\mathfrak{g}}$ is the Lie bracket on \mathfrak{g} .

A sub-Lie-Leibniz triple of $(\mathfrak{g}, V, \Theta)$ is a Lie-Leibniz triple $(\mathfrak{t}, U, \Theta|_U)$ such that \mathfrak{t} (resp. U) is a Lie subalgebra of \mathfrak{g} (resp. Leibniz subalgebra of V).

Remark 1. What physicists call linear constraint, or representation constraint, is a condition on Θ that singles out the \mathfrak{g} sub-module of $\text{Hom}(V, \mathfrak{g})$ to which Θ belongs. In the physics literature, the linear constraint sometimes appears under the form that the symmetric bracket $\{.,.\}$ factors through a \mathfrak{g} sub-module of $S^2(V)$, see e.g. Equation (6) in [21]. This linear constraint can be interpreted as a weakening of one of the axioms of the notion of Lie algebra crossed module (Equation (40) in Lemma 11).

The Leibniz algebra structure on V is uniquely defined from the action of the Lie algebra \mathfrak{g} on V and Equation (13). This linear constraint shows that the embedding tensor is a *lift* of the adjoint action along the representation ρ :

$$\begin{array}{ccc} & & \mathfrak{g} \\ & \nearrow \Theta & \downarrow \rho \\ V & \xrightarrow{\text{ad}} & \text{End}(V) \end{array}$$

It is thus redundant to assume in Definition 5 that V is a Leibniz algebra beforehand. The same redundancy occurs in the definition of Lie algebra crossed modules. Moreover, Equations (13) and (14) are similar to those defining Lie algebra crossed modules. We will investigate in subsection 2.2 the extent of the precise relationship between the former and the latter notions.

In supergravity models, \mathfrak{g} is often taken to be the split real form of a semi-simple Lie algebra and V is the smallest-dimensional faithful representation of \mathfrak{g} [43]. The term *embedding tensor* has been introduced in the 2000s and refers to the fact that the image of the map Θ defines a Lie sub-algebra of \mathfrak{g} : this can be straightforwardly read from Equation (14). This Lie sub-algebra $\text{Im}(\Theta) \subset \mathfrak{g}$ plays the role of the gauge algebra in supergravity theories. For the particular role that this Lie subalgebra plays in this paper, we denote it by \mathfrak{h} . Throughout this section and the next one, major emphasis will be on the respective roles of \mathfrak{g} and of \mathfrak{h} in a general Lie-Leibniz triple, as well as on the relationship between ad and Θ . The quadratic constraint (14) additionally implies two things: that $\text{Ker}(\Theta)$ is an ideal of V and that $\mathcal{I} \subset \text{Ker}(\Theta)$. The linear constraint then implies that this ideal is central. Thus we have the following successive inclusions:

$$\mathcal{I} \subset \text{Ker}(\Theta) \subset \mathcal{Z} \tag{15}$$

The embedding tensor Θ then induces a Lie algebra isomorphism between $V/\text{Ker}(\Theta)$ and $\mathfrak{h} = \text{Im}(\Theta)$. The equality $\text{Ker}(\Theta) = \mathcal{Z}$ is satisfied only when V is a faithful representation of \mathfrak{g} .

Example 3. Let (V, \circ) be a Leibniz algebra and take $\mathfrak{g} = \text{End}(V)$. Let $\Theta = \text{ad}$ be the map that sends any element $x \in V$ to the adjoint endomorphism $\text{ad}_x : y \mapsto x \circ y$. The gauge algebra $\mathfrak{h} = \text{Im}(\Theta)$ consists of the inner derivations of V . Then $(\text{End}(V), V, \text{ad})$ is a Lie-Leibniz triple.

Another way of understanding the quadratic constraint is to note that it implies that Θ is $\text{Im}(\Theta)$ -equivariant (but not necessarily \mathfrak{g} -equivariant). This can be shown as follows: $\text{Hom}(V, \mathfrak{g})$ is equipped with a \mathfrak{g} -module structure through the induced action of \mathfrak{g} on V^* on the one hand and through the adjoint action of \mathfrak{g} on itself on the other hand. This defines a map:

$$\begin{aligned} \eta : \mathfrak{g} &\longrightarrow \text{End}(\text{Hom}(V, \mathfrak{g})) \\ a &\longmapsto \eta(a; -) : \Xi \longmapsto \left(x \mapsto [a, \Xi(x)]_{\mathfrak{g}} - \Xi(a \cdot x) \right) \end{aligned}$$

We also write $a \cdot \Theta$ for $\eta(a; \Theta)$, and more generally in the present paper, in order to represent the action of an element $a \in \mathfrak{g}$ on a vector r of any \mathfrak{g} -module R , we write $a \cdot r$. Then, for any Lie subalgebra \mathfrak{t} of the Lie algebra \mathfrak{g} , we say that Θ is \mathfrak{t} -equivariant if

$$\forall a \in \mathfrak{t} \quad a \cdot \Theta = 0 \quad (16)$$

Then, considering how \mathfrak{g} acts on V and on itself, Equation (14) can be rewritten:

$$\forall x \in V \quad \Theta(x) \cdot \Theta = 0 \quad (17)$$

In other words, the embedding tensor is \mathfrak{h} -equivariant since $\mathfrak{h} = \text{Im}(\Theta)$. We emphasize that Θ needs not be \mathfrak{g} -equivariant. The difference between the \mathfrak{h} -equivariance versus the \mathfrak{g} -equivariance of the embedding tensor Θ plays a central role in gauging procedures in supergravity.

Let us now investigate the relationship between the adjoint map and the embedding tensor Θ . Given a Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$, the adjoint map induced by the Leibniz product is an element of $\text{Hom}(V, \text{End}(V))$. Since V is a representation of \mathfrak{g} , the spaces $\text{End}(V)$ and $\text{Hom}(V, \text{End}(V))$ canonically inherit a structure of \mathfrak{g} -module. Then, since Equation (13) establishes that the embedding tensor is the lift of the adjoint action, Equation (10) implies that:

$$\Theta(x) \cdot \text{ad}_y - \text{ad}_{\Theta(x) \cdot y} = 0 \quad (18)$$

for every $x, y \in V$. In turn, this is equivalent to:

$$a \cdot \text{ad} = 0 \quad (19)$$

for every $a \in \mathfrak{h}$. That is to say: the map ad is \mathfrak{h} -equivariant, as is the embedding tensor. However, it is not necessarily \mathfrak{g} -equivariant, i.e. the action of \mathfrak{g} on ad may not be trivial. This discussion justifies the following definition:

Definition 6. *We say that a Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ is semi-strict if the adjoint map $\text{ad} : V \rightarrow \text{End}(V)$ is \mathfrak{g} -equivariant, and we say that it is strict if the embedding tensor Θ is \mathfrak{g} -equivariant.*

The semi-strictness and the strictness conditions are equivariance properties satisfied by, respectively, the adjoint action and its lift, the embedding tensor. The relationship between these two conditions relies on the relationship between these two maps. As the name indicates, a strict Lie-Leibniz triple is semi-strict but the converse is not necessarily true, as we will now show using another equivalent characterization of semi-strictness:

Lemma 7. *The Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ is semi-strict if and only if the Lie algebra morphism $\rho : \mathfrak{g} \rightarrow \text{End}(V)$ defining the \mathfrak{g} -module structure on V takes values in $\text{Der}(V)$.*

Proof. The fact that the map $\text{ad} \in \text{Hom}(V, \text{End}(V))$ is \mathfrak{g} -equivariant translates as Equation (19) for every $a \in \mathfrak{g}$. Then, following the reverse reasoning than led to Equation (19), one finds that this equation can be equivalently written as:

$$a \cdot (x \circ y) - x \circ (a \cdot y) - (a \cdot x) \circ y = 0 \quad (20)$$

for every $a \in \mathfrak{g}$ and $x, y \in V$. This equation characterizes the fact that $\rho : \mathfrak{g} \rightarrow \text{End}(V)$ takes values in $\text{Der}(V)$, hence the result. \square

Going back to our discussion: using the linear constraint (13), one can rewrite Equation (20) as:

$$a \cdot (\Theta(x) \cdot y) - \Theta(x) \cdot (a \cdot y) - \Theta(a \cdot x) \cdot y = 0 \quad (21)$$

The left-hand side can be rewritten as $(a \cdot \Theta)(x) \cdot y$, hence the property that $(\mathfrak{g}, V, \Theta)$ is semi-strict is given by the following condition:

$$\forall a \in \mathfrak{g}, \quad \forall x, y \in V \quad (a \cdot \Theta)(x) \cdot y = 0 \quad (22)$$

Thus, if the Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ is strict, i.e. if $a \cdot \Theta = 0$ for every $a \in \mathfrak{g}$, then it is semi-strict, but the converse is not true, for Equation (22) does not necessarily imply that Θ is \mathfrak{g} -equivariant, unless $\rho : \mathfrak{g} \rightarrow \text{End}(V)$ is injective, i.e. unless the representation V is faithful.

Remark 2. The notion of strict Lie-Leibniz triple had already been introduced by Loday and Pirashvili in Proposition 3.2 of [34], as well as by Kinyon and Weinstein in Definition 2.3. of [27], although it had not been given a name at the time. In [14], it is called a Loday-Pirashvili module.

Example 4. Let $(\mathfrak{g}, V, \Theta)$ be a Lie-Leibniz triple, and set $\mathfrak{h} = \text{Im}(\Theta)$. Then, by Equation (14), the Lie-Leibniz triple $(\mathfrak{h}, V, \Theta)$ is strict.

Example 5. By Lemma 7, the Lie-Leibniz triple $(\text{Der}(V), V, \text{ad})$ is semi-strict. Since the equality $f(\text{ad}_x(-)) = \text{ad}_{f(x)}(-) + \text{ad}_x(f(-))$ holds for every derivation f of V and $x \in V$, we deduce that the embedding tensor ad is $\text{Der}(V)$ -equivariant. This means that $(\text{Der}(V), V, \text{ad})$ is actually a strict Lie-Leibniz triple.

Example 6. We take the canonical example of [35] generalizing the construction of a Lie algebra associated to any associative algebra. Let (A, \cdot) be a \mathbb{R} associative algebra equipped with an endomorphism $D : A \rightarrow A$ which is not necessarily an algebra homomorphism but which satisfies the following identities:

$$D(x \cdot Dy) = Dx \cdot Dy = D(Dx \cdot y) \quad (23)$$

Then the bilinear product $\circ : A \times A \rightarrow A$ defined as:

$$x \circ y := Dx \cdot y - y \cdot Dx \quad (24)$$

is a Leibniz product on A . If $D = \text{id}_A$, then we obtain the canonical Lie algebra structure associated to A . Let us denote the corresponding Lie bracket by $[\cdot, \cdot]_A$ – not to be confused with the skew-symmetric part of the Leibniz product (24) – so that Equation (24) then reads:

$$x \circ y = [Dx, y]_A \quad (25)$$

and we say that \circ is a derived bracket [29].

With these data we can define a canonical Lie-Leibniz triple associated to any associative algebra A : \mathfrak{g} is the Lie algebra $(A, [\cdot, \cdot]_A)$, the Leibniz algebra V is (A, \circ) and the embedding tensor Θ is the endomorphism D . The action of the Lie algebra \mathfrak{g} on V is mediated through the adjoint action of the Lie bracket $[\cdot, \cdot]_A : x \cdot y = [x, y]_A$. The linear constraint is precisely Equation (25), while the quadratic constraint is a consequence of properties (23):

$$D(x \circ y) = D(Dx \cdot y) - D(y \cdot Dx) = Dx \cdot Dy - Dy \cdot Dx = [Dx, Dy]_A \quad (26)$$

Moreover, the gauge algebra \mathfrak{h} is precisely the image of the endomorphism D , itself a subspace of A . Now, since the Leibniz product is a derived bracket, the Jacobi identity on \mathfrak{g} together with properties (23) imply that we have, for any $x, y, z \in A$:

$$[x, y \circ z]_A = [x, [Dy, z]_A]_A \quad (27)$$

$$= [[x, Dy]_A, z]_A + [Dy, [x, z]_A]_A \quad (28)$$

$$= -[y \circ x, z]_A + y \circ [x, z]_A \quad (29)$$

$$= [x, y]_A \circ z + y \circ [x, z]_A - [y \circ x - D([y, x]_A), z]_A \quad (30)$$

So, the (semi)-strictness of the Lie-Leibniz triple $((A, [\cdot, \cdot]_A), (A, \circ), D)$ associated to the associative algebra A depends on the endomorphism D . The Lie-Leibniz triple is semi-strict if the bracket $[y \circ x - D([y, x]_A), z]_A$ vanishes for every x, y, z while the strictness condition – the \mathfrak{g} -equivariance of the embedding tensor – here reads $[x, Dy]_A = D[x, y]_A$ and is precisely equivalent to the vanishing of $y \circ x - D([y, x]_A)$. Let us apply these observations to the following two cases, that were introduced in [35]:

1. D is an algebra homomorphism and it is idempotent: $D^2 = D$;
2. D is a differential: $D(x \cdot y) = Dx \cdot y + x \cdot Dy$ and $D^2 = 0$.

In the first case, we first deduce from the homomorphism property that $D([x, y]_A) = [Dx, Dy]_A$. By Equation (26) we then conclude that $D([x, y]_A) = D(x \circ y)$ so the Lie-Leibniz triple associated to the associative algebra A and the endomorphism D is strict if and only if $x \circ y = D(x \circ y)$. In the second case, we deduce from the derivation property that $[Dx, y]_A = D([x, y]_A) + [Dy, x]_A$ so we obtain the skew-symmetric bracket of the Leibniz product:

$$[x, y] = \frac{1}{2}D([x, y]_A) \quad (31)$$

This has the following consequences: first, $D(x \circ y) = D([x, y]) = 0$ because $D^2 = 0$, and second: $y \circ x - D([y, x]_A) = x \circ y$ so the corresponding Lie-Leibniz triple is strict if and only if $x \circ y = 0$, i.e. the Leibniz algebra structure is zero.

Let us now observe some consequences of semi-strictness on the symmetric bracket $\{.,.\}$ entailed by the Leibniz algebra structure \circ on V . This bracket defines a map from $S^2(V)$ to V , also denoted $\{.,.\}$. This map, as an element of $\text{Hom}(S^2(V), V)$ is subject to the action of \mathfrak{g} induced from the \mathfrak{g} -module structure on V :

$$a \cdot (\{.,.\})(x, y) = a \cdot \{x, y\} - \{a \cdot x, y\} - \{x, a \cdot y\} \quad (32)$$

for every $a \in \mathfrak{g}$ and $x, y \in V$. The \mathfrak{g} -equivariance of the symmetric bracket is equivalent to the vanishing of the right hand side, which is a priori not granted. However, thanks to the linear constraint (13), the left-hand side of Equation (32) can be written as the following:

$$a \cdot (\{.,.\})(x, y) = \frac{1}{2}((a \cdot \Theta)(x)) \cdot y + \frac{1}{2}((a \cdot \Theta)(y)) \cdot x \quad (33)$$

Notice that, due to Equation (17), this term is always 0 when $a \in \mathfrak{h}$, thus implying that the symmetric bracket is always \mathfrak{h} -equivariant, but there is no guarantee that it is \mathfrak{g} -equivariant. However we deduce from Equation (22) that a sufficient condition for the symmetric bracket to be \mathfrak{g} -equivariant is that the Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ is semi-strict, so that the right-hand side of Equation (33) vanishes. These observations imply that, for any Lie-Leibniz triple, the space $\text{Ker}(\{.,.\}) \subset S^2(V)$ is always a \mathfrak{h} -module but not necessarily a \mathfrak{g} -module. This latter fact is central in the construction of the tensor hierarchy associated to a Lie-Leibniz triple.

2.2 A relationship between Lie-Leibniz triples and Lie algebra crossed modules

In order to study in a more consistent way the notion of Lie-Leibniz triple with respect to what is already known for Lie algebra crossed modules, we make use of categories. That is why we now introduce the correct notion for morphisms of Lie-Leibniz triples:

Definition 8. A morphism between two Lie-Leibniz triples $(\mathfrak{g}, V, \Theta)$ and $(\mathfrak{g}', V', \Theta')$ is a pair (φ, χ) consisting of a Lie algebra morphism $\varphi : \mathfrak{g} \rightarrow \mathfrak{g}'$ and a Leibniz algebra morphism $\chi : V \rightarrow V'$, satisfying the following compatibility conditions:

$$\Theta'(\chi(x)) = \varphi(\Theta(x)) \quad (34)$$

$$\chi(a \cdot x) = \varphi(a) \cdot \chi(x) \quad (35)$$

for every $a \in \mathfrak{g}$ and $x \in V$. That is to say, the following prism is commutative:

$$\begin{array}{ccccc}
& & \mathfrak{g} \otimes V & \xrightarrow{\varphi \otimes \chi} & \mathfrak{g}' \otimes V' \\
& \nearrow \Theta \otimes \text{id} & \downarrow \rho & & \nearrow \Theta' \otimes \text{id} \\
V \otimes V & \xrightarrow{\chi \otimes \chi} & V' \otimes V' & & \\
& \searrow \circ & \downarrow \chi & & \searrow \circ' \\
& & V & \xrightarrow{\chi} & V' \\
& & & & \downarrow \rho'
\end{array}$$

where ρ (resp. ρ') denotes the action of \mathfrak{g} (resp. \mathfrak{g}') on V (resp. V'). We say that (φ, χ) is an isomorphism of Lie-Leibniz triples when both φ and χ are isomorphisms in their respective categories.

Remark 3. Let (V, \circ) be a Leibniz algebra, then any Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ whose embedding tensor generates the Leibniz product \circ as given in Equation (13) induces a canonical Lie-Leibniz triple morphism (ρ, id_V) from $(\mathfrak{g}, V, \Theta)$ to $(\text{End}(V), V, \text{ad})$. Moreover, if $(\mathfrak{g}, V, \Theta)$ is semi-strict, the image of the Lie-Leibniz triple morphism (ρ, id_V) is a sub-Lie-Leibniz triple of $(\text{Der}(V), V, \text{ad})$.

A particular case of Lie-Leibniz triple morphisms are those for which the linear map $\chi \odot \chi : S^2 V \rightarrow S^2 V'$ – here, \odot symbolizes the symmetric product – preserves a particular submodule of $\text{Ker}(\{.,.\}) \subset S^2(V)$. Assume that we have a morphism (φ, χ) between two Lie-Leibniz triples $(\mathfrak{g}, V, \Theta)$ and $(\mathfrak{g}', V', \Theta')$. The kernel of the symmetric bracket on V is a subspace $\text{Ker}(\{.,.\})$ of $S^2(V)$ and a \mathfrak{h} -module but not necessarily a \mathfrak{g} -module. However, it admits \mathfrak{g} -submodules, and among them one is the biggest, that we denote by K . The same occurs for the symmetric bracket on V' and the biggest \mathfrak{g}' -submodule of $\text{Ker}(\{.,.\}) \subset S^2(V')$ is denoted K' . Since χ is a morphism of Leibniz algebras, we have that:

$$\chi \odot \chi(K) \subset \text{Ker}(\{.,.\})' \quad (36)$$

By Equation (35), the vector space $\chi \odot \chi(K)$ is only a $\varphi(\mathfrak{g})$ -module, and not a \mathfrak{g}' -module. So a priori, there is no reason that $\chi \odot \chi(K)$ would be a subspace of K' . We decide to give a name to morphisms that precisely have this property:

Definition 9. We say that a morphism of Lie-Leibniz triples $(\varphi, \chi) : (\mathfrak{g}, V, \Theta) \rightarrow (\mathfrak{g}', V', \Theta')$ is compatible if

$$\chi \odot \chi(K) \subset K' \quad (37)$$

This notion of compatible morphism allows us to define several categories: we call **Lie-Leib** the category of Lie-Leibniz triples with their associated morphisms given in Definition 8. We call **compLie-Leib** the wide subcategory of **Lie-Leib** whose morphisms are the compatible morphisms of Lie-Leibniz triples. In other words, $\text{Ob}(\mathbf{compLie-Leib}) = \text{Ob}(\mathbf{Lie-Leib})$, while $\text{Mor}(\mathbf{compLie-Leib})$ is contained in $\text{Mor}(\mathbf{Lie-Leib})$. The semi-strict (resp. strict) Lie-Leibniz triples form a full subcategory of both **compLie-Leib** and **Lie-Leib**, denoted **semLie-Leib** (resp. **strLie-Leib**). This can be seen from the fact that in semi-strict Lie-Leibniz triples, the biggest \mathfrak{g} -submodule of $\text{Ker}(\{.,.\}) \subset S^2(V)$ is the kernel itself. So, for $K = \text{Ker}(\{.,.\})$ and $K' = \text{Ker}(\{.,.\})'$, inclusion (36), which is valid for any morphism of Lie-Leibniz triples, becomes $\chi \odot \chi(K) \subset K'$. Hence any morphism of semi-strict Lie-Leibniz triples is compatible.

These inclusions of subcategories will allow us to be more precise when studying the relationship between Lie-Leibniz triples and Lie algebra crossed modules. Recall the definition of the latter notion:

Definition 10. A Lie algebra crossed module consists of the following data: two Lie algebras $(\mathfrak{g}, [\cdot, \cdot]_{\mathfrak{g}})$ and $(\mathfrak{c}, [\cdot, \cdot]_{\mathfrak{c}})$, a linear map $\Theta : \mathfrak{c} \rightarrow \mathfrak{g}$ and a Lie algebra morphism $\rho : \mathfrak{g} \rightarrow \text{Der}(\mathfrak{c})$, satisfying the following axioms:

$$\rho(\Theta(x); y) = [x, y]_{\mathfrak{c}} \quad (38)$$

$$\Theta(\rho(a; x)) = [a, \Theta(x)]_{\mathfrak{g}} \quad (39)$$

for every $x, y \in \mathfrak{c}$ and $a \in \mathfrak{g}$. We denote by **Lie \times Mod** the category of Lie algebra crossed modules.

Notice that the definition of ρ implies that \mathfrak{c} is a \mathfrak{g} -module, equipped with a Lie algebra structure entirely defined by Equation (38). Moreover, Equation (39) for $a = \Theta(z)$ implies that Θ is a Lie algebra morphism as well: $\Theta([z, x]_{\mathfrak{c}}) = [\Theta(z), \Theta(x)]_{\mathfrak{g}}$. These observations imply that we have an alternative, equivalent definition of Lie algebra crossed modules that relies on the following data:

Lemma 11. *A Lie algebra crossed module consists of the following data:*

1. a Lie algebra \mathfrak{g} ,
2. a \mathfrak{g} -module \mathfrak{c} , and
3. a \mathfrak{g} -equivariant linear map $\Theta : \mathfrak{c} \rightarrow \mathfrak{g}$ satisfying:

$$\Theta(x) \cdot y = -\Theta(y) \cdot x \quad (40)$$

Proof. Because of Equation (40), the vector space \mathfrak{c} can be equipped with a skew-symmetric bracket $[x, y]_{\mathfrak{c}} = \Theta(x) \cdot y$. This definition, together with the fact that \mathfrak{c} is a \mathfrak{g} -module has two consequences: on the one hand, the bracket $[\cdot, \cdot]_{\mathfrak{c}}$ satisfies the Jacobi identity and on the other hand, the representation map $\rho : \mathfrak{g} \rightarrow \text{End}(\mathfrak{c})$ takes values in $\text{Der}(\mathfrak{c})$. Eventually, Equation (38) is the definition of the bracket on \mathfrak{c} while Equation (39) corresponds to the \mathfrak{g} -equivariance of Θ . \square

The criteria established in Lemma 11 are reminiscent of the definition of a Lie-Leibniz triples. Items 1. and 2. establish that one can equip \mathfrak{c} with a Leibniz product, as in Equation (13), but Equation (40) implies that this product is skew-symmetric, hence a Lie bracket. Moreover, the \mathfrak{g} -equivariance of Θ that is constitutive of Lie algebra crossed modules straightforwardly implies the quadratic constraint (14). We then have the following relationship between Lie algebra crossed modules and Lie-Leibniz triples:

Lemma 12. *The category $\mathbf{Lie} \times \mathbf{Mod}$ forms a full subcategory of $\mathbf{strLie-Leib}$.*

Proof. Let $\mathfrak{c} \xrightarrow{\Theta} \mathfrak{g}$ be a Lie algebra crossed module. Then, Equations (38) and (39) imply Equations (13) and (14), under the following assumptions:

1. $\mathfrak{g} = \mathfrak{g}$ and $V = \mathfrak{c}$;
2. the Leibniz product \circ on \mathfrak{c} corresponds to the Lie algebra structure $[\cdot, \cdot]_{\mathfrak{c}}$.

This makes $(\mathfrak{g}, \mathfrak{c}, \Theta)$ a Lie-Leibniz triple. Moreover, Equation (39) tells us that the embedding tensor Θ is \mathfrak{g} -equivariant, i.e. that the Lie-Leibniz triple is strict. Morphisms of Lie algebra crossed modules are precisely the morphisms of the corresponding Lie-Leibniz triples, hence the result. \square

This lemma characterizes Lie algebra crossed modules as the strict Lie-Leibniz triples $(\mathfrak{g}, V, \Theta)$ for which V is a Lie algebra and for which the embedding tensor is \mathfrak{g} -equivariant. It moreover implies that we have the following inclusions of categories:

$$\mathbf{Lie} \times \mathbf{Mod} \subset \mathbf{strLie-Leib} \subset \mathbf{semLie-Leib} \subset \mathbf{compLie-Leib} \subset \mathbf{Lie-Leib}$$

where each inclusion is full in the sense of categories, except on the right-most inclusion where $\mathbf{compLie-Leib}$ is a wide subcategory of $\mathbf{Lie-Leib}$. The similarity between Lie algebra crossed modules and Lie-Leibniz triples is striking: one can see the latter as natural generalizations of the former, in which some stringent conditions (a Lie algebra structure on \mathfrak{c} and \mathfrak{g} -equivariance of Θ) have been relaxed.

Let us now push the analogy further by analyzing the differential graded Lie algebra structure of Lie algebra crossed modules and its natural extension to Lie-Leibniz triples. Indeed, an important property of Lie algebra crossed modules is that they can be considered as 2-term differential graded Lie algebras concentrated in degrees 0 and 1 [1] or, equivalently, in degrees 0 and -1 (our convention). This correspondence is derived

from the equivalence of categories between the category of crossed modules of Lie groups and that of strict Lie 2-groups [7, 25]. We recall here how the identification works, since it will be a cornerstone of the present section.

Given a Lie algebra crossed module $\mathfrak{c} \xrightarrow{\Theta} \mathfrak{g}$, assign degree -1 to \mathfrak{c} and degree 0 to \mathfrak{g} . Then, define the graded Lie bracket on $\mathfrak{c} \oplus \mathfrak{g}$ as follows:

$$[[a, b]] = [a, b]_{\mathfrak{g}} \quad (41)$$

$$[[x, y]] = 0 \quad (42)$$

$$[[a, x]] = a \cdot x \quad \text{and} \quad [[x, a]] = -a \cdot x \quad (43)$$

for every $a, b \in \mathfrak{g}$ and $x, y \in \mathfrak{c}$. This bracket and the differential Θ are indeed compatible, in the sense that together they form a differential graded Lie algebra structure. The first non-trivial compatibility condition between the graded Lie bracket $[[\cdot, \cdot]]$ and the differential Θ is:

$$\Theta([a, x]) = [[a, \Theta(x)]] \quad (44)$$

This is nothing but Equation (39). The other non-trivial compatibility condition is:

$$\Theta([x, y]) = [[\Theta(x), y]] - [[x, \Theta(y)]] \quad (45)$$

Notice that the left hand side vanishes identically because of Equation (42), and then the right hand side just tells us that $\Theta(x) \cdot y = -\Theta(y) \cdot x$ (Equation (40)). However this is a tautology because by Equation (38), $\Theta(x) \cdot y = [x, y]_{\mathfrak{c}}$ and we know that the Lie bracket on \mathfrak{c} is skew-symmetric. Eventually, the bracket defined in Equations (41)-(43) satisfies the Jacobi identity as the only non-obvious identity to satisfy is the following one:

$$[[a, [b, x]]] - [[b, [a, x]]] = [[a, b], x] \quad (46)$$

This is an alternative, equivalent rewriting of the fact that $\rho : \mathfrak{g} \rightarrow \text{Der}(\mathfrak{c})$ is a Lie algebra morphism. Thus, a Lie algebra crossed module is equivalent to a 2-term differential graded Lie algebra concentrated in degrees 0 and -1 .

Let us now present an alternative, equivalent formulation of this observation by using the cyclic representation generated by Θ . Let $(\mathfrak{g}, V, \Theta)$ be a Lie-Leibniz triple and let $\eta : \mathfrak{g} \rightarrow \text{End}(\text{Hom}(V, \mathfrak{g}))$ denote the induced action of \mathfrak{g} on $\text{Hom}(V, \mathfrak{g})$ as defined in Section 2.1. As an element of $\text{Hom}(V, \mathfrak{g})$, the embedding tensor Θ generates a cyclic \mathfrak{g} -submodule of the \mathfrak{g} -module $\text{Hom}(V, \mathfrak{g})$; we call it R_{Θ} . More precisely, noting

$$\Theta_{a_1 a_2 \dots a_n} := a_1 \cdot (a_2 \cdot (\dots (a_n \cdot \Theta) \dots)) = \eta(a_1; \eta(a_2; \dots (\eta(a_n; \Theta)) \dots)) \quad (47)$$

for any $a_1, a_2, \dots, a_n \in \mathfrak{g}$, we have:

$$R_{\Theta} := \text{Span}(\Theta, \Theta_{a_1 a_2 \dots a_m} \mid a_1, a_2, \dots, a_m \in \mathfrak{g}) \subset \text{Hom}(V, \mathfrak{g}) \quad (48)$$

In the case where the Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ is a Lie algebra crossed module – i.e. when $\mathfrak{g} = \mathfrak{g}$ and $V = \mathfrak{c}$ – then Θ being \mathfrak{g} -equivariant means that R_{Θ} is the one-dimensional irreducible trivial representation of \mathfrak{g} and can be identified with \mathbb{R} . Then, the differential graded Lie algebra structure on $\mathfrak{c} \oplus \mathfrak{g}$ presented earlier can be straightforwardly extended to a 3-term differential graded Lie algebra structure on the following cochain complex:

$$\mathfrak{c} \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{0} \mathbb{R}$$

One consider that elements of \mathbb{R} carry a degree $+1$, so that the grading on the complex is increasing from left to right. Actually, the above sequence is a particular case of the following observation:

Proposition 13. *To every Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ in which V is a Lie algebra, there corresponds a unique 3-term differential graded Lie algebra concentrated in degrees $-1, 0$ and 1^2 :*

$$V[1] \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{-\eta(-; \Theta)} R_\Theta[-1]$$

which canonically extends the graded Lie algebra structure on $V[1] \oplus \mathfrak{g}$ defined by Equations (41)-(43).

Proof. Given a Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ in which V is assumed to be a Lie algebra (i.e. the Leibniz product \circ is skew-symmetric), then one defines the following brackets (the first four being those of Equations (41)-(43)):

$$[[a, b]] = [a, b]_{\mathfrak{g}}, \quad [[x, y]] = 0, \quad (49)$$

$$[[a, x]] = a \cdot x, \quad [[x, a]] = -a \cdot x, \quad (50)$$

$$[[a, \Theta]] = a \cdot \Theta, \quad [[\Theta, a]] = -a \cdot \Theta, \quad (51)$$

$$[[\Theta, x]] = \Theta(x) \quad \text{and} \quad [[x, \Theta]] = \Theta(x), \quad (52)$$

for every $a, b \in \mathfrak{g}$ and $x, y \in V[1]$. Since R_Θ is generated by the successive actions of \mathfrak{g} on Θ , one extends the above brackets to the whole of $R_\Theta[-1]$ by the following formulas:

$$[[\Theta_{a_1 a_2 \dots a_m}, u]] = a_1 \cdot [[\Theta_{a_2 \dots a_m}, u]] - [[\Theta_{a_2 \dots a_m}, a_1 \cdot u]] \quad (53)$$

$$[[u, \Theta_{a_1 a_2 \dots a_m}]] = -(-1)^{|u|} [[\Theta_{a_1 a_2 \dots a_m}, u]] \quad (54)$$

$$0 = [[R_\Theta[-1], R_\Theta[-1]]] \quad (55)$$

for any homogeneous element $u \in \mathfrak{g} \oplus V[1]$ of degree $|u|$, and where $\Theta_{a_1 a_2 \dots a_m} = a_1 \cdot (a_2 \cdot (\dots (a_m \cdot \Theta) \dots))$ for any $a_1, a_2, \dots, a_m \in \mathfrak{g}$. Moreover setting $\partial_1 = -\eta(-, \Theta)$ and $\partial_0 = \Theta$, one has $\partial_1(\partial_0(x)) = 0$ by Equation (17). Actually, Lines (51) and (52) imply that the induced differential ∂ coincides with the adjoint action of Θ :

$$\partial = [[\Theta, -]] \quad (56)$$

Using the properties of Lie-Leibniz triples, one may check that the above brackets and differentials satisfy the graded Jacobi identities and Leibniz rule. We have already shown several of them: the Jacobi identity on $\mathfrak{g} \oplus V[1]$ is Equation (46), while the Leibniz rule for any pair of element $x \in V[1]$ and $a \in \mathfrak{g}$ can be found in Equations (44) and (45). We will now show that their validity extends to $R_\Theta[-1]$. In the following, we assume that $a, b, a_1, \dots, a_m \in \mathfrak{g}$, and $x, y \in V$. We first turn to the Leibniz rules.

Using Equation (53) for $[[\Theta_a, b]]$ and reorganizing the terms, one has:

$$\partial([a, b]) = [[\Theta, [a, b]_{\mathfrak{g}}]] = a \cdot [[\Theta, b]] - [[\Theta_a, b]] \quad (57)$$

$$= [[a, [[\Theta, b]]]] + [[[[\Theta, a], b]] = [[a, \partial(b)]] + [[\partial(a), b]] \quad (58)$$

This proves the Leibniz rule for the bracket $[[a, b]]$. Next, because of the degree, one knows that any Leibniz rule involving an element of $R_\Theta[-1]$ and an element of \mathfrak{g} is automatically zero. This leaves us with only one Leibniz rule to check: the one involving one element $x \in V[1]$ and every generator of $R_\Theta[-1]$. The Leibniz rule between x and Θ should read:

$$\partial([[\Theta, x]]) = -[[\Theta, \partial(x)]]] \quad (59)$$

because $\partial(R_\Theta[-1]) = 0$ by construction. But Equation (56) implies that the left-hand side is zero, while the right-hand side is $\eta(\partial(x), \Theta) = \Theta(x) \cdot \Theta$ which also vanishes by the quadratic constraint (17). This proves that the Leibniz rules is satisfied on the pair of elements x and Θ .

²Our convention for suspension is the following: for any graded vector space $U_\bullet = \bigoplus_{i \in \mathbb{Z}} U_i$, the *suspension operator* $[-1] : U \rightarrow U[-1]$ shifts the degree of every elements in U by 1, i.e. it is such that $(U[-1])_i = U_{i-1}$. The *desuspension operator* $[1]$ has the reverse effect.

It is now sufficient to prove it on any other choice of generator $\Theta_{a_1 \dots a_m}$ of R_Θ . While one can deduce from Equation (53) that $\partial(\llbracket \Theta_a, x \rrbracket) = -\llbracket \Theta_a, \partial(x) \rrbracket$, the same does not hold for generators of $R_\Theta[-1]$ of the form $\Theta_{a_1 \dots a_m}$ for $m \geq 2$ (see the discussion following Equation (148)). Then, as a further assumption, one is allowed to additionally require that:

$$\partial(\llbracket \Theta_{a_1 \dots a_m}, x \rrbracket) = -\llbracket \Theta_{a_1 \dots a_m}, \partial(x) \rrbracket \quad (60)$$

This is precisely the Leibniz rule for x and $\Theta_{a_1 \dots a_m}$. One concludes that the Leibniz rule holds for every pair of elements taken from $V[1]$ and $R_\Theta[-1]$, and thus on the whole of $V[1] \oplus \mathfrak{g} \oplus R_\Theta[-1]$.

We can now turn to proving the remaining Jacobi identities. We already know by Equation (46) and because \mathfrak{g} is a Lie algebra that the graded Jacobi identity is satisfied on $V[1] \oplus \mathfrak{g}$. Moreover it is trivially satisfied on $R_\Theta[-1]$ because the bracket is zero on this space. Then we need only check the cases for which one or two terms belong to $R_\Theta[-1]$ and the other one or two belong to $V[1] \oplus \mathfrak{g}$. The proof is made by induction and, in order to avoid unnecessary technical computation at this stage of the paper, we postpone it to the second part of the proof of Proposition 23 (see in particular the discussion below Equation (163)). This concludes the proof of Proposition 13. \square

This result shows that the assignment to a Lie algebra crossed module of a differential graded Lie algebra can be extended to specific cases of Lie-Leibniz triples: those for which the Leibniz algebra is a Lie algebra, without further assumption on the \mathfrak{g} -equivariance of the embedding tensor. One objective of the paper is to show that one can generalize Proposition 13 to every Lie-Leibniz triples. However, since Lie-Leibniz triples involve Leibniz algebras and not just Lie algebras, we expect new intricate structures to emerge. In particular, one would expect to associate a bigger differential graded Lie algebra to every Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ whose Leibniz algebra V is not a Lie algebra:

$$\dots \longrightarrow V[1] \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{-\eta(-; \Theta)} R_\Theta[-1]$$

More precisely, let \overline{G} be the functor associating to each Lie algebra crossed module the corresponding unique 3-term differential graded Lie algebra:

$$\begin{aligned} \overline{G}: \mathbf{Lie} \times \mathbf{Mod} &\longrightarrow \mathbf{DGLie}_{\leq 1} \\ (\mathfrak{c} \xrightarrow{\Theta} \mathfrak{g}) &\longmapsto (\mathfrak{c} \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{0} \mathbb{R}[-1]) \end{aligned}$$

This functor canonically extends to the category of Lie-Leibniz triples for which V is a Lie algebra – this is the content of Proposition 13. Moreover, it turns out that this functor satisfies Theorem 1, which the rest of the paper is dedicated to proving.

We will show that, in the presence of a symmetric component in the Leibniz product of a Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$, the cochain complex of Proposition 13 actually extends to the left:

$$\dots \xrightarrow{\partial_{-3}} T_{-3} \xrightarrow{\partial_{-2}} T_{-2} \xrightarrow{\partial_{-1}} V[1] \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{-\eta(-; \Theta)} R_\Theta[-1] \quad (61)$$

The existence of a tower of spaces $T_\bullet = \bigoplus_{i \geq 1} T_{-i}$ beyond degree -1 is thus the sign that the Leibniz algebra V is *not* a Lie algebra. The \mathfrak{g} -module structure of V is key in defining T_{-2} , as it is used in its very definition. The other T_{-i} for $i \geq 3$ are built by induction as quotients of sums of products of other T_{-j} for $1 \leq j \leq i-1$ so that the property of being a \mathfrak{g} -module is inherited by induction. The action of \mathfrak{g} on T_{-i} is moreover necessary to define the differential on T_\bullet , showing the centrality of the Lie algebra \mathfrak{g} in the construction. As proved in Proposition 23, the total space $\mathbb{T} = T_\bullet \oplus \mathfrak{g} \oplus R_\Theta[-1]$ then canonically inherits a differential graded Lie algebra structure, which restricts to that of Proposition 13 when V is a Lie algebra.

This construction improves and simplifies the one presented in [31]. In that paper, the author associated a differential graded Lie algebra $((T_{-i})_{i \geq 0}, \partial, [\cdot, \cdot])$ to any Lie-Leibniz triple such that $T_{-1} = V[1]$, $T_0 = \mathfrak{h} = \text{Im}(\Theta)$ and $\partial_0 = \Theta$:

$$\dots \xrightarrow{\partial_{-3}} T_{-3} \xrightarrow{\partial_{-2}} T_{-2} \xrightarrow{\partial_{-1}} V[1] \xrightarrow{\Theta} \mathfrak{h} \quad (62)$$

Here, the spaces T_{-i} and the linear maps ∂_{-i} are the same as in the cochain complex (61). The construction presented in [31] used a dual point of view which implied rather cumbersome computations and had the serious drawback that the resulting differential graded Lie algebra lost important information about the Lie algebra \mathfrak{g} (for example it did not appear in the resulting cochain complex (62)). In the present paper, we propose to offer a novel and direct construction of the cochain complex (61), drawing on theoretical work done by physicists, among others [10, 20]. Moreover, we shall prove that each map ∂_{-i} , as an element of $\text{Hom}(T_{-i-1}, T_{-i})$ on which \mathfrak{g} acts, generates a cyclic \mathfrak{g} -submodule of $\text{Hom}(T_{-i-1}, T_{-i})$ which is isomorphic to (a sub-module of) R_Θ . This result, although having been conjectured and used for a long time by physicists, is new and is needed precisely in order to extend the cochain complex (62) to the right as in diagram (61).

3 The tensor hierarchy associated to a Lie-Leibniz triple

Subsection 3.1 is dedicated to defining the particular graded vector space T_\bullet and equip it with a graded Lie bracket. Subsection 3.2 is dedicated to defining the differential, while subsection 3.3 proves the compatibility of the bracket with the differential, turning the total space $\mathbb{T} = T_\bullet \oplus \mathfrak{g} \oplus R_\Theta[-1]$ into a differential graded Lie algebra. This algebra is the one defining the function G in Theorem 1. We refer the reader to Example 7 in order to fully appreciate the details of the first steps of the construction of the algebra. Finally, subsection 3.4 is devoted to showing that the restriction of G to the subcategory **compLie-Leib** is a faithful functor.

3.1 Associating a graded Lie algebra to a Lie-Leibniz triple

The main result of the current subsection shows that any Lie-Leibniz triple gives rise to a graded Lie algebra generalizing the \mathfrak{g} -module \mathfrak{c} that appears in any Lie algebra crossed module $\mathfrak{c} \xrightarrow{\Theta} \mathfrak{g}$:

Proposition 14. *To any Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ one can associate a negatively graded Lie algebra $(T_\bullet = (T_{-i})_{i \geq 1}, [\cdot, \cdot])$, which satisfies the following three conditions:*

1. T_\bullet is a graded \mathfrak{g} -module, and the corresponding representation $\rho : \mathfrak{g} \rightarrow \text{End}(T_\bullet)$ takes values in $\text{Der}(T_\bullet)$, the space of derivations of $[\cdot, \cdot]$;
2. $T_\bullet = V[1] \oplus [T_\bullet, T_\bullet]$;
3. $T_\bullet = T_{-1} = V[1]$ if and only if V is a Lie algebra.

The first item of the proposition means that for every $i \geq 1$, T_{-i} is a \mathfrak{g} -module under the action of a Lie algebra morphism $\rho_{-i} : \mathfrak{g} \rightarrow \text{End}(T_{-i})$, so that ρ is the unique (graded) Lie algebra morphism restricting to ρ_{-i} on $\text{End}(T_{-i})$. Moreover for every $a \in \mathfrak{g}$, $x \in T_{-i}$ and $y \in T_{-j}$, $\rho(a; -)$ is a derivation of the graded Lie bracket $[x, y]$:

$$\rho_{-i-j}(a; [x, y]) = [\rho_{-i}(a; x), y] + [x, \rho_{-j}(a; y)] \quad (63)$$

The second item can be reformulated as follows: $T_{-1} = V[1]$ and for every $i \geq 2$, $T_{-i} = \sum_{j=1}^{i-1} [T_{-j}, T_{-i+j}]$. Eventually, the last item is the condition that $T_{-i} = 0$ for every $i \geq 2$ if and only if V is a Lie algebra. In other words, when V is a Lie algebra, the tower of vector spaces reduces to $T_\bullet = T_{-1} = V[1]$ and the graded Lie bracket $[\cdot, \cdot]$ is identically zero, whereas when V is a Leibniz algebra whose Leibniz product is not fully skew-symmetric, then the spaces of lower degrees exist and satisfy item 2. of Proposition 14.

The proof is split in two parts. First, we will define the T_{-i} recursively, starting by setting $T_{-1} = V[1]$, and then by taking the quotient of the free graded Lie algebra generated by $V[1]$ by a particular graded ideal $K_{\bullet} = \bigoplus_{i \geq 2} K_{-i}$ which is constructed by induction. The quotient then is a (possibly not bounded) graded vector space $T_{\bullet} = \bigoplus_{i=1}^{\infty} T_{-i}$ which has the following properties:

1. Every vector space T_{-i} is a \mathfrak{g} -module;
2. $T_{-1} = V[1]$;
3. $T_{-i} = 0$ for every $i \geq 2$ if and only if V is a Lie algebra.

This represents the biggest part of the proof, and it crucially relies on key Lemmas 15 and 16. Then, we will define a graded Lie bracket on T_{\bullet} , so that items 1. and 2. of Proposition 14 are satisfied by construction.

Remark 4. *The idea behind the proof (quotienting a graded ideal out of a free graded Lie algebra) in the context of building a tensor hierarchy has first been given in [10], where the authors use a partition function in order to compute the \mathfrak{g} -modules that form the building blocks of the hierarchy. On the other hand, our proof relies on an inductive argument that is algebraic and that enables us to make explicit the definition of the graded Lie bracket. The relationship between the graded Lie algebra structure presented in [10] and ours has still to be investigated, but we expect it to be very close if not identical.*

Let us first give a brief account of our conventions and a technical reminder of graded algebra. Given a finite dimensional vector space X , one can define the free Lie algebra $\text{Free}(X)$ as the Lie algebra generated by X , imposing only the defining relations of skew-symmetry and the Jacobi identity. It can be thought of as the vector space generated by the successive commutators of a basis of X . The vector space $F_{\bullet} = \text{Free}(X)$ is graded by the length of the commutators, e.g. $[\cdots [x_1, x_2], x_3], \cdots x_i]$, for any $x_1, \dots, x_i \in X$, has length i . There exist various formulas for a basis of $F_{\bullet} = \bigoplus_{k \geq 1} F_k$ [22, 36, 40]. The construction also works for graded vector spaces X , hence defining free graded Lie algebras, also called free Lie superalgebras when the emphasis is on the parity [15, 41].

First, let us recall that for a graded vector space X , the free graded skew-commutative algebra generated by X is denoted by $\Lambda^{\bullet}(X)$. That is, the algebra spanned by elements $x_1 \wedge x_2 \wedge \cdots \wedge x_n$. With the degree of x denoted $|x|$, we have:

$$x \wedge y = -(-1)^{|x||y|} y \wedge x \quad (64)$$

For example, pick a vector space V and define $X = V[1]$ to be its desuspension, that is to say: X is isomorphic to V as a vector space, but its elements are considered to carry a grading defined to be -1 . Then, due to the shifted degree and Equation (64), the vector space $\Lambda^2(X)$ is isomorphic to $S^2(V)$ (understood in the category of vector spaces).

Thus, the free graded Lie algebra $F_{\bullet} = \text{Free}(X)$ generated by the degree -1 vector space $F_{-1} = X = V[1]$ admits $\Lambda^2(X) \simeq S^2(V)$ as the desired space F_{-2} , where \simeq indicates that the two spaces are isomorphic. Notice that for $x, y \in F_{-1}$, we have $[x, y] = [y, x]$, where here the bracket is the free graded Lie bracket on F_{\bullet} . The next vector space F_{-3} has a basis of some elements of the form $[[x, y], z]$ for $x, y, z \in F_{-1}$, subject to the Jacobi identity. Thus F_{-3} is isomorphic to the quotient of $F_{-1} \otimes \Lambda^2(F_{-1}) \simeq V \otimes S^2(V)$ by $\Lambda^3(F_{-1}) \simeq S^3(V)$. More generally, the negative grading on $F_{\bullet} = \bigoplus_{k \geq 1} F_{-k}$ is defined as follows: the number $k \geq 1$ labelling F_{-k} indicates the number of elements $x_1, \dots, x_k \in X = V[1]$ used in the brackets $[\cdots [x_1, x_2], x_3], \dots, x_k]$ defining the generators of F_{-k} .

In this section, we will consider the total exterior algebra $\Lambda(F_{\bullet}) = \bigoplus_{p=0}^{\infty} \Lambda^p F_{\bullet}$ of a free graded Lie algebra of the form $F_{\bullet} = \text{Free}(V[1])$. The exterior algebra inherits a grading from F_{\bullet} which is such that the component in total degree $-i$ is denoted $\Lambda(F_{\bullet})_{-i} = \bigoplus_{p \geq 0} \Lambda^p(F_{\bullet})|_{-i}$. The space $\Lambda(F_{\bullet})_{-i}$ at degree $-i$ is a finite dimensional graded vector space. This exterior algebra is obtained as the quotient of the tensor algebra $T(F_{\bullet})$ by the (graded) ideal generated by elements of the form:

$$x \otimes y + (-1)^{|x||y|} y \otimes x \quad (65)$$

for every two homogeneous elements $x, y \in F_\bullet$. The resulting quotient is isomorphic to a graded sub-vector space of the tensor algebra, and is a graded algebra on its own, when equipped with the wedge product.

Let us discuss the first few spaces appearing in the decomposition of $\Lambda^2(F_\bullet)$. As a vector space, $\Lambda^2(F_\bullet)|_{-2}$ is canonically isomorphic to $F_{-1} \wedge F_{-1}$, while $\Lambda^2(F_\bullet)|_{-3}$ is canonically isomorphic to $F_{-1} \otimes F_{-2}$, whereas $\Lambda^2(F_\bullet)|_{-4}$ is canonically isomorphic to $F_{-1} \otimes F_{-3} \oplus F_{-2} \wedge F_{-2}$. More generally, $\Lambda^2(F_\bullet)_{-i}$ (for $i \geq 3$) is canonically isomorphic to the following vector spaces depending on the parity of i :

$$\text{if } i \text{ is even} \quad F_{-\frac{i}{2}} \wedge F_{-\frac{i}{2}} \quad \oplus \quad \bigoplus_{1 \leq j \leq \frac{i-2}{2}} F_{-j} \otimes F_{-i+j} \quad (66)$$

$$\text{if } i \text{ is odd} \quad \bigoplus_{1 \leq j \leq \frac{i-1}{2}} F_{-j} \otimes F_{-i+j} \quad (67)$$

When $j \neq \frac{i}{2}$, what we denote by $F_{-j} \wedge F_{-i+j}$ is the vector space isomorphic to $F_{-j} \otimes F_{-i+j}$, understood as a subspace of $\Lambda^2(F_\bullet)_{-i}$. In particular, if $x \in F_{-j}$ and $y \in F_{-i+j}$ then we will write $x \wedge y$ instead of $x \otimes y$ when we want to emphasize that we work in $F_{-j} \wedge F_{-i+j}$. Regarding $F_{-j} \wedge F_{-i+j}$ as isomorphic to both $F_{-j} \otimes F_{-i+j}$ and $F_{-i+j} \otimes F_{-j}$, we then have $x \wedge y$ represented by $x \otimes y \in F_{-j} \otimes F_{-i+j}$ or equally by $-(-i)^{j(i-j)} y \otimes x \in F_{-i+j} \otimes F_{-j}$.

The Chevalley-Eilenberg homology of a graded Lie algebra \mathfrak{g} is the homology of their chain complex $CE_\bullet(\mathfrak{g})$ with $CE_n(\mathfrak{g}) = \Lambda^n \mathfrak{g}$ and differential $d = \{d_n : CE_n(\mathfrak{g}) \rightarrow CE_{n-1}(\mathfrak{g})\}$ given by ([19] p. 471):

$$d_n(x_1 \wedge \dots \wedge x_n) = \sum_{1 \leq i < j \leq n} (-1)^{i+j+|x_j|\eta_j+|x_i|\eta_i-|x_i||x_j|} [x_i, x_j] \wedge x_1 \wedge \dots \wedge x_{i-1} \wedge x_{i+1} \wedge \dots \wedge x_{j-1} \wedge x_{j+1} \wedge \dots \wedge x_n \quad (68)$$

where $\eta_m = \sum_{1 \leq k \leq m-1} |x_k|$. Degrees of the elements appear in the formula because one has to permute x_i and x_j to the first and second positions in order to eventually take the Lie bracket of x_i and x_j . There exists another, identical but more compact formula, which uses unshuffles:

$$d_n(x_1 \wedge \dots \wedge x_n) = - \sum_{\sigma \in \text{Un}(2, n-2)} \epsilon^\sigma [x_{\sigma(1)}, x_{\sigma(2)}] \wedge x_{\sigma(3)} \wedge \dots \wedge x_{\sigma(n)} \quad (69)$$

where $\text{Un}(2, n-2)$ are the $(2, n-2)$ -unshuffles, and where ϵ^σ is the sign that is induced by the permutation σ on $x_1 \wedge \dots \wedge x_n$. More precisely, a permutation $\sigma \in S_n$ is called a $(p, n-p)$ -unshuffle if $\sigma(1) < \dots < \sigma(p)$ and $\sigma(p+1) < \dots < \sigma(n)$, and ϵ^σ is defined through the following formula:

$$x_1 \wedge \dots \wedge x_n = \epsilon^\sigma x_{\sigma(1)} \wedge \dots \wedge x_{\sigma(n)} \quad (70)$$

One may check that Equation (68) is the same as Equation (69), and that they correspond to the common definition of the Chevalley-Eilenberg differential (in homology) when \mathfrak{g} is concentrated in degree 0 (see [28] p. 284). Then the following result holds:

Lemma 15. *For a free graded Lie algebra $F_\bullet = (F_{-i})_{i \geq 1}$, the following sequences are exact:*

$$0 \longrightarrow \Lambda^2(F_{-1}) \xrightarrow{d_2} F_{-2} \longrightarrow 0 \quad (71)$$

$$\Lambda^3(F_{-1}) \xrightarrow{d_3} F_{-1} \wedge F_{-2} \xrightarrow{d_2} F_{-3} \longrightarrow 0 \quad (72)$$

and more generally, for every $i \geq 4$:

$$\Lambda^3\left(\bigoplus_{j=1}^{i-2} F_{-j}\right)\Big|_{-i} \xrightarrow{d_3} \Lambda^2\left(\bigoplus_{j=1}^{i-1} F_{-j}\right)\Big|_{-i} \xrightarrow{d_2} F_{-i} \longrightarrow 0 \quad (73)$$

Proof. For a free graded Lie algebra F_\bullet , one has $H_1(F_\bullet) = 0$ and $H_2(F_\bullet) = 0$. The first equality is justified because on the one hand, the map $d_1 : F_\bullet \rightarrow \mathbb{R}$ is the zero map and, on the other hand, $d_2 : \Lambda^2(F_\bullet) \rightarrow F_{\leq -2}$ is by definition surjective: F_\bullet is a free graded Lie algebra so each F_{-i} (for $i \geq 2$) is spanned by (iterated) brackets. The second equation, $H_2(F_\bullet) = 0$, is deduced from Corollary 6.5 in [19] by keeping track of the natural graduation of F_\bullet . Then it suffices to write the equations $H_1(F_\bullet) = 0$ and $H_2(F_\bullet) = 0$ degree-wise, starting at $n = 2$, to obtain the exact sequences (71), (72) and (73). \square

The proof of Proposition 14 then relies on the use of Lemma 15 in conjunction with the following Lemma:

Lemma 16. *Let \mathcal{A} be an abelian category and assume that the following diagram in \mathcal{A} is commutative:*

$$\begin{array}{ccccccc}
 0 & \longrightarrow & A & \xrightarrow{\text{id}} & A & \longrightarrow & 0 \\
 & & \downarrow u & & \downarrow k & & \\
 W & \xrightarrow{a} & X & \xrightarrow{b} & Y & \xrightarrow{c} & Z \\
 & & \downarrow & & \downarrow & & \\
 W & \xrightarrow{\alpha} & X/u(A) & \xrightarrow{\beta} & Y/k(A) & \xrightarrow{\gamma} & Z
 \end{array}$$

where α, β and γ are the linear maps canonically induced from a, b and c . Then exactness of the second row implies exactness of the third.

Remark 5. Here, given a morphism $\chi \in \text{Hom}_{\mathcal{A}}(P, Q)$, the quotient notation $P/\chi(Q)$ denotes the cokernel of the map χ , that is: $P/\chi(Q) := \text{coker}(\chi : P \rightarrow Q)$. In the category of R -modules, this reduces to the usual quotient of R -modules.

Proof. Exactness of the sequence:

$$W \xrightarrow{a} X \xrightarrow{b} Y \xrightarrow{c} Z$$

is equivalent to the existence of two isomorphisms:

$$\text{Coker}(a) \simeq \text{Im}(b) \quad \text{and} \quad \text{Coker}(b) \simeq \text{Im}(c)$$

Now we have the following isomorphisms:

$$\text{Im}(\gamma) \simeq \text{Im}(c) \simeq \text{Coker}(b) \simeq \text{Coker}(\beta)$$

which implies that the sequence:

$$W \xrightarrow{\alpha} X/u(A) \xrightarrow{\beta} Y/k(A) \xrightarrow{\gamma} Z$$

is exact at $Y/k(A)$. Exactness at $X/u(A)$ is implied by the following sequences of isomorphisms:

$$\text{Coker}(\alpha) \simeq \text{Coker}(A \rightarrow \text{Coker}(a)) \simeq \text{Coker}(A \rightarrow \text{Im}(b)) \simeq \text{Im}(\beta)$$

where the first and last equalities are obtained by definition of the maps α and β . This concludes the proof. \square

Proof of Proposition 14. The first part of the proof relies on constructing the graded vector space T_{\bullet} by taking the quotient of the free graded Lie algebra $F_{\bullet} = \text{Free}(V[1])$ by a graded ideal $K_{\bullet} = \bigoplus_{i \geq 2} K_{-i}$. More precisely, for every $i \geq 2$:

$$T_{-i} = F_{-i} / K_{-i}$$

and $T_{-1} = V[1]$ of course. This will be done by using Lemma 16, where we justify that the lines are exact by invoking Lemma 15.

Given this strategy, let us first define T_{-2} as a quotient of $\Lambda^2(F_{-1}) \simeq S^2(V)$. The kernel $\text{Ker}(\{ \cdot, \cdot \})$ of the symmetric bracket on V is a subspace of $S^2(V)$; as seen from Equation (33), it is a \mathfrak{h} -module but not necessarily a \mathfrak{g} -module. However, it admits \mathfrak{g} -submodules, and among them one is the biggest, that we denote by K . We then set T_{-2} as the following quotient:

$$T_{-2} := (S^2(V) / K) [2]$$

Here, the desuspension operator [1] is applied twice so that elements of T_{-2} are considered to carry degree -2 . Identifying $\Lambda^2(F_{-1})$ and $S^2(V)[2]$, and setting $K_{-2} = K[2]$, the vector space T_{-2} can moreover be identified with:

$$T_{-2} = F_{-2}/K_{-2}$$

since $\Lambda^2(F_{-1})$ is by construction canonically isomorphic to F_{-2} . This vector space is a \mathfrak{g} -module, inheriting this structure from the canonical quotient map $p : F_{-2} \rightarrow T_{-2}$. Recalling that $F_{-1} = T_{-1} = V[1]$, let us define $q_{-2} : \Lambda^2(T_{-1}) \rightarrow T_{-2}$ to be the following composite map, which is by construction \mathfrak{g} -equivariant:

$$\begin{array}{ccc} & & F_{-2} \\ & \nearrow -d_2 & \downarrow 2p \\ \Lambda^2(T_{-1}) & \xrightarrow{q_{-2}} & T_{-2} \end{array}$$

Remark 6. *This choice of T_{-2} satisfies all required conditions: it is a \mathfrak{g} -module and it is zero if and only if V is a Lie algebra, since then $\{\cdot, \cdot\} = 0$ and $K = S^2(V)$. In particular, it is precisely for this last reason that we did not pick up any \mathfrak{g} -submodule of $\text{Ker}(\{\cdot, \cdot\})$, but only the biggest such.*

Let us now use Lemmas 15 and 16 to define T_{-3} . Since K_{-2} is a submodule of F_{-2} , there is a canonical embedding $u_{-3} : F_{-1} \otimes K_{-2} \rightarrow F_{-1} \wedge F_{-2}$. This makes the following diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & F_{-1} \otimes K_{-2} & \xrightarrow{\text{id}} & F_{-1} \otimes K_{-2} & \longrightarrow & 0 \\ & & \downarrow u_{-3} & & \downarrow k_{-3} & & \\ \Lambda^3(F_{-1}) & \xrightarrow{-d_3} & F_{-1} \wedge F_{-2} & \xrightarrow{-d_2} & F_{-3} & \longrightarrow & 0 \end{array}$$

commutative with exact rows (the bottom row is exact by Lemma 15). We appended a minus sign in front of the linear maps d_2 and d_3 because the definition of the Chevalley-Eilenberg differential d_2 is minus the Lie bracket defining F_{-3} (see Equation (69)). The map $k_{-3} : F_{-1} \otimes K_{-2} \rightarrow F_{-3}$ is the composite $-d_2 \circ u_{-3}$, making this diagram commutative. As a composition of \mathfrak{g} -equivariant morphisms, the map k_{-3} is itself \mathfrak{g} -equivariant, so that $\text{Im}(k_{-3})$ defines a \mathfrak{g} -submodule K_{-3} of F_{-3} .

Since $T_{-1} = F_{-1}$ the quotient of $F_{-1} \wedge F_{-2}$ by $F_{-1} \wedge K_{-2}$ is $T_{-1} \wedge T_{-2}$, so by Lemma 16 we deduce that:

$$\Lambda^3(F_{-1}) \longrightarrow T_{-1} \wedge T_{-2} \longrightarrow F_{-3}/K_{-3} \longrightarrow 0 \quad (74)$$

is an exact sequence. Here, we adopted the same convention for $T_{-1} \wedge T_{-2}$ as we did for $F_{-1} \wedge F_{-2}$, so that it can be identified with $T_{-1} \otimes T_{-2}$. We set:

$$T_{-3} := F_{-3}/K_{-3}$$

and we denote by $q_{-3} : T_{-1} \wedge T_{-2} \rightarrow T_{-3}$ the map induced from $-d_2|_{F_{-1} \wedge F_{-2}}$ by Lemma 16, i.e. the unique map such that the following diagram is commutative, with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & F_{-1} \otimes K_{-2} & \xrightarrow{\text{id}} & F_{-1} \otimes K_{-2} & \longrightarrow & 0 \\ & & \downarrow u_{-3} & & \downarrow k_{-3} & & \\ \Lambda^3(F_{-1}) & \xrightarrow{-d_3} & F_{-1} \wedge F_{-2} & \xrightarrow{-d_2} & F_{-3} & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \\ \Lambda^3(F_{-1}) & \longrightarrow & T_{-1} \wedge T_{-2} & \xrightarrow{q_{-3}} & T_{-3} & \longrightarrow & 0 \end{array}$$

The bottom vertical rows are the quotient maps induced by u_{-3} and k_{-3} . Elements of T_{-3} are considered to carry homogeneous degree -3 . By construction, T_{-3} is a \mathfrak{g} -module and q_{-3} is \mathfrak{g} -equivariant. Finally, if V is a Lie algebra, then $T_{-2} = 0$ which implies in turn that $T_{-3} = 0$ as well, as required by item 3. of Proposition 14.

The next step is not as straightforward and illustrates what happens for higher degrees; for this reason, it is worth its own treatment. Since K_{-2} (resp. K_{-3}) is a well-defined subspace of F_{-2} (resp. F_{-3}), one can define a linear map $u_{-4} : F_{-1} \otimes K_{-3} \oplus F_{-2} \otimes K_{-2} \rightarrow \Lambda^2(F_{\bullet})|_{-4}$ as the composition of the following two maps:

$$(F_{-1} \otimes K_{-3}) \oplus (F_{-2} \otimes K_{-2}) \hookrightarrow (F_{-1} \otimes F_{-3}) \oplus (F_{-2} \otimes F_{-2}) \oplus (F_{-3} \otimes F_{-1}) \longrightarrow \Lambda^2(F_{\bullet})|_{-4}$$

The first arrow is the canonical inclusion, whereas the second arrow is the quotient map defining the exterior algebra. The action of u_{-4} can be explicitly given by the following definition:

$$u_{-4}(x \otimes y) = x \wedge y \quad \text{for every } x \otimes y \in (F_{-1} \otimes K_{-3}) \oplus (F_{-2} \otimes K_{-2})$$

Then, by construction, the map u_{-4} is injective. This implies in turn that it is \mathfrak{g} -equivariant because K_{-2} and K_{-3} are \mathfrak{g} -modules. We have the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & F_{-1} \otimes K_{-3} \oplus F_{-2} \otimes K_{-2} & \xrightarrow{\text{id}} & F_{-1} \otimes K_{-3} \oplus F_{-2} \otimes K_{-2} & \longrightarrow & 0 \\ & & \downarrow u_{-4} & & \downarrow k_{-4} & & \\ \Lambda^3(F_{-1} \oplus F_{-2})|_{-4} & \xrightarrow{-d_3} & F_{-1} \wedge F_{-3} \oplus \Lambda^2(F_{-2}) & \xrightarrow{-d_2} & F_{-4} & \longrightarrow & 0 \end{array}$$

Notice that the bottom row is exact by Lemma 15. The dashed vertical arrow k_{-4} is the composite $-d_2 \circ u_{-4}$ making this diagram commutative. As a composition of \mathfrak{g} -equivariant morphisms, this map is \mathfrak{g} -equivariant. Its image is thus a \mathfrak{g} -submodule of F_{-4} , denoted K_{-4} . By construction, this space is uniquely defined. Then, recalling that $T_{-1} = F_{-1}$, one can quotient the bottom row by the respective images of u_{-4} and k_{-4} and obtain:

$$\Lambda^3(F_{-1} \oplus F_{-2})|_{-4} \longrightarrow T_{-1} \wedge T_{-3} \oplus \Lambda^2(T_{-2}) \longrightarrow F_{-4}/K_{-4} \longrightarrow 0 \quad (75)$$

We set:

$$T_{-4} := F_{-4}/K_{-4}$$

so that elements of T_{-4} are considered to carry homogeneous degree -4 . We call $q_{-4} : T_{-1} \wedge T_{-3} \oplus \Lambda^2(T_{-2}) \rightarrow T_{-4}$ the map induced from $-d_2|_{\Lambda^2(F_{\bullet})|_{-4}}$ by the quotient. T_{-4} inherits the quotient \mathfrak{g} -module structure and q_{-4} is by construction \mathfrak{g} -equivariant. By Lemma 16 the sequence (75) is exact. It implies in particular that q_{-4} is surjective. Moreover, if V is a Lie algebra, then $T_{-2} = 0$ and $T_{-3} = 0$, which implies that $T_{-4} = 0$ as desired in that case.

Now, let $n \geq 4$ and assume that the spaces T_{-i} , for $4 \leq i \leq n$, have been defined as quotients:

$$T_{-i} = F_{-i}/K_{-i}$$

where K_{-i} is the \mathfrak{g} -submodule of F_{-i} (of degree $-i$) corresponding to the canonical image of $\bigoplus_{j=1}^{i-2} F_{-j} \otimes K_{-i+j}$ in F_{-i} , as the following commutative diagram (with exact rows) shows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \bigoplus_{j=1}^{i-2} F_{-j} \otimes K_{-i+j} & \xrightarrow{\text{id}} & \bigoplus_{j=1}^{i-2} F_{-j} \otimes K_{-i+j} & \longrightarrow & 0 \\ & & \downarrow u_{-i} & & \downarrow k_{-i} & & \\ \Lambda^3\left(\bigoplus_{j=1}^{i-2} F_{-j}\right)|_{-i} & \xrightarrow{-d_3} & \Lambda^2\left(\bigoplus_{j=1}^{i-1} F_{-j}\right)|_{-i} & \xrightarrow{-d_2} & F_{-i} & \longrightarrow & 0 \end{array}$$

where the (injective) map u_{-i} is the composition of the two following maps:

$$\bigoplus_{j=1}^{i-2} F_{-j} \otimes K_{-i+j} \hookrightarrow \bigoplus_{j=1}^{i-1} F_{-j} \otimes F_{-i+j} \longrightarrow \Lambda^2(F_\bullet)|_{-i}$$

and where k_{-i} is the unique linear making the diagram commutative. We can assume that both u_{-i} and k_{-i} are \mathfrak{g} -equivariant. Moreover, if V is a Lie algebra, assume that every $T_{-i} = 0$, for $2 \leq i \leq n$.

Let us show that these data uniquely define a vector space T_{-n-1} of degree $-(n+1)$ satisfying the same conditions, one level higher. We will then use this induction to conclude the proof of Proposition (14). The finite direct sum $\bigoplus_{j=1}^{n-1} F_{-j} \otimes K_{-(n+1)+j}$ is a subspace of $\bigoplus_{j=1}^{n-1} F_{-j} \otimes F_{-(n+1)+j}$ and then, this induces a unique linear map $u_{-n-1} : \bigoplus_{j=1}^{n-1} F_{-j} \otimes K_{-(n+1)+j} \rightarrow \Lambda^2(F_\bullet)|_{-n-1}$ through the composition:

$$\bigoplus_{j=1}^{n-1} F_{-j} \otimes K_{-(n+1)+j} \hookrightarrow \bigoplus_{j=1}^{n-1} F_{-j} \otimes F_{-(n+1)+j} \longrightarrow \Lambda^2\left(\bigoplus_{i=1}^n F_{-i}\right)|_{-n-1}$$

The map u_{-n-1} is injective and a morphism of \mathfrak{g} -modules. One can then define a commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \bigoplus_{j=1}^{n-1} F_{-j} \otimes K_{-(n+1)+j} & \xrightarrow{\text{id}} & \bigoplus_{j=1}^{n-1} F_{-j} \otimes F_{-(n+1)+j} & \longrightarrow & 0 \\ & & \downarrow u_{-n-1} & & \downarrow k_{-n-1} & & \\ \Lambda^3\left(\bigoplus_{i=1}^{n-1} F_{-i}\right)|_{-n-1} & \xrightarrow{-d_3} & \Lambda^2\left(\bigoplus_{i=1}^n F_{-i}\right)|_{-n-1} & \xrightarrow{-d_2} & F_{-n-1} & \longrightarrow & 0 \end{array}$$

The dashed vertical line on the right is defined as the unique linear map – denoted k_{-n-1} – making the diagram commutative. As a composition of two \mathfrak{g} -equivariant morphisms, it is \mathfrak{g} -equivariant. The image of this map uniquely defines a \mathfrak{g} -submodule of F_{-n-1} that we call K_{-n-1} . Since by Lemma 15 the bottom row of the above diagram is exact, Lemma 16 implies that its quotient by the above row is an exact sequence:

$$\Lambda^3\left(\bigoplus_{i=1}^{n-1} F_{-i}\right)|_{-n-1} \longrightarrow \Lambda^2\left(\bigoplus_{i=1}^n T_{-i}\right)|_{-n-1} \xrightarrow{q_{-(n+1)}} F_{-n-1}/K_{-n-1} \longrightarrow 0 \quad (76)$$

We set:

$$T_{-n-1} := F_{-n-1}/K_{-n-1}$$

and elements of T_{-n-1} are considered to carry homogeneous degree $-(n+1)$. We call $q_{-n-1} : \Lambda^2(\bigoplus_{i=1}^n T_{-i})|_{-n-1} \rightarrow T_{-n-1}$ the map induced from $-d_2|_{\Lambda^2(F_\bullet)|_{-n-1}}$ by taking the quotient. By construction, T_{-n-1} is a \mathfrak{g} -module and q_{-n-1} is \mathfrak{g} -equivariant. Exactness of the sequence (76) implies that q_{-n-1} is surjective. Thus, if V is a Lie algebra, since by induction $T_{-i} = 0$ for every $2 \leq i \leq n$, we obtain that $T_{-n-1} = 0$ as well. We then have proved that the induction hypothesis is true at level $n+1$.

Remark 7. Notice that exactness of the sequence (76) is preserved if one changes each F_{-i} by T_{-i} in the term at the extreme left, so that the following sequence is also exact:

$$\Lambda^3\left(\bigoplus_{i=1}^{n-1} T_{-i}\right)|_{-n-1} \longrightarrow \Lambda^2\left(\bigoplus_{i=1}^n T_{-i}\right)|_{-n-1} \longrightarrow T_{-n-1} \longrightarrow 0 \quad (77)$$

This fact will be used in the proof of Lemma 19.

Continuing the induction provides us with a (possibly infinite) graded vector space $T_\bullet = \bigoplus_{i=1}^\infty T_{-i}$ which has the following properties:

1. Every vector space T_{-i} is a \mathfrak{g} -module;
2. $T_{-1} = V[1]$;

3. $T_{-i} = 0$ for every $i \geq 2$ if and only if V is a Lie algebra.

So, items 1. and 2. of Proposition 14 are partially shown, whereas item 3. is proved. Let us now define a graded Lie bracket on T_\bullet so that $T_\bullet = V[1] \oplus \llbracket T_\bullet, T_\bullet \rrbracket$. The graded vector space T_\bullet can be seen as a quotient of the free graded Lie algebra $F_\bullet = \text{Free}(V[1])$ by the graded \mathfrak{g} -submodule $K_\bullet = \bigoplus_{i=2}^\infty K_{-i}$. Additionally, by construction, this graded module is a Lie ideal in the free graded Lie algebra F_\bullet . This implies in particular that the graded vector space T_\bullet inherits a graded Lie algebra structure that descends from the free graded Lie algebra structure on F_\bullet .

The corresponding graded Lie bracket $\llbracket \cdot, \cdot \rrbracket$ can be made explicit: let $q : \Lambda^2(T_\bullet) \rightarrow T_\bullet$ be the degree 0 map whose restriction to $\Lambda^2(T_\bullet)_{-i}$ coincides with q_{-i} for every $i \geq 2$. This condition uniquely defines the map q . This bilinear map, being induced from the Chevalley-Eilenberg differential, satisfies the cohomological condition:

$$q^2 = 0 \quad (78)$$

Additionally, it is surjective and \mathfrak{g} -equivariant, by construction. Then, for every two homogeneous elements $x, y \in T_\bullet$, one has:

$$\llbracket x, y \rrbracket = q(x, y) \quad (79)$$

This bracket is a (degree 0) graded bracket – i.e. $\llbracket x, y \rrbracket \in F_{|x|+|y|}$, and it is graded skew symmetric:

$$\llbracket x, y \rrbracket = -(-1)^{|x||y|} \llbracket y, x \rrbracket \quad (80)$$

Moreover, it satisfies the Jacobi identity – which is equivalent to q satisfying Equation (78). Since q is surjective on T_{-i} for every $i \geq 2$, the condition $T_\bullet = V[1] \oplus \llbracket T_\bullet, T_\bullet \rrbracket$ is automatically satisfied. Item 2. of Proposition 14 is proved, there remains the last part of item 1.

Let us call $\rho_{-i} : \mathfrak{g} \otimes T_{-i} \rightarrow T_{-i}$ the map representing the inherited action of \mathfrak{g} on T_{-i} , and let $\rho : \mathfrak{g} \otimes T_\bullet \rightarrow T_\bullet$ be the degree 0 map restricting to ρ_{-i} on $\mathfrak{g} \otimes T_{-i}$. Alternatively, ρ can of course be seen as a degree 0 graded Lie algebra morphism from \mathfrak{g} to $\text{End}(T_\bullet)$. Then, since q is \mathfrak{g} -equivariant, it means that the action of \mathfrak{g} on T_\bullet is a derivation of the Lie bracket, i.e. the representation $\rho : \mathfrak{g} \rightarrow \text{End}(T_\bullet)$ actually takes values in $\text{Der}(T_\bullet)$. This concludes the proof of Proposition 14. \square

Remark 8. *The graded Lie algebra structure defined on T_\bullet is not independent of the Leibniz algebra structure on V since we defined T_{-2} as a quotient of $S^2(V)$ by the biggest \mathfrak{g} -submodule K of $\text{Ker}(\{\cdot, \cdot\}) \subset S^2(V)$. By induction, the dependence on the Leibniz structure has thus been propagated implicitly along the whole structure of T_\bullet , in particular through the maps q_{-i} . In particular, semi-strict Lie-Leibniz triples satisfy:*

$$T_{-2} = \left(S^2(V) / \text{Ker}(\{\cdot, \cdot\}) \right) [2]$$

Remark 9. *The graded Lie algebra obtained in Proposition 14 coincides with the one defined in Lemma 3.13 in [31]. This follows from: 1. the choice for T_{-1} and T_{-2} are identical in both constructions and 2. the map q in the present section is dual to the map π defined in [31]. The surjectivity of q hence corresponds to the injectivity of π , then the spaces of both hierarchies coincide. Notice that a typo has unfortunately been left in the final, published version of the afore mentioned paper [31]: in Definition 2.13, one should read $H_{\text{CE}}^2(\mathfrak{g}) = \wedge^2(\mathfrak{g}_{-1})^* / d_{\text{CE}}(\mathfrak{g}_{-2})^*$.*

3.2 Existence and uniqueness of the differential

From now on, in order to avoid any misunderstanding, we will denote by $T_{\leq -1}$ the graded vector space T_\bullet constructed in Proposition 14, and more generally we will write $T_{\leq -i} = \bigoplus_{k \geq i} T_{-k}$. The direct sum $T_\bullet \oplus \mathfrak{g}$ is thus denoted $T_{\leq 0}$ while $T_\bullet \oplus \mathfrak{g} \oplus R_\Theta[-1]$ is denoted $T_{\leq +1}$, where the space $T_{+1} = R_\Theta[-1]$ is understood as the space R_Θ , shifted by degree +1. Additionally, for brevity sometimes we will write \mathbb{T} in order to denote $T_{\leq +1}$.

This subsection is dedicated to building a (non-trivial) differential $\partial = (\partial_{-i})_{i \geq 1}$ on the graded vector space $T_{\leq -1}$ so that it eventually extends to the right as a cochain complex:

$$\dots \xrightarrow{\partial_{-3}} T_{-3} \xrightarrow{\partial_{-2}} T_{-2} \xrightarrow{\partial_{-1}} V[1] \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{-\eta(-;\Theta)} R_{\Theta}[-1]$$

The differential will be constructed precisely so that it is compatible with the graded Lie algebra structure on $T_{\leq -1}$ defined in Proposition 14. However we postpone to subsection 3.3 the proof that such objects equip the graded vector space \mathbb{T} with a differential graded Lie algebra structure. The following proposition then provides existence and unicity of the desired differential:

Proposition 17. *Let $(\mathfrak{g}, V, \Theta)$ be a Lie-Leibniz triple, and let $(T_{\leq -1}, \llbracket \cdot, \cdot \rrbracket)$ be the negatively graded Lie algebra associated to $(\mathfrak{g}, V, \Theta)$ by Proposition 14. Then there exists a unique family of degree +1 \mathfrak{h} -equivariant linear maps $\partial = (\partial_{-i} : T_{-i-1} \rightarrow T_{-i})_{i \geq 1}$ and satisfying the following conditions:*

$$\partial(\llbracket u, v \rrbracket) = 2\{u, v\} \quad (81)$$

$$\partial(\llbracket u, x \rrbracket) = \rho(\Theta(u); (x)) - \llbracket u, \partial(x) \rrbracket \quad (82)$$

$$\partial(\llbracket x, y \rrbracket) = \llbracket \partial(x), y \rrbracket + (-1)^{|x|} \llbracket x, \partial(y) \rrbracket \quad (83)$$

for every $u, v \in T_{-1}$ and $x, y \in T_{\leq -2}$. Moreover, the degree +1 linear map $\partial = (\partial_{-i} : T_{-i-1} \rightarrow T_{-i})_{i \geq 1}$ is a differential, i.e. for every $i \geq 1$,

$$\partial_{-i} \circ \partial_{-i-1} = 0. \quad (84)$$

Remark 10. *The differential defined in Proposition 17 does not turn $T_{\leq -1}$ into a differential graded Lie algebra, but when one appends $T_0 = \mathfrak{g}$ and $T_{+1} = R_{\Theta}[-1]$ to $T_{\leq -1}$ with adequate graded Lie brackets, the total graded vector space $\mathbb{T} = T_{\leq +1}$ becomes a differential graded Lie algebra, as is shown in Proposition 23.*

Proof. The proof is build on three lemmas 18, 19 and 20 involving a bilinear map $m : T_{-1} \wedge T_{\leq -1} \rightarrow T_{\leq -1}$ that is a convenient rewriting of the bilinear map $\rho(\Theta(-); -)$ appearing in the right-hand side of Equation (82). We define it component-wise, so that for every $i \geq 2$, let $m_{-i} : T_{-1} \wedge T_{-i} \rightarrow T_{-i}$ be the linear map defined by composing ρ_{-i} with Θ :

$$\begin{array}{ccc} T_{-1} \wedge T_{-i} & \xrightarrow{m_{-i}} & T_{-i} \\ \Theta \otimes \text{id} \downarrow & \nearrow \rho_{-i} & \\ \mathfrak{g} \otimes T_{-i} & & \end{array}$$

As for $i = 1$, we take $m_{-1} = 2\{\cdot, \cdot\} : \Lambda^2(T_{-1}) \rightarrow T_{-1}$, as we have the following identity:

$$m_{-1}(x, y) = \rho_{-1}(\Theta(x); y) + \rho_{-1}(\Theta(y); x) = 2\{x, y\} \quad (85)$$

where ρ_{-1} denotes the action of \mathfrak{g} on $T_{-1} = V[1]$. Then, the map m_{-i} lifts the action of \mathfrak{g} on T_{-i} to an action of V on T_{-i} and the quadratic constraint (17) implies that each m_{-i} is \mathfrak{h} -equivariant.

Let $m : \Lambda^2(T_{\leq -1}) \rightarrow T_{\leq -1}$ be the unique morphism whose restriction to $\Lambda^2(T_{\leq -1})|_{-i-1}$ is m_{-i} (for every $i \geq 1$). Therefore, on any subspace of $\Lambda^2(T_{\leq -1})|_{-i-1}$ other than $T_{-1} \wedge T_{-i}$ (for $i \geq 2$), the morphism m is the zero morphism. Both the bilinear map q defined at the end of the proof of Proposition 14 and the bilinear map m can be extended to multilinear maps on $\Lambda^p(T_{\leq -1})$ (for every $p \geq 3$) as follows:

$$q(x_1 \wedge \dots \wedge x_p) = \sum_{\sigma \in \text{Un}(2, p-2)} \epsilon^\sigma q(x_{\sigma(1)} \wedge x_{\sigma(2)}) \wedge x_{\sigma(3)} \wedge \dots \wedge x_{\sigma(p)} \quad (86)$$

$$m(x_1 \wedge \dots \wedge x_p) = \sum_{\sigma \in \text{Un}(2, p-2)} \epsilon^\sigma m(x_{\sigma(1)} \wedge x_{\sigma(2)}) \wedge x_{\sigma(3)} \wedge \dots \wedge x_{\sigma(p)} \quad (87)$$

where $\text{Un}(2, p-2)$ is the set of $(2, p-2)$ -unshuffles, and ϵ^σ has been defined in Equation (70). Then, the first Lemma is the following:

Lemma 18. *The following identity holds:*

$$m \circ q + q \circ m = 0 \quad (88)$$

That is: the following diagram is commutative, for every $i \geq 1$:

$$\begin{array}{ccc} & \Lambda^2(T_{\leq -1})_{-i-1} & \xrightarrow{q} T_{-i-1} \\ & \nearrow m & \nearrow -m \\ \Lambda^3(T_{\leq -1})_{-i-2} & \xrightarrow{q} \Lambda^2(T_{\leq -1})_{-i-2} & \end{array}$$

Proof. Let $x \wedge y \wedge z \in \Lambda^3(T_{\leq -1})_{-i-2}$. If x , y , and z have degree strictly lower than -1 , then (88) is trivially satisfied (because m and $-m$ both vanish). Then, one can assume that one element is in $T_{-1} = V[1]$, say x , which is then of degree -1 . In particular $|y| + |z| = -i - 1$. The action of q on $x \wedge y \wedge z$ is defined through formula (86):

$$q(x \wedge y \wedge z) = q(x \wedge y) \wedge z - (-1)^{|y||z|} q(x \wedge z) \wedge y + (-1)^{-|y|-|z|} q(y \wedge z) \wedge x \quad (89)$$

In the following we will define Θ to be 0 on any element whose degree is lower than -1 . This will allow us to keep track of those terms even if they formally vanish. Keeping in mind that $|x| = -1$, then applying m on both sides of Equation (89) implies that:

$$m(q(x \wedge y \wedge z)) \quad (90)$$

$$= m\left(-(-1)^{|z|(|y|-1)} z \wedge q(x \wedge y) + (-1)^{2|y||z|-|y|} y \wedge q(x \wedge z) - (-1)^{-|y|-|z|+i+1} x \wedge q(y \wedge z)\right) \quad (91)$$

$$= -(-1)^{|z|(|y|-1)} \Theta(z) \cdot q(x \wedge y) + (-1)^{|y|} \Theta(y) \cdot q(x \wedge z) - \Theta(x) \cdot q(y \wedge z) \quad (92)$$

$$= -(-1)^{|z|(|y|-1)} q\left((\Theta(z) \cdot x) \wedge y + x \wedge (\Theta(z) \cdot y)\right) \quad (93)$$

$$+ (-1)^{|y|} q\left((\Theta(y) \cdot x) \wedge z + x \wedge (\Theta(y) \cdot z)\right) \quad (94)$$

$$- q\left((\Theta(x) \cdot y) \wedge z + y \wedge (\Theta(x) \cdot z)\right) \quad (95)$$

$$= -q\left(m(x \wedge y) \wedge z - (-1)^{|y||z|} m(x \wedge z) \wedge y + (-1)^{|y|+|z|} m(y \wedge z) \wedge x\right) \quad (96)$$

$$= -q(m(x \wedge y \wedge z)) \quad (97)$$

as desired (where we used the fact that q is \mathfrak{g} -equivariant, thus in particular \mathfrak{h} -equivariant). To pass from the antepenultimate line to the penultimate one, we used the fact that $m(y, z) \neq 0$ only if neither y nor z have simultaneously their degrees strictly lower than -1 . Obviously, Equation (88) extends to the whole of $\Lambda(F_\bullet)$. \square

Going back to the proof of Proposition 17, the construction of the degree $+1$ linear endomorphism $\partial = (\partial_{-i} : T_{-i-1} \rightarrow T_{-i})_{i \geq 1}$ is made by induction. First, for $i = 1$, the definition of ∂_{-1} is a by-product of the observation that m_{-1} factors through T_{-2} :

$$\begin{array}{ccc}
& & T_{-1} \\
& \nearrow m_{-1} = 2\{\cdot, \cdot\} & \uparrow \partial_{-1} \\
\Lambda^2(T_{-1}) & \xrightarrow{q_{-2}} & T_{-2}
\end{array}$$

The map ∂_{-1} is defined as the unique linear map making the above triangle commutative, i.e. such that

$$\partial_{-1} \circ q(u, v) = m(u, v) \quad (98)$$

for every $u, v \in T_{-1}$. In particular, it is surjective on the ideal of squares \mathcal{I} and it is \mathfrak{h} -equivariant because $\{\cdot, \cdot\}$ has both of these properties. A priori ∂_{-1} is not \mathfrak{g} -equivariant because the symmetric bracket $\{\cdot, \cdot\}$ – equivalently m_{-1} – is not. By construction, Equation (98) is equivalent to Equation (81).

Now assume that we have constructed a family of \mathfrak{h} -equivariant linear maps $\partial_{(-n)} = (\partial_{-i} : T_{-i-1} \rightarrow T_{-i})_{1 \leq i \leq n}$ up to order $n \geq 1$ satisfying the following conditions: $\partial_{(-n)}$ is the unique linear degree +1 endomorphism of $T_{\leq -1}$ restricting to ∂_{-i} on T_{-i-1} for every $1 \leq i \leq n$ (in particular it vanishes on T_{-1}), extending as a graded derivation on $\Lambda^\bullet(\bigoplus_{i=1}^{n+1} T_{-i})$, and satisfying the following equation on $\Lambda^\bullet(\bigoplus_{i=1}^{n+1} T_{-i})$:

$$\partial_{(-n)} \circ q = m + q \circ \partial_{(-n)} \quad (99)$$

To define the map $\partial_{-n-1} : T_{-n-2} \rightarrow T_{-n-1}$ such that it extends Equation (99) one step further, we first define a degree +1 linear map $j_{-n-1} : \Lambda^2(T_{\leq -1})|_{-n-2} \rightarrow T_{-n-1}$ as the sum:

$$j_{-n-1} := m_{-n-1} + q_{-n-1} \circ \partial_{(-n)} \quad (100)$$

By construction the map j_{-n-1} is \mathfrak{h} -equivariant, and allows us to prove the existence and uniqueness of the linear map ∂_{-n-1} :

Lemma 19. *The map j_{-n-1} factors through T_{-n-2} ; in fact, there exists a unique \mathfrak{h} -equivariant linear map*

$$\partial_{-n-1} : T_{-n-2} \rightarrow T_{-n-1}$$

such that the following triangle is commutative:

$$\begin{array}{ccc}
& & T_{-n-1} \\
& \nearrow j_{-n-1} & \uparrow \partial_{-n-1} \\
\Lambda^2(T_{\leq -1})|_{-n-2} & \xrightarrow{q_{-n-2}} & T_{-n-2}
\end{array}$$

Proof. Let $x \in T_{-n-2}$. By surjectivity of q_{-n-2} , there exists $y \in \Lambda^2(T_{\leq -1})|_{-n-2}$ such that $x = q_{-n-2}(y)$. Then one would define:

$$\partial_{-n-1}(x) = j_{-n-1}(y) \quad (101)$$

if guaranteed that any other choice of pre-image of x does not change the result, i.e. that $j_{-n-1}(\text{Ker}(q_{-n-2})) = 0$. Thus, let $w \in \text{Ker}(q_{-n-2})$. Exactness of the sequence (77):

$$\Lambda^3(T_{\leq -1})|_{-n-2} \xrightarrow{q} \Lambda^2(T_{\leq -1})|_{-n-2} \xrightarrow{q_{-n-2}} T_{-n-2} \longrightarrow 0 \quad (102)$$

implies the equality $\text{Ker}(q_{-n-2}) = q(\Lambda^3(T_{\leq -1})|_{-n-2})$. Then there exists $\alpha \in \Lambda^3(T_{\leq -1})|_{-n-2}$ such that $q(\alpha) = w$, and by Lemma 18, we have:

$$m_{-n-1}(w) = -q \circ m(\alpha) \quad (103)$$

But by the induction hypothesis, m satisfies Equation (99) on $\Lambda^3(T_{\leq -1})|_{-n-2}$, so we obtain that:

$$m_{-n-1}(w) = -q_{-n-1} \circ \partial_{(-n)} \circ q(\alpha) + q_{-n-1} \circ q \circ \partial_{(-n)}(\alpha) = -q_{-n-1} \circ \partial_{(-n)}(w) \quad (104)$$

where we used Equation (78) to get rid of the second term on the right hand side. Hence the result: $j_{-n-1}(w) = 0$. \mathfrak{h} -equivariance of ∂_{-n-1} is straightforward because q_{-n-2} is \mathfrak{g} -equivariant and j_{-n-1} is \mathfrak{h} -equivariant, so for any $a \in \mathfrak{h}$ one has:

$$\partial_{-n-1}(a \cdot x) = \partial_{-n-1}(a \cdot q_{-n-2}(y)) = \partial_{-n-1}(q_{-n-2}(a \cdot y)) = j_{-n-1}(a \cdot y) = a \cdot (j_{-n-1}(y)) = a \cdot \partial_{-n-1}(x) \quad (105)$$

Notice that a priori ∂_{-n-1} is not \mathfrak{g} -equivariant. \square

The linear map ∂_{-n-1} defined in Lemma 19 is unique and has the advantage of extending Equation (99) to the level $-(n+1)$, by construction. That is to say, setting $\partial_{(-(n+1))}$ to be the unique linear degree +1 endomorphism of $T_{\leq -1}$ that restricts to ∂_{-i} on T_{-i-1} for every $1 \leq i \leq n+1$, it extends as a graded derivation on $\Lambda^\bullet(\bigoplus_{i=1}^{n+2} T_{-i})$, and satisfies the following equation on $\Lambda^\bullet(\bigoplus_{i=1}^{n+2} T_{-i})$:

$$\partial_{(-(n+1))} \circ q = m + q \circ \partial_{(-(n+1))} \quad (106)$$

By induction, this demonstrates the existence and the uniqueness of a degree +1 map ∂ satisfying the following identity on $\Lambda(T_{\leq -1})$:

$$\partial \circ q = m + q \circ \partial \quad (107)$$

This equation, when restricted to $\Lambda^2(T_{\leq -1})$, is equivalent to the three Equations (81), (82) and (83). Indeed, if two elements of T_{-1} are involved in Equation (107), then the last term disappears and $m = m_{-1}$ so that we are left with Equation (98), which is equivalent to Equation (81); if only one element of T_{-1} is involved in Equation (107), then we obtain (82), while if no element of T_{-1} is involved, $m = 0$ and we obtain Equation (83). This argument proves the first part of Proposition 17.

Now we will show that this degree +1 linear map ∂ is a differential. The proof is made by induction and relies on the following lemma:

Lemma 20. *The following identity holds:*

$$m \circ \partial + \partial \circ m = 0 \quad (108)$$

That is: the following diagram is commutative for every $i \geq 1$:

$$\begin{array}{ccc} & & T_{-i} \\ & \nearrow^{-m_{-i}} & \uparrow \partial_{-i} \\ \Lambda^2(T_{\leq -1})|_{-i-1} & & T_{-i-1} \\ \uparrow \partial_{(-i)} & \nearrow m_{-i-1} & \\ \Lambda^2(T_{\leq -1})|_{-i-2} & & \end{array}$$

Proof. Let $i \geq 1$ and let $x \wedge y \in \Lambda^2(T_{\leq -1})_{-i-2}$. If neither x nor y belong to T_{-1} , then $m(x \wedge y) = 0$ and $m(\partial(x) \wedge y + (-1)^{|x|} x \wedge \partial(y)) = 0$ (because either Θ vanishes on x, y or $\Theta \circ \partial = 0$), so that commutativity of the diagram is trivially satisfied. Thus, we can assume that at least one element, say x , belongs to T_{-1} . In that case $|y| = -i - 1 < -1$. As usual we assume that Θ vanishes on elements of degree lower than -1 , and that $\partial(T_{-1}) = 0$. Then, one has:

$$-m \circ \partial(x \wedge y) = m(x \wedge \partial(y)) \quad (109)$$

$$= \Theta(x) \cdot \partial(y) - (-1)^{-i} \Theta(\partial(y)) \cdot x \quad (110)$$

$$= \partial(\Theta(x) \cdot y) \quad (111)$$

$$= \partial \circ m(x \wedge y) \quad (112)$$

as desired. Notice that we used the \mathfrak{h} -equivariance of ∂ to pass from the antepenultimate line to the penultimate one. \square

The proof of the second part of Proposition 17 is then straightforward. First, let us show that $\partial_{-1} \circ \partial_{-2} = 0$. Let $x \in T_{-3}$, then by construction there exists $y \in T_{-1} \wedge T_{-2}$ such that $\partial_{-2}(x) = j_{-2}(y)$. Then, we get:

$$\partial_{-1} \circ j_{-2}(y) = \partial_{-1} \circ m_{-2}(y) + \partial_{-1} \circ q_{-2} \circ \partial_{(-1)}(y) \quad (113)$$

$$= \partial_{-1} \circ m_{-2}(y) + m_{-1} \circ \partial_{(-1)}(y) \quad (114)$$

$$= 0 \quad (115)$$

where we used Lemma 20 to make the penultimate line vanish.

Secondly, let $n \geq 1$ and assume that Equation (84) has been shown for every $1 \leq i \leq n$. In particular the following equation is valid:

$$\partial_{(-i)} \circ \partial_{(-i-1)} = 0 \quad (116)$$

for every $1 \leq i \leq n$, where $\partial_{(-i)}$ is the unique linear degree +1 endomorphism of $T_{\leq -1}$ that restricts to ∂_{-j} on T_{-j-1} for every $1 \leq j \leq i$. We want to show that:

$$\partial_{-n-1} \circ \partial_{-n-2} = 0 \quad (117)$$

Let $x \in T_{-n-3}$. By construction, we have $\partial_{-n-2}(x) = j_{-n-2}(y)$ for some $y \in \Lambda^2(T_{\leq -1})_{-n-3}$. Then, using Lemma 20, we have:

$$\partial_{-n-1} \circ \partial_{-n-2}(x) = \partial_{-n-1} \circ j_{-n-2}(y) \quad (118)$$

$$= \partial_{-n-1} \circ (m_{-n-2} + q_{-n-2} \circ \partial_{(-n-1)})(y) \quad (119)$$

$$= (-m_{-n-1} \circ \partial_{(-n-1)} + \partial_{-n-1} \circ q_{-n-1} \circ \partial_{(-n-1)})(y) \quad (120)$$

$$= (-m_{-n-1} + \partial_{-n-1} \circ q_{-n-2}) \circ \partial_{(-n-1)}(y) \quad (121)$$

$$= q_{-n-1} \circ \partial_{(-n)} \circ \partial_{(-n-1)}(y) \quad (122)$$

$$= 0 \quad (123)$$

where we used Equation (99) between the fourth and fifth line, and the induction hypothesis Equation (116) for $i = n$ at the penultimate line. This concludes the proof of Proposition 17. \square

Before going further, we need to investigate the relationship between the linear maps $(\partial_{-i})_{i \geq 1}$, and the embedding tensor Θ . As Θ , each of the linear map ∂_{-i} is \mathfrak{h} -equivariant, but certainly not \mathfrak{g} -equivariant. The cyclic module generated by each linear map ∂_{-i} is a \mathfrak{g} -submodule of $\text{Hom}(T_{-i-1}, T_{-i})$, that we denote R_{-i} :

$$R_{-i} := \text{Span}(\partial_{-i}, a_1 \cdot (a_2 \cdot (\dots (a_m \cdot \partial_{-i}) \dots)) \mid a_1, a_2, \dots, a_m \in \mathfrak{g}) \quad (124)$$

Recall that R_Θ is itself a cyclic \mathfrak{g} -submodule of $\text{Hom}(V, \mathfrak{g})$ (see Equation (48)). Then the R_{-i} are characterized by the following observation:

Proposition 21. *For every $i \geq 1$, there exists a surjective morphism of \mathfrak{g} -modules $\mu_{-i} : R_{\Theta} \rightarrow R_{-i}$, such that:*

$$\partial_{-i} = \mu_{-i}(\Theta) \quad (125)$$

Proof. The proof is made by induction. We will show that the R_{-i} are quotients of the cyclic module R_{Θ} , so that the μ_{-i} are the quotient maps. First let us observe that, for every $i \geq 2$, as an element of $\text{Hom}(T_{-1} \wedge T_{-i}, T_{-i})$, the map m_{-i} generates a \mathfrak{g} -submodule M_{-i} of $\text{Hom}(T_{-1} \wedge T_{-i}, T_{-i})$ through the successive actions of \mathfrak{g} on m_{-i} :

$$M_{-i} := \text{Span}\left(m_{-i}, a_1 \cdot (a_2 \cdot (\dots (a_m \cdot m_{-i}) \dots)) \mid a_1, a_2, \dots, a_m \in \mathfrak{g}\right) \quad (126)$$

The representation $\rho_{-i} : \mathfrak{g} \otimes T_{-i} \rightarrow T_{-i}$ induces a morphism of \mathfrak{g} -modules $\overline{\rho_{-i}}$ from $\text{Hom}(T_{-1} \wedge T_{-i}, \mathfrak{g} \otimes T_{-i})$ to $\text{Hom}(T_{-1} \wedge T_{-i}, T_{-i})$. By construction, this homomorphism satisfies:

$$m_{-i} = \overline{\rho_{-i}}(\Theta \wedge \text{id}_{T_{-i}}) \quad (127)$$

Here, the right hand side is the composition of $\overline{\rho_{-i}}$ with the following injective linear map:

$$\begin{array}{ccc} \text{Hom}(T_{-1}, \mathfrak{g}) & \longrightarrow & \text{Hom}(T_{-1} \wedge T_{-i}, \mathfrak{g} \otimes T_{-i}) \\ \alpha & \longmapsto & \alpha \wedge \text{id}_{T_{-i}} \end{array}$$

This composition of maps yields a morphism of \mathfrak{g} -modules $\nu_{-i} = \overline{\rho_{-i}} \circ \alpha : \text{Hom}(T_{-1}, \mathfrak{g}) \rightarrow \text{Hom}(T_{-1} \wedge T_{-i}, T_{-i})$ sending Θ onto m_{-i} :

$$m_{-i} = \nu_{-i}(\Theta) \quad (128)$$

In particular, since ν_{-i} is \mathfrak{g} -equivariant, the action of \mathfrak{g} on m_{-i} depends only on the action of \mathfrak{g} on Θ , through ν_{-i} . In particular, it sends R_{Θ} onto M_{-i} , which is then isomorphic to a quotient of R_{Θ} . This line of argument is still valid for $i = 1$, where $m_{-1} = 2\{\cdot, \cdot\}$.

Initialization: Let us now turn to the maps ∂_{-i} : assume that we have built the differential $\partial = (\partial_{-i} : T_{-i-1} \rightarrow T_{-i})_{i \geq 1}$ as in Proposition 17. Given that the map $q_{-2} : T_{-1} \wedge T_{-1} \rightarrow T_{-2}$ is surjective and \mathfrak{g} -equivariant, and that $m_{-1} = \partial_{-1} \circ q_{-2}$, it follows that q_{-2} induces a morphism of \mathfrak{g} -modules $\overline{q_{-2}} : M_{-1} \rightarrow R_{-1}$, sending m_{-1} onto ∂_{-1} . Pre-composing it with ν_{-1} induces a homomorphism of \mathfrak{g} -modules $\mu_{-1} : R_{\Theta} \rightarrow R_{-1}$:

$$\begin{array}{ccccc} R_{\Theta} & \xrightarrow{\nu_{-1}} & M_{-1} & \xrightarrow{\overline{q_{-2}}} & R_{-1} \\ & \searrow & & \nearrow & \\ & & \mu_{-1} & & \end{array}$$

In particular the composition of maps implies that:

$$\partial_{-1} = \mu_{-1}(\Theta) \quad (129)$$

which, by definition of R_{-1} , implies that μ_{-1} is onto. Moreover, since $\text{Ker}(\mu_{-1})$ is a \mathfrak{g} -submodule of R_{Θ} then R_{-1} is isomorphic, as a \mathfrak{g} -module, to the quotient $R_{\Theta} / \text{Ker}(\mu_{-1})$.

Inductive step: Let $n \geq 1$ and assume that for every $1 \leq i \leq n$ there exists a surjective homomorphism of \mathfrak{g} -modules $\mu_{-i} : R_{\Theta} \rightarrow R_{-i}$, such that:

$$\partial_{-i} = \mu_{-i}(\Theta) \quad (130)$$

Let us prove that this is still the case one level higher.

The bilinear map j_{-n-1} , as an element of $\text{Hom}(\Lambda^2(T_{\leq -1})|_{-n-2}, T_{-n-1})$, generates a \mathfrak{g} -submodule of $\text{Hom}(\Lambda^2(T_{\leq -1})|_{-n-2}, T_{-n-1})$ that we call S_{-n-1} :

$$S_{-n-1} := \text{Span}\left(j_{-n-1}, a_1 \cdot (a_2 \cdot (\dots (a_m \cdot j_{-n-1}) \dots))\right) \Big|_{a_1, a_2, \dots, a_m \in \mathfrak{g}} \quad (131)$$

By the induction hypothesis and Equation (128) at level $-n-1$, the right hand side of Equation (100) consists of a set of terms, each of which is the image of Θ through a \mathfrak{g} -equivariant map. Then it means that there exists a morphism of \mathfrak{g} -modules $s_{-n-1} : R_\Theta \rightarrow S_{-n-1}$ such that:

$$j_{-n-1} = s_{-n-1}(\Theta) \quad (132)$$

By definition of S_{-n-1} , this morphism is surjective.

Now given that the map $q_{-n-2} : \Lambda^2(T_{\leq -1})|_{-n-2} \rightarrow T_{-n-2}$ is surjective and \mathfrak{g} -equivariant, and that $\partial_{-n-1} \circ q_{-n-2} = j_{-n-1}$ by Lemma 19, it induces a morphism of \mathfrak{g} -modules $\overline{q_{-n-2}} : S_{-n-1} \rightarrow R_{-n-1}$, such that $\overline{q_{-n-2}}(j_{-n-1}) = \partial_{-n-1}$. Pre-composing it with s_{-n-1} induces in turn a homomorphism of \mathfrak{g} -modules $\mu_{-n-1} : R_\Theta \rightarrow R_{-n-1}$:

$$\begin{array}{ccccc} R_\Theta & \xrightarrow{s_{-n-1}} & S_{-n-1} & \xrightarrow{\overline{q_{-n-2}}} & R_{-n-1} \\ & \searrow & & \nearrow & \\ & & \mu_{-n-1} & & \end{array}$$

The composition of maps implies that:

$$\partial_{-n-1} = \mu_{-n-1}(\Theta) \quad (133)$$

which, by definition of R_{-n-1} , implies that μ_{-n-1} is onto. Moreover, since $\text{Ker}(\mu_{-n-1})$ is a \mathfrak{g} -submodule of R_Θ then R_{-n-1} is isomorphic, as a \mathfrak{g} -module, to the quotient $R_\Theta / \text{Ker}(\mu_{-n-1})$. This concludes the proof of Proposition 21. \square

Recall what is it we know so far. Let $(\mathfrak{g}, V, \Theta)$ be a Lie-Leibniz triple. Proposition 14 uniquely associates to this Lie-Leibniz triple a negatively graded vector space $T_{\leq -1}$ equipped with a graded Lie algebra structure. Then, Proposition 17 has shown that the complex $T_{\leq -1}$ can be equipped with a differential $\partial = (\partial_{-i} : T_{-i-1} \rightarrow T_{-i})_{i \geq 1}$ which is almost compatible with the graded Lie bracket, see Equations (81)- (83). We will now extend this differential and the Lie bracket to the graded vector space $\mathbb{T} = T_{\leq -1} \oplus \mathfrak{g} \oplus R_\Theta[-1]$, making it a differential graded Lie algebra. The construction is precisely made so that when the Lie-Leibniz triple is a Lie algebra crossed module or more generally when V is a Lie algebra, one finds the differential graded Lie algebra associated to it as in Proposition 13.

3.3 The differential graded Lie algebra structure on \mathbb{T}

The present subsection is dedicated to showing that the graded Lie bracket on $T_{\leq -1}$ defined in Proposition 14 and the differential defined in Proposition 17 can be extended to the whole of $\mathbb{T} = T_{\leq -1} \oplus \mathfrak{g} \oplus R_\Theta[-1]$ in such a way that it forms a differential graded Lie algebra.

Let $(\mathfrak{g}, V, \Theta)$ be a Lie-Leibniz triple, and let $T_{\leq -1}$ be the negatively graded Lie algebra associated to $(\mathfrak{g}, V, \Theta)$ by Proposition 14. Recall that $T_0 = \mathfrak{g}$ and $T_{+1} = R_\Theta[-1]$. The differential defined on $T_{\leq -1}$ by Proposition 17 straightforwardly extends to T_{-1} and T_0 by setting:

$$\partial_0 = \Theta : T_{-1} \rightarrow T_0 \quad \text{and} \quad \partial_{+1} = -\eta(-, \Theta) : T_0 \rightarrow T_{+1} \quad (134)$$

We indeed have $\partial_0 \circ \partial_{-1} = 0$ because the subspace $\text{Im}(\partial_{-1}) = \mathcal{I}$ is by construction included in $\text{Ker}(\Theta)$, while $\partial_{+1} \circ \partial_0 = 0$ because Θ is $\text{Im}(\Theta)$ -equivariant, by the quadratic constraint (17). For convenience, we denote the

extended differential ∂ as well. Now, let us extend the graded Lie algebra structure defined in Proposition 14 to the higher levels $T_0 = \mathfrak{g}$ and $T_{+1} = R_\Theta[-1]$. For every $a, b \in \mathfrak{g}$ and $x \in T_{\leq -1}$, let us set:

$$[[a, x]] = a \cdot x \quad \text{and} \quad [[x, a]] = -a \cdot x, \quad (135)$$

$$[[a, b]] = [a, b], \quad (136)$$

where $[\cdot, \cdot]$ is the Lie bracket on \mathfrak{g} . The first result is the following:

Lemma 22. *$(T_{\leq 0}, [[\cdot, \cdot]])$ is a graded Lie algebra.*

Proof. The Jacobi identity for the Lie bracket on T_0 is automatically satisfied because \mathfrak{g} is a Lie algebra. Then, one need only check that the bracket defined in Equation (135) and the graded Lie bracket defined in Equation (79) are compatible, in the sense that they satisfy the two Jacobi identities:

$$[[a, [[b, x]]]] = [[[[a, b], x]] + [[b, [[a, x]]]] \quad (137)$$

$$[[a, [[x, y]]]] = [[[[a, x], y]] + [[x, [[a, y]]]] \quad (138)$$

for any $a, b \in \mathfrak{g}$ and $x, y \in T_{\leq -1}$. The first equation encodes the fact that $T_{|x|}$ is a \mathfrak{g} -module:

$$a \cdot (b \cdot x) = [a, b] \cdot x + b \cdot (a \cdot x) \quad (139)$$

so it is automatically satisfied. Recalling that on $T_{\leq -1}$ the bracket corresponds to the family of maps $q = (q_{-i} : \Lambda^2(T_{\leq -1})|_{-i} \rightarrow T_{-i})_{i \leq -2}$, Equation (138) reads:

$$a \cdot q(x, y) = q(a \cdot x, y) + q(x, a \cdot y) \quad (140)$$

which corresponds to the \mathfrak{g} -equivariance of the map q , which is satisfied by construction. \square

Remark 11. *The graded Lie algebra $(T_{\leq 0}, [[\cdot, \cdot]])$, when equipped with the differential $\partial = (\partial_{-i} : T_{-i-1} \rightarrow T_{-i})_{i \geq 0}$ is not a differential graded Lie algebra. Indeed, for any $a \in \mathfrak{g}$ and $x \in T_{\leq -1}$, the Leibniz rule reads:*

$$\partial([a, x]) = [[\partial(a), x]] + [[a, \partial(x)]] = [[a, \partial(x)]] \quad (141)$$

as the restriction of the differential on \mathfrak{g} is (in the present case) the zero map. The difference between the right-hand side and the left-hand side is thus zero, meaning that we have the following identity:

$$(a \cdot \partial)(x) = [[a, \partial(x)]] - \partial([a, x]) = 0 \quad (142)$$

for every $a \in \mathfrak{g}$. But this is in contradiction with the fact that the maps $\partial_{-i} : T_{-i-1} \rightarrow T_{-i}$ are certainly not \mathfrak{g} -equivariant. Equation (142) would be only valid on \mathfrak{h} , implying in turn that $T_{\leq -1} \oplus \mathfrak{h}$ is a differential graded Lie algebra. However, this latter partial result [31], is quite unsatisfying since one loses most data from \mathfrak{g} .

We will now extend the graded Lie algebra structure to $T_{+1} = R_\Theta[-1]$ and check at the same time that it forms a graded Lie algebra structure on \mathbb{T} which is compatible with the differential ∂ . Recalling that the generators of R_Θ are of the form $\Theta_{a_1 a_2 \dots a_m} = a_1 \cdot (a_2 \cdot (\dots (a_m \cdot \Theta) \dots))$ for some $a_1, a_2, \dots, a_m \in \mathfrak{g}$, we set the following definitions for the bracket involving $T_{+1} = R_\Theta[-1]$:

$$[[\Theta, x]] = \partial(x) \quad \text{and} \quad [[x, \Theta]] = -(-1)^{|x|} \partial(x) \quad (143)$$

$$[[\Theta_{a_1 a_2 \dots a_m}, x]] = a_1 \cdot [[\Theta_{a_2 \dots a_m}, x]] - [[\Theta_{a_2 \dots a_m}, a_1 \cdot x]] \quad (144)$$

$$[[x, \Theta_{a_1 a_2 \dots a_m}]] = -(-1)^{|x|} [[\Theta_{a_1 a_2 \dots a_m}, x]] \quad (145)$$

$$[[T_{+1}, T_{+1}]] = 0 \quad (146)$$

for every homogeneous element $x \in T_{\leq 0}$ and every $a_1, \dots, a_m \in \mathfrak{g}$. As a side remark, be aware that physicists do not necessarily impose that $\llbracket T_{+1}, T_{+1} \rrbracket = 0$, see Section 3.4. of [4].

Notice that Equations (143) and (144) need a bit of explanation. Indeed, Θ is a linear map of vector spaces whereas ∂ is a family of linear maps, which do not necessarily even belong to the same \mathfrak{g} -modules. However, we have seen in the proof of Proposition 21 that each cyclic \mathfrak{g} -module R_{-i} is a quotient of R_Θ , through a quotient map μ_{-i} sending Θ on ∂_{-i} , see Equation (125). We can straightforwardly extend Proposition 21 to ∂_0 and ∂_{+1} . Since $\partial_0 = \Theta$, we know that $R_0 = R_\Theta$ so that we set $\mu_0 = \text{id}_{R_\Theta}$. Next, we define R_{+1} to be the cyclic \mathfrak{g} -submodule of $\text{Hom}(\mathfrak{g}, R_\Theta)$ generated by ∂_{+1} , so that the \mathfrak{g} -equivariant linear map $-\eta$ sending Θ to ∂_{+1} induces a morphism of \mathfrak{g} -modules $\mu_{+1} : R_\Theta \rightarrow R_{+1}$ such that $\mu_{+1}(\Theta) = \partial_{+1}$. Then one can make sense of Equation (143) by understanding that $\llbracket \Theta, - \rrbracket$, evaluated on a homogeneous element x , actually corresponds to $\mu_{|x|+1}(\Theta)(x) = \partial_{|x|+1}(x)$. Moreover, \mathfrak{g} -equivariance of $\mu_{|x|+1}$ allows to make sense of Equation (144) as:

$$\llbracket \Theta_{a_1 a_2 \dots a_m}, x \rrbracket = a_1 \cdot (a_2 \cdot (\dots (a_m \cdot \partial) \dots))(x) \quad (147)$$

The right-hand side of Equation (147) could have been alternatively found by iterating Equation (144) m times on the left-hand side of Equation (147).

From Equations (143)-(146), we can deduce the following identities:

$$\partial(\llbracket \Theta_a, x \rrbracket) = -\llbracket \Theta_a, \partial(x) \rrbracket \quad \text{and} \quad a \cdot (\partial)(\llbracket \Theta, x \rrbracket) = -\llbracket \Theta, a \cdot (\partial)(x) \rrbracket \quad (148)$$

Then, Equations (148) imply that the following identities are equivalent:

$$\partial(\llbracket \Theta_{ab}, x \rrbracket) = -\llbracket \Theta_{ab}, \partial(x) \rrbracket \quad \Longleftrightarrow \quad a \cdot (\partial)(\llbracket \Theta_b, x \rrbracket) = -\llbracket \Theta_b, a \cdot (\partial)(x) \rrbracket \quad (149)$$

$$\Longleftrightarrow \quad b \cdot (a \cdot (\partial))(\llbracket \Theta, x \rrbracket) = -\llbracket \Theta, b \cdot (a \cdot (\partial))(x) \rrbracket \quad (150)$$

However, none of them is deducible from existing equations. Then, one is free to chose these identities to hold, implying that at the next level the following are equivalent:

$$\partial(\llbracket \Theta_{abc}, x \rrbracket) = -\llbracket \Theta_{abc}, \partial(x) \rrbracket \quad \Longleftrightarrow \quad a \cdot (\partial)(\llbracket \Theta_{bc}, x \rrbracket) = -\llbracket \Theta_{bc}, a \cdot (\partial)(x) \rrbracket \quad (151)$$

$$\Longleftrightarrow \quad b \cdot (a \cdot (\partial))(\llbracket \Theta_c, x \rrbracket) = -\llbracket \Theta_c, b \cdot (a \cdot (\partial))(x) \rrbracket \quad (152)$$

$$\Longleftrightarrow \quad c \cdot (b \cdot (a \cdot (\partial))) (\llbracket \Theta, x \rrbracket) = -\llbracket \Theta, c \cdot (b \cdot (a \cdot (\partial)))(x) \rrbracket \quad (153)$$

being understood that none of them is deducible from existing equations. Assuming that any – equivalently, all – of them hold, we then find another set of equivalent identities at the next level, etc. By induction, we are thus free (and led) to assume the following compatibility condition between the bracket and the differential, generalizing the left-hand sides of Lines (148), (149) and (151) to any level:

$$\forall u \in T_{+1}, v \in T_{\leq -1} \quad \partial(\llbracket u, v \rrbracket) = -\llbracket u, \partial(v) \rrbracket \quad (154)$$

It is now time to check the compatibility of the extended bracket and the extended differential on \mathbb{T} . Recall that the graded bracket $\llbracket \cdot, \cdot \rrbracket$ is defined by Proposition 14 and Equations (135), (136), (143), (144), (145) and (146), while the differential is defined in Proposition 17 and extended to T_0 and T_{+1} by Equations (134). Additionally, we have required that these two objects satisfy Equation (154). Then, under these assumptions, we have the following important result, which generalizes Proposition 13 to any Lie-Leibniz triple:

Proposition 23. $(\mathbb{T}, \llbracket \cdot, \cdot \rrbracket, \partial)$ is a differential graded Lie algebra.

Proof. The proof is not much different than that of Proposition 13 because most problems only arise in the interaction between elements at level $+1$ and 0 . We already know that $\partial^2 = 0$ by construction, and that the

Jacobi identity is satisfied on $T_{\leq 0}$ by Lemma 22. First let us prove that the bracket and the differential are compatible, i.e. that they satisfy the Leibniz rule:

$$\partial(\llbracket u, v \rrbracket) = \llbracket \partial(u), v \rrbracket + (-1)^{|u|} \llbracket u, \partial(v) \rrbracket \quad (155)$$

for every $u, v \in \mathbb{T}$. We have several cases to handle:

1. If $u, v \in T_{+1}$, or if $u \in T_{+1}$ and $v \in T_0$ (or conversely) the identity is trivial as a consequence of Equation (146) and because $\partial|_{T_{+1}}$ is supposed to be the zero map.
2. Let $u \in T_{+1}$ and $v \in T_{\leq -1}$. Then, the Leibniz rule for u and v is nothing but Equation (154).
3. If $u, v \in T_0 = \mathfrak{g}$ then, recalling that $\partial|_{T_0} = -\eta(-; \Theta)$, Equation (155) is equivalent to the following:

$$-[u, v] \cdot \Theta = v \cdot (u \cdot \Theta) - u \cdot (v \cdot \Theta) \quad (156)$$

which corresponds to the fact that R_Θ is a \mathfrak{g} -module.

4. If $u \in T_0$ and $v \in T_{-1}$, then Equation (155) becomes:

$$\partial(u \cdot v) = \llbracket -u \cdot \Theta, v \rrbracket + u \cdot \partial(v) \quad (157)$$

This equation is equivalent to:

$$\llbracket u \cdot \Theta, v \rrbracket = u \cdot (\partial(v)) - \partial(u \cdot v) \quad (158)$$

which is nothing but Equation (144) for $\Theta_u = u \cdot \Theta$.

5. If $u, v \in T_{\leq -1}$, then we have several subcases (we will implicitly use Equation (79) in this part):
- 5.a. If $u, v \in T_{-1}$, recalling that $\partial_0 = \Theta$, Equation (155) reads:

$$\partial_{-1}(q_{-2}(u, v)) = \Theta(u) \cdot v + \Theta(v) \cdot u \quad (159)$$

This equation is nothing but Equation (81).

- 5.b. If $u \in T_{-1}$ and $v \in T_{\leq -2}$, then Equation (155) becomes:

$$\partial(q(u, v)) = \Theta(u) \cdot v - q(u, \partial(v)) \quad (160)$$

This equation can be rewritten as:

$$\Theta(u) \cdot v = \partial(q(u, v)) + q(u, \partial(v)) \quad (161)$$

which, given the definition of $m_{|v|}$, is nothing but Equation (82).

- 5.c. If $u, v \in T_{\leq -2}$, then Equation (155) can be rewritten as:

$$\partial(q(u, v)) = q \circ \partial(u \wedge v) \quad (162)$$

which is, once the term on the right hand side is transported to the left hand side, nothing but Equation (83). Hence the three subcases of item 5., gathered together, are equivalent to the identities (81), (82), and (83). This result was expected since by construction the bilinear map $m : \Lambda^2(T_{\leq -1}) \rightarrow T_{\leq -1}$ is the composition of ∂_0 with the graded Lie bracket. To conclude, the Leibniz rule is thus satisfied on the whole of \mathbb{T} .

Now let us check the Jacobi identity for the graded Lie bracket. For notational purposes, recall that for $u, v, w \in \mathbb{T}$, the Jacobi identity reads:

$$\llbracket u, \llbracket v, w \rrbracket \rrbracket = \llbracket \llbracket u, v \rrbracket, w \rrbracket + (-1)^{|u||v|} \llbracket v, \llbracket u, w \rrbracket \rrbracket \quad (163)$$

We already know by Lemma 22 that the graded Jacobi identity is satisfied on $T_{\leq 0}$. Moreover it is trivially satisfied on T_{+1} because the bracket is zero on this space. Then we need only check the cases for which one

or two terms belong to T_{+1} , that is to say: **I.** $u \in T_{+1}$ and $v, w \in T_{\leq 0}$, and **II.** $u, v \in T_{+1}$ and $w \in T_{\leq 0}$. It is sufficient to check both of these Jacobi identities on the generators of T_{+1} .

I. First notice that if v or w belongs to $T_0 = \mathfrak{g}$ then Equation (163) is Equation (144). Thus one can restrict oneself to the situation where $v, w \in T_{\leq -1}$. The proof is made by induction. First assume that $u = \Theta$. Then, since $[\Theta, -] = \partial$ (see Equation (143)), Equation (163) becomes:

$$\partial([v, w]) = [\partial(v), w] + (-1)^{|v|}[v, \partial(w)] \quad (164)$$

which is Equation (155) for $v, w \in T_{\leq -1}$, and has been proved to be satisfied in item 5. above. Therefore, assume that $u = \Theta_a$ for some $a \in \mathfrak{g}$. Then, using Equation (144), one obtains:

$$[\Theta_a, [v, w]] = a \cdot [\Theta, [v, w]] - [\Theta, a \cdot [v, w]] \quad (165)$$

$$= a \cdot ([[\Theta, v], w] + (-1)^{|v|}[v, [\Theta, w]]) - [\Theta, [a \cdot v, w] - [v, a \cdot w]] \quad (166)$$

$$= [a \cdot [\Theta, v], w] + [[\Theta, v], a \cdot w] + (-1)^{|v|}([a \cdot v, [\Theta, w]] + [v, a \cdot [\Theta, w]]) \quad (167)$$

$$\begin{aligned} & - [[\Theta, a \cdot v], w] - (-1)^{|v|}[a \cdot v, [\Theta, w]] - [[\Theta, v], a \cdot w] - (-1)^{|v|}[v, [\Theta, a \cdot w]] \\ & = [[\Theta_a, v]w] + (-1)^{|v|}[v, [\Theta_a, w]] \end{aligned} \quad (168)$$

where, from the first line to the second line we used Equation (164) on the term on the left, and Lemma 22 on the term on the right; from the second to the third line, we used Lemma 22 on the first term, and Equation (164) on the second one; the transition from the third line to the fourth and last one results from cancellation of four terms, and the use of Equation (144).

Now let $n \geq 1$ and assume that Equation (163) has been shown for every term $u \in T_{+1}$ of the form $\Theta_{a_1 \dots a_n}$, and every $v, w \in T_{\leq -1}$. Chose $n+1$ elements $a_1, \dots, a_{n+1} \in \mathfrak{g}$, then the proof of the following identity:

$$[\Theta_{a_1 \dots a_{n+1}}, [v, w]] = [[\Theta_{a_1 \dots a_{n+1}}, v]w] + (-1)^{|v|}[v, [\Theta_{a_1 \dots a_{n+1}}, w]] \quad (169)$$

relies on the same step as for Equations (165)-(168), except that one uses the induction hypothesis in place of Equation (164). This proves the first case.

II. Now assume that $u, v \in T_{+1}$. Then, by Equation (146), the graded Jacobi identity reduces to:

$$[u, [v, w]] = -[v, [u, w]] \quad (170)$$

Notice that if $w \in T_0 = \mathfrak{g}$, then it is automatically satisfied because both $[v, w]$ and $[u, w]$ belong to T_{+1} . Hence one can assume that $w \in T_{\leq -1}$. The proof is made by induction as well. It is sufficient to check Equation (170) on the generators of T_{+1} . Let v be a fixed element of T_{+1} . If $u = \Theta$ then Equation (170) corresponds to Equation (154). Next, let $u = \Theta_a$ for some $a \in \mathfrak{g}$. Then, following the same lines of argument as for Equations (165)-(167) and recalling that $[T_{+1}, T_{+1}] = 0$, one obtains the following identity:

$$[\Theta_a, [v, w]] = -([a \cdot v, [\Theta, w]] + [v, a \cdot [\Theta, w]]) + [a \cdot v, [\Theta, w]] + [v, [\Theta, a \cdot w]] \quad (171)$$

The right hand side can be recast as $[v, [\Theta_a, w]]$, which is the desired result. Notice that in the present context, where both u and v belong to T_{+1} , the use of Equation (154) was crucial to pass from the left-hand side of Equation (171) to the right-hand side.

Now let $n \geq 1$ and assume that Equation (170) has been proved for every u of the form $\Theta_{a_1 \dots a_n}$. Then pick some $a_1, \dots, a_{n+1} \in \mathfrak{g}$ so, using Equation (144), one has:

$$[\Theta_{a_1 \dots a_{n+1}}, [v, w]] = a_1 \cdot [\Theta_{a_2 \dots a_{n+1}}, [v, w]] - [\Theta_{a_2 \dots a_{n+1}}, a_1 \cdot [v, w]] \quad (172)$$

$$= a_1 \cdot (-[v, [\Theta_{a_2 \dots a_{n+1}}, w]]) - [\Theta_{a_2 \dots a_{n+1}}, [a_1 \cdot v, w] - [v, a_1 \cdot w]] \quad (173)$$

$$= -[a_1 \cdot v, [\Theta_{a_2 \dots a_{n+1}}, w]] - [v, a_1 \cdot [\Theta_{a_2 \dots a_{n+1}}, w]] \quad (174)$$

$$\begin{aligned} & + [a_1 \cdot v, [\Theta_{a_2 \dots a_{n+1}}, w]] + [v, [\Theta_{a_2 \dots a_{n+1}}, a_1 \cdot w]] \\ & = -[v, [\Theta_{a_1 \dots a_{n+1}}, w]] \end{aligned} \quad (175)$$

where we used three times the induction hypothesis, applied to $\Theta_{a_2 \dots a_{n+1}}$. This proves that Equation (170) is satisfied at level $n + 1$. Notice that this result could have been straightforwardly obtained from the generalization of equivalences (149)-(153) at every level, justified by the fact that we assumed Equation (154) to hold. By induction this proves that Equation (170) is satisfied over the generators of T_{+1} , and thus for every $u, v \in T_{+1}$. This proves that $[\![\cdot, \cdot]\!]$ is a graded Lie bracket on \mathbb{T} . \square

The fact that the maps $(\partial_{-i})_{i \geq -1}$ generate \mathfrak{g} -modules $(R_{-i})_{i \geq -1}$ satisfying Proposition 21 is of crucial importance in the proof of Proposition (23). Indeed this fact allows us to identify the action of the differential ∂ and the adjoint action of the embedding tensor Θ , thus proving the Jacobi identities that the graded bracket has to satisfy. Without this identification, one cannot simply extend the graded Lie algebra structure defined on $T_{\leq -1}$ in Proposition 14 to the whole complex $\mathbb{T} = T_{\leq -1} \oplus \mathfrak{g} \oplus R_\Theta[-1]$, and should rather limit oneself to extend it to $T_{\leq -1} \oplus \mathfrak{h}$ only, as was emphasized in Remark 11. By overcoming the problem established in this remark, Proposition (23) is thus a great improvement to the situation, since we do not lose any data in the process. This justifies the following characterization:

Definition 24. *Given a Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$, the (differential) graded Lie algebra $(\mathbb{T}, [\![\cdot, \cdot]\!], \partial)$ defined in Proposition 23 is called the tensor hierarchy associated to $(\mathfrak{g}, V, \Theta)$.*

Remark 12. *The tensor hierarchy here defined should not be confused with the ‘tensor hierarchy algebra’ defined in [38], which a priori is not a differential graded Lie algebra, but a \mathbb{Z} -graded Lie superalgebra with a subspace at degree +1 (in our convention) accommodating all possible embedding tensors satisfying the representation constraint. Moreover, in Section 3.4 of [4], the authors explain that although the condition $[\![R_\Theta, R_\Theta]\!] = 0$ defines a completely consistent mathematical model, it is however not general enough for gauged supergravity.*

When applying this construction to specific cases of Lie-Leibniz triples we obtain the following result, whose proof is immediate:

Proposition 25. *The tensor hierarchy associated to a Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ for which V is a Lie algebra is precisely the 3-term differential graded Lie algebra obtained in Proposition 13. In particular the tensor hierarchy associated to a Lie algebra crossed module $\mathfrak{c} \xrightarrow{\Theta} \mathfrak{g}$ is the 3-term differential graded Lie algebra:*

$$\mathfrak{c} \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{0} \mathbb{R}[-1]$$

Example 7. *We take a particular case of the general Example 6. Let A be the associative algebra of 2-by-2 real matrices, i.e. $A = \mathcal{M}_{2 \times 2}(\mathbb{R})$ and the associative product \cdot is the matrix product \times . The associative algebra structure on A induces a Lie algebra structure on A where the Lie bracket $[\cdot, \cdot]_A$ is the commutator of matrices; this is $\mathfrak{gl}_2(\mathbb{R})$. Let $D : A \rightarrow A$ be the linear projection on the diagonal matrices:*

$$D : \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \longmapsto \begin{pmatrix} a_{11} & 0 \\ 0 & a_{22} \end{pmatrix} \quad (176)$$

This linear map satisfies Equation (23). Then, we can define a Leibniz product on A via Equation (25):

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \circ \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} = \begin{pmatrix} 0 & (a_{11} - a_{22})b_{12} \\ -(a_{11} - a_{22})b_{21} & 0 \end{pmatrix} \quad (177)$$

From now on, when we want to emphasize the algebraic structures equipping A , we denote by \mathfrak{g} the Lie algebra structure $(A, [\cdot, \cdot]_A)$ and by V the Leibniz algebra structure (A, \circ) . The action of \mathfrak{g} on V (equivalently, on A) is materialized via the adjoint action of the Lie bracket. As the image of D consists of the Lie subalgebra \mathfrak{h} of diagonal 2-by-2 matrices, Equation (268) shows that Equation (26) is automatically satisfied. Together with Equation (25), it implies that the triple (\mathfrak{g}, V, D) is a Lie-Leibniz triple. Since $[a \circ b - D([a, b]_A), c]_A \neq 0$,

by the discussion at the end of Example 6, this Lie-Leibniz triple is not semi-strict. Then, let us compute how \mathfrak{g} acts on the embedding tensor D . Let $a, b \in A$, if we denote by $\eta : \mathfrak{g} \rightarrow \text{End}(\text{End}(A))$ the representation of \mathfrak{g} on $\text{End}(A)$ induced by the adjoint action of \mathfrak{g} on A , then we have:

$$\eta(a; D)(b) = [a, Db]_A - D([a, b]_A) = -(b \circ a - D([b, a]_A)) = \begin{pmatrix} b_{12}a_{21} - a_{12}b_{21} & -(b_{11} - b_{22})a_{12} \\ (b_{11} - b_{22})a_{21} & b_{21}a_{12} - a_{21}b_{12} \end{pmatrix} \quad (178)$$

In particular it confirms that the embedding tensor is not \mathfrak{g} -equivariant, but that it is \mathfrak{h} -equivariant, i.e. $\text{Im}(D)$ -equivariant (the right-hand side of Equation (270) vanishes when a is a diagonal matrix).

As a consequence, the symmetric bracket $\{.,.\} : S^2A \rightarrow A$ is not \mathfrak{g} -equivariant and its kernel is not a \mathfrak{g} -module. Let us decompose S^2A into \mathfrak{g} -modules to find the biggest \mathfrak{g} -submodule K of $\text{Ker}(\{.,.\})$. A basis of A is given by the four matrices:

$$E_{11} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad E_{12} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad E_{21} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad E_{22} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \quad (179)$$

The \mathfrak{g} -action of a matrix $a = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ on each of them is given by the adjoint action $[a, .]_A$:

$$[a, E_{11}]_A = -a_{12}E_{12} + a_{21}E_{21}, \quad [a, E_{12}]_A = -a_{21}E_{11} + (a_{11} - a_{22})E_{12} + a_{21}E_{22} \quad (180)$$

$$[a, E_{22}]_A = a_{12}E_{12} - a_{21}E_{21}, \quad [a, E_{21}]_A = a_{12}E_{11} - (a_{11} - a_{22})E_{21} - a_{12}E_{22} \quad (181)$$

The matrices E_{ij} induce a basis of the 10-dimensional space S^2A via the symmetric product $E_{ij} \odot E_{kl}$. We can alternatively describe this space with degree 2 polynomials of four variables by assigning the variable X to E_{11} , Y to E_{12} , Z to E_{21} and T to E_{22} . Then the action (180), (181) of \mathfrak{g} on the matrices E_{ij} induces an action on the monomials X, Y, Z, T :

$$a \cdot X = -a_{12}Y + a_{21}Z, \quad a \cdot Y = -a_{21}X + (a_{11} - a_{22})Y + a_{21}T \quad (182)$$

$$a \cdot T = a_{12}Y - a_{21}Z, \quad a \cdot Z = a_{12}X - (a_{11} - a_{22})Z - a_{12}T \quad (183)$$

The action of $\mathfrak{g} = \mathfrak{gl}_2(\mathbb{R})$ on S^2A is induced from Equations (182) and (183) by derivation.

Since $\mathfrak{gl}_2(\mathbb{R})$ is reductive, the symmetric space $S^2A = S^2(\mathfrak{gl}_2(\mathbb{R}))$ is completely reducible into four irreducible $\mathfrak{gl}_2(\mathbb{R})$ -submodules (equivalently: $\mathfrak{sl}_2(\mathbb{R})$ -modules):

$$\begin{aligned} U_1 &= \langle XT - YZ \rangle \\ U_2 &= \langle X^2 + T^2 + XT + YZ \rangle \\ U_3 &= \langle X^2 - T^2, XY + YT, XZ + ZT \rangle \\ U_4 &= \langle Y^2, Z^2, X^2 + T^2 - 2(XT + YZ), XY - YT, XZ - ZT \rangle \end{aligned}$$

The generators of the modules form an alternative basis for S^2A because one can pass from the standard basis of S^2A to the generators of U_1, U_2, U_3 and U_4 by the following formula:

$$\begin{aligned} a_{X^2}X^2 + a_{XY}XY + a_{XZ}XZ + a_{XT}XT + a_{Y^2}Y^2 \\ + a_{YZ}YZ + a_{YT}YT + a_{Z^2}Z^2 + a_{ZT}ZT + a_{T^2}T^2 = \end{aligned} \quad (184)$$

$$\begin{aligned} & \frac{a_{XT} - a_{YZ}}{2}(XT - YZ) + \frac{2(a_{X^2} + a_{T^2}) + a_{XT} + a_{YZ}}{6}(X^2 + T^2 + XT + YZ) \\ & + \frac{a_{X^2} - a_{T^2}}{2}(X^2 - T^2) + \frac{a_{XY} + a_{YT}}{2}(XY + YT) + \frac{a_{XZ} + a_{ZT}}{2}(XZ + ZT) \\ & + a_{Y^2}Y^2 + a_{Z^2}Z^2 + \frac{a_{X^2} + a_{T^2} - a_{XT} - a_{YZ}}{6}(X^2 + T^2 - 2(XT + YZ)) \\ & + \frac{a_{XY} - a_{YT}}{2}(XY - YT) + \frac{a_{XZ} - a_{ZT}}{2}(XZ - ZT) \end{aligned}$$

We then have two equivalent ways of parametrizing S^2A , whose respective usefulness is relative to the given computation.

The symmetric product $a \odot b \in S^2A$ of two matrices can be decomposed on the polynomials in X, Y, Z, T as the following:

$$\begin{aligned} a \odot b = & a_{11}b_{11}X^2 + \frac{1}{2}(a_{11}b_{12} + b_{11}a_{12})XY + \frac{1}{2}(a_{11}b_{21} + b_{11}a_{21})XZ + \frac{1}{2}(a_{11}b_{22} + b_{11}a_{22})XT + a_{12}b_{12}Y^2 \\ & + \frac{1}{2}(a_{12}b_{21} + b_{12}a_{21})YZ + \frac{1}{2}(a_{12}b_{22} + b_{12}a_{22})YT + a_{21}b_{21}Z^2 + \frac{1}{2}(a_{21}b_{22} + b_{21}a_{22})ZT + a_{22}b_{22}T^2 \end{aligned} \quad (185)$$

By comparing the anti-diagonal matrix elements of the matrix on the right-hand side of the symmetrized version of Equation (268) to the decomposition (185), we deduce that the symmetric bracket $\{.,.\} : S^2A \rightarrow A$ is defined on an element of S^2A as:

$$\begin{aligned} & a_{X^2}X^2 + a_{XY}XY + a_{XZ}XZ + a_{XT}XT + a_{Y^2}Y^2 \\ & + a_{YZ}YZ + a_{YT}YT + a_{Z^2}Z^2 + a_{ZT}ZT + a_{T^2}T^2 \end{aligned} \xrightarrow{\{.,.\}} \begin{pmatrix} 0 & a_{XY} - a_{YT} \\ -a_{XZ} + a_{ZT} & 0 \end{pmatrix} \quad (186)$$

The elements of the left-hand side of Equation (186) that do not appear on the right-hand side live in the kernel of the symmetric bracket, which is then the 8-dimensional subspace of S^2A generated by the following vectors:

$$\text{Ker}(\{.,.\}) = \langle X^2, Y^2, Z^2, T^2, XT, YZ, XY + YT, XZ + ZT \rangle$$

The three first \mathfrak{g} -submodules U_1, U_2, U_3 are contained in $\text{Ker}(\{.,.\})$ while for the last module, only the first three generators belong to the kernel:

$$\text{Ker}(\{.,.\}) = U_1 \oplus U_2 \oplus U_3 \oplus \langle Y^2, Z^2, X^2 + T^2 - 2(XT + YZ) \rangle$$

As the action of \mathfrak{g} on the remaining generators $Y^2, Z^2, X^2 + T^2 - 2(XT + YZ)$ leaves the kernel, the biggest \mathfrak{g} -submodule of the kernel is the 5-dimensional subspace $K = U_1 \oplus U_2 \oplus U_3$. We can then identify $T_{-2} = (S^2(A)/K)[2]$, the degree -2 space of the tensor hierarchy, with $U_4[2]$ (U_4 shifted by -2).

The projection $p : S^2A \rightarrow U_4$ then induces the graded Lie bracket on $T_{-1} = A[1]$ taking values in T_{-2} by Equation (79), that is to say $\llbracket a, b \rrbracket = q_{-2}(a \odot b) = 2p(a \odot b)$. We can give an explicit formula of this bracket, thanks to the decomposition (185) and the correspondence (184):

$$\begin{aligned} \left[\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}, \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} \right] = & \frac{1}{6} \begin{pmatrix} 2a_{11}b_{11} - (a_{11}b_{22} + b_{11}a_{21}) \\ +2a_{22}b_{22} - (a_{12}b_{21} + b_{12}a_{21}) \end{pmatrix} (X^2 + T^2 - 2(XT + YZ)) \\ & + \frac{1}{2} \left((a_{11}b_{12} + b_{11}a_{12}) - (a_{12}b_{22} + b_{12}a_{22}) \right) (XY - YT) \\ & + \frac{1}{2} \left((a_{11}b_{21} + b_{11}a_{21}) - (a_{21}b_{22} + b_{21}a_{22}) \right) (XZ - ZT) \\ & + 2a_{12}b_{12}Y^2 + 2a_{21}b_{21}Z^2 \end{aligned} \quad (187)$$

The linear map $\partial_{-1} : T_{-2} \rightarrow T_{-1}$ is then defined as:

$$\begin{aligned} & pY^2 + qZ^2 \\ & + r(X^2 + T^2 - 2(XT + YZ)) \\ & + s(XY - YT) + t(XZ - ZT) \end{aligned} \xrightarrow{\partial_{-1}} \begin{pmatrix} 0 & 2s \\ -2t & 0 \end{pmatrix} \quad (188)$$

so that we indeed have $\partial_{-1}(\llbracket a, b \rrbracket) = 2\{a, b\}$. The map ∂_{-1} has a 3-dimensional kernel. The tensor hierarchy associated to the Lie-Leibniz triple (\mathfrak{g}, V, D) then begins with the following sequence:

$$\dots \longrightarrow U_4[2] \xrightarrow{\partial_{-1}} A[1] \xrightarrow{D} A \xrightarrow{-\eta(-;D)} R_D[-1]$$

where $R_D \subset \text{End}(A)$ is the \mathfrak{g} -submodule generated by D .

In order to define T_{-3} , one need to decompose the following diagram in terms of irreducible $\mathfrak{sl}_2(\mathbb{R})$ -modules (because they are the irreducible $\mathfrak{gl}_2(\mathbb{R})$ -modules):

$$\begin{array}{ccccccc}
0 & \longrightarrow & A \otimes (U_1 \oplus U_2 \oplus U_3) & \xrightarrow{\text{id}} & A \otimes (U_1 \oplus U_2 \oplus U_3) & \longrightarrow & 0 \\
& & \downarrow u_{-3} & & \downarrow k_{-3} & & \\
S^3(A) & \xrightarrow{-d_3} & A \otimes S^2(A) & \xrightarrow{-d_2} & F_{-3} & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \\
S^3(A) & \longrightarrow & A \otimes U_4 & \xrightarrow{q_{-3}} & T_{-3} & \longrightarrow & 0
\end{array} \tag{189}$$

Here, recall that lines and columns are exact sequences. For any $k \geq 0$, let us denote by V_k the irreducible $\mathfrak{sl}_2(\mathbb{R})$ -module of highest weight k , so in particular it is a $k+1$ -dimensional vector space. With this convention, V_2 denotes the 3-dimensional adjoint representation of $\mathfrak{sl}_2(\mathbb{R})$, and $U_4 = V_4$. Given the discussion above and since $\mathfrak{gl}_2(\mathbb{R}) = \mathfrak{sl}_2(\mathbb{R}) \oplus \langle I_2 \rangle = V_2 \oplus V_0$, the decomposition of the second symmetric power of $A = \mathfrak{gl}_2(\mathbb{R})$ in terms of the $\mathfrak{sl}_2(\mathbb{R})$ -modules U_1, U_2, U_3, U_4 can be rewritten as:

$$S^2(\mathfrak{gl}_2(\mathbb{R})) = S^2(V_2 \oplus V_0) = V_0 \oplus V_0 \oplus V_2 \oplus V_4 \tag{190}$$

Then, using the classical formulas for reducibility of (symmetric) powers of $\mathfrak{sl}_2(\mathbb{R})$ -modules, we have:

$$S^3(\mathfrak{gl}_2(\mathbb{R})) = 2V_0 \oplus 2V_2 \oplus V_4 \oplus V_6 \tag{191}$$

$$\mathfrak{gl}_2(\mathbb{R}) \otimes S^2(\mathfrak{gl}_2(\mathbb{R})) = 3V_0 \oplus 5V_2 \oplus 3V_4 \oplus V_6 \tag{192}$$

where the integers in front of each module denote their multiplicities. By exactness of the lines of Diagram (189), the difference between Lines (192) and (191) gives:

$$F_{-3} = V_0 \oplus 3V_2 \oplus 2V_4 \tag{193}$$

As every space in the present example is completely reducible, the module T_{-3} can be understood as a sub-module of both F_{-3} and $A \otimes U_4 = \mathfrak{gl}_2(\mathbb{R}) \otimes V_4 = V_2 \oplus 2V_4 \oplus V_6$. From this observation we deduce that:

$$T_{-3} \subset F_{-3} \cap (A \otimes U_4) = V_2 \oplus 2V_4 \tag{194}$$

Since $T_{-3} \simeq A \otimes U_4 / S^3(A)$, by carefully analyzing the intersection of $A \otimes U_4$ with $S^3(\mathfrak{gl}_2(\mathbb{R}))$, we can deduce the explicit decomposition of T_{-3} . For example, as we have the inclusion $A \otimes U_4 \subset \mathfrak{gl}_2(\mathbb{R}) \otimes S^2(\mathfrak{sl}_2)$ and since the latter tensor product contains $S^3(\mathfrak{sl}_2) = V_2 \oplus V_6$, we deduce that the term $V_2 \oplus V_6$ in $A \otimes U_4$ corresponds to $S^3(\mathfrak{sl}_2)$ so, by exactness of the bottom line in Diagram (189), V_2 cannot appear in T_{-3} . Now, there is only one module V_4 in $S^3(\mathfrak{gl}_2(\mathbb{R}))$, while there are two in $A \otimes U_4$ so T_{-3} contains at least one module V_4 , if not two. The question is then if either one of the two modules V_4 in the decomposition of $A \otimes U_4$ lies in the image of $S^3(\mathfrak{gl}_2(\mathbb{R}))$. Since the two modules V_4 in $A \otimes U_4$ are precisely those of $\mathfrak{gl}_2(\mathbb{R}) \otimes S^2(\mathfrak{sl}_2(\mathbb{R}))$, it is equivalent to asking if either one of the two submodules V_4 of $\mathfrak{gl}_2(\mathbb{R}) \otimes S^2(\mathfrak{sl}_2(\mathbb{R}))$ lies in the image of $S^3(\mathfrak{gl}_2(\mathbb{R}))$.

The following, speculative argument suggests that neither of them lie in this image. First, notice that:

$$S^3(\mathfrak{gl}_2(\mathbb{R})) = S^3(\mathfrak{sl}_2(\mathbb{R})) \oplus S^2(\mathfrak{gl}_2(\mathbb{R})) \tag{195}$$

$$\mathfrak{gl}_2(\mathbb{R}) \otimes S^2(\mathfrak{gl}_2(\mathbb{R})) = (\mathfrak{gl}_2(\mathbb{R}) \otimes S^2(\mathfrak{sl}_2(\mathbb{R}))) \oplus (\mathfrak{gl}_2(\mathbb{R}) \otimes \mathfrak{gl}_2(\mathbb{R})) \tag{196}$$

Then, comparing the decompositions (195) and (196), we deduce that the submodule $S^2(\mathfrak{gl}_2(\mathbb{R}))$ of $S^3(\mathfrak{gl}_2(\mathbb{R}))$ sits in the submodule $\mathfrak{gl}_2(\mathbb{R}) \otimes \mathfrak{gl}_2(\mathbb{R})$ of $\mathfrak{gl}_2(\mathbb{R}) \otimes S^2(\mathfrak{gl}_2(\mathbb{R}))$. Then, since $S^3(\mathfrak{sl}_2) = V_2 \oplus V_6$, the submodule

V_4 of $S^3(\mathfrak{gl}_2(\mathbb{R}))$ is necessarily in $S^2(\mathfrak{gl}_2(\mathbb{R}))$, so it seems to be sent to the submodule V_4 of $\mathfrak{gl}_2(\mathbb{R}) \otimes \mathfrak{gl}_2(\mathbb{R})$ in Line (196). From this, we conjecture that none of the two submodules V_4 of $\mathfrak{gl}_2(\mathbb{R}) \otimes S^2(\mathfrak{sl}_2(\mathbb{R}))$ – and thus, of $A \otimes U_4$ – lies in the image of $S^3(\mathfrak{gl}_2(\mathbb{R}))$. Then, Diagram (189) can be recast in the following form, where the colors mark the correspondence between the domain and the range of each map (and where the purple color means that it is both red and blue):

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \textcolor{red}{3} V_0 \oplus \textcolor{red}{4} V_2 \oplus V_4 & \xrightarrow{\text{id}} & \textcolor{red}{3} V_0 \oplus \textcolor{red}{4} V_2 \oplus V_4 & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \\
 & & \textcolor{purple}{2} V_0 \oplus \textcolor{red}{V}_0 & & & & \\
 \textcolor{blue}{2} V_0 \oplus \textcolor{blue}{2} V_2 & \longrightarrow & \oplus \textcolor{blue}{V}_2 \oplus \textcolor{blue}{V}_2 \oplus \textcolor{red}{3} V_2 & \longrightarrow & \textcolor{red}{V}_0 \oplus \textcolor{red}{3} V_2 \oplus 2 V_4 & \longrightarrow & 0 \\
 \oplus \textcolor{blue}{V}_4 \oplus \textcolor{blue}{V}_6 & & \oplus \textcolor{blue}{V}_4 \oplus 2 V_4 & & \downarrow & & \\
 & & \oplus \textcolor{blue}{V}_6 & & & & \\
 & & \downarrow & & \downarrow & & \\
 \textcolor{blue}{2} V_0 \oplus \textcolor{blue}{2} V_2 & \longrightarrow & \textcolor{blue}{V}_2 \oplus 2 V_4 \oplus \textcolor{blue}{V}_6 & \longrightarrow & 2 V_4 & \longrightarrow & 0 \\
 \oplus \textcolor{blue}{V}_4 \oplus \textcolor{blue}{V}_6 & & & & & &
 \end{array}$$

From this we deduce that $T_{-3} = (V_4 \oplus V_4)[3]$. The Lie bracket between $T_{-2} = V_4[2]$ and $T_{-1} = A[1]$ is uniquely defined as the bottom horizontal line, while the linear map $\partial_{-2} : T_{-3} \rightarrow T_{-2}$ is also uniquely defined by Lemma 19. This gives the next level of the tensor hierarchy associated to the Lie-Leibniz triple (\mathfrak{g}, V, D) :

$$\dots \longrightarrow (V_4 \oplus V_4)[3] \xrightarrow{\partial_{-2}} V_4[2] \xrightarrow{\partial_{-1}} A[1] \xrightarrow{D} A \xrightarrow{-\eta(-;D)} R_D[-1]$$

We will not provide the next space of the hierarchy as the computations become more cumbersome, but in principle there is no mathematical obstruction to this task.

3.4 Proof of Theorem 1 and open questions

Since the tensor hierarchy \mathbb{T} associated to a given Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ includes \mathfrak{g} and V as T_0 and T_{-1} , we immediately have the following result, which is the first part of Theorem 1:

Proposition 26. *The function $G : \mathbf{Lie-Leib} \rightarrow \mathbf{DGLie}_{\leq 1}$ associating a tensor hierarchy to a given Lie-Leibniz triple is injective-on-objects.*

Thus, tensor hierarchies can be seen as Lie-ifications of Lie-Leibniz triples, where no information has been lost, generalizing what occurs for Lie algebra crossed modules. We now show that, when restricted to a particular subcategory of **Lie-Leib**, this function is a functor. The following Proposition is the second part of Theorem 1:

Proposition 27. *The restriction of the function $G : \mathbf{Lie-Leib} \rightarrow \mathbf{DGLie}_{\leq 1}$ to **compLie-Leib** is a faithful functor. Moreover, **compLie-Leib** is the biggest wide subcategory of **Lie-Leib** (with respect to inclusion) such that this functorial property holds.*

Proof. Let us first prove that the assignment G is functorial on **compLie-Leib**, i.e. that a compatible morphism of Lie-Leibniz triples (see Definition 9) induces a dgLa morphism between their tensor hierarchies. Faithfulness will follow from the definition of the action of G on morphisms.

Let $(\mathfrak{g}, V, \Theta)$ and $(\mathfrak{g}', V', \Theta')$ be Lie-Leibniz triples, and let $(\mathbb{T}, \llbracket \cdot, \cdot \rrbracket, \partial)$ and $(\mathbb{T}', \llbracket \cdot, \cdot \rrbracket', \partial')$ be their associated tensor hierarchies. Let us introduce some notation: for every $i \geq 1$, let $T_{(-i)}$ be the direct sum $\bigoplus_{0 \leq j \leq i} T_{-j}$, the same convention applying to \mathbb{T}' . In the proof, ϕ_{-i} will always denote a degree 0 linear map

from T_{-i} to T'_{-i} , and $\phi_{(-i)}$ will denote the unique degree 0 linear map from $T_{(-i)}$ to $T'_{(-i)}$ that restricts to ϕ_{-i} on T_{-i} . It can be straightforwardly extended to $\Lambda^\bullet(T_{(-i)})$ as a morphism of graded algebras.

Definition of ϕ_{+1} : Let $(\varphi, \chi) : (\mathfrak{g}, V, \Theta) \rightarrow (\mathfrak{g}', V', \Theta')$ be any (compatible) morphism of Lie-Leibniz triples, and let us denote $\phi_0 = \varphi$ and $\phi_{-1} = \chi : V[1] \rightarrow V'[1]$. The compatibility of the respective graded Lie brackets on \mathbb{T} and \mathbb{T}' with ϕ_0 and ϕ_{-1} is deduced from the fact that ϕ_0 is a Lie algebra morphism and by Equation (35):

$$\llbracket \phi_0(a), \phi_0(b) \rrbracket = \phi_0(\llbracket a, b \rrbracket) \quad (197)$$

$$\llbracket \phi_0(a), \phi_{-1}(x) \rrbracket = \phi_{-1}(\llbracket a, x \rrbracket) \quad (198)$$

for every $a, b \in T_0$ and $x \in T_{-1}$. Now, let $R_\Theta \subset \text{Hom}(V, \mathfrak{g})$ and $R_{\Theta'} \subset \text{Hom}(V', \mathfrak{g}')$ the cyclic \mathfrak{g} -modules generated by Θ and Θ' , respectively (see Equation (48)). Then we define the linear map $\phi_{+1} : R_\Theta[-1] \rightarrow R_{\Theta'}[-1]$ to be the unique morphism of cyclic \mathfrak{g} -modules sending Θ to Θ' and compatible with ϕ_0 . That is to say, since Θ (resp. Θ') generates R_Θ (resp. $R_{\Theta'}$), we have:

$$\phi_{+1}(\Theta_{a_1 \dots a_n}) = \Theta'_{\varphi(a_1) \dots \varphi(a_n)} \quad (199)$$

where $\Theta_{a_1 \dots a_n} = a_1 \cdot (a_2 \cdot (\dots (a_n \cdot \Theta) \dots))$, for every $a_1, \dots, a_n \in \mathfrak{g}$. Moreover, since R_{-i} (resp. R'_{-i}) is generated by ∂_{-i} (resp. ∂'_{-i}) – see Equation (124) – the fact that $\phi_{+1}(\Theta) = \Theta'$ implies that the map ϕ_{+1} induces a morphism of cyclic \mathfrak{g} -modules from R_{-i} to R'_{-i} – also denoted ϕ_{+1} for convenience – such that:

$$\phi_{+1}(\partial_{-i}) = \partial'_{-i} \quad (200)$$

So far, there are two compatibility equations involving ϕ_{+1} and the graded Lie bracket. The first one is of the form:

$$\llbracket \phi_{+1}(\Theta), \phi_{-1}(x) \rrbracket = \phi_0(\llbracket \Theta, x \rrbracket) \quad (201)$$

for any $x \in T_{-1}$. It is automatically satisfied because it is a rewriting of Equation (34), under the convention that $\partial = \llbracket \Theta, - \rrbracket$. The second equation is:

$$\llbracket \phi_{+1}(\Theta), \phi_0(a) \rrbracket = \phi_{+1}(\llbracket \Theta, a \rrbracket) \quad (202)$$

for every $a \in T_0$. It is also satisfied because it can be equivalently rewritten as

$$-\varphi(a) \cdot \Theta' = \phi_{+1}(-a \cdot \Theta) \quad (203)$$

which is a particular case of Equation (199). Using successively Equations (199), (144) and (35), together with the fact that φ is a morphism of Lie algebras, one can show that Equations (201) and (202) imply that the following two equations hold for any generator $\Theta_{a_1 \dots a_m}$ of $T_{+1} = R_\Theta[-1]$:

$$\llbracket \phi_{+1}(\Theta_{a_1 \dots a_m}), \phi_{-1}(x) \rrbracket = \phi_0(\llbracket \Theta_{a_1 \dots a_m}, x \rrbracket) \quad (204)$$

$$\llbracket \phi_{+1}(\Theta_{a_1 \dots a_m}), \phi_0(a) \rrbracket = \phi_{+1}(\llbracket \Theta_{a_1 \dots a_m}, a \rrbracket) \quad (205)$$

Hence, the first three maps ϕ_{+1} , ϕ_0 and ϕ_{-1} are so far compatible with the graded Lie brackets on \mathbb{T} and \mathbb{T}' , respectively. Now, because of Equations (143) and (200), one can check that the identity $\partial'_0 \circ \phi_{-1} = \phi_0 \circ \partial_0$ corresponds to Equation (201), and that $\partial'_{+1} \circ \phi_0 = \phi_{+1} \circ \partial_{+1}$ corresponds to Equation (202).

Definition of ϕ_{-i} , $i \geq 2$: The proof is made by induction. Let $T_{-2} = (S^2(V)/K)[2]$ and $T'_{-2} = (S^2(V')/K')[2]$. Since the morphism $(\varphi, \chi) : (\mathfrak{g}, V, \Theta) \rightarrow (\mathfrak{g}', V', \Theta')$ is a compatible morphism, Equation (37) implies that $\chi \odot \chi$ passes to the quotient, i.e. that it induces a well-defined map $\phi_{-2} : T_{-2} \rightarrow T'_{-2}$ that renders the following prism commutative:

$$\begin{array}{ccccc}
& & T_{-2} & \xrightarrow{\phi_{-2}} & T'_{-2} \\
& \nearrow q_{-2} & \downarrow \partial_{-1} & & \nearrow q'_{-2} \\
\Lambda^2(T_{-1}) & \xrightarrow{\phi_{-1} \wedge \phi_{-1}} & \Lambda^2(T'_{-1}) & & \\
& \searrow m_{-1} & \downarrow \partial'_{-1} & & \searrow m'_{-1} \\
& & T_{-1} & \xrightarrow{\phi_{-1}} & T'_{-1}
\end{array}$$

where we have written χ under its current denomination $\phi_{-1} : T_{-1} \rightarrow T'_{-1}$. This is precisely here that we use the compatibility property of the morphism. We refer to the end of the proof for a discussion of what happens when the morphism is not compatible.

The commutative diagram proves the following two equations:

$$\partial'_{-1} \circ \phi_{-2} = \phi_{-1} \circ \partial_{-1} \quad (206)$$

$$\phi_{-2}(\llbracket x, y \rrbracket) = \llbracket \phi_{-1}(x), \phi_{-1}(y) \rrbracket \quad (207)$$

for every $x, y \in T_{-1}$. The fact that χ satisfies Equation (35) implies that ϕ_{-2} in turn satisfies:

$$\phi_{-2}(\llbracket a, x \rrbracket) = \llbracket \phi_0(a), \phi_{-2}(x) \rrbracket \quad (208)$$

for every $a \in \mathfrak{g}$ and $x \in T_{-2}$. The last compatibility equations to check are:

$$\llbracket \phi_{+1}(\Theta), \phi_{-2}(x) \rrbracket = \phi_{-1}(\llbracket \Theta, x \rrbracket) \quad (209)$$

$$\llbracket \phi_{+1}(\Theta_{a_1 \dots a_m}), \phi_{-2}(x) \rrbracket = \phi_{-1}(\llbracket \Theta_{a_1 \dots a_m}, x \rrbracket) \quad (210)$$

for every $x \in T_{-2}$. By Equations (143) and (200), the first equation is an alternative form of Equation (206). Equation (210) can be proved by using Equations (35), (144), (199), (208), as well as Equation (209). For example, for $m = 1$, one has:

$$\llbracket \phi_{+1}(\Theta_a), \phi_{-2}(x) \rrbracket = \llbracket \Theta'_{\varphi(a)}, \phi_{-2}(x) \rrbracket \quad (211)$$

$$= \varphi(a) \cdot \llbracket \Theta', \phi_{-2}(x) \rrbracket - \llbracket \Theta', \varphi(a) \cdot \phi_{-2}(x) \rrbracket \quad (212)$$

$$= \varphi(a) \cdot \phi_{-1}(\llbracket \Theta, x \rrbracket) - \llbracket \Theta', \phi_{-2}(a \cdot x) \rrbracket \quad (213)$$

$$= \phi_{-1}(a \cdot \llbracket \Theta, x \rrbracket) - \phi_{-1}(\llbracket \Theta, a \cdot x \rrbracket) \quad (214)$$

$$= \phi_{-1}(\llbracket \Theta_a, x \rrbracket) \quad (215)$$

Lets now turn to the inductive step. Assume that the linear map $\phi_{(-i)} : T_{(-i)} \rightarrow T'_{(-i)}$ is constructed up to order $i \geq 2$, and that it satisfies:

$$\partial' \circ \phi_{(-i)}(x) = \phi_{(-i)} \circ \partial(x) \quad (216)$$

$$\phi_{(-i)}(\llbracket x, y \rrbracket) = \llbracket \phi_{(-i)}(x), \phi_{(-i)}(y) \rrbracket \quad (217)$$

for every $x, y \in T_{\leq 0}$ such that $-i \leq |x| + |y| \leq 0$. Moreover, notice that for $a \in T_0 = \mathfrak{g}$, and $u \in T_{\leq 0}$, Equation (217) implies the following result:

$$\phi_{(-i)}(a \cdot u) = \phi_0(a) \cdot \phi_{(-i)}(u) \quad (218)$$

This equation extends to every element $y \in \Lambda(T_{\leq -1})$. The only remaining non-trivial equations are:

$$\llbracket \phi_{+1}(\Theta), \phi_{-i}(x) \rrbracket = \phi_{-i+1}(\llbracket \Theta, x \rrbracket) \quad (219)$$

$$\llbracket \phi_{+1}(\Theta_{a_1 \dots a_m}), \phi_{-i}(x) \rrbracket = \phi_{-i+1}(\llbracket \Theta_{a_1 \dots a_m}, x \rrbracket) \quad (220)$$

for every $x \in T_{-i}$. By Equations (143) and (200), Equation (219) is automatically satisfied because it is equivalent to Equation (216), while Equation (220) is a consequence of the former as well as (218), as can be shown by using the same kind of arguments as for Equations (211)-(215).

Let us construct $\phi_{-i-1} : T_{-i-1} \rightarrow T'_{-i-1}$ such that Equations (216) and (217) are valid one step higher, i.e. such that i is replaced by $i+1$. Then, as explained above, Equations (219) and (220) would be obtained as a mere consequence of the former. Since, by Equation (79), the graded Lie bracket $\llbracket \cdot, \cdot \rrbracket$ (resp. $\llbracket \cdot, \cdot \rrbracket'$) coincides with q (resp. q'), Equation (217) can be rewritten as:

$$\phi_{(-i)} \circ q = q' \circ \phi_{(-i)} \quad (221)$$

and straightforwardly extended to $\Lambda^3(T_{\leq -1})|_{-i-1}$ (in particular, see Equation (86)), making the following diagram commutative :

$$\begin{array}{ccc} \Lambda^3(T_{\leq -1})|_{-i-1} & \xrightarrow{\phi_{(-i)}} & \Lambda^3(T'_{\leq -1})|_{-i-1} \\ q \downarrow & & \downarrow q' \\ \Lambda^2(T_{\leq -1})|_{-i-1} & \xrightarrow{\phi_{(-i)}} & \Lambda^2(T'_{\leq -1})|_{-i-1} \end{array}$$

where it is understood that $\phi_{(-i)}$ acts as an algebra morphism. Commutativity of the above diagram, together with the fact that $\text{Im}(q|_{\Lambda^3(T_{\leq -1})}) = \text{Ker}(q|_{\Lambda^2(T_{\leq -1})})$ (and a similar equality for q'), implies in turn that:

$$\phi_{(-i)}(\text{Ker}(q|_{\Lambda^2(T_{\leq -1})|_{-i-1}})) \subset \text{Ker}(q'|_{\Lambda^2(T'_{\leq -1})|_{-i-1}})$$

Since $\Lambda^2(T_{\leq -1})|_{-i-1} / \text{Ker}(q|_{\Lambda^2(T_{\leq -1})|_{-i-1}}) \simeq T_{-i-1}$ (and a similar identity for T'_{-i-1}), this implies that $\phi_{(-i)}$ passes to the quotient, and induces a linear map $\phi_{-i-1} : T_{-i-1} \rightarrow T'_{-i-1}$ which makes the following diagram commutative:

$$\begin{array}{ccc} \Lambda^2(T_{\leq -1})|_{-i-1} & \xrightarrow{\phi_{(-i)}} & \Lambda^2(T'_{\leq -1})|_{-i-1} \\ q \downarrow & & \downarrow q' \\ T_{-i-1} & \xrightarrow{\phi_{-i-1}} & T'_{-i-1} \end{array}$$

Replacing back q by $\llbracket \cdot, \cdot \rrbracket$, commutativity of the diagram is equivalent to the following equation:

$$\phi_{-i-1}(\llbracket x, y \rrbracket) = \llbracket \phi_{(-i)}(x), \phi_{(-i)}(y) \rrbracket \quad (222)$$

for every $x, y \in T_{\leq -1}$ such that $|x| + |y| = -i - 1$.

The similar condition involving an element of \mathfrak{g} and an element of T_{-i-1} comes from the fact that $q : \Lambda^2(T_{\leq -1}) \rightarrow T_{\leq -2}$ (resp. q') is a surjective \mathfrak{g} -equivariant (resp. \mathfrak{g}' -equivariant) map. More precisely, let $a \in \mathfrak{g}$ and $x \in T_{-i-1}$, then there exists $u \in \Lambda^2(T_{\leq -1})|_{-i-1}$ such that $x = q(u)$. But then, one obtains:

$$\phi_{-i-1}(\llbracket a, x \rrbracket) = \phi_{-i-1}(a \cdot q(u)) \quad (223)$$

$$= \phi_{-i-1}(q(a \cdot u)) = q'(\phi_{(-i)}(a \cdot u)) \quad (224)$$

$$= q'(\phi_0(a) \cdot \phi_{(-i)}(u)) = \phi_0(a) \cdot q'(\phi_{(-i)}(u)) \quad (225)$$

$$= \phi_0(a) \cdot \phi_{-i-1}(q(u)) = \llbracket \phi_0(a), \phi_{-i-1}(x) \rrbracket \quad (226)$$

where we used successively Equation (135), the \mathfrak{g} -equivariance of q , Equation (222), Equation (218), \mathfrak{g}' -equivariance of q' , Equation (222) again, and finally Equation (135). The proof does not depend on the choice of preimage u we make because $\text{Ker}(q|_{\Lambda^2(T_{\leq -1})|_{-i-1}})$ is stable under the action of \mathfrak{g} . Thus, we have extended Equation (217) one level higher:

$$\phi_{(-i-1)}(\llbracket x, y \rrbracket) = \llbracket \phi_{(-i-1)}(x), \phi_{(-i-1)}(y) \rrbracket \quad (227)$$

for every $x, y \in T_{\leq 0}$ such that $-i-1 \leq |x| + |y| \leq 0$.

There are two more compatibility equations between ϕ_{-i-1} and the graded Lie bracket to check, that is:

$$\phi_{-i}(\llbracket \Theta, x \rrbracket) = \llbracket \phi_{+1}(\Theta), \phi_{-i-1}(x) \rrbracket \quad (228)$$

$$\phi_{-i}(\llbracket \Theta_{a_1 \dots a_m}, x \rrbracket) = \llbracket \phi_{+1}(\Theta_{a_1 \dots a_m}), \phi_{-i-1}(x) \rrbracket \quad (229)$$

for every $x \in T_{-i-1}$ and every $a_1, \dots, a_m \in \mathfrak{g}$. Equation (228) can be rewritten in the form of Equation (216), at level $-i-1$:

$$\phi_{-i}(\partial(x)) = \partial'(\phi_{-i-1}(x)) \quad (230)$$

This is indeed satisfied; by choosing a pre-image $u \in \Lambda^2(T_{\leq -1})_{-i-1}$ of x , one has the following result:

$$\partial'(\phi_{-i-1}(x)) = \partial'(\phi_{-i-1}(q(u))) = \partial'(q'(\phi_{(-i)}(u))) = q'(\partial'(\phi_{(-i)}(u))) + m'(\phi_{(-i)}(u)) \quad (231)$$

$$= q'(\phi_{(-i)}(\partial(u))) + \phi_{(-i)}(m(u)) = \phi_{(-i)}(q(\partial(u)) + m(u)) = \phi_{-i}(\partial(x)) \quad (232)$$

where we have successively used Equations (222), (82) (or (83)) (216), (217), and Equation (82) (or (83)) again. Notice that we passed from $m'(\phi_{(-i)}(u))$ to $\phi_{(-i)}(m(u))$ by using Equation (217) and the definition of m (resp. m'). Proving Equation (229) then relies on using Equations (227) (on $T_0 \wedge T_{-i-1}$) and (228), as well as the same kind of arguments as for Equations (211)-(215). This concludes the proof of the inductive step.

The induction provides us with a linear application $\phi : \mathbb{T} \rightarrow \mathbb{T}'$ satisfying the following conditions:

$$\partial' \circ \phi(a) = \phi \circ \partial(a) \quad (233)$$

$$\phi(\llbracket a, b \rrbracket) = \llbracket \phi(a), \phi(b) \rrbracket \quad (234)$$

Moreover, this differential graded Lie algebra morphism between \mathbb{T} and \mathbb{T}' is such that $\phi|_{T_{-1}} = \chi$, $\phi|_{T_0} = \varphi$, and $\phi_{+1}(\Theta) = \Theta'$. The morphism ϕ is the image through the function $G : \mathbf{Lie-Leib} \rightarrow \mathbf{DGLie}_{\leq 1}$ of the compatible morphism of Lie-Leibniz triples (φ, χ) :

$$G(\varphi, \chi) = \phi \quad (235)$$

Then, when restricted to the wide subcategory **compLie-Leib**, the injective-on-objects function G is a functor. Moreover, since by construction $G(\varphi, \chi) = G(\varphi', \chi')$ if only if $\varphi = \varphi'$ and $\chi = \chi'$, G is faithful.

Finally, let us explain why **compLie-Leib** is the biggest wide subcategory of **Lie-Leib** (with respect to inclusion) such that G is a (faithful) functor. The idea is that in the general case where the morphism of Lie-Leibniz triples (φ, χ) is not compatible in the sense of Definition 9, an obstruction arises. Indeed, let (φ, χ) be any morphism of Lie-Leibniz triples between $(\mathfrak{g}, V, \Theta)$ and $(\mathfrak{g}', V', \Theta')$. Then, as seen in the first step of the induction, the couple (φ, χ) would induce a morphism of differential graded Lie algebras between the associated tensor hierarchies \mathbb{T} and \mathbb{T}' if and only if there exists a map $\tau : S^2(V)/K \rightarrow S^2(V')/K'$ making the following diagram commutative:

$$\begin{array}{ccc} S^2(V) & \xrightarrow{\chi \odot \chi} & S^2(V') \\ p \downarrow & & \downarrow p' \\ S^2(V)/K & \xrightarrow{\tau} & S^2(V')/K' \end{array}$$

However, the existence of such a map τ is dependent on the fact that $\chi \odot \chi(K) \subset K'$, namely that (φ, χ) is a compatible morphism. \square

Remark 13. *The equivalence of 2-categories between **Lie** \times **Mod** and that of strict Lie 2-algebras [1] does not seem to extend to **compLie-Leib**, as it is not obvious how to define a homotopy between two morphisms of tensor hierarchies $(\phi_{-i})_i$ and $(\phi'_{-i})_i$ from the data of a homotopy between morphisms of Lie-Leibniz triples.*

We conclude this section and the paper by some discussion on the algebraic structures involved and open questions. First, what is the significance of only partial functoriality in Theorem 1? While the construction of tensor hierarchies from Lie-Leibniz triples was inspired by theoretical physical considerations, the question of functoriality was motivated by the mathematical relevance of characterizing the injective-on-objects function G . With respect to this latter motivation, the result established in Theorem 1 seems to be the best one can hope for. It hints at the possibility that the more relevant notion of morphism in the category of Lie-Leibniz triples is that of *compatible morphisms* (see Definition 9).

Second, what is the meaning of the cohomology of the tensor hierarchy associated to a given Lie-Leibniz triple? In supergravity theories, the tensor hierarchy appears as a tool to cope with gauge theories in which the gauge algebra is not Lie [16, 18, 43]. The cohomology of the tensor hierarchy then parametrizes the p -form gauge fields which cannot be obtained from $p+1$ -form gauge fields, thus measuring some sort of independence in the gauge parameters. In the algebraic construction described in the present paper, the meaning of the cohomology is not so clear as the relationship with the original physical inspiration has vanished. However, the following argument may shed some light on this question. For every Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ for which:

$$K = \text{Ker}(\{ \cdot, \cdot \}) \quad (236)$$

the map $\partial_{-1} : T_{-2} \rightarrow T_{-1}$ is injective because $T_{-2} = S^2(V)/\text{Ker}(\{ \cdot, \cdot \})[2]$ is isomorphic to the ideal of squares \mathcal{I} , the image of ∂_{-1} . Moreover, one can then show that Condition (236) implies that $\partial_{-2} = 0$, as would be expected from $T_{\leq -1}$ being a cochain complex. The question arises: what about the higher cohomology spaces? We have some reason to think that the following conjecture holds:

Conjecture 1. *For a Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ satisfying condition (236) (so in particular for semi-strict Lie-Leibniz triples), the associated tensor hierarchy induces an exact sequence of the ideal of squares \mathcal{I} :*

$$\dots \xrightarrow{\partial_{-4}} T_{-4} \xrightarrow{\partial_{-3}} T_{-3} \xrightarrow{\partial_{-2}} T_{-2} \xrightarrow{\partial_{-1}} \mathcal{I}[1] \longrightarrow 0$$

This would be plausible since in this case, the role of \mathfrak{g} in the construction of the tensor hierarchy is not as salient as compared to the general case (e.g. in defining T_{-2}).

Third, the motivation behind this axiomatic construction of tensor hierarchies goes well beyond supergravity theories (the following argument is taken from [3]). In classical gauge theories, gauge transformations are interpreted as smooth transformations of the fields ϕ^i leaving the action invariant:

$$0 = \delta_\epsilon(S[\phi^i]) = \int \frac{\delta S}{\delta \phi^i} \delta_\epsilon \phi^i \quad (237)$$

where ϕ^i denote all the fields and ϵ denotes an arbitrary vector in the space of gauge parameters V . Invariance of the action under the commutator of two gauge transformations $[\delta_\epsilon, \delta_{\epsilon'}]$ implies that the space of gauge transformations forms a Lie algebra \mathfrak{g} . From this, physicists deduce that the space of gauge parameters V is itself a Lie algebra and $\delta : V \rightarrow \mathfrak{g}$ is a Lie algebra morphism. But as was noticed in [3, 23], it could well be that V is a Leibniz algebra and δ is an embedding tensor. As is currently established in higher gauge theories, if the gauge algebra is not Lie, we expect some *higher form of Lie algebras* to appear, as these are the ones integrating to (higher) symmetry groups. In the case at hand, the corresponding higher gauge algebra would be the tensor hierarchy, or an alternative, equivalent higher structure [6]. The present article thus provides a rigorous construction of the needed material.

Finally, the question arises as to how the statement of Theorem 1 generalizes to pairs of algebras over other operads. As explained earlier the motivation behind the construction of the tensor hierarchy associated to a Lie-Leibniz triple comes from theoretical physical considerations. We did not (yet) consider the possibility of extending this construction to other kinds of algebras, with suitable compatibility conditions. This question may be answered once we have a more categorical approach to the construction of tensor hierarchies, departing from the technical, computational proofs presented in Section 3. The present paper only provides an explicit, constructive proof of Theorem 1, and hints at the existence of a proof using more operadic and categorical tools. Actually, this kind of proof may be of great help in order to generalize Theorem 1 to the category of Leibniz algebroids.

A Corrigendum (in collaboration with Jakob Palmkvist)

In [33], Theorem 1, $\mathbf{DGLie}_{\leq 1}$ should be replaced by \mathbf{DGLie} . The reason is that when the embedding tensor is not \mathfrak{g} -equivariant, consistency of the differential graded Lie algebra structure at level $+1$ may sometimes require the introduction of spaces of higher degrees (see Proposition 13' and Sequence (252)). This observation had been missed in [33] and is based on an example inspired by supergravity theories contradicting Proposition 13 in [33] which has been provided by one of the authors of the present Corrigendum (see Example 8). Likewise, the part of the proof of Proposition 23 in [33] involving the space of degree $+1$ is modified accordingly. These modifications imply that $\mathbf{DGLie}_{\leq 1}$ should be replaced by \mathbf{DGLie} in Propositions 26 and 27, and thus in Theorem 1 too. Table 1 summarizes how this updated result applies to the most distinguished cases.

I. First, Proposition 13 in [33] should be rewritten:

Proposition 13'. *To every Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ in which V is a Lie algebra, there corresponds a differential graded Lie algebra:*

$$V[1] \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{-\eta(-; \Theta)} R_\Theta[-1] \xrightarrow{[\Theta, \cdot]} T_2 \xrightarrow{[\Theta, \cdot]} \dots$$

which canonically extends the graded Lie algebra structure on $V[1] \oplus \mathfrak{g}$ defined by Equations (41)-(43).

Remark 14. *The error in the proof of Proposition 13 in [33] was that Equation (60) cannot be enforced, as both sides of the equation are determined independently from existing data, and so they may not coincide.*

Likewise, one cannot assume Equation (154). Moreover, Equation (55) and Equation (146) in [33] should be replaced by $[\Theta, \Theta] = 0$ (which is equivalent to the cohomological equation $\partial \circ \partial = 0$).

Proof. The proof of Proposition 13 in [33] is valid, until Equation (60). The construction of the spaces of degrees higher than one is similar to that appearing in [37] (where the grading convention is reversed). We generate the spaces T_i , $i \geq 2$ by induction, from the commutators of the adjoint action of the elements at level $1 \leq k \leq i - 1$. Given an element $\chi \in T_1 = R_\Theta[-1]$, the adjoint action of χ is defined from Equations (53) and (54) in [33] by $\text{ad}_\chi(x) = [\chi, x]$.

Then, one defines T_2 to be the vector space generated by elements of the form $(\text{ad}_\chi \circ \text{ad}_\psi + \text{ad}_\psi \circ \text{ad}_\chi)|_V$, which correspond to the graded commutator $[\text{ad}_\chi, \text{ad}_\psi]|_V$. If T_2 is zero (see Example 9), the process stops and the differential graded Lie algebra obtained through Proposition 13' is $V[1] \rightarrow \mathfrak{g} \rightarrow T_1$. Otherwise, we represent the element $[\text{ad}_\chi, \text{ad}_\psi]|_V$ of T_2 by the bracket $[\chi, \psi]$ and we assign to it a degree +2. The evaluation of $[\chi, \psi]$ on an element $x \in V$ is symbolized by the nested bracket $[[\chi, \psi], x]$ or, alternatively, by the notation $\text{ad}_{[\chi, \psi]}(x)$. By construction, we then have the following Jacobi identity, for every $x \in V[1]$:

$$[[\chi, \psi], x] = [\chi, [\psi, x]] + [\psi, [\chi, x]] \quad (238)$$

Moreover, the bracket $[\chi, \psi]$ is graded skew-symmetric, i.e. $[\chi, \psi] = -[\psi, \chi]$ and represents the action of T_1 on T_1 , taking values in T_2 . The action of \mathfrak{g} on T_2 is then defined by:

$$a \cdot [\chi, \psi] \stackrel{\text{def}}{=} [a \cdot \chi, \psi] + [\chi, a \cdot \psi] \quad (239)$$

Therefore, if $\chi = \psi = \Theta$, we have:

$$0 = a \cdot [\Theta, \Theta] = 2[\Theta, a \cdot \Theta] \quad (240)$$

which is consistent with Equation (53) in [33]. As $a \cdot \Theta = \eta(a, \Theta)$, where $\eta : \mathfrak{g} \rightarrow \text{Hom}(V, \mathfrak{g})$ is the representation of \mathfrak{g} on $\text{Hom}(V, \mathfrak{g})$, Equation (240) means that $[\Theta, \eta(a, \Theta)] = 0$, implying in turn that the linear operator $\partial_2 = [\Theta, \cdot] : T_1 \rightarrow T_2$ extends the chain complex $V[1] \rightarrow \mathfrak{g} \rightarrow R_\Theta[-1]$ to the right.

One defines T_3 to be the vector space generated by elements of the form:

$$[\text{ad}_\phi, \text{ad}_{[\chi, \psi]}]|_V \stackrel{\text{def}}{=} (\text{ad}_\phi \circ \text{ad}_{[\chi, \psi]} - \text{ad}_{[\chi, \psi]} \circ \text{ad}_\phi)|_V \quad (241)$$

Such elements are represented in T_3 by the nested bracket $[\phi, [\chi, \psi]]$, inheriting the graded skew-symmetry property of the commutator from Equation (241):

$$[\phi, [\chi, \psi]] = -[[\chi, \psi], \phi] \quad (242)$$

This property is consistent with the grading of ϕ and $[\chi, \psi]$. To elements of T_3 are attributed a degree +3.

The definition of the nested bracket $[\phi, [\chi, \psi]]$ corresponds to the following Jacobi identity:

$$[[\phi, [\chi, \psi]], x] = [\phi, [[\chi, \psi], x]] - [[\chi, \psi], [\phi, x]] \quad (243)$$

characterizing the action of T_3 on every element $x \in V[1]$. Using Equation (238), the first term on the right-hand side reads:

$$[\phi, [[\chi, \psi], x]] = [\phi, [\chi, [\psi, x]]] + [\phi, [\psi, [\chi, x]]] \quad (244)$$

while using Equation (239) shows that the second term of the right-hand side of Equation (243) reads:

$$-[[\chi, \psi], [\phi, x]] = -[\chi, [\psi, [\phi, x]]] - [\psi, [\chi, [\phi, x]]] \quad (245)$$

Summing both right-hand sides of Equations (244) and (245), gathering terms by using Equation (238) and eventually using Equation (243), one deduces that, as elements of T_3 :

$$[\phi, [\chi, \psi]] = [[[\phi, \chi], \psi] - [\chi, [\phi, \psi]]] \quad (246)$$

This is the Jacobi identity for three elements of degree +1, and can be understood as how T_2 acts on T_1 (or conversely). In particular, for $\phi = \chi = \Theta$, one has:

$$[[\Theta, [\Theta, \psi]] = \underbrace{[[\Theta, \Theta], \psi]}_{=0} - [[\Theta, [\Theta, \psi]]] \quad (247)$$

From this, one deduces that $\partial_3 = [[\Theta, \cdot]] : T_2 \rightarrow T_3$ extends the chain complex:

$$V[1] \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{-\eta(-;\Theta)} R_\Theta[-1] \xrightarrow{[[\Theta, \cdot]]} T_2$$

to the right. The action of \mathfrak{g} on T_3 is defined in a similar fashion as in Equation (239) and meets no obstruction.

If T_3 is zero-dimensional, then the process stops because T_4 could only be generated by elements of the form $[\text{ad}_\xi, \text{ad}_{[[\chi, \psi]]}]|_V$ or $[\text{ad}_{[[\xi, \zeta]], \text{ad}_{[[\chi, \psi]]}}]|_V$ (where the commutator is graded), but the latter kind of elements can be rewritten using terms of the former kind. Indeed, by using Equations (243) and (246), together with the knowledge of the action of \mathfrak{g} on T_3 , one finds:

$$[\text{ad}_{[[\xi, \zeta]], \text{ad}_{[[\chi, \psi]]}}](x) = [\text{ad}_\xi, \text{ad}_{[[\zeta, [[\chi, \psi]]]}](x) + [\text{ad}_\zeta, \text{ad}_{[[\xi, [[\chi, \psi]]]}](x) \quad (248)$$

for every $x \in V$. If T_3 is not zero-dimensional, the induction continues, with the space at level $k \geq 4$ being defined as:

$$T_k = \text{Span} \left([\text{ad}_{T_1}, \text{ad}_{T_{k-1}}]|_V \right)$$

where the commutator is again graded. If $T_i = 0$ for some i then $T_j = 0$ for every $j \geq i$. The graded Lie bracket between two elements Ξ and Ω whose degrees sum up to k comes from the very definition of T_k , while the graded Jacobi identity between elements of various degrees can be obtained from a similar technique performed through Equations (239) and (243)–(246). At each step, the chain complex

$$V[1] \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{-\eta(-;\Theta)} R_\Theta[-1] \xrightarrow{[[\Theta, \cdot]]} T_2 \xrightarrow{[[\Theta, \cdot]]} T_3 \xrightarrow{[[\Theta, \cdot]]} \dots \xrightarrow{[[\Theta, \cdot]]} T_{k-1}$$

can be extended to the right by the linear map $\partial_k = [[\Theta, \cdot]] : T_{k-1} \rightarrow T_k$. Eventually, the compatibility between the graded Lie bracket and the differential is guaranteed by construction. \square

II. We modify the proof of Proposition 23 in [33] following the same lines as in heading **I.**. The only remaining data to be introduced for the differential graded Lie algebra structure to be consistent is the action of the positively graded spaces T_k , for $k \geq 2$, on the negatively graded spaces T_{-i} , for $i \geq 2$. This is done by noticing that 1. the action of T_k on V is properly defined by the proof of Proposition 13', and 2. that the graded Lie bracket on $T_{\leq -1}$ is surjective on $T_{\leq -2}$ (this is item 2 of Proposition 14 in [33]). Then, we proceed by induction and we set, to begin with, for any $x, y \in V[1]$ and $\Xi \in T_k$:

$$[\Xi, [x, y]] \stackrel{\text{def}}{=} [[\Xi, x], y] + (-1)^k [x, [\Xi, y]] \quad (249)$$

Assume that the action of T_k has been defined on T_{-j} for $1 \leq j \leq i-1$, then for any $u \in T_{|u|}$ and $v \in T_{|v|}$ – where $|u|$ symbolizes the (necessarily negative) degree of u – such that $-i-1 \leq |u|, |v| \leq -1$ and $|u| + |v| = -i$, we set:

$$[\Xi, [u, v]] \stackrel{\text{def}}{=} [[\Xi, u], v] + (-1)^{k|u|} [u, [\Xi, v]] \quad (250)$$

In order for the definition of the left-hand side to be consistent – e.g. it should not depend on the choice of representant of $[u, v]$ in the free graded Lie algebra of $V[1]$ – one should check by induction that Equation (250) respects the graded Jacobi identity, in the following sense:

$$[[\Xi, u], [v, w]] + (-1)^{k|u|} [u, [[\Xi, v], w]] + (-1)^{k(|u|+|v|)} [u, [v, [\Xi, w]]] + \text{cyclic} = 0 \quad (251)$$

where the arrow symbolizes graded circular permutation. This statement has been justified and proved (for the non-graded case) in Section 2.2 in [39].

Definitions (249) and (250), being consistent with the existing graded Lie bracket and differential, turn the following chain complex:

$$\dots \xrightarrow{\partial_{-3}} T_{-3} \xrightarrow{\partial_{-2}} T_{-2} \xrightarrow{\partial_{-1}} V[1] \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{-\eta(\cdot; \Theta)} R_{\Theta}[-1] \xrightarrow{\llbracket \Theta, \cdot \rrbracket} T_2 \xrightarrow{\llbracket \Theta, \cdot \rrbracket} T_3 \xrightarrow{\llbracket \Theta, \cdot \rrbracket} \dots \quad (252)$$

into a differential graded Lie algebra, called the *tensor hierarchy* associated to the Lie-Leibniz triple $(\mathfrak{g}, V, \Theta)$ in [33]. We present in Table 1 the tensor hierarchies associated to several distinguished Lie-Leibniz triples.

| Lie-Leibniz triple | corresponding dgLa |
|--|---|
| Lie alg. crossed module $V \xrightarrow{\Theta} \mathfrak{g}$ V is a Lie algebra Θ is \mathfrak{g} -equivariant | $V[1] \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{0} \mathbb{R}[-1]$ |
| augmented Leibniz alg. [5] $V \xrightarrow{\Theta} \mathfrak{g}$ Θ is \mathfrak{g} -equivariant | $\dots \xrightarrow{\partial_{-3}} T_{-2} \xrightarrow{\partial_{-1}} V[1] \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{0} \mathbb{R}[-1]$ |
| Proposition 13' $V \xrightarrow{\Theta} \mathfrak{g}$ V is a Lie algebra | $V[1] \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{-\eta(\cdot; \Theta)} R_{\Theta}[-1] \xrightarrow{\llbracket \Theta, \cdot \rrbracket} T_2 \xrightarrow{\llbracket \Theta, \cdot \rrbracket} \dots$ |
| general Lie-Leibniz triple $V \xrightarrow{\Theta} \mathfrak{g}$ | $\dots \xrightarrow{\partial_{-3}} T_{-2} \xrightarrow{\partial_{-1}} V[1] \xrightarrow{\Theta} \mathfrak{g} \xrightarrow{-\eta(\cdot; \Theta)} R_{\Theta}[-1] \xrightarrow{\llbracket \Theta, \cdot \rrbracket} T_2 \xrightarrow{\llbracket \Theta, \cdot \rrbracket} \dots$ |

Table 1: Description of the injective-on-object function $\mathbf{LieLeib} \rightarrow \mathbf{DGLie}$ in various cases.

Remark 15. *It is conjectured that the negatively graded part of these tensor hierarchies are isomorphic to the positively graded part of the Borchers algebras considered in [11, 37]. This isomorphism will be explored elsewhere.*

III. As a consequence of heading **II.**, $\mathbf{DGLie}_{\leq 1}$ should be replaced by \mathbf{DGLie} in Propositions 26 and 27 in [33]. The proof of the former is not impacted by this change however, as it relies on the non-positively graded part of the algebra only. On the other hand, the functorial property described in Proposition 27 is still valid under the modifications brought in by heading **II.** This is because the linear map $\phi_1 : T_1 \rightarrow T'_1$ straightforwardly extends by induction to a graded Lie algebra morphism from $T_{1 \leq i}$ to $T'_{1 \leq i}$, as we now show.

To start with, one sets:

$$\phi_2(\llbracket \chi, \psi \rrbracket) \stackrel{\text{def}}{=} \llbracket \phi_1(\chi), \phi_1(\psi) \rrbracket' \quad (253)$$

for every $\chi, \psi \in T_1 = R_{\Theta}[-1]$, and more generally, for every $\chi_1, \dots, \chi_i \in T_1$:

$$\phi_i(\llbracket \chi_1, \llbracket \chi_2, \llbracket \dots \llbracket \chi_{i-1}, \chi_i \rrbracket \dots \rrbracket \rrbracket) \stackrel{\text{def}}{=} \llbracket \phi_1(\chi_1), \llbracket \phi_1(\chi_2), \llbracket \dots \llbracket \phi_1(\chi_{i-1}), \phi_1(\chi_i) \rrbracket' \dots \rrbracket' \rrbracket' \quad (254)$$

Since the graded Lie bracket of two elements $\Xi \in T_{j_1}$, $\Omega \in T_{j_2}$, $j_1, j_2 \geq 1$, can be written as a sum of nested brackets of elements of T_1 , one deduces from Equation (254) that:

$$\phi_{j_1+j_2}(\llbracket \Xi, \Omega \rrbracket) = \llbracket \phi_{j_1}(\Xi), \phi_{j_2}(\Omega) \rrbracket' \quad (255)$$

This implies that the collection of linear maps $(\phi_i)_{i \geq 1}$ forms a graded Lie algebra morphism from $T_{\geq 1}$ to $T'_{\geq 1}$.

These linear maps are moreover compatible with their non-positively graded counterpart, in the sense that Equations (204), (205), (209), (210) and (228), (229) in [33] are still valid for elements of degree higher than

1. More precisely, these equations, together with Equations (250) and (255), imply by induction that for any $\Xi \in T_i$ ($i \geq 2$) and $u \in T_{-j}$ ($j \geq 0$) one has:

$$\phi_{i-j}(\llbracket \Xi, u \rrbracket) = \llbracket \phi_i(\Xi), \phi_{-j}(u) \rrbracket \quad (256)$$

Let $\phi : \mathbb{T} \rightarrow \mathbb{T}'$ be the unique degree 0 linear map restricting to ϕ_i on T_i , $i \in \mathbb{Z}$. From the proof of Proposition 27 in [33] together with the above discussion, we obtain that ϕ is a differential graded Lie algebra morphism. As a consequence, $\mathbf{DGLie}_{\leq 1}$ should be replaced by \mathbf{DGLie} in Proposition 27. As Propositions 26 and 27 are constitutive of Theorem 1, we conclude that $\mathbf{DGLie}_{\leq 1}$ should be replaced by \mathbf{DGLie} in Theorem 1 in [33].

IV. Example 8 was proposed by Jakob Palmkvist as a counter-example to Proposition 13 in [33]. It has been inspired by the treatment of tensor hierarchies in gauged supergravity, from which the notion originates, and in extended geometry. We refer to [13, 37] and references within for details and more examples on this perspective. The graded Lie algebras appearing in Examples 8 and 9 are the Lie superalgebras of Cartan type denoted by $W(3)$ and $W(2)$ in [26]. The relation between these Lie superalgebras and the more general tensor hierarchy algebras (originally introduced in the gauged supergravity context) was explained in [9].

Example 8. *The data for the Lie-Leibniz triple is $\mathfrak{g} = \mathfrak{gl}_3(\mathbb{R})$, $V = \mathbb{R}^3$ is the defining representation, and the embedding tensor is defined on the basis vectors e_1, e_2, e_3 as:*

$$\Theta(e_1) = -E_{22}, \quad \Theta(e_2) = E_{21} \quad \text{and} \quad \Theta(e_3) = 0 \quad (257)$$

where E_{ij} is the 3×3 matrix with 1 at the i -th line and the j -th column. The Leibniz product is null, except in the following two cases:

$$e_1 \circ e_2 = -e_2 \quad \text{and} \quad e_2 \circ e_1 = e_2 \quad (258)$$

Thus, the Leibniz product is actually a Lie bracket. The quadratic constraint is verified.

Let $\chi = E_{13} \cdot \Theta$ and $\psi = E_{32} \cdot \Theta$. We can compute their action on the basis vectors of V , using the following generic formula, valid for any vector of $T_{+1} = R_\Theta[-1]$:

$$(a \cdot \chi)(x) = [a, \chi(x)] - \chi(a \cdot x) \quad (259)$$

for any $x \in V$. It gives:

$$\chi(e_i) = [E_{13}, -E_{22}\delta_i^1 + E_{21}\delta_i^2] - \Theta(e_1\delta_i^3) = -E_{23}\delta_i^2 + E_{22}\delta_i^3 \quad (260)$$

$$\psi(e_i) = [E_{32}, -E_{22}\delta_i^1 + E_{21}\delta_i^2] - \Theta(e_3\delta_i^2) = -E_{32}\delta_i^1 + E_{31}\delta_i^2 \quad (261)$$

where δ_i^j is the Kronecker delta. Then, we compute the right-hand side of Equation (238) in order to prove that the left-hand side is not zero, hence proving that Proposition 13 in [33] was erroneous. We have:

$$\llbracket \chi, \llbracket \psi, e_i \rrbracket \rrbracket + \llbracket \psi, \llbracket \chi, e_i \rrbracket \rrbracket = \delta_i^1 E_{32} \cdot \chi - \delta_i^2 E_{31} \cdot \chi + \delta_i^2 E_{23} \cdot \psi - \delta_i^3 E_{22} \cdot \psi \quad (262)$$

The first term cannot be canceled by any other and is not identically null since, by Equation (259):

$$E_{32} \cdot \chi(e_i) = [E_{32}, -E_{23}\delta_i^2 + E_{22}\delta_i^3] - \chi(e_3\delta_i^2) = -E_{33}\delta_i^2 + E_{22}\delta_i^2 + E_{32}\delta_i^3 \quad (263)$$

This means that $\llbracket \chi, \llbracket \psi, e_1 \rrbracket \rrbracket + \llbracket \psi, \llbracket \chi, e_1 \rrbracket \rrbracket \neq 0$, proving that $\llbracket \chi, \psi \rrbracket \neq 0$. Notice that in that particular example, although T_2 is not zero, it turns out that T_3 is, so every space of degree higher than 3 is identically zero.

Example 9. *This example illustrates that it is possible that $\dim(R_\Theta) > 1$ while at the same time $T_i = 0$ for every $i \geq 2$. Let $\mathfrak{g} = \mathfrak{gl}_2(\mathbb{R})$ and $V = \mathbb{R}^2$ be the defining representation of \mathfrak{g} . As embedding tensor, we set:*

$$\Theta(e_1) = -E_{12} \quad \text{and} \quad \Theta(e_2) = E_{11} \quad (264)$$

The kernel of the embedding tensor is zero, so the Leibniz product induced on V is actually Lie: $e_1 \circ e_2 = -e_1$ and $e_2 \circ e_1 = e_1$, and there are no spaces in degrees lower than -1 . The image of the embedding tensor in $\mathfrak{gl}_2(\mathbb{R})$ – that is to say the subspace spanned by E_{12} and E_{11} – acts trivially on Θ . From Equation (259), one deduces the action of the remaining two matrices on the embedding tensor:

$$E_{21} \cdot \Theta(e_i) = -E_{22}\delta_i^1 + E_{21}\delta_i^2 \quad (265)$$

$$E_{22} \cdot \Theta(e_i) = E_{12}\delta_i^1 - E_{11}\delta_i^2 \quad (266)$$

One notices that $E_{22} \cdot \Theta = -\Theta$, while $\psi = E_{21} \cdot \Theta : V \rightarrow \mathfrak{g}$ is another embedding tensor as it satisfies the quadratic constraint. The only non-trivial actions of matrices of \mathfrak{g} on ψ are the following: $E_{11} \cdot \psi = -\psi$ and $E_{12} \cdot \psi = \Theta$. The space R_Θ is thus two-dimensional and we only have to compute the nested bracket involving Θ and ψ :

$$[[\Theta, [\psi, e_i]]] + [[\psi, [\Theta, e_i]]] = -\delta_i^1 \Theta - \delta_i^2 \psi + \delta_i^1 \Theta + \delta_i^2 \psi = 0 \quad (267)$$

There is thus no need to introduce the degree 2 bracket $[[\Theta, \psi]]$ so T_2 , and more generally every T_i for $i > 1$, is zero. Finally, notice that Equation (267) is nothing but Equation (240), as seen from the present example.

Example 10. This example provides a case where the Lie algebra is the same as that of 9, but where the representation V is different, thus generating another orbit at level $+1$ and at higher levels as well. This example is a continuation of Example 7 of [33], where the spaces of positive degree had not been computed. In the present case, the Lie algebra is $\mathfrak{g} = \mathfrak{gl}_2(\mathbb{R})$, while the Leibniz algebra is $V = \mathcal{M}_{2 \times 2}(\mathbb{R})$, equipped with the following Leibniz product:

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \circ \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} = \begin{pmatrix} 0 & (a_{11} - a_{22})b_{12} \\ -(a_{11} - a_{22})b_{21} & 0 \end{pmatrix} \quad (268)$$

The action of \mathfrak{g} on V occurs through the matrix commutator. The embedding tensor is given by:

$$D : \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \longmapsto \begin{pmatrix} a_{11} & 0 \\ 0 & a_{22} \end{pmatrix} \quad (269)$$

Using Equation (259), one deduces the action of \mathfrak{g} on the embedding tensor D . This action was originally given by Equation (270) in [33] as:

$$\eta(a; D)(b) = [a, Db]_A - D([a, b]) = \begin{pmatrix} b_{12}a_{21} - a_{12}b_{21} & -(b_{11} - b_{22})a_{12} \\ (b_{11} - b_{22})a_{21} & b_{21}a_{12} - a_{21}b_{12} \end{pmatrix} \quad (270)$$

where $[a, b]$ denotes the commutator of matrices.

Equation (270) implies two things. On the one hand, D is invariant under the action of the two matrices E_{11} and E_{22} , which translates the fact that D is $\text{Im}(D)$ -equivariant, as it should be. On the other hand, the actions of E_{12} and E_{21} on D define two linear maps $D^+ : V \rightarrow \mathfrak{g}$ and $D^- : V \rightarrow \mathfrak{g}$ whose actions are:

$$D^+(a) = \begin{pmatrix} -a_{21} & a_{22} - a_{11} \\ 0 & a_{21} \end{pmatrix} \quad \text{and} \quad D^-(a) = \begin{pmatrix} a_{12} & 0 \\ a_{11} - a_{22} & -a_{12} \end{pmatrix} \quad (271)$$

Using Equation (259), one observes that the actions of E_{12} and E_{21} generate three other linear operators D^{++} , D^{--} and D^0 from V to \mathfrak{g} :

$$D^{++}(a) = E_{12} \cdot D^+(a) = 4 \begin{pmatrix} 0 & a_{21} \\ 0 & 0 \end{pmatrix}, \quad D^{--}(a) = E_{21} \cdot D^-(a) = 4 \begin{pmatrix} 0 & 0 \\ a_{12} & 0 \end{pmatrix} \quad (272)$$

$$D^0(a) = E_{21} \cdot D^+(a) = E_{12} \cdot D^-(a) = 2 \begin{pmatrix} a_{11} - a_{22} & -a_{12} \\ -a_{21} & a_{22} - a_{11} \end{pmatrix} \quad (273)$$

Each one of these linear maps from V to \mathfrak{g} are eigenvectors of the actions of the diagonal matrices (the image of D). The \mathfrak{g} -orbit $R_D \subset \text{Hom}(V, \mathfrak{g})$ is then spanned by the linear maps $D, D^+, D^-, D^{++}, D^{--}$ and D^0 , so it is 6-dimensional.

By computing the actions of every matrix of $\mathfrak{g} = \mathfrak{gl}_2(\mathbb{R})$ on the generators D^+, D^-, D^{++}, D^{--} and D^0 via Equation (259), one deduces that they form an irreducible representation of $\mathfrak{sl}_2(\mathbb{R})$. Indeed, the vectors:

$$v_0 = D^{++}, \quad v_1 = 4D^+, \quad v_2 = 2D^0, \quad v_3 = 4D^- \quad \text{and} \quad v_4 = D^{--}, \quad (274)$$

satisfy the usual properties of the 5-dimensional irreducible representation of $\mathfrak{sl}_2(\mathbb{R})$, denoted V_4 . One can check that the operator $\Omega = 6D - D^0$ generates the one-dimensional trivial representation of $\mathfrak{sl}_2(\mathbb{R})$ – denoted V_0 – so the \mathfrak{g} -orbit R_D is fully reducible: $R_D = V_0 \oplus V_4$. In particular, the embedding tensor D has one leg in each of the representations.

Then, by denoting $[[\chi, a]] = \chi(a)$ for every $\chi \in R_D$, one observes that:

$$[[D^+, [[D^-, a]]] + [[D^-, [[D^+, a]]] = -2a_{12}D^+ - 2a_{21}D^- \quad (275)$$

The right-hand side being non-zero for a general matrix $a \in V$, one deduces that $[[D^+, D^-]]$ cannot be zero, and rather should be set to act on matrices as the right-hand side of Equation (275). The space R_D being 6-dimensional and the bracket of two elements of R_D being symmetric, one should compute 21 brackets in total to exhaust all the possibilities and find a basis for the space T_2 . The dimension of the latter is strictly lower than 21 however, since some of the brackets vanish, as can be seen from the two following examples:

$$[[D^{++}, [[D^{++}, a]]] = -4a_{21}E_{12} \cdot D^{++} = 0 \quad \text{and} \quad [[D^{--}, [[D^{--}, a]]] = -4a_{12}E_{21} \cdot D^{--} = 0 \quad (276)$$

Acknowledgements:

The first author wants to thank the University of Pennsylvania, where the project started, for hosting him during one month. He is also grateful to Georges Skandalis and Stéphane Vassout for having funded most of his research under the ANR project ‘Analysis on singular and non compact spaces. A C^* -algebra approach’. The final stage of this research could not have happened without the financial sustainability of the French social welfare system, as well as scholarship No. 075-15-2019-1620 from the government of the Russian Federation. Many thanks to Christian Saemann and Henning Samtleben for their helpful comments during the preparation of the paper. We are indebted to Jakob Palmkvist for his insightful ideas on taking a quotient of the free graded Lie algebra and to the anonymous referees who encouraged us to ponder several questions that led us to improve the first version of this paper. Thanks to Marco Manetti and Mario Trigiante whose comments benefited the current version. We thank Juan Moreno and Jonathan Wise for providing the elegant Lemma 16, which is essential to the construction of the tensor hierarchy. We finally would like to thank Johannes Huebschmann for his careful reading and remarks in the final stage of redaction. Concerning the appendix, Sylvain Lavau and Jakob Palmkvist are grateful to the Mainz Institute for Theoretical Physics (MITP) of the DFG Cluster of Excellence PRISMA* (Project ID 39083149), for its hospitality and support during the workshop "Higher Structures, Gravity and Fields" in January 2023, when this work was completed. We would also like to thank the organizers Martin Cederwall and Henning Samtleben for insightful comments and discussions.

References

- [1] BAEZ, J. C., AND CRANS, A. S. Higher-dimensional algebra VI: Lie 2-algebras. *Theor. Appl. Categor.* 12 (2004), 492–528.
- [2] BLOKH, A. A generalization of the concept of a Lie algebra. *Dokl. Akad. Nauk SSSR* 165, 3 (1965), 471–473.
- [3] BONEZZI, R., AND HOHM, O. Leibniz gauge theories and infinity structures. *Commun. Math. Phys.* 377, 3 (2020), 2027–2077.
- [4] BONEZZI, R., AND HOHM, O. Duality hierarchies and differential graded Lie algebras. *Commun. Math. Phys.* 382, 1 (2021), 277–315.
- [5] BORDEMANN, M., AND WAGEMANN, F. Global integration of Leibniz algebras. *J. Lie Theory* 27, 2 (2017), 555–567.
- [6] BORSTEN, L., KIM, H., AND SAEMANN, C. EL_∞ -algebras, generalized geometry, and tensor hierarchies. *arXiv e-prints* (May 2021), arXiv:2106.00108.
- [7] BROWN, R., HIGGINS, P. J., AND SIVERA, R. *Nonabelian Algebraic Topology*, vol. 15 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2011.
- [8] CAGNACCI, Y., CODINA, T., AND MARQUES, D. L_∞ algebras and tensor hierarchies in exceptional field theory and gauged supergravity. *JHEP* 01 (2019), 117.
- [9] CARBONE, L., CEDERWALL, M., AND PALMKVIST, J. Generators and relations for Lie superalgebras of Cartan type. *J. Phys. A: Math. Theor.* 52, 5 (2019), 055203.
- [10] CEDERWALL, M., AND PALMKVIST, J. Superalgebras, constraints and partition functions. *JHEP* 08 (2015), 36.
- [11] CEDERWALL, M., AND PALMKVIST, J. Superalgebras, constraints and partition functions. *JHEP* 2015, 8 (2015), 36.
- [12] CEDERWALL, M., AND PALMKVIST, J. Extended geometries. *JHEP* 02 (2018), 071.
- [13] CEDERWALL, M., AND PALMKVIST, J. Tensor hierarchy algebras and extended geometry. Part I. Construction of the algebra. *J. High Energ. Phys.* 2020, 2 (2020), 144.
- [14] CHEN, Z., QIAO, Y., XIANG, M., AND ZHANG, T. Dg Loday-Pirashvili modules over Lie algebras. *arXiv e-prints* (Oct. 2021), arXiv:2110.11623.
- [15] CHIBRIKOV, E. S. The right-normed basis for a free Lie superalgebra and Lyndon-Shirshov words. *Algebr. Logic* 45, 4 (2006), 261–276.
- [16] DE WIT, B., NICOLAI, H., AND SAMTLEBEN, H. Gauged supergravities, tensor hierarchies, and M-theory. *JHEP* 02 (2008), 44.
- [17] DE WIT, B., AND SAMTLEBEN, H. Gauged maximal supergravities and hierarchies of non-abelian vector-tensor systems. *Fortschr. Phys.* 53, 5-6 (2005), 442–449.
- [18] DE WIT, B., AND SAMTLEBEN, H. The end of the p -form hierarchy. *JHEP* 08 (2008), 15.

- [19] GARCIA-MARTINEZ, X., KHMALADZE, E., AND LADRA, M. Non-abelian tensor product and homology of Lie superalgebras. *J. Algebra* 440 (2015), 464 – 488.
- [20] GOMIS, J., KLEINSCHMIDT, A., AND PALMKVIST, J. Symmetries of M-theory and free Lie superalgebras. *JHEP* 03 (2019), 160.
- [21] GREITZ, J., HOWE, P., AND PALMKVIST, J. The tensor hierarchy simplified. *Class. Quant. Grav.* 31, 1 (2014), 087001.
- [22] HALL, M. A basis for free Lie rings and higher commutators in free groups. *Proc. Amer. Math. Soc.* 1 (1950), 575–581.
- [23] HOHM, O., AND SAMTLEBEN, H. Higher gauge structures in double and exceptional field theory. *Fortsch. Phys.* 67, 8-9 (2019), 1910008.
- [24] HUEBSCHMANN, J. On the history of Lie brackets, crossed modules, and Lie-Rinehart algebras. *J. Geom. Mech.* 13, 3 (2021), 385–402.
- [25] JANELIDZE, G. Internal crossed modules. *Georgian Math. J.* 10, 1 (2003), 99–114.
- [26] KAC, V. G. Lie superalgebras. *Adv. Math.* 26, 1 (1977), 8–96.
- [27] KINYON, M. K., AND WEINSTEIN, A. Leibniz algebras, Courant algebroids, and multiplications on reductive homogeneous spaces. *Am. J. of Math.* 123, 3 (2001), 525–550.
- [28] KNAPP, A. W. *Lie Groups, Lie Algebras, and Cohomology*. No. 34 in Mathematical Notes. Princeton University Press, Princeton, 1988.
- [29] KOSMANN-SCHWARZBACH, Y. Derived brackets. *Lett. Math. Phys.* 69, 1 (2004), 61–87.
- [30] KOTOV, A., AND STROBL, T. The embedding tensor, Leibniz–Loday algebras, and their higher gauge theories. *Commun. Math. Phys.* 376, 1 (2020), 235–258.
- [31] LAVAU, S. Tensor hierarchies and Leibniz algebras. *J. Geom. Phys.* 144 (2019), 147–189.
- [32] LAVAU, S., PALMKVIST, J., AND STASHEFF, J. Corrigendum to “From Lie algebra crossed modules to tensor hierarchies” [J. Pure Appl. Algebra 227 (6) (2023) 107311]. *J. Pure Appl. Algebra* 227, 12 (2023), 107428.
- [33] LAVAU, S., AND STASHEFF, J. From Lie algebra crossed modules to tensor hierarchies. *J. Pure Appl. Algebra* 227, 6 (2023), 107311.
- [34] LODAY, J., AND PIRASHVILI, T. The tensor category of linear maps and Leibniz algebras. *Georgian Math. J.* 5, 3 (1998), 263–276.
- [35] LODAY, J.-L., AND PIRASHVILI, T. Universal enveloping algebras of Leibniz algebras and (co)homology. *Math. Ann.* 296, 1 (1993), 139–158.
- [36] LYNDON, R. C. On Burnside’s problem. *Trans. Amer. Math. Soc.* 77, 2 (1954), 202–215.
- [37] PALMKVIST, J. The tensor hierarchy algebra. *J. Math. Phys.* 55, 1 (2014), 011701.
- [38] PALMKVIST, J. The tensor hierarchy algebra. *J. Math. Phys.* 55 (2014), 011701.
- [39] PALMKVIST, J. Nonlinear realizations of Lie superalgebras. *Commun. Algebra* 50, 11 (2022), 4917–4936.

- [40] SHIRSHOV, A. I. On the bases of a free Lie algebra. In *Selected Works of A.I. Shirshov*, L. Bokut, I. Shestakov, V. Latyshev, and E. Zelmanov, Eds., Contemporary Mathematicians. Birkhäuser, Basel, 2009, pp. 113–118.
- [41] SHTERN, A. S. Free Lie superalgebras. *Sib. Math. J.* 27, 1 (1986), 136–140.
- [42] STROBL, T. Leibniz-Yang-Mills gauge theories and the 2-Higgs mechanism. *Phys. Rev. D* 99, 11 (2019), 115026.
- [43] TRIGIANTE, M. Gauged supergravities. *Phys. Rept.* 680 (2017), 1–175.