

On the saturation mechanism of the fluctuation dynamo at $\text{Pr}_M \geq 1$

Amit Seta,^{1,2,*} Paul J. Bushby,² Anvar Shukurov,² and Toby S. Wood²

¹*Research School of Astronomy and Astrophysics, Australian National University, Canberra, ACT 2611, Australia*

²*School of Mathematics, Statistics and Physics, Newcastle University, Newcastle Upon Tyne, NE1 7RU, UK*

(Dated: December 5, 2021)

The presence of magnetic fields in many astrophysical objects is due to dynamo action, whereby a part of the kinetic energy is converted into magnetic energy. A turbulent dynamo that produces magnetic field structures on the same scale as the turbulent flow is known as the fluctuation dynamo. We use numerical simulations to explore the nonlinear, statistically steady state of the fluctuation dynamo in driven turbulence. We demonstrate that as the magnetic field growth saturates, its amplification and diffusion are both affected by the back-reaction of the Lorentz force upon the flow. The amplification of the magnetic field is reduced due to stronger alignment between the velocity field, magnetic field, and electric current density. Furthermore, we confirm that the amplification decreases due to a weaker stretching of the magnetic field lines. The enhancement in diffusion relative to the field line stretching is quantified by a decrease in the computed local value of the magnetic Reynolds number. Using the Minkowski functionals, we quantify the shape of the magnetic structures produced by the dynamo as magnetic filaments and ribbons in both kinematic and saturated dynamos and derive the scalings of the typical length, width, and thickness of the magnetic structures with the magnetic Reynolds number. We show that all three of these magnetic length scales increase as the dynamo saturates. The magnetic intermittency, strong in the kinematic dynamo (where the magnetic field strength grows exponentially) persists in the statistically steady state, but intense magnetic filaments and ribbons are more volume-filling.

I. INTRODUCTION

Magnetic fields are observed in a variety of astrophysical objects, including stars, galaxies and galaxy clusters, where they play an important role in various physical processes. Based on length and time scales, astrophysical magnetic fields can be divided into two types: the large-scale or mean field, which is coherent over scales comparable to the size of the system, and the small-scale or fluctuating field, whose correlation length is of the order of the driving scale of the underlying turbulent flow. The driving scale of turbulence, l_0 , is of the order of 0.1 kpc in spiral galaxies [1–3], and 10 kpc in galaxy clusters [4, 5]. The fluctuating magnetic field is believed to evolve over the eddy turnover timescale, which is considerably shorter than the corresponding evolution timescale for the large-scale field (which is typically of the order of 10^8 yr in spiral galaxies, comparable to the rotation period). For spiral galaxies, the mean and fluctuating fields have comparable magnitudes and thus both kinds of fields are equally important for the galactic dynamics [6]. There are a number of reviews covering the theoretical, numerical, and observational aspects of the subject [7–12].

The evolution and maintenance of magnetic fields is generally explained by dynamo action, a process by which kinetic energy is converted to magnetic energy. Astrophysical flows leading to dynamo action are typically turbulent; such flows may be driven by convection in stars, supernovae in galaxies, and merger shocks, motion of galaxies and AGN outflows in galaxy clusters. Magnetic field amplification by turbulent motions has also been observed in laboratory experiments [13]. Depending upon the magnetic fields that they produce, such dynamos are generally categorized as either

mean-field or fluctuation (or “small-scale”) dynamos. Mean-field dynamos produce large-scale magnetic fields, whereas the fluctuation dynamo generates the small-scale component of the field via random stretching of field lines by the turbulent velocity [14, 15] (as conceptually explained by the stretch-twist-fold mechanism [16, 17]). Fluctuation dynamo action plays a crucial role not only in spiral galaxies [7, 10, 18–21], elliptical galaxies [22, 23] and galaxy clusters [24–27], but also in stars such as the Sun [28–31], making it a general type of astrophysical process. Fluctuation dynamos naturally produce intermittent magnetic fields [32–34], characterised by the presence of intense, localised field structures. In the galactic context, a better understanding of these structures is needed for cosmic ray propagation studies [35, 36] and in the galaxy cluster context for the interpretation of radio observations [37]. The initial stages of magnetic field growth, when the Lorentz force is negligible, have been thoroughly studied [9, 32], so here we focus on the nonlinear states of the fluctuation dynamo, for which it is possible to consider relatively simple idealised flows (i.e., homogeneous, isotropic turbulence). A mean-field dynamo would require additional physics, such as rotation, velocity shear and density stratification; such effects can be safely ignored over the length and time scales that will be of interest here.

In a fluctuation dynamo, the root mean square (rms) magnetic field grows exponentially if the magnetic Reynolds number Re_M (quantifying the efficiency of inductive effects compared to magnetic diffusion) exceeds its critical value $\text{Re}_M^{(\text{crit})}$, which depends on the properties of the flow. When the magnetic energy is low in comparison to the turbulent kinetic energy, the flow dynamics are not influenced by the magnetic field (the kinematic stage). For an isotropic, incompressible, mirror-symmetric, homogeneous and Gaussian random velocity field, which is also δ -correlated in time, it can be shown that the magnetic field power spectrum M_k in the kinematic stage follows a power-law (at low wave numbers) with

* amit.seta@anu.edu.au

slope $3/2$ [9, 14]. However, an exponentially growing magnetic field also leads to the exponential growth of the Lorentz force, which eventually makes the problem nonlinear. This slows down the growth and finally leads to the saturation of the dynamo (the saturated stage). The nonlinear problem is mostly studied via numerical simulations, in which the Navier-Stokes and induction equations are solved simultaneously [e.g., 26, 28, 30, 33, 38–47]. Our aim in this paper is to explore the saturation mechanism of the fluctuation dynamo and to characterize the magnetic structures it generates.

For fluctuation dynamos driven by homogeneous and isotropic turbulence, the following three quantities are prescribed: the driving scale of the turbulent flow l_0 , the fluid viscosity ν , and the magnetic resistivity η . Based on the magnetic Prandtl number Pr_M (defined to be the ratio of viscosity to resistivity, $\text{Pr}_M = \nu/\eta$), fluctuation dynamos can be divided into small and large Pr_M cases. Pr_M is greater than unity ($\eta < \nu$) for hot diffuse plasma (interstellar and intergalactic medium) and Pr_M is much smaller than unity ($\eta > \nu$) for dense plasma (planets, stars and liquid metal dynamo experiments). The critical magnetic Reynolds number $\text{Re}_M^{(\text{crit})}$, which is a threshold for dynamo action to occur, increases with decreasing Pr_M [48–52]. We focus upon the $\text{Pr}_M \geq 1$ regime, fixing the underlying flow (i.e., fixing Re) and then varying Re_M in order to study the sensitivity of the magnetic structures of nonlinear dynamo states to the magnetic Reynolds number.

This paper is structured as follows. In Section II, we introduce the basic equations and describe the numerical setup and provide parameters of the simulations. In Section III, we discuss magnetic field intermittency and in Section IV, we examine possible saturation mechanisms. Then, in Section V, we use Minkowski functionals to quantify the magnetic field structures (as a function of the magnetic Reynolds number) in both the kinematic and nonlinear regimes. Finally, in Section VI, we conclude with a discussion and propose some future directions of research.

II. BASIC EQUATIONS AND NUMERICAL MODELLING

To study the fluctuation dynamo action in a turbulent flow driven by a prescribed random force, we solve the equations of magnetohydrodynamics, using the Pencil code [53]. The computational domain is a triply-periodic cubic box of non-dimensional width $L = 2\pi$, with 256^3 or 512^3 grid points. The equations are solved with sixth-order finite differences in space and a third-order Runge–Kutta scheme for the temporal evolution. The governing equations are

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) = 0, \quad (1)$$

$$\frac{\partial \mathbf{b}}{\partial t} = \nabla \times (\mathbf{u} \times \mathbf{b}) + \eta \nabla^2 \mathbf{b}, \quad (2)$$

$$\begin{aligned} \frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} = & -\frac{\nabla p}{\rho} + \frac{\mathbf{j} \times \mathbf{b}}{c\rho} \\ & + \nu \left(\nabla^2 \mathbf{u} + \frac{1}{3} \nabla (\nabla \cdot \mathbf{u}) + 2\mathbf{S} \cdot \nabla \ln \rho \right) + \mathbf{F}, \end{aligned} \quad (3)$$

where \mathbf{u} is the velocity field, \mathbf{b} is the magnetic field, ρ is the fluid density, p is the pressure, η is the magnetic diffusivity, $\mathbf{j} = (c/4\pi) \nabla \times \mathbf{b}$ is the electric current density, c is the speed of light, ν is the viscosity, $S_{ij} = \frac{1}{2} (u_{i,j} + u_{j,i} - \frac{2}{3} \delta_{ij} \nabla \cdot \mathbf{u})$ is the rate-of-strain tensor, and \mathbf{F} is the forcing function (defined below). We use an isothermal equation of state, $p = c_s^2 \rho$, where the constant c_s is the sound speed. Eq. (2) is solved in terms of the magnetic vector potential to ensure that the magnetic field remains divergence free.

We drive the flow with a mirror-symmetric and δ -correlated in time forcing [39] of the form

$$\mathbf{F}(\mathbf{x}, t) = \text{Re}\{N \mathbf{F}_{\mathbf{k}(t)} \exp[i\mathbf{k}(t) \cdot \mathbf{x} + i\phi(t)]\}, \quad (4)$$

where \mathbf{k} is the wave vector, \mathbf{x} is the position vector and $-\pi < \phi \leq \pi$ is a random phase. To ensure that the forcing is nearly δ -correlated in time, \mathbf{k} and ϕ are changed at each time step δt . Also, to ensure that the time-integrated force is independent of the chosen time step δt , the normalization is $N = F_0 c_s (|\mathbf{k}| c_s / \delta t)^{1/2}$, where F_0 is the non-dimensional forcing amplitude chosen such that the maximum Mach number is small enough ($u_{\text{rms}}/c_s \lesssim 0.1$) to avoid strong compressibility. We select many random wave vectors \mathbf{k} , each of magnitude k (a multiple of $2\pi/L$ to make sure that the flow is periodic) in a given range. Then we select an arbitrary unit vector \mathbf{e} (neither parallel nor anti-parallel to \mathbf{k}) and set

$$\mathbf{F}_{\mathbf{k}} = \frac{\mathbf{k} \times \mathbf{e}}{|\mathbf{k} \times \mathbf{e}|}. \quad (5)$$

The form of Eq. (5) ensures that the forcing is solenoidal, i.e. $\nabla \cdot \mathbf{F} = 0$ by construction. The average wave number at which the flow is driven is denoted by k_F . Even when the flow is periodic, k_F need not be a multiple of $2\pi/L$. Physically, $2\pi/k_F$ represents the driving scale of the turbulent flow, l_0 , in the system.

The turbulent plasma is characterized by the hydrodynamic Reynolds number Re and magnetic Reynolds number Re_M , defined in terms of the rms velocity u_{rms} and the forcing scale k_F [54], as

$$\text{Re} = \frac{u_{\text{rms}}}{\nu} \frac{2\pi}{k_F}, \quad \text{Re}_M = \frac{u_{\text{rms}}}{\eta} \frac{2\pi}{k_F}. \quad (6)$$

We use non-dimensional units with lengths in units of the domain size $L = 2\pi$, speed in units of the isothermal sound speed c_s , time in units of the eddy turnover time $t_0 = 2\pi/u_{\text{rms}} k_F$, density in units of the initial density ρ_0 and the magnetic field in units of $(4\pi\rho_0 c_s^2)^{1/2}$. Initially, the density is constant everywhere and $\mathbf{u} = \mathbf{0}$, whilst there is a weak random, seed magnetic field with zero net flux across the domain.

For the first set of simulations, with parameters given in Table I, the turbulent motions are driven at the wave numbers $2\pi/L$ and $2(2\pi/L)$ at equal intensities, which implies that $k_F \approx 1.5(2\pi/L)$. The magnetic field grows for $\text{Re}_M \geq \text{Re}_M^{(\text{crit})}$, with $\text{Re}_M^{(\text{crit})} \approx 220$ for $\text{Pr}_M = 1$ [39]. The evolution of the rms velocity field, u_{rms} , and magnetic field, b_{rms} , is shown in Fig. 1 for $\text{Re}_M = 1122$. The flow speed is controlled by the forcing function and thus remains nearly constant. The

TABLE I. Summary of fluctuation dynamo simulations in a numerical domain of size $(L = 2\pi)^3$ with 256^3 mesh points. In all cases, the forcing scale k_F is approximately equal to $1.5(2\pi/L)$, the forcing amplitude $F_0 = 0.02$, the magnetic Prandtl number $\text{Pr}_M = 1$ and the rms velocity in the saturated state is $u_{\text{rms}}/c_s \approx 0.11$. For each simulation, we quote the Reynolds number, the magnetic Reynolds number, the rms magnetic field in the saturated state b_{rms} , the ratio of magnetic to kinetic energy in the saturated state $\varepsilon_M/\varepsilon_K = b_{\text{rms}}^2/u_{\text{rms}}^2$, the correlation length of the velocity and magnetic field in the kinematic stage $l_{u\text{kin}}$ and $l_{b\text{kin}}$, and similarly in the saturated stage $l_{u\text{sat}}$ and $l_{b\text{sat}}$.

η, ν	Re_M, Re	b_{rms}	$\varepsilon_M/\varepsilon_K$	$l_{u\text{kin}}$	$l_{b\text{kin}}$	$l_{u\text{sat}}$	$l_{b\text{sat}}$
10×10^{-4}	449	0.033	0.08	3.14	1.82	3.77	1.95
5×10^{-4}	898	0.042	0.14	3.20	1.26	3.45	1.76
4×10^{-4}	1122	0.048	0.20	3.01	0.94	3.64	1.76
3×10^{-4}	1496	0.049	0.21	3.01	0.88	3.39	1.57
2.5×10^{-4}	1796	0.054	0.25	2.95	0.75	3.58	1.57
2×10^{-4}	2244	0.055	0.26	2.95	0.69	3.33	1.56




FIG. 1. Root mean square (rms) velocity field u_{rms} (red) and magnetic field b_{rms} (blue) as functions of normalized time t/t_0 (where $t_0 = 2\pi/u_{\text{rms}}k_F$ is the eddy turnover time) for $\text{Re} = \text{Re}_M = 1122$. During the kinematic stage (area shaded in light red), the black dashed line corresponds to the exponential growth. As the magnetic field grows, the dynamo passes through a transitional stage (area shaded in light green), before reaching a statistically steady saturated state (area shaded in light blue).

magnetic field first decays until it reaches an eigenstate of the induction equation. Then it grows exponentially in the kinematic stage at the growth rate of $0.4 u_{\text{rms}} k_F / 2\pi$ in dimensional units. As it becomes stronger, the Lorentz force affects the flow and slows down the exponential increase. Finally, when the magnetic field becomes strong enough, the dynamo reaches a statistically steady state in the saturated stage. The exponential growth and then saturation of the magnetic field occurs in all of the runs shown in Table I.

The shell-averaged (one-dimensional) power spectra, for various stages of the magnetic field evolution, are shown in




FIG. 2. The shell-averaged (one-dimensional) kinetic E_k (dashed) and magnetic M_k (solid) energy spectra in the kinematic (red) and saturated (blue) stages for $\text{Re} = \text{Re}_M = 1122$. The kinetic energy spectrum is close to the Kolmogorov spectrum, $E_k \propto k^{-5/3}$ (dotted, black) in the main part of the wave number range. The magnetic spectrum is initially of the form $M_k \propto k^{3/2}$ (dashed, black) at smaller wave numbers. As the magnetic field saturates, its power shifts to smaller wave numbers and the magnetic spectrum flattens.

Fig. 2. At all times, the kinetic energy spectrum is close to the Kolmogorov spectrum, $E_k \propto k^{-5/3}$, in the range $3 \leq kL/2\pi \leq 20$ (flow is driven at $k = 2\pi/L$ and $k = 2(2\pi/L)$), which suggest that the velocity field is turbulent in nature. The magnetic spectrum in the kinematic stage has a broad maximum at large wave numbers and its slope agrees with the Kazantsev model, $M_k \propto k^{3/2}$, in the range $2 \leq kL/2\pi \leq 10$ with maximum power at approximately $kL/2\pi = 10$. Kazantsev's theory assumes that the turbulent flow is δ -correlated in time. Whilst we have used a δ -correlated forcing in the Navier-Stokes equation (term \mathbf{F} in Eq. (3)), the flow that it drives is not δ -correlated, especially at high Re . However, it is known that the slope of the spectrum in the kinematic stage remains the same even when the flow has a finite but small correlation time [55, 56], which explains why we recover the Kazantsev result in these simulations. As the magnetic field grows, the spectral maximum shifts to smaller wave numbers and the spectrum becomes much flatter with a broad maximum in the range $2 \leq kL/2\pi \leq 5$.

III. MAGNETIC INTERMITTENCY

Intermittency in a random field can manifest itself via heavy tails in its probability distribution function (PDF) and leads to an increased kurtosis in comparison with the Gaussian distribution. For the random velocity field \mathbf{u} with zero mean, the kurtosis is defined by

$$K(\mathbf{u}) = \frac{\langle \mathbf{u}^4 \rangle}{\langle \mathbf{u}^2 \rangle^2}, \quad (7)$$




FIG. 3. The PDF of the normalized velocity field component u_x / u_{rms} for $\text{Re}_M = 1122$ and $\text{Re}_M = 2244$ in the kinematic (dashed) and saturated (solid) stages for the value of Re_M given in the legend. The PDF of the single component of the velocity field is roughly Gaussian (dashed, black) in both the stages for both Re_M . Here only u_x / u_{rms} is shown but similar behaviour is exhibited by all three velocity components.

with angular brackets denoting the volume average. A useful diagnostic of the spatial structure is the correlation length of the field, l_u , which is calculated from the power spectrum E_k as

$$l_u = \frac{\pi}{2} \frac{\int_0^\infty 2\pi k^{-1} E_k dk}{\int_0^\infty E_k dk}. \quad (8)$$

Here, using such tools, we discuss the spatial intermittency of the velocity and magnetic fields in nonlinear fluctuation dynamos.

Fig. 3 shows the PDF of a single component of the velocity field u_x / u_{rms} in the kinematic and saturated dynamo stages for $\text{Re}_M = 1122$ and $\text{Re}_M = 2244$. The PDF is nearly Gaussian in both the kinematic and saturated stages. This is generally true for homogenous turbulence [57]. The velocity PDFs remain Gaussian even in the case of supersonic turbulence with a compressible forcing [e.g., Fig. A1. in 58]. For all cases of Table I, the kurtosis of the velocity field is very close to $K = 3$, which is the value for a Gaussian distribution. The correlation length of the velocity field l_u , also given in Table I, is about half of the periodic domain size $L = 2\pi$, as can also be seen from Fig. 4. It decreases slightly as Re increases and is slightly larger in the saturated stage than in the kinematic stage for all Re_M . The velocity field thus becomes more volume filling as the magnetic field saturates. This is directly attributable to the dynamical effects of the magnetic fields.

Even though the velocity field statistics are nearly Gaussian, the magnetic field in both the kinematic and saturated stages is spatially intermittent and strongly non-Gaussian. This can be seen from the PDFs of a normalized component of the magnetic field b_x / b_{rms} in Fig. 5. The distribution is far from a Gaussian one and has long, heavy tails. The non-

linearity truncates the most extreme relative magnetic field strengths above $|b_x| / b_{\text{rms}} \approx 3$. The magnetic field intermittency is further demonstrated in Fig. 6 which shows the PDF of b / b_{rms} for $\text{Re}_M = 1122$ and $\text{Re}_M = 2244$ in the kinematic and saturated stages. The PDF of the kinematic magnetic field strength follows a lognormal distribution and it has heavier tails in comparison to that of the saturated magnetic field. Thus the magnetic field is intermittent in both the kinematic and the saturated stages, but the level of intermittency decreases as the field saturates. It should be noted that this conclusion is consistent with that of Schekochihin et al. [33] (Fig. 6 in this paper is similar to their Fig. 27), who studied a closely related system. This confirms that this finding is robust to small variations in the model setup and parameters.

Magnetic intermittency can also be quantified by measuring the quantity $Q_{nl} = \langle (b/b_{\text{rms}})^{nl} \rangle^{(1/nl)}$ and its rate of change as nl changes (for example, Q_{nl}/Q_{nl-1}). Higher Q_{nl} and Q_{nl}/Q_{nl-1} is a signature of a larger degree of intermittency. Fig. 7 shows Q_{nl} and Q_{nl}/Q_{nl-1} for the magnetic field in the kinematic and saturated stages for $\text{Re}_M = 1122$ for $nl = 1, 2, 3, \dots, 50$. Q_{nl} and its rate of change are higher for the kinematic stage as compared to the saturated stage. This further demonstrates that the magnetic field in the saturated stage is less intermittent than that in the kinematic stage. We further compare both terms with the corresponding Gaussian versions obtained by randomizing phases in Fourier space [keeping the exact same magnetic field spectrum but destroying intermittent structures, as done in 35, 36, 59]. Q_{nl} and Q_{nl}/Q_{nl-1} are higher for the dynamo generated field in comparison to its randomized Gaussian versions in both the kinematic and saturated stages. Thus, the dynamo generated field is always spatially intermittent and the degree of intermittency decreases as the field saturates due to nonlinearity.

The two-dimensional vector plots of the magnetic fields in Fig. 8 also show larger structures in the saturated stage. This can be further seen in Fig. 9, which shows the isosurfaces of magnetic fields in the kinematic and saturated stages. The kurtosis of the kinematic magnetic field for $\text{Re}_M = 1122$ is 5.29 but is 3.32 in the saturated stage. This also suggests that the magnetic field in the kinematic stage is more intermittent than the saturated stage. The magnetic field correlation length l_b is calculated using Eq. (8) by replacing E_k with M_k , the magnetic field power spectrum. The magnetic field correlation length in the kinematic $l_{b\text{kin}}$ and saturated $l_{b\text{sat}}$ stages is given in Table I. The magnetic field correlation length decreases as Re_M increases, both for the kinematic and saturated stages (see Section V for further details). Thus, the magnetic field intermittency increases during both kinematic and saturated dynamo stages as Re_M increases. It is also clear that $l_{b\text{sat}} > l_{b\text{kin}}$ for all Re_M which confirms again that the magnetic field in the kinematic stage is less volume filling. The increase in the correlation length due to magnetic field saturation is true regardless of the choice of Re_M and agrees with previous numerical studies [26, 40].




FIG. 4. A 2D cut in the xy -plane with vectors $(u_x/u_{\text{rms}}, u_y/u_{\text{rms}})$ and colour showing the magnitude of u_z/u_{rms} in the kinematic (a) and saturated (b) stages with $\text{Re}_M = 2244$. The velocity field in both the stages looks qualitatively the same. The structures span approximately half of the domain.




FIG. 5. The PDF of the normalized magnetic field component b_x/b_{rms} for $\text{Re}_M = 1122$ and $\text{Re}_M = 2244$ in the kinematic (dashed) and saturated (solid) stages for the values of Re_M given in the legend. The magnetic field for both Re_M in both stages is far from a Gaussian (dashed, black). It has heavy tails, which is a sign of intermittency. Here only b_x/b_{rms} is shown but similar behaviour can be observed in all three magnetic field components.

IV. SATURATION OF THE FLUCTUATION DYNAMO

Several mechanisms have previously been considered to explain the saturation of the fluctuation dynamo, including a reduction in magnetic field line stretching due to the suppression of the Lagrangian chaos in the velocity field [60, 61], changes in the mutual alignment of the velocity and magnetic field lines [43], the folded structure of magnetic fields and energy equipartition between magnetic and velocity fields for




FIG. 6. The PDF of the normalized magnetic field strength b/b_{rms} for $\text{Re}_M = 1122$ and $\text{Re}_M = 2244$ in the kinematic (dashed) and saturated (solid) stages for the values of Re_M given in the legend. The PDF of the magnetic field in the kinematic state follows a lognormal distribution (dashed, black). The magnetic field is more intermittent in the kinematic stage than in the saturated stage.

$\text{Pr}_M \gg 1$ [33, 62], enhancement in diffusion due to additional nonlinear velocity drift [63, 64] and selective dissipation of the turbulent kinetic energy [65, 66]. From the induction equation (2), there are two type of processes that could lead to the saturation: a decrease in the induction term $(\nabla \times (\mathbf{u} \times \mathbf{b}))$ or an increase in the dissipation term $(\eta \nabla^2 \mathbf{b})$. We explore each scenario here.




FIG. 7. $Q_{nl} = \langle (b/b_{\text{rms}})^{nl} \rangle^{(1/nl)}$ (a) and Q_{nl}/Q_{nl-1} (b) as function of nl for the kinematic (red, solid) and saturated (blue, solid) stages for $\text{Re}_M = 1122$. The corresponding quantities for randomized fields which have almost Gaussian statistics (dashed) are also plotted. The dynamo generated magnetic field is always intermittent with the degree of intermittency being higher in the kinematic stage.




FIG. 8. As Fig. 4 but for the magnetic field. The magnetic field in the kinematic stage (a) is intermittent with random magnetic structures. In the saturated stage (b), the field remains intermittent but the structures are larger.

A. Alignment of velocity field, magnetic field and electric current density

We first examine how the induction term is affected when the field becomes stronger. The rms magnitude of both the velocity and magnetic fields are statistically steady, as shown in Fig. 1. Thus, we consider the alignment of the magnetic field with the velocity field as a possible mechanism for the saturation. Such an alignment has been studied in the context of convectively driven fluctuation dynamos [43, 67], MHD turbulence in the presence of a strong guide field [68] and decaying isotropic MHD turbulence [69]. For the numerical simulations described in Table I, we calculate the angle between the velocity \mathbf{u} and magnetic field \mathbf{b} , and between the current

density \mathbf{j} and \mathbf{b} ,

$$\cos(\theta)_{\mathbf{u}, \mathbf{b}} = \frac{\mathbf{u} \cdot \mathbf{b}}{|\mathbf{u}||\mathbf{b}|}, \quad \text{and} \quad \cos(\theta)_{\mathbf{j}, \mathbf{b}} = \frac{\mathbf{j} \cdot \mathbf{b}}{|\mathbf{j}||\mathbf{b}|}, \quad (9)$$

respectively. An increase in the level of alignment between \mathbf{u} and \mathbf{b} implies a decrease in the effectiveness of magnetic induction. On the other hand, an increase in the level of alignment between \mathbf{j} and \mathbf{b} leads to a decrease in the Lorentz force, i.e., the field becomes more force-free.

Fig. 10 and Fig. 11 show the probability density functions of the cosines in the kinematic and saturated stages for $\text{Re}_M = 1122$ and $\text{Re}_M = 1496$. Since both angles are symmetric about $\mathbf{b} = \mathbf{0}$, we show PDFs of the absolute value of their cosines. For both values of Re_M , the cosine of the angle




FIG. 9. Isosurfaces of $b^2/b_{\text{rms}}^2 = 4$ (blue) and $b^2/b_{\text{rms}}^2 = 5$ (yellow) for the magnetic fields in the kinematic (a) and saturated stages (b) for $\text{Re}_M = 2244$. The structures in the saturated stage are larger in size as compared to that in the kinematic stage.

between the velocity and magnetic field, $|\cos(\theta)_{\mathbf{u}, \mathbf{b}}|$, tends to be larger in the saturated stage than in the kinematic stage. The better alignment between \mathbf{u} and \mathbf{b} decreases the induction term $\nabla \times (\mathbf{u} \times \mathbf{b})$ and thus reduces the amplification of the magnetic field. To put this another way, the enhanced alignment between \mathbf{u} and \mathbf{b} implies a decrease in the energy transfer from the flow to the magnetic field (which is a process that has been studied in some detail in the context of shell models of magnetohydrodynamic turbulence [70–73]). However, there is a significant fraction of the volume where the two fields are not aligned and so the amplification is not completely suppressed. This minimum level of amplification is required to balance the magnetic diffusion. The cosine of the angle between the current density and magnetic field $\cos(\theta)_{\mathbf{j}, \mathbf{b}}$ is also statistically larger by magnitude in the saturated stage. Thus, the field becomes closer to a force-free form as it saturates. This also implies that the morphology of magnetic field changes on saturation, which motivates us to study the morphology of magnetic structures in Section V. Overall, because of the enhanced local alignment between the velocity and magnetic field, the field amplification rate decreases. At the same time, due to the increase in the local alignment between the current density and magnetic field, the field becomes more force-free.

Similar broad conclusions apply when we consider conditional PDFs that focus exclusively upon the regions of stronger field (higher b/b_{rms} in Fig. 10 and Fig. 11). However, the level of alignment between the velocity and magnetic field is higher in the strong field regions in both the kinematic and saturated stages. This suggests that the strong field regions require a larger reduction in amplification by alignment. The distribution of $\cos(\theta)_{\mathbf{j}, \mathbf{b}}$ in the kinematic stage shows some dependence upon the field strength but in the saturated stage the

difference is less pronounced. In the kinematic stage, alignment is weakest in the relatively strong field regions, suggesting that in the strong field regions, not only because of its higher strength (as the Lorentz force is proportional to the strength of the field) but also because of the lower level of alignment, the field produces a stronger back reaction on the flow.

Another important question is whether the alignment between the velocity and magnetic fields and the magnetic field and current density occur in the same spatial region. To answer this, we show the cross-correlation between the two angles in Fig. 12 which suggests that the velocity, magnetic field and current density are always nearly aligned to each other at same spatial positions. It is difficult to see any further difference between the kinematic and saturated stages in Fig. 12a and Fig. 12b. Fig. 12c and Fig. 12d show the same correlation but only for strong field regions, $b/b_{\text{rms}} > 1.5$. In Fig. 12c, the kinematic stage shows higher correlation in regions with high $\cos(\theta)_{\mathbf{u}, \mathbf{b}}$ and low $\cos(\theta)_{\mathbf{j}, \mathbf{b}}$, which is absent in the saturated stage. The larger misalignment of \mathbf{j} and \mathbf{b} , especially in the strong field regions, enhances the work done on the magnetic field by the flow. This promotes growth of the magnetic field. Once the field saturates, the larger correlation at high $\cos(\theta)_{\mathbf{u}, \mathbf{b}}$ and low $\cos(\theta)_{\mathbf{j}, \mathbf{b}}$ disappears in Fig. 12d. This implies a statistical decrease in the back-reaction of the magnetic field on the flow as the field saturates.

To summarize, the alignment between the velocity and magnetic field vectors and the magnetic field and current density vectors is statistically enhanced as the dynamo saturates. The alignment does not completely inhibit the amplification, so there is always some field generated to balance the resistive decay. This in turn also implies that the back reaction of the




FIG. 10. The total and conditional probability distribution functions of the cosines of the angles between \mathbf{u} and \mathbf{b} , $\cos(\theta)_{\mathbf{u}, \mathbf{b}}$ (a) and between \mathbf{j} and \mathbf{b} , $\cos(\theta)_{\mathbf{j}, \mathbf{b}}$ (b) for $\text{Re}_M = 1122$ in the kinematic (red) and saturated (blue) states. The magnetic field in the saturated stage is more aligned with the velocity field (reducing the induction effects) as compared to the kinematic stage. The magnetic field also becomes better aligned with the electric current density, reducing the back reaction on the velocity field.




FIG. 11. As Fig. 10 but for $\text{Re}_M = 1496$.

Lorentz force always remains significant.

B. Magnetic field stretching

To explore another mechanism by which magnetic field amplification can be suppressed, we consider the stretching of the magnetic field lines by the turbulent velocity. For this, we consider the alignment of the magnetic field with the eigenvectors of the rate of strain tensor. Neglecting the rather weak divergence of the flow, the symmetric 3×3 matrix $S_{ij} = \frac{1}{2} (u_{i,j} + u_{j,i})$ is calculated at each point in the domain using sixth-order finite differences, and its eigenvalues and eigenvectors are calculated. The eigenvalues are arranged in an increasing order, $\lambda_1 < \lambda_2 < \lambda_3$. The corresponding eigenvectors are $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$. The sum of the eigenvalues is close

to zero since the flow is nearly incompressible. λ_1 is always negative and the vector \mathbf{e}_1 corresponds to the direction of local compression of magnetic field, λ_3 is always positive and the vector \mathbf{e}_3 corresponds to the direction of local stretching, whereas λ_2 can be obtained from $\lambda_1 + \lambda_2 + \lambda_3 \approx 0$. The direction \mathbf{e}_2 (sometimes referred to as the ‘null’ direction [33, 74]) can correspond to either local stretching or compression depending on the sign of λ_2 . We then quantify the alignment with the magnetic field \mathbf{b} of the vectors \mathbf{e}_1 and \mathbf{e}_3 by considering

$$\cos(\theta)_{\mathbf{e}_1, \mathbf{b}} = \frac{\mathbf{e}_1 \cdot \mathbf{b}}{|\mathbf{e}_1| |\mathbf{b}|} \quad \text{and} \quad \cos(\theta)_{\mathbf{e}_3, \mathbf{b}} = \frac{\mathbf{e}_3 \cdot \mathbf{b}}{|\mathbf{e}_3| |\mathbf{b}|}. \quad (10)$$

Fig. 13 shows the PDF of the cosines in the kinematic and saturated stages for $\text{Re}_M = 1796$. In most of the volume, the direction of the magnetic field is perpendicular to the direction




FIG. 12. The cross correlation of $\cos(\theta)_{u,b}$ and $\cos(\theta)_{j,b}$ in the kinematic (a,c) and saturated (b,d) stages for $Re_M = 1122$. Panels (a) and (b) refer to the whole domain and the difference between them is not significant. Panels (c) and (d) refers to only the strong field regions ($b/b_{rms} \geq 1.5$). The yellow patch close to low $\cos(\theta)_{j,b}$ and high $\cos(\theta)_{u,b}$ in the kinematic stage vanishes for the saturated stage. The peak in the count is always at high $\cos(\theta)_{u,b}$ and high $\cos(\theta)_{j,b}$, which implies significant alignment between magnetic field, velocity field and current density.

of the local compression (Fig. 13a), which leads to the amplification of magnetic field, and this trend is slightly stronger in the kinematic stage. The PDF of the angle between the direction of local stretching and the magnetic field $\cos(\theta)_{e_3,b}$ has maxima at $\cos(\theta)_{e_3,b} = 0$ and $\cos(\theta)_{e_3,b} = 1$ in the kinematic stage. In the saturated stage, however, all angles are nearly equiprobable. This change in behaviour is more pronounced in the strong field regions, $b/b_{rms} \geq 1$. In Fig. 14a, we also show the PDF of $\cos(\theta)_{e_2,b}$, $\cos(\theta)_{e_1,b}$ and $\cos(\theta)_{e_3,b}$. The forms of the PDF for $\cos(\theta)_{e_2,b}$ are different from that of $\cos(\theta)_{e_1,b}$ and $\cos(\theta)_{e_3,b}$ in the kinematic and saturated stages. The magnetic field is less aligned to the direction e_2 in the kinematic stage as compared to the saturated stage and its effect, locally on the magnetic field, is decided by the sign of the eigenvalue λ_2 (dashed lines in Fig. 14b). Fig. 14b shows the PDF of all three eigenvalues in the kinematic and saturated stages. All three eigenvalues are statistically lower in

magnitude in the saturated stage as compared to the kinematic stage. However, as can be seen in Fig. 13 and Fig. 14a, the difference between the PDFs in the kinematic and saturated stages, whilst statistically significant, is not very strong. This suggest that a small reduction in the local stretching and compression of magnetic field contributes towards the saturation of the fluctuation dynamo.

Before concluding this section, we note that some of these conclusions are similar to those reached independently in the PhD thesis of Denis St-Onge [74], albeit for a different model setup.

C. Local magnetic energy balance

We now directly consider the equation for magnetic energy evolution and calculate its local growth and dissipation terms.




FIG. 13. The total and conditional PDFs of the cosine of the angle between the direction of local field line compression and the magnetic field $\cos(\theta)_{\mathbf{e}_1, \mathbf{b}}$ (a) and between the direction of local field line stretching and the magnetic field $\cos(\theta)_{\mathbf{e}_3, \mathbf{b}}$ (b) for $\text{Re}_M = 1796$ in the kinematic (red) and saturated (blue) stages.




FIG. 14. The PDFs of the cosine of the angle between three eigenvectors ($\mathbf{e}_1, \mathbf{e}_2$, and \mathbf{e}_3) with the local magnetic field direction (a) and three eigenvalues (λ_1, λ_2 , and λ_3) normalized by u_{rms} (b) for $\text{Re}_M = 1796$ in the kinematic (red) and saturated (blue) stages.

For an incompressible flow in a periodic domain, the magnetic energy evolution equation can be written as [75]

$$\frac{dE_M}{dt} = \int_V b_i b_j S_{ij} dV - \eta \int_V (\nabla \times \mathbf{b})^2 dV, \quad (11)$$

where $E_M = \frac{1}{2} \int_V \mathbf{b}^2 dV$ and summation over repeated indices is understood. The term contributing to the energy growth, $b_i b_j S_{ij}$, is calculated at each point in the volume as follows. First, we project the magnetic field vector \mathbf{b} on to each of the eigenvectors of the rate of strain tensor, $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$. Let these be $\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3$, and then the local growth term $b_i b_j S_{ij} = \lambda_1 \mathbf{b}_1^2 + \lambda_2 \mathbf{b}_2^2 + \lambda_3 \mathbf{b}_3^2$ at each position. This term can be positive or negative ($\lambda_1 < 0$ and $\lambda_3 > 0$). A negative local growth term leads to a decrease in the magnetic energy, whilst a positive value leads to an increase. The term contributing to the decay in energy is calculated by computing $(\nabla \times \mathbf{b})^2$ ($\eta = \text{constant}$)

at each point in space.

Fig. 15 and Fig. 16 show the total and conditional PDFs of the local growth and dissipation terms in the kinematic and saturated stages for $\text{Re}_M = 1122$ and $\text{Re}_M = 1796$ respectively. Fig. 15a and Fig. 16a show that the local growth term decreases on saturation and this is equally true of the strong and weak field regions. This confirms that the stretching of the magnetic field line reduces, which in turn decreases the amplification. Numerically, this can be quantified by calculating the skewness of the local growth term distribution in the kinematic and saturated stages (solid red and blue lines in Fig. 15a and Fig. 16a). The skewness is defined for a quantity X as $\langle (X - \langle X \rangle)^3 \rangle / \langle (X - \langle X \rangle)^2 \rangle^{3/2}$, where $\langle \dots \rangle$ refers to the mean. The skewness of the local growth term distribution in the kinematic (solid red line in Fig. 15a) and saturated (solid blue line in Fig. 15a) stage for $\text{Re}_M = 1122$ are 0.4 and




FIG. 15. The total and conditional PDFs of the local growth term $\lambda_1(\mathbf{b}_1/b_{\text{rms}})^2 + \lambda_2(\mathbf{b}_2/b_{\text{rms}})^2 + \lambda_3(\mathbf{b}_3/b_{\text{rms}})^2$ (a) and the local dissipation term $(\nabla \times \mathbf{b})^2$ (b) in the kinematic (red) and saturated (blue) stages for $\text{Re}_M = 1122$. The skewness of the local growth term distribution (solid red line in (a)) is 0.4 in the kinematic stage and 0.1 in the saturated stage, so the tendency of this term to promote growth decreases on saturation, as could be expected. The local dissipation term (b) also decreases statistically as the field saturates. These conclusions hold in both the weak and strong field regions, except for the local dissipation term, which increases in the weak field regions.




FIG. 16. As Fig. 15 but for $\text{Re}_M = 1796$. The skewness of the local growth term distribution (solid red line in (a)) is 0.9 in the kinematic stage and 0.4 in the saturated stage. The conclusions remain the same here as for $\text{Re}_M = 1122$ in Fig. 15.

0.1 respectively. The corresponding values for $\text{Re}_M = 1796$ (Fig. 16a) in the kinematic and saturated stages are 0.9 and 0.4 respectively. The local growth term always has a positive skewness implying continuous magnetic field generation. The skewness decreases on saturation, where the growth is only required to compensate the dissipation. The dissipation term also exhibits an overall decrease on saturation as shown in Fig. 15b and Fig. 16b, but its behaviour differs in the strong and weak field regions, where the dissipation increases in the latter regions.

To calculate the overall decrease or increase in the magnetic energy at each point in the domain, we calculate the local magnetic Reynolds number. This helps us to explore the behaviour of the diffusion term $(\eta \nabla^2 \mathbf{b})$ in the induction equation

(Eq. (2)) as the dynamo saturates. Both terms in Eq. (11) are calculated at each point in the volume, and the local magnetic Reynolds number is derived at each position as

$$(\text{Re}_M)_{\text{loc}} = \frac{b_i b_j S_{ij}}{\eta (\nabla \times \mathbf{b})^2}, \quad (12)$$

providing a measure of the local dynamo efficiency. The local magnetic Reynolds number can be positive or negative, signifying the locally increasing or decreasing magnetic field strength, respectively. Fig. 17 and Fig. 18 show the total and conditional PDFs of the local magnetic Reynolds number in the kinematic and saturated stages for $\text{Re}_M = 1122$ and $\text{Re}_M = 1796$. $(\text{Re}_M)_{\text{loc}}$ varies from values much less than to those much greater than $\text{Re}_M^{(\text{crit})}$ in both the kinematic and sat-




FIG. 17. The total and conditional PDFs of the local magnetic Reynolds number $(\text{Re}_M)_{\text{loc}}$ in the kinematic (red) and saturated (blue) stages with $\text{Re}_M = 1122$. The purple dashed line shows the critical magnetic Reynolds number $\text{Re}_M^{(\text{crit})} = 220$ and the black dashed line shows Re_M for this run.




FIG. 18. As Fig. 17 but for $\text{Re}_M = 1796$.

urated stages. Thus, magnetic field grows and decays in different parts of the volume but remains in a statistically steady state overall in the saturated stage. On saturation, both Fig. 17 and Fig. 18 show that $(\text{Re}_M)_{\text{loc}}$ decreases statistically. The mean of $(\text{Re}_M)_{\text{loc}}$ for $\text{Re}_M = 1122$ in the kinematic stage is 808 and that in the saturated stage is 595. Thus, the mean value of the local magnetic Reynolds number over the entire domain decreases on saturation (to a value close to but not exactly equal to the critical value, $\text{Re}_M^{(\text{crit})} \approx 220$). This effectively implies a relative enhancement in the local diffusion in comparison to the local stretching, which also contributes towards the saturation of the fluctuation dynamo.

To summarize, the fluctuation dynamo saturates due to both reduction in stretching and altered diffusion. The alignment between the velocity and magnetic fields increases as the field saturates, signifying reduced amplification. Furthermore, the

TABLE II. Four Minkowski functionals (MF) V_0, V_1, V_2 and V_3 , their geometrical interpretation and definitions in three dimensions. dV is the volume element, dS is the surface element, and κ_1 and κ_2 are the principle curvatures of the surface of a structure.

MF	Geometric interpretation	Expression
V_0	Volume	$\iiint dV$
V_1	Surface area	$(1/6) \iint dS$
V_2	Integral mean curvature	$(1/6\pi) \iint (\kappa_1 + \kappa_2) dS$
V_3	Euler characteristic	$(1/4\pi) \iint (\kappa_1 \kappa_2) dS$

current density and magnetic field are also statistically better aligned in the saturated stage, which implies a trend towards a force-free field. The local growth term statistically decreases (the skewness of the distribution, though remaining positive, decreases on saturation), which implies that the reduced magnetic field stretching reduces the amplification, which contributes towards the saturation of the fluctuation dynamo. The local magnetic Reynolds number, though varying over a wide range from values much less than to much higher than the critical value, decreases on average. This further implies relative enhancement in the local dissipation compared to the local stretching, which also contributes towards the saturation of the fluctuation dynamo.

V. MORPHOLOGY OF MAGNETIC STRUCTURES

As shown in Section III, magnetic field generated by a fluctuation dynamo is intermittent as it is concentrated in filaments, sheets and ribbons (Fig. 8 and Fig. 9). To characterize the magnetic structures, we use the Minkowski functionals [76]. Minkowski functionals have been used in studying morphology of structures in a number of numerical simulations [34, 77–82] and observations [83–86]. The morphology of a d -dimensional structure can be described by $d + 1$ Minkowski functionals. In three dimensions, there are four Minkowski functionals, as described in Table II. We calculate the Minkowski functionals using Crofton's formulae [87, 88] and then calculate the representative length scales (l_1, l_2, l_3) of magnetic structures (defined by isosurfaces at a fixed value of the magnetic field strength, e.g., see Fig. 9) as [77, 89]

$$l_1 = \frac{V_0}{2V_1}, \quad l_2 = \frac{2V_1}{\pi V_2}, \quad l_3 = \frac{3V_2}{4V_3}. \quad (13)$$

We associate the smallest of these length scales with the thickness T of the structures, the next largest with the width W and the largest length scale with the length L , i.e., if $l_1 \leq l_2 \leq l_3$, then $T = l_1, W = l_2$ and $L = l_3$. The thickness, width and length can be further used to obtain dimensionless measures of the structure shape: planarity p and filamentarity f , given by

$$p = \frac{W - T}{W + T}, \quad f = \frac{L - W}{L + W}. \quad (14)$$

By definition, $0 \leq p \leq 1$ and $0 \leq f \leq 1$; $p = 0$ and $f = 1$ for a perfect filament, $p = 0$ and $f = 0$ for a sphere, and $p = 1$

TABLE III. Parameters of various runs for the nonlinear fluctuation dynamo in a numerical domain size of $(2\pi)^3$ with 512^3 mesh points. In all cases, the forcing scale is approximately $L/5$, the forcing amplitude is $F_0 \approx 0.02$ and the hydrodynamic viscosity is $\nu = 4 \times 10^{-4}$. The magnetic diffusivity η , the rms velocity in the saturated stage u_{rms} , the Reynolds number Re , the magnetic Reynolds number Re_M , the magnetic Prandtl number Pr_M and the critical magnetic Reynolds number $\text{Re}_M^{(\text{crit})} (\approx 220\text{Pr}_M^{-1/2})$ are given.

η	u_{rms}	Re	Re_M	Pr_M	$\text{Re}_M^{(\text{crit})}$
4×10^{-4}	0.11	346	346	1.00	220
3×10^{-4}	0.11	346	461	1.33	191
2×10^{-4}	0.10	314	628	2.00	156
1×10^{-4}	0.09	283	1131	4.00	110
7.5×10^{-5}	0.09	283	1508	5.33	95
5×10^{-5}	0.09	283	2261	8.00	78

and $f = 0$ for a sheet. The planarity and filamentarity are not sensitive to the size of the structures but quantify the shape. It is useful to remember that, unlike the Minkowski functionals, p and f are not additive.

To explore the morphology of magnetic structures for a range of Re_M values, we use simulations with parameters given in Table III. We keep Re about the same for all runs, vary Re_M (making sure $\text{Pr}_M \geq 1$), and choose $k_F \approx 5(2\pi/L)$, so there is a sufficient number of magnetic correlation cells within the volume (with 5^3 velocity correlation cells).

Fig. 19a shows the thickness, width and length of magnetic structures obtained by averaging over 30 values of magnetic field strengths ranging from $b/b_{\text{rms}} = 2.5$ to 4. The lower limit of the magnetic field strength is chosen to ensure that the structures represent the tail of the PDF (e.g., see Fig. 6), whilst the upper limit is chosen to ensure a sufficient number of points within each structure. The computed values of planarity and filamentarity also remain roughly constant within this selected range of magnetic field strengths. For the kinematic stage, we expect that the largest length scale L will be independent of Re_M . This is because the length of the structures is controlled by the correlation length of the flow since the magnetic correlation function of the fastest growing dynamo mode decreases exponentially after that scale [32]. As seen in Fig. 19a, the length remains roughly constant but then increases slightly after $\text{Re}_M \approx 600$ and again remains roughly constant. This variation is likely to be due to the decrease in the Reynolds number Re (Table III). The other two scales (W and T) decrease as $\text{Re}_M^{-0.5}$. This scaling can be obtained by balancing the rate of magnetic dissipation with the local shearing rate [90], $\eta/W^2 \approx u_{\text{rms}}/l_0$, where η is the magnetic resistivity, u_{rms} is the rms turbulent velocity and l_0 is the driving scale of the turbulence. This gives $W \approx l_0(\eta/u_{\text{rms}}l_0)^{1/2} = l_0\text{Re}_M^{-0.5}$. This means that the shape of the magnetic structures becomes more filamentary ($L \gg W \approx T$) and ribbon-like ($T \lesssim W \ll L$) as Re_M increases, but the filamentarity is always larger than the planarity, so the filaments dominate among the magnetic structures [91]. The differences in Re_M scalings with the previ-

ous work [34] is probably due to the following reasons. First, they have a prescribed velocity field with forcing at a range of scales, whereas we force the flow at two scales ($k = 4$ and $k = 6$) and then let it evolve via the Navier-Stokes equation. Second, our simulations are at a higher resolution (512^3) as compared to theirs (128^3) and thus magnetic structures, especially at higher Re_M , are better resolved in our case. Last and most importantly, they consider values of Re_M which are both lower and higher than $\text{Re}_M^{(\text{crit})}$, whereas we only consider $\text{Re}_M > \text{Re}_M^{(\text{crit})}$. This is because we strongly believe that those two regimes ($\text{Re}_M < \text{Re}_M^{(\text{crit})}$ and $\text{Re}_M \geq \text{Re}_M^{(\text{crit})}$) are physically different and must not be considered together to characterize the length scales of magnetic structures as functions of Re_M .

All three scales are larger in the saturated stage than in the kinematic stage. Thus, the magnetic structures become larger as the magnetic field saturates. This is also the reason that the magnetic field correlation length scale increases as the field saturates (as shown in Table I). The increase in the length (the largest length scale) of magnetic structures on saturation is consistent with the finding by Schekochihin et al. [33]. The Re_M scaling for all three scales is roughly the same for both the kinematic and saturated stages.

Fig. 19b shows the planarity and filamentarity of magnetic structures as functions of Re_M . The filamentarity is always higher than the planarity and thus the magnetic structures are more like filaments in both the kinematic and saturated stages. The dependence of these morphological measures on Re_M is the same for the kinematic and saturated stages.

VI. CONCLUSIONS AND DISCUSSION

It is important to understand the saturated state of the fluctuation dynamo because the saturated state seeds the mean field dynamo, controls the small-scale magnetic field structure and decides the magnetic field length scales in the system where the mean field dynamo is absent (for example, elliptical galaxies). Moreover, it is crucial to understand the physics of the saturation mechanism because numerical simulations, at present, are at much lower values of Re_M than their estimated values ($\text{Re}_M \approx 10^{18}$ for spiral galaxies, $\text{Re}_M \approx 10^{22}$ for elliptical galaxies and $\text{Re}_M \approx 10^{29}$ for galaxy clusters).

Using numerical simulations of driven nearly incompressible turbulence, we have explored the saturation mechanism of the fluctuation dynamo. We find that the dynamo saturates because both the amplification and diffusion are affected by the action of the Lorentz force on the flow. Most previously suggested mechanisms hinted at changes in either of those two and thus required significant changes in the properties of the velocity and magnetic fields from the kinematic stage. For example, if only the enhancement in diffusion is responsible, it would require the effective Re_M in the saturated state to reduce from hugely supercritical levels to values close to $\text{Re}_M^{(\text{crit})}$ ($\approx 10^2$ – 10^3 [14]). And, if only the decrease in amplification is responsible for saturating the dynamo, it would require a drastic decrease in the Lyapunov exponents (which




FIG. 19. (a) Average length (L), thickness (T) and width (W) of magnetic structures in the kinematic (dashed, color) and saturated stages (solid, color) of the nonlinear fluctuation dynamo as functions of Re_M . The width and thickness of the magnetic structures both decrease as $Re_M^{-0.5}$. The Re_M dependence of the size of the structures is approximately the same in both the kinematic and saturated stages. (b) Planarity (p) and filamentarity (f) of the magnetic structures, as functions of Re_M , for the kinematic (dashed, color) and saturated (solid, color) stages. As Re_M increases, the filamentarity increases and the planarity decreases but they seem to approach an asymptotic value after $Re_M \approx 1200$. The trend with respect to Re_M is the same for both the kinematic and saturated stages.

are a measure of chaotic properties of the flow) [60]. We suggest that both occur and thus such a dramatic change is not necessary. We confirm that the amplification decreases by reduction in the stretching of magnetic field lines. The local magnetic Reynolds number $(Re_M)_{\text{loc}}$, which is suggested as a measure of the local magnetic diffusion, decreases slightly. This confirms that the local diffusion of magnetic field relative to field line stretching is enhanced, which is also responsible for saturating the dynamo.

The fluctuation dynamo-generated magnetic field is spatially intermittent. So, we studied the morphology of the magnetic structures in the kinematic and saturated stages. In both cases, the largest length scale is roughly independent of Re_M and the other two scales decrease as $Re_M^{-0.5}$. We find that the structures are of a larger size (all three length scales increase) in the saturated stage as compared to the kinematic stage. This agrees with the results in Table I, where we find that the correlation length is higher for the saturated magnetic field. This also aligns with the conclusion in the Section III (also shown in [33]) that the magnetic field is less intermittent in the saturated stage as compared to the kinematic stage. However, the Re_M dependence is the same for both the stages and thus the overall shape of magnetic structures produced by the fluctuation dynamo is not affected by the Lorentz force to any significant extent (all three length scales increase but in a very similar way).

The study explores physical effects over a range of Re_M for $Pr_M \geq 1$. However, for $Pr_M > 1$ at very high Re_M ($\gtrsim 10^3$), the fields might be unstable to fast magnetic reconnection [12]. This might change the morphology of magnetic fields, locally affect velocity fields and thus might alter the saturated state of the fluctuation dynamo. However, the effect of fast, stochastic magnetic reconnection on the dynamo is not very well under-

stood yet [92] and would require high-resolution numerical simulations over a number of very high Re_M values to study the effect of fast magnetic reconnection on the fluctuation dynamo saturation mechanism.

The study can be extended in several ways. An immediate extension would be to repeat the entire analysis for dynamos in a stratified medium [42, 46, 47, 93], which is more relevant for young galaxies and star-forming gas clouds. We have performed the analysis for $Pr_M \geq 1$ which is of relevance to fluctuation dynamo in the interstellar and intergalactic medium but this should be extended to the $Pr_M < 1$ regime which is important for stars, planets and liquid metal experiments [94, 95]. We have adopted the MHD approximation but plasma effects might also play an important role. It would also be interesting to compare our results with those of the plasma dynamo [96, 97] and see how the relationship between velocity and magnetic fields and the magnetic field structure change when plasma effects are considered. Plasma effects might be particularly important for the weakly collisional gas in galaxy clusters. We aim to consider such problems in our future work.

ACKNOWLEDGMENTS

We thank Kandaswamy Subramanian and Christoph Federath for useful discussions and comments on the paper. We acknowledge financial support of the STFC (ST/N000900/1, Project 2) and the Leverhulme Trust (RPG-2014-427). We thank one of the referees for highlighting Denis St-Onge's PhD thesis [74], which independently reports results similar to those found in Section IV B. In fact, Fig. 14 was added following the initial review so as to facilitate a direct comparison

between the two studies.

-
- [1] B. M. Gaensler, M. Haverkorn, L. Staveley-Smith, J. M. Dickey, N. M. McClure-Griffiths, J. R. Dickel, and M. Wolleben, “The Magnetic Field of the Large Magellanic Cloud Revealed Through Faraday Rotation,” *Science* **307**, 1610–1612 (2005), [astro-ph/0503226](#).
- [2] A. Fletcher, R. Beck, A. Shukurov, E. M. Berkhuijsen, and C. Horellou, “Magnetic fields and spiral arms in the galaxy M51,” *Mon. Not. R. Astron. Soc.* **412**, 2396–2416 (2011).
- [3] M. Houde, A. Fletcher, R. Beck, R. H. Hildebrand, J. E. Vaillancourt, and J. M. Stil, “Characterizing Magnetized Turbulence in M51,” *Astrophys. J.* **766**, 49 (2013), [arXiv:1301.7508](#).
- [4] Federica Govoni and Luigina Feretti, “Magnetic Fields in Clusters of Galaxies,” *International Journal of Modern Physics D* **13**, 1549–1594 (2004).
- [5] A. A. Schekochihin and S. C. Cowley, “Turbulence, magnetic fields, and plasma physics in clusters of galaxies,” *Physics of Plasmas* **13**, 056501–056501 (2006).
- [6] R. Beck, “Magnetic fields in spiral galaxies,” *Ann. Rev. Astron. Astrophys.* **24**, 4 (2016), [arXiv:1509.04522](#).
- [7] R. Beck, A. Brandenburg, D. Moss, A. Shukurov, and D. Sokoloff, “Galactic Magnetism: Recent Developments and Perspectives,” *Ann. Rev. Astron. Astrophys.* **34**, 155–206 (1996).
- [8] L. M. Widrow, “Origin of galactic and extragalactic magnetic fields,” *Reviews of Modern Physics* **74**, 775–823 (2002), [astro-ph/0207240](#).
- [9] A. Brandenburg and K. Subramanian, “Astrophysical magnetic fields and nonlinear dynamo theory,” *Phys. Rep.* **417**, 1–209 (2005).
- [10] Russell M. Kulsrud and Ellen G. Zweibel, “On the origin of cosmic magnetic fields,” *Reports on Progress in Physics* **71** (2008), [10.1088/0034-4885/71/4/046901](#).
- [11] Christoph Federrath, “Magnetic field amplification in turbulent astrophysical plasmas,” *Journal of Plasma Physics* **82**, 535820601 (2016).
- [12] François Rincon, “Dynamo theories,” *Journal of Plasma Physics* **85**, 205850401 (2019), [arXiv:1903.07829 \[physics.plasm-ph\]](#).
- [13] P. Tzeferacos, A. Rigby, A. F. A. Bott, A. R. Bell, R. Bingham, A. Casner, F. Cattaneo, E. M. Churazov, J. Emig, F. Fiuza, C. B. Forest, J. Foster, C. Graziani, J. Katz, M. Koenig, C. K. Li, J. Meinecke, R. Petrasso, H. S. Park, B. A. Remington, J. S. Ross, D. Ryu, D. Ryutov, T. G. White, B. Reville, F. Miniati, A. A. Schekochihin, D. Q. Lamb, D. H. Froula, and G. Gregori, “Laboratory evidence of dynamo amplification of magnetic fields in a turbulent plasma,” *Nature Communications* **9**, 591 (2018), [arXiv:1702.03016 \[physics.plasm-ph\]](#).
- [14] A. P. Kazantsev, “Enhancement of a Magnetic Field by a Conducting Fluid,” *Soviet Journal of Experimental and Theoretical Physics* **26**, 1031 (1968).
- [15] Ya. B. Zel’dovich, A. A. Ruzmaikin, S. A. Molchanov, and D. D. Sokoloff, “Kinematic dynamo problem in a linear velocity field,” *Journal of Fluid Mechanics* **144**, 1–11 (1984).
- [16] S. Childress and A. D. Gilbert, *The Fast Dynamo*, XI, 406 pp., Springer-Verlag Berlin Heidelberg New York. Also *Lecture Notes in Physics*, volume 37 (1995).
- [17] A. Seta, P. Bhat, and K. Subramanian, “Saturation of Zeldovich stretch-twist-fold map dynamos,” *Journal of Plasma Physics* **81**, 395810503 (2015), [arXiv:1410.8455](#).
- [18] A. A. Ruzmaikin, D. D. Sokoloff, and A. M. Shukurov, eds., *Astrophysics and Space Science Library*, Astrophysics and Space Science Library, Vol. 133 (1988).
- [19] R. M. Kulsrud and S. W. Anderson, “The spectrum of random magnetic fields in the mean field dynamo theory of the Galactic magnetic field,” *Astrophys. J.* **396**, 606–630 (1992).
- [20] A. Shukurov and D. Sokoloff, “Astrophysical dynamos,” in *Les Houches, Session LXXXVIII, Dynamos*, Vol. 88, edited by P. Cardin and L. F. Cugliandolo (Amsterdam: Elsevier, 2007) pp. 251–299.
- [21] Rüdiger Pakmor, Facundo A. Gómez, Robert J. J. Grand, Federico Marinacci, Christine M. Simpson, Volker Springel, David J. R. Campbell, Carlos S. Frenk, Thomas Guillet, Christoph Pfrommer, and Simon D. M. White, “Magnetic field formation in the Milky Way like disc galaxies of the Auriga project,” *Mon. Not. R. Astron. Soc.* **469**, 3185–3199 (2017).
- [22] A. H. Minter and S. R. Spangler, “Observation of Turbulent Fluctuations in the Interstellar Plasma Density and Magnetic Field on Spatial Scales of 0.01 to 100 Parsecs,” *Astrophys. J.* **458**, 194 (1996).
- [23] Amit Seta, “Magnetic fields in elliptical galaxies,” (2019).
- [24] A. Ruzmaikin, D. Sokoloff, and A. Shukurov, “The dynamo origin of magnetic fields in galaxy clusters,” *Mon. Not. R. Astron. Soc.* **241**, 1–14 (1989).
- [25] K. Subramanian, A. Shukurov, and N. E. L. Haugen, “Evolving turbulence and magnetic fields in galaxy clusters,” *Mon. Not. R. Astron. Soc.* **366**, 1437–1454 (2006), [astro-ph/0505144](#).
- [26] Pallavi Bhat and Kandaswamy Subramanian, “Fluctuation dynamos and their Faraday rotation signatures,” *Mon. Not. R. Astron. Soc.* **429**, 2469–2481 (2013).
- [27] F. Vazza, G. Brunetti, M. Brüggen, and A. Bonafede, “Resolved magnetic dynamo action in the simulated intracluster medium,” *Mon. Not. R. Astron. Soc.* **474**, 1672–1687 (2018).
- [28] Fausto Cattaneo, “On the Origin of Magnetic Fields in the Quiet Photosphere,” *Astrophys. J.* **515**, L39–L42 (1999).
- [29] Jonathan Pietarila Graham, Robert Cameron, and Manfred Schüssler, “Turbulent Small-Scale Dynamo Action in Solar Surface Simulations,” *Astrophys. J.* **714**, 1606–1616 (2010), [arXiv:1002.2750 \[astro-ph.SR\]](#).
- [30] P. J. Bushby and B. Favier, “Mesogranulation and small-scale dynamo action in the quiet Sun,” *Astron. Astrophys.* **562**, A72 (2014).
- [31] M. Rempel, “Numerical Simulations of Quiet Sun Magnetism: On the Contribution from a Small-scale Dynamo,” *Astrophys. J.* **789**, 132 (2014), [arXiv:1405.6814 \[astro-ph.SR\]](#).
- [32] Ya. B. Zeldovich, A. A. Ruzmaikin, and D. D. Sokoloff, *The Almighty Chance* (World Scientific, Singapore, 1990).
- [33] A. A. Schekochihin, S. C. Cowley, S. F. Taylor, J. L. Maron, and J. C. McWilliams, “Simulations of the Small-Scale Turbulent Dynamo,” *Astrophys. J.* **612**, 276–307 (2004), [astro-ph/0312046](#).
- [34] S. L. Wilkin, C. F. Barenghi, and A. Shukurov, “Magnetic Structures Produced by the Small-Scale Dynamo,” *Phys. Rev. Lett.* **99**, 134501 (2007).
- [35] A. Shukurov, A. P. Snodin, A. Seta, P. J. Bushby, and T. S. Wood, “Cosmic Rays in Intermittent Magnetic Fields,” *Astrophys. J. Lett.* **839**, L16 (2017), [arXiv:1702.06193 \[astro-ph\]](#).

- ph.HE].
- [36] A. Seta, A. Shukurov, T. S. Wood, P. J. Bushby, and A. P. Snodin, “Relative distribution of cosmic rays and magnetic fields,” *Mon. Not. R. Astron. Soc.* **473**, 4544–4557 (2018), arXiv:1708.07499.
 - [37] T. A. Enßlin and C. Vogt, “Magnetic turbulence in cool cores of galaxy clusters,” *Astron. Astrophys.* **453**, 447–458 (2006), arXiv:astro-ph/0505517 [astro-ph].
 - [38] M. Meneguzzi, U. Frisch, and A. Pouquet, “Helical and non-helical turbulent dynamos,” *Physical Review Letters* **47**, 1060–1064 (1981).
 - [39] N. E. Haugen, A. Brandenburg, and W. Dobler, “Simulations of nonhelical hydromagnetic turbulence,” *Phys. Rev. E* **70**, 016308 (2004), astro-ph/0307059.
 - [40] Jungyeon Cho and Dongsu Ryu, “Characteristic Lengths of Magnetic Field in Magnetohydrodynamic Turbulence,” *Astrophys. J.* **705**, L90–L94 (2009), arXiv:0908.0610 [astro-ph.CO].
 - [41] Fausto Cattaneo and Steven M. Tobias, “Dynamo properties of the turbulent velocity field of a saturated dynamo,” *Journal of Fluid Mechanics* **621**, 205 (2009).
 - [42] C. Federrath, G. Chabrier, J. Schober, R. Banerjee, R. S. Klessen, and D. R. G. Schleicher, “Mach Number Dependence of Turbulent Magnetic Field Amplification: Solenoidal versus Compressive Flows,” *Phys. Rev. Lett.* **107**, 114504 (2011), arXiv:1109.1760 [physics.flu-dyn].
 - [43] B. Favier and P. J. Bushby, “Small-scale dynamo action in rotating compressible convection,” *Journal of Fluid Mechanics* **690**, 262–287 (2012), arXiv:1110.0374.
 - [44] Sharanya Sur, Christoph Federrath, Dominik R. G. Schleicher, Robi Banerjee, and Ralf S. Klessen, “Magnetic field amplification during gravitational collapse - influence of turbulence, rotation and gravitational compression,” *Mon. Not. R. Astron. Soc.* **423**, 3148–3162 (2012), arXiv:1202.3206 [astro-ph.SR].
 - [45] A. Beresnyak, “Universal Nonlinear Small-Scale Dynamo,” *Phys. Rev. Lett.* **108**, 035002 (2012), arXiv:1109.4644 [astro-ph.GA].
 - [46] Christoph Federrath, Jennifer Schober, Stefano Bovino, and Dominik R. G. Schleicher, “The Turbulent Dynamo in Highly Compressible Supersonic Plasmas,” *Astrophys. J.* **797**, L19 (2014), arXiv:1411.4707 [astro-ph.GA].
 - [47] Sharanya Sur, Pallavi Bhat, and Kandaswamy Subramanian, “Faraday rotation signatures of fluctuation dynamos in young galaxies,” *Mon. Not. R. Astron. Soc.* **475**, L72–L76 (2018).
 - [48] Stanislav Boldyrev and Fausto Cattaneo, “Magnetic-Field Generation in Kolmogorov Turbulence,” *Phys. Rev. Lett.* **92**, 144501 (2004), arXiv:astro-ph/0310780 [astro-ph].
 - [49] A. A. Schekochihin, N. E. L. Haugen, A. Brandenburg, S. C. Cowley, J. L. Maron, and J. C. McWilliams, “The Onset of a Small-Scale Turbulent Dynamo at Low Magnetic Prandtl Numbers,” *Astrophys. J.* **625**, L115–L118 (2005), arXiv:astro-ph/0412594 [Astrophysics].
 - [50] A. A. Schekochihin, A. B. Iskakov, S. C. Cowley, J. C. McWilliams, M. R. E. Proctor, and T. A. Yousef, “Fluctuation dynamo and turbulent induction at low magnetic Prandtl numbers,” *New Journal of Physics* **9**, 300 (2007), arXiv:0704.2002.
 - [51] A. B. Iskakov, A. A. Schekochihin, S. C. Cowley, J. C. McWilliams, and M. R. E. Proctor, “Numerical Demonstration of Fluctuation Dynamo at Low Magnetic Prandtl Numbers,” *Phys. Rev. Lett.* **98**, 208501 (2007), arXiv:astro-ph/0702291 [astro-ph].
 - [52] A. Brandenburg, N. E. L. Haugen, Xiang-Yu Li, and K. Subramanian, “Varying the forcing scale in low Prandtl number dynamos,” *Mon. Not. R. Astron. Soc.* **479**, 2827–2833 (2018), arXiv:1805.01249 [physics.flu-dyn].
 - [53] Website: <https://github.com/pencil-code>.
 - [54] It is also common to define the hydrodynamic and magnetic Reynolds number with respect to the forcing wave number instead of the driving length scale. Then the Reynolds numbers are smaller by a factor 2π than the values we quote.
 - [55] Pallavi Bhat and Kandaswamy Subramanian, “Fluctuation Dynamo at Finite Correlation Times and the Kazantsev Spectrum,” *Astrophys. J.* **791**, L34 (2014).
 - [56] Pallavi Bhat and Kandaswamy Subramanian, “Fluctuation dynamos at finite correlation times using renewing flows,” *Journal of Plasma Physics* **81**, 395810502 (2015), arXiv:1411.0885 [astro-ph.GA].
 - [57] A. Vincent and M. Meneguzzi, “The spatial structure and statistical properties of homogeneous turbulence,” *Journal of Fluid Mechanics* **225**, 1–20 (1991).
 - [58] Christoph Federrath, “On the universality of supersonic turbulence,” *Mon. Not. R. Astron. Soc.* **436**, 1245–1257 (2013).
 - [59] A. H. Waelkens, A. A. Schekochihin, and T. A. Enßlin, “Probing magnetic turbulence by synchrotron polarimetry: statistics and structure of magnetic fields from Stokes correlators,” *Mon. Not. R. Astron. Soc.* **398**, 1970–1988 (2009), arXiv:0903.3056.
 - [60] F. Cattaneo, D. W. Hughes, and E.-J. Kim, “Suppression of chaos in a simplified nonlinear dynamo model,” *Phys. Rev. Lett.* **76**, 2057–2060 (1996).
 - [61] Eun-jin Kim, “Nonlinear dynamo in a simplified statistical model,” *Physics Letters A* **259**, 232–239 (1999).
 - [62] A. A. Schekochihin, S. C. Cowley, G. W. Hammett, J. L. Maron, and J. C. McWilliams, “A model of nonlinear evolution and saturation of the turbulent MHD dynamo,” *New Journal of Physics* **4**, 84 (2002).
 - [63] K. Subramanian, “Unified Treatment of Small- and Large-Scale Dynamos in Helical Turbulence,” *Phys. Rev. Lett.* **83**, 2957–2960 (1999).
 - [64] Kandaswamy Subramanian, “Hyperdiffusion in Nonlinear Large- and Small-Scale Turbulent Dynamos,” *Phys. Rev. Lett.* **90**, 245003 (2003).
 - [65] S. I. Braginskii, “Transport Processes in a Plasma,” *Reviews of Plasma Physics* **1**, 205 (1965).
 - [66] Leonid Malyshkin and Russell M. Kulsrud, “Magnetized Turbulent Dynamos in Protogalaxies,” *Astrophys. J.* **571**, 619–637 (2002).
 - [67] A. Brandenburg, R. L. Jennings, Å. Nordlund, M. Rieutord, R. F. Stein, and I. Tuominen, “Magnetic structures in a dynamo simulation,” *Journal of Fluid Mechanics* **306**, 325–352 (1996).
 - [68] Joanne Mason, Fausto Cattaneo, and Stanislav Boldyrev, “Dynamic Alignment in Driven Magnetohydrodynamic Turbulence,” *Phys. Rev. Lett.* **97**, 255002 (2006).
 - [69] S. Servidio, W. H. Matthaeus, and P. Dmitruk, “Depression of Nonlinearity in Decaying Isotropic MHD Turbulence,” *Phys. Rev. Lett.* **100**, 095005 (2008).
 - [70] Mahendra K. Verma, “Statistical theory of magnetohydrodynamic turbulence: recent results,” *Phys. Rep.* **401**, 229–380 (2004), arXiv:nlin/0404043 [nlin.CD].
 - [71] Rohit Kumar, Mahendra K. Verma, and Ravi Samtaney, “Energy transfers and magnetic energy growth in small-scale dynamo,” *EPL (Europhysics Letters)* **104**, 54001 (2013).
 - [72] Franck Plunian, Rodion Stepanov, and Peter Frick, “Shell models of magnetohydrodynamic turbulence,” *Phys. Rep.* **523**, 1–60 (2013), arXiv:1209.3844 [physics.flu-dyn].
 - [73] Mahendra K. Verma and Rohit Kumar, “Dynamos at extreme magnetic Prandtl numbers: insights from shell models,” *Journal of Turbulence* **17**, 1112–1141 (2016), arXiv:1510.00122 [physics.flu-dyn].
 - [74] Denis A. St-Onge, “Fluctuation Dynamo in Collisionless and

- Weakly Collisional Magnetized Plasmas,” arXiv e-prints , arXiv:1912.11072 (2019), [arXiv:1912.11072 \[physics.plasm-ph\]](#).
- [75] P. H. Roberts, *An Introduction to Magnetohydrodynamics*, by P. H. Roberts. Textbook published by Longmans, Green and Co Ltd, London, 1967 (1967).
- [76] Hermann Minkowski, “Volumen und oberflche,” *Mathematische Annalen* **57**, 447–495 (1903).
- [77] J. Schmalzing, T. Buchert, A. L. Melott, V. Sahni, B. S. Sathyaprakash, and S. F. Shandarin, “Disentangling the Cosmic Web. I. Morphology of Isodensity Contours,” *Astrophys. J.* **526**, 568–578 (1999), [astro-ph/9904384](#).
- [78] T. Leung, N. Swaminathan, and P. A. Davidson, “Geometry and interaction of structures in homogeneous isotropic turbulence,” *Journal of Fluid Mechanics* **710**, 453–481 (2012).
- [79] V. Zhdankin, S. Boldyrev, J. C. Perez, and S. M. Tobias, “Energy Dissipation in Magnetohydrodynamic Turbulence: Coherent Structures or “Nanoflares”?” *Astrophys. J.* **795**, 127 (2014), [arXiv:1409.3285 \[astro-ph.SR\]](#).
- [80] Akanksha Kapahtia, Pravabati Chingangbam, Stephen Appleby, and Changbom Park, “A novel probe of ionized bubble shape and size statistics of the epoch of reionization using the contour Minkowski Tensor,” *J. Cosmol. Astropart. Phys.* **2018**, 011 (2018), [arXiv:1712.09195 \[astro-ph.CO\]](#).
- [81] Satadru Bag, Rajesh Mondal, Prakash Sarkar, Somnath Bhargadwaj, and Varun Sahni, “The shape and size distribution of H II regions near the percolation transition,” *Mon. Not. R. Astron. Soc.* **477**, 1984–1992 (2018), [arXiv:1801.01116 \[astro-ph.CO\]](#).
- [82] Satadru Bag, Rajesh Mondal, Prakash Sarkar, Somnath Bhargadwaj, Tirthankar Roy Choudhury, and Varun Sahni, “Studying the morphology of HI isodensity surfaces during reionization using Shapefinders and percolation analysis,” *Mon. Not. R. Astron. Soc.* , 527 (2019), [arXiv:1809.05520 \[astro-ph.CO\]](#).
- [83] J. Schmalzing and K. M. Gorski, “Minkowski functionals used in the morphological analysis of cosmic microwave background anisotropy maps,” *Mon. Not. R. Astron. Soc.* **297**, 355–365 (1998), [astro-ph/9710185](#).
- [84] S. Bhargadwaj, V. Sahni, B. S. Sathyaprakash, S. F. Shandarin, and C. Yess, “Evidence for Filamentarity in the Las Campanas Redshift Survey,” *Astrophys. J.* **528**, 21–29 (2000), [astro-ph/9904406](#).
- [85] I. Makarenko, A. Fletcher, and A. Shukurov, “3D morphology of a random field from its 2D cross-section,” *Mon. Not. R. Astron. Soc.* **447**, L55–L59 (2015), [arXiv:1407.4048 \[physics.flu-dyn\]](#).
- [86] P. K. Joby, Pravabati Chingangbam, Tuhin Ghosh, Vidhya Ganesan, and C. D. Ravikumar, “Search for anomalous alignments of structures in Planck data using Minkowski Tensors,” *Journal of Cosmology and Astro-Particle Physics* **2019**, 009 (2019), [arXiv:1807.01306 \[astro-ph.CO\]](#).
- [87] M. W. Crofton, “On the Theory of Local Probability, Applied to Straight Lines Drawn at Random in a Plane; The Methods Used Being Also Extended to the Proof of Certain New Theorems in the Integral Calculus,” *Philosophical Transactions of the Royal Society of London Series I* **158**, 181–199 (1868).
- [88] D. Legland, K. Ki  u, and M-F. Devaux, “COMPUTATION OF MINKOWSKI MEASURES ON 2D AND 3D BINARY IMAGES,” *Image Analysis & Stereology* **2**, 83–92 (2011).
- [89] V. Sahni, B. S. Sathyaprakash, and S. F. Shandarin, “Shapefinders: A New Shape Diagnostic for Large-Scale Structure,” *Astrophys. J. Lett.* **495**, L5–L8 (1998), [astro-ph/9801053](#).
- [90] Kandaswamy Subramanian, “Can the turbulent galactic dynamo generate large-scale magnetic fields?” *Mon. Not. R. Astron. Soc.* **294**, 718–728 (1998).
- [91] For $\text{Pr}_M > 1$, fast magnetic reconnection at very high Re_M might alter the shape of the structures [12], see Section VI for further discussion.
- [92] Gregory L. Eyink, “Stochastic flux freezing and magnetic dynamo,” *Phys. Rev. E* **83**, 056405 (2011).
- [93] Nils Erland L. Haugen, Axel Brandenburg, and Antony J. Mee, “Mach number dependence of the onset of dynamo action,” *Mon. Not. R. Astron. Soc.* **353**, 947–952 (2004), [arXiv:astro-ph/0405453 \[astro-ph\]](#).
- [94] Axel Brandenburg, “Nonlinear Small-scale Dynamos at Low Magnetic Prandtl Numbers,” *Astrophys. J.* **741**, 92 (2011), [arXiv:1106.5777 \[astro-ph.SR\]](#).
- [95] Ganapati Sahoo, Prasad Perlekar, and Rahul Pandit, “Systematics of the magnetic-Prandtl-number dependence of homogeneous, isotropic magnetohydrodynamic turbulence,” *New Journal of Physics* **13**, 013036 (2011), [arXiv:1006.5585 \[nlin.CD\]](#).
- [96] Fran  ois Rincon, Francesco Califano, Alexander A. Schekochihin, and Francesco Valentini, “Turbulent dynamo in a collisionless plasma,” *Proceedings of the National Academy of Science* **113**, 3950–3953 (2016).
- [97] Denis A. St-Onge and Matthew W. Kunz, “Fluctuation Dynamo in a Collisionless, Weakly Magnetized Plasma,” *Astrophys. J.* **863**, L25 (2018).