

Overcharging problem and thermodynamics in five-dimensional black holes

Deyou Chen and Shan Zeng

School of Science, Xihua University, Chengdu 610039, China
E-mails: deyouchen@hotmail.com, shanzengxhu@hotmail.com

Abstract: In this paper, the overcharging problem and thermodynamics in the extended phase spaces of the five-dimensional spherically symmetric topological black holes are investigated by absorptions of scalar particles and fermions. The cosmological constant is regarded as a variable related to pressure and its conjugate quantity is a thermodynamic volume. The first law of thermodynamics is recovered. The second law is violated in the extended phase space of the extremal and near-extremal black holes. The overcharging problem is tested by the existence of the event horizons. The event horizon is determined by the metric component $f(r)$. The minimal values of the metric component at the final stage show that the extremal and near-extremal black holes can not be overcharged.

1 Introduction

It is well known that a spacetime singularity can be formed at the end of gravitational collapse. Near the singularity, physical laws fail. To avoid the physical damage caused by the singularity, Penrose pointed out that under normal material properties and initial conditions, naked singularities cannot be formed in the real physical process [1]. This is the weak form of the cosmic censorship conjecture. It shows that singularities are not naked and hide behind event horizons. This cosmic censorship conjecture plays an important role in black hole physics, and its validity is a necessary condition to ensure the predictability of physical laws. Although its correctness is widely accepted, there is no complete evidence to prove it.

An effective method to test the weak cosmic censorship conjecture (WCCC) is the Gedanken experiment designed by Wald [2]. In this experiment, a particle with energy, large enough angular momentum and charge is thrown into a charged rotating black hole to test the evolution of the black hole. The evolution result is described by the new spacetime metric. If the final state of the black hole evolution is no longer a black hole, there is no event horizon and the singularity is naked. The solution of the event horizon is determined by the metric components. Therefore, one only needs to check the existence of the horizon to judge whether the singularity is naked. The energy, charge and angular momentum of the black hole can also be gotten by the interaction between the black hole and a field. The field has a finite energy (i.e. it contains a wave packet). At first, the field does not exist, there is only the black hole. Then, the field comes from infinity and interacts with the black hole. Due to the interaction, energy, charge and angular momentum are transferred between the black hole and field. Part of the field is absorbed by the black hole and the other part is reflected back to infinity. Finally, the field decays, and the energy, charge and angular momentum of

the black hole change. Using this field view, Semiz first adopted a complex scalar field to test the WCCC and found that the dyonic Kerr-Newman black hole could not be overcharged [3]. This result was supported by the work of Toth [4]. Based on the seed work of Wald, the validity of the WCCC in the various spacetimes has been tested [5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28]. In the relevant discussions, the self-force effect, tunnelling effect, radiation reaction were taken into account [29, 30, 31, 32]. However, there is no consensus. In these researches, the black holes are assumed to exist in advance. The formation of the naked singularity was also discussed by the numerical method [33]. In the discussion, it was assumed that a celestial body before particles being absorbed is not a black hole, but a general dense star.

The cosmological constant used to be a fixed constant. Recently, it was seen as a variable in the work of Kastor and others [34, 35, 36, 37]. One of the reasons for this consideration is that the cosmological constant, as a variable, can reconcile the inconsistency between the first law of thermodynamics of black holes and the Smarr relation derived from the scaling method. The second reason is that if people consider "more fundamental" theories, physical constants, such as the gauge coupling constant, Yukawa coupling constant, Newtonian constant or cosmological constant arise as vacuum expectation values are not fixed in advance and can vary [38]. The cosmological constant was seen as a pressure, and its conjugate quantity is a thermodynamic volume. Using the standard thermodynamics, Kubiznak and Mann studied the critical behavior of the charged anti-de Sitter black hole. They found that its critical index was consistent with that of van der Waals system [37]. The original first law of black hole thermodynamics was modified to

$$dM = TdS + \phi dQ + VdP + \Omega dJ. \quad (1)$$

The corresponding Smarr relation is

$$(n-3)M = (n-2)TS + (n-2)\Omega J + (n-3)\phi Q - 2VP. \quad (2)$$

where the mass M is expressed as the enthalpy and n is the dimension of the space-time. Introducing the thermodynamic pressure and volume into the black hole system, one can use the standard thermodynamic formulae to calculate various thermodynamic quantities and discuss various thermodynamic phenomena. In the extended phase space, many interesting thermodynamic phenomena have been found, including the reentrant phase transition, repulsive interaction in the high temperature black holes, multiphase transition in the quasi topological gravity and Lovelock gravity, and other thermodynamic phenomena [39, 40, 41, 42, 43, 44, 45].

In the recent work, Gwak studied the WCCC in the extended phase space of the rotating black hole, where the pressure and volume were introduced [46]. He found that the black hole could not be overspun by throwing a particle with energy and large enough angular momentum into the hole. This result is in consistence with that gotten in the new version Gedanken experiment [47, 48, 49, 50, 51, 52]. In the

extremal and near-extremal cases, the entropies decreased. This means the violation of the second law of thermodynamics in the extended phase space. The same result was gotten in the subsequent work [53, 54, 55, 56, 57, 58, 59, 60]. In these work, an important relation between the momentum, energy and angular momentum or charge of the particle was gotten by the Hamilton-Jacobi equation.

In this paper, we investigate the WCCC and thermodynamics in extended phase spaces of five-dimensional charged topological black holes by absorptions of scalar particles and fermions. These black holes are a spacial case of a $(n+1)$ -dimensional Einstein-Maxwell-dilaton gravity solution [61], where the dilaton $\alpha = 0$. The test particles in this paper are scalar particles and fermions, which satisfy the Klein-Gordon equation and Dirac equation in curved spacetime, respectively. The first law of thermodynamics is recovered by the absorptions of the particles. In the extremal and near-extremal cases, the second law is violated in the extended phase spaces. The overcharging problem is tested by the existence of the event horizon. The minimal values of the metric component $f(r)$ at the final stages show that the extremal and near-extremal black holes can not be overcharged.

The rest is organized as follows. The five-dimensional topological black hole is given and the thermodynamics are discussed in the next section. In section 3, we investigate the absorptions of the scalar particle and fermion by the black hole. The relation between the momentum, energy and charge of the particle is gotten. In section 4, the thermodynamics in the extended phase space are investigated by the absorptions of the particles. In section 5, the overcharging problem is tested by throwing a particle in the near-extremal and extremal black holes. The last section is devoted to our discussion and conclusion.

2 The black hole solution

The action described the $(n+1)$ -dimensional Einstein-Maxwell-dilaton gravity is [62]

$$\mathcal{S} = \frac{1}{16\pi} \int dx^{n+1} \sqrt{-g} \left[\mathcal{R} - \frac{4}{n-1} (\nabla\Phi)^2 - V(\Phi) - e^{-\frac{4\alpha\Phi}{n-1}} F_{\mu\nu} F^{\mu\nu} \right], \quad (3)$$

where $n \geq 3$, \mathcal{R} is the Ricci scalar curvature, Φ is a dilaton field, and $V(\Phi)$ is a potential for Φ ,

$$V(\Phi) = 2\Lambda e^{\frac{4\alpha\Phi}{n-1}} + \frac{k(n-1)(n-2)\alpha^2}{b^2(\alpha^2-1)} e^{\frac{4\Phi}{(n-1)\alpha}}. \quad (4)$$

Λ is a negative cosmological constant. α is a constant determining the coupling strength of the scalar and electromagnetic fields. $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the electromagnetic field tensor and A_μ denotes the electromagnetic potential. The black hole solutions in the Einstein-Maxwell-dilaton gravity with the dilaton coupling parameter α are given by [61]

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2 R^2(r) d\Omega_{k,n-1}^2, \quad (5)$$

where

$$f(r) = -\frac{k(n-2)(\alpha^2+1)^2b^{-2\gamma}r^{2\gamma}}{(\alpha^2-1)(\alpha^2+n-2)} - \frac{m}{r^{(n-1)(1-\gamma)-1}} + \frac{2q^2(\alpha^2+1)^2b^{-2(n-2)\gamma}r^{2(n-2)(\gamma-1)}}{(\alpha^2-1)(\alpha^2+n-2)} + \frac{2\Lambda(\alpha^2+1)^2b^{2\gamma}r^{2(1-\gamma)}}{(n-1)(\alpha^2-n)}, \quad (6)$$

$R(r) = e^{\frac{2\alpha\Phi}{n-1}}$, $\Phi(r) = \frac{(n-1)\alpha}{2(1+\alpha^2)} \ln\left(\frac{b}{r}\right)$, $\gamma = \frac{\alpha^2}{\alpha^2+1}$ and b is an arbitrary constant. $d\Omega_{k,n-1}^2$ describes a $(n-1)$ -dimensional subspace with $n(n-1)k$ constant curvature and volume ω_{n-1} . k is a constant and characterizes the hypersurface. Without loss of generality, the value of k is $-1, 0, 1$, which correspond to a negative (hyperbolic), zero (flat), positive (elliptic) constant curvature hypersurface in the metric (5), respectively. m and q are integration constants. According to the definition of mass due to Abbott and Deser [63, 64], the mass takes the form

$$M = \frac{(n-1)b^{(n-1)\gamma}m\omega_{n-1}}{16\pi(\alpha^2+1)}. \quad (7)$$

The charge is

$$Q = \frac{q\omega_{n-1}}{4\pi}. \quad (8)$$

The electric potential is $A_t = -\frac{qb^{(3-n)\gamma}}{\lambda r^\lambda}$, where $\lambda = (n-3)(1-\gamma) + 1$. The thermodynamic properties in the extended phase space of the $(n+1)$ -dimensional dilaton black holes were deeply discussed in [65, 66, 67, 68, 69, 70].

In this paper, our interest is focused on the overcharging problem and thermodynamics in the extended phase spaces of the five-dimensional black holes. Therefore, we let $n = 4$ in the metric (5). Considering that the cosmological constant is a variable related to pressure, $P = -\frac{\Lambda}{8\pi} = \frac{n(n-1)}{16\pi l^2}$, we replace the cosmological constant with the pressure in the following discussions. In the absence of the non-trivial dilaton ($\alpha = 0$), the solution (6) is reduced to

$$f(r) = k - \frac{M}{r^2} \frac{16\pi}{3\omega_3} + \frac{Q^2}{r^4} \frac{16\pi^2}{3\omega_3^2} + \frac{4}{3}\pi r^2 P, \quad (9)$$

Thus the metric (5) describes a five-dimensional anti-de Sitter topological spacetime where $R(r) = 0$. Now the electric potential is given by

$$A_t = -\frac{2\pi Q}{\omega_3 r^2}. \quad (10)$$

The Hawking temperature and entropy are

$$T = \frac{f'(r_+)}{4\pi}, \quad S = \frac{\omega_3 r_+^3}{4}, \quad (11)$$

respectively. r_+ is the event horizon radius derived by $f(r) = 0$. The mass is expressed by the entropy, charge and pressure

$$M = \frac{3k\omega_3}{16\pi} \left(\frac{4S}{\omega_3}\right)^{\frac{2}{3}} + \frac{\pi Q^2}{\omega_3} \left(\frac{\omega_3}{4S}\right)^{\frac{2}{3}} + \frac{\omega_3 P}{4} \left(\frac{4S}{\omega_3}\right)^{\frac{4}{3}}. \quad (12)$$

Here, S , Q and P are regarded as the extensive parameters for the mass M . Their conjugate quantities are the temperature, electric potential and volume, respectively.

$$\begin{aligned} T &= \left(\frac{\partial M}{\partial S}\right)_{Q,P} = \frac{k\omega_3}{8\pi} \left(\frac{4}{\omega_3}\right)^{\frac{2}{3}} S^{-\frac{1}{3}} - \frac{2\pi Q^2}{3\omega_3} \left(\frac{\omega_3}{4}\right) S^{-\frac{5}{3}} + \frac{\omega_3 P}{3} \left(\frac{4}{\omega_3}\right)^{\frac{4}{3}} S^{\frac{1}{3}}, \\ \phi &= \left(\frac{\partial M}{\partial Q}\right)_{P,S} = \frac{2\pi Q}{\omega_3 r_+^2}, \quad V = \left(\frac{\partial M}{\partial P}\right)_{S,Q} = \frac{\omega_3 r_+^4}{4}. \end{aligned} \quad (13)$$

After a straightforward calculation, it is easily found that these thermodynamic quantities obey the first law of thermodynamics

$$dM = TdS + \phi dQ + VdP, \quad (14)$$

and the Smarr relation

$$2M = 3TS + 2\phi Q - 2VP. \quad (15)$$

In what follows the thermodynamics are investigated by the absorptions of the scalar particle and fermion and the first law of thermodynamics is recovered.

3 Particles' absorption

3.1 Scalar particle's absorption

In this subsection, we discuss the absorption of the scalar particle in the five-dimensional spacetime. In the curved spacetime, the motion of the charged scalar particle obeys the Klein-Gordon equation

$$\frac{1}{\sqrt{-g}} \left(\frac{\partial}{\partial x^\mu} - \frac{iq_e}{\hbar} A_\mu \right) \left[\sqrt{-g} g^{\mu\nu} \left(\frac{\partial}{\partial x^\nu} - \frac{iq_e}{\hbar} A_\nu \right) \right] \Psi_S - \frac{m_e^2}{\hbar^2} \Psi_S = 0, \quad (16)$$

where m_e and q_e are the mass and charge of the particle, respectively. Ψ_S is the scalar field. Following the ansatz [71, 72, 73], we use the *WKB* approximation and adopt the following expression of the wave function

$$\Psi_S = \exp\left(\frac{i}{\hbar}I_s + I_1 + \mathcal{O}(\hbar)\right). \quad (17)$$

Inserting the above ansatz and the contravariant metric components of the five-dimensional black hole into the Klein-Gordon equation yields an equation about the action I_s . Dividing by the exponential term and taking the leading contribution of \hbar , we get

$$f^{-1}(\partial_t I_s - q_e A_t)^2 - f(\partial_r I_s)^2 - g^{22}(\partial_{\theta_2} I_s)^2 - g^{33}(\partial_{\theta_3} I_s)^2 - g^{44}(\partial_{\theta_4} I_s)^2 + m_e^2 = 0. \quad (18)$$

Here we use $\theta_2, \theta_3, \theta_4$ to express the coordinates x^2, x^3, x^4 , respectively. And g^{22}, g^{33} and g^{44} are the corresponding contravariant metric components in the five-dimensional metric. Their expressions are not specified here, since they do not affect the later calculation result. This is different from the previous discussion of tunneling radiation with consideration of quantum gravity effects [74]. Taking into account the symmetries of the spacetime, we carry out the separation of variables to the action,

$$I_s = -\omega t + W(r) + \Theta(\theta_2, \theta_3, \theta_4). \quad (19)$$

In the above equation, ω is the energy of the absorbed scalar particle. Substituting the separated action into Eq. (18) and solving it yield

$$\partial_r W = \pm \frac{\sqrt{\left(\omega - q_e \frac{2\pi Q}{\omega_3 r^2}\right)^2 + [m_e^2 - g^{22}(\partial_{\theta_2} \Theta)^2 - g^{33}(\partial_{\theta_3} \Theta)^2 - g^{44}(\partial_{\theta_4} \Theta)^2] f}}{f}, \quad (20)$$

where $+$ ($-$) denote the cases of the ingoing (outgoing) particles. We assume that the particle is completely absorbed by the black hole. Then the negative sign in the above equation is neglected. Defining $p_r = \partial_r I_s$, we get

$$\begin{aligned} p^r &= g^{rr} p_r \\ &= \sqrt{\left(\omega - q_e \frac{2\pi Q}{\omega_3 r^2}\right)^2 + [m_e^2 - g^{22}(\partial_{\theta_2} \Theta)^2 - g^{33}(\partial_{\theta_3} \Theta)^2 - g^{44}(\partial_{\theta_4} \Theta)^2] f}, \end{aligned} \quad (21)$$

Near the event horizon, we have $f(r) \rightarrow 0$ and the above equation is simplified to

$$p^r = \omega - q_e \phi, \quad (22)$$

where $\phi = \frac{2\pi Q}{\omega_3 r_+^2}$ represents the electric potential at the event horizon. The condition of the superradiation is that the boundary condition of the scalar field should be in the asymptotic region and $\omega < q_e \phi$. In this paper, the effects of superradiation are not taken into account, which leads to $\omega > q_e \phi$ and $p^r > 0$. The relation (22) plays an important role in the derivation of thermodynamics and is recovered by the absorption of the fermion in the next subsection.

3.2 Fermion's absorption

In this subsection, we use a charged fermion with spin-1/2 to discuss its absorption at the event horizon. The motion of the fermion obeys the Dirac equation in curved spacetime

$$i\gamma^\mu \left(\partial_\mu + \Omega_\mu + \frac{i}{\hbar} q_e A_\mu \right) \Psi_F + \frac{m_e}{\hbar} \Psi_F = 0, \quad (23)$$

where m_e and q_e denote the mass and charge of the fermion, respectively. Ψ_F is the wave function. $\Omega_\mu \equiv \frac{i}{2} \omega_\mu{}^{ab} \Sigma_{ab}$, $\omega_\mu{}^{ab}$ is the spin connection defined by the ordinary connection and the tetrad $e^\lambda{}_b$, namely, $\omega_\mu{}^a{}_b = e_\nu{}^a e^\lambda{}_b \Gamma_{\mu\lambda}^\nu - e^\lambda{}_b \partial_\mu e_\lambda{}^a$. The Greek indices are raised and lowered by the curved metric $g_{\mu\nu}$. The Latin indices are governed by the flat metric η_{ab} . The construction of the tetrad needs to use the following definitions

$$g_{\mu\nu} = e_\mu{}^a e_\nu{}^b \eta_{ab}, \quad \eta_{ab} = g_{\mu\nu} e^\mu{}_a e^\nu{}_b, \quad e^\mu{}_a e_\nu{}^a = \delta_\nu^\mu, \quad e^\mu{}_a e_\mu{}^b = \delta_a^b. \quad (24)$$

The Lorentz spinor generators are defined by $\Sigma_{ab} = \frac{i}{4} [\gamma^a, \gamma^b]$ and γ^a satisfy $\{\gamma^a, \gamma^b\} = 2\eta^{ab}$. We can readily construct the γ^μ 's in curved spacetime as

$$\gamma^\mu = e^\mu{}_a \gamma^a, \quad \{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}. \quad (25)$$

For a spin-1/2 fermion, there are two states corresponding to spin up and spin down. First of all, we discuss the spin up state. The wave function takes the form

$$\Psi_{F\uparrow} = \begin{pmatrix} A \\ 0 \\ B \\ 0 \end{pmatrix} \exp\left(\frac{i}{\hbar} I_\uparrow\right), \quad (26)$$

where A , B and I_\uparrow are the functions of $(t, r, \theta_2, \theta_3, \theta_4)$. To solve the wave function, we should construct the γ^μ matrices. There are many choices to construct them. For the five-dimensional metric, we first choose the tetrad

$$e_\mu{}^a = \text{diag}\left(\sqrt{f}, 1/\sqrt{f}, \sqrt{g^{22}}, \sqrt{g^{33}}, \sqrt{g^{44}}\right). \quad (27)$$

Then the γ^μ matrices are

$$\begin{aligned} \gamma^t &= \frac{1}{\sqrt{f(r)}} \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}, & \gamma^{\theta_2} &= \sqrt{g^{22}} \begin{pmatrix} 0 & \sigma^1 \\ \sigma^1 & 0 \end{pmatrix}, \\ \gamma^r &= \sqrt{f(r)} \begin{pmatrix} 0 & \sigma^3 \\ \sigma^3 & 0 \end{pmatrix}, & \gamma^{\theta_3} &= \sqrt{g^{33}} \begin{pmatrix} 0 & \sigma^2 \\ \sigma^2 & 0 \end{pmatrix}, \\ \gamma^{\theta_4} &= \sqrt{g^{44}} \begin{pmatrix} -I & 0 \\ 0 & I \end{pmatrix}. \end{aligned} \quad (28)$$

σ^1 , σ^2 and σ^3 are the Pauli matrices and I is a identity matrix. Inserting the spin connection, wave function and gamma matrices into the Dirac equation, dividing the exponential term and multiplying by \hbar yield four equations to the leading order in \hbar . They are

$$-B\frac{1}{\sqrt{f}}(\partial_t I_\uparrow - q_e A_t) - B\sqrt{f}\partial_r I_\uparrow + A\sqrt{g^{44}}\partial_{\theta_4} I_\uparrow + Am_e = 0, \quad (29)$$

$$A\frac{1}{\sqrt{f}}(\partial_t I_\uparrow - q_e A_t) - A\sqrt{f}\partial_r I_\uparrow - B\sqrt{g^{44}}\partial_{\theta_4} I_\uparrow + Bm_e = 0, \quad (30)$$

$$-B\left[\sqrt{g^{22}}\partial_{\theta_2} I_\uparrow + i\sqrt{g^{33}}\partial_{\theta_3} I_\uparrow\right] = 0, \quad (31)$$

$$-A\left[\sqrt{g^{22}}\partial_{\theta_2} I_\uparrow + i\sqrt{g^{33}}\partial_{\theta_3} I_\uparrow\right] = 0. \quad (32)$$

The last two equations are simplified to an equation and yield $g^{22}(\partial_{\theta_2} I_\uparrow)^2 + g^{33}(\partial_{\theta_3} I_\uparrow)^2 = 0$. Previous discussions have shown that the contribution of the angular part does not affect the results of the tunneling radiation when quantum gravity effects are not taken into account [74]. Therefore, we focus on the first two equations. Due to the same symmetry in the black hole spacetime, we still adopt the separation of variables in the above subsection

$$I_\uparrow = -\omega t + W(r) + \Theta(\theta_2, \theta_3, \theta_4). \quad (33)$$

Now ω denotes the energy of the absorbed fermion. Combining these two equations and eliminating A and B , we get an equation about the action I_\uparrow . Substituting the separated variables into this equation and solving it, we get

$$\partial_r W = \pm \frac{\sqrt{\left(\omega - q_e \frac{2\pi Q}{\omega_3 r^2}\right)^2 + [m_e^2 - g^{44}\partial_{\theta_4} I_\uparrow] f}}{f}, \quad (34)$$

where $+$ ($-$) denote the cases of the ingoing (outgoing) fermion. As addressed in the above subsection, the particle is assumed to be completely absorbed. Therefore, there is no outgoing fermion. Using the definition $p_r = \partial_r I_\uparrow$, we get $p^r = g^{rr}\partial_r I_\uparrow$. Near the event horizon,

$$p^r = \omega - q_e \phi, \quad (35)$$

where ϕ is the electric potential at the event horizon given by Eq. (13). It recovers the relation gotten by the absorption of the scalar particle. For the spin down, the calculation is parrel and the same relation (35) is gotten.

4 Thermodynamics and particles' absorption

Thermodynamics of black holes were deeply discussed through particles' tunnelling across event horizons. In this section, it is shown that the thermodynamics in the extended phase space of the five-dimensional black hole can be recovered by the absorptions of the scalar particle and fermion. The relation $p^r = \omega - q_e \phi$ is used in the derivation. Therefore, the particle can be either a scalar particle or a fermion. When a particle with energy and charge falls into the black hole, the energy and charge of the black hole increase. Let $f(M, Q, P, r_+) = f(r_+)$ represent the metric component of the original state of the black hole and $f(M + dM, Q + dQ, P + dP, r_+ + dr_+)$ be the metric component of the final state after the particle being absorbed. These metric components satisfy

$$\begin{aligned} f(M + dM, Q + dQ, P + dP, r_+ + dr_+) - f(M, Q, P, r_+) = \\ \left. \frac{\partial f(M, Q, P, r)}{\partial M} \right|_{r=r_+} dM + \left. \frac{\partial f(M, Q, P, r)}{\partial Q} \right|_{r=r_+} dQ + \left. \frac{\partial f(M, Q, P, r)}{\partial P} \right|_{r=r_+} dP \\ + \left. \frac{\partial f(M, Q, P, r)}{\partial r} \right|_{r=r_+} dr_+, \end{aligned} \quad (36)$$

where

$$\begin{aligned} \left. \frac{\partial f(M, Q, P, r)}{\partial M} \right|_{r=r_+} &= -\frac{16\pi}{3\omega_3 r_+^2}, & \left. \frac{\partial f(M, Q, P, r)}{\partial Q} \right|_{r=r_+} &= \frac{32\pi^2 Q}{3\omega_3^2 r_+^4}, \\ \left. \frac{\partial f(M, Q, P, r)}{\partial P} \right|_{r=r_+} &= \frac{4\pi r_+^2}{3}, & \left. \frac{\partial f(M, Q, P, r)}{\partial r} \right|_{r=r_+} &= 4\pi T. \end{aligned} \quad (37)$$

Following Gwak's work, we assume that the final state is still a black hole. This implies $f(M + dM, Q + dQ, P + dP, r_+ + dr_+) = f(M, Q, P, r_+) = 0$. From Eqs. (36) and (37), the increase of the horizon radius takes the form

$$dr_+ = \frac{4p^r}{3\omega_3 r_+^2 T - 4\omega_3 r_+^3 P}. \quad (38)$$

Using the entropy relation $S = \frac{1}{4}\omega_3 r_+^3$ yields

$$dS = \frac{3p^r}{3T - 4r_+ P}, \quad (39)$$

which shows that the increase of the entropy is related to the Hawking temperature, pressure and p^r . For the non-extremal black hole or the fixed cosmological constant, we readily get $dS > 0$, which implies that the entropy increase when the particle is absorbed by the black hole. In the extremal and near-extremal cases, $dS < 0$. This implies that the entropy decreases when the particle falls into the black hole. It

violates the second law of thermodynamics which shows the entropy never decreases in the clockwise direction.

We associate the particle's energy and charge with the increases of the internal energy and charge of the black hole,

$$\omega = dU, \quad q_e = dQ. \quad (40)$$

Since the cosmological constant is regarded as the thermodynamic pressure, the black hole mass is seen as the enthalpy. The relationship between the mass and internal energy is

$$U = M - PV. \quad (41)$$

Combining Eqs. (39), (40) and (41) yields

$$dM = TdS + \phi dQ + VdP, \quad (42)$$

which is the first law of thermodynamics recovered by the absorptions of the scalar particle and fermion. This derivation is inevitable, since the final state of the black hole after the particle being absorbed was assumed to be still a black hole.

5 Overcharging problem in the extremal and near-extremal black holes

In this section, we test the WCCC in the extremal and near-extremal five-dimensional black holes by the absorptions of the scalar particle and fermion. Here the final states of the black holes are unknown. An effective way to test the WCCC is to check whether a event horizon exists when a particle with energy and large enough charge or angular momentum falls into a black hole. The event horizon is determined by the metric component $f(r)$. If the minimal value of $f(r)$ is negative or zero, the horizon exists. Otherwise, the horizon does not exist.

We use $f(M, Q, P, r_0)$ to represent the metric component of the original state and $f(M + dM, Q + dQ, P + dP, r_0 + dr_0)$ to denote the metric component of the final state, where r_0 is the location of the minimal value. The relation between these two metric components is

$$f(M + dM, Q + dQ, P + dP, r_0 + dr_0) = f(M, Q, P, r_0) + \left. \frac{\partial f(M, Q, P, r)}{\partial r} \right|_{r=r_0} dr_0 + \left. \frac{\partial f(M, Q, P, r)}{\partial M} \right|_{r=r_0} dM + \left. \frac{\partial f(M, Q, P, r)}{\partial Q} \right|_{r=r_0} dQ + \left. \frac{\partial f(M, Q, P, r)}{\partial P} \right|_{r=r_0} dP, \quad (43)$$

where

$$\begin{aligned}
f(M, Q, P, r_0) &= k - \frac{M}{r_0^2} \frac{16\pi}{3\omega_3} + \frac{Q^2}{r_0^4} \frac{16\pi^2}{3\omega_3^2} + \frac{4}{3}\pi r_0^2 P, \\
\left. \frac{\partial f(M, Q, P, r)}{\partial r} \right|_{r=r_0} &= 0, \quad \left. \frac{\partial f(M, Q, P, r)}{\partial P} \right|_{r=r_0} = \frac{4\pi r_0^2}{3}, \\
\left. \frac{\partial f(M, Q, P, r)}{\partial M} \right|_{r=r_0} &= -\frac{16\pi}{3\omega_3 r_0^2}, \quad \left. \frac{\partial f(M, Q, P, r)}{\partial Q} \right|_{r=r_0} = \frac{32\pi^2 Q}{3\omega_3^2 r_0^4}, \quad (44)
\end{aligned}$$

For the extremal black hole, its Hawking temperature is zero. The location of the event horizon coincides with that of the minimal value, namely, $r_+ = r_0$. Thus,

$$\begin{aligned}
&f(M + dM, Q + dQ, P + dP, r_0 + dr_0) \\
&= f(M, Q, P, r_+) + \frac{16\pi}{3\omega_3 r_+^2} [dM - V dP - \phi dQ] \\
&= 0, \quad (45)
\end{aligned}$$

where $f(M, Q, P, r_+) = 0$ and the first law of thermodynamics for the extremal black hole were used to derive the second equal sign. It shows that the final state of the extremal black hole is still a extremal one with new mass and charge. Therefore, the extremal black hole can not be overcharged.

In the near-extremal case, $f(M, P, Q, r_0) < 0$. We order $r_0 = r_+ - \epsilon$, where $0 < \epsilon \ll 1$. Eq. (43) is reduced to

$$\begin{aligned}
f(M + dM, Q + dQ, P + dP, r_0 + dr_0) &= f(M, Q, P, r_0) + \mathcal{O}(\epsilon) \\
&- \frac{16\pi}{3\omega_3 r_+^2} \left(1 + \frac{4\epsilon}{r_+}\right) \frac{3T p^r}{3T - 4r_+ P} + \frac{2\epsilon}{r_+} \left(\frac{16\pi}{3\omega_3 r_+^2} dM - 4\pi r_+^2 dP \right). \quad (46)
\end{aligned}$$

The near-extremity leads to $T \rightarrow 0$. Considering $r_+ \sim l \gg 1 \gg \epsilon$, we neglect the second term of the second line in the above equation. Thus we get $f(M + dM, Q + dQ, P + dP, r_0 + dr_0) < 0$. It shows that the final state of the near-extremal black hole is still a near-extremal one with the new mass and charge.

Therefore, the extremal and near-extremal five-dimensional black holes can not be overcharged by absorbing the scalar particle or fermion with mass and large enough charge.

6 Discussion and Conclusion

In this paper, we investigated the overcharging problem and thermodynamics in the extended phase spaces of the five-dimensional spherically symmetric topological black holes by the absorptions of the scalar particle and fermion. In the investigation, the cosmological constant was regarded as a variable related to pressure and its conjugate quantity is the thermodynamic volume. The calculations of the absorptions of the

scalar particle and fermion reduced to the same relation $p^r = \omega - q_e \phi$. This relation is full in consistence with that gotten by the Hamilton-Jacobi equation [46]. Maybe the reason is that the Hamilton-Jacobi equation can be derived by the Klein-Gordon equation and Dirac equation. This derivation is referred to Ref. [75]. We used this relation to discuss the thermodynamics and recovered the first law of thermodynamics. This result is inevitable because the final state of the black hole after the particles being absorbed was assumed to be still a black hole. But the second law of thermodynamics was violated in the near-extremal and extremal black holes. The event horizon is determined by the metric component $f(r)$. To check the existence of the horizons at the final stages of the extremal and near-extremal black holes, we estimate the minimal values of the metric component. For the extremal black hole, its minimal value is zero, which shows that the final state is still a extremal black hole. For the near-extremal black hole, its minimal value is a negative value, which corresponds to a near-extremal black hole with new mass and charge. Therefore, the extremal and near-extremal black holes can not be overcharged.

7 Acknowledgments

This work is supported by the NSFC (Grant No. 11205125).

References

- [1] R. Penrose, *Gravitational collapse: the role of general relativity*, *Riv. Nuovo Cim.* **1** (1969) 252.
- [2] R.M. Wald, *Gedanken experiments to destroy a black hole*, *Ann. Phys.* **82** (1974) 548.
- [3] I. Semiz, *Dyonic Kerr-Newman black holes, complex scalar field and cosmic censorship*, *Gen. Relat. Grav.* **43** (2011) 833.
- [4] G.Z. Toth, *Test of the weak cosmic censorship conjecture with a charged scalar field and dyonic Kerr-Newman black holes*, *Gen. Relat. Grav.* **44** (2012) 2019.
- [5] J. Crisostomo and R. Olea, *Hamiltonian treatment of the gravitational collapse of thin shells*, *Phys. Rev. D* **69** (2004) 104023.
- [6] G.E.A. Matsas and A.R.R. da Silva, *Overspinning a nearly extreme charged black hole via a quantum tunneling process*, *Phys. Rev. Lett.* **99** (2007) 181301.
- [7] S. Hod, *Weak cosmic censorship: as strong as ever*, *Phys. Rev. Lett.* **100** (2008) 121101.
- [8] M. Richartz and A. Saa, *Overspinning a nearly extreme black hole and the weak cosmic censorship conjecture*, *Phys. Rev. D* **78** (2008) 081503(R).
- [9] T. Jacobson and T.P. Sotiriou, *Over-spinning a black hole with a test body*, *Phys. Rev. Lett.* **103** (2009) 141101.
- [10] A. Saa and R. Santarelli, *Destroying a near-extremal Kerr-Newman black hole*, *Phys. Rev. D* **84** (2011) 027501.

- [11] S. Isoyama, N. Sago and T. Tanaka, *Cosmic censorship in overcharging a Reissner-Nordstrom black hole via charged particle absorption*, *Phys. Rev. D* **84** (2011) 124024.
- [12] S. Hod, *Cosmic censorship: formation of a shielding horizon around a fragile horizon*, *Phys. Rev. D* **87** (2013) 024037.
- [13] S. Gao and Y. Zhang, *Destroying extremal Kerr-Newman black holes with test particles*, *Phys. Rev. D* **87** (2013) 044028.
- [14] K. Düztaş and I. Semiz, *Cosmic censorship, black holes and integer-spin test fields*, *Phys. Rev. D* **88** (2013) 064043.
- [15] H.M. Siahhaan, *Destroying Kerr-Sen black holes*, *Phys. Rev. D* **93** (2016) 064028.
- [16] K. Düztaş, *Overspinning BTZ black holes with test particles and fields*, *Phys. Rev. D* **94** (2016) 124031.
- [17] K. Düztaş, *Can test fields destroy the event horizon in the Kerr-Taub-NUT spacetime?* *Class. Quant. Gravi.* **35** (2018) 045008.
- [18] V. Husain and S. Singh, *On the Penrose inequality in anti-de Sitter space*, *Phys. Rev. D* **96** (2017) 104055.
- [19] K.S. Revelar and I. Vega, *Overcharging higher-dimensional black holes with point particles*, *Phys. Rev. D* **96** (2017) 064010.
- [20] C. Fairoos, A. Ghosh and S. Sarkar, *Massless charged particles: cosmic censorship, and third law of black hole mechanics*, *Phys. Rev. D* **96** (2017) 084013.
- [21] B. Gwak, *Weak cosmic censorship conjecture in Kerr-(Anti-)de Sitter black hole with scalar field*, *JHEP* **1809** (2018) 081.
- [22] Y. Gim and B. Gwak, *Charged black hole in gravity's rainbow: violation of weak cosmic censorship*, *Phys. Lett. B* **794** (2019) 122.
- [23] B. Gwak, *Thermodynamics and cosmic censorship conjecture in Kerr-Newman-de Sitter black hole*, *Entropy* **20** (2018) 855.
- [24] T. Crisford, G.T. Horowitz and J.E. Santos, *Testing the weak gravitycosmic censorship connection*, *Phys. Rev. D* **97** (2018) 066005.
- [25] T.Y. Yu and W.Y. Wen, *Cosmic censorship and weak gravity conjecture in the Einstein-Maxwell-dilaton theory*, *Phys. Lett. B* **781** (2018) 713.
- [26] B. Liang, S.W. Wei and Y.X. Liu, *Weak cosmic censorship conjecture in Kerr black holes of modified gravity*, *Mod. Phys. Lett. A* **34** (2019) 1950037.
- [27] J. Natario, L. Queimada and R. Vicente *Test fields cannot destroy extremal black holes*, *Class. Quant. Grav.* **33** (2016) 175002.
- [28] J. Natario and R. Vicente *Test fields cannot destroy extremal de Sitter black holes*, *Gen. Rel. Grav.* **52** (2020) 5.
- [29] M. Colleoni and L. Barack, *Overspinning a Kerr black hole: The effect of the self-force*, *Phys. Rev. D* **91** (2015) 104024.

- [30] M. Colleoni, L. Barack, Abhay G. Shah, and M. van de Meent, *Self-force as a cosmic censor in the Kerr overspinning problem*, *Phys. Rev. D* **92** (2015) 084044.
- [31] E. Barausse, V. Cardoso and G. Khanna, *Test bodies and naked singularities: Is the self-force the cosmic censor?* *Phys. Rev. Lett.* **105** (2010) 261102.
- [32] P. Zimmerman, I. Vega, E. Poisson and R. Haas, *Selfforce as a cosmic censor*, *Phys. Rev. D* **87** (2013) 041501.
- [33] T. Moon and Y.S. Myung, *Gregory-Laflamme instability of BTZ black hole in new massive gravity*, *Phys. Rev. D* **88** (2013) 124014.
- [34] D. Kastor, S. Ray and J. Traschen, *Enthalpy and the mechanics of AdS black holes*, *Class. Quantum Gravity* **26** (2009) 195011.
- [35] D. Kastor, S. Ray and J. Traschen, *Smarr formula and an extended first law for lovelock gravity*, *Class. Quantum Gravity* **27** (2010) 235014.
- [36] H. Lu, Y. Pang, C.N. Pope and J.F. Vazquez-Poritz, *AdS and Lifshitz black holes in conformal and Einstein-Weyl gravities*, *Phys. Rev. D* **86** (2012) 044011.
- [37] D. Kubiznak and R.B. Mann, *P-V criticality of charged AdS black holes*, *JHEP* **1207** (2012) 033.
- [38] J. Creighton and R.B. Mann, *Quasilocal thermodynamics of dilaton gravity coupled to gauge fields*, *Phys. Rev. D* **52** (1995) 4569.
- [39] A.M. Frassino, D. Kubiznak, R.B. Mann and F. Simovic, *Multiple reentrant phase transitions and triple points in Lovelock thermodynamics*, *JHEP* **1409** (2014) 080.
- [40] R.A. Hennigar, W.G. Brenna and R.B. Mann, *P-V criticality in quasitopological gravity*, *JHEP* **1507** (2015) 077.
- [41] R.G. Cai, L.M. Cao, L. Li and R.Q. Yang, *P-V criticality in the extended phase space of GB black holes in AdS space*, *JHEP* **1309** (2013) 005.
- [42] S.W. Wei, Y.X. Liu and R.B. Mann, *Repulsive interactions and universal properties of charged anti-de Sitter black hole microstructures*, *Phys. Rev. Lett.* **123** (2019) 071103.
- [43] S.W. Wei and Y.X. Liu, *Insight into the microscopic structure of an AdS black hole from thermodynamical phase transition*, *Phys. Rev. Lett.* **115** (2015) 111302, Erratum: *Phys. Rev. Lett.* **116** (2016) 169903.
- [44] D.C. Zou, S.J. Zhang and B. Wang, *Critical behavior of Born-Infeld AdS black holes in the extended phase space*, *Phys. Rev. D* **89** (2013) 044002.
- [45] C. Niu, Y. Tian and X.N. Wu, *Critical phenomena and thermodynamic geometry of RN-AdS black holes*, *Phys. Rev. D* **85** (2012) 024017.
- [46] B. Gwak, *Thermodynamics with pressure and volume under charged particle absorption*, *JHEP* **1711** (2017) 129.
- [47] J. Sorce and R.M. Wald, *Gedanken experiments to destroy a black hole II: Kerr-Newman black holes cannot be over-charged or over-spun*, *Phys. Rev. D* **96** (2017) 104014.

- [48] R.M. Wald, *Kerr-Newman black holes cannot be over-charged or over-spun*, *Int. J. Mod. Phys. D* **27** (2018) 1843003.
- [49] J.C. An, J.R. Shan, H.B. Zhang and S.T. Zhao, *Five dimensional Myers-Perry black holes cannot be over-spun by gedanken experiments*, *Phys. Rev. D* **97** (2018) 104007.
- [50] B.Y. Chen, F.L. Lin and B. Ning, *Gedanken experiments to destroy a BTZ black hole*, *Phys. Rev. D* **100** (2019) 044043.
- [51] A.K. Mishra and S. Sarkar, *Overcharging a multi black hole system and cosmic censorship*, *Phys. Rev. D* **100** (2019) 024030.
- [52] J. Jiang, B.L. Deng, Z.Z. Chen, *Static charged dilaton black cannot be over-charged by gedanken experiments*, *Phys. Rev. D* **100** (2019) 066024.
- [53] B. Gwak, *Weak cosmic censorship with pressure and volume in charged anti-de Sitter black hole under charged scalar field*, *JCAP* **1908** (2019) 016.
- [54] X.X. Zeng, X.Y. Hu and K.J. He, *Weak cosmic censorship conjecture with pressure and volume in the Gauss-Bonnet AdS black hole*, *Nucl. Phys. B* **949** (2019) 114823
- [55] W. Hong, B.R. Mu and J. Tao, *Thermodynamics and weak cosmic censorship conjecture in the charged RN-AdS black hole surrounded by quintessence under the scalar field*, *Nucl. Phys. B* **949** (2019) 114826.
- [56] P. Wang, H.W. Wu and H. Yang, *Thermodynamics of nonlinear electrodynamics black holes and the validity of weak cosmic censorship at charged particle absorption*, *Eur. Phys. J. C* **79** (2019) 572.
- [57] P. Wang, H.W. Wu and S.X. Ying, *Validity of thermodynamic laws and weak cosmic censorship for AdS black holes and black holes in a cavity*, [arXiv:2002.12233[gr-qc]]
- [58] B.R. Mu and J. Tao, *Minimal length effect on thermodynamics and weak cosmic censorship conjecture in anti-de Sitter black holes via charged particle absorption*, *Advances in High Energy Physics*, **2020** (2020) 2612946.
- [59] Y.W. Han, X.X. Zeng and Y. Hong, *Thermodynamics and weak cosmic censorship conjecture of the torus-like black hole*, *Eur. Phys. J. C* **79** (2019) 252.
- [60] X.X. Zeng and H.Q. Zhang, *Thermodynamics and weak cosmic censorship conjecture in Born-Infeld-anti-de Sitter black holes*, [arXiv:1901.04247 [hep-th]]
- [61] A. Sheykhi, *Thermodynamics of charged topological dilaton black holes*, *Phys. Rev. D* **76** (2007) 124025.
- [62] K.C.K. Chan, J.H. Horne and R.B. Mann, *Charged dilaton black holes with unusual asymptotics*, *Nucl. Phys. B* **447** (1995) 441.
- [63] L.F. Abbott and S. Deser, *Stability of gravity with a cosmological constant*, *Nucl. Phys. B* **195** (1982) 76.
- [64] G. Kofinas, R. Olea, *Vacuum energy in Einstein-Gauss-Bonnet AdS gravity*, *Phys. Rev. D* **74** (2006) 084035

- [65] R. Zhao, H.H. Zhao, M.S. Ma and L.C. Zhang, *On the critical phenomena and thermodynamics of charged topological dilaton AdS black holes*, *Eur. Phys. J. C* **73** (2013) 2645.
- [66] M.H. Dehghani, S. Kamrani and A. Sheykhi, *P-V criticality of charged dilatonic black holes*, *Phys. Rev. D* **90** (2014) 104020.
- [67] S.H. Hendi, A. Sheykhi, S. Panahiyan and B. Eslam Panah, *Phase transition and thermodynamic geometry of Einstein-Maxwell-dilaton black holes*, *Phys. Rev. D* **92** (2015) 064028.
- [68] A. Dehyadegari, A. Sheykhi and A. Montakhab, *Novel phase transition in charged dilaton black holes*, *Phys. Rev. D* **96** (2017) 084012.
- [69] S.H. Hendi, M. Faizal, B. Eslam Panah and S. Panahiyan, *Charged dilatonic black holes in gravity's rainbow*, *Eur. Phys. J. C* **76** (2016) 296.
- [70] S.H. Hendi, B. Eslam Panah, S. Panahiyan and M. Momennia, *Dilatonic black holes in gravity's rainbow with a nonlinear source: the effects of thermal fluctuations*, *Eur. Phys. J. C* **77** (2017) 647.
- [71] U.A. Gillani, M. Rehman and K. Saifullah, *Hawking radiation of scalar particles from accelerating and rotating black holes*, *JCAP* **1106** (2011) 016.
- [72] A. Ejaz, H. Gohar, H. Lin, K. Saifullah and S.T. Yau, *Quantum tunneling from three-dimensional black holes*, *Phys. Lett. B* **726** (2013) 827.
- [73] I. Sakalli and H. Gursel, *Quantum tunneling from rotating black holes with scalar hair in three dimensions*, *Eur. Phys. J. C* **76** (2016) 318.
- [74] D.Y. Chen, H. Yang and X.T. Zu, *Hawking radiation of black holes in the $z = 4$ Horava-Lifshitz gravity*, *Phys. Lett. B* **681** (2009) 463.
- [75] H. Erbin and V. Lahoche, *Universality of tunnelling particles in Hawking radiation*, *Phys. Rev. D* **98** (2018) 104001.