

Palatini quadratic gravity: spontaneous breaking of gauged scale symmetry and inflation

D. M. Ghilencea *

Department of Theoretical Physics, National Institute of Physics
and Nuclear Engineering, Bucharest 077125, Romania

Abstract

We study quadratic gravity $R^2 + R^2_{[\mu\nu]}$ in the Palatini formalism where the connection and the metric are independent. This action has a *gauged* scale symmetry (also known as Weyl gauge symmetry) of Weyl gauge field $v_\mu = (\tilde{\Gamma}_\mu - \Gamma_\mu)/2$, with $\tilde{\Gamma}_\mu$ (Γ_μ) the trace of the Palatini (Levi-Civita) connection, respectively. The underlying geometry is non-metric due to the $R^2_{[\mu\nu]}$ term acting as a gauge kinetic term for v_μ . We show that this theory has an elegant spontaneous breaking of gauged scale symmetry and mass generation in the absence of matter, where the necessary scalar field (ϕ) is not added ad-hoc to this purpose but is “extracted” from the R^2 term. The gauge field becomes massive by absorbing the derivative term $\partial_\mu \ln \phi$ of the Stueckelberg field (“dilaton”). In the broken phase one finds the Einstein-Proca action of v_μ of mass proportional to the Planck scale $M \sim \langle \phi \rangle$, and a positive cosmological constant. Below this scale v_μ decouples, the connection becomes Levi-Civita and metricity and Einstein gravity are recovered. These results remain valid in the presence of non-minimally coupled scalar field (Higgs-like) with Palatini connection and the potential is computed. In this case the theory gives successful inflation and a specific prediction for the tensor-to-scalar ratio $0.007 \leq r \leq 0.01$ for current spectral index n_s (at 95%CL) and $N = 60$ efolds. This value of r is mildly larger than in inflation in Weyl quadratic gravity of similar symmetry, due to different non-metricity. This establishes a connection between non-metricity and inflation predictions and enables us to test such theories by future CMB experiments.

*E-mail: dimitru.ghilencea@cern.ch

1 Introduction

At a fundamental level gravity may be regarded as a theory of connections. An example is the “Palatini approach” to gravity due to Einstein [1,2], hereafter called EP approach [3,4]. In this case the “Palatini connection” ($\tilde{\Gamma}$) is apriori independent of the metric ($g_{\alpha\beta}$) and is actually determined by its equations of motion, from the action considered. For simple actions, $\tilde{\Gamma}$ plays an auxiliary role only, with no dynamics. For example, for an Einstein action in the EP approach the variation principle gives that $\tilde{\Gamma}$ is actually equal to the Levi-Civita connection (Γ). With this solution for $\tilde{\Gamma}$, one then recovers Einstein gravity - the metric formulation and EP approach are equivalent.

However, this equivalence is not true in general, for complicated actions, with matter present, etc [5–20]. For example, for quadratic gravity actions of the type studied here in the EP approach, the equations of motion of $\tilde{\Gamma}$ become complicated *second-order* differential equations; further, some components of $\tilde{\Gamma}$ even become dynamical in a sense discussed shortly, etc. The question remains, however, if such general actions in the EP formalism and *in the absence of matter* can recover *dynamically* the Levi-Civita connection and Einstein gravity. If true, this would be similar to the original Weyl quadratic gravity theory [21–24] as we showed recently in [25,26]. The main goal of this paper is to answer this question.

To address this question we study a gravity action in the EP approach with *gauged* scale symmetry also called Weyl gauge symmetry, see [25,26] for an example¹. This symmetry, first present in Weyl gravity [21–23] is important for mass generation, hence our interest. This symmetry demands us to consider quadratic gravity actions, with no dimensionful parameters. For such action we shall: **1)** explain the spontaneous breaking of this symmetry and the emergence of Levi-Civita connection, Einstein gravity and Planck scale in the broken phase, even in the absence of matter. This answers the above question; **2)** study the relation of this action to Weyl theory [21–23] of similar symmetry; **3)** study its inflation predictions.

In Section 2 we first review the $R(\tilde{\Gamma}, g)^2$ gravity in the EP approach, where $R(\tilde{\Gamma}, g)$ denotes the scalar curvature in this formalism. This action is local scale invariant. The connection is shown to be conformally related to the Levi-Civita connection. When “fixing the gauge” of this symmetry, the “auxiliary” scalar field ϕ introduced to “linearise” the R^2 term decouples. As a result, one finds that $\tilde{\Gamma} = \Gamma$ and Einstein action is obtained.

In Sections 3 and 4 we study the quadratic action $R(\tilde{\Gamma}, g)^2 + R_{[\mu\nu]}(\tilde{\Gamma})^2$ in the EP approach, hereafter called “EP quadratic gravity”. Here we used the notation $R_{[\mu\nu]} \equiv (R_{\mu\nu} - R_{\nu\mu})/2$. In this action the trace $\tilde{\Gamma}_\mu$ of the Palatini connection (assumed symmetric) is *dynamical* in the sense that $R_{[\mu\nu]}(\tilde{\Gamma})^2$ is a gauge kinetic term for $\tilde{\Gamma}_\mu$ or, more exactly, for the vector field² $v_\mu \sim \tilde{\Gamma}_\mu - \Gamma_\mu$, ($\tilde{\Gamma}_\mu \equiv \tilde{\Gamma}_{\mu\alpha}^\alpha$, $\Gamma_\mu \equiv \Gamma_{\mu\alpha}^\alpha$). With $\tilde{\Gamma}$ independent of $g_{\mu\nu}$, one notices that the local scale symmetry of this action is actually a *gauged* scale symmetry, of gauge field v_μ .

A consequence of the gauged scale symmetry is that EP quadratic gravity is non-metric³ i.e. $\tilde{\nabla}_\mu g_{\alpha\beta} \neq 0$. This is due to a dynamical $v_\mu \sim \tilde{\Gamma}_\mu$ and v_μ is the non-metricity field, also

¹See e.g. [21–46] for models with gauged scale symmetry and [47–68] for conformal or global scale symmetry.

²Unlike $\tilde{\Gamma}_\mu$ and Γ_μ , $v_\mu \propto \tilde{\Gamma}_\mu - \Gamma_\mu$ is indeed a vector (see Appendix). We assume $\tilde{\Gamma}_{\mu\nu}^\alpha = \tilde{\Gamma}_{\nu\mu}^\alpha$ (no torsion).

³Non-metricity means that under parallel transport along a curve a vector changes its norm i.e. is path dependent; therefore, in realistic theories with matter present (as here) it must be suppressed by a large scale (e.g. Planck) to avoid atomic spectral lines changes that it would otherwise induce [21]. In the absence of matter non-metricity can be traded for torsion in related R^2 theory [69].

called Weyl gauge field. Further, we find that the equations of motion for $\tilde{\Gamma}$ are *second-order* differential equations. In this case the usual EP approach in $f(R)$ theories to solve algebraically for $\tilde{\Gamma}$ [4] does not work, due to local scale symmetry and non-metricity. Nevertheless, we compute $\tilde{\Gamma}$ and find that EP quadratic gravity with $\tilde{\Gamma}$ *onshell* is equivalent to a ghost-free second-order gauged scale invariant theory with an *additional* dynamical field (“dilaton”). (Expressed in terms of this field the differential equations of $\tilde{\Gamma}$ simplify considerably and this is how they are solved).

The main result of this work (Section 3) is that the gauged scale invariance of the above action is broken spontaneously by a new mechanism [25, 26] valid even in the absence of matter; in this, the necessary scalar field (ϕ) is *not* added ad-hoc to this purpose (as usually done), but is “extracted” from the R^2 term in the action; ϕ is thus of geometric origin. After a Stueckelberg mechanism [70–72] the gauge field v_μ becomes massive, of mass m_v near the Planck scale $M \sim \langle \phi \rangle$, by “absorbing” the derivative $\partial_\mu \ln \phi$ of the Stueckelberg field (also referred to as “dilaton”). Near the Planck scale we obtain the Einstein-Proca action of v_μ . Further, below the scale $m_v \propto M$, the field v_μ decouples and we recover metricity, Levi-Civita connection and Einstein gravity; the Planck scale M is then an *emergent* scale where this symmetry is broken. These results remain true if the theory also has matter fields (higgs, etc) non-minimally coupled with *Palatini connection*, while respecting gauged scale invariance (Section 4). Briefly, the EP quadratic gravity is a gauged scale invariant theory broken à la Stueckelberg, even in the absence of matter, to an Einstein-Proca action with a positive cosmological constant and a potential for the scalar fields - if present. This answers the main goal of the paper.

Another theory where the connection is not determined by the metric itself is the original Weyl quadratic gravity of gauged scale invariance [21–23] (also [24]). With hindsight, it is then not too surprising that the above results are similar to those in [25–27] for Weyl theory. This theory came under early criticism from Einstein [21] for its non-metricity implying e.g. changes of the atomic spectral lines, in contrast to experiment; however, if the Weyl “photon” (v_μ) of non-metricity is actually massive (mass $\sim M$) by the same Stueckelberg mechanism, metricity and Einstein gravity are recovered below its decoupling scale (\sim Planck scale). Non-metricity effects are then strongly suppressed by a large M (their current lower bound seems low [73, 74]). Hence, the long-held criticisms that have implicitly assumed v_μ be massless are actually avoided and Weyl gravity is then viable [25–27]. As outlined, in this work we obtain similar results in EP quadratic gravity, up to different non-metricity effects.

We also study inflation in EP quadratic gravity (Section 5). We consider this theory with an extra scalar field (Higgs-like) with perturbative non-minimal coupling and Palatini connection, that plays the role of the inflaton. We compute the potential after the gauged scale symmetry breaking. With the Planck scale a simple phase transition scale in our theory, field values above M are natural. Interestingly, the inflaton potential is similar to that in Weyl quadratic gravity [27], up to couplings and field redefinitions (due to a different non-metricity of the theory). Inflation in EP quadratic gravity has a specific prediction for the tensor-to-scalar ratio $0.007 \leq r \leq 0.010$ for the current spectral index n_s at 95% CL. This range of r is distinct from that predicted by inflation in Weyl gravity [27, 46] and will soon be reached by CMB experiments [75–77]. The Conclusions are presented in Section 6 followed by an Appendix.

2 Palatini R^2 gravity

For later reference we first review R^2 gravity in the EP formalism [78, 79]. As discussed below, the action is local scale invariant (unlike its Riemannian counterpart):

$$L_1 = \sqrt{g} \frac{\xi_0}{4!} R(\tilde{\Gamma}, g)^2, \quad \xi_0 > 0, \quad (1)$$

where

$$R(\tilde{\Gamma}, g) = g^{\mu\nu} R_{\mu\nu}(\tilde{\Gamma}), \quad R_{\mu\nu}(\tilde{\Gamma}) = \partial_\lambda \tilde{\Gamma}_{\mu\nu}^\lambda - \partial_\mu \tilde{\Gamma}_{\lambda\nu}^\lambda + \tilde{\Gamma}_{\rho\lambda}^\lambda \tilde{\Gamma}_{\mu\nu}^\rho - \tilde{\Gamma}_{\rho\mu}^\lambda \tilde{\Gamma}_{\nu\lambda}^\rho \quad (2)$$

$R_{\mu\nu}(\tilde{\Gamma})$ is the metric-independent Ricci tensor in the EP formalism. Our conventions are as in [80] with metric $(+, -, -, -)$, $g \equiv |\det g_{\mu\nu}|$ and we assume there is no torsion i.e. $\tilde{\Gamma}_{\mu\nu}^\rho = \tilde{\Gamma}_{\nu\mu}^\rho$.

There is an equivalent “linearised” version of L_1 , found by using an auxiliary field ϕ

$$L_1 = \sqrt{g} \frac{\xi_0}{4!} \left\{ -2\phi^2 R(\tilde{\Gamma}, g) - \phi^4 \right\}. \quad (3)$$

Indeed, (1) is recovered if we use in (3) the solution $\phi^2 = -R(\tilde{\Gamma}, g)$ of the equation of motion of the scalar field ϕ . With the connection $\tilde{\Gamma}$ independent of the metric, (3) and (1) have local scale symmetry i.e. are invariant under a Weyl transformation $\Omega = \Omega(x)$ with⁴

$$\hat{g}_{\mu\nu} = \Omega^2 g_{\mu\nu}, \quad \sqrt{\hat{g}} = \Omega^4 \sqrt{g}, \quad \hat{\phi} = \frac{1}{\Omega} \phi, \quad \hat{R}(\tilde{\Gamma}, \hat{g}) = \frac{1}{\Omega^2} R(\tilde{\Gamma}, g). \quad (4)$$

Unlike in the metric case, $R_{\mu\nu}(\tilde{\Gamma})$ is invariant under (4) while $R(\tilde{\Gamma}, g)$ transforms covariantly, hence (1), (3) are invariant. L_1 has a shift symmetry: $\ln \phi \rightarrow \ln \phi - \ln \Omega$. In global cases $\ln \phi$ is the dilaton field generating a mass scale from its vev (assumed to be non-zero); here, $\ln \phi$ is similar to a would-be Goldstone, as seen if we “gauge” symmetry (4) (see later, eq.(18)).

Let us solve the equation of motion for $\tilde{\Gamma}$, then find the action for $\tilde{\Gamma}$ *onshell*⁵. The change of $R_{\mu\nu}(\tilde{\Gamma})$ under a variation of the connection is $\delta R_{\mu\nu}(\tilde{\Gamma}) = \tilde{\nabla}_\lambda (\delta \tilde{\Gamma}_{\mu\nu}^\lambda) - \tilde{\nabla}_\nu (\delta \tilde{\Gamma}_{\mu\lambda}^\lambda)$, where the operator $\tilde{\nabla}$ is defined with connection $\tilde{\Gamma}$. Then from (3) the equation of motion of $\tilde{\Gamma}_{\mu\nu}^\lambda$ gives

$$\tilde{\nabla}_\lambda (\sqrt{g} g^{\mu\nu} \phi^2) - \frac{1}{2} \left[\tilde{\nabla}_\rho (\sqrt{g} g^{\rho\mu} \phi^2) \delta_\lambda^\nu + (\mu \leftrightarrow \nu) \right] = 0. \quad (5)$$

Setting $\nu = \lambda$ and then summing over, then⁶

$$\tilde{\nabla}_\rho (\sqrt{g} g^{\rho\mu} \phi^2) = 0. \quad (6)$$

⁴From $\phi^2 = -R(\tilde{\Gamma}, g)$, ϕ^2 transforms under metric rescaling like $R(\tilde{\Gamma}, g)$, as expected for a scalar field.

⁵Obviously, with $\Omega^2 = \xi_0 \phi^2 / (6M^2)$ (with M the Planck scale), one can set ϕ to a constant (fix the “gauge” of local scale symmetry). L_1 becomes $L_1 = \sqrt{-g} \left\{ -(1/2)M^2 R(\tilde{\Gamma}, g) - 3/(2\xi_0)M^4 \right\}$. This is the Palatini formulation of Einstein action; via eqs motion then $\tilde{\nabla}_\mu g_{\alpha\beta} = 0$ where $\tilde{\nabla}_\mu$ is computed with $\tilde{\Gamma}$. Hence $\tilde{\Gamma}$ is a Levi-Civita connection. However, this approach obscures the role of local scale symmetry, relevant later.

⁶If we use $\phi^2 = -R(\tilde{\Gamma}, g)$ in (5), (6), one recovers the equation of motion of $\tilde{\Gamma}$ found directly from (1) [78].

To simplify notation, introduce an auxiliary dimensionful “metric” $h_{\mu\nu} \equiv \phi^2 g_{\mu\nu}$, then

$$\tilde{\nabla}_\lambda(\sqrt{h} h^{\mu\nu}) = 0. \quad (7)$$

This means that in terms of $h_{\mu\nu}$, the connection is Levi-Civita⁷

$$\tilde{\Gamma}_{\mu\nu}^\alpha(h) = (1/2) h^{\alpha\lambda} (\partial_\mu h_{\lambda\nu} + \partial_\nu h_{\lambda\mu} - \partial_\lambda h_{\mu\nu}), \quad (9)$$

or, in terms of $g_{\mu\nu}$

$$\tilde{\Gamma}_{\mu\nu}^\alpha = \Gamma_{\mu\nu}^\alpha(g) + (1/2) (\delta_\nu^\alpha u_\mu + \delta_\mu^\alpha u_\nu - g^{\alpha\lambda} g_{\mu\nu} u_\lambda), \quad u_\mu \equiv \partial_\mu \ln \phi^2, \quad (10)$$

with Levi-Civita $\Gamma_{\mu\nu}^\alpha(g) = (1/2) g^{\alpha\lambda} (\partial_\mu g_{\lambda\nu} + \partial_\nu g_{\lambda\mu} - \partial_\lambda g_{\mu\nu})$. Next, if we use the equation of motion of ϕ of solution $\phi^2 = -R(\tilde{\Gamma}, g)$, eq.(10) for $\tilde{\Gamma}$ (also (5), (6)) becomes a *second-order* differential equation since $\partial\phi^2 \sim \partial R \sim \partial^2 \tilde{\Gamma}$, and it is difficult to solve (and since solution $\tilde{\Gamma}$ of (10) involves $\partial g_{\mu\nu}$ from $\Gamma(g)$ then for $\tilde{\Gamma}$ onshell action (1) is a *four-derivative* theory in $g_{\mu\nu}$). An easy way out is to keep ϕ an independent variable hereafter (no use of its eq of motion), then eqs.(5), (6) have solution $\tilde{\Gamma}$ given by the rhs of (10). For this solution, then

$$R(\tilde{\Gamma}, g) = R(g) - 3\nabla_\mu u^\mu - \frac{3}{2} g^{\mu\nu} u_\mu u_\nu, \quad (11)$$

with the Ricci scalar $R(g)$ for $g_{\mu\nu}$ while ∇ is defined with the Levi-Civita connection (Γ). Using (11) in (3) of the same metric, we find for $\tilde{\Gamma}$ onshell⁸

$$L_1 = \sqrt{g} \left\{ \frac{\xi_0}{2} \left[-\frac{1}{6} \phi^2 R(g) - (\partial_\mu \phi)^2 - \frac{1}{12} \phi^4 \right] \right\}. \quad (12)$$

L_1 is a second order theory with an additional dynamical variable demanded by symmetry (4) and is equivalent to action (1) which for $\tilde{\Gamma}$ onshell is a four-derivative theory, as noticed.

Lagrangian (12) has local scale symmetry so one may like to “fix the gauge”. We choose the Einstein or unitarity gauge reached by a ϕ -dependent transformation $\Omega^2 = \phi^2 / \langle \phi \rangle^2$ that is gauge-fixing ϕ to a *constant* (vev); in this gauge $M^2 = \xi_0 \langle \phi \rangle^2 / 6$ is the Planck mass. From (12)

$$L_1 = \sqrt{\hat{g}} \left\{ \frac{-1}{2} M^2 \hat{R}(\hat{g}) - \frac{3}{2\xi_0} M^4 \right\}. \quad (13)$$

Hence Einstein action (13) is recovered *as a gauge fixed form* of (12); symmetry (4) is now spontaneously broken and ϕ decouples⁹ [81]; this may be expected since the local scale symmetry current of (12) is vanishing [82–84] (this will change in Section 3.3). With ϕ “gauge fixed” to a constant, eqs.(7),(10) give $h_{\mu\nu} \propto g_{\mu\nu}$ and $\tilde{\Gamma} = \Gamma$ so the theory is metric^{10,11}.

⁷One shows $\tilde{\nabla} h_{\mu\nu} = 0$ by using $\tilde{\nabla}_\lambda h^{\mu\nu} = -h^{\mu\sigma} h^{\nu\rho} \tilde{\nabla}_\lambda h_{\sigma\rho}$, and $\tilde{\nabla}_\lambda \sqrt{h} = (1/2) \sqrt{h} h^{\alpha\beta} \tilde{\nabla}_\lambda h_{\alpha\beta}$. (8)

⁸From (12) the equation of motion for $g^{\mu\nu}$ and its trace give that on the ground state $\phi^2 = -R$ and $R(R_{\mu\nu} - 1/4 g_{\mu\nu} R) = 0$ which is similar to that found directly from equivalent action (1), see also footnote 6.

⁹There is no gauge field here to “eat” would-be Goldstone $\ln \phi$ but in Section 3, v_μ of (18) will “eat” $\ln \phi$.

¹⁰This result is also valid for Palatini $f(R)$ action instead of (1); we do not consider it here since it violates Weyl scale symmetry, but (unlike here) the trace of the equation of motion of $g^{\mu\nu}$ is non-trivial, giving $f'(R) = \text{constant}$, which fixes R , then $h_{\mu\nu} \propto g_{\mu\nu}$, $\tilde{\Gamma} = \Gamma$ so metricity/Einstein action is recovered[3, 4].

¹¹This situation is different from the Riemannian $R(g)^2$ gravity [85] (eq.2.11), also [86], where in the Einstein frame a kinetic term for ϕ remains present in (13) (being absent in eq.(12)).

3 Palatini quadratic gravity with gauged scale symmetry

3.1 The Lagrangian and its expression for onshell $\tilde{\Gamma}$

Consider now the following EP quadratic gravity, with $\alpha=\text{constant}$ and $R_{[\mu\nu]} \equiv (R_{\mu\nu} - R_{\nu\mu})/2$

$$L_2 = \sqrt{g} \left\{ \frac{\xi_0}{4!} R(\tilde{\Gamma}, g)^2 - \frac{1}{4\alpha^2} R_{[\mu\nu]}(\tilde{\Gamma}) R^{\mu\nu}(\tilde{\Gamma}) \right\}. \quad (14)$$

With $R_{\mu\nu}(\tilde{\Gamma})$ from eq.(2) and $\tilde{\Gamma}_{\mu\nu}^\alpha$ symmetric in (μ, ν) , L_2 has a more intuitive form

$$L_2 = \sqrt{g} \left\{ \frac{\xi_0}{4!} R(\tilde{\Gamma}, g)^2 - \frac{1}{4\alpha^2} F_{\mu\nu}(\tilde{\Gamma}) F^{\mu\nu}(\tilde{\Gamma}) \right\}. \quad (15)$$

This is a natural extension of L_1 of eq.(1), with the second term above indicating we now have a dynamical trace ($\tilde{\Gamma}_\mu$) of the Palatini connection, as seen from the notation below:

$$F_{\mu\nu}(\tilde{\Gamma}) = \tilde{\nabla}_\mu v_\nu - \tilde{\nabla}_\nu v_\mu; \quad v_\mu = (1/2)(\tilde{\Gamma}_\mu - \Gamma_\mu(g)), \quad (16)$$

with $\tilde{\Gamma}_\mu \equiv \tilde{\Gamma}_{\mu\lambda}^\lambda$ and $\Gamma_\mu \equiv \Gamma_{\mu\lambda}^\lambda$. Since $\tilde{\Gamma}_{\mu\nu}^\alpha = \tilde{\Gamma}_{\nu\mu}^\alpha$ and $\tilde{\nabla}_\mu v_\nu = \partial_\mu v_\nu - \tilde{\Gamma}_{\mu\nu}^\alpha v_\alpha$, then we have $F_{\mu\nu} = \partial_\mu v_\nu - \partial_\nu v_\mu = (\partial_\mu \tilde{\Gamma}_\nu - \partial_\nu \tilde{\Gamma}_\mu)/2 = -R_{[\mu\nu]}$, and eqs.(14), (15) are equivalent. While $\Gamma_\mu(g)$ does not contribute to $F_{\mu\nu}(\tilde{\Gamma})^2$, it is needed to ensure that v_μ is a vector under coordinate transformation (which is not true for $\tilde{\Gamma}_\mu$ or Γ_μ , see Appendix). v_μ is the Weyl field¹² and measures the trace of the deviation of the Palatini connection $\tilde{\Gamma}$ from Levi-Civita connection $\Gamma(g)$. L_2 is quadratic in R but for $\tilde{\Gamma}$ offshell resembles a second order theory.

As in previous section, write L_2 in an equivalent “linearised” form useful later on

$$L_2 = \sqrt{g} \left\{ -\frac{\xi_0}{12} \phi^2 R(\tilde{\Gamma}, g) - \frac{1}{4\alpha^2} F_{\mu\nu}(\tilde{\Gamma})^2 - \frac{\xi_0}{4!} \phi^4 \right\}. \quad (17)$$

The equation of motion for ϕ has solution $\phi^2 = -R(\tilde{\Gamma}, g)$ which replaced in L_2 recovers (15).

Since $\tilde{\Gamma}$ does not transform under (4) and with $\Gamma_\mu(g) = \partial_\mu \ln \sqrt{g}$ that follows from the definition of Levi-Civita connection, then L_2 is invariant under (4) extended by

$$\hat{v}_\lambda = v_\lambda - \partial_\lambda \ln \Omega^2. \quad (18)$$

The invariance of L_2 under transformations (4), (18), is referred to as *gauged* scale invariance or Weyl gauge symmetry, with a (dilatation) group isomorphic to \mathbf{R}^+ , as in Weyl gravity.

Let us then compute the connection $\tilde{\Gamma}_{\mu\nu}^\lambda$ from its equation of motion which is

$$\tilde{\nabla}_\lambda(\sqrt{g} g^{\mu\nu} \phi^2) - \left\{ \frac{1}{2} \delta_\lambda^\nu \left[\tilde{\nabla}_\rho(\sqrt{g} g^{\mu\rho} \phi^2) - \frac{6\sqrt{g}}{\alpha^2 \xi_0} \nabla_\rho F^{\rho\mu} \right] + (\mu \leftrightarrow \nu) \right\} = 0. \quad (19)$$

¹²Definition (16) of gauge field v_μ is general, it also applies to Weyl gravity of similar symmetry (Appendix).

Here $\tilde{\nabla}_\mu$ and ∇_μ are evaluated with the Palatini ($\tilde{\Gamma}$) and Levi-Civita (Γ) connections, respectively. Setting $\lambda = \nu$ and summing over gives (compare against eq.(6))

$$\tilde{\nabla}_\rho(\sqrt{g} g^{\mu\rho} \phi^2) = \frac{10}{\alpha^2} \frac{1}{\xi_0} \sqrt{g} \nabla_\rho F^{\rho\mu}, \quad (20)$$

which is an equation of motion for the trace $\tilde{\Gamma}_\mu \sim v_\mu$. Replacing (20) back in (19) leads to

$$\tilde{\nabla}_\lambda(\sqrt{g} g^{\mu\nu} \phi^2) - \frac{1}{5} \left\{ \delta_\lambda^\nu \tilde{\nabla}_\rho(\sqrt{g} g^{\rho\mu} \phi^2) + (\mu \leftrightarrow \nu) \right\} = 0. \quad (21)$$

Therefore, the set of eqs.(19) is equivalent to the combined set of eqs.(21) and (20)¹³.

Let us find $\tilde{\Gamma}_{\mu\nu}^\lambda$ from (21). Note that if one used the equation of motion of ϕ of solution $\phi^2 = -R(\tilde{\Gamma}, g)$, then (21) would be a *second-order* differential equation for $\tilde{\Gamma}_{\mu\nu}^\alpha$, since $\tilde{\nabla}_\lambda \phi^2 \sim \partial \phi^2 \sim \partial R(\tilde{\Gamma}, g) \sim \partial^2 \tilde{\Gamma}$, with further complications. It is however easier to simply regard ϕ hereafter as an independent variable¹⁴ (i.e. no use of its equation of motion) in terms of which one then easily computes $\tilde{\Gamma}$ algebraically, as we do below. To find a solution to (21) we first introduce, based on an approach of [7]:

$$\tilde{\nabla}_\lambda(\sqrt{g} g^{\mu\nu} \phi^2) = (-2)\sqrt{g} \phi^2 (\delta_\lambda^\mu V^\nu + \delta_\lambda^\nu V^\mu), \quad (22)$$

where V_μ is some arbitrary vector field (to be determined later). V_μ is introduced since, due to underlying symmetry, eq.(21) with $\lambda = \nu$ summed over is automatically respected for fixed μ ($= 0, 1, 2, 3$); this is leaving four undetermined components, accounted for by V_μ . Further, if in eq.(21) one replaces $\tilde{\nabla}(\dots)$ terms by the rhs of (22) one easily shows that (21) is indeed verified. Hence, instead of finding $\tilde{\Gamma}$ from (21), it is sufficient to compute $\tilde{\Gamma}$ from (22)¹⁵, which is easier. To this end, multiply (22) by $g_{\mu\nu}$ and use $g_{\mu\nu} \tilde{\nabla}_\lambda g^{\mu\nu} = -2\tilde{\nabla}_\lambda \ln \sqrt{g}$, to find that

$$V_\lambda = -(1/2) \tilde{\nabla}_\lambda \ln(\sqrt{g} \phi^4). \quad (23)$$

From (22), (23)

$$\tilde{\nabla}_\lambda(\phi^2 g_{\mu\nu}) = (-2) (g_{\mu\nu} V_\lambda - g_{\mu\lambda} V_\nu - g_{\nu\lambda} V_\mu) \phi^2, \quad (24)$$

so the theory is non-metric. From (24) we find the solution¹⁶ $\tilde{\Gamma}$ to (21) *in terms of* V_λ :

$$\tilde{\Gamma}_{\mu\nu}^\alpha = \Gamma_{\mu\nu}^\alpha(\phi^2 g) - (3g_{\mu\nu} V_\lambda - g_{\nu\lambda} V_\mu - g_{\lambda\mu} V_\nu) g^{\lambda\alpha}, \quad (25)$$

$$\text{with } \Gamma_{\mu\nu}^\alpha(\phi^2 g) = \Gamma_{\mu\nu}^\alpha(g) + 1/2 (\delta_\nu^\alpha \partial_\mu + \delta_\mu^\alpha \partial_\nu - g^{\alpha\lambda} g_{\mu\nu} \partial_\lambda) \ln \phi^2.$$

¹³Unlike in (19), setting $\lambda = \nu$ in (21) brings no information - this was “moved” into 4 eqs in (20).

¹⁴The consequence of doing so is that ϕ acquires a kinetic term and becomes dynamical (see also Section 2).

¹⁵One cannot solve algebraically (22) as done in Palatini $f(R)$ theories [3, 4] due to non-vanishing rhs (dynamical $\tilde{\Gamma}_\mu$) and to the conformal symmetry of L_2 , absent in $f(R)$ theories, see discussion in [87], p.5-6.

¹⁶Use that $\tilde{\nabla}_\lambda g_{\mu\nu} = \partial_\lambda g_{\mu\nu} - \tilde{\Gamma}_{\mu\lambda}^\rho g_{\rho\nu} - \tilde{\Gamma}_{\nu\lambda}^\rho g_{\mu\rho}$, for cyclic permutations of indices and combine them.

$\Gamma_{\mu\nu}^\alpha(g)$ is Levi-Civita connection of $g_{\mu\nu}$. From (25), $\tilde{\Gamma}_\lambda = \Gamma_\lambda(\phi^2 g) + 2V_\lambda$ and with (16), (23)

$$v_\lambda = -(1/2) \tilde{\nabla}_\lambda \ln \sqrt{g}, \quad (26)$$

and finally, $V_\lambda = v_\lambda - \partial_\lambda \ln \phi^2$. With this relation between V_λ and v_λ , the solution $\tilde{\Gamma}$ in (25) is finally expressed as a function of v_λ , ϕ , and will be used shortly to compute the action for $\tilde{\Gamma}$ onshell (see eq.(29) below¹⁷). Notice that solution $\tilde{\Gamma}$ of (25) and also (24), are invariant under transformations (4), (18) for any $\Omega(x)$ since $\phi^2 g_{\mu\nu}$, V_λ , $\sqrt{g}\phi^4$ are invariant.

As expected, v_λ is the Weyl field of non-metricity defined as $Q_{\lambda\mu\nu} \equiv \tilde{\nabla}_\lambda g_{\mu\nu}$, since from (26) the trace $Q_{\lambda\mu}^\mu = -4v_\lambda$. Non-metricity is a consequence of the dynamical v_λ , see (20). Eq.(26) is similar to that in Weyl quadratic gravity of same symmetry (e.g.[32]).

Finally, from solution (25) and (2) we compute $R_{\mu\nu}(\tilde{\Gamma})$ and scalar curvature¹⁸ $R(\tilde{\Gamma}, g)$

$$R(\tilde{\Gamma}, g) = R(g) - 6g^{\mu\nu} \nabla_\mu \nabla_\nu \ln \phi - 6(\nabla_\mu \ln \phi)^2 - 12(\nabla_\lambda V^\lambda + V^\lambda \partial_\lambda \ln \phi^2) - 6V_\mu V^\mu. \quad (28)$$

$R(g)$ is here the usual Ricci scalar and $V_\lambda = w_\lambda - \partial_\lambda \ln \phi^2$. Using (28) in (17), then finally

$$L_2 = \sqrt{g} \left\{ -\frac{\xi_0}{12} [\phi^2 R(g) + 6(\partial_\mu \phi)^2] + \frac{\xi_0}{2} \phi^2 (v_\mu - \partial_\mu \ln \phi^2)^2 - \frac{1}{4\alpha^2} F_{\mu\nu}^2 - \frac{\xi_0}{4!} \phi^4 \right\}. \quad (29)$$

This is the “onshell” Lagrangian of EP quadratic gravity of eq.(14) and is gauged scale invariant. L_2 is a second-order scalar-vector-tensor theory of gravity which is ghost-free according to [89] for a torsion-free connection as here (this is also obvious from (30) below). This is relevant since initial action (14) which (offshell) was of second order is actually a four-derivative theory in the metric¹⁹ for $\tilde{\Gamma}$ onshell; indeed, $R(\tilde{\Gamma}, g)^2$ in (14) with replacement (28) contains the higher derivative term $R^2(g) + \dots$; this four-derivative theory has an equivalent second-order formulation with additional ϕ , as shown in eq.(29). Finally, if $v_\mu = \partial_\mu \ln \phi^2$ (“pure gauge”), the model is Weyl integrable and (29) recovers (12).

Lagrangian (29) (also initial (15)) is similar to that of Weyl quadratic gravity [25,26], up to a Weyl tensor-squared term not included here. However, unlike in Weyl theory, here $\tilde{\Gamma}$ is ϕ -dependent; also, in Weyl theory non-metricity follows from the underlying Weyl conformal geometry, while here it emerges after we determine $\tilde{\Gamma}$ from its equation of motion.

¹⁷As a remark, recall that eqs.(19) for $\tilde{\Gamma}$ were shown to be equivalent to the combined set of (21), (20) and we solved (21) with solution $\tilde{\Gamma}_{\mu\nu}^\alpha$ in (25) *expressed in terms of* $V_\mu \sim \tilde{\Gamma}_\mu$. We have four remaining equations in (20) for $\tilde{\Gamma}_\mu$ itself, that could in principle be used to also “fix” $\tilde{\Gamma}_\mu \sim V_\mu$ or equivalently v_μ , since $V_\mu = v_\mu - \partial_\mu \ln \phi^2$; however we do not do this step since v_μ is a *massless, dynamical* gauge field enforcing gauged scale symmetry (18) of initial action (14), (17). Then what information does (20) bring? With eq.(22) for $\lambda=\nu$, eq.(20) is actually $\tilde{\nabla}_\rho F^{\rho\mu} + \alpha^2 \xi_0 \phi^2 V^\mu = 0$, which is just the equation of motion of v_λ ; this may be seen from final Lagrangian (29) which has all $\tilde{\Gamma}_{\mu\nu}^\alpha$ onshell (expressed in terms of $\tilde{\Gamma}_\mu$) but $\tilde{\Gamma}_\mu$ is kept offshell, for the reason mentioned; v_λ may be integrated out after becoming massive, see later.

¹⁸ $R_{\mu\nu}(\tilde{\Gamma})$ has the following expression (which by contraction with $g^{\mu\nu}$ gives $R(\tilde{\Gamma}, g)$ of (28)):

$$\begin{aligned} R_{\mu\nu}(\tilde{\Gamma}) &= R_{\mu\nu}(g) - 3g_{\mu\nu}(\nabla^\lambda V_\lambda + V^\lambda \partial_\lambda \ln \phi^2) - (\nabla_\mu V_\nu - \nabla_\nu V_\mu) - 6V_\mu V_\nu + 1/2(\partial_\mu \theta)(\partial_\nu \theta) \\ &- 1/2 g_{\mu\nu} g^{\alpha\beta}(\partial_\alpha \theta)(\partial_\beta \theta) - 1/2 g_{\mu\nu} \nabla^\lambda \partial_\lambda \theta + 1/2 \nabla_\nu \partial_\mu \theta - 3/2 \nabla_\mu \partial_\nu \theta, \quad \theta \equiv \ln \phi^2. \end{aligned} \quad (27)$$

¹⁹This agrees with e.g.[18] that in general in a Palatini model its metric part leads to a fourth order theory.

3.2 Stueckelberg breaking to Einstein-Proca action

Given L_2 in (29) with gauged scale symmetry we would like to “fix the gauge”. We choose the Einstein gauge obtained from (29) by transformations (4), (18) of a special $\Omega^2 = \xi_0 \phi^2 / (6M^2)$ fixing ϕ to a *constant* ($\langle \phi \rangle \neq 0$). After removing the hats ($\hat{}$) on transformed g, v_μ, R , we find

$$L_2 = \sqrt{g} \left\{ -\frac{1}{2} M^2 R(g) + 3 M^2 v_\mu v_\nu g^{\mu\nu} - \frac{1}{4\alpha^2} F_{\mu\nu}^2 - \frac{3}{2\xi_0} M^4 \right\}. \quad (30)$$

This is the Einstein-Proca action for the gauge field v_μ with a positive cosmological constant, in which we identified M with the Planck scale (M) as seen from eq.(29)

$$M^2 \equiv \xi_0 \langle \phi \rangle^2 / 6. \quad (31)$$

The initial gauged scale invariance is broken by a gravitational Stueckelberg mechanism [70–72]: the massless ϕ is not part of the action anymore, but v_μ has become massive, after “absorbing” the derivative $\partial_\mu(\ln \phi)$ of the Stueckelberg field (dilaton) in eq.(29). Note that $\partial_\mu(\ln \phi)$ is actually the Goldstone of special conformal symmetry - this Goldstone is not independent but is determined by the derivative of the dilaton [88]. The number of degrees of freedom (dof) other than graviton is conserved in going from (29) to (30), as it should be for spontaneous breaking: massless v_μ and dynamical ϕ are replaced by massive v_μ (dof=3). The mass of v_μ is $m_v^2 = 6\alpha^2 M^2$ which is near Planck scale M (unless one fine-tunes $\alpha \ll 1$).

Using the same transformation Ω , from (24)

$$\tilde{\nabla}_\lambda g_{\mu\nu} = (-2)(g_{\mu\nu} v_\lambda - g_{\mu\lambda} v_\nu - g_{\nu\lambda} v_\mu). \quad (32)$$

This has a solution $\tilde{\Gamma}$ that is immediate from (25) for ϕ constant and V_λ replaced by v_λ . Finally, after the massive field v_μ decouples, metricity is recovered below m_v , so $\tilde{\nabla}_\lambda g_{\mu\nu} = 0$ and $\tilde{\Gamma} = \Gamma(g)$. Briefly, Einstein action is a “low energy” limit of Einstein-Palatini quadratic gravity, and the Planck scale $M \sim \langle \phi \rangle$ is a phase transition scale (up to coupling α)²⁰.

For comparison, in Weyl quadratic gravity e.g. [25, 26], non-metricity is different²¹

$$\tilde{\nabla}_\lambda g_{\mu\nu} = -g_{\mu\nu} v_\lambda. \quad (33)$$

Interestingly the different non-metricity of these theories (giving different $\tilde{\Gamma}$) has phenomenological impact, see Section 5. In both theories the non-metricity scale is $m_v \sim$ Planck scale and is large enough (current bounds [73, 74] are low \sim TeV) to suppresses unwanted effects e.g. atomic spectral lines spacing. Past critiques of non-metricity assumed a massless v_μ .

Finally, let us remark that the above spontaneous symmetry breaking mechanism for initial action (14) is special since it takes place *in the absence of matter*. Indeed, the necessary scalar (Stueckelberg) field $\ln \phi$ was not added ad-hoc to this purpose, as usually done in the literature; instead, this field was “extracted” from the R^2 term in the initial, symmetric action (14) and is thus of geometric origin. This situation is similar to Weyl quadratic gravity where this mechanism was first noticed [25, 26].

²⁰A special case: consider (17) with $\phi^2 = 6M^2/\xi_0 = \text{constant}$, i.e. a different initial action with no symmetry! then (28), (29) simplify; we still find (30) but there is no dynamical ϕ and thus no Stueckelberg mechanism.

²¹Contracting (32), (33) by $g^{\mu\nu}$ gives the same non-metricity trace, justifying our normalization of v_μ eq(16).

3.3 Conserved current

Eqs.(20) and (22) show there is now a non-trivial current due to dynamical $v_\mu \sim \tilde{\Gamma}_\mu$

$$J^\mu = \sqrt{g} g^{\rho\mu} \phi (\partial_\rho - 1/2 v_\rho) \phi, \quad \nabla_\mu J^\mu = 0, \quad (34)$$

This is conserved since $F_{\mu\nu}$ in (20) is anti-symmetric. To obtain (34) we used that the lhs of (20) and of (22) (with $\lambda = \nu$) are equal and replaced $V_\lambda = v_\lambda - \partial_\lambda \ln \phi^2$. The current J^μ is the same as that in Weyl quadratic gravity [25] (eq.18) which has similar symmetry but different non-metricity. The presence of this conserved current extends to the case of the *gauged* scale symmetry a similar conservation for a global scale symmetry [61]. For a Friedmann-Robertson-Walker metric with ϕ only t -dependent such current conservation in the global case naturally leads to $\phi = \text{constant}$ [61] and a breaking of scale symmetry. In our case, since eq.(30) has $\phi = \text{constant}$ (assumed $\langle \phi \rangle \neq 0$), then from (34) one has $\nabla_\mu v^\mu = 0$ which is a condition similar to that for a Proca (massive) gauge field, leaving 3 degrees of freedom for v_μ in (30).

4 Palatini quadratic gravity: adding matter

In this section we re-do the previous analysis in the presence of a scalar χ which can be the SM Higgs, with non-minimal coupling with *Palatini connection* to the EP quadratic gravity.

The general Lagrangian of the field χ , with gauged scale invariance, eqs.(4),(18) is

$$L_3 = \sqrt{g} \left[\frac{\xi_0}{4!} R(\tilde{\Gamma}, g)^2 - \frac{1}{4\alpha^2} F_{\mu\nu}^2 - \frac{1}{12} \xi_1 \chi^2 R(\tilde{\Gamma}, g) + \frac{1}{2} (\tilde{D}_\mu \chi)^2 - \frac{\lambda_1}{4!} \chi^4 \right], \quad (35)$$

with the potential dictated by this symmetry and with

$$\tilde{D}_\mu \chi = (\partial_\mu - 1/2 v_\mu) \chi. \quad (36)$$

Under (4), (18) the Weyl-covariant derivative transforms as $\hat{\tilde{D}}_\mu \hat{\chi} = (1/\Omega) \tilde{D}_\mu \chi$. As in previous sections, replace $R(\tilde{\Gamma}, g)^2 \rightarrow -2\phi^2 R(\tilde{\Gamma}, g) - \phi^4$ to find an equivalent “linearised” L_3

$$L_3 = \sqrt{g} \left[-\frac{1}{2} \rho^2 R(\tilde{\Gamma}, g) - \frac{1}{4\alpha^2} F_{\mu\nu}^2 + \frac{1}{2} (\tilde{D}_\mu \chi)^2 - \mathcal{V}(\chi, \rho) \right], \quad (37)$$

where

$$\mathcal{V}(\chi, \rho) \equiv \frac{1}{4!} \left[\frac{1}{\xi_0} (6\rho^2 - \xi_1 \chi^2)^2 + \lambda_1 \chi^4 \right], \quad \text{and} \quad \rho^2 = \frac{1}{6} (\xi_1 \chi^2 + \xi_0 \phi^2). \quad (38)$$

Notice that we also replaced the scalar field ϕ by the new, radial direction field ρ ; $\ln \rho$ transforms as $\ln \rho \rightarrow \ln \rho - \ln \Omega$ and acts as the (would-be) Goldstone of the symmetry.

The equation of motion for $\tilde{\Gamma}_{\mu\nu}^\lambda$ is similar to (19) but with a replacement $\phi \rightarrow \rho$ and with an additional contribution from the kinetic term of χ . Following the same steps as in the previous section, we eliminate the contributions of the kinetic terms of χ and v_μ to the equation of $\tilde{\Gamma}$ and find an equation similar to (21) with $\phi \rightarrow \rho$:

$$\tilde{\nabla}_\lambda(\sqrt{g}g^{\mu\nu}\rho^2) - \frac{1}{5}\left\{\delta_\lambda^\nu \tilde{\nabla}_\sigma(\sqrt{g}g^{\sigma\mu}\rho^2) + (\mu \leftrightarrow \nu)\right\} = 0. \quad (39)$$

This gives (see previous section):

$$\tilde{\nabla}_\lambda(\rho^2 g_{\mu\nu}) = (-2)\rho^2(g_{\mu\nu} V_\lambda - g_{\mu\lambda} V_\nu - g_{\nu\lambda} V_\mu), \quad (40)$$

where $V_\mu = (-1/2)\tilde{\nabla}_\mu \ln(\sqrt{g}\rho^4) = v_\mu - \partial_\mu \ln \rho^2$. From (40) one finds the solution for Palatini connection $\tilde{\Gamma}_{\mu\nu}^\alpha$ in terms of $v_\mu \sim \tilde{\Gamma}_\mu$, with a result similar to (25) but with $\phi \rightarrow \rho$. We use this solution for the connection back in the action and find for $\tilde{\Gamma}$ onshell²²

$$L_3 = \sqrt{g}\left\{\frac{-1}{2}\left[\rho^2 R(g) + 6(\partial_\mu \rho)^2\right] + 3\rho^2(v_\mu - \partial_\mu \ln \rho^2)^2 - \frac{1}{4\alpha^2} F_{\mu\nu}^2 + \frac{1}{2}(\tilde{D}_\mu \chi)^2 - \mathcal{V}(\chi, \rho)\right\}. \quad (41)$$

L_3 has a gauged scale symmetry and extents (29) in the presence of scalar field χ .

Finally, we choose the Einstein gauge by using transformation (4),(18) of a particular $\Omega = \rho/M$ which essentially sets $\hat{\rho}$ to a *constant* (vev). In terms of the new variables (with a hat) we find

$$L_3 = \sqrt{\hat{g}}\left\{-\frac{1}{2}M^2 R(\hat{g}) + 3M^2 \hat{v}_\mu \hat{v}^\mu - \frac{1}{4\alpha^2} \hat{F}_{\mu\nu}^2 + \frac{1}{2}(\hat{D}_\mu \hat{\chi})^2 - \mathcal{V}(\hat{\chi}, M)\right\}, \quad (42)$$

with $\hat{D}_\mu \hat{\chi} = (\partial_\mu - 1/2 \hat{v}_\mu) \hat{\chi}$ and we identify M with the Planck scale ($M = \langle \hat{\rho} \rangle$). As in the absence of matter, we obtained the Einstein-Proca action of a gauge field that became massive after Stueckelberg mechanism of “absorbing” the derivative term $\partial_\mu(\ln \rho)$. A canonical kinetic term of $\hat{\chi}$ remained present in the action, since only one degree of freedom (radial direction ρ) was “eaten” by v_μ . The mass of v_μ is $m_v^2 = 6\alpha^2 M^2$. The potential becomes

$$\mathcal{V} = \frac{3M^4}{2\xi_0}\left[1 - \frac{\xi_1 \hat{\chi}^2}{6M^2}\right]^2 + \frac{\lambda_1}{4!} \hat{\chi}^4. \quad (43)$$

For a “standard” kinetic term for $\hat{\chi}$, similar to a “unitary gauge” in electroweak case, we remove the coupling $\hat{v}^\mu \partial_\mu \hat{\chi}$ in the Weyl-covariant derivative in (42) by a field redefinition

$$\hat{v}'_\mu = \hat{v}_\mu - \partial_\mu \ln \cosh^2 \left[\frac{\sigma}{2M\sqrt{6}} \right], \quad \hat{\chi} = 2M\sqrt{6} \sinh \left[\frac{\sigma}{2M\sqrt{6}} \right], \quad (44)$$

which replaces $\hat{\chi} \rightarrow \sigma$. After some algebra, we find the final Lagrangian

²²As in Section 3, the trace $\tilde{\Gamma}_\mu \sim v_\mu$ is kept offshell since we do not integrate out *massless* dynamical v_μ .

$$L_3 = \sqrt{\hat{g}} \left\{ -\frac{1}{2} M^2 \hat{R} + 3M^2 \cosh^2 \left[\frac{\sigma}{2M\sqrt{6}} \right] \hat{v}'_\mu \hat{v}'^\mu - \frac{1}{4\alpha^2} \hat{F}_{\mu\nu}^2 + \frac{\hat{g}^{\mu\nu}}{2} \partial_\mu \sigma \partial_\nu \sigma - \hat{\mathcal{V}}(\sigma) \right\} \quad (45)$$

with

$$\hat{\mathcal{V}}(\sigma) = \hat{\mathcal{V}}_0 \left\{ \left[1 - 4\xi_1 \sinh^2 \frac{\sigma}{2M\sqrt{6}} \right]^2 + 16 \lambda_1 \xi_0 \sinh^4 \frac{\sigma}{2M\sqrt{6}} \right\}, \quad \hat{\mathcal{V}}_0 \equiv \frac{3}{2} \frac{M^4}{\xi_0}. \quad (46)$$

In (45) one finally rescales $\hat{v}'_\mu \rightarrow \alpha \hat{v}'_\mu$ for a canonical gauge kinetic term.

For *small* field values, $\sigma \ll M$, then $\hat{\chi} \approx \sigma$ (up to $\mathcal{O}(\sigma^3/M^2)$) and a SM Higgs-like potential is recovered²³, see eq.(43). For $\xi_1 > 0$ it has spontaneous breaking of the symmetry carried by σ i.e. electroweak (EW) symmetry if σ is the Higgs; this is triggered by the non-minimal coupling to gravity ($\xi_1 \neq 0$) and Stueckelberg mechanism. The negative mass term originates in (38) due to the ϕ^4 term (itself induced by \tilde{R}^2). The mass $m_\sigma^2 \propto \xi_1 M^2 / \xi_0$ may be small enough, near the EW scale by tuning $\xi_1 \ll \xi_0$. It may be interesting to study if the gauged scale symmetry brings some “protection” to m_σ at the quantum level.

L_3 of (45) is similar to that in Weyl quadratic gravity with a non-minimally coupled scalar/Higgs field [25–27]²⁴, up to a rescaling of the couplings (ξ_1, λ_1) and fields (σ). This difference is due to the different non-metricity of the two theories, eqs.(32), (33). Both cases provide a gauged scale invariant theory of quadratic gravity coupled to matter. They both recover Einstein gravity in their broken phase, see eq.(45), and also metricity below the scale $m_v \sim \alpha M$ ($\alpha \leq 1$). This result may be more general - it may apply to other theories with this symmetry and can be used for model building.

To conclude, mass generation (Planck scale, v_μ mass) and Einstein gravity emerge naturally from spontaneous breaking of gauged scale symmetry in Einstein-Palatini theories, even in the absence of matter. Actions (14), (35) were inspired by Weyl quadratic gravity of similar breaking [26]; but in a more general case, additional operators may be present in (14), (35); for a list of all quadratic operators and a complementary study see [13]. The mechanism of symmetry breaking should remain valid in their presence if one includes the terms in (14): R^2 that ‘supplied’ the scalar field and $R_{[\mu\nu]}^2$ generating the symmetry and non-metricity. However, in such general case it is unclear that one can still solve algebraically the second-order differential equations of motion of $\tilde{\Gamma}$ (eqs.(19)) without simplifying assumptions, since these equations acquire new terms of different indices structure and new states will be present (ghosts, etc).

5 Palatini R^2 inflation

In this section we consider an application to inflation of the action in the previous section.

For *large* field values, the potential in (46) can also be used for inflation (hereafter Palatini R^2 inflation), with σ as the inflaton²⁵. For a Friedmann-Robertson-Walker metric (FRW) $(1, -a^2(t), -a^2(t), -a^2(t))$ and compatible background $v_\mu(t) = (v_0(t), 0, 0, 0)$ the gauge fixing condition $\nabla_\mu v^\mu = 0$ gives that $v_\mu(t)$ redshifts to zero $v_\mu(t) \sim 1/a^3(t)$. Then the

²³We shall see shortly that inflation “prefers” ultraweak or vanishing values for λ_1 in (38) and (43).

²⁴For comparison to Weyl gravity Lagrangian see e.g. eqs.(39)-(41) in [25] and eqs.(21), (22) in [26].

²⁵Unlike in Starobinsky models, there is no scalaron here, its counterpart was “eaten” by massive v_μ .

coupling $v_\mu - \sigma$ in (45) is vanishing and therefore $v_\mu(t)$ cannot affect inflation; this means we have single-field inflation of potential (46) and standard slow-roll formulae can be used. Further, since M is just a phase transition scale, field values $\sigma \geq M$ are natural. $\hat{\mathcal{V}}(\sigma)$ is similar to that in Weyl gravity R^2 -inflation, see [27, 46] for a detailed analysis²⁶; however, as mentioned, the couplings and field normalization in the potential differ (for same initial couplings and non-metricity trace); hence the spectral index n_s and tensor-to-scalar ratio r are different, too, and need to be analyzed separately.

The potential is shown in Figure 1 for perturbative values of the couplings relevant for successful inflation. This demands $\lambda_1 \xi_0 \ll \xi_1^2 \ll 1$, with the first relation from demanding that the initial energy be larger than at the end of inflation $\hat{\mathcal{V}}_0 > \hat{\mathcal{V}}_{\min}$, respected by choosing a small enough λ_1 for given $\xi_{0,1}$. Therefore, we shall work in the leading order in $(\lambda_1 \xi_0)$.

The slow-roll parameters are:

$$\epsilon = \frac{M^2}{2} \left\{ \frac{\hat{\mathcal{V}}'(\sigma)}{\hat{\mathcal{V}}(\sigma)} \right\}^2 = \frac{4}{3} \xi_1^2 \sinh^2 \frac{\sigma}{M\sqrt{6}} + \mathcal{O}(\xi_1^3) \quad (47)$$

$$\eta = M^2 \frac{\hat{\mathcal{V}}''(\sigma)}{\hat{\mathcal{V}}(\sigma)} = -\frac{2}{3} \xi_1 \cosh \frac{\sigma}{M\sqrt{6}} + \mathcal{O}(\xi_1^2) \quad (48)$$

Then

$$n_s = 1 + 2\eta_* - 6\epsilon_* = 1 - \frac{4}{3} \xi_1 \cosh \frac{\sigma_*}{M\sqrt{6}} + \mathcal{O}(\xi_1^2), \quad (49)$$

where σ_* is the value of σ at the horizon exit. With $r = 16\epsilon_*$ we have²⁷

$$r = 12(1 - n_s)^2 + \mathcal{O}(\xi_1^2) \quad (50)$$

The contribution of ϵ is subleading for small ξ_1 considered here. The slope of the curves in the plane (n_s, r) , shown in leading order in (50), is steeper than in Weyl R^2 inflation [27] (or Starobinsky model) where $r = 3(1 - n_s)^2 + \mathcal{O}(\xi_1^2)$.

The exact numerical results for (n_s, r) in our model, for different e-folds number N , are shown in Figure 1. From experimental data $n_s = 0.9670 \pm 0.0037$ (68% CL) and $r < 0.07$ (95% CL) from Planck 2018 (TT, TE, EE + low E + lensing + BK14 + BAO) [101]. Using this data, Figure 1 (right plot) shows that a specific, small range for r is predicted in our model for the current range for n_s at 95% CL:

$$N = 60, \quad 0.007 \leq r \leq 0.010, \quad [\text{Palatini } R^2 \text{ inflation}]. \quad (51)$$

Similar values for r can be read from Figure 1 for $55 \leq N \leq 65$. The lower bound on r comes from that for n_s while the upper one corresponds to a saturation limit, $\xi_1 \rightarrow 0$, with values $\xi_1 < 10^{-3}$ having similar (n_s, r) . One should also respect the constraint $\lambda_1 \leq \xi_1^2/\xi_0$, giving $\lambda_1 \sim 10^{-12}$ or smaller (with the CMB anisotropy constraint $\xi_0 \geq 6.89 \times 10^8$).

²⁶For related works on inflation in the Einstein-Palatini formalism see e.g. [91–100].

²⁷There is also a constraint on the parametric space from the normalization of CMB anisotropy $\mathcal{V}_0/(24\pi^2 M^4 \epsilon_*) = \kappa_0$, $\kappa_0 = 2.1 \times 10^{-9}$ and $r = 16\epsilon_*$ with $r < 0.07$ [101] then $\xi_0 = 1/(\pi^2 r \kappa) \geq 6.89 \times 10^8$. The aforementioned condition $\lambda_1 \xi_0 \ll \xi_1^2$ is then respected for perturbative ξ_1 , $1/\xi_0$ by choosing small $\lambda_1 \ll \xi_1^2/\xi_0$.




Figure 1: **Left plot:** The potential $\hat{V}(\sigma)/\hat{V}_0$ for $\lambda_1 \xi_0 = 10^{-10} \ll \xi_1^2$ with different $\xi_1 \ll 1$. For larger $\lambda_1 \xi_0$ the curves move to the left while the minimum of the rightmost ones is lifted. Larger values of $\lambda_1 \xi_0$ are allowed, but inflation becomes less likely when $\lambda_1 \xi_0 \sim \xi_1^2$. The flat region is wide for a large range of σ , with the width controlled by $1/\sqrt{\xi_1}$ while its height is $\hat{V}_0 \propto 1/\xi_0$. We have $\hat{V}/\hat{V}_{\min} \propto \xi_1^2/(\lambda_1 \xi_0)$. **Right plot:** The values of (n_s, r) for different values of ξ_1 that enable values of $n_s = 0.9670 \pm 0.0037$ at 68% CL (blue band) and 95% CL (light blue region). For each curve $N = 60$ e-folds is marked by a red point and the dark blue interval corresponds to $55 \leq N \leq 65$. Curves of $\xi_1 < 10^{-3}$ are degenerate with the red one while those with $\xi_1 > 2.5 \times 10^{-2}$ have $N > 65$.

For comparison, in Weyl R^2 -inflation for same n_s at 95% CL one has a smaller r [27, 46]

$$N = 60, \quad 0.00257 \leq r \leq 0.00303, \quad [\text{Weyl } R^2 \text{ inflation}]. \quad (52)$$

The different range for r in eq.(51) versus eq.(52) is important since it enables us to distinguish these two inflation models based on gauged scale invariance, and is due to their different non-metricity^{28,29}. Such values for $r \sim 10^{-3}$ will soon be reached by various CMB experiments [75–77] that will then be able to test both models. This establishes an interesting connection between non-metricity and testable inflation predictions.

Similar values for r were found in other recent inflation models in Palatini R^2 gravity [98–100] but these are not gauged scale invariant. In the absence of this symmetry, other successful models (e.g. Starobinsky model [102]) have corrections to r from higher curvature operators (R^4 , etc) of unknown coefficients [104]. Such operators (and their corrections) are not allowed here because they must be suppressed by some effective scale whose presence would violate scale invariance³⁰. Another advantage is that due to the *gauged* scale symmetry Palatini R^2 inflation is allowed by black-hole physics (similarly for Weyl R^2 inflation [27]), in contrast to models of inflation with *global* scale symmetry³¹.

²⁸For a more detailed comparison of Einstein-Palatini R^2 -inflation to Weyl R^2 -inflation see [90].

²⁹In the Starobinsky model [102] for similar n_s one has $r \sim \mathcal{O}(10^{-3})$, e.g. $r = 0.0034$ ($N = 55$) [103].

³⁰The dilaton field cannot suppress them itself since it is “eaten” to all orders by the gauge field v_μ .

³¹A global symmetry is broken since global charges can be eaten by black holes which then evaporate [105].

6 Conclusions

At a fundamental level gravity may be regarded as a theory of connections. An example is the Einstein-Palatini (EP) approach to gravity where the connection ($\tilde{\Gamma}$) is apriori independent of the metric, and is determined by its equation of motion, from the action. For simple actions $\tilde{\Gamma}$ plays an auxiliary role (no dynamics) and can be solved algebraically. In particular, for Einstein action in the EP approach one finds that the connection is actually equal to the Levi-Civita connection (of the metric formulation); then Einstein gravity is recovered, so the metric and EP approaches are equivalent. However, this equivalence is not true in general, for complicated actions, etc. In this work we considered quadratic gravity actions in the EP approach, with the goal to show that, while this equivalence does not hold true, one can still find actions that recover *dynamically* the Levi-Civita connection, metricity, Einstein gravity and Planck mass in some “low-energy” limit, even in the *absence* of matter.

We studied EP quadratic gravity given by $R(\tilde{\Gamma}, g)^2 + R_{[\mu\nu]}(\tilde{\Gamma})^2$ which has local scale symmetry. $R_{[\mu\nu]}(\tilde{\Gamma})^2$ can be regarded as a gauge kinetic term for the vector field $v_\mu \sim \tilde{\Gamma}_\mu - \Gamma_\mu$ where $\tilde{\Gamma}_\mu$ (Γ_μ) denotes the trace of the Palatini (Levi-Civita) connections, respectively. Hence this theory actually has a *gauged* scale symmetry, with v_μ the Weyl gauge field. A consequence of this symmetry is that the theory is non-metric i.e. $\tilde{\nabla}_\mu g_{\alpha\beta} \neq 0$ (due to dynamical $v_\mu \sim \tilde{\Gamma}_\mu$). At the same time, the equations of motion of the connection ($\tilde{\Gamma}$) become complicated second-order differential equations and we showed how to solve them *algebraically* in terms of an auxiliary scalar ϕ that “linearises” the $R(\tilde{\Gamma}, g)^2$ term. While initially the action appears to be of second order, for $\tilde{\Gamma}$ onshell it is a *higher derivative* theory since $R(\tilde{\Gamma}, g)^2$ contains a (four-derivative) metric contribution $R(g)^2 + \dots$. We showed that for $\tilde{\Gamma}$ onshell, the action is equivalent to a second-order theory in which the initial auxiliary field ϕ has become dynamical, while preserving the symmetry of the theory.

The main result is that our EP quadratic gravity action has an elegant spontaneous breaking mechanism of gauged scale invariance and mass generation valid even in the absence of matter; in this, the necessary scalar field (ϕ) was not added ad-hoc to this purpose (as usually done), but was “extracted” from the R^2 term, as mentioned, being of geometric origin. The derivative $\partial_\mu \ln \phi$ of this field acting as a Stueckelberg field is “eaten” by v_μ which becomes massive, of mass m_v proportional to the Planck scale $M \sim \langle \phi \rangle$. One obtains the Einstein-Proca action for the gauge field v_μ and a positive cosmological constant. This is a “low-energy” broken phase of the initial action. Below the scale $m_v \sim M$, the Proca field v_μ decouples and metricity and the Einstein action are recovered. Non-metricity effects are strongly suppressed by a large scale ($\propto M$), which is important for the theory to be viable.

The above results remain valid in the presence of scalar matter (Higgs, etc) with a (perturbative) non-minimal coupling to this theory with a *Palatini connection*; in such case and following the Stueckelberg mechanism, the scalar potential also has a breaking of the symmetry under which this scalar is charged, e.g. electroweak symmetry in the higgs case. This is relevant for building models with this symmetry for physics beyond the SM.

To summarise, Einstein-Palatini quadratic gravity $R(\tilde{\Gamma}, g)^2 + R_{[\mu\nu]}^2(\tilde{\Gamma})$ is a gauged theory of scale invariance that is spontaneously broken to the Einstein-Proca action for the Weyl field with a positive cosmological constant; if initial action also contains (non-minimally coupled) scalar fields with Palatini connection, a scalar potential is also present.

This picture is similar to a recent analysis for the original Weyl quadratic gravity, despite the different non-metricity of these two theories. With hindsight, this is not too surprising, since in both theories there is a gauged scale symmetry and the connection is not fixed by the metric, except that in Weyl gravity non-metricity is present from the onset (due to underlying Weyl conformal geometry) while here it emerges for $\tilde{\Gamma}$ onshell. It is worth studying further the relation of these two theories, by including any remaining operators (on the Einstein-Palatini side) that can have this symmetry.

There are also interesting predictions from inflation. While the scalar potential is Higgs-like for small field values ($\ll M$), for large field values it can be used for inflation. With the Planck scale M a simple phase transition scale, field values above M are natural. The inflaton potential is similar to that in Weyl quadratic gravity, up to couplings and field redefinitions (due to different non-metricity of the two theories). We find a specific prediction for the tensor-to-scalar ratio, $0.007 \leq r \leq 0.01$, for the current value of the spectral index at 95% CL. This value of r is mildly larger than that predicted by inflation in Weyl gravity. This enables us to distinguish and test these two theories by future CMB experiments that will reach such values of r . It also establishes an interesting connection between non-metricity and inflation predictions.

Acknowledgements: The author thanks Graham Ross for helpful discussions on this topic at an early stage of this work.

Appendix

For a self-contained presentation, we include here basic aspects of the Palatini formalism used in the text. In the (pseudo-)Riemannian geometry, the Levi-Civita connection $\Gamma(g)$ is determined by the metric. In general, however, the connection can be introduced without reference to $g_{\mu\nu}$. In the Palatini approach the connection $\tilde{\Gamma}$ is apriori independent of the metric and is determined by the equations of motion. To ensure the covariant derivatives transform under coordinate change ($x \rightarrow x'(x)$) as true tensors, Palatini connection has a transformation law

$$\tilde{\Gamma}'_{\mu\nu}{}^{\lambda} = \frac{\partial x^{\alpha}}{\partial x'^{\mu}} \frac{\partial x^{\beta}}{\partial x'^{\nu}} \frac{\partial x'^{\lambda}}{\partial x^{\rho}} \tilde{\Gamma}_{\alpha\beta}{}^{\rho} + \frac{\partial x'^{\lambda}}{\partial x^{\sigma}} \frac{\partial^2 x^{\sigma}}{\partial x'^{\mu} \partial x'^{\nu}} \quad (53)$$

with $\tilde{\Gamma}' = \tilde{\Gamma}'(x')$, $\tilde{\Gamma} = \tilde{\Gamma}(x)$. In the text we also assumed $\tilde{\Gamma}_{\mu\nu}{}^{\rho} = \tilde{\Gamma}_{\nu\mu}{}^{\rho}$ (no torsion). The Levi-Civita connection $\Gamma_{\mu\nu}^{\lambda}(g)$ has a similar transformation

$$\Gamma'_{\mu\nu}{}^{\lambda}(g) = \frac{\partial x^{\alpha}}{\partial x'^{\mu}} \frac{\partial x^{\beta}}{\partial x'^{\nu}} \frac{\partial x'^{\lambda}}{\partial x^{\rho}} \Gamma_{\alpha\beta}{}^{\rho}(g) + \frac{\partial x'^{\lambda}}{\partial x^{\sigma}} \frac{\partial^2 x^{\sigma}}{\partial x'^{\mu} \partial x'^{\nu}}. \quad (54)$$

Note that the difference of these connections transforms as a tensor

$$\tilde{\Gamma}'_{\mu\nu}{}^{\lambda} - \Gamma'_{\mu\nu}{}^{\lambda}(g) = \frac{\partial x^{\alpha}}{\partial x'^{\mu}} \frac{\partial x^{\beta}}{\partial x'^{\nu}} \frac{\partial x'^{\lambda}}{\partial x^{\rho}} (\tilde{\Gamma}_{\alpha\beta}{}^{\rho}(x) - \Gamma_{\alpha\beta}{}^{\rho}(g)). \quad (55)$$

Setting $\lambda = \nu$ and with the notation $\tilde{\Gamma}_\mu \equiv \tilde{\Gamma}_{\mu\nu}^\nu$, $\Gamma_\mu \equiv \Gamma_{\mu\nu}^\nu$, etc, then

$$\tilde{\Gamma}'_\mu - \Gamma'_\mu(g) = \frac{\partial x^\alpha}{\partial x'^\mu} (\tilde{\Gamma}_\alpha - \Gamma_\alpha(g)) \quad (56)$$

and therefore v_μ introduced in Section 3 transforms as a covariant vector

$$v'_\mu(x') = \frac{\partial x^\alpha}{\partial x'^\mu} v_\alpha(x) \quad (57)$$

Further, the covariant derivatives used in the text are

$$\tilde{\nabla}_\nu v_\mu = \partial_\nu v_\mu - \tilde{\Gamma}_{\mu\nu}^\lambda v_\lambda, \quad \tilde{\nabla}_\nu v^\mu = \partial_\nu v^\mu + \tilde{\Gamma}_{\lambda\nu}^\mu v^\lambda. \quad (58)$$

One also has $\tilde{\nabla}_\lambda g_{\mu\nu} = \partial_\lambda g_{\mu\nu} - \tilde{\Gamma}_{\mu\lambda}^\rho g_{\rho\nu} - \tilde{\Gamma}_{\nu\lambda}^\rho g_{\rho\mu}$, also used in the text.

In Section 3 we introduced the gauge field v_μ in eq.(16). This is general. For example, in Weyl gravity of similar gauged scale symmetry, an identical formula exists for the gauge field. To see this, note that in Weyl gravity [25,26], see also eq.(33) in the text, non-metricity is different from EP quadratic gravity: $\tilde{\nabla}_\lambda g_{\mu\nu} = -v_\lambda g_{\mu\nu}$. Contracting this equation with $g^{\mu\nu}$ and using $\tilde{\nabla}_\lambda \sqrt{g} = (1/2)\sqrt{g} g^{\mu\nu} \tilde{\nabla}_\lambda g_{\mu\nu}$ we find $v_\lambda = (-1/2)\tilde{\nabla}_\lambda \ln \sqrt{g}$. This is similar to Palatini case, eq.(26) in the text, although the connection is different. From these last two equations, by writing the action of $\tilde{\nabla}_\lambda$ on $g_{\mu\nu}$ one immediately finds

$$v_\mu = (1/2)(\tilde{\Gamma}_\mu - \Gamma_\mu(g)) \quad (59)$$

with the trace of Levi-Civita connection $\Gamma_\mu(g) = \partial_\mu \ln \sqrt{g}$. Eq.(59) was used as a definition for the gauge field in Einstein-Palatini quadratic gravity, eq.(16).

References

- [1] A. Einstein, “Einheitliche Feldtheories von Gravitation und Electrizaritat”, Sitzungber Preuss Akad. Wiss (1925) 414-419.
- [2] M. Ferraris, M. Francaviglia and C. Reina, “Variational formulation of general relativity from 1915 to 1925, “Palatini’s method” discovered by Einstein in 1925”, Gen. Rel. Grav. 14 (1982) 243-254.
- [3] For a review and references, see G. J. Olmo, “Palatini Approach to Modified Gravity: f(R) Theories and Beyond,” Int. J. Mod. Phys. D **20** (2011) 413 [arXiv:1101.3864 [gr-qc]].
- [4] Another review is: T. P. Sotiriou and S. Liberati, “Metric-affine f(R) theories of gravity,” Annals Phys. **322** (2007) 935 [gr-qc/0604006]. T. P. Sotiriou and V. Faraoni, “f(R) Theories Of Gravity,” Rev. Mod. Phys. **82** (2010) 451 [arXiv:0805.1726 [gr-qc]].
- [5] H. A. Buchdahl, “Representation Of The Einstein-proca Field By An A(4)*,” J. Phys. A **12** (1979) 1235.
- [6] H. A. Buchdahl, “Non-linear Lagrangians and cosmological theory,” Mon. Not. Roy. Astron. Soc. **150** (1970) 1.

- [7] D. N. Vollick, “Einstein-Maxwell and Einstein-Proca theory from a modified gravitational action,” gr-qc/0601016.
- [8] D. N. Vollick, “Born-Infeld-Einstein theory with matter,” Phys. Rev. D **72** (2005) 084026 [gr-qc/0506091].
- [9] S. Cotsakis, J. Miritzis and L. Querella, “Variational and conformal structure of nonlinear metric connection gravitational Lagrangians,” J. Math. Phys. **40** (1999) 3063 [gr-qc/9712025].
- [10] B. Shahid-Saless, “First Order Formalism Treatment of $R + R^{**2}$ Gravity,” Phys. Rev. D **35** (1987) 467.
- [11] E. E. Flanagan, “Palatini form of $1/R$ gravity,” Phys. Rev. Lett. **92** (2004) 071101 [astro-ph/0308111].
- [12] E. E. Flanagan, “Higher order gravity theories and scalar tensor theories,” Class. Quant. Grav. **21** (2003) 417 [gr-qc/0309015].
- [13] M. Borunda, B. Janssen and M. Bastero-Gil, “Palatini versus metric formulation in higher curvature gravity,” JCAP **0811** (2008) 008 [arXiv:0804.4440 [hep-th]].
- [14] L. Järv, M. Rünkla, M. Saal and O. Vilson, “Nonmetricity formulation of general relativity and its scalar-tensor extension,” Phys. Rev. D **97** (2018) no.12, 124025 [arXiv:1802.00492 [gr-qc]].
- [15] R. Percacci and E. Sezgin, “New class of ghost- and tachyon-free metric affine gravities,” Phys. Rev. D **101** (2020) no.8, 084040 [arXiv:1912.01023 [hep-th]].
- [16] L. Querella, “Variational principles and cosmological models in higher order gravity,” gr-qc/9902044.
- [17] V. Vitagliano, T. P. Sotiriou and S. Liberati, “The dynamics of generalized Palatini Theories of Gravity,” Phys. Rev. D **82** (2010) 084007 [arXiv:1007.3937 [gr-qc]].
- [18] G. Allemandi, A. Borowiec, M. Francaviglia and S. D. Odintsov, “Dark energy dominance and cosmic acceleration in first order formalism,” Phys. Rev. D **72** (2005), 063505 [arXiv:gr-qc/0504057 [gr-qc]].
- [19] A. Kozak and A. Borowiec, “Palatini frames in scalar-tensor theories of gravity,” Eur. Phys. J. C **79** (2019) no.4, 335 [arXiv:1808.05598 [hep-th]].
- [20] J. Annala, “Higgs inflation and higher-order gravity in Palatini formulation”, PhD thesis, Univ of Helsinki, <https://inspirehep.net/literature/1799417>
- [21] Hermann Weyl, Gravitation und elektrizität, Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin (1918), pp.465; Einstein’s critical comment appended (on atomic spectral lines changes).
- [22] Hermann Weyl “Eine neue Erweiterung der Relativitätstheorie” (“A new extension of the theory of relativity”), Ann. Phys. (Leipzig) (4) 59 (1919), 101-133.
- [23] Hermann Weyl “Raum, Zeit, Materie”, vierte erweiterte Auflage. Julius Springer, Berlin 1921 “Space-time-matter”, translated from German by Henry L. Brose, 1922, Methuen & Co Ltd, London.

- [24] For review and references on Weyl’s theory, see E. Scholz, “The unexpected resurgence of Weyl geometry in late 20-th century physics,” *Einstein Stud.* **14** (2018) 261 [arXiv:1703.03187 [math.HO]];
- [25] D. M. Ghilencea, “Spontaneous breaking of Weyl quadratic gravity to Einstein action and Higgs potential,” *JHEP* **1903** (2019) 049 [arXiv:1812.08613 [hep-th]].
- [26] D. M. Ghilencea, “Stueckelberg breaking of Weyl conformal geometry and applications to gravity,” *Phys. Rev. D* **101** (2020) no.4, 045010 [arXiv:1904.06596 [hep-th]].
- [27] D. M. Ghilencea, “Weyl R^2 inflation with an emergent Planck scale,” *JHEP* **1910** (2019) 209 [arXiv:1906.11572 [gr-qc]].
- [28] P. A. M. Dirac, “Long range forces and broken symmetries,” *Proc. Roy. Soc. Lond. A* **333** (1973) 403. doi:10.1098/rspa.1973.0070
- [29] L. Smolin, “Towards a Theory of Space-Time Structure at Very Short Distances,” *Nucl. Phys. B* **160** (1979) 253.
- [30] H. Cheng, “The Possible Existence of Weyl’s Vector Meson,” *Phys. Rev. Lett.* **61** (1988) 2182.
- [31] I. Quiros, “Scale invariant theory of gravity and the standard model of particles,” E-print arXiv:1401.2643 [gr-qc].
- [32] T. Fulton, F. Rohrlich and L. Witten, “Conformal invariance in physics,” *Rev. Mod. Phys.* **34** (1962) 442.
- [33] J. T. Wheeler, “Weyl geometry,” *Gen. Rel. Grav.* **50** (2018) no.7, 80 [arXiv:1801.03178 [gr-qc]].
- [34] M. de Cesare, J. W. Moffat and M. Sakellariadou, “Local conformal symmetry in non-Riemannian geometry and the origin of physical scales,” *Eur. Phys. J. C* **77** (2017) no.9, 605 [arXiv:1612.08066 [hep-th]].
- [35] H. C. Ohanian, “Weyl gauge-vector and complex dilaton scalar for conformal symmetry and its breaking,” *Gen. Rel. Grav.* **48** (2016) no.3, 25 [arXiv:1502.00020 [gr-qc]].
- [36] D. M. Ghilencea and H. M. Lee, “Weyl symmetry and its spontaneous breaking in Standard Model and inflation,” arXiv:1809.09174 [hep-th].
- [37] A. Barnaveli, S. Lucat and T. Prokopec, “Inflation as a spontaneous symmetry breaking of Weyl symmetry,” arXiv:1809.10586 [gr-qc].
- [38] J. W. Moffat, “Scalar-tensor-vector gravity theory,” *JCAP* **0603** (2006) 004 [gr-qc/0506021].
- [39] L. Heisenberg, “Scalar-Vector-Tensor Gravity Theories,” arXiv:1801.01523 [gr-qc].
- [40] J. Beltran Jimenez, L. Heisenberg, T. S. Koivisto, “Cosmology for quadratic gravity in generalized Weyl geometry,” *JCAP* **1604** (2016) no.04, 046 [arXiv:1602.07287 [hep-th]].
- [41] J. Beltran Jimenez, T. S. Koivisto, “Spacetimes with vector distortion: Inflation from generalised Weyl geometry,” *Phys. Lett. B* **756** (2016) 400 [arXiv:1509.02476 [gr-qc]].
- [42] C. T. Hill, “Inertial Symmetry Breaking,” arXiv:1803.06994 [hep-th].
- [43] E. Scholz, “Higgs and gravitational scalar fields together induce Weyl gauge,” *Gen. Rel. Grav.* **47** (2015) no.2, 7 [arXiv:1407.6811 [gr-qc]].

- [44] W. Drechsler and H. Tann, “Broken Weyl invariance and the origin of mass,” *Found. Phys.* **29** (1999) 1023 [gr-qc/9802044].
- [45] S. Dengiz, “A Note on Noncompact and Nonmetricit Quadratic Curvature Gravity Theories,” *Turk. J. Phys.* **42** (2018) no.1, 70 [arXiv:1404.2714 [hep-th]].
- [46] P. G. Ferreira, C. T. Hill, J. Noller and G. G. Ross, “Scale-independent R^2 inflation,” *Phys. Rev. D* **100** (2019) no.12, 123516 [arXiv:1906.03415 [gr-qc]].
- [47] G. ’t Hooft, “Local conformal symmetry: The missing symmetry component for space and time,” *Int. J. Mod. Phys. D* **24** (2015) no.12, 1543001.
- [48] I. Bars, P. Steinhardt, N. Turok, “Local Conformal Symmetry in Physics and Cosmology,” *Phys. Rev. D* **89** (2014) no.4, 043515 [arXiv:1307.1848 [hep-th]].
- [49] G. ’t Hooft, “Imagining the future, or how the Standard Model may survive the attacks,” *Int. J. Mod. Phys.* **31** (2016) no.16, 1630022.
- [50] G. ’t Hooft, “Local conformal symmetry in black holes, standard model, and quantum gravity,” *Int. J. Mod. Phys. D* **26** (2016) no.03, 1730006.
- [51] M. Shaposhnikov and D. Zenhausern, “Quantum scale invariance, cosmological constant and hierarchy problem,” *Phys. Lett. B* **671** (2009) 162 [arXiv:0809.3406 [hep-th]].
- [52] R. Armillis, A. Monin and M. Shaposhnikov, “Spontaneously Broken Conformal Symmetry: Dealing with the Trace Anomaly,” *JHEP* **1310** (2013) 030 [arXiv:1302.5619 [hep-th]].
- [53] F. Bezrukov, G. K. Karananas, J. Rubio and M. Shaposhnikov, “Higgs-Dilaton Cosmology: an effective field theory approach,” *Physical Review D* **87** (2013) no.9, 096001 [arXiv:1212.4148 [hep-ph]].
- [54] F. Gretsche and A. Monin, “Perturbative conformal symmetry and dilaton,” *Phys. Rev. D* **92** (2015) no.4, 045036 [arXiv:1308.3863 [hep-th]].
- [55] D. M. Ghilencea, “Quantum implications of a scale invariant regularization,” *Phys. Rev. D* **97** (2018) no.7, 075015 [arXiv:1712.06024 [hep-th]]; “Manifestly scale-invariant regularization and quantum effective operators,” *Phys. Rev. D* **93** (2016) no.10, 105006 [arXiv:1508.00595 [hep-ph]]; “One-loop potential with scale invariance and effective operators,” *PoS CORFU 2015* (2016) 040 [arXiv:1605.05632 [hep-ph]].
- [56] D. M. Ghilencea, Z. Lalak and P. Olszewski, “Two-loop scale-invariant scalar potential and quantum effective operators,” *Eur. Phys. J. C* **76** (2016) no.12, 656 [arXiv:1608.05336 [hep-th]].
- [57] D. M. Ghilencea, Z. Lalak and P. Olszewski, “Standard Model with spontaneously broken quantum scale invariance,” *Phys. Rev. D* **96** (2017) no.5, 055034 [arXiv:1612.09120 [hep-ph]].
- [58] J. Kubo, M. Lindner, K. Schmitz and M. Yamada, “Planck mass and inflation as consequences of dynamically broken scale invariance,” *Phys. Rev. D* **100** (2019) no.1, 015037 [arXiv:1811.05950 [hep-ph]].
- [59] R. Foot, A. Kobakhidze, K. L. McDonald and R. R. Volkas, “Poincaré protection for a natural electroweak scale,” *Phys. Rev. D* **89** (2014) no.11, 115018 [arXiv:1310.0223 [hep-ph]].
- [60] P. G. Ferreira, C. T. Hill and G. G. Ross, “Scale-Independent Inflation and Hierarchy Generation,” *Phys. Lett. B* **763** (2016) 174 [arXiv:1603.05983 [hep-th]].

- [61] P. G. Ferreira, C. T. Hill and G. G. Ross, “Inertial Spontaneous Symmetry Breaking and Quantum Scale Invariance,” arXiv:1801.07676 [hep-th].
- [62] P. G. Ferreira, C. T. Hill and G. G. Ross, “Weyl Current, Scale-Invariant Inflation and Planck Scale Generation,” Phys. Rev. D **95** (2017) no.4, 043507 [arXiv:1610.09243 [hep-th]].
- [63] E. J. Chun, S. Jung and H. M. Lee, “Radiative generation of the Higgs potential,” Phys. Lett. B **725** (2013) 158 Erratum: [Phys. Lett. B **730** (2014) 357] [arXiv:1304.5815 [hep-ph]].
- [64] O. Lebedev and H. M. Lee, “Higgs Portal Inflation,” Eur. Phys. J. C **71** (2011) 1821 [arXiv:1105.2284 [hep-ph]].
- [65] Z. Lalak and P. Olszewski, “Vanishing trace anomaly in flat spacetime,” Phys. Rev. D **98** (2018) no.8, 085001 [arXiv:1807.09296 [hep-th]].
- [66] E. Elizalde, S. D. Odintsov and A. Romeo, “Manifestations of quantum gravity in scalar QED phenomena,” Phys. Rev. D **51** (1995), 4250-4253 [arXiv:hep-th/9410028 [hep-th]].
- [67] I. Buchbinder, S. Odintsov and I. Shapiro, “Effective action in quantum gravity,” Published in Bristol, UK: IOP (1992) 413 p.
- [68] A. Salvio and A. Strumia, “Agravity,” JHEP **1406** (2014) 080 [arXiv:1403.4226 [hep-ph]].
A. Salvio and A. Strumia, “Agravity up to infinite energy,” Eur. Phys. J. C **78** (2018) no.2, 124 [arXiv:1705.03896 [hep-th]].
- [69] D. Iosifidis, A. C. Petkou and C. G. Tsagas, “Torsion/non-metricity duality in f(R) gravity,” Gen. Rel. Grav. **51** (2019) no.5, 66 [arXiv:1810.06602 [gr-qc]].
- [70] E. C. G. Stueckelberg, “Interaction forces in electrodynamics and in the field theory of nuclear forces,” Helv. Phys. Acta **11** (1938) 299.
- [71] R. Percacci, “Gravity from a Particle Physicists’ perspective,” PoS ISFTG (2009) 011 [arXiv:0910.5167 [hep-th]].
- [72] R. Percacci, “The Higgs phenomenon in quantum gravity,” Nucl. Phys. B **353** (1991) 271 [arXiv:0712.3545 [hep-th]].
- [73] A. D. I. Latorre, G. J. Olmo and M. Ronco, “Observable traces of non-metricity: new constraints on metric-affine gravity,” Phys. Lett. B **780** (2018) 294 [arXiv:1709.04249 [hep-th]].
- [74] I. P. Lobo and C. Romero, “Experimental constraints on the second clock effect,” Phys. Lett. B **783** (2018) 306 [arXiv:1807.07188 [gr-qc]].
- [75] K. N. Abazajian *et al.* [CMB-S4 Collaboration], “CMB-S4 Science Book, First Edition,” arXiv:1610.02743 [astro-ph.CO]. <https://cmb-s4.org/>
- [76] J. Errard, S. M. Feeney, H. V. Peiris and A. H. Jaffe, “Robust forecasts on fundamental physics from the foreground-obscured, gravitationally-lensed CMB polarization,” JCAP **1603** (2016) no.03, 052 [arXiv:1509.06770 [astro-ph.CO]].
- [77] A. Suzuki *et al.*, “The LiteBIRD Satellite Mission - Sub-Kelvin Instrument,” J. Low. Temp. Phys. **193** (2018) no.5-6, 1048 [arXiv:1801.06987 [astro-ph.IM]].
- [78] For an early study of this theory, see F. Englert, E. Gunzig, C. Truffin and P. Windey, “Conformal Invariant General Relativity with Dynamical Symmetry Breakdown,” Phys. Lett. **57B** (1975) 73.

- [79] P. W. Higgs, “Quadratic lagrangians and general relativity,” *Nuovo Cim.* **11** (1959) no.6, 816-820
- [80] D. Gorbunov, V. Rubakov, “Introduction to the theory of the early Universe”, World Scientific, 2011.
- [81] A. Edery and Y. Nakayama, “Palatini formulation of pure R^2 gravity yields Einstein gravity with no massless scalar,” *Phys. Rev. D* **99** (2019) no.12, 124018 [arXiv:1902.07876 [hep-th]].
- [82] R. Jackiw and S. Y. Pi, “Fake Conformal Symmetry in Conformal Cosmological Models,” *Phys. Rev. D* **91** (2015) no.6, 067501 [arXiv:1407.8545 [gr-qc]].
- [83] R. Jackiw and S. Y. Pi, “New Setting for Spontaneous Gauge Symmetry Breaking?,” *Fundam. Theor. Phys.* **183** (2016) 159 [arXiv:1511.00994 [hep-th]].
- [84] G. ’t Hooft, “Local Conformal Symmetry: the Missing Symmetry Component for Space and Time,” arXiv:1410.6675 [gr-qc]. (Essay written for the Gravity Research Foundation - 2015 Awards for Essays on Gravitation).
- [85] C. Kounnas, D. Lüst and N. Toubas, “ R^2 inflation from scale invariant supergravity and anomaly free superstrings with fluxes,” *Fortsch. Phys.* **63** (2015) 12 [arXiv:1409.7076 [hep-th]].
- [86] L. Alvarez-Gaume, A. Kehagias, C. Kounnas, D. Lüst and A. Riotto, “Aspects of Quadratic Gravity,” *Fortsch. Phys.* **64** (2016) no.2-3, 176 [arXiv:1505.07657 [hep-th]].
- [87] D. N. Vollick, “Modified Palatini action that gives the Einstein-Maxwell theory,” *Phys. Rev. D* **93** (2016) no.4, 044061 [arXiv:1612.05829 [gr-qc]].
- [88] K. i. Kobayashi and T. Uematsu, “Nonlinear Realization of Superconformal Symmetry,” *Nucl. Phys. B* **263** (1986) 309. doi:10.1016/0550-3213(86)90119-7
- [89] J. Beltrán Jiménez and A. Delhom, “Ghosts in metric-affine higher order curvature gravity,” *Eur. Phys. J. C* **79** (2019) no.8, 656 [arXiv:1901.08988 [gr-qc]].
- [90] D. M. Ghilencea, “Gauging scale symmetry and inflation: Weyl versus Palatini gravity,” [arXiv:2007.14733 [hep-th]].
- [91] F. Bauer and D. A. Demir, “Higgs-Palatini Inflation and Unitarity,” *Phys. Lett. B* **698** (2011) 425 [arXiv:1012.2900 [hep-ph]].
- [92] F. Bauer and D. A. Demir, “Inflation with Non-Minimal Coupling: Metric versus Palatini Formulations,” *Phys. Lett. B* **665** (2008) 222 [arXiv:0803.2664 [hep-ph]].
- [93] T. Koivisto and H. Kurki-Suonio, “Cosmological perturbations in the palatini formulation of modified gravity,” *Class. Quant. Grav.* **23** (2006) 2355 [astro-ph/0509422].
- [94] S. Rasanen and P. Wahlman, “Higgs inflation with loop corrections in the Palatini formulation,” *JCAP* **1711** (2017) 047 [arXiv:1709.07853 [astro-ph.CO]].
- [95] V. M. Enckell, K. Enqvist, S. Rasanen and E. Tomberg, “Higgs inflation at the hilltop,” *JCAP* **1806** (2018) 005 [arXiv:1802.09299 [astro-ph.CO]].
- [96] T. Markkanen, T. Tenkanen, V. Vaskonen and H. Veermäe, “Quantum corrections to quartic inflation with a non-minimal coupling: metric vs. Palatini,” *JCAP* **1803** (2018) 029 [arXiv:1712.04874 [gr-qc]].

- [97] L. Järv, A. Racioppi and T. Tenkanen, “Palatini side of inflationary attractors,” *Phys. Rev. D* **97** (2018) no.8, 083513 [arXiv:1712.08471 [gr-qc]].
- [98] I. Antoniadis, A. Karam, A. Lykkas and K. Tamvakis, “Palatini inflation in models with an R^2 term,” *JCAP* **1811** (2018) 028 [arXiv:1810.10418 [gr-qc]].
- [99] V. M. Enckell, K. Enqvist, S. Rasanen and L. P. Wahlman, “Inflation with R^2 term in the Palatini formalism,” *JCAP* **1902** (2019) 022 [arXiv:1810.05536 [gr-qc]].
- [100] I. D. Gialamas and A. Lahanas, “Reheating in R^2 Palatini inflationary models,” *Phys. Rev. D* **101** (2020) no.8, 084007 [arXiv:1911.11513 [gr-qc]].
- [101] Y. Akrami *et al.* [Planck Collaboration], “Planck 2018 results. X. Constraints on inflation,” arXiv:1807.06211 [astro-ph.CO].
- [102] A. A. Starobinsky “A New Type of Isotropic Cosmological Models Without Singularity,” *Phys. Lett. B* **91** (1980) 99 [*Phys. Lett.* **91B** (1980) 99] [*Adv. Ser. Astrophys. Cosmol.* **3** (1987) 130].
- [103] C. Patrignani *et al.* [Particle Data Group], “Review of Particle Physics,” *Chin. Phys. C* **40** (2016) no.10, 100001.
- [104] J. Edholm, “UV completion of the Starobinsky model, tensor-to-scalar ratio, and constraints on nonlocality,” *Phys. Rev. D* **95** (2017) no.4, 044004 [arXiv:1611.05062 [gr-qc]] and references therein.
- [105] R. Kallosh, A. D. Linde, D. A. Linde and L. Susskind, “Gravity and global symmetries,” *Phys. Rev. D* **52** (1995) 912 [hep-th/9502069].