

FIBONACCI REPRESENTATIONS OF SEQUENCES IN HILBERT SPACES

J. SEDGHI MOGHADDAM, A. NAJATI AND Y. KHEDMATI

ABSTRACT. Dynamical sampling deals with frames of the form $\{T^n\varphi\}_{n=0}^\infty$, where $T \in B(\mathcal{H})$ belongs to certain classes of linear operators and $\varphi \in \mathcal{H}$. The purpose of this paper is to investigate a new representation, namely, Fibonacci representation of sequences $\{f_n\}_{n=1}^\infty$ in a Hilbert space \mathcal{H} ; having the form $f_{n+2} = T(f_n + f_{n+1})$ for all $n \geq 1$ and a linear operator $T : \text{span}\{f_n\}_{n=1}^\infty \rightarrow \text{span}\{f_n\}_{n=1}^\infty$. We apply this kind of representations for complete sequences and frames. Finally, we present some properties of Fibonacci representation operators.

1. INTRODUCTION

We recall some definitions and standard results from frame theory.

Definition 1.1. Consider a sequence $F = \{f_i\}_{i=1}^\infty$ in \mathcal{H} .

(i) F is called a *frame* for \mathcal{H} , if there exist two constants $A_F, B_F > 0$ such that

$$A_F \|f\|^2 \leq \sum_{i=1}^{\infty} |\langle f, f_i \rangle|^2 \leq B_F \|f\|^2, \quad f \in \mathcal{H}.$$

(ii) F is called a *Bessel sequence* with Bessel bound B_F if at least the upper frame condition holds.

(iii) F is called *complete* in \mathcal{H} if $\overline{\text{span}}\{f_i\}_{i=1}^\infty = \mathcal{H}$, i.e., $\text{span}\{f_i\}_{i=1}^\infty$ is dense in \mathcal{H} .

(iv) F is called *linearly independent* if $\sum_{k=1}^m c_k f_k = 0$ for some scalar coefficients $\{c_k\}_{k=1}^m$, then $c_k = 0$ for all $k = 1, \dots, m$. We say F is *linearly dependent* if F is not linearly independent.

Theorem 1.2. [4, Theorem 5.5.1] *A sequence $F = \{f_i\}_{i=1}^\infty \subseteq \mathcal{H}$ is a frame for \mathcal{H} if and only if*

$$T_F : \ell^2 \rightarrow \mathcal{H}, \quad T_F(\{c_i\}_{i=1}^\infty) = \sum_{i=1}^{\infty} c_i f_i,$$

is a well-defined mapping from ℓ^2 onto \mathcal{H} . Moreover, the adjoint of T_F is given by

$$T_F^* : \mathcal{H} \rightarrow \ell^2, \quad T_F^* f = \{\langle f, f_i \rangle\}_{i=1}^\infty.$$

It is well known that if $F = \{f_i\}_{i=1}^\infty \subseteq \mathcal{H}$ is a Bessel sequence for \mathcal{H} , then T_F is well-defined and bounded.

In [2], Aldroubi et al. introduced dynamic sampling which it deals with frame properties of sequences of the form $\{T^n\varphi\}_{n=0}^\infty$, where $T \in B(\mathcal{H})$ belongs to certain classes of linear operators (such as diagonalizable normal operators) and $\varphi \in \mathcal{H}$. Various characterizations of frames having the form $\{f_k\}_{k \in I} = \{T^k\varphi\}_{k \in I}$, where T is a linear (not necessarily bounded) operator can be found in [5, 6, 7, 8].

Proposition 1.3. [6, Proposition 2.3] *Consider a frame sequence $F = \{f_i\}_{i=1}^\infty$ in a Hilbert space \mathcal{H} which spans an infinite-dimensional subspace. The following are equivalent:*

2010 *Mathematics Subject Classification.* 42C15, 47B40.

Key words and phrases. Frame, Operator representation, Fibonacci representation, Basis, Perturbation.

- (i) F is linearly independent.
- (ii) There exists a linear operator $T : \text{span}\{f_i\}_{i=1}^\infty \rightarrow \mathcal{H}$ such that $Tf_i := f_{i+1}$.

Theorem 1.4. [7, Theorem 2.1] *Consider a frame $F = \{f_i\}_{i=1}^\infty$ in \mathcal{H} . Then the following are equivalent:*

- (i) $F = \{T^{i-1}f_1\}_{i=1}^\infty$ for some $T \in B(\mathcal{H})$.
- (ii) For some dual frame $G = \{g_i\}_{i=1}^\infty$ (and hence all)

$$f_{j+1} = \sum_{i=1}^{\infty} \langle f_j, g_i \rangle f_{i+1}, \forall j \in \mathbb{N}.$$

- (iii) The $\ker T_F$ is invariant under the right-shift operator $\mathcal{T} : \ell^2 \rightarrow \ell^2$ defined by

$$\mathcal{T}(c_1, c_2, \dots) = (0, c_1, c_2, \dots).$$

2. SPECIAL SEQUENCES

It is well known, cf. [4, Example 5.4.6] that if $\{e_n\}_{n=1}^\infty$ is an orthonormal basis $\{e_n\}_{n=1}^\infty$ for \mathcal{H} , then $\{e_n + e_{n+1}\}_{n=1}^\infty$ is complete and a Bessel sequence but not a frame. This motivates us to investigate some results concerning the sequences $F = \{f_n\}_{n=1}^\infty$, $M = \{f_n + f_{n+1}\}_{n=1}^\infty$ and $N = \{f_n - f_{n+1}\}_{n=1}^\infty$ in a Hilbert space \mathcal{H} .

Proposition 2.1. *Let α and β be nonzero scalars and $F = \{f_n\}_{n=1}^\infty \subseteq \mathcal{H}$. Then*

- (i) F is a Bessel sequence for \mathcal{H} , if and only if $M = \{\alpha f_n + \beta f_{n+1}\}_{n=1}^\infty$ and $N = \{\alpha f_n - \beta f_{n+1}\}_{n=1}^\infty$ are Bessel sequences for \mathcal{H} .
- (ii) Suppose that F is a Bessel sequence for \mathcal{H} . Then F is complete, if and only if $M = \{\alpha f_n + \beta f_{n+1}\}_{n=1}^\infty$ is complete, whenever $|\alpha| \geq |\beta|$.

Proof. (i) Assume that $\{f_n\}_{n=1}^\infty$ is a Bessel sequence with Bessel bound B_F and $\mu = \max\{|\alpha|^2, |\beta|^2\}$. Then for $f \in \mathcal{H}$, we have

$$\begin{aligned} & \sum_{n=1}^{\infty} |\langle f, \alpha f_n + \beta f_{n+1} \rangle|^2 + \sum_{n=1}^{\infty} |\langle f, \alpha f_n - \beta f_{n+1} \rangle|^2 \\ &= 2|\alpha|^2 \sum_{n=1}^{\infty} |\langle f, f_n \rangle|^2 + 2|\beta|^2 \sum_{n=1}^{\infty} |\langle f, f_{n+1} \rangle|^2 \\ &\leq 4\mu \sum_{n=1}^{\infty} |\langle f, f_n \rangle|^2 \leq 4\mu B_F \|f\|^2. \end{aligned}$$

Then M and N are Bessel sequences. For the opposite implication, let B_M and B_N be Bessel bounds for sequences M and N , respectively. Then

$$\begin{aligned} 2|\alpha|^2 \sum_{n=1}^{\infty} |\langle f, f_n \rangle|^2 &\leq 2 \sum_{n=1}^{\infty} |\langle f, \alpha f_n \rangle|^2 + 2 \sum_{n=1}^{\infty} |\langle f, \beta f_{n+1} \rangle|^2 \\ &= \sum_{n=1}^{\infty} |\langle f, \alpha f_n + \beta f_{n+1} \rangle|^2 + \sum_{n=1}^{\infty} |\langle f, \alpha f_n - \beta f_{n+1} \rangle|^2 \\ &\leq (B_M + B_N) \|f\|^2, \quad f \in \mathcal{H}. \end{aligned}$$

- (ii) Suppose that F is complete and $f \in \mathcal{H}$ such that $\langle f, \alpha f_n + \beta f_{n+1} \rangle = 0$ for all $n \in \mathbb{N}$. Then $\overline{\alpha} \langle f, f_n \rangle = -\overline{\beta} \langle f, f_{n+1} \rangle$ for all $n \in \mathbb{N}$. Since $|\alpha| \geq |\beta|$ and

$$|\langle f, f_1 \rangle|^2 \sum_{n=0}^{\infty} \left| \frac{\alpha}{\beta} \right|^n = \sum_{n=1}^{\infty} |\langle f, f_n \rangle|^2 \leq B_F \|f\|^2,$$

we get $\langle f, f_1 \rangle = 0$ and consequently $\langle f, f_n \rangle = 0$ for $n \in \mathbb{N}$. Hence $f = 0$ and this shows that $\{\alpha f_n + \beta f_{n+1}\}_{n=1}^{\infty}$ is complete. In order to show the other implication, assume that M is complete and $f \in \mathcal{H}$ such that $\langle f, f_n \rangle = 0$ for all $n \in \mathbb{N}$. Since

$$\langle f, \alpha f_n + \beta f_{n+1} \rangle = \overline{\alpha} \langle f, f_n \rangle + \overline{\beta} \langle f, f_{n+1} \rangle = 0, \quad n \in \mathbb{N},$$

we conclude that $f = 0$ and therefore F is complete. \square

Proposition 2.2. *Let $F = \{f_n\}_{n=1}^{\infty}$, $M = \{\alpha f_n + \beta f_{n+1}\}_{n=1}^{\infty}$ and $N = \{\alpha f_n - \beta f_{n+1}\}_{n=1}^{\infty}$ be sequences in a Hilbert space \mathcal{H} and $\alpha \neq 0$. Then F is a frame for \mathcal{H} , if and only if $M \cup N$ is a frame for \mathcal{H} .*

Proof. Let $\mu = \max\{|\alpha|^2, |\beta|^2\}$. Then the result follows from

$$\begin{aligned} |\alpha|^2 \sum_{n=1}^{\infty} |\langle f, f_n \rangle|^2 &\leq 2 \sum_{n=1}^{\infty} |\langle f, \alpha f_n \rangle|^2 + 2 \sum_{n=1}^{\infty} |\langle f, \beta f_{n+1} \rangle|^2 \\ &= \sum_{n=1}^{\infty} |\langle f, \alpha f_n + \beta f_{n+1} \rangle|^2 + \sum_{n=1}^{\infty} |\langle f, \alpha f_n - \beta f_{n+1} \rangle|^2 \\ &= 2|\alpha|^2 \sum_{n=1}^{\infty} |\langle f, f_n \rangle|^2 + 2|\beta|^2 \sum_{n=1}^{\infty} |\langle f, f_{n+1} \rangle|^2 \\ &\leq 4\mu \sum_{n=1}^{\infty} |\langle f, f_n \rangle|^2, \quad f \in \mathcal{H}. \end{aligned}$$

\square

Theorem 2.3. *Let $M = \{f_n + f_{n+1}\}_{n=1}^{\infty}$ and $N = \{f_n - f_{n+1}\}_{n=1}^{\infty}$ be frames for \mathcal{H} . Then $F = \{f_n\}_{n=1}^{\infty}$ is a frame for \mathcal{H} and*

$$(2.1) \quad 4S_F f = S_M f + S_N f + 2\langle f, f_1 \rangle f_1, \quad f \in \mathcal{H},$$

where S_F, S_M and S_N are frame operators for F, M and N , respectively.

Proof. By Proposition 2.1, F is a Bessel sequence for \mathcal{H} . Let A_M and A_N be lower frame bounds for M and N , respectively. Then we have

$$\begin{aligned} (A_M + A_N) \|f\|^2 &\leq \sum_{n=1}^{\infty} |\langle f, f_n + f_{n+1} \rangle|^2 + \sum_{n=1}^{\infty} |\langle f, f_n - f_{n+1} \rangle|^2 \\ &= 2 \sum_{n=1}^{\infty} |\langle f, f_n \rangle|^2 + 2 \sum_{n=1}^{\infty} |\langle f, f_{n+1} \rangle|^2 \\ &\leq 4 \sum_{n=1}^{\infty} |\langle f, f_n \rangle|^2, \quad f \in \mathcal{H}. \end{aligned}$$

Therefore, F is a frame for \mathcal{H} . Furthermore, since

$$\begin{aligned} & \sum_{n=1}^{\infty} |\langle f, f_n + f_{n+1} \rangle|^2 + \sum_{n=1}^{\infty} |\langle f, f_n - f_{n+1} \rangle|^2 \\ &= 2 \sum_{n=1}^{\infty} |\langle f, f_n \rangle|^2 + 2 \sum_{n=1}^{\infty} |\langle f, f_{n+1} \rangle|^2, \quad f \in \mathcal{H}, \end{aligned}$$

we obtain

$$\begin{aligned} \langle S_M f, f \rangle + \langle S_N f, f \rangle &= 4 \langle S_F f, f \rangle - 2 |\langle f, f_1 \rangle|^2 \\ &= 4 \langle S_F f, f \rangle - 2 \langle \langle f, f_1 \rangle f_1, f \rangle, \quad f \in \mathcal{H}. \end{aligned}$$

Then we obtain (2.1). \square

Theorem 2.4. Let $M = \{f_n + f_{n+1}\}_{n=1}^{\infty}$, $N = \{f_n - f_{n+1}\}_{n=1}^{\infty}$ and $F = \{f_n\}_{n=1}^{\infty}$ be Bessel sequences for \mathcal{H} and $\ker T_F$ be invariant under \mathcal{T}_R . Then $\ker T_F = \ker T_M \cap \ker T_N$.

Proof. Let $\{c_n\}_{n=1}^{\infty} \in \ker T_F$. Since $\ker T_F$ is invariant under \mathcal{T}_R , we get $\{0, c_1, c_2, \dots\} \in \ker T_F$. Therefore $\{c_1, c_1 + c_2, c_2 + c_3, \dots\}, \{c_1, c_2 - c_1, c_3 - c_2, \dots\} \in \ker T_F$. Hence

$$\begin{aligned} \sum_{n=1}^{\infty} c_n (f_n + f_{n+1}) &= \sum_{n=1}^{\infty} (c_n + c_{n+1}) f_{n+1} + c_1 f_1 = 0, \\ \sum_{n=1}^{\infty} c_n (f_n - f_{n+1}) &= \sum_{n=1}^{\infty} (c_{n+1} - c_n) f_{n+1} + c_1 f_1 = 0. \end{aligned}$$

Then we conclude $\{c_n\}_{n=1}^{\infty} \in \ker T_M \cap \ker T_N$. On the other hand, if $\{c_n\}_{n=1}^{\infty} \in \ker T_M \cap \ker T_N$, then we have

$$0 = \sum_{n=1}^{\infty} c_n (f_n - f_{n+1}) + \sum_{n=1}^{\infty} c_n (f_n + f_{n+1}) = 2 \sum_{n=1}^{\infty} c_n f_n.$$

Therefore, $\{c_n\}_{n=1}^{\infty} \in \ker T_F$. \square

3. FIBONACCI REPRESENTATION

In this section we want to consider representation of a sequence $\{f_n\}_{n=1}^{\infty} \subseteq \mathcal{H}$ on the form $f_n = T(f_{n-1} + f_{n-2})$ for $n \geq 3$, where T is a linear operator defined on an appropriate subspace of \mathcal{H} .

Definition 3.1. We say that a sequence $F = \{f_n\}_{n=1}^{\infty}$ has a *Fibonacci representation* if there is a linear operator $T : \text{span}\{f_n\}_{n=1}^{\infty} \rightarrow \text{span}\{f_n\}_{n=1}^{\infty}$ such that $f_n = T(f_{n-1} + f_{n-2})$ for $n \geq 3$. In the affirmative case, we say that F is represented by T , and T is called a *Fibonacci representation operator* with respect to F .

Throughout this segment, \mathcal{H} denotes a Hilbert space and $\{e_n\}_{n=1}^{\infty}$ is an orthonormal basis for \mathcal{H} .

Example 3.2. It is clear that $F = \{f_n\}_{n=1}^{\infty} = \{e_1, e_1, e_2, \dots\}$ is a frame for \mathcal{H} . We define the linear operator $T : \text{span}\{e_n\}_{n=1}^{\infty} \rightarrow \text{span}\{e_n\}_{n=1}^{\infty}$ by

$$T e_1 = \frac{e_2}{2}, \quad T e_n = \sum_{i=0}^{n-2} (-1)^i e_{n-i+1} + (-1)^{n+1} \frac{e_2}{2}, \quad n \geq 2.$$

Then F is represented by T . Indeed, $Te_1 + Te_2 = e_3$ and for $n \geq 4$ we have

$$\begin{aligned}
 Te_{n-2} + Te_{n-1} &= \sum_{i=0}^{n-4} (-1)^i e_{n-i-1} + (-1)^{n-1} \frac{e_2}{2} + \sum_{i=0}^{n-3} (-1)^i e_{n-i} + (-1)^n \frac{e_2}{2} \\
 &= \sum_{i=0}^{n-4} (-1)^i e_{n-i-1} + e_n + \sum_{i=1}^{n-3} (-1)^i e_{n-i} \\
 &= \sum_{i=0}^{n-4} (-1)^i e_{n-i-1} + \sum_{i=0}^{n-4} (-1)^{i+1} e_{n-i-1} + e_n \\
 &= e_n.
 \end{aligned}$$

Note that F is not linearly independent, and so by [6, Proposition 2.3], there does not exist a linear operator $S : \text{span}\{e_n\}_{n=1}^\infty \rightarrow \text{span}\{e_n\}_{n=1}^\infty$ such that $Se_1 = e_1$ and $Se_{n-1} = e_n$, $n \geq 2$.

Example 3.3. The frame $F = \{f_n\}_{n=1}^\infty = \{e_1, e_2, e_3, e_1, e_4, e_5, e_6, \dots\}$ is represented by T , where $T : \text{span}\{f_n\}_{n=1}^\infty \rightarrow \text{span}\{f_n\}_{n=1}^\infty$ is defined by

$$\begin{aligned}
 Te_1 &= \frac{1}{2}(e_4 + e_3 - e_1), & Te_2 &= \frac{1}{2}(-e_4 + e_3 + e_1), & Te_3 &= \frac{1}{2}(e_4 - e_3 + e_1), \\
 Te_4 &= e_5 - Te_1, & Te_n &= e_{n+1} - Te_{n-1}, & n &\geq 5.
 \end{aligned}$$

Proposition 3.4. *A sequence $F = \{f_n\}_{n=1}^\infty$ is represented by T , if and only if $M = \{f_n + f_{n+1}\}_{n=1}^\infty$ and $N = \{f_n - f_{n+1}\}_{n=1}^\infty$ are represented by T .*

Proof. First, let F be represented by T . For every $n \in \mathbb{N}$, we have

$$\begin{aligned}
 T((f_n + f_{n+1}) + (f_{n+1} + f_{n+2})) &= T(f_n + f_{n+1}) + T(f_{n+1} + f_{n+2}) \\
 &= f_{n+2} + f_{n+3}, \\
 T((f_n - f_{n+1}) + (f_{n+1} - f_{n+2})) &= T(f_n + f_{n+1}) - T(f_{n+1} + f_{n+2}) \\
 &= f_{n+2} - f_{n+3},
 \end{aligned}$$

Then M and N are represented by T . Conversely, if M and N are represented by T , then

$$\begin{aligned}
 T(f_n + f_{n+1}) &= \frac{1}{2}T(f_n + f_{n+1} + f_n - f_{n+1} + f_{n+1} + f_{n+2} + f_{n+1} - f_{n+2}) \\
 &= \frac{1}{2}(T(f_n + f_{n+1} + f_{n+1} + f_{n+2}) + T(f_n - f_{n+1} + f_{n+1} - f_{n+2})) \\
 &= \frac{1}{2}(f_{n+2} + f_{n+3} + f_{n+2} - f_{n+3}) = f_{n+2}, \quad n \in \mathbb{N}.
 \end{aligned}$$

Hence F is represented by T . □

A frame may have more than one Fibonacci representation and a frame may not have any.

Example 3.5. The frame $G = \{f_n\}_{n=1}^\infty = \{e_1, e_2, e_1, e_3, e_4, \dots\}$ does not have any Fibonacci representations. Indeed, if G is represented by T , then

$$Te_1 + Te_2 = e_1, \quad Te_2 + Te_1 = e_3,$$

which is a contradiction. We note that $\{f_n + f_{n+1}\}_{n=1}^\infty$ is not linearly independent.

Example 3.6. Consider the frame $E = \{e_n\}_{n=1}^\infty \subseteq \mathcal{H}$ and let $T, S : \text{span}\{e_n\}_{n=1}^\infty \rightarrow \text{span}\{e_n\}_{n=1}^\infty$ be linear operators defined by

$$Te_1 = Te_2 = \frac{1}{2}e_3, \quad Te_n = \frac{(-1)^n}{n}e_3 - \sum_{i=4}^{n+1} (-1)^{n+i}e_i, \quad n \geq 4,$$

$$Se_1 = 0, \quad Se_2 = e_3, \quad Se_3 = e_4 - e_3, \quad Se_n = e_3 - \sum_{i=4}^{n+1} (-1)^{n+i}e_i, \quad n \geq 4.$$

Then it is easy to see that E is represented by T and S . We note that $\{e_n + e_{n+1}\}_{n=1}^\infty$ is linearly independent.

In general if $\{f_n\}_{n=1}^\infty \subseteq \mathcal{H}$ is linearly independent with a Fibonacci representation T , then for each $g \in \text{span}\{f_n\}_{n=1}^\infty$ the linear operator $S : \text{span}\{f_n\}_{n=1}^\infty \rightarrow \text{span}\{f_n\}_{n=1}^\infty$ defined by

$$S\left(\sum_{i=1}^k c_i f_i\right) = \sum_{i=1}^k c_i T f_i + \sum_{i=1}^k (-1)^i c_i g$$

is a Fibonacci representation for $\{f_n\}_{n=1}^\infty$.

Now, we want to get a sufficient condition for a frame $F = \{f_n\}_{n=1}^\infty$ to have a Fibonacci representation. We need the following lemma.

Lemma 3.7. *Consider a sequence $\{f_n\}_{n=1}^\infty$ in \mathcal{H} . Then the following hold:*

(i) *For $n \geq 2$, we have*

$$f_n = \sum_{i=0}^{m-1} (-1)^i (f_{n-i-1} + f_{n-i}) + (-1)^m f_{n-m}, \quad 1 \leq m \leq n-1.$$

(ii) $\text{span}\{f_n\}_{n=1}^\infty = \text{span}\{\{f_1\} \cup \{f_n + f_{n+1}\}_{n=1}^\infty\}$.

(iii) *If $\{f_n\}_{n=1}^\infty$ is linearly independent, then $\{f_n + f_{n+1}\}_{n=1}^\infty$ is linearly independent.*

(iv) *If $\{f_1\} \cup \{f_n + f_{n+1}\}_{n=1}^\infty$ is linearly independent, then $\{f_n\}_{n=1}^\infty$ is linearly independent.*

Proof. (i) Let $n \geq 2$ and $1 \leq m \leq n-1$. Then we have

$$\begin{aligned} \sum_{i=0}^{m-1} (-1)^i (f_{n-i-1} + f_{n-i}) &= \sum_{i=0}^{m-1} (-1)^i f_{n-i-1} + \sum_{i=0}^{m-1} (-1)^i f_{n-i} \\ &= \sum_{i=1}^m (-1)^{i-1} f_{n-i} + \sum_{i=0}^{m-1} (-1)^i f_{n-i} \\ &= \sum_{i=1}^{m-1} (-1)^{i-1} f_{n-i} + (-1)^{m-1} f_{n-m} + f_n + \sum_{i=1}^{m-1} (-1)^i f_{n-i} \\ &= (-1)^{m-1} f_{n-m} + f_n. \end{aligned}$$

For the proof of (ii), it is clear that $\text{span}\{\{f_n + f_{n+1}\}_{n=1}^\infty \cup \{f_1\}\} \subseteq \text{span}\{f_n\}_{n=1}^\infty$. On the other hand, by (i) (for $m = n-1$) we infer $\text{span}\{f_n\}_{n=1}^\infty \subseteq \text{span}\{\{f_n + f_{n+1}\}_{n=1}^\infty \cup \{f_1\}\}$. This proves (ii). To prove (iii), let $\{c_n\}_{n=1}^k \subseteq \mathbb{C}$ such that $\sum_{n=1}^k c_n (f_n + f_{n+1}) = 0$. Then we have

$$c_1 f_1 + \sum_{n=2}^k (c_{n-1} + c_n) f_n + c_k f_{k+1} = 0.$$

Since $\{f_n\}_{n=1}^\infty$ is linearly independent, we get $c_1 = c_k = 0$ and $c_{n-1} + c_n = 0$ for all $2 \leq n \leq k$. Therefore, $c_n = 0$ for all $1 \leq n \leq k$. This completes the proof of (iii).

To prove (iv), let $\{c_n\}_{n=1}^N \subseteq \mathbb{C}$ such that $\sum_{n=1}^N c_n f_n = 0$. Then by (i), we have

$$\begin{aligned}
0 &= \sum_{n=1}^N c_n f_n = c_1 f_1 + \sum_{n=2}^N c_n f_n \\
&= c_1 f_1 + \sum_{n=2}^N c_n \left(\sum_{i=0}^{n-2} (-1)^i (f_{n-i-1} + f_{n-i}) + (-1)^{n-1} f_1 \right) \\
&= \left(c_1 + \sum_{n=2}^N c_n (-1)^{n-1} \right) f_1 + \sum_{n=2}^N c_n \sum_{i=0}^{n-2} (-1)^i (f_{n-i-1} + f_{n-i}) \\
&= \left(c_1 + \sum_{n=2}^N c_n (-1)^{n-1} \right) f_1 + \sum_{i=0}^{N-2} \sum_{n=i+2}^N c_n (-1)^i (f_{n-i-1} + f_{n-i}) \\
&= \left(c_1 + \sum_{n=2}^N c_n (-1)^{n-1} \right) f_1 + \sum_{i=2}^N \sum_{n=i}^N c_n (-1)^{i-2} (f_{n-i+1} + f_{n-i+2}) \\
&= \left(c_1 + \sum_{n=2}^N c_n (-1)^{n-1} \right) f_1 + \sum_{i=2}^N \sum_{n=0}^{N-i} c_{i+n} (-1)^i (f_{n+1} + f_{n+2}) \\
&= \left(c_1 + \sum_{n=2}^N c_n (-1)^{n-1} \right) f_1 + \sum_{n=0}^{N-2} \left(\sum_{i=2}^{N-n} c_{i+n} (-1)^i \right) (f_{n+1} + f_{n+2}).
\end{aligned}$$

Since $\{f_1\} \cup \{f_n + f_{n+1}\}_{n=1}^\infty$ is linearly independent, we get

$$c_1 + \sum_{k=2}^N c_k (-1)^{k-1} = 0, \quad \sum_{i=2}^{N-n} c_{i+n} (-1)^i = 0, \quad 0 \leq n \leq N-2.$$

Hence we conclude that $c_n = 0$ for all $n = 1, 2, \dots, N$. Then $\{f_n\}_{n=1}^\infty$ is linearly independent. \square

In the following, we give a sufficient condition for a sequence $F = \{f_n\}_{n=1}^\infty$ to have a Fibonacci representation.

Theorem 3.8. *Let $F = \{f_n\}_{n=1}^\infty$ be a sequence in \mathcal{H} . If $\{f_n + f_{n+1}\}_{n=1}^\infty$ is linearly independent, then F has a Fibonacci representation.*

Proof. First we assume that $f_1 \in \text{span}\{f_n + f_{n+1}\}_{n=1}^\infty$. Then by (ii) of Lemma 3.7, we have $\text{span}\{f_n + f_{n+1}\}_{n=1}^\infty = \text{span}\{f_n\}_{n=1}^\infty$. We define a linear operator $T : \text{span}\{f_n\}_{n=1}^\infty \rightarrow \text{span}\{f_n\}_{n=1}^\infty$ by

$$(3.1) \quad T(f_n + f_{n+1}) = f_{n+2}; \quad n \geq 2.$$

Since $\{f_n + f_{n+1}\}_{n=1}^\infty$ is linearly independent sequence, T is well-defined and F is represented by T . If $f_1 \notin \text{span}\{f_n + f_{n+1}\}_{n=1}^\infty$, then $\{f_1\} \cup \{f_n + f_{n+1}\}_{n=1}^\infty$ is linearly independent and so by Lemma 3.7 (iv), $\{f_n\}_{n=1}^\infty$ is linearly independent. Hence we can define a linear operator $T : \text{span}\{f_n\}_{n=1}^\infty \rightarrow \text{span}\{f_n\}_{n=1}^\infty$ by

$$Tf_n = \sum_{i=0}^n (-1)^i f_{n+1-i}, \quad n \in \mathbb{N}.$$

We show that F is represented by T . Indeed,

$$\begin{aligned}
 Tf_n + Tf_{n+1} &= \sum_{i=0}^n (-1)^i f_{n+1-i} + \sum_{i=0}^{n+1} (-1)^i f_{n+2-i} \\
 &= \sum_{i=0}^n (-1)^i f_{n+1-i} + \sum_{i=1}^{n+1} (-1)^i f_{n+2-i} + f_{n+2} \\
 &= \sum_{i=0}^n (-1)^i f_{n+1-i} + \sum_{i=0}^n (-1)^{i+1} f_{n+1-i} + f_{n+2} = f_{n+2}.
 \end{aligned}$$

□

The following example shows that the converse of Theorem 3.8 is not satisfied in general.

Example 3.9. The frame $F = \{f_n\}_{n=1}^\infty = \{e_1, e_2, e_3, e_2, e_2, e_4, e_5, e_6, \dots\}$ is represented by the linear operator $T : \text{span}\{f_n\}_{n=1}^\infty \rightarrow \text{span}\{f_n\}_{n=1}^\infty$ given by

$$\begin{aligned}
 Te_1 &= e_3 - \frac{e_4}{2}, \quad Te_2 = \frac{e_4}{2}, \quad Te_3 = e_2 - \frac{e_4}{2}, \\
 Te_n &= \sum_{i=0}^{n-4} (-1)^i e_{n-i+1} + (-1)^{n-3} \frac{e_4}{2}, \quad n \geq 4.
 \end{aligned}$$

But $\{f_n + f_{n+1}\}_{n=1}^\infty = \{e_1 + e_2, e_2 + e_3, e_3 + e_2, 2e_2, \dots\}$ is not linearly independent.

Corollary 3.10. *Let $F = \{f_n\}_{n=1}^\infty$ be a linear independent sequence in \mathcal{H} . Then F has a Fibonacci representation.*

Proof. It follows from Lemma 3.7 (iii) and Theorem 3.8. □

Now, we provide sufficient conditions to make the converse of Theorem 3.8 satisfy.

Theorem 3.11. *Let $F = \{f_n\}_{n=1}^\infty$ be a complete sequence in an infinite dimensional Hilbert space \mathcal{H} which is represented by T . If there exists $m \in \mathbb{N}$ such that $f_{m+1}, Tf_1 \in \text{span}\{f_n\}_{n=1}^m$, then $\{f_n + f_{n+1}\}_{n=1}^\infty$ is linearly independent.*

Proof. Suppose that $\{f_n + f_{n+1}\}_{n=1}^\infty$ is not linearly independent. Then there exists $n_0 \in \mathbb{N}$ such that $f_{n_0} + f_{n_0+1} = \sum_{n=1}^{n_0-1} c_n(f_n + f_{n+1})$. Hence

$$(3.2) \quad f_{n_0+2} = T(f_{n_0} + f_{n_0+1}) = \sum_{n=1}^{n_0-1} c_n f_{n+2} \in \text{span}\{f_n\}_{n=1}^{n_0+1}.$$

Let $V = \text{span}\{f_n\}_{n=1}^l$, where $l = \max\{n_0 + 1, m\}$. By (3.2) and $f_{m+1} \in \text{span}\{f_n\}_{n=1}^m$, we get $f_{l+1} \in V$. We show V is invariant under T . Suppose that $f = \sum_{n=1}^l c_n f_n \in V$. By using (i) of

Lemma 3.7, we have

$$\begin{aligned}
Tf &= T\left(\sum_{n=1}^l c_n f_n\right) = c_1 T f_1 + \sum_{n=2}^l c_n T f_n \\
&= c_1 T f_1 + \sum_{n=2}^l c_n T \left(\sum_{i=0}^n (-1)^i (f_{n-i-1} + f_{n-i}) + (-1)^{n-1} f_1 \right) \\
&= c_1 T f_1 + \sum_{n=2}^l c_n \left(\sum_{i=0}^n (-1)^i T(f_{n-i-1} + f_{n-i}) + (-1)^{n-1} T f_1 \right) \\
&= \left(c_1 + \sum_{n=2}^l c_n (-1)^{n-1} \right) T f_1 + \sum_{n=2}^l c_n \sum_{i=0}^n (-1)^i f_{n-i+1}.
\end{aligned}$$

Since $T f_1 \in \text{span}\{f_n\}_{n=1}^m \subseteq V$ and $f_{l+1} \in V$, the above argument proves that V is invariant under T . Therefore $f_n \in V$ for all $n \geq l+1$ and consequently $\text{span}\{f_n\}_{n=1}^\infty = V$. Since $\{f_n\}_{n=1}^\infty$ is complete in \mathcal{H} , we have $\mathcal{H} = \overline{\text{span}}\{f_n\}_{n=1}^\infty = \overline{V} = V$ which is in contradiction to $\dim \mathcal{H} = \infty$. \square

Proposition 3.12. *Let $\{f_n\}_{n=1}^\infty$ be a complete and linearly dependent sequence with $f_1 \neq 0$ in an infinite dimensional Hilbert space \mathcal{H} . Then there exists $m \geq 2$ such that $f_m \in \text{span}\{f_n\}_{n=1}^{m-1}$ and $f_{m+1} \notin \text{span}\{f_n\}_{n=1}^m$.*

Proof. Since $\{f_n\}_{n=1}^\infty$ is linearly dependent, there exists $k \geq 2$ such that $f_k \in \text{span}\{f_n\}_{n=1}^{k-1}$. We claim that there exists an integer $l > k$ such that $f_l \notin \text{span}\{f_n\}_{n=1}^{l-1}$. If $f_l \in \text{span}\{f_n\}_{n=1}^{l-1}$ for each $l > k$, then $f_{k+1} \in \text{span}\{f_n\}_{n=1}^{k-1}$ because $f_k \in \text{span}\{f_n\}_{n=1}^{k-1}$ and $f_{k+1} \in \text{span}\{f_n\}_{n=1}^k$. Hence by induction we get $f_l \in \text{span}\{f_n\}_{n=1}^{k-1}$ for each $l > k$. Therefore $\text{span}\{f_n\}_{n=1}^\infty = \text{span}\{f_n\}_{n=1}^{k-1}$. Since $\{f_n\}_{n=1}^\infty$ is complete and $\dim \mathcal{H} = \infty$, the contradiction is achieved. Now, let $i \in \mathbb{N}$ be the smallest number such that $f_{k+i} \notin \text{span}\{f_n\}_{n=1}^{k+i-1}$. Putting $m = k + i - 1$, we get $f_m \in \text{span}\{f_n\}_{n=1}^{m-1}$ and $f_{m+1} \notin \text{span}\{f_n\}_{n=1}^m$. \square

Proposition 3.13. *Let $F = \{f_n\}_{n=1}^\infty$ be a sequence in \mathcal{H} which is represented by T . Suppose that $f_m \in \text{span}\{f_n\}_{n=1}^{m-1}$ and $f_{m+1} \notin \text{span}\{f_n\}_{n=1}^m$ for some integer $m \geq 2$. Then $T f_i \in \text{span}\{f_n\}_{n=3}^{m+1}$ for $1 \leq i \leq m$.*

Proof. By the assumption, we have $f_m = \sum_{n=1}^{m-1} c_n f_n$, so

$$\begin{aligned}
& \sum_{n=1}^{m-2} \left(\sum_{i=0}^{n-1} (-1)^i c_{n-i} \right) (f_n + f_{n+1}) \\
&= \sum_{n=1}^{m-2} \left(\sum_{i=0}^{n-1} (-1)^i c_{n-i} \right) f_n + \sum_{n=1}^{m-2} \left(\sum_{i=0}^{n-1} (-1)^i c_{n-i} \right) f_{n+1} \\
&= \sum_{n=1}^{m-2} \left(\sum_{i=0}^{n-1} (-1)^i c_{n-i} \right) f_n + \sum_{n=2}^{m-1} \left(\sum_{i=0}^{n-2} (-1)^i c_{n-i-1} \right) f_n \\
&= c_1 f_1 + \sum_{n=2}^{m-2} \left(\sum_{i=0}^{n-1} (-1)^i c_{n-i} \right) f_n \\
&\quad + \sum_{n=2}^{m-2} \left(\sum_{i=0}^{n-2} (-1)^i c_{n-i-1} \right) f_n + \left(\sum_{i=0}^{m-3} (-1)^i c_{m-i-2} \right) f_{m-1} \\
&= c_1 f_1 + \sum_{n=2}^{m-2} \left(\sum_{i=0}^{n-1} (-1)^i c_{n-i} + \sum_{i=0}^{n-2} (-1)^i c_{n-i-1} \right) f_n \\
&\quad + \left(\sum_{i=0}^{m-3} (-1)^i c_{m-i-2} \right) f_{m-1} \\
&= c_1 f_1 + \sum_{n=2}^{m-2} \left(c_n + \sum_{i=1}^{n-1} (-1)^i c_{n-i} + \sum_{i=1}^{n-1} (-1)^{i-1} c_{n-i} \right) f_n \\
&\quad + \left(\sum_{i=0}^{m-3} (-1)^i c_{m-i-2} \right) f_{m-1} \\
&= c_1 f_1 + \sum_{n=2}^{m-2} c_n f_n + \left(\sum_{i=0}^{m-3} (-1)^i c_{m-i-2} \right) f_{m-1} \\
&= f_m + \left(-c_{m-1} + \sum_{i=0}^{m-3} (-1)^i c_{m-i-2} \right) f_{m-1} \\
&= f_m + \left(\sum_{i=0}^{m-2} (-1)^{i-1} c_{m-i-1} \right) f_{m-1},
\end{aligned}$$

thus

$$(3.3) \quad f_{m-1} + f_m = \sum_{n=1}^{m-2} \left(\sum_{i=0}^{n-1} (-1)^i c_{n-i} \right) (f_n + f_{n+1}) + \left(1 - \sum_{i=0}^{m-2} (-1)^{i-1} c_{m-i-1} \right) f_{m-1}.$$

Since F is represented by T , the equality (3.3) implies that

$$(3.4) \quad f_{m+1} = \sum_{n=1}^{m-2} \left(\sum_{i=0}^{n-1} (-1)^i c_{n-i} \right) f_{n+2} + \left(1 - \sum_{i=0}^{m-2} (-1)^{i-1} c_{m-i-1} \right) T f_{m-1}.$$

If $1 - \sum_{i=0}^{m-2} (-1)^{i-1} c_{m-i-1} = 0$, then $f_{m+1} \in \text{span}\{f_n\}_{n=3}^m \subseteq \text{span}\{f_n\}_{n=1}^{m-1}$ which is a contradiction. Hence (3.4) implies that

$$(3.5) \quad Tf_{m-1} = \frac{f_{m+1} - \sum_{n=1}^{m-2} \left(\sum_{i=0}^{n-1} (-1)^i c_{n-i} \right) f_{n+2}}{1 - \sum_{i=0}^{m-2} (-1)^{i-1} c_{m-i-1}} \in \text{span}\{f_n\}_{n=3}^{m+1}.$$

Also, by (i) of Lemma 3.7, for $1 \leq j \leq m-1$, we have

$$\begin{aligned} f_{m+1} &= Tf_m + Tf_{m-1} = T \left(\sum_{i=0}^{j-1} (-1)^i (f_{m-i-1} + f_{m-i}) + (-1)^j f_{m-j} \right) + Tf_{m-1} \\ &= \sum_{i=0}^{j-1} (-1)^i f_{m-i+1} + (-1)^j Tf_{m-j} + Tf_{m-1}. \end{aligned}$$

Therefore

$$(3.6) \quad Tf_{m-j} = (-1)^j (f_{m+1} - Tf_{m-1} - \sum_{i=0}^{j-1} (-1)^i f_{m-i+1}).$$

Hence it follows from (3.5) and (3.6) that $Tf_i \in \text{span}\{f_n\}_{n=3}^{m+1}$ for each $1 \leq i \leq m-1$. \square

Corollary 3.14. *Let $F = \{f_n\}_{n=1}^\infty$ be a sequence in \mathcal{H} which is represented by T . Suppose that $f_m \in \text{span}\{f_n\}_{n=1}^{m-1}$ and $f_{m+1} \notin \text{span}\{f_n\}_{n=1}^{m-1}$ for some $m \in \mathbb{N}$. Then, $Tf_{m+i} \in \text{span}\{f_n\}_{n=3}^{m+i+1}$ for each $i \in \mathbb{N}$.*

Proof. Since $Tf_{m+i} = f_{m+i+1} - Tf_{m+i-1}$, the result follows by induction on i and Proposition 3.13. \square

Corollary 3.15. *Let $F = \{f_n\}_{n=1}^\infty$ be a complete sequence in an infinite dimensional Hilbert space \mathcal{H} .*

- (i) *If F is linearly independent, then it has a Fibonacci representation T such that $\mathcal{R}(T) = \text{span}\{f_n\}_{n=3}^\infty$.*
- (ii) *If F is linearly dependent, then for any Fibonacci representation T of F we have $\mathcal{R}(T) = \text{span}\{f_n\}_{n=3}^\infty$.*

Proof. First we note that if F is represented by T , then $f_n = T(f_{n-1} + f_{n-2}) \in \mathcal{R}(T)$ for every $n \geq 3$, and consequently $\text{span}\{f_n\}_{n=3}^\infty \subseteq \mathcal{R}(T)$.

To prove (i), consider the linear operator $T : \text{span}\{f_n\}_{n=1}^\infty \rightarrow \text{span}\{f_n\}_{n=1}^\infty$ defined by

$$Tf_1 = Tf_2 = \frac{1}{2}f_3, \quad Tf_n = \sum_{i=0}^{n-3} (-1)^i f_{n+1-i} + \frac{(-1)^n}{2}f_3, \quad n \geq 3.$$

Then

$$\begin{aligned} Tf_1 + Tf_2 &= f_3, \quad Tf_2 + Tf_3 = f_4, \\ Tf_n + Tf_{n+1} &= \sum_{i=0}^{n-3} (-1)^i f_{n-i+1} + \sum_{i=0}^{n-2} (-1)^i f_{n-i+2} = f_{n+2}, \quad n \geq 3. \end{aligned}$$

Hence F is represented by T and it is obvious that $\mathcal{R}(T) \subseteq \text{span}\{f_n\}_{n=3}^\infty$. In order to prove (ii), by Proposition 3.12 there exists $m \geq 2$ such that $f_m \in \text{span}\{f_n\}_{n=1}^{m-1}$ and $f_{m+1} \notin \text{span}\{f_n\}_{n=1}^{m-1}$. If F is represented by T , then by Proposition 3.13 and Corollary 3.14 we have $\mathcal{R}(T) \subseteq \text{span}\{f_n\}_{n=3}^\infty$. \square

In Theorem 3.11, we showed that $\{f_n + f_{n+1}\}_{n=1}^{\infty}$ is linearly independent under some conditions. In the following, we show that (under some conditions) by removing finitely many elements of $\{f_n + f_{n+1}\}_{n=1}^{\infty}$ the remaining elements will be linearly independent.

Theorem 3.16. *Let $F = \{f_n\}_{n=1}^{\infty}$ be a complete sequence in an infinite dimensional Hilbert space \mathcal{H} which is represented by T . Then there exists $m \in \mathbb{N}$ such that $\{f_{m+n} + f_{m+n+1}\}_{n=1}^{\infty}$ is linearly independent.*

Proof. If F is linearly independent, then the result follows by (iii) of Lemma 3.7. Suppose that F is linearly dependent. Then by Proposition 3.12 and Proposition 3.13, there exists $m \geq 2$ such that $f_m \in \text{span}\{f_n\}_{n=1}^{m-1}$, $f_{m+1} \notin \text{span}\{f_n\}_{n=1}^{m-1}$ and $Tf_1 \in \text{span}\{f_n\}_{n=3}^{m+1}$. We prove $\{f_{m+n} + f_{m+n+1}\}_{n=1}^{\infty}$ is linearly independent. Suppose by contradiction that $\{f_{m+n} + f_{m+n+1}\}_{n=1}^{\infty}$ is not linearly independent. Then there exists $j \in \mathbb{N}$ such that $f_{m+j} + f_{m+j+1} = \sum_{n=1}^{j-1} c_n(f_{m+n} + f_{m+n+1})$. Hence we have

$$(3.7) \quad f_{m+j+2} = T(f_{m+j} + f_{m+j+1}) = \sum_{n=1}^{j-1} c_n f_{m+n+2} \in \text{span}\{f_n\}_{n=1}^{m+j+1}.$$

Let $V = \text{span}\{f_n\}_{n=1}^{m+j+1}$. We show that V is invariant under T . Let $f = \sum_{n=1}^{m+j+1} c_n f_n \in V$. Then by (i) of Lemma 3.7, we have

$$\begin{aligned} Tf &= c_1 Tf_1 + \sum_{n=2}^{m+j+1} c_n Tf_n \\ &= c_1 Tf_1 + \sum_{n=2}^{m+j+1} c_n T \left(\sum_{i=0}^{n-2} (-1)^i (f_{n-i-1} + f_{n-i}) + (-1)^{n-1} f_1 \right) \\ &= c_1 Tf_1 + \sum_{n=2}^{m+j+1} c_n \left(\sum_{i=0}^{n-2} (-1)^i f_{n-i+1} + (-1)^{n-1} Tf_1 \right) \\ &= \left(c_1 + \sum_{n=2}^{m+j+1} c_n (-1)^{n-1} \right) Tf_1 + \sum_{n=2}^{j+m+1} c_n \sum_{i=0}^{n-2} (-1)^i f_{n-i+1}. \end{aligned}$$

Using $Tf_1 \in \text{span}\{f_n\}_{n=3}^{m+1} \subseteq V$ and (3.7), we get $Tf \in V$. Then we conclude $f_n \in V$ for all $n \geq m+j+2$. Thus, $\text{span}\{f_n\}_{n=1}^{\infty} = V$ and since $\{f_n\}_{n=1}^{\infty}$ is complete in \mathcal{H} , we have $\mathcal{H} = \overline{\text{span}}\{f_n\}_{n=1}^{\infty} = \overline{V} = V$ which is a contradiction. \square

4. FIBONACCI REPRESENTATION OPERATORS

In a frame that indeed has the form $\{T^n \varphi\}_{n=0}^{\infty}$, where $T \in B(\mathcal{H})$ and $\varphi \in \mathcal{H}$, all sequence members are represented by iterative actions of T on φ . In the case where $\{f_n\}_{n=1}^{\infty}$ has a Fibonacci representation operator T , we expect (Theorem 4.2) all members of the sequence $\{f_n\}_{n=1}^{\infty}$ to be identified in terms of iterative actions of T on elements f_1 and f_2 . In this section, we present some results concerning Fibonacci representation operators. One of the results characterizes types of frame which can be represented in terms of a bounded operator T .

Notation 4.1. $[x]$ denotes the integer part of $x \in \mathbb{R}$ and $\binom{n}{k} := \frac{n!}{k!(n-k)!}$ for integers $0 \leq k \leq n$.

We let $\binom{n}{k} := 0$ when $k > n$ or $k < 0$.

Theorem 4.2. *Let $T : \text{span}\{f_n\}_{n=1}^\infty \rightarrow \text{span}\{f_n\}_{n=1}^\infty$ be a linear operator, then the following statements are equivalent:*

- (i) $F = \{f_n\}_{n=1}^\infty$ is represented by T .
- (ii) $Tf_1 + Tf_2 = f_3$ and

$$(4.1) \quad f_n = \sum_{i=a_n}^{2a_n} \left(\binom{i+b_n}{2i-2a_n+b_n} T^{i+b_n} f_2 + \binom{i+b_n}{2i-2a_n+b_n+1} T^{i+b_n+1} f_1 \right), \quad n \geq 4,$$

where $a_n = \lfloor \frac{n-1}{2} \rfloor$, $b_n = n - 2a_n - 2$.

Proof. (i) \Rightarrow (ii) We prove (4.1) by induction on n . For $n = 4$, we have $a_4 = 1$ and $b_4 = 0$. Then

$$\begin{aligned} & \binom{1}{0} T f_2 + \binom{1}{1} T^2 f_1 + \binom{2}{2} T^2 f_2 + \binom{2}{3} T^3 f_1 \\ &= T f_2 + T(T f_1 + T f_2) = T f_2 + T f_3 = f_4. \end{aligned}$$

Now, assume that $k > 4$ and (4.1) holds for all $n \leq k$ and we prove (4.1) for $n = k+1$. If $k+1$ is even, then $b_k = -1$, $b_{k-1} = b_{k+1} = 0$ and $a_{k+1} = a_k = 1 + a_{k-1}$. Hence

$$\begin{aligned} f_{k+1} &= T f_k + T f_{k-1} \\ &= \sum_{i=a_k}^{2a_k} \left(\binom{i-1}{2i-2a_k-1} T^i f_2 + \binom{i-1}{2i-2a_k} T^{i+1} f_1 \right) \\ &\quad + \sum_{i=a_{k-1}}^{2a_{k-1}} \left(\binom{i}{2i-2a_{k-1}} T^{i+1} f_2 + \binom{i}{2i-2a_{k-1}+1} T^{i+2} f_1 \right) \\ &= \sum_{i=a_k}^{2a_k} \left(\binom{i-1}{2i-2a_k-1} T^i f_2 + \binom{i-1}{2i-2a_k} T^{i+1} f_1 \right) \\ &\quad + \sum_{i=a_k}^{2a_k-1} \left(\binom{i-1}{2i-2a_k} T^i f_2 + \binom{i-1}{2i-2a_k+1} T^{i+1} f_1 \right) \\ &= \sum_{i=a_k}^{2a_k} \left(\binom{i-1}{2i-2a_k-1} T^i f_2 + \binom{i-1}{2i-2a_k} T^{i+1} f_1 \right) \\ &\quad + \sum_{i=a_k}^{2a_k} \left(\binom{i-1}{2i-2a_k} T^i f_2 + \binom{i-1}{2i-2a_k+1} T^{i+1} f_1 \right) \\ &= \sum_{i=a_k}^{2a_k} \left(\binom{i}{2i-2a_k} T^i f_2 + \binom{i}{2i-2a_k+1} T^{i+1} f_1 \right). \end{aligned}$$

Since $b_{k+1} = 0$ and $a_{k+1} = a_k$, we obtain (4.1). If $k+1$ is odd, then $b_k = 0$, $b_{k-1} = b_{k+1} = -1$ and $1 + a_{k-1} = 1 + a_k = a_{k+1}$. Hence

$$\begin{aligned}
f_{k+1} &= Tf_k + Tf_{k-1} \\
&= \sum_{i=a_k}^{2a_k} \left(\binom{i}{2i-2a_k} T^{i+1}f_2 + \binom{i}{2i-2a_k+1} T^{i+2}f_1 \right) \\
&\quad + \sum_{i=a_{k-1}}^{2a_{k-1}} \left(\binom{i-1}{2i-2a_{k-1}-1} T^i f_2 + \binom{i-1}{2i-2a_{k-1}} T^{i+1}f_1 \right) \\
&= \sum_{i=1+a_{k+1}}^{2a_{k+1}} \left(\binom{i-2}{2i-2a_{k+1}-2} T^{i-1}f_2 + \binom{i-2}{2i-2a_{k+1}-1} T^i f_1 \right) \\
&\quad + \sum_{i=a_{k+1}}^{2a_{k+1}-1} \left(\binom{i-2}{2i-2a_{k+1}-1} T^{i-1}f_2 + \binom{i-2}{2i-2a_{k+1}} T^i f_1 \right) \\
&= \sum_{i=a_{k+1}}^{2a_{k+1}} \left(\binom{i-2}{2i-2a_{k+1}-2} T^{i-1}f_2 + \binom{i-2}{2i-2a_{k+1}-1} T^i f_1 \right) \\
&\quad + \sum_{i=a_{k+1}}^{2a_{k+1}} \left(\binom{i-2}{2i-2a_{k+1}-1} T^{i-1}f_2 + \binom{i-2}{2i-2a_{k+1}} T^i f_1 \right) \\
&= \sum_{i=a_{k+1}}^{2a_{k+1}} \left(\binom{i-1}{2i-2a_{k+1}-1} T^{i-1}f_2 + \binom{i-1}{2i-2a_{k+1}} T^i f_1 \right).
\end{aligned}$$

Hence we get (4.1). To prove (ii) \Rightarrow (i), there are two possibilities. If $n > 4$ is odd, then $b_n = -1$, $b_{n-1} = b_{n+1} = 0$ and $a_{n+1} = a_n = 1 + a_{n-1}$. Hence

$$\begin{aligned}
Tf_n &= \sum_{i=a_n}^{2a_n} \left(\binom{i-1}{2i-2a_n-1} T^i f_2 + \binom{i-1}{2i-2a_n} T^{i+1}f_1 \right), \\
Tf_{n-1} &= \sum_{i=a_{n-1}}^{2a_{n-1}} \left(\binom{i}{2i-2a_{n-1}} T^{i+1}f_2 + \binom{i}{2i-2a_{n-1}+1} T^{i+2}f_1 \right) \\
&= \sum_{i=a_n}^{2a_n-1} \left(\binom{i-1}{2i-2a_n} T^i f_2 + \binom{i-1}{2i-2a_n+1} T^{i+1}f_1 \right) \\
&= \sum_{i=a_n}^{2a_n} \left(\binom{i-1}{2i-2a_n} T^i f_2 + \binom{i-1}{2i-2a_n+1} T^{i+1}f_1 \right).
\end{aligned}$$

Therefore,

$$Tf_n + Tf_{n-1} = \sum_{i=a_n}^{2a_n} \left(\binom{i}{2i-2a_n} T^i f_2 + \binom{i}{2i-2a_n+1} T^{i+1}f_1 \right) = f_{n+1}.$$

If $n \geq 4$ is even, the argument is similar to the previous case. \square

Remark 4.3. We recall that

$$\ell^2(\mathcal{H}) := \left\{ \{f_n\}_{n=1}^\infty \subseteq \mathcal{H} : \sum_{n=1}^\infty \|f_n\|^2 < \infty \right\},$$

and $\mathcal{T}_L, \mathcal{T}_R : \ell^2(\mathcal{H}) \rightarrow \ell^2(\mathcal{H})$ are bounded linear operators defined by

$$\mathcal{T}_L\{f_n\}_{n=1}^\infty = \{f_{n+1}\}_{n=1}^\infty, \quad \mathcal{T}_R\{f_n\}_{n=1}^\infty = \{0, f_1, f_2, f_3, \dots\}.$$

Proposition 4.4. *Let $F = \{f_n\}_{n=1}^\infty$ be a Bessel sequence in \mathcal{H} which is represented by T_0 and let $M = \{f_n + f_{n+1}\}_{n=1}^\infty$ be a frame for \mathcal{H} . Then $\ker T_M \subseteq \ker T_{\mathcal{T}_L^2 F}$ if and only if $T := T_0|_{\text{span}\{f_n + f_{n+1}\}_{n=1}^\infty}$ is bounded with $\|T\| \leq \sqrt{\frac{B_F}{A_M}}$, where B_F is a Bessel bound for F and A_M is a lower frame bound for M .*

Proof. Let $\ker T_M \subseteq \ker T_{\mathcal{T}_L^2 F}$ and $f = \sum_{n=1}^k c_n(f_n + f_{n+1})$, where $\{c_n\}_{n=1}^\infty \in \ell^2$ with $c_n = 0$ for $n \geq k+1$. Then

$$\begin{aligned} Tf &= T\left(\sum_{n=1}^k c_n(f_n + f_{n+1})\right) = \sum_{n=1}^k c_n T(f_n + f_{n+1}) \\ &= \sum_{n=1}^\infty c_n f_{n+2} \\ &= \sum_{n=1}^\infty d_n f_{n+2} + \sum_{n=1}^\infty r_n f_{n+2}, \end{aligned}$$

where $\{d_n\}_{n=1}^\infty \in \ker T_M \subseteq \ker T_{\mathcal{T}_L^2 F}$ and $\{r_n\}_{n=1}^\infty \in (\ker T_M)^\perp$. Since $\{d_n\}_{n=1}^\infty \in \ker T_{\mathcal{T}_L^2 F}$, we have $\sum_{n=1}^\infty d_n f_{n+2} = 0$ and consequently $Tf = \sum_{n=1}^\infty r_n f_{n+2}$. Therefore by applying [4, Lemma 5.5.5], we have

$$\begin{aligned} \|Tf\|^2 &= \left\| \sum_{n=1}^\infty r_n f_{n+2} \right\|^2 \leq B_F \sum_{n=1}^\infty |r_n|^2 \leq \frac{B_F}{A_M} \left\| \sum_{n=1}^\infty r_n (f_n + f_{n+1}) \right\|^2 \\ &= \frac{B_F}{A_M} \left\| \sum_{n=1}^\infty r_n (f_n + f_{n+1}) + \sum_{n=1}^\infty d_n (f_n + f_{n+1}) \right\|^2 \\ &= \frac{B_F}{A_M} \left\| \sum_{n=1}^k c_n (f_n + f_{n+1}) \right\|^2 = \frac{B_F}{A_M} \|f\|^2. \end{aligned}$$

For the other implication, let $\{c_n\}_{n=1}^\infty \in \ker T_M$. Since $\sum_{n=1}^\infty c_n (f_n + f_{n+1}) = 0$ and T is bounded, we have

$$0 = T\left(\sum_{n=1}^\infty c_n (f_n + f_{n+1})\right) = \sum_{n=1}^\infty c_n f_{n+2}.$$

This means $\{c_n\}_{n=1}^\infty \in \ker T_{\mathcal{T}_L^2 F}$. □

Remark 4.5. In Theorem 1.4, the invariance of $\ker T_F$ under the right-shift operator \mathcal{T} is sufficient condition for the boundedness of T . It is obvious that the invariance of $\ker T_F$ under \mathcal{T} is equivalent with $\ker T_F \subseteq \ker T_{\mathcal{T}_L F}$. In fact, for $\{c_n\}_{n=1}^\infty \in \ell^2$ we have $\mathcal{T}(\{c_n\}_{n=1}^\infty) \in \ker T_F$ if and only if $\{c_n\}_{n=1}^\infty \in \ker T_{\mathcal{T}_L F}$.

Proposition 4.6. *Let $F = \{f_n\}_{n=1}^\infty$ be a Bessel sequence in \mathcal{H} which is represented by $T \in B(\mathcal{H})$ and $M = \{f_n + f_{n+1}\}_{n=1}^\infty$ be a frame for \mathcal{H} . Then T is injective if and only if $\ker T_{\mathcal{T}_L^2 F} \subseteq \ker T_M$*

Proof. Let T be injective and $\{c_n\}_{n=1}^\infty \in \ker T_{\mathcal{T}_L^2 F}$. Then

$$T\left(\sum_{n=1}^\infty c_n(f_n + f_{n+1})\right) = \sum_{n=1}^\infty c_n f_{n+2} = 0.$$

Since T is injective, we get $\sum_{n=1}^\infty c_n(f_n + f_{n+1}) = 0$ and consequently $\{c_n\}_{n=1}^\infty \in \ker T_M$. Conversely, assume that $f \in \mathcal{H}$ and $Tf = 0$. Since $M = \{f_n + f_{n+1}\}_{n=1}^\infty$ is a frame for \mathcal{H} , we have $f = \sum_{n=1}^\infty c_n(f_n + f_{n+1})$ for some $\{c_n\}_{n=1}^\infty \in \ell^2$. Then

$$\sum_{n=1}^\infty c_n f_{n+2} = T\left(\sum_{n=1}^\infty c_n(f_n + f_{n+1})\right) = Tf = 0.$$

Hence $\{c_n\}_{n=1}^\infty \in \ker T_{\mathcal{T}_L^2 F}$ and consequently $\{c_n\}_{n=1}^\infty \in \ker T_M$. This means $f = \sum_{n=1}^\infty c_n(f_n + f_{n+1}) = 0$ and the proof is completed. \square

Proposition 4.7. *Let $\{f_n\}_{n=1}^\infty$ be represented by T . Then the following hold:*

- (i) *If $K \in B(\mathcal{H})$ is injective and has closed range, then $\{Kf_n\}_{n=1}^\infty$ has a Fibonacci representation.*
- (ii) *If $K \in B(\mathcal{H})$ is surjective, then $\{K^*f_n\}_{n=1}^\infty$ and $\{KK^*f_n\}_{n=1}^\infty$ have Fibonacci representations.*

Proof. (i) By Open Mapping Theorem, there exists a bonded linear operator $S : \mathcal{R}(K) \rightarrow \mathcal{H}$ such that $SK = I_{\mathcal{H}}$. Therefore

$$KTS(Kf_n + Kf_{n-1}) = KT(f_n + f_{n-1}) = Kf_{n+1}, \quad n \geq 2.$$

To prove (ii), by [4, Lemma 2.4.1], K^* is injective and has closed range. Also KK^* is invertible. Then (i) implies (ii). \square

Proposition 4.8. *Let $\{f_n\}_{n=1}^\infty$ be a frame for \mathcal{H} and represented by $T \in B(\mathcal{H})$. If $Tf_1 \in \text{span}\{f_n\}_{n=3}^\infty$, then $\mathcal{R}(T)$ is closed and $\mathcal{R}(T) = \overline{\text{span}}\{Tf_n\}_{n=1}^\infty = \overline{\text{span}}\{f_n\}_{n=3}^\infty$.*

Proof. For each $f \in \mathcal{H}$, there exists $\{c_n\}_{n=1}^\infty \in \ell^2$ such that $f = \sum_{n=1}^\infty c_n f_n$. Then $Tf = \sum_{n=1}^\infty c_n Tf_n \in \overline{\text{span}}\{Tf_n\}_{n=1}^\infty$, and therefore $\mathcal{R}(T) \subseteq \overline{\text{span}}\{Tf_n\}_{n=1}^\infty$. On the other hand, for $g \in \overline{\text{span}}\{f_n\}_{n=3}^\infty$ there exists $\{c_n\}_{n=1}^\infty \in \ell^2$ such that

$$g = \sum_{n=1}^\infty c_n f_{n+2} = \sum_{n=1}^\infty c_n T(f_n + f_{n+1}) = T\left(\sum_{n=1}^\infty c_n(f_n + f_{n+1})\right) \in \mathcal{R}(T).$$

Then $\overline{\text{span}}\{f_n\}_{n=3}^\infty \subseteq \mathcal{R}(T)$. Since $Tf_1 \in \text{span}\{f_n\}_{n=3}^\infty$, we can now apply (ii) of Lemma 3.7 to conclude that

$$\text{span}\{Tf_n\}_{n=1}^\infty = \text{span}\left\{\{Tf_1\} \cup \{Tf_n + Tf_{n+1}\}_{n=1}^\infty\right\} = \text{span}\{f_n\}_{n=3}^\infty.$$

Therefore $\mathcal{R}(T) = \overline{\text{span}}\{Tf_n\}_{n=1}^\infty = \overline{\text{span}}\{f_n\}_{n=3}^\infty$. \square

Proposition 4.9. *Let $\{f_n\}_{n=1}^\infty$ be a linearly dependent frame sequence represented by $T \in B(\mathcal{K})$, where $\mathcal{K} = \overline{\text{span}}\{f_n\}_{n=1}^\infty$ is an infinite dimensional Hilbert space. Then $\mathcal{R}(T)$ is closed and $\mathcal{R}(T) = \overline{\text{span}}\{f_n\}_{n=3}^\infty$.*

Proof. Let T_0 be the restriction of T on $\text{span}\{f_n\}_{n=1}^\infty$. Then by (ii) of Corollary 3.15, we have $\mathcal{R}(T_0) = \text{span}\{f_n\}_{n=3}^\infty$, and therefore $\mathcal{R}(T) \subseteq \overline{\text{span}}\{f_n\}_{n=3}^\infty$. On the other hand, Since $\{f_n\}_{n=1}^\infty$ is a frame sequence, for each $f \in \overline{\text{span}}\{f_n\}_{n=3}^\infty$, there exists $\{c_n\}_{n=1}^\infty \in \ell^2$ such that

$$f = \sum_{n=1}^\infty c_n f_{n+2} = \sum_{n=1}^\infty c_n (Tf_n + Tf_{n+1}) = T\left(\sum_{n=1}^\infty c_n(f_n + f_{n+1})\right) \in \mathcal{R}(T).$$

Hence $\overline{\text{span}}\{f_n\}_{n=3}^\infty \subseteq \mathcal{R}(T)$ and this completes the proof. \square

Theorem 4.10. *Let $\{f_n\}_{n=1}^\infty$ be represented by T and S . If $Tf_1 = Sf_1$, then $T = S$.*

Proof. Since $Tf_1 = Sf_1$ and $T(f_n + f_{n+1}) = f_{n+2} = S(f_n + f_{n+1})$ for all $n \in \mathbb{N}$, we get $Tf_n = Sf_n$ for all $n \in \mathbb{N}$ (we can use (i) of Lemma 3.7). This proves $T = S$ on $\text{span}\{f_n\}_{n=1}^\infty$. \square

Corollary 4.11. *Let $\{f_n\}_{n=1}^\infty$ be represented by T and S . If $f_1 \in \text{span}\{f_n + f_{n+1}\}_{n=k}^\infty$ for some $k \in \mathbb{N}$, then $T = S$.*

Proof. Since $f_1 \in \text{span}\{f_n + f_{n+1}\}_{n=k}^\infty$, we have $f_1 = \sum_{n=k}^m c_n(f_n + f_{n+1})$ for some scalars c_k, \dots, c_m . Then

$$Tf_1 = T\left(\sum_{n=k}^m c_n(f_n + f_{n+1})\right) = \sum_{n=k}^m c_n T(f_n + f_{n+1}) = \sum_{n=k}^m c_n f_{n+2} = \sum_{n=k}^m c_n S(f_n + f_{n+1}) = Sf_1.$$

Therefore $T = S$ by Theorem 4.10. \square

REFERENCES

- [1] A. Aldroubi, C. Cabrelli, A. F. Cakmak, U. Molter and A. Petrosyan, *Iterative actions of normal operators*, J. Funct. Anal. **272** (2017), 1121–1146.
- [2] A. Aldroubi, C. Cabrelli, U. Molter and S. Tang, *Dynamical sampling*, Appl. Comput. Harmon. Anal. **42** (2017), 378–401.
- [3] A. Aldroubi and A. Petrosyan, *Dynamical sampling and systems from iterative actions of operators*, Frames and other bases in abstract and function spaces, 15–26, Appl. Numer. Harmon. Anal., Birkh user/Springer, Cham, 2017.
- [4] Ole Christensen, *An introduction to frames and Riesz bases*. Second expanded edition. Birkh user (2016).
- [5] O. Christensen and M. Hasannasab, *Operator representations of frames: boundedness, duality, and stability*, Integr. Equ. Oper. Theory **88** (2017), 483–499.
- [6] O. Christensen and M. Hasannasab, *Frames property of systems arising via iterative actions of operators*, Appl. Comput. Harmon. Anal., **46** (2019), 3, 664–673.
- [7] O. Christensen, M. Hasannasab, and E. Rashidi, *Dynamical sampling and frame representations with bounded operators*, J. Math. Anal. Appl. **463** (2018), 634–644.
- [8] E. Rashidi, A. Najati, and E. Osgoei, *Dynamical sampling: mixed frame operators, representations and perturbations*, arXiv:1904.00697v1.

DEPARTMENT OF MATHEMATICS

FACULTY OF SCIENCES

UNIVERSITY OF MOHAGHEGH ARDABILI

ARDABIL, IRAN

E-mail address: j.smoghaddam@uma.ac.ir, a.nejati@yahoo.com, khedmati.y@uma.ac.ir