

# THE RIGOROUS DERIVATION OF THE $\mathbb{T}^2$ FOCUSING CUBIC NLS FROM 3D

SHUNLIN SHEN

**ABSTRACT.** We derive rigorously the 2D periodic focusing cubic NLS as the mean-field limit of the 3D focusing quantum many-body dynamics describing a dilute Bose gas with periodic boundary condition in the  $x$ -direction and a well of infinite-depth in the  $z$ -direction. Physical experiments for these systems are scarce. We find that, to fulfill the empirical requirement for observing NLS dynamics in experiments, namely, the kinetic energy dominates the potential energy, it is necessary to impose an extra restriction on the system parameters. This restriction gives rise to an unusual coupling constant.

## 1. INTRODUCTION

Boes-Einstein condensate (BEC) is the phenomenon that occurs when particles the same quantum state. The first experimental observation of BEC in an interacting atomic gas occurred in 1995, using laser cooling techniques [2, 29].

Let  $t \in \mathbb{R}$  be the time variable and  $\mathbf{r}_N = (r_1, r_2, \dots, r_N) \in \mathbb{R}^{3N}$  be the position vector of  $N$  particles in  $\mathbb{R}^3$ . Then BEC naively means that the  $N$ -body wave function  $\psi_N(t, \mathbf{r}_N)$  satisfies

$$\psi_N(t, \mathbf{r}_N) \sim \prod_{j=1}^N \varphi(t, r_j)$$

up to a phase factor solely depending on  $t$ , for some one particle state  $\varphi$ . That is, every particle takes the same quantum state. Equivalently, there is the Penrose-Onsager formulation of BEC: if we take  $\gamma_N^{(k)}$  be the  $k$ -particle marginal densities associated with  $\psi_N$  by

$$(1.1) \quad \gamma_N^{(k)}(t, \mathbf{r}_k, \mathbf{r}'_k) = \int \psi_N(t, \mathbf{r}_k, \mathbf{r}_{N-k}) \overline{\psi}_N(t, \mathbf{r}'_k, \mathbf{r}_{N-k}) d\mathbf{r}_{N-k}, \quad \mathbf{r}_k, \mathbf{r}'_k \in \mathbb{R}^{3k}.$$

Then BEC equivalently means

$$(1.2) \quad \gamma_N^{(k)}(t, \mathbf{r}_k, \mathbf{r}'_k) \sim \prod_{j=1}^k \varphi(t, r_j) \overline{\varphi}(t, r'_j).$$

It is widely believed that the cubic nonlinear Schrödinger equation (NLS)

$$(1.3) \quad i\partial_t \varphi = -\Delta \varphi + \mu |\varphi|^2 \varphi,$$

which is called focusing if  $\mu < 0$  and defocusing  $\mu > 0$ , describes BEC in the sense that  $\varphi$  satisfies NLS. In this paper, we are interested in the focusing case. There have been many physical experiments [28, 30, 46, 66] and mathematical results [21, 23, 24, 52, 54, 55, 58] regarding the focusing case. However, from the experiment [28], one infers that not only it is very difficult to prove the 3D focusing NLS as the mean-field limit of a 3D focusing quantum many-body dynamic, but such a limit also may not be true. Thus, in focusing settings, both physical experiments and mathematical results emphasize one dimensional and two dimensional behaviours. To our knowledge, physical experiments regarding the two dimensional behavior in the real-world three dimensional setting are limited and the corresponding mathematical research only studies the two dimensional behaviour in 2D. Therefore, we turn our attention to the derivation of 2D focusing NLS from 3D. Interestingly, our analysis produces an unusual microscopic-to-macroscopic coupling constant and might provide some suggestions to the experiment. To expect a two-dimensional behaviour, we should confine a large number of bosons inside a trap with strong confinement in one direction. We consider a simple physical model, namely, quantum many-body dynamics with periodic boundary condition in the  $x$ -direction

and a well of infinite-depth in the  $z$ -direction<sup>1</sup>. Such model with strong restriction in one direction was first considered by Schnee and Yngvason [62] for the defocusing time-independent problem. Then, the defocusing time-dependent 3D-to-2D program was studied by X. Chen and Holmer in [19], in which they used the quadratic potential  $|\cdot|^2$  to represent the trap. Here, we model the trap by using a well of infinite-depth in the  $z$ -direction. That is, we consider the Hamiltonian (see [62])

$$(1.4) \quad H_{N,L,a} = \sum_{j=1}^N (-\Delta_{r_j} + L^{-2}V^\perp(z_j/L)) + \sum_{1 \leq i < j \leq N} \frac{1}{a^{3\beta-1}} V\left(\frac{r_i - r_j}{a^\beta}\right),$$

where

$$V^\perp(z) = \begin{cases} 1, & z \in (-\pi/2, \pi/2), \\ \infty, & z \notin (-\pi/2, \pi/2). \end{cases}$$

For parameter  $a$ , the scaling of the interaction potential, we consider the case, which is called Region I in [62]. Schnee and Yngvason define  $g = g(N, L, a)$  as follows

$$g \sim (-\ln(L^2\bar{\rho}) + L/a)^{-1},$$

where  $\bar{\rho}$  is the mean density. The Gross-Pitaevskii limit means  $Ng \sim 1$  and hence  $\bar{\rho} \sim N$ . Then, the term  $a/L$  dominates in the definition of  $g$ . Therefore,

$$1 \sim Ng \sim Na/L \iff a \sim L/N.$$

For mathematical convenience, we take  $a = L/N$  and consider the Hamiltonian

$$(1.5) \quad H_{N,L} = \sum_{j=1}^N -\Delta_{r_j} + \frac{L}{N-1} \sum_{1 \leq i < j \leq N} V_{N,L}(r_i - r_j).$$

acting on the Hilbert space  $L_s^2(\Omega_L^{\otimes N})$ , the subspace of  $L^2(\Omega_L^{\otimes N})$  consisting of functions that are symmetric with respect to permutations of the  $N$  particles, where

$$V_{N,L}(r_i - r_j) = (N/L)^{3\beta} V((N/L)^\beta(r_i - r_j))$$

and the domain<sup>2</sup>  $\Omega_L = (-\pi, \pi)^2 \times (-L\pi/2, L\pi/2)$ . As  $L \rightarrow 0$ , we see that the particles are strongly confined in the  $z$ -direction. For more detailed analysis of system parameters, also see [19].

We take the periodic boundary condition<sup>3</sup> in the  $x$ -direction and Dirichlet boundary condition in the  $z$ -direction. We will derive rigorously  $\mathbb{T}^2$  focusing cubic NLS from the 3D quantum many-body dynamic. For simplicity, we take  $\cos_L(z) = (2/\pi)^{1/2} \cos(z/L)/L^2$ , which is the normalized ground state eigenfunction. With the lowest energy, we notice that, as  $L \rightarrow 0$ ,  $\cos_L(z)$  has infinite energy. Thus, our main theorem is better to be stated regarding the renormalization.

Let  $\psi_{N,L}(t, \cdot) = e^{itH_{N,L}}\psi_{N,L}(0, \cdot)$  denote the evolution of this initial data corresponding to the Hamiltonian operator (1.5). Define the rescaled solution

$$(1.6) \quad \tilde{\psi}_{N,L}(t, \mathbf{r}_N) \stackrel{def}{=} L^{N/2} \psi_{N,L}(t, \mathbf{x}_N, L\mathbf{z}_N), \quad \mathbf{r}_N \in \mathbb{T}^{2N} \times (-\pi/2, \pi/2)^N,$$

and the rescaled Hamiltonian

$$(1.7) \quad \tilde{H}_{N,L} = \sum_{j=1}^N \left( -\Delta_{x_j} - \frac{1}{L^2} \partial_{z_j}^2 \right) + \frac{1}{N-1} \sum_{1 \leq i < j \leq N} \tilde{V}_{N,L}(r_i - r_j),$$

where

$$(1.8) \quad \tilde{V}_{N,L}(r) = L(N/L)^{3\beta} V((N/L)^\beta x, L(N/L)^\beta z).$$

Then

$$\left( \tilde{H}_{N,L} \tilde{\psi}_{N,L} \right) (t, \mathbf{x}_N, \mathbf{z}_N) = L^{N/2} (H_{N,L} \psi_{N,L}) (t, \mathbf{x}_N, L\mathbf{z}_N),$$

<sup>1</sup>Our exact proof also works for the case in which we put  $\mathbb{R}^2$  in the  $x$ -direction. We choose  $\mathbb{T}^2$  here, considering all of these limits problem originated from the thermodynamic limit on  $\mathbb{T}^3$  (see a survey in [4]).

<sup>2</sup>When  $L = 1$ , we take  $\Omega = \Omega_1$  for convenience.

<sup>3</sup>To match the periodic condition,  $V_{N,L}(r)$  is considered as the periodic extension in the  $x$ -direction of the rescaled  $V$  which is compactly supported on  $\Omega$ .

and hence, we have

$$(1.9) \quad \tilde{\psi}_{N,L}(t, \mathbf{r}_N) = e^{it\tilde{H}_{N,L}} \tilde{\psi}_{N,L}(0, \mathbf{r}_N).$$

**Definition 1.1.** We denote  $C_{gn}$  the sharp constant of the 2D inhomogeneous Gagliardo-Nirenberg estimate<sup>4</sup> on torus:

$$(1.10) \quad \|\phi\|_{L^4(\mathbb{T}^2)} \leq C_{gn} \|\phi\|_{L^2(\mathbb{T}^2)}^{\frac{1}{2}} \|\sqrt{1-\Delta}\phi\|_{L^2(\mathbb{T}^2)}^{\frac{1}{2}}.$$

**Theorem 1.2.** Assume  $L(N/L)^\beta \rightarrow 1^-$  and the pair interaction  $V$  is an even nonpositive smooth function compactly support on  $\Omega$  such that  $\|V\|_{L_z^\infty L_x^1} \leq \frac{2\alpha}{C_{gn}^4}$  for some  $\alpha \in (0, 1)$ . Let  $\{\tilde{\gamma}_{N,L}^{(k)}(t, \mathbf{r}_k, \mathbf{r}'_k)\}$  be the family of marginal densities associated with the 3D rescaled Hamiltonian evolution  $\tilde{\psi}_{N,L}(t) = e^{it\tilde{H}_{N,L}} \tilde{\psi}_{N,L}(0)$  for  $\beta \in (0, 3/7)$ . Suppose the initial datum  $\tilde{\psi}_{N,L}(0)$  satisfies the following:

- (i)  $\tilde{\psi}_{N,L}(0)$  is normalized, that is,  $\|\tilde{\psi}_{N,L}(0)\|_{L^2} = 1$ ,
- (ii)  $\tilde{\psi}_{N,L}(0)$  is asymptotically factorized in the sense that

$$\lim_{N, 1/L \rightarrow \infty} \text{Tr} \left| \tilde{\gamma}_{N,L}^{(1)}(0, x_1, z_1; x'_1, z'_1) - \frac{2}{\pi} \phi_0(x_1) \overline{\phi_0}(x'_1) \cos(z_1) \cos(z'_1) \right| = 0,$$

for some one particle state  $\phi_0 \in H^1(\mathbb{T}^2)$ ,

- (iii) Away from the  $z$ -direction ground state energy,  $\tilde{\psi}_{N,L}(0)$  has finite energy per particle

$$\sup_{N,L} \langle \tilde{\psi}_{N,L}(0), (N^{-1} \tilde{H}_{N,L} - 1/L^2) \tilde{\psi}_{N,L}(0) \rangle \leq C.$$

Then  $\forall k \geq 1, t \geq 0$ , we have the convergence in trace norm that

$$\lim_{\substack{N, 1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} \text{Tr} \left| \tilde{\gamma}_{N,L}^{(k)}(t, \mathbf{x}_k, \mathbf{z}_k; \mathbf{x}'_k, \mathbf{z}'_k) - \prod_{j=1}^k \frac{2}{\pi} \phi(t, x_j) \overline{\phi}(t, x'_j) \cos(z_j) \cos(z'_j) \right| = 0,$$

where  $\phi(t, x)$  solves the 2D periodic focusing cubic NLS with coupling constant

$$g_0 = \frac{4}{\pi^2} \int \int V(x, z_1 - z_2) dx |\cos(z_1) \cos(z_2)|^2 dz_1 dz_2 < 0,$$

that is

$$(1.11) \quad i\partial_t \phi = -\Delta_x \phi + g_0 |\phi|^2 \phi,$$

with initial condition  $\phi(0, x) = \phi_0(x)$ .

It is well-known that Theorem 1.2 is equivalent to Theorem 1.3 by the method of Erdős, Schlein, and Yau [32, 33, 34, 35, 36].

**Theorem 1.3.** Assume  $L(N/L)^\beta \rightarrow 1^-$  and the pair interaction  $V$  is an even nonpositive smooth function compactly support on  $\Omega$  such that  $\|V\|_{L_z^\infty L_x^1} \leq \frac{2\alpha}{C_{gn}^4}$  for some  $\alpha \in (0, 1)$ . Let  $\{\tilde{\gamma}_{N,L}^{(k)}(t, \mathbf{r}_k, \mathbf{r}'_k)\}$  be the family of marginal densities associated with the 3D rescaled Hamiltonian evolution  $\tilde{\psi}_{N,L}(t) = e^{it\tilde{H}_{N,L}} \tilde{\psi}_{N,L}(0)$  for  $\beta \in (0, 3/7)$ . Suppose the initial datum  $\tilde{\psi}_{N,L}(0)$  is normalized asymptotically factorized and satisfies the energy condition that

- (iii') there is a constant  $C > 0$  such that

$$\sup_{N,L} \langle \tilde{\psi}_{N,L}(0), (N^{-1} \tilde{H}_{N,L} - 1/L^2)^k \tilde{\psi}_{N,L}(0) \rangle \leq C^k, \quad \forall k \geq 1.$$

Then  $\forall k \geq 1, t \geq 0$ , we have the convergence in trace norm that

$$\lim_{\substack{N, 1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} \text{Tr} \left| \tilde{\gamma}_{N,L}^{(k)}(t, \mathbf{x}_k, \mathbf{z}_k; \mathbf{x}'_k, \mathbf{z}'_k) - \prod_{j=1}^k \frac{2}{\pi} \phi(t, x_j) \overline{\phi}(t, x'_j) \cos(z_j) \cos(z'_j) \right| = 0,$$

<sup>4</sup>There are many versions of the Gagliardo-Nirenberg inequalities on  $\mathbb{T}^2$ . Our proof works more or less the same.

<sup>5</sup>We use the notation  $L(N/L)^\beta \rightarrow 1^-$  to denote  $L(N/L)^\beta \leq 1$  and  $L(N/L)^\beta \rightarrow 1$ .

where  $\phi(t, x)$  solves the 2D periodic focusing cubic NLS with the coupling constant

$$g_0 = \frac{4}{\pi^2} \int \int V(x, z_1 - z_2) dx |\cos(z_1) \cos(z_2)|^2 dz_1 dz_2 < 0,$$

that is

$$(1.12) \quad i\partial_t \phi = -\Delta_x \phi + g_0 |\phi|^2 \phi,$$

with initial condition  $\phi(0, x) = \phi_0(x)$ .

For  $\beta < 1/3$ , our exact proof also works for the case in which we put  $\mathbb{R}^2$  in the  $x$ -direction.

**Theorem 1.4.** *Under the same condition of Theorem 1.2 with  $\beta \in (0, 1/3)$ , then  $\forall k \geq 1$ ,  $t \geq 0$ , we have the convergence in trace norm that*

$$\lim_{\substack{N, 1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} \text{Tr} \left| \tilde{\gamma}_{N,L}^{(k)}(t, \mathbf{x}_k, \mathbf{z}_k; \mathbf{x}'_k, \mathbf{z}'_k) - \prod_{j=1}^k \frac{2}{\pi} \phi(t, x_j) \bar{\phi}(t, x'_j) \cos(z_j) \cos(z'_j) \right| = 0,$$

where  $\phi(t, x)$  solves the 2D focusing cubic NLS with the coupling constant

$$g_0 = \frac{4}{\pi^2} \int \int V(x, z_1 - z_2) dx |\cos(z_1) \cos(z_2)|^2 dz_1 dz_2 < 0,$$

that is

$$(1.13) \quad i\partial_t \phi = -\Delta_x \phi + g_0 |\phi|^2 \phi,$$

with initial condition  $\phi(0, x) = \phi_0(x) \in H^1(\mathbb{R}^2)$ .

We notice that Theorems 1.2 – 1.3 carry an extra requirement  $L(N/L)^\beta \rightarrow 1^-$  and a different coupling constant<sup>6</sup>, if compared to the previous work, for example [19, 23, 24], in which the constant is usually  $\int V$  or the scattering length of  $V$ . It emerges from the empirical requirement for observing NLS dynamics in experiments, namely, the kinetic energy dominates the potential energy. We will certainly explain it in detail during the course of the proof. Due to the requirement, the limit of  $\tilde{V}_{N,L}$  defined by (1.8) is not a 3D  $\delta$ -function, though it scales like one.

There are two well-developed schemes to deal with this type of procedure. One is the Fock space method, while the other is the hierarchy approach. We take the hierarchy approach here<sup>7</sup>. The BBGKY hierarchy associated with  $\tilde{\psi}_{N,L}$  is

$$(1.14) \quad i\partial_t \tilde{\gamma}_{N,L}^{(k)} = \sum_{j=1}^k \left[ -\Delta_{x_j}, \tilde{\gamma}_{N,L}^{(k)} \right] + \frac{1}{L^2} \sum_{j=1}^k \left[ -\partial_{z_j}^2, \tilde{\gamma}_{N,L}^{(k)} \right] \\ + \frac{1}{N-1} \sum_{1 \leq i < j \leq k} \left[ \tilde{V}_{N,L}(r_i - r_j), \tilde{\gamma}_{N,L}^{(k)} \right] + \frac{N-k}{N-1} \sum_{j=1}^k \text{Tr}_{r_{k+1}} \left[ \tilde{V}_{N,L}(r_j - r_{k+1}), \tilde{\gamma}_{N,L}^{(k+1)} \right].$$

It was Erdős, Schlein, and Yau who first rigorously derived the 3D cubic defocusing NLS from a 3D quantum many-body dynamic in their fundamental papers [32, 33, 34, 35, 36]. They proved a-priori  $L_T^\infty H_x^1$  bound to establish the compactness of BBGKY with respect to a topology on the trace class operators. Then, they showed that the limit point satisfies GP hierarchy. Finally, the proof for the uniqueness of GP hierarchy was the principal part and also surprisingly dedicate due to the fact that it is a system of infinitely many coupled equations over an unbounded number of variables. It motivated a large amount of works [1, 8, 9, 10, 11, 12, 13, 15, 16, 17, 18, 19, 25, 47, 48, 60, 67].

Subsequently, with imposing an additional a-priori condition on space-time norm, Klainerman and Machedon [48] gave an another proof of the uniqueness of GP hierarchy in a different space of density matrices defined by Hilbert-Schmidt type Sobolev norms. Later, the approach of Klainerman and Machedon was used by Kirkpatrick, Schlein and Staffilani [47] to derived the 2D cubic defocusing NLS from the 2D quantum many-body dynamic both on  $\mathbb{R}^2$  and  $\mathbb{T}^2$ ; by T. Chen and Pavlović [10] to derive the quintic NLS for  $d = 1$ ,

<sup>6</sup>This extra requirement and the coupling constant certainly give rises to a density condition for the gas. We do not compute this density as it is not our main goal here.

<sup>7</sup>We believe the Fock space method will reach the same result. We just prefer a  $H^1$  result here. In fact, some techniques we used come from the Fock space literatures [52, 55].

2; by X. Chen [18] to investigate the trapping problem in 2D and 3D; and by X. Chen and Holmer [19] to derive 2D cubic defocusing NLS from the 3D quantum many-body dynamic.

Later on, T. Chen, Hainzl, Pavlović and Seiringer [8], using the quantum de Finetti theorem from [53], provided a simplified proof of the  $L_T^\infty H_x^1$ -type 3D cubic uniqueness theorem in [33]. This method in [8] inspired the study for refined uniqueness theorems, such as [26, 45, 64].

Using Fock space methods to study the convergence rate has also been worked on by many authors, for example, see [3, 7, 39, 40, 50, 51, 58, 61], and the references within.

For the focusing setting, which is a natural continuation of the defocusing problem, X. Chen and Holmer [21] first derived the 1D focusing cubic NLS and later a 3D-to-1D reduction in [23]. But the 2D cubic case did not see any process until [54], in which Lewin, Nam, and Rougerie used a quantitative version of the quantum de Finetti theorem [27] to show that the ground state energy of the 2D  $N$ -body was described by a NLS ground state energy. Using the finite-dimensional quantum de Finetti theorem in [54], X. Chen and Holmer [24] derived 2D focusing cubic NLS from the 2D quantum many-body dynamic for  $\beta \in (0, 1/6)$ . For higher  $\beta$ , Lewin, Nam, and Rougerie [55] used a bootstrapping argument to improve  $\beta$ , which, together with the approach in [24], implied the convergence of the quantum many-body dynamics to the focusing NLS for  $\beta \in (0, 3/4)$ . In the 2D focusing case, the stability of the second kind which is the energy bound when  $k = 1$  was improved to  $\beta < 1$  in [59]. Besides the convergence of density matrix, the convergence rate is also of interest and extended to lower dimension in both focusing and defocusing cases by Nam and Napiórkowski in [58] using  $H^4$  regularity.

The derivation of 2D defocusing cubic NLS from 3D was first by X. Chen and Holmer in [19] and then by Boßmann in [5] for the regime  $\beta \in (0, 1]$ . To our knowledge, the derivation of 2D focusing cubic NLS from 3D has not been completed before. In this paper, we follow the lead of the aforementioned focusing works [21, 23, 24, 52, 54, 55] and pursue the treatment of 2D case from the 3D physical setting.

**1.1. Outline of the Proof of Theorem 1.2.** We first establish in Section 2, under the assumption  $L(N/L)^\beta \rightarrow 1^-$ , that the renormalized kinetic energy controls the potential energy and hence yield an  $H^1$  regularity bound to make the other parts of the paper work.

In section 2.1, we use scaling arguments to show why we choose the uncommon mixed norm  $\|V\|_{L_z^\infty L_x^1}$  and we are bounded by the extra restriction  $L(N/L)^\beta \rightarrow 1^-$ . In fact, a similar requirement would also show up in the harmonic well case studied by X. Chen and Holmer [19] if one wants the renormalized kinetic energy to bound the potential energy instead of dropping it. Subsequently we prove the energy bound with  $\beta < 1/2$  when  $k = 1$ , which is divided into two parts. First in Section 2.2, we get to  $\beta < 1/3$ . Instead of taking the approach in X. Chen and Holmer [19, 23, 24], our proof improvises from Lewin, Nam, and Rougerie [55] and Lewin [52]. This proof does not use the finite-dimensional quantum de Finetti theorem and thus can be applied to the  $\mathbb{R}^2$  case as well. Hence, the main Theorem 1.2 works the same for  $\mathbb{R}^2$  with  $\beta < 1/3$ . Second in Section 2.3, we adapt the bootstrapping argument in [55] to reach  $\beta < 1/2$ . Then in Section 2.4, we complete Theorem 2.1 when  $k > 1$ , where the restriction  $\beta < 3/7$  is required. In the defocusing case [19], if we require a similar requirement, the index  $\beta$  can be improved to  $3/7$  as well.

In Section 3, we show the compactness of the BBGKY sequence. Then, we use a modified version of the approximation of identity type lemma to show that limit points satisfy the GP hierarchy with the unusual coupling constant  $g_0$ . The uniqueness for GP hierarchy on  $\mathbb{T}^2$  has been well studied by Kirkpatrick, Schlein and Staffilani [47], Herr and Sohinger [42, 43]. We use their uniqueness theorems to conclude our proof.

## 2. FOCUSING ENERGY ESTIMATES

In this section, we prove focusing energy estimates. Define

$$S_j := (1 - \Delta_{r_j} - 1/L^2)^{1/2},$$

and write

$$S^{(k)} = \prod_{j=1}^k S_j.$$

**Theorem 2.1.** Assume  $L(N/L)^\beta \rightarrow 1^-$ ,  $\beta < \frac{3}{7}$ , and  $\|V\|_{L_z^\infty L_x^1} \leq \frac{2\alpha}{C_{gn}^4}$  for some  $\alpha \in (0, 1)$ , then let  $c_0 = \min\left(\frac{1-\alpha}{\sqrt{2}}, \frac{1}{2}\right)$ , we have  $\forall k \geq 0$ , there exists an  $N_0(k) > 0$  such that

$$(2.1) \quad \langle \psi_{N,L}, (2 + N^{-1}H_{N,L} - 1/L^2)^k \psi_{N,L} \rangle \geq c_0^k \|S^{(k)} \psi_{N,L}\|_{L^2}^2$$

for all  $N > N_0$  and for all  $\psi_{N,L} \in L_s^2(\Omega_L^{\otimes N})$ .

*Proof.* For smoothness of presentation, we postpone the proof of Theorem 2.1 to Section 2.2 – 2.4.  $\square$

Now we convert the conclusions of Theorem 2.1 into the statement about the rescaled solution, which we will use in the remainder of the paper.

Let  $\tilde{P}_0$  denote the orthogonal projection onto the ground state of  $-\partial_z^2 - 1$  on the region  $(-\pi/2, \pi/2)$  with Dirichlet boundary condition and  $\tilde{P}_{\geq 1} = I - \tilde{P}_0$ . We define  $\tilde{P}_0^j$  and  $\tilde{P}_1^j$  to be respectively  $\tilde{P}_0$  and  $\tilde{P}_{\geq 1}$  acting on the  $z_j$ -variable, and

$$(2.2) \quad \tilde{P}_\alpha = \tilde{P}_{\alpha_1}^1 \cdots \tilde{P}_{\alpha_k}^k$$

for a  $k$ -tuple  $\alpha = (\alpha_1, \dots, \alpha_k)$  with  $\alpha_j \in \{0, 1\}$  and adopt the notation  $|\alpha| = \alpha_1 + \cdots + \alpha_k$ . Then

$$(2.3) \quad I = \sum_{\alpha} \tilde{P}_\alpha,$$

where  $I : L^2(\Omega^k) \rightarrow L^2(\Omega^k)$ .

*Corollary 2.2.* Define

$$\tilde{S}_j = (1 - \Delta_{x_j} - \partial_{z_j}^2 / L^2 - 1/L^2)^{1/2},$$

and write

$$\tilde{S}^{(k)} = \prod_{j=1}^k \tilde{S}_j, \quad \langle \nabla \rangle^{(k)} = \prod_{j=1}^k \sqrt{1 - \Delta_{r_j}}.$$

Assume  $L(N/L)^\beta \rightarrow 1^-$ . Let  $\tilde{\psi}_{N,L}(t) = e^{it\tilde{H}_{N,L}} \tilde{\psi}_{N,L}(0)$  and  $\{\tilde{\gamma}_{N,L}^{(k)}(t)\}$  be the associated marginal densities. Then for all  $k \geq 0$ , we have the uniform-in-time bound

$$(2.4) \quad Tr \tilde{S}^{(k)} \tilde{\gamma}_{N,L}^{(k)} \tilde{S}^{(k)} = \|\tilde{S}^{(k)} \tilde{\psi}_{N,L}(t)\|_{L^2}^2 \leq C^k.$$

Consequently,

$$(2.5) \quad Tr \langle \nabla \rangle^{(k)} \tilde{\gamma}_{N,L}^{(k)} \langle \nabla \rangle^{(k)} = \|\langle \nabla \rangle^{(k)} \tilde{\psi}_{N,L}(t)\|_{L^2}^2 \leq C^k,$$

and

$$(2.6) \quad \|\tilde{P}_\alpha \tilde{\psi}_{N,L}\|_{L^2} \leq C^k L^{|\alpha|}, \quad |Tr \tilde{P}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{P}_\beta| \leq C^k L^{|\alpha|+|\beta|},$$

where  $\tilde{P}_\alpha$  and  $\tilde{P}_\beta$  are defined as in (2.2).

*Proof.* We notice that

$$\begin{aligned} (\tilde{S}_j^2 \tilde{\psi}_{N,L})(t, \mathbf{x}_N, \mathbf{z}_N) &= L^{N/2} (S_j^2 \psi_{N,L})(t, \mathbf{x}_N, L\mathbf{z}_N), \\ (\tilde{H}_{N,L} \tilde{\psi}_{N,L})(t, \mathbf{x}_N, \mathbf{z}_N) &= L^{N/2} (H_{N,L} \psi_{N,L})(t, \mathbf{x}_N, L\mathbf{z}_N), \end{aligned}$$

where  $\tilde{\psi}_{N,L}$  is defined by (1.6). Thus, we have

$$\begin{aligned} \|\tilde{S}^{(k)} \tilde{\psi}_{N,L}\|_{L^2}^2 &= \|S^{(k)} \psi_{N,L}\|_{L^2}^2, \\ \langle \tilde{\psi}_{N,L}, (2 + N^{-1} \tilde{H}_{N,L} - 1/L^2)^k \tilde{\psi}_{N,L} \rangle &= \langle \psi_{N,L}, (2 + N^{-1} H_{N,L} - 1/L^2)^k \psi_{N,L} \rangle. \end{aligned}$$

From estimate (2.1) in Theorem 2.1, we obtain

$$\|\tilde{S}^{(k)} \tilde{\psi}_{N,L}(t)\|_{L^2}^2 \leq C^k \langle \tilde{\psi}_{N,L}(t), (2 + N^{-1} \tilde{H}_{N,L} - 1/L^2)^k \tilde{\psi}_{N,L}(t) \rangle.$$

The term on the right-hand side is conserved, so

$$\|\tilde{S}^{(k)} \tilde{\psi}_{N,L}(t)\|_{L^2}^2 \leq C^k \langle \tilde{\psi}_{N,L}(0), (2 + N^{-1} \tilde{H}_{N,L} - 1/L^2)^k \tilde{\psi}_{N,L}(0) \rangle.$$

Applying the binomial theorem twice,

$$\begin{aligned} \|\tilde{S}^{(k)}\tilde{\psi}_{N,L}(t)\|_{L^2}^2 &\leq C^k \sum_{j=0}^k \binom{k}{j} 2^j \langle \tilde{\psi}_{N,L}(0), (N^{-1}\tilde{H}_{N,L} - 1/L^2)^{k-j} \tilde{\psi}_{N,L}(0) \rangle \\ &\leq C^k \sum_{j=0}^k \binom{k}{j} 2^j C^{k-j} \\ &= C^k (2 + C)^k \leq \tilde{C}^k, \end{aligned}$$

where we used initial condition in the second-to-last line. So we have established (2.4). Combining (2.4) and (A.31), estimate (2.5) then follows. By (2.4) and (A.30), we obtain the first inequality of (2.6). By Lemma A.7,

$$\text{Tr} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta = \langle \tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}, \tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L} \rangle,$$

so the second inequality of (2.6) follows by Cauchy-Schwarz inequality.  $\square$

**2.1. Explanations on the assumptions.** We will explain the idea that we choose the mixed norm  $\|V\|_{L_z^\infty L_x^1}$  and the relationship  $L(N/L)^\beta \rightarrow 1^-$ , both of which are different from the previous work, such as [19, 23, 24]. In fact, to derive the 2D focusing NLS equations, the key point is that the interaction energy can be controlled by the kinetic energy, which is described by Theorem 2.1 when  $k = 1$ . By a scaling, we can see that the mixed norm  $\|V\|_{L_z^\infty L_x^1}$  is reasonable and  $L(N/L)^\beta$  should be bounded.

2.1.1. We begin by setting up some notations for simplicity. Let

$$H_{ij} = S_i^2 + S_j^2 + H_{Iij},$$

and

$$H_{Iij} = LV_{N,L}(r_i - r_j),$$

where the subscript  $I$  represents the interaction energy. Then, we can rewrite

$$(2.7) \quad 1 + N^{-1}H_{N,L} - 1/L^2 = \frac{1}{N(N-1)} \sum_{1 \leq i < j \leq N} H_{ij}.$$

When we take  $\psi_{N,L} = \phi_L^{\otimes N}$  with  $\|\phi_L\|_{L^2} = 1$ , we find

$$(2.8) \quad \langle \psi_{N,L}, (1 + N^{-1}H_{N,L} - 1/L^2) \psi_{N,L} \rangle = \frac{1}{2} \langle \phi_L^{\otimes 2}, H_{12} \phi_L^{\otimes 2} \rangle.$$

If  $H_{12} \geq 0$ , we can deduce that

$$(2.9) \quad \begin{aligned} -L \int_{\Omega_L^2} V_{N,L}(r_1 - r_2) |\phi_L(r_1) \phi_L(r_2)|^2 dr_1 dr_2 &\leq C(V) \langle S_1^2 \phi_L(r_1) \phi_L(r_2), \phi_L(r_1) \phi_L(r_2) \rangle \\ &\leq C(V) \langle (1 - \Delta_{r_1}) \phi_L(r_1) \phi_L(r_2), \phi_L(r_1) \phi_L(r_2) \rangle, \end{aligned}$$

where  $C(V)$  depends on  $V$ . Moreover, if we assume  $C(V) = C\|V\|_X$  where  $\|\cdot\|_X$  is a norm, by a scaling argument, it should satisfy

$$\|V\|_X \leq \lambda^{-2} \|V(\cdot/\lambda)\|_X, \quad \forall \lambda \in (0, 1).$$

Indeed, we take  $V^\lambda(\cdot) = V(\cdot/\lambda)$  and  $\phi_L^\lambda(\cdot) = \lambda^{-3/2} \phi_L(\cdot/\lambda)$  to replace  $V$  and  $\phi_L$  respectively. Since we take the periodic condition in the  $x$ -direction, a scaling argument can only be used for the function supported in the interior of the domain. Thus, we consider the test function  $\phi_L \in C_c^\infty(\Omega_L)$ , the space of smooth functions compactly supported in  $(-\pi, \pi)^2 \times (-L\pi/2, L\pi/2)$ . For every  $\phi_L \in C_c^\infty(\Omega_L)$ , we have

$$(2.10) \quad -L \int_{\Omega_L^2} V_{N,L}(r_1 - r_2) |\phi_L(r_1) \phi_L(r_2)|^2 dr_1 dr_2 = -L \int_{\Omega_L^2} V_{N,L}^\lambda(r_1 - r_2) |\phi_L^\lambda(r_1) \phi_L^\lambda(r_2)|^2 dr_1 dr_2,$$

and

$$(2.11) \quad C\|V^\lambda\|_X \langle (1 - \Delta_{r_1}) \phi_L^\lambda(r_1) \phi_L^\lambda(r_2), \phi_L^\lambda(r_1) \phi_L^\lambda(r_2) \rangle \leq C\|V\|_X \lambda^{-2} \langle (1 - \Delta_{r_1}) \phi_L(r_1) \phi_L(r_2), \phi_L(r_1) \phi_L(r_2) \rangle.$$

If there exists a  $\lambda_0 \in (0, 1)$  such that  $\lambda_0^{-2} \|V(\cdot/\lambda_0)\|_X \leq q_0 \|V\|_X$  for some  $q_0 \in (0, 1)$ , we take  $\lambda = \lambda_0$ . Putting (2.9) (2.10) and (2.11) together, we get

$$-L \int_{\Omega_L^2} V_{N,L}(r_1 - r_2) |\phi_L(r_1) \phi_L(r_2)|^2 dr_1 dr_2 \leq q_0 C \|V\|_X \langle (1 - \Delta_{r_1}) \phi_L(r_1) \phi_L(r_2), \phi_L(r_1) \phi_L(r_2) \rangle,$$

for all  $\phi_L \in C_c^\infty(\Omega_L)$ . Iterating the process, it will lead to a contradiction for  $q_0 < 1$ .

On the one hand, the common norm  $\|\cdot\|_{L^1}$  cannot satisfy the above requirement, since

$$\|V\|_{L^1} = \lambda^{-3} \|V^\lambda\|_{L^1} \geq \lambda^{-2} \|V^\lambda\|_{L^1}, \quad \forall \lambda \in (0, 1).$$

On the other hand, we note that

$$\|V\|_{L_z^\infty L_x^1} = \lambda^{-2} \|V(\cdot/\lambda)\|_{L_z^\infty L_x^1}, \quad \forall \lambda \in (0, 1).$$

That is, the mixed norm  $L_z^\infty L_x^1$  satisfies the requirement. Indeed, we can establish a general Lemma 2.7 in Section 2.2.

2.1.2. To derive the relationship between  $N$  and  $L$ , the key point is also that the interaction energy can be controlled by the kinetic energy. More precisely, let us consider the rescaled system. We take the test function  $\tilde{\phi}(r) = f(x)\tilde{g}(z)$ , where  $f(x) \in C_c^\infty((-\pi, \pi)^2)$ ,  $\tilde{g}(z) \in C_c^\infty(-\pi/2, \pi/2)$  and

$$\|f\|_{L^2} = \|\tilde{g}\|_{L^2} = 1.$$

Define

$$(2.12) \quad f_\varepsilon(x) = \frac{1}{\varepsilon} f(x/\varepsilon) \in C_c^\infty((-\pi, \pi)^2), \quad \varepsilon \in (0, 1),$$

$$(2.13) \quad \tilde{g}_\lambda(z) = \frac{1}{\sqrt{\lambda}} \tilde{g}(z/\lambda) \in C_c^\infty(-\pi/2, \pi/2), \quad \lambda \in (0, 1),$$

$$(2.14) \quad \tilde{\phi}_{\varepsilon,\lambda}(r) = f_\varepsilon(x) \tilde{g}_\lambda(z),$$

$$(2.15) \quad \tilde{\psi}_{\varepsilon,\lambda}(r_1, r_2) = \tilde{\phi}_{\varepsilon,\lambda}(r_1) \tilde{\phi}_{\varepsilon,\lambda}(r_2).$$

where  $f_\varepsilon(x)$  should be considered as a periodic extension and  $f_\varepsilon(x) \in C^\infty(\mathbb{T}^2)$ .

The interaction energy is

$$(2.16) \quad \begin{aligned} & \int \tilde{V}_{N,L}(r_1 - r_2) |f_\varepsilon(x_1) \tilde{g}_\lambda(z_1) f_\varepsilon(x_2) \tilde{g}_\lambda(z_2)|^2 dr_1 dr_2 \\ &= \int \tilde{V}_{N,L}(\varepsilon(x_1 - x_2), \lambda(z_1 - z_2)) |f(x_1) \tilde{g}(z_1) f(x_2) \tilde{g}(z_2)|^2 dr_1 dr_2. \end{aligned}$$

The kinetic energy is

$$(2.17) \quad \begin{aligned} & \langle \tilde{S}_1^2 \tilde{\psi}_{\varepsilon,\lambda}, \tilde{\psi}_{\varepsilon,\lambda} \rangle \\ &= \|f_\varepsilon\|_{L^2}^2 \|\tilde{g}_\lambda\|_{L^2}^2 \left( \langle (1 - \Delta_x) f_\varepsilon(x) \tilde{g}_\lambda(z), f_\varepsilon(x) \tilde{g}_\lambda(z) \rangle + \frac{1}{L^2} \langle f_\varepsilon(x) (-\partial_z^2 - 1) \tilde{g}_\lambda(z), f_\varepsilon(x) \tilde{g}_\lambda(z) \rangle \right) \\ &= \|f_\varepsilon\|_{L^2}^2 \|\tilde{g}_\lambda\|_{L^2}^2 \left( \|\nabla_x f_\varepsilon\|_{L^2}^2 \|\tilde{g}_\lambda\|_{L^2}^2 + \|f_\varepsilon\|_{L^2}^2 \|\tilde{g}_\lambda\|_{L^2}^2 + \frac{1}{L^2} \|f_\varepsilon\|_{L^2}^2 \|\partial_z \tilde{g}_\lambda\|_{L^2}^2 - \frac{1}{L^2} \|f_\varepsilon\|_{L^2}^2 \|\tilde{g}_\lambda\|_{L^2}^2 \right) \\ &= \|f\|_{L^2}^2 \|\tilde{g}\|_{L^2}^4 \left( \frac{\|\nabla_x f\|_{L^2}^2}{\varepsilon^2} + \|f\|_{L^2}^2 \right) + \frac{1}{L^2} \|f\|_{L^2}^4 \|\tilde{g}\|_{L^2}^2 \left( \frac{\|\partial_z \tilde{g}\|_{L^2}^2}{\lambda^2} - \|\tilde{g}\|_{L^2}^2 \right). \end{aligned}$$

When we take  $\lambda^{-1} = L(N/L)^\beta$ ,  $\varepsilon^{-1} = (N/L)^\beta$ , the interaction energy is equal to

$$L(N/L)^{3\beta} \int V(x_1 - x_2, z_1 - z_2) |f(x_1) \tilde{g}(z_1) f(x_2) \tilde{g}(z_2)|^2 dr_1 dr_2$$

and the kinetic energy is controlled by

$$(N/L)^{2\beta} + \frac{L^2(N/L)^{2\beta}}{L^2} = 2(N/L)^{2\beta}.$$

Since  $L(N/L)^\beta \rightarrow \infty$ , the interaction energy cannot be controlled by the kinetic energy. Therefore, it implies that we should consider the case  $L(N/L)^\beta \leq C$ . On the other hand, to make the limit of  $\tilde{V}_{N,L}$  exist, we should take  $L(N/L)^\beta$  to be a constant or tend to 0. For the case  $L(N/L)^\beta \rightarrow 0$ , we note that the limit of



$\tilde{V}_{N,L}$  equals to 0, which is not sufficient to derive the cubic NLS equation. Hence, we only consider the case  $L(N/L)^\beta \rightarrow 1$  and it works the same for  $L(N/L)^\beta \rightarrow R_0$ .

*Remark 2.3.* A similar argument can also apply to [19] in the focusing setting. To control the interaction energy instead of dropping it like the defocusing case, it also needs an extra condition  $(N\sqrt{\omega})^\beta \omega^{-1/2} \leq C$ .

## 2.2. Focusing energy estimates when $k = 1$ .

**Theorem 2.4.** Assume  $L(N/L)^\beta \rightarrow 1^-$ ,  $\beta < \frac{1}{2}$ , and  $\|V\|_{L_z^\infty L_x^1} \leq \frac{2\alpha}{C_{gn}^4}$  for some  $\alpha \in (0, 1)$ , then let  $c_0 = \min\left(\frac{1-\alpha}{\sqrt{2}}, \frac{1}{2}\right)$ , we have  $\forall C_0 > 0$ , there exists an  $N_0 > 0$  such that

$$(2.18) \quad \langle \psi_{N,L}, (C_0 + 1 + N^{-1}H_{N,L} - 1/L^2)\psi_{N,L} \rangle \geq c_0 \|S_1 \psi_{N,L}\|_{L^2}^2,$$

for all  $N > N_0$  and for all  $\psi_{N,L} \in L_s^2(\Omega_L^{\otimes N})$ .

We divide the proof of Theorem 2.4 into two parts. The first is Theorem 2.5 where the energy bound holds for  $\beta < 1/3$  and the second is Theorem 2.15 which implies Theorem 2.4. In this section, we prove Theorem 2.5 with  $\beta < 1/3$ .

**Theorem 2.5.** Assume  $L(N/L)^\beta \rightarrow 1^-$ ,  $\beta < \frac{1}{3}$ , and  $\|V\|_{L_z^\infty L_x^1} \leq \frac{2\alpha}{C_{gn}^4}$  for some  $\alpha \in (0, 1)$ , then  $\forall C_0 > 0$ , there exists an  $N_0 > 0$  such that

$$(2.19) \quad \langle \psi_{N,L}, (C_0 + 1 + N^{-1}H_{N,L} - 1/L^2)\psi_{N,L} \rangle \geq (1 - \alpha) \|S_1 \psi_{N,L}\|_{L^2}^2,$$

for all  $N > N_0$  and for all  $\psi_{N,L} \in L_s^2(\Omega_L^{\otimes N})$ .

The key of the proof of Theorem 2.5 is the following theorem.

**Theorem 2.6.** Assume  $L(N/L)^\beta \rightarrow 1^-$ ,  $\beta < \frac{1}{3}$ , and  $\|V\|_{L_z^\infty L_x^1} \leq \frac{2\alpha}{C_{gn}^4}$  for some  $\alpha \in (0, 1)$ , define the operator

$$(2.20) \quad H_{ij,\alpha} = \alpha S_i^2 + \alpha S_j^2 + H_{Iij}.$$

Then  $\forall C_0 > 0$ , there exists an  $N_0 > 0$  such that

$$\langle \psi_{N,L}, (2C_0 + H_{12,\alpha})\psi_{N,L} \rangle \geq 0,$$

for all  $N > N_0$  and for all  $\psi_{N,L} \in L_s^2(\Omega_L^{\otimes N})$ .

**Proof of Theorem 2.5 assuming Theorem 2.6.** Using formula (2.7) and the symmetry of  $\psi_{N,L}$ , we have

$$\begin{aligned} & \langle \psi_{N,L}, (C_0 + 1 + N^{-1}H_{N,L} - 1/L^2)\psi_{N,L} \rangle \\ &= \frac{1}{N(N-1)} \sum_{1 \leq i < j \leq N} \langle \psi_{N,L}, (2C_0 + H_{ij})\psi_{N,L} \rangle \\ &= \frac{1}{2} \langle \psi_{N,L}, (2C_0 + H_{12})\psi_{N,L} \rangle \\ &= \frac{1}{2} \langle \psi_{N,L}, (2C_0 + H_{12,\alpha})\psi_{N,L} \rangle + \frac{1-\alpha}{2} \langle \psi_{N,L}, (S_1^2 + S_2^2)\psi_{N,L} \rangle \\ &\geq (1-\alpha) \|S_1 \psi_{N,L}\|_{L^2}^2. \end{aligned}$$

□

Next, we turn our attention onto the proof of Theorem 2.6. Under the assumption  $L(N/L)^\beta \rightarrow 1^-$ , the renormalized kinetic energy can control the potential energy.

**Lemma 2.7.** Assume  $L(N/L)^\beta \rightarrow 1^-$ ,  $M \geq 1$ , then for all  $\psi_{M,L} \in L_s^2(\Omega_L^{\otimes M})$  with  $\|\psi_{M,L}\|_{L^2} = 1$ , we have

$$(2.21) \quad L \int_{\Omega_L^2} |V_{N,L}(r_1 - r_2)| \rho_{M,L}(r_1) \rho_{M,L}(r_2) dr_1 dr_2 \leq C_{gn}^4 \|V\|_{L_z^\infty L_x^1} \langle S_1^2 \psi_{M,L}, \psi_{M,L} \rangle,$$

where density function  $\rho_{M,L}(r_1) := \int \cdots \int |\psi_{M,L}|^2(r_1, \dots, r_M) dr_2 \cdots dr_M$ . Especially, if  $\psi_{M,L} = \phi_L^{\otimes M}$  with  $\|\phi_L\|_{L^2} = 1$ , then

$$(2.22) \quad L \int_{\Omega_L^2} |V_{N,L}(r_1 - r_2)| \phi_L(r_1) \phi_L(r_2) dr_1 dr_2 \leq C_{gn}^4 \|V\|_{L_z^\infty L_x^1} \langle S_1^2 \phi_L, \phi_L \rangle.$$

*Proof.* Using Cauchy-Schwarz inequality and Young's convolution inequality, we get

$$(2.23) \quad L \int_{\Omega_L^2} |V_{N,L}(r_1 - r_2)| \rho_{M,L}(r_1) \rho_{M,L}(r_2) dr_1 dr_2 \leq L \|V_{N,L} * \rho_{M,L}\|_{L_z^\infty L_x^2} \|\rho_{M,L}\|_{L_z^1 L_x^2} \\ \leq L \|V_{N,L}\|_{L_z^\infty L_x^1} \|\rho_{M,L}\|_{L_z^1 L_x^2}^2.$$

For  $\|\rho_{M,L}\|_{L_z^1 L_x^2}^2$  on the right-hand side of (2.23), we use 2D inhomogeneous Gagliardo-Nirenberg inequality (1.10),

$$(2.24) \quad \|\rho_{M,L}\|_{L_z^1 L_x^2}^2 = \|\sqrt{\rho_{M,L}}\|_{L_z^2 L_x^4}^4 \leq C_{gn}^4 \|\sqrt{\rho_{M,L}}\|_{L_z^2 L_x^2}^2 \|\sqrt{1 - \Delta_x} \sqrt{\rho_{M,L}}\|_{L_z^2 L_x^2}^2 \\ = C_{gn}^4 \|\sqrt{\rho_{M,L}}\|_{L_z^2 L_x^2}^2 \left( \|\sqrt{\rho_{M,L}}\|_{L_z^2 L_x^2}^2 + \|\nabla_x \sqrt{\rho_{M,L}}\|_{L_z^2 L_x^2}^2 \right).$$

By the Hoffman-Ostenhof inequality (A.1), we have

$$(2.25) \quad \|\nabla_x \sqrt{\rho_{M,L}}\|_{L^2}^2 \leq \langle -\Delta_{x_1} \psi_{M,L}, \psi_{M,L} \rangle.$$

Now, putting (2.25) into (2.24) and (2.24) into (2.23), we have

$$(2.26) \quad L \int_{\Omega_L^2} |V_{N,L}(r_1 - r_2)| \rho_{M,L}(r_1) \rho_{M,L}(r_2) dr_1 dr_2 \leq C_{gn}^4 L \|V_{N,L}\|_{L_z^\infty L_x^1} \langle (1 - \Delta_{x_1}) \psi_{M,L}, \psi_{M,L} \rangle.$$

Noting that  $\|V_{N,L}\|_{L_z^\infty L_x^1} = (N/L)^\beta \|V\|_{L_z^\infty L_x^1}$ , with the assumption  $L(N/L)^\beta \rightarrow 1^-$ , we conclude that

$$(2.27) \quad L \int_{\Omega_L^2} |V_{N,L}(r_1 - r_2)| \rho_{M,L}(r_1) \rho_{M,L}(r_2) dr_1 dr_2 \leq C_{gn}^4 \|V\|_{L_z^\infty L_x^1} \langle (1 - \Delta_{x_1}) \psi_{M,L}, \psi_{M,L} \rangle.$$

Since  $S_1^2 \geq (1 - \Delta_{x_1})$ , we arrive at the estimate (2.21) from estimate (2.27).  $\square$

The following lemma is used to estimate the two-body interaction energy by a one-body term.

**Lemma 2.8.** *If  $V \in C_c^\infty(\mathbb{T}^2 \times \mathbb{R})$  and has a positive Fourier transform  $\widehat{V} \geq 0$ , then for all real function  $\eta \in L^1(\mathbb{T}^2 \times \mathbb{R})$*

$$(2.28) \quad \sum_{1 \leq j < k \leq N} V(r_j - r_k) \geq \frac{1}{(2\pi)^3} \sum_{j=1}^N \eta * V(r_j) - \frac{1}{2(2\pi)^6} \int_{\mathbb{T}^2 \times \mathbb{R}} \int_{\mathbb{T}^2 \times \mathbb{R}} V(r_1 - r_2) \eta(r_1) \eta(r_2) dr_1 dr_2 - \frac{N}{2} V(0).$$

*Proof.* With the Fourier inversion formula,

$$(2.29) \quad \sum_{1 \leq j < k \leq N} V(r_j - r_k) \\ = \int \sum_{1 \leq j < k \leq N} e^{i\xi \cdot (r_j - r_k)} \widehat{V}(\xi) d\xi \\ = \frac{1}{2} \int \left| \sum_{j=1}^N e^{i\xi \cdot r_j} - \widehat{\eta}(\xi) \right|^2 \widehat{V}(\xi) d\xi + \frac{1}{(2\pi)^3} \sum_{j=1}^N \eta * V(r_j) \\ - \frac{1}{2(2\pi)^6} \int_{\mathbb{T}^2 \times \mathbb{R}} \int_{\mathbb{T}^2 \times \mathbb{R}} V(r_1 - r_2) \eta(r_1) \eta(r_2) dr_1 dr_2 - \frac{N}{2} V(0) \\ \geq \frac{1}{(2\pi)^3} \sum_{j=1}^N \eta * V(r_j) - \frac{1}{2(2\pi)^6} \int_{\mathbb{T}^2 \times \mathbb{R}} \int_{\mathbb{T}^2 \times \mathbb{R}} V(r_1 - r_2) \eta(r_1) \eta(r_2) dr_1 dr_2 - \frac{N}{2} V(0),$$

where  $\xi = (n_1, n_2, \tau)$  and  $\int \cdot d\xi$  is short for  $\int_{\mathbb{R}} \sum_{n_1, n_2} \cdot d\tau$ .  $\square$

*Remark 2.9.* In our setting, the integral region is  $\Omega_L$ . To use Lemma 2.8,  $V_{N,L}$  should be understood as the periodic extension in the  $x$ -direction and zero extension in the  $z$ -direction of the rescaled  $V$  which is compactly supported on  $\Omega$ . That is,  $V_{N,L} \in C_c^\infty(\mathbb{T}^2 \times \mathbb{R})$ . Similarly,  $\rho_{N,L}(r)$  and  $\psi_{N,L}$  should be seen as  $\rho_{N,L}(r) 1_{\Omega_L}(r) \in L^1(\mathbb{T}^2 \times \mathbb{R})$  and  $\psi_{N,L} 1_{\Omega_L^{\otimes N}}$  respectively.

In case  $\widehat{V}_{N,L} \geq 0$ , if we take  $V = V_{N,L}$  and  $\eta = (2\pi)^3 N \rho_{N,L}$ , we obtain

$$\begin{aligned}
 (2.30) \quad & \sum_{1 \leq j < k \leq N} \langle \psi_{N,L}, V_{N,L}(r_j - r_k) \psi_{N,L} \rangle \\
 & \geq N \sum_{j=1}^N \int \rho_{N,L} * V_{N,L}(r_j) |\psi_{N,L}|^2(\mathbf{r}_N) d\mathbf{r}_N - \frac{N^2}{2} \langle \rho_{N,L} * V_{N,L}, \rho_{N,L} \rangle - \frac{N}{2} V_{N,L}(0) \\
 & = \frac{N^2}{2} \int \rho_{N,L}(r_1) \rho_{N,L}(r_2) V_{N,L}(r_1 - r_2) dr_1 dr_2 - \frac{N(N/L)^{3\beta} V(0)}{2} \\
 & \geq -\frac{N(N/L)^{3\beta} V(0)}{2},
 \end{aligned}$$

where we have used

$$\int \rho_{N,L}(r_1) \rho_{N,L}(r_2) V_{N,L}(r_1 - r_2) dr_1 dr_2 = (2\pi)^6 \int |\widehat{\rho}_{N,L}(\xi)|^2 \widehat{V}_{N,L}(\xi) d\xi$$

and  $\widehat{V}_{N,L} \geq 0$  in the last inequality.

By estimate (2.30), we have

$$\langle \psi_{N,L}, (2C_0 + H_{12,\alpha}) \psi_{N,L} \rangle \geq 2C_0 - \frac{(N/L)^{3\beta} V(0)}{(N-1)/L} \geq 0,$$

as long as  $\beta < 1/3$  and  $N/L$  is large enough.

Hence, we have established Theorem 2.6 if  $\widehat{V}_{N,L} \geq 0$ . Next, we will use Lemma 2.8 to deal with a general interaction function  $V$ .

**Proof of Theorem 2.6.** For general  $V$ , we consider  $N = 2M$  particles which we split into two groups of  $M$ . For the case  $N = 2M + 1$ , the proof works the same if we split the system into two groups of  $M$  and  $M + 1$ . We denote the first  $M$  variables by  $r_1, \dots, r_M$  and the others by  $r'_1 = r_{M+1}, \dots, r'_M = r_{2M}$ . We decompose  $V_{N,L} = V_{N,L}^+ - V_{N,L}^-$  on the Fourier side where  $\widehat{V_{N,L}^+} = (\widehat{V_{N,L}})_+ \geq 0$  and  $\widehat{V_{N,L}^-} = (\widehat{V_{N,L}})_- \geq 0$ . By its symmetry in the  $2M$  variables, we rewrite

$$\begin{aligned}
 & \frac{1}{2} \langle \psi_{N,L}, H_{12} \psi_{N,L} \rangle \\
 & = \frac{L}{2M(2M-1)} \left\langle \psi_{2M,L}, \sum_{1 \leq j < k \leq 2M} V_{N,L}(r_j - r_k) \psi_{2M,L} \right\rangle \\
 & = \frac{L}{M(M-1)} \left\langle \psi_{2M,L}, \sum_{1 \leq j < k \leq M} V_{N,L}^+(r_j - r_k) \psi_{2M,L} \right\rangle \\
 & \quad + \frac{L}{M(M-1)} \left\langle \psi_{2M,L}, \sum_{1 \leq l < m \leq M} V_{N,L}^-(r'_l - r'_m) \psi_{2M,L} \right\rangle - \frac{L}{M^2} \left\langle \psi_{2M,L}, \sum_{j=1}^M \sum_{l=1}^M V_{N,L}^-(r_j - r'_l) \psi_{2M,L} \right\rangle.
 \end{aligned}$$

This means that

$$(2.31) \quad \frac{1}{2} \langle \psi_{2M,L}, H_{12} \psi_{2M,L} \rangle = \langle \psi_{2M,L}, I_{M,L} \psi_{2M,L} \rangle,$$

where

$$\begin{aligned}
 (2.32) \quad I_{M,L} & = \frac{L}{M(M-1)} \sum_{1 \leq j < k \leq M} V_{N,L}^+(r_j - r_k) + \frac{L}{M(M-1)} \sum_{1 \leq l < m \leq M} V_{N,L}^-(r'_l - r'_m) \\
 & \quad - \frac{L}{M^2} \sum_{j=1}^M \sum_{l=1}^M V_{N,L}^-(r_j - r'_l).
 \end{aligned}$$

Then we have

$$(2.33) \quad \langle \psi_{2M,L}, (2C_0 + H_{12,\alpha}) \psi_{2M,L} \rangle = \langle \psi_{2M,L}, (2C_0 + 2I_{M,L} + \alpha(S_1^2 + S_2^2)) \psi_{2M,L} \rangle.$$

Thus, in order to bound the interaction  $2C_0 + H_{12,\alpha}$  from below, it suffices to consider  $I_{M,L}$ . We fix the variables  $r'_1, \dots, r'_M$  in the second group. For simplicity, we use the notation  $\langle \cdot, \cdot \rangle_{\mathbf{r}_M}$  to denote the integral only in the variables  $\mathbf{r}_M := (r_1, \dots, r_M)$ . We denote the one-particle density by

$$(2.34) \quad \rho_{M,L}(r, \mathbf{r}'_M) := \int_{\Omega_L} \cdots \int_{\Omega_L} |\psi_{2M,L}|^2(r, r_2, \dots, r_M, \mathbf{r}'_M) dr_2 \cdots dr_M.$$

Our goal is to get

$$(2.35) \quad \langle \psi_{2M,L}, (2C_0 + 2I_{M,L} + \alpha(S_1^2 + S_2^2)) \psi_{2M,L} \rangle_{\mathbf{r}_M} \\ \geq (2\alpha - C_{gn}^4 \|V\|_{L_z^\infty L_x^1}) \langle \psi_{2M,L}, S_1^2 \psi_{2M,L} \rangle_{\mathbf{r}_M} + \int \rho_{M,L}(r, \mathbf{r}'_M) dr \left( 2C_0 - \frac{LV_{N,L}^+(0) + LV_{N,L}^-(0)}{(M-1)} \right).$$

First, we may assume  $\int \rho_{M,L}(r, \mathbf{r}'_M) dr > 0$ . The case  $\int \rho_{M,L}(r, \mathbf{r}'_M) dr = 0$  is easier and will be presented later. By using Lemma 2.8 with

$$V = V_{N,L}^+, \quad \eta = \frac{(2\pi)^3 M \rho_{M,L}(r, \mathbf{r}'_M)}{\int \rho_{M,L}(r, \mathbf{r}'_M) dr},$$

we have

$$(2.36) \quad \langle \psi_{2M,L}, I_{M,L} \psi_{2M,L} \rangle_{\mathbf{r}_M} \\ \geq \frac{ML}{2(M-1) \int \rho_{M,L}(r, \mathbf{r}'_M) dr} \int V_{N,L}^+(r_1 - r_2) \rho_{M,L}(r_1, \mathbf{r}'_M) \rho_{M,L}(r_2, \mathbf{r}'_M) dr_1 dr_2 \\ - \frac{LV_{N,L}^+(0) \int \rho_{M,L}(r, \mathbf{r}'_M) dr}{2(M-1)} + \frac{L}{M(M-1)} \sum_{1 \leq l < m \leq M} V_{N,L}^-(r'_l - r'_m) \int \rho_{M,L}(r, \mathbf{r}'_M) dr \\ - \frac{L}{M} \sum_{l=1}^M \rho_{M,L} * V_{N,L}^-(r'_l).$$

Next we use again Lemma 2.8 with

$$V = V_{N,L}^-, \quad \eta = \frac{(2\pi)^3 (M-1) \rho_{M,L}(r, \mathbf{r}'_M)}{\int \rho_{M,L}(r, \mathbf{r}'_M) dr},$$

and obtain

$$(2.37) \quad \frac{L}{M(M-1)} \sum_{1 \leq l < m \leq M} V_{N,L}^-(r'_l - r'_m) \int \rho_{M,L}(r, \mathbf{r}'_M) dr - \frac{L}{M} \sum_{l=1}^M \rho_{M,L} * V_{N,L}^-(r'_l) \\ \geq - \frac{(M-1)}{2M} \frac{L}{\int \rho_{M,L}(r, \mathbf{r}'_M) dr} \int V_{N,L}^-(r_1 - r_2) \rho_{M,L}(r_1, \mathbf{r}'_M) \rho_{M,L}(r_2, \mathbf{r}'_M) dr_1 dr_2 \\ - \frac{LV_{N,L}^-(0) \int \rho_{M,L}(r, \mathbf{r}'_M) dr}{2(M-1)}.$$

Thus, we have

$$(2.38) \quad \langle \psi_{2M,L}, I_{M,L} \psi_{2M,L} \rangle_{\mathbf{r}_M} \\ \geq \int \rho_{M,L}(r, \mathbf{r}'_M) dr \int \frac{L}{2} V_{N,L}(r_1 - r_2) \frac{\rho_{M,L}(r_1, \mathbf{r}'_M)}{\int \rho_{M,L}(r, \mathbf{r}'_M) dr} \frac{\rho_{M,L}(r_2, \mathbf{r}'_M)}{\int \rho_{M,L}(r, \mathbf{r}'_M) dr} dr_1 dr_2 \\ - \int \rho_{M,L}(r, \mathbf{r}'_M) dr \frac{LV_{N,L}^+(0) + LV_{N,L}^-(0)}{2(M-1)}.$$

By Lemma 2.7, we obtain

$$(2.39) \quad \langle \psi_{2M,L}, (2C_0 + 2I_{M,L} + \alpha(S_1^2 + S_2^2)) \psi_{2M,L} \rangle_{\mathbf{r}_M} \\ \geq (2\alpha - C_{gn}^4 \|V\|_{L_z^\infty L_x^1}) \langle \psi_{2M,L}, S_1^2 \psi_{2M,L} \rangle_{\mathbf{r}_M} + \int \rho_{M,L}(r, \mathbf{r}'_M) dr \left( 2C_0 - \frac{LV_{N,L}^+(0) + LV_{N,L}^-(0)}{(M-1)} \right).$$

We arrive at the estimate (2.35) for the case  $\int \rho_{M,L}(r, \mathbf{r}'_M) dr > 0$ .

Next, if  $\int \rho_{M,L}(r, \mathbf{r}'_M) dr = 0$ , we can deduce that  $\rho_{M,L}(r, \mathbf{r}'_M) = 0$  due to the nonnegativity and smoothness of  $\rho_{M,L}$ . Then, we have

$$\begin{aligned}
(2.40) \quad & \langle \psi_{2M,L}, I_{M,L} \psi_{2M,L} \rangle_{\mathbf{r}_M} \\
&= \frac{L}{M(M-1)} \sum_{1 \leq j < k \leq M} \int V_{N,L}^+(r_j - r_k) |\psi_{M,L}(\mathbf{r}_M, \mathbf{r}'_M)|^2 d\mathbf{r}_M \\
&\quad + \frac{L}{M(M-1)} \sum_{1 \leq l < m \leq M} V_{N,L}^-(r'_l - r'_m) \int \rho_{M,L}(r, \mathbf{r}'_M) dr \\
&\quad - \frac{L}{M^2} \sum_{j=1}^M \sum_{l=1}^M \int V_{N,L}^-(r_j - r'_l) \rho_{M,L}(r_j, \mathbf{r}'_M) dr_j \\
&= I + II + III.
\end{aligned}$$

Since  $V_{N,L}^+$  has a positive Fourier transform, we have  $I \geq 0$ . By the fact that  $\rho_{M,L}(r, \mathbf{r}'_M) = 0$ , we obtain  $II = III = 0$ . That is, the estimate (2.35) still holds.

Hence, when  $\|V\|_{L_z^\infty L_x^1} \leq \frac{2\alpha}{C_{gn}^4}$ , we have

$$\begin{aligned}
(2.41) \quad & \langle \psi_{2M,L}, (2C_0 + H_{12,\alpha}) \psi_{2M,L} \rangle = \langle \psi_{2M,L}, (2C_0 + 2I_{M,L} + \alpha(S_1^2 + S_2^2)) \psi_{2M,L} \rangle \\
& \geq 2C_0 - \frac{L(N/L)^{3\beta} (V^+(0) + V^-(0))}{2(M-1)}.
\end{aligned}$$

Since  $\beta < \frac{1}{3}$ , there exists an  $N_0 > 0$  such that

$$(2.42) \quad \langle \psi_{2M,L}, (2C_0 + H_{12,\alpha}) \psi_{2M,L} \rangle \geq 0,$$

for all  $N = 2M \geq N_0$ . □

The proof also works for the case in which we put  $\mathbb{R}^2$  in the  $x$ -direction and hence the main theorem works the same for  $\mathbb{R}^2$  with  $\beta < 1/3$ .

**2.3. Bootstrapping argument.** <sup>8</sup> In [55], Lewin, Nam, and Rougerie can improve the index  $\beta$  by a bootstrapping argument for 2D case. Here, such a method also works for  $\beta < 1/2$  with a starting point  $\eta_0 \leq (22\beta - 4)/5$  in our setting. Let us define

$$(2.43) \quad E_{N,L} := \inf_{\|\psi_{N,L}\|_{L^2}=1} \langle \psi_{N,L}, (1 + N^{-1}H_{N,L} - 1/L^2) \psi_{N,L} \rangle,$$

$$(2.44) \quad E_{N,L,\varepsilon} := \inf_{\|\psi_{N,L}\|_{L^2}=1} \langle \psi_{N,L}, \left( 1 + N^{-1}H_{N,L} - 1/L^2 - \varepsilon N^{-1} \sum_{j=1}^N S_j^2 \right) \psi_{N,L} \rangle,$$

where  $E_{N,L}$  denotes the many-body ground state energy per particle. From the definition (2.44), estimate (2.19) in Theorem 2.5, is equivalent to prove  $E_{N,L,1-\alpha} \geq -C_0$  for  $N \geq N_0$ . Indeed, if  $E_{N,L,1-\alpha} \geq -C_0$  for  $N \geq N_0$ , it means that

$$C_0 + 1 + N^{-1}H_{N,L} - 1/L^2 - (1 - \alpha)S_1^2 \geq 0,$$

for  $N \geq N_0$ , which is the estimate (2.19).

Thus, our goal is to bound  $E_{N,L,\varepsilon}$  from below. We note that  $E_{N,L,\varepsilon} = (1 - \varepsilon)E_{N,L}^\varepsilon$ , where  $E_{N,L}^\varepsilon$  is the ground state energy with interaction function  $V^\varepsilon = (1 - \varepsilon)^{-1}V$ . So we only need to deal with  $E_{N,L}^\varepsilon$  or  $E_{N,L}$ . A main tool is the finite-dimensional quantum de Finetti theorem (Lemma 2.11). Then we can give a lower bound on the Hamiltonian energy in Lemma 2.13, that is,

$$(2.45) \quad \left\langle \psi_{N,L}, (1 + N^{-1}H_{N,L} - 1/L^2) \psi_{N,L} \right\rangle \geq C(V, N, L, \lambda_x, \lambda_z, \|S_1 \psi_{N,L}\|_{L^2}, \|S_1 S_2 \psi_{N,L}\|_{L^2}),$$

where  $\lambda_x$  and  $\lambda_z$  are cut-off parameters.

---

<sup>8</sup>The finite dimensional quantum de Finetti theorem is only used in this section. In Section 2.2, we have already reached  $\beta < 1/3$  without it. Hence, the main theorem works the same for  $\mathbb{R}^2$  with  $\beta < 1/3$ .

Subsequently in Lemma 2.14, we will control  $\|S_1\psi_{N,L}\|_{L^2}$  and  $\|S_1S_2\psi_{N,L}\|_{L^2}$  for the ground state  $\psi_{N,L}$ . More precisely,

$$(2.46) \quad \text{Tr} \left( S_1^2 \gamma_{N,L}^{(1)} \right) \lesssim C_{up} \frac{1 + |E_{N,L,\theta}|}{\theta},$$

$$(2.47) \quad \text{Tr} \left( S_1^2 S_2^2 \gamma_{N,L}^{(2)} \right) \lesssim C_{up}^2 \left( \frac{1 + |E_{N,L,\theta}|}{\theta} \right)^2,$$

where  $C_{up}$  is an upper bound constant defined by (2.50).

With Lemma 2.13 and 2.14, we arrive at a closed control relationship, namely,

$$|E_{N,L}| \leq C(V, N, L, \lambda_x, \lambda_z, |E_{N,L,\theta}|).$$

Thus we can use the bootstrapping argument to bound  $|E_{N,L,\varepsilon}|$  as long as there exists a starting point.

Now, we present the above procedure in detail. First, we take  $\psi_{N,L} = \phi_L^{\otimes N}$  with  $\|\phi_L\|_{L^2} = 1$  and obtain the NLS energy functional

$$(2.48) \quad \begin{aligned} \mathcal{E}_{N,L}(\phi_L) &= \langle \phi_L^{\otimes N}, (1 + N^{-1}H_{N,L} - 1/L^2) \phi_L^{\otimes N} \rangle \\ &= \frac{1}{2} \langle \phi_L^{\otimes 2}, H_{12} \phi_L^{\otimes 2} \rangle = \langle S_1^2 \phi_L, \phi_L \rangle + \frac{L}{2} \int_{\Omega_L^2} V_{N,L}(r_1 - r_2) |\phi_L(r_1) \phi_L(r_2)|^2 dr_1 dr_2. \end{aligned}$$

Define

$$(2.49) \quad e_{N,L} := \inf_{\|\phi_L\|_{L^2}=1} \mathcal{E}_{N,L}(\phi_L),$$

where  $e_{N,L}$  stands for the ground state energy of the NLS energy functional. From the above definition, we know  $e_{N,L} \geq E_{N,L} \geq E_{N,L,\varepsilon}$ . To bound  $E_{N,L,\varepsilon}$  from below, it is necessary to bound  $e_{N,L}$  from below. Here, we first give a sufficient condition to bound  $e_{N,L}$  as follows.

**Lemma 2.10.** *Assume  $L(N/L)^\beta \rightarrow 1^-$  and  $\|V\|_{L_z^\infty L_x^1} \leq \frac{2}{C_{gn}^4}$ , then  $0 \leq e_{N,L} \leq C_{up}$ , where*

$$(2.50) \quad C_{up} = 1 + \frac{C_{gn}^4 \|V\|_{L_z^\infty L_x^1}}{2}.$$

*Proof.* For the lower bound, it suffices to prove  $\mathcal{E}_{N,L}(\phi_L) \geq 0$ . From estimate (2.22) in Lemma 2.7, we have

$$(2.51) \quad \begin{aligned} \mathcal{E}_{N,L}(\phi_L) &= \langle S_1^2 \phi_L, \phi_L \rangle + \frac{L}{2} \int_{\Omega_L^2} V_{N,L}(r_1 - r_2) |\phi_L(r_1) \phi_L(r_2)|^2 dr_1 dr_2 \\ &\geq \langle S_1^2 \phi_L, \phi_L \rangle - \frac{C_{gn}^4 \|V\|_{L_z^\infty L_x^1}}{2} \langle S_1^2 \phi_L, \phi_L \rangle \geq 0, \end{aligned}$$

as long as  $\|V\|_{L_z^\infty L_x^1} \leq \frac{2}{C_{gn}^4}$ .

For the upper bound, we use estimate (2.22) again and obtain

$$(2.52) \quad \begin{aligned} \mathcal{E}_{N,L}(\phi_L) &= \langle S_1^2 \phi_L, \phi_L \rangle + \frac{L}{2} \int_{\Omega_L^2} V_{N,L}(r_1 - r_2) |\phi_L(r_1) \phi_L(r_2)|^2 dr_1 dr_2 \\ &\leq \langle S_1^2 \phi_L, \phi_L \rangle + \frac{C_{gn}^4 \|V\|_{L_z^\infty L_x^1}}{2} \langle S_1^2 \phi_L, \phi_L \rangle. \end{aligned}$$

When we take  $\phi_L(x, z) = \cos_L(z)$ , which is the  $L^2$  normalized ground state wave function of  $S_1^2$ , then  $\langle S_1^2 \phi_L, \phi_L \rangle = 1$ . Hence, we have

$$(2.53) \quad e_{N,L} \leq 1 + \frac{C_{gn}^4 \|V\|_{L_z^\infty L_x^1}}{2}.$$

□

To establish the lower bound estimate for  $E_{N,L}$ , we need the finite-dimensional quantum de Finetti theorem. We define the Littlewood-Paley projectors (eigenspace projector) by

$$(2.54) \quad P_{x_j, m} = \chi_{\{m\}}(-\Delta_{x_j}), \quad m \geq 0,$$

$$(2.55) \quad P_{x_j, \leq \lambda} = \chi_{[0, \lambda^2]}(-\Delta_{x_j}), \quad P_{z_j, \leq \lambda} = \chi_{[1/L^2, \lambda^2/L^2]}(-\partial_{z_j}^2),$$

$$(2.56) \quad P_{x_j, > \lambda} = I - P_{x_j, \leq \lambda}, \quad P_{z_j, > \lambda} = I - P_{z_j, \leq \lambda}.$$

From the definition, we notice that  $P_{z_j, \leq \lambda}$  and  $P_{z_j, > \lambda}$  are  $L$  dependent. However, we omit it for simplicity.

**Lemma 2.11** (Finite-dimensional quantum de Finetti[54]). *Assume  $\left\{\gamma_{N,L}^{(k)}\right\}_{k=1}^N$  is the marginal density generated by an  $N$ -body wave function  $\psi_{N,L} \in L^2_s(\Omega_L^N)$  and  $P_L$  be a finite-rank orthogonal projector with*

$$\dim(P_L(L^2(\Omega_L))) = d < \infty,$$

*where  $d$  can be independent of  $L$ . Then, there is a positive Borel measure  $d\mu_{N,L}$  supported on the unit sphere  $SP_L(L^2(\Omega_L))$  such that*

$$(2.57) \quad \text{Tr} \left| P_L^{\otimes 2} \gamma_{N,L}^{(2)} P_L^{\otimes 2} - \int_{S(P_L(L^2(\Omega_L)))} |\phi_L^{\otimes 2}\rangle \langle \phi_L^{\otimes 2}| d\mu_{N,L}(\phi_L) \right| \leq \frac{8d}{N}.$$

Moreover, we will need operator inequalities for two-body interaction as follows.

**Lemma 2.12.** *Assume  $L(N/L)^\beta \rightarrow 1^-$ . For  $\delta \in (0, 1)$ , the multiplication operator  $V_{N,L}(r_1 - r_2)$  on  $L^2(\Omega_L^{\otimes 2})$  satisfies*

$$(2.58) \quad L|V_{N,L}(r_1 - r_2)| \leq C_\delta (N/L)^\delta \|V\|_{L_z^\infty L_x^{1+\delta}} (1 - \Delta_{x_1}),$$

$$(2.59) \quad L|V_{N,L}(r_1 - r_2)| \leq C_\delta \|V\|_{L_z^\infty L_x^1} (1 - \Delta_{x_1})^{1/2+\delta} (1 - \Delta_{x_2})^{1/2+\delta},$$

$$(2.60) \quad \|\langle \nabla_{x_1} \rangle^{-1} \langle \nabla_{x_2} \rangle^{-1} V(r_1 - r_2) \langle \nabla_{x_1} \rangle^{-1} \langle \nabla_{x_2} \rangle^{-1}\|_{op} \leq C \|V\|_{L_z^\infty L_x^1},$$

$$(2.61) \quad S_1^2 L V_{N,L}(r_1 - r_2) + L V_{N,L}(r_1 - r_2) S_1^2 \geq -C_\delta(V) (N/L)^{\beta+\delta} S_1^2 S_2^2,$$

where  $C_\delta(V)$  is dependent on  $V$ .

*Proof.* For smoothness of presentation, we put the proof in the Appendix. □

Now, along with Lemma 2.11, we can establish the following lower bound estimate.

**Lemma 2.13.** *Assume  $L(N/L)^\beta \rightarrow 1^-$  and  $\|V\|_{L_z^\infty L_x^1} \leq \frac{2}{C_{gn}^4}$ . Then for every  $\delta \in (0, 1/2)$ , there exists a constant  $C_\delta > 0$  such that for all  $N \geq 2$ ,  $\lambda_z \geq 1$ ,  $\lambda_x \geq 0$  and for all wave functions  $\psi_{N,L}$*

$$\begin{aligned} & \left\langle \psi_{N,L}, \left(1 + N^{-1} H_{N,L} - 1/L^2\right) \psi_{N,L} \right\rangle \\ & \gtrsim -C_\delta \|V\|_{L_z^\infty L_x^1} (1 + \lambda_x^2)^{1+2\delta} \frac{d}{N} \\ & \quad - \frac{C_\delta \|V\|_{L_z^\infty L_x^1}}{\min\{1 + \lambda_x^2, 1 + (\lambda_z^2 - 1)/L^2\}^{1/4-\delta/2}} \left( \text{Tr} \left( S_1^2 \gamma_{N,L}^{(1)} \right) \right)^{1/4-\delta/2} \left( \text{Tr} \left( S_1^2 S_2^2 \gamma_{N,L}^{(2)} \right) \right)^{1/2+\delta}, \end{aligned}$$

where  $d = \dim(P_{x, \leq \lambda_x} P_{z, \leq \lambda_z} L^2(\Omega_L))$ .

*Proof.* For simplicity, we adopt the notation

$$\begin{aligned} P &:= P_{x, \leq \lambda_x} P_{z, \leq \lambda_z}, \quad Q = 1 - P, \\ P^j &:= P_{x_j, \leq \lambda_x} P_{z_j, \leq \lambda_z}, \quad Q^j := 1 - P^j, \\ P^{(k)} &:= \prod_{j=1}^k P_{x_j, \leq \lambda_x}, \quad P_{z_j, \leq \lambda_z} = \prod_{j=1}^k P^j. \end{aligned}$$

Recall  $H_{12} = S_1^2 + S_2^2 + L V_{N,L}(r_1 - r_2)$  and  $S_1^2 = 1 - \Delta_{r_1} - 1/L^2$ . Using Lemma 2.11, we write

$$\begin{aligned} \left\langle \psi_{N,L}, \left(1 + N^{-1} H_{N,L} - 1/L^2\right) \psi_{N,L} \right\rangle &= \frac{1}{2} \langle \psi_{N,L}, H_{12} \psi_{N,L} \rangle \\ &\geq \langle \psi_{N,L}, (1 - \Delta_{x_1} + \frac{1}{2} L V_{N,L}(r_1 - r_2)) \psi_{N,L} \rangle \\ &= I + II + III, \end{aligned}$$

where

$$(2.62) \quad I = \int \langle \phi_L^{\otimes 2}, (1 - \Delta_{x_1} + \frac{1}{2}LV_{N,L}(r_1 - r_2))\phi_L^{\otimes 2} \rangle d\mu_{N,L}(\phi_L)$$

$$(2.63) \quad II = Tr \left( (1 - \Delta_{x_1} + \frac{1}{2}LV_{N,L}(r_1 - r_2)) \left[ \gamma_{N,L}^{(2)} - P^{(2)}\gamma_{N,L}^{(2)}P^{(2)} \right] \right)$$

$$(2.64) \quad III = Tr \left( (1 - \Delta_{x_1} + \frac{1}{2}LV_{N,L}(r_1 - r_2)) \left[ P^{(2)}\gamma_{N,L}^{(2)}P^{(2)} - \int_{S(P(L^2(\Omega_L)))} |\phi_L^{\otimes 2}\rangle \langle \phi_L^{\otimes 2}| d\mu_{N,L}(\phi_L) \right] \right)$$

We only need to bound these terms from below.

We first handle  $I$ . By estimate (2.27) with  $\psi_{M,L} = \phi_L^{\otimes 2}$ , we have

$$\langle \phi_L^{\otimes 2}, (1 - \Delta_{x_1} + \frac{1}{2}LV_{N,L}(r_1 - r_2))\phi_L^{\otimes 2} \rangle \geq 0.$$

Noting that  $\mu_{N,L}$  is a positive measure, we obtain  $I \geq 0$ . Hence we can discard it.

We now deal with  $II$  and rewrite

$$(2.65) \quad II = Tr \left( (1 - \Delta_{x_1}) \left[ \gamma_{N,L}^{(2)} - P^{(2)}\gamma_{N,L}^{(2)}P^{(2)} \right] \right) + \frac{1}{2}Tr \left( LV_{N,L}(r_1 - r_2) \left[ \gamma_{N,L}^{(2)} - P^{(2)}\gamma_{N,L}^{(2)}P^{(2)} \right] \right) \\ = II_K + II_V.$$

Since  $(1 - \Delta_{x_1}) \geq (1 - \Delta_{x_1})P^{(2)}$ , with Lemma A.7, we obtain

$$(2.66) \quad II_K = Tr \left( (1 - \Delta_{x_1}) \left[ \gamma_{N,L}^{(2)} - P^{(2)}\gamma_{N,L}^{(2)}P^{(2)} \right] \right) \\ = \langle [(1 - \Delta_{x_1}) - (1 - \Delta_{x_1})P^{(2)}] \psi_{N,L}, \psi_{N,L} \rangle \geq 0.$$

For  $II_V$ , we expand

$$(2.67) \quad 2 \left( LV_{N,L}(r_1 - r_2) - P^{(2)}LV_{N,L}(r_1 - r_2)P^{(2)} \right) \\ = LV_{N,L}(r_1 - r_2)(1 - P^{(2)}) + (1 - P^{(2)})LV_{N,L}(r_1 - r_2) \\ + (1 - P^{(2)})LV_{N,L}(r_1 - r_2)P^{(2)} + P^{(2)}LV_{N,L}(r_1 - r_2)(1 - P^{(2)}).$$

By Lemma A.7 and Cauchy-Schwarz inequality,

$$(2.68) \quad \frac{1}{2} \left| Tr \left( (LV_{N,L}(r_1 - r_2)(1 - P^{(2)}) + (1 - P^{(2)})LV_{N,L}(r_1 - r_2)) \gamma_{N,L}^{(2)} \right) \right| \\ = \frac{1}{2} \left| \langle LV_{N,L}(r_1 - r_2)(1 - P^{(2)})\psi_{N,L}, \psi_{N,L} \rangle + \langle LV_{N,L}(r_1 - r_2)\psi_{N,L}, (1 - P^{(2)})\psi_{N,L} \rangle \right| \\ \leq L \| |V_{N,L}(r_1 - r_2)|^{1/2} \psi_{N,L} \|_{L^2} \| |V_{N,L}(r_1 - r_2)|^{1/2} (1 - P^{(2)})\psi_{N,L} \|_{L^2}.$$

Computing in the same way, we have

$$(2.69) \quad \frac{1}{2} \left| Tr \left( (1 - P^{(2)})LV_{N,L}(r_1 - r_2)P^{(2)} + P^{(2)}LV_{N,L}(r_1 - r_2)(1 - P^{(2)}) \right) \gamma_{N,L}^{(2)} \right| \\ = \frac{1}{2} \left| \langle LV_{N,L}(r_1 - r_2)P^{(2)}\psi_{N,L}, (1 - P^{(2)})\psi_{N,L} \rangle + \langle LV_{N,L}(r_1 - r_2)(1 - P^{(2)})\psi_{N,L}, P^{(2)}\psi_{N,L} \rangle \right| \\ \leq L \| |V_{N,L}(r_1 - r_2)|^{1/2} P^{(2)}\psi_{N,L} \|_{L^2} \| |V_{N,L}(r_1 - r_2)|^{1/2} (1 - P^{(2)})\psi_{N,L} \|_{L^2}.$$

Combining estimates (2.68) and (2.69), we obtain

$$(2.70) \quad |II_V| \leq L \left( \| |V_{N,L}(r_1 - r_2)|^{1/2} \psi_{N,L} \|_{L^2} + \| |V_{N,L}(r_1 - r_2)|^{1/2} P^{(2)}\psi_{N,L} \|_{L^2} \right) \\ \times \| |V_{N,L}(r_1 - r_2)|^{1/2} (1 - P^{(2)})\psi_{N,L} \|_{L^2}.$$

Next, we need to bound the right side terms. From estimate (2.59), we obtain

$$|LV_{N,L}(r_1 - r_2)| \leq C_\delta \|V\|_{L_z^\infty L_x^1} ((1 - \Delta_{x_1})(1 - \Delta_{x_2}))^{1/2+\delta} \\ \leq C_\delta \|V\|_{L_z^\infty L_x^1} \left( \eta^{-1}(1 - \Delta_{x_1})(1 - \Delta_{x_2}) + \eta^{\frac{1+2\delta}{1-2\delta}} \right),$$



where we have used the interpolation inequality for fractional powers in Lemma A.3 in the last line. By optimizing over  $\eta > 0$ , we have

$$(2.71) \quad L \| |V_{N,L}(r_1 - r_2)|^{1/2} \psi_{N,L} \|_{L^2}^2 = \langle L |V_{N,L}(r_1 - r_2)| \psi_{N,L}, \psi_{N,L} \rangle \\ \leq C_\delta \|V\|_{L_z^\infty L_x^1} (\langle (1 - \Delta_{x_1})(1 - \Delta_{x_2}) \psi_{N,L}, \psi_{N,L} \rangle)^{1/2+\delta}.$$

Similarly,

$$(2.72) \quad L \| |V_{N,L}(r_1 - r_2)|^{1/2} P^{(2)} \psi_{N,L} \|_{L^2}^2 \leq C_\delta \|V\|_{L_z^\infty L_x^1} (\langle (1 - \Delta_{x_1})(1 - \Delta_{x_2}) \psi_{N,L}, \psi_{N,L} \rangle)^{1/2+\delta}.$$

Using estimate (2.59) again, we get

$$(2.73) \quad (1 - P^{(2)}) |LV_{N,L}(r_1 - r_2)| (1 - P^{(2)}) \\ \leq C_\delta \|V\|_{L_z^\infty L_x^1} (1 - P^{(2)}) ((1 - \Delta_{x_1})(1 - \Delta_{x_2}))^{1/2+\delta} (1 - P^{(2)}) \\ = C_\delta \|V\|_{L_z^\infty L_x^1} (1 - P^{(2)}) ((1 - \Delta_{x_1})(1 - \Delta_{x_2}))^{1/2+\delta},$$

where we used  $(1 - \Delta_{x_j})P^{(2)} = P^{(2)}(1 - \Delta_{x_j})$  in the last equality.

Since  $1 - P^{(2)} \leq Q_1 + Q_2$ , we use Property 3 in Lemma A.6 to get

$$(2.74) \quad (1 - P^{(2)}) ((1 - \Delta_{x_1})(1 - \Delta_{x_2}))^{1/2+\delta} \leq (Q_1 + Q_2) ((1 - \Delta_{x_1})(1 - \Delta_{x_2}))^{1/2+\delta}.$$

By  $\min\{1 + \lambda_x^2, 1 + (\lambda_z^2 - 1)/L^2\} Q \leq S^2 Q$ , we obtain

$$(2.75) \quad (Q_1 + Q_2) ((1 - \Delta_{x_1})(1 - \Delta_{x_2}))^{1/2+\delta} \\ \leq (Q_1 + Q_2) (S_1^2 S_2^2)^{1/2+\delta} \\ \leq \frac{1}{\min\{1 + \lambda_x^2, 1 + (\lambda_z^2 - 1)/L^2\}^{1/2-\delta}} \left( S_1^2 (S_2^2)^{1/2+\delta} + (S_1^2)^{1/2+\delta} S_2^2 \right) \\ \leq \frac{1}{\min\{1 + \lambda_x^2, 1 + (\lambda_z^2 - 1)/L^2\}^{1/2-\delta}} \left[ \eta^{-1} S_1^2 S_2^2 + \eta^{\frac{1+2\delta}{1-2\delta}} (S_1^2 + S_2^2) \right].$$

where we have used the interpolation inequality for fractional powers in Lemma A.3 in the last line.

Putting estimates (2.73) (2.74) and (2.75) together, we have

$$(2.76) \quad (1 - P^{(2)}) |LV_{N,L}(r_1 - r_2)| (1 - P^{(2)}) \\ \leq \frac{C_\delta \|V\|_{L_z^\infty L_x^1}}{\min\{1 + \lambda_x^2, 1 + (\lambda_z^2 - 1)/L^2\}^{1/2-\delta}} \left[ \eta^{-1} S_1^2 S_2^2 + \eta^{\frac{1+2\delta}{1-2\delta}} (S_1^2 + S_2^2) \right].$$

By optimizing over  $\eta$ , we deduce that

$$(2.77) \quad L \| |V_{N,L}(r_1 - r_2)|^{1/2} (1 - P^{(2)}) \psi_{N,L} \|_{L^2}^2 \\ = \langle L |V_{N,L}(r_1 - r_2)| (1 - P^{(2)}) \psi_{N,L}, (1 - P^{(2)}) \psi_{N,L} \rangle \\ \leq \frac{C_\delta \|V\|_{L_z^\infty L_x^1}}{\min\{1 + \lambda_x^2, 1 + (\lambda_z^2 - 1)/L^2\}^{1/2-\delta}} \left( \text{Tr} \left( S_1^2 \gamma_{N,L}^{(1)} \right) \right)^{1/2-\delta} \left( \text{Tr} \left( S_1^2 S_2^2 \gamma_{N,L}^{(2)} \right) \right)^{1/2+\delta}.$$

Combining estimates (2.70) (2.71) (2.72) and (2.77), we get

$$(2.78) \quad II \geq - \frac{C_\delta \|V\|_{L_z^\infty L_x^1}}{\min\{1 + \lambda_x^2, 1 + (\lambda_z^2 - 1)/L^2\}^{1/4-\delta/2}} \left( \text{Tr} \left( S_1^2 \gamma_{N,L}^{(1)} \right) \right)^{1/4-\delta/2} \left( \text{Tr} \left( S_1^2 S_2^2 \gamma_{N,L}^{(2)} \right) \right)^{1/2+\delta}.$$

For III, we rewrite

$$(2.79) \quad III = \text{Tr} (1 - \Delta_{x_1}) \left[ P^{(2)} \gamma_{N,L}^{(2)} P^{(2)} - \int_{SP(L^2(\Omega_L))} |\phi_L^{\otimes 2}\rangle \langle \phi_L^{\otimes 2}| d\mu_{N,L}(\phi_L) \right] \\ + \frac{1}{2} \text{Tr} \left( LV_{N,L}(r_1 - r_2) \left[ P^{(2)} \gamma_{N,L}^{(2)} P^{(2)} - \int |\phi_L^{\otimes 2}\rangle \langle \phi_L^{\otimes 2}| d\mu_{N,L}(\phi_L) \right] \right) \\ = III_K + III_V.$$

For the first term  $III_K$ , we can use the inequality  $|TrAB| \leq \|A\|_{op} Tr|B|$  to get

$$(2.80) \quad |III_K| \leq \|(1 - \Delta_x)P\|_{op} Tr \left| P^{(2)} \gamma_{N,L}^{(2)} P^{(2)} - \int_{SP(L^2(\Omega_L))} |\phi_L^{\otimes 2}\rangle \langle \phi_L^{\otimes 2}| d\mu_{N,L}(\phi_L) \right|.$$

Using  $(1 - \Delta_x)P \leq (1 + \lambda_x^2)P$  and Lemma 2.11, we have

$$(2.81) \quad |III_K| \lesssim (1 + \lambda_x^2) \frac{d}{N}.$$

For the second term  $III_V$ , we use the inequality  $|TrAB| \leq \|A\|_{op} Tr|B|$  again to get

$$(2.82) \quad |III_V| \leq \|P^{(2)}|LV_{N,L}(r_1 - r_2)|P^{(2)}\|_{op} Tr \left| P^{(2)} \gamma_{N,L}^{(2)} P^{(2)} - \int_{SP(L^2(\Omega_L))} |\phi_L^{\otimes 2}\rangle \langle \phi_L^{\otimes 2}| d\mu_{N,L}(\phi_L) \right|.$$

By estimate (2.59) and  $P^j(1 - \Delta_{x_j}) \leq (1 + \lambda_x^2)P^j$ , we have

$$(2.83) \quad \begin{aligned} P^{(2)}|LV_{N,L}(r_1 - r_2)|P^{(2)} &\leq C_\delta \|V\|_{L_z^\infty L_x^1} \left( (P^1(1 - \Delta_{x_1})) \otimes (P^2(1 - \Delta_{x_2})) \right)^{1/2+\delta} \\ &\leq C_\delta \|V\|_{L_z^\infty L_x^1} (1 + \lambda_x^2)^{1+2\delta} P^{(2)}. \end{aligned}$$

Noting that  $[P^{(2)}|LV_{N,L}(r_1 - r_2)|P^{(2)}, P^{(2)}] = 0$ , by Property 2 in Lemma A.6, we deduce that

$$\|P^{(2)}|LV_{N,L}(r_1 - r_2)|P^{(2)}\|_{op} \leq C_\delta \|V\|_{L_z^\infty L_x^1} (1 + \lambda_x^2)^{1+2\delta}.$$

Therefore, using Lemma 2.11 again, we get

$$(2.84) \quad \begin{aligned} &\left| Tr \left( LV_{N,L}(r_1 - r_2) \left[ P^{(2)} \gamma_{N,L}^{(2)} P^{(2)} - \int_{SP(L^2(\Omega_L))} |\phi_L^{\otimes 2}\rangle \langle \phi_L^{\otimes 2}| d\mu_{N,L}(\phi_L) \right] \right) \right| \\ &\lesssim C_\delta \|V\|_{L_z^\infty L_x^1} (1 + \lambda_x^2)^{1+2\delta} \frac{d}{N}. \end{aligned}$$

Thus, for all  $\delta > 0$ , combining estimates (2.81) and (2.84), we obtain

$$(2.85) \quad III \gtrsim -C_\delta \|V\|_{L_z^\infty L_x^1} (1 + \lambda_x^2)^{1+2\delta} \frac{d}{N}.$$

Combining estimates (2.78) and (2.85), we get the desired lower bound, that is,

$$\begin{aligned} &\left\langle \psi_{N,L}, (1 + N^{-1}H_{N,L} - 1/L^2) \psi_{N,L} \right\rangle \\ &\gtrsim -C_\delta \|V\|_{L_z^\infty L_x^1} (1 + \lambda_x^2)^{1+2\delta} \frac{d}{N} \\ &\quad - \frac{C_\delta \|V\|_{L_z^\infty L_x^1}}{\min \{1 + \lambda_x^2, 1 + (\lambda_z^2 - 1)/L^2\}^{1/4-\delta/2}} \left( Tr \left( S_1^2 \gamma_{N,L}^{(1)} \right) \right)^{1/4-\delta/2} \left( Tr \left( S_1^2 S_2^2 \gamma_{N,L}^{(2)} \right) \right)^{1/2+\delta}. \end{aligned}$$

□

From Lemma 2.13, to get the lower bound estimate for  $E_{N,L}$ , we are left to control  $Tr(S_1^2 \gamma_{N,L}^{(1)})$  and  $Tr(S_1^2 S_2^2 \gamma_{N,L}^{(2)})$  for the ground state  $\psi_{N,L}$ .

**Lemma 2.14.** *Assume  $L(N/L)^\beta \rightarrow 1^-$ ,  $\beta < \frac{1}{2}$ . Let  $\psi_{N,L}$  be a ground state of  $1 + H_{N,L} - 1/L^2$ . For all  $\theta \in (0, 1)$ , we have*

$$(2.86) \quad Tr \left( S_1^2 \gamma_{N,L}^{(1)} \right) \lesssim C_{up} \frac{1 + |E_{N,L,\theta}|}{\theta},$$

$$(2.87) \quad Tr \left( S_1^2 S_2^2 \gamma_{N,L}^{(2)} \right) \lesssim C_{up}^2 \left( \frac{1 + |E_{N,L,\theta}|}{\theta} \right)^2.$$

*Proof.* For estimate (2.86), from the definition of  $E_{N,L,\theta}$ , we have

$$(2.88) \quad 1 + N^{-1}H_{N,L} - 1/L^2 - \theta N^{-1} \sum_{j=1}^N S_j^2 \geq E_{N,L,\theta}.$$

Let  $\psi_{N,L}$  be a ground state of  $1 + N^{-1}H_{N,L} - 1/L^2$ . By taking the expectation against  $\psi_{N,L}$  on both sides of (2.88), we obtain

$$Tr \left( S_1^2 \gamma_{N,L}^{(1)} \right) \leq \frac{1}{\theta} \left( \left\langle \psi_{N,L}, (1 + N^{-1}H_{N,L} - 1/L^2) \psi_{N,L} \right\rangle - E_{N,L,\theta} \right) = \frac{E_{N,L} - E_{N,L,\theta}}{\theta}.$$

From Lemma 2.10, we see  $C_{up} \geq e_{N,L} \geq E_{N,L} \geq E_{N,L,\theta}$ . Therefore,  $|E_{N,L}| \leq C_{up} + |E_{N,L,\theta}|$  and

$$(2.89) \quad Tr \left( S_1^2 \gamma_{N,L}^{(1)} \right) \leq \frac{2C_{up}(1 + |E_{N,L,\theta}|)}{\theta}.$$

For estimate (2.87), we notice that

$$(2.90) \quad Tr \left( S_1^2 S_2^2 \gamma_{N,L}^{(2)} \right) \leq \frac{2}{N^2} \left\langle \psi_{N,L}, \left( \sum_{j=1}^N S_j^2 \right)^2 \psi_{N,L} \right\rangle.$$

Thus, it needs only to control the right term of (2.90). We rewrite

$$(2.91) \quad \begin{aligned} \frac{2}{N^2} \left( \sum_{j=1}^N S_j^2 \right)^2 &= (1 + N^{-1}H_{N,L} - 1/L^2) \frac{1}{N} \sum_{j=1}^N S_j^2 + \frac{1}{N} \sum_{j=1}^N S_j^2 (1 + N^{-1}H_{N,L} - 1/L^2) \\ &\quad - \frac{1}{N^2(N-1)} \sum_{i=1}^N \sum_{j < k} (S_i^2 LV_{N,L}(r_j - r_k) + LV_{N,L}(r_j - r_k) S_i^2). \end{aligned}$$

We need to control the right side terms of (2.91).

For the first and second terms, by using the ground state  $\psi_{N,L}$  and estimate (2.89), we have

$$(2.92) \quad \begin{aligned} &\left\langle \psi_{N,L}, \left( (1 + N^{-1}H_{N,L} - 1/L^2) \frac{1}{N} \sum_{j=1}^N S_j^2 + \frac{1}{N} \sum_{j=1}^N S_j^2 (1 + N^{-1}H_{N,L} - 1/L^2) \right) \psi_{N,L} \right\rangle \\ &= \frac{2E_{N,L}}{N} \left\langle \psi_{N,L}, \sum_{j=1}^N S_j^2 \psi_{N,L} \right\rangle \leq \frac{4C_{up}^2(1 + |E_{N,L,\theta}|)^2}{\theta}. \end{aligned}$$

For the third term, we decompose it into two cases. On the one hand, we consider the case  $j \neq i$  and  $k \neq i$ . By estimate (2.58), we have

$$(2.93) \quad \begin{aligned} &\frac{1}{N(N-1)} \sum_{i \neq j < k \neq i} LV_{N,L}(r_j - r_k) \\ &= 1 + N^{-1}H_{N,L} - 1/L^2 - \theta N^{-1} \sum_{j=1}^N S_j^2 - (1 - \theta) N^{-1} \sum_{j=1}^N S_j^2 - \frac{1}{N(N-1)} \sum_{j \neq i} LV_{N,L}(r_i - r_j) \\ &\geq E_{N,L,\theta} - \left( \frac{1 - \theta}{N} + \frac{C_\delta(N/L)^\delta \|V\|_{L_z^\infty L_x^{1+\delta}}}{N^2} \right) \sum_{j=1}^N S_j^2. \end{aligned}$$

Since  $[V_{N,L}(r_j - r_k), S_i^2] = 0$ , by summing over  $i$ , we obtain

$$(2.94) \quad \begin{aligned} &\frac{1}{N^2(N-1)} \sum_{i=1}^N \sum_{i \neq j < k \neq i} (S_i^2 LV_{N,L}(r_j - r_k) + LV_{N,L}(r_j - r_k) S_i^2) \\ &\geq \frac{2}{N} \sum_{i=1}^N S_i^2 \left( E_{N,L,\theta} - \left( \frac{1 - \theta}{N} + \frac{C_\delta(N/L)^\delta \|V\|_{L_z^\infty L_x^{1+\delta}}}{N^2} \right) \sum_{j=1}^N S_j^2 \right) \\ &= \frac{2E_{N,L,\theta}}{N} \sum_{j=1}^N S_j^2 - \frac{2}{N} \left( \frac{1 - \theta}{N} + \frac{C_\delta(N/L)^\delta \|V\|_{L_z^\infty L_x^{1+\delta}}}{N^2} \right) \left( \sum_{j=1}^N S_j^2 \right)^2. \end{aligned}$$

On the other hand, when  $j = i$  or  $k = i$ , by estimate (2.61), we have

$$S_j^2 LV_{N,L}(r_j - r_k) + LV_{N,L}(r_j - r_k) S_j^2 \geq -C_\delta(V)(N/L)^{\beta+\delta} S_j^2 S_k^2.$$

Therefore,

$$(2.95) \quad \begin{aligned} & \frac{1}{N^2(N-1)} \sum_{j \neq k} (S_j^2 LV_{N,L}(r_j - r_k) + LV_{N,L}(r_j - r_k) S_j^2) \\ & \geq -C_\delta(V) \frac{(N/L)^{\beta+\delta}}{N^3} \left( \sum_{j=1}^N S_j^2 \right)^2. \end{aligned}$$

Putting estimates (2.94) and (2.95) together, we obtain

$$(2.96) \quad \begin{aligned} & \frac{1}{N^2(N-1)} \sum_{i=1}^N \sum_{j < k} (S_i^2 LV_{N,L}(r_j - r_k) + LV_{N,L}(r_j - r_k) S_i^2) \\ & \geq -\frac{2}{N^2} \left( 1 - \theta + \frac{C_\delta(V)(N/L)^{\beta+\delta}}{N} \right) \left( \sum_{j=1}^N S_j^2 \right)^2 + \frac{2E_{N,L,\theta}}{N} \sum_{j=1}^N S_j^2. \end{aligned}$$

Taking the expectation against the ground state  $\psi_{N,L}$  on both sides of (2.96), we obtain

$$(2.97) \quad \begin{aligned} & \frac{1}{N^2(N-1)} \sum_{i=1}^N \sum_{j < k} \langle \psi_{N,L}, (S_i^2 LV_{N,L}(r_j - r_k) + LV_{N,L}(r_j - r_k) S_i^2) \psi_{N,L} \rangle \\ & \geq -\frac{2}{N^2} \left( 1 - \theta + \frac{C_\delta(V)(N/L)^{\beta+\delta}}{N} \right) \left\langle \psi_{N,L}, \left( \sum_{j=1}^N S_j^2 \right)^2 \psi_{N,L} \right\rangle + \frac{2E_{N,L,\theta}}{N} \sum_{j=1}^N \langle \psi_{N,L}, S_j^2 \psi_{N,L} \rangle \\ & \geq -\frac{2}{N^2} \left( 1 - \theta + \frac{C_\delta(V)(N/L)^{\beta+\delta}}{N} \right) \left\langle \psi_{N,L}, \left( \sum_{j=1}^N S_j^2 \right)^2 \psi_{N,L} \right\rangle - \frac{2C_{up}(1 + |E_{N,L,\theta}|)^2}{\theta}, \end{aligned}$$

where we used estimate (2.89) in the last line.

Taking the expectation against the ground state  $\psi_{N,L}$  on both sides of (2.91), we use estimates (2.92) and (2.97) to obtain

$$(2.98) \quad \begin{aligned} & \frac{2}{N^2} \left\langle \psi_{N,L}, \left( \sum_{j=1}^N S_j^2 \right)^2 \psi_{N,L} \right\rangle \\ & \leq \frac{8C_{up}^2(1 + |E_{N,L,\theta}|)^2}{\theta} + \frac{2}{N^2} \left( 1 - \theta + \frac{C_\delta(V)(N/L)^{\beta+\delta}}{N} \right) \left\langle \psi_{N,L}, \left( \sum_{j=1}^N S_j^2 \right)^2 \psi_{N,L} \right\rangle. \end{aligned}$$

Equivalently, we have

$$(2.99) \quad \frac{2}{N^2} \left( \theta - \frac{C_\delta(V)(N/L)^{\beta+\delta}}{N} \right) \left\langle \psi_{N,L}, \left( \sum_{j=1}^N S_j^2 \right)^2 \psi_{N,L} \right\rangle \leq \frac{8C_{up}^2(1 + |E_{N,L,\theta}|)^2}{\theta}.$$

Since  $\beta < 1/2$ , we can take  $\delta$  such that  $2\beta + \delta < 1$ . Then, with  $L(N/L)^\beta \rightarrow 1^-$ , we deduce that

$$\theta - \frac{C_\delta(V)(N/L)^{\beta+\delta}}{N} \geq \theta - \frac{C_\delta(V)(N/L)^{2\beta+\delta}}{N/L} \geq \theta/2,$$

for large  $N$  and  $1/L$ . With estimate (2.99), we conclude

$$Tr \left( S_1^2 S_2^2 \gamma_{N,L}^{(2)} \right) \leq \frac{2}{N^2} \left\langle \left( \sum_{j=1}^N S_j^2 \right)^2 \psi_{N,L}, \psi_{N,L} \right\rangle \lesssim \frac{C_{up}^2(1 + |E_{N,L,\theta}|)^2}{\theta^2}.$$

□

Then, we prove the following theorem with a bootstrapping argument.

**Theorem 2.15.** *Assume  $L(N/L)^\beta \rightarrow 1^-$ ,  $\beta < \frac{1}{2}$ , and  $\|V\|_{L_z^\infty L_x^1} \leq \frac{2\alpha}{C_{gn}^4}$  for some  $\alpha \in (0, 1)$ , then we have*

$$\liminf_{\substack{N, 1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} E_{N,L,\varepsilon} \geq 0$$

for  $\varepsilon \in [0, 1 - \alpha)$ .

*Proof.* From Lemma 2.13 and Lemma 2.14, we have

$$(2.100) \quad E_{N,L} \gtrsim -C_\delta \|V\|_{L_z^\infty L_x^1} (1 + \lambda_x^2)^{1+2\delta} \frac{d}{N} - \frac{C_\delta \|V\|_{L_z^\infty L_x^1}}{\min\{1 + \lambda_x^2, 1 + (\lambda_z^2 - 1)/L^2\}^{1/4-\delta/2}} \left( \frac{C_{up}(1 + |E_{N,L,\theta}|)}{\theta} \right)^{5/4+3\delta/2}.$$

Similarly, we use  $e_{N,L}^\varepsilon$ ,  $E_{N,L}^\varepsilon$ ,  $E_{N,L,\theta}^\varepsilon$  to denote the ground state energy and  $C_{up}^\varepsilon$  to denote the upper bound in Lemma 2.10 with interaction function  $V^\varepsilon = (1 - \varepsilon)^{-1}V$ . Then, we obtain

$$(2.101) \quad E_{N,L}^\varepsilon \gtrsim C_\delta \|V^\varepsilon\|_{L_z^\infty L_x^1} (1 + \lambda_x^2)^{1+2\delta} \frac{d}{N} - \frac{C_\delta \|V^\varepsilon\|_{L_z^\infty L_x^1}}{\min\{1 + \lambda_x^2, 1 + (\lambda_z^2 - 1)/L^2\}^{1/4-\delta/2}} \left( \frac{C_{up}^\varepsilon(1 + |E_{N,L,\theta}^\varepsilon|)}{\theta} \right)^{5/4+3\delta/2},$$

Noting that  $E_{N,L,\varepsilon} = (1 - \varepsilon)E_{N,L}^\varepsilon$  and  $d \lesssim \lambda_x^2 \lambda_z$ , we take  $\theta = (\varepsilon' - \varepsilon)/(1 - \varepsilon)$  to obtain

$$(2.102) \quad E_{N,L,\varepsilon} \geq -C(V, \delta, \varepsilon, \varepsilon') \left( (1 + \lambda_x^2)^{1+2\delta} \frac{\lambda_x^2 \lambda_z}{N} + \frac{|E_{N,L,\varepsilon'}|^{5/4+3\delta/2}}{\min\{1 + \lambda_x^2, 1 + (\lambda_z^2 - 1)/L^2\}^{1/4-\delta/2}} \right).$$

for all  $0 < \varepsilon < \varepsilon' < 1/2$  and  $\delta \in (0, 1/2)$ ,  $N \geq N(\delta, \varepsilon, \varepsilon')$ .

We make the induction hypothesis (labeled  $I_\eta$ )

$$(2.103) \quad \limsup_{\substack{N, 1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} \frac{|E_{N,L,\varepsilon}|}{1 + (N/L)^\eta} < \infty \text{ for all } 0 < \varepsilon < 1 - \alpha.$$

We may assume  $I_\eta$  holds for  $\eta = \eta_0$  as a start point. Under the assumption  $L(N/L)^\beta \rightarrow 1^-$ , we have

$$N^{-1} \sim (N/L)^{\beta-1}.$$

Then, by taking  $\lambda_z = 2$  and  $\lambda_x = (N/L)^\tau$  with  $\tau \leq \beta$ , we deduce that  $I_{\eta'}$  holds provided that

$$(2.104) \quad \eta' > \max\{4\tau + \beta - 1, 5\eta_0/4 - \tau/2\}.$$

With the optimal choice  $\tau = 5\eta_0/18 + 2(1 - \beta)/9$ , we get

$$(2.105) \quad \eta' > \eta_0 + \frac{\eta_0 + \beta - 1}{9}.$$

Noting that  $\tau \leq \beta$ , we have to require

$$(2.106) \quad \eta_0 \leq \frac{22\beta - 4}{5}.$$

By Theorem 2.5, we can choose  $\eta_0 = 3\beta - 1$ , which satisfies (2.106). With  $\beta < 1/2$ , we choose a constant  $c$  such that

$$0 < c < \frac{2 - 4\beta}{9}$$

and  $I_{\eta'}$  holds with  $\eta' = \eta_0 - c$ . Repeating the process, we finally deduce that  $I_0$  holds. It means that  $|E_{N,L,\varepsilon}| \leq C_\varepsilon$  for  $N \geq N_\varepsilon$ . Then by taking  $\lambda_z = 2$  and  $\lambda_x = (N/L)^\tau$  with  $\tau$  small enough, we use estimate (2.102) to obtain

$$\liminf_{\substack{N, 1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} E_{N,L,\varepsilon} \geq 0.$$

□

By taking  $\varepsilon = \frac{1-\alpha}{\sqrt{2}}$  in Theorem 2.15, we obtain Theorem 2.4.

**2.4. High Energy estimates when  $k > 1$ .** Assuming (2.1) holds for  $k$ , we now prove it for  $k+2$ . By the induction hypothesis, we have

$$\begin{aligned} & \frac{1}{c_0^{k+2}} \langle \psi_{N,L}, (2 + N^{-1}H_{N,L} - 1/L^2)^{k+2} \psi_{N,L} \rangle \\ & \geq \frac{1}{c_0^2} \|S^{(k)}(2 + N^{-1}H_{N,L} - 1/L^2)\psi_{N,L}\|_{L^2}^2 \\ & = MS + E_C + E_P \end{aligned}$$

where the main sum  $MS$  is

$$MS = \frac{1}{c_0^2 N^2 (N-1)^2} \sum_{\substack{1 \leq i_1 < j_1 \leq N \\ 1 \leq i_2 < j_2 \leq N \\ i_1 > k, i_2 > k}} \langle S^{(k)}(2 + H_{i_1 j_1})\psi_{N,L}, S^{(k)}(2 + H_{i_2 j_2})\psi_{N,L} \rangle,$$

the cross error term  $E_C$  is

$$E_C = \frac{1}{c_0^2 N^2 (N-1)^2} \sum_{\substack{1 \leq i_1 < j_1 \leq N \\ 1 \leq i_2 < j_2 \leq N \\ i_1 \leq k, i_2 > k}} 2\operatorname{Re} \langle S^{(k)}(2 + H_{i_1 j_1})\psi_{N,L}, S^{(k)}(2 + H_{i_2 j_2})\psi_{N,L} \rangle,$$

and the nonnegative error term  $E_P$  is

$$\begin{aligned} E_P &= \frac{1}{c_0^2 N^2 (N-1)^2} \sum_{\substack{1 \leq i_1 < j_1 \leq N \\ 1 \leq i_2 < j_2 \leq N \\ i_1 \leq k, i_2 \leq k}} \langle S^{(k)}(2 + H_{i_1 j_1})\psi_{N,L}, S^{(k)}(2 + H_{i_2 j_2})\psi_{N,L} \rangle \\ &= \frac{1}{c_0^2 N^2 (N-1)^2} \left\langle \sum_{\substack{1 \leq i < j \leq N \\ i \leq k}} S^{(k)}(2 + H_{ij})\psi_{N,L}, \sum_{\substack{1 \leq i < j \leq N \\ i \leq k}} S^{(k)}(2 + H_{ij})\psi_{N,L} \right\rangle \\ &\geq 0. \end{aligned}$$

**2.4.1. Handling the main sum.** Commuting  $(1 + H_{i_1 j_1})$  and  $(1 + H_{i_2 j_2})$  with  $S^{(k)}$  in  $MS$ , we get

$$MS = M_1 + M_2 + M_3,$$

where  $M_1$  consists of the terms with

$$\{i_1, j_1\} \cap \{i_2, j_2\} = \emptyset,$$

$M_2$  consists of the terms with

$$|\{i_1, j_1\} \cap \{i_2, j_2\}| = 1,$$

and  $M_3$  consists of the terms with

$$|\{i_1, j_1\} \cap \{i_2, j_2\}| = 2.$$

By symmetric of  $\psi_{N,L}$ , we have

$$\begin{aligned} M_1 &= \frac{1}{4c_0^2} \langle S^{(k)}(2 + H_{(k+1)(k+2)})\psi_{N,L}, S^{(k)}(2 + H_{(k+3)(k+4)})\psi_{N,L} \rangle, \\ M_2 &= \frac{1}{2c_0^2} N^{-1} \langle S^{(k)}(2 + H_{(k+1)(k+2)})\psi_{N,L}, S^{(k)}(2 + H_{(k+2)(k+3)})\psi_{N,L} \rangle, \\ M_3 &= \frac{1}{2c_0^2} N^{-2} \langle S^{(k)}(2 + H_{(k+1)(k+2)})\psi_{N,L}, S^{(k)}(2 + H_{(k+1)(k+2)})\psi_{N,L} \rangle, \end{aligned}$$

up to an unimportant combination number.

Since  $M_3 \geq 0$ , we drop it. By the fact that

$$[2 + H_{(k+1)(k+2)}, 2 + H_{(k+3)(k+4)}] = 0,$$

we have

$$M_1 \geq \frac{4(1-\alpha)^2}{4c_0^2} \langle S^{(k)} \psi_{N,L}, S_{k+1}^2 S_{k+2}^2 S^{(k)} \psi_{N,L} \rangle$$

using Theorem 2.5 and Lemma A.6. Recall  $c_0 = \min\left(\frac{1-\alpha}{\sqrt{2}}, \frac{1}{2}\right)$ , hence

$$(2.107) \quad M_1 \geq 2 \langle S^{(k+2)} \psi_{N,L}, S^{(k+2)} \psi_{N,L} \rangle = 2 \|S^{(k+2)} \psi_{N,L}\|_{L^2}^2.$$

To deal with  $M_2$ , we expand

$$M_2 = M_{21} + M_{22} + M_{23},$$

where

$$\begin{aligned} M_{21} &= \frac{N^{-1}}{2c_0^2} \langle (2 + S_{k+1}^2 + S_{k+2}^2) S^{(k)} \psi_{N,L}, (2 + S_{k+2}^2 + S_{k+3}^2) S^{(k)} \psi_{N,L} \rangle, \\ M_{22} &= \frac{N^{-1}}{c_0^2} \text{Re} \langle (2 + S_{k+1}^2 + S_{k+2}^2) S^{(k)} \psi_{N,L}, LV_{N,L}(r_{k+2} - r_{k+3}) S^{(k)} \psi_{N,L} \rangle, \\ M_{23} &= \frac{N^{-1}}{2c_0^2} \langle LV_{N,L}(r_{k+1} - r_{k+2}) S^{(k)} \psi_{N,L}, LV_{N,L}(r_{k+2} - r_{k+3}) S^{(k)} \psi_{N,L} \rangle. \end{aligned}$$

We keep only the  $S_{k+2}^4$  terms inside  $M_{21}$ , which is the main contribution. That is,

$$(2.108) \quad \begin{aligned} M_{21} &\geq \frac{N^{-1}}{2c_0^2} \langle S_{k+2}^2 S_{k+2}^2 S^{(k)} \psi_{N,L}, S^{(k)} \psi_{N,L} \rangle \geq 2N^{-1} \langle S_{k+1}^4 S^{(k)} \psi_{N,L}, S^{(k)} \psi_{N,L} \rangle \\ &= 2N^{-1} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2. \end{aligned}$$

For  $M_{22}$ , we expand

$$\begin{aligned} M_{22} &= \frac{2N^{-1}}{c_0^2} \langle S^{(k)} \psi_{N,L}, LV_{N,L}(r_{k+2} - r_{k+3}) S^{(k)} \psi_{N,L} \rangle \\ &\quad + \frac{N^{-1}}{c_0^2} \langle S^{(k+1)} \psi_{N,L}, LV_{N,L}(r_{k+2} - r_{k+3}) S^{(k+1)} \psi_{N,L} \rangle \\ &\quad + \frac{N^{-1}}{c_0^2} \langle S^{(k)} \psi_{N,L}, S_{k+2}^2 LV_{N,L}(r_{k+2} - r_{k+3}) S^{(k)} \psi_{N,L} \rangle \\ &= M_{221} + M_{222} + M_{223}. \end{aligned}$$

By estimate (2.58)

$$(2.109) \quad |M_{221}| + |M_{222}| \lesssim \frac{L(N/L)^{\beta+}}{N} \left( \|S^{(k+1)} \psi_{N,L}\|_{L^2}^2 + \|S^{(k+2)} \psi_{N,L}\|_{L^2}^2 \right).$$

By estimate (2.61),

$$(2.110) \quad |M_{223}| \lesssim \frac{(N/L)^{\beta+}}{N} \|S^{(k+2)} \psi_{N,L}\|_{L^2}^2.$$

This requires  $\beta < \frac{1}{2}$ .

For  $M_{23}$ , with Hölder inequality, we have

$$(2.111) \quad \begin{aligned} |M_{23}| &\leq N^{-1} \|LV_{N,L}(r_{k+1} - r_{k+2})\|_{L_{z_{k+1}}^\infty L_{x_{k+1}}^{1+}} \|LV_{N,L}(r_{k+2} - r_{k+3})\|_{L_{z_{k+3}}^\infty L_{x_{k+3}}^{1+}} \|S^{(k)} \psi_{N,L}\|_{L^2 L_{x_{k+1}}^\infty L_{x_{k+3}}^\infty}^2 \\ &\lesssim N^{-1} L^2 (N/L)^{2\beta+2\delta} \|S^{(k+2)} \psi_{N,L}\|_{L^2}^2. \end{aligned}$$

Putting (2.107)–(2.111) together, with the assumption  $L(N/L)^\beta \rightarrow 1^-$ , we arrive at the following estimate for  $MS$ :

$$(2.112) \quad MS \geq (2 - C(N/L)^{2\beta-1+}) \left( \|S^{(k+2)} \psi_{N,L}\|_{L^2}^2 + N^{-1} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2 \right).$$

**2.4.2. Handling the cross error term.** Next, we turn our attention to estimate  $E_C$ . We will prove that

$$(2.113) \quad E_C \geq -C \max \left( N^{-\frac{3}{2}} (N/L)^{2\beta+}, N^{-1} (N/L)^{\beta+} \right) (\|S^{(k+2)} \psi_{N,L}\|_{L^2}^2 + N^{-1} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2).$$

Since  $L(N/L)^\beta \rightarrow 1^-$ , (2.113) is equivalent to

$$(2.114) \quad E_C \geq -C(N/L)^{(7\beta-3)/2} (\|S^{(k+2)} \psi_{N,L}\|_{L^2}^2 + N^{-1} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2).$$

That is, when  $\beta < 3/7$ ,  $E_C$  can be absorbable if added into (2.112).

We assume  $k \geq 1$ , since  $E_C = 0$  when  $k = 0$ . We decompose the sum into three parts

$$E_C = E_1 + E_2 + E_3,$$

where  $E_1$  contains the terms with  $j_1 \leq k$ ,  $E_2$  contains the terms with  $j_1 > k$  and  $j_1 \in \{i_2, j_2\}$  and  $E_3$  contains those term with  $j_1 > k$ ,  $j_1 \neq j_2$ .

Since  $H_{ij} = H_{ji}$ , by symmetry of  $\psi_{N,L}$ , we have

$$\begin{aligned} E_1 &= N^{-2} \langle S^{(k)} (2 + H_{12}) \psi_{N,L}, S^{(k)} (2 + H_{(k+1)(k+2)}) \psi_{N,L} \rangle, \\ E_2 &= N^{-2} \langle S^{(k)} (2 + H_{1(k+1)}) \psi_{N,L}, S^{(k)} (2 + H_{(k+1)(k+2)}) \psi_{N,L} \rangle, \\ E_3 &= N^{-1} \langle S^{(k)} (2 + H_{1(k+1)}) \psi_{N,L}, S^{(k)} (2 + H_{(k+2)(k+3)}) \psi_{N,L} \rangle, \end{aligned}$$

up to an unimportant combination number.

when  $k = 1$ ,  $E_1 = 0$ . Therefore, we address  $E_1$  for  $k \geq 2$ .

$$E_1 = E_{11} + E_{12} + E_{13} + E_{14},$$

where

$$\begin{aligned} E_{11} &= N^{-2} \langle S^{(k)} (2 + S_1^2 + S_2^2) \psi_{N,L}, S^{(k)} (2 + S_{k+1}^2 + S_{k+2}^2) \psi_{N,L} \rangle, \\ E_{12} &= N^{-2} \langle S^{(k)} (2 + S_1^2 + S_2^2) \psi_{N,L}, S^{(k)} LV_{N,L}(r_{k+1} - r_{k+2}) \psi_{N,L} \rangle, \\ E_{13} &= N^{-2} \langle S^{(k)} LV_{N,L}(r_1 - r_2) \psi_{N,L}, S^{(k)} (2 + S_{k+1}^2 + S_{k+2}^2) \psi_{N,L} \rangle, \\ E_{14} &= N^{-2} \langle S^{(k)} LV_{N,L}(r_1 - r_2) \psi_{N,L}, S^{(k)} LV_{N,L}(r_{k+1} - r_{k+2}) \psi_{N,L} \rangle. \end{aligned}$$

Since  $E_{11} \geq 0$ , we discard it. For  $E_{12}$ , by symmetry of  $\psi_{N,L}$ , we need to only consider

$$N^{-2} \langle S^{(k)} S_1^2 \psi_{N,L}, S^{(k)} LV_{N,L}(r_{k+1} - r_{k+2}) \psi_{N,L} \rangle.$$

Since  $[V_{N,L}(r_{k+1} - r_{k+2}), S_1] = 0$ , we use estimate (2.58) to obtain

$$\begin{aligned} &N^{-2} \langle S^{(k)} S_1^2 \psi_{N,L}, S^{(k)} LV_{N,L}(r_{k+1} - r_{k+2}) \psi_{N,L} \rangle \\ &= N^{-2} \langle S^{(k)} S_1 \psi_{N,L}, LV_{N,L}(r_{k+1} - r_{k+2}) S^{(k)} S_1 \psi_{N,L} \rangle \\ &\lesssim N^{-2} L(N/L)^{\beta+} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2. \end{aligned}$$

Hence

$$(2.115) \quad |E_{12}| \lesssim N^{-2} L(N/L)^{\beta+} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2.$$

For  $E_{13}$ , we decompose

$$E_{13} = E_{131} + E_{132} + E_{133},$$

where

$$\begin{aligned} E_{131} &= N^{-2} \langle S^{(k)} LV_{N,L}(r_1 - r_2) \psi_{N,L}, S^{(k)} S_{k+1}^2 \psi_{N,L} \rangle, \\ E_{132} &= N^{-2} \langle S^{(k)} LV_{N,L}(r_1 - r_2) \psi_{N,L}, S^{(k)} S_{k+2}^2 \psi_{N,L} \rangle, \\ E_{133} &= N^{-2} \langle S^{(k)} LV_{N,L}(r_1 - r_2) \psi_{N,L}, 2S^{(k)} \psi_{N,L} \rangle. \end{aligned}$$



For  $E_{131}$ , we expand

$$\begin{aligned}
E_{131} &= N^{-2} \langle LV_{N,L}(r_1 - r_2) \prod_{j=3}^{k+1} S_j \psi_{N,L}, S_1^2 S_2^2 \prod_{j=3}^{k+1} S_j P_{z_1, >1} \psi_{N,L} \rangle \\
&= N^{-2} \langle LV_{N,L}(r_1 - r_2) \prod_{j=3}^{k+1} S_j \psi_{N,L}, (1 - \Delta_{r_1} - 1/L^2) S_2^2 \prod_{j=3}^{k+1} S_j P_{z_1, >1} \psi_{N,L} \rangle \\
&= E_{1311} + E_{1312},
\end{aligned}$$

where

$$\begin{aligned}
E_{1311} &= N^{-2} \langle LV_{N,L}(r_1 - r_2) \prod_{j=3}^{k+1} S_j \psi_{N,L}, -\Delta_{r_1} S_2^2 \prod_{j=3}^{k+1} S_j P_{z_1, >1} \psi_{N,L} \rangle, \\
E_{1312} &= N^{-2} \langle LV_{N,L}(r_1 - r_2) \prod_{j=3}^{k+1} S_j \psi_{N,L}, (1 - 1/L^2) S_2^2 \prod_{j=3}^{k+1} S_j P_{z_1, >1} \psi_{N,L} \rangle.
\end{aligned}$$

Using integration by parts for  $E_{1311}$ ,

$$\begin{aligned}
E_{1311} &= N^{-2} \langle \nabla_{r_1} (LV_{N,L}(r_1 - r_2) \prod_{j=3}^{k+1} S_j \psi_{N,L}), \nabla_{r_1} S_2^2 \prod_{j=3}^{k+1} S_j P_{z_1, >1} \psi_{N,L} \rangle \\
&= N^{-2} L(N/L)^\beta \langle (\nabla_{r_1} V)_{N,L}(r_1 - r_2) \prod_{j=3}^{k+1} S_j \psi_{N,L}, \nabla_{r_1} S_2^2 \prod_{j=3}^{k+1} S_j P_{z_1, >1} \psi_{N,L} \rangle \\
&\quad + N^{-2} L \langle V_{N,L}(r_1 - r_2) \nabla_{r_1} \prod_{j=3}^{k+1} S_j \psi_{N,L}, \nabla_{r_1} S_2^2 \prod_{j=3}^{k+1} S_j P_{z_1, >1} \psi_{N,L} \rangle.
\end{aligned}$$

Using Hölder and Sobolev inequality,

$$\begin{aligned}
|E_{1311}| &\leq N^{-2} L(N/L)^\beta \|(\nabla V)_{N,L}\|_{L_z^\infty L_x^{2+}} \left\| \prod_{j=3}^{k+1} S_j \psi_{N,L} \right\|_{L^2 L_{x_2}^\infty} \left\| \nabla_{r_1} S_2^2 \prod_{j=3}^{k+1} S_j P_{z_1, >1} \psi_{N,L} \right\|_{L^2} \\
&\quad + N^{-2} L \|V_{N,L}\|_{L_z^\infty L_x^{2+}} \left\| \nabla_{r_1} \prod_{j=3}^{k+1} S_j \psi_{N,L} \right\|_{L^2 L_{x_2}^\infty} \left\| \nabla_{r_1} S_2^2 \prod_{j=3}^{k+1} S_j P_{z_1, >1} \psi_{N,L} \right\|_{L^2} \\
&\lesssim N^{-2} L(N/L)^{3\beta+} \left\| \prod_{j=2}^{k+1} S_j \psi_{N,L} \right\|_{L^2} \left\| \nabla_{r_1} S_2^2 \prod_{j=3}^{k+1} S_j P_{z_1, >1} \psi_{N,L} \right\|_{L^2} \\
&\quad + N^{-2} L(N/L)^{2\beta+} \left\| \nabla_{r_1} \prod_{j=2}^{k+1} S_j \psi_{N,L} \right\|_{L^2} \left\| \nabla_{r_1} S_2^2 \prod_{j=2}^{k+1} S_j P_{z_1, >1} \psi_{N,L} \right\|_{L^2}
\end{aligned}$$

By estimates (A.28) and (A.29), with  $L(N/L)^\beta \rightarrow 1^-$ , we have

$$\begin{aligned}
(2.116) \quad |E_{1311}| &\lesssim N^{-2} (N/L)^{2\beta+} \left\| \prod_{j=2}^{k+1} S_j \psi_{N,L} \right\|_{L^2} \|S_2 S^{(k+1)} \psi_{N,L}\|_{L^2} \\
&\quad + N^{-2} (N/L)^{2\beta+} \|S^{(k+1)} \psi_{N,L}\|_{L^2} \|S_2 S^{(k+1)} \psi_{N,L}\|_{L^2} \\
&\leq N^{-\frac{3}{2}} (N/L)^{2\beta+} (\|S^{(k+1)} \psi_{N,L}\|_{L^2}^2 + N^{-1} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2).
\end{aligned}$$

For  $E_{1312}$ , with Hölder and Sobolev inequality, we obtain

$$\begin{aligned} |E_{1312}| &\leq N^{-2}L^{-1}\|V_{N,L}\|_{L_z^\infty L_x^{2+}}\left\|\prod_{j=3}^{k+1}S_j\psi_{N,L}\right\|_{L^2L_{x_1}^\infty}\left\|S_2^2\prod_{j=3}^{k+1}S_jP_{z_1,>1}\psi_{N,L}\right\|_{L^2} \\ &\lesssim N^{-2}L^{-1}(N/L)^{2\beta+}\|S_1\prod_{j=3}^{k+1}S_j\psi_{N,L}\|_{L^2}\|S_2^2\prod_{j=3}^{k+1}S_jP_{z_1,>1}\psi_{N,L}\|_{L^2} \end{aligned}$$

By  $L^{-2}P_{z_1,>1} \leq S_1^2P_{z_1,>1}$ , we get

$$\begin{aligned} (2.117) \quad |E_{1312}| &\lesssim N^{-2}(N/L)^{2\beta+}\|S^{(k)}\psi_{N,L}\|_{L^2}\|S_1S^{(k+1)}\psi_{N,L}\|_{L^2} \\ &\leq N^{-\frac{3}{2}}(N/L)^{2\beta+}(\|S^{(k)}\psi_{N,L}\|_{L^2}^2 + N^{-1}\|S_1S^{(k+1)}\psi_{N,L}\|_{L^2}^2). \end{aligned}$$

Estimated in the same way as  $E_{131}$ ,

$$(2.118) \quad |E_{132}| \lesssim N^{-\frac{3}{2}}(N/L)^{2\beta+}(\|S^{(k)}\psi_{N,L}\|_{L^2}^2 + N^{-1}\|S_1S^{(k+1)}\psi_{N,L}\|_{L^2}^2),$$

$$(2.119) \quad |E_{133}| \lesssim N^{-\frac{3}{2}}(N/L)^{2\beta+}(\|S^{(k)}\psi_{N,L}\|_{L^2}^2 + N^{-1}\|S_1S^{(k+1)}\psi_{N,L}\|_{L^2}^2).$$

For  $E_{14}$ , we decompose

$$\begin{aligned} E_{14} &= N^{-2}\langle LV_{N,L}(r_1 - r_2) \prod_{j=3}^k S_j\psi_{N,L}, S_1^2S_2^2LV_{N,L}(r_{k+1} - r_{k+2}) \prod_{j=3}^k S_jP_{z_1,>1}\psi_{N,L} \rangle \\ &= E_{141} + E_{142}, \end{aligned}$$

where

$$\begin{aligned} E_{141} &= N^{-2}\langle LV_{N,L}(r_1 - r_2) \prod_{j=3}^k S_j\psi_{N,L}, -\Delta_{r_1}LV_{N,L}(r_{k+1} - r_{k+2})S_2^2\prod_{j=3}^k S_jP_{z_1,>1}\psi_{N,L} \rangle, \\ E_{142} &= N^{-2}\langle LV_{N,L}(r_1 - r_2) \prod_{j=3}^k S_j\psi_{N,L}, (1 - 1/L^2)LV_{N,L}(r_{k+1} - r_{k+2})S_2^2\prod_{j=3}^k S_jP_{z_1,>1}\psi_{N,L} \rangle. \end{aligned}$$

For  $E_{141}$ , using integration by parts, we obtain

$$\begin{aligned} E_{141} &= N^{-2}L(N/L)^\beta \langle \nabla_{r_1}(LV_{N,L}(r_1 - r_2) \prod_{j=3}^k S_j\psi_{N,L}), \nabla_{r_1}LV_{N,L}(r_{k+1} - r_{k+2})S_2^2\prod_{j=3}^k S_jP_{z_1,>1}\psi_{N,L} \rangle, \\ &= N^{-2}L(N/L)^\beta \langle (\nabla_{r_1}V)_{N,L}(r_1 - r_2) \prod_{j=3}^k S_j\psi_{N,L}, \nabla_{r_1}LV_{N,L}(r_{k+1} - r_{k+2})S_2^2\prod_{j=3}^k S_jP_{z_1,>1}\psi_{N,L} \rangle \\ &\quad + N^{-2}L\langle V_{N,L}(r_1 - r_2)\nabla_{r_1}\prod_{j=3}^k S_j\psi_{N,L}, \nabla_{r_1}LV_{N,L}(r_{k+1} - r_{k+2})S_2^2\prod_{j=3}^k S_jP_{z_1,>1}\psi_{N,L} \rangle. \end{aligned}$$

Using Hölder and Sobolev inequality,

$$\begin{aligned} &|E_{141}| \\ &\leq N^{-2}L^2(N/L)^\beta \|(\nabla_r V)_{N,L}\|_{L_z^\infty L_x^{2+}} \left\|\prod_{j=3}^k S_j\psi_{N,L}\right\|_{L^2L_{x_2}^\infty L_{x_{k+1}}^\infty} \|V_{N,L}\|_{L_z^\infty L_x^{1+}} \|\nabla_{r_1}S_2^2\prod_{j=3}^k S_jP_{z_1,>1}\psi_{N,L}\|_{L^2L_{x_{k+1}}^\infty} \\ &\quad + N^{-2}L^2\|V_{N,L}\|_{L_z^\infty L_x^{2+}} \|\nabla_{r_1}\prod_{j=3}^k S_j\psi_{N,L}\|_{L^2L_{x_2}^\infty L_{x_{k+1}}^\infty} \|V_{N,L}\|_{L_z^\infty L_x^{1+}} \|\nabla_{r_1}S_2^2\prod_{j=3}^k S_jP_{z_1,>1}\psi_{N,L}\|_{L^2L_{x_{k+1}}^\infty} \\ &\lesssim N^{-2}L^2(N/L)^{4\beta+} \left\|\prod_{j=2}^{k+1} S_j\psi_{N,L}\right\|_{L^2} \|\nabla_{r_1}S_2^2\prod_{j=3}^{k+1} S_jP_{z_1,>1}\psi_{N,L}\|_{L^2} \\ &\quad + N^{-2}L^2(N/L)^{3\beta+} \|\nabla_{r_1}\prod_{j=2}^{k+1} S_j\psi_{N,L}\|_{L^2} \|\nabla_{r_1}S_2^2\prod_{j=3}^{k+1} S_jP_{z_1,>1}\psi_{N,L}\|_{L^2}. \end{aligned}$$

By estimates (A.28) and (A.29), with  $L(N/L)^\beta \rightarrow 1^-$ , we have

$$(2.120) \quad \begin{aligned} |E_{141}| &\lesssim N^{-2}(N/L)^{2\beta+} \left\| \prod_{j=2}^{k+1} S_j \psi_{N,L} \right\|_{L^2} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2} \\ &\quad + N^{-2}(N/L)^{2\beta+} \|S^{(k+1)} \psi_{N,L}\|_{L^2} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2} \\ &\leq N^{-\frac{3}{2}}(N/L)^{2\beta+} (\|S^{(k+1)} \psi_{N,L}\|_{L^2}^2 + N^{-1} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2). \end{aligned}$$

For  $E_{142}$ , with Hölder and Sobolev inequality, we obtain

$$(2.121) \quad \begin{aligned} |E_{142}| &\lesssim N^{-2} \|V_{N,L}\|_{L_z^\infty L_x^{1+}} \left\| \prod_{j=3}^k S_j \psi_{N,L} \right\|_{L^2 L_{x_1}^{\infty-} L_{x_{k+1}}^{\infty-}} \|V_{N,L}\|_{L_z^\infty L_x^{1+}} \|S_2^2 \prod_{j=3}^k S_j P_{z_1, >1} \psi_{N,L}\|_{L^2 L_{x_1}^{\infty-} L_{x_{k+1}}^{\infty-}} \\ &\lesssim N^{-2}(N/L)^{2\beta+} \|S^{(k)} \psi_{N,L}\|_{L^2} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2} \\ &\leq N^{-\frac{3}{2}}(N/L)^{2\beta+} (\|S^{(k+1)} \psi_{N,L}\|_{L^2}^2 + N^{-1} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2). \end{aligned}$$

Hence, we obtain

$$(2.122) \quad E_1 \geq -CN^{-\frac{3}{2}}(N/L)^{2\beta+} (\|S^{(k+1)} \psi_{N,L}\|_{L^2}^2 + N^{-1} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2).$$

Next, we deal with  $E_2$ . We write

$$E_2 = E_{21} + E_{22} + E_{23} + E_{24},$$

where

$$\begin{aligned} E_{21} &= N^{-2} \langle S^{(k)} (2 + S_1^2 + S_{k+1}^2) \psi_{N,L}, S^{(k)} (2 + S_{k+1}^2 + S_{k+2}^2) \psi_{N,L} \rangle, \\ E_{22} &= N^{-2} \langle S^{(k)} (2 + S_1^2 + S_{k+1}^2) \psi_{N,L}, S^{(k)} LV_{N,L}(r_{k+1} - r_{k+2}) \psi_{N,L} \rangle, \\ E_{23} &= N^{-2} \langle S^{(k)} LV_{N,L}(r_1 - r_{k+1}) \psi_{N,L}, S^{(k)} (2 + S_{k+1}^2 + S_{k+2}^2) \psi_{N,L} \rangle, \\ E_{24} &= N^{-2} \langle S^{(k)} LV_{N,L}(r_1 - r_{k+1}) \psi_{N,L}, S^{(k)} LV_{N,L}(r_{k+1} - r_{k+2}) \psi_{N,L} \rangle. \end{aligned}$$

Since  $E_{21} \geq 0$ , we can discard it. For  $E_{22}$ , we decompose

$$E_{22} = E_{221} + E_{222} + E_{223},$$

where

$$\begin{aligned} E_{221} &= 2N^{-2} \langle S^{(k)} \psi_{N,L}, S^{(k)} LV_{N,L}(r_{k+1} - r_{k+2}) \psi_{N,L} \rangle, \\ E_{222} &= N^{-2} \langle S^{(k)} S_1^2 \psi_{N,L}, S^{(k)} LV_{N,L}(r_{k+1} - r_{k+2}) \psi_{N,L} \rangle, \\ E_{223} &= N^{-2} \langle S^{(k)} S_{k+1}^2 \psi_{N,L}, S^{(k)} LV_{N,L}(r_{k+1} - r_{k+2}) \psi_{N,L} \rangle. \end{aligned}$$

By estimate (2.58), we obtain

$$(2.123) \quad |E_{221}| \lesssim N^{-2} L(N/L)^{\beta+} \|S^{(k+1)} \psi_{N,L}\|_{L^2}^2,$$

$$(2.124) \quad |E_{222}| \lesssim N^{-2} L(N/L)^{\beta+} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2.$$

For  $E_{223}$ , by Hölder and Sobolev inequality, we have

$$(2.125) \quad \begin{aligned} |E_{223}| &\leq N^{-2} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2} \|LV_{N,L}\|_{L_z^\infty L_x^{2+}} \|S^{(k)} \psi_{N,L}\|_{L^2 L_{x_{k+1}}^{\infty-}} \\ &\lesssim N^{-2} L(N/L)^{2\beta+} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2. \end{aligned}$$

For  $E_{23}$ , we expand

$$E_{23} = E_{231} + E_{232} + E_{233},$$

where

$$\begin{aligned} E_{231} &= N^{-2} \langle S^{(k)} LV_{N,L}(r_1 - r_{k+1}) \psi_{N,L}, S^{(k)} S_{k+2}^2 \psi_{N,L} \rangle, \\ E_{232} &= N^{-2} \langle S^{(k)} LV_{N,L}(r_1 - r_{k+1}) \psi_{N,L}, S^{(k)} S_{k+1}^2 \psi_{N,L} \rangle, \\ E_{233} &= N^{-2} \langle S^{(k)} LV_{N,L}(r_1 - r_{k+1}) \psi_{N,L}, 2S^{(k)} \psi_{N,L} \rangle. \end{aligned}$$

For  $E_{231}$ , we expand

$$\begin{aligned} E_{231} &= N^{-2} \left\langle \prod_{j=2}^k S_j S_{k+2} L V_{N,L}(r_1 - r_{k+1}) \psi_{N,L}, S_1^2 \prod_{j=2}^k S_j S_{k+2} P_{z_1, >1} \psi_{N,L} \right\rangle \\ &= E_{2311} + E_{2312}, \end{aligned}$$

where

$$\begin{aligned} E_{2311} &= N^{-2} \left\langle \prod_{j=2}^k S_j S_{k+2} L V_{N,L}(r_1 - r_{k+1}) \psi_{N,L}, -\Delta_{r_1} \prod_{j=2}^k S_j S_{k+2} P_{z_1, >1} \psi_{N,L} \right\rangle, \\ E_{2312} &= N^{-2} \left\langle \prod_{j=2}^k S_j S_{k+2} L V_{N,L}(r_1 - r_{k+1}) \psi_{N,L}, (1 - 1/L^2) \prod_{j=2}^k S_j S_{k+2} P_{z_1, >1} \psi_{N,L} \right\rangle. \end{aligned}$$

For  $E_{2311}$ , using integration by parts, we have

$$\begin{aligned} E_{2311} &= N^{-2} \left\langle \nabla_{r_1} (L V_{N,L}(r_1 - r_{k+1}) \prod_{j=2}^k S_j S_{k+2} \psi_{N,L}), \nabla_{r_1} \prod_{j=2}^k S_j S_{k+2} P_{z_1, >1} \psi_{N,L} \right\rangle \\ &= N^{-2} L(N/L)^\beta \left\langle (\nabla V)_{N,L}(r_1 - r_{k+1}) \prod_{j=2}^k S_j S_{k+2} \psi_{N,L}, \nabla_{r_1} \prod_{j=2}^k S_j S_{k+2} P_{z_1, >1} \psi_{N,L} \right\rangle \\ &\quad + N^{-2} L \left\langle V_{N,L}(r_1 - r_{k+1}) \nabla_{r_1} \prod_{j=2}^k S_j S_{k+2} \psi_{N,L}, \nabla_{r_1} \prod_{j=2}^k S_j S_{k+2} P_{z_1, >1} \psi_{N,L} \right\rangle. \end{aligned}$$

Using Hölder and Sobolev inequality, we obtain

$$\begin{aligned} |E_{2311}| &\leq N^{-2} L(N/L)^\beta \|(\nabla_{r_1} V)_{N,L}\|_{L_z^\infty L_x^{1+}} \left\| \prod_{j=2}^k S_j S_{k+2} \psi_{N,L} \right\|_{L^2 L_{x_{k+1}}^{\infty-}} \left\| \nabla_{r_1} \prod_{j=2}^k S_j S_{k+2} P_{z_1, >1} \psi_{N,L} \right\|_{L^2 L_{x_{k+1}}^{\infty-}} \\ &\quad + N^{-2} L \|V_{N,L}\|_{L_z^\infty L_x^{1+}} \left\| \nabla_{r_1} \prod_{j=2}^k S_j S_{k+2} \psi_{N,L} \right\|_{L^2 L_{x_{k+1}}^{\infty-}} \left\| \nabla_{r_1} \prod_{j=2}^k S_j S_{k+2} P_{z_1, >1} \psi_{N,L} \right\|_{L^2 L_{x_{k+1}}^{\infty-}} \\ &\lesssim N^{-2} L(N/L)^{2\beta+} \left\| \prod_{j=2}^{k+2} S_j \psi_{N,L} \right\|_{L^2} \left\| \nabla_{r_1} \prod_{j=2}^{k+2} S_j P_{z_1, >1} \psi_{N,L} \right\|_{L^2} \\ &\quad + N^{-2} L(N/L)^{\beta+} \left\| \nabla_{r_1} \prod_{j=2}^{k+2} S_j \psi_{N,L} \right\|_{L^2} \left\| \nabla_{r_1} \prod_{j=2}^{k+2} S_j P_{z_1, >1} \psi_{N,L} \right\|_{L^2}. \end{aligned}$$

By estimates (A.28) and (A.29), with  $L(N/L)^\beta \rightarrow 1^-$ , we have

$$(2.126) \quad |E_{2311}| \lesssim N^{-2} (N/L)^{\beta+} \|S^{(k+2)} \psi_{N,L}\|_{L^2}^2.$$

For  $E_{2312}$ , with Hölder and Sobolev inequality, we have

$$\begin{aligned} |E_{2312}| &\lesssim N^{-2} L^{-2} \|L V_{N,L}\|_{L_z^\infty L_x^{1+}} \left\| \prod_{j=2}^k S_j S_{k+2} \psi_{N,L} \right\|_{L^2 L_{x_{k+1}}^{\infty-}} \left\| \prod_{j=2}^k S_j S_{k+2} P_{z_1, >1} \psi_{N,L} \right\|_{L^2 L_{x_{k+1}}^{\infty-}} \\ &\lesssim N^{-2} L^{-1} (N/L)^{\beta+} \left\| \prod_{j=2}^{k+2} S_j \psi_{N,L} \right\|_{L^2} \left\| \prod_{j=2}^{k+2} S_j P_{z_1, >1} \psi_{N,L} \right\|_{L^2}. \end{aligned}$$

By  $L^{-2} P_{z_1, >1} \leq S_1^2 P_{z_1, >1}$ , we get

$$(2.127) \quad |E_{2312}| \lesssim N^{-2} (N/L)^{\beta+} \|S^{(k+1)} \psi_{N,L}\|_{L^2} \|S^{(k+2)} \psi_{N,L}\|_{L^2}$$

Estimated in the same way as  $E_{131}$ ,

$$(2.128) \quad |E_{232}| \lesssim N^{-\frac{3}{2}} (N/L)^{2\beta+} (\|S^{(k+1)} \psi_{N,L}\|_{L^2}^2 + N^{-1} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2).$$

Estimated in the same way as  $E_{231}$ ,

$$(2.129) \quad |E_{233}| \lesssim N^{-2} (N/L)^{\beta+} \|S^{(k+2)} \psi_{N,L}\|_{L^2}^2.$$

For  $E_{24}$ , we decompose

$$\begin{aligned} E_{24} &= N^{-2} \langle LV_{N,L}(r_1 - r_{k+1}) \prod_{j=2}^k S_j \psi_{N,L}, S_1^2 LV_{N,L}(r_{k+1} - r_{k+2}) \prod_{j=2}^k S_j P_{z_1, >1} \psi_{N,L} \rangle \\ &= E_{241} + E_{242}, \end{aligned}$$

where

$$\begin{aligned} E_{241} &= N^{-2} \langle LV_{N,L}(r_1 - r_{k+1}) \prod_{j=2}^k S_j \psi_{N,L}, -\Delta_{r_1} LV_{N,L}(r_{k+1} - r_{k+2}) \prod_{j=2}^k S_j P_{z_1, >1} \psi_{N,L} \rangle, \\ E_{242} &= N^{-2} \langle LV_{N,L}(r_1 - r_{k+1}) \prod_{j=2}^k S_j \psi_{N,L}, (1 - 1/L^2) LV_{N,L}(r_{k+1} - r_{k+2}) \prod_{j=2}^k S_j P_{z_1, >1} \psi_{N,L} \rangle. \end{aligned}$$

For  $E_{241}$ , with Hölder and Sobolev inequality, we obtain

$$\begin{aligned} |E_{241}| &\leq N^{-2} \|LV_{N,L}\|_{L_z^\infty L_{x_1}^{2+}} \left\| \prod_{j=2}^k S_j \psi_{N,L} \right\|_{L^2 L_{x_{k+1}}^{\infty-} L_{x_1}^{\infty-}} \|LV_{N,L}\|_{L_z^\infty L_{x_{k+1}}^{2+}} \|\Delta_{r_1} \prod_{j=2}^k S_j P_{z_1, >1} \psi_{N,L}\|_{L^2} \\ &\lesssim N^{-2} L^2 (N/L)^{4\beta+} \|S^{(k+1)} \psi_{N,L}\|_{L^2} \|\Delta_{r_1} \prod_{j=2}^k S_j P_{z_1, >1} \psi_{N,L}\|_{L^2}. \end{aligned}$$

By estimate (A.29), with  $L(N/L)^\beta \rightarrow 1^-$ , we have

$$\begin{aligned} (2.130) \quad |E_{241}| &\leq N^{-2} (N/L)^{2\beta+} \|S^{(k+1)} \psi_{N,L}\|_{L^2} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2} \\ &\leq N^{-\frac{3}{2}} (N/L)^{2\beta+} (\|S^{(k+1)} \psi_{N,L}\|_{L^2}^2 + N^{-1} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2). \end{aligned}$$

For  $E_{242}$ , with Hölder and Sobolev inequality, we have

$$\begin{aligned} (2.131) \quad |E_{242}| &\lesssim N^{-2} \|V_{N,L}\|_{L_z^\infty L_{x_1}^{1+}} \left\| \prod_{j=2}^k S_j \psi_{N,L} \right\|_{L^2 L_{x_{k+1}}^{\infty-} L_{x_1}^{\infty-}} \|V_{N,L}\|_{L_z^\infty L_{x_{k+1}}^{1+}} \left\| \prod_{j=2}^k S_j P_{z_1, >1} \psi_{N,L} \right\|_{L^2 L_{x_{k+1}}^{\infty-} L_{x_1}^{\infty-}} \\ &\lesssim N^{-2} (N/L)^{2\beta+} \|S^{(k+1)} \psi_{N,L}\|_{L^2}^2. \end{aligned}$$

Hence, we get

$$(2.132) \quad E_2 \gtrsim -\max \left( N^{-\frac{3}{2}} (N/L)^{2\beta+}, N^{-2} (N/L)^{2\beta+} \right) (\|S^{(k+2)} \psi_{N,L}\|_{L^2}^2 + N^{-1} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2).$$

Finally, we handle  $E_3$  and expand

$$E_3 = E_{31} + E_{32} + E_{33},$$

where

$$\begin{aligned} E_{31} &= N^{-1} \langle S^{(k)} (2 + S_1^2 + S_{k+1}^2) \psi_{N,L}, S^{(k)} (2 + H_{(k+2)(k+3)}) \psi_{N,L} \rangle, \\ E_{32} &= N^{-1} \langle S^{(k)} LV_{N,L}(r_1 - r_{k+1}) \psi_{N,L}, S^{(k)} (2 + S_{k+2}^2 + S_{k+3}^2) \psi_{N,L} \rangle, \\ E_{33} &= N^{-1} \langle S^{(k)} LV_{N,L}(r_1 - r_{k+1}) \psi_{N,L}, S^{(k)} LV_{N,L}(r_{k+2} - r_{k+3}) \psi_{N,L} \rangle. \end{aligned}$$

We first discard  $E_{31}$ , since  $E_{31} \geq 0$  by Theorem 2.5 and Lemma A.6. For  $E_{32}$ , we expand

$$E_{32} = E_{321} + E_{322} + E_{323},$$

where

$$\begin{aligned} E_{321} &= N^{-1} \langle S^{(k)} LV_{N,L}(r_1 - r_{k+1}) \psi_{N,L}, S^{(k)} S_{k+2}^2 \psi_{N,L} \rangle, \\ E_{322} &= N^{-1} \langle S^{(k)} LV_{N,L}(r_1 - r_{k+1}) \psi_{N,L}, S^{(k)} S_{k+3}^2 \psi_{N,L} \rangle, \\ E_{323} &= N^{-1} \langle S^{(k)} LV_{N,L}(r_1 - r_{k+1}) \psi_{N,L}, 2S^{(k)} \psi_{N,L} \rangle. \end{aligned}$$

Estimated in the same way as  $E_{231}$ ,

$$(2.133) \quad |E_{321}| \lesssim N^{-1}(N/L)^{\beta+} \|S^{(k+2)}\psi_{N,L}\|_{L^2}^2,$$

$$(2.134) \quad |E_{322}| \lesssim N^{-1}(N/L)^{\beta+} \|S^{(k+2)}\psi_{N,L}\|_{L^2}^2,$$

$$(2.135) \quad |E_{323}| \lesssim N^{-1}(N/L)^{\beta+} \|S^{(k+2)}\psi_{N,L}\|_{L^2}^2.$$

For  $E_{33}$ , we expand

$$\begin{aligned} E_{33} &= N^{-1} \langle LV_{N,L}(r_1 - r_{k+1}) \prod_{j=2}^k S_j \psi_{N,L}, S_1^2 LV_{N,L}(r_{k+2} - r_{k+3}) \prod_{j=2}^k S_j \psi_{N,L} \rangle \\ &= E_{331} + E_{332}, \end{aligned}$$

where

$$\begin{aligned} E_{331} &= N^{-1} \langle LV_{N,L}(r_1 - r_{k+1}) \prod_{j=2}^k S_j \psi_{N,L}, -\Delta_{r_1} LV_{N,L}(r_{k+2} - r_{k+3}) \prod_{j=2}^k S_j P_{z_1, >1} \psi_{N,L} \rangle, \\ E_{332} &= N^{-1} \langle LV_{N,L}(r_1 - r_{k+1}) \prod_{j=2}^k S_j \psi_{N,L}, (1 - 1/L^2) LV_{N,L}(r_{k+2} - r_{k+3}) \prod_{j=2}^k S_j P_{z_1, >1} \psi_{N,L} \rangle. \end{aligned}$$

Using integration by parts for  $E_{331}$ ,

$$\begin{aligned} E_{331} &= N^{-1} \langle \nabla_{r_1} LV_{N,L}(r_1 - r_{k+1}) \prod_{j=2}^k S_j \psi_{N,L}, \nabla_{r_1} LV_{N,L}(r_{k+2} - r_{k+3}) \prod_{j=2}^k S_j P_{z_1, >1} \psi_{N,L} \rangle \\ &= N^{-1} \langle (\nabla_{r_1} V)_{N,L}(r_1 - r_{k+1}) \prod_{j=2}^k S_j \psi_{N,L}, \nabla_{r_1} LV_{N,L}(r_{k+2} - r_{k+3}) \prod_{j=2}^k S_j P_{z_1, >1} \psi_{N,L} \rangle \\ &\quad + N^{-1} L \langle V_{N,L}(r_1 - r_{k+1}) \nabla_{r_1} \prod_{j=2}^k S_j \psi_{N,L}, \nabla_{r_1} LV_{N,L}(r_{k+2} - r_{k+3}) \prod_{j=2}^k S_j P_{z_1, >1} \psi_{N,L} \rangle. \end{aligned}$$

Using Hölder and Sobolev inequality,

$$\begin{aligned} &|E_{331}| \\ &\leq N^{-1} L \|(\nabla_r V)_{N,L}\|_{L_z^\infty L_x^{1+}} \left\| \prod_{j=2}^k S_j \psi_{N,L} \right\|_{L^2 L_{x_{k+1}}^{\infty-} L_{x_{k+2}}^{\infty-}} \|V_{N,L}\|_{L_z^\infty L_x^{1+}} \|\nabla_{r_1} \prod_{j=2}^k S_j P_{z_1, >1} \psi_{N,L}\|_{L^2 L_{x_{k+1}}^{\infty-} L_{x_{k+2}}^{\infty-}} \\ &\quad + N^{-1} L^2 \|V_{N,L}\|_{L_z^\infty L_x^{1+}} \|\nabla_{r_1} \prod_{j=2}^k S_j \psi_{N,L}\|_{L^2 L_{x_{k+1}}^{\infty-} L_{x_{k+2}}^{\infty-}} \|V_{N,L}\|_{L_z^\infty L_x^{1+}} \|\nabla_{r_1} \prod_{j=2}^k S_j P_{z_1, >1} \psi_{N,L}\|_{L^2 L_{x_{k+1}}^{\infty-} L_{x_{k+2}}^{\infty-}} \\ &\lesssim N^{-1} L (N/L)^{2\beta+} \left\| \prod_{j=2}^{k+2} S_j \psi_{N,L} \right\|_{L^2} \|\nabla_{r_1} \prod_{j=2}^{k+2} S_j P_{z_1, >1} \psi_{N,L}\|_{L^2} \\ &\quad + N^{-1} L^2 (N/L)^{2\beta+} \|\nabla_{r_1} \prod_{j=2}^{k+2} S_j \psi_{N,L}\|_{L^2} \|\nabla_{r_1} \prod_{j=2}^{k+2} S_j P_{z_1, >1} \psi_{N,L}\|_{L^2}. \end{aligned}$$

By estimates (A.28) and (A.29), with  $L(N/L)^\beta \rightarrow 1^-$ , we have

$$(2.136) \quad \begin{aligned} |E_{331}| &\lesssim N^{-1} (N/L)^{\beta+} \|S^{(k+1)}\psi_{N,L}\|_{L^2} \|S^{(k+2)}\psi_{N,L}\|_{L^2} \\ &\quad + N^{-1} (N/L)^{\beta+} \|S^{(k+2)}\psi_{N,L}\|_{L^2} \|S^{(k+2)}\psi_{N,L}\|_{L^2}. \end{aligned}$$

For  $E_{332}$ , with Hölder and Sobolev inequality, we obtain

$$\begin{aligned} |E_{332}| &\lesssim N^{-1} \|V_{N,L}\|_{L_z^\infty L_x^{1+}} \left\| \prod_{j=2}^k S_j \psi_{N,L} \right\|_{L^2 L_{x_{k+1}}^{\infty-} L_{x_{k+2}}^{\infty-}} \|V_{N,L}\|_{L_z^\infty L_x^{1+}} \left\| \prod_{j=2}^k S_j P_{z_1, >1} \psi_{N,L} \right\|_{L^2 L_{x_{k+1}}^{\infty-} L_{x_{k+2}}^{\infty-}} \\ &\lesssim N^{-1} (N/L)^{2\beta+} \left\| \prod_{j=2}^{k+2} S_j \psi_{N,L} \right\|_{L^2} \left\| \prod_{j=2}^{k+2} S_j P_{z_1, >1} \psi_{N,L} \right\|_{L^2}. \end{aligned}$$

By  $L^{-2} P_{z_1, >1} \leq S_1^2 P_{z_1, >1}$ , with  $L(N/L)^\beta \rightarrow 1^-$ , we get

$$(2.137) \quad |E_{332}| \leq N^{-1} (N/L)^{\beta+} \|S^{(k+1)} \psi_{N,L}\|_{L^2} \|S^{(k+2)} \psi_{N,L}\|_{L^2}.$$

That is

$$(2.138) \quad E_3 \gtrsim -N^{-1} (N/L)^{\beta+} \|S^{(k+2)} \psi_{N,L}\|_{L^2}^2.$$

Putting (2.122), (2.132) and (2.138) together, we obtain the estimate for the cross error term

$$(2.139) \quad E_C \geq -C \max \left( N^{-\frac{3}{2}} (N/L)^{2\beta+}, N^{-1} (N/L)^{\beta+} \right) (\|S^{(k+2)} \psi_{N,L}\|_{L^2}^2 + N^{-1} \|S_1 S^{(k+1)} \psi_{N,L}\|_{L^2}^2).$$

Hence we have proved for all  $k$  and established Theorem 2.1.

### 3. COMPACTNESS, CONVERGENCE, AND UNIQUENESS

To work on compactness, convergence and uniqueness, we introduce an appropriate topology on the density matrices, as was previously done in [10, 16, 19, 23, 24, 32, 33, 34, 35, 36, 47, 64]. Denote the spaces of compact operators and trace class operators on  $L^2(\Omega^{\otimes k})$  as  $\mathcal{K}_k$  and  $\mathcal{L}_k^1$ , respectively. Then  $(\mathcal{K}_k)' = \mathcal{L}_k^1$ . By the fact that  $\mathcal{K}_k$  is separable, we select a dense countable subset  $\{J_i^{(k)}\}_{i \geq 1} \subset \mathcal{K}_k$  in the unit ball of  $\mathcal{K}_k$  (so  $\|J_i^{(k)}\|_{op} \leq 1$  where  $\|\cdot\|_{op}$  is the operator norm). For  $\gamma^{(k)}, \tilde{\gamma}^{(k)} \in \mathcal{L}_k^1$ , we then define a metric  $d_k$  on  $\mathcal{L}_k^1$  by

$$d_k(\gamma^{(k)}, \tilde{\gamma}^{(k)}) = \sum_{i=1}^{\infty} 2^{-i} \left| \text{Tr} J_i^{(k)} (\gamma^{(k)} - \tilde{\gamma}^{(k)}) \right|.$$

A uniformly bounded sequence  $\tilde{\gamma}_{N,L}^{(k)} \in \mathcal{L}_k^1$  converges to  $\tilde{\gamma}^{(k)}$  with respect to the weak\* topology if and only if

$$\lim_{N, 1/L \rightarrow \infty} d_k(\tilde{\gamma}_{N,L}^{(k)}, \tilde{\gamma}^{(k)}) = 0.$$

For fixed  $T > 0$ , let  $C([0, T]; \mathcal{L}_k^1)$  be the space of functions of  $t \in [0, T]$  with values in  $\mathcal{L}_k^1$  that are continuous with respect to the metric  $d_k$ . On  $C([0, T]; \mathcal{L}_k^1)$ , we define the metric

$$(3.1) \quad \hat{d}_k(\gamma^{(k)}(\cdot), \tilde{\gamma}^{(k)}(\cdot)) = \sup_{t \in [0, T]} d_k(\gamma^{(k)}(t), \tilde{\gamma}^{(k)}(t)),$$

and denote by  $\tau_{prod}$  the topology on the space  $\bigotimes_{k \geq 1} C([0, T]; \mathcal{L}_k^1)$  given by the product of topologies generated by the metrics  $\hat{d}_k$  on  $C([0, T], \mathcal{L}_k^1)$ .

#### 3.1. Compactness of the BBGKY sequence.

**Theorem 3.1.** *Assume  $L(N/L)^\beta \rightarrow 1^-$ . Then the sequence*

$$(3.2) \quad \left\{ \Gamma_{N,L}(t) = \left\{ \tilde{\gamma}_{N,L}^{(k)} \right\}_{k=1}^N \right\} \subset \bigotimes_{k \geq 1} C([0, T]; \mathcal{L}_k^1),$$

which satisfies the BBGKY hierarchy, is compact with respect to the product topology  $\tau_{prod}$ . For any limit point  $\Gamma(t) = \{\tilde{\gamma}^{(k)}(t)\}_{k=1}^\infty$ , we have  $\tilde{\gamma}^{(k)}$  is a symmetric nonnegative trace class operator with trace bounded by 1.

*Proof.* By the standard diagonalization argument, it suffices to show the compactness of  $\tilde{\gamma}_{N,L}^{(k)}$  for fixed  $k$  with respect to the metric  $\hat{d}_k$ . By the Arzelà-Ascoli theorem, this is equivalent to the equicontinuity of  $\gamma_{N,L}^{(k)}$ , and by, this is equivalent to the statement that for every observable  $J^{(k)}$  from a dense subset of  $\mathcal{K}_k$  and for every  $\varepsilon > 0$ , there exists  $\delta(J^{(k)}, \varepsilon)$  such that for all  $t_1, t_2 \in [0, T]$  with  $|t_1 - t_2| \leq \delta$ , we have

$$(3.3) \quad \sup_{N,L} \left| \text{Tr} J^{(k)} \tilde{\gamma}_{N,L}^{(k)}(t_1) - \text{Tr} J^{(k)} \tilde{\gamma}_{N,L}^{(k)}(t_2) \right| \leq \varepsilon.$$

We assume that compact operators  $J^{(k)}$  have been cutoff in Lemma A.8. Since the observable  $J^{(k)}$  can be written as a sum of a self-adjoint operator and an anti-self-adjoint operator, we may assume  $J^{(k)}$  is self-adjoint. Inserting the decomposition (2.3) on the left and right sides of  $\tilde{\gamma}_{N,L}^{(k)}$ , we obtain

$$\tilde{\gamma}_{N,L}^{(k)} = \sum_{\alpha, \beta} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta,$$

where the sum is taken over all  $k$ -tuples  $\alpha$  and  $\beta$ .

To establish (3.3), it suffices to prove that, for each  $\alpha$  and  $\beta$ , we have

$$(3.4) \quad \sup_{N,L} \left| \text{Tr} J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta(t_1) - \text{Tr} J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta(t_2) \right| \leq \varepsilon.$$

To this end, we establish the estimate

$$(3.5) \quad \begin{aligned} & \left| \text{Tr} J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta(t_1) - \text{Tr} J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta(t_2) \right| \\ & \leq C |t_2 - t_1| \left( 1_{\{\alpha=0, \beta=0\}} + \max \left\{ 1, L^{|\alpha|+|\beta|-2} \right\} 1_{\{\alpha \neq 0 \text{ or } \beta \neq 0\}} \right). \end{aligned}$$

By (3.5), we can directly establish (3.4) except for the case  $|\alpha| + |\beta| = 1$ . However, from Corollary 2.2, we can also get a bound

$$(3.6) \quad \begin{aligned} & \left| \text{Tr} J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta(t_1) - \text{Tr} J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta(t_2) \right| \\ & \leq 2 \sup_t |\langle J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}(t), \tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L}(t) \rangle| \\ & \leq 2 \|J^{(k)}\|_{op} \|\tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}(t)\|_{L^2} \|\tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L}(t)\|_{L^2} \\ & \lesssim L^{|\alpha|+|\beta|}. \end{aligned}$$

By averaging (3.5) and (3.6) in the case  $|\alpha| + |\beta| = 1$ , we obtain

$$\left| \text{Tr} J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta(t_1) - \text{Tr} J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta(t_2) \right| \lesssim |t_2 - t_1|^{1/2},$$

which suffices to establish (3.4).

Thus, we are left to prove (3.5). The BBGKY hierarchy (1.14) yields

$$(3.7) \quad \partial_t \text{Tr} J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta = I + II + III + IV,$$

where

$$\begin{aligned} I &= -i \sum_{j=1}^k \text{Tr} J^{(k)} \left[ -\Delta_{x_j}, \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta \right], \\ II &= -i \frac{1}{L^2} \sum_{j=1}^k \text{Tr} J^{(k)} \left[ -\partial_{z_j}^2, \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta \right], \\ III &= -i \frac{1}{N-1} \sum_{1 \leq i < j \leq k} \text{Tr} J^{(k)} \tilde{\mathcal{P}}_\alpha \left[ \tilde{V}_{N,L}(r_i - r_j), \tilde{\gamma}_{N,L}^{(k)} \right] \tilde{\mathcal{P}}_\beta, \\ IV &= -i \frac{N-k}{N-1} \sum_{j=1}^k \text{Tr} J^{(k)} \text{Tr}_{r_{k+1}} \tilde{\mathcal{P}}_\alpha \left[ \tilde{V}_{N,L}(r_j - r_{k+1}), \tilde{\gamma}_{N,L}^{(k+1)} \right] \tilde{\mathcal{P}}_\beta. \end{aligned}$$



First, we handle  $I$ . By Lemma A.7 and integration by parts, we have

$$\begin{aligned} I &= i \sum_{j=1}^k \left( \langle J^{(k)} \Delta_{x_j} \tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}, \tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L} \rangle - \langle J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}, \tilde{\mathcal{P}}_\beta \Delta_{x_j} \tilde{\psi}_{N,L} \rangle \right) \\ &= i \sum_{j=1}^k \left( \langle J^{(k)} \Delta_{x_j} \tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}, \tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L} \rangle - \langle \Delta_{x_j} J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}, \tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L} \rangle \right). \end{aligned}$$

Hence

$$(3.8) \quad |I| \leq \sum_{j=1}^k \left( \|J^{(k)} \Delta_{x_j}\|_{op} + \|\Delta_{x_j} J^{(k)}\|_{op} \right) \|\tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}\|_{L^2} \|\tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L}\|_{L^2} \leq C_{k,J^{(k)}},$$

where in the last step we used the energy estimate.

Next, we consider  $II$ . When  $\alpha = \beta = 0$ , we have

$$II = -i \frac{1}{L^2} \sum_{j=1}^k Tr J^{(k)} \left[ -\partial_{z_j}^2 - 1, \tilde{\mathcal{P}}_0 \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_0 \right] = 0,$$

where we used  $[1, \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta] = 0$  in the first equality.

When  $|\alpha| + |\beta| \geq 1$ , applying Lemma A.7 and integration by parts again, we have

$$\begin{aligned} II &= i \frac{1}{L^2} \sum_{j=1}^k \left( \langle J^{(k)} \partial_{z_j}^2 \tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}, \tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L} \rangle - \langle J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}, \partial_{z_j}^2 \tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L} \rangle \right) \\ &= i \frac{1}{L^2} \sum_{j=1}^k \left( \langle J^{(k)} \partial_{z_j}^2 \tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}, \tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L} \rangle - \langle \partial_{z_j}^2 J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}, \tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L} \rangle \right). \end{aligned}$$

Hence

$$|II| \leq \frac{1}{L^2} \sum_{j=1}^k \left( \|J^{(k)} \partial_{z_j}^2\|_{op} + \|\partial_{z_j}^2 J^{(k)}\|_{op} \right) \|\tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}\|_{L^2} \|\tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L}\|_{L^2}.$$

By the energy estimate (2.6),

$$(3.9) \quad |II| = 0, \quad |\alpha| + |\beta| = 0,$$

$$(3.10) \quad |II| \lesssim C_{k,J^{(k)}} L^{|\alpha|+|\beta|-2}, \quad |\alpha| + |\beta| \geq 1.$$

Next, we consider  $III$ . Similarly,

$$\begin{aligned} III &= \frac{-i}{N-1} \sum_{1 \leq i < j \leq k} \langle J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{V}_{N,L}(r_i - r_j) \tilde{\psi}_{N,L}, \tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L} \rangle \\ &\quad + \frac{i}{N-1} \sum_{1 \leq i < j \leq k} \langle J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}, \tilde{\mathcal{P}}_\beta \tilde{V}_{N,L}(r_i - r_j) \tilde{\psi}_{N,L} \rangle \\ &= \frac{-i}{N-1} \sum_{1 \leq i < j \leq k} \langle J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{V}_{N,L}(r_i - r_j) \tilde{\psi}_{N,L}, \tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L} \rangle \\ &\quad + \frac{i}{N-1} \sum_{1 \leq i < j \leq k} \langle \tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}, J^{(k)} \tilde{\mathcal{P}}_\beta \tilde{V}_{N,L}(r_i - r_j) \tilde{\psi}_{N,L} \rangle. \end{aligned}$$

Let

$$W_{ij} = \langle \nabla_{r_i} \rangle^{-1} \langle \nabla_{r_j} \rangle^{-1} \tilde{V}_{N,L}(r_i - r_j) \langle \nabla_{r_i} \rangle^{-1} \langle \nabla_{r_j} \rangle^{-1}.$$

Hence

$$\begin{aligned} |III| &\lesssim N^{-1} \sum_{1 \leq i < j \leq k} \|J^{(k)} \langle \nabla_{r_i} \rangle \langle \nabla_{r_j} \rangle\|_{op} \|W_{ij}\|_{op} \|\langle \nabla_{r_i} \rangle \langle \nabla_{r_j} \rangle \tilde{\psi}_{N,L}\|_{L^2} \|\tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L}\|_{L^2} \\ &\quad + N^{-1} \sum_{1 \leq i < j \leq k} \|\langle \nabla_{r_i} \rangle \langle \nabla_{r_j} \rangle J^{(k)}\|_{op} \|W_{ij}\|_{op} \|\langle \nabla_{r_i} \rangle \langle \nabla_{r_j} \rangle \tilde{\psi}_{N,L}\|_{L^2} \|\tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}\|_{L^2}. \end{aligned}$$

Since  $\|W_{ij}\|_{op} \lesssim \|\tilde{V}_{N,L}\|_{L_z^\infty L_x^1} \leq \|V\|_{L_z^\infty L_x^1}$  by (2.60), the energy estimates (2.5) (2.6) imply that

$$(3.11) \quad |III| \lesssim \frac{C_{k,J^{(k)}}}{N}.$$

For  $IV$ , we have

$$\begin{aligned} IV &= -i \frac{N-k}{N-1} \sum_{j=1}^k \langle J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{V}_{N,L}(r_j - r_{k+1}) \psi_{N,L}, \tilde{\mathcal{P}}_\beta \psi_{N,L} \rangle \\ &\quad + i \frac{N-k}{N-1} \sum_{j=1}^k \langle J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{V}_{N,L}(r_j - r_{k+1}) \psi_{N,L}, \tilde{\mathcal{P}}_\beta \psi_{N,L} \rangle. \end{aligned}$$

Then, since  $J^{(k)} \langle \nabla_{r_{k+1}} \rangle = \langle \nabla_{r_{k+1}} \rangle J^{(k)}$ ,

$$\begin{aligned} IV &= -i \frac{N-k}{N-1} \sum_{j=1}^k \langle J^{(k)} \tilde{\mathcal{P}}_\alpha \langle \nabla_{r_j} \rangle W_{j(k+1)} \langle \nabla_{r_j} \rangle \langle \nabla_{r_{k+1}} \rangle \tilde{\psi}_{N,L}, \tilde{\mathcal{P}}_\beta \langle \nabla_{r_{k+1}} \rangle \tilde{\psi}_{N,L} \rangle \\ &\quad + i \frac{N-k}{N-1} \sum_{j=1}^k \langle \langle \nabla_{r_j} \rangle J^{(k)} \tilde{\mathcal{P}}_\alpha \langle \nabla_{r_{k+1}} \rangle \tilde{\psi}_{N,L}, \tilde{\mathcal{P}}_\beta W_{j(k+1)} \langle \nabla_{r_j} \rangle \langle \nabla_{r_{k+1}} \rangle \tilde{\psi}_{N,L} \rangle. \end{aligned}$$

Hence

$$|IV| \lesssim \sum_{j=1}^k \left( \|J^{(k)} \langle \nabla_{r_j} \rangle\|_{op} + \|\langle \nabla_{r_j} \rangle J^{(k)}\|_{op} \right) \|W_{j(k+1)}\|_{op} \|\langle \nabla_{r_j} \rangle \langle \nabla_{r_{k+1}} \rangle \tilde{\psi}_{N,L}\|_{L^2} \|\langle \nabla_{r_{k+1}} \rangle \tilde{\psi}_{N,L}\|_{L^2}.$$

By energy estimate (2.5),

$$(3.12) \quad |IV| \lesssim C_{k,J^{(k)}}.$$

Integrating (3.7) from  $t_1$  to  $t_2$  and putting (3.8), (3.9), (3.10), (3.11) and (3.12) together, we obtain (3.5).  $\square$

*Corollary 3.2.* Let  $\Gamma(t) = \{\tilde{\gamma}^{(k)}\}_{k=1}^\infty$  be a limit point of  $\left\{ \Gamma_{N,L}(t) = \left\{ \tilde{\gamma}_{N,L}^{(k)} \right\}_{k=1}^N \right\}$ , with respect to the product topology  $\tau_{prod}$ . Then  $\tilde{\gamma}^{(k)}$  satisfies the priori bound

$$(3.13) \quad Tr \langle \nabla \rangle^{(k)} \tilde{\gamma}^{(k)} \langle \nabla \rangle^{(k)} \leq C^k,$$

and takes the structure

$$(3.14) \quad \tilde{\gamma}^{(k)}(t, (\mathbf{x}_k, \mathbf{z}_k); (\mathbf{x}'_k, \mathbf{z}'_k)) = \tilde{\gamma}_x^{(k)}(t, \mathbf{x}_k; \mathbf{x}'_k) \left( \prod_{j=1}^k \frac{2}{\pi} \cos(z_j) \cos(z'_j) \right),$$

where  $\tilde{\gamma}_x^{(k)} = Tr_z \tilde{\gamma}^{(k)}$ .

*Proof.* The estimate (3.13) follows by (2.5) in Corollary 2.2 and Theorem 3.1. To establish the formula (3.14), it suffices to prove  $\tilde{\mathcal{P}}_\alpha \tilde{\gamma}^{(k)} \tilde{\mathcal{P}}_\beta = 0$  if either  $\alpha \neq 0$  or  $\beta \neq 0$ . This is equivalent to the statement that for any  $J^{(k)} \in \mathcal{K}_k$ ,  $Tr J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}^{(k)} \tilde{\mathcal{P}}_\beta = 0$ . By Corollary (2.2), we obtain

$$(3.15) \quad Tr J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}^{(k)} \tilde{\mathcal{P}}_\beta = \lim_{\substack{N, 1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} Tr J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta.$$

By Lemma A.7,

$$Tr J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L}^{(k)} \tilde{\mathcal{P}}_\beta = \langle J^{(k)} \mathcal{P}_\alpha \tilde{\psi}_{N,L}, \tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L} \rangle,$$

and by Cauchy-Schwarz and (2.6),

$$\begin{aligned} \left| Tr J^{(k)} \tilde{\mathcal{P}}_\alpha \tilde{\gamma}_{N,L} \tilde{\mathcal{P}}_\beta \right| &\leq \|J^{(k)}\|_{op} \|\tilde{\mathcal{P}}_\alpha \tilde{\psi}_{N,L}\|_{L^2} \|\tilde{\mathcal{P}}_\beta \tilde{\psi}_{N,L}\|_{L^2} \\ &\leq C^k L^{|\alpha|+|\beta|}. \end{aligned}$$

Hence the right side of (3.15) is 0.  $\square$

**Theorem 3.3.** *The sequence*

$$\Gamma_{x,N,L} = \left\{ \tilde{\gamma}_{x,N,L} = \text{Tr}_z \tilde{\gamma}_{N,L}^{(k)} \right\}_{k=1}^N \in \bigotimes_{k \geq 1} C([0, T]; \mathcal{L}_k^1(\mathbb{T}^{2k}))$$

is compact with respect to the 2D version of the product topology  $\tau_{\text{prod}}$  used in Theorem 3.1.

*Proof.* The proof is similar to the 3D case and we omit it. Also see [19, Theorem 5].  $\square$

**3.2. Limit points satisfy GP hierarchy.** To prove the limit points satisfy the GP hierarchy, a technical tool we need is the approximation of identity type lemma, which is used to compare the  $\delta$ -function and its approximation. Since we request  $L(N/L)^\beta \rightarrow 1^-$ , we see that  $\tilde{V}_{N,L}(x, z)$  defined by (1.8) formally converges to  $\delta(x) \int V(x, z) dx$ . Thus, we need a modified version of this type lemma as follows.

**Lemma 3.4.** *Let  $\rho \in L^1(\Omega)$  be a function compactly supported on  $\Omega$  such that*

$$\sup_z \int |\rho(x, z)| |x| dx < \infty$$

and define  $\rho_{\varepsilon, \lambda}(x, z) = \varepsilon^{-2} \lambda^{-1} \rho(x/\varepsilon, z/\lambda)$  and  $g(z) = \int \rho(x, z) dx$ . Then, for every  $\kappa \in [0, 1)$ , there exists  $C > 0$ , such that

$$\begin{aligned} & \left| \text{Tr} J^{(k)}(\rho_{\varepsilon, \lambda}(r_j - r_{k+1}) - \delta(x_j - x_{k+1})g(z_j - z_{k+1}))\gamma^{(k+1)} \right| \\ & \leq C\varepsilon^\kappa \sup_z \int |\rho(x, z)| |x|^\kappa dx \left( \|\langle \nabla_{x_j} \rangle^{-1} J^{(k)} \langle \nabla_{x_j} \rangle\|_{\text{op}} + \|\langle \nabla_{x_j} \rangle J^{(k)} \langle \nabla_{x_j} \rangle^{-1}\|_{\text{op}} \right) \text{Tr} \langle \nabla_{x_j} \rangle^2 \langle \nabla_{x_{k+1}} \rangle^2 \gamma^{(k+1)} \\ & \quad + C_J \|\rho_{1, \lambda} - \rho_{1, 1}\|_{L^1} \text{Tr} \langle \nabla_{r_j} \rangle^2 \langle \nabla_{r_{k+1}} \rangle^2 \gamma^{(k+1)} \end{aligned}$$

for all nonnegative  $\gamma^{(k+1)} \in \mathcal{L}_{k+1}^1$ .

*Proof.* We will give a proof for Lemma 3.4 in the Appendix.  $\square$

**Theorem 3.5.** *Assume  $L(N/L)^\beta \rightarrow 1^-$ , and let  $\Gamma(t) = \{\tilde{\gamma}^{(k)}\}_{k=1}^\infty$  be a limit point of*

$$\left\{ \Gamma_{N,L}(t) = \left\{ \tilde{\gamma}_{N,L}^{(k)}(t) \right\}_{k=1}^N \right\}$$

with respect to the product topology  $\tau_{\text{prod}}$ . Then  $\left\{ \tilde{\gamma}_x^{(k)} = \text{Tr}_z \tilde{\gamma}^{(k)} \right\}_{k=1}^\infty$  is a solution to the coupled focusing Gross-Pitaevskii hierarchy subject to initial data  $\tilde{\gamma}_x^{(k)}(0) = |\phi_0\rangle \langle \phi_0|^{\otimes k}$ , which, rewritten in integral form, is

$$\tilde{\gamma}_x^{(k)} = U^{(k)}(t) \tilde{\gamma}_x^{(k)}(0) - i \sum_{j=1}^k \int_0^t U^{(k)}(t-s) \text{Tr}_{x_{k+1}} \text{Tr}_z \left[ \delta(x_j - x_{k+1}) \int V(x, z_j - z_{k+1}) dx, \tilde{\gamma}^{(k+1)}(s) \right] ds,$$

where  $U^{(k)}(t) = \prod_{j=1}^k e^{it\Delta_{x_j}} e^{-it\Delta_{x'_j}}$ .

*Proof.* Passing to subsequences if necessary, we have

$$(3.17) \quad \lim_{\substack{N, 1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} \sup_t \text{Tr} J^{(k)} \left( \tilde{\gamma}_{N,L}^{(k)} - \tilde{\gamma}^{(k)}(t) \right) = 0, \quad \forall J^{(k)} \in \mathcal{K}(L^2(\Omega^k))$$

$$(3.18) \quad \lim_{\substack{N, 1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} \sup_t \text{Tr} J_x^{(k)} \left( \tilde{\gamma}_{x,N,L}^{(k)} - \tilde{\gamma}_x^{(k)}(t) \right) = 0, \quad \forall J_x^{(k)} \in \mathcal{K}(L^2(\mathbb{T}^{2k}))$$

from Theorem 3.1 and 3.3.

It suffices to test the limit point against the test function  $J_x^{(k)} \in \mathcal{K}(L^2(\mathbb{T}^{2k}))$ . We will prove that the limit point satisfies

$$(3.19) \quad \text{Tr} J_x^{(k)} \tilde{\gamma}_x^{(k)}(0) = \text{Tr} J_x^{(k)} |\phi_0\rangle \langle \phi_0|^{\otimes k},$$

and

$$(3.20) \quad \begin{aligned} Tr J_x^{(k)} \tilde{\gamma}_x^{(k)}(t) = & Tr J_x^{(k)} U^{(k)}(t) \tilde{\gamma}_x^{(k)}(0) \\ & + i \sum_{j=1}^k \int_0^t U^{(k)}(t-s) Tr_{x_{k+1}} Tr_z \left[ \delta(x_j - x_{k+1}) g(z_j - z_{k+1}), \tilde{\gamma}^{(k+1)}(s) \right] ds, \end{aligned}$$

where we adopt the notation  $g(z) = \int V(x, z) dx$  for simplicity.

To end this, we use the coupled focusing BBGKY hierarchy, which is

$$(3.21) \quad Tr J_x^{(k)} \tilde{\gamma}_{x,N,L}^{(k)}(t) = A - \frac{i}{N-1} \sum_{i < j}^k B - i \left( 1 - \frac{k+1}{N-1} \right) \sum_{j=1}^k D,$$

where

$$\begin{aligned} A &= Tr J_x^{(k)} U^{(k)}(t) \tilde{\gamma}_{x,N,L}^{(k)}(0), \\ B &= \int_0^t Tr J_x^{(k)} U^{(k)}(t-s) \left[ \tilde{V}_{N,L}(r_i - r_j), \tilde{\gamma}_{N,L}^{(k)}(s) \right] ds, \\ D &= \int_0^t Tr J_x^{(k)} U^{(k)}(t-s) \left[ \tilde{V}_{N,L}(r_j - r_{k+1}), \tilde{\gamma}_{N,L}^{(k+1)}(s) \right] ds. \end{aligned}$$

By (3.18), we have

$$(3.22) \quad \lim_{\substack{N,1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} Tr J_x^{(k)} \tilde{\gamma}_{x,N,L}^{(k)}(t) = Tr J_x^{(k)} \tilde{\gamma}_x^{(k)}(t),$$

$$(3.23) \quad \lim_{\substack{N,1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} Tr J_x^{(k)} U^{(k)}(t) \tilde{\gamma}_{x,N,L}^{(k)}(0) = Tr J_x^{(k)} U^{(k)}(t) \tilde{\gamma}_x^{(k)}(0).$$

By the argument in [56], we know, from assumption (ii) in Theorem 1.2,

$$\tilde{\gamma}_{N,L}^{(1)}(0) \rightarrow \frac{2}{\pi} \phi_0(x_1) \overline{\phi_0}(x'_1) \cos(z_1) \cos(z'_1), \quad \text{strongly in trace norm};$$

that is,

$$\tilde{\gamma}_{N,L}^{(k)}(0) \rightarrow \prod_{j=1}^k \frac{2}{\pi} \phi_0(x_j) \overline{\phi_0}(x'_j) \cos(z_j) \cos(z'_j), \quad \text{strongly in trace norm}.$$

Thus we have checked (3.19), the left-hand side of (3.20), and the first term on the right-hand side of (3.20) for the limit point. We are left to prove that

$$(3.24) \quad \lim_{\substack{N,1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} \frac{B}{N-1} = 0,$$

$$(3.25) \quad \lim_{\substack{N,1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} \left( 1 - \frac{k+1}{N-1} \right) D = \int_0^t Tr J_x^{(k)} U^{(k)}(t-s) \left[ \delta(x_j - x_{k+1}) g(z_j - z_{k+1}), \tilde{\gamma}^{(k+1)}(s) \right] ds.$$

First, we will show the boundedness of  $|B|$  and  $|D|$  for every finite time  $t$ . Noting that  $[U^{(k)}, \langle \nabla_{r_i} \rangle] = 0$ , we have

$$\begin{aligned}
|B| &\leq \int_0^t |Tr J_x^{(k)} U^{(k)}(t-s) [\tilde{V}_{N,L}(r_i - r_j), \tilde{\gamma}_{N,L}^{(k)}(s)]| ds \\
&= \int_0^t ds |Tr \langle \nabla_{r_i} \rangle^{-1} \langle \nabla_{r_j} \rangle^{-1} J_x^{(k)} \langle \nabla_{r_i} \rangle \langle \nabla_{r_j} \rangle U^{(k)}(t-s) W_{ij} \langle \nabla_{r_i} \rangle \langle \nabla_{r_j} \rangle \tilde{\gamma}_{N,L}^{(k)}(s) \langle \nabla_{r_i} \rangle \langle \nabla_{r_j} \rangle \\
&\quad - Tr \langle \nabla_{r_i} \rangle \langle \nabla_{r_j} \rangle J_x^{(k)} \langle \nabla_{r_i} \rangle^{-1} \langle \nabla_{r_j} \rangle^{-1} U^{(k)}(t-s) \langle \nabla_{r_i} \rangle \langle \nabla_{r_j} \rangle \tilde{\gamma}_{N,L}^{(k)} \langle \nabla_{r_i} \rangle \langle \nabla_{r_j} \rangle W_{ij}| \\
&\leq \int_0^t ds \|\langle \nabla_{r_i} \rangle^{-1} \langle \nabla_{r_j} \rangle^{-1} J_x^{(k)} \langle \nabla_{r_i} \rangle \langle \nabla_{r_j} \rangle\|_{op} \|U^{(k)}\|_{op} \|W_{ij}\|_{op} Tr \langle \nabla_{r_i} \rangle^2 \langle \nabla_{r_j} \rangle^2 \tilde{\gamma}_{N,L}^{(k)}(s) \\
&\quad + \int_0^t ds \|\langle \nabla_{r_i} \rangle \langle \nabla_{r_j} \rangle J_x^{(k)} \langle \nabla_{r_i} \rangle^{-1} \langle \nabla_{r_j} \rangle^{-1}\|_{op} \|U^{(k)}\|_{op} \|W_{ij}\|_{op} Tr \langle \nabla_{r_i} \rangle^2 \langle \nabla_{r_j} \rangle^2 \tilde{\gamma}_{N,L}^{(k)}(s) \\
&\leq C_J t.
\end{aligned}$$

$|D|$  can be estimated in the same way as  $|B|$  and hence

$$|D| \leq C_J t.$$

That is,

$$\lim_{\substack{N, 1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} \frac{B}{N-1} = \lim_{\substack{N, 1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} \frac{(k+1)D}{N-1} = 0.$$

Next, we will use Lemma 3.4 to prove

$$(3.26) \quad \lim_{\substack{N, 1/L \rightarrow \infty \\ L(N/L)^\beta \rightarrow 1^-}} D = \int_0^t Tr J_x^{(k)} U^{(k)}(t-s) \left[ \delta(x_j - x_{k+1}) g(z_j - z_{k+1}), \tilde{\gamma}^{(k+1)}(s) \right] ds$$

Let  $\eta \in L^1(\mathbb{T}^2)$  be a smooth probability density function compactly supported on  $(-\pi, \pi)^2$  and define  $\eta_\varepsilon(x) = \varepsilon^{-2} \eta(x/\varepsilon)$ . For simplicity, we adopt the notation  $M_{s,t}^{(k)} = J_x^{(k)} U^{(k)}(t-s)$ . Then, we expand

$$\begin{aligned}
&Tr J_x^{(k)} U^{(k)}(t-s) \left( \tilde{V}_{N,L}(r_j - r_{k+1}) \tilde{\gamma}_{N,L}^{(k+1)}(s) - \delta(x_j - x_{k+1}) g(z_j - z_{k+1}) \tilde{\gamma}^{(k+1)}(s) \right) \\
&= I + II + III + IV,
\end{aligned}$$

where

$$\begin{aligned}
I &= Tr M_{s,t}^{(k)} \left( \tilde{V}_{N,L}(r_j - r_{k+1}) - \delta(x_j - x_{k+1}) g(z_j - z_{k+1}) \right) \tilde{\gamma}_{N,L}^{(k+1)}(s), \\
II &= Tr M_{s,t}^{(k)} (\delta(x_j - x_{k+1}) - \eta_\varepsilon(x_j - x_{k+1})) g(z_j - z_{k+1}) \tilde{\gamma}_{N,L}^{(k+1)}(s), \\
III &= Tr M_{s,t}^{(k)} \eta_\varepsilon(x_j - x_{k+1}) g(z_j - z_{k+1}) \left( \tilde{\gamma}_{N,L}^{(k+1)}(s) - \tilde{\gamma}^{(k+1)}(s) \right), \\
IV &= Tr M_{s,t}^{(k)} (\eta_\varepsilon(x_j - x_{k+1}) - \delta(x_j - x_{k+1})) g(z_j - z_{k+1}) \tilde{\gamma}^{(k+1)}(s).
\end{aligned}$$

It needs only to prove  $I - IV$  converge to 0 as  $N, 1/L \rightarrow \infty$ . By Lemma 3.4, we have

$$\begin{aligned}
(3.27) \quad |I| &= \left| Tr M_{s,t}^{(k)} \left( \tilde{V}_{N,L}(r_j - r_{k+1}) - \delta(x_j - x_{k+1}) g(z_j - z_{k+1}) \right) \tilde{\gamma}_{N,L}^{(k+1)}(s) \right| \\
&\leq \frac{C}{(N/L)^{\beta\kappa}} \sup_z \int |V(x, z)| |x|^\kappa dx \left( \left\| \langle \nabla_{x_j} \rangle J_x^{(k)} \langle \nabla_{x_j} \rangle^{-1} \right\|_{op} + \left\| \langle \nabla_{x_j} \rangle^{-1} J_x^{(k)} \langle \nabla_{x_j} \rangle \right\|_{op} \right) \\
&\quad \times Tr \langle \nabla_{x_j} \rangle \langle \nabla_{x_{k+1}} \rangle \tilde{\gamma}_{N,L}^{(k+1)}(s) \langle \nabla_{x_{k+1}} \rangle \langle \nabla_{x_j} \rangle + C_J \|\tilde{V}_{1,\lambda} - \tilde{V}_{1,1}\|_{L^1} Tr \langle \nabla_{r_j} \rangle^2 \langle \nabla_{r_{k+1}} \rangle^2 \gamma_{N,L}^{(k+1)} \\
&\leq C_J \left[ \frac{\sup_z \int |V(x, z)| |x|^\kappa dx}{(N/L)^{\beta\kappa}} + \|L(N/L)^\beta \tilde{V}(x, L(N/L)^\beta z) - \tilde{V}(x, z)\|_{L^1} \right].
\end{aligned}$$

Similarly, for  $II$  and  $IV$ , via Lemma 3.4, we have

$$\begin{aligned}
 (3.28) \quad |II| &= |Tr M_{s,t}^{(k)} (\delta(x_j - x_{k+1}) - \eta_\varepsilon(x_j - x_{k+1})) g(z_j - z_{k+1}) \tilde{\gamma}_{N,L}^{(k+1)}(s)| \\
 &\leq C\varepsilon^\kappa \|V\|_{L_z^\infty L_x^1} \int \eta(x) |x|^\kappa dx \left( \left\| \langle \nabla_{x_j} \rangle J_x^{(k)} \langle \nabla_{x_j} \rangle^{-1} \right\|_{op} + \left\| \langle \nabla_{x_j} \rangle^{-1} J_x^{(k)} \langle \nabla_{x_j} \rangle \right\|_{op} \right) \\
 &\quad \times Tr \langle \nabla_{x_j} \rangle \langle \nabla_{x_{k+1}} \rangle \tilde{\gamma}_{N,L}^{(k+1)}(s) \langle \nabla_{x_{k+1}} \rangle \langle \nabla_{x_j} \rangle \\
 &\leq C_J \varepsilon^\kappa,
 \end{aligned}$$

where the last inequality follows from Corollary 2.2, and

$$\begin{aligned}
 (3.29) \quad |IV| &= |Tr M_{s,t}^{(k)} (\eta_\varepsilon(x_j - x_{k+1}) - \delta(x_j - x_{k+1})) g(z_j - z_{k+1}) \tilde{\gamma}_{N,L}^{(k+1)}(s)| \\
 &\leq C\varepsilon^\kappa \|V\|_{L_z^\infty L_x^1} \int \eta(x) |x|^\kappa dx \left( \left\| \langle \nabla_{x_j} \rangle J_x^{(k)} \langle \nabla_{x_j} \rangle^{-1} \right\|_{op} + \left\| \langle \nabla_{x_j} \rangle^{-1} J_x^{(k)} \langle \nabla_{x_j} \rangle \right\|_{op} \right) \\
 &\quad \times Tr \langle \nabla_{x_j} \rangle \langle \nabla_{x_{k+1}} \rangle \tilde{\gamma}_{N,L}^{(k+1)}(s) \langle \nabla_{x_{k+1}} \rangle \langle \nabla_{x_j} \rangle \\
 &\leq C_J \varepsilon^\kappa,
 \end{aligned}$$

where the last inequality follows from Corollary 3.2. That is,

$$|II| \leq C_J \varepsilon^\kappa \text{ and } |IV| \leq C_J \varepsilon^\kappa.$$

Hence  $II$  and  $IV$  converge to 0 as  $\varepsilon \rightarrow 0$ , uniformly in  $N$  and  $L$ .

For  $III$ ,

$$\begin{aligned}
 (3.30) \quad |III| &= \left| Tr M_{s,t}^{(k)} \eta_\varepsilon(x_j - x_{k+1}) g(z_j - z_{k+1}) \left( \tilde{\gamma}_{N,L}^{(k+1)}(s) - \tilde{\gamma}^{(k+1)}(s) \right) \right| \\
 &\leq \left| Tr M_{s,t}^{(k)} \eta_\varepsilon(x_j - x_{k+1}) g(z_j - z_{k+1}) \frac{1}{1 + \theta \langle \nabla_{r_{k+1}} \rangle} \left( \tilde{\gamma}_{N,L}^{(k+1)}(s) - \tilde{\gamma}^{(k+1)}(s) \right) \right| \\
 &\quad + \left| Tr M_{s,t}^{(k)} \eta_\varepsilon(x_j - x_{k+1}) g(z_j - z_{k+1}) \frac{\theta \langle \nabla_{r_{k+1}} \rangle}{1 + \theta \langle \nabla_{r_{k+1}} \rangle} \left( \tilde{\gamma}_{N,L}^{(k+1)}(s) - \tilde{\gamma}^{(k+1)}(s) \right) \right|.
 \end{aligned}$$

The first term in the above estimate goes to zero as  $N, 1/L \rightarrow \infty$  for every  $\theta > 0$ , since we have condition (3.17) and  $M_{s,t}^{(k)} \eta_\varepsilon(x_j - x_{k+1}) g(z_j - z_{k+1}) \frac{1}{1 + \theta \langle \nabla_{r_{k+1}} \rangle}$  is a compact operator. Due to the energy bounds on  $\tilde{\gamma}_{N,L}^{(k+1)}$  and  $\tilde{\gamma}^{(k+1)}$ , the second term tends to zero as  $\theta \rightarrow 0$ , uniformly in  $N$  and  $L$ .

Putting together the estimates for  $I - IV$ , we have established (3.26). Hence, we have obtained Theorem 3.5.  $\square$

Combining Corollary 3.2 and Theorem 3.5, we see that  $\tilde{\gamma}_x^{(k)}$  solves the 2D Gross-Pitaevskii hierarchy with the desired coupling constant  $g_0 = \frac{4}{\pi^2} \iint V(x, z_1 - z_2) dx |\cos(z_1) \cos(z_2)|^2 dz_1 dz_2$ .

*Corollary 3.6.* Assume  $L(N/L)^\beta \rightarrow 1^-$ , and let  $\Gamma(t) = \{\tilde{\gamma}^{(k)}\}_{k=1}^\infty$  be a limit point of

$$\left\{ \Gamma_{N,L}(t) = \left\{ \tilde{\gamma}_{N,L}^{(k)}(t) \right\}_{k=1}^N \right\}$$

with respect to the product topology  $\tau_{prod}$ . Then  $\left\{ \tilde{\gamma}_x^{(k)} = Tr_z \tilde{\gamma}^{(k)} \right\}_{k=1}^\infty$  is a solution to the 2D Gross-Pitaevskii hierarchy subject to initial data  $\tilde{\gamma}_x^{(k)}(0) = |\phi_0\rangle \langle \phi_0|^{\otimes k}$  with coupling constant

$$g_0 = \frac{4}{\pi^2} \iint V(x, z_1 - z_2) dx |\cos(z_1) \cos(z_2)|^2 dz_1 dz_2,$$

which, rewritten in integral form, is

$$(3.31) \quad \tilde{\gamma}_x^{(k)}(t) = U^{(k)}(t) \tilde{\gamma}_x^{(k)}(0) - i g_0 \sum_{j=1}^k \int_0^t U^{(k)}(t-s) Tr_{x_{k+1}} \left[ \delta(x_j - x_{k+1}), \tilde{\gamma}_x^{(k+1)}(s) \right] ds.$$

*Proof.* The inhomogeneous term in hierarchy (3.16) is

$$\begin{aligned}
& i \int_0^t U^{(k)}(t-s) Tr_{x_{k+1}} Tr_z \left[ \delta(x_j - x_{k+1}) \int V(x, z_j - z_{k+1}) dx, \tilde{\gamma}^{(k+1)}(s) \right] ds \\
&= i \int_0^t U^{(k)}(t-s) Tr_{x_{k+1}} Tr_z \left( \delta(x_j - x_{k+1}) \int V(x, z_j - z_{k+1}) dx \tilde{\gamma}^{(k+1)}(s) \right) ds \\
&\quad - i \int_0^t U^{(k)}(t-s) Tr_{x_{k+1}} Tr_z \left( \delta(x'_j - x'_{k+1}) \int V(x, z'_j - z'_{k+1}) dx \tilde{\gamma}^{(k+1)}(s) \right) ds \\
&= I - II.
\end{aligned}$$

From Corollary 3.2, we have

$$\begin{aligned}
I &= i \int_0^t U^{(k)}(t-s) Tr_{x_{k+1}} \left( \delta(x_j - x_{k+1}) \tilde{\gamma}_x^{(k+1)}(s) Tr_z \left( \int V(x, z_j - z_{k+1}) dx \prod_{j=1}^{k+1} \frac{2}{\pi} \cos(z_j) \cos(z'_j) \right) \right) ds \\
&= i g_0 \int_0^t U^{(k)}(t-s) Tr_{x_{k+1}} \left( \delta(x_j - x_{k+1}) \tilde{\gamma}_x^{(k+1)}(s) \right) ds.
\end{aligned}$$

In the same manner we can see that

$$II = i g_0 \int_0^t U^{(k)}(t-s) Tr_{x_{k+1}} \left( \delta(x'_j - x'_{k+1}) \tilde{\gamma}_x^{(k+1)}(s) \right) ds.$$

In summary, we have

$$\tilde{\gamma}_x^{(k)}(t) = U^{(k)}(t) \tilde{\gamma}_x^{(k)}(0) - i g_0 \sum_{j=1}^k \int_0^t U^{(k)}(t-s) Tr_{x_{k+1}} \left[ \delta(x_j - x_{k+1}), \tilde{\gamma}_x^{(k+1)}(s) \right] ds.$$

□

**3.3. Uniqueness of the 2D GP Hierarchy.** By Bourgain [6], as we are below the Gagliardo-Nirenberg threshold here, we have the  $H^1$  global wellposedness for the  $\mathbb{T}^2$  focusing cubic NLS (1.11). Thus, when  $\tilde{\gamma}_x^{(k)}(0) = |\phi_0\rangle\langle\phi_0|^{\otimes k}$ , we know one solution to the focusing GP hierarchy (1.14), namely  $|\phi\rangle\langle\phi|^{\otimes k}$ , where  $\phi$  solves (1.11).

**Theorem 3.7** ([42, 43, 47]). *There is at most one nonnegative operator sequence*

$$\left\{ \gamma_x^{(k)} \right\}_{k=1}^{\infty} \in \bigotimes_{k \geq 1} C([0, T], \mathcal{L}_k^1(\mathbb{T}^2))$$

*that solves the 2D Gross-Pitaevskii hierarchy (3.31) subject to the energy condition*

$$(3.32) \quad Tr \left( \prod_{j=1}^k (1 - \Delta_{x_j}) \right) \gamma_x^{(k)} \leq C^k.$$

From Theorem 3.7, we conclude that the compact sequence

$$\left\{ \Gamma_{N,L}(t) = \left\{ \tilde{\gamma}_{N,L}^{(k)} \right\}_{k=1}^N \right\}$$

has only one  $L(N/L)^\beta \rightarrow 1^-$  limit point, namely

$$\tilde{\gamma}^{(k)} = \prod_{j=1}^k \frac{2}{\pi} \phi(t, x_j) \bar{\phi}(t, x'_j) \cos(z_j) \cos(z'_j).$$

We then infer that as trace class operators

$$\tilde{\gamma}_{N,L}^{(k)} \rightarrow \prod_{j=1}^k \frac{2}{\pi} \phi(t, x_j) \bar{\phi}(t, x'_j) \cos(z_j) \cos(z'_j) \text{ weak}^*.$$

Since the limit point  $\tilde{\gamma}^{(k)}$  is an orthogonal projection, the well-known argument in [35,  $P_{296}$ ] upgrades the weak\* convergence to strong, by using Gr\"{u}mm's convergence theorem [63, Theorem 2.19].

**Acknowledgements** The author would like to thank his PhD advisors Xuwen Chen and Zhifei Zhang. The author received partial support from the Graduate School of Peking University.

#### APPENDIX A. BASIC OPERATOR FACTS AND SOBOLEV-TYPE LEMMAS

**Lemma A.1** (Hoffman-Ostenhof inequality[44]). *For  $\psi_N \in H^1(\mathbb{T}^{dN})$ , we have*

$$(A.1) \quad \|\nabla_x \sqrt{\rho_N}\|_{L^2(\mathbb{T}^d)} \leq \|\nabla_{x_1} \psi_N\|_{L^2(\mathbb{T}^{dN})}.$$

with the one-particle density

$$\rho_N(x) = \int_{\mathbb{T}^d} \cdots \int_{\mathbb{T}^d} |\psi_N(x, x_2, \dots, x_N)|^2 dx_2 \cdots dx_N.$$

*Proof.* We may assume  $\psi_N$  is a test function, So  $\rho_N(x) \in C^\infty(\mathbb{T}^d)$ . By Cauchy-Schwarz inequality, we have

$$\begin{aligned} \|\nabla_x \sqrt{\rho_N + \varepsilon}\|_{L^2}^2 &= \int \left| \frac{\nabla_x \rho_N}{2\sqrt{\rho_N + \varepsilon}} \right|^2 dx \\ &= \int \frac{|\int \psi_N \nabla_x \psi_N dx_2 \cdots dx_N|^2}{\rho_N(x) + \varepsilon} dx \\ &\leq \int |\nabla_x \psi_N|^2 dx_2 \cdots dx_N \frac{\rho_N(x)}{\rho_N(x) + \varepsilon} dx \\ &\leq \int |\nabla_x \psi_N|^2 dx dx_2 \cdots dx_N. \end{aligned}$$

Thus,  $\sqrt{\rho_N + \varepsilon}$  is uniform bounded in  $H^1(\mathbb{T}^d)$  norm. We note that  $-\Delta_x$  is a self-adjoint operator on  $H^1(\mathbb{T}^d)$  and  $\sqrt{\rho_N + \varepsilon}$  converges to  $\sqrt{\rho_N}$  in  $L^2(\mathbb{T}^d)$  norm as  $\varepsilon \rightarrow 0$ . From the definition of adjoint operator, we deduce that  $\sqrt{\rho_N} \in H^1(\mathbb{T}^d)$  and

$$\|\nabla_x \sqrt{\rho_N}\|_{L^2}^2 \leq \int |\nabla_x \psi_N|^2 dx dx_2 \cdots dx_N.$$

□

**Lemma A.2.** *Assume  $L(N/L)^\beta \rightarrow 1^-$ . For  $\delta \in (0, 1)$ , the multiplication operator  $V_{N,L}(r_1 - r_2)$  on  $L^2(\Omega_L^{\otimes 2})$  satisfies*

$$(A.2) \quad L|V_{N,L}(r_1 - r_2)| \leq C_\delta (N/L)^\delta \|V\|_{L_x^\infty L_x^{1+\delta}} (1 - \Delta_{x_1}),$$

$$(A.3) \quad L|V_{N,L}(r_1 - r_2)| \leq C_\delta \|V\|_{L_x^\infty L_x^1} (1 - \Delta_{x_1})^{1/2+\delta} (1 - \Delta_{x_2})^{1/2+\delta},$$

$$(A.4) \quad \|\langle \nabla_{x_1} \rangle^{-1} \langle \nabla_{x_2} \rangle^{-1} V(r_1 - r_2) \langle \nabla_{x_1} \rangle^{-1} \langle \nabla_{x_2} \rangle^{-1}\|_{op} \leq C \|V\|_{L_x^\infty L_x^1},$$

$$(A.5) \quad S_1^2 LV_{N,L}(r_1 - r_2) + LV_{N,L}(r_1 - r_2) S_1^2 \geq -C_\delta(V) (N/L)^{\beta+\delta} S_1^2 S_2^2,$$

where  $C_\delta(V)$  is dependent on  $V$ .

*Proof.* For estimate (A.2), by H\"{o}lder and Sobolev inequality, we get

$$\begin{aligned} (A.6) \quad \langle LV_{N,L}(r_1 - r_2) \varphi_L, \varphi_L \rangle &\leq \|LV_{N,L}\|_{L_x^\infty L_x^{1+}} \|\varphi_L\|_{L^2 L_{x_1}^\infty}^2 \\ &\leq C_\delta L(N/L)^{\beta+\delta} \|V\|_{L_x^\infty L_x^{1+\delta}} \|\sqrt{1 - \Delta_{x_1}} \varphi_L\|_{L^2}^2 \\ &= C_\delta L(N/L)^{\beta+\delta} \|V\|_{L_x^\infty L_x^{1+\delta}} \langle (1 - \Delta_{x_1}) \varphi_L, \varphi_L \rangle. \end{aligned}$$

With  $L(N/L)^\beta \rightarrow 1^-$ , we obtain estimate (A.2).

For estimate (A.3), we recall Littlewood-Paley projectors defined by (2.54) and decompose  $\varphi_L$  that

$$(A.7) \quad \varphi_L = \varphi_{L,1} + \varphi_{L,2},$$

where

$$\varphi_{L,1} = \sum_{0 \leq m_1 < m_2} P_{x_1, m_1} P_{x_2, m_2} \varphi_L,$$



and

$$\varphi_{L,2} = \sum_{0 \leq m_2 \leq m_1} P_{x_1, m_1} P_{x_2, m_2} \varphi_L.$$

Then we have

$$(A.8) \quad \|\sqrt{1 - \Delta_{x_1}} \varphi_{L,1}\|_{L^2} \leq \|\sqrt{1 - \Delta_{x_2}} \varphi_{L,1}\|_{L^2},$$

$$(A.9) \quad \|\sqrt{1 - \Delta_{x_2}} \varphi_{L,2}\|_{L^2} \leq \|\sqrt{1 - \Delta_{x_1}} \varphi_{L,2}\|_{L^2}.$$

By estimates (A.8), (A.9) and Sobolev inequality, we find that

$$(A.10) \quad \begin{aligned} & |\langle LV_{N,L}(r_1 - r_2) \varphi_L, \varphi_L \rangle| \\ & \leq \int L |V_{N,L}(r_1 - r_2)| (|\varphi_{L,1}|^2 + |\varphi_{L,2}|^2) dr_1 dr_2 \\ & \leq L \|V_{N,L}\|_{L_z^\infty L_x^1} (\|\varphi_{L,1}\|_{L^2 L_{x_1}^\infty}^2 + \|\varphi_{L,2}\|_{L^2 L_{x_2}^\infty}^2) \\ & \leq C_\delta L (N/L)^\beta \|V\|_{L_z^\infty L_x^1} \left( \|(\sqrt{1 - \Delta_{x_1}})^{1+2\delta} \varphi_{L,1}\|_{L^2}^2 + \|(\sqrt{1 - \Delta_{x_2}})^{1+2\delta} \varphi_{L,2}\|_{L^2}^2 \right) \\ & \leq C_\delta L (N/L)^\beta \|V\|_{L_z^\infty L_x^1} \langle (1 - \Delta_{x_1})^{1/2+\delta} (1 - \Delta_{x_2})^{1/2+\delta} \varphi_L, \varphi_L \rangle. \end{aligned}$$

With  $L(N/L)^\beta \rightarrow 1^-$ , we obtain estimate (A.3).

For estimate (A.4), in the same manner as estimate (A.3), we get

$$(A.11) \quad \langle |V(r_1 - r_2)| \varphi, \varphi \rangle \leq C \|V\|_{L_z^\infty L_x^1} \langle (1 - \Delta_{x_1})(1 - \Delta_{x_2}) \varphi, \varphi \rangle,$$

which is equivalent to

$$(A.12) \quad \langle \nabla_{x_1} \rangle^{-1} \langle \nabla_{x_2} \rangle^{-1} |V(r_1 - r_2)| \langle \nabla_{x_1} \rangle^{-1} \langle \nabla_{x_2} \rangle^{-1} \leq C \|V\|_{L_z^\infty L_x^1}.$$

By Property 2 in Lemma A.6, we obtain estimate (A.4).

For estimate (A.5), we decompose

$$(A.13) \quad \begin{aligned} & \langle S_1^2 LV_{N,L}(r_1 - r_2) \varphi_L, \varphi_L \rangle \\ & = \langle (1 - \Delta_{x_1})(LV_{N,L}(r_1 - r_2) \varphi_L), \varphi_L \rangle + \langle (-\partial_{z_1}^2 - 1/L^2)(LV_{N,L}(r_1 - r_2) \varphi_L), \varphi_L \rangle \\ & = A + B. \end{aligned}$$

It needs only to control  $A$  and  $B$ . Using the identity  $\nabla_{x_1} V_{N,L}(r_1 - r_2) = -\nabla_{x_2} V_{N,L}(r_1 - r_2)$  and integration by parts, we get

$$\begin{aligned} A & = \langle \nabla_{x_1} (LV_{N,L}(r_1 - r_2) \varphi_L), \nabla_{x_1} \varphi_L \rangle \\ & = \langle LV_{N,L}(r_1 - r_2) \nabla_{x_1} \varphi_L, \nabla_{x_1} \varphi_L \rangle + \langle L(\nabla_{x_1} V_{N,L})(r_1 - r_2) \varphi_L, \nabla_{x_1} \varphi_L \rangle \\ & = \langle LV_{N,L}(r_1 - r_2) \nabla_{x_1} \varphi_L, \nabla_{x_1} \varphi_L \rangle - \langle L(\nabla_{x_2} V_{N,L})(r_1 - r_2) \varphi_L, \nabla_{x_1} \varphi_L \rangle \\ & = \langle LV_{N,L}(r_1 - r_2) \nabla_{x_1} \varphi_L, \nabla_{x_1} \varphi_L \rangle + \int LV_{N,L}(r_1 - r_2) \nabla_{x_2} \varphi_L \nabla_{x_1} \bar{\varphi}_L dr_1 dr_2 \\ & \quad + \int LV_{N,L}(r_1 - r_2) \varphi_L \nabla_{x_2} \nabla_{x_1} \bar{\varphi}_L dr_1 dr_2 \\ & = A_1 + A_2 + A_3. \end{aligned}$$

By Hölder and Sobolev inequality, we have

$$(A.14) \quad \begin{aligned} |A_1| & \leq L \|V_{N,L}\|_{L_z^\infty L_x^2} \|\nabla_{x_1} \varphi_L\|_{L^2 L_{x_2}^4}^2 \\ & \leq L (N/L)^{2\beta} \|V\|_{L_z^\infty L_x^2} \|\sqrt{1 - \Delta_{x_2}} \nabla_{x_1} \varphi_L\|_{L^2} \|\nabla_{x_1} \varphi_L\|_{L^2} \\ & \leq L (N/L)^{2\beta} \|V\|_{L_z^\infty L_x^2} \langle S_1^2 S_2^2 \varphi_L, \varphi_L \rangle. \end{aligned}$$

Estimated in the same way as  $A_1$ ,

$$(A.15) \quad \begin{aligned} |A_2| & \leq \int L |V_{N,L}(r_1 - r_2)| (|\nabla_{x_1} \varphi_L|^2 + |\nabla_{x_2} \varphi_L|^2) dr_1 dr_2 \\ & \leq L (N/L)^{2\beta} \|V\|_{L_z^\infty L_x^2} \langle S_1^2 S_2^2 \varphi_L, \varphi_L \rangle. \end{aligned}$$

For  $A_3$ , we have

$$(A.16) \quad \begin{aligned} |A_3| &\leq \|LV_{N,L}(r_1 - r_2)\varphi_L\|_{L^2} \|\nabla_{x_1} \nabla_{x_2} \varphi_L\|_{L^2} \\ &\leq CL(N/L)^{2\beta} \|V\|_{L_z^\infty L_x^2} \langle (1 - \Delta_{x_1})(1 - \Delta_{x_2})\varphi_L, \varphi_L \rangle, \end{aligned}$$

where we have used estimate (A.3) with  $\delta = 1/2$  and  $V$  replaced by  $V^2$  in the last inequality.

Hence we have

$$(A.17) \quad |A| \leq CL(N/L)^{2\beta} \|V\|_{L_z^\infty L_x^2} \langle (1 - \Delta_{x_1})(1 - \Delta_{x_2})\varphi_L, \varphi_L \rangle.$$

Next, we decompose  $B$  into two terms

$$(A.18) \quad \begin{aligned} B &= \langle (-\partial_{z_1}^2 - 1/L^2)LV_{N,L}(r_1 - r_2)\varphi_L, P_{z_1, >1}\varphi_L \rangle \\ &= \langle -\partial_{z_1}^2 (LV_{N,L}(r_1 - r_2)\varphi_L), P_{z_1, >1}\varphi_L \rangle - L^{-2} \langle LV_{N,L}(r_1 - r_2)\varphi_L, P_{z_1, >1}\varphi_L \rangle \\ &= B_1 + B_2. \end{aligned}$$

For  $B_1$ , we expand

$$(A.19) \quad B_1 = B_{11} + B_{12},$$

where

$$\begin{aligned} B_{11} &= \langle L(\partial_{z_1} V_{N,L}(r_1 - r_2))\varphi_L, \partial_{z_1} P_{z_1, >1}\varphi_L \rangle, \\ B_{12} &= \langle LV_{N,L}(r_1 - r_2)\partial_{z_1}\varphi_L, \partial_{z_1} P_{z_1, >1}\varphi_L \rangle \end{aligned}$$

For  $B_{11}$ , applying Hölder inequality at  $x_2$ , we obtain

$$(A.20) \quad \begin{aligned} |B_{11}| &\leq \|L\partial_{z_1} V_{N,L}(r_1 - r_2)\|_{L_{z_2}^\infty L_{x_2}^{1+}} \|\varphi_L\|_{L^2 L_{x_2}^\infty} \|\partial_{z_1} P_{z_1, >1}\varphi_L\|_{L^2 L_{x_2}^\infty} \\ &\leq C_\delta L(N/L)^{2\beta+\delta} \|\partial_z V\|_{L_z^\infty L_x^{1+\delta}} \|S_2 \varphi_L\|_{L^2} \|S_2 \partial_{z_1} P_{z_1, >1}\varphi_L\|_{L^2} \\ &\leq C_\delta L(N/L)^{2\beta+\delta} \|\partial_z V\|_{L_z^\infty L_x^{1+\delta}} \langle S_1^2 S_2^2 \varphi_L, \varphi_L \rangle. \end{aligned}$$

where we have used estimate (A.29) in the last inequality.

Computing in the same way, we have

$$(A.21) \quad \begin{aligned} |B_{12}| &\leq \|LV_{N,L}\|_{L_{z_2}^\infty L_{x_2}^{1+}} \|\partial_{z_1} \varphi_L\|_{L^2 L_{x_2}^\infty} \|\partial_{z_1} P_{z_1, >1}\varphi_L\|_{L^2 L_{x_2}^\infty} \\ &\leq \|LV_{N,L}\|_{L_{z_2}^\infty L_{x_2}^{1+}} \|S_2 \partial_{z_1} \varphi_L\|_{L^2} \|S_2 \partial_{z_1} P_{z_1, >1}\varphi_L\|_{L^2} \\ &\leq C_\delta (N/L)^{\beta+\delta} \|V\|_{L_z^\infty L_x^{1+\delta}} \|S_2 S_1 \varphi_L\|_{L^2}^2, \end{aligned}$$

where we have used (A.28) and (A.29) in the last inequality.

Hence, with  $L(N/L)^\beta \rightarrow 1^-$ , we have

$$(A.22) \quad |B_1| \leq C_\delta(V) L(N/L)^{2\beta+\delta} \|S_1 S_2 \varphi_L\|_{L^2}^2.$$

For  $B_2$ , we have

$$(A.23) \quad \begin{aligned} |B_2| &\leq L^{-2} \|LV_{N,L}\|_{L_{z_2}^\infty L_{x_2}^{1+}} \|\varphi_L\|_{L^2 L_{x_2}^\infty} \|P_{z_1, >1}\varphi_L\|_{L^2 L_{x_2}^\infty} \\ &\leq C_\delta L^{-1} (N/L)^{\beta+\delta} \|V\|_{L_z^\infty L_x^{1+\delta}} \|S_2 \varphi_L\|_{L^2} \|S_2 P_{z_1, >1}\varphi_L\|_{L^2} \\ &\leq C_\delta (N/L)^{\beta+\delta} \|V\|_{L_z^\infty L_x^{1+\delta}} \langle S_1^2 S_2^2 \varphi_L, \varphi_L \rangle, \end{aligned}$$

where we used  $L^{-2} P_{z_1, >1} \leq S_1^2 P_{z_1, >1}$  in the last inequality.

Putting (A.17) (A.22) and (A.23) together, with  $L(N/L)^\beta \rightarrow 1^-$ , we obtain estimate (A.5).  $\square$

**Lemma A.3.** *Let  $A : D(A) \mapsto H$ , be a positive selfadjoint operator in the Hilbert space  $H$ , and let  $0 < \alpha < 1$ . Then*

$$A^\alpha \leq (1 - \alpha)\eta^{-1}A + \alpha\eta^{\frac{\alpha}{1-\alpha}},$$

for  $\eta \in (0, 1)$ .

*Proof.* By the spectral representation and the inequality

$$\lambda^\alpha \leq \alpha \eta^{-1} \lambda + (1 - \alpha) \eta^{\frac{\alpha}{1-\alpha}}, \quad \forall \lambda \in (0, \infty),$$

we obtain

$$\begin{aligned} \langle A^\alpha u, u \rangle_H &= \int_0^\infty \lambda^\alpha d\|E_\lambda u\|_H^2 \\ &\leq (1 - \alpha) \eta^{-1} \int_0^\infty \lambda d\|E_\lambda u\|_H^2 + \alpha \eta^{\frac{\alpha}{1-\alpha}} \int_0^\infty d\|E_\lambda u\|_H^2 \\ &= (1 - \alpha) \eta^{-1} \langle Au, u \rangle_H + \alpha \eta^{\frac{\alpha}{1-\alpha}} \langle u, u \rangle_H. \end{aligned}$$

□

Recall Lemma 3.4, we now give the proof as follows.

**Lemma A.4.** *Let  $\rho \in L^1(\Omega)$  be a function compactly supported on  $\Omega$  such that*

$$\sup_z \int |\rho(x, z)| |x| dx < \infty$$

*and define  $\rho_{\varepsilon, \lambda}(x, z) = \varepsilon^{-2} \lambda^{-1} \rho(x/\varepsilon, z/\lambda)$  and  $g(z) = \int \rho(x, z) dx$ . Then, for every  $\kappa \in [0, 1)$ , there exists  $C > 0$ , such that*

$$\begin{aligned} &\left| \text{Tr} J^{(k)}(\rho_{\varepsilon, \lambda}(r_j - r_{k+1}) - \delta(x_j - x_{k+1})g(z_j - z_{k+1})) \gamma^{(k+1)} \right| \\ &\leq C \varepsilon^\kappa \sup_z \int |\rho(x, z)| |x|^\kappa dx \left( \|\langle \nabla_{x_j} \rangle^{-1} J^{(k)} \langle \nabla_{x_j} \rangle\|_{op} + \|\langle \nabla_{x_j} \rangle J^{(k)} \langle \nabla_{x_j} \rangle^{-1}\|_{op} \right) \text{Tr} \langle \nabla_{x_j} \rangle^2 \langle \nabla_{x_{k+1}} \rangle^2 \gamma^{(k+1)} \\ &\quad + C_J \|\rho_{1, \lambda} - \rho_{1, 1}\|_{L^1} \text{Tr} \langle \nabla_{r_j} \rangle^2 \langle \nabla_{r_{k+1}} \rangle^2 \gamma^{(k+1)} \end{aligned}$$

*for all nonnegative  $\gamma^{(k+1)} \in \mathcal{L}_{k+1}^1$ .*

*Proof.* We present a proof by modifying the proof in [47]. Such a method has been used by various authors, for example [19]. It suffices to prove the estimate for  $k = 1$ . Since the observable  $J^{(1)}$  can be written as a sum of a self-adjoint operator and an anti-self-adjoint operator, we may assume  $J^{(1)}$  is self-adjoint. We represent  $\gamma^{(2)}$  by  $\gamma^{(2)} = \sum_j \lambda_j |\varphi_j\rangle \langle \varphi_j|$ , where  $\varphi_j \in L^2(\Omega^2)$  and  $\lambda_j \geq 0$ . Then, we have

$$\begin{aligned} &\text{Tr} J^{(1)}(\rho_{\varepsilon, \lambda}(r_1 - r_2) - \delta(x_1 - x_2)g(z_1 - z_2)) \gamma^{(2)} \\ &= \sum_j \lambda_j \langle \varphi_j, J^{(1)}(\rho_{\varepsilon}(r_1 - r_2) - \delta(x_1 - x_2)g(z_1 - z_2)) \varphi_j \rangle \\ &= \sum_j \lambda_j \langle \psi_j, (\rho_{\varepsilon, \lambda}(r_1 - r_2) - \delta(x_1 - x_2)g(z_1 - z_2)) \varphi_j \rangle, \end{aligned}$$

where  $\psi_j = (J^{(1)} \otimes 1) \varphi_j$ . Then, we decompose

$$\langle \psi_j, (\rho_{\varepsilon, \lambda}(r_1 - r_2) - \delta(x_1 - x_2)g(z_1 - z_2)) \varphi_j \rangle = A_j + B_j,$$

where

$$\begin{aligned} A_j &= \langle \psi_j, (\rho_{\varepsilon, 1}(r_1 - r_2) - \delta(x_1 - x_2)g(z_1 - z_2)) \varphi_j \rangle, \\ B_j &= \langle \psi_j, (\rho_{\varepsilon, \lambda}(r_1 - r_2) - \rho_{\varepsilon, 1}(r_1 - r_2)) \varphi_j \rangle. \end{aligned}$$

For  $A_j$ , switching to Fourier space in the  $x$ -direction, we find

$$\begin{aligned} |A_j| &= \left| \int \overline{\psi_j}(n_1, n_2; z_1, z_2) \widehat{\varphi_j}(m_1, m_2; z_1, z_2) \rho_{1, \lambda}(x, z_1 - z_2) (e^{i\varepsilon x \cdot (n_1 - m_1)} - 1) \right. \\ &\quad \times \delta(n_1 + n_2 - m_1 - m_2) dx dz_1 dz_2 dn_1 dn_2 dm_1 dm_2 \Big| \\ &\leq \int |\widehat{\psi_j}(n_1, n_2; z_1, z_2)| |\widehat{\varphi_j}(m_1, m_2; z_1, z_2)| \delta(n_1 + n_2 - m_1 - m_2) \\ &\quad \times \left| \int \rho(x, z_1 - z_2) (e^{i\varepsilon x \cdot (n_1 - m_1)} - 1) dx \right| dz_1 dz_2 dn_1 dn_2 dm_1 dm_2. \end{aligned}$$

Using the inequality that  $\forall \kappa \in (0, 1)$

$$\begin{aligned} \left| e^{i\varepsilon x \cdot (n_1 - m_1)} - 1 \right| &\leq \varepsilon^\kappa |x|^\kappa |n_1 - m_1|^\kappa \\ &\leq \varepsilon^\kappa |x|^\kappa (|n_1|^\kappa + |m_1|^\kappa), \end{aligned}$$

we get

$$\begin{aligned} &|\langle \psi_j, (\rho_{\varepsilon,1}(r_1 - r_2) - \delta(x_1 - x_2)g(z_1 - z_2))\varphi_j \rangle| \\ &\leq \varepsilon^\kappa \sup_z \int |\rho(x, z)| |x|^\kappa dx \int \delta(n_1 + n_2 - m_1 - m_2) \\ &\quad (|n_1|^\kappa + |m_1|^\kappa) |\widehat{\psi}_j(n_1, n_2; z_1, z_2)| |\widehat{\varphi}_j(m_1, m_2; z_1, z_2)| dz_1 dz_2 dn_1 dn_2 dm_1 dm_2 \\ &= \varepsilon^\kappa \sup_z \int |\rho(x, z)| |x|^\kappa dx (I + II), \end{aligned}$$

where

$$\begin{aligned} I &= \int \delta(n_1 + n_2 - m_1 - m_2) |n_1|^\kappa |\widehat{\psi}_j(n_1, n_2; z_1, z_2)| |\widehat{\varphi}_j(m_1, m_2; z_1, z_2)| dz_1 dz_2 dn_1 dn_2 dm_1 dm_2, \\ II &= \int \delta(n_1 + n_2 - m_1 - m_2) |m_1|^\kappa |\widehat{\psi}_j(n_1, n_2; z_1, z_2)| |\widehat{\varphi}_j(m_1, m_2; z_1, z_2)| dz_1 dz_2 dn_1 dn_2 dm_1 dm_2. \end{aligned}$$

The estimates for  $I$  and  $II$  are similar, so we only deal with  $I$  explicitly.

$$\begin{aligned} I &\leq \int \delta(n_1 + n_2 - m_1 - m_2) \frac{\langle n_1 \rangle \langle n_2 \rangle}{\langle m_1 \rangle \langle m_2 \rangle} |\widehat{\psi}_j(n_1, n_2; z_1, z_2)| \\ &\quad \times \frac{\langle m_1 \rangle \langle m_2 \rangle}{\langle n_1 \rangle^{1-\kappa} \langle n_2 \rangle} |\widehat{\varphi}_j(m_1, m_2; z_1, z_2)| dz_1 dz_2 dn_1 dn_2 dm_1 dm_2 \\ &\leq \theta \int \delta(n_1 + n_2 - m_1 - m_2) \frac{\langle n_1 \rangle^2 \langle n_2 \rangle^2}{\langle m_1 \rangle^2 \langle m_2 \rangle^2} |\widehat{\psi}_j(n_1, n_2; z_1, z_2)|^2 dz_1 dz_2 dn_1 dn_2 dm_1 dm_2 \\ &\quad + \theta^{-1} \int \delta(n_1 + n_2 - m_1 - m_2) \frac{\langle m_1 \rangle^2 \langle m_2 \rangle^2}{\langle n_1 \rangle^{2(1-\kappa)} \langle n_2 \rangle^2} |\widehat{\varphi}_j(m_1, m_2; z_1, z_2)|^2 dz_1 dz_2 dn_1 dn_2 dm_1 dm_2 \\ &\leq \theta \langle \psi_j, (1 - \Delta_{x_1})(1 - \Delta_{x_2})\psi_j \rangle \sup_n \sum_{m_2} \frac{1}{\langle m_2 - n \rangle^2 \langle m_2 \rangle^2} \\ &\quad + \theta^{-1} \langle \varphi_j, (1 - \Delta_{x_1})(1 - \Delta_{x_2})\varphi_j \rangle \sup_n \sum_{m_2} \frac{1}{\langle m_2 - n \rangle^{2(1-\kappa)} \langle m_2 \rangle^2}. \end{aligned}$$

Since

$$\sup_n \sum_{m \in \mathbb{Z}^2} \frac{1}{\langle m - n \rangle^{2(1-\kappa)} \langle m \rangle^2} < \infty$$

for  $\kappa \in [0, 1)$ , we have

$$(A.24) \quad |A_j| \leq \theta \langle \psi_j, (1 - \Delta_{x_1})(1 - \Delta_{x_2})\psi_j \rangle + \theta^{-1} \langle \varphi_j, (1 - \Delta_{x_1})(1 - \Delta_{x_2})\varphi_j \rangle.$$

Hence, we obtain

(A.25)

$$\begin{aligned}
& \left| Tr J^{(1)}(\rho_{\varepsilon,1}(r_1 - r_2) - \delta(x_1 - x_2)g(z_1 - z_2))\gamma^{(2)} \right| \\
& \leq C\varepsilon^\kappa \sup_z \int |\rho(x, z)| |x|^\kappa dx \left( \theta Tr \langle \nabla_{x_1} \rangle^2 \langle \nabla_{x_2} \rangle^2 \gamma^{(2)} + \theta^{-1} Tr J^{(1)} \langle \nabla_{x_1} \rangle^2 \langle \nabla_{x_2} \rangle^2 J^{(1)} \gamma^{(2)} \right) \\
& = C\varepsilon^\kappa \sup_z \int |\rho(x, z)| |x|^\kappa dx \left( \theta Tr \langle \nabla_{x_1} \rangle^2 \langle \nabla_{x_2} \rangle^2 \gamma^{(2)} \right. \\
& \quad \left. + \theta^{-1} Tr \langle \nabla_{x_1} \rangle^{-1} \langle \nabla_{x_2} \rangle^{-1} J^{(1)} \langle \nabla_{x_1} \rangle^2 J^{(1)} \langle \nabla_{x_1} \rangle^{-1} \langle \nabla_{x_2} \rangle^2 \gamma^{(2)} \langle \nabla_{x_1} \rangle \langle \nabla_{x_2} \rangle \right) \\
& \leq C\varepsilon^\kappa \sup_z \int |\rho(x, z)| |x|^\kappa dx \left( \theta + \theta^{-1} \| \langle \nabla_{x_1} \rangle^{-1} J^{(1)} \langle \nabla_{x_1} \rangle \|_{op} \| \langle \nabla_{x_1} \rangle J^{(1)} \langle \nabla_{x_1} \rangle^{-1} \|_{op} \right) Tr \langle \nabla_{x_1} \rangle^2 \langle \nabla_{x_2} \rangle^2 \gamma^{(2)} \\
& = C\varepsilon^\kappa \sup_z \int |\rho(x, z)| |x|^\kappa dx \left( \| \langle \nabla_{x_1} \rangle^{-1} J^{(1)} \langle \nabla_{x_1} \rangle \|_{op} + \| \langle \nabla_{x_1} \rangle J^{(1)} \langle \nabla_{x_1} \rangle^{-1} \|_{op} \right) Tr \langle \nabla_{x_1} \rangle^2 \langle \nabla_{x_2} \rangle^2 \gamma^{(2)},
\end{aligned}$$

where we have taken  $\theta = \| \langle \nabla_{x_1} \rangle J^{(1)} \langle \nabla_{x_1} \rangle^{-1} \|_{op}$  in the last line.

For  $B_j$ , we use the operator inequality (also see [31, (A.63)])

$$|V(r_1 - r_2)| \leq C \|V\|_{L^1} (1 - \Delta_{r_1})(1 - \Delta_{r_2}),$$

which can be estimated in the same way as estimate (A.4). Then, we get

$$\begin{aligned}
(A.26) \quad |B_j| & \leq \theta \langle \psi_i, |(\rho_{\varepsilon,\lambda} - \rho_{\varepsilon,1})(r_1 - r_2)| \psi_i \rangle + \theta^{-1} \langle \varphi_i, |(\rho_{\varepsilon,\lambda} - \rho_{\varepsilon,1})(r_1 - r_2)| \varphi_i \rangle \\
& \leq C \| \rho_{\varepsilon,\lambda} - \rho_{\varepsilon,1} \|_{L_z^1 L_x^1} \left( \theta \langle \psi_i, (1 - \Delta_{r_1})(1 - \Delta_{r_2}) \psi_i \rangle + \theta^{-1} \langle \varphi_i, (1 - \Delta_{r_1})(1 - \Delta_{r_2}) \varphi_i \rangle \right) \\
& = C \| \rho_{1,\lambda} - \rho_{1,1} \|_{L_z^1 L_x^1} \left( \theta \langle \psi_i, (1 - \Delta_{r_1})(1 - \Delta_{r_2}) \psi_i \rangle + \theta^{-1} \langle \varphi_i, (1 - \Delta_{r_1})(1 - \Delta_{r_2}) \varphi_i \rangle \right).
\end{aligned}$$

Similarly, we have

$$(A.27) \quad |Tr J^{(1)}(\rho_{\varepsilon,\lambda}(r_1 - r_2) - \rho_{\varepsilon,\lambda}(r_1 - r_2))\gamma^{(2)}| \leq C_J \| \rho_{1,\lambda} - \rho_{1,1} \|_{L^1} Tr \langle \nabla_{r_1} \rangle^2 \langle \nabla_{r_2} \rangle^2 \gamma^{(2)}.$$

Together with estimate (A.25) and (A.27), we complete the proof.  $\square$

**Lemma A.5.** *Recall*

$$\begin{aligned}
S^2 &= 1 - \Delta_r - 1/L^2, \\
\tilde{S}^2 &= 1 - \Delta_x - \partial_z^2/L^2 - 1/L^2.
\end{aligned}$$

We have

$$(A.28) \quad (1 - \Delta_r) \leq 2L^{-2} S^2,$$

$$(A.29) \quad (1 - \Delta_r) P_{z,>1} \leq 2S^2 P_{z,>1},$$

$$(A.30) \quad \tilde{S}^2 \tilde{P}_1 \geq L^{-2} \tilde{P}_1,$$

$$(A.31) \quad 2\tilde{S}^2 \geq 1 - \Delta_r$$

*Proof.* For (A.28), we have

$$1 - \Delta_r = S^2 + 1/L^2 \leq 2L^{-2} S^2.$$

For (A.29), we note that

$$L^{-2} P_{z,>1} \leq (-\partial_z^2 - 1/L^2) P_{z,>1} \leq S^2 P_{z,>1}.$$

Thus, we obtain

$$(1 - \Delta_r) P_{z,>1} = (S^2 + 1/L^2) P_{z,>1} \leq 2S^2 P_{z,>1}.$$

For (A.30), we have

$$L^{-2} \tilde{P}_1 \leq (-\partial_z^2/L^2 - 1/L^2) \tilde{P}_1 \leq \tilde{S}^2 \tilde{P}_1.$$

For (A.31), we note that

$$(A.32) \quad 2(-\partial_z^2 - 1) \tilde{P}_1 = -\partial_z^2 \tilde{P}_1 + (-\partial_z^2 - 2) \tilde{P}_1 \geq -\partial_z^2 \tilde{P}_1.$$

Then we have

$$\begin{aligned}
 (A.33) \quad 2\tilde{S}^2 &\geq 2(1 - \Delta_x) + 2L^{-2}(-\partial_z^2 - 1)\tilde{P}_1 \\
 &\geq 2(1 - \Delta_x) - L^{-2}\partial_z^2\tilde{P}_1 \\
 &\geq 2(1 - \Delta_x) - \partial_z^2\tilde{P}_1.
 \end{aligned}$$

Noting that  $-\partial_z^2\tilde{P}_0 = \tilde{P}_0$ , we obtain

$$(A.34) \quad 1 - \Delta_x \geq 1 - \partial_z^2\tilde{P}_0.$$

Putting (A.33) and (A.34) together, we establish (A.31).  $\square$

We need the following facts as well. The proofs are elementary and we omit them.

- Lemma A.6.** 1. Suppose that  $A \geq 0$ ,  $P_j = P_j^*$ , and  $I = P_0 + P_1$ . Then  $A \leq 2P_0AP_0 + 2P_1AP_1$ .  
 2. If  $A \geq B \geq 0$ , and  $AB = BA$ , then  $A^\alpha \geq B^\alpha$  for any  $\alpha > 0$ . Especially, if  $\alpha = 2$ , then  $\|A\|_{op} \geq \|B\|_{op}$ .  
 3. If  $A_1 \geq A_2 \geq 0$ ,  $B_1 \geq B_2 \geq 0$  and  $A_iB_j = B_jA_i$  for all  $1 \leq i, j \leq 2$ , then  $A_1B_1 \geq A_2B_2$ .  
 4. If  $A \geq 0$  and  $AB = BA$ , then  $A^{1/2}B = BA^{1/2}$ .

**Lemma A.7.** Suppose  $\sigma : L^2(\Omega^k) \mapsto L^2(\Omega^k)$  has kernel

$$\sigma(\mathbf{r}_k, \mathbf{r}'_k) = \int \psi(\mathbf{r}_k, \mathbf{r}_{N-k}) \overline{\psi}(\mathbf{r}'_k, \mathbf{r}_{N-k}) d\mathbf{r}_{N-k},$$

for some  $\psi \in L^2(\Omega^k)$ , and let  $A, B : L^2(\Omega^k) \mapsto L^2(\Omega^k)$ . Then the composition  $A\sigma B$  has kernel

$$(A\sigma B)(\mathbf{r}_k, \mathbf{r}'_k) = \int (A\psi)(\mathbf{r}_k, \mathbf{r}_{N-k}) \overline{(B^*\psi)}(\mathbf{r}'_k, \mathbf{r}_{N-k}) d\mathbf{r}_{N-k}.$$

It follows that

$$Tr A\sigma B = \langle A\psi, B^*\psi \rangle.$$

**Lemma A.8.** Let  $P_M^j$  be the orthogonal projection onto the sum of the first  $M$  eigenspaces with respect to the spectral decomposition of  $L^2(\Omega)$  to the operator  $-\Delta_{r_j}$  and

$$P_M^{(k)} = \prod_{j=1}^k P_M^j.$$

1. Suppose that  $J^{(k)}$  is a compact operator. Then  $J_M^{(k)} := P_M^{(k)} J^{(k)} P_M^{(k)}$  converges to  $J^{(k)}$  in the operator norm.
2.  $\Delta_{r_j} J_M^{(k)}$  and  $J_M^{(k)} \Delta_{r_j}$  are bounded operators.
3. There exists a countable dense subset  $\{T_i\}$  of the closed unit ball in the space of bounded operators on  $L^2(\Omega^k)$  such that  $T_i$  is compact and in fact for each  $i$  there exists  $M$  (depending on  $i$ ) such that  $T_i = P_M^{(k)} T_i P_M^{(k)}$ .

## REFERENCES

- [1] R. Adami, F. Golse, and A. Teta. Rigorous derivation of the cubic NLS in dimension one. *J. Stat. Phys.*, 127(6):1193–1220, 2007.
- [2] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell. Observation of bose-einstein condensation in a dilute atomic vapor. *Science*, 269(5221):198–201, 1995.
- [3] N. Benedikter, G. de Oliveira, and B. Schlein. Quantitative derivation of the Gross-Pitaevskii equation. *Comm. Pure Appl. Math.*, 68(8):1399–1482, 2015.
- [4] C. Boccato, C. Brennecke, S. Cenatiempo, and B. Schlein. Bogoliubov theory in the Gross-Pitaevskii limit. *Acta Math.*, 222(2):219–335, 2019.
- [5] L. Boßmann. Derivation of the 2d Gross-Pitaevskii equation for strongly confined 3d bosons. *arXiv preprint arXiv:1907.04547*, 2019.
- [6] J. Bourgain. Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. I. Schrödinger equations. *Geom. Funct. Anal.*, 3(2):107–156, 1993.
- [7] L. Chen, J. O. Lee, and B. Schlein. Rate of convergence towards Hartree dynamics. *J. Stat. Phys.*, 144(4):872–903, 2011.
- [8] T. Chen, C. Hainzl, N. Pavlović, and R. Seiringer. Unconditional uniqueness for the cubic Gross-Pitaevskii hierarchy via quantum de Finetti. *Comm. Pure Appl. Math.*, 68(10):1845–1884, 2015.
- [9] T. Chen and N. Pavlović. On the Cauchy problem for focusing and defocusing Gross-Pitaevskii hierarchies. *Discrete Contin. Dyn. Syst.*, 27(2):715–739, 2010.

- [10] T. Chen and N. Pavlović. The quintic NLS as the mean field limit of a boson gas with three-body interactions. *J. Funct. Anal.*, 260(4):959–997, 2011.
- [11] T. Chen and N. Pavlović. A new proof of existence of solutions for focusing and defocusing Gross-Pitaevskii hierarchies. *Proc. Amer. Math. Soc.*, 141(1):279–293, 2013.
- [12] T. Chen and N. Pavlović. Derivation of the cubic NLS and Gross-Pitaevskii hierarchy from manybody dynamics in  $d = 3$  based on spacetime norms. *Ann. Henri Poincaré*, 15(3):543–588, 2014.
- [13] T. Chen and N. Pavlović. Higher order energy conservation and global well-posedness of solutions for Gross-Pitaevskii hierarchies. *Comm. Partial Differential Equations*, 39(9):1597–1634, 2014.
- [14] T. Chen, N. Pavlović, and N. Tzirakis. Energy conservation and blowup of solutions for focusing Gross-Pitaevskii hierarchies. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 27(5):1271–1290, 2010.
- [15] T. Chen and K. Taliaferro. Derivation in strong topology and global well-posedness of solutions to the Gross-Pitaevskii hierarchy. *Comm. Partial Differential Equations*, 39(9):1658–1693, 2014.
- [16] X. Chen. Collapsing estimates and the rigorous derivation of the 2d cubic nonlinear Schrödinger equation with anisotropic switchable quadratic traps. *J. Math. Pures Appl.* (9), 98(4):450–478, 2012.
- [17] X. Chen. Second order corrections to mean field evolution for weakly interacting bosons in the case of three-body interactions. *Arch. Ration. Mech. Anal.*, 203(2):455–497, 2012.
- [18] X. Chen. On the rigorous derivation of the 3D cubic nonlinear Schrödinger equation with a quadratic trap. *Arch. Ration. Mech. Anal.*, 210(2):365–408, 2013.
- [19] X. Chen and J. Holmer. On the rigorous derivation of the 2D cubic nonlinear Schrödinger equation from 3D quantum many-body dynamics. *Arch. Ration. Mech. Anal.*, 210(3):909–954, 2013.
- [20] X. Chen and J. Holmer. Correlation structures, many-body scattering processes, and the derivation of the Gross-Pitaevskii hierarchy. *Int. Math. Res. Not. IMRN*, (10):3051–3110, 2016.
- [21] X. Chen and J. Holmer. Focusing quantum many-body dynamics: the rigorous derivation of the 1D focusing cubic nonlinear Schrödinger equation. *Arch. Ration. Mech. Anal.*, 221(2):631–676, 2016.
- [22] X. Chen and J. Holmer. On the Klainerman-Machedon conjecture for the quantum BBGKY hierarchy with self-interaction. *J. Eur. Math. Soc. (JEMS)*, 18(6):1161–1200, 2016.
- [23] X. Chen and J. Holmer. Focusing quantum many-body dynamics, II: The rigorous derivation of the 1D focusing cubic nonlinear Schrödinger equation from 3D. *Anal. PDE*, 10(3):589–633, 2017.
- [24] X. Chen and J. Holmer. The rigorous derivation of the 2D cubic focusing NLS from quantum many-body evolution. *Int. Math. Res. Not. IMRN*, (14):4173–4216, 2017.
- [25] X. Chen and J. Holmer. The derivation of the  $T^3$  energy-critical NLS from quantum many-body dynamics. *Invent. Math.*, 217(2):433–547, 2019.
- [26] X. Chen and P. Smith. On the unconditional uniqueness of solutions to the infinite radial Chern-Simons-Schrödinger hierarchy. *Anal. PDE*, 7(7):1683–1712, 2014.
- [27] M. Christandl, R. König, G. Mitchison, and R. Renner. One-and-a-half quantum de Finetti theorems. *Comm. Math. Phys.*, 273(2):473–498, 2007.
- [28] S. L. Cornish, N. R. Claussen, J. L. Roberts, E. A. Cornell, and C. E. Wieman. Stable Rb-85 Bose-Einstein Condensates with Widely Tunable Interactions. *Phys. Rev. Lett.*, 85:1795–1798, 2000.
- [29] K. B. Davis, M. O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle. Bose-Einstein Condensation in a Gas of Sodium Atoms. *Phys. Rev. Lett.*, 75:3969–3973, Nov 1995.
- [30] E. A. Donley, N. R. Claussen, S. L. Cornish, J. L. Roberts, E. A. Cornell, and C. E. Wieman. Dynamics of collapsing and exploding Bose-Einstein condensates. *Nature*, 412(6844):295–299, 2001.
- [31] A. Elgart, L. Erdős, B. Schlein, and H.-T. Yau. Gross-Pitaevskii equation as the mean field limit of weakly coupled bosons. *Arch. Ration. Mech. Anal.*, 179(2):265–283, 2006.
- [32] L. Erdős, B. Schlein, and H.-T. Yau. Derivation of the Gross-Pitaevskii hierarchy for the dynamics of Bose-Einstein condensate. *Comm. Pure Appl. Math.*, 59(12):1659–1741, 2006.
- [33] L. Erdős, B. Schlein, and H.-T. Yau. Derivation of the cubic non-linear Schrödinger equation from quantum dynamics of many-body systems. *Invent. Math.*, 167(3):515–614, 2007.
- [34] L. Erdős, B. Schlein, and H.-T. Yau. Rigorous derivation of the Gross-Pitaevskii equation with a large interaction potential. *J. Amer. Math. Soc.*, 22(4):1099–1156, 2009.
- [35] L. Erdős, B. Schlein, and H.-T. Yau. Derivation of the Gross-Pitaevskii equation for the dynamics of Bose-Einstein condensate. *Ann. of Math.* (2), 172(1):291–370, 2010.
- [36] L. Erdős, B. Schlein, and H.-T. Yau. Rigorous Derivation of the Gross-Pitaevskii Equation. *Phys. Rev. Lett.*, 98:040404, Jan 2007.
- [37] J. Fröhlich, A. Knowles, and S. Schwarz. On the mean-field limit of bosons with Coulomb two-body interaction. *Comm. Math. Phys.*, 288(3):1023–1059, 2009.
- [38] P. Gressman, V. Sohinger, and G. Staffilani. On the uniqueness of solutions to the periodic 3D Gross-Pitaevskii hierarchy. *J. Funct. Anal.*, 266(7):4705–4764, 2014.
- [39] M. Grillakis and M. Machedon. Pair excitations and the mean field approximation of interacting bosons, I. *Comm. Math. Phys.*, 324(2):601–636, 2013.
- [40] M. Grillakis and M. Machedon. Pair excitations and the mean field approximation of interacting bosons, II. *Comm. Partial Differential Equations*, 42(1):24–67, 2017.
- [41] K. Hepp. The classical limit for quantum mechanical correlation functions. *Comm. Math. Phys.*, 35:265–277, 1974.
- [42] S. Herr and V. Sohinger. The Gross-Pitaevskii hierarchy on general rectangular tori. *Arch. Ration. Mech. Anal.*, 220(3):1119–1158, 2016.

- [43] S. Herr and V. Sohinger. Unconditional uniqueness results for the nonlinear Schrödinger equation. *Commun. Contemp. Math.*, 21(7):1850058, 33, 2019.
- [44] M. Hoffmann-Ostenhof and T. Hoffmann-Ostenhof. “Schrödinger inequalities” and asymptotic behavior of the electron density of atoms and molecules. *Phys. Rev. A* (3), 16(5):1782–1785, 1977.
- [45] Y. Hong, K. Taliaferro, and Z. Xie. Unconditional uniqueness of the cubic Gross-Pitaevskii hierarchy with low regularity. *SIAM J. Math. Anal.*, 47(5):3314–3341, 2015.
- [46] L. Khaykovich, F. Schreck, G. Ferrari, T. Bourdel, J. Cubizolles, L. D. Carr, Y. Castin, and C. Salomon. Formation of a matter-wave bright soliton. *Science*, 296(5571):1290–1293, 2002.
- [47] K. Kirkpatrick, B. Schlein, and G. Staffilani. Derivation of the two-dimensional nonlinear Schrödinger equation from many body quantum dynamics. *Amer. J. Math.*, 133(1):91–130, 2011.
- [48] S. Klainerman and M. Machedon. On the uniqueness of solutions to the Gross-Pitaevskii hierarchy. *Comm. Math. Phys.*, 279(1):169–185, 2008.
- [49] A. Knowles and P. Pickl. Mean-field dynamics: singular potentials and rate of convergence. *Comm. Math. Phys.*, 298(1):101–138, 2010.
- [50] E. Kuz. Rate of convergence to mean field for interacting bosons. *Comm. Partial Differential Equations*, 40(10):1831–1854, 2015.
- [51] E. Kuz. Exact evolution versus mean field with second-order correction for bosons interacting via short-range two-body potential. *Differential Integral Equations*, 30(7-8):587–630, 2017.
- [52] M. Lewin. Mean-Field limit of Bose systems: rigorous results. In: *Proceedings of the International Congress of Mathematical Physics*, 2015.
- [53] M. Lewin, P. T. Nam, and N. Rougerie. Derivation of Hartree’s theory for generic mean-field Bose systems. *Adv. Math.*, 254:570–621, 2014.
- [54] M. Lewin, P. T. Nam, and N. Rougerie. The mean-field approximation and the non-linear Schrödinger functional for trapped Bose gases. *Trans. Amer. Math. Soc.*, 368(9):6131–6157, 2016.
- [55] M. Lewin, P. T. Nam, and N. Rougerie. A note on 2D focusing many-boson systems. *Proc. Amer. Math. Soc.*, 145(6):2441–2454, 2017.
- [56] E. H. Lieb and R. Seiringer. Proof of Bose-Einstein Condensation for Dilute Trapped Gases. *Phys. Rev. Lett.*, 88:170409, Apr 2002.
- [57] E. H. Lieb, R. Seiringer, J. P. Solovej, and J. Yngvason. *The mathematics of the Bose gas and its condensation*, volume 34 of *Oberwolfach Seminars*. Birkhäuser Verlag, Basel, 2005.
- [58] P. T. Nam and M. Napiórkowski. Norm approximation for many-body quantum dynamics: focusing case in low dimensions. *Adv. Math.*, 350:547–587, 2019.
- [59] P. T. Nam and N. Rougerie. Improved stability for 2D attractive Bose gases. *J. Math. Phys.*, 61(2):021901, 8, 2020.
- [60] P. Pickl. Derivation of the time dependent Gross-Pitaevskii equation without positivity condition on the interaction. *J. Stat. Phys.*, 140(1):76–89, 2010.
- [61] I. Rodnianski and B. Schlein. Quantum fluctuations and rate of convergence towards mean field dynamics. *Comm. Math. Phys.*, 291(1):31–61, 2009.
- [62] K. Schnee and J. Yngvason. Bosons in disc-shaped traps: from 3D to 2D. *Comm. Math. Phys.*, 269(3):659–691, 2007.
- [63] B. Simon. *Trace ideals and their applications*, volume 120 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, second edition, 2005.
- [64] V. Sohinger. A rigorous derivation of the defocusing cubic nonlinear Schrödinger equation on  $\mathbb{T}^3$  from the dynamics of many-body quantum systems. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 32(6):1337–1365, 2015.
- [65] H. Spohn. Kinetic equations from Hamiltonian dynamics: Markovian limits. *Rev. Modern Phys.*, 52(3):569–615, 1980.
- [66] K. E. Strecker, G. B. Partridge, A. G. Truscott, and R. G. Hulet. Formation and propagation of matter-wave soliton trains. *Nature*, 417(6885):150–153, 2002.
- [67] Z. Xie. Derivation of a nonlinear Schrödinger equation with a general power-type nonlinearity in  $d = 1, 2$ . *Differential Integral Equations*, 28(5-6):455–504, 2015.

SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING, 100871, CHINA & DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER, ROCHESTER, NY 14627, USA

E-mail address: sshen20@ur.rochester.edu