

THE STEFAN PROBLEM FOR THE FISHER-KPP EQUATION WITH UNBOUNDED INITIAL RANGE

WEIWEI DING[†], YIHONG DU[‡] AND ZONGMING GUO[‡]

ABSTRACT. We consider the nonlinear Stefan problem

$$\begin{cases} u_t - d\Delta u = au - bu^2 & \text{for } x \in \Omega(t), t > 0, \\ u = 0 \text{ and } u_t = \mu|\nabla_x u|^2 & \text{for } x \in \partial\Omega(t), t > 0, \\ u(0, x) = u_0(x) & \text{for } x \in \Omega_0, \end{cases}$$

where $\Omega(0) = \Omega_0$ is an unbounded smooth domain in \mathbb{R}^N , $u_0 > 0$ in Ω_0 and u_0 vanishes on $\partial\Omega_0$. When Ω_0 is bounded, the long-time behavior of this problem has been rather well-understood by [7, 8, 12, 13]. Here we reveal some interesting different behavior for certain unbounded Ω_0 . We also give a unified approach for a weak solution theory to this kind of free boundary problems with bounded or unbounded Ω_0 .

1. INTRODUCTION

In this paper, we investigate the following nonlinear Stefan problem

$$\begin{cases} u_t - d\Delta u = au - bu^2 & \text{for } x \in \Omega(t), t > 0, \\ u = 0 \text{ and } u_t = \mu|\nabla_x u|^2 & \text{for } x \in \Gamma(t), t > 0, \\ u(0, x) = u_0(x) & \text{for } x \in \Omega_0, \end{cases} \quad (1.1)$$

where $\Omega(t) \subset \mathbb{R}^N$ ($N \geq 2$) is a varying domain with boundary $\Gamma(t)$, a, b, μ and d are given positive constants, u_0 is positive in Ω_0 and vanishes on $\partial\Omega_0$. We are interested in the case that $\Omega(0) = \Omega_0$ is an unbounded smooth domain, which induces extra difficulties from the case of bounded Ω_0 , and gives rise to some interesting new phenomena (to be explained below).

Problem (1.1) is an analogue of the classical one-phase Stefan problem but with a logistic type nonlinear source term on the right side of the diffusive equation. Such a diffusive equation is often called a Fisher-KPP equation due to the pioneering works of Fisher [17] and Kolmogorov, Petrovski and Piskunov [22], and has been widely used in the study of propagation questions. In [7, 8, 12, 13], (1.1) with a bounded Ω_0 was used to describe the spreading of a new or invasive species with population density $u(t, x)$ and population range $\Omega(t)$. The evolution of the spreading front is determined by the free boundary $\Gamma(t)$, governed by the equation $u_t = \mu|\nabla_x u|^2$ on $\Gamma(t)$, which means the velocity of the movement of a point $x \in \Gamma(t)$ is given by $\mu|\nabla_x u|\nu_x$, where ν_x denotes the unit outward normal vector of $\Omega(t)$ at x (a deduction of this condition from ecological

Date: March 14, 2024.

1991 Mathematics Subject Classification. MSC: primary 35K20, 35R35 secondary 35J60.

Key words and phrases. Free boundary, Stefan problem, Fisher-KPP equation, weak solutions, unbounded initial range.

Y. Du would like to thank Prof Changfeng Gui for inspiring discussions on this problem. This work was supported by the Australian Research Council, the National Natural Science Foundation of China (11171092, 11571093) and the Innovation Scientists and Technicians Troop Construction Projects of Henan Province (114200510011).

[†] School of Mathematical Sciences, South China Normal University, Guangzhou 510631, China.

[‡] School of Science and Technology, University of New England, Armidale, NSW 2351, Australia.

[‡] Department of Mathematics, Henan Normal University, Xinxiang, 453007, P.R. China.

considerations can be found in [2]). In one space dimension, such a problem was first considered in [9].

Problem (1.1) is closely related to the following Cauchy problem:

$$\begin{cases} U_t - d\Delta U = aU - bU^2 & \text{for } x \in \mathbb{R}^N, t > 0, \\ U(0, x) = u_0(x) & \text{for } x \in \mathbb{R}^N, \end{cases} \quad (1.2)$$

where $u_0(x)$ is given in (1.1) but extended to \mathbb{R}^N with value 0 outside Ω_0 . Indeed, when Ω_0 is bounded, it has been shown in [8] that as $\mu \rightarrow \infty$, for any fixed $t > 0$, $\Omega(t) \rightarrow \mathbb{R}^N$ and $u(t, x) \rightarrow U(t, x)$, where u and U are the unique solutions of (1.1) and (1.2), respectively. (This holds true for unbounded Ω_0 as well; see Theorem 2.11 below.) Problem (1.2) has been studied extensively as a model for propagation (see, for example, [1, 17, 22]).

The basic feature of (1.1) with a bounded Ω_0 is given by the following theorem of [13]:

Theorem A.

- (i) $\Omega(t)$ is expanding: $\bar{\Omega}_0 \subset \Omega(t) \subset \Omega(s)$ if $0 < t < s$;
- (ii) $\Gamma(t) \setminus (\text{convex hull of } \bar{\Omega}_0)$ is smooth (e.g., C^2);
- (iii) $\Omega_\infty := \cup_{t>0} \Omega(t)$ is either the entire space \mathbb{R}^N , or it is a bounded set;
- (iv) When Ω_∞ is bounded, $\lim_{t \rightarrow \infty} \|u(t, \cdot)\|_{L^\infty(\Omega(t))} = 0$;
- (v) When $\Omega_\infty = \mathbb{R}^N$, for all large t , $\Gamma(t)$ is a smooth closed hypersurface in \mathbb{R}^N , and there exists a continuous function $M(t)$ such that

$$\Gamma(t) \subset \left\{ x : M(t) - \frac{\pi}{2}d_0 \leq |x| \leq M(t) \right\},$$

where d_0 is the diameter of Ω_0 .

Moreover, it follows from results of [8] that there exists $c_* = c_*(\mu) \in (0, 2\sqrt{ad})$ such that, in case (v) of Theorem A,

$$\lim_{t \rightarrow \infty} M(t)/t = c_*,$$

and for any $\epsilon > 0$,

$$\lim_{t \rightarrow \infty} \sup_{|x| \leq (c_* - \epsilon)t} \left| u(t, x) - \frac{a}{b} \right| = 0.$$

The number c_* is usually called the spreading speed, which is the unique positive value of c such that the following problem has a unique solution q :

$$-dq'' + cq' = aq - bq^2, \quad q > 0 \text{ in } (0, \infty), \quad q(0) = 0, \quad q'(0) = c/\mu. \quad (1.3)$$

The main purpose of this paper is to reveal some rather different and interesting behavior of (1.1) for the case that Ω_0 is unbounded. It turns out that such a case is much more complicated than the Ω_0 bounded case. To keep the paper at a reasonable length, we will only examine some very simple situations of unbounded Ω_0 for the long-time behavior of (1.1).

As in [8], a solution to (1.1) will be understood in a certain weak sense (to be made precise in section 2 below). We will show that (1.1) has a unique weak solution defined for all $t > 0$. As for Ω_0 , we will assume that there exist two parallel circular cones Λ_1 and Λ_2 , with the same axis, such that

$$\Lambda_1 \subset \Omega_0 \subset \Lambda_2.$$

Without loss of generality, we may assume that the common axis of Λ_1 and Λ_2 is the x_N axis in \mathbb{R}^N . If we denote

$$e_N := (0, \dots, 0, 1) \in \mathbb{R}^N,$$

then there exist $\phi \in (0, \pi)$ and $\xi_1, \xi_2 \in \mathbb{R}^1$ so that

$$\Lambda_i = \left\{ x \in \mathbb{R}^N : \frac{x - \xi_i e_N}{|x - \xi_i e_N|} \cdot e_N > \cos \phi \right\}, \quad i = 1, 2.$$

In other words, Λ_i has vertex at $\xi_i e_N$ and opening angle 2ϕ . Figure 1 below illustrates a case that $\phi \in (\pi/2, \pi)$ with $\theta = \pi - \phi \in (0, \pi/2)$.




FIGURE 1. $\Lambda^\phi + \xi_1 e_N \subset \Omega_0 \subset \Lambda^\phi + \xi_2 e_N$ with $\phi = \pi - \theta \in (\pi/2, \pi)$

For clarity of notations, for $\phi \in (0, \pi)$ and $\xi \in \mathbb{R}^1$, we define

$$\Lambda^\phi := \left\{ x \in \mathbb{R}^N \setminus \{0\} : \frac{x}{|x|} \cdot e_N > \cos \phi \right\}$$

and

$$\Lambda^\phi + \xi e_N := \left\{ x \in \mathbb{R}^N : x - \xi e_N \in \Lambda^\phi \right\}.$$

So Λ^ϕ and $\Lambda^\phi + \xi e_N$ are parallel cones with vertices at the origin and ξe_N , respectively.

We are now able to describe the main results of this paper.

Theorem 1.1. *Suppose that there exist $\phi \in (\pi/2, \pi)$ and $\xi_1 > \xi_2$ such that*

$$\Lambda^\phi + \xi_1 e_N \subset \Omega_0 \subset \Lambda^\phi + \xi_2 e_N \quad (1.4)$$

and

$$u_0 \in C^1(\overline{\Omega}_0) \cap L^\infty(\Omega_0), \quad \liminf_{d(x, \partial\Omega_0) \rightarrow \infty} u_0(x) > 0.$$

Then for any given small $\epsilon > 0$, there exists $T = T(\epsilon) > 0$ such that, for all $t > T$,

$$\Lambda^\phi - \left(\frac{c_*}{\sin \phi} - \epsilon \right) t e_N \subset \Omega(t) \subset \Lambda^\phi - \left(\frac{c_*}{\sin \phi} + \epsilon \right) t e_N, \quad (1.5)$$

where c_* is the spreading speed determined by (1.3).

The above result indicates that as $t \rightarrow \infty$, the free boundary $\Gamma(t) = \partial\Omega(t)$ propagates to infinity in the direction $-e_N$ at roughly the speed $c_*/\sin \phi$, and the shape of the free boundary can be roughly approximated by the boundary of the cone Λ^ϕ .

For any $R > 0$, if we denote the R -neighbourhood of Λ^ϕ by $N[\Lambda^\phi, R]$, namely

$$N[\Lambda^\phi, R] := \{x \in \mathbb{R}^N : d(x, \Lambda^\phi) < R\},$$

then, it is easily seen that when $\phi \in [\pi/2, \pi)$, we have

$$N[\Lambda^\phi, R] = \Lambda^\phi - \frac{R}{\sin \phi} e_N,$$

i.e., $N[\Lambda^\phi, R]$ is a shift of Λ^ϕ in the direction of $-e_N$ by $R/\sin \phi$. Thus (1.5) is equivalent to

$$N[\Lambda^\phi, (c_* - \tilde{\epsilon})t] \subset \Omega(t) \subset N[\Lambda^\phi, (c_* + \tilde{\epsilon})t] \text{ for } t > T, \text{ with } \tilde{\epsilon} := \epsilon \sin \phi. \quad (1.6)$$

In sharp contrast, if $\phi \in (0, \pi/2)$, then $N[\Lambda^\phi, R]$ has smooth boundary for all $R > 0$, and for large $R > 0$, $N[\Lambda^\phi, R]$ becomes very different from Λ^ϕ geometrically. Indeed, for such ϕ , it is easily verified that $\partial N[\Lambda^\phi, R]$ can be decomposed into two parts: a spherical part S_R on the sphere $\partial B_R(0)$, and the remaining part C_R on the boundary of the cone $\Lambda^\phi - \frac{R}{\sin \phi} e_N$, as follows

$$\partial N[\Lambda^\phi, R] = S_R \cup C_R, \quad S_R := \partial B_R(0) \setminus \Lambda^{\phi+\frac{\pi}{2}}, \quad C_R := \left(\partial \Lambda^\phi - \frac{R}{\sin \phi} e_N \right) \cap \Lambda^{\phi+\frac{\pi}{2}}.$$

The following theorem shows that (1.6) also holds when $\phi \in (0, \pi/2)$, though the geometric implications of (1.6) are very different from that in Theorem 1.1 now.

Theorem 1.2. *In Theorem 1.1 above, if we replace $\phi \in (\pi/2, \pi)$ by $\phi \in (0, \pi/2)$, then for any $\epsilon > 0$, there exists $\tilde{T} = \tilde{T}(\epsilon) > 0$ such that*

$$N[\Lambda^\phi, (c_* - \epsilon)t] \subset \Omega(t) \subset N[\Lambda^\phi, (c_* + \epsilon)t] \text{ for all } t \geq \tilde{T}. \quad (1.7)$$

Let us note that (1.7) implies that for all large t , $\partial \Omega(t)$ is roughly approximated in shape by $\partial N[\Lambda^\phi, R]$, which is very different from $\partial \Lambda^\phi$; in particular the propagation of $\partial \Omega(t)$ in the directions ν satisfying $\nu \cdot e_N \leq \cos(\phi + \frac{\pi}{2})$ is roughly at speed c_* .

Remark 1.3. (i) *If (1.4) holds with $\phi = \pi/2$, then much better results than those in Theorems 1.1 and 1.2 can be obtained; it can be shown that the free boundary $\partial \Omega(t)$ converges to a moving hyperplane, and the solution converges to the corresponding planar semi-wave. These require very different techniques and will be considered elsewhere.*
(ii) *In the case of Theorem 1.1, it is natural to conjecture that the solution u converges to a traveling wave with a V-shaped front (free boundary). This and related questions will be investigated in a future work ([6]).*

Let us mention here that the existence of “V-shaped” non-planar traveling wave solutions for the Cauchy problem (1.2) are well known by now (see e.g., [18, 19, 20, 24, 25]), which lend support to the validity of our conjecture above, but the existence of V-shaped semi-wave with free boundary is technically much more demanding. Note also that the nonlinearities in [19, 20, 24, 25] are of bistable type, very different from the Fisher-KPP type here.

The rest of the paper is organized as follows. In Section 2, we prove the existence and uniqueness of a weak solution to (1.1) by following the approach of [8], where the case of bounded Ω_0 was treated. For unbounded Ω_0 , some extra difficulties arise. In view of possible future applications, here we give a unified approach for a much more general problem than (1.1), with Ω_0 either bounded or unbounded; see (2.1).

In Section 3, we study the long-time dynamical behavior of problem (1.1). We consider a special type of unbounded Ω_0 , namely (1.4) holds for some $\phi \in (0, \pi)$. The main results are Theorems 3.1, 3.3 and 3.11. In particular, the conclusions stated in Theorem 1.1 above follow from Theorem 3.3, and the statements in Theorem 1.2 are consequences of Theorem 3.11.

The Appendix (Section 4) is concerned with the existence and uniqueness of classical solutions for an auxiliary radially symmetric free boundary problem with initial range the exterior of a ball. These conclusions are used in the proof of Theorem 3.3.

2. WEAK SOLUTIONS

In this section, we extend the weak solution theory in [8] to cover the case that the initial population range Ω_0 is unbounded. To our knowledge, little is known for problem (1.1) with a general unbounded Ω_0 . For future applications, we consider the following more general problem

$$\begin{cases} u_t - d\Delta u = g(x, u) & \text{for } x \in \Omega(t), t > 0, \\ u = 0, \quad u_t = \mu|\nabla_x u|^2 & \text{for } x \in \Gamma(t), t > 0, \\ u(0, x) = u_0(x) & \text{for } x \in \Omega_0, \end{cases} \quad (2.1)$$

where Ω_0 is a domain in \mathbb{R}^N with Lipschitz boundary $\partial\Omega_0$, the initial function $u_0(x)$ satisfies

$$u_0 \in C(\overline{\Omega_0}) \cap H_{loc}^1(\Omega_0) \cap L^\infty(\Omega_0), \quad u_0 > 0 \text{ in } \Omega_0, \quad u_0 = 0 \text{ on } \partial\Omega_0, \quad (2.2)$$

and the reaction term $g(x, u)$ is assumed to satisfy

$$\left. \begin{aligned} \text{(i)} \quad & g(x, u) \text{ is continuous for } (x, u) \in \mathbb{R}^N \times [0, \infty), \\ \text{(ii)} \quad & g(x, u) \text{ is locally Lipschitz in } u \text{ uniformly for } x \in \mathbb{R}^N, \\ \text{(iii)} \quad & g(x, 0) \equiv 0, \\ \text{(iv)} \quad & \text{there exists } K > 0 \text{ such that } g(x, u) \leq Ku \text{ for all } x \in \mathbb{R}^N \text{ and } u \geq 0. \end{aligned} \right\} \quad (2.3)$$

To describe the weak formulation of (2.1), it is convenient to start by considering $0 < t \leq T$ for some arbitrarily given $T \in (0, \infty)$. As in [8], the idea is to consider the extended u in the bigger region $H_T := [0, T] \times \mathbb{R}^N$ by defining $u(t, x) = 0$ for $x \in \mathbb{R}^N \setminus \Omega(t)$, $0 \leq t \leq T$, and regard it as a weak solution of an associated equation with certain jumping discontinuity. Throughout this section, we denote

$$\alpha(w) = \begin{cases} w & \text{if } w > 0, \\ w - d\mu^{-1} & \text{if } w \leq 0, \end{cases}$$

and

$$\tilde{u}_0(x) = \begin{cases} u_0(x) & \text{for } x \in \Omega_0, \\ 0 & \text{for } x \in \mathbb{R}^N \setminus \Omega_0. \end{cases}$$

Definition 2.1. Suppose that u_0 satisfies (2.2) and g satisfies (2.3). A nonnegative function $u \in H_{loc}^1(H_T) \cap L^\infty(H_T)$ is called a weak solution of (2.1) over H_T if

$$\int_0^T \int_{\mathbb{R}^N} \left[d\nabla_x u \cdot \nabla_x \phi - \alpha(u) \phi_t \right] dx dt - \int_{\mathbb{R}^N} \alpha(\tilde{u}_0) \phi(0, x) dx = \int_0^T \int_{\mathbb{R}^N} g(x, u) \phi dx dt \quad (2.4)$$

for every function $\phi \in C^\infty(H_T)$ such that ϕ has compact support¹ and $\phi = 0$ on $\{T\} \times \mathbb{R}^N$.

Correspondingly, if u satisfies (2.4) with “=” replaced by “ \geq ” (resp. “ \leq ”) for every test function $\phi \in C^\infty(H_T)$ satisfying $\phi \geq 0$ in H_T , $\phi = 0$ on $\{T\} \times \mathbb{R}^N$ and ϕ has compact support, then we call it a weak supersolution (resp. weak subsolution) of (2.1) over H_T .

Moreover, as in [8, 16], for each weak solution (or weak supersolution, or weak subsolution) $u(t, x)$, the function $\alpha(u(t, x))$ is defined as $u(t, x)$ if $u(t, x) > 0$; at points where $u(t, x) = 0$ the function $\alpha(u(t, x))$ is only required to satisfy $-d\mu^{-1} \leq \alpha(u(t, x)) \leq 0$ and to be such that it is altogether a measurable function over H_T . However, if $w(x)$ is continuous and positive in Ω_0 and identically zero in $\mathbb{R}^N \setminus \Omega_0$, then we understand that $\alpha(w) = -d\mu^{-1}$ on $\mathbb{R}^N \setminus \Omega_0$.

¹This means that there exists a ball B_R in \mathbb{R}^N such that $\phi(t, x) = 0$ for $t \in [0, T]$ and $x \in \mathbb{R}^N \setminus B_R$.

Remark 2.2. *Definition 2.1 is an extension of [8, Definition 2.1] for weak solution of problem (2.1) with bounded Ω_0 to the case that Ω_0 is generally unbounded. Indeed, the choice of the test function in (2.4) implies that if Ω_0 is bounded and if u is a weak solution to (2.1) over H_T by Definition 2.1, then the restriction of u to $G \times [0, T]$ is a weak solution of (2.1) over $G \times [0, T]$ by [8, Definition 2.1], where G is any given bounded domain in \mathbb{R}^N such that $\Omega(t)$ stays inside G for $0 \leq t \leq T$.*

By a *classical solution* of problem (2.1) for $0 < t \leq T < \infty$, we mean a pair $(u(t, x), \Omega(t))$ such that $\bigcup_{0 \leq t \leq T} \partial\Omega(t)$ is a C^1 hypersurface in \mathbb{R}^{N+1} , $u, \nabla_x u$ are continuous in $\bigcup_{0 \leq t \leq T} \overline{\Omega(t)}$, and $u_t, u_{x_i x_j}$ are continuous in $\bigcup_{0 < t \leq T} \Omega(t)$, and all the equations in (2.1) are satisfied in the classical sense. In this case, there exists $\Phi \in C^1\left(\overline{\bigcup_{0 < t \leq T} \Omega(t)}\right)$ such that

$$\Phi(t, x) = 0 \text{ on } \Gamma(t), \quad \nabla_x \Phi(x, t) \neq 0 \text{ on } \Gamma(t), \quad \Phi(t, x) < 0 \text{ in } \Omega(t),$$

and it follows from

$$u = 0 \text{ and } u_t = \mu |\nabla_x u|^2 \text{ on } \Gamma(t)$$

that

$$\Phi_t = \mu \nabla_x u \cdot \nabla_x \Phi \text{ for } x \in \Gamma(t).$$

Theorem 2.3. (i) *Assume that $(u(t, x), \Omega(t))$ is a classical solution of (2.1) for $0 < t \leq T$. Then*

$$w(t, x) := \begin{cases} u(t, x) & \text{for } x \in \Omega(t), 0 < t \leq T, \\ 0 & \text{for } x \in \mathbb{R}^N \setminus \Omega(t), 0 < t \leq T \end{cases}$$

is a weak solution of (2.1) over H_T .

(ii) *Let w be a weak solution of (2.1) on H_T . Assume that there exists a C^1 function Φ in H_T satisfying*

$$\Omega(t) := \{x \in \mathbb{R}^N : w(t, x) > 0\} = \{x \in \mathbb{R}^N : \Phi(t, x) < 0\}$$

with $\Omega(0) = \Omega_0$, and

$$\nabla_x \Phi \neq 0 \text{ on } \Gamma(t) := \partial\Omega(t).$$

Setting $u = w$ in $\bigcup_{0 < t \leq T} \overline{\Omega(t)}$, and assume that $u, \nabla_x u$ are continuous in $\bigcup_{0 \leq t \leq T} \overline{\Omega(t)}$ and that $\nabla_x^2 u, u_t$ are continuous in $\bigcup_{0 < t \leq T} \Omega(t)$. Then $(u(t, x), \Omega(t))$ is a classical solution of (2.1) for $t \in (0, T]$.

Proof. The proof follows that of [8, Theorem 2.3] with some obvious modifications, and we omit the details. \square

Next we prove the existence and uniqueness of weak solution to problem (2.1). The strategies of the proof are adapted from the approximation method of [21] as used in [8, 16], but new techniques are required to handle difficulties arising from the unboundedness of Ω_0 . We begin with the uniqueness part, since the existence proof will use some of the arguments in the uniqueness proof.

Theorem 2.4. *Suppose that $\mu_1 \geq \mu_2 > 0$, u_1 is a weak supersolution of (2.1) over H_T with $\mu = \mu_1$ and u_2 is a weak subsolution with $\mu = \mu_2$, where the initial data u_0 and Ω_0 are shared. Then $u_1 \geq u_2$ a.e. in H_T . In particular, problem (2.1) can have at most one weak solution over H_T .*

Proof. We complete the proof in three steps.

Step 1: *Setup of the approximation method.*

With u_1 and u_2 as given in the statement of the theorem, we define

$$\ell(t, x) = \begin{cases} \frac{g(x, u_2(t, x)) - g(x, u_1(t, x))}{u_2(t, x) - u_1(t, x)} & \text{if } u_2(t, x) \neq u_1(t, x), \\ 0 & \text{if } u_2(t, x) = u_1(t, x), \end{cases}$$

and for $i = 1, 2$, let $\alpha_i(u)$ denote $\alpha(u)$ with $\mu = \mu_i$, and

$$e(t, x) = \begin{cases} \frac{u_2(t, x) - u_1(t, x)}{\alpha_2(u_2(t, x)) - \alpha_1(u_1(t, x))} & \text{if } u_2(t, x) \neq u_1(t, x), \\ 0 & \text{if } u_2(t, x) = u_1(t, x). \end{cases}$$

We then have

$$\int_0^T \int_{\mathbb{R}^N} [\alpha_2(u_2) - \alpha_1(u_1)] (\phi_t + de\Delta\phi + e\ell\phi) dxdt \geq d(\mu_2^{-1} - \mu_1^{-1}) \int_{\mathbb{R}^N \setminus \overline{\Omega_0}} \phi(0, x) dx \quad (2.5)$$

for every nonnegative $\phi \in C^\infty(H_T)$ such that ϕ has compact support, and $\phi = 0$ on $\{T\} \times \mathbb{R}^N$.

It is easily checked that if we write

$$\alpha_2(u_2(t, x)) - \alpha_1(u_1(t, x)) = \bar{\alpha}(t, x) [u_2(t, x) - u_1(t, x)]$$

when $u_2(t, x) \neq u_1(t, x)$, then $\bar{\alpha}(t, x) \geq 1$ for almost all $(t, x) \in H_T$. And hence,

$$0 \leq e(t, x) \leq 1 \text{ for almost all } (t, x) \in H_T. \quad (2.6)$$

Moreover, by the condition (2.3) on g and the fact that $u_1, u_2 \in L^\infty(H_T)$, it follows that $\ell \in L^\infty(H_T)$. We then approximate e and ℓ by smooth functions $e_m \in C^\infty(H_T)$ and $\ell_m \in C^\infty(H_T)$, respectively, such that for any $R > 0$,

$$\|e - e_m\|_{L^2([0, T] \times B_R)} \rightarrow 0, \quad \|\ell - \ell_m\|_{L^2([0, T] \times B_R)} \rightarrow 0 \text{ as } m \rightarrow \infty, \quad (2.7)$$

and that

$$\inf_{H_T} e_m \geq \frac{1}{m}, \quad \|e_m\|_{L^\infty(H_T)} \leq C_1, \quad \|\ell_m\|_{L^\infty(H_T)} \leq C_1, \quad (2.8)$$

for some positive constant C_1 independent of m . Here and in what follows, we use B_R to denote the ball with center the origin and radius R .

Now, let $R_0 > 0$ be fixed, and choose a nonnegative function $f \in C^\infty(H_T)$ with $f(t, x) = 0$ for all $|x| \geq R_0$ and $0 \leq t \leq T$. For any $m \geq 1$ and $R > R_0 + 1$, we consider the following backward parabolic equation

$$\begin{cases} \frac{\partial \phi}{\partial t} + de_m \Delta \phi + e_m \ell_m \phi = f & \text{in } [0, T) \times B_R, \\ \phi = 0 & \text{on } \{T\} \times B_R, \\ \phi = 0 & \text{on } [0, T] \times \partial B_R. \end{cases} \quad (2.9)$$

This is a nondegenerate problem and it has a unique smooth solution ϕ_m (see e.g., [23]). Furthermore, the parabolic maximum principle (applied to $\phi_m(T - t, x)$) implies that

$$\phi_m \leq 0 \text{ in } [0, T] \times B_R.$$

Next, for $0 < \epsilon \ll 1$, we take a cutoff function $\xi_\epsilon \in C_0^\infty(\mathbb{R}^N)$ such that

$$0 \leq \xi_\epsilon \leq 1 \text{ in } \mathbb{R}^N, \quad \xi_\epsilon = 1 \text{ in } B_{R-2\epsilon}, \quad \xi_\epsilon = 0 \text{ in } \mathbb{R}^N \setminus B_{R-\epsilon}, \quad (2.10)$$

and that

$$\|\nabla \xi_\epsilon\|_{L^\infty(\mathbb{R}^N)} \leq C_2 \epsilon^{-1}, \quad \|\Delta \xi_\epsilon\|_{L^\infty(\mathbb{R}^N)} \leq C_2 \epsilon^{-2} \quad (2.11)$$

for some positive constant C_2 independent of ϵ . Then for any $m > 1$ and $0 < \epsilon \ll 1$, define

$$\psi_m^\epsilon(t, x) = \begin{cases} \phi_m(t, x) \xi_\epsilon(x) & \text{in } [0, T] \times B_R, \\ 0 & \text{in } [0, T] \times (\mathbb{R}^N \setminus B_R). \end{cases}$$

Clearly, $\psi_m^\epsilon \leq 0$, it belongs to $C^\infty(H_T)$, and vanishes in $(\{T\} \times \mathbb{R}^N) \cup ([0, T] \times (\mathbb{R}^N \setminus B_R))$.

We may now take $-\psi_m^\epsilon$ as a test function in (2.5) to obtain, due to $\mu_1 \geq \mu_2 > 0$,

$$\int_0^T \int_{\mathbb{R}^N} [\alpha_2(u_2) - \alpha_1(u_1)] \left((\psi_m^\epsilon)_t + de\Delta\psi_m^\epsilon + e\ell\psi_m^\epsilon \right) dxdt \leq 0.$$

Since $\xi_\epsilon f \equiv f$ in H_T , it then follows that

$$\begin{aligned} & \int_0^T \int_{\mathbb{R}^N} [\alpha_2(u_2) - \alpha_1(u_1)] f dxdt \\ &= \int_0^T \int_{B_R} [\alpha_2(u_2) - \alpha_1(u_1)] \left(\frac{\partial \phi_m}{\partial t} + de_m \Delta \phi_m + e_m \ell_m \phi_m \right) \xi_\epsilon dxdt \\ &\leq I_m + J_m + K_m, \end{aligned} \tag{2.12}$$

where

$$\begin{aligned} I_m &= I_m(\epsilon) := \int_0^T \int_{B_R} [\alpha_2(u_2) - \alpha_1(u_1)] d\xi_\epsilon (e_m - e) \Delta \phi_m dxdt, \\ J_m &= J_m(\epsilon) := \int_0^T \int_{B_R} [\alpha_2(u_2) - \alpha_1(u_1)] \xi_\epsilon (e_m \ell_m - e\ell) \phi_m dxdt, \end{aligned}$$

and

$$K_m = K_m(\epsilon) := \int_0^T \int_{B_R} [\alpha_2(u_2) - \alpha_1(u_1)] \left(-de\phi_m \Delta \xi_\epsilon - 2de \nabla \phi_m \cdot \nabla \psi_\epsilon \right) dxdt.$$

Our aim is to show, through suitable estimates on I_m , J_m and K_m , that

$$\int_0^T \int_{\mathbb{R}^N} [\alpha_2(u_2) - \alpha_1(u_1)] f dxdt \leq 0,$$

and the conclusions of the theorem would then follow easily from this inequality.

Step 2: *Estimates of I_m , J_m and K_m .*

We first consider I_m . Let

$$C_3 := \|u_1\|_{L^\infty(H_T)} + \|u_2\|_{L^\infty(H_T)} + d\mu_1^{-1} + d\mu_2^{-1}.$$

By the Hölder inequality, we have

$$\begin{aligned} I_m &\leq dC_3 \int_0^T \int_{B_R} |e_m - e| |\Delta \phi_m| dxdt \\ &\leq dC_3 \left(\int_0^T \int_{B_R} \frac{|e_m - e|^2}{|e_m|} dxdt \right)^{\frac{1}{2}} \left(\int_0^T \int_{B_R} |e_m| |\Delta \phi_m|^2 dxdt \right)^{\frac{1}{2}} \\ &\leq dC_3 \|e_m - e\|_{L^2([0,T] \times B_R)}^{\frac{1}{2}} \left(\int_0^T \int_{B_R} \frac{|e_m - e|^2}{|e_m|^2} dxdt \right)^{\frac{1}{4}} \left(\int_0^T \int_{B_R} |e_m| |\Delta \phi_m|^2 dxdt \right)^{\frac{1}{2}}. \end{aligned}$$

It follows from the proof of [8, Lemma 3.7] that there is a positive constant $C_4 = C_4(T, f, R)$ such that

$$\left(\int_0^T \int_{B_R} |e_m| |\Delta \phi_m|^2 dxdt \right)^{\frac{1}{2}} \leq C_4.$$

Moreover, by the same analysis as that used in the proof of [3, Lemma 5], we may require that the approximation sequence e_m satisfies additionally

$$\left\| \frac{e}{e_m} \right\|_{L^2([0,T] \times B_R)} \leq C_5$$

for some positive constant $C_5 = C_5(T, R)$. We thus obtain

$$I_m \leq C_6 \|e_m - e\|_{L^2([0,T] \times B_R)}^{\frac{1}{2}}, \tag{2.13}$$

with $C_6 = d(T|B_R| + C_5^2)^{1/4}C_3C_4$ independent of m .

Next, it is easily seen that

$$J_m \leq C_3 \|\phi_m\|_{L^\infty([0,T] \times B_R)} \left(\|e_m\|_{L^2([0,T] \times B_R)} \|\ell_m - \ell\|_{L^2([0,T] \times B_R)} + \|\ell\|_{L^2([0,T] \times B_R)} \|e_m - e\|_{L^2([0,T] \times B_R)} \right).$$

By the comparison arguments used in [8, Lemma 3.6], we have

$$\|\phi_m\|_{L^\infty([0,T] \times B_R)} \leq C_7,$$

for some positive constant $C_7 = C_7(T, f)$. Hence, by setting $C_8 = (T|B_R|)^{1/2}C_1C_3C_7$, we obtain

$$J_m \leq C_8 \left(\|\ell_m - \ell\|_{L^2([0,T] \times B_R)} + \|e_m - e\|_{L^2([0,T] \times B_R)} \right). \quad (2.14)$$

It remains to estimate K_m . Making use of (2.6), (2.10) and (2.11), we have

$$\begin{aligned} K_m &\leq dC_3 \int_0^T \int_{B_R} \left(|\phi_m \Delta \xi_\epsilon| + 2|\nabla \phi_m \cdot \nabla \xi_\epsilon| \right) dx dt \\ &\leq dC_2C_3 \left(\int_0^T \int_{B_{R-\epsilon} \setminus B_{R-2\epsilon}} \left(\frac{|\phi_m|}{\epsilon^2} + \frac{2|\nabla \phi_m|}{\epsilon} \right) dx dt \right). \end{aligned} \quad (2.15)$$

Since $\phi_m(t, x) = 0$ on $[0, T] \times \partial B_R$, it follows by letting $\epsilon \rightarrow 0$ in (2.15) that

$$\limsup_{\epsilon \rightarrow 0} K_m(\epsilon) \leq C_9 R^{N-1} \sup_{0 \leq t \leq T, x \in \partial B_R} \left| \frac{\partial \phi_m(t, x)}{\partial \nu_x} \right|, \quad (2.16)$$

where $C_9 = C_9(T, C_2, C_3)$ is a positive constant independent of m and R , and ν_x is the outward unit normal vector of B_R at $x \in \partial B_R$.

We next estimate $\partial \phi_m(t, x)/\partial \nu_x$ at the boundary ∂B_R by making use of barrier functions inspired by the proof of [4, Theorem 2.1]. Let

$$v(t, x) := -C_{10}e^{-\beta_1 t}(1 + |x|^2)^{-\beta_2} \text{ for } (t, x) \in [0, T] \times B_R,$$

where C_{10} , β_1 and β_2 are positive constants independent of R , to be chosen later. It is straightforward to verify that

$$v_t = -\beta_1 v \quad \text{and} \quad \Delta v \geq 4(\beta_2^2 + \beta_2)v \quad \text{for } (t, x) \in [0, T] \times B_R,$$

and thus,

$$v_t + de_m \Delta v + e_m \ell_m v \geq -v \left(\beta_1 - 4(\beta_2^2 + \beta_2)de_m - e_m \ell_m \right) \text{ for } (t, x) \in [0, T] \times B_R.$$

Due to (2.8) and the fact that $f(t, x) = 0$ for all $|x| \geq R_0$ and $0 \leq t \leq T$, we may choose

$$\beta_1 = 4(\beta_2^2 + \beta_2)dC_1 + C_1^2 + 1,$$

and

$$C_{10} = e^{\beta_1 T} (1 + R_0^2)^{\beta_2} \max_{(t, x) \in [0, T] \times B_{R_0}} |f(t, x)|,$$

and thus conclude that

$$\begin{cases} v_t + de_m \Delta v + e_m \ell_m v \geq f & \text{in } [0, T] \times B_R, \\ v < 0 & \text{on } \{T\} \times B_R, \\ v < 0 & \text{on } [0, T] \times \partial B_R. \end{cases}$$

It then follows from the parabolic maximum principle applied to problem (2.9) that

$$v(t, x) \leq \phi_m(t, x) \leq 0 \text{ for } (t, x) \in [0, T] \times B_R. \quad (2.17)$$

We next consider the function

$$v^*(t, x) := \frac{-C_{10}e^{-\beta_1 t}\sigma(|x|)}{\sigma(R-1)(1+(R-1)^2)^{\beta_2}} \quad \text{for } (t, x) \in [0, T] \times B_R \setminus B_{R-1},$$

where

$$\sigma(\rho) = \begin{cases} \rho^{2-N} - R^{2-N} & \text{if } N \geq 3, \\ \log(R/\rho) & \text{if } N = 2. \end{cases}$$

The function $v^*(t, x)$ satisfies $\Delta v^* = 0$ and

$$v_t^* + e_m \ell_m v^* = -v^*(\beta_1 - e_m \ell_m) \geq 0 \quad \text{in } [0, T] \times B_R \setminus B_{R-1}.$$

Since $f(t, x) = 0$ for all $0 \leq t \leq T$ and $x \in B_R \setminus B_{R-1}$ (note that $R > R_0 + 1$), it then follows that

$$v_t^* + de_m \Delta v^* + e_m \ell_m v^* \geq f \quad \text{in } [0, T] \times B_R \setminus B_{R-1}.$$

On the other hand, we have, in view of (2.17),

$$\begin{cases} v^* < 0 & \text{on } \{T\} \times B_R \setminus B_{R-1}, \\ v^* = 0 & \text{on } [0, T] \times \partial B_R, \\ v^* = -C_{10}e^{-\beta_1 t}(1+(R-1)^2)^{-\beta_2} = v \leq \phi_m & \text{on } [0, T] \times \partial B_{R-1}. \end{cases}$$

It then follows from the parabolic maximum principle again that

$$v^*(t, x) \leq \phi_m(t, x) \leq 0 \quad \text{for } (t, x) \in [0, T] \times B_R \setminus B_{R-1}.$$

Since both ϕ_m and v^* vanish on ∂B_R , we thus obtain

$$\frac{\partial v^*(t, x)}{\partial \nu_x} \geq \frac{\partial \phi_m(t, x)}{\partial \nu_x} \geq 0 \quad \text{for } (t, x) \in [0, T] \times \partial B_R.$$

This together with the estimate (2.16) implies that

$$\begin{aligned} \limsup_{\epsilon \rightarrow 0} K_m(\epsilon) &\leq C_9 R^{N-1} \sup_{0 \leq t \leq T, x \in \partial B_R} \left| \frac{\partial v^*(t, x)}{\partial \nu_x} \right| \\ &= \frac{C_9 C_{10} R^{N-1} |\sigma'(R)|}{\sigma(R-1)(1+(R-1)^2)^{\beta_2}}. \end{aligned}$$

Thus, if we take $\beta_2 = N/2$, then there exists some positive constant $C_{11} = C_{11}(N, C_9, C_{10})$ independent of R and m such that

$$\limsup_{\epsilon \rightarrow 0} K_m(\epsilon) \leq C_{11} R^{-1}. \quad (2.18)$$

Step 3: Completion of the proof.

Combining the estimates in (2.13), (2.14) and (2.18), we obtain from (2.12) that

$$\begin{aligned} &\int_0^T \int_{\mathbb{R}^N} [\alpha_2(u_2) - \alpha_1(u_1)] f dx dt \\ &\leq C_6 \|e_m - e\|_{L^2([0, T] \times B_R)}^{\frac{1}{2}} + C_8 \left(\|\ell_m - \ell\|_{L^2([0, T] \times B_R)} + \|e_m - e\|_{L^2([0, T] \times B_R)} \right) + C_{11} R^{-1}. \end{aligned}$$

In view of (2.7) and the facts that C_6, C_8 depend on R but not on m and that C_{11} is independent of m and R , passing to the limit as $m \rightarrow \infty$ followed by letting $R \rightarrow \infty$ in the above inequality, we obtain

$$\int_0^T \int_{\mathbb{R}^N} [\alpha_2(u_2) - \alpha_1(u_1)] f dx dt \leq 0.$$

Since it holds for all nonnegative smooth function $f(t, x)$ with compact support in x , it follows that $\alpha_2(u_2) \leq \alpha_1(u_1)$, and hence in the a.e. sense $u_1(t, x) > 0$ whenever $u_2(t, x) > 0$, and $u_1 \geq u_2$ a.e. in H_T . The proof of Theorem 2.4 is now complete. \square

We now consider the existence of weak solutions of problem (2.1) over H_T .

Theorem 2.5. *There exists a unique weak solution w of (2.1) over H_T .*

The proof of this theorem also follows the approximation arguments used in [8, 16] and we only provide the details where considerable changes are required. Before giving the proof, we first introduce some notations and approximation functions. Let $\{\alpha_m(w)\}_{m \in \mathbb{N}}$ be a sequence of smooth functions such that

$$\alpha_m(w) \rightarrow \alpha(w) \text{ uniformly in any compact subset of } \mathbb{R}^1 \setminus \{0\},$$

and

$$\alpha_m(0) \rightarrow -d\mu^{-1}, \quad w - d\mu^{-1} \leq \alpha_m(w) \leq w \text{ for all } w \in \mathbb{R}^1.$$

We may choose $\alpha_m(w)$ in such a way that

$$\alpha'_m(w) \geq 1. \quad (2.19)$$

We now consider the following sequence of approximating problems:

$$\begin{cases} \frac{\partial \alpha_m(w)}{\partial t} - d\Delta w = g(x, w) & \text{in } (0, T] \times \mathbb{R}^N, \\ w(0, x) = \tilde{u}_0(x) & \text{in } \mathbb{R}^N. \end{cases} \quad (2.20)$$

For any fixed $m \in \mathbb{N}$, we call a function w a *bounded supersolution* to problem (2.20) if $w \in C^{1,2}((0, T] \times \mathbb{R}^N) \cap C(H_T) \cap L^\infty(H_T)$ and it satisfies

$$\begin{cases} \frac{\partial \alpha_m(w)}{\partial t} - d\Delta w \geq g(x, w) & \text{in } (0, T] \times \mathbb{R}^N, \\ w(0, x) \geq \tilde{u}_0(x) & \text{in } \mathbb{R}^N. \end{cases} \quad (2.21)$$

Such a w is called a *bounded subsolution* to problem (2.20) if the reversed inequalities in (2.21) are satisfied. We have the following comparison result.

Lemma 2.6. *Let $m \in \mathbb{N}$ be fixed. Assume that \bar{w} and \underline{w} are bounded supersolution and subsolution of problem (2.20), respectively. Then $\bar{w} \geq \underline{w}$ in H_T .*

Proof. It is easily verified from (2.20) that \bar{w} satisfies

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^N} [d\nabla_x \bar{w} \cdot \nabla_x \phi - \alpha_m(\bar{w})\phi_t] dx dt &= \int_{\mathbb{R}^N} \alpha_m(\bar{w}(0, x))\phi(0, x) dx \\ &\geq \int_0^T \int_{\mathbb{R}^N} g(x, \bar{w})\phi dx dt \end{aligned} \quad (2.22)$$

for every nonnegative function $\phi \in C^\infty(H_T)$ such that ϕ has compact support and $\phi = 0$ on $\{T\} \times \mathbb{R}^N$. Analogously, \underline{w} satisfies (2.22) with the inequality sign reversed. Subtracting these two inequalities, due to (2.19) and the fact that $\bar{w}(0, x) \geq \underline{w}(0, x)$ in \mathbb{R}^N , we obtain

$$\int_0^T \int_{\mathbb{R}^N} [\alpha_m(\underline{w}) - \alpha_m(\bar{w})] (\phi_t + d\tilde{\ell}\Delta\phi + \tilde{\ell}\tilde{\ell}\phi) dx dt \geq 0,$$

where

$$\tilde{\ell}(t, x) = \begin{cases} \frac{g(x, \underline{w}(t, x)) - g(x, \bar{w}(t, x))}{\underline{w}(t, x) - \bar{w}(t, x)} & \text{if } \underline{w}(t, x) \neq \bar{w}(t, x), \\ 0 & \text{if } \underline{w}(t, x) = \bar{w}(t, x), \end{cases}$$

and

$$\tilde{e}(t, x) = \begin{cases} \frac{\underline{w}(t, x) - \bar{w}(t, x)}{\alpha_m(\underline{w}(t, x)) - \alpha_m(\bar{w}(t, x))} & \text{if } \underline{w}(t, x) \neq \bar{w}(t, x), \\ 0 & \text{if } \underline{w}(t, x) = \bar{w}(t, x). \end{cases}$$

Since $\bar{w}, \underline{w} \in L^\infty(H_T)$ and since $\alpha'_m(w) \geq 1$ by (2.19), it then follows from the same arguments as those used in the proof of Theorem 2.4 that

$$\alpha_m(\underline{w}(t, x)) - \alpha_m(\bar{w}(t, x)) \leq 0 \text{ for almost all } (t, x) \in H_T.$$

This immediately gives

$$\underline{w}(t, x) \leq \bar{w}(t, x) \text{ for all } (t, x) \in H_T,$$

as $\alpha_m(w)$, $\underline{w}(t, x)$ and $\bar{w}(t, x)$ are all continuous functions, and $\alpha_m(w)$ is increasing in w . The proof of Lemma 2.6 is thus complete. \square

Clearly, $\underline{w} \equiv 0$ is a subsolution of (2.20). On the other hand, by the condition (2.3) on g and the property (2.19) of α_m , we can conclude that the function $\bar{w}(t)$ is a bounded supersolution of (2.20), where

$$\bar{w}(t) = \|\tilde{u}_0\|_{L^\infty(\mathbb{R}^N)} e^{Kt} \text{ for } 0 \leq t \leq T \quad (2.23)$$

is the unique solution to the problem

$$\frac{d\bar{w}}{dt} = K\bar{w} \text{ for } 0 < t < T, \quad \bar{w}(0) = \|\tilde{u}_0\|_{L^\infty(\mathbb{R}^N)}.$$

We are ready to show the existence and uniqueness of classical solution to (2.20).

Lemma 2.7. *Let $m \in \mathbb{N}$ be fixed and $\bar{w}(t)$ be given as in (2.23). Then problem (2.20) admits a unique solution $w_m \in C^{1,2}((0, T] \times \mathbb{R}^N) \cap C(H_T) \cap L^\infty(H_T)$. Moreover,*

$$0 \leq w_m(t, x) \leq \bar{w}(t) \text{ for } (t, x) \in H_T, \quad (2.24)$$

and for any $R > 0$, there exists a positive constant $\tilde{C}_1 = \tilde{C}_1(T, R)$ independent of m such that

$$\|w_m\|_{H^1([0, T] \times B_R)} \leq \tilde{C}_1 \text{ for all } m \in \mathbb{N}. \quad (2.25)$$

Proof. The existence part must be a known result, but we failed to find a proof in the literature. So we include a proof here. Note that the uniqueness part follows directly from Lemma 2.6.

For each $k \in \mathbb{N}$, let $B_k \in \mathbb{R}^N$ be the ball with center at 0 and radius k . Consider the following initial boundary value problem

$$\begin{cases} \frac{\partial \alpha_m(v)}{\partial t} - d\Delta v = g(x, v) & \text{in } (0, T] \times B_k, \\ v(t, x) = 0 & \text{on } (0, T] \times \partial B_k, \\ v(0, x) = \tilde{u}_0(x) & \text{in } B_k. \end{cases} \quad (2.26)$$

It is well known that, for each $k \in \mathbb{N}$, problem (2.26) admits a unique solution $v_k \in C([0, T] \times B_k) \cap C^{1,2}((0, T] \times \bar{B}_k)$ (see e.g., [23]). Moreover, it follows from the comparison result given in [8, Lemma 3.2] that

$$0 \leq v_k(t, x) \leq \bar{w}(t) \text{ for all } (t, x) \in [0, T] \times B_k.$$

It thus follows that

$$0 \leq v_k(t, x) \leq \tilde{C}_2 := \|\tilde{u}_0\|_{L^\infty(\mathbb{R}^N)} e^{KT} \text{ for all } (t, x) \in [0, T] \times B_k.$$

The comparison argument also gives $v_k(t, x) \leq v_{k+1}(t, x)$ for $t \in [0, T]$ and $x \in B_k$. Thus

$$v(t, x) := \lim_{k \rightarrow \infty} v_k(t, x) \text{ exists for every } x \in \mathbb{R}^N \text{ and } t \in [0, T],$$

and $0 \leq v(t, x) \leq \bar{w}(t)$. By a standard regularity consideration, one sees that v solves (2.20), and (2.24) holds. To mark its dependence on m , we will denote $v = w_m$.

Next, we show the estimate (2.25). Clearly, we have

$$0 \leq w_m(t, x) \leq \tilde{C}_2 \text{ for } (t, x) \in H_T. \quad (2.27)$$

For any fixed $R > 0$, let $\xi(x)$ be a smooth function such that

$$0 \leq \xi \leq 1 \text{ in } \mathbb{R}^N, \quad \xi \equiv 1 \text{ in } B_R, \text{ and } \xi \equiv 0 \text{ in } \mathbb{R}^N \setminus B_{2R}.$$

If we multiply the differential equation in (2.20) by $\alpha_m(w_m)\xi^2$, then by following the proof of [8, Lemma 5.1], we obtain

$$\int_0^T \int_{B_R} |\nabla_x w_m|^2 \leq \tilde{C}_3,$$

for some positive constant $\tilde{C}_3 = \tilde{C}_3(R, T)$ independent of m . Furthermore, if we multiply the differential equation in (2.20) by $\frac{\partial w_m}{\partial t} \xi^2$ and estimate the resulting equation over $[0, T] \times B_{2R}$, then the proof of [8, Lemma 5.2] implies that there exists a positive constant $\tilde{C}_4 = \tilde{C}_4(R, T)$ such that

$$\int_0^T \int_{B_R} \left| \frac{\partial w_m}{\partial t} \right|^2 \leq \tilde{C}_4.$$

Therefore, combining the above, we obtain (2.25). The proof of Lemma 2.7 is now complete. \square

To complete the proof of Theorem 2.5, we shall need the following estimate.

Lemma 2.8. *For any $R > 0$, there is a positive constant \tilde{C}_5 depending on R and T but independent of m such that*

$$\int_{B_{R/2}} \left| \nabla_x w_m(\sigma, x) \right|^2 dx \leq \tilde{C}_5, \text{ for all } m \in \mathbb{N}, \sigma \in [0, T].$$

Proof. Let $\xi(x)$ be a smooth function such that

$$0 \leq \xi \leq 1 \text{ in } \mathbb{R}^N, \quad \xi \equiv 1 \text{ in } B_{R/2}, \text{ and } \xi \equiv 0 \text{ in } \mathbb{R}^N \setminus B_R.$$

We multiply both sides of the differential equation in (2.20) by $\frac{\partial w_m}{\partial t} \xi^2$ and then integrate the resulting equation over $[0, \sigma] \times B_R$. After suitable integration by parts, we obtain

$$\begin{aligned} \int_0^\sigma \int_{B_R} \alpha'_m(w_m) \left(\frac{\partial w_m}{\partial t} \right)^2 \xi^2 dx dt &+ d \int_0^\sigma \int_{B_R} \nabla_x w_m \cdot \nabla_x \left(\frac{\partial w_m}{\partial t} \xi^2 \right) dx dt \\ &= \int_0^\sigma \int_{B_R} g(x, w_m) \frac{\partial w_m}{\partial t} \xi^2 dx dt. \end{aligned} \quad (2.28)$$

Moreover, using the assumption (2.3) on g and the estimate (2.27), we have

$$\int_0^\sigma \int_{B_R} g(x, w_m) \frac{\partial w_m}{\partial t} \xi^2 dx dt \leq \frac{1}{4} \int_0^\sigma \int_{B_R} \left(\frac{\partial w_m}{\partial t} \right)^2 \xi^2 dx dt + \tilde{C}_6(T, R).$$

On the other hand, making use of the estimate (2.25), we deduce that

$$\begin{aligned}
& d \int_0^\sigma \int_{B_R} \nabla_x w_m \cdot \nabla_x \left(\frac{\partial w_m}{\partial t} \xi^2 \right) dx dt \\
&= d \int_0^\sigma \int_{B_R} \xi^2 \nabla_x w_m \cdot \nabla_x \left(\frac{\partial w_m}{\partial t} \right) dx dt + 2d \int_0^\sigma \int_{B_R} \xi \frac{\partial w_m}{\partial t} \nabla_x w_m \cdot \nabla_x \xi dx dt \\
&\geq \frac{d}{2} \int_{B_R} |\nabla_x w_m(\sigma, x)|^2 \xi^2(x) dx - \frac{d}{2} \int_{B_R} |\nabla_x \tilde{u}_0(x)|^2 \xi^2(x) dx \\
&\quad - 16d^2 \int_0^\sigma \int_{B_R} (\nabla_x w_m \cdot \nabla_x \xi)^2 dx dt - \frac{1}{4} \int_0^\sigma \int_{B_R} \left(\frac{\partial w_m}{\partial t} \right)^2 \xi^2 dx dt \\
&\geq \frac{d}{2} \int_{B_R} |\nabla_x w_m(\sigma, x)|^2 \xi^2(x) dx - \frac{d}{2} \int_{B_R} |\nabla_x \tilde{u}_0(x)|^2 \xi^2(x) dx \\
&\quad - \frac{1}{4} \int_0^\sigma \int_{B_R} \left(\frac{\partial w_m}{\partial t} \right)^2 \xi^2 dx dt - \tilde{C}_7(T, R).
\end{aligned}$$

Substituting the above estimates into (2.28), and recalling that $\alpha'_m(w) \geq 1$ from (2.19), we obtain

$$\begin{aligned}
& \frac{1}{2} \int_0^\sigma \int_{B_R} \left(\frac{\partial w_m}{\partial t} \right)^2 \xi^2 dx dt + \frac{d}{2} \int_{B_R} |\nabla_x w_m(\sigma, x)|^2 \xi^2(x) dx \\
&\leq \frac{d}{2} \int_{B_R} |\nabla_x \tilde{u}_0(x)|^2 \xi^2(x) dx + \tilde{C}_6(T, R) + \tilde{C}_7(T, R).
\end{aligned}$$

Let

$$\tilde{C}_5 := d \int_{B_R} |\nabla_x \tilde{u}_0(x)|^2 dx + 2\tilde{C}_6(T, R) + 2\tilde{C}_7(T, R).$$

It finally follows that

$$\int_{B_{R/2}} |\nabla_x w_m(\sigma, x)|^2 dx \leq \int_{B_R} |\nabla_x w_m(\sigma, x)|^2 \xi^2(x) dx \leq \tilde{C}_5 \quad \text{for all } m \in \mathbb{N}.$$

This completes the proof. \square

We are now ready to give the proof of Theorem 2.5.

Proof of Theorem 2.5. In what follows we will select various subsequences from $\{w_m\}$ and, to avoid inundation by subscripts, we will always denote the subsequence again by $\{w_m\}$. By the estimate (2.25), and by Rellich's Lemma and a standard diagonal argument, there is a subsequence of $\{w_m\}$ (denoted by itself) and a function $w \in H_{loc}^1(H_T)$ such that, for any $R > 0$,

$$w_m \rightarrow w \text{ weakly in } H^1([0, T] \times B_R) \text{ and strongly in } L^2([0, T] \times B_R) \text{ as } m \rightarrow \infty.$$

In particular, $w_m \rightarrow w$ as $m \rightarrow \infty$ and $w \geq 0$ almost everywhere in H_T . Moreover, in view of (2.27), we have $0 \leq w \leq \tilde{C}_2$ in H_T , and hence $w \in H_{loc}^1(H_T) \cap L^\infty(H_T)$. Furthermore, Lemma 2.8 implies that

$$\int_{B_{R/2}} |\nabla_x w(t, x)|^2 dx \leq \tilde{C}_5 \quad \text{for a.e. } t \in [0, T],$$

since the set

$$\left\{ v \in H^1([0, T] \times B_{R/2}) : \int_{B_{R/2}} |\nabla_x v(t, x)|^2 dx \leq \tilde{C}_5 \text{ for a.e. } t \in [0, T] \right\}$$

is closed and convex in $H^1([0, T] \times B_{R/2})$, and such sets are closed under the weak limit.

With the above preparations, and noting that the test function ϕ in Definition 2.1 has compact support, we can follow the same lines as those used in the proof of [8, Theorem 3.1] to verify that

w is a weak solution of (2.1) over H_T . We do not repeat the details here. The proof of Theorem 2.5 is thereby complete. \square

Note that since $T > 0$ is arbitrary in Theorem 2.5, the weak solution $u(t, x)$ of (2.1) given over H_T can be extended to all $t > 0$, and it is unique due to Theorem 2.4.

Next, we present a comparison principle which will be used frequently in the subsequent sections. Suppose that μ and $\hat{\mu}$ are two positive constants, g and \hat{g} both satisfy the assumption (2.3), Ω_0 and $\hat{\Omega}_0$ are smooth domains in \mathbb{R}^N , u_0 satisfies (2.2) and \hat{u}_0 satisfies (2.2) with Ω_0 replaced by $\hat{\Omega}_0$. Let u be a weak subsolution of (2.1) corresponding to (Ω_0, u_0, g, μ) , and \hat{u} be a weak supersolution corresponding to $(\hat{\Omega}_0, \hat{u}_0, \hat{g}, \hat{\mu})$, respectively.

Theorem 2.9. *Suppose that $\Omega_0 \subset \hat{\Omega}_0$, $u_0 \leq \hat{u}_0$, $\mu \leq \hat{\mu}$ and $g \leq \hat{g}$. Then $u \leq \hat{u}$ a.e. in H_T .*

Proof. Let w and \hat{w} be the weak solutions of (2.1) corresponding to (Ω_0, u_0, g, μ) and $(\hat{\Omega}_0, \hat{u}_0, \hat{g}, \hat{\mu})$ over H_T , respectively. As a direct application of Theorem 2.4, we have

$$u \leq w, \quad \hat{u} \geq \hat{w} \quad \text{a.e. in } H_T. \quad (2.29)$$

On the other hand, let w_m be the solution of problem (2.20), and \hat{w}_m be the solution determined by (2.20) with reaction term \hat{g} , and the initial function is obtained by extending \hat{u}_0 to \mathbb{R}^N . By the comparison result Lemma 2.6, we clearly have $w_m \leq \hat{w}_m$ in H_T . Furthermore, from the proof of Theorem 2.5 and the uniqueness of the weak solution, we know that

$$w_m \rightarrow w, \quad \hat{w}_m \rightarrow \hat{w} \quad \text{a.e. in } H_T \text{ as } m \rightarrow \infty,$$

and hence $w \leq \hat{w}$ a.e. in H_T . This together with (2.29) completes the proof. \square

The next result is useful for constructing supersolutions and subsolutions to (2.1).

Theorem 2.10. *Assume that there exists a C^1 function Φ over H_T such that, with*

$$\Omega(t) := \left\{ x \in \mathbb{R}^N : \Phi(t, x) < 0 \right\},$$

one has

$$\Omega(0) = \Omega_0 \text{ and } |\nabla_x \Phi| \neq 0 \text{ on } \partial\Omega(t).$$

Suppose that $w(t, x)$ and $\nabla_x w(t, x)$ are continuous in $\bigcup_{0 \leq t \leq T} \overline{\Omega(t)}$, $\nabla_x^2 w(t, x)$, $w_t(t, x)$ are continuous in $\bigcup_{0 < t \leq T} \Omega(t)$. Then $w(t, x)$ (extended by 0 for $x \notin \Omega(t)$) is a weak supersolution of (2.1) for $t \in (0, T]$, provided that

$$\begin{cases} w_t - d\Delta w \geq g(x, w), \quad w > 0 & \text{for } x \in \Omega(t), \quad 0 < t \leq T, \\ w = 0, \quad \Phi_t \leq \mu \nabla_x w \cdot \nabla_x \Phi & \text{for } x \in \partial\Omega(t), \quad 0 < t \leq T, \\ w(0, x) \geq u_0(x) & \text{for } x \in \Omega_0. \end{cases} \quad (2.30)$$

If all the inequalities in (2.30) are reversed, then w is a weak subsolution to (2.1).

Proof. We only prove the case for weak supersolution, as the proof for the weak subsolution case is similar. We need to show that

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^N} \left[d \nabla_x w \cdot \nabla_x \phi - \alpha(w) \phi_t \right] dx dt - \int_{\mathbb{R}^N} \alpha(\tilde{u}_0(x)) \phi(0, x) dx \\ \geq \int_0^T \int_{\mathbb{R}^N} g(x, w) \phi dx dt \end{aligned} \quad (2.31)$$

for every nonnegative function $\phi \in C^\infty(H_T)$ with compact support and satisfying $\phi = 0$ on $\{T\} \times \mathbb{R}^N$.

For each test function ϕ as described above, we use the divergence theorem to calculate the following integral

$$\int_0^T \int_{\Omega(t)} \operatorname{div} \Psi \, dxdt \quad \text{with } \Psi(t, x) = (\phi(t, x), 0, \dots, 0) \in \mathbb{R}^{N+1},$$

and obtain, with $S := \bigcup_{0 < t < T} \partial\Omega(t)$,

$$\begin{aligned} \int_0^T \int_{\Omega(t)} \phi_t \, dxdt &= - \int_S \phi \frac{\Phi_t}{|(\Phi_t, \nabla_x \Phi)|} d\sigma - \int_{\Omega(0)} \phi(0, x) \, dxdt \\ &= - \int_0^T \int_{\partial\Omega(t)} \phi \frac{\Phi_t}{|\nabla_x \Phi|} dS_x dt - \int_{\Omega(0)} \phi(0, x) dxdt. \end{aligned}$$

Since the unit outward normal of $\Omega(t)$ at $x \in \partial\Omega(t)$ is given by $\nu = \nu(t, x) := \nabla_x \Phi(t, x) / |\nabla_x \Phi(t, x)|$, by (2.30) we have

$$\frac{\Phi_t}{|\nabla_x \Phi|} \leq \mu \nabla_x w \cdot \nu \quad \text{for } x \in \partial\Omega(t),$$

and hence

$$- \frac{d}{\mu} \int_0^T \int_{\Omega(t)} \phi_t \, dxdt \leq d \int_0^T \int_{\partial\Omega(t)} \phi \nabla_x w \cdot \nu dS_x dt + \frac{d}{\mu} \int_{\Omega(0)} \phi(0, x) dxdt. \quad (2.32)$$

On the other hand, we multiply both sides of the first inequality in (2.30) by ϕ and integrate the resulting inequality over $\bigcup_{0 < t < T} \Omega(t)$, making use of integration by parts and the last inequality in (2.30), and obtain

$$\begin{aligned} \int_0^T \int_{\Omega(t)} [\phi_t w - d \nabla_x \phi \nabla_x w] \, dxdt &+ \int_0^T \int_{\Omega(t)} g(x, w) \phi \, dxdt \\ &\leq -d \int_0^T \int_{\partial\Omega(t)} \phi \nabla_x w \cdot \nu dS_x dt - \int_{\Omega(0)} u_0(x) \phi(0, x) dxdt. \end{aligned}$$

By this inequality, (2.32), the fact that $w(t, x) = 0$ for $x \notin \Omega(t)$, and the definition of α , we immediately obtain (2.31). \square

We now consider the asymptotic behavior of the weak solution to (2.1) as $\mu \rightarrow \infty$. To emphasize its dependence on μ , we denote by u_μ the unique weak solution, and denote $\Omega_\mu(t) = \{x : u_\mu(t, x) > 0\}$.

Theorem 2.11. *Let u_μ be the unique weak solution to problem (2.1). Then*

$$\lim_{\mu \rightarrow \infty} \Omega_\mu(t) = \mathbb{R}^N, \quad \forall t > 0,$$

and

$$u_\mu \rightarrow U \quad \text{in } C_{loc}^{\frac{1+\nu_0}{2}, 1+\nu_0}((0, \infty) \times \mathbb{R}^N) \quad \text{as } \mu \rightarrow \infty,$$

where ν_0 can be any number in $(0, 1)$ and $U(t, x)$ is the unique solution of the Cauchy problem

$$\begin{cases} U_t - d \Delta U = g(x, U) & \text{in } (0, \infty) \times \mathbb{R}^N, \\ U(0, x) = \tilde{u}_0(x) & \text{in } \mathbb{R}^N. \end{cases} \quad (2.33)$$

Moreover, $u_\mu(t, x) \leq U(t, x)$ for all $t > 0$ and $x \in \Omega_\mu(t)$.

Proof. Making use of Theorem 2.9, the proof is almost identical to that of [8, Theorem 5.4], and we omit the details. \square

Remark 2.12. *Although our main interest of this paper is for the case $\Omega_0 \subset \mathbb{R}^N$ with $N \geq 2$, it is easily seen from their proofs that all the results in this section remain valid when $N = 1$.*

3. SPREADING PROFILE OF THE FISHER-KPP EQUATION

In this section, we study the long-time behavior of the weak solution to problem (1.1), for some special unbounded Ω_0 . More precisely, we assume that there exist $\phi \in (0, \pi)$ and $\xi_1 > \xi_2$ such that (1.4) holds, namely

$$\Lambda^\phi + \xi_1 e_N \subset \Omega_0 \subset \Lambda^\phi + \xi_2 e_N. \quad (3.1)$$

Our first result on the long-time behavior of (1.1) is the following theorem.

Theorem 3.1. *Let $u(t, x)$ be the unique weak solution of problem (1.1) with Ω_0 satisfying (3.1). Denote $\Omega(t) = \{x : u(t, x) > 0\}$. Then*

$$\lim_{t \rightarrow \infty} \Omega(t) = \mathbb{R}^N, \quad (3.2)$$

and

$$\lim_{t \rightarrow \infty} u(t, x) = \frac{a}{b} \text{ locally uniformly in } x \in \mathbb{R}^N. \quad (3.3)$$

Proof. We first prove (3.2). Since Ω_0 satisfies (3.1), we can find a ball $B_{r_0}(x_0) \subset \Omega_0$ with radius

$$r_0 \geq R^* := \sqrt{\frac{d}{a} \lambda_1}, \quad (3.4)$$

where λ_1 is the first eigenvalue of the following eigenvalue problem

$$-\Delta \phi = \lambda \phi \text{ in } B_1(0); \quad \phi = 0 \text{ on } \partial B_1(0).$$

We then choose a C^2 radial function $v_0(r)$ ($r = |x - x_0|$) satisfying

$$v_0(|x - x_0|) \leq u_0(x) \text{ for } |x - x_0| < r_0, \quad v_0(r) > 0 \text{ for } r \in [0, r_0) \text{ and } v_0'(0) = v_0(r_0) = 0,$$

and consider the following radially symmetric problem

$$\begin{cases} v_t - d\Delta v = av - bv^2, & t > 0, \quad 0 < r < k(t), \\ v_r(t, 0) = 0, \quad v(t, k(t)) = 0, & t > 0, \\ k'(t) = -\mu v_r(t, k(t)), & t > 0, \\ k(0) = r_0, \quad v(0, r) = v_0(r), & 0 \leq r \leq r_0. \end{cases}$$

It follows from [7, Theorem 2.1] that this problem admits a (unique) classical solution $(v(t, r), k(t))$ defined for all $t > 0$ such that $k'(t) > 0$, $v(t, r) > 0$ for $0 \leq r < k(t)$, $t > 0$. Moreover, by [7, Theorem 2.5], we have

$$\lim_{t \rightarrow \infty} k(t) = \infty. \quad (3.5)$$

Denote

$$\mathcal{G}(t) = \{x \in \mathbb{R}^N : |x - x_0| < k(t)\},$$

and

$$V(t, x) = v(t, |x - x_0|).$$

We also extend $V(t, \cdot)$ to be zero outside $\mathcal{G}(t)$. Clearly, for any given $T > 0$, (V, \mathcal{G}) is a classical solution of the free boundary problem

$$\begin{cases} u_t - d\Delta u = au - bu^2 & \text{for } x \in \mathcal{G}(t), \quad 0 < t \leq T, \\ u = 0, \quad u_t = \mu |\nabla_x u|^2 & \text{for } x \in \partial \mathcal{G}(t), \quad 0 < t \leq T, \\ u(0, x) = v_0(|x - x_0|) & \text{for } x \in \mathcal{G}(0). \end{cases} \quad (3.6)$$

By Theorems 2.3 and 2.4, V is the unique weak solution of (3.6) over H_T . It then follows from Theorem 2.9 that $u \geq V$ in H_T , and hence by the arbitrariness of T , we have $\mathcal{G}(t) \subset \Omega(t)$ for all $t \geq 0$. This together with (3.5) implies (3.2).

It remains to prove (3.3). We claim that

$$V(t, x) \leq u(t, x) \leq U(t, x) \text{ in } [0, \infty) \times \mathbb{R}^N, \quad (3.7)$$

where $U(t, x)$ is the unique solution to the Cauchy problem (2.33). Indeed, the first inequality in (3.7) follows directly from Theorem 2.9, while the second inequality is a consequence of Theorem 2.11.

Applying [8, Theorem 6.2] to the equation of V , we obtain

$$\lim_{t \rightarrow \infty} V(t, x) = \frac{a}{b} \text{ locally uniformly in } x \in \mathbb{R}^N. \quad (3.8)$$

On the other hand, it is well known that

$$\lim_{t \rightarrow \infty} U(t, x) = \frac{a}{b} \text{ locally uniformly in } x \in \mathbb{R}^N. \quad (3.9)$$

By (3.7), (3.8) and (3.9), we immediately obtain (3.3), and the proof of Theorem 3.1 is thus complete. \square

Next we examine the long-time profile of the free boundary $\partial\Omega(t)$. For convenience, we first recall the following result from [2].

Proposition 3.2. *Let $d > 0$, $a > 0$ and $b > 0$ be given constants. For any $k \in [0, 2\sqrt{ad})$, the problem*

$$-dZ'' + kZ' = aZ - bZ^2 \text{ in } (0, \infty), \quad Z(0) = 0 \quad (3.10)$$

admits a unique positive solution $Z = Z_k$, and it satisfies $Z_k(r) \rightarrow a/b$ as $r \rightarrow \infty$. Moreover, $Z'_k(r) > 0$ for $r \geq 0$, $Z'_{k_1}(0) > Z'_{k_2}(0)$, $Z_{k_1}(r) > Z_{k_2}(r)$ for $r > 0$ and $k_1 < k_2$, and for each $\mu > 0$, there is a unique $c_ = c_*(\mu, a, b, d)$ such that $\mu Z'_k(0) = k$ when $k = c_*$. Furthermore, $c_*(\mu, a, b, d)$ depends continuously on its arguments, is increasing in μ and $\lim_{\mu \rightarrow \infty} c_*(\mu, a, b, d) = 2\sqrt{ad}$.*

In our discussion below, since a, b, d and μ are always fixed, we use c_* to denote $c_*(\mu, a, b, d)$.

Theorem 3.3. *Let $u(t, x)$ be the unique weak solution of problem (1.1) with Ω_0 satisfying (3.1) for some $\phi \in (\pi/2, \pi)$, and u_0 satisfying (2.2) and*

$$\liminf_{d(x, \partial\Omega_0) \rightarrow \infty} u_0(x) > 0. \quad (3.11)$$

Denote $\Omega(t) = \{x : u(t, x) > 0\}$. Then for any $\epsilon > 0$, there exists $T = T(\epsilon) > 0$ such that for all $t > T$ we have

$$\Lambda^\phi - \left(\frac{c_*}{\sin \phi} - \epsilon \right) t e_N \subset \Omega(t) \subset \Lambda^\phi - \left(\frac{c_*}{\sin \phi} + \epsilon \right) t e_N. \quad (3.12)$$

Moreover,

$$\lim_{t \rightarrow \infty} \left[\sup_{x \in \Lambda^\phi - \left(\frac{c_*}{\sin \phi} - \epsilon \right) t e_N} \left| u(t, x) - \frac{a}{b} \right| \right] = 0. \quad (3.13)$$

We prove this theorem by a series of lemmas. We first show

$$\Lambda^\phi - \left(\frac{c_*}{\sin \phi} - \epsilon \right) t e_N \subset \Omega(t) \text{ for all large } t. \quad (3.14)$$

By assumption (3.11), we can find a one-dimensional function $w_0 \in C^2([0, \infty))$ such that

$$0 < w_0 \leq \sigma_0 := \frac{1}{2} \liminf_{d(x, \partial\Omega_0) \rightarrow \infty} u_0(x) \text{ in } (0, \infty), \quad w_0(0) = 0 \quad \text{and} \quad w'_0 > 0 \text{ in } [0, \infty). \quad (3.15)$$

Then we consider the following one space dimension free boundary problem

$$\begin{cases} w_t - dw_{yy} = aw - bw^2, & t > 0, \rho(t) < y < \infty, \\ w(t, \rho(t)) = 0, & t > 0, \\ \rho'(t) = -\mu w_y(t, \rho(t)), & t > 0, \\ \rho(0) = 0, \quad w(0, y) = w_0(y), & 0 \leq y < \infty. \end{cases} \quad (3.16)$$

It follows from [5, Theorem 2.11] that, (3.16) admits a (unique) classical solution $(w(t, y), \rho(t))$ defined for all $t > 0$ and $\rho'(t) < 0$, $w(t, y) > 0$ for $\rho(t) \leq y < \infty$, $t > 0$. Moreover, we have the following comparison result.

Lemma 3.4. *For any given $\tilde{T} \in (0, \infty)$, suppose $\tilde{\rho} \in C^1([0, \tilde{T}])$ and $\tilde{w} \in C^{1,2}(\tilde{D}_{\tilde{T}})$ with $\tilde{D}_{\tilde{T}} = \{(t, y) : 0 \leq t \leq \tilde{T}, \tilde{\rho}(t) \leq y < \infty\}$. If*

$$\begin{cases} \tilde{w}_t - d\tilde{w}_{yy} \geq \tilde{w}(a - b\tilde{w}), & 0 < t \leq \tilde{T}, \tilde{\rho}(t) < y < \infty, \\ \tilde{w}(t, \tilde{\rho}(t)) = 0, \quad \tilde{\rho}'(t) \leq -\mu\tilde{w}_y(t, \tilde{\rho}(t)), & 0 < t \leq \tilde{T}, \end{cases}$$

and

$$\tilde{\rho}(0) \leq 0 \quad \text{and} \quad w_0(y) \leq \tilde{w}(0, y) \quad \text{in } [0, \infty),$$

then the solution (w, ρ) of problem (3.16) satisfies

$$\tilde{\rho}(t) \leq \rho(t) \quad \text{in } (0, \tilde{T}] \quad \text{and} \quad w(t, y) \leq \tilde{w}(t, y) \quad \text{for } 0 < t \leq \tilde{T}, \rho(t) \leq y < \infty.$$

Proof. If for any $0 \leq t \leq \tilde{T}$, we extend $w(t, y)$ (resp. $\tilde{w}(t, y)$) to be zero for $y < \rho(t)$ (resp. $y < \tilde{\rho}(t)$), then it is easily checked that w (reps. \tilde{w}) is a weak solution (resp. weak supersolution) of the free boundary problem induced from (3.16) over $H_{\tilde{T}}$, and hence the desired comparison result follows from Theorem 2.9. (One could also prove the result directly along the lines of [9].) \square

We next show the following estimate of (w, ρ) .

Lemma 3.5. *For any $\epsilon > 0$, there exists $T_1 = T_1(\epsilon) > 0$ such that*

$$\rho(t) \leq -\left(c_* - \frac{2}{3}\epsilon\right)t \quad \text{for all } t \geq T_1, \quad (3.17)$$

and

$$\liminf_{t \rightarrow \infty} \left[\inf_{y \geq -(c_* - \frac{2}{3}\epsilon)t} w(t, y) \right] \geq \frac{a}{b}. \quad (3.18)$$

Proof. The proof follows from [9, Theorem 4.2] with some modifications. For the sake of completeness, and also for the convenience of later applications, we include the details below.

We first claim that for any given small $\delta > 0$, there exists $t_1 = t_1(\delta) > 0$ such that

$$w(t_1, y) \geq \frac{a - \delta}{b + \delta} \quad \text{for all } y \geq 0. \quad (3.19)$$

Indeed, applying the proof of Theorem 3.1 to the one-dimensional problem (3.16), we easily obtain $\lim_{t \rightarrow \infty} \rho(t) = -\infty$ and

$$\lim_{t \rightarrow \infty} w(t, y) = \frac{a}{b} \quad \text{locally uniformly for } y \in \mathbb{R}.$$

Since $w_0(y)$ is nondecreasing in $y \in [0, \infty)$, it follows from the comparison result stated in Lemma 3.4 and the uniqueness of solution to problem (3.16) that, for any fixed $t > 0$, $w(t, y)$ is nondecreasing in $y \in [\rho(t), \infty)$. We thus obtain

$$\liminf_{t \rightarrow \infty} w(t, y) \geq \frac{a}{b} \quad \text{uniformly for } y \geq 0,$$

which clearly implies (3.19).

Next, we construct a subsolution to problem (3.16). To do this, we need a few more notations. For any small $\delta > 0$, denote

$$c_\delta := c_*(\mu, a - \delta, b + \delta, d)$$

and denote by $Z_\delta(r)$ the solution of (3.10) with k, a, b replaced by $c_\delta, a - \delta, b + \delta$, respectively. Set

$$\eta(t) = \eta_\delta(t) := -(1 - \delta)^2 c_\delta t \text{ for } t > 0,$$

and

$$\underline{w}(t, y) = \underline{w}_\delta(t, y) := (1 - \delta)^2 Z_\delta(y - \eta_\delta(t)) \text{ for } \eta_\delta(t) \leq y < \infty, t > 0.$$

It is straightforward to verify that

$$\underline{w}(t, \eta(t)) = 0, \quad \eta'(t) = -\mu \underline{w}_y(t, \eta(t)) \text{ for } t > 0.$$

Moreover, since $Z'_\delta > 0$ in $[0, \infty)$, direct calculations yield

$$\begin{aligned} \underline{w}_t - d \underline{w}_{yy} &= (1 - \delta)^4 c_\delta Z'_\delta - d(1 - \delta)^2 Z''_\delta \\ &\leq (1 - \delta)^2 (Z'_\delta - d Z''_\delta) \\ &= (1 - \delta)^2 [(a - \delta) Z_\delta - (b + \delta) Z_\delta^2] \\ &\leq (a - \delta) \underline{w} - (b + \delta) \underline{w}^2 \end{aligned}$$

for $\eta(t) < y < \infty, t > 0$. By Proposition 3.2, we have

$$\underline{w}(0, y) = (1 - \delta)^2 Z_\delta(y) < (1 - \delta)^2 \frac{a - \delta}{b + \delta} \text{ for } 0 \leq y < \infty.$$

This together with (3.19) implies

$$w(t_1, y) \geq \underline{w}(0, y) \text{ for } 0 \leq y < \infty.$$

It then follows from Lemma 3.4 that

$$\rho(t + t_1) \leq \eta(t), \quad w(t + t_1, y) \geq \underline{w}(t, y) \text{ for } \eta(t) \leq y < \infty, t > 0. \quad (3.20)$$

Since $\lim_{\delta \rightarrow 0} (1 - \delta)^2 c_\delta = c_* := c_*(\mu, a, b, d)$, for any $\epsilon > 0$, we can find some $\delta_\epsilon \in (0, \epsilon)$ such that

$$|(1 - \delta_\epsilon)^2 c_{\delta_\epsilon} - c_*| \leq \epsilon/2. \quad (3.21)$$

We now fix $\delta = \delta_\epsilon$ in Z_δ, η_δ and $t_1(\delta)$. Then (3.20) implies

$$\rho(t) \leq -(1 - \delta_\epsilon)^2 c_{\delta_\epsilon} (t - t_1) \leq -\left(c_* - \frac{2}{3}\epsilon\right)t - \frac{\epsilon}{6}t + (1 - \delta_\epsilon)^2 c_{\delta_\epsilon} t_1 \text{ for } t \geq t_1.$$

Thus (3.17) holds with $T_1 = 6\epsilon^{-1}(1 - \delta_\epsilon)^2 c_{\delta_\epsilon} t_1$.

It remains to prove (3.18). With $\delta = \delta_\epsilon$ chosen as above, it follows from Proposition 3.2 that there exists $y_0 > 0$ sufficiently large such that

$$Z_{\delta_\epsilon}(y) \geq \frac{a - 2\delta_\epsilon}{b + 2\delta_\epsilon} \text{ for all } y \geq y_0. \quad (3.22)$$

On the other hand, by (3.21), we clearly have

$$y - \eta(t - t_1) \geq y + \left(c_* - \frac{2}{3}\epsilon\right)t + \frac{\epsilon}{6}t - (1 - \delta_\epsilon)^2 c_{\delta_\epsilon} t_1 \text{ for } t \geq t_1.$$

Thus, if we choose $\tilde{t}_1 = 6\epsilon^{-1}[y_0 + (1 - \delta_\epsilon)^2 c_{\delta_\epsilon} t_1]$, then

$$y - \eta(t - t_1) \geq y_0 \text{ for all } y \geq -\left(c_* - \frac{2}{3}\epsilon\right)t, t \geq \tilde{t}_1.$$

This together with (3.20) and (3.22) implies

$$w(t, y) \geq \underline{w}(t - t_1, y) \geq (1 - \delta_\epsilon)^2 \frac{a - 2\delta_\epsilon}{b + 2\delta_\epsilon} \text{ for all } y \geq -\left(c_* - \frac{2}{3}\epsilon\right)t, t \geq \tilde{t}_1.$$

Since

$$(1 - \delta_\epsilon)^2 \frac{a - 2\delta_\epsilon}{b + 2\delta_\epsilon} \rightarrow \frac{a}{b} \text{ as } \epsilon \rightarrow 0,$$

this gives (3.18), and the proof of Lemma 3.5 is now complete. \square

Remark 3.6. *The conclusions in Lemma 3.5 can be considerably sharpened (though they are not needed in this paper). It is possible to modify the method of [14] to show that, as $t \rightarrow \infty$,*

$$\rho(t) - c_* t \rightarrow C \in \mathbb{R}^1, \quad \sup_{y \in [\rho(t), \infty)} |w(t, y) - Z_{c_*}(y - \rho(t))| \rightarrow 0,$$

where Z_{c_*} is given in Proposition 3.2.

Lemma 3.7. *Let $u(t, x)$ and $\Omega(t)$ be given in the statement of Theorem 3.3. Then for any $\epsilon > 0$, there exists $T_2 = T_2(\epsilon) > 0$ such that (3.14) holds for all $t \geq T_2$, and*

$$\liminf_{t \rightarrow \infty} \left[\inf_{x \in \Lambda^\phi - \left(\frac{c_*}{\sin \phi} - \epsilon\right) t e_N} u(t, x) \right] \geq \frac{a}{b}. \quad (3.23)$$

Proof. By the assumption (3.11), there exists $r_1 > 0$ sufficiently large such that

$$u_0(x) \geq \sigma_0 := \frac{1}{2} \liminf_{d(\tilde{x}, \partial\Omega_0) \rightarrow \infty} u_0(\tilde{x}) \text{ for all } x \in \Omega_0 \text{ with } d(x, \partial\Omega_0) \geq r_1.$$

Then, due to the assumption (3.1), we have

$$u_0(x) \geq \sigma_0 \text{ for all } x \in \Lambda^\phi + \left(\xi_1 + \frac{r_1}{\sin \theta}\right) e_N, \quad (3.24)$$

where $\theta = \pi - \phi$ (and so $\sin \theta = \sin \phi$).

Let (w, ρ) be the unique solution to problem (3.16) with initial function w_0 satisfying (3.15). For convenience of notation, we write

$$z_1 := \left(\xi_1 + \frac{r_1}{\sin \theta}\right) e_N \quad \text{and} \quad \Lambda_{z_1} := \Lambda^\phi + z_1.$$

For any fixed $z \in \partial\Lambda_{z_1} \setminus \{z_1\}$, let ν_z be the inward unit normal vector of Λ_{z_1} at z , and define

$$\Omega_z(t) = \left\{ x : x \cdot \nu_z \geq \rho(t) + r_1 + \xi_1 \sin \theta \right\}$$

($\Omega_z(0)$ is illustrated in Figure 2), and

$$w_z(t, x) = w(t, x \cdot \nu_z - r_1 - \xi_1 \sin \theta).$$

We also extend $w_z(t, \cdot)$ to be zero outside $\Omega_z(t)$ for $t \geq 0$. Clearly, (w_z, Ω_z) is a classical solution of the following problem

$$\begin{cases} w_t - d\Delta w = w(a - bw) & \text{for } x \in \Omega(t), t > 0, \\ w = 0, w_t = \mu |\nabla_x w|^2 & \text{for } x \in \partial\Omega(t), t > 0, \\ w(0, x) = w_0(x \cdot \nu_z - r_1 - \xi_1 \sin \theta) & \text{for } x \in \Omega_z(0), \end{cases}$$

and hence by Theorems 2.3 and 2.4, it is the unique weak solution.

By Lemma 3.5, for any $\epsilon > 0$, there exists $T_1 = T_1(\epsilon) > 0$ such that


$$\Omega_z(t) \supset \left\{ x : x \cdot \nu_z \geq -\left(c_* - \frac{2}{3}\epsilon\right)t + r_1 + \xi_1 \sin \theta \right\} \text{ for all } t \geq T_1,$$

and

$$\liminf_{t \rightarrow \infty} \left[\inf_{x \cdot \nu_z - r_1 - \xi_1 \sin \theta \geq -(c_* - \frac{2}{3}\epsilon)t} w_z(t, x) \right] \geq \frac{a}{b}.$$

Therefore, if we choose

$$T_2 := \max \left\{ T_1, \frac{3|r_1 + \xi_1 \sin \theta|}{\epsilon} \right\},$$

FIGURE 2. The domain $\Omega_z(0)$ with given $z \in \partial\Lambda_{z_1} \setminus \{z_1\}$

then

$$\Omega_z(t) \supset \left\{ x : x \cdot \nu_z \geq -(c_* - \epsilon)t \right\} \text{ for all } t \geq T_2,$$

and

$$\liminf_{t \rightarrow \infty} \left[\inf_{x \cdot \nu_z \geq -(c_* - \epsilon)t} w_z(t, x) \right] \geq \frac{a}{b}.$$

On the other hand, by the choice of w_0 in (3.15) and the property (3.24), we have

$$\Omega_z(0) \subset \Lambda_{z_1} \subset \Omega(0),$$

and

$$w_0(x \cdot \nu_z - r_1 - \xi_1 \sin \theta) \leq u_0(x) \text{ for } x \in \Omega_z(0).$$

Hence we can use Theorem 2.9 to compare u and w_z and then obtain

$$w_z(t, x) \leq u(t, x) \text{ in } [0, \infty) \times \mathbb{R}^N,$$

which clearly implies $\Omega_z(t) \subset \Omega(t)$ for $t \geq 0$.

Thus, we obtain

$$\left\{ x : x \cdot \nu_z \geq -(c_* - \epsilon)t \right\} \subset \Omega(t) \text{ for all } t \geq T_2,$$

and

$$\liminf_{t \rightarrow \infty} \left[\inf_{x \cdot \nu_z \geq -(c_* - \epsilon)t} u(t, x) \right] \geq \frac{a}{b}.$$

Finally, by the arbitrariness of $z \in \partial\Lambda_{z_1} \setminus \{z_1\}$, we obtain

$$\Lambda^\phi - \frac{c_* - \epsilon}{\sin \phi} t e_N = \bigcup_{z \in \partial\Lambda_{z_1} \setminus \{z_1\}} \left\{ x : x \cdot \nu_z \geq -(c_* - \epsilon)t \right\} \subset \Omega(t) \text{ for all } t \geq T_2.$$

The desired results then follow if we replace ϵ by $\tilde{\epsilon} := \epsilon / \sin \phi$. □

Next we prove

$$\Omega(t) \subset \Lambda^\phi - \left(\frac{c_*}{\sin \phi} + \epsilon \right) t e_N \text{ for all large } t$$

by constructing a suitable weak supersolution to problem (1.1). We do this with several lemmas.

Lemma 3.8. *Let $u(t, x)$ and $\Omega(t)$ be given in the statement of Theorem 3.3. Then for any $\delta > 0$, there exist $t_2 = t_2(\delta) > 0$ and $r_2 = r_2(\delta) > 0$ such that*

$$u(t, x) \leq \frac{a + \delta}{b - \delta} \text{ for } t \geq t_2, x \in \mathbb{R}^N, \quad (3.25)$$

and

$$\Omega(t_2) \subset \Lambda_{z_2} := \Lambda^\phi + \left(\xi_2 - \frac{r_2}{\sin \phi} \right) e_N. \quad (3.26)$$

Proof. Let $u^*(t)$ be the unique solution of the problem

$$\frac{du^*}{dt} = u^*(a - bu^*) \text{ for } t > 0; \quad u^*(0) = \max \left\{ \frac{a}{b}, \|u_0\|_{L^\infty(\Omega_0)} \right\}.$$

Clearly, we have

$$u^*(t) \geq \frac{a}{b} \text{ for all } t \geq 0 \text{ and } \lim_{t \rightarrow \infty} u^*(t) = \frac{a}{b}.$$

Moreover, it follows from Theorem 2.11 and the parabolic comparison principle that

$$u(t, x) \leq U(t, x) \leq u^*(t) \text{ for } t \geq 0, x \in \mathbb{R}^N,$$

where $U(t, x)$ is the unique solution of the Cauchy problem (2.33). As a consequence, for any $\delta > 0$, there exists $t_2 = t_2(\delta) > 0$ such that

$$u(t, x) \leq u^*(t) \leq \frac{a + \delta}{b - \delta} \text{ for all } t \geq t_2, x \in \mathbb{R}^N,$$

which clearly gives (3.25).

It remains to prove (3.26). Let $t_2 > 0$ be determined as above. It then follows from Proposition 4.1 in the Appendix below that there exists $R_0 > 1$ depending on t_2 such that, for any given radially symmetric function $\hat{v}_0 \in C^2([0, \infty))$ satisfying

$$0 < \hat{v}_0(r) < 2\|u_0\|_{L^\infty(\Omega_0)} \text{ for } r \in (0, \infty), \quad \hat{v}_0(0) = 0, \quad \|\hat{v}_0\|_{C^1([0, \infty))} \leq 3\|u_0\|_{L^\infty(\Omega_0)}, \quad (3.27)$$

the following free boundary problem

$$\begin{cases} \hat{v}_t - d\Delta \hat{v} = a\hat{v} - b\hat{v}^2, & 0 < t < t_2, \hat{h}(t) < r < \infty, \\ \hat{v}(t, \hat{h}(t)) = 0, & 0 < t < t_2, \\ \hat{h}'(t) = -\mu \hat{v}_r(t, \hat{h}(t)), & 0 < t < t_2, \\ \hat{h}(0) = R_0, \quad \hat{v}(0, r) = \hat{v}_0(r - R_0), & R_0 \leq r < \infty, \end{cases} \quad (3.28)$$

admits a unique classical solution (\hat{v}, \hat{h}) defined for $0 < t \leq t_2$ with $\hat{v}(t, r) > 0$, $\hat{h}'(t) < 0$ for $0 < t \leq t_2$, $\hat{h}(t) < r < \infty$, and

$$\hat{h}(t_2) \geq R_0/2, \quad (3.29)$$

where due to the radial symmetry, $\Delta \hat{v} = \hat{v}_{rr} + \frac{N-1}{r} \hat{v}_r$.

Set

$$r_2 := R_0 + 1 \quad \text{and} \quad \Lambda_{z_2} := \Lambda^\phi + \left(\xi_2 - \frac{r_2}{\sin \phi} \right) e_N.$$

Clearly

$$B_{r_2}(\hat{x}_0) \cap (\Lambda^\phi + \xi_2 e_N) = \emptyset \text{ for all } \hat{x}_0 \in \mathbb{R}^N \setminus \Lambda_{z_2}.$$

Since $\Omega_0 \subset (\Lambda^\phi + \xi_2 e_N)$, it is easily seen that $|x - \hat{x}_0| - R_0 > 1$ for all $x \in \Omega_0$, $\hat{x}_0 \in \mathbb{R}^N \setminus \Lambda_{z_2}$. In view of this, we may require that, in addition to the constraint (3.27), \hat{v}_0 also satisfies

$$\hat{v}_0(|x - \hat{x}_0| - R_0) \geq u_0(x) \text{ for } x \in \Omega_0, \hat{x}_0 \in \mathbb{R}^N \setminus \Lambda_{z_2}. \quad (3.30)$$

Indeed, this can be ensured by requiring \hat{v}_0 to satisfy $\hat{v}_0(r) \geq \|u_0\|_{L^\infty(\Omega_0)}$ for $r \geq 1$.

We now define, for each $\hat{x}_0 \in \mathbb{R}^N \setminus \Lambda_{z_2}$,

$$\hat{V}(t, x) = \hat{v}(t, |x - \hat{x}_0|) \text{ for } |x - \hat{x}_0| \geq \hat{h}(t),$$

and extend it to zero for $|x - \hat{x}_0| < \hat{h}(t)$ ($0 \leq t \leq t_2$), then it is easily seen that \hat{V} is the unique weak solution of the free boundary problem induced from (3.28) over $[0, t_2] \times \mathbb{R}^N$ with initial function $\hat{v}_0(|x - \hat{x}_0| - R_0)$. Furthermore, due to (3.30), we conclude from Theorem 2.9 that

$$u(t, x) \leq \hat{V}(t, x) \text{ in } [0, t_2] \times \mathbb{R}^N.$$

This together with (3.29) clearly implies

$$\Omega(t_2) \subset \left\{ x : |x - \hat{x}_0| \geq \hat{h}(t_2) \right\} \subset \left\{ x : |x - \hat{x}_0| \geq R_0/2 \right\} \text{ for all } \hat{x}_0 \in \mathbb{R}^N \setminus \Lambda_{z_2}.$$

It follows that

$$\Omega(t_2) \subset \bigcap_{\hat{x}_0 \in \mathbb{R}^N \setminus \Lambda_{z_2}} \left\{ x : |x - \hat{x}_0| \geq R_0/2 \right\} \subset \Lambda_{z_2}.$$

The proof of Lemma 3.8 is now complete. \square

We are now ready to construct a weak supersolution of problem (1.1). For any given small $\delta > 0$, denote

$$c^\delta := c_*(\mu, a + \delta, b - \delta, d),$$

and denote by $Z^\delta(r)$ the solution of (3.10) with k, a, b replaced by $c^\delta, a + \delta, b - \delta$, respectively. For $R > 0$, we define

$$\xi_R(t) := \xi_2 - \frac{R + r_2}{\sin \phi} - \left[(1 - \delta)^{-2} \frac{c^\delta}{\sin \phi} \right] t \text{ for } t \geq 0,$$

with r_2 given in Lemma 3.8 depending on $\delta > 0$, and

$$\Omega_R(t) := \Lambda_R^\phi + \xi_R(t)e_N \text{ for } t \geq 0,$$

with

$$\Lambda_R^\phi := \{x \in \Lambda^\phi : d(x, \partial \Lambda^\phi) > R\}.$$

Then define

$$\bar{u}(t, x) = \bar{u}_R(t, x) := u_R(x - \xi_R(t)e_N) \text{ for } t \geq 0, x \in \mathbb{R}^N,$$

with

$$u_R(x) := \begin{cases} (1 - \delta)^{-2} Z^\delta(d(x, \partial \Lambda_R^\phi)), & x \in \Lambda_R^\phi, \\ 0, & x \notin \Lambda_R^\phi. \end{cases}$$

We are going to show that for suitably chosen R , $\bar{u}_R(t, x)$ is a weak supersolution to the equation satisfied by $u(t_2 + t, x)$; the desired result then easily follows.

Lemma 3.9. *Let \bar{u} be given as above. Then there exists $R = R(\delta)$ sufficiently large such that \bar{u} is a weak supersolution of (1.1) with $u_0(x)$ replaced by $\bar{u}(0, x)$.*

Proof. Let us observe that $\partial \Lambda_R^\phi$ is smooth and it can be decomposed into two parts, a spherical part

$$\Sigma_R^1 := \partial B_R(0) \cap \Lambda^{\phi - \frac{\pi}{2}},$$

and part of the surface of the cone $\Lambda^\phi + \frac{R}{\sin \theta} e_N$ (recall $\theta = \pi - \phi$):

$$\Sigma_R^2 := \left(\partial \Lambda^\phi + \frac{R}{\sin \theta} e_N \right) \setminus \Lambda^{\phi - \frac{\pi}{2}}.$$

Correspondingly, we can decompose Λ_R^ϕ into two parts:

$$\Lambda_R^\phi := \Lambda_{R,1}^\phi \cup \Lambda_{R,2}^\phi \text{ with } \Lambda_{R,1}^\phi := \Lambda_R^\phi \cap \Lambda^{\phi - \frac{\pi}{2}} \text{ and } \Lambda_{R,2}^\phi := \Lambda_R^\phi \setminus \Lambda^{\phi - \frac{\pi}{2}}$$

($\Lambda_{R,1}^\phi$ and $\Lambda_{R,2}^\phi$ are illustrated in Figure 3).




FIGURE 3. The domains $\Lambda_{R,1}^\phi$ and $\Lambda_{R,2}^\phi$

In a similar way, for each $t \geq 0$, we can write

$$\Omega_R(t) = \Omega_R^1(t) \cup \Omega_R^2(t),$$

with

$$\Omega_R^1(t) := \Lambda_{R,1}^\phi + \xi_R(t) \quad \text{and} \quad \Omega_R^2(t) := \Lambda_{R,2}^\phi + \xi_R(t).$$

Clearly

$$d(x, \partial\Lambda_R^\phi) = d(x, \Sigma_R^1) = |x| - R \quad \text{if } x \in \Lambda_{R,1}^\phi,$$

and, by some simple geometrical calculations, for $x = (x', x_N) \in \Lambda_{R,2}^\phi$,

$$d(x, \partial\Lambda_R^\phi) = d(x, \Sigma_R^2) = |x'| \cos \theta + x_N \sin \theta - R.$$

It is straightforward to check that \bar{u} and $\nabla_x \bar{u}$ are continuous in $\bigcup_{t \geq 0} \overline{\Omega_R(t)}$, that $\nabla_x^2 \bar{u}$, \bar{u}_t are continuous in $\bigcup_{t > 0} \overline{\Omega_R(t)}$.

Next, we show that for $R > 0$ sufficiently large,

$$\bar{u}_t - d\Delta \bar{u} \geq a\bar{u} - b\bar{u}^2 \quad \text{for } x \in \Omega_R(t), \quad t > 0. \quad (3.31)$$

Denote $z := x - \xi_R(t)e_N$; direct calculation shows that, for $x \in \Omega_R^1(t)$ and $t > 0$,

$$\bar{u}_t = (1 - \delta)^{-4} (Z^\delta)'(z) \frac{z_N}{|z|} \frac{c^\delta}{\sin \phi},$$

and

$$d\Delta \bar{u} = (1 - \delta)^{-2} \left[d(Z^\delta)''(z) + \frac{d(N-1)}{|z|} (Z^\delta)'(z) \right].$$

Due to $|z| > R$ and $z_N \geq |z| \sin \theta = |z| \sin \phi$ for $x \in \Omega_R^1(t)$ (i.e., $z \in \Lambda_{R,1}^\phi$), and $(Z^\delta)' > 0$, we have

$$\begin{aligned} \bar{u}_t - d\Delta \bar{u} &\geq (1-\delta)^{-2} \left[(1-\delta)^{-2} c^\delta (Z^\delta)' - \frac{d(N-1)}{|z|} (Z^\delta)' - d(Z^\delta)'' \right] \\ &\geq (1-\delta)^{-2} \left(\left[(1-\delta)^{-2} c^\delta - \frac{d(N-1)}{R} \right] (Z^\delta)' - d(Z^\delta)'' \right) \\ &\geq (1-\delta)^{-2} \left(c^\delta (Z^\delta)' + \left[\delta c^\delta - \frac{d(N-1)}{R} \right] (Z^\delta)' - d(Z^\delta)'' \right). \end{aligned}$$

Therefore, if we choose

$$R \geq \frac{d(N-1)}{\delta c^\delta}, \quad (3.32)$$

then for $x \in \Omega_R^1(t)$,

$$\bar{u}_t - d\Delta \bar{u} \geq (1-\delta)^{-2} [c^\delta (Z^\delta)' - d(Z^\delta)'] \geq (a+\delta)\bar{u} - (b-\delta)\bar{u}^2 \geq a\bar{u} - b\bar{u}^2.$$

For $x \in \Omega_R^2(t)$ and $t > 0$, it follows from a direct calculation that

$$\bar{u}_t = (1-\delta)^{-4} c^\delta (Z^\delta)',$$

and

$$d\Delta \bar{u} = (1-\delta)^{-2} \left[d(Z^\delta)'' + \frac{d(N-2) \cos \theta (Z^\delta)'}{|x'|} \right],$$

where $x' := (x_1, \dots, x_{N-1}) \in \mathbb{R}^{N-1}$.

It is easily checked that for $z = (z', z_N) \in \Lambda_{R,2}^\phi$, we always have $|z'| \geq R \cos \theta$. Thus using $z = x - \xi_R(t)e_N \in \Lambda_{R,2}^\phi$, we obtain

$$\frac{\cos \theta}{|x'|} \leq \frac{1}{R}.$$

Thus for $R > 0$ satisfying (3.32) and $x \in \Omega_R^2(t)$, we have

$$\begin{aligned} \bar{u}_t - d\Delta \bar{u} &\geq (1-\delta)^{-2} \left[(1-\delta)^{-2} c^\delta (Z^\delta)' - \frac{d(N-2)}{R} (Z^\delta)' - d(Z^\delta)'' \right] \\ &\geq (1-\delta)^{-2} \left[c^\delta (Z^\delta)' + \left(\delta c^\delta - \frac{d(N-2)}{R} \right) (Z^\delta)' - d(Z^\delta)'' \right] \\ &\geq (1-\delta)^{-2} [c^\delta (Z^\delta)' - d(Z^\delta)'] \\ &\geq (a+\delta)\bar{u} - (b-\delta)\bar{u}^2, \end{aligned}$$

and hence,

$$\bar{u}_t - d\Delta \bar{u} \geq a\bar{u} - b\bar{u}^2.$$

We have thus proved that (3.31) holds for all R satisfying (3.32). We henceforth fix such an R .

We next define

$$\Phi(t, x) = R - d(x, \partial\Omega_R(t)) = \begin{cases} R - |x - \xi_R(t)e_N|, & x \in \Omega_R^1(t), \\ R - [|x'| \cos \theta + (x_N - \xi_R(t)) \sin \theta], & x \in \Omega_R^2(t). \end{cases}$$

Clearly, Φ is smooth, $\Omega_R(t) = \{x : \Phi(t, x) < 0\}$ and $|\nabla_x \Phi| \neq 0$ for $x \in \partial\Omega_R(t)$. We next show that

$$\Phi_t \leq \mu \nabla_x \bar{u} \cdot \nabla_x \Phi \quad \text{for } x \in \partial\Omega_R(t), \quad t > 0. \quad (3.33)$$

It is straightforward to calculate that, for $x \in \partial\Omega_R(t)$ and $t > 0$,

$$\nabla_x \Phi \cdot \nabla_x \bar{u} = -(1-\delta)^{-2} (Z^\delta)'(0),$$

and

$$\Phi_t(t, x) = \begin{cases} -(1-\delta)^{-2} \frac{x_N - \xi_R(t)}{|x - \xi_R(t)e_N|} \frac{c^\delta}{\sin \theta} & \text{if } x \in \Omega_R^1(t), \\ -(1-\delta)^{-2} c^\delta & \text{if } x \in \Omega_R^2(t). \end{cases}$$

On the other hand, it is easily seen that for any $z \in \Lambda_{R,1}^\phi$, $z_N \geq |z| \sin \theta$. It then follows that

$$\frac{x_N - \xi_R(t)}{|x - \xi_R(t)e_N|} \geq \sin \theta \quad \text{for } x \in \Omega_R^1(t),$$

and hence

$$\Phi_t(t, x) \leq -(1-\delta)^{-2} c^\delta \quad \text{for } x \in \Omega_R(t).$$

From this and $\mu(Z^\delta)'(0) = c^\delta$, we deduce (3.33). We may now apply Theorem 2.10 to conclude that \bar{u} is a weak supersolution of (1.1) with $u_0(x)$ replaced by $\bar{u}(0, x)$. \square

Lemma 3.10. *Let $u(t, x)$ and $\Omega(t)$ be given in the statement of Theorem 3.3. Then for any $\epsilon > 0$, there exists $T_3 = T_3(\epsilon) > 0$ such that*

$$\Omega(t) \subset \Lambda^\phi - \left(\frac{c_*}{\sin \phi} + \epsilon \right) t e_N \quad \text{for all } t \geq T_3.$$

Proof. For any small $\delta > 0$, let $t_2 = t_2(\delta)$ and $R = R(\delta)$ be given in Lemma 3.8 and Lemma 3.9, respectively. By Proposition 3.2, there exists $r_3 = r_3(\delta) > 0$ such that

$$Z^\delta(r) \geq (1-\delta) \frac{a+\delta}{b-\delta} \quad \text{for all } r \geq r_3. \quad (3.34)$$

We first claim that

$$\Omega(t_2) \subset \Omega_R(0) \quad (3.35)$$

and

$$u(t_2, x) \leq \bar{u}(0, x + \tilde{r}_3 e_N) \quad \text{for } x \in \Omega(t_2), \quad (3.36)$$

where

$$\tilde{r}_3 := \frac{r_3}{\sin \theta}.$$

Indeed, it is easily seen from the definition that

$$\Omega_R(0) \supset \Lambda_{z_2} = \Lambda^\phi + \left(\xi_2 - \frac{r_2}{\sin \phi} \right) e_N.$$

Thus, (3.35) is a consequence of $\Omega(t_2) \subset \Lambda_{z_2}$ proved in Lemma 3.8.

We now prove (3.36). For any $x \in \Omega(t_2)$, due to $\Omega(t_2) \subset \Lambda_{z_2} \subset \Omega_R(0)$, we obtain

$$\begin{aligned} d\left(x + \frac{r_3}{\sin \theta} e_N, \partial \Omega_R(0)\right) &\geq d\left(x + \frac{r_3}{\sin \theta} e_N, \partial \Lambda_{z_2}\right) \\ &= d\left(x, \partial \Lambda_{z_2} - \frac{r_3}{\sin \theta} e_N\right) \\ &\geq r_3 + d(x, \partial \Lambda_{z_2}) \geq r_3. \end{aligned}$$

Thus, for $x \in \Omega(t_2)$, due to $(Z^\delta)'(r) > 0$ in $(0, \infty)$, we have

$$\begin{aligned} \bar{u}(0, x + \tilde{r}_3 e_N) &= (1-\delta)^{-2} Z^\delta \left(d\left(x + \frac{r_3}{\sin \theta} e_N - \xi_R(0) e_N, \partial \Lambda_R^\phi\right) \right) \\ &= (1-\delta)^{-2} Z^\delta \left(d\left(x + \frac{r_3}{\sin \theta} e_N, \partial \Omega_R(0)\right) \right) \\ &\geq (1-\delta)^{-2} Z^\delta(r_3). \end{aligned}$$

This together with (3.25) and (3.34) implies that

$$\bar{u}(0, x + \tilde{r}_3 e_N) \geq (1 - \delta)^{-1} \frac{a + \delta}{b - \delta} \geq u(t_2, x) \quad \text{for } x \in \Omega(t_2),$$

which proves (3.36).

By Lemma 3.9, $\bar{u}(t, x + \tilde{r}_3 e_N)$ is a weak supersolution of problem (1.1) with u_0 replaced by $\bar{u}(0, x + \tilde{r}_3 e_N)$, and since $u(t + t_2, x)$ is a weak solution of (1.1) with u_0 replaced by $u(t_2, x)$, it follows from (3.35), (3.36) and Theorem 2.9 that

$$u(t + t_2, x) \leq \bar{u}(t, x + \tilde{r}_3 e_N) \quad \text{in } [0, \infty) \times \mathbb{R}^N,$$

and hence,

$$\Omega(t) \subset \Omega_R(t - t_2) - \tilde{r}_3 e_N = \Lambda_R^\phi + [\xi_R(t - t_2) - \tilde{r}_3] e_N \quad \text{for } t \geq t_2.$$

Since $\Lambda_R^\phi \subset \Lambda^\phi$, we thus obtain

$$\Omega(t) \subset \Lambda^\phi + [\xi_R(t - t_2) - \tilde{r}_3] e_N \quad \text{for } t \geq t_2.$$

Clearly,

$$\xi_R(t - t_2) - \tilde{r}_3 = M - (1 - \delta)^{-2} \frac{c^\delta}{\sin \theta} t,$$

with

$$M = M_\delta := \xi_2 - \frac{R + r_2 + r_3}{\sin \theta} + (1 - \delta)^{-2} \frac{c^\delta}{\sin \theta} t_2.$$

Since

$$\lim_{\delta \rightarrow 0} (1 - \delta)^{-2} c^\delta = c_*,$$

for any small $\epsilon > 0$, we can find some $\delta'_\epsilon \in (0, \epsilon)$ such that

$$-(1 - \delta'_\epsilon)^{-2} \frac{c^{\delta'_\epsilon}}{\sin \theta} \geq -\frac{c_*}{\sin \theta} - \frac{\epsilon}{2}.$$

We now fix $\delta = \delta'_\epsilon$ and obtain

$$\xi_R(t - t_2) - \tilde{r}_3 \geq M - \left(\frac{c_*}{\sin \theta} + \frac{\epsilon}{2} \right) t = M + \frac{\epsilon}{2} t - \left(\frac{c_*}{\sin \theta} + \epsilon \right) t \geq - \left(\frac{c_*}{\sin \theta} + \epsilon \right) t$$

for $t \geq T_3$ with

$$T_3 = T_3(\epsilon) := \max \left\{ t_2, \frac{2}{\epsilon} |M| \right\}.$$

Therefore,

$$\Omega(t) \subset \Lambda^\phi - \left(\frac{c_*}{\sin \theta} + \epsilon \right) t e_N \quad \text{for } t \geq T_3,$$

as desired. \square

It is easily seen that (3.12) in Theorem 3.3 follows from Lemmas 3.7 and 3.10, while (3.13) is a direct consequence of (3.23) and (3.25). Thus, Theorem 3.3 is now proved.

Theorem 3.3 implies that if Ω_0 satisfies (3.1) with $\phi \in (\pi/2, \pi)$, then for all large time, the free boundary $\partial\Omega(t)$ propagates to infinity in the negative x_N -direction with speed $c_*/\sin \phi$. Moreover, given any direction $\nu \in \mathbb{S}^{N-1}$ pointing outward of Λ^ϕ , if we denote

$$\psi := \arccos[(-e_N) \cdot \nu] \quad (\text{and so } \psi \in (0, \theta) = (0, \pi - \phi)),$$

then the spreading of $\Omega(t)$ in the direction ν is roughly at speed $c_*/(\sin \phi \cos \psi)$. In sharp contrast, we will show in the following theorem, that when Ω_0 satisfies (3.1) with $\phi \in (0, \pi/2)$, the spreading of $\Omega(t)$ in a set of directions $\nu \in \mathbb{S}^{N-1}$ pointing outward of Λ^ϕ , including $\nu = -e_N$, is roughly at the speed c_* . (This set of directions ν is given by Σ_ϕ below.)

Theorem 3.11. *Let $u(t, x)$ be the unique weak solution of problem (1.1) with Ω_0 satisfying (3.1) for some $\phi \in (0, \pi/2)$, and u_0 satisfying (2.2) and (3.11). Denote $\Omega(t) = \{x : u(t, x) > 0\}$. Then for any $\epsilon > 0$, there exists $\tilde{T} = \tilde{T}(\epsilon) > 0$ such that*

$$N[\Lambda^\phi, (c_* - \epsilon)t] \subset \Omega(t) \subset N[\Lambda^\phi, (c_* + \epsilon)t] \text{ for all } t \geq \tilde{T}. \quad (3.37)$$

Moreover, we have

$$\lim_{t \rightarrow \infty} \left[\sup_{x \in N[\Lambda^\phi, (c_* - \epsilon)t]} \left| u(t, x) - \frac{a}{b} \right| \right] = 0. \quad (3.38)$$

Proof. We prove this theorem by two steps.

Step 1: *Proof of the first relation of (3.37) and (3.38).*

Due to the assumptions (3.1) and (3.11), we can find $\xi_0 > \xi_1$ such that

$$u_0(x) \geq \sigma_0 := \frac{1}{2} \liminf_{d(\tilde{x}, \partial\Omega_0) \rightarrow \infty, \tilde{x} \in \Omega_0} u_0(\tilde{x}) \text{ for all } x \in \Lambda^\phi + \xi_0 e_N.$$

Let $R^* > 0$ be the positive constant given in (3.4). Then we choose a radial function $\tilde{v}_0 \in C^2([0, R^*])$ such that

$$0 < \tilde{v}_0(x) \leq \sigma_0 \text{ in } [0, R^*), \quad \tilde{v}_0'(0) = \tilde{v}_0(R^*) = 0.$$

Next, for any fixed $x_0 \in \mathbb{R}^N$ with $B_{R^*}(x_0) \subset \Lambda^\phi + \xi_0 e_N$, set $r = |x - x_0|$, and consider the following radially symmetric free boundary problem

$$\begin{cases} \tilde{v}_t - d\Delta\tilde{v} = \tilde{v}(a - b\tilde{v}), & t > 0, 0 < r < \tilde{k}(t), \\ \tilde{v}_r(t, 0) = 0, \quad \tilde{v}(t, \tilde{k}(t)) = 0, & t > 0, \\ \tilde{k}'(t) = -\mu v_r(t, \tilde{k}(t)), & t > 0, \\ \tilde{k}(0) = R^*, \quad \tilde{v}(0, r) = \tilde{v}_0(r), & 0 \leq r \leq R^*. \end{cases} \quad (3.39)$$

It follows from [7, Theorems 2.1, 2.5 and Corollary 3.7] that problem (3.39) admits a unique classical solution $(\tilde{v}(t, r), \tilde{k}(t))$ defined for all $t \geq 0$, and

$$\lim_{t \rightarrow \infty} \frac{\tilde{k}(t)}{t} = c_*. \quad (3.40)$$

Furthermore, applying [8, Theorem 6.4] to this problem, we have

$$\lim_{t \rightarrow \infty} \max_{r \leq (c_* - \epsilon/2)t} \left| \tilde{v}(t, r) - \frac{a}{b} \right| = 0 \quad (3.41)$$

for every small $\epsilon > 0$.

On the other hand, by our choices of \tilde{v}_0 and ξ_0 , we have

$$\tilde{v}_0(|x - x_0|) \leq u_0(x) \text{ for all } x \in B_{R^*}(x_0).$$

Then extending $\tilde{v}(t, |x - x_0|)$ to be zero for $|x - x_0| > \tilde{k}(t)$ and applying the comparison principle Theorem 2.9, we obtain

$$\tilde{v}(t, |x - x_0|) \leq u(t, x) \text{ for all } t \geq 0, x \in \mathbb{R}^N, \quad (3.42)$$

and hence,

$$\left\{ x \in \mathbb{R}^N : |x - x_0| \leq \tilde{k}(t) \right\} \subset \Omega(t) \text{ for all } t \geq 0.$$

This together with (3.40) implies that, for any $\epsilon > 0$, there exists $\tilde{T}_1 = \tilde{T}_1(\epsilon) > 0$ such that

$$\left\{ x \in \mathbb{R}^N : |x - x_0| \leq \left(c_* - \frac{\epsilon}{2}\right)t \right\} \subset \Omega(t) \text{ for all } t \geq \tilde{T}_1.$$

Note that the above analysis remains valid if x_0 replaced by any point \tilde{x}_0 such that $B_{R^*}(\tilde{x}_0) \subset \Lambda^\phi + \xi_0 e_N$, and that the constant \tilde{T}_1 is independent of the choice of such \tilde{x}_0 . It then follows that

$$\bigcup_{B_{R^*}(x_0) \subset \Lambda^\phi + \xi_0 e_N} \left\{ x \in \mathbb{R}^N : |x - x_0| \leq (c_* - \frac{\epsilon}{2})t \right\} \subset \Omega(t) \quad \text{for all } t \geq \tilde{T}_1.$$

Furthermore, it is easily seen that there exists $\tilde{T}_2 = \tilde{T}_2(\epsilon) \geq \tilde{T}_1$ such that, for all $t \geq \tilde{T}_2$,

$$N[\Lambda^\phi, (c_* - \epsilon)t] \subset \bigcup_{B_{R^*}(x_0) \subset \Lambda^\phi + \xi_0 e_N} \left\{ x \in \mathbb{R}^N : |x - x_0| \leq (c_* - \frac{\epsilon}{2})t \right\}. \quad (3.43)$$

We thus obtain the first relation of (3.37).

To complete the proof of this step, it remains to show (3.38). On the one hand, by (3.42) and (3.43), we have

$$\inf_{x \in N[\Lambda^\phi, (c_* - \epsilon)t]} u(t, x) \geq \inf \left\{ \min_{|x - x_0| \leq (c_* - \epsilon/2)t} \tilde{v}(t, |x - x_0|) : B_{R^*}(x_0) \subset \Lambda^\phi + \xi_0 e_N \right\}$$

for all $t \geq \tilde{T}_2$. It further follows from (3.41) that

$$\liminf_{t \rightarrow \infty} \inf_{x \in N[\Lambda^\phi, (c_* - \epsilon)t]} u(t, x) \geq \frac{a}{b}.$$

On the other hand, since u_0 is bounded, by the arguments used at the beginning of the proof of Lemma 3.8, we have

$$\limsup_{t \rightarrow \infty} \sup_{x \in \mathbb{R}^N} u(t, x) \leq \frac{a}{b}.$$

Combining the above, we immediately obtain (3.38).

Step 2: *Proof of the second relation of (3.37).*

Choose a one-dimensional function $\tilde{w}_0 \in C^2((-\infty, 1]) \cap L^\infty((-\infty, 1])$ such that

$$\tilde{w}_0(y) \geq \|u_0\|_{L^\infty(\Omega_0)} \text{ in } (-\infty, 0], \quad \tilde{w}_0(x) > 0 \text{ in } (0, 1) \quad \text{and} \quad \tilde{w}_0(1) = 0.$$

Then we consider the following one-dimensional free boundary problem

$$\begin{cases} \tilde{w}_t - d\tilde{w}_{yy} = \tilde{w}(a - b\tilde{w}), & t > 0, \quad -\infty < y < \tilde{\rho}(t), \\ \tilde{w}(t, \tilde{\rho}(t)) = 0, & t > 0, \\ \tilde{\rho}'(t) = -\mu\tilde{w}_y(t, \tilde{\rho}(t)), & t > 0, \\ \tilde{\rho}(0) = 1, \quad \tilde{w}(0, y) = \tilde{w}_0(y), & -\infty < y \leq 1. \end{cases} \quad (3.44)$$

It follows from [5, Theorem 2.11] that, (3.44) admits a (unique) classical solution $(\tilde{w}(t, y), \tilde{\rho}(t))$ defined for all $t > 0$ and $\tilde{\rho}'(t) > 0$, $\tilde{w}(t, y) > 0$ for $-\infty < y < \tilde{\rho}(t)$, $t > 0$.

For any

$$\nu \in \Sigma_\phi := \left\{ x \in \mathbb{S}^{N-1} : \arccos(x \cdot e_N) \in [\phi + \frac{\pi}{2}, \pi] \right\},$$

define

$$\tilde{\Omega}_\nu(t) = \left\{ x \in \mathbb{R}^N : x \cdot \nu \leq \tilde{\rho}(t) \right\} \quad \text{and} \quad \tilde{w}_\nu(t, x) = \tilde{w}(t, x \cdot \nu).$$

Clearly, we have $\Omega_0 - \xi_2 e_N \subset \tilde{\Omega}_\nu(0)$, and $u_0(\cdot + \xi_2 e_N) \leq \tilde{w}_\nu(0, \cdot)$ in $\Omega_0 - \xi_2 e_N$. Then by similar comparison arguments as those used in the proof of Lemma 3.7, we obtain

$$\Omega(t) - \xi_2 e_N \subset \tilde{\Omega}_\nu(t) \quad \text{for all } t \geq 0, \quad \nu \in \Sigma_\phi. \quad (3.45)$$

Furthermore, it follows from the proof of [9, Theorem 4.2] with similar modifications as those given in the proof of Lemma 3.5 that, for the given $\epsilon > 0$, there exists $\tilde{T}_3 = \tilde{T}_3(\epsilon) > 0$ such that

$$\tilde{\rho}(t) \leq (c_* + \frac{\epsilon}{2})t \quad \text{for all } t \geq \tilde{T}_3.$$

This together with (3.45) implies

$$\Omega(t) - \xi_2 e_N \subset \left\{ x \in \mathbb{R}^N : x \cdot \nu \leq \left(c_* + \frac{\epsilon}{2}\right)t \right\} \text{ for all } t \geq \tilde{T}_3, \nu \in \Sigma_\phi.$$

Since

$$N[\Lambda^\phi, (c_* + \frac{\epsilon}{2})t] = \bigcup_{\nu \in \Sigma_\phi} \left\{ x \in \mathbb{R}^N : x \cdot \nu \leq \left(c_* + \frac{\epsilon}{2}\right)t \right\},$$

by enlarging \tilde{T}_3 if necessary (depending on ξ_2 and ϵ), we obtain

$$\Omega(t) \subset N[\Lambda^\phi, (c_* + \epsilon)t] \text{ for all } t \geq \tilde{T}_3.$$

which clearly gives the second relation of (3.37). \square

Remark 3.12. *The estimates in Theorem 3.11 can be improved by making use of sharp estimates for the spreading speed for one space dimension free boundary problems in [14] and for radially symmetric free boundary problems in [15]. We leave the details to the interested reader.*

4. APPENDIX

This appendix is concerned with the existence and uniqueness of classical solutions to an auxiliary radially symmetric problem with initial range the exterior of a ball. These results have been used to construct the weak supersolution for problem (1.1) in the proof of Theorem 3.3, and here we consider a more general problem which might have other applications.

More precisely, for any given $T > 0$, $C_1 > 0$ and $C_2 > 0$, we consider the following radially symmetric free boundary problem

$$\begin{cases} v_t - d\Delta v = \tilde{g}(r, v), & 0 < t < T, h(t) < r < \infty, \\ v(t, h(t)) = 0, & 0 < t < T, \\ h'(t) = -\mu v_r(t, h(t)), & 0 < t < T, \\ h(0) = R_0, \quad v(0, r) = v_0(r - R_0), & R_0 \leq r < \infty, \end{cases} \quad (4.1)$$

where $\Delta v = v_{rr} + \frac{N-1}{r}v_r$, $R_0 > 1$ is a constant to be determined by T , C_1 , C_2 later, and v_0 is a given function in $C^2([0, \infty))$ satisfying

$$0 < v_0(r) \leq C_1 \text{ for } r \in (0, \infty), \quad v_0(0) = 0, \quad \|v_0\|_{C^1([0, \infty))} \leq C_2. \quad (4.2)$$

Here we assume that $\tilde{g}(r, v)$ is a continuous function defined over $\mathbb{R}^+ \times \mathbb{R}^+$ satisfying

$$\left. \begin{aligned} \text{(i)} \quad & \tilde{g}(r, v) \text{ is Hölder continuous in } r \geq 0, v \geq 0, \\ \text{(ii)} \quad & \tilde{g}(r, v) \text{ is locally Lipschitz in } v \text{ uniformly for } r \geq 0, \\ \text{(iii)} \quad & \tilde{g}(r, 0) \equiv 0, \\ \text{(iv)} \quad & \text{there exists } K > 0 \text{ such that } \tilde{g}(r, v) \leq Kv \text{ for all } r \geq 0 \text{ and } v \geq 0. \end{aligned} \right\} \quad (4.3)$$

Proposition 4.1. *Assume that (4.3) is satisfied. For any $T > 0$, $C_1 > 0$ and $C_2 > 0$, there exists a constant $R_0 > 1$ depending on T , C_1 and C_2 such that for any v_0 satisfying (4.2), problem (4.1) admits a unique solution $(v(t, r), h(t))$ with $h \in C^1([0, T])$, $v \in C^{1,2}(D_T)$, where $D_T = \{(t, r) : t \in [0, T], r \in [h(t), \infty)\}$. Moreover, $v(t, r) > 0$, $h'(t) < 0$ for $0 < t \leq T$, $h(t) < r < \infty$, and*

$$h(T) \geq \frac{R_0}{2}. \quad (4.4)$$

Proof. For given $R_0 > 1$, following the proof of [7, Theorem 2.1] we can show that (4.1) has a unique solution for some small $T > 0$. The proof involves the straightening of the free boundary, and different from [7], the resulting problem here is over an unbounded interval for the new space variable. However, the estimates easily carry over (by using suitable interior estimates, similar to a

related situation treated in [10]) and so we obtain the local existence and uniqueness all the same. Moreover, all the stated properties in the proposition, except (4.4), also hold.

Furthermore, the solution can be extended as long as $h(t) > 0$. Let $T_\infty = T_\infty(R_0)$ be the maximal existence time of the solution. If $h(t) \geq R_0/2$ for all $t \in (0, T_\infty)$, then necessarily $T_\infty = \infty$ and thus $h(T) > R_0/2$, and there is nothing left to prove.

Suppose now $h(t_0) < R_0/2$ for some $t_0 \in (0, T_\infty)$. Since $h(0) = R_0$, we can find $T_0 = T_0(R_0) \in (0, t_0)$ such that $h(T_0) = R_0/2$. We are going to show that $T_0(R_0) > T$ provided that R_0 is sufficiently large, which clearly implies $h(T) > R_0/2$, as desired. We use an indirect argument and assume that $T_0(R_0) \leq T$ for all $R_0 > 1$.

By the assumption (4.3), it follows from the parabolic comparison principle that $v(t, r) \leq \bar{v}(t)$ for $r > h(t)$ and $0 \leq t \leq T_0$, where $\bar{v}(t)$ is the solution to

$$\frac{d\bar{v}}{dt} = K\bar{v} \text{ for } t > 0; \quad \bar{v}(0) = \|v_0\|_{L^\infty([0, \infty))}.$$

Clearly, $\bar{v}(t) = \|v_0\|_{L^\infty([0, \infty))} e^{Kt}$ for $t \geq 0$. Thus, we have

$$v(t, r) \leq C_3 := C_1 e^{KT} \text{ for } r > h(t), 0 \leq t \leq T_0.$$

Next we prove that there exists a positive constant C_4 independent of R_0 such that

$$-C_4 \leq h'(t) < 0 \text{ for } t \in (0, T_0]. \quad (4.5)$$

This would lead to a contradiction, since it follows that

$$R_0/2 = h(T_0) \geq h(0) - C_4 T_0 \geq R_0 - C_4 T > R_0/2 \text{ for all large } R_0.$$

Therefore to complete the proof, it suffices to show (4.5). To this end, for $M > 0$ to be determined later, we define

$$\Omega = \Omega_M := \left\{ (t, r) : 0 < t \leq T_0, \quad h(t) < r < h(t) + M^{-1} \right\}$$

and construct an auxiliary function

$$w(t, r) := C_3 [2M(r - h(t)) - M^2(r - h(t))^2] \in (0, C_3) \text{ for } (t, r) \in \Omega.$$

We will show that for some suitable choice of $M > 0$, $w(t, r) \geq v(t, r)$ holds over Ω .

Direct calculations give, for $(t, r) \in \Omega$,

$$w_t = 2C_3 M(-h'(t)) [1 - M(r - h(t))] \geq 0,$$

and

$$-w_r = -2C_3 M [1 - M(r - h(t))] \geq -2C_3 M, \quad -w_{rr} = 2C_3 M^2.$$

Making use of $1/2 \leq R_0/2 \leq h(t)$ for $t \in (0, T_0]$, we obtain, for $(t, r) \in \Omega$,

$$w_t - d \left(w_{rr} + \frac{N-1}{r} w_r \right) \geq 2dC_3 M^2 - 2dC_3 M \frac{N-1}{r} \geq 2dC_3 [M^2 - 2(N-1)M].$$

Thus, if we choose

$$M \geq (N-1) + \sqrt{\frac{K}{2d} + (N-1)^2},$$

where K is given in (4.3), then

$$w_t - d \left(w_{rr} + \frac{N-1}{r} w_r \right) \geq KC_3 \geq \tilde{g}(r, w) \text{ for } (t, r) \in \Omega.$$

Let us also note that for $t \in (0, T_0]$,

$$w(t, h(t) + M^{-1}) = C_3 \geq v(t, h(t) + M^{-1}),$$

and

$$w(t, h(t)) = 0 = v(t, h(t)).$$

Thus, if our choice of M also ensures

$$v(0, r) \leq w(0, r) \text{ for } r \in [R_0, R_0 + M^{-1}],$$

then we can apply the maximum principle to $w - v$ over Ω to deduce that $v(t, r) \leq w(t, r)$ for $(t, r) \in \Omega$. It would then follow that

$$v_r(t, h(t)) \leq w_r(t, h(t)) = 2MC_3,$$

and so

$$h'(t) = -\mu v_r(t, h(t)) \geq -C_4 := -2MC_3\mu,$$

as we wanted.

To complete the proof, we calculate

$$w_r(0, r) = 2C_3M[1 - M(r - R_0)] \geq C_3M \text{ for } r \in [R_0, R_0 + (2M)^{-1}].$$

Therefore, upon choosing

$$M := \max \left\{ (N-1) + \sqrt{\frac{K}{2d} + (N-1)^2}, \frac{2C_2}{C_3} \right\},$$

we will have

$$w_r(0, r) \geq 2C_2 > v'_0(r) = v_r(0, r) \text{ for } [R_0, R_0 + (2M)^{-1}].$$

Since $w(0, R_0) = v(0, R_0) = 0$, the above inequality implies

$$w(0, r) \geq v(0, r) \text{ for } r \in [R_0, R_0 + (2M)^{-1}].$$

For $r \in [R_0 + (2M)^{-1}, R_0 + M^{-1}]$, we have $w(0, r) \geq C_3/2$ and

$$v(0, r) \leq \|v(0, \cdot)\|_{C^1([R_0, R_0 + M^{-1}])} M^{-1} \leq C_2 M^{-1} \leq \frac{C_3}{2},$$

which clearly gives $v(0, r) \leq w(0, r)$. Since M is independent of R_0 , this completes the proof of (4.5). \square

REFERENCES

- [1] D. G. Aronson, H. F. Weinberger, *Multidimensional nonlinear diffusions arising in population genetics*, Adv. Math. **30** (1978), 33-76.
- [2] G. Bunting, Y. Du and K. Krakowski, *Spreading speed revisited: Analysis of a free boundary model, Networks and Heterogeneous Media* (special issue dedicated to H. Matano), **7** (2012), 583-603.
- [3] J. R. Cannon, C. D. Hill, *On the movement of a chemical reaction interface*, Indiana Univ. Math. J. **20** (1970), 429-454.
- [4] A. De Pablo, J. L. Vazquez, *Travelling waves and finite propagation in a reaction-diffusion equation*, J. Differential Equations **93** (1991), 19-61.
- [5] W. Ding, Y. Du and X. Liang, *Spreading in space-time periodic media governed by a monostable equation with free boundaries, Part 1: Continuous initial functions*, J. Diff. Eqns. **262** (2017), 4988-5021.
- [6] Y. Du, C. Gui, K. Wang and M. Zhou, *Semi-waves with λ -shaped free boundary for nonlinear Stefan problems*, in preparation.
- [7] Y. Du, Z. Guo, *Spreading-Vanishing dichotomy in the diffusive logistic model with a free boundary II*, J. Differential Equations **250** (2011), 4336-4366.
- [8] Y. Du, Z. Guo, *The Stefan problem for the Fisher-KPP equation*, J. Differential Equations **253** (2012), 996-1035.
- [9] Y. Du, Z. Lin, *Spreading-vanishing dichotomy in the diffusive logistic model with a free boundary*, SIAM J. Math. Anal. **42** (2010), 377-405.
- [10] Y. Du and Z. Lin, *The diffusive competition model with a free boundary: Invasion of a superior or inferior competitor*, Discrete Cont. Dyn. Syst.-B **19** (2014), 3105-3132.
- [11] Y. Du, B. Lou, *Spreading and vanishing in nonlinear diffusion problems with free boundaries*, J. Eur. Math. Soc. **17** (2015), 2673-2724.
- [12] Y. Du, B. Lou and M. Zhou, *Spreading and vanishing for nonlinear Stefan problems in high space dimensions*, J. Ellip. Par. Eqns. **2** (2016), 297-321.

- [13] Y. Du, H. Matano and K. Wang, *Regularity and asymptotic behavior of nonlinear Stefan problems*, Arch. Rational Mech. Anal. **212** (2014), 957-1010.
- [14] Y. Du, H. Matsuzawa and M. Zhou, *Sharp estimate of the spreading speed determined by nonlinear free boundary problems*, SIAM J. Math. Anal. **46** (2014), 375-396.
- [15] Y. Du, H. Matsuzawa and M. Zhou, *Spreading speed and profile for nonlinear Stefan problems in high space dimensions*, J. Math. Pures Appl. **103** (2015), 741-787.
- [16] A. Friedman, *The Stefan problem in several space variables*, Trans. Amer. Math. Soc. **132** (1968), 51-87. (Correction, Trans. Amer. Math. Soc. **142** (1969), 557.)
- [17] R.A. Fisher, *The wave of advance of advantageous genes*, Ann. Eugenics **7** (1937), 335-369.
- [18] F. Hamel, N. Nadirashvili, *Traveling waves and entire solutions of the Fisher-KPP equation in \mathbb{R}^N* , Arch. Ration. Mech. Anal. **157** (2001), 91-163.
- [19] F. Hamel, R. Monneau, J.-M. Roquejoffre, *Existence and qualitative properties of multidimensional conical bistable fronts*, Discrete Contin. Dyn. Syst. **13** (2005), 1069-1096.
- [20] F. Hamel, R. Monneau, J.-M. Roquejoffre, *Asymptotic properties and classification of bistable fronts with Lipschitz level sets*, Discrete Contin. Dyn. Syst. **14** (2006), 75-92.
- [21] S.L. Kamenomostskaja, *On Stefan's problem*, Math. Sb. **53** (1965), 485-514.
- [22] A.N. Kolmogorov, I.G. Petrovski and N.S. Piskunov, *Étude de l'équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique*, Bull. Univ. Moscou Sér. Internat. A 1 (1937), 1-26; English transl. in: P. Pelcé (Ed.), Dynamics of Curved Fronts, Academic Press, 1988, 105-130.
- [23] O.A. Ladyzenskaja, V.A. Solonnikov, N.N. Ural'tseva, *Linear and Quasilinear Equations of Parabolic Type*, Amer. Math. Soc. Providence, RI, 1968.
- [24] H. Ninomiya, M. Taniguchi, *Existence and global stability of traveling curved fronts in the Allen-Cahn equations*, J. Differential Equations **213** (2005), 204-233.
- [25] M. Taniguchi, *An $(N - 1)$ -dimensional convex compact set gives an N -dimensional traveling front in the Allen-Cahn equation*, SIAM J. Math. Anal. **47** (2015), no. 1, 455-476.