

# Model Comparison tests of modified gravity from the Eöt-Wash experiment

Aditi Krishak<sup>a</sup> and Shantanu Desai,<sup>b,1</sup>

<sup>a</sup>Department of Physics, Indian Institute of Science Education and Research, Bhopal, Madhya Pradesh 462066, India

<sup>b</sup>Department of Physics, Indian Institute of Technology, Hyderabad, Telangana-502285, India

E-mail: [aditi16@iiserb.ac.in](mailto:aditi16@iiserb.ac.in), [shntn05@gmail.com](mailto:shntn05@gmail.com)

**Abstract.** Perivolaropoulos et al [1] (P19) have argued that the residual torque data in the Eöt-Wash experiment is consistent with an oscillating signal. This could either be a signature of non-local modified gravity theories or some other systematic error in the data. We independently assess the viability of such an oscillating signal in the same data using Bayesian and information theoretical criterion, to complement the frequentist analysis in P19. We fit this data to three different parametrizations (an offset Newtonian, Yukawa, and an oscillating model), and assess the significance of the oscillating model using AIC, BIC, WAIC, and Bayes factor. All these techniques provide decisive evidence for the oscillating model compared to the Newtonian model, provided the phase is fixed at the same value as P19. If the phase is allowed to vary, then significance from BIC, WAIC, and Bayes based tests reduces to strong evidence, whereas only AIC still shows decisive evidence. Our analysis codes have been made publicly available.

---

<sup>1</sup>Corresponding author.

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Summary of Eöt-Wash results and P19</b>	<b>2</b>
<b>3</b>	<b>Model comparison techniques</b>	<b>3</b>
<b>4</b>	<b>Analysis and Results</b>	<b>5</b>
4.1	Model Comparison using $\mathbf{AIC_c}$ and BIC	7
4.2	Model Comparison using WAIC	7
4.3	Bayesian Model Comparison	7
<b>5</b>	<b>Conclusions</b>	<b>8</b>

---

## 1 Introduction

Despite being more than a century old, General relativity (GR) currently agrees with all tests using solar system, binary pulsar, and gravitational wave based observations [2–9]. In spite of this, a large number of modified theories of gravity have been explored since the inception of GR. Most of the recent resurgence in these alternate theories of gravity has been driven by the need to address problems in Cosmology such as Dark matter, Dark Energy, Inflation, Baryogenesis, and data driven cosmological tensions [10–23]. Apart from this, a number of alternatives have also been proposed to resolve conceptual issues, such as the Big-Bang singularity [24], arrow of time [25, 26], or the quantization of gravity [27–29].

One of the most stringent probes of gravity at very short distance scales are torsion balance experiments. For over three decades, the Eöt-Wash group at the University of Washington <sup>1</sup> has been conducting a series of such high precision torsion balance based tests of gravity at sub-millimeter scales to look for departures from Newtonian gravity. Their own analysis has not revealed any signs of new physics. However, other authors [1] (P19 hereafter) have independently analyzed the data from these experiments [30–33], and have argued that the residual data show signatures of an oscillating signal. They concluded that one possible explanation for the observed oscillations in the data, could be that this is a potential signature of non-local theories of gravity [34–40]. Other possible reasons include statistical fluctuations of the data in one of the experiments conducted, or a periodic distance-dependent systematic feature of the data [41].

In this work, we independently assess the statistical significance of this result using multiple model selection methods motivated from Bayesian and information theoretic considerations. This complements the frequentist model comparison analysis done in P19. We have previously used these same techniques to address a number of model selection problems in Astrophysics and Cosmology [42–47].

The outline of this manuscript is as follows. We recap the details of the Eöt-Wash experiment and analysis of their data by P19 in Sect. 2. We provide an abridged summary of different model comparison techniques used in Sect. 3. We present the results of our

---

<sup>1</sup><https://www.npl.washington.edu/eotwash/node/1>

analysis in Sect. 4. We conclude in Sect. 5. Our analysis codes are publicly available at [https://github.com/aditikrishak/EotWash\\_analysis](https://github.com/aditikrishak/EotWash_analysis).

## 2 Summary of Eöt-Wash results and P19

P19 independently analyzed the data from the Eöt-Wash experiment [30]. The Eöt-Wash group used a torsion pendulum detector to look for departures from Newton’s law of gravity at sub-millimeter scales. The torsion pendulum was suspended by a thin tungsten fiber. The gravitational interactions were measured via a torque that developed between the holes in the torsion pendulum and similar holes machined on a molybdenum detector ring below the detector. A schematic description of the Eöt-Wash experiment can be found in Fig. 1 of Ref. [30] or Fig. 15 of P19. The Eöt-Wash group conducted three different experiments (which they named as Experiment I, II and III), consisting of the same detector ring and upper attractor disk, but using different sizes for the lower attractor disks. This feature was introduced to discriminate between potential new Physics and systematic errors. For each of these three setups, residuals between the measured torques and the expected Newtonian values were published. Based on these differences, 95% c.l. upper limits were set on possible Yukawa interactions [30–33].

P19 (and also Ref. [41]) fitted the 87 residual torque data points from these experiments ( $\delta\tau$ ) to three different functions:

$$\delta\tau_1(\alpha', m', r) = \alpha' \quad (2.1)$$

$$\delta\tau_2(\alpha', m', r) = \alpha' e^{-m'r} \quad (2.2)$$

$$\delta\tau_3(\alpha', m', r) = \alpha' \cos(m'r + \phi) \quad (2.3)$$

These correspond to an offset Newtonian potential (which can be considered as white noise for the purpose of fitting), a Yukawa potential, and an oscillating potential respectively. Here,  $\alpha'$  and  $m'$  are two ad-hoc parameters, and the relations between these parameters and the corresponding parameters of any modified gravity theory depend on the details of the experimental setup [41]. In Eq. 2.3,  $\phi$  fixed to  $3\pi/4$  in P19, as this provided the best fit compared to other phases and all model comparison was done using this premise. For each of these functions, P19 carried out  $\chi^2$  minimization to obtain the best-fit parameters. Among the three models, they found that the oscillating function Eq. 2.3 has the smallest  $\chi^2$  (cf. Table IV of P19), with  $\Delta\chi^2 \sim -15$  with respect to the other models. They assess the statistical significance based on this difference in  $\chi^2$  with respect to the other models to be  $3\sigma$ . P19 also carried out Monte-Carlo simulations of the null hypothesis and showed that the probability for the oscillation signal to be a statistical fluctuation is about 10%. Therefore, P19 concluded from this frequentist model comparison technique that what they have found is one of three possibilities: either a statistical fluctuation in one of the experiments causing this anomaly, or a periodic distance-dependent systematic endemic to such laboratory-based experiments; or a signature for a short-distance modification to GR. P19 also pointed out that such an oscillating potential at short distances is a characteristic of many non-local gravity theories [34–40], involving infinite derivatives of the Lagrangian. These theories have been constructed to solve the black hole and big-bang singularity problem [24].

Given the potential ramification for the third possibility, we independently do a fit to the same residual torque data and assess the statistical significance of the oscillating parametrization using Bayesian and information theory-based model selection techniques, which we have

previously used to address multiple problems in Astrophysics and Cosmology [42–47]. This complements the frequentist analysis in P19.

### 3 Model comparison techniques

There are basically two distinct ways used to compare or rank two models used to fit a given dataset [48–50]. The Bayesian analysis compares the probability of the model given the data, whereas the frequentist method compares the expected predictive accuracy of the two models for future data. The information theory techniques are a distinct class and have both frequentist and Bayesian interpretations [50]. More details comparing and contrasting these techniques, including the pitfalls and assumptions in each of them can be found in Refs. [48–54]. A comparison of these techniques within the statistics community can be found in Ref. [55].

We briefly summarize the different selection techniques used to rank between multiple models. More details can be found in specialized reviews [48–50, 52–55] or in some of our previous works [45–47]. In this work we only use information theory and Bayesian techniques to compare the different models, since a frequentist model comparison technique has already been carried out in P19.

- **AIC and BIC:** The Akaike Information Criterion (AIC) as well as the Bayesian Inference Criterion (BIC) are used to penalize for any free parameters to avoid overfitting. AIC is an approximate minimization of Kullback-Leibler information entropy, which estimates the distance between two probability distributions [50]. In this work we use a variant of AIC corrected for small sample sizes (called AIC<sub>c</sub>) and is given by [50]:

$$\text{AIC}_c = -2 \ln L_{\max} + 2p + \frac{2p(p+1)}{N-p-1}, \quad (3.1)$$

where  $p$  is the number of free parameters;  $L_{\max}$  is the maximum likelihood which is also used for parameter estimation.

For comparing two models using AIC, the model with the lower value is the preferred model. To assess the significance, one can apply the following “strength of evidence” rules for difference in AIC between two models  $\Delta\text{AIC} = \text{AIC}_i - \text{AIC}_{\min}$  [51, 56]:

$0 - 2$	Substantial
$4 - 7$	Considerably Less
$> 10$	Essentially None

BIC is an approximation to the Bayes factor and is given by [49]:

$$\text{BIC} = -2 \ln L_{\max} + p \ln N, \quad (3.2)$$

where all the parameters have the same interpretation as in Eq. 3.1. We note that one assumption in applying BIC is that the posterior pdf is Gaussian and this may not always be satisfied.

For the BIC, one can apply the following “strength of evidence”, where  $\Delta\text{BIC} = \text{BIC}_i - \text{BIC}_{\min}$  [51, 56]:

$\Delta\text{BIC}$	Evidence against Model i
0 – 2	Not Worth More Than A Bare Mention
2 – 6	Positive
6 – 10	Strong
> 10	Very Strong

Therefore, for both AIC and BIC, the difference between two models must be greater than 10, for the model with smaller AIC/BIC to be decisively favored.

- **WAIC:** The Widely Applicable Information Criterion (WAIC) (also known as Watanabe-Akaike information criterion) [57] uses a Bayesian approach to quantitatively describe the ability of a model to predict new data. The model with a higher predictive power is the one with a smaller WAIC value. WAIC has some advantages compared to AIC or BIC. WAIC uses the Bayesian predictive density as opposed to AIC which uses the point estimate for the parameter [48, 55, 58, 59]. If a model is singular, AIC and BIC do not work well, whereas WAIC works well for such cases. WAIC is also invariant under reparametrization and in the asymptotic limit of large sample size, WAIC is equivalent to leave-one-out cross-validation [48].

The WAIC for a given model is defined as [48, 57, 58]:

$$WAIC = -2(lppd - p_{WAIC}) \quad (3.3)$$

where  $lppd$  (log pointwise predictive density) is given by [58]:

$$lppd = \sum_{i=1}^N \ln \left( \frac{1}{S} \sum_{s=1}^S P(y_i | \theta^s) \right), \quad (3.4)$$

where  $S$  is the total number of posterior samples;  $P(y_i | \theta^s)$  is the likelihood for the data-point  $y_i$ , given the parameter vector  $\theta^s$ ; and  $N$  is the total number of data points. We note that in order to calculate  $lppd$ , one first needs to estimate the average likelihood for each data point over all the parameter samples, and then sum its logarithm over all the data points. On the other hand, in parameter estimation and also for calculating AIC/BIC, what is calculated is the sum of log likelihood over all data points *for a fixed parameter set*.  $p_{WAIC}$  in Eq. 3.3 is given by:

$$p_{WAIC} = \sum_{i=1}^N \frac{1}{S-1} \sum_{s=1}^S (\ln P(y_i | \theta^s) - \langle \ln P(y_i | \theta^s) \rangle)^2, \quad (3.5)$$

where  $\langle \ln P(y_i | \theta^s) \rangle$  is the average value of  $P(y_i | \theta^s)$  over all parameter samples. We note that  $p_{WAIC}$  is an estimate of the effective number of free parameters in the model and can be interpreted as a penalty term, which compensates for overfitting [58, 59]. In lieu of  $p_{WAIC}$ , an alternate parameterization has also been considered to compensate for the free parameters [48, 55, 57]<sup>2</sup>. Between the two, Gelman et al [55] have recommended the use of  $p_{WAIC}$  (from Eq. 3.5), as its series expansion is closer to leave-one out cross-validation.

Similar to AIC and BIC, the model with the smaller value of WAIC is the preferred model and the significance can be obtained using the same “strength of evidence” rules as used for AIC and BIC.

---

<sup>2</sup>See Eq. 12 in Ref. [55]

Parametrization	$\alpha'$ (fN · m)	$m'$ (mm <sup>-1</sup> )	$\phi$
Newtonian	$0.0001^{+0.0007}_{-0.0007}$	-	-
Yukawa	$(7.2 \pm 2.9) \times 10^7$	$366.1^{+64}_{-47}$	-
Oscillating ( $\phi$ fixed)	$0.0042^{+0.0020}_{-0.0020}$	$65.29^{+0.93}_{-0.84}$	-
Oscillating ( $\phi$ free)	$0.0042^{+0.0012}_{-0.0015}$	$65.28^{+0.20}_{-0.21}$	$(0.756^{+0.15}_{-0.15})\pi$

**Table 1.** The best-fit parameters obtained for the three parametrizations (Eq. 2.1, Eq. 2.2, Eq. 2.3) by  $\chi^2$  minimization.

- **Bayesian Model Selection:** The Bayesian model selection technique used to compare two models ( $M_1$  and  $M_2$ ) is based on the calculation of the Bayesian odds ratio, given by:

$$O_{21} = \frac{P(M_2|D)}{P(M_1|D)}, \quad (3.6)$$

where  $P(M_2|D)$  is the posterior probability for  $M_2$  given data  $D$ , and similarly for  $P(M_1|D)$ . The posterior probability for a general model  $M$  is given by

$$P(M|D) = \frac{P(D|M)P(M)}{P(D)}, \quad (3.7)$$

where  $P(M)$  is the prior probability for the model  $M$ ,  $P(D)$  is the probability for the data  $D$  and  $P(D|M)$  is the marginal likelihood or Bayesian evidence for model  $M$  and is given by:

$$P(D|M) = \int P(D|M, \theta)P(\theta|M)d\theta, \quad (3.8)$$

where  $\theta$  is a vector of parameters, encapsulating the model  $M$ . If the prior probabilities of the two models are equal, the odds ratio can be written as:

$$B_{21} = \frac{\int P(D|M_2, \theta_2)P(\theta_2)d\theta_2}{\int P(D|M_1, \theta_1)P(\theta_1)d\theta_1}, \quad (3.9)$$


The quantity  $B_{21}$  in Eq. 3.9 is known as the Bayes factor. We shall compute this quantity in order to obtain a Bayesian estimate of the statistical significance. Note that unlike the frequentist and AIC/BIC based tests, the Bayesian test does not use the best-fit values of the parameters.

To assess the significance, a qualitative criterion based on Jeffreys' scale is used to interpret the odds ratio or Bayes factor [52, 53]. A value  $> 10$  represents strong evidence in favor of  $M_2$ , and a value  $> 100$  represents decisive evidence [52].

## 4 Analysis and Results

For each of the three models, we find the best-fit parameters by  $\chi^2$  minimization, where  $\chi^2$  is defined as follows:

$$\chi^2 = \sum_{i=1}^N \left( \frac{\delta\tau_i - \delta\tau_{model}}{\sigma_{\delta\tau_i}} \right)^2, \quad (4.1)$$



**Figure 1.** Plots showing the data points for the residual torque (obtained from P19) from the Eöt-Wash experiment, along with the best fits obtained for the three models using  $\chi^2$  minimization. The best-fit values for each of the three models are shown in Table 1. We note that for the best-fit Yukawa model, except for the first few points, the best fit is indistinguishable by eye compared to the corresponding Newtonian fit.

Parameter	Prior	Minimum	Maximum
$\alpha'$ (Eq. 2.1)	Uniform	-0.05	0.07
$\alpha'$ (Eq. 2.2)	Uniform	0	$1.0 \times 10^8$
$\alpha'$ (Eq. 2.3)	Uniform	0	0.01
$m'$ (Yukawa)	Uniform	0	500
$m'$ (Oscillating)	Uniform	60	70
$\phi$ (Oscillating with phase free)	Uniform	0	$2\pi$

**Table 2.** List of priors used to evaluate the Bayesian evidence for each of the three different models. The units for  $\alpha'$  and  $m'$  are in  $\text{fN} \cdot \text{m}$  and  $\text{mm}^{-1}$  respectively.

where  $\delta_{\tau_i}$  denotes the data for the residual torques (provided to us by L. Kazantidis) and  $\sigma_{\delta_{\tau_i}}$  indicates its associated error;  $\delta_{\tau_{\text{model}}}$  encapsulates the three model functions (defined in Eqs 2.1, 2.2, 2.3) used to fit the residual torque. We use the **Nelder-Mead** simplex algorithm [60] as coded in the **scipy** Python module to carry out the optimization. For the oscillating model we did two fits: with the phase fixed to  $3\pi/4$  and also keeping the phase ( $\phi$ ) as a free parameter. The best-fit parameters for each of these models are shown in Table 1 and the corresponding plots with all the three functions superposed on the data are shown in Fig. 1. We note that for the Yukawa model, we get very large values of  $\alpha'$  and  $m'$ , compared to P19. However, our  $\chi^2$  value of 82.1 for the Yukawa model is smaller than the value of 85.4 obtained in P19. Except for the first few points, the Yukawa best-fit is indistinguishable by eye compared to the Newtonian model. The best-fit for  $\phi$  in the oscillating parametrization we obtain is approximately the same as that in P19. We now present results from all the

different model comparison tests.

#### 4.1 Model Comparison using $AIC_c$ and BIC

We calculate the  $AIC_c$  and BIC values for each of the three models in Eqs. 2.1, 2.2, 2.3 using equations 3.1 and 3.2. Note that  $AIC_c$  and BIC are computed from the maximum value of the Gaussian likelihood using the best-fit parameters, obtained by  $\chi^2$  minimization. These values are listed in Table 3. We find that the difference in  $AIC_c$  and BIC between Yukawa model and the Newtonian one is negligible. So neither model is preferred.

However, when we compare the oscillating model with the offset Newtonian model when the phase is fixed at  $3\pi/4$ , we find that both  $\Delta AIC_c$  and  $\Delta BIC$  are  $> 10$ . Therefore, with the phase fixed, the oscillating model is decisively favored as compared to the Newtonian offset model. When the phase is a free parameter, only  $AIC_c$  still shows decisive evidence for the oscillating model, whereas BIC now shows only strong evidence. This is because BIC harshly penalizes models with additional free parameters compared to AIC [48].

#### 4.2 Model Comparison using WAIC

Using the equations 3.3, 3.4, and 3.5 we calculate the WAIC values for each of the three models with all the samples in each case. The posterior samples for each model have been obtained by nested sampling using the `Nestle` package in `Python`. The priors used to calculate the posterior samples are the same as for Bayesian model comparison, to be described in Sect. 4.3.

The values of WAIC obtained for the three models are listed in Table 3, along with the  $\Delta WAIC$  values with respect to the null hypothesis (Newtonian model). We find that the results broadly agree with  $AIC_c$  and BIC. The oscillating model has the lowest value of WAIC, implying that it has the highest predictive power among the three models. The values of WAIC for the Newtonian and Yukawa model are almost comparable. When the phase in the oscillating model is fixed at  $3\pi/4$ , the difference in WAIC between the Newtonian and oscillating hypothesis is greater than 10, implying that using this test also, the oscillating parametrization is decisively favored compared to the offset Newtonian model. When the phase is kept free, the difference in WAIC falls marginally below 10 and therefore, the oscillating model is strongly favored, but not decisively.

#### 4.3 Bayesian Model Comparison

Considering the Newtonian model to be the null hypothesis, we calculate the Bayes factor for the Yukawa model as well as the oscillating model in comparison with the constant Newtonian offset as the null hypothesis. We chose a Gaussian likelihood, using the data, model and experimental errors per data point. We chose uniform priors for all the parameters. A tabular summary of the priors used for each of the parametrizations can be found in Table 2. We note that for the oscillating model, we have only used positive definite priors for  $\alpha'$  as compared to the other two models. This is because the cosine term in the oscillating model can take both positive and negative values, so a negative value for the amplitude would be redundant. Also, as noticed in P19, there are multiple minima in  $(\alpha', m')$  parameter space. In this work, we wanted to hone in on the best-fit parameter space found by P19. Therefore, for  $m'$ , we also used a narrow range of priors near the maximum value. A tabular summary of the priors used for each of the parametrizations can be found in Table 2.

To calculate the Bayesian evidence for each of the hypotheses, We used the `Nestle`<sup>3</sup> package in `Python`, which uses the Nested sampling algorithm [61]. The Bayes factor was

---

<sup>3</sup><http://kylebarbary.com/nestle/>

	Newtonian	Yukawa	Oscillating ( $\phi = 3\pi/4$ )	Oscillating ( $\phi$ free)
<b>AIC<sub>c</sub> values</b>	-571.3	-572.6	-584.0	-582.1
$\Delta\text{AIC}_c = \text{AIC}_c(\text{Newt.}) - \text{AIC}_c$	-	1.2	12.7	10.5
<b>BIC values</b>	-568.9	-567.6	-579.2	-574.7
$\Delta\text{BIC} = \text{BIC}(\text{Newt.}) - \text{BIC}$	-	-1.2	10.3	5.9
<b>WAIC values</b>	-571.4	-574.3	-583.6	-579.4
$\Delta\text{WAIC} = \text{WAIC}(\text{Newt.}) - \text{WAIC}$	-	2.9	12.2	8.0
<b>Bayes Factor</b>	-	29	544	83

**Table 3.** Summary of results for the model comparison tests using Bayesian and information theoretic techniques. The Bayes factor (in this table) is the ratio of Bayesian evidence for the Yukawa or oscillating model to the same for Newtonian model.  $\Delta\text{AIC}_c$  (also BIC and WAIC) indicates the difference in  $\text{AIC}_c$ /BIC between the Newtonian model and the Yukawa/oscillating model. When the phase ( $\phi$ ) is fixed at  $3\pi/4$  similar to P19, all the tests used here, (viz.  $\text{AIC}_c$ , BIC, WAIC, and Bayes factor) decisively favor the oscillating model compared to the Newtonian model. When  $\phi$  is a free parameter, only  $\text{AIC}_c$  shows decisive evidence for the oscillating model; whereas BIC, WAIC, and Bayes factor now only show strong evidence for the same.

calculated using Eq. 3.9 for both the Yukawa and the oscillating models in comparison with the null hypothesis. The values for the Bayes factor are shown in Table 3. We find that the value for the Bayes factor is about 29 for the Yukawa model and 544 (83) for the oscillating model, depending on whether the phase is fixed to  $3\pi/4$  or is allowed to be a free parameter (cf. Table 3). According to the Jeffreys’ scale [52], Bayesian model comparison provides “strong” evidence for the Yukawa model and “decisive”/“strong” evidence in favor of the oscillating model, depending on whether the phase is fixed at  $3\pi/4$  or kept as a free parameter. Therefore, the results from Bayesian model comparison result concur with BIC/WAIC based information theory tests regarding the significance of the oscillating model as compared to the Newtonian model.

## 5 Conclusions

In a series of papers, Perivolaropoulos et al [1, 41] independently analyzed the residual torque data from the Eöt-Wash experiment [30–33], wherein the data was obtained after subtracting the torques due to a Newtonian potential. They argued that an oscillating parametrization with a fixed phase of  $3\pi/4$  provides a better fit to this data as compared to a constant term (equivalent to an offset Newtonian potential). If the oscillating fit is the true description of the data, one possible implication is that this could be a signature of a non-local modified theory of gravity [34–37].

To further investigate the viability and statistical significance of this claim, we independently analyze this same residual torque data, and carry out a fit to the same three functions used in Refs. [1, 41]: offset Newtonian, Yukawa and oscillating models. To discern the relative significance of each of the models, we carry out model comparison techniques using four different model comparison tests:  $\text{AIC}_c$  (a variant of AIC to account for small number of samples), BIC and WAIC based information theory test, and finally a Bayesian model comparison technique based on a calculation of the Bayes factor.

Our results from these model comparison analyses are summarized in Table 3. When we compare the Yukawa and Newtonian parametrizations, we find that the difference between the

models is marginal. The Bayesian test on the other hand prefers the Yukawa parametrization, where according to Jeffreys’ criterion, the support for the Yukawa model is strong, but not decisive. However, all the four tests decisively favor the oscillating model with the phase fixed to the same value as in P19. When the phase is kept as a free parameter, the oscillating model is now decisively favored using only the AIC test. Using all other tests, this model is strongly offered compared to the offset Newtonian model.

Therefore, we agree with P19 that our complementary suite of statistical tests also favor a spatially oscillating signal in the Eöt-Wash dataset, if the phase is fixed. But once we vary the phase, the significance reduces to “strong evidence” using three of the four tests used, with only AIC still showing decisive evidence. However, our tests cannot discern whether this spatial oscillation is a signature of modified gravity or other systematic effects which could explain this oscillating pattern [41]. Further analysis of data from other experiments is needed to distinguish between these possibilities.

To improve transparency in data analysis, we have made our analysis codes and datasets analyzed publicly accessible. These can be found at [https://github.com/aditikrishak/EotWash\\_analysis](https://github.com/aditikrishak/EotWash_analysis).

## Acknowledgments

Aditi Krishak is supported by a DST-INSPIRE fellowship. We are grateful to Lavrentios Kazantzidis for providing us the data from P19 for our analysis. We also acknowledge useful discussions with Sanjib Sharma about WAIC.

## References

- [1] L. Perivolaropoulos and L. Kazantzidis, *Hints of modified gravity in cosmos and in the lab?*, *International Journal of Modern Physics D* **28** (2019) 1942001 [[1904.09462](#)].
- [2] C. M. Will, *The Confrontation between General Relativity and Experiment*, *Living Reviews in Relativity* **17** (2014) 4 [[1403.7377](#)].
- [3] S. G. Turyshev, *Experimental Tests of General Relativity*, *Annual Review of Nuclear and Particle Science* **58** (2008) 207 [[0806.1731](#)].
- [4] I. H. Stairs, *Testing General Relativity with Pulsar Timing*, *Living Reviews in Relativity* **6** (2003) 5 [[astro-ph/0307536](#)].
- [5] N. Yunes, K. Yagi and F. Pretorius, *Theoretical physics implications of the binary black-hole mergers GW150914 and GW151226*, *Phys. Rev. D* **94** (2016) 084002 [[1603.08955](#)].
- [6] LIGO SCIENTIFIC, VIRGO collaboration, *Tests of General Relativity with the Binary Black Hole Signals from the LIGO-Virgo Catalog GWTC-1*, *Phys. Rev. D* **100** (2019) 104036 [[1903.04467](#)].
- [7] LIGO SCIENTIFIC, VIRGO collaboration, *Tests of General Relativity with GW170817*, *Phys. Rev. Lett.* **123** (2019) 011102 [[1811.00364](#)].
- [8] E. O. Kahya and S. Desai, *Constraints on frequency-dependent violations of Shapiro delay from GW150914*, *Physics Letters B* **756** (2016) 265 [[1602.04779](#)].
- [9] S. Boran, S. Desai, E. O. Kahya and R. P. Woodard, *GW170817 falsifies dark matter emulators*, *Phys. Rev. D* **97** (2018) 041501 [[1710.06168](#)].
- [10] C. M. Will, *Theory and Experiment in Gravitational Physics*. Mar., 1993.

- [11] T. Clifton, P. G. Ferreira, A. Padilla and C. Skordis, *Modified gravity and cosmology*, *Physics Reports* **513** (2012) 1 [[1106.2476](#)].
- [12] K. Koyama, *Cosmological tests of modified gravity*, *Reports on Progress in Physics* **79** (2016) 046902 [[1504.04623](#)].
- [13] A. Joyce, B. Jain, J. Khoury and M. Trodden, *Beyond the cosmological standard model*, *Physics Reports* **568** (2015) 1 [[1407.0059](#)].
- [14] R. Woodard, *Avoiding Dark Energy with 1/R Modifications of Gravity*, in *The Invisible Universe: Dark Matter and Dark Energy*, L. Papantonopoulos, ed., vol. 720 of *Lecture Notes in Physics*, Berlin Springer Verlag, p. 403, 2007, [[astro-ph/0601672](#)].
- [15] A. Joyce, L. Lombriser and F. Schmidt, *Dark Energy Versus Modified Gravity*, *Annual Review of Nuclear and Particle Science* **66** (2016) 95 [[1601.06133](#)].
- [16] J. Martin, C. Ringeval and V. Vennin, *Encyclopædia Inflationaris*, *Physics of the Dark Universe* **5** (2014) 75 [[1303.3787](#)].
- [17] S. Alexander and N. Yunes, *Chern-Simons modified general relativity*, *Physics Reports* **480** (2009) 1 [[0907.2562](#)].
- [18] N. J. Poplawski, *Cosmological consequences of gravity with spin and torsion*, *The Astronomical Review* **8** (2013) 108 [[1106.4859](#)].
- [19] K. Bamba, S. Capozziello, S. Nojiri and S. D. Odintsov, *Dark energy cosmology: the equivalent description via different theoretical models and cosmography tests*, *Astrophys. Space Sci.* **342** (2012) 155 [[1205.3421](#)].
- [20] I. Debono and G. F. Smoot, *General Relativity and Cosmology: Unsolved Questions and Future Directions*, *Universe* **2** (2016) 23 [[1609.09781](#)].
- [21] P. G. Ferreira, *Cosmological Tests of Gravity*, *Ann. Rev. Astron. Astrophys.* **57** (2019) 335 [[1902.10503](#)].
- [22] M. Ishak, *Testing general relativity in cosmology*, *Living Reviews in Relativity* **22** (2019) 1 [[1806.10122](#)].
- [23] S. Desai, *Limit on graviton mass from galaxy cluster Abell 1689*, *Physics Letters B* **778** (2018) 325 [[1708.06502](#)].
- [24] S. W. Hawking and R. Penrose, *The Singularities of Gravitational Collapse and Cosmology*, *Proceedings of the Royal Society of London Series A* **314** (1970) 529.
- [25] G. F. R. Ellis, *The arrow of time and the nature of spacetime*, *Studies in the History and Philosophy of Modern Physics* **44** (2013) 242 [[1302.7291](#)].
- [26] T. Padmanabhan, *Why does the universe expand?*, *General Relativity and Gravitation* **42** (2010) 2743 [[1001.3380](#)].
- [27] S. Carlip, *Quantum gravity: a progress report*, *Reports on Progress in Physics* **64** (2001) 885 [[gr-qc/0108040](#)].
- [28] A. Ashtekar, *Gravity and the quantum*, *New Journal of Physics* **7** (2005) 198 [[gr-qc/0410054](#)].
- [29] R. P. Woodard, *How far are we from the quantum theory of gravity?*, *Reports on Progress in Physics* **72** (2009) 126002 [[0907.4238](#)].
- [30] D. J. Kapner, T. S. Cook, E. G. Adelberger, J. H. Gundlach, B. R. Heckel, C. D. Hoyle et al., *Tests of the gravitational inverse-square law below the dark-energy length scale*, *Phys. Rev. Lett.* **98** (2007) 021101 [[hep-ph/0611184](#)].
- [31] C. D. Hoyle, D. J. Kapner, B. R. Heckel, E. G. Adelberger, J. H. Gundlach, U. Schmidt et al., *Submillimeter tests of the gravitational inverse-square law*, *Phys. Rev. D* **70** (2004) 042004 [[hep-ph/0405262](#)].

- [32] C. D. Hoyle, U. Schmidt, B. R. Heckel, E. G. Adelberger, J. H. Gundlach, D. J. Kapner et al., *Submillimeter tests of the gravitational inverse square law: a search for 'large' extra dimensions*, *Phys. Rev. Lett.* **86** (2001) 1418 [[hep-ph/0011014](#)].
- [33] D. J. Kapner, *A short-range test of Newton's gravitational inverse-square law*, Ph.D. thesis, University of Washington, Nov., 2005.
- [34] J. Edholm, A. S. Koshelev and A. Mazumdar, *Behavior of the Newtonian potential for ghost-free gravity and singularity free gravity*, *Phys. Rev. D.* **94** (2016) 104033 [[1604.01989](#)].
- [35] A. Kehagias and M. Maggiore, *Spherically symmetric static solutions in a nonlocal infrared modification of General Relativity*, *Journal of High Energy Physics* **2014** (2014) 29 [[1401.8289](#)].
- [36] V. P. Frolov and A. Zelnikov, *Head-on collision of ultrarelativistic particles in ghost-free theories of gravity*, *Phys. Rev. D.* **93** (2016) 064048 [[1509.03336](#)].
- [37] M. Maggiore and M. Mancarella, *Nonlocal gravity and dark energy*, *Phys. Rev. D.* **90** (2014) 023005 [[1402.0448](#)].
- [38] A. Accioly, B. L. Giacchini and I. L. Shapiro, *Low-energy effects in a higher-derivative gravity model with real and complex massive poles*, *Phys. Rev. D.* **96** (2017) 104004 [[1610.05260](#)].
- [39] B. L. Giacchini, *On the cancellation of Newtonian singularities in higher-derivative gravity*, *Physics Letters B* **766** (2017) 306 [[1609.05432](#)].
- [40] B. L. Giacchini and T. de Paula Netto, *Weak-field limit and regular solutions in polynomial higher-derivative gravities*, *European Physical Journal C* **79** (2019) 217 [[1806.05664](#)].
- [41] L. Perivolaropoulos, *Submillimeter spatial oscillations of Newton's constant: Theoretical models and laboratory tests*, *Phys. Rev.* **D95** (2017) 084050 [[1611.07293](#)].
- [42] S. Desai and D. W. Liu, *A search for evidence of solar rotation in Super-Kamiokande solar neutrino dataset*, *Astroparticle Physics* **82** (2016) 86 [[1604.06758](#)].
- [43] S. Desai, *Frequentist model comparison tests of sinusoidal variations in measurements of Newton's gravitational constant*, *EPL (Europhysics Letters)* **115** (2016) 20006 [[1607.03845](#)].
- [44] S. Kulkarni and S. Desai, *Classification of gamma-ray burst durations using robust model-comparison techniques*, *Astrophys. and Space Science* **362** (2017) 70 [[1612.08235](#)].
- [45] S. Ganguly and S. Desai, *Statistical significance of spectral lag transition in GRB 160625B*, *Astroparticle Physics* **94** (2017) 17 [[1706.01202](#)].
- [46] A. Krishak and S. Desai, *An independent assessment of significance of annual modulation in COSINE-100 data*, *Open J. Astrophys.* (2019) [[1907.07199](#)].
- [47] A. Krishak, A. Dantuluri and S. Desai, *Robust model comparison tests of DAMA/LIBRA annual modulation*, *JCAP* **2002** (2020) 007 [[1906.05726](#)].
- [48] S. Sharma, *Markov Chain Monte Carlo Methods for Bayesian Data Analysis in Astronomy*, *Ann. Rev. Astron. Astrophys.* **55** (2017) 213 [[1706.01629](#)].
- [49] A. R. Liddle, *How many cosmological parameters?*, *Mon. Not. R. Astron. Soc.* **351** (2004) L49 [[astro-ph/0401198](#)].
- [50] A. R. Liddle, *Information criteria for astrophysical model selection*, *Mon. Not. R. Astron. Soc.* **377** (2007) L74 [[astro-ph/0701113](#)].
- [51] K. Shi, Y. F. Huang and T. Lu, *A comprehensive comparison of cosmological models from the latest observational data*, *Mon. Not. R. Astron. Soc.* **426** (2012) 2452 [[1207.5875](#)].
- [52] R. Trotta, *Bayesian Methods in Cosmology*, *arXiv e-prints* (2017) arXiv:1701.01467 [[1701.01467](#)].

- [53] Ž. Ivezić, A. Connolly, J. Vanderplas and A. Gray, *Statistics, Data Mining and Machine Learning in Astronomy*. Princeton University Press, 2014.
- [54] M. Kerscher and J. Weller, *On model selection in cosmology*, *SciPost Physics Lecture Notes* **9** (2019) [[1901.07726](#)].
- [55] A. Gelman, J. Hwang and A. Vehtari, *Understanding predictive information criteria for bayesian models*, *Statistics and computing* **24** (2014) 997.
- [56] F. J. Fabozzi, S. M. Focardi, S. T. Rachev and B. G. Arshanapalli, *The basics of financial econometrics: Tools, concepts, and asset management applications*. John Wiley & Sons, 2014.
- [57] S. Watanabe, *Asymptotic equivalence of bayes cross validation and widely applicable information criterion in singular learning theory*, *Journal of Machine Learning Research* **11** (2010) 3571.
- [58] P. Ranalli, E. Koulouridis, I. Georgantopoulos, S. Fotopoulou, L. T. Hsu, M. Salvato et al., *The 2-10 keV unabsorbed luminosity function of AGN from the LSS, CDFS, and COSMOS surveys*, *Astron. & Astrophys.* **590** (2016) A80 [[1512.05563](#)].
- [59] P. Ranalli, D. Hobbs and L. Lindegren, *Astrometry and exoplanets in the Gaia era: a Bayesian approach to detection and parameter recovery*, *Astron. & Astrophys.* **614** (2018) A30 [[1704.02493](#)].
- [60] W. H. Press, S. A. Teukolsky, W. T. Vetterling and B. P. Flannery, *Numerical recipes in FORTRAN. The art of scientific computing*. 1992.
- [61] P. Mukherjee, D. Parkinson and A. R. Liddle, *A Nested Sampling Algorithm for Cosmological Model Selection*, *Astrophys. J. Lett.* **638** (2006) L51 [[astro-ph/0508461](#)].