

The topological period–index conjecture

Benjamin Antieau and Ben Williams

January 6, 2022

Abstract

We prove the topological analogue of the period–index conjecture in each dimension away from a small set of primes.¹

Résumé

Nous prouvons l’analogie topologique de la conjecture exposant et indice dans chaque dimension en dehors d’un petit ensemble de nombres premiers.

1 Introduction

We prove the following theorem, partly solving the so-called topological period–index conjecture of [AW14b]. For background, see §2.

Theorem A. *Let X be a finite $2d$ -dimensional CW complex² and let $\alpha \in \mathrm{Br}(X) = \mathrm{H}^3(X, \mathbb{Z})_{\mathrm{tors}}$ be a Brauer class. Setting $n = \mathrm{per}(\alpha)$, we have*

$$\mathrm{ind}(\alpha) | n^{d-1} \prod_{p|n} p^{v_p((d-1)!)},$$

where the product ranges over the prime divisors of n and where v_p denotes the p -adic valuation.

Away from $(d-1)!$, the result simplifies.

Corollary B. *If X is a finite $2d$ -dimensional CW complex, $\alpha \in \mathrm{Br}(X)$, and $\mathrm{per}(\alpha)$ is prime to $(d-1)!$, then $\mathrm{ind}(\alpha) | \mathrm{per}(\alpha)^{d-1}$.*

The corollary is an exact topological analogue, away from some small primes, of the well-known period–index conjecture for division algebras over function fields (see [CT06, Section 2.4]).

Conjecture (Function field period–index conjecture). *Let k be algebraically closed and let K be of transcendence degree d over k . If $\alpha \in \mathrm{Br}(K)$, then*

$$\mathrm{ind}(\alpha) | \mathrm{per}(\alpha)^{d-1}.$$

¹**Key Words;** Brauer groups, period–index problems, twisted K -theory. **Mathematics Subject Classification 2010:** 14F22, 19L50, 55R40.

²More precisely, what we require is for X to have the homotopy type of a retract of a finite CW complex and for $\mathrm{H}^i(X, A) = 0$ for $i > 2d$ and any abelian group A .

Tsen's theorem implies that $\mathrm{Br}(K) = 0$ when K has transcendence degree 1 over an algebraically closed field (see [GS17]). In [dJ04], de Jong proved the conjecture when $d = 2$ for Brauer classes of order relatively prime to the characteristic. Lieblich removed this restriction in [Lie08] thus establishing the $d = 2$ case in full generality. There are no other cases known of the conjecture. More precisely, the period–index conjecture for function fields is not known for a single function field of transcendence degree $d > 2$ over an algebraically closed field.

Nevertheless, work of de Jong and Starr [SdJ10] has reduced the conjecture above to the following special cases.

Conjecture (Global period–index conjecture). *Let k be an algebraically closed field and let X be a smooth projective k -scheme of dimension d . If $\alpha \in \mathrm{Br}(X) \subseteq \mathrm{Br}(k(X))$, then*

$$\mathrm{ind}(\alpha) \mid \mathrm{per}(\alpha)^{d-1}.$$

In other words, to prove the period–index conjecture for function fields, it is enough to prove it for unramified classes. We therefore view the period–index conjecture as a global problem. If $k = \mathbb{C}$, then the space of complex points $X(\mathbb{C})$ of a smooth projective \mathbb{C} -scheme of dimension d admits the structure of a $2d$ -dimensional CW complex. Thus, Corollary B provides evidence for this conjecture.

The topological period–index problem was introduced by the authors in [AW14] where weak lower bounds were given and where the $d \leq 2$ cases were solved. The $d = 3$ case of the topological period–index problem was settled in [AW14b], where it was proved moreover that the bound appearing in Theorem A is sharp in that case. The case of $d = 4$ was solved by Gu in [Gu19, Gu18], where the upper bound of Theorem A was found independently (in the $d = 4$ case) and where it is shown that the bound appearing in the theorem is sharp for square-free classes. The best possible bound for $d = 4$ and $n = \mathrm{per}(\alpha)$, as proved by Gu, is

$$\mathrm{ind}(\alpha) \mid \begin{cases} e_3(n)n^3 & \text{if } 4 \mid n \text{ and} \\ e_2(n)e_3(n)n^3 & \text{otherwise,} \end{cases}$$

where $e_p(n) = p$ if $p \mid n$ and 1 otherwise. In other words, there exist 8-dimensional finite CW complexes where these upper bounds on the index are achieved.

The fact that for $d = 3$ there exist 6-dimensional finite CW complexes with Brauer classes α having $\mathrm{per}(\alpha) = 2$ and $\mathrm{ind}(\alpha) = 8$ leads to the natural question in [AW14b] of whether the global period–index conjecture might be false at the prime 2 for threefolds over the complex numbers. However, recent work of Crowley–Grant [CG18] proves (a) that these topological examples can be found among closed orientable 6-manifolds but that (b) these examples cannot be found among closed 6-dimensional Spin^c -manifolds and hence they cannot be found among closed 6-manifolds of the form $X(\mathbb{C})$ for a smooth projective complex 3-fold X . The question of what happens for $d = 4$ is the subject of ongoing work of Crowley–Gu–Haesemeyer who prove in [CGH19] that for closed orientable 8-manifolds X one has $\mathrm{ind}(\alpha) \mid \mathrm{per}(\alpha)^3$ for $\alpha \in \mathrm{Br}(X)$ unless $\mathrm{per}(\alpha) \equiv 2 \pmod{4}$ in which case $\mathrm{ind}(\alpha) \mid 2\mathrm{per}(\alpha)^3$.

Note that Gu's bound $\mathrm{ind}(\alpha) \mid e_3(n)n^3$ if $4 \mid n$ is better than the bound $\mathrm{ind}(\alpha) \mid e_2(n)e_3(n)n^3$ arising from Theorem A. We do not further address in this paper the sharpness of the bounds in Theorem A except to make the following conjecture.

Conjecture C. *The bounds of Corollary B are the best possible. That is, for every $d \geq 1$ and every natural number n prime to $(d-1)!$, there exists a finite $2d$ -dimensional CW complex X and a Brauer class $\alpha \in \mathrm{Br}(X)$ such that $\mathrm{per}(\alpha) = n$ and $\mathrm{ind}(\alpha) = n^{d-1}$.*

The bounds in the period–index conjecture for function fields are known to be sharp, for example by Gabber's appendix to [CT02].

Acknowledgments. We would like to thank Diarmuid Crowley, Mark Grant, Xing Gu, and Christian Haesemeyer for their interest in this project and for many conversations over the years about the topological period–index problem. The first author was supported by NSF Grant DMS-1552766.

2 Background and strategy

We quickly review the period–index problem in three settings.

Period–index for fields. The period–index problem originated in the domain of division algebras over fields. Specifically, for a field K , we have the Brauer group $\mathrm{Br}(K) = H_{\mathrm{\acute{e}t}}^2(\mathrm{Spec} K, \mathbb{G}_m)$. This group is isomorphic to the set of isomorphism classes of finite dimensional division K -algebras with center exactly K . Given $\alpha \in \mathrm{Br}(K)$, we have two numbers: $\mathrm{per}(\alpha)$, which is the order of α in the torsion abelian group $\mathrm{Br}(K)$, and $\mathrm{ind}(\alpha)$ which is the unique positive integer such that $\mathrm{ind}(\alpha)^2 = \dim_K D$ where D is a division algebra with Brauer class $[D] = \alpha$. It is not hard to see that

$$\mathrm{per}(\alpha) | \mathrm{ind}(\alpha)$$

and Noether proved that these two numbers have the same prime divisors. It follows that there is some integer e_α such that $\mathrm{ind}(\alpha) | \mathrm{per}(\alpha)^{e_\alpha}$.

The period–index problem is to find for a fixed field K a number e such that

$$\mathrm{ind}(\alpha) | \mathrm{per}(\alpha)^e$$

for all $\alpha \in \mathrm{Br}(K)$ and, this done, to find the smallest such number. For example, the Albert–Brauer–Hasse–Noether theorem says that if K is a number field, then $\mathrm{ind}(\alpha) = \mathrm{per}(\alpha)$ so that $e = 1$ works (see [GS17, Remark 6.5.6]).

The period–index conjecture for function fields can be rephrased as saying that if K has transcendence degree d over an algebraically closed field, then $e = d - 1$ is the solution. For $d \geq 3$, it is not yet known that there is any e that works for all Brauer classes, but some results are known one prime at a time [Mat16].

Period–index for schemes. We introduced the period–index problem in other settings in [AW14]. For instance, if X is a quasicompact scheme, then the Brauer group

$$\mathrm{Br}(X) \subseteq H_{\mathrm{\acute{e}t}}^2(X, \mathbb{G}_m)_{\mathrm{tors}} \subseteq H_{\mathrm{\acute{e}t}}^2(X, \mathbb{G}_m)$$

of Azumaya algebras of Grothendieck [Gro68] is a torsion abelian group. Given $\alpha \in \mathrm{Br}(X)$, we again let $\mathrm{per}(\alpha)$ be the order of α in $\mathrm{Br}(X)$. We define

$$\mathrm{ind}(\alpha) = \gcd\{\deg(\mathcal{A}) : \mathcal{A} \text{ is an Azumaya algebra with } [\mathcal{A}] = \alpha\}.$$

In [AW14c], we showed that even on smooth schemes over the complex numbers, it is necessary to take the greatest common divisor to obtain a good theory. In this setting, we have $\mathrm{per}(\alpha) | \mathrm{ind}(\alpha)$ and the numbers have the same prime divisors by [AW15]. Thus, one can formulate the period–index problem for X .

Period–index for topological spaces. Finally, if X is a topological space, we have the Brauer group $\mathrm{Br}(X) \subseteq H^3(X, \mathbb{Z})_{\mathrm{tors}} \subseteq H^3(X, \mathbb{Z})$ of topological Azumaya algebras, also introduced in [Gro68]. We define $\mathrm{per}(\alpha)$ and $\mathrm{ind}(\alpha)$ as for schemes. Again, $\mathrm{per}(\alpha) | \mathrm{ind}(\alpha)$ and we proved that these have the same prime divisors if X is a finite CW complex in [AW14] and in general in [AW15].

The topological period–index problem of [AW14] asks the following. Given $d \geq 1$, find bounds e such that if X is a finite $2d$ -dimensional CW complex and $\alpha \in \text{Br}(X)$, then $\text{ind}(\alpha) | \text{per}(\alpha)^e$. We proposed $e = d - 1$ as a straw man in [AW14b], where we immediately proved that for $d = 3$ this fails in general when $2 | \text{per}(\alpha)$. Gu has proved this fails for $d = 4$ if 2 or 3 divides $\text{per}(\alpha)$. But, in these low-dimensional cases, these small primes are the only obstruction. We prove in Theorem A that this pattern continues in higher dimensions.

The topological results reveal a pattern which has not yet been discovered in algebra: a dependence on the prime divisors of the period and their relationship to d . This dependence comes, as we will see, from the ‘jumps’ in the cohomology of the Eilenberg–MacLane spaces $K(\mathbb{Z}/(n), 2)$.

Strategy. If X is a topological space and $\alpha \in H^3(X, \mathbb{Z})$, Atiyah and Segal constructed in [AS04] an α -twisted form of complex K -theory $KU(X)_\alpha$ and in [AS06] an α -twisted Atiyah–Hirzebruch spectral sequence

$$E_2^{s,t} = H^s(X, \mathbb{Z}(\tfrac{t}{2})) \Rightarrow KU^{s+t}(X)_\alpha,$$

where $\mathbb{Z}(\tfrac{t}{2}) \cong \mathbb{Z}$ if t is even and $\mathbb{Z}(\tfrac{t}{2}) = 0$ if t is odd. The differentials d_r^α have bidegree $(r, 1 - r)$. We proved in [AW14] that if X is a connected finite CW complex, then $\text{ind}(\alpha)$ generates the group

$$E_\infty^{0,0} \subseteq H^0(X, \mathbb{Z}) \cong \mathbb{Z}$$

of permanent cycles. Thus, to bound the index, one attempts to bound the orders of the differentials

$$d_{2r+1} : E_r^{0,0} \rightarrow E_r^{2r+1, -2r}.$$

There is a universal case to consider for all order m topological Brauer classes, namely the space $K(\mathbb{Z}/(m), 2)$ and a generator α of $H^3(K(\mathbb{Z}/(m), 2), \mathbb{Z}) \cong \mathbb{Z}/(m)$. By studying the orders of the differentials in this particular case, we prove Theorem A.

3 The cohomology of $K(\mathbb{Z}/(n), 2)$

We recall some results of Cartan [Car54] on the cohomology of Eilenberg–MacLane spaces in the special case of $K(\mathbb{Z}/(n), 2)$, which we will use in the next section to give upper bounds on the index of period n classes. We claim no originality in our presentation here, but we hope its inclusion will be useful to the reader.

Write

$$n = p_1^{r_1} \cdots p_k^{r_k},$$

and pick a generator u_i of the subgroup $\mathbb{Z}/(p_i^{r_i})$ of $\mathbb{Z}/(n)$ for $1 \leq i \leq k$. Let $v_i = p_i^{r_i-1} u_i$. Cartan [Car54, Théorème 1] gives a recipe for computing the integral homology (Pontryagin) ring of $K(\mathbb{Z}/(n), 2)$.

For each prime p and positive integer f , consider certain words in the 4 symbols:

$$\sigma, \quad \gamma_p, \quad \phi_p, \quad \psi_{pf}$$

The symbol ψ_{pf} , if it appears, is the last symbol in a word. The *height* of a word α is the total number of σ , ϕ_p and ψ_{pf} appearing. The *degree* of α is defined recursively by letting $\deg(\emptyset) = 0$ and

$$\deg(\sigma\alpha) = 1 + \deg(\alpha), \quad \deg(\gamma_p\alpha) = p \deg(\alpha), \quad \deg(\phi_p\alpha) = 2 + p \deg(\alpha), \quad \deg(\psi_{pf}) = 2$$

For each prime p , an *admissible* p -word α is a word on 3 symbols σ , γ_p , and ϕ_p such that α is non-empty, the first and last letters of α are σ or ϕ_p , and for each letter γ_p or ϕ_p , the number of letters σ appearing to the right in α is even. In addition to the admissible words, we will use the auxiliary words $\sigma^{h-1}\psi_{pf}$, of height h and degree $h + 1$.

Let $E(x, 2q-1)$ denote the exterior graded algebra over \mathbb{Z} with generator x of degree $2q-1$ endowed with the trivial dg-algebra structure $dx = 0$. Let $P(x, 2q)$ be the divided power polynomial algebra over \mathbb{Z} with generator x of degree $2q$ and given the trivial dg algebra structure with $dx = 0$. Cartan calls $E(x, 2q-1)$ and $P(x, 2q)$ *elementary complexes of the first type*. Define tensor dg algebras $E(x, 2q-1) \otimes_{\mathbb{Z}} P(y, 2q)$ by $dx = 0$ and $dy = hx$ for some integer h and $P(x, 2q) \otimes_{\mathbb{Z}} E(y, 2q+1)$ by $dx = 0$ and $dy = hx$ (the integer h is part of the data, even though it is not specified in the notation). These are the *elementary complexes of the second type*. The positive-degree homology groups of $E(x, 2q-1) \otimes_{\mathbb{Z}} P(x, 2q)$ are

$$H_{2q-1+2qk}(E(x, 2q-1) \otimes_{\mathbb{Z}} P(x, 2q)) = \mathbb{Z}/h \cdot x\gamma_k(y)$$

for $k \geq 0$ and 0 otherwise, where γ_k is the k th divided power operation. For $P(x, 2q) \otimes_{\mathbb{Z}} E(y, 2q+1)$, we get

$$H_{2qk}(P(x, 2q) \otimes_{\mathbb{Z}} E(y, 2q+1)) = \mathbb{Z}/hk \cdot \gamma_k(x)$$

for $k \geq 0$ and all other homology groups are 0.

The height-2 admissible or auxiliary p -words are

$$\sigma^2, \quad \sigma\gamma_p^k\phi_p, \quad \phi_p\gamma_p^k\phi_p, \quad \sigma\psi_{p^f}$$

for $k \geq 0$ and $f \geq 1$. These are of degrees 2 , $1 + 2p^k$, $2 + 2p^{k+1}$, and 3 , respectively. Below, the symbols u_i and v_i , are just formal indeterminates to keep track of generators for different dg algebras.

For each p_i , we define a dg algebra X_{p_i} as the following tensor product of elementary complexes of the second type

$$X_{p_i} = P(\sigma^2 u_i, 2) \otimes E(\sigma\psi_{p_i^{r_i}} u_i, 3) \otimes \bigotimes_{k=0}^{\infty} E(\sigma\gamma_{p_i}^{k+1} \phi_{p_i} v_i, 1 + 2p_i^{k+1}) \otimes P(\phi_{p_i} \gamma_{p_i}^k \phi_{p_i} v_i, 2 + 2p_i^{k+1}).$$

The differentials are

$$d(\sigma\psi_{p_i^{r_i}} u_i) = p_i^{r_i} \sigma^2 u_i \quad \text{and} \quad d(\phi_{p_i} \gamma_{p_i}^k \phi_{p_i} v_i) = p_i \sigma \gamma_{p_i}^{k+1} \phi_{p_i} v_i,$$

i.e., $h = p_i^{r_i}$ or $h = p_i$, respectively. (Here the r_i are the exponents appearing in the prime decomposition of n .) Let

$$X = X_{p_1} \otimes \cdots \otimes X_{p_k}.$$

Then, Cartan [Car54, Théorème 1] gives a surjection, which depends on the choice of the u_i ,

$$H_k(X) \rightarrow H_k(K(\mathbb{Z}/(n), 2), \mathbb{Z}),$$

and the kernel is described. This map induces for each i a surjection

$$H_k(X_{p_i}) \rightarrow H_k(K(\mathbb{Z}/(n), 2), \mathbb{Z})\{p_i\},$$

the p_i -primary part of homology. The largest possible torsion in $H_{2k}(X_{p_i})$ comes from the first term $P(\sigma^2 u_i, 2) \otimes E(\sigma\psi_{p_i^{r_i}} u_i, 3)$. Specifically, by using the Künneth theorem, the exponent of $H_{2k}(X_{p_i})$ is the same as the exponent of $H_{2k}(P(\sigma^2 u_i, 2) \otimes E(\sigma\psi_{p_i^{r_i}} u_i, 3))$, namely

$$p_i^{r_i} k.$$

It follows that the exponent of $H_{2k}(X)$ is at most nk . Write $k = k'a$, where a is largest integer dividing k and prime to m . Since $H_{2k}(K(\mathbb{Z}/(n), 2), \mathbb{Z})$ is n -primary torsion, we see that nk' kills all of

$H_{2k}(K(\mathbb{Z}/(n), 2), \mathbb{Z})$; in fact, the exponent is exactly nk' , as follows from Cartan's description of the kernel of $H_{2k}(X) \rightarrow H_{2k}(K(\mathbb{Z}/(n), 2), \mathbb{Z})$, but we will not need this fact.

Since we will be able to argue prime-by-prime in a moment, it is helpful to record the p_i -primary part of the exponent of $H_{2k}(K(\mathbb{Z}/(n), 2), \mathbb{Z})$ for a given prime p_i . If $\xi \in H_{2k}(K(\mathbb{Z}/(n), 2), \mathbb{Z})$ is an element of p_i -primary order, then the order of ξ divides $p_i^{r_i + v_{p_i}(k)}$.

4 Proof of the main theorem

If X is a topological space with finitely many connected components, then the topological Brauer group $\text{Br}(X)$ is a subgroup of $\text{Br}'(X) = H^3(X, \mathbb{Z})_{\text{tors}}$. Serre showed that if X is compact, then $\text{Br}(X) = \text{Br}'(X)$ (see [Gro68]). This will be the main setting of this paper. We want to make a general remark on a different version of the index to which our methods apply whether or not X is compact.

We may weaken the hypothesis on the finiteness of X in Theorem A at the expense of using the K -theoretic index rather than the topological index. Recall from [AW14] that the K -theoretic index $\text{ind}_K(\alpha)$ is defined as the (positive) generator of the image of the rank map $\text{KU}^0(X)_\alpha \rightarrow \mathbb{Z}$, where $\text{KU}^0(X)_\alpha$ denotes the α -twisted K -theory group. When X is finite-dimensional (and connected), one computes $\text{ind}_K(\alpha)$ as the generator of the group

$$E_\infty^{0,0} \subseteq E_2^{0,0} \cong \mathbb{Z}$$

by the convergence of the α -twisted Atiyah–Hirzebruch spectral sequence; hence the differentials $d_{2k+1}: E_{2k+1}^{0,0} \rightarrow E_{2k+1}^{2k+1, -2k}$ of the twisted Atiyah–Hirzebruch spectral sequence control the K -theoretic index. It is this crucial fact we exploit below.

In general, $\text{per}(\alpha) | \text{ind}_K(\alpha) | \text{ind}(\alpha)$ and if X is compact, then $\text{ind}_K(\alpha) = \text{ind}(\alpha)$. Thus, Theorem A follows from the following theorem.

Theorem 4.1. *Let X be a $2d$ -dimensional CW complex, and let $\alpha \in \text{Br}'(X)$ have period $m = p_1^{r_1} \cdots p_k^{r_k}$ where the p_i are distinct primes. Then,*

$$|\text{ind}_K(\alpha)| \prod_{i=1}^k p_i^{(d-1)r_i + v_{p_i}(2) + \cdots + v_{p_i}(d-1)} = m^{d-1} \prod_{i=1}^k p_i^{v_{p_i}((d-1)!)},$$

where v_{p_i} is the p_i -adic valuation.

Proof. According to the main result of [AW17], it is enough to prove the theorem when m is a prime power. So, suppose that $m = p_1^{r_1} = p^r$. Let β be a generator of $H^3(K(\mathbb{Z}/(p^r), 2), \mathbb{Z}) \cong \mathbb{Z}/(p^r)$. There is some map $\sigma: X \rightarrow K(\mathbb{Z}/(p^r), 2)$ such that $\sigma^*\beta = \alpha$. The twisted Atiyah–Hirzebruch spectral sequence is functorial, so we obtain—in particular—a map on the $(0, 0)$ -terms in each page:

$$E_j^{0,0}(K(\mathbb{Z}/(p^r), 2)) \rightarrow E_j^{0,0}(X).$$

An elementary induction argument shows that this map is a monomorphism on each page, and so by cellular approximation, $\text{ind}_K(\alpha)$ is bounded above by ind_K for the restriction of β to a $2d$ -skeleton of $K(\mathbb{Z}/(n), 2)$.

There is an isomorphism

$$\tilde{H}^{2j+1}(K(\mathbb{Z}/(p^r), 2), \mathbb{Z}) \cong \tilde{H}_{2j}(K(\mathbb{Z}/(p^r), 2), \mathbb{Z}).$$

We have seen in Section 3 that the latter group is $p^r p^{v_p(j)}$ -torsion. In the Atiyah–Hirzebruch spectral sequence for the Bockstein β on a $2d$ -skeleton of $K(\mathbb{Z}/(p^r), 2)$, only the differentials d_{2j+1}^β for $1 \leq j \leq d-1$

are possibly non-zero—in particular, the cohomology group in degree $2d$ of the skeleton, which may differ from that of $K(\mathbb{Z}/(p^r), 2)$, plays no part in the calculation. We are interested in the order of the image of

$$d_{2j+1}^\beta : E_{2j+1}^{0,0} \rightarrow E_{2j+1}^{2j+1,-2j},$$

where the latter group is a subquotient of $H^{2j+1}(K(\mathbb{Z}/(p^r), 2), \mathbb{Z})$, and hence the order of the image divides $p^{r+v_p(j)}$. This gives the result. \square

References

- [AS04] M. Atiyah and G. Segal, *Twisted K-theory*, Ukr. Mat. Visn. **1** (2004), no. 3, 287–330. MR2172633 [↑2](#)
- [AS06] ———, *Twisted K-theory and cohomology*, Inspired by S. S. Chern, 2006, pp. 5–43. MR2307274 [↑2](#)
- [AW14a] B. Antieau and B. Williams, *The period-index problem for twisted topological K-theory*, Geom. Topol. **18** (2014), no. 2, 1115–1148. MR3190609 [↑1](#), [2](#), [2](#), [2](#), [4](#)
- [AW14b] ———, *The topological period-index problem over 6-complexes*, J. Topol. **7** (2014), no. 3, 617–640. MR3252958 [↑1](#), [2](#)
- [AW14c] ———, *Unramified division algebras do not always contain Azumaya maximal orders*, Invent. Math. **197** (2014), no. 1, 47–56. MR3219514 [↑2](#)
- [AW15] ———, *The prime divisors of the period and index of a Brauer class*, J. Pure Appl. Algebra **219** (2015), no. 6, 2218–2224. MR3299728 [↑2](#), [2](#)
- [AW17] ———, *Prime decomposition for the index of a Brauer class*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **17** (2017), no. 1, 277–285. MR3676049 [↑4.1](#)
- [Car54] H. Cartan, *Détermination des algèbres $H_*(\pi, n; z)$* , Séminaire Henri Cartan **7** (1954), no. 1, 1–24. [↑3](#)
- [CG18] D. Crowley and M. Grant, *The topological period-index conjecture for $Spin^c$ 6-manifolds*, arXiv preprint arXiv:1802.01296 (2018). [↑1](#)
- [CGH19] D. Crowley, X. Gu, and C. Haesemeyer, *On the topological period-index problem over 8-manifolds*, arXiv preprint arXiv:1911.07206 (2019). [↑1](#)
- [CT02] J.-L. Colliot-Thélène, *Exposant et indice d’algèbres simples centrales non ramifiées*, Enseign. Math. (2) **48** (2002), no. 1-2, 127–146. With an appendix by Ofer Gabber. MR1923420 [↑1](#)
- [CT06] ———, *Algèbres simples centrales sur les corps de fonctions de deux variables (d’après A. J. de Jong)*, 2006, pp. Exp. No. 949, ix, 379–413. Séminaire Bourbaki. Vol. 2004/2005. MR2296425 [↑1](#)
- [dJ04] A. J. de Jong, *The period-index problem for the Brauer group of an algebraic surface*, Duke Math. J. **123** (2004), no. 1, 71–94. MR2060023 [↑1](#)
- [Gro68] A. Grothendieck, *Le groupe de Brauer. I. Algèbres d’Azumaya et interprétations diverses*, Dix exposés sur la cohomologie des schémas, 1968, pp. 46–66. MR244269 [↑2](#), [2](#), [4](#)
- [GS17] P. Gille and T. Szamuely, *Central simple algebras and Galois cohomology*, Cambridge Studies in Advanced Mathematics, vol. 165, Cambridge University Press, Cambridge, 2017. Second edition of [MR2266528]. MR3727161 [↑1](#), [2](#)
- [Gu18] X. Gu, *The topological period-index problem over 8-complexes, II*, arXiv preprint arXiv:1803.05100 (2018). [↑1](#)
- [Gu19] ———, *The topological period-index problem over 8-complexes, I*, Journal of Topology **12** (2019), no. 4, 1368–1395. [↑1](#)
- [Lie08] M. Lieblich, *Twisted sheaves and the period-index problem*, Compos. Math. **144** (2008), no. 1, 1–31. MR2388554 [↑1](#)
- [Mat16] E. Matzri, *Symbol length in the Brauer group of a field*, Trans. Amer. Math. Soc. **368** (2016), no. 1, 413–427. MR3413868 [↑2](#)
- [SdJ10] J. Starr and J. de Jong, *Almost proper GIT-stacks and discriminant avoidance*, Doc. Math. **15** (2010), 957–972. MR2745688 [↑1](#)