

# POTENTIAL THEORY FOR DRIFT DIFFUSION EQUATIONS WITH CRITICAL DIFFUSION AND APPLICATIONS TO THE DISSIPATIVE SQG EQUATION

QUOC-HUNG NGUYEN AND YANNICK SIRE

ABSTRACT. The purpose of this paper is to prove new regularity results for drift diffusion equations under minimal assumptions on the divergence free drift term. These results are presented as potential estimates and these estimates do not on the drift, allowing for applications to the dissipative SQG equation. We deal with quite general equations with non-homogeneous right hand sides.

## CONTENTS

1. Introduction	1
2. Proof of Theorem 1.1	6
3. Proof of Theorem 1.2	13
4. The case of bounded domains	21
5. Appendix	22
References	23

## 1. INTRODUCTION

The goal of this paper is to derive new regularity estimates for drift diffusion equations with a critical dissipative term in  $\mathbb{R}^d$ . Our goal is actually to derive finer estimates by means of Parabolic Riesz potentials independent of the vector-field in the drift term, except via its *a priori* regularity (see [Kis10] for a survey and references therein). This allows us to obtain new estimates for the critical dissipative SQG equation. Such equations has been introduced in [CMT94] as a toy model for the regularity of Navier-Stokes and several seminal works in the last years have led to major breakthrough on this topic [CV10, KNV07, KN09].

Let  $d \geq 2$  and consider the following drift-diffusion equation in  $\mathbb{R}^d$

$$(1.1) \quad \partial_t u + \mathbf{b} \cdot \nabla u + (-\Delta)^{1/2} u = f \quad \text{in } \mathbb{R} \times \mathbb{R}^d,$$

---

ShanghaiTech University, 393 Middle Huaxia Road, Pudong, Shanghai, 201210, China,  
Email address: [qhnguyen@shanghaitech.edu.cn](mailto:qhnguyen@shanghaitech.edu.cn).

Department of Mathematics, Johns Hopkins University, Baltimore, MD 21218, United States, Email address: [sire@math.jhu.edu](mailto:sire@math.jhu.edu).

for a given divergence free vector field  $\mathbf{b}$  and a datum  $f \in \mathcal{M}(\mathbb{R}^{d+1})$ , the set of Radon measures on  $\mathbb{R}^{d+1}$ . The operator  $(-\Delta)^{1/2}$  stands for the half-laplacian of symbol  $|\xi|$ , alternatively given by the integral formula

$$(-\Delta)^{1/2}u(x) = P.V. \int_{\mathbb{R}^d} \frac{u(x) - u(x+h)}{|h|^{d+1}} dh.$$

To introduce our results, we first define the following Parabolic Riesz potentials: let  $\mu \in \mathcal{M}(\mathbb{R}^{d+1})$ , be a locally finite Radon measure and consider  $Q_\rho(x, t) = B_\rho(x) \times (t - \rho, t)$ , a parabolic cylinder. Then we denote  $\mathbf{P}_s^r[\mu]$  the parabolic Riesz potential of  $\mu$  defined by

$$\mathbf{P}_s^r[\mu](x, t) = \int_0^r \frac{|\mu|(Q_\rho(x, t)) d\rho}{\rho^{d+1-s} \rho}.$$

One could consider two natural generalizations of (1.1). A first one would be

$$(1.2) \quad \partial_t u + \mathbf{b} \cdot \nabla u + (-\Delta)^\alpha u = f \quad \text{in } \mathbb{R} \times \mathbb{R}^d,$$

where  $\alpha \in (0, 1)$  and one has

$$(-\Delta)^\alpha u(x) = P.V. \int_{\mathbb{R}^d} \frac{u(x) - u(x+h)}{|h|^{d+2\alpha}} dh.$$

In view of the scaling, one gets a sub-critical problem if  $\alpha > 1/2$  and a supercritical for  $\alpha < 1/2$ . Very little is known on the super-critical problem when the drift  $\mathbf{b}$  is considered in this generality. See e.g. [CW99].

A second generalization would be to consider any integral operator  $\mathcal{L}u$  in place  $(-\Delta)^{1/2}u$  of the form

$$\mathcal{L}u(x) = P.V. \int_{\mathbb{R}^d} (u(x) - u(x+h)) K(h) dh,$$

where one has the bounds for  $c, C > 0$

$$\frac{c}{|h|^{d+1}} \leq K(h) \leq \frac{C}{|h|^{d+1}}.$$

In this case, because of our proof, the potential estimates of Theorem 1.3 hold using the same argument. The whole point would be to prove an analogue of Theorem 1.2. We plan to address this problem in a future work.

One can find in the literature (see e.g. [CI16]) a version of the critical dissipative SQG equation in a bounded domain with homogeneous Dirichlet conditions where the previously defined operator  $(-\Delta)^{1/2}$  is replaced by  $(-\Delta)_D^{1/2}$ . The latter is defined spectrally in the following way: let  $\{\varphi_k\}_{k=1}^\infty$  denote an orthonormal basis of  $L^2(\Omega)$  consisting of eigenfunctions of  $-\Delta$  in  $\Omega$  with homogeneous Dirichlet boundary conditions, associated to the eigenvalues  $\{\mu_k\}_{k=1}^\infty$ . Namely,  $0 < \mu_1 < \mu_2 \leq \mu_3 \leq \dots \leq \mu_k \rightarrow +\infty$ ,  $\int_{B_1} \varphi_j \varphi_k dx = \delta_{j,k}$  and

$$\begin{cases} -\Delta \varphi_k = \mu_k \varphi_k & \text{in } \Omega \\ \varphi_k = 0 & \text{on } \partial\Omega. \end{cases}$$

The operator  $(-\Delta)_D^{\frac{1}{2}}$  is defined for any  $u \in C_c^\infty(\Omega)$  by

$$(1.3) \quad (-\Delta)_D^{\frac{1}{2}}u = \sum_{k=1}^{\infty} \mu_k^{\frac{1}{2}} u_k \varphi_k,$$

where

$$u = \sum_{k=1}^{\infty} u_k \varphi_k, \quad \text{and} \quad u_k = \int_{\Omega} u \varphi_k \, dx.$$

This operator can be extended by density for  $u$  in the Hilbert space  $H = H_{00}^{1/2}(\Omega)$ , the Lions-Magenes space (see [CT10]). In this case, the drift-diffusion equation writes

$$(1.4) \quad \partial_t u + \mathbf{b} \cdot \nabla u + (-\Delta)_D^{1/2} u = f \quad \text{in } \mathbb{R} \times \Omega.$$

**Definition 1.** Let  $f \in L^q(\mathbb{R}^{n+1})$  for some  $q > 1$ . A function  $u \in L_{loc}^1((0, \infty) \times \mathbb{R}^d)$  is a (very) weak solution to (1.1) if for all test function  $\varphi \in C_0^\infty((0, \infty) \times \mathbb{R}^d)$  with

$$(1.5) \quad -\varphi_t - \mathbf{b} \cdot \nabla \varphi + (-\Delta)^{1/2} \varphi \in L^\infty,$$

we have

$$(1.6) \quad \int_0^\infty \int_{\mathbb{R}^d} u \left[ -\varphi_t - \mathbf{b} \cdot \nabla \varphi + (-\Delta)^{\frac{1}{2}} \varphi \right] = \int_0^\infty \int_{\mathbb{R}^d} \varphi f.$$

We will be considering depending on the statement a more natural (energy-type) notion of solutions, which are given by a standard Galerkin method, providing a (weak) solution  $u \in L^\infty(\mathbb{R}; L^2(\mathbb{R}^d)) \cap L^2(\mathbb{R}; H^{1/2}(\mathbb{R}^d))$ .

**Definition 2.** We define the contribution at infinity of a function  $v \in L_{loc}^1(\mathbb{R}^{d+1})$  by

$$\text{Tail}(v, x, t, r) = \inf_{\lambda \in \mathbb{R}} \int_{t-r}^t \int_{|x-y|>r} \frac{|v(y, s) - \lambda|}{|x-y|^{d+1}} \, dy \, ds.$$

**Main results.** Our first result describes a Hölder estimate for (1.1) under minimal assumptions on  $\mathbf{b}$  and  $f$ . In the following we *always* assume that  $\mathbf{b}$  is divergence free almost-everywhere in time  $t \in \mathbb{R}$ .

**Theorem 1.1.** Let  $\mathbf{b} : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}^d$  be such that

$$\sup_{(x,t) \in \mathbb{R}^{d+1}} \left| \int_{B_1(x)} \mathbf{b}(y, t) \, dy \right| \leq M_1 < \infty, \quad \|\mathbf{b}\|_{L^\infty(\mathbb{R}; BMO(\mathbb{R}^d))} \leq M_2 < \infty.$$

for some  $M_1, M_2 \geq 1$ . Let  $f \in (L^1 \cap L^q)(\mathbb{R}^{d+1})$  for  $q > d+1$ . Assume that  $u$  is a very weak solution to (1.1) satisfying additionally  $u \in L^\infty(\mathbb{R}; L^2(\mathbb{R}^d)) \cap L^2(\mathbb{R}; H^{1/2}(\mathbb{R}^d))$ . Then

$$(1.7) \quad \sup_{t,x} \|u\|_{C^\alpha(Q_{\delta_0/M_1^2}(x,t))} + \|u\|_{L_{t,x}^\infty} \lesssim_{M_2} \|f\|_{L_{t,x}^1} + \|f\|_{L_{t,x}^q},^1$$

for some  $\alpha \in (0, \frac{q-d-1}{q})$  and  $\delta_0$  only depending on  $M_2, q, d$ .

<sup>1</sup>We denote by  $L_{t,x}^p$  the space  $L_{t,x}^p(\mathbb{R} \times \mathbb{R}^d)$  for  $1 \leq p \leq \infty$  and  $Q_r(x, t) = (-r+t, t) \times B_r(x)$ . We use the expression  $A \lesssim_C B$  to mean that there exists a universal constant  $C$  depending only on  $c$  such that  $A \leq CB$ .

We now derive finer estimates, by means of parabolic potentials.

**Theorem 1.2.** *Let  $\omega$  be a bounded Radon measure in  $\mathbb{R}^{d+1}$  and  $\sigma \in BV(\mathbb{R}^d)$ . Assume that  $f = \omega(x, t) + \delta_{t=0} \otimes \sigma(x)$  and*

$$\|\mathbf{b}\|_{L^\infty} \leq M_3 < \infty.$$

Let  $\alpha \in (0, 1)$  be as in Theorem 1.1 and  $q_0 = 1 + \frac{1}{100d}$ . The following pointwise estimate holds

$$(1.8) \quad |u(x, t)| \lesssim_{M_3} \left( \int_{Q_{2r}(x, t)} |u|^{q_0} \right)^{1/q_0} + \mathbf{P}_1^r[\omega](x, t) + \mathbf{P}_2^r[\delta_{t=0} \otimes |\nabla \sigma|](x, t) + Tail(u, x, t, r).$$

Furthermore, for any  $\beta \in (0, \alpha)$  and  $|x_1 - x_2| + |t_1 - t_2| < r/2$

$$(1.9) \quad \frac{|u(x_1, t_1) - u(x_2, t_2)|}{(|x_1 - x_2| + |t_1 - t_2|)^\beta} \lesssim_{M_3} r^{-\beta} \left( \sum_{i=1,2} \left( \int_{Q_r(x_i, t_i)} |u|^{q_0} \right)^{1/q_0} + \mathbf{P}_{1-\beta}^r[\omega](x_i, t_i) + \mathbf{P}_{2-\beta}^r[\delta_{t=0} \otimes |\nabla \sigma|](x_i, t_i) + Tail(u, x_i, t_i, r) \right).$$

**Theorem 1.3.** *Assume that*

$$\sup_{(x, t) \in \mathbb{R}^{d+1}} \left| \int_{B_1(x)} \mathbf{b}(y, t) dy \right| \leq M_1 < \infty, \quad \|\mathbf{b}\|_{L^\infty(\mathbb{R}; BMO(\mathbb{R}^d))} \leq M_2 < \infty,$$

for some  $M_1, M_2 \geq 1$ . Let  $\alpha \in (0, 1)$  be as in Theorem 1.1 with  $q = 2d$  and  $q_0 = 1 + \frac{1}{100d}$ . The following pointwise estimate holds for any  $0 < r \leq 10^{-2}$

(1.10)

$$|u(x, t)| \lesssim_{M_1, M_2} |\log(r)|^{\frac{3d}{q_0}} \left( \int_{Q_r(x, t)} |u|^{q_0} \right)^{1/q_0} + \int_0^r \frac{|\omega|(Q_r(x, t))}{r^d |\log(r)|^{-3(d+1)}} \frac{dr}{r} + \int_0^r \frac{|\delta_{t=0} \otimes \nabla \sigma|(Q_r(x, t))}{r^d |\log(r)|^{-3(d+1)}} \frac{dr}{r} + Tail(u, x, t, r).$$

Furthermore, for any  $\beta \in (0, \alpha)$  and  $|x_1 - x_2| + |t_1 - t_2| < r/2 \leq 10^{-2}/2$ ,

(1.11)

$$(1.12) \quad \frac{|u(x_1, t_1) - u(x_2, t_2)|}{(|x_1 - x_2| + |t_1 - t_2|)^\beta} \lesssim_{M_1, M_2} \left( \sum_{i=1,2} r^{-\beta} |\log(r)|^{\beta_1} \left( \int_{Q_r(x_i, t_i)} |u|^{q_0} \right)^{1/q_0} + \int_0^{r_0} |\log(r)|^{\beta_2} \frac{|\omega|(Q_r(x_i, t_i))}{r^{d+\beta}} \frac{dr}{r} + \int_0^{r_0} |\log(r)|^{\beta_2} \frac{|\delta_{t=0} \otimes \nabla \sigma|(Q_r(x_i, t_i))}{r^{d-1+\beta}} \frac{dr}{r} + r^{-\beta} |\log(r)|^{\frac{3d}{q_0}} Tail(u, x_i, t_i, r) \right),$$

where  $\beta_1 = \frac{3\alpha}{\alpha-\beta}(\beta + \frac{d}{q_0}) + \frac{3d}{q_0}$  and  $\beta_2 = \frac{3\alpha(d+1+\beta)}{\alpha-\beta} + \frac{3d}{q_0}$ .

The previous theorems allow us to get results for the SQG equation, namely

$$(1.13) \quad \partial_t u + \mathcal{K}u \cdot \nabla u + (-\Delta)^{1/2} u = f \quad \text{in } \mathbb{R} \times \mathbb{R}^d,$$

where  $\mathcal{K}$  is a vectorial Calderon-Zygmund operator such that  $\operatorname{div} u = 0$ . In the case  $d = 2$ , one has  $\mathcal{K}u = \nabla^\perp(-\Delta)^{-1/2}u$  a rotation of the vectorial Riesz transform.

**Corollary 1.1.** *Let  $u$  be a weak solution of (1.13) with  $f \in \mathcal{M}(\mathbb{R}^{d+1})$ .*

- i:** *If  $f \in L^1 \cap L^q(\mathbb{R}^{d+1})$  for  $q > d + 1$ , then  $u$  is Hölder continuous in space and time*
- ii:** *If  $f(t, x) = \omega(t, x) + \delta_{t=0} \otimes \sigma(x)$  for  $\omega \in L^q, q > d + 1$  and  $\sigma \in L^\infty \cap BV(\mathbb{R}^d)$  satisfying*

$$\sup_{x,t} \int_0^1 \frac{|\delta_{t=0} \otimes \nabla \sigma|(Q_r(x, t))}{r^{d-1+\beta_0}} \frac{dr}{r} < \infty,$$

*for some  $\beta_0 > 0$ , then  $u$  is also Hölder continuous in space and time.*

*Proof.* For (1.13), one has that  $\mathbf{b} = \nabla^\perp(-\Delta)^{1/2}u$ . Since  $\mathbf{b}$  is divergence free, Proposition 2.1 gives that  $u(t) \in L^\infty(\mathbb{R}^d)$ ; hence by Calderon-Zygmund estimate one has  $\mathbf{b} \in L_t^\infty BMO_x$ . Applying Theorem (1.3) gives actually that  $u$  is  $C^\alpha$ . The proof is complete.  $\square$

**Space-time integrability.** In the following, we mention some consequences of the potential estimates. Indeed, it is well-known (see e.g. [DM11] and the references therein) that potential estimates lead to space-time estimates for solutions, using the mapping properties of the Riesz Parabolic potentials in Lorentz/Marcinkiewicz spaces for the solutions. The lemma below provides the general estimates. It is classical but we provide a proof for sake of completeness.

**Lemma 1.1.** *The following weak estimate holds for any positive Radon measure  $\mu$*

$$(1.14) \quad \mathcal{L}^{d+1}(\{(x, t) : \mathbf{P}_a[\mu](x, t) > \lambda\}) \leq \frac{C}{\lambda^{\frac{d+1}{d+1-a}}} \int_{\mathbb{R}^{d+1}} d\mu, \quad \forall \lambda > 0,$$

and

$$(1.15) \quad \|\mathbf{P}_a[\mu]\|_{L^{\frac{p(d+1)}{d+1-pa}}(\mathbb{R}^{d+1})} \leq C \|\mu\|_{L^p(\mathbb{R}^{d+1})} \quad \forall 1 < p < \frac{d+1}{a}.$$

*Proof.* Define the maximal function  $M\mu(x, t) := \sup_{r>0} \int_{Q_r(x,t)} d\mu$ . Standard estimates give

$$(1.16) \quad \|M\|_{\mathcal{M}_b \rightarrow w-L^1} + \|M\|_{L^q \rightarrow L^q} \leq C(d, \alpha, q) \quad \forall q > 1.$$

It is not hard to check ( see [Ngu14] ) that

$$\mathbf{P}_a[\mu](x, t) \leq C |\omega|(\mathbb{R}^{d+1})^{\frac{a}{d+1}} (M\mu(x, t))^{\frac{d+1-a}{d+1}},$$

and

$$\mathbf{P}_a[\mu](x, t) \leq C \|\mu\|_{L^p(\mathbb{R}^{d+1})}^{\frac{ap}{d+1}} (M\mu(x, t))^{\frac{d+1-ap}{d+1}}.$$

Combining these with (1.16) yields (1.14) and (1.15). The proof is complete.  $\square$

A direct application of this lemma along the lines of [DM11] gives straightforward results in "weak" spaces.

**Acknowledgements.** Quoc-Hung Nguyen's research was supported by the ShanghaiTech University startup funds.

## 2. PROOF OF THEOREM 1.1

This section is devoted to the proof of Theorem 1.1 following the approach by Caffarelli and Vasseur [CV10].

**Proposition 2.1.** *Under the assumptions of Theorem 1.1, there exists a unique weak solution of (1.1) such that*

$$(2.1) \quad \sup_{t \in \mathbb{R}} \int_{\mathbb{R}^d} u(t)^2 dx + \int_{\mathbb{R}^{d+1}} |(-\Delta)^{1/4} u|^2 dx dt \lesssim \|f\|_{L^{\frac{2(d+1)}{d+2}}}^2,$$

$$(2.2) \quad \|u\|_{L^{\frac{d+1}{d}, \infty}} \lesssim \|f\|_{L^1},$$

$$(2.3) \quad \|u\|_{L^\infty(\mathbb{R}^{d+1})} \lesssim_{d,q} \|f\|_{L^{\frac{2(q-d-1)}{2(q-d-1)+qd}}(\mathbb{R}^{d+1})} \|f\|_{L^q(\mathbb{R}^{d+1})}^{\frac{qd}{2(q-d-1)+qd}}.$$

*Proof.* A standard argument ensures existence and uniqueness of weak solutions for (1.1) satisfying (2.1). As proof of Lemma 3.1 in Section 3, we have (2.2). To prove (2.3), we only need to show that

$$(2.4) \quad \|u\|_{L^\infty([1, \infty) \times \mathbb{R}^d)} \leq C,$$

with  $\|f\|_{L^{\frac{2(d+1)}{d+2}}(\mathbb{R}^{d+1})} + \|f\|_{L^q(\mathbb{R}^{d+1})} \leq 1$ .

For this purpose we use a De Giorgi iteration. Set  $T_k = 1 - 2^{-k}$  and  $\lambda_k = \lambda(1 - 2^{-k})$ ,  $u_k = (u - \lambda_k)_+$  and the level set of energy:

$$U_k = \sup_{t \geq T_k} \int_{\mathbb{R}^d} u_k(t)^2 dx + \int_{T_k}^{\infty} \int_{\mathbb{R}^d} |(-\Delta)^{1/4} u_k|^2 dx dt$$

We have

$$U_k \leq 2 \int_{\mathbb{R}^d} u_k(s)^2 dx + 2 \int_{T_{k-1}}^{\infty} \int_{\mathbb{R}^d} u_k |f| dx dt.$$

Taking the mean value in  $s$  on  $[T_{k-1}, T_k]$  we find,

$$U_k \leq 2^{k+1} \|u_k\|_{L^2([T_{k-1}, \infty) \times \mathbb{R}^d)}^2 + 2 \|u_k\|_{L^{q'}([T_{k-1}, \infty) \times \mathbb{R}^d)} \|f\|_{L^q(\mathbb{R}^{d+1})},$$

where  $q' = \frac{q}{q-1} < \frac{d+1}{d}$ .

It is not hard to check (see [CV10]) that

$$\begin{aligned} \|u_k\|_{L^2([T_{k-1}, \infty) \times \mathbb{R}^d)}^2 &\leq \left(\frac{2^k}{\lambda}\right)^{\frac{2}{d}} \|u_{k-1}\|_{L^{\frac{2(d+1)}{d}}([T_{k-1}, \infty) \times \mathbb{R}^d)}^{\frac{2(d+1)}{d}}, \\ \|u_k\|_{L^{q'}([T_{k-1}, \infty) \times \mathbb{R}^d)} &\leq \left(\frac{2^k}{\lambda}\right)^{\frac{2(d+1)}{dq'}-1} \|u_{k-1}\|_{L^{\frac{2(d+1)}{d}}([T_{k-1}, \infty) \times \mathbb{R}^d)}^{\frac{2(d+1)}{d}}, \\ \|u_{k-1}\|_{L^{\frac{2(d+1)}{d}}([T_{k-1}, \infty) \times \mathbb{R}^d)} &\leq C U_{k-1}^{1/2}. \end{aligned}$$

Then, we get

$$U_k \leq C \frac{2^{k(1+2/d)}}{\lambda^{2/d}} U_{k-1}^{\frac{d+1}{d}} + C \left(\frac{2^k}{\lambda}\right)^{\frac{2(d+1)}{dq'}-1} U_{k-1}^{\frac{d+1}{d}} \quad \forall k \geq 1.$$

Thanks to (2.1), one has  $U_0 \leq C$ . Since  $\frac{d+1}{d}$  and  $\frac{d+1}{dq} > 1$ , so for  $\lambda > 1$  large enough, we get  $\lim_{k \rightarrow \infty} U_k = 0$ . This means  $u \leq \lambda$  almost everywhere in  $[1, \infty) \times \mathbb{R}^d$ , similarly, we also obtain  $u \geq -\lambda$ . So, we get (2.4). The proof is complete.  $\square$

**2.1. Local energy inequality.** A technical tool, now wildly used in the present framework, is the harmonic extension of an  $\dot{H}^{1/2}(\mathbb{R}^d)$  function. If  $u$  is such a function on  $\mathbb{R}^d$ , we denote by  $u^*$  its extension to the half-space  $\mathbb{R}_+^{d+1}$ .

**Proposition 2.2.** *Let  $t_1, t_2$  be such that  $t_1 < t_2$  and denote  $B_r^* = B_r \times [0, r)$ . Let  $u \in L^\infty(t_1, t_2; L^2(\mathbb{R}^d))$  with  $(-\Delta)^{1/2}u \in L^2((t_1, t_2) \times \mathbb{R}^d)$  be a weak solution of (1.1). For every  $t_1 \leq t \leq t_2$  and cut-off function  $\varphi$  such that the restriction of  $\eta[u^*]_+$  on  $B_2^*$  is compactly supported in  $B_2 \times (-2, 2)$ , it holds*

(2.5)

$$\begin{aligned} & \int_{B_2} \eta^2 [u]_+^2(t_2) dx + \int_{t_1}^{t_2} \int_{B_2^*} |\nabla(\eta[u^*]_+)|^2 dx dz dt \\ & \lesssim \int_{B_2} \eta^2 [u]_+^2(t_1) dx + \int_{t_1}^{t_2} \int_{B_2^*} |\nabla \eta|^2 [u^*]_+^2 dx dz dt \\ & + \|\mathbf{b}\|_{L^\infty(t_1, t_2; L^{2d}(B_2))}^2 \int_{t_1}^{t_2} \int_{\mathbb{R}^d} |\nabla \eta|^2 [u]_+^2 dx ds + \int_{t_1}^{t_2} \int_{B_2} \eta^2 [u]_+ |g| dx ds. \end{aligned}$$

*Proof.* We have for every  $t_1 < t < t_2$ :

$$\begin{aligned} 0 &= \int_{B_2^*} \eta^2 [u^*]_+ \Delta u^* = - \int_{B_2^*} \nabla(\eta^2 [u^*]_+) \nabla u^* + \int_{B_2} \eta^2 [u]_+ (-\Delta)^{1/2} u \\ &= - \int_{B_2^*} |\nabla(\eta[u^*]_+)|^2 + \int_{B_2^*} |\nabla \eta|^2 [u^*]_+^2 + \int_{B_2} \eta^2 [u]_+ (-\Delta)^{1/2} u. \end{aligned}$$

Using  $\eta^2 [u^*]_+$  as test function in (1.1), one has

$$\frac{d}{dt} \int_{B_2} \eta^2 \frac{[u]_+^2}{2} dx - \int_{B_2} \nabla \eta^2 \cdot \mathbf{b} \frac{[u]_+^2}{2} dx + \int_{B_2} \eta^2 [u]_+ (-\Delta)^{1/2} u = \int_{B_2} \eta^2 [u]_+ f.$$

This implies

$$\begin{aligned} & \int_{B_2} \eta^2 \frac{[u]_+^2(t_2)}{2} dx + \int_{t_1}^{t_2} \int_{B_2^*} |\nabla(\eta[u^*]_+)|^2 dx dz dt = \\ & \int_{B_2} \eta^2 \frac{[u]_+^2(t_1)}{2} dx + \int_{t_1}^{t_2} \int_{B_2^*} |\nabla \eta|^2 [u^*]_+^2 dx dz dt \\ & + \int_{t_1}^{t_2} \int_{B_2} \nabla \eta^2 \cdot \mathbf{b} \frac{[u]_+^2}{2} dx dt + \int_{t_1}^{t_2} \int_{B_2} \eta^2 [u]_+ f dx dt. \end{aligned}$$

By Sobolev inequality,

$$\begin{aligned} \|\eta[u]_+\|_{L^{\frac{2d}{d-1}}(B_2)}^2 &\leq C \|\mathbf{1}_{B_2} \eta[u]_+\|_{H^{1/2}(\mathbb{R}^d)}^2 = C \int_0^\infty |\nabla(\mathbf{1}_{B_2} \eta[u]_+)^*|^2 dx dz \\ &\leq C \int_0^\infty |\nabla(\mathbf{1}_{B_2^*} \eta[u^*]_+)|^2 dx dz = C \int_{B_2^*} |\nabla(\eta[u^*]_+)|^2 dx dz. \end{aligned}$$

Thus, using Holder's inequality we get

$$\begin{aligned} \left| \int_{t_1}^{t_2} \int_{B_2} \nabla \eta^2 \cdot \mathbf{b} \frac{[u]_+^2}{2} dx dt \right| &\leq \varepsilon \int_{t_1}^{t_2} \|\eta[u]_+\|_{L^{\frac{2d}{d-1}}}^2 dt + \\ &C_\varepsilon \|\mathbf{b}\|_{L^\infty(t_1, t_2; L^{2d}(B_2))}^2 \int_{t_1}^{t_2} \int_{\mathbb{R}^d} |\nabla \eta|^2 [u]_+^2 dx ds, \end{aligned}$$

which implies (2.5).  $\square$

The following lemma describes the oscillation reduction.

**Lemma 2.1.** *Let  $q > d + 1$ . Assume that*

$$(2.6) \quad \|\mathbf{b}\|_{L^\infty(t_1, t_2; L^{2d}(B_2))} \leq M_0.$$

*There exists  $\varepsilon_0 > 0$  (depending only on  $d, M_0$ ) and  $\lambda > 0$  (depending only on  $d$ ) such that for every  $u$  solution to (1.1), there holds if*

$$u^* \leq 2 \text{ in } [-4, 0] \times B_4^*,$$

and

$$(2.7) \quad \int_{-4}^0 \int_{B_4^*} (u^*)_+^2 dx dz ds + \int_{-4}^0 \int_{B_4} (u)_+^2 dx ds + \|f\|_{L^q([-4, 0] \times B_4)}^2 \leq \varepsilon_0,$$

then:

$$(2.8) \quad (u)_+ \leq 2 - \lambda \text{ on } [-1, 0] \times B_1.$$

*Proof.* We follow proof of [CV10, Lemma 6]. First, let  $E_1$  be solution of  $\Delta E_1 = 0$  in  $B_4^*$ ,  $E_1 = 2$  in  $\partial B_4^* \setminus \{z = 0\}$ , and  $E_1 = 0$  in  $\partial B_4^* \cap \{z = 0\}$ . Let  $E_2$  be solution of  $\Delta E_2(s, z) = 0$  in  $[0, \infty) \times [0, 1]$ ,  $E_2(0, z) = 2$  in  $[0, 1]$ , and  $E_2(s, 0) = E_2(s, 1) = 0$  in  $(0, \infty)$ . There exists  $\lambda \in (0, 1/2)$  and  $c_1 > 0$  such that

$$(2.9) \quad E_1(x, z) \leq 2 - 4\lambda \text{ in } B_2^*,$$

$$(2.10) \quad |E_2(s, z)| \leq c_1 e^{-s/2} \text{ in } [0, \infty) \times [0, 1].$$

We set  $u_k = (u - C_k)_+$ ,  $u_k^* = (u^* - C_k)_+$ , with  $C_k = 2 - \lambda(1 + 2^{-k})$ . Let  $\eta_k$  be a cut off function in  $x$  only such that:  $\mathbf{1}_{B_{1+2^{-k-1/2}}} \leq \eta_k \leq \mathbf{1}_{B_{1+2^{-k}}}$  and  $|\nabla \eta_k| \leq C2^k$ . We denote

$$A_k = \int_{-1-2^{-k}}^0 \int_0^{\delta^k} \int_{\mathbb{R}^d} |\nabla(\eta_k u_k^*)|^2 dx dz dt + \sup_{t \in [-1-2^k, 0]} \int_{\mathbb{R}^d} (\eta_k u_k)^2 dx.$$

As in the proof of [CV10, Lemma 7], there exist  $0 < \delta < 1$  and  $M > 1$  such that for every  $k \geq 2d$ :

$$2dc_1 e^{-\frac{2^{-k}}{4(\sqrt{2}+1)\delta^k}} \leq \lambda 2^{-k-2}, \quad \frac{M^{-k/2}}{\delta^{\frac{d(k+1)}{2}}} \|P\|_{L^2(\mathbb{R}^d)} \leq \lambda 2^{-k-2}, \quad M^{-k} \geq c_2^k M^{-\frac{d+1}{dq'}(k-3)}.$$

where  $q' = \frac{q}{q-1} < \frac{d+1}{d}$  and we denote by  $P$  the Poisson kernel, which satisfies  $P(1)(x) = P(x, 1)$  for any  $x \in \mathbb{R}^d$  and  $c_2$  will be chosen in later.

We want to prove that for every  $k \geq 0$ :

$$(2.11) \quad A_k \leq M^{-k},$$



$$(2.12) \quad \eta_k u_k^* = 0 \text{ for } \delta^k \leq z \leq \min\{2, \delta^{k-1}\}.$$

We use that energy inequality (2.5) with cut-off function  $\eta_k(x)\psi(z)$  where  $\psi$  is a fixed cut-off function in  $z$  only. It follows from (2.5) that for any  $t_1 \in [-4, -2]$

$$\begin{aligned} A_k &\leq \sup_{t_2 \in [-1-2^{-k}, 0]} \left( \int_{B_2} (\eta_k \psi)^2 [u]_+^2(t_2) dx + \int_{t_1}^{t_2} \int_{B_2^*} |\nabla(\eta_k \psi [u^*]_+)|^2 dx dz dt \right) \\ &\leq \int_{B_2} (\eta_k \psi)^2 [u]_+^2(t_1) dx + C \int_{-4}^0 \int_{B_2^*} |\nabla(\eta_k \psi)|^2 [u^*]_+^2 dx dz dt \\ &\quad + CM_0 \int_{-4}^0 \int_{\mathbb{R}^d} |\nabla(\eta_k \psi)|^2 [u]_+^2 dx ds + C \left( \int_{-4}^0 \int_{B_2} \eta_k^2 [u]_+^2 dx ds \right)^{1/2} \|f\|_{L^2([-4, 0] \times B_4)}. \end{aligned}$$

Thus, using the fact that  $|\nabla \eta_k| \leq C2^{24d}$  for  $0 \leq k \leq 12d$ , we get

$$A_k \leq C(1 + M_0)2^{24d} \varepsilon_0.$$

Choosing  $\varepsilon_0 \in (0, 1/100)$  such that  $C(1 + M_0)2^{24d} \varepsilon_0 \leq M^{-12d}$ , this gives that (2.11) is verified for  $0 \leq k \leq 12d$ . Moreover, following the proof of [CV10, Lemma 6] we also obtain (2.12) at  $k = 0$  for  $\varepsilon_0 > 0$  small enough. Now assume that (2.11) and (2.12) are true for any  $k \leq m$ . We will prove that (2.11) and (2.12) are true at  $k = m + 1$ . As in the proof of [CV10, Lemma 6], we have that (2.12) is true  $k = m + 1$ . Thus, it is enough to show that

$$(2.13) \quad A_m \leq c_2^m (A_{m-2}^{\frac{d+1}{d}} + A_{m-2}^{\frac{d+1}{dq'}}).$$

Indeed, repeating proof of [CV10, (15) in Lemma 6] we have

$$\begin{aligned} A_{m+1} &\leq \frac{C2^{2(m+1)(1+1/d)}}{\lambda^{2/d}} A_{m-2}^{1+\frac{1}{d}} + C \int_{1-2^{-m-1}}^0 \int_{\mathbb{R}^d} \eta_m^2 u_{m+1} |f| dx ds, \\ \|\eta_{m-1} u_m\|_{L^{\frac{2(d+1)}{d}}}^2 &\leq CA_{m-2}. \end{aligned}$$

Using Holder's inequality

$$\begin{aligned} \int_{1-2^{-m-1}}^0 \int_{\mathbb{R}^d} \eta_m^2 u_{m+1} |f| dx ds &\leq \|f\|_{L^q([-4, 0] \times B_4)} \left( \int_{1-2^{-m-1}}^0 \int_{\mathbb{R}^d} (\eta_m u_{m+1})^{q'} dx ds \right)^{\frac{1}{q'}} \\ &\leq C\varepsilon_0^{1/2} \left( \frac{2^{m+1}}{\lambda} \right)^{\frac{2(d+1)}{dq'} - 1} A_{m-2}^{\frac{d+1}{dq'}}, \end{aligned}$$

we have used the fact that  $(\eta_m u_{m+1})^{q'} \leq C \left( \frac{2^{m+1}}{\lambda} \right)^{\frac{2(d+1)}{d} - q'} (\eta_{m-1} u_m)^{\frac{2(d+1)}{d}}$ . Therefore, we get (2.13) for some  $c_2 > 0$ .  $\square$

In the following, we denote  $Q_r = B_r \times [-r, 0]$  and  $Q^* = B_r^* \times [-r, 0]$ . As in [CV10, Lemma 8], we also have

**Lemma 2.2.** *Let  $q > d + 1$ . Let*

$$(2.14) \quad \|\mathbf{b}\|_{L^\infty(t_1, t_2; L^{2d}(B_2))} \leq M_0.$$

*Assume that  $u^* \leq 2$  in  $Q_4^*$  and*

$$\mathcal{L}^{d+2}(\{(x, z, t) \in Q_4^* : u^*(x, z, t) \leq 0\}) \geq \mathcal{L}^{d+2}(Q_4^*)/2.$$

for every  $\varepsilon_1 > 0$ , there exists  $\varepsilon_0 = \varepsilon_0(d, M_0) > 0$  and  $\delta_1 = \delta_1(d, M_0, \varepsilon_1)$  such that if  $\|f\|_{L^q([-4,0] \times B_4)}^2 \leq \varepsilon_0$  and

$$\mathcal{L}^{d+2}(\{(x, z, t) \in Q_4^* : 0 < u^*(x, s, t) < 1\}) \leq \delta_1$$

then

$$(2.15) \quad \int_{Q_1} (u-1)_+^2 dxdt + \int_{Q_1^*} (u^*-1)_+^2 dx dz dt \leq C(d, M_0) \sqrt{\varepsilon_1}$$

The previous two lemmas give the usual dichotomy.

**Theorem 2.1.** *Let  $q > d + 1$ . Let*

$$(2.16) \quad \|\mathbf{b}\|_{L^\infty(t_1, t_2; L^{2d}(B_2))} \leq M_0.$$

There exist  $\lambda^* = \lambda^*(d, M_0) > 0$  and  $\varepsilon_0 = \varepsilon_0(d, M_0)$  such that for every weak solution  $u$  of (1.1) with

$$(2.17) \quad \|f\|_{L^q([-4,0] \times B_4)}^2 \leq \varepsilon_0,$$

if

$$|u^*| \leq 2 \quad \text{in } Q_1^*,$$

then

$$(2.18) \quad \text{osc}_{Q_{1/4}^*} u^* \leq 4 - \lambda^*.$$

*Proof.* Let  $\varepsilon_0$  be as in Lemma 2.1. Then,  $\varepsilon_1 > 0$  satisfies  $4C\sqrt{\varepsilon_1} = \varepsilon_0$ , where  $C$  is the constant in Lemma 2.2. Let  $\delta_1$  be in Lemma 2.2 associated to  $\varepsilon_1$ . It is enough to show that if

$$\mathcal{L}^{d+2}(\{(x, z, t) \in Q_1^* : u^*(x, z, t) \leq 0\}) \geq \mathcal{L}^{d+2}(Q_1^*)/2,$$

then

$$(2.19) \quad u^* \leq 2 - \lambda^*/2 \quad \text{in } Q_{1/16}^*.$$

For every  $k \in \mathbb{N}$ ,  $k \leq K = [1/\delta_1 + 1]$ , we define

$$\bar{u}_k = 2(\bar{u}_{k-1} - 1) \quad \text{with } \bar{u}_0 = 0.$$

So, we have  $\bar{u}_k = 2^k(u - 2) + 2$ . Note that for every  $k$ ,  $\bar{u}_k$  is a solution of (1.1) with data  $2^k f$  and

$$\mathcal{L}^{d+2}(\{(x, z, t) \in Q_1^* : \bar{u}_k^*(x, z, t) \leq 0\}) \geq \mathcal{L}^{d+2}(Q_1^*)/2.$$

Assume that for every  $k \leq K$ ,  $\mathcal{L}^{d+2}(\{(x, z, t) \in Q_1^* : 0 < \bar{u}_k^*(x, z, t) < 1\}) \geq \delta_1$ . Then for every  $k \leq K$

$$\mathcal{L}^{d+2}(\{Q_1^* : \bar{u}_k^* < 0\}) = \mathcal{L}^{d+2}(\{Q_1^* : \bar{u}_{k-1}^* < 1\}) \geq \delta_1 + \mathcal{L}^{d+2}(\{Q_1^* : \bar{u}_{k-1}^* < 0\})$$

Thus,  $\mathcal{L}^{d+2}(\{Q_1^* : \bar{u}_K^* < 0\}) \geq 1$  and  $\bar{u}_K^* \leq 0$  almost everywhere, which means  $u^* \leq 2 - 2^{-K}$ . Otherwise, there exists  $0 \leq k_0 \leq K$  such that  $\mathcal{L}^{d+2}(\{(x, z, t) \in Q_1^* : 0 < \bar{u}_{k_0}^*(x, z, t) < 1\}) \leq \delta_1$ . Applying Lemma 2.2 for  $\bar{u}_{k_0+1}$  with data  $2^{k_0+1} f$ , we get  $\bar{u}_{k_0+1} \leq 2 - \lambda$  for  $\varepsilon_0$  small enough. Thus,

$$u \leq 2 - 2^{-k_0-1} \lambda \leq 2 - 2^{-K} \lambda \quad \text{in } Q_{1/8}.$$

Then, as in the proof of [CV10, Proposition 9], we get (2.19).  $\square$

*Proof of Theorem 1.1.* It is enough to show that

$$(2.20) \quad \sup_{|t_1-t_2|+|x_1-x_2|<\rho_0} \frac{|u(t_1, x_1) - u(t_2, x_2)|}{(|t_1 - t_2| + |x_1 - x_2|)^\alpha} \leq C,$$

with

$$\|f\|_{L^1(\mathbb{R}^{d+1})} + \|f\|_{L^q(\mathbb{R}^{d+1})} \leq \varepsilon_0.$$

and  $\varepsilon_0 > 0$  small enough and  $\rho_0 = 10^{-21} M_1^{-2}$  and for some  $\delta \in (0, 1)$ . By (2.1) and (2.3), one has

$$(2.21) \quad \sup_{t \in \mathbb{R}} \int_{\mathbb{R}^d} u(t)^2 dx + \int_{\mathbb{R}^{d+1}} |(-\Delta)^{1/4} u|^2 dx dt + \|u\|_{L^\infty(\mathbb{R}^{d+1})}^2 \leq C \varepsilon_0^2$$

We define

$$F_0(y, s) = u(s/4, (y - x_0(s))/4), \quad f_0(y, s) = \frac{1}{4} f(s/4, (y - x_0(s))/4)$$

where  $x_0(s)$  is solution to

$$\dot{x}_0(s) = \frac{1}{\mathcal{L}^d(B_4)} \int_{B_4} \mathbf{b}(s/4, (y - x_0(s))/4) dy, \quad x_0(0) = 0.$$

We set

$$\tilde{u}_0^* = \frac{4}{\sup_{Q_4^*} F_0^* - \inf_{Q_4^*} F_0^* + 1} \left( F_0^* - \frac{\sup_{Q_4^*} F_0^* + \inf_{Q_4^*} F_0^*}{2} \right),$$

$$B_0(y, s) = \mathbf{b}(s/4, (y - x_0(s))/4) - \dot{x}_0(s),$$

and then for every  $k > 0$

$$F_k(y, s) = F_{k-1}(\sigma s, \sigma(y - x_k(s))), \quad f_k(y, s) = \sigma f_{k-1}(\sigma s, \sigma(y - x_k(s))),$$

$$\tilde{u}_k^* = \frac{4}{\sup_{Q_4^*} F_k^* - \inf_{Q_4^*} F_k^* + \sigma^{\frac{k(q-d-1)}{q}}} \left( F_k^* - \frac{\sup_{Q_4^*} F_k^* + \inf_{Q_4^*} F_k^*}{2} \right),$$

with

$$\dot{x}_k(s) = \frac{1}{\mathcal{L}^d(B_4)} \int_{B_4} B_{k-1}(\sigma s, \sigma(y - x_k(s))) dy, \quad x_k(0) = 0,$$

and

$$B_k(y, s) = B_{k-1}(\sigma s, \sigma(y - x_k(s))) - \dot{x}_k(s).$$

Clearly,  $\tilde{u}_k(y, s) = \tilde{u}_k^*(s, y, 0)$  is solution to

$$\partial_t \tilde{u}_k + B_k \cdot \nabla \tilde{u}_k + (-\Delta)^{1/2} \tilde{u}_k = \frac{4}{\sup_{Q_4^*} F_k^* - \inf_{Q_4^*} F_k^* + \sigma^{\frac{k(q-d-1)}{q}}} f_k,$$

for any  $(x, t) \in (-\infty, +\infty) \times \mathbb{R}^d$  with

$$\tilde{u}_k^* \leq 2 \quad \text{in } Q_4^*,$$

and

$$\operatorname{div} B_k = 0, \quad \int_{B_4} B_k(y, s) dy = 0, \quad \|B_k\|_{BMO} = \|\mathbf{b}\|_{BMO} \leq M_2.$$

And, for any  $m \geq 2$ ,

$$(2.22) \quad \left| \int_{B_{2^m}} B_k dy \right| \leq C m \|B_k\|_{BMO}.$$

We will prove that

$$(2.23) \quad \sup_{s \in [0,1]} |x_0(s)| + \sup_{s \in [0,1]} |\dot{x}_0(s)| \leq M_1,$$

and

$$(2.24) \quad \sup_{s \in [0,1]} |x_k(s)| + \sup_{s \in [0,1]} |\dot{x}_k(s)| \leq C(1 + M_2^2)\sigma^{-1/4} \forall k \geq 1, \sigma \in (0, 1/10).$$

Indeed, since  $|\dot{x}_0(s)| \leq M_1$  and  $x_0(0) = 0$ , one obtains (2.23). By an interpolation inequality in [DDN18] and (2.22), one has for any  $R \geq 1$ ,  $p \geq 2$ ,  $k \geq 1$

$$\|B_k\|_{L_t^\infty L^p(B_R)} \leq C(p, d)R^{d/p} \log(R+1)^{1/p} M_2.$$

This implies for  $k \geq 1$ ,

$$\begin{aligned} |\dot{x}_k(s)| &\leq C\sigma^{-1/4} \|B_{k-1}\|_{L_t^\infty L^{4d}(B_{\sigma(-\sigma x_k(s))})} \leq C\sigma^{-1/4} \|B_{k-1}\|_{L^{4d}(B_{\sigma(1+|x_k(s)|)})} \\ &\leq C\sigma^{-1/4} M_2 + C(2 + |x_k(s)|)^{\frac{1}{2}} |\log(\sigma)|^{1/(4d)} M_2. \end{aligned}$$

So, since  $x_k(0) = 0$

$$\sup_{s \in [0,1]} |x_k(s)| \leq C\sigma^{-1/4} M_2 + C|\log(\sigma)|^{1/(4d)} M_2 + C \sup_{s \in [0,1]} |x_k(s)|^{1/2} |\log(\sigma)|^{1/(4d)} M_2.$$

Thanks to Holder's inequality, one gets (2.24).

So, for  $0 \leq s \leq 1$ , and  $y \in B_4$ , we have

$$|\sigma(y - x_k(s))| \leq \sigma(4 + C(1 + M_2^2)\sigma^{-1/4}) \forall k \geq 1.$$

For  $\sigma \leq (C + 10 + M_2)^{-4}$ , then  $|\sigma(y - x_k(s))| \leq 10^{-2}$  for any  $k \geq 1$ .

Since  $f_k(y, s) = \frac{\sigma^k}{4} f(\frac{\sigma^k}{4}s, \frac{\sigma^k}{4}y - \sum_{m=0}^k \sigma^{k-m} x_m(s)/4)$

$$\left\| \frac{4}{\sup_{Q_4^*} F_k^* - \inf_{Q_4^*} F_k^* + \sigma^{\frac{k(q-d-1)}{q}}} f_k \right\|_{L^q(\mathbb{R}^{d+1})} \leq 4^{\frac{d+1}{q}} \|f\|_{L^q(\mathbb{R}^{d+1})} \leq 4^{\frac{d+1}{q}} \varepsilon_0, \quad \forall k \geq 0.$$

Thanks to Lemma 2.1, we have

$$(2.25) \quad \text{osc}_{Q_{1/4}^*} \tilde{u}_k^* \leq 4 - \lambda^*,$$

for some  $\lambda^* = \lambda^*(d, M_2)$  and  $\varepsilon_0 = \varepsilon_0(d, M_2)$  small enough. This implies,

$$\frac{4}{\text{osc}_{Q_4^*} F_k^* + \sigma^{\frac{k(q-d-1)}{q}}} \text{osc}_{Q_{1/4}^*} F_k^* \leq 4 - \lambda^*.$$

So,

$$(2.26) \quad \text{osc}_{Q_{1/4}^*} F_k^* \leq (1 - \lambda^*/4) \text{osc}_{Q_{1/4}^*} F_{k-1}^* + (1 - \lambda^*/4) \sigma^{\frac{k(q-d-1)}{q}}.$$

Thus,

$$(2.27) \quad \text{osc}_{Q_{1/4}^*} F_k^* \leq (1 - \lambda^*/4)^k \text{osc}_{Q_{1/4}^*} F_0^* + \sum_{m=0}^k (1 - \lambda^*/4)^m \sigma^{(k-m)\frac{q-d-1}{q}} \leq C(1 - \lambda^*/4)^k.$$

Here we have used the fact that

$$|\text{osc}_{Q_{1/4}^*} F_0^*| \leq 2\|F_0^*\|_{L^\infty} \leq 2\|u^*\|_{L^\infty} \leq 2\|u\|_{L^\infty} \leq C.$$

By (2.23) and (2.24) for  $s \leq \sigma^{2k}$  and  $\sigma \leq (C + 10 + M_2)^{-4}$ ,

$$\left| \sum_{m=0}^k \sigma^{k-m} x_m(s)/4 \right| \leq \sigma^{2k} M_1 + C(1 + M_2^2) \sigma^{2k-1/4} \sum_{m=1}^k \sigma^{k-m} \leq \left( \frac{1}{100} + \sigma^k M_1 \right) \sigma^k.$$

It follows from (2.27) and  $F_k^*(y, s) = u(\frac{\sigma^k}{4}s, \frac{\sigma^k}{4}y - \sum_{m=0}^k \sigma^{k-m} x_m(s)/4)$  that for  $\sigma = (C + 10 + M_2)^{-4}$

$$(2.28) \quad \text{osc}_{[-\sigma^{2k}, 0] \times B_{\sigma^{2k}}^*} u^* \leq C(1 - \lambda^*/2)^k = C\sigma^{\frac{|\log(1-\lambda^*/2)|}{|\log(\sigma)|} k}$$

for any  $k \geq k_0 := \frac{\log(100M_1+2)}{|\log(\sigma)|}$ . This gives

$$\sup_{|t_1-t_2|+|x_1-x_2|<\rho_0} \frac{|u(t_1, x_1) - u(t_2, x_2)|}{(|t_1 - t_2| + |x_1 - x_2|)^{\frac{|\log(1-\lambda^*/2)|}{2|\log(\sigma)|}}} \leq C$$

with  $\rho_0 = 10^{-21} M_1^{-2}$ . Hence (2.20) follows and the proof is complete.  $\square$

### 3. PROOF OF THEOREM 1.2

We now come to the potential estimates. Following the approach in [DM11], we first derive a comparison estimate. Notice that here we do not use the fact that the half-laplacian is the Dirichlet-to-Neumann operator of the harmonic extension of a given function. In particular, once one has Hölder regularity of the solution  $u$  translated into decay estimates, the following results work for any integral operator as already mentioned in the introduction.

We first start out with a comparison estimate. Consider  $u$  a weak solution of (1.1) and let  $w$  be a solution of

$$(3.1) \quad \begin{cases} w_t(x, t) + \mathbf{b}(x, t) \cdot \nabla w(x, t) + (-\Delta)^{\frac{1}{2}} w(x, t) = 0 & \text{in } Q_1, \\ w(-1, x) = \bar{u}(-1, x) & \forall x \in \mathbb{R}^d, \\ w(x, t) = \bar{u}(x, t) & \forall x \in B_1(0)^c, t \in (-1, 0). \end{cases}$$

We will rely on the explicit integral expression of the square root of the laplacian, namely  $\int_{\mathbb{R}^d} \frac{w(x, t) - w(y, t)}{|x-y|^{d+1}} dy$ .

**Lemma 3.1.** *Let  $w$  be a solution to (3.1) with  $\bar{u} = u$ . There holds,*

$$(3.2) \quad \left( \int_{Q_1} |u - w|^p \right)^{1/p} \leq C \int_{Q_1} |f|,$$

for any  $p < \frac{d+1}{d}$ .

*Proof.* Set  $v = u - w$ . One has

$$(3.3) \quad \begin{aligned} & \int_{-1}^t \int_{B_1} \phi(x, s) \partial_s v(x, s) dx ds + \int_{-1}^t \int_{B_1} \mathbf{b}(x, s) \cdot \nabla v(x, s) \phi(x, s) dx ds \\ & + \int_{-1}^t \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(v(x, s) - v(y, s))(\phi(x, s) - \phi(y, s))}{|x-y|^{d+1}} dx dy \\ & = \int_{-1}^t \int_{B_1(0)} f(x, s) \phi(x, s) dx ds. \end{aligned}$$

for any  $t \in (-1, 0)$ . By approximation, we can take  $\phi = T_k(v) = \min\{|v|, k\}\text{sign}(v)$  and get

$$(3.4) \quad \sup_{-1 < t < 0} \int_{\mathbb{R}^d} \bar{T}_k(v(x, t)) dx + \int_{-1}^0 \int_{\mathbb{R}^d} \frac{|T_k(v(x, s)) - T_k(v(y, s))|^2}{|x - y|^{d+1}} \leq 2k \int_{Q_1} |f|,$$

where  $\bar{T}_k(\rho) = \int_0^\rho T_k(\rho') d\rho'$ , here we have used the fact that  $v(-1, x) = 0$  for any  $x \in \mathbb{R}^d$ ,

$$\int_{-1}^t \int_{B_1} b(x, s) \cdot \nabla v(x, s) T_k(v(x, s)) dx ds = 0,$$

and

$$(y_1 - y_2)(T_k(y_1) - T_k(y_2)) \geq (T_k(y_1) - T_k(y_2))^2 \quad \forall y_1, y_2 \in \mathbb{R}.$$

Thanks to  $\lim_{k \rightarrow 0} \bar{T}_k(\rho) k^{-1} = |\rho|$  and Sobolev inequality

$$\|f\|_{L^{\frac{2d}{d-1}}(\mathbb{R}^d)}^2 \leq C \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{|f(x) - f(y)|^2}{|x - y|^{d+1}} dx dy \quad \forall f \in \dot{H}^{1/2}(\mathbb{R}^d).$$

We get from (3.4) that

$$(3.5) \quad \sup_{-1 < t < 0} \int_{B_1} |v(x, t)| dx + \frac{1}{k} \int_{-1}^0 \|T_k(v(t))\|_{L^{\frac{2d}{d-1}}(Q_1)}^2 dt \leq C \int_{Q_1} |f| \quad \forall k > 0.$$

Thus, for  $k > 0$ ,

$$\begin{aligned} \mathcal{L}^{d+1}(Q_1 \cap \{|v| > k\}) &\leq \left[ \sup_{-1 < t < 0} \int_{B_1} \mathbf{1}_{|v(x, t)| > k} dx \right]^{\frac{1}{d}} \int_{-1}^0 \left[ \int_{B_1} \mathbf{1}_{|v(x, t)| > k} dx \right]^{\frac{d-1}{d}} dt \\ &\leq C \left[ k^{-1} \sup_{-1 < t < 0} \int_{B_1} |v(x, t)| dx \right]^{\frac{1}{d}} \int_{-1}^0 \|T_k(v)\|_{L^{\frac{2d}{d-1}}(Q_1)}^2 dt k^{-2} \\ &\leq C \left( k^{-1} \int_{Q_1} |f| \right)^{1 + \frac{1}{d}}. \end{aligned}$$

This gives (3.2). The proof is complete.  $\square$

In the following we denote

$$(3.6) \quad \mathbf{I}(u, x, t, \rho) = \left( \int_{Q_\rho(x, t)} |u - (u)_{Q_\rho(x, t)}|^{q_0} \right)^{1/q_0}$$

for  $q_0 = 1 + \frac{1}{100d}$ .

**Lemma 3.2.** *Let  $q_0 = 1 + \frac{1}{100d}$ . Assume that*

$$\sup_{(x, t) \in \mathbb{R}^{d+1}} \left| \int_{B_1(x)} \mathbf{b}(y, t) dy \right| \leq M_1, \quad \|\mathbf{b}\|_{L^\infty(\mathbb{R}; BMO(\mathbb{R}^d))} \leq M_2,$$

for some  $M_1, M_2 \geq 1$ . Let  $\alpha \in (0, 1)$  be as in Theorem 1.1. There holds

$$(3.7) \quad \|w\|_{C^\alpha(Q_{1/2})} + \|w\|_{L^\infty(Q_{1/2})} \lesssim_{M_2} M_1^{3\alpha} \left( \|w\|_{L^{q_0}(Q_1)} + \int_{-1}^0 \int_{|y| > 1} \frac{|\bar{u}(y, t)|}{|y|^{d+1}} \right).$$

In particular, we have for all  $\varepsilon \in (0, 1/2)$

$$(3.8) \quad \mathbf{I}(w, 0, 0, \varepsilon) \lesssim_{M_2} M_1^{3\alpha} \varepsilon^\alpha (\mathbf{I}(w, 0, 0, 1) + \text{Tail}(u, 0, 0, 1)).$$

*Proof of Lemma 3.2.* The proof goes by several steps.

*Step 1:* We have,

$$\partial_t(\phi w)(x, t) + \mathbf{b}(x, t) \cdot \nabla(\phi w)(x, t) + \int_{\mathbb{R}^d} \frac{(\phi w)(x, t) - (\phi w)(y, t)}{|x - y|^{d+1}} dy = F(x, t)$$

where  $\phi \in C_c^\infty(Q_{3/4})$  with  $\mathbf{1}_{Q_{5/8}} \leq \phi \leq \mathbf{1}_{Q_{3/4}}$  and

$$\begin{aligned} F(x, t) &= w \partial_t \phi + w \mathbf{b} \cdot \nabla \phi + \mathbf{1}_{B_1} \int_{\mathbb{R}^d} \frac{(\phi(x, t) - \phi(y, t))(w(y, t) - w(x, t))}{|x - y|^{d+1}} dy \\ &\quad + \mathbf{1}_{B_1} \int_{\mathbb{R}^d} \frac{\phi(x, t) - \phi(y, t)}{|x - y|^{d+1}} dy w(x, t) - \mathbf{1}_{|x| > 1} \int_{B_1} \frac{\phi(y, t) w(y, t)}{|x - y|^{d+1}} dy. \end{aligned}$$

Applying Theorem 1.1, one has

$$\sup_{(x, t) \in Q_{1/2}} \|w\|_{C^\alpha(Q_{\delta_0/(M_1+2)^2}(x, t))} + \|w\|_{L^\infty(Q_{1/2})} \lesssim_{M_2} \|F\|_{L^1} + \|F\|_{L^{d+2}}$$

for some  $\delta_0 = \delta_0(M_2, d)$  and  $\alpha \in (0, 1/10)$ . Thanks to

$$\|\mathbf{b}(t)\|_{L^p(B_1)} \lesssim_{p, M_2} M_1^{1/p} \quad \forall p \geq 2,$$

we obtain for any  $\delta_1 \in (0, 10^{-1})$ ,

$$(3.9)$$

$$\begin{aligned} &\|w\|_{C^\alpha(Q_{1/2})} + \|w\|_{L^\infty(Q_{1/2})} \\ &\lesssim M_1^{2\alpha} \left( \mathbf{I} + \|w\|_{L^{d+3}(Q_{\frac{3}{4}})} + \left( \int_{-1}^0 \int_{|x| < \frac{7}{8}} \left( \int_{|y| < \frac{3}{4}} \frac{|w(x, t) - w(y, t)|}{|x - y|^d} dy \right)^{d+2} dx dt \right)^{\frac{1}{d+2}} \right), \end{aligned}$$

where

$$\mathbf{I} = \|w\|_{L^{q_0}(Q_1)} + \int_{-1}^0 \int_{|y| > 1} \frac{|\bar{u}(y, s)|}{|y|^{d+1}} dy ds.$$

*Step 2:* Let  $\varphi \in C_c^\infty(B_r)$  for  $\frac{3}{4} < r < 1$  and  $q > 0$ . Using  $|w|^{q-1} w \varphi^{q+1}$  as a test function for (3.1), one gets

$$\begin{aligned} &\frac{1}{q+1} \frac{d}{dt} \int |w(t)|^{q+1} \varphi^{q+1} + \int \int \frac{(w(x, t) - w(y, t))(|w|^{q-1} w \varphi^{q+1}(x, t) - |w|^{q-1} w \varphi^{q+1}(y, t))}{|x - y|^{d+1}} \\ &= \frac{1}{q+1} \int \mathbf{b} \cdot \nabla(\varphi^{q+1}) |w|^{q+1}. \end{aligned}$$

Thanks to (5.1), we have

$$\begin{aligned} &\frac{d}{dt} \int |w(t) \varphi|^{q+1} + c \| |w \varphi|^{\frac{q+1}{2}} \|_{H^{1/2}}^2 \lesssim \int |\mathbf{b}| |\nabla \varphi| \varphi^p |w|^{q+1} + \int_{B_r \times B_r} \frac{(|w(x, t)| + |w(y, t)|)^{q+1}}{|x - y|^{d+1}} G \\ &+ \left( \sup_{x \in \text{supp} \varphi} \int_{\mathbb{R}^d \setminus B_r} \frac{|w(y, t)|}{|x - y|^{d+1}} \right) \int_{B_r} |w|^q \varphi^{q+1} dx + \int_{B_r} \frac{|w(x, t) \varphi(x)|^{q+1}}{r - |x|} dx, \end{aligned}$$

where

$$G(x, y) = \mathbf{1}_{q \geq 1} (|\varphi(x)| + |\varphi(y)|)^{q-1} |\varphi(x) - \varphi(y)|^2 + \mathbf{1}_{q < 1} |\varphi(x) - \varphi(y)|^{q+1}.$$

This implies that for  $0 < R_1 < R_2$ ,

$$\begin{aligned}
& \sup_{-R_1 < t < 0} \int |w(t)\varphi|^{q+1} + \int_{-R_1}^0 \| |w(t)\varphi(t)|^{\frac{q+1}{2}} \|_{H^{1/2}}^2 \lesssim \frac{1}{R_2 - R_1} \int_{-R_2}^{-R_1} \int |w(t)\varphi|^{q+1} \\
& + \int_{-R_2}^0 \int |\mathbf{b}| |\nabla \varphi| \varphi^p |w|^{q+1} + \int_{-R_2}^0 \int_{B_r} \int_{B_r} \frac{(|w(x,t)| + |w(y,t)|)^{q+1}}{|x-y|^{d+1}} G(x,y) \\
(3.10) \quad & + \int_{-R_2}^0 \left( \sup_{x \in \text{supp} \varphi} \int_{\mathbb{R} \setminus B_r} \frac{|w(y,t)|}{|x-y|^{d+1}} \right) \int_{B_r} |w|^q \varphi^{q+1} dx + \int_{-R_2}^0 \int_{B_r} \frac{|w(x,t)\varphi(x)|^{q+1}}{r-|x|} dx dt
\end{aligned}$$

Using Sobolev inequality,

$$\begin{aligned}
& \| |w\varphi| \|_{L^{\frac{(d+1)(q+1)}{d}}((-R_1,0) \times \mathbb{R}^d)}^{q+1} \lesssim \frac{1}{R_2 - R_1} \int_{-R_2}^{-R_1} \int |w(t)\varphi|^{q+1} + \int_{-R_2}^0 \int |\mathbf{b}| |\nabla \varphi| \varphi^p |w|^{q+1} \\
& + \int_{-R_2}^0 \int_{B_r} \int_{B_r} \frac{(|w(x,t)| + |w(y,t)|)^{q+1}}{|x-y|^{d+1}} G \\
& + \int_{-R_2}^0 \left( \sup_{x \in \text{supp} \varphi} \int_{\mathbb{R} \setminus B_r} \frac{|w(y,t)|}{|x-y|^{d+1}} \right) \int_{B_r} |w|^q \varphi^{q+1} dx + \int_{-R_2}^0 \int_{B_r} \frac{|w(x,t)\varphi(x)|^{q+1}}{r-|x|} dx dt.
\end{aligned}$$

Therefore, by a bootstrap argument, we obtain for any  $\delta_1 > 0$

$$(3.11) \quad \|w\|_{L^{d+3}(Q_{\frac{3}{4}})^+} \left( \int_{-1}^0 \int_{|x| < \frac{7}{8}} \left( \int_{|y| < \frac{3}{4}} \frac{|w(x,t) - w(y,t)|}{|x-y|^d} dy dt \right)^{d+2} dx \right)^{\frac{1}{d+2}} \lesssim_{\delta_1, M_2} M_1^{\delta_1} \mathbf{I}.$$

Combining this with (3.9) yields (3.7). The proof is complete.  $\square$

Lemma 3.1 implies

**Lemma 3.3.** *Assume that*

$$\sup_{(x,t) \in \mathbb{R}^{d+1}} \left| \int_{B_1(x)} \mathbf{b}(y,t) dy \right| \leq M_1, \quad \|\mathbf{b}\|_{L^\infty(\mathbb{R}; BMO(\mathbb{R}^d))} \leq M_2$$

for some  $M_1, M_2 \geq 1$ . Let  $\alpha \in (0, 1)$  be as in Theorem 1.1. There holds for any  $\varepsilon \in (0, 1/2)$

$$(3.12) \quad \mathbf{I}(u, 0, 0, \varepsilon) \lesssim_{M_2} M_1^{3\alpha} \varepsilon^\alpha \mathbf{I}(u, 0, 0, 1) + \varepsilon^{-d-1} \int_{Q_1} |f| + M_1^{3\alpha} \varepsilon^\alpha \text{Tail}(u, 0, 0, 1).$$

*Proof.* We have the chain of inequalities

$$\begin{aligned}
\mathbf{I}(u, 0, 0, \varepsilon) & \leq \mathbf{I}(w, 0, 0, \varepsilon) + 2 \left( \int_{Q_\varepsilon} |u - w|^{q_0} \right)^{1/q_0} \\
& \stackrel{(3.8)}{\lesssim} M_1^{3\alpha} \varepsilon^\alpha \mathbf{I}(w, 0, 0, 1) + \varepsilon^{-d-1} \left( \int_{Q_1} |u - w|^{q_0} \right)^{1/q_0} + M_1^{3\alpha} \varepsilon^\alpha \text{Tail}(u, 0, 0, 1) \\
& \lesssim M_1^{3\alpha} \varepsilon^\alpha \mathbf{I}(u, 0, 0, 1) + \varepsilon^{-d-1} \left( \int_{Q_1} |u - w|^{q_0} \right)^{1/q_0} + M_1^{3\alpha} \varepsilon^\alpha \text{Tail}(u, 0, 0, 1)
\end{aligned}$$

Applying (3.2) with  $p = q_0$ , we get (3.8). The proof is complete.  $\square$



The following lemma is a simple iteration of the previous one. Notice that this is the place where we use the fact that the drift  $\mathbf{b}$  is bounded in  $L_t^\infty L_x^\infty$ . This is necessary to ensure that under the iteration one keeps the bound uniform. With more work, one could obtain the same lemma assuming that the drift is BMO in space, but at the price of losing integrability on the data  $f$ . However, in view of the applications to SQG, the boundedness in  $L^\infty$  is enough and we then refrain to go into such a generality.

**Lemma 3.4.** *Assume that*

$$\|\mathbf{b}\|_{L^\infty} \leq M_3 < \infty.$$

*There holds for any  $\varepsilon \in (0, 1/2)$*

$$\mathbf{I}(u, x, t, \varepsilon r) \lesssim_{M_3} \varepsilon^\alpha \mathbf{I}(u, x, t, r) + \varepsilon^{-d-1} \frac{\int_{Q_r(x,t)} |f|}{r^d} + \varepsilon^\alpha \text{Tail}(u, x, t, r).$$

**Lemma 3.5.** *Assume that*

$$\sup_{(x,t) \in \mathbb{R}^{d+1}} \left| \int_{B_1(x)} \mathbf{b}(y, t) dy \right| \leq M_1 < \infty, \quad \|\mathbf{b}\|_{L^\infty(\mathbb{R}; BMO(\mathbb{R}^d))} \leq M_2 < \infty$$

*for some  $M_1, M_2 \geq 1$ . There holds for any  $\varepsilon \in (0, 1/2)$  and  $r \in (0, 1/2)$*

(3.13)

$$\mathbf{I}(u, x, t, \varepsilon r) \lesssim_{M_1, M_2} \varepsilon^\alpha |\log(r)|^{3\alpha} \mathbf{I}(u, x, t, r) + \varepsilon^{-d-1} \frac{\int_{Q_r(x,t)} |f|}{r^d} + \varepsilon^\alpha |\log(r)|^{3\alpha} \text{Tail}(u, x, t, r).$$

*In particular, for any  $\varepsilon \in (0, 10^{-1})$  and  $r \in (0, 1/2)$*

$$\begin{aligned} \mathbf{I}(u, x, t, \varepsilon |\log(r)|^{-3} r) &\lesssim_{M_1, M_2} \varepsilon^\alpha \mathbf{I}(u, x, t, r) \\ &+ \varepsilon^{-d-1} |\log(r)|^{3(d+1)} \frac{\int_{Q_r(x,t)} |f|}{r^d} + \varepsilon^\alpha \text{Tail}(u, x, t, r). \end{aligned}$$

*Proof.* Since  $\sup_{x,t} \left| \int_{B_r(x)} \mathbf{b}(y, t) dy \right| \lesssim |\log(r)| M_2 + M_1 \lesssim_{M_1, M_2} |\log(r)|$  for any  $r \leq 1/2$ , (3.13) follows from (3.12).  $\square$

The following is a consequence of the previous 2 Lemma:

**Corollary 3.1.** *Assume  $f(x, t) = \omega(x, t) + \delta_{t=0} \otimes \sigma(x)$  where  $\omega$  is a bounded Radon measure in  $\mathbb{R}^{d+1}$  and  $\nabla \sigma$  is a bounded Radon measure in  $\mathbb{R}^d$ . Set*

$$\mathbf{E}(\omega, \sigma, x, t, r) = \frac{|\omega|(Q_r(x, t))}{r^d} + \frac{|\delta_{t=0} \otimes \nabla \sigma|(Q_r(x, t))}{r^{d-1}}.$$

*Assume that*

$$\sup_{(x,t) \in \mathbb{R}^{d+1}} \left| \int_{B_1(x)} \mathbf{b}(y, t) dy \right| \leq M_1 < \infty, \quad \|\mathbf{b}\|_{L^\infty(\mathbb{R}; BMO(\mathbb{R}^d))} \leq M_2 < \infty$$

*for some  $M_1, M_2 \geq 1$ . Then, it holds, for any  $\varepsilon \in (0, 10^{-1})$  and  $r \in (0, 1/2)$*

(3.14)

$$\begin{aligned} \mathbf{I}(u, x, t, \varepsilon |\log(r)|^{-3} r) &\lesssim_{M_1, M_2} \varepsilon^\alpha \mathbf{I}(u, x, t, r) \\ &+ \varepsilon^{-d-1} |\log(r)|^{3(d+1)} \mathbf{E}(\omega, \sigma, x, t, r) + \varepsilon^\alpha \text{Tail}(u, x, t, r). \end{aligned}$$

*Moreover, if  $\|\mathbf{b}\|_{L^\infty} \leq M_3 < \infty$ , then for any  $\varepsilon \in (0, 10^{-1})$  and  $r \in (0, 1/2)$*

(3.15)

$$\mathbf{I}(u, x, t, \varepsilon r) \lesssim_{M_3} \varepsilon^\alpha \mathbf{I}(u, x, t, r) + \varepsilon^{-d-1} \mathbf{E}(\omega, \sigma, x, t, r) + \varepsilon^\alpha \text{Tail}(u, x, t, r).$$

*Proof.* Let  $\varrho_n$  be the standard sequence of mollifiers in  $\mathbb{R}$  and  $u_n$  be a solution of

$$(3.16) \quad \partial_t u_n + \mathbf{b} \cdot \nabla u_n + (-\Delta)^{1/2} u_n = \omega(x, t) + \varrho_n(t) \sigma(x) \quad \text{in } \mathbb{R} \times \mathbb{R}^d.$$

Define  $v_n(x, t) = u_n(x, t) - g_n(t)$  with

$$g_n(t) = \int_{n^{-1/4}}^t \varrho_n(s) ds \int_{B_r(x_0)} \sigma(y) dy.$$

The function  $v_n$  solves

$$\partial_t v_n + \mathbf{b} \cdot \nabla v_n + (-\Delta)^{1/2} v_n = \omega(x, t) + \varrho_n(t) \left( \sigma(x) - \int_{B_r(x_0)} \sigma(y) dy \right) \quad \text{in } \mathbb{R} \times \mathbb{R}^d.$$

and  $v_n \rightarrow u$  in  $L_{\text{loc}}^p$  for any  $p < 1 + 1/d$ . By Lemma 3.4, (3.5) and Poincaré's inequality, we have for  $Q_\rho = Q_\rho(t_0, x_0)$

$$\begin{aligned} \mathbf{I}(v_n, x, t, \varepsilon r) &\lesssim_{M_3} \varepsilon^\alpha \mathbf{I}(v_n, x, t, r) \\ &+ \varepsilon^{-d-1} \left( \frac{|\omega|(Q_r(x, t))}{r^d} + \frac{\int_{Q_r} |\varrho_n(t)| |\nabla \sigma| dy dt}{r^{d-1}} \right) + \varepsilon^\alpha \text{Tail}(v_n, x_0, t_0, r), \end{aligned}$$

and

$$\begin{aligned} \mathbf{I}(u_n, x, t, \varepsilon |\log(r)|^{-3} r) &\lesssim_{M_1, M_2} \varepsilon^\alpha \mathbf{I}(u_n, x_0, t_0, r) \\ &+ \varepsilon^{-d-1} |\log(r)|^{3(d+1)} \left( \frac{|\omega|(Q_r(x_0, t_0))}{r^d} + \frac{\int_{Q_r} |\varrho_n(t)| |\nabla \sigma| dy dt}{r^{d-1}} \right) + \varepsilon^\alpha \text{Tail}(u_n, x_0, t_0, r). \end{aligned}$$

Letting  $n \rightarrow \infty$ , we get (3.14) and (3.15). The proof is complete.  $\square$

*Proof of Theorem 1.2.* We can assume that  $u(x, t) = \lim_{\rho \rightarrow 0} \int_{Q_\rho(x, t)} u$ . It follows from (3.15) that

$$\begin{aligned} \int_0^{\varepsilon r_0} \mathbf{I}(u, x, t, r) \frac{dr}{r} &\leq C \varepsilon^\alpha \int_0^{r_0} \mathbf{I}(u, x, t, r) \frac{dr}{r} + C_\varepsilon \int_0^{r_0} \mathbf{E}(\omega, \sigma, x, t, r) \frac{dr}{r} \\ &+ C \varepsilon^\alpha \int_0^{r_0} \text{Tail}(u, x, t, r) \frac{dr}{r}, \end{aligned}$$

for  $C = C(d, M_3)$ .

Choosing  $\varepsilon > 0$  such that  $C \varepsilon^\alpha \leq 1/2$ , we find

$$\begin{aligned} \int_0^{r_0} \mathbf{I}(u, x, t, r) \frac{dr}{r} &\leq C \left( \int_{Q_{r_0}} |u|^{q_0} \right)^{1/q_0} + C_\varepsilon \int_0^{r_0} \mathbf{E}(\omega, \sigma, x, t, r) \frac{dr}{r} \\ &+ C \alpha^{\alpha_0} \int_0^{r_0} \text{Tail}(u, x, t, r) \frac{dr}{r}. \end{aligned}$$

It is not hard to show that

$$\int_0^{r_0} \text{Tail}(u, x, t, r) \frac{dr}{r} \lesssim \int_0^{r_0} \mathbf{I}(u, x, t, r) \frac{dr}{r} + \inf_\lambda \int_{t-2r_0}^t \int_{|x-y|>r_0/2} \frac{|u(y, s) - \lambda|}{|x-y|^{d+1}} dy ds.$$

Thus, letting  $\varepsilon$  be small enough, we obtain

$$\int_0^{r_0} \mathbf{I}(u, x, t, r) \frac{dr}{r} \leq C \left( \int_{Q_{2r_0}} |u|^{q_0} \right)^{1/q_0} + C \int_0^{r_0} \mathbf{E}(\omega, \sigma, x, t, r) \frac{dr}{r} + C \text{Tail}(u, x, t, r_0).$$

Therefore, using the fact that

$$|u(x, t)| \lesssim \int_0^{r_0} \mathbf{I}(u, x, t, r) \frac{dr}{r} + \left( \int_{Q_{r_0}} |u|^{q_0} \right)^{1/q_0},$$

we get (1.8).

To prove (1.9), we apply (1.8) to  $u - a$  with  $a = (u)_{Q_s(x_1, t_1)}$ ,  $s = |x_1 - x_2| + |t_1 - t_2| < r$  to obtain

$$|u(x_i, t_i) - a| \lesssim_{M_3} \left( \int_{Q_s(x_i, t_i)} |u - a|^{q_0} \right)^{1/q_0} + \int_0^{2s} \mathbf{E}(\omega, \sigma, x_i, t_i, \rho) \frac{d\rho}{\rho} + \text{Tail}(u, x_i, t_i, s).$$

Thus,

(3.17)

$$\begin{aligned} & |u(x_1, t_1) - u(x_2, t_2)| \\ & \leq C \sum_{i=1,2} \left( \mathbf{I}(u, x_i, t_i, 3s) + \int_0^{2s} \mathbf{E}(\omega, \sigma, x_i, t_i, \rho) \frac{d\rho}{\rho} + \text{Tail}(u, x_i, t_i, s) \right), \end{aligned}$$

where we have used the fact that

$$\sum_{i=1,2} \left( \int_{Q_s(x_i, t_i)} |u - a|^{q_0} \right)^{1/q_0} \leq C \mathbf{I}(u, x_1, t_1, 3s).$$

So, we get that

(3.18)

$$\begin{aligned} \frac{|u(x_1, t_1) - u(x_2, t_2)|}{(|x_1 - x_2| + |t_1 - t_2|)^\beta} & \lesssim s^{-\beta} \mathbf{I}(u, x_1, t_1, 3s) \\ & + \sum_{i=1,2} \int_0^{2s} \mathbf{E}(\omega, \sigma, x_i, t_i, \rho) \frac{d\rho}{\rho^{1+\beta}} + s^{-\beta} \text{Tail}(u, x_i, t_i, s). \end{aligned}$$

We now estimate the terms  $s^{-\beta} \mathbf{I}(u, x, t, s)$  and  $s^{-\beta} \text{Tail}(u, x, t, s)$  in terms of potentials. Similarly, we also can show that

$$\begin{aligned} \varepsilon^\beta \int_0^{\varepsilon r_0} \mathbf{I}(u, x, t, r) \frac{dr}{r^{1+\beta}} & \lesssim \varepsilon^\alpha \int_0^{r_0} \mathbf{I}(u, x, t, r) \frac{dr}{r^{1+\beta}} + C_\varepsilon \int_0^{r_0} \mathbf{E}(\omega, \sigma, x, t, \rho) \frac{d\rho}{\rho^{1+\beta}} \\ & + \varepsilon^\alpha \int_0^{r_0} \text{Tail}(u, x, t, r) \frac{dr}{r^{1+\beta}}. \end{aligned}$$

Thus, we get

$$\begin{aligned} & \int_0^{r_0} \mathbf{I}(u, x, t, r) \frac{dr}{r^{1+\beta}} + \int_0^{r_0} \text{Tail}(u, x, t, r) \frac{dr}{r^{1+\beta}} \\ & \lesssim r_0^{-\beta} \left( \int_{Q_{2r_0}} |u|^{q_0} \right)^{1/q_0} + \int_0^{r_0} \mathbf{E}(\omega, \sigma, x, t, \rho) \frac{d\rho}{\rho^{1+\beta}} + r_0^{-\beta} \text{Tail}(u, x, t, r_0). \end{aligned}$$

Combining this with (3.18) we find (1.9).

*Proof of Theorem 1.3.* In this proof, the constant  $C$  will depend on  $d, M_1, M_2$ . We can assume that  $u(x, t) = \lim_{\rho \rightarrow 0} \mathcal{f}_{Q_\rho(x, t)} u$ . It follows from (3.14) that for any  $r_0 \leq 10^{-10}$ ,

$$\begin{aligned} \int_0^{r_0} \mathbf{I}(u, x, t, \varepsilon |\log(r)|^{-3} r) \frac{dr}{r} &\leq C \varepsilon^\alpha \int_0^{r_0} \mathbf{I}(u, x, t, r) \frac{dr}{r} \\ &+ C \varepsilon^{-d-1} \int_0^{r_0} |\log(r)|^{3(d+1)} \mathbf{E}(\omega, \sigma, x, t, r) \frac{dr}{r} + C \varepsilon^\alpha \int_0^{r_0} \text{Tail}(u, x, t, r) \frac{dr}{r}. \end{aligned}$$

So,

$$\begin{aligned} \int_0^{\varepsilon r_0 |\log(r_0)|^{-3}} \mathbf{I}(u, x, t, r) \frac{dr}{r} &\leq C \varepsilon^\alpha \int_0^{r_0} \mathbf{I}(u, x, t, r) \frac{dr}{r} \\ &+ C \varepsilon^{-d-1} \int_0^{r_0} |\log(r)|^{3(d+1)} \mathbf{E}(\omega, \sigma, x, t, r) \frac{dr}{r} + C \varepsilon^\alpha \int_0^{r_0} \text{Tail}(u, x, t, r) \frac{dr}{r}. \end{aligned}$$

As proof of Theorem 1.2, we have

$$\begin{aligned} |u(x, t)| + \int_0^{r_0} \mathbf{I}(u, x, t, r) \frac{dr}{r} &\leq C |\log(r_0)|^{\frac{3d}{q_0}} \left( \int_{Q_{2r_0}} |u|^{q_0} \right)^{1/q_0} \\ &+ C \int_0^{r_0} |\log(r)|^{3(d+1)} \mathbf{E}(\omega, \sigma, x, t, r) \frac{dr}{r} + C \text{Tail}(u, x, t, r_0). \end{aligned}$$

This gives (1.10).

To prove (1.11), as proof of Theorem 1.2 we apply (1.8) to  $u - (u)_{Q_s(x, t)}$  with  $s = |x_1 - x_2| + |t_1 - t_2| < r/2$  to obtain

$$\begin{aligned} (3.19) \quad \frac{|u(x_1, t_1) - u(x_2, t_2)|}{(|x_1 - x_2| + |t_1 - t_2|)^\beta} &\leq C s^{-\beta} |\log(s)|^{\frac{3d}{q_0}} \mathbf{I}(u, x_1, t_1, 3s) \\ &+ \sum_{i=1,2} C \int_0^{2s} |\log(r)|^{3(d+1)} \mathbf{E}(\omega, \sigma, x_i, t_i, r) \frac{dr}{r^{1+\beta}} + C s^{-\beta} \text{Tail}(u, x_i, t_i, s). \end{aligned}$$

We now estimate the terms  $s^{-\beta} |\log(s)|^{\frac{3d}{q_0}} \mathbf{I}(u, x, t, s)$  and  $s^{-\beta} |\log(s)|^{\frac{3d}{q_0}} \text{Tail}(u, x, t, s)$  in terms of potentials. Apply (3.14) to  $\varepsilon = \varepsilon_0 |\log(r)|^{-\frac{3\beta}{\alpha-\beta}}$  yields

$$\begin{aligned} \mathbf{I}(u, x, t, \varepsilon_0 |\log(r)|^{-\frac{3\alpha}{\alpha-\beta}} r) &\leq C \varepsilon_0^\alpha |\log(r)|^{-\frac{3\alpha\beta}{\alpha-\beta}} \mathbf{I}(u, x, t, r) \\ &+ C \varepsilon_0^{-d-1} |\log(r)|^{\frac{3\alpha(d+1)}{\alpha-\beta}} \mathbf{E}(\omega, \sigma, x, t, r) + C \varepsilon_0^\alpha |\log(r)|^{-\frac{3\alpha\beta}{\alpha-\beta}} \text{Tail}(u, x, t, r). \end{aligned}$$

So, for any  $r < \varepsilon_0^2 < 10^{-20}$

$$\begin{aligned} C^{-1} \int_0^{\varepsilon_0 |\log(r_0)|^{-\frac{3\alpha}{\alpha-\beta}} r_0} \frac{\mathbf{I}(u, x, t, r)}{r^\beta |\log(r)|^{-\frac{3d}{q_0}} r} \frac{dr}{r} &\leq \int_0^{r_0} \frac{\mathbf{I}(u, x, t, \varepsilon_0 |\log(r)|^{-\frac{3\alpha}{\alpha-\beta}} r)}{(\varepsilon_0 |\log(r)|^{-\frac{3\alpha}{\alpha-\beta}} r)^\beta |\log(r)|^{-\frac{3d}{q_0}} r} \frac{dr}{r} \\ &\leq C \varepsilon_0^{\alpha-\beta} \int_0^{r_0} \frac{\mathbf{I}(u, x, t, r)}{r^\beta |\log(r)|^{-\frac{3d}{q_0}} r} \frac{dr}{r} + C \varepsilon_0^{-d-1-\beta} \int_0^{r_0} |\log(r)|^{\frac{3\alpha(d+1+\beta)}{\alpha-\beta} + \frac{3d}{q_0}} \mathbf{E}(\omega, \sigma, x, t, r) \frac{dr}{r^{1+\beta}} \\ &+ C \varepsilon_0^{\alpha-\beta} \int_0^{r_0} \frac{\text{Tail}(u, x, t, r)}{r^\beta |\log(r)|^{-\frac{3d}{q_0}} r} \frac{dr}{r}. \end{aligned}$$

As proof of Theorem 1.2, we get

$$\begin{aligned} & \int_0^{r_0} \frac{\mathbf{I}(u, x, t, r)}{r^\beta |\log(r)|^{-\frac{3d}{q_0}} r} dr + \int_0^{r_0} \frac{\text{Tail}(u, x, t, r)}{r^\beta |\log(r)|^{-\frac{3d}{q_0}} r} dr \leq Cr_0^{-\beta} |\log(r_0)|^{\beta_1} \left( \int_{Q_{2r_0}} |u|^{q_0} \right)^{1/q_0} \\ & + C \int_0^{r_0} |\log(r)|^{\beta_2} \mathbf{E}(\omega, \sigma, x, t, r) \frac{dr}{r^{1+\beta}} + Cr_0^{-\beta} |\log(r_0)|^{\frac{3d}{q_0}} \text{Tail}(u, x, t, r_0), \end{aligned}$$

where  $\beta_1 = \frac{3\alpha}{\alpha-\beta}(\beta + \frac{d}{q_0}) + \frac{3d}{q_0}$  and  $\beta_2 = \frac{3\alpha(d+1+\beta)}{\alpha-\beta} + \frac{3d}{q_0}$ .

Combining this with (3.19) we find (1.11). The proof is complete.

#### 4. THE CASE OF BOUNDED DOMAINS

This section is devoted to the investigation of similar results for (1.4). In the case of bounded domains the main mathematical issue is the boundary regularity (see e.g. [SV19] and references therein). The methods we developed in the previous sections adapt straightforwardly to the *interior* regularity. We state below those results in the case of the critical SQG system, i.e.

$$(4.1) \quad \partial_t u + \mathcal{K}u \cdot \nabla u + (-\Delta)_D^{1/2} u = f \quad \text{in } \mathbb{R} \times \Omega.$$

where  $\mathcal{K}u = \nabla^\perp (-\Delta)_D^{-1/2} u$ , the vectorial Riesz transform defined via functional calculus.

**Theorem 4.1.** *Let  $u$  be a weak solution of (4.1) and  $f$  satisfying  $f \in L^q(\Omega)$  for some  $q > d + 1$ . Then  $u$  is locally Hölder continuous in space and time and satisfies the parabolic potential estimates in Theorem 1.2 and 1.3.*

We explain below the main amendments to the previous proofs. The technical tool is the extension in one more dimension in a cylinder-like geometry. The extended problem is set in the cylinder  $\Omega \times (0, \infty)$  and it will be convenient to use the following notation:  $x \in \mathbb{R}^d$ ,  $y > 0$ , and  $X = (x, y) \in \mathbb{R}_+^{d+1} := \mathbb{R}^d \times (0, \infty)$ ; likewise, we denote by  $\mathcal{C}$  the cylinder  $\Omega \times (0, \infty)$  and by  $\partial_L \mathcal{C}$  its lateral boundary, i.e.  $\partial\Omega \times (0, \infty)$ . The ambient space  $H_{0,L}^s(\mathcal{C})$  is defined as the completion of

$$C_{0,L}^{\frac{1}{2}}(\mathcal{C}) := \{U \in C^\infty(\overline{\mathcal{C}}) : U = 0 \text{ on } \partial_L \mathcal{C}\}$$

with respect to the norm

$$(4.2) \quad \|U\|_{\mathcal{C}} = \left( \int_{\mathcal{C}} |\nabla U|^2 \right)^{1/2}.$$

This is a Hilbert space endowed with the natural inner product. We point out that if  $\Omega$  is a smooth bounded domain then

$$H(\Omega) = \{u = \text{trace}|_{\Omega \times \{0\}} U : U \in H_{0,L}^s(\mathcal{C})\}.$$

The extension problem is the following: given  $u \in H(\Omega)$ , we solve

$$(4.3) \quad \begin{cases} \text{div}(\nabla u^*) = 0 & \text{in } \mathcal{C}, \\ u^* = 0 & \text{on } \partial_L \mathcal{C}, \\ u^* = u & \text{on } \Omega, \end{cases}$$

for  $u^* \in H_{0,L}^s(\mathcal{C})$ . Then, up to a multiplicative constant,

$$(4.4) \quad (-\Delta)_D^{\frac{1}{2}} u = - \lim_{y \rightarrow 0} \partial_y u^*.$$

See [CT10] for more details. The previous extended problem allows to run the De Giorgi method exactly as in Theorem 1.1 to obtain interior Hölder regularity.

To deal with the potential estimates, we rely on the kernel representation of  $(-\Delta)_D^{\frac{1}{2}}$  which can be found in [SV03], i.e.

$$(4.5) \quad (-\Delta)_D^{\frac{1}{2}} u(x) = \int_{\Omega} (u(x) - u(y)) K(x, y) dy + B(x)u(x)$$

where  $B \sim \frac{1}{\text{dist}(x, \partial\Omega)}$  and  $K$  is supported in  $\bar{\Omega} \times \bar{\Omega}$  and satisfies

$$K(x, y) \sim \frac{1}{|x - y|^{d+1}} \left( \frac{\text{dist}(x, \partial\Omega)}{|x - y|} \wedge 1 \right) \left( \frac{\text{dist}(y, \partial\Omega)}{|x - y|} \wedge 1 \right).$$

The potential estimates being localized away from the boundary, the zero order term in (4.5) does not introduce any difficulty. Concerning the integral term in (4.5), the same computations we developed before, choosing an interior cut-off functions, leads to the same estimates.

## 5. APPENDIX

We prove here a technical lemma used for the proof of the potential estimates.

**Lemma 5.1.** *Let  $q > 0$  and  $\theta \in (0, \min\{q, 1\})$ . For any  $x, y \in \mathbb{R}$  and  $a, b \in \mathbb{R}$  such that  $b \geq a > 0$ , define*

$$M = 1_{q < 1} (|x| + |y|)^{q+1} |a - b|^{q+1} + 1_{q \geq 1} (|x| + |y|)^{q+1} (|a| + |b|)^{q-1} |a - b|^2.$$

*Then there exists a constant  $C$  such that*

$$(5.1) \quad (x - y)(|x|^{q-1} x a^{q+1} - |y|^{q-1} y b^{q+1}) \geq \frac{1}{2} (x a - y b)(|x a|^{q-1} x a - |y b|^{q-1} y b) - C M$$

*Proof.* We have

$$\begin{aligned} & (x - y)(|x|^{q-1} x a^{q+1} - |y|^{q-1} y b^{q+1}) \\ &= (x b - y b)(|x a|^{q-1} x a - |y b|^{q-1} y b) + (x b - y b)|x a|^{q-1} x a (a/b - 1) \\ &= (x a - y b)(|x a|^{q-1} x a - |y b|^{q-1} y b) + x(b - a)(|x a|^{q-1} x a - |y b|^{q-1} y b) \\ & \quad + (x a - y b)|x a|^{q-1} x a (a/b - 1) + x(b - a)|x a|^{q-1} x a (a/b - 1) \end{aligned}$$

Using the fact that

$$(s - 1)(|s|^{q-1} s - 1) \sim (|s| + 1)^{q-1} |s - 1|^2 \sim (|s| + 1)^{1-q} ||s|^{q-1} s - 1|^2 \quad \forall s \in \mathbb{R},$$

we have

$$\begin{aligned} & (x - y)(|x|^{q-1} x a^{q+1} - |y|^{q-1} y b^{q+1}) \geq \frac{1}{2} (x a - y b)(|x a|^{q-1} x a - |y b|^{q-1} y b) \\ & \quad + \left[ \frac{c}{2} (|x a| + |y b|)^{1-q} ||x a|^{q-1} x a - |y b|^{q-1} y b|^2 + x(b - a)(|x a|^{q-1} x a - |y b|^{q-1} y b) \right] \\ & \quad + \left[ \frac{c}{2} (|x a| + |y b|)^{q-1} |x a - y b|^2 + (x a - y b)|x a|^{q-1} x a (a/b - 1) \right] - \frac{1}{b} (b - a)^2 |x|^{q+1} a^q. \end{aligned}$$

By Holder's inequality, we get the result.  $\square$

#### REFERENCES

- [CI16] Peter Constantin and Mihaela Ignatova. Critical SQG in bounded domains. *Ann. PDE*, 2(2):Art. 8, 42, 2016.
- [CMT94] Peter Constantin, Andrew J. Majda, and Esteban Tabak. Formation of strong fronts in the 2-D quasigeostrophic thermal active scalar. *Nonlinearity*, 7(6):1495–1533, 1994.
- [CT10] Xavier Cabré and Jिंगgang Tan. Positive solutions of nonlinear problems involving the square root of the Laplacian. *Adv. Math.*, 224(5):2052–2093, 2010.
- [CV10] Luis A. Caffarelli and Alexis Vasseur. Drift diffusion equations with fractional diffusion and the quasi-geostrophic equation. *Ann. of Math. (2)*, 171(3):1903–1930, 2010.
- [CW99] Peter Constantin and Jiahong Wu. Behavior of solutions of 2D quasi-geostrophic equations. *SIAM J. Math. Anal.*, 30(5):937–948, 1999.
- [DDN18] Nguyen Anh Dao, Jesus Ildefonso Díaz, and Quoc-Hung Nguyen. Generalized gagliardo-nirenberg inequalities using lorentz spaces, bmo, holder spaces and fractional sobolev spaces. *Nonlinear Analysis*, 173:146–153, 2018.
- [DM11] Frank Duzaar and Giuseppe Mingione. Gradient estimates via non-linear potentials. *Amer. J. Math.*, 133(4):1093–1149, 2011.
- [Kis10] A. Kiselev. Regularity and blow up for active scalars. *Math. Model. Nat. Phenom.*, 5(4):225–255, 2010.
- [KN09] A. Kiselev and F. Nazarov. A variation on a theme of Caffarelli and Vasseur. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 370(Kraevye Zadachi Matematicheskoi Fiziki i Smezhnye Voprosy Teorii Funktsii. 40):58–72, 220, 2009.
- [KNV07] A. Kiselev, F. Nazarov, and A. Volberg. Global well-posedness for the critical 2D dissipative quasi-geostrophic equation. *Invent. Math.*, 167(3):445–453, 2007.
- [Ngu14] Quoc-Hung Nguyen. Potential estimates and quasilinear parabolic equations with measure data. *Submitted for publication.*, 7, 2014.
- [SV03] Renming Song and Zoran Vondraček. Potential theory of subordinate killed Brownian motion in a domain. *Probab. Theory Related Fields*, 125(4):578–592, 2003.
- [SV19] Logan Stokols and Alexis Vasseur. Hölder regularity up to the boundary for critical sqg on bounded domains. *Preprint*, 2019.