

On the period of Lehn, Lehn, Sorger, and van Straten's symplectic eightfold

Nicolas Addington and Franco Giovenzana

Abstract

For the irreducible holomorphic symplectic eightfold Z associated to a cubic fourfold Y not containing a plane, we show that a natural Abel–Jacobi map from $H_{\text{prim}}^4(Y)$ to $H_{\text{prim}}^2(Z)$ is a Hodge isometry. We describe the full $H^2(Z)$ in terms of the Mukai lattice of the K3 category \mathcal{A} of Y . We give numerical conditions for Z to be birational to a moduli space of sheaves on a K3 surface or to $\text{Hilb}^4(\text{K3})$. We propose a conjecture on how to use Z to produce equivalences from \mathcal{A} to the derived category of a K3 surface.

Introduction

Beauville and Donagi [6] showed that if $Y \subset \mathbb{P}_{\mathbb{C}}^5$ is a smooth cubic fourfold, then the variety F of lines on Y is an irreducible holomorphic symplectic fourfold, deformation-equivalent to the Hilbert scheme of two points on a K3 surface. They showed moreover that the universal line $P \subset Y \times F$ induces a Hodge isometry

$$[P]_*: H_{\text{prim}}^4(Y, \mathbb{Z}) \xrightarrow{\sim} H_{\text{prim}}^2(F, \mathbb{Z}),$$

with the intersection pairing on the left-hand side and the opposite of the Beauville–Bogomolov–Fujiki pairing on the right.

The first author later refined this result by describing the full $H^2(F, \mathbb{Z})$ in terms of the Mukai lattice of the K3 category of Y , and characterized the cubics for which F is birational to the Hilbert scheme of two points on a K3 surface, or more generally to a moduli space of sheaves on a K3 surface [1].

In his survey paper [11], Hassett remarked that it would be useful to have analogous results for the symplectic eightfold Z constructed from twisted cubics on Y by Lehn, Lehn, Sorger, and van Straten [19]. This is the subject of the present paper. We begin with a statement on primitive cohomology.

Theorem 1. *Let Y be a smooth cubic fourfold not containing a plane. Let*

$$u: M \rightarrow Z$$

be the contraction from the ten-dimensional space of generalized twisted cubics on Y to the LLSvS symplectic eightfold. Let $C \subset Y \times M$ be the universal curve. Then the pullback

$$u^*: H^2(Z, \mathbb{Z}) \rightarrow H^2(M, \mathbb{Z})$$

is injective, and the map

$$[C]_*: H^4(Y, \mathbb{Z}) \rightarrow H^2(M, \mathbb{Z}) \tag{1}$$

restricts to a Hodge isometry

$$[C]_*: H_{\text{prim}}^4(Y, \mathbb{Z}) \xrightarrow{\sim} u^*(H_{\text{prim}}^2(Z, \mathbb{Z})), \tag{2}$$

with the intersection pairing on the left-hand side and the opposite of the Beauville–Bogomolov–Fujiki pairing on the right.

For a statement about the full $H^2(Z, \mathbb{Z})$, we must recall Kuznetsov’s K3 category

$$\mathcal{A} = \langle \mathcal{O}_Y, \mathcal{O}_Y(h), \mathcal{O}_Y(2h) \rangle^\perp \subset D^b(Y),$$

the Mukai lattice introduced in [3, §2],

$$K_{\text{top}}(\mathcal{A}) = \{[\mathcal{O}_Y], [\mathcal{O}_Y(h)], [\mathcal{O}_Y(2h)]\}^\perp \subset K_{\text{top}}(Y),$$

and the two special classes $\lambda_1, \lambda_2 \in K_{\text{top}}(\mathcal{A})$. For a recent account of the lattice theory of cubic fourfolds, see Huybrechts’ survey paper [16].

Theorem 2. *Continue the notation of Theorem 1. Let*

$$\Phi^K: K_{\text{top}}(\mathcal{A}) \subset K_{\text{top}}(Y) \rightarrow K_{\text{top}}(M)$$

be the map on topological K -theory induced by the Fourier–Mukai kernel $I_C^\vee(-3h) \in D^b(Y \times M)$. Then the map

$$c_1 \circ \Phi^K: K_{\text{top}}(\mathcal{A}) \rightarrow H^2(M, \mathbb{Z}) \tag{3}$$

restricts to a Hodge isometry

$$\langle \lambda_2 - \lambda_1 \rangle^\perp \xrightarrow{\sim} u^*(H^2(Z, \mathbb{Z})), \tag{4}$$

with the Euler pairing on the left-hand side and the opposite of the Beauville–Bogomolov–Fujiki pairing on the right.

One of the main ideas of [2] is that Z is a moduli space of complexes in \mathcal{A} containing the projections of \mathcal{O}_y for points $y \in Y$; the K-theory class of these complexes is $\lambda_2 - \lambda_1$, so (4) is natural in light of O’Grady’s description of the period of a moduli space of sheaves on a K3 surface [25]. Li, Pertusi, and Zhao developed this idea further using Bridgeland stability conditions in [21]; note that their class $2\lambda_1 + \lambda_2$ is related to our $\lambda_2 - \lambda_1$ by an auto-equivalence of \mathcal{A} . Statements similar to Theorem 2 appear in [21, Prop. 5.2] and [4, Thm. 29.2(2)]. We offer the present paper nonetheless because our proofs require less machinery, the statement in Theorem 1 is very classical and explicit, and we thought it worthwhile to articulate Theorem 3 and Conjecture 4 below.¹

Next we recall Hassett’s Noether–Lefschetz divisors \mathcal{C}_d in the moduli space \mathcal{C} of cubic fourfolds, indexed by an integer d satisfying

$$d > 6 \text{ and } d \equiv 0 \text{ or } 2 \pmod{6}. \quad (*)$$

Again we refer to the survey papers of Huybrechts [16] and Hassett [11]. From [3], [4], and [1] we know that the K3 category \mathcal{A} is equivalent to the derived category of coherent sheaves on a K3 surface, and the fourfold F is birational to a moduli space of sheaves on a K3 surface, if and only if the cubic $Y \in \mathcal{C}_d$ for some d satisfying

$$d/2 \text{ is not divisible by } 9 \text{ or any prime } p \equiv 2 \pmod{3}, \quad (**)$$

or equivalently,

$$d \text{ divides } 2n^2 + 2n + 2 \text{ for some } n \in \mathbb{Z}. \quad (**)$$

From [15] and [4] we know that the K3 category \mathcal{A} is equivalent to the derived category of *twisted* sheaves on a K3 surface, and F is birational to a moduli space of twisted sheaves, if and only if d satisfies the weaker condition

$$\begin{aligned} &\text{In the prime factorization of } d/2, \\ &\text{primes } p \equiv 2 \pmod{3} \text{ appear with even exponents.} \quad (**') \end{aligned}$$

On the other hand, from [1] we know that F is birational to the Hilbert scheme of two points on a K3 surface if and only if d satisfies the stronger condition

$$d \text{ is of the form } \frac{2n^2 + 2n + 2}{a^2} \text{ for some } n, a \in \mathbb{Z}. \quad (***)$$

¹After the first version of this paper was posted, Li, Pertusi, and Zhao informed us that the submitted version of [21] includes a statement analogous to our Theorem 3(c).

Let us introduce a new condition

$$d \text{ is of the form } \frac{6n^2 + 6n + 2}{a^2} \text{ for some } n, a \in \mathbb{Z}. \quad (***)'$$

It is strictly stronger than $(**)$ but incomparable to $(***)$: the first few ds satisfying $(***)$ are 14, 26, 38, 42, 62, and 86, whereas the first few ds satisfying $(***)'$ are 14, 38, 62, 74, and 86.

Theorem 3. *Let Y be a cubic fourfold not containing a plane — that is, not lying in \mathcal{C}_8 — so the symplectic eightfold Z is defined.*

- (a) *Z is birational to a moduli space of sheaves on a K3 surface if and only if $Y \in \mathcal{C}_d$ for some d satisfying $(**)$.*
- (b) *Z is birational to a moduli space of twisted sheaves on a K3 surface if and only if $Y \in \mathcal{C}_d$ for some d satisfying $(**')$.*
- (c) *Z is birational to the Hilbert scheme of four points on a K3 surface if and only if $Y \in \mathcal{C}_d$ for some d satisfying $(***)$.*

From [5, Thm. 1.2(c)] we know that if Z is birational to a moduli space of sheaves on a K3 surface, then Z is *isomorphic* to a moduli space of stable complexes on the same surface for some Bridgeland stability condition. This suggests an alternative approach to proving that \mathcal{A} is equivalent to the derived category of a K3 surface if and only if d satisfies $(**)$:²

Conjecture 4. *Suppose that $Y \in \mathcal{C}_d$ for some d satisfying $(**)$, so Z is isomorphic to a moduli space of Bridgeland-stable complexes on some K3 surface S . Let $\alpha \in \text{Br}(Z)$ be the obstruction to the existence of a universal complex on $Z \times S$, and let $U \in D^b(Z \times S, \alpha \boxtimes 1)$ be a twisted universal complex. Recall that Y is naturally embedded in Z , and consider the restriction $U|_{Y \times S}$, which can be untwisted because $\text{Br}(Y) = 0$. Then the functor $\mathcal{A} \rightarrow D^b(S)$ induced by $U|_{Y \times S}$ is an equivalence.*

We also expect a similar statement with $(**')$ and twisted K3 surfaces. To prove Conjecture 4 would require a careful analysis of the “wrong-way slices” $U|_{Z \times x}$ for $x \in S$, and of their restrictions to Y .

²On the other hand, if one assumes this fact (that \mathcal{A} is geometric if and only if d satisfies $(**)$), then Conjecture 4 seems to follow from [21, Prop. 5.7].

Acknowledgments. N.A. thanks the organizers of the Japanese–European Symposium on Symplectic Varieties at RIMS, Kyoto in October 2015, where these ideas first took shape. We thank M. Lehn, Ch. Lehn, G. Mongardi, J. C. Ottem, R. Takahashi, L. Giovenzana, and D. C. Veniani for valuable conversations, and A. Kuznetsov for detailed comments on an earlier version of the paper. N.A. was partially supported by NSF grant no. DMS-1902213. F.G. was supported by DFG research grants Le 3093/2-1 and Le 3093/2-2.

1 Proof of Theorems 1 and 2

1.1 Recollections and reductions

For a smooth cubic fourfold Y not containing a plane, Lehn, Lehn, Sorger, and van Straten considered the irreducible component $M \subset \mathrm{Hilb}^{3n+1}(Y)$ containing twisted cubics, which is a smooth tenfold [19, Thm. A]. They produced a contraction $u: M \rightarrow Z$ onto an irreducible holomorphic symplectic eightfold, and a copy of Y naturally embedded in Z [ibid., Thm. B]. Addington and Lehn showed that Z is deformation-equivalent to the Hilbert scheme of four points on a K3 surface [2].

Lemma 5. *The pullback map $u^*: H^2(Z, \mathbb{Z}) \rightarrow H^2(M, \mathbb{Z})$ is injective.*

Proof. The map u factors as the blowup $Z' \rightarrow Z$ along $Y \subset Z$, and a \mathbb{P}^2 -fibration $M \rightarrow Z'$, both of which induce injections on cohomology. Alternatively, u is surjective, so $u^*: H^*(Z, \mathbb{Q}) \rightarrow H^*(M, \mathbb{Q})$ is injective by [28, Lem. 7.28], and $H^2(Z, \mathbb{Z})$ is torsion-free because Z is simply-connected. \square

The map (1) is clearly a map of Hodge structures; let us argue that (3) is as well. For basics about the map on topological K-theory induced by a Fourier–Mukai kernel, and its compatibility with the more usual map on rational cohomology, we refer to [3, §2.1]. The Hodge structure on $K_{\mathrm{top}}(\mathcal{A}) \subset K_{\mathrm{top}}(Y)$ is pulled back via the Mukai vector $v: K_{\mathrm{top}}(Y) \rightarrow \bigoplus H^{2i}(Y, \mathbb{Q})$. We have a commutative diagram

$$\begin{array}{ccc} K_{\mathrm{top}}(\mathcal{A}) & \xrightarrow{\Phi^K} & K_{\mathrm{top}}(M) \\ v \downarrow & & v \downarrow \\ H^*(Y, \mathbb{Q}) & \xrightarrow{\Phi^H} & H^*(M, \mathbb{Q}), \end{array} \tag{5}$$

where Φ^K and Φ^H are the maps induced by $I_C^\vee(-3h) \in D^b(Y \times M)$. Thus we see that $v \circ \Phi^K$ is a map of Hodge structures. To get the map $c_1 \circ \Phi^K$ of (3), we first do $v \circ \Phi^K$, then multiply by $(\mathrm{td} M)^{-1/2}$ which preserves Hodge type, then take the degree-2 part.

Now it is a topological question whether the restrictions (2) and (4) take values in the subgroups claimed, and whether they are isometries, so it is enough to prove the claims for a single cubic Y . In a bit more detail, let $U \subset \mathbb{P}(H^0(\mathcal{O}_{\mathbb{P}^5}(3))) = \mathbb{P}^{55}$ be the Zariski open set of cubic fourfolds that are smooth and do not contain a plane. The construction of M and Z works in families, as discussed in the introduction to [19], so we get smooth families $\mathcal{Y} \rightarrow U$, $\mathcal{M} \rightarrow \mathcal{Z} \rightarrow U$, and $\mathcal{C} \subset \mathcal{M} \times_U \mathcal{Y}$. Then (2) and (4) are specializations of maps of local systems over U , so to prove the claims about them it is enough to do so at one point $Y \in U$.

In §§1.2–1.3 we do this for (4) when Y is a Pfaffian cubic fourfold whose associated K3 surface X has Picard rank 1. In §1.4 we do it for (2) when Y does not belong to any Noether–Lefschetz divisor.

1.2 Proof of Theorem 2 for a very general Pfaffian cubic

To a generic choice of six two-forms on \mathbb{C}^6 , Beauville and Donagi [6] associate a “Pfaffian” cubic fourfold $Y \subset \mathbb{P}^5$ and a K3 surface $X \subset \mathbb{P}^8$ that parametrizes a complete family of quartic scrolls on Y . In [2] the family is called $\Gamma \subset X \times Y$; it is generically 4-to-1 over Y . We will recall more details in the next section, but we do not need them yet.

Assuming that Y does not contain a plane, we continue to let M be the space of generalized twisted cubics on Y , $u: M \rightarrow Z$ the contraction onto the LLSvS eightfold, and $C \subset Y \times M$ the universal curve.

We will finish proving Theorem 2 under the assumption that X has Picard rank 1, which by our discussion above is enough to prove the theorem in general. We will construct a commutative diagram of Fourier–Mukai functors between $D^b(X)$ and $D^b(M)$, and deduce from it a commutative diagram of maps between lattices, most of which are isomorphisms, and ultimately compare the map (3) with the composition

$$[I_\xi]^\perp \subset K_{\text{top}}(X) \xrightarrow{I_\Xi^\vee} K_{\text{top}}(X^{[4]}) \xrightarrow{c_1} H^2(X^{[4]}, \mathbb{Z}),$$

which is known to be a Hodge isometry; here $\xi \subset X$ is a subscheme of length 4 and $\Xi \subset X \times X^{[4]}$ is the universal subscheme. For standard facts about Fourier–Mukai kernels, their adjoints, induced maps on cohomology, etc. we refer to Huybrechts’ book [13, Ch. 5].

We consider the convolution

$$T := I_\Gamma \circ I_C(2h) \in D^b(X \times M).$$

and recall some facts proved in [2, §3], with a few improvements:

- (a) There is a non-empty, Zariski open set $M_0 \subset M$ such that the restriction $T[1]|_{X \times M_0}$ is quasi-isomorphic to an M_0 -flat family of ideal sheaves of length-4 subschemes of X . We take M_0 as big as possible.
- (b) There is an open set $Z_0 \subset Z$ such that $M_0 = u^{-1}(Z_0)$.

Proof: From [2, Prop. 2] we know that M_0 is a union of fibers of u , so $M_0 = u^{-1}(Z_0)$ for some $Z_0 \subset Z$, maybe not open *a priori*. But u is surjective, so $Z \setminus Z_0 = u(M \setminus M_0)$, and u is proper, hence closed, so $Z \setminus Z_0$ is closed.

- (c) The classifying map $t': M_0 \rightarrow X^{[4]}$ descends to an open immersion $t: Z_0 \rightarrow X^{[4]}$.

Proof: From [2, §3] we know that t' descends to a map t that is injective. But an injective holomorphic map between complex manifolds of the same dimension is an open immersion by the proof of [10, Prop. on p. 19]; see also [26]. (So the smaller open set $Z_1 \subset Z_0$ of [2, §3] is unnecessary.)

The key to the present proof will be the following fact, whose proof we postpone to the next section:

Proposition 6. *If the K3 surface X has Picard rank 1, then the open set Z_0 contains Y , and its complement $Z \setminus Z_0$ has codimension at least 2 in Z .*

Thus $M \setminus M_0$ has codimension at least 2 in M as well, because u is a \mathbb{P}^2 -fibration over $Z \setminus Y$; and thus the inclusions $Z_0 \hookrightarrow Z$ and $M_0 \hookrightarrow M$ induce isomorphisms on H^2 .

The reader may wonder why we need Proposition 6, when we can say immediately that because Z and $X^{[4]}$ have nef canonical bundles, the birational map $Z \dashrightarrow X^{[4]}$ is biregular on an open set W containing Z_0 whose complement has codimension at least 2 [17, Cor. 3.54]. The reason is that we would not know anything about the restriction of T to $X \times u^{-1}(W)$, which we need in what follows.

Consider the functor $D^b(M) \rightarrow D^b(Y)$ induced by $I_C \otimes \mathcal{O}_Y(2h)$, the functor $D^b(Y) \rightarrow D^b(X)$ induced by I_Γ , their composition

$$D^b(M) \xrightarrow{I_C(2h)} D^b(Y) \xrightarrow{I_\Gamma} D^b(X), \quad (6)$$

and its left adjoint

$$D^b(X) \xrightarrow{I_\Gamma^\vee[2]} D^b(Y) \xrightarrow{I_C^\vee(-5h)[4]} D^b(M), \quad (7)$$

which it will later be convenient to rewrite as

$$D^b(X) \xrightarrow{I_\Gamma^\vee(-2h)[2]} D^b(Y) \xrightarrow{I_C^\vee(-3h)[4]} D^b(M). \quad (7')$$

We post-compose with the restriction

$$D^b(X) \xrightarrow{I_\Gamma^\vee(-2h)[2]} D^b(Y) \xrightarrow{I_C^\vee(-3h)[4]} D^b(M) \xrightarrow{i^*} D^b(M_0). \quad (8)$$

The first composition (6) is induced by T , so the left adjoint (7) or (7') is induced by $T^\vee[2]$, so the restriction (8) is induced by $T^\vee|_{X \times M_0}[2]$.

Because $T[1]|_{X \times M_0}$ is an M_0 -flat family of ideal sheaves with classifying map $t': M_0 \rightarrow X^{[4]}$, there is a line bundle \mathcal{L} on M_0 such that

$$T[1]|_{X \times M_0} = (1 \times t')^* I_\Xi \otimes \mathcal{L},$$

where again $\Xi \subset X \times X^{[4]}$ is the universal subscheme.³ So the composition (8) agrees with

$$D^b(X) \xrightarrow{I_\Xi^\vee[3]} D^b(X^{[4]}) \xrightarrow{t'^*} D^b(M_0) \xrightarrow{\otimes \mathcal{L}^\vee} D^b(M_0).$$

Because $t' = t \circ u$, this is

$$D^b(X) \xrightarrow{I_\Xi^\vee[3]} D^b(X^{[4]}) \xrightarrow{t^*} D^b(Z_0) \xrightarrow{u^*} D^b(M_0) \xrightarrow{\otimes \mathcal{L}^\vee} D^b(M_0).$$

Passing to topological K-theory [3, §2.1], and taking first Chern classes where shown, we get a diagram

$$\begin{array}{ccccccc} K_{\text{top}}(X) & \xrightarrow{-I_\Xi^\vee} & K_{\text{top}}(X^{[4]}) & \xrightarrow{t^*} & K_{\text{top}}(Z_0) & \xrightarrow{u^*} & K_{\text{top}}(M_0) \xrightarrow{[\mathcal{L}^\vee]} K_{\text{top}}(M_0) \\ & & c_1 \downarrow & & c_1 \downarrow & & c_1 \downarrow \\ & & H^2(X^{[4]}, \mathbb{Z}) & \xrightarrow{t^*} & H^2(Z_0, \mathbb{Z}) & \xrightarrow{u^*} & H^2(M_0, \mathbb{Z}) \dashrightarrow H^2(M_0, \mathbb{Z}), \end{array} \quad (9)$$

where there is no well-defined map to fill in the dashed arrow, and the injectivity of the lower u^* follows from Lemma 5 and Proposition 6.

The map $D^b(X^{[4]}) \xrightarrow{I_\Xi^\vee} D^b(X)$ takes the skyscraper sheaf of a point to I_ξ , where $\xi \subset X$ is a subscheme of length 4, so the left adjoint $I_\Xi^\vee[2]$ and its induced map on K-theory

$$K_{\text{top}}(X) \xrightarrow{I_\Xi^\vee} K_{\text{top}}(X^{[4]})$$

³For a careful proof that Hilbert schemes are moduli spaces of stable sheaves when $H^1(\mathcal{O}) = 0$, see [18, Lem. B.5.6].

takes classes in $[I_\xi]^\perp$ to classes of rank 0. Thus in the big diagram (9), if we replace $K_{\text{top}}(X)$ with the subspace $[I_\xi]^\perp$ then multiplying by $[\mathcal{L}^\vee]$ does not affect the first Chern class, so we can forget the last column of the diagram.

The composition

$$[I_\xi]^\perp \subset K_{\text{top}}(X) \xrightarrow{I_\Gamma^\vee} K_{\text{top}}(X^{[4]}) \xrightarrow{c_1} H^2(X^{[4]}, \mathbb{Z})$$

is a Hodge isometry by O'Grady's classic calculation [25]. Moreover the map t^* is an isomorphism on H^2 , so the big diagram (9) gives a Hodge isometry from $[I_\xi]^\perp \subset K_{\text{top}}(X)$ to $u^*(H^2(Z_0, \mathbb{Z})) \subset H^2(M_0, \mathbb{Z})$. Because $M \setminus M_0$ and $Z \setminus Z_0$ have codimension at least 2, this is the same as $u^*(H^2(Z, \mathbb{Z})) \subset H^2(M, \mathbb{Z})$.

If we take (8), pass to topological K-theory, and include first Chern classes, we get a diagram

$$\begin{array}{ccccccc} K_{\text{top}}(X) & \xrightarrow{I_\Gamma^\vee(-2h)} & K_{\text{top}}(Y) & \xrightarrow{I_C^\vee(-3h)} & K_{\text{top}}(M) & \xrightarrow{i^*} & K_{\text{top}}(M_0) \\ & & & & \downarrow c_1 & & \downarrow c_1 \\ & & & & H^2(M, \mathbb{Z}) & \xrightarrow{i^*} & H^2(M_0, \mathbb{Z}). \end{array} \quad (10)$$

In [2, Prop. 3] it is proved that the functor

$$D^b(X) \xrightarrow{I_\Gamma^\vee(-3h)[4]} D^b(Y)$$

is fully faithful, and its image is $\langle \mathcal{O}_Y(-h), \mathcal{O}_Y, \mathcal{O}_Y(h) \rangle^\perp$. In this paper we are taking $\mathcal{A} = \langle \mathcal{O}_Y, \mathcal{O}_Y(h), \mathcal{O}_Y(2h) \rangle^\perp$, so we should use the equivalence

$$D^b(X) \xrightarrow{I_\Gamma^\vee(-2h)[4]} \mathcal{A}.$$

We claim that the induced map on topological K-theory takes $[I_\xi]$ to our class $\lambda_2 - \lambda_1$. To see this, first observe that the functor $D^b(Y) \xrightarrow{I_\Gamma(-h)} D^b(X)$ takes the skyscraper sheaf of a point \mathcal{O}_y to some I_ξ , so its right adjoint $D^b(X) \xrightarrow{I_\Gamma^\vee(-2h)[4]} D^b(Y)$ takes I_ξ to the projection of \mathcal{O}_y into \mathcal{A} . In K-theory we have $[\mathcal{O}_y] = [\mathcal{O}_\ell(2)] - [\mathcal{O}_\ell(1)]$, where ℓ is a line on Y , so the class of the projection of \mathcal{O}_y is $\lambda_2 - \lambda_1$.

Thus in our second big diagram (10), the first map takes $[I_\xi]^\perp$ isometrically to $(\lambda_2 - \lambda_1)^\perp \subset K_{\text{top}}(\mathcal{A}) \subset K_{\text{top}}(Y)$. We conclude that the map (3) takes $(\lambda_2 - \lambda_1)^\perp$ isometrically to $u^*(H^2(Z, \mathbb{Z}))$, as desired.

1.3 Proof of Proposition 6

Now we recall the construction of the Pfaffian cubic fourfold Y , the K3 surface X , and the correspondence $\Gamma \subset X \times Y$ in detail. Fix a vector space $V \cong \mathbb{C}^6$ and a generic 6-dimensional subspace $L \subset \Lambda^2 V^*$. The cubic is

$$Y = \left\{ [\varphi] \in \mathbb{P}(L) \mid \text{rank}(\varphi) = 4 \right\} = \left\{ [\varphi] \in \mathbb{P}(L) \mid \varphi^3 = 0 \right\},$$

the K3 surface is

$$X = \left\{ [P] \in \text{Gr}(2, V) \mid \varphi|_P = 0 \text{ for all } \varphi \in L \right\},$$

and the correspondence is

$$\Gamma = \left\{ ([P], [\varphi]) \in X \times Y \mid P \cap \ker \varphi \neq 0 \right\}.$$

For a generic choice of L , both X and Y are smooth, X does not contain a line, and Y does not contain a plane.

In [2, Lem. 4] it is proved that Γ is generically 4-to-1 over Y . We improve this as follows:

Lemma 7. *If X does not contain a (-2) -curve then Γ is flat over Y .*

Proof. Let $\Gamma_\varphi \subset X$ be the scheme-theoretic fiber of Γ over a point $[\varphi] \in Y$. We will argue that Γ_φ is zero-dimensional; then by the proof of [2, Lem. 4] it has length 4.

Consider the Schubert cycle

$$\Sigma_\varphi = \left\{ [P] \in \text{Gr}(2, V) \mid P \cap \ker \varphi \neq 0 \right\}$$

of which Γ_φ is a linear section, and the normalization

$$\tilde{\Sigma}_\varphi = \left\{ (l, P) \in \mathbb{P}(\ker \varphi) \times \text{Gr}(2, V) \mid l \subset P \right\}.$$

We observe that $\tilde{\Sigma}_\varphi$ is a \mathbb{P}^4 -bundle over $\mathbb{P}(\ker \varphi) = \mathbb{P}^1$. First we claim that the preimage of Γ_φ in $\tilde{\Sigma}_\varphi$ meets each \mathbb{P}^4 fiber in at most one point, even scheme-theoretically: to see this, note that the \mathbb{P}^4 s are embedded linearly in $\text{Gr}(2, V) \subset \mathbb{P}(\Lambda^2 V)$, so if a linear section contained more than one point of a \mathbb{P}^4 it would contain a line, contradicting our hypothesis on X .

Next we claim that the preimage of Γ_φ in $\tilde{\Sigma}_\varphi$ meets at most finitely many \mathbb{P}^4 fibers. Otherwise it would meet them all, hence would give a section of this \mathbb{P}^4 -bundle over \mathbb{P}^1 , hence a smooth rational curve on X , again contradicting our hypothesis.

Thus the preimage of Γ_φ in $\tilde{\Sigma}_\varphi$ is zero-dimensional, so Γ_φ is as well. \square

Lemma 8. *Suppose that X has Picard rank 1, so in particular X contains no (-2) -curve and thus Γ induces a regular map $j: Y \rightarrow X^{[4]}$. Let \tilde{H} and B be the basis for $\text{Pic}(X^{[4]})$ used in [5, §13]. Then*

$$\begin{aligned} j^*B &= 9h \\ j^*\tilde{H} &= 14h \end{aligned}$$

where $h \in \text{Pic}(Y)$ is the hyperplane class.

Proof. By construction of $j: Y \rightarrow X^{[4]}$ we have a Cartesian diagram

$$\begin{array}{ccc} \Gamma & \longrightarrow & \Xi \\ \downarrow p & & \downarrow p' \\ Y & \xrightarrow{j} & X^{[4]}. \end{array}$$

By [20, Lem. 3.7] we have $B = -c_1(p'_*\mathcal{O}_\Xi)$, and thus

$$j^*B = -c_1(p_*\mathcal{O}_\Gamma).$$

Applying the Grothendieck–Riemann–Roch formula to the embedding $i: Y \hookrightarrow \mathbb{P}^5$, we get

$$i_* \text{ch}(p_*\mathcal{O}_\Gamma) = \text{ch}(i_*p_*\mathcal{O}_\Gamma) \cdot \text{td}(\mathcal{O}_{\mathbb{P}^5}(3)). \quad (11)$$

Let $0 \rightarrow \mathcal{P} \rightarrow \mathcal{O}_X \otimes V \rightarrow \mathcal{Q} \rightarrow 0$ be the restriction to X of the tautological bundle sequence on $\text{Gr}(2, V)$. In [2, §2] it is argued that $(1 \times i)(\Gamma) \subset X \times \mathbb{P}^5$ is the degeneracy locus of a map $\mathcal{P} \boxtimes \mathcal{O}_{\mathbb{P}^5} \rightarrow \mathcal{Q}^\vee \boxtimes \mathcal{O}_{\mathbb{P}^5}(1)$, of the expected dimension. Thus with the Eagon–Northcott resolution of $(1 \times i)_*\mathcal{O}_\Gamma$ and a lot of Schubert calculus, we can compute the right-hand side of (11). We carried out this computation by hand and checked it with the Schubert2 package of Macaulay2 [9]; we include our code in the ancillary file `Gamma.m2`. The result is

$$i_* \text{ch}(p_*\mathcal{O}_\Gamma) = 12h - 27h^2 + \frac{65}{2}h^3 - \frac{33}{2}h^4 + \frac{19}{8}h^5 \in H^*(\mathbb{P}^5, \mathbb{Q}).$$

Because $i_*(h) = 3h^2$, we read off $c_1(p_*\mathcal{O}_\Gamma) = -9h$, so

$$j^*B = 9h \in H^2(Y, \mathbb{Q}).$$

To find $j^*\tilde{H}$, let $H \subset X$ be a hyperplane section, and let $q': \Xi \rightarrow X$ be the natural projection. Then $p'_*q'^*\mathcal{O}_H$ is a sheaf supported on the locus of 4-tuples of which one is contained in H , that is, on \tilde{H} ; and this sheaf

has generic rank 1 on its support, so its first Chern class is \tilde{H} . Using the resolution

$$0 \rightarrow \mathcal{O}_X(-H) \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_H \rightarrow 0$$

we find that

$$i_* \operatorname{ch}(p_*(\mathcal{O}_\Gamma \otimes q^* \mathcal{O}_H)) = 42h^2 - 91h^3 + 56h^4 - \frac{35}{4}h^5.$$

Again the code is included in `Gamma.m2`. Taking the coefficient of h^2 and dividing by 3, we get $j^*(\tilde{H}) = 14h$ as desired. \square

Lemma 9. *Suppose that X has Picard rank 1, so again Γ induces a regular map $j: Y \rightarrow X^{[4]}$. Then every effective divisor on $X^{[4]}$ meets $j(Y)$.*

Proof. First we calculate the movable cone of $X^{[4]}$, applying Bayer and Macrì's [5, Prop. 13.1] with $d = 7$ and $n = 4$. Then $d(n - 1) = 21$ is not a perfect square; and $3X^2 - 7Y^2 = 1$ has no solution, as we see by reducing mod 3; so we are in case (c). The minimal positive solution to $X^2 - 21Y^2 = 1$ is $X = 55$, $Y = 12$. This satisfies $X \equiv 1 \pmod{3}$ rather than $X \equiv -1 \pmod{3}$, contradicting [5, eq. (33)], but the latter contains a typo which Debarre corrects in [8, Example 3.20]. Thus the movable cone is spanned by

$$\tilde{H} \quad \text{and} \quad 55\tilde{H} - 84B.$$

The same information can in principle be obtained from work of Markman [23], which does not use Bridgeland stability conditions, but [5] is more user-friendly.

Next we need to know that if D is an effective divisor on $X^{[4]}$ and M is movable, then $q(D, M) \geq 0$, where q is the Beauville–Bogomolov–Fujiki pairing. This follows from the proof of [12, Thm. 7]. In fact a result of Boucksom [7, Prop. 4.4] implies that the pseudo-effective cone is dual to the closure of the movable cone, but we do not need the full strength of this.

Now if an effective divisor D on $X^{[4]}$ does not meet $j(Y)$, then $j^*(D) = 0$, so from Lemma 8 we know that D is a multiple of $9\tilde{H} - 14B$. But we have

$$q(\tilde{H}, \tilde{H}) = 14 \quad q(\tilde{H}, B) = 0 \quad q(B, B) = -6,$$

and thus

$$q(9\tilde{H} - 14B, \tilde{H}) = 126 \quad q(9\tilde{H} - 14B, 55\tilde{H} - 84B) = -126,$$

so no multiple of $9\tilde{H} - 14B$ pairs non-negatively with both walls of the movable cone. \square

We conclude this section by deducing Proposition 6 from the lemmas above. From [2, §3] we know that the open immersion $t: Z_0 \rightarrow X^{[4]}$, restricted to $Y_0 := Y \cap Z_0$, agrees with the map $j: Y_0 \rightarrow X^{[4]}$ induced by I_Γ . If X has Picard rank 1 then $Y_0 = Y$ by Lemma 7. The open immersion $t: Z_0 \rightarrow X^{[4]}$ gives a birational map $Z \dashrightarrow X^{[4]}$. Because Z and $X^{[4]}$ are minimal models, this birational map is biregular on an open set W containing Z_0 whose complement has codimension at least 2 in both Z and $X^{[4]}$. If $Z \setminus Z_0$ has any component of codimension 1, then so too does $W \setminus Z_0$, and taking its closure in $X^{[4]}$ we get an effective divisor that does not meet $j(Y)$, contradicting Lemma 9.

1.4 Proof of Theorem 1 for a very general cubic

By our discussion in §1.1, to prove Theorem 1 it is enough to show that for a single cubic Y , the map (1) takes $H_{\text{prim}}^4(Y, \mathbb{Z})$ into $u^*(H_{\text{prim}}^2(Z, \mathbb{Z}))$ and respects the pairings. We deduce this from Theorem 2 when Y is not in any Noether–Lefschetz divisor.

Because Y is Noether–Lefschetz general, the transcendental lattice of $\langle \lambda_2 - \lambda_1 \rangle^\perp \subset K_{\text{top}}(\mathcal{A})$ is $\langle \lambda_1, \lambda_2 \rangle^\perp$. The transcendental lattice of $H^2(Z, \mathbb{Z})$ is contained in $H_{\text{prim}}^2(Z, \mathbb{Z})$, so the Hodge isometry (4) takes $\langle \lambda_1, \lambda_2 \rangle^\perp$ into $u^*(H_{\text{prim}}^2(Z, \mathbb{Z}))$; but they are primitive sublattices of the same rank, so (4) gives an isomorphism between them.

So we need to compare the action of (3) on $\langle \lambda_1, \lambda_2 \rangle^\perp$ with the action of (1) on $H_{\text{prim}}^4(Y, \mathbb{Z})$. In the diagram (5), we know from [3, Prop. 2.3] that v takes $\langle \lambda_1, \lambda_2 \rangle^\perp \subset K_{\text{top}}(\mathcal{A})$ isometrically to $H_{\text{prim}}^4(Y, \mathbb{Z}) \subset H^*(Y, \mathbb{Q})$. To finish proving Theorem 1, it remains to show that the maps

$$H_{\text{prim}}^4(Y, \mathbb{Z}) \rightarrow H^2(M, \mathbb{Q})$$

induced by $v(I_C^\vee(-3h)) \in H^*(Y \times M, \mathbb{Q})$ and $[C] \in H^6(Y \times M, \mathbb{Z})$ are the same. To see this, observe that

$$\begin{aligned} \text{ch}(I_C) &= 1 - [C] + \text{higher-order terms}, \\ \text{ch}(I_C^\vee) &= 1 + [C] + \text{higher-order terms}. \end{aligned}$$

What happens when we map a class $\alpha \in H_{\text{prim}}^4(Y)$ to $H^*(M, \mathbb{Q})$ using $v(I_C^\vee(-3h))$? First we multiply by $(\text{td } Y)^{1/2}$ and $\text{ch}(\mathcal{O}_Y(-3h))$, which are polynomials in the hyperplane class $h \in H^2(Y, \mathbb{Z})$ and thus have no effect on α . Then we pull up to $H^*(Y \times M, \mathbb{Q})$, multiply with $\text{ch}(I_C^\vee)$, and push down to $H^*(M, \mathbb{Q})$, which yields $[C]_*\alpha + \text{higher-order terms}$. Then we multiply by $(\text{td } M)^{1/2}$, which does not affect the leading term $[C]_*\alpha$. Then we take the degree-2 part, leaving $[C]_*\alpha$ as desired.

2 Proof of Theorem 3

We adapt an argument developed in [1] and [15, Prop. 4.1] and polished in [16, §3.2], staying especially close to the latter reference.

From Verbitsky's hyperkähler Torelli theorem [27, 14, 22] and Markman's work of the monodromy of manifolds of $\mathrm{K3}^{[n]}$ -type [23, §9], we know that if $n - 1$ is a prime power, then two manifolds M and M' of $\mathrm{K3}^{[n]}$ -type are birational (or bimeromorphic) if and only if there is a Hodge isometry $H^2(M, \mathbb{Z}) \cong H^2(M', \mathbb{Z})$. If $n - 1$ has two or more prime factors there is a more subtle statement, but this does not concern us since the eightfold Z is of $\mathrm{K3}^{[4]}$ -type.

By [16, Prop. 1.24 and Prop. 1.13], a cubic fourfold Y is in \mathcal{C}_d for some d satisfying $(**)$ if and only if the Mukai lattice $K_{\mathrm{top}}(\mathcal{A})$ is Hodge-isometric to the Mukai lattice $\tilde{H}(S, \mathbb{Z})$ of a K3 surface, and similarly with $(**')$ and the Mukai lattice $\tilde{H}(S, \alpha, \mathbb{Z})$ of a twisted K3 surface.

If $\varphi: K_{\mathrm{top}}(\mathcal{A}) \rightarrow \tilde{H}(S, \mathbb{Z})$ is a Hodge isometry, let $v = \varphi(\lambda_2 - \lambda_1)$, which is a primitive vector satisfying $v^2 = 6$. Then for a v -generic polarization $h \in \mathrm{Pic}(S)$, the moduli space $M := M_h(v)$ of h -stable sheaves with Mukai vector v is a smooth variety of $\mathrm{K3}^{[4]}$ -type, and satisfies $H^2(M, \mathbb{Z}) \cong v^\perp \subset \tilde{H}(S, \mathbb{Z})$. Thus

$$H^2(Z, \mathbb{Z}) \cong \langle \lambda_2 - \lambda_1 \rangle^\perp \cong v^\perp \cong H^2(M, \mathbb{Z}).$$

Conversely, any Hodge isometry $\langle \lambda_2 - \lambda_1 \rangle^\perp \cong v^\perp$ extends to a Hodge isometry $K_{\mathrm{top}}(\mathcal{A}) \rightarrow \tilde{H}(S, \mathbb{Z})$ taking $\lambda_2 - \lambda_1$ to $\pm v$ by [24, Cor. 1.5.2]: the discriminant group of the lattice v^\perp is $\mathbb{Z}/6$ whose automorphism group is ± 1 . With twisted K3 surfaces the argument is similar. This proves parts (i) and (ii) of Theorem 3.

To prove part (iii), by [1, Prop. 5] or by the proof of [16, Prop. 3.4(i)] we see that Z is birational to $S^{[4]}$ for some K3 surface S if and only if the algebraic lattice $K_{\mathrm{num}}(\mathcal{A}) \subset K_{\mathrm{top}}(\mathcal{A})$ contains a copy of the hyperbolic plane U which contains $\lambda_2 - \lambda_1$; or equivalently, if and only if there is a vector $w \in K_{\mathrm{num}}(\mathcal{A})$ with $\chi(w, w) = 0$ and $\chi(w, \lambda_2 - \lambda_1) = 1$.⁴

⁴Huybrechts [16] and many other authors define the Mukai pairing on $K_{\mathrm{top}}(\mathcal{A})$ as having the opposite sign from the Euler pairing, in order to agree with the Beauville–Bogomolov pairing on H^2 of various hyperkähler varieties, whereas Addington and Thomas [3, 1] define it having the same sign as the Euler pairing. In effect, the first convention regards $K_{\mathrm{top}}(\mathcal{A})$ as a Hodge structure of weight 2, the second as a Hodge structure of weight 0. In the calculation that follows, to avoid confusion, we only use Euler pairing and do not mention the Mukai pairing.

If there is such a w , let $L = \langle \lambda_1, \lambda_2, w \rangle \subset K_{\text{num}}(\mathcal{A})$, and let $n = \chi(w, \lambda_1)$, so the Gram matrix of the Euler pairing on L is

$$\begin{pmatrix} -2 & 1 & n \\ 1 & -2 & n+1 \\ n & n+1 & 0 \end{pmatrix}$$

Thus $\text{disc}(L) = 6n^2 + 6n + 2$. Let M be the saturation of L in $K_{\text{num}}(\mathcal{A})$, let a be the index of L in M , and let $d = \text{disc}(M)$; then $a^2 d = \text{disc}(L)$, so $d = (6n^2 + 6n + 2)/a^2$. By [3, Prop. 2.5] we have $Y \in \mathcal{C}_d$.

Conversely, suppose that $Y \in \mathcal{C}_d$ for some d of the form $(6n^2 + 6n + 2)/a^2$. We claim that $a \equiv 1 \pmod{6}$: to see this, observe that $6n^2 + 6n + 2$ is the norm of the primitive vector $(n, -n-1)$ in the A_2 lattice, hence satisfies $(**)$ by [16, Prop. 1.13(iii)], so a must be a product of primes $p \equiv 1 \pmod{3}$. Thus $d \equiv 2 \pmod{6}$; write $d = 6k + 2$ and $a = 3m + 1$. By [1, Lem. 9] there is an element $\tau \in K_{\text{num}}(\mathcal{A})$ such that $\langle \lambda_1, \lambda_2, \tau \rangle$ is a primitive sublattice on which the Gram matrix of the Euler pairing is

$$\begin{pmatrix} -2 & 1 & 0 \\ 1 & -2 & 1 \\ 0 & 1 & 2k \end{pmatrix}.$$

Then the vector

$$w := (m - n)\lambda_1 + (2m - n)\lambda_2 + a\tau$$

satisfies $\chi(w, w) = 0$ and $\chi(w, \lambda_2 - \lambda_1) = 1$, as desired.

References

- [1] N. Addington. On two rationality conjectures for cubic fourfolds. *Math. Res. Lett.*, 23(1):1–13, 2016. Also [arXiv:1405.4902](#).
- [2] N. Addington and M. Lehn. On the symplectic eightfold associated to a Pfaffian cubic fourfold. *J. reine angew. Math.*, 731:129–137, 2017. Also [arXiv:1404.5657](#).
- [3] N. Addington and R. Thomas. Hodge theory and derived categories of cubic fourfolds. *Duke Math. J.*, 163(10):1885–1927, 2014. Also [arXiv:1211.3758](#).
- [4] A. Bayer, M. Lahoz, E. Macrì, H. Nuer, A. Perry, and P. Stellari. Stability conditions in families. Preprint, [arXiv:1902.08184](#).

- [5] A. Bayer and E. Macrì. MMP for moduli of sheaves on K3s via wall-crossing: nef and movable cones, Lagrangian fibrations. *Invent. Math.*, 198(3):505–590, 2014. Also [arXiv:1301.6968](#).
- [6] A. Beauville and R. Donagi. La variété des droites d’une hypersurface cubique de dimension 4. *C. R. Acad. Sci. Paris Sér. I Math.*, 301(14):703–706, 1985. Also [math1.unice.fr/~beauvill/pubs/bd.pdf](#).
- [7] S. Boucksom. Divisorial Zariski decompositions on compact complex manifolds. *Ann. Sci. École Norm. Sup. (4)*, 37(1):45–76, 2004. Also [math/0204336](#).
- [8] O. Debarre. Hyperkähler manifolds. Preprint, [arXiv:1810.02087](#).
- [9] D. Grayson and M. Stillman. Macaulay2, a software system for research in algebraic geometry. Available at [www.math.uiuc.edu/Macaulay2/](#).
- [10] P. Griffiths and J. Harris. *Principles of algebraic geometry*. Wiley-Interscience [John Wiley & Sons], New York, 1978. Pure and Applied Mathematics.
- [11] B. Hassett. Cubic fourfolds, K3 surfaces, and rationality questions. In *Rationality problems in algebraic geometry*, volume 2172 of *Lecture Notes in Math.*, pages 29–66. Springer, Cham, 2016. Also [arXiv:1601.05501](#).
- [12] B. Hassett and Yu. Tschinkel. Moving and ample cones of holomorphic symplectic fourfolds. *Geom. Funct. Anal.*, 19(4):1065–1080, 2009. Also [arXiv:0710.0390](#), although there are significant corrections in the published version.
- [13] D. Huybrechts. *Fourier-Mukai transforms in algebraic geometry*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, Oxford, 2006.
- [14] D. Huybrechts. A global Torelli theorem for hyperkähler manifolds [after M. Verbitsky]. *Astérisque*, 348:Exp. No. 1040, x, 375–403, 2012. Séminaire Bourbaki: Vol. 2010/2011. Exposés 1027–1042. Also [arXiv:1106.5573](#).
- [15] D. Huybrechts. The K3 category of a cubic fourfold. *Compos. Math.*, 153(3):586–620, 2017. Also [1505.01775](#).

- [16] D. Huybrechts. Hodge theory of cubic fourfolds, their Fano varieties, and associated K3 categories. In *Lecture Notes of the Unione Matematica Italiana*, volume 26, pages 165–198. Springer, 2019. Also [arXiv:1811.02876](#).
- [17] J. Kollár and S. Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998.
- [18] A. Kuznetsov, Yu. Prokhorov, and C. Shramov. Hilbert schemes of lines and conics and automorphism groups of Fano threefolds. *Jpn. J. Math.*, 13(1):109–185, 2018. Also [arXiv:1605.02010](#).
- [19] Ch. Lehn, M. Lehn, Ch. Sorger, and D. van Straten. Twisted cubics on cubic fourfolds. *J. reine angew. Math.*, 731:87–128, 2017. Also [arXiv:1305.0178](#).
- [20] M. Lehn. Chern classes of tautological sheaves on Hilbert schemes of points on surfaces. *Invent. Math.*, 136(1):157–207, 1999. Also [math/9803091](#).
- [21] C. Li, L. Pertusi, and X. Zhao. Twisted cubics on cubic fourfolds and stability conditions. Preprint, [arXiv:1802.01134](#).
- [22] E. Looijenga. Teichmüller spaces and Torelli theorems for hyperkähler manifolds. *Math. Z.*, published online ahead of print, 2020. Also [arXiv:1912.02550](#).
- [23] E. Markman. A survey of Torelli and monodromy results for holomorphic-symplectic varieties. In *Complex and differential geometry*, volume 8 of *Springer Proc. Math.*, pages 257–322. Springer, Heidelberg, 2011. Also [arXiv:1101.4606](#).
- [24] V. Nikulin. Integer symmetric bilinear forms and some of their geometric applications. *Izv. Akad. Nauk SSSR Ser. Mat.*, 43(1):111–177, 238, 1979.
- [25] K. O’Grady. The weight-two Hodge structure of moduli spaces of sheaves on a K3 surface. *J. Algebraic Geom.*, 6(4):599–644, 1997. Also [alg-geom/9510001](#).
- [26] J.-P. Rosay. Injective holomorphic mappings. *Amer. Math. Monthly*, 89(8):587–588, 1982.

- [27] M. Verbitsky. Mapping class group and a global Torelli theorem for hyperkähler manifolds. *Duke Math. J.*, 162(15):2929–2986, 2013. Appendix A by Eyal Markman. Also [arXiv:0908.4121](#).
- [28] C. Voisin. *Hodge theory and complex algebraic geometry. I*, volume 76 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2002. Translated from the French original by Leila Schneps.

Nicolas Addington
 Department of Mathematics
 University of Oregon
 Eugene, OR 97403
 United States
adding@uoregon.edu

Franco Giovenzana
 Fakultät für Mathematik
 Technische Universität Chemnitz
 Reichenhainer Strasse 39
 09126 Chemnitz
 Germany
franco.giovenzana@mathematik.tu-chemnitz.de