

# Nonperturbative Flavor Breaking in Topological Susceptibility at Chiral Crossover

Mamiya Kawaguchi\*

*Department of Physics and Center for Field Theory and Particle Physics,  
Fudan University, 220 Handan Road, 200433 Shanghai, China*

Shinya Matsuzaki†

*Center for Theoretical Physics and College of Physics, Jilin University, Changchun, 130012, China*

Akio Tomiya‡

*RIKEN BNL Research center, Brookhaven National Laboratory, Upton, NY, 11973, USA*

We demonstrate that the QCD topological susceptibility nonperturbatively gets a significant contribution signaled by flavor-nonuniversal quark condensates at around the pseudo-critical temperature of the chiral crossover. It implies a remarkable flavor breaking in the axial anomaly as well as the QCD theta vacuum in high temperature QCD, which are almost flavor universal in the vacuum. A nontrivial flavor breaking is triggered by nonperturbative thermal loop corrections at around the chiral crossover, which is different from the trivial flavor violation just scaled by the quark mass ratio, observed at asymptotically high temperatures. This critical flavor violation cannot be dictated by the chiral perturbation theory with that lattice QCD usually compares, or the dilute instanton gas approximation based on that its astrophysical implications have conventionally been made. This would give an impact on the thermal history and the cosmological evolution of QCD axion including the estimate of the relic abundance as a dark matter candidate.

PACS numbers:

## I. INTRODUCTION

The topological susceptibility is a crucial probe in studying the QCD  $\theta$  vacuum structure and the axial anomaly. It is also important for QCD axion, which is postulated as an elegant solution to so-called the strong CP problem. In particular, when the QCD axion potential and mass are evaluated in the thermal history, the relic abundance as a cold dark matter today and the cosmological evolution of axion field in our Universe will be subject mainly to the temperature dependence of the topological susceptibility around the QCD phase transition epoch [1]. It would furthermore be a key quantity for hot QCD phenomena relevant to ongoing and designated heavy-ion collision experiments, such as physics induced by the topological charge fluctuation (i.e. susceptibility) closely tied with the presence of the QCD sphaleron [2–6]. Thus the QCD topological susceptibility has extensively been analyzed in multi-point of views with field theoretical, cosmological and astrophysical concerns.

The topological susceptibility  $\chi_{\text{top}}$  is defined as the curvature of the free energy of QCD with respect to the  $\theta$  at the QCD vacuum with  $\theta = 0$ . To our best knowledge, the key role for the  $\chi_{\text{top}}$  in real-life QCD is provided by an approximate chiral symmetry: with the approximate chiral symmetry for quarks, we can evaluate the  $\chi_{\text{top}}$  directly and generically from the QCD generating

functional, in relation to the quark condensates  $\langle \bar{q}q \rangle$  and the current quark mass  $m_q$  #1

$$\chi_{\text{top}} = \bar{m}^2(q) \sum_q \frac{\langle \bar{q}q \rangle}{m_q}, \quad \frac{1}{\bar{m}(q)} \equiv \sum_q \frac{1}{m_q}, \quad (1)$$

at the leading order of  $m_q$  (omitting the higher-order pseudoscalar susceptibility terms). The relation in Eq. (1) works even at finite temperature, because it is based on the Ward-Takahashi identity with the flavor singlet nature of the axial anomaly (as for the derivation of Eq. (1), see e.g. [8]). This is the intriguing formula having the nonperturbative correlation in QCD between the axial anomaly along with the  $\theta$  (dictated by the left hand side) and the chiral symmetry breaking (right hand side). The mediator for this correlation, the effective mass  $\bar{m}(q)$  reflects the flavor-singlet nature of the axial anomaly in QCD [9]: the axial anomaly detected by the  $\chi_{\text{top}}$  goes away if either of quarks get massless.

In the QCD vacuum with the lightest three flavors ( $u, d, s$  having the mass well below the intrinsic QCD scale  $\sim \mathcal{O}(1)$  GeV), the quark condensates are well degenerated ( $\langle \bar{u}u \rangle \simeq \langle \bar{d}d \rangle \simeq \langle \bar{s}s \rangle \equiv \Sigma$ ) as observed in the recent lattice simulation,  $\langle \bar{s}s \rangle / \langle \bar{l}l \rangle = 1.08 \pm 0.16$  ( $l = u, d$ ) [10]. In that case, the generic formula in Eq.(1) can be reduced to the Leutwyler-Smiluga (LS) relation [11]:  $\chi_{\text{top}}|_{\text{LS}} = \bar{m}(u, d, s)\Sigma$ , which is derived at the leading order in the chiral perturbation theory (ChPT). This LS relation indeed works well in the QCD

\*kawaguchi@fudan.edu.cn

†synya@jlu.edu.cn

‡akio.tomiya@riken.jp

#1 One may for instance refer to Appendix of the literature [7].

vacuum since the size of the three-flavor breaking is of  $\mathcal{O}(\frac{m_l}{m_s} \frac{\langle \bar{s}s \rangle}{\langle \bar{l}l \rangle}) = \mathcal{O}(10^{-2})$ , which supports the success of the ChPT also for this topological sector. The lattice QCD, with a physical pion mass realized and the continuum limit taken, have also shown consistency with the ChPT prediction [12–14].

The situation would get complicated at finite temperatures: at low temperatures enough, where the strange quark thermally decouples, the generic formula Eq.(1) implies the  $\chi_{\text{top}}$  (normalized by the vacuum value) will keep almost constant reflecting the residual two-flavor ( $u, d$ ) symmetry. This will be well described by the two-flavor ChPT. Given Eq.(1), going to a fairly high temperature regime, where all the three quarks may decouple from gluons, so the  $\chi_{\text{top}}$  (normalized as such) could asymptotically get damped with quadratic power in temperature, just like the ideal quark gas picture.

Highly nontrivial is, in particular, the response of the  $\chi_{\text{top}}$  at around the chiral phase transition: As has been observed in several analyses on hot lattice QCD [15, 16], thermal loop effects would cause a partial restoration of the chiral symmetry (chiral crossover), where only the lightest  $l$  quark condensates  $\langle \bar{l}l \rangle$  drop, but the strange quark's  $\langle \bar{s}s \rangle$  still keeps nonzero (or damps more slowly than the  $\langle \bar{l}l \rangle$ ), as will be seen in the panel (b) of Fig. 1.

This implies a nonperturbative flavor breaking in the quark condensates, hence would lead to a nonperturbative flavor breaking for the  $\chi_{\text{top}}$  at the hot QCD via the generic formula Eq.(1), which surely cannot be captured by the ChPT, or the ideal quark gas picture, or even the semi-classical dilute instanton gas picture [17, 18].

In fact, the recent lattice simulations [13, 14] indicate a substantial deviation from the ChPT prediction on the temperature dependence of the  $\chi_{\text{top}}$  around and/or above the chiral-crossover transition point. This discrepancy should be understood by some nonperturbative analysis based on the generic formula Eq.(1), though no definite implications to the  $\chi_{\text{top}}$  have been made in a sense of the quark-flavor symmetry at the crossover, even on the lattice QCD and chiral-effective model approaches.

In this Letter, we show that the QCD topological susceptibility indeed nonperturbatively gets a significant flavor violation signaled by a sizable strange-quark condensate-contribution at around the pseudo-critical temperature of the chiral crossover. It implies a non-trivial flavor breaking in the axial anomaly as well as the QCD theta vacuum at the chiral crossover.

We employ a nonperturbative analysis in a linear sigma model with the lightest three flavors, based on the Cornwall-Jackiw-Tomboulis (CJT) formalism [19], and compute thermal corrections to the effective potential coming from meson loops. The linear sigma model description allows us to go across the chiral phase boundary, in which the role of the chiral order parameter is played by sigma fields, as will be seen below.

Some nonperturbative analyses on the QCD topological susceptibility at finite temperatures have so far been done based on chiral effective models [20–22]. However,

no discussion on the correlation with quark condensates was made because their topological susceptibilities do not hold the flavor singlet form as in Eq.(1) (see also Eq.(2), for the flavor singlet condition), hence it seems to have been impossible to find the nonperturbative flavor breaking as addressed in the present Letter <sup>#2</sup>.

## II. CHIRAL EFFECTIVE MODEL DESCRIPTION

We begin by introducing a linear sigma model with the lightest three flavors, which has recently been proposed by including a possible axial-anomaly induced-flavor breaking term [24]. We parameterize the scalar- and pseudoscalar-meson nonets by a  $3 \times 3$  matrix field  $\Phi$  as  $\Phi = (\sigma_a + i\pi_a)T_a$ , where  $\sigma_a$  are the scalar fields and  $\pi_a$  are the pseudoscalar fields.  $T_a = \lambda_a/2$  ( $a = 0, 1, \dots, 8$ ) are the generators of  $U(3)$  normalized by  $\text{tr}[T_a T_b] = 1/2\delta_{ab}$ , where  $\lambda_{a=1, \dots, 8}$  are the Gell-Mann matrices with  $\lambda_0 = \sqrt{2/3}\mathbf{1}_{3 \times 3}$ . Under the chiral  $SU(3)_L \times SU(3)_R \times U(1)_A$  symmetry,  $\Phi$  transforms as  $\Phi \rightarrow g_A \cdot g_L \cdot \Phi \cdot g_R^\dagger$ , where  $g_{L,R} \in SU(3)_{L,R}$  and  $g_A \in U(1)_A$ . The three-flavor linear sigma model is thus written as [24],  $\mathcal{L} = \text{tr}[\partial_\mu \Phi \partial^\mu \Phi^\dagger] - V(\Phi)$ , with  $V(\Phi) = V_0 + V_{\text{anom}} + V_{\text{anom}} + V_{\text{SB}} + V_{\text{SB-anom}} \cdot V_0$  is an invariant part under the  $SU(3)_L \times SU(3)_R \times U(1)_A$  symmetry,  $V_0 = \mu^2 \text{tr}[(\Phi^\dagger \Phi)] + \lambda_1 \text{tr}[(\Phi^\dagger \Phi)^2] + \lambda_2 (\text{tr}[(\Phi^\dagger \Phi)])^2$ . The chiral  $SU(3)_L \times SU(3)_R$  invariant, but  $U(1)_A$  anomalous part is incorporated in  $V_{\text{anom}}$ :  $V_{\text{anom}} = -B (\det[\Phi] + \det[\Phi^\dagger])$ .

$V_{\text{SB}}$  denotes the explicit chiral symmetry breaking term, which is originated from the current quark mass matrix  $\mathcal{M} = \text{diag}[m_u, m_d, m_s]$  in the underlying QCD Lagrangian. This  $\mathcal{M}$  acts as a spurion field that transforms in the same way as the  $\Phi$  does. At the leading order of expansion in  $m_q$ , the  $V_{\text{SB}}$  thus takes the form  $V_{\text{SB}} = -\text{ctr}[\mathcal{M}\Phi^\dagger + \mathcal{M}^\dagger\Phi]$ . The parameter  $c$  necessarily comes along with the current quark masses ( $\mathcal{M}$ ) in physical observables, which reflects the renormalization scale ambiguity in defining quark condensates.

$V_{\text{SB-anom}}$  is the axial-anomaly induced-flavor breaking term. In the minimal flavor violation limit where single  $\mathcal{M}$  is only allowed to be inserted, the  $V_{\text{SB-anom}}$  is cast into the form  $V_{\text{SB-anom}} = -kc [\epsilon_{abc} \epsilon^{def} \mathcal{M}_d^a \Phi_e^b \Phi_f^c + \text{h.c.}]$ . This non-standard axial-anomaly interaction induced by the flavor violation potentially makes  $a_0(980)$  heavier than  $K_0^*(700)$  because by construction, it supplies the strange quark mass only for  $a_0(980)$ . Thus, this  $k$  coupling term plays a crucial role to realize the inverse mass hierarchy for scalar mesons below 1 GeV:  $m_{a_0(980)} \simeq m_{f_0(980)} > m_{K_0^*(700)} > m_{f_0(500)}$ , as has been shown in [24]. Moreover, intriguingly, the ax-

<sup>#2</sup> Another nonperturbative flavor breaking has been discussed based on Dyson–Schwinger analysis [23].

ial anomaly-induced flavor breaking directly contributes to the quark condensates (see Eq.(5)), and then provides the nonperturbative flavor violation for the chiral crossover.

In the QCD generating functional, the  $\theta$ -term can be rotated away by the  $U(1)_A$  rotation with the rotation angle  $\theta_q$  for quark fields. Then the  $\theta$ -dependence is fully transferred into the quark mass matrix,  $\bar{q}_L \mathcal{M} q_R$  ( $\bar{q}_R \mathcal{M}^\dagger q_L$ ) with  $\mathcal{M} = \text{diag}[m_u e^{i\theta_u}, m_d e^{i\theta_d}, m_s e^{i\theta_s}]$ . It is important to note here that the  $\theta_q$  is constrained by the flavor singlet condition (for  $\theta \ll 1$ ),

$$\theta_q = \frac{\bar{m}(u, d, s)}{m_q} \theta, \quad (2)$$

so that the  $U(1)_A$  anomaly keeps the flavor singlet nature [9]. We shall make a matching of this QCD generating functional with the one corresponding to the present linear sigma model. We are thus allowed to eliminate the  $\theta$ -dependence in the sigma field  $\Phi$  ( $\Phi^\dagger$ ), which has been introduced as the interpolating field of quark bilinear  $\bar{q}_R q_L$  ( $\bar{q}_L q_R$ ), so that the  $\theta$ -parameter should be entered in  $V_{\text{SB}}$  and  $V_{\text{SB-anom}}$  only via the quark mass matrix  $\mathcal{M}$ .

We assume the isospin symmetry,  $m_l = m_u = m_d \neq m_s$ , so that the vacuum expectation values are taken as  $\langle \Phi \rangle = \text{diag}[\bar{\Phi}_1, \bar{\Phi}_2, \bar{\Phi}_3]$  with  $\bar{\Phi}_1 = \bar{\Phi}_2 \neq \bar{\Phi}_3$ .

All the introduced parameters  $\mu^2, \lambda_{1,2}, B, cm_{l,s}, k$ , and  $\theta$  are taken to be real and positive.

### III. NONPERTURBATIVE FLAVOR BREAKING IN TOPOLOGICAL SUSCEPTIBILITY

To perform nonperturbative calculation of thermal loop contributions, we work on the CJT formalism [19] by using the imaginary time formalism. The meson propagators contributing to thermal loops are then treated as full propagators determined by stationary conditions for the CJT effective potential ( $V_{\text{CJT}}$ ), simultaneously with the vacuum expectation values  $\bar{\Phi}_{1,3}$ . Details are reported in [8], instead, we shall here just show the result on the numerical analysis of the CJT formalism.

The CJT potential is given by

$$V_{\text{CJT}} = V_{\text{tree}}(\bar{\Phi}_{1,3}) + V_{1\text{PI}}(\bar{\Phi}_{1,3}, S, P) + V_{2\text{PI}}(\bar{\Phi}_{1,3}, S, P), \quad (3)$$

where  $V_{\text{tree}}(\bar{\Phi}_{1,3})$  is the tree-level potential and  $V_{1(2)\text{PI}}(\bar{\Phi}_{1,3}, S, P)$  arise from one (two)-particle irreducible diagrams, in which all meson loop lines are drawn by the dressed-full propagators, denoted as  $S$  (for scalar mesons) and  $P$  (pseudoscalar mesons). In the present study we work in the Hartree approximation, corresponding to the leading order approximation in the large  $N$  expansion (only double-bubble diagrams survive for  $V_{2\text{PI}}(\bar{\Phi}_{1,3}, S, P)$ ). Since the CJT effective potential depends on the theta parameter  $\theta$  through the quark mass

matrix  $\mathcal{M}$  as noted above, the topological susceptibility is straightforwardly computed as

$$\chi_{\text{top}} = -\frac{\partial^2 V_{\text{CJT}}}{\partial \theta^2} \Big|_{\theta=0} = \left( \frac{2\langle \bar{l}l \rangle}{m_l} + \frac{\langle \bar{s}s \rangle}{m_s} \right) \bar{m}^2(u, d, s), \quad (4)$$

which is precisely in accordance with Eq.(1). Terms suppressed by the higher order in  $m_q$  have been omitted, as in Eq.(1). The quark condensates are calculated also through the  $V_{\text{CJT}}$  as

$$\begin{aligned} \langle \bar{l}l \rangle &= \frac{\partial V_{\text{CJT}}}{\partial m_l} = -2c(\bar{\Phi}_1 + 2k\bar{\Phi}_1\bar{\Phi}_3), \\ \langle \bar{s}s \rangle &= \frac{\partial V_{\text{CJT}}}{\partial m_s} = -2c(\bar{\Phi}_3 + 2k\bar{\Phi}_1^2), \end{aligned} \quad (5)$$

in which we will only take into account thermal correction parts <sup>#3</sup>.

Input parameters have been chosen by following the literature [24] as  $\mu^2 = 1.02 \times 10^4 \text{ MeV}^2$ ,  $\lambda_1 = 11.8$ ,  $\lambda_2 = 20.4$ ,  $cm_l = 6.11 \times 10^5 \text{ MeV}^3$ ,  $cm_s = 198 \times 10^5 \text{ MeV}^3$ ,  $B = 3.85 \times 10^3 \text{ MeV}$ ,  $k = 3.40 \text{ GeV}^{-1}$ , which is a set of the best-fit values and well reproduces the scalar and pseudoscalar spectroscopy up to the mass scale of 1 GeV at the vacuum [24].

As to the vacuum value of  $\chi_{\text{top}}$ , the literature predicts  $\chi_{\text{top}}(T=0) = 0.019(9)/\text{fm}^4$  [13], and  $\chi_{\text{top}}(T=0) = 0.0245(24)_{\text{stat}}(03)_{\text{flow}}(12)_{\text{cont}}/\text{fm}^4$  [14]. For the latter the first error is statistical, the second error is systematic error and the third error comes from changing the upper limit of the lattice spacing range in the fit. Our prediction  $\chi_{\text{top}}(T=0) \simeq 0.0263/\text{fm}^4$  is in agreement with both those lattice observations with their systematic errors taken into account.

By increasing temperature, we find the crossover “phase transition” for the chiral symmetry in [8]. Similar crossover phenomenon has also been observed in other three-flavor models based on the CJT formalism [26, 27]. The pseudo-critical temperature  $T_{pc}^*$  can simply be identified by  $d^2\langle \bar{l}l \rangle(T)/dT^2|_{T=T_{pc}^*} = 0$ , to be  $\simeq 215 \text{ MeV}$  in the present analysis. This crossover phenomenon is in a qualitative sense consistent with the current result of the lattice QCD with 2 + 1 flavors yielding the pseudo critical temperature  $\simeq 155 \text{ MeV}$  [28–31]. Note that the definition of our pseudo-critical temperature  $T_{pc}^*$  is different from the lattice QCD’s and has been estimated to

<sup>#3</sup> Although the vacuum part has the ultraviolet divergences and is subject to renormalization schemes, the thermal contribution to the loop integral is independent of the ultraviolet divergence and renormalization. Actually, the qualitative results obtained based on the CJT analysis at finite temperatures are fairly insensitive to renormalization schemes, as was studied in [25–27]. Thus, as far as qualitative features deduced from nonperturbative corrections are concerned, we may be almost free from the ultraviolet sensitivity, allowing to just work on the thermal corrections from the meson loops.

be larger than the lattice QCD value [28–31]: by construction of the present chiral effective model with single current quark mass matrix  $\mathcal{M}$  only included in operators, we cannot evaluate the chiral susceptibility, through which the pseudo-critical temperature in the lattice simulation is defined. However the temperature at which the maximum of the chiral susceptibility should reach is usually compatible with the inflection point of the chiral condensate in lattice QCD data, i.e. the point where  $d^2\langle\bar{l}l\rangle(T)/dT^2|_{T=T_{pc}} = 0$ . So, we may have quantitative comparison between them, to find about 30% deviation for almost whole relevant temperature regime. See the panel (a) of Fig. 1.

In the panel (b) of Fig. 1 we compare the  $\langle\bar{s}s\rangle(T)/\langle\bar{l}l\rangle(T)$  with the one corresponding to the free theory of quarks, in which the quarks behave non-interacting free-particles. The ratio of free-quark condensates  $\langle\bar{s}s\rangle_{\text{FT}}(T)/\langle\bar{l}l\rangle_{\text{FT}}(T)$  is perturbatively obtained from the thermal one-loop calculation. For the low temperature regions where  $T < m_s \sim 100$  MeV, the free-theory (FT) quark-condensate ratio  $\langle\bar{s}s\rangle_{\text{FT}}(T)/\langle\bar{l}l\rangle_{\text{FT}}(T)$  keeps the value below  $m_s/m_l \simeq 27$ , because of the naive Boltzmann suppression for the strange quark contribution. After passing  $T \simeq m_s \sim 100$  MeV, the  $\langle\bar{s}s\rangle_{\text{FT}}(T)/\langle\bar{l}l\rangle_{\text{FT}}(T)$  asymptotically approaches  $m_s/m_l$ , which merely reflects the trivial and overall flavor-breaking just by quark masses:  $\langle\bar{q}q\rangle \sim m_q T^2$ . In contrast, in whole low-temperature regions  $T < T_{pc}^* \simeq 215$  MeV the CJT analysis exhibits a gigantic suppression

for the quark-condensate ratio more than the Boltzmann suppression, to respect the flavor symmetry, i.e., the vacuum value  $\sim 1$ . This flavor symmetric behavior is consistent with the conventional three-flavor ChPT observation, in which the vectorial  $SU(3)$  flavor symmetry cannot explicitly be violated at the leading order of the chiral expansion based on the nonlinear-sigma model setup, so that  $\langle\bar{s}s\rangle_{\text{ChPT}}(T)/\langle\bar{l}l\rangle_{\text{ChPT}}(T) \simeq 1$  even including the next-to leading order corrections. As the temperature further increases, the CJT analysis non-perturbatively undergoes the chiral crossover around the pseudo-critical temperature  $T_{pc}^* \simeq 215$  MeV, where the light quark condensate  $\langle\bar{l}l\rangle(T)$  starts to drop more efficiently than the strange quark condensate  $\langle\bar{s}s\rangle(T)$ . Consequently, the quark-condensate ratio rapidly starts to grow from  $T \simeq T_{pc}^*$ . This is a nontrivial flavor breaking, essentially different from the trivial FT flavor violation just by  $m_s/m_l$ . In the end, at around  $T \simeq 600$  MeV, the quark-condensate ratio asymptotically merges with the FT yielding the trivial-flavor breaking value  $m_s/m_l$ . This would imply that the present linear sigma model would converge to an ideal-quark gas picture consistently with the asymptotic free nature of the underlying QCD. Thus we can conclude that what we call a nonperturbative flavor breaking is certainly a nonperturbative output in association with the characteristic chiral crossover phenomenon.




FIG. 1: The temperature dependence of the quark condensates. (a): comparison of the subtracted chiral condensate  $\Delta_{l,s}(T) = \langle\bar{l}l\rangle - \frac{2m_l}{m_s}\langle\bar{s}s\rangle$  with the lattice QCD data [32]; (b): the ratio of quark condensates as a function of the temperature,  $\langle\bar{s}s\rangle(T)/\langle\bar{l}l\rangle(T)$ , which implies the significance of the flavor breaking in the quark condensates as the temperature gets higher than the pseudo-critical temperature  $T_{pc}^* \simeq 215$  MeV while  $\langle\bar{s}s\rangle(0)/\langle\bar{l}l\rangle(0) \simeq 1.18$  [24].

To extract the strange quark contribution in the topological susceptibility, in Fig. 2 we show the temperature dependence of  $\chi_{\text{top}}(T)$  normalized to the one in the three-

flavor universal limit,  $\chi_{\text{top}}^{\text{3fl}} = \left(\frac{2}{m_l} + \frac{1}{m_s}\right)^{-1} \langle\bar{l}l\rangle(T)$ . We see that due to the sizable strange quark condensate as seen from the panel (b) of Fig. 1, the topological sus-




FIG. 2: The temperature dependence of the  $\chi_{top}(T)/\chi_{top}^{3fl}(T)$  compared with the free-quark theory,  $\chi_{top}^{(FT)}(T)/\chi_{top}^{(FT)3fl}(T)$ .

ceptibility is rapidly enhanced from  $T \simeq T_{pc}^*$ . Eventually, after arriving at the high temperature regions where  $\langle \bar{s}s \rangle(T)/\langle \bar{l}l \rangle(T)$  reaches the trivial-flavor breaking value  $m_s/m_l$ , the  $\chi_{top}(T)/\chi_{top}^{3fl}(T)$  merges with the quark-FT regime, to asymptotically converge to  $\chi_{top}(T)/\chi_{top}^{3fl}(T) \simeq 3/(2 + m_l/m_s) \simeq 1.5$ . Thus, the topological susceptibility gets the nonperturbative flavor breaking at around the chiral crossover, which is manifestly different from the trivial-flavor breaking as seen in the quark-FT.

Actually, it is obvious from the robust formula in Eq.(1) that the strange quarks contribute to the  $\chi_{top}$ , and, at high enough temperatures  $T \gg T_{pc}^*$ , all flavors exhibit an equal importance as also deduced from the FT scaling in Fig. 1. At low temperatures  $T < T_{pc}^*$ , on the other hand, Eq.(1) and the ChPT tells us that the contribution of the strange quarks is suppressed by a factor of approximately  $m_s/m_l (\sim 27)$ . What is nontrivial is that strange quarks start to become important in the  $\chi_{top}$  for  $T \gtrsim T_{pc}^*$ , as one can expect from Fig. 1 together with Eq.(1), that has presently been made clarified in Fig. 2, and we have dubbed the nonperturbative flavor violation in the  $\chi_{top}$ .

The panel (a) of Fig. 3 shows the  $\chi_{top}$  normalized to the vacuum value, in comparison with the ChPT prediction up to the next-leading order (one loop) [40] and the recent lattice data with  $2 + 1$  (+1) flavors having a physical pion mass and the continuum limit being taken [13, 14]. Since the linear sigma model including only mesons tends to systematically give a larger  $T_{pc}^*$  than theories with quarks and mesons including lattice QCD, comparison with the lattice data has been made by normalizing  $T$  by  $T_{pc}^*$ , so that such a systematic uncertainty is reduced. Note also that the linear-sigma model picture is thought to break down at high  $T$ , where degrees of freedom of quarks (and gluons) become important. Such a  $T$  may be interpreted as  $T \sim m_q^{\text{constituent}} \sim 330$  MeV, where the latter denotes mass of constituent up and down

quarks (given roughly by one-third of proton mass). In terms of the normalized  $T$ , it corresponds to  $T/T_{pc}^* \sim 1.5$ . Therefore, the present model prediction may be operative only up until  $T/T_{pc}^* \sim 1.5$ , above which quarks and gluons would govern the system. As seen from Figs.1 and 2, however, it suffices for  $T/T_{pc}^* \lesssim 1.5$  to explore the nonperturbative flavor violation in the  $\chi_{top}$ , which starts to be eminent at around the chiral crossover  $T/T_{pc}^* \gtrsim 1$ .

Remarkably, the predicted  $T/T_{pc}^*$  dependence (denoted by “CJT” in the figure) is actually slower-damping, and is overall consistent with both two lattice QCD data [13, 14] #4, which is not realized by the ChPT. This would manifest the importance of nonperturbative thermal contribution including the enhanced flavor breaking by  $\langle \bar{s}s \rangle(T)/\langle \bar{l}l \rangle(T) \gg 1$  above  $T_{pc}^*$ , as depicted in Fig. 1.

The CJT result favors the ChPT at  $T/T_{pc}^* \lesssim 1$ , which implies that the chiral symmetry for light quarks might be essential in the low-temperature region, as discussed in the literature [40] #5. Beyond the ChPT-governed domain, above  $T_{pc}$  the strange quark condensate would serve as an important source to develop the topological susceptibility, as the consequence of the nonperturbative flavor breaking.

The predicted curve also coincides with the dilute instanton gas approximation [17, 18] for  $T/T_{pc}^* \lesssim 1.5$ , with which the lattice result in Ref.[14] is shown to be fully consistent. A substantial deviation starts in higher  $T$ , which might be because of lack of quarks (and gluons) in the present model.

In the panel (b) of Fig. 3, we also plot the temperature dependence of an unnormalized susceptibility  $\chi_{top}^{1/4}(\text{MeV})$ , where comparison with some recent lattice simulations with the extrapolation to the continuum limit is available for  $T/T_{pc}^* \leq 1.5$  [14, 42] #6. We see that at around the chiral crossover  $T/T_{pc}^* \sim 1 - 1.5$  the CJT result is in quite good agreement with those continuum extrapolated data. Although qualitatively having agreement, for  $T/T_{pc}^* \gtrsim 1.5$  the CJT result tends to predict a somewhat larger  $\chi_{top}^{1/4}(\text{MeV})$ . This discrepancy gets more larger as  $T/T_{pc}^*$  gets larger and larger, which agrees with the expected validity of the present linear sigma model, as noted above.

To reconcile the gap observed when  $T/T_{pc}^* \gtrsim 1.5$ , we may note that model parameters in the chiral effective model can actually have the intrinsic-temperature dependence, which could mimic a part of nondecoupling effects from integrating out quarks and/or gluons. For instance, as noted in [44], the instanton study predicts

#4 Note that in Ref. [13], data have been taken only for two different lattice spacings. So, continuum extrapolation in Ref. [13] may not reliably be performed to give predictions.

#5 Note that the ChPT-governed domain is intact even if one includes the next-to-next-to-leading order (NNLO) correction in the ChPT analysis [41].

#6 Another continuum extrapolated result in Ref. [43] was provided for  $T/T_{pc}^* \gtrsim 3$ .

the parameter  $B$  in the  $U(1)_A$  anomalous part to have an intrinsic-temperature dependence, which is relevant to the QCD topological structure at high temperatures. Thus, such an intrinsic-temperature dependence might pull the CJT result down, to coincide with the lattice results in the continuum limit [14, 42, 43].




FIG. 3: (a): The comparison of the  $\chi_{\text{top}}(T)/\chi_{\text{top}}(T=0)$  (labeled as “CJT” in the plot) with the ChPT prediction up to the next-to-leading order (NLO) (one-loop) [40] and the lattice QCD data [13, 14]. We have taken  $T_{\text{pc}}^* = 215$  MeV for the CJT result, and  $T_{\text{pc}} = 155$  MeV for the lattice simulations and the ChPT prediction. The band corresponds to the continuum extrapolation of the lattice QCD data [13], which is estimated by the function  $\chi(a, T)/\chi(a, T=0) = D_0(1 + D_1 a^2)(T/T_c)^{D_2}$  with  $D_0 = 1.17$ ,  $D_1 = 0$ ,  $D_2 = -2.71$  and  $T_c = 155$  MeV. (b): The unnormalized topological susceptibility,  $\chi_{\text{top}}^{1/4}$  (MeV), versus temperature, in comparison with the continuum extrapolated results of the recent lattice simulations. The band corresponds to the continuum extrapolation of the lattice QCD data [42].

#### IV. CONCLUSION

In conclusion, we showed that the strange quark contribution to the topological susceptibility becomes eminent around the chiral crossover, which cannot be detected by the chiral perturbation theory and can clearly be distinguished from the trivial flavor breaking in high enough temperatures, just set by the quark mass weight. Our

finding would be crucial to understand the hot QCD with the theta parameter, at around the chiral crossover as well as the associated  $U(1)_A$  anomaly. In particular, the lattice QCD measurement of strange quark condensate at around and over the pseudo critical temperature would be crucial to check how the nonperturbative flavor breaking is critically operative in the topological susceptibility. This critical phenomenon should directly be observed in the future lattice QCD with appropriate accuracy.

Another remark is that the topological susceptibility dumps slowly due to the nondecoupled nonperturbative strange quark condensate which implies the delay of the effective  $U(1)_A$  restoration. Thus the role of this nondecoupled strange quark condensate may account for the tension in the effective restoration of the  $U(1)_A$  symmetry currently observed on lattices with the two-flavor [33–36] and the 2+1 flavor [37–39] near the chiral limit.

Our proposal has been explicitized using some specific chiral effective model, especially including an axial anomaly-induced flavor breaking term. We have checked that even without this speciality, our finding is substantially unchanged, which in details is reported in [8].

The nonperturbative flavor breaking, especially, the significance of the strange quark condensate, in the topological susceptibility at around the pseudo critical temperature would give an impact on applications to QCD axion cosmological models. The epoch for QCD axion to start rolling and oscillate as well as the position in the potential is crucial to estimate the relic abundance of the axion as a dark matter today. This is subject to the temperature (or time) dependence of the  $\chi_{\text{top}}$  (which corresponds to the potential height) at around and/or above the chiral crossover boundary. As recently investigated in [45], below and above this crossover boundary, it has simply been assumed so far to apply the two-flavor ChPT and dilute instanton gas descriptions, separately, in evaluating the  $\chi_{\text{top}}$  (with imposing a continuity condition between two domains through macroscopic thermodynamics quantities). Our present work would improve or refine this existing approach by including the strange quark contribution with use of the linear sigma model description, instead of the dilute instanton gas, at around and/or above the chiral crossover, and would provide a complementary evaluation of the QCD axion cosmology, with higher reliability. Detailed study deserves to another publication.

#### Acknowledgements

We are grateful to Massimo D’Elia, Xu-Guang Huang and Robert Pisarski for useful comments. This work was supported in part by the National Science Foundation of China (NSFC) under Grant No. 11747308 and 11975108 and the Seeds Funding of Jilin University (S.M.). M.K. and A.T. thank for the hospitality of Center for Theoretical Physics and College of Physics, Jilin University where the present work has been partially done.



The work of A.T. was supported by the RIKEN Special Postdoctoral Researcher program.

- 
- [1] For a review, e.g., see the following paper:  
J. E. Kim and G. Carosi, *Rev. Mod. Phys.* **82**, 557 (2010)  
Erratum: [*Rev. Mod. Phys.* **91**, no. 4, 049902 (2019)]  
doi:10.1103/RevModPhys.91.049902, 10.1103/RevMod-  
Phys.82.557 [arXiv:0807.3125 [hep-ph]], and references  
therein. One may refer also to related recent papers that  
cite this paper.
- [2] N. S. Manton, *Phys. Rev. D* **28**, 2019 (1983);  
F. R. Klinkhamer and N. S. Manton, *Phys. Rev. D* **30**,  
2212 (1984).
- [3] L. D. McLerran, E. Mottola and M. E. Shaposhnikov,  
*Phys. Rev. D* **43**, 2027 (1991).
- [4] P. Arnold, D. Son and L. G. Yaffe, *Phys. Rev. D* **55**, 6264  
(1997) [arXiv:hep-ph/9609481].
- [5] P. Huet and D. T. Son, *Phys. Lett. B* **393**, 94 (1997)  
[arXiv:hep-ph/9610259].
- [6] D. Bodeker, *Phys. Lett. B* **426**, 351 (1998) [arXiv:hep-  
ph/9801430].
- [7] Y. Y. Mao *et al.* [TWQCD Collaboration], *Phys. Rev. D*  
**80**, 034502 (2009) doi:10.1103/PhysRevD.80.034502  
[arXiv:0903.2146 [hep-lat]].
- [8] M. Kawaguchi, S. Matsuzaki and A. Tomiya,  
[arXiv:2005.07003 [hep-ph]].
- [9] V. Baluni, *Phys. Rev. D* **19**, 2227 (1979).  
doi:10.1103/PhysRevD.19.2227
- [10] C. McNeile, A. Bazavov, C. T. H. Davies, R. J. Dow-  
dall, K. Hornbostel, G. P. Lepage and H. D. Trot-  
tier, *Phys. Rev. D* **87**, no. 3, 034503 (2013)  
doi:10.1103/PhysRevD.87.034503 [arXiv:1211.6577 [hep-  
lat]].
- [11] H. Leutwyler and A. V. Smilga, *Phys. Rev. D* **46**, 5607  
(1992). doi:10.1103/PhysRevD.46.5607
- [12] S. Aoki *et al.* [Flavour Lattice Averaging Group], *Eur.*  
*Phys. J. C* **80**, no. 2, 113 (2020) doi:10.1140/epjc/s10052-  
019-7354-7 [arXiv:1902.08191 [hep-lat]].
- [13] C. Bonati, M. D'Elia, M. Mariti, G. Martinelli,  
M. Mesiti, F. Negro, F. Sanfilippo and G. Villadoro,  
*JHEP* **1603**, 155 (2016) doi:10.1007/JHEP03(2016)155  
[arXiv:1512.06746 [hep-lat]].
- [14] S. Borsanyi *et al.*, *Nature* **539**, no. 7627, 69 (2016)  
doi:10.1038/nature20115 [arXiv:1606.07494 [hep-lat]].
- [15] M. Cheng *et al.*, *Phys. Rev. D* **77**, 014511 (2008)  
doi:10.1103/PhysRevD.77.014511 [arXiv:0710.0354 [hep-  
lat]].
- [16] A. Gómez Nicola and J. Ruiz de Elvira, *JHEP* **1603**, 186  
(2016) doi:10.1007/JHEP03(2016)186 [arXiv:1602.01476  
[hep-ph]].
- [17] D. J. Gross, R. D. Pisarski and L. G. Yaffe, *Rev. Mod.*  
*Phys.* **53**, 43 (1981). doi:10.1103/RevModPhys.53.43
- [18] T. Schäfer and E. V. Shuryak, *Rev. Mod. Phys.*  
**70**, 323 (1998) doi:10.1103/RevModPhys.70.323 [hep-  
ph/9610451].
- [19] J. M. Cornwall, R. Jackiw and E. Tomboulis, *Phys. Rev. D*  
**10**, 2428 (1974). doi:10.1103/PhysRevD.10.2428
- [20] Y. Jiang, T. Xia and P. Zhuang, *Phys. Rev. D* **93**,  
no. 7, 074006 (2016) doi:10.1103/PhysRevD.93.074006  
[arXiv:1511.06466 [hep-ph]].
- [21] K. Fukushima, K. Ohnishi and K. Ohta, *Phys. Rev. C*  
**63**, 045203 (2001) doi:10.1103/PhysRevC.63.045203  
[nucl-th/0101062].
- [22] Y. Jiang and P. Zhuang, *Phys. Rev. D* **86**, 105016 (2012)  
doi:10.1103/PhysRevD.86.105016 [arXiv:1209.0507 [hep-  
ph]].
- [23] D. Horvatić, D. Kekez and D. Klabučar, *Uni-  
verse* **5**, no.10, 208 (2019) doi:10.3390/universe5100208  
[arXiv:1909.09879 [hep-ph]].
- [24] Y. Kuroda, M. Harada, S. Matsuzaki and D. Jido,  
arXiv:1910.09146 [hep-ph].
- [25] J. T. Lenaghan and D. H. Rischke, *J. Phys. G*  
**26**, 431 (2000) doi:10.1088/0954-3899/26/4/309 [nucl-  
th/9901049].
- [26] J. T. Lenaghan, D. H. Rischke and J. Schaffner-  
Bielich, *Phys. Rev. D* **62**, 085008 (2000)  
doi:10.1103/PhysRevD.62.085008 [nucl-th/0004006].
- [27] D. Roder, J. Ruppert and D. H. Rischke, *Phys. Rev. D*  
**68**, 016003 (2003) doi:10.1103/PhysRevD.68.016003  
[nucl-th/0301085].
- [28] Y. Aoki, G. Endrodi, Z. Fodor, S. D. Katz and K. K. Szabo,  
*Nature* **443**, 675 (2006) doi:10.1038/nature05120  
[hep-lat/0611014].
- [29] S. Borsanyi *et al.* [Wuppertal-Budapest Collaboration],  
*J. Phys. Conf. Ser.* **316**, 012020 (2011) doi:10.1088/1742-  
6596/316/1/012020 [arXiv:1109.5032 [hep-lat]].
- [30] H. T. Ding, F. Karsch and S. Mukherjee, *Int.*  
*J. Mod. Phys. E* **24**, no. 10, 1530007 (2015)  
doi:10.1142/S0218301315300076 [arXiv:1504.05274 [hep-  
lat]].
- [31] H. T. Ding, arXiv:2002.11957 [hep-lat].
- [32] Y. Aoki, S. Borsanyi, S. Durr, Z. Fodor, S. D. Katz,  
S. Krieg and K. K. Szabo, *JHEP* **06**, 088 (2009)  
doi:10.1088/1126-6708/2009/06/088 [arXiv:0903.4155  
[hep-lat]].
- [33] G. Cossu, S. Aoki, H. Fukaya, S. Hashimoto, T. Kaneko,  
H. Matsufuru and J. I. Noaki, *Phys. Rev. D* **87**,  
no.11, 114514 (2013) [erratum: *Phys. Rev. D* **88**,  
no.1, 019901 (2013)] doi:10.1103/PhysRevD.87.114514  
[arXiv:1304.6145 [hep-lat]].
- [34] A. Tomiya, G. Cossu, S. Aoki, H. Fukaya, S. Hashimoto,  
T. Kaneko and J. Noaki, *Phys. Rev. D* **96**,  
no.3, 034509 (2017) doi:10.1103/PhysRevD.96.034509  
[arXiv:1612.01908 [hep-lat]].
- [35] K. Suzuki *et al.* [JLQCD], *PoS CD2018*, 085 (2019)  
doi:10.22323/1.317.0085 [arXiv:1908.11684 [hep-lat]].
- [36] K. Suzuki *et al.* [JLQCD], *PoS LATTICE2019*, 178  
(2020) doi:10.22323/1.363.0178 [arXiv:2001.07962 [hep-  
lat]].
- [37] A. Bazavov *et al.* [HotQCD], *Phys. Rev. D* **86**,  
094503 (2012) doi:10.1103/PhysRevD.86.094503  
[arXiv:1205.3535 [hep-lat]].
- [38] M. I. Buchoff, M. Cheng, N. H. Christ, H. T. Ding,  
C. Jung, F. Karsch, Z. Lin, R. D. Mawhinney,  
S. Mukherjee and P. Petreczky, *et al.* *Phys. Rev. D* **89**,  
no.5, 054514 (2014) doi:10.1103/PhysRevD.89.054514  
[arXiv:1309.4149 [hep-lat]].
- [39] H. T. Ding, P. Hegde, O. Kaczmarek, F. Karsch,  
A. Lahiri, S. T. Li, S. Mukherjee, H. Ohno, P. Petreczky

- and C. Schmidt, *et al.* Phys. Rev. Lett. **123**, no.6, 062002 (2019) doi:10.1103/PhysRevLett.123.062002 [arXiv:1903.04801 [hep-lat]].
- [40] G. Grilli di Cortona, E. Hardy, J. Pardo Vega and G. Villadoro, JHEP **1601**, 034 (2016) doi:10.1007/JHEP01(2016)034 [arXiv:1511.02867 [hep-ph]].
- [41] A. Gómez Nicola, J. Ruiz De Elvira and A. Vioque-Rodríguez, JHEP **1911**, 086 (2019) doi:10.1007/JHEP11(2019)086 [arXiv:1907.11734 [hep-ph]].
- [42] P. Petreczky, H. P. Schadler and S. Sharma, Phys. Lett. B **762**, 498-505 (2016) doi:10.1016/j.physletb.2016.09.063 [arXiv:1606.03145 [hep-lat]].
- [43] C. Bonati, M. D'Elia, G. Martinelli, F. Negro, F. Sanfilippo and A. Todaro, JHEP **11**, 170 (2018) doi:10.1007/JHEP11(2018)170 [arXiv:1807.07954 [hep-lat]].
- [44] R. D. Pisarski and F. Rennecke, Phys. Rev. D **101**, no.11, 114019 (2020) doi:10.1103/PhysRevD.101.114019 [arXiv:1910.14052 [hep-ph]].
- [45] J. E. Kim and S. J. Kim, Phys. Lett. B **783**, 357 (2018) doi:10.1016/j.physletb.2018.07.020 [arXiv:1804.05173 [hep-ph]].