

SOME APPLICATIONS OF UNIFORMLY p -CONVERGENT SETS

M. ALIKHANI.

ABSTRACT. In this paper, we introduce a new class of subsets of bounded linear operators between Banach spaces which is p -version of the uniformly completely continuous sets. Then, we study the relationship between these sets with the equicontact sets. Moreover, we introduce the concept of weakly p -sequentially continuous differentiable mappings and obtain some characterizations of these mappings. Finally, we give a factorization result for differentiable mappings through p -convergent operators.

1. INTRODUCTION

Let X and Y be Banach spaces. The study uniformly completely continuous sets in the class of all bounded linear operators between Banach spaces have been obtained in recent years by several authors. The research works of Cilia et al. [9], shows that if $U \subseteq X$ is an open convex and $f : U \rightarrow Y$ is a differentiable mapping whose derivative f' is uniformly continuous on U -bounded subsets of U ($f \in C^{1u}(U, Y)$), then f takes weakly Cauchy U -bounded sequences into norm convergent sequences (in short, $f \in C_{wsc}(U, Y)$), if and only if f' takes Rosenthal and U -bounded subsets of U into uniformly completely continuous subsets of $L(X, Y)$. For more information in this kinds of researches, we refer the reader to [2, 5, 10, 11, 13, 15, 17, 18, 19, 20] and references therein. Recently, Chen et al. [16], by introducing the notion p -(V) sets, showed that an operator $T : X \rightarrow Y$ is p -convergent if and only if its adjoint $T^* : Y^* \rightarrow X^*$ takes bounded subsets of Y^* into p -(V) subset X^* . Motivated by the above work and the research works of Cilia et al [9, 10, 11], we give similar results for differentiable mappings. Here, we introduce the notion uniformly p -convergent sets and we try answer to the following interesting questions:

1991 *Mathematics Subject Classification.* Primary: 46B25; Secondary: 46E50, 46G05.

Key words and phrases. uniformly completely continuous subsets; weakly sequentially continuous differentiable mappings; Dunford-Pettis property of order p ;

- For given a differentiable mapping $f : U \rightarrow Y$, under which conditions its derivative f' takes U -bounded weakly p -precompact subsets of U into uniformly p -convergent subsets?
- For given a differentiable mapping $f : U \rightarrow Y$, under which conditions its derivative f' takes U -bounded sets into uniformly p -convergent sets?

This paper deals with the p -version of uniformly completely continuous sets and weakly sequentially continuous differentiable mappings. In Section 2 of this article provides a wide range of definitions and concepts in Banach spaces. These concepts are mostly well known, but we need them in the sequel.

In Section 3, we define the concepts of uniformly p -convergent sets and weakly p -sequentially continuous differentiable mappings. Also, we apply the notion weakly equicontact sets in order to find a characterization for those Banach spaces in which the double dual of them have the p -Schur property. Finally, we find some equivalent conditions for all $f \in C^{1u}(U, Y)$ such that f' takes U -bounded and weakly p -precompact subsets of U into uniformly p -convergent subsets of the class of all bounded linear operators from X to Y .

In the Section 4, we apply the concept uniformly p -convergent sets in order to find a factorization result for a differentiable mapping through a p -convergent operator.

2. NOTIONS AND DEFINITIONS

Throughout this paper X, Y and Z will always denote real Banach spaces and U is an open convex subset of X . We denote the spaces of all bounded linear operators, compact operators and weakly compact operators from X into Y by $L(X, Y)$, $K(X, Y)$ and $W(X, Y)$, respectively. The topological dual of X is denoted by X^* and the adjoint of an operator T is denoted by T^* . Also we use $\langle x^*, x \rangle$ or $x^*(x)$ for the duality between $x \in X$ and $x^* \in X^*$. We denote the closed unit ball of X and the identity operator on X by B_X and id_X respectively. p^* will always denote the conjugate number of p for $1 \leq p < \infty$; if $p = 1$, ℓ_{p^*} plays the role of c_0 . The unit coordinate vector in ℓ_p (resp. c_0 or ℓ_∞) is denoted by e_n^p (resp. e_n). In this paper $1 \leq p < \infty$, except for the cases where we consider other assumptions.

To state our results, we need to recall some definitions. A sequence $(x_n)_n$ in X is called weakly p -summable, if $(x^*(x_n))_n \in \ell_p$ for each $x^* \in X^*$. We denote by $\ell_p^w(X)$ the space of all weakly p -summable sequences in X ; see [14]. A bounded subset K of X is relatively weakly- p -compact, if every sequence in K has a

weakly- p -convergent subsequence with limit in X . A sequence $(x_n)_n$ in X is called weakly p -Cauchy, provided that $(x_{m_k} - x_{n_k})_k \in \ell_p^w(X)$ for any increasing sequences $(m_k)_k$ and $(n_k)_k$ of positive integers; see [8]. A subset K of X is said to be weakly p -precompact, provided that every sequence from K has a weakly p -Cauchy subsequence; see [8]. Note that the weakly ∞ -precompact sets are precisely the weakly precompact sets or Rosenthal sets. An operator $T \in L(X, Y)$ is said to be weakly p -precompact, if $T(B_X)$ is weakly p -precompact. An operator $T \in L(X, Y)$ is called p -convergent, if $\lim_{n \rightarrow \infty} \|T(x_n)\| = 0$ for all $(x_n)_n \in \ell_p^w(X)$. We denote the space of all p -convergent operators from X into Y , by $C_p(X, Y)$; see [6]. If the identity operator on X is p -convergent, we say that a Banach space X has the p -Schur property, which is equivalent to every weakly p -compact subset of X is norm compact; see [12]. A Banach space X is said to have the Dunford-Pettis property of p (in short, (DPP_p)), if for any Banach space Y , every weakly compact operator $T : X \rightarrow Y$ is p -convergent; see [8]. A bounded subset K of X^* is a p -(V) set, if $\lim_{n \rightarrow \infty} \sup_{x^* \in K} |x^*(x_n)| = 0$, for every $(x_n)_n \in \ell_p^w(X)$; see [16]. Given $x, y \in X$, the segment with bounds x and y denoted by $I(x, y)$. A set $B \subset U$ is U -bounded, if it is bounded and the distance between B and the boundary of U is strictly positive; see [10]. The space of all differentiable mappings $f : U \rightarrow Y$ whose derivative $f' : U \rightarrow L(X, Y)$ is uniformly continuous on U -bounded subsets of U will be denoted by $C^{1u}(U, Y)$; see [9]. A set $M \subset K(X, Y)$ is equicontact, if there exists a null sequence $(x_n^*)_n$ in X^* so that $\|T(x)\| \leq \sup_n |x_n^*(x)|$ for all $x \in X$ and all $T \in M$, which is equivalent to every bounded sequence $(x_n)_n$ in X has a subsequence $(x_{k_n})_n$ such that $(Tx_{k_n})_n$ is uniformly convergent for $T \in M$; see [18]. For given a mapping $f : U \rightarrow Y$ and a class \mathcal{M} of subsets of U such that every singleton belongs to \mathcal{M} , the mapping f is \mathcal{M} -differentiable at $x \in U$, if there exists an operator $f'(x) \in L(X, Y)$ such that

$$\lim_{\varepsilon \rightarrow 0} \frac{f(x + \varepsilon y) - f(x) - f'(x)(\varepsilon y)}{\varepsilon} = 0$$

uniformly to y on each member of \mathcal{M} . In this case, we write $f \in D_{\mathcal{M}}(x, Y)$; see [15]. We say that a mapping f is Gâteaux differentiable at x , if $f \in D_{\mathcal{M}}(x, Y)$ where \mathcal{M} is the class of all single-point subsets of X . We also, say that f is Fréchet differentiable at x if $f \in D_{\mathcal{M}}(x, Y)$, where \mathcal{M} is the class of all bounded subsets of X .

3. WEAKLY p -SEQUENTIALLY CONTINUOUS DIFFERENTIABLE MAPPINGS

Here, we introduce the notion uniformly p -convergent sets in $L(X, Y)$ and give some properties of these sets. Then, we study the weakly p -sequentially continuous differentiable mappings.

Definition 3.1. *Let $K \subset L(X, Y)$. We say that K is a uniformly p -convergent set, if every $(x_n)_n \in \ell_p^w(X)$ converges uniformly on K , that is,*

$$\limsup_n \sup_{T \in K} \|T(x_n)\| = 0.$$

Note that, the uniformly ∞ -convergent sets in $L(X, Y)$ are precisely the uniformly completely continuous sets; see [18]. Also, every uniformly q -convergent subset of $L(X, Y)$ is uniformly p -convergent, whenever $1 \leq p < q \leq \infty$. It would be interesting to obtain conditions under which every uniformly p -convergent set in $L(X, Y)$ is uniformly q -convergent. In particular, we obtain a characterization for those Banach spaces in which uniformly p -convergent sets in X^* are uniformly q -convergent; see [4].

The following example shows that, there exists a uniformly p -convergent subset of $L(\ell_2, Y)$ so that it is not uniformly q -convergent.

Example 3.1. *Let $X = \ell_2$ and Y be an arbitrary Banach space. Since ℓ_2 does not have 2-Schur property, B_{ℓ_2} is not a 2-(V) set in ℓ_2 . Therefore B_{ℓ_2} is not uniformly 2-convergent subsets of ℓ_2 . On the other hand, ℓ_2 contain no copy of c_0 . Therefore, ℓ_2 has the 1-Schur property; see [12]. Hence, B_{ℓ_2} is a 1-(V) set and so, B_{ℓ_2} is a uniformly 1-convergent subsets of ℓ_2 . Now, let $0 \neq y_0 \in B_Y$ and $S : \mathbb{R} \rightarrow Y$ be the operator given by $S(\lambda) := \lambda y_0$ ($\lambda \in \mathbb{R}$). Define an operator $T : \ell_2 \rightarrow L(\ell_2, Y)$ by $T(\phi)(h) := \phi(h)y_0$, for $\phi \in \ell_2^* = \ell_2$ and $h \in \ell_2$. Then*

$$\|T(\phi)\| = \sup_{h \in B_{\ell_2}} \|T(\phi)(h)\| = \sup_{h \in B_{\ell_2}} \|\phi(h)y_0\| = \|\phi\|.$$

Since, uniformly p -convergent sets are stable under isometry, there exists a uniformly 1-convergent subset of $L(\ell_2, Y)$ such that it is not uniformly 2-convergent.

In the following result, we give some properties of uniformly p -convergent sets.

Proposition 3.1. (i) *Every subset of a uniformly p -convergent set in $L(X, Y)$ is uniformly p -convergent.*

(ii) Absolutely closed convex hull of a uniformly p -convergent set in $L(X, Y)$ is uniformly p -convergent.

(iii) If K_1, \dots, K_n are uniformly p -convergent sets in $L(X, Y)$, then $\bigcup_{i=1}^n K_i$ and

$\sum_{i=1}^n K_i$ are uniformly p -convergent sets in $L(X, Y)$.

(iv) Every relatively compact subset of $C_p(X, Y)$ is uniformly p -convergent.

(v) If $K \subset L(X, Y)$ is a uniformly p -convergent set, then $K \subset C_p(X, Y)$.

Theorem 3.2. *Let X be a Banach space. If there exists a Banach space Y so that every uniformly p -convergent set of $K(X, Y)$ is equicontact, then $C_p(X, Y) = K(X, Y)$.*

Proof. Since the p -(V) sets in X^* coincides with the uniformly p -convergent subsets of X^* , it is enough to show that every uniformly p -convergent subset M of X^* is relatively compact; see ([4, Theorem 2.4]). For this purpose, consider $y_0 \in S_Y$ and put $H = M \otimes y_0$. Obviously, H is a uniformly p -convergent subset of $K(X, Y)$. Hence, by the hypothesis, H is equicontact, which yields the equicontactness of M as a subset of $K(X, \mathbb{R})$. Therefore, an application of ([19, Lemma 2.1]) yields the result. \square

As an immediate consequence of Proposition 2.2 of [18], one can conclude the following result.

Proposition 3.2. *If B_X is weakly p -precompact, then a bounded subset M of $K(X, Y)$ is equicontact if and only if M is uniformly p -convergent.*

Example 3.3. *For each $b = (b_n)_n \in \ell_2$, Define the operator $T_b : \ell_2 \rightarrow \ell_1$ by $T_b(a_n) := (a_n b_n)$. Obviously, $M := \{T_b : b \in B_{\ell_2}\}$ is not equicontact in $K(\ell_2, \ell_1)$. So, Proposition 3.2 implies that M is not a uniformly p -convergent subset of $K(\ell_2, \ell_1)$.*

A subset M of $K(X, Y)$ is said to be collectively compact, if $\bigcup_{T \in M} T(B_X)$ is a relatively compact set. Recall that $M \subset K(X, Y)$ is equicontact if and only if $M^* = \{T^* : T \in M\}$ is collectively compact; see [18].

Proposition 3.3. *Let $S : X \rightarrow Z$ be a weakly p -precompact operator. If for each Banach space Y , every $N \subset C_p(Z, Y)$ is uniformly p -convergent, then the set $N \circ S := \{T \circ S : T \in N\}$ is equicontact.*

Proof. We prove that $S^* \circ N^*$ is collectively compact. Consider a sequence $((S^* \circ T_n^*)y_n^*)_n$ in $\bigcup_{T \in N} S^* \circ T^*(B_Y)$ and put $A := \{T_n^*y_n^* : n \in \mathbb{N}\}$. It is easy to verify that, A is a uniformly p -convergent set in Z^* . Indeed, if $(z_n)_n \in \ell_p^w(Z)$, we have

$$\lim_{n \rightarrow \infty} \sup_m |\langle z_n, T_m^*(y_m^*) \rangle| \leq \lim_{n \rightarrow \infty} \sup_m \|T_m(z_n)\| = 0.$$

Let $(z_n^*)_n \subset A$ and let $(z_n)_n \in \ell_p^w(Z)$. Consider an operator $S_1 : Z \rightarrow \ell_\infty$ defined by $S_1(z) := (z_n^*(z))$. Since A is uniformly p -convergent set in Z^* , $\lim_n \|S_1(z_n)\| = \lim_n \sup_i |z_i^*(z_n)| = 0$, and so S_1 is p -convergent. Hence, the operator $S_1 S : X \rightarrow \ell_\infty$ is compact, since $S : X \rightarrow Z$ is a weakly p -precompact operator. Thus $S^* \circ S_1^*$ is compact and so, $S^*(z_n^*)_n = (S^*(S_1^*(e_n^1)))_n$ is relatively compact, where (e_n^1) is the unit basis of ℓ_1 . Hence, $S^*(A)$ is a relatively compact set and so, $((S^* \circ T_n^*)y_n^*)_n$ has a convergent subsequence. \square

In [19], the authors defined weakly equicontact sets as those subsets M of $W(X, Y)$ satisfying that, for every bounded sequence $(x_n)_n$ in X , there exists a subsequence $(x_{k_n})_n$ such that $(T(x_{k_n}))_n$ is weakly uniformly convergent for $T \in M$.

Proposition 3.4. *If B_X is weakly p -precompact, then the following statements are equivalent for a set $M \subset W(X, Y)$.*

- (i) M is weakly equicontact.
- (ii) $M^*(y^*)$ is a uniformly p -convergent set in X for every $y^* \in Y^*$.
- (iii) $T(x_n) \xrightarrow{w} 0$ uniformly for $T \in M$ whenever $(x_n)_n \in \ell_p^w(X)$.

Proof. Since the assertions (i) \Rightarrow (ii) is straightforward of ([19, Corollary 2.3]) and (ii) \Rightarrow (iii) is obvious, we only have to show that (iii) \Rightarrow (i).

Let $(x_n)_n$ be a bounded sequence in X . Since B_X is weakly p -precompact, we can suppose that $(x_n)_n$ is weakly p -Cauchy. Assume that M is not weakly equicontact. Thus, $(T(x_n))_n$ is not weakly Cauchy uniformly for $T \in M$. So, there exist $\varepsilon > 0$, $y^* \in Y^*$, strictly increasing sequences $(p_n)_n \subset \mathbb{N}$ and $(q_n)_n \subset \mathbb{N}$, a sequence $(T_n)_n$ in M such that:

$$|\langle x_{p_n} - x_{q_n}, T_n^*(y^*) \rangle| = |\langle T_n(x_{p_n}) - T_n(x_{q_n}), y^* \rangle| \geq \frac{\varepsilon}{2} \quad \text{for all } n \in \mathbb{N},$$

which is a contradiction. \square

In the following example, we show that the hypothesis about X cannot be omitted in Proposition 3.4.

Example 3.4. Suppose $M := B_{\ell_\infty} \otimes B_{\ell_1}$. An easy verification shows that $M \subset W(\ell_1, \ell_1)$ satisfies conditions (ii) and (iii) of Proposition 3.4 while, M is not weakly equicontact.

Here, we obtain a characterization of double dual of Banach space X with the p -Schur property.

Theorem 3.5. *If X is a Banach space, then the following statements are equivalent:*

- (i) *For every Banach space Y , if $M \subset W(X^*, Y)$ is relatively weakly p -compact, then it is weakly equicontact.*
- (ii) *For some Banach space $Y \neq \{0\}$, if $M \subset W(X^*, Y)$ is relatively weakly p -compact, then it is weakly equicontact.*
- (iii) *X^{**} has the p -Schur property.*

Proof. The assertion (i) \Rightarrow (ii) is straightforward. Therefore, we only prove that the assertions (ii) \Rightarrow (iii) and (iii) \Rightarrow (i).

(ii) \Rightarrow (iii) Let K be a relatively weakly p -compact set in X^{**} . We claim that K is relatively norm compact. For this purpose, consider $M = K \otimes y_0$, so that $y_0 \in Y - \{0\}$. One can see that, M is a relatively weakly p -compact set in $W(X^*, Y)$. By the hypothesis, M is weakly equicontact, which yields the weakly equicontactness of K . Hence, ([19, Lemma 2.1]) implies that K is relatively norm compact.

(iii) \Rightarrow (i) Suppose that $M \subset W(X^*, Y)$ is relatively weakly p -compact and $(x_n^*)_n \in \ell_p^w(X^*)$. If $T(x_n^*) \xrightarrow{w} 0$, so that the convergence is not uniform for $T \in M$, then there exist $y^* \in Y^*$, $\varepsilon > 0$, strictly increasing sequences $(p_n)_n \subset \mathbb{N}$ and $(q_n)_n \subset \mathbb{N}$, and a sequence $(T_n)_n$ in M such that:

$$|\langle x_{p_n}^* - x_{q_n}^*, T_n^*(y^*) \rangle| = |\langle T_n(x_{p_n}^*) - T_n(x_{q_n}^*), y^* \rangle| \geq \frac{\varepsilon}{2} \quad \text{for all } n \in \mathbb{N},$$

On the other hand, $(x_{p_n}^* - x_{q_n}^*)_n$ is weakly p -Cauchy and $(T_n^*(y^*))_n$ admits a weakly convergent subsequence. Since X^{**} has the p -Schur property, Theorem 2.8 of [4] implies that $X^{**} \in (DPP_p)$. Hence, by using Theorem 3.1 in [8], we have $\lim_{n \rightarrow \infty} |\langle x_{p_n}^* - x_{q_n}^*, T_n^*(y^*) \rangle| = 0$, which is a contradiction. \square

Proposition 3.5. *Let X be a Banach space. If there exists a Banach space $Y \neq \{0\}$ such that every uniformly p -convergent set of $W(X, Y)$ is weakly equicontact, then $C_p(X, Y) = K(X, Y)$.*

Proof. It suffices to show that every uniformly p -convergent set $K \subset X^*$ is relatively compact; see ([4, Theorem 2.4]). Choose $y_0 \in Y$ and $y_0^* \in Y^*$ such that $\langle y_0^*, y_0 \rangle = 1$. Clearly, $M = K \otimes y_0$ is a uniformly p -convergent set in $W(X, Y)$ and so, by the hypothesis, M is weakly equicontact. Hence, by using Proposition 2.2 of [19], $K = \langle y_0^*, y_0 \rangle K = M^*(y_0^*)$ is relatively compact. \square

Definition 3.2. Let $U \subset X$ be an open convex and $1 \leq p \leq \infty$. We say that $f : U \rightarrow Y$ is a weakly p -sequentially continuous map, if it takes U -bounded and weakly p -Cauchy sequences of U into norm convergent sequences in Y . We denote the space of all such mappings by $C_{wsc}^p(U, Y)$.

The class of all weakly ∞ -sequentially continuous mappings is precisely the class of all weakly sequentially continuous mappings; see [9]. Also, note that $C_{wsc}^q(U, Y) \subseteq C_{wsc}^p(U, Y)$, whenever $1 \leq p < q \leq \infty$. But, we do not have any example of a mapping $f \in C^{1u}(U, Y) \cap C_{wsc}^p(U, Y)$ which does not belong to $C_{wsc}^q(U, Y)$.

Proposition 3.6. Let $U \subset X$ be an open convex and $1 \leq p \leq \infty$. If f is compact and takes U -bounded and weakly p -Cauchy sequences into weakly Cauchy sequences, then $f \in C_{wsc}^p(U, Y)$.

Proposition 3.7. Let $U \subset X$ be an open convex and $1 \leq p \leq \infty$. If $f \in C^{1u}(U, Y)$ so that $f' \in C_{wsc}^p(U, C_p(X, Y))$, then $f \in C_{wsc}^p(U, Y)$.

Proof. Let $(x_n)_n$ be a U -bounded and weakly p -Cauchy sequence. By the Mean Value Theorem ([7, Theorem 6.4]), we have

$$\|f(x_n) - f(x_m)\| \leq \|f'(c_{n,m})(x_n - x_m)\|$$

for some $c_{n,m} \in I(x_n, x_m)$. Since the sequence $(c_{n,m})$ is U -bounded and weakly p -Cauchy, the sequence $(f'(c_{n,m}))$ norm converges to some $T \in C_p(X, Y)$. Therefore we have:

$$\lim_{n,m \rightarrow \infty} \|f'(c_{n,m})(x_n - x_m) - T(x_n - x_m)\| + \|T(x_n - x_m)\|$$

$$\leq \lim_{n,m \rightarrow \infty} \|f'(c_{n,m})(x_n - x_m) - T(x_n - x_m)\| + \lim_{n,m \rightarrow \infty} \|T(x_n - x_m)\| = 0.$$

So, $\lim_{n,m \rightarrow \infty} \|f'(c_{n,m})(x_n - x_m)\| = 0$. Hence, the sequence $(f(x_n))_n$ is norm convergent. \square

Proposition 3.8. *Let $U \subset X$ be an open convex and $1 \leq p \leq \infty$. If $f : U \rightarrow Y$ is a differentiable mapping such that for every U -bounded set K , $f'(K)$ is a uniformly p -convergent set in $L(X, Y)$, then $f \in C_{wsc}^p(U, Y)$.*

Proof. Let $(x_n)_n$ be a U -bounded and weakly p -Cauchy sequence. By the Mean Value Theorem ([7, Theorem 6.4]), for all $n, m \in \mathbb{N}$, there is $c_{i,j} \in I(x_n, x_m)$ such that

$$\|f(x_n) - f(x_m)\| \leq \|f'(c_{i,j})(x_n - x_m)\| \leq \sup_{i,j} \|f'(c_{i,j})(x_n - x_m)\|$$

Obviously, the set $K := \{c_{i,j} : i, j \in \mathbb{N}\}$ is contained in the convex hull of all x_n and then in U , since U is a convex set. Moreover K is still a U -bounded set. By the hypothesis, $f'(K)$ is a uniformly p -convergent set in $L(X, Y)$. Since $(x_n - x_m) \in \ell_p^w(X)$, it follows that $\limsup_{n,m} \sup_{i,j} \|f'(c_{i,j})(x_n - x_m)\| = 0$. Therefore,

$$\|f(x_n) - f(x_m)\| \rightarrow 0. \quad \square$$

Now by using the same argument of ([9, Theorem 2.1]), we find a method to get uniformly p -convergent subsets of $L(X, Y)$.

Theorem 3.6. *Let $U \subseteq X$ be an open convex subset and $1 \leq p < \infty$. If $f \in C^{1u}(U, Y)$, then the following assertions are equivalent:*

- (i) $f \in C_{wsc}^p(U, Y)$;
- (ii) *For every U -bounded and weakly p -Cauchy sequence (x_n) and every weakly p -Cauchy sequence $(h_n) \subset X$, the sequence $(f'(x_n)(h_n))_n$ norm converges in Y ;*
- (iii) *For every U -bounded weakly p -Cauchy sequence $(x_n)_n$ and every weakly p -summable sequence $(h_n)_n \subset X$, we have*

$$\limsup_n \sup_m \|f'(x_m)(h_n)\| = 0;$$

- (iv) *For every U -bounded and weakly p -Cauchy sequence $(x_n)_n$ and every weakly p -summable sequence $(h_n)_n \subset X$, we have*

$$\lim_n f'(x_n)(h_n) = 0;$$

- (v) *f' takes U -bounded and weakly p -precompact subsets of U into uniformly p -convergent subsets of $L(X, Y)$.*

Proof. (i) \Rightarrow (ii) Let $(x_n)_n$ be a U -bounded weakly p -Cauchy sequence and let $(h_n)_n$ be a weakly p -Cauchy sequence in X . Without loss of generality, we

assume that $(h_n)_n$ is bounded. Consider $B := \{x_n : n \in \mathbb{N}\}$ and let $d := \min\{1, \text{dist}(B, \partial U)\}$. It is easy to show that the set

$$B' := B + \frac{d}{2}B_X \subset U$$

is also U -bounded. Since $f \in C^{1u}(U, Y)$, f' is uniformly continuous on B' . Hence, for given $\varepsilon > 0$, there exists $0 < \delta < \frac{d}{4}$ such that if $t_1, t_2 \in B'$ satisfy $\|t_1 - t_2\| < 2\delta$, then

$$(1) \quad \|f'(t_1) - f'(t_2)\| < \frac{\varepsilon}{4}.$$

If $c \in I(x_n, x_n + \delta h_n)$ for some $n \in \mathbb{N}$, then

$$\|c - x_n\| \leq \delta \|h_n\| < \delta < 2\delta < \frac{d}{2},$$

and so,

$$c = x_n + (c - x_n) \in B' = B + \frac{d}{2}B_X$$

As an immediate consequence of the Mean Value Theorem ([7, Theorem 6.4]), and formula (1), we obtain

$$\begin{aligned} & \|f'(x_n)(\delta h_n) - f(x_n + \delta h_n) + f(x_n)\| \\ & \leq \sup_{c \in I(x_n, x_n + \delta h_n)} \|f'(c) - f'(x_n)\| \|\delta h_n\| \leq \frac{\varepsilon \delta}{4}. \end{aligned}$$

Similarly,

$$\begin{aligned} & \|f(x_m + \delta h_m) - f(x_m) - f'(x_m)(h_m)\| \\ & \leq \sup_{c \in I(x_m, x_m + \delta h_m)} \|f'(c) - f'(x_m)\| \|\delta h_m\| \leq \frac{\varepsilon \delta}{4}. \end{aligned}$$

On the other hand, the sequences $(x_n + \delta h_n)_n$ and $(x_n)_n$ are U -bounded and weakly p -Cauchy in U . Hence, by the hypothesis the sequences $(f(x_n + \delta h_n))_n$ and $(f(x_n))_n$ are norm convergent in Y . Hence, we can find $n_0 \in \mathbb{N}$ so that for $n, m > n_0$:

$$\|f(x_n + \delta h_n) - f(x_m + \delta h_m)\| < \frac{\varepsilon \delta}{4}, \quad \|f(x_n) - f(x_m)\| < \frac{\varepsilon \delta}{4}$$

So, for $n, m > n_0$, we have

$$\| f'(x_n)(h_n) - f'(x_m)(h_m) \| < \varepsilon.$$

(ii) \Rightarrow (iii) Let $(x_n)_n$ be a U -bounded weakly p -Cauchy sequence and let $(h_n)_n$ be a weakly p -summable sequence in X . By the part (ii), for every $h \in X$, the set $\{f'(x_n)(h) : n \in \mathbb{N}\}$ is bounded in Y . On the other hand, there exists a subsequence $(x_{m_k})_k$ of $(x_m)_m$ in U such that

$$\| f'(x_{m_k})(h_k) \| \geq \sup_m \| f'(x_m)(h_k) \| - \frac{1}{k} \quad (k \in \mathbb{N}).$$

Since the sequences $(x_{m_k})_k$ in U and $(h_1, 0, h_2, 0, h_3, 0, \dots)$ in X are weakly p -Cauchy, the sequence

$$(f'(x_{m_1})(h_1), 0, f'(x_{m_2})(h_2), 0, f'(x_{m_3})(h_3), 0, \dots)$$

converges in Y . Therefore, $\lim_k f'(x_{m_k})(h_k) = 0$. Hence, we have:

$$\limsup_k \sup_m \| f'(x_m)(h_k) \| = 0.$$

(iii) \Rightarrow (iv) is obvious.

(iv) \Rightarrow (v) Let K be a weakly p -precompact and U -bounded set. It is clear that, for every $h \in X$, the set $f'(K)(h)$ is bounded in Y . Let $(h_n)_n$ be a weakly p -summable sequence in X . If $(h_{n_k})_k$ is a subsequence of $(h_n)_n$, then for every $k \in \mathbb{N}$, there exists $a_k \in K$ such that

$$\sup_{a \in K} \| f'(a)(h_{n_k}) \| < \| f'(a_k)(h_{n_k}) \| + \frac{1}{k}.$$

Since K is a weakly p -precompact set, the sequence $(a_k)_k$ admits a weakly p -Cauchy subsequence $(a_{k_r})_r$. Hence, by the hypothesis,

$$\lim_r \| f'(a_{k_r})(h_{n_{k_r}}) \| = 0.$$

Therefore we have, $\limsup_r \sup_{a \in K} \| f'(a)(h_{n_{k_r}}) \| = 0$. Hence, every subsequence of $(\sup_{a \in K} \| f'(a)(h_n) \|)_n$ has a subsequence converging to 0. Therefore, the sequence itself converges to 0, that is, $\limsup_n \sup_{a \in K} \| f'(a)(h_n) \| = 0$.

(v) \Rightarrow (i) Let $(x_n)_n$ be a U -bounded and weakly p -Cauchy sequence. Since U is

convex, the segment $I(x_n, x_m)$ is contained in U for all $n, m \in \mathbb{N}$. By the Mean Value Theorem ([7, Theorem 6.4]), there exists $c_{nm} \in I(x_n, x_m)$ such that

$$\|f(x_n) - f(x_m)\| \leq \|f'(c_{n,m})(x_n - x_m)\| \leq \sup_{i,j \in \mathbb{N}} \|f'(c_{i,j})(x_n - x_m)\|.$$

Since $(c_{i,j})_{i,j}$ is a weakly p -Cauchy and U -bounded sequence, the part (v) implies that

$$\limsup_{n,m} \sup_{i,j \in \mathbb{N}} \|f'(c_{i,j})(x_n - x_m)\| = 0.$$

Therefore, $\lim_{n,m \rightarrow \infty} \|f(x_n) - f(x_m)\| = 0$. \square

Example 3.7. Let $h \in C^1(\mathbb{R})$ and $1 < p < 2$. Define $f : \ell_{p^*} \rightarrow \mathbb{R}$ by $f((x_n)_n) = \sum_{n=1}^{\infty} \frac{h(x_n)}{2^n}$. The same argument as in the ([11, Example 2.4]), shows that f is differentiable so that $f'((x_n)_n) = (\frac{h'(x_n)}{2^n})_n \in \ell_p$. By Pitt's Theorem ([11, Theorem 2.1.4]), $f' : \ell_{p^*} \rightarrow \ell_p$ is compact and so, $f'(B_{\ell_{p^*}})$ is a relatively compact set in $L(\ell_{p^*}, \mathbb{R}) = C_p(\ell_{p^*}, \mathbb{R})$. Therefore, the part (iv) of Proposition 3.1, yields that $f'(B_{\ell_{p^*}})$ is a uniformly p -convergent set in $L(\ell_{p^*}, \mathbb{R})$. Hence, Proposition 3.8 implies that f is weakly p -sequentially continuous. On the other hand, $\frac{1}{2}B_{\ell_{p^*}}$ is a $B_{\ell_{p^*}}$ -bounded set. Hence, by Theorem 3.6 $f'(\frac{1}{2}B_{\ell_{p^*}})$ is a p -(V) set in ℓ_p .

4. FACTORIZATION THEOREM THROUGH A p -CONVERGENT OPERATOR

Here, for given a mapping $f : U \rightarrow Y$, we show that f is differentiable so that f' takes U -bounded sets into uniformly p -convergent sets if and only if it happens $f = g \circ S$, where S is a p -convergent operator from X into a suitable Banach space Z and $g : S(U) \rightarrow Y$ is a Gâteaux differentiable mapping with some additional properties.

Theorem 4.1. Let $U \subset X$ be an open convex and $1 \leq p \leq \infty$. If $f : U \rightarrow Y$ is a mapping, then the following assertions are equivalent:

- (a) f is differentiable so that f' takes U -bounded sets into uniformly p -convergent sets and f is weakly p -sequentially continuous.
- (b) There exist a Banach space Z , an operator $S \in C_p(X, Z)$ and a mapping $g : S(U) \rightarrow Y$ such that:
 - (i) $f(x) = g(S(x))$ for all $x \in U$.
 - (ii) $g \in D_{\mathcal{M}}(S(x), Y)$ for every $x \in U$, where

$$\mathcal{M} := \{S(B) : B \text{ is a } U\text{-bounded subset of } X\}.$$

(iii) g' is bounded on $S(B)$ for every U -bounded subset $B \subset X$.

Moreover, if this factorization holds, f is weakly p -sequentially continuous.

Proof. (a) \Rightarrow (b) For every $n \in \mathbb{N}$, put

$$W_n = \{x \in U : d(x, \partial U) > \frac{1}{n}\} \cap nB_X.$$

By the hypothesis for every $r \in \mathbb{N}$, $f'(\frac{W_r}{r\|f'\|_{W_r}})$ is a uniformly p -convergent set. Now, we define $K := \bigcup_{r=1}^{\infty} \frac{f'(W_r)}{r\|f'\|_{W_r}}$. We claim that, K is a uniformly

p -convergent set. Indeed, for every $N \in \mathbb{N}$, we define $A_N := \bigcup_{r \leq N} \frac{f'(W_r)}{r\|f'\|_{W_r}}$

and $B_N := \bigcup_{r > N} \frac{f'(W_r)}{r\|f'\|_{W_r}}$. Since A_N is a uniformly p -convergent set, it is

enough to show that B_N is a uniformly p -convergent set. For this purpose, let $(x_n)_n \in \ell_p^w(X)$ and $M = \sup_n \|x_n\|$. If $T \in B_N$, then there are $r > N$ and

$x \in W_r$ so that $T = \frac{f'(x)}{r\|f'\|_{W_r}}$. It is clear that, $\|T\| \leq \frac{1}{r} < \frac{1}{N}$. Hence, for each $N \in \mathbb{N}$ we have:

$$\lim_{n \rightarrow \infty} \sup_{T \in K} \|T(x_n)\| = \max\{\lim_{n \rightarrow \infty} \sup_{T \in A_N} \|T(x_n)\|, \lim_{n \rightarrow \infty} \sup_{T \in B_N} \|T(x_n)\|\} \leq \frac{M}{N}.$$

Therefore, $\lim_{n \rightarrow \infty} \sup_{T \in K} \|T(x_n)\| = 0$ and so, K is a uniformly p -convergent set.

Now, as in the proof of ([10, Theorem 2.1]), let

$$V_K := \{x \in X : \sup_{\phi \in K} \|\phi(x)\|_Y = 0\}$$

and $G := \frac{X}{V_K}$. If $S : X \rightarrow G$ is the quotient map G , then G is normed space respect the norm $\|S(x)\| = \sup_{\phi \in K} \|\phi(x)\|_Y$, $\forall x \in X$. Suppose that Z is

the completion of G . Let $(x_n)_n \in \ell_p^w(X)$, since K is a uniformly p -convergent set, $\|S(x_n)\| = \sup_{\phi \in K} \|\phi(x_n)\| \rightarrow 0$. Hence, $S \in C_p(X, Z)$. Now we define

$g : S(U) \rightarrow Y$ by $g(S(x)) = f(x)$, $x \in U$. In the first, we proved that g is well defined. Suppose that $\|S(x - y)\| = 0$. Since the span of K contains $f'(U)$,

we have

$$\| f'(c)(x - y) \| = 0 \quad (c \in U).$$

By using the Mean Value Theorem ([7, Theorem 6.4]),

$$\sup_{c \in I(x,y)} \| f(x) - f(y) \| \leq \| f'(c)(x - y) \| = 0,$$

and so $f(x) = f(y)$. Therefore g well defined. Now, we show that g is Gâteaux differentiable. For given $x, y \in U$,

$$(2) \quad \lim_{t \rightarrow 0} \frac{g(S(x) + tS(y)) - g(S(x))}{t} = \lim_{t \rightarrow 0} \frac{f(x + ty) - f(x)}{t} = f'(x)(y)$$

where $|t|$ is sufficiently small so that $x + ty \in U$. For $x \in U$ fixed, the mapping $g'(S(x)) : G \rightarrow Y$ given by $g'(S(x))(S(y)) = f'(x)(y)$ ($y \in X$) is linear. Choosing $r \in \mathbb{N}$ so that $x \in W_r$, we have

$$\begin{aligned} \| g'(S(x))(S(y)) \| &= \\ \| f'(x)(y) \| &\leq r \| f' \|_{W_r} \| S(y) \|. \end{aligned}$$

Hence $g'(S(x))$ is continuous and may be extended to the completion Z of G . Hence g is Gâteaux differentiable. Moreover, since f is differentiable, for every U -bounded set B , the limit in (2) exists uniformly to $S(y)$ in $S(B)$. So, $g \in D_{\mathcal{M}}(S(x), Y)$ for every $x \in U$, where

$$\mathcal{M} = \{S(B) : B \text{ is a } U\text{-bounded subset of } X\}$$

and (ii) is proved.

From the inequality (ii), we have

$$\| g'(S(x)) \| = \sup_{\| S(y) \| \leq 1} \| g'(S(x))(S(y)) \| \leq r \| f' \|_{W_r}, \quad (x \in W_r)$$

and this implies (iii).

(b) \Rightarrow (a). Assume that there exists a Banach space Z and $S \in C_p(X, Z)$, and a mapping $g : S(U) \rightarrow Y$ satisfying (b). Obviously, f is differentiable. We claim that f' takes U -bounded sets into uniformly p -convergent sets. For this purpose, suppose that B is a U -bounded set and $(x_n)_n \in \ell_p^w(X)$. Since $S \in C_p(X, Z)$, we have

$$\begin{aligned} \sup_{x \in B} \| f'(x)(x_n) \| &= \sup_{x \in B} \| g'(S(x))(S(x_n)) \| \leq \\ &\sup_{x \in B} \| g'(S(x)) \| \| S(x_n) \| \rightarrow 0. \end{aligned}$$

So, $f'(B)$ is a uniformly p -convergent set. Given $r \in \mathbb{N}$, the mapping g is uniformly continuous on $S(W_r)$. Indeed, for $x, y \in W_r$, we have

$$\begin{aligned} \|g(S(x)) - g(S(y))\| &= \|f(x) - f(y)\| \leq \sup_{c \in W_r} \|f'(c)(x - y)\| \\ &\leq r \|f'\|_{W_r} \sup_{\phi \in K} \|\phi(x - y)\| = r \|f'\|_{W_r} \|S(x - y)\| \end{aligned}$$

where we have used that W_r is a convex set. Hence, if $(x_n)_n$ is a U -bounded weakly p -Cauchy sequence, then the sequence $(S(x_n))_n$ in $S(W_r)$, for a suitable index r , is norm Cauchy and so, $(f(x_n))_n = (g(S(x_n)))_n$ is also norm Cauchy. Hence, f is weakly p -sequentially continuous. \square

Example 4.2. Let $h \in C^1(\mathbb{R})$. Define $f : c_0 \rightarrow \mathbb{R}$ by $f((x_n)_n) = \sum_{n=1}^{\infty} \frac{h(x_n)}{2^n}$. It is easy to verify that f is differentiable so that $f'((x_n)_n) = (\frac{h'(x_n)}{2^n})_n \in \ell_1$. Since $f' : c_0 \rightarrow L(c_0, \mathbb{R}) = C_p(c_0, \mathbb{R})$ is compact, $f'(B_{c_0})$ is a relatively compact set in $C_p(c_0, \mathbb{R})$ and so, $f'(B_{c_0})$ is a uniformly p -convergent set in ℓ_1 . Therefore, Proposition 3.8 implies that f is weakly p -sequentially continuous. Hence, there exist a Banach space Z , an operator $S \in C_p(c_0, Z)$ and a Gâteaux differentiable mapping $g : S(B_{c_0}) \rightarrow \mathbb{R}$ such that $f = g \circ S$ with some additional properties.

REFERENCES

- [1] F. Albiac and N. J. Kalton, Topics in Banach Space Theory, Graduate Texts in Mathematics, 233, Springer, New York, 2006.
- [2] R. M. Aron, Weakly uniformly continuous and weakly sequentially continuous entire functions, Advances in Holomorphy, in: Math. Stud., vol. 34, North-Holland, Amsterdam, 1979, pp. 47-66.
- [3] R. M. Aron, C. Hervs, M. Valdivia, Weakly continuous mappings on Banach spaces, J. Funct. Anal. **52** (1983) 189-204.
- [4] M. Alikhani, M. Fakhar and J. Zafarani, p -convergent operators and p -Schur property, Analysis Mathematica (Accept), (2019).
- [5] F. Bombal, J. M. Gutiérrez, I. Villanueva, Derivative and factorization of holomorphic functions, J. Math. Anal. Appl. **348** (2008) 444-453.
- [6] J. M. F. Castillo and F. Sánchez, Dunford-Pettis-like properties of continuous function vector spaces, Rev. Mat. Univ. Complut. Madrid **6** (1993), 43-59.
- [7] S. B. Chae, Holomorphy and Calculus in Normed Spaces, Monogr. Textbooks Pure Appl. Math. 92, Dekker, New York 1985.
- [8] D. Chen, J. Alejandro Chávez-Domínguez and L. Li, p -converging operators and Dunford-Pettis property of order p , J. Math. Anal. Appl. **461** (2018), 1053-1066.

- [9] R. Cilia, J. M. Gutiérrez, Weakly sequentially continuous differentiable mappings. *J. Math. Anal. Appl.* **360** (2009), 609-623.
- [10] R. Cilia, J. M. Gutiérrez, Factorization of weakly continuous differentiable mappings. *Bull. Braz. Math. Soc. (N.S.)* **40** (2009), 371-380.
- [11] R. Cilia, J. M. Gutiérrez, and G. Saluzzo, Compact factorization of differentiable mappings. *Proc. Amer. Math. Soc.* **137** (2009), 1743-1752.
- [12] M. Dehghani and S. M. Moshtaghioun, On p -Schur property of Banach spaces, *Ann. Funct. Anal.* **9** (2018), 123-136.
- [13] R. Deville, E. Matheron, Pyramidal vectors and smooth functions on Banach spaces, *Proc. Amer. Math. Soc.* **128** (2000) 3601-3608.
- [14] J. Diestel, H. Jarchow and A. Tonge, *Absolutely Summing Operators*, Cambridge Univ. Press, (1995).
- [15] M. González and J. M. Gutiérrez, Factorization of weakly continuous holomorphic mappings, *Studia Math.* **118** (1996), 117-133.
- [16] L. Li, D. Chen and J. Alejandro Chávez-Domínguez, Pelczyński's property (V^*) of order p and its quantification, *Math. Nachr.* **291** (2018) 420-442.
- [17] F. Mayoral, Compact sets of compact operators in absence of ℓ_1 , *Proc. Amer. Math. Soc.* **129** (2001) 79-82.
- [18] E. Serrano, C. Piñeiro, and J. M. Delgado, Equicompact sets of operators defined on Banach spaces, *Proc. Amer. Math. Soc.* **134** (2006), 689-695.
- [19] E. Serrano, C. Piñeiro, and J. M. Delgado, Weakly Equicompact sets of operators defined on Banach spaces, *Arch. Math.* **86** (2006), 231-240.
- [20] E. Serrano, C. Piñeiro, and J. M. Delgado, Some properties and applications of equicompact sets of operators, *Studia. Math.* **181** (2) (2007).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ISFAHAN
E-mail address: m2020alikhani@ yahoo.com