

Variability of paths and differential equations with BV -coefficients

Michael Hinz* and Jonas M. Tölle† and Lauri Viitasaari‡

March 27, 2020

Abstract

We define compositions $\varphi(X)$ of Hölder paths X in \mathbb{R}^n and functions of bounded variation φ under a relative condition involving the path and the gradient measure of φ . We show the existence and properties of generalized Lebesgue-Stieltjes integrals of compositions $\varphi(X)$ with respect to a given Hölder path Y . These results are then used, together with Doss' transform, to obtain existence and, in a certain sense, uniqueness results for differential equations in \mathbb{R}^n driven by Hölder paths and involving coefficients of bounded variation. Examples include equations with discontinuous coefficients driven by paths of two-dimensional fractional Brownian motions.

Mathematics Subject Classification (2010): 31B10, 34A12, 34A34 (primary); 26A33, 26A42, 26B30, 26B35, 28A78, 31B99, 60G22 (secondary).

Keywords: functions of bounded variation; generalized Lebesgue-Stieltjes integrals; occupation measure; Hölder path; Riesz potential; systems of nonlinear differential equations.

Contents

1 Introduction	2
1.1 Related literature	3
1.2 Brief description of our approach	4
1.3 Structure of the article	5

*Fakultät für Mathematik, Universität Bielefeld, Germany, mhinz@math.uni-bielefeld.de. Research supported in part by the DFG IRTG 2235 ‘Searching for the regular in the irregular: Analysis of singular and random systems’ and by the DFG CRC 1283, ‘Taming uncertainty and profiting from randomness and low regularity in analysis, stochastics and their applications’.

†Department of Mathematics and Statistics, University of Helsinki, Finland, jonas.tolle@helsinki.fi. JMT would like to thank Panu Lahti and Mario Santilli for sharing their insight into BV functions. Financial support by the Dean’s Office, Faculty of Mathematics and Economics, Universität Ulm is gratefully acknowledged.

‡Department of Information and Management Science, Aalto University School of Business, Finland, lauri.viitasaari@iki.fi.

2	Stieltjes integrals and BV-coefficients	6
2.1	Compositions of paths with BV -functions	6
2.2	Existence and properties of Stieltjes integrals	8
3	Systems of differential equations	12
3.1	Upper regularity and Hölder continuity of coefficients	13
3.2	Solutions in dimension one	14
3.3	Doss' transformation	15
3.4	Solutions in arbitrary dimensions	18
3.5	A uniqueness result	20
4	Variability and consequences	21
4.1	Riesz potentials and occupation measures	21
4.2	Upper regularity and bounded potentials	24
4.3	Fourier transform and trading of regularity	29
4.4	Probabilistic examples	29
4.5	A brief discussion of variability versus irregularity	33
4.6	Compositions of BV_{loc} -functions and Hölder paths	34
4.7	Proof of existence and regularity of the integral	40
4.8	Interpretation as currents	42
4.9	Proof of the change of variable formula	43
5	Existence and uniqueness proofs	45
5.1	Invertibility and BV -regularity	45
5.2	Solutions to the deterministic equation	46
5.3	Existence of solutions in the invertible case	48
5.4	Uniqueness of solutions in the invertible case	50
A	Convolution of Riesz kernels	56
B	Mollification results	57
C	Some properties of maximal functions	59
D	Elements of fractional calculus	61

1 Introduction

We prove new results on the existence and regularity of generalized Lebesgue-Stieltjes integrals

$$\int_0^t \varphi(X_u) dY_u, \quad t \in [0, T], \quad (1.1)$$

as in [73, 90, 91], where $X : [0, T] \rightarrow \mathbb{R}^n$ and $Y : [0, T] \rightarrow \mathbb{R}$ are Hölder continuous functions with sum of Hölder orders greater than one and $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}$ is a function locally of bounded variation, [5, 94], possibly discontinuous. We then employ these results to study equations in \mathbb{R}^n of form

$$X_t = \dot{x} + \int_0^t \sigma(X_u) dY_u, \quad t \in [0, T], \quad (1.2)$$

where Y is a given path in \mathbb{R}^n , Hölder of order $\gamma > \frac{1}{2}$, and σ is a (bounded) matrix valued function of locally bounded variation. We implement a Doss transform, [24, 87], and use it to construct Hölder continuous solutions X to (1.2), unique in a certain class. This produces novel first results for discontinuous coefficients σ in dimensions $n \geq 2$. The main difficulties are to provide a meaningful definition of the compositions $\varphi(X)$ (resp. $\sigma(X)$) and to show they are regular enough for the integrals in (1.1) or (1.2), respectively, to make sense and for the Doss transformation method [24, 87] to work. Our main tool is a quantitative condition which ensures that X spends little time in regions where the gradient measure of φ (resp. σ) is very concentrated.

1.1 Related literature

To study equations of type (1.2) for deterministic integrators Y of low regularity or for probabilistic integrators Y lacking semimartingale or other good distributional properties, the use of Stieltjes type integrals, [73, 89, 90, 91], and much more generally, the theory of rough paths, [41, 65, 66, 67], have become established tools. However, rather little is known about equations with irregular diffusion coefficients σ , and we are only aware of the few references mentioned below. Stochastic differential equations with respect to Brownian motion involving non-Lipschitz (drift or diffusion) coefficients can be discussed in several different ways, [75]. Most classical and recent results for singular, irregular, or degenerate coefficients, and notions of uniqueness such as [22, 30, 34, 56, 61, 83, 93], and results specific to the one-dimensional case, [26, 27, 60, 71], are built upon the connection to diffusion theory and partial differential equations. For equations (1.2) driven by rough deterministic or fractional Gaussian signals Y , such tools are not available.

For an integrator Y that is Hölder of order $\gamma > \frac{1}{2}$, Peano existence for solutions to (1.2) in \mathbb{R}^n is well known for coefficients σ that are s -Hölder continuous provided that $\gamma > \frac{1}{1+s}$. Moreover, Picard existence and uniqueness holds if the coefficient is $\mathcal{C}^{1,s}$ (with the same s), [41, 67]. See [68, 73, 90, 91] for the more classical Lipschitz resp. \mathcal{C}^2 -cases. In [62], new results have been obtained for equations (1.2) in \mathbb{R}^n for the case $\gamma < \frac{1}{1+s}$. For $n = 1$ and continuous coefficients σ whose reciprocal is integrable on compact intervals around zero, the authors of [62] constructed solutions to (1.2) by means of a Lamperti transform [58], see [62, Theorem 3.7]. For $n \geq 1$ they can solve (1.2) if the components of the coefficient σ are bounded from below by $|x|^s$, their gradients are Hölder continuous of order $\frac{1}{\gamma} - 1$ away from zero, and the integral is understood in terms of Riemann sum approximations, [62, Theorem 4.15]. The first results on the existence of Stieltjes integrals with discontinuous coefficients (in the case $n = 1$) were obtained in [18] (see also [17, Chapter 5]). There the authors proved the existence of (1.1) if φ is of locally finite variation and X is a sufficiently active path, [18, Theorem 3.1 and Remark 3.3]. For random X an integrability assumptions on its probability densities ensures this condition. They also prove a change of variable formula and several results on the approximation of (1.1) by Riemann-Stieltjes sums. A first study for differential equations (1.2) was provided in [38], where the authors prove existence and uniqueness of solutions to (1.2) for $n = 1$ if Y is a fractional Brownian motion with Hurst index greater $\frac{1}{2}$ and φ

is a (scaled) Heaviside step function. The authors of [38] used a Lamperti transform and smoothing arguments. Merging the assumptions from [18] and the transform used in [38], the authors of [81] were able to prove existence and uniqueness for (1.2) in the case $n = 1$ and in a probabilistic setup. Finally, we mention [88], where an alternative existence proof for integrals of type (1.1) was given using Riemann-Stieltjes approximations and suitable controls (avoiding fractional calculus), extending the results of [18].

1.2 Brief description of our approach

In the present paper we use a quantitative condition on the given individual path X w.r.t. the given coefficient φ (or σ), which we call (s, p) -variability, Definition 2.1. It may be seen as an deterministic version of the probabilistic Assumption 2.1 of [81], and as a higher dimensional analog of a condition in [88, Corollary 3]. Our first main result is Theorem 2.13, where we state that the composition $\varphi(X)$ of φ with a Hölder path that is $(s, 1)$ -variable w.r.t. φ is well defined and a member of a certain fractional Sobolev space, ensuring the existence of (1.1). The stronger assumption of (s, p) -variability with large p guarantees that (1.1) is Hölder continuous. A key step to obtain these results is a multiplicative estimate for Gagliardo seminorms of $\varphi(X)$, Proposition 4.27, that can be viewed as a generalization of [18, Proposition 4.6] ([81, Proposition 4.1]) to higher space dimensions. To obtain this estimate, one bounds differences of type $|\varphi(X_t) - \varphi(X_u)|$ in terms of a fractional maximal functions of the total variation $\|D\varphi\|$ of the gradient measure of φ , Proposition C.1; this is a fractional version of a prominent argument, [20, Lemma A.3]. Then one estimates further using the fact that the fractional maximal functions of order $1 - s$ are trivially bounded by Riesz potentials of $\|D\varphi\|$ of order $(1 - s)$, evaluated at X_t (resp. X_u). The (s, p) -variability condition just means that these functions have the desired integrability in time. The $(s, 1)$ -variability of X w.r.t. φ is tantamount to saying that the total variation $\|D\varphi\|$ of the gradient measure of φ and the occupation measure $\mu_X^{[0, T]}$ on $[0, T]$ have a finite mutual Riesz energy of order $1 - s$, see Proposition 4.2. Phenomenologically this means that these two measures are sufficiently disperse with respect to each other to make the singular (repulsive) interaction kernel of order $-n + 1 - s$ integrable, which is a polarized version of well known arguments, see [28, 59, 69] for background and [42] for a related application. Mutual Riesz energies are not necessarily easy to handle, but they encode dimensional properties in a neat way and this permits to easily connect to the well known scaling properties of gradient measures of BV -functions, [5, Section 3.9], and well known scaling properties of fractal curves, [28], such as realizations of prominent stochastic processes, [7, 28, 50, 86]. If in a certain region of space φ has a jump or strong oscillation so that $\|D\varphi\|$ is too concentrated, this can be compensated if X is so fast moving in that part of space that the Hausdorff dimension of $\mu_X^{[0, T]}$ is sufficiently high to guarantee sufficient integrability, see Corollary 4.15. The motivation that increased activity of a path can compensate low coefficient regularity is also central in regularization by noise, [16, 22, 32, 34, 56], see Remark 4.3 below and also Subsection 4.1, where we compare $(s, 1)$ -variability to a condition used in [16]. We begin our discussion of differential equations (1.2) by

showing that a uniform boundedness condition on the Riesz potential of order $1 - s$ of the total variations of the gradient measures naturally takes us back to the case of γ -Hölder coefficients with $\gamma > \frac{1}{1-s}$, so that a well known Peano existence argument applies, Theorem 3.6. This is the extreme case, where the activity of the solution path is not used. To obtain existence results taking into account the activity of the solution path, we implement a Doss transform for BV -coefficients σ under the main assumptions that σ is invertible, Assumption 3.12, its inverse has curl free columns, Assumption 3.15, and a bounded distortion condition holds. Of course in particular the curl free condition is quite restrictive, but as in classical implementations of the Doss method, [87], it is inevitable. However, in lack of other existence results for equations with BV -coefficients it seems reasonable to establish a BV -variant of Doss' transformation under these assumptions. They guarantee the existence of a Lipschitz function f so that, roughly speaking, a solution X is obtained as an image of the driver Y under f . This uses the fact that dimensional lower bounds for Y are stable under Lipschitz transformations and produces our second main result, Theorem 3.24, which states the existence of Hölder continuous solutions X to (1.2) with BV -coefficients σ . A one-dimensional version of this result under less restrictive assumptions is formulated in Theorem 3.8. These theorems are purely deterministic, the regularization effect of the irregular path is rather mild, and the dimensional hypotheses are too restrictive to allow satisfactory result in cases where Y is taken to be, say, a fixed path of a fractional Brownian motion in \mathbb{R}^n when $n \geq 2$. In our third main result, Theorem 3.25, we assume that Y is a stochastic process satisfying a specific bound for the mean value of a Riesz potential of its occupation measure and that the gradient measures of the coefficient obey a specific moment condition, and obtain Hölder continuous solutions for almost every realization of Y . One can regard Theorem 3.25 as (a partial) extension of the probabilistic [81, Theorem 2.1]. Our fourth main result is a related uniqueness result, Theorem 3.28. It shows that Assumptions 3.12 and 3.15 guarantee uniqueness in the class of variability solutions.

1.3 Structure of the article

The structure of the article is as follows: In Section 2 we introduce the notion of (s, p) -variability, define compositions $\varphi(X)$ ($\sigma(X)$, respectively), and state our results on existence and properties of (1.1). We also provide a change of variable formula and a result on Riemann sum approximation. Section 3 contains our results on existence and uniqueness of variability solutions to (1.2). In Section 4 we provide a systematic discussion of $(s, 1)$ -variability, some of its immediate consequences, conditions sufficient to ensure it, and some probabilistic examples. The mentioned multiplicative estimate, the properties of (1.1) and the change of variable formula, are verified, and we point out a connection to currents. The Doss transformation and the claimed existence and uniqueness results for (1.2) are proved in Section 5. Basic facts on Riesz kernels, mollification, maximal functions, and fractional calculus are collected in Appendices.

By $|\cdot|$ we denote the Euclidean norm in \mathbb{R}^n . We write $B(x, r)$ for the open ball of radius $r > 0$ centered at $x \in \mathbb{R}^n$. The symbol \mathcal{L}^n stands for the n -dimensional Lebesgue measure and the symbol \mathcal{H}^d for the d -dimensional Hausdorff measure on \mathbb{R}^n . For spaces

of \mathbb{R}^m -valued functions we use notations like $BV(\mathbb{R}^n)^m$ (to stay close to reference [5]) or $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ (because it is more practical for other function spaces). For $m = 1$, we suppress \mathbb{R}^m from notation and write $L^1_{\text{loc}}(\mathbb{R}^n)$.

2 Stieltjes integrals and BV -coefficients

2.1 Compositions of paths with BV -functions

Recall that a function $\varphi \in L^1_{\text{loc}}(\mathbb{R}^n)$ is of *locally bounded variation*, denoted $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$, if its distributional partial derivatives $D_i\varphi$ are signed Radon measures, $i = 1, \dots, n$. We write $D\varphi = (D_1\varphi, \dots, D_n\varphi)$ for its \mathbb{R}^n -valued gradient measure, and $\|D\varphi\|$ for the total variation of $D\varphi$. If $\varphi \in L^1(\mathbb{R}^n)$ and $\|D\varphi\|(\mathbb{R}^n) < +\infty$, then φ is said to be of *bounded variation*, $\varphi \in BV(\mathbb{R}^n)$.

Let $T > 0$. We consider continuous paths from $[0, T]$ into \mathbb{R}^n , that is, continuous functions $X = (X^1, \dots, X^n) : [0, T] \rightarrow \mathbb{R}^n$. The following definition is our key tool to provide a meaningful and sufficiently regular definition of the composition $\varphi \circ X$ of a BV_{loc} -function φ and a path X . As usual, $L^p(0, T)$ denotes the Lebesgue space of classes of p -integrable functions on $(0, T)$.

Definition 2.1. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$, $p \in [1, +\infty]$ and $s \in (0, 1)$. We say that a path $X = (X^1, \dots, X^n) : [0, T] \rightarrow \mathbb{R}^n$ is (s, p) -variable with respect to φ if there is a relatively compact open neighborhood \mathcal{U} of $X([0, T])$ such that*

$$\int_{\mathcal{U}} |X - z|^{-n+1-s} \|D\varphi\|(dz) \in L^p(0, T). \quad (2.1)$$

We write $V(\varphi, s, p)$ for the class of paths X that are (s, p) -variable w.r.t. φ and use the short notation $V(\varphi, s) := V(\varphi, s, 1)$.

Note that $V(\varphi, s, p) \subset V(\varphi, s, q)$ for any $q < p$ and $V(\varphi, s, p) \subset V(\varphi, r, p)$ for any $r < s$. The (s, p) -variability condition (2.1) is a quantitative and relative condition on the path X and the function φ . Roughly speaking, it ensures that X varies sufficiently around sites where φ has strong oscillations or jumps, encoded in the requirement that the Riesz potential of order $1 - s$ of the restriction of $\|D\varphi\|$ to \mathcal{U} is in $L^p(0, T)$, see Section 4 for a systematic discussion. The use of an open neighborhood \mathcal{U} of $X([0, T])$ in (2.1) allows mollification arguments.

We admit a component-wise point of view upon functions with values in \mathbb{R}^m .

Definition 2.2. *Let $\varphi = (\varphi_1, \dots, \varphi_m) \in BV_{\text{loc}}(\mathbb{R}^n)^m$, $p \in [1, +\infty]$ and $s \in (0, 1)$. We say that a path $X : [0, T] \rightarrow \mathbb{R}^n$ is (s, p) -variable with respect to φ if it is (s, p) -variable with respect to each component φ_i of φ . That is, $V(\varphi, s, p) := \bigcap_{i=1}^m V(\varphi_i, s, p)$. Similarly as before, we write $V(\varphi, s) := V(\varphi, s, 1)$.*

Recall the following classical definition.

Definition 2.3. A function $\varphi \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ is said to have an approximate limit at $x \in \mathbb{R}^n$ if there exists $\lambda_\varphi(x) \in \mathbb{R}^m$ such that

$$\lim_{r \rightarrow 0} \frac{1}{\mathcal{L}^n(B(x, r))} \int_{B(x, r)} |\varphi(y) - \lambda_\varphi(x)| dy = 0.$$

In this situation, the unique value $\lambda_\varphi(x)$ is called the approximate limit of φ at x . The set of points $x \in \mathbb{R}^n$ for which this property does not hold is called approximate discontinuity set (or exceptional set) and is denoted by S_φ .

The set S_φ does not depend on the choice of the representative for φ . If $\tilde{\varphi}$ is a representative of $\varphi \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ then a point $x \notin S_\varphi$ with $\tilde{\varphi}(x) = \lambda_\varphi(x)$ is called a *Lebesgue point* of $\tilde{\varphi}$, and the set of all Lebesgue points of $\tilde{\varphi}$ is called the *Lebesgue set* of φ . See for instance [5, Definition 3.63]. The set S_φ is Borel and of zero Lebesgue measure, [5, Proposition 3.64]. If $\varphi = (\varphi_1, \dots, \varphi_m) \in BV(\mathbb{R}^n)^m$ then by the Federer-Vol'pert theorem, [5, Theorem 3.78], the set S_φ is countably \mathcal{H}^{n-1} -rectifiable.

We say that a Borel function $\tilde{\varphi} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a *Lebesgue representative* of $\varphi = (\varphi_1, \dots, \varphi_m) \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ if

$$\tilde{\varphi}(x) = \lambda_\varphi(x), \quad x \in \mathbb{R}^n \setminus S_\varphi. \quad (2.2)$$

Using Definition 2.3 and the equivalence of norms on \mathbb{R}^n it is easy to see that

$$S_\varphi = \bigcup_{i=1}^m S_{\varphi_i}. \quad (2.3)$$

In particular, if for any i the function $\tilde{\varphi}_i : \mathbb{R}^n \rightarrow \mathbb{R}$ is a Lebesgue representative of φ_i then $(\tilde{\varphi}_1, \dots, \tilde{\varphi}_m)$ is a Lebesgue representative of φ , and we refer to such representatives as *component-wise Lebesgue representatives*.

The following observation will be proved in Section 4.

Lemma 2.4. Let $\varphi = (\varphi_1, \dots, \varphi_m) \in BV_{\text{loc}}(\mathbb{R}^n)^m$ and $X \in V(\varphi, s)$ for some $s \in (0, 1)$. Then for any component-wise Lebesgue representatives $\tilde{\varphi}^{(1)}$ and $\tilde{\varphi}^{(2)}$ of φ we have

$$\tilde{\varphi}^{(1)}(X_t) = \tilde{\varphi}^{(2)}(X_t)$$

at \mathcal{L}^1 -a.e. $t \in [0, T]$.

Lemma 2.4 could be rephrased by saying that under the $(s, 1)$ -variability condition the equivalence class φ has a well defined trace on the range $X([0, T]) \subset \mathbb{R}^n$ of X , endowed with a suitable measure. See [2] or [49] for pointwise redefinitions of functions and traces to closed subsets of \mathbb{R}^n in other contexts.

Definition 2.5. Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)^m$ and suppose that $X \in V(\varphi, s)$ for some $s \in (0, 1)$. We define the composition $\varphi \circ X$ to be the \mathcal{L}^1 -equivalence class of $t \mapsto \tilde{\varphi}(X_t)$ on $[0, T]$, where $\tilde{\varphi}$ is a component-wise Lebesgue representative of φ . Given $p \in [1, +\infty]$ we say that φ is p -integrable w.r.t. X , in symbols $\varphi \in L^p(X, \mathbb{R}^m)$, if $\varphi \circ X$ is an element of $L^p(0, T, \mathbb{R}^m)$.

Thanks to Lemma 2.4 the composition and the notion of p -integrability w.r.t. X are well defined. The component-wise choice of representatives is not essential, but it is convenient in conjunction with Definitions 2.1 and 2.2.

We discuss $(s, 1)$ -variability in some examples.

Example 2.6. If $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}$ is locally Lipschitz, then for any $s \in (0, 1)$ any path X is in $V(\varphi, s, \infty)$. This follows from the fact that $\|D\varphi\| = |\nabla f| \cdot \mathcal{L}^n$ with $|\nabla f| \in L^\infty(\mathcal{U})$ on any relatively compact open set $\mathcal{U} \subset \mathbb{R}^n$.

Remark 2.7. Note that saying *any* path includes the constant path.

Example 2.8. Let $\mathcal{C} \subset [0, 1]$ be the classical middle third Cantor set and $\nu_{\mathcal{C}}$ the unique self-similar probability measure with support \mathcal{C} , see [28]. Let $\varphi_{\mathcal{C}} : \mathbb{R}^n \rightarrow [0, 1]$ be the function that satisfies $\varphi_{\mathcal{C}}(x) = \nu_{\mathcal{C}}((0, x_1))$ for all $x = (x_1, x_2, \dots, x_n) \in [0, 1] \times \mathbb{R}^{n-1}$, $\varphi_{\mathcal{C}}(x) = 0$ for all $x \in (-\infty, 0) \times \mathbb{R}^{n-1}$ and $\varphi_{\mathcal{C}}(x) = 1$ for all $x \in (1, +\infty) \times \mathbb{R}^{n-1}$. Then $\varphi_{\mathcal{C}} \in BV_{\text{loc}}(\mathbb{R}^n)$, and on $[0, 1]^n$ we have $\|D\varphi_{\mathcal{C}}\| = D\varphi_{\mathcal{C}} = \nu_{\mathcal{C}} \otimes \mathcal{H}^{n-1}$. Writing $d_{\mathcal{C}} = \frac{\log 2}{\log 3}$ for the Hausdorff dimension of \mathcal{C} , we find that for $s \in (0, d_{\mathcal{C}})$ any path X in \mathbb{R}^n is in $V(\varphi_{\mathcal{C}}, s, \infty)$. Now suppose $s \in (d_{\mathcal{C}}, 1)$. The constant path $X \equiv (\frac{1}{2}, 0, \dots, 0)$ in \mathbb{R}^n is in $V(\varphi_{\mathcal{C}}, s, \infty)$, but the constant path $X \equiv (0, 0, \dots, 0)$ is not in $V(\varphi_{\mathcal{C}}, s)$. For $n = 1$ any smooth function $X : (0, T) \rightarrow (0, 1)$ with a finite number of critical points is in $V(\varphi_{\mathcal{C}}, s, \infty)$. For $n = 2$ a smooth curve $X : (0, T) \rightarrow (0, 1)^2$, parametrized to have unit speed, does not have to be in $V(\varphi_{\mathcal{C}}, s)$. On the other hand, a path of Brownian motion is in $V(\varphi_{\mathcal{C}}, s, \infty)$ with probability one. For $n \geq 3$ paths of fractional Brownian motions with Hurst index $H \in (0, \frac{1}{n-1+s})$ are in $V(\varphi_{\mathcal{C}}, s, \infty)$ with probability one. See Example 4.17 and Subsection 4.4 for details.

The function φ in Example 2.8 is Hölder continuous. The next example discussed variability with respect to discontinuous functions.

Example 2.9. Suppose that $D \subset \mathbb{R}^n$ is a smooth domain with $\mathcal{H}^{n-1}(\partial D) < +\infty$. By [5, Proposition 3.62] the function $\mathbf{1}_D$ is in $BV(\mathbb{R}^n)$ and D has finite perimeter $\mathcal{P}(D) = \|D\mathbf{1}_D\| < +\infty$. Let $s \in (0, 1)$ be arbitrary. If a smooth unit speed curve $X : [0, T] \rightarrow \mathbb{R}^n$ hits ∂D in finitely many points then we have $X \in V(\mathbf{1}_D, s)$, but if $n \geq 2$ and X spends \mathcal{L}^1 -positive time in ∂D then it cannot be an element of $V(\mathbf{1}_D, s)$, see Example 4.18. For $n = 1$ or $n = 2$ the path of a Brownian motion is in $V(\mathbf{1}_D, s, \infty)$ with probability one. For arbitrary $n \geq 1$, the path of a fractional Brownian motion with Hurst index $H \in (0, \frac{1}{n-1+s})$ is in $V(\mathbf{1}_D, s, \infty)$ with probability one. For arbitrary $n \geq 1$ it also follows that if $H \in (0, \frac{1}{s})$ and the fractional Brownian motion is started in $(\partial D)^c$ then it is in $V(s, 1)$ with probability one, see Subsection 4.4.

2.2 Existence and properties of Stieltjes integrals

As mentioned, we are interested in generalized Lebesgue-Stieltjes integrals defined in terms of fractional calculus, [76], introduced in [90, 91] and used e.g. in [73].

We introduce suitable function spaces to discuss the existence and the continuity properties of the integral. Let $p \in [1, +\infty)$ and $0 < \theta < 1$. The Gagliardo seminorm of

order θ with exponent p of a measurable function $f : (0, T) \rightarrow \mathbb{R}^m$ is defined as

$$[f]_{\theta,p} = \left(\int_0^T \int_0^T \frac{|f(t) - f(u)|^p}{|t - u|^{1+\theta p}} du dt \right)^{\frac{1}{p}}. \quad (2.4)$$

By $W^{\theta,p}(0, T, \mathbb{R}^m)$ we denote the space of measurable functions $f : (0, T) \rightarrow \mathbb{R}^m$ such that

$$\|f\|_{W^{\theta,p}(0,T,\mathbb{R}^m)} := \|f\|_{L^p(0,T,\mathbb{R}^m)} + [f]_{\theta,p} < +\infty.$$

Recall that for $m = 1$ we agreed to suppress \mathbb{R}^m from notation and simply write $W^{\theta,p}(0, T)$, which we do similarly for the spaces in the sequel. The Hölder seminorm of order $0 < \theta < 1$ of a measurable function $f : [0, T] \rightarrow \mathbb{R}^m$ is denoted by

$$[f]_{\theta,\infty} = \sup_{0 \leq u < t \leq T} \frac{|f(t) - f(u)|}{|t - u|^\theta},$$

and we write $\mathcal{C}^\theta([0, T], \mathbb{R}^m)$ to denote the space of Hölder continuous functions $f : [0, T] \rightarrow \mathbb{R}^m$, endowed with the norm

$$\|f\|_{\mathcal{C}^\theta([0,T],\mathbb{R}^m)} := \sup_{t \in [0,T]} |f(t)| + [f]_{\theta,\infty}.$$

Remark 2.10. The spaces $W^{\theta,p}(0, T)$ and $\mathcal{C}^\theta([0, T])$ are classical Besov spaces of type $B_{p,p}^\theta$ and $B_{\infty,\infty}^\theta$, see for instance [82].

Because they appear naturally in connection with Stieltjes integrals we also consider the following more specific types of spaces, which (in this or a similar form) were introduced in [73]. Accepting a slight abuse of notation we use the symbol $W^{\theta,\infty}(0, T, \mathbb{R}^m)$ to denote the space of all measurable $f : (0, T) \rightarrow \mathbb{R}^m$ such that

$$\|f\|_{W^{\theta,\infty}(0,T,\mathbb{R}^m)} := \|f\|_{L^\infty(0,T,\mathbb{R}^m)} + \operatorname{ess\,sup}_{t \in [0,T]} \int_0^t \frac{|f(t) - f(u)|}{|t - u|^{1+\theta}} du < +\infty.$$

By $W_T^{\theta,\infty}(0, T, \mathbb{R}^m)$ we denote the space of measurable functions $f : (0, T) \rightarrow \mathbb{R}^m$ such that $f(T-) \in \mathbb{R}^m$ exists and

$$\|f\|_{W_T^{\theta,\infty}(0,T,\mathbb{R}^m)} = \sup_{t \in (0,T)} \frac{|f(T-) - f(t)|}{(T-t)^\theta} + \sup_{t \in (0,T)} \int_t^T \frac{|f(t) - f(u)|}{|t - u|^{1+\theta}} du < \infty.$$

It is well known and easily seen that

$$\mathcal{C}^{\theta+\varepsilon}([0, T], \mathbb{R}^m) \subset W^{\theta,\infty}(0, T, \mathbb{R}^m) \quad \text{and} \quad \mathcal{C}^{\theta+\varepsilon}([0, T], \mathbb{R}^m) \subset W_T^{\theta,\infty}(0, T, \mathbb{R}^m). \quad (2.5)$$

We write $W_0^{\theta,p}(0, T, \mathbb{R}^m)$ for the space of measurable functions $f : (0, T) \rightarrow \mathbb{R}^m$ such that

$$\|f\|_{W_0^{\theta,p}(0,T,\mathbb{R}^m)} := \int_0^T \frac{|f(t)|^p}{t^{\theta p}} dt + [f]_{\theta,p} < +\infty.$$

We emphasize that in the present paper the symbols $W^{\theta,\infty}(0,T)$ and $W_0^{\theta,p}(0,T)$ do not have the standard meaning.

The following definition is due to [90, 91], see also [73]. By D_{0+}^θ and $D_{T-}^{1-\theta}$ we denote the (left and right sided) fractional Weyl-Marchaud derivatives of orders θ and $1-\theta$, respectively, see [76, 73, 90, 91] or Appendix D for details.

Definition 2.11. *Let $f \in W_0^{\theta,1}(0,T)$ and $g \in W_T^{1-\theta,\infty}(0,T)$ for some $\theta \in (0,1)$. Then we define the integral by*

$$\int_0^T f_t dg_t = (-1)^\theta \int_0^T D_{0+}^\theta f(t) D_{T-}^{1-\theta}(g - g(T-))(t) dt. \quad (2.6)$$

This definition is correct, that is, independent of the choice of θ , [90, Proposition 2.1]. If f and g are as in the definition and g has bounded variation, then (2.6) equals to the classical Lebesgue-Stieltjes integral of f w.r.t. g , [90, Theorem 2.4]. The following duality estimate and restriction property are well known, see [90, 91] or [73]. By Γ we denote the Euler Gamma function.

Proposition 2.12. *Assume that $f \in W_0^{\theta,1}(0,T)$ and $g \in W_T^{1-\theta,\infty}(0,T)$ for some $\theta \in (0,1)$. Then the integral in (2.6) admits the bound*

$$\left| \int_0^T f_t dg_t \right| \leq \frac{\|f\|_{W_0^{\theta,1}} \|g\|_{W_T^{1-\theta,\infty}}}{\Gamma(\theta)\Gamma(1-\theta)}. \quad (2.7)$$

For every $t \in [0,T]$ the restriction $\mathbf{1}_{[0,t]}f$ belongs to $W_0^{\theta,1}(0,T)$ and the integral

$$\int_0^t f_u dg_u = \int_0^T \mathbf{1}_{[0,t]}(u) f_u dg_u$$

is well-defined.

Our first new contribution is the following Theorem 2.13 on the regularity and integrability of the composition $\varphi \circ X$ of a Hölder path X with a BV -function φ , the existence of the generalized Lebesgue-Stieltjes integral

$$\int_0^t \varphi(X_u) dY_u \quad (2.8)$$

of $\varphi(X)$ with respect to a given Hölder path Y in the sense of (2.6), and the Hölder regularity of this integral, seen as a function of $t \in [0,T]$.

Theorem 2.13. *Suppose that $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ and that $X \in \mathcal{C}^\alpha([0,T], \mathbb{R}^n)$ is $(s,1)$ -variable with respect to φ for some $s \in (0,1)$.*

- (i) *For any $0 < \beta < \alpha s$ the composition $\varphi \circ X$ is an element of $W_0^{\beta,1}(0,T)$.*
- (ii) *If in addition $Y \in \mathcal{C}^\gamma([0,T], \mathbb{R})$ and $1 - \alpha s < \gamma$ then for any $t \in [0,T]$ the integral (2.8) exists.*

(iii) Moreover, if X is (s, p) -variable with respect to φ for some $p \in (1, +\infty]$, then for any $0 < \beta < \alpha s$ we have $\varphi \circ X \in W^{\beta, p}(0, T)$. If in addition $Y \in \mathcal{C}^\gamma([0, T], \mathbb{R})$ with $\frac{1}{p} < 1 - \beta < \gamma$, where we write $\frac{1}{+\infty} := 0$, then

$$\int_0^\cdot \varphi(X_u) dY_u \in \mathcal{C}^{1-\beta-1/p}([0, T], \mathbb{R})$$

and

$$\left\| \int_0^\cdot \varphi(X) dY \right\|_{\mathcal{C}^{1-\beta-1/p}([0, T])} \leq c \|\varphi \circ X\|_{W^{\beta, p}(0, T)} \|Y\|_{\mathcal{C}^\gamma([0, T])}. \quad (2.9)$$

In Subsections 4.6 and 4.7 of Section 4 we provide a proof of Theorem 2.13, along with quantitative estimates for the integral (2.8) involving (2.1).

Variability permits the following change of variable formula.

Theorem 2.14. *Let $F \in W_{\text{loc}}^{1,1}(\mathbb{R}^n)$ be such that $\partial_k F \in BV_{\text{loc}}(\mathbb{R}^n)$ for $k = 1, \dots, n$. If $X \in \mathcal{C}^\alpha([0, T], \mathbb{R}^n)$ with $\alpha > \frac{1}{2}$ is a path which is $(s, 1)$ -variable w.r.t. each $\partial_k F$ for some $s \in (\frac{1-\alpha}{\alpha}, 1)$, then we have*

$$F(X_t) = F(\dot{x}) + \sum_{k=1}^n \int_0^t \partial_k F(X_u) dX_u^k \quad (2.10)$$

for \mathcal{L}^1 -a.e. $t \in [0, T]$, provided that $\dot{x} \in \mathbb{R}^n \setminus S_F$. If in addition F is continuous, then (2.10) holds for all $t \in [0, T]$ and no matter where $\dot{x} \in \mathbb{R}^n$ is located.

The proof of this theorem can be found in Subsection 4.9 of Section 4.

The following result on the coincidence of (2.8) and the corresponding Riemann-Stieltjes integral under the (s, p) -variability condition with large enough p is immediate from Proposition 4.27 and Lemma 4.29 below together with [90, Theorems 4.1.1 and 4.2.1]. We provide this result for systematic reasons while it will not be used in the sequel.

Theorem 2.15. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$, let $X \in \mathcal{C}^\alpha([0, T], \mathbb{R}^n)$ be a path which is (s, p) -variable with respect to φ for some $s \in (0, 1)$ and $p \in (\frac{1}{\alpha s}, +\infty]$. Then $\varphi \circ X$ is Hölder continuous of any order less than $\alpha s - \frac{1}{p}$. If in addition $Y \in \mathcal{C}^\gamma([0, T], \mathbb{R})$ for some $\gamma > 1 - \alpha s + \frac{1}{p}$, where $\frac{1}{+\infty} := 0$, then both the generalized Lebesgue-Stieltjes integral $\int_0^T \varphi(X_u) dY_u$ as in (2.8) and the Riemann-Stieltjes integral of $\varphi(X)$ w.r.t. Y over $[0, T]$ exist and agree.*

Moreover, if in this case we are given $0 < \varepsilon < \alpha s - 1/p - 1 + \gamma$, a refining sequence $(\pi_k)_{k \geq 1}$ of finite partitions $\pi_k = \{0 = t_0^{(k)} < t_1^{(k)} < \dots < t_{N_k}^{(k)} = T\}$ of $[0, T]$ and points $\xi_i^{(k)} \in [t_{i-1}^{(k)}, t_i^{(k)}]$, then we have

$$\left| \int_0^T \varphi(X) dY - \sum_{i=1}^{N_k} \varphi(X_{\xi_i^{(k)}}) (Y_{t_i^{(k)}} - Y_{t_{i-1}^{(k)}}) \right| \leq c |\pi_k|^{\alpha s - 1/p - 1 + \gamma - \varepsilon}$$

for all k , where $|\pi_k| = \max_i (t_i^{(k)} - t_{i-1}^{(k)})$ denotes the mesh of the partition π_k and $c > 0$ is a constant depending on α, γ, s, p and Y .

3 Systems of differential equations

In this section we discuss equations of form

$$X_t = \hat{x} + \int_0^t \sigma(X_u) dY_u, \quad t \in [0, T], \quad (3.1)$$

where $T > 0$, $\hat{x} \in \mathbb{R}^n$, $\sigma : \mathbb{R}^n \mapsto \mathbb{R}^{n \times n}$ is a coefficient function $\sigma = (\sigma_{jk})_{1 \leq j, k \leq n}$ and $Y = (Y^1, \dots, Y^n) : [0, T] \rightarrow \mathbb{R}^n$ is a given Hölder path. As usual, (3.1) is to be understood in the sense that all components X^j of $X = (X^1, \dots, X^n) : [0, T] \rightarrow \mathbb{R}^n$ satisfy the equations

$$X_t^j = \hat{x}_j + \sum_{k=1}^n \int_0^t \sigma_{jk}(X_u^1, \dots, X_u^n) dY_u^k, \quad (3.2)$$

where $\hat{x} = (\hat{x}_1, \dots, \hat{x}_n)$. Each integral in these sums is a generalized Lebesgue-Stieltjes integral as in (2.8).

We employ Definition 2.2 for matrix coefficients: Given $\sigma = (\sigma_{jk})_{1 \leq j, k \leq n}$ with $\sigma_{jk} \in BV_{\text{loc}}(\mathbb{R}^n)$ for all j and k and parameters $s \in (0, 1)$ and $p \in [1, +\infty]$, a path $X = (X^1, \dots, X^n) : [0, T] \mapsto \mathbb{R}^n$ is called (s, p) -variable with respect to σ if it is (s, p) -variable with respect to all σ_{jk} . We consider the following notion of solution.

Definition 3.1. *Let $\sigma = (\sigma_{jk})_{1 \leq j, k \leq n}$ be such that $\sigma_{jk} \in BV_{\text{loc}}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ for all j and k and let $Y \in \mathcal{C}^\gamma([0, T], \mathbb{R}^n)$. A path $X : [0, T] \rightarrow \mathbb{R}^n$ is called a variability solution for σ and Y started at $\hat{x} \in \mathbb{R}^n$ if*

- (i) $X_0 = \hat{x}$
- (ii) the path X is in $\mathcal{C}^\alpha([0, T], \mathbb{R}^n)$ and also in $V(\sigma, s)$ for some $s \in (\frac{1-\gamma}{\alpha}, 1)$
- (iii) X satisfies (3.1).

Examples 3.2. Let $n = 1$ and let $\sigma = \varphi_\ell$ be as in Example 2.8 (with $n = 1$). If Y is a typical realization of fractional Brownian motion with Hurst index $H > \frac{\log 3}{\log 3 + \log 2}$ then by Theorem 3.6 below, variability solutions for σ and Y started at zero exist. Also the zero function $X \equiv 0$ is a variability solution.

Not every solution is a variability solution as the following example shows.

Examples 3.3. If $n = 1$, $\kappa \in (0, 1)$, $\sigma(x) = |x|^\kappa$ for $x \in (-1, 1)$ and $\sigma(x) \equiv 1$ for $x \in \mathbb{R} \setminus (-1, 1)$ and (nowhere constant) $Y \in \mathcal{C}^\gamma([0, T])$ then by Theorem 3.8 there are (non-constant) variability solutions for σ and Y started at zero. Also the zero function $X \equiv 0$ is a solution, but it can be a variability solution only if $\gamma + \kappa > 1$: Since in a neighborhood of zero $\|D\sigma\|(dx) = |x|^{\kappa-1} dx$ the zero solution is a variability solution only if we can find $s \in (1 - \gamma, 1)$ such that $|x|^{-s+\kappa-1}$ is integrable around the origin.

Remark 3.4. In [62] the authors considered monotone and continuous σ having power-type non-linearity at the origin similarly as in the preceding example. The assumptions of [62, Theorem 3.6] ensure that their solution candidate is actually an $(s, 1)$ -variable solution. As pointed out in [62], one cannot expect unique solutions, because also the zero function is a solution.

3.1 Upper regularity and Hölder continuity of coefficients

To put our results in perspective we begin with a case that encodes a known result.

Recall that, given $d \geq 0$, a Borel measure μ on \mathbb{R}^n is said to be *upper d -regular on a Borel set $B \subset \mathbb{R}^n$* if there are constants $c > 0$ and $r_0 > 0$ such that

$$\mu(B(x, r)) \leq cr^d, \quad x \in B \cap \text{supp } \mu, \quad 0 < r < r_0.$$

We call a function $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ *upper d -regular on B* if $\|D\varphi\|$ is *upper d -regular on B* . We say that a function $\varphi = (\varphi_1, \dots, \varphi_m) \in (BV_{\text{loc}}(\mathbb{R}^n))^m$ is *upper d -regular on B* if each of its components φ_i is. If $B = \mathbb{R}^n$ we simply say *upper d -regular*.

The following is verified in Proposition 4.8.

Proposition 3.5. *Let $\sigma = (\sigma_{jk})_{1 \leq j, k \leq n}$ be such that $\sigma_{jk} \in BV_{\text{loc}}(\mathbb{R}^n)$ for all j and k , let $s \in (0, 1)$, and assume that σ is upper d -regular with $d > n - 1 + s$. Then σ is Hölder continuous of order s . Moreover, any path is (s, ∞) -variable w.r.t. σ .*

Concerning (3.1) we are led back to a well known Peano type existence result.

Theorem 3.6. *Let $\sigma = (\sigma_{jk})_{1 \leq j, k \leq n}$ be such that $\sigma_{jk} \in BV_{\text{loc}}(\mathbb{R}^n)$ for all j and k , let $s \in (0, 1)$, and assume that σ is upper d -regular with $d > n - 1 + s$. If $Y \in \mathcal{C}^\gamma([0, T], \mathbb{R}^n)$ for some $\gamma \in (\frac{1}{1+s}, 1)$, then for any $\dot{x} \in \mathbb{R}^n$ there exists a variability solution $X \in \mathcal{C}^\gamma([0, T], \mathbb{R}^n)$ for σ and Y starting at \dot{x} , and $X \in V(\sigma, s, \infty)$.*

For a proof, see for instance [41, Proposition 5] or, in a slightly different formulation, [67, Theorem 1.20].

We also consider upper regularity conditions for paths. If for given $s > 0$ and $B \subset \mathbb{R}^n$ Borel a path $Y : [0, T] \rightarrow \mathbb{R}^n$ satisfies

$$\sup_{x \in Y([0, T]) \cap B} \int_0^T |Y_t - x|^{-s} dt < +\infty, \quad (3.3)$$

then there are constants $c > 0$ and $r_0 > 0$ such that

$$\mathcal{L}^1(\{t \in [0, T] : |Y_t - Y_u| < r\}) \leq cr^s, \quad u \in \{\tau \in [0, T] : Y_\tau \in B\}, \quad 0 < r < r_0. \quad (3.4)$$

If (3.4) holds with some $d > s$ in place of s , then (3.3) holds. These facts are standard, see Proposition 4.11. Condition (3.4) says that the occupation measure of Y on $[0, T]$ is upper s -regular on B , and for $B = \mathbb{R}^n$ this encodes a lower bound on its Hausdorff dimension, see Remark 4.10. If (3.4) holds, we also say that Y is *upper s -regular on B* and *upper s -regular* if $B = \mathbb{R}^n$.

Remark 3.7.

- (i) No non-constant $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ can be upper d -regular with $d > n$, as a Lebesgue differentiation argument shows.

- (ii) Suppose $n = 1$. Then we trivially have $\dim_H Y([0, T]) \leq 1$ for any path Y , with equality if Y is non-constant. For a constant path clearly $\dim_H Y([0, T]) = 0$. If Y is upper d -regular, then $d \leq \dim_H Y([0, T])$, [28, Theorem 4.13]. A path Y constant on some nonempty open interval cannot be upper d -regular for any $d > 0$. A nowhere constant Lipschitz path Y on $[0, T]$ that is upper d -regular for a given number $d \in (0, 1]$ but not upper d' -regular for any $d' > d$ is $Y_t = t^{\frac{1}{d}}$, $t \in [0, T]$. In this case we have $\mu_Y^{[0, T]}([0, r]) = r^d$ for any $0 < r < T^{\frac{1}{d}}$.

3.2 Solutions in dimension one

For the case $n = 1$ we obtain the following slight modification of the constructive existence results [38, Theorem 3.3] and [81, Theorem 2.1], see Example 3.9 below. It allows to compensate the failure of σ to be Hölder of sufficiently high order in some space region by sufficient activity of Y in some other space region, stated in terms of (3.3).

Theorem 3.8. *Let $\sigma \in BV_{\text{loc}}(\mathbb{R}) \cap L^\infty(\mathbb{R})$ be nonnegative \mathcal{L}^1 -a.e. and such that $\frac{1}{\sigma} \in L^1_{\text{loc}}(\mathbb{R})$.*

- (i) *The function $g(x) = \int_0^x \frac{dz}{\sigma(z)}$, $x \in \mathbb{R}$, is absolutely continuous and strictly increasing on \mathbb{R} . Its inverse $f := g^{-1}$ is Lipschitz and satisfies $\sigma(f) = f'$ \mathcal{L}^1 -a.e. on \mathbb{R} .*
- (ii) *Let $s \in (0, 1)$, $\gamma \in (\frac{1}{1+s}, 1)$, $Y \in \mathcal{C}^\gamma([0, T])$ with $Y_0 = 0$ and $\hat{x} \in \mathbb{R}$. Let $-\infty \leq a < b \leq +\infty$. Suppose that Y satisfies (3.3) for $B = (g(a), g(b))$ or that σ is upper d -regular on (a, b) with $d > s$, and similarly for $\mathbb{R} \setminus (g(a), g(b))$ and $\mathbb{R} \setminus (a, b)$, respectively. Then the function*

$$X_t = f(Y_t + g(\hat{x})), \quad t \in [0, T],$$

is a variability solution $X \in \mathcal{C}^\gamma([0, T]) \cap V(\sigma, s)$ for σ and Y started at \hat{x} .

In the case that in Theorem 3.8 no upper regularity of σ is used but Y is assumed to be upper d_Y -regular, the Hölder continuity and upper regularity of Y together can produce sufficient regularity: The hypotheses in Theorem 3.8 force the condition

$$\frac{1}{\gamma} < 1 + d_Y, \quad (3.5)$$

and for $s \in (\frac{1}{\gamma} - 1, d_Y)$ the theorem yields solutions. If the Hölder order γ of Y is higher (lower), lower (higher) upper regularity of Y is needed.

Example 3.9. A real valued fractional Brownian motion $(B_t^H)_{t \in [0, T]}$ of arbitrary Hurst index $H \in (0, 1)$ is almost surely γ -Hölder for any $\gamma < H$. On the other hand, it is upper 1-regular almost surely, as can for instance be concluded from the joint continuity of its local times, see [11, p. 1271], [12, Theorem] or [84], which entails their local boundedness, see Proposition 4.13 below. If $H > \frac{1}{2}$ then for any $\gamma \in (\frac{1}{2}, H)$ condition (3.5) holds and for almost every realization (seen as a deterministic path) in place of Y , Theorem 3.8 yields variability solutions for any starting point \hat{x} and no matter whether

σ is upper regular of any order or not. This recovers the fractional Brownian motion special case of [81, Theorem 2.1], which roughly speaking made use of the facts that, for any $\gamma < H$, the paths of B^H are γ -Hölder almost surely and that

$$\sup_{x \in \mathbb{R}} \mathbb{E} \int_0^T |Y_t - x|^{-\frac{1}{H}} dt < +\infty, \quad (3.6)$$

[81, Assumption 2.1 and Example 2.1]. Condition (3.6) implies that with probability one we have $Y \in V(\sigma, \frac{1}{H}, 1)$ for any BV -function σ .

Example 3.10. Functions $Y : [0, 1] \rightarrow \mathbb{R}$ having Hölder and upper regularity properties satisfying the assumptions of Theorem 3.8 can be found in the class of self-affine functions [51, Definition 1]. A self-affine function of order $H \in (0, 1]$ is Hölder continuous of order H but of no order $H' > H$, see [51, Theorem 1] or [53]. For any choice of integers $2 \leq b < a$ one can construct self-affine functions Y of order $H = \frac{\log b}{\log a}$ which have bounded local times L_1^Y , see [13, Proposition 7 et sa preuve], and for any such Y Theorem 3.8 yields variability solutions no matter whether σ is upper regular of any order or not. Although Theorem 3.8 does not apply to this case, it is interesting to note that one can find self-affine functions of order $H = \frac{1}{2}$ having square integrable local times, [13, p. 438, Remarques], what ensures Y is upper $\frac{1}{2}$ -regular, Proposition 4.13.

Example 3.11. Suppose that $\sigma(x) = \varphi_{\mathcal{E}}(\frac{x}{3})\mathbf{1}_{[0,3]}(x) + 4 \cdot \mathbf{1}_{(3,+\infty)}(x)$, $x \geq 0$, where $\varphi_{\mathcal{E}}$ is as in Example 2.8, and $\sigma(x) := \sigma(-x)$, $x < 0$. Then $\sigma(0) = 0$ and σ is $d_{\mathcal{E}}$ -Hölder except at $x = \pm 3$, where it jumps. Since $\nu_{\mathcal{E}}$ is lower $d_{\mathcal{E}}$ -regular, we have $\varphi_{\mathcal{E}}(r) \geq cr^{d_{\mathcal{E}}}$ for small $r > 0$, which implies that σ^{-1} is locally integrable. Since $\nu_{\mathcal{E}}$ is upper $d_{\mathcal{E}}$ -regular, σ is upper $d_{\mathcal{E}}$ -regular on $(-2, 2)$. Let $d, \delta \in (0, 1]$ and let $Z : [0, 2] \rightarrow \mathbb{R}$ be the function defined by $Z_t = t^{\frac{1}{d}}$ for $t \in (0, 1]$ and $Z_t = (1 - (1 - t))^{\delta}$ for $t \in (1, 2]$. Let g be as in Theorem 3.8 and suppose that Y is such that on each of the intervals $[0, \frac{1}{2}g(1)]$, $[\frac{1}{2}g(1), \frac{3}{4}g(1)]$, $[\frac{3}{4}g(1), \frac{7}{8}g(1)]$ etc. it equals a scaled down copy of Z , and on $[g(1), +\infty)$ it equals a typical path of fractional Brownian motion with Hurst index H started at zero. Then Y is Hölder of any order $\gamma < H \wedge \delta$, upper d -regular on $(g(-1), g(1))$ and upper 1-regular on $\mathbb{R} \setminus (g(-1), g(1))$. If $d = \frac{1}{2}$ and $\delta = H = \frac{3}{4}$ then Theorem 3.8 yields variability solutions for all starting points, note that on $(-2, 2)$ we can use the upper d_{σ} -regularity of σ and on $\mathbb{R} \setminus (-g(1), g(1))$ the upper 1-regularity of Y .

3.3 Doss' transformation

To understand analogs of Theorem 3.8 for $n \geq 2$ we implement a multidimensional Doss transformation, [24, 87]. Let $\sigma : \mathbb{R}^n \rightarrow \mathbb{R}^{n \times n}$ be a matrix coefficient as before, and let $\sigma_j = (\sigma_{j1}, \dots, \sigma_{jn})$ denote its j th row. Suppose that $f = (f_1, \dots, f_n) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a function with components $f_k : \mathbb{R}^n \mapsto \mathbb{R}$ which satisfies the deterministic equation

$$\nabla f = \sigma(f), \quad (3.7)$$

where we have used the symbol

$$\nabla f = (\partial_k f_j)_{1 \leq j, k \leq n}$$

for the Jacobian matrix of f to avoid confusion with the symbol for gradient measures. The j th row of ∇f is the gradient $\nabla f_j = (\partial_1 f_j, \dots, \partial_n f_j)$ of f_j , and (3.7) states that $\nabla f_j = \sigma_j(f)$ for all j . If we set

$$X_t^j = f_j(Y_t^1, \dots, Y_t^n),$$

then $X = (X^1, \dots, X^n)$ is a solution to (3.1) since

$$\begin{aligned} X_t^j &= f_j(Y_0^1, \dots, Y_0^n) + \sum_{k=1}^n \int_0^t \partial_k f_j(Y_u^1, \dots, Y_u^n) dY_u^k \\ &= f_j(Y_0^1, \dots, Y_0^n) + \sum_{k=1}^n \int_0^t \sigma_{jk}(f_1(Y_u^1, \dots, Y_u^n), \dots, f_n(Y_u^1, \dots, Y_u^n)) dY_u^k \quad (3.8) \\ &= X_0^j + \sum_{k=1}^n \int_0^t \sigma_{jk}(X_u^1, \dots, X_u^n) dY_u^k, \end{aligned}$$

provided that we can justify the use of the change of variable formula (2.10) with f_j in place of F and (3.7). As in the classical case, [87], one needs strong assumptions to find solutions f to (3.7), and in our case an additional difficulty is that the components of σ are BV_{loc} only.

Our first main assumption on σ is as follows.

Assumption 3.12.

- (i) $\sigma = (\sigma_{jk})_{1 \leq j, k \leq n}$ with components $\sigma_{jk} \in BV_{\text{loc}}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ for all j and k ,
- (ii) $\det(\sigma) > \varepsilon$ \mathcal{L}^n -a.e. on \mathbb{R}^n for some $\varepsilon > 0$.

Remark 3.13. Since all paths are continuous and T is finite, it suffices to have σ with the respective properties defined on a bounded domain in \mathbb{R}^n . To save notation and have shorter proofs we formulate Assumption 3.12 as stated.

The following lemma is a direct consequence of Assumption 3.12 and the Cayley-Hamilton theorem, see Subsection 5.1.

Lemma 3.14. *Let Assumption 3.12 be satisfied. Then there exists a matrix $\hat{\sigma} = (\hat{\sigma}_{jk})_{1 \leq j, k \leq n}$ of functions $\hat{\sigma}_{jk} \in BV_{\text{loc}}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ so that*

$$\sigma \hat{\sigma} = \hat{\sigma} \sigma = I \quad \mathcal{L}^n\text{-a.e.}$$

and this matrix is unique up to \mathcal{L}^n -a.e. equivalence. Moreover, there exists some $\hat{\varepsilon} > 0$ such that $\det(\hat{\sigma}) > \hat{\varepsilon}$ \mathcal{L}^n -a.e. on \mathbb{R}^n .

By Assumption 3.12 and Lemma 3.14 our second main assumption makes sense. We write D_i to denote the partial differentiation in direction of the i th coordinate in the sense of tempered distributions.

Assumption 3.15. For all i, j , and k we have

$$D_i \hat{\sigma}_{kj} - D_j \hat{\sigma}_{ki} = 0 \quad (3.9)$$

in the sense of tempered distributions.

Example 3.16. Let σ be a diagonal matrix with entries $\sigma_{ii} \in BV(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ such that $\sigma_{ii} > \varepsilon$ \mathcal{L}^1 -a.e. on \mathbb{R} for some $\varepsilon > 0$ and for all $i = 1, 2, \dots, n$. Then Assumption 3.12 is clearly satisfied. If also Assumption 3.15 is satisfied, it follows that σ_{ii} depends only on x_i . Consequently, we are led back to one-dimensional equations as discussed in Subsection 3.2.

Example 3.17. Let $n = 2$, $c > 1$ and consider

$$\sigma(x_1, x_2) = \begin{pmatrix} c & 1 \\ \mathbf{1}_{\{cx_1 < x_2\}} & c \end{pmatrix}, \quad (x_1, x_2) \in \mathbb{R}^2.$$

The function σ is bounded and in $BV_{\text{loc}}(\mathbb{R}^2)$, and it jumps across the straight line $J = \{x_2 = cx_1\}$. Obviously $\det \sigma \geq c^2 - 1$, and we find

$$\hat{\sigma}(x_1, x_2) = \frac{\mathbf{1}_{\{cx_1 < x_2\}}}{c^2 - 1} \begin{pmatrix} c & -1 \\ -1 & c \end{pmatrix} + \frac{\mathbf{1}_{\{cx_1 \geq x_2\}}}{c^2} \begin{pmatrix} c & -1 \\ 0 & c \end{pmatrix}.$$

Since $\hat{\sigma}_{11}$, $\hat{\sigma}_{12}$ and $\hat{\sigma}_{22}$ are constant and by [5, (3.90)] the measure $D\hat{\sigma}_{21}$ is a constant multiple of $\mathcal{H}^1|_J$, (3.9) trivially holds.

Assumptions 3.12 and 3.15 allow the following result for solutions to (3.7).

Proposition 3.18. *Suppose that σ satisfies Assumption 3.12. For $n \geq 2$ suppose also that it satisfies Assumption 3.15 and that there exists $\delta > -1$ such that for \mathcal{L}^n -a.e. $x \in \mathbb{R}^n$ we have*

$$\langle \xi, \sigma(x) \xi \rangle \geq \delta |\sigma(x) \xi| |\xi|, \quad \xi \in \mathbb{R}^n. \quad (3.10)$$

Then there exists a bi-Lipschitz function $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ which solves (3.7).

Condition (3.10) ensures the applicability of a global inversion result, Proposition 5.4, which yields a global Lipschitz solution f to (3.7).

Remark 3.19. Let $n \geq 2$. Then $\sigma \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^{n \times n})$ satisfies (3.10) if and only if $\text{spec}(\sigma(x)) \cap (-\infty, 0) = \emptyset$ for \mathcal{L}^n -a.e. $x \in \mathbb{R}^n$, where $\text{spec}(\sigma(x))$ denotes the spectrum of the square matrix $\sigma(x)$, see [54, p. 3], [6] and [64, Proposition 2.1]. This holds in particular if σ has an \mathcal{L}^n -a.e. nonnegative definite symmetric part.

Example 3.20. The coefficient σ in Example 3.17 is positive definite, and hence (3.10) is immediate. Moreover, the function

$$f(y_1, y_2) = (cy_1 + y_2, y_1 \mathbf{1}_{\{y_1 < 0\}} + cy_2), \quad (y_1, y_2) \in \mathbb{R}^2,$$

solves (3.7) for this σ .

Example 3.21. Let $n = 2$, $0 < a < b$ and consider the cone $C = \{\frac{a}{b}x_1 < x_2 < \frac{b}{a}x_1\}$. The function

$$\sigma(x_1, x_2) = \begin{pmatrix} b & a\mathbf{1}_C(x_1, x_2) \\ a\mathbf{1}_C(x_1, x_2) & b \end{pmatrix}, \quad (x_1, x_2) \in \mathbb{R}^2,$$

satisfies Assumptions 3.12 and 3.15 and (3.10). A solution to (3.7) is given by

$$f(y_1, y_2) = (by_1 + ay_2\mathbf{1}_{\{y_1 > 0, y_2 > 0\}}, ay_1\mathbf{1}_{\{y_1 > 0, y_2 > 0\}} + by_2), \quad (y_1, y_2) \in \mathbb{R}^2.$$

Example 3.22. Let $n = 2$, let $\varphi_{\mathcal{C}} : \mathbb{R} \rightarrow \mathbb{R}$ be as in Example 2.8 (with $n = 1$), and $\Phi(z) = \int_0^z (1 + \varphi_{\mathcal{C}}(z))dz$, $z \in \mathbb{R}$. The function

$$\begin{aligned} \sigma(x_1, x_2) = \mathbf{1}_{\{x_1 \leq 0\}} & \begin{pmatrix} 1 & 0 \\ 0 & 1 + \varphi_{\mathcal{C}}(\Phi^{-1}(x_2)) \end{pmatrix} \\ & + \mathbf{1}_{\{x_1 > 0\}} \begin{pmatrix} 1 & 0 \\ 1 & 1 + \varphi_{\mathcal{C}}(\Phi^{-1}(x_2 - x_1)) \end{pmatrix}, \quad (x_1, x_2) \in \mathbb{R}^2, \end{aligned}$$

satisfies Assumptions 3.12 and 3.15 and (3.10), the fact that $\sigma_{22} \in BV_{\text{loc}}(\mathbb{R}^n)$ follows using [5, Theorem 3.16]. For this choice of σ the function

$$f(y_1, y_2) = (y_1, y_1\mathbf{1}_{\{y_1 > 0\}} + \Phi(y_2)), \quad (y_1, y_2) \in \mathbb{R}^2,$$

solves (3.7).

Remark 3.23. The existence of almost everywhere defined local inverses of Sobolev functions is treated in [37, 36]. In [8, Section 5] fine properties of a.e. local inverses are studied under the requirement that $\det(\hat{\sigma}(x)) > 0$ for \mathcal{L}^n -a.e. $x \in \mathbb{R}^n$. The local inverse becomes an everywhere defined local homeomorphism on open subsets where the determinant of $\hat{\sigma}$ equals a positive constant, see [36, Corollary 6.3].

3.4 Solutions in arbitrary dimensions

The following is a version of Theorem 3.8 for arbitrary $n \geq 1$.

Theorem 3.24. *Suppose that σ is as in Assumption 3.12 and $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a bi-Lipschitz function which solves (3.7). Let $s \in (0, 1)$, $\gamma \in (\frac{1}{1+s}, 1)$, $Y \in \mathcal{C}^\gamma([0, T], \mathbb{R}^n)$ with $Y_0 = 0$, $B \subset \mathbb{R}^n$ is a Borel and $\hat{x} \in \mathbb{R}^n$. Suppose that σ is upper d -regular on B with $d > n - 1 + s$ or that*

$$\sup_{x \in Y([0, T]) \cap (f^{-1}(B) - f^{-1}(\hat{x}))} \int_0^T |Y_t - x|^{-n+1-s} dt < +\infty, \quad (3.11)$$

and suppose the same is true for B^c in place of B . Then the path

$$X_t = f(Y_t + f^{-1}(\hat{x})), \quad t \in [0, T], \quad (3.12)$$

is a variability solution $X \in \mathcal{C}^\gamma([0, T], \mathbb{R}^n) \cap V(\sigma, s)$ for σ and Y started at \hat{x} .

By Proposition 3.18, the first sentence in Theorem 3.24 could be replaced by requiring σ to satisfy Assumptions 3.12 and, if $n \geq 2$, also Assumption 3.15 and (3.10). Note that for $n = 1$, Theorem 3.8 gives the same result under less restrictive assumptions on σ .

If only the upper regularity of σ is used, Theorem 3.24 complements Theorem 3.6 by constructing a solution. If also the upper regularity of Y is used one can, in some cases, obtain solutions for discontinuous σ . However, the upper regularity condition on Y is quite restrictive: Already if $n = 2$ and σ has jumps one cannot hope to use Theorem 3.24 to obtain solutions to (3.1) when Y is a typical path of a fractional Brownian motion B^H with Hurst index $H > \frac{1}{2}$: On the one hand, B^H is almost surely γ -Hölder of any order $\gamma < H$, what forces $\frac{1}{H} < 1 + s$. On the other hand, the Hausdorff dimension $\dim_H B^H([0, T])$ of $B^H([0, T])$ is known to equal $\frac{1}{H}$ almost surely, see [28, Chapter 11] or [50, Section 18.3, Theorem 1 and Corollary], so that we would run into the contradictive chain of inequalities $s < d_Y \leq \dim_H Y([0, T]) = \frac{1}{H} < 1 + s$. For $n = 1$ one has $\dim_H B^H([0, T]) = 1$ almost surely, and this contradiction does not occur.

In comparison with (3.6) condition (3.3) is quite restrictive. It ensures a rather weak regularizing effect solely due to a dimensional lower bound. Condition (3.6) encodes a strong additional regularization by randomness. The following result may be seen as a generalization of [38, Theorem 3.3], and [81, Theorem 2.1]. It uses a weighted version of condition (3.3) together with a moment condition on the gradient measures of the coefficient.

Theorem 3.25. *Suppose that σ is as in Assumption 3.12 and $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a bi-Lipschitz function which solves (3.7). Let $s \in (0, 1)$, $\gamma \in (\frac{1}{1+s}, 1)$, let $Y = (Y_t)_{t \in [0, T]}$ be an \mathbb{R}^n -valued stochastic process with $Y_0 = 0$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with paths \mathbb{P} -a.s. Hölder continuous of order γ , and let $\hat{x} \in \mathbb{R}^n$. Suppose that there are $\varepsilon \in (0, 1 - s)$, $c > 0$, and $\delta \in (0, n - 1 + s - \varepsilon)$ such that*

$$\mathbb{E} \int_0^T |Y_t - x|^{-n+\varepsilon} dt < c|x|^{-n+\delta}, \quad x \in \mathbb{R}^n, \quad (3.13)$$

and for all j and k we have

$$\int_{\mathbb{R}^n} |x - \hat{x}|^{-n+1-s-\varepsilon+\delta} \|D\sigma_{jk}\| (dx) < +\infty. \quad (3.14)$$

Then for \mathbb{P} -a.e. $\omega \in \Omega$ the path

$$X_t(\omega) = f(Y_t(\omega) + f^{-1}(\hat{x})), \quad t \in [0, T], \quad (3.15)$$

is a variability solution $X \in \mathcal{C}^\gamma([0, T], \mathbb{R}^n) \cap V(\sigma, s)$ for σ and Y started at \hat{x} .

At any time $t > 0$ the averaging effect of the process Y lead to a regularization. The moment condition (3.14) excludes a too bad behavior of σ at the starting point \hat{x} at time $t = 0$ when this effect is not yet present.

Example 3.26. Let $n = 2$ and let $Y = B^H$ be a fractional Brownian motion with Hurst index $H > \frac{1}{2}$. It satisfies (3.13) with $0 < \varepsilon < 1 - \frac{1}{H}$ and $\delta = \frac{1}{H} + \varepsilon$, see Example 4.24. If σ satisfies Assumptions 3.12 and 3.15, (3.10), and

$$\int_{\mathbb{R}^n} |x - \hat{x}|^{-n+1} \|D\sigma_{jk}\| (dx) < +\infty, \quad j, k = 1, 2, \quad (3.16)$$

then for \mathbb{P} -a.s. realization of Y the path X as in Theorem 3.25 is a variability solution for σ and Y started at \hat{x} , and $X \in \mathcal{C}^\gamma([0, T], \mathbb{R}^2) \cap V(\sigma, \frac{1}{H})$ for any $\gamma < H$. If σ is as in Example 3.17, then (3.16) holds for all starting points $\hat{x} \in \mathbb{R}^n \setminus J$ off the line J on which σ jumps. If σ is as in Example 3.21, then (3.16) holds for all starting points $\hat{x} \in \mathbb{R}^2 \setminus \partial C$, where ∂C is the boundary of the cone C . If σ is as in Example 3.22 and $\hat{x} = (\hat{x}_1, \hat{x}_2)$ is such that $\hat{x}_1 < 0$ and $\hat{x}_2 \in \mathcal{C}$, where \mathcal{C} is the middle third Cantor set, then (3.16) holds.

Remark 3.27. Theorem 3.25 is stated in the above form because our main examples are probabilistic. Basically the same proof yields a deterministic counterpart: Suppose that σ and f are as in Theorem 3.25, $s \in (0, 1)$, $\gamma \in (\frac{1}{1+s}, 1)$, $Y \in \mathcal{C}^\gamma([0, T], \mathbb{R}^n)$ is a path with $Y_0 = 0$ and $\hat{x} \in \mathbb{R}^n$. If ε , c , and δ are as in Theorem 3.25,

$$\int_0^T |Y_t - x|^{-n+\varepsilon} dt < c|x|^{-n+\delta}, \quad x \in \mathbb{R}^n,$$

and (3.14) holds, then (3.12) is a variability solution $X \in \mathcal{C}^\gamma([0, T], \mathbb{R}^n) \cap V(\sigma, s)$ for σ and Y started at \hat{x} .

3.5 A uniqueness result

We establish a uniqueness result for variability solutions.

Theorem 3.28. *Suppose that $\sigma = (\sigma_{jk})_{1 \leq j, k \leq n}$ satisfies Assumptions 3.12 and 3.15, $Y \in \mathcal{C}^\gamma([0, T], \mathbb{R}^n)$ for some $\gamma \in (0, 1)$, and $\hat{x} \in \mathbb{R}^n$. Then there exists at most one variability solution of Hölder order greater $\frac{1}{2}$ for σ and Y started at \hat{x} .*

The proof of Theorem 3.28 shows that the only solution candidate is (3.12). We can conclude the following results.

Corollary 3.29. *Suppose that Assumptions 3.12 and 3.15 hold and let f be the solution to (3.7). If assumptions of Theorem 3.24 (or Theorem 3.8 in the case $n = 1$) are satisfied, then for any $\hat{x} \in \mathbb{R}^n$ there exists a unique variability solution $X \in \mathcal{C}^\gamma([0, T], \mathbb{R}^n)$ for σ and Y starting at \hat{x} , given by (3.12).*

Corollary 3.30. *Suppose that Assumptions 3.12 and 3.15 hold and let f be the solution to (3.7). If assumptions of Theorem 3.25 are satisfied, then for any $\hat{x} \in \mathbb{R}^n$ and \mathbb{P} -a.e. $\omega \in \Omega$ there exists a unique variability solution $X(\omega) \in \mathcal{C}^\gamma([0, T], \mathbb{R}^n)$ for σ and $Y(\omega)$ starting at \hat{x} , given by (3.15).*

Remark 3.31. Following the ideas of [81], we could replace $\det(\sigma) > \varepsilon$ in Assumption 3.12 with $\det(\sigma) \geq 0$. In this case we would obtain a uniqueness result similar to Theorem 3.28 up to the first time when $\det(\sigma(X_t)) = 0$. However, the existence of the solution becomes more complicated.

4 Variability and consequences

We provide a potential theoretic interpretation of the $(s, 1)$ -variability. This allows to prove that $\varphi \circ X$ is a well defined class in $W_0^{\beta,1}(0, T)$.

4.1 Riesz potentials and occupation measures

The *Riesz potential of order* $0 < \gamma < n$ of a non-negative Borel measure ν on \mathbb{R}^n is defined by

$$U^\gamma \nu(x) := \int_{\mathbb{R}^n} |x - y|^{-n+\gamma} \nu(dy), \quad x \in \mathbb{R}^n, \quad (4.1)$$

see for instance [59, Chapter I, Section 3] or [15, Section V.4]. The *mutual Riesz energy of order* $0 < \gamma < n$ of two non-negative Borel measures ν_1 and ν_2 on \mathbb{R}^n is defined by

$$I^\gamma(\nu_1, \nu_2) := \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |x - y|^{-n+\gamma} \nu_1(dy) \nu_2(dx). \quad (4.2)$$

This quantity takes values in $[0, +\infty]$, is symmetric in ν_1 and ν_2 , and we have

$$I^\gamma(\nu_1, \nu_2) = \int_{\mathbb{R}^n} U^\gamma \nu_1(x) \nu_2(dx),$$

see [59, Chapter I, Section 4]. We write $I^\gamma \nu := I^\gamma(\nu, \nu)$.

Given a path $X = (X_t)_{t \in [0, T]}$ we can define a finite non-negative Radon measure $\mu_X^{[0, T]}$ on \mathbb{R}^n by the identity

$$\int_{\mathbb{R}^n} g(y) \mu_X^{[0, T]}(dy) = \int_0^T g(X_t) dt, \quad (4.3)$$

valid for any bounded Borel function $g : \mathbb{R}^n \rightarrow \mathbb{R}$. To $\mu_X^{[0, T]}$ one usually refers as the *occupation measure of X* and to (4.3) as the *occupation time formula*. The measure $\mu_X^{[0, T]}$ is supported on the compact set $X([0, T])$. Clearly the occupation measure depends on the parametrization of the path X .

Remark 4.1.

- (i) Discussions of absolutely continuous occupation measures in terms of their densities, referred to as *local times*, are a classical subject in probability theory, [10, 11, 39, 86]. As mentioned in [39, Section 3], applications of occupation densities to nonrandom functions are much less discussed. Results on possibly singular occupation measures are rather sparse and have mainly been used to obtain results on dimensions of images or graphs of stochastic processes, see for instance [28, Section 16] and [85, Section 4] and the references cited there.
- (ii) In the special case that X is an absolutely continuous curve parametrized to have unit speed the right hand side of (4.3) is just the line integral of g along X , and

formula (4.3) may be seen as a special case of the area formula for the path X , [29, Theorem 3.2.6]. Another interpretation of formulas of type (4.3) in the spirit of geometric measure theory have been established in [14]. There the authors proved an occupation time formula for \mathbb{R}^n -valued semimartingales which has features of a coarea formula for $\mathcal{C}^2(\mathbb{R}^n)$ -functions. They observed that the fluctuations of the semimartingale can lead to the existence of certain ‘transversal’ (in a Rokhlin sense) densities which may be seen as generalizations of local times of one-dimensional semimartingales, see [14, Theorems 1 and 3].

The definitions (4.1) and (4.2) and the identity (4.3) now immediately allow to rewrite (2.1) in terms of Riesz energies. Given a function $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ and a set $\mathcal{U} \subset \mathbb{R}^n$ we use

$$\mu_{\varphi, \mathcal{U}} := \|D\varphi\| \llcorner_{\mathcal{U}}$$

to abbreviate the restriction of $\|D\varphi\|$ to \mathcal{U} .

Proposition 4.2. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$, $p \in [1, +\infty]$ and $s \in (0, 1)$. A path X is (s, p) -variable with respect to φ if and only if there is a relatively compact open neighborhood \mathcal{U} of $X([0, T])$ such that*

$$U^{1-s} \mu_{\varphi, \mathcal{U}} \in L^p(\mathbb{R}^n, \mu_X^{[0, T]}). \quad (4.4)$$

In particular, X is $(s, 1)$ -variable with respect to φ if and only if there is a relatively compact open neighborhood of $X([0, T])$ such that the mutual Riesz energy of order $1 - s$ of $\mu_{\varphi, \mathcal{U}}$ and $\mu_X^{[0, T]}$ is finite,

$$\int_{\mathbb{R}^n} U^{1-s} \mu_{\varphi, \mathcal{U}}(x) \mu_X^{[0, T]}(dx) < +\infty. \quad (4.5)$$

Remark 4.3.

- (i) Both ideas, the variability of curves and the use of Riesz energies, also appear in connection with regularization properties of operators in harmonic analysis, see for instance [80] for the first idea and [40, 43] for the second.
- (ii) Occupation measures appear naturally in connection with *regularization by noise*, [16, 22, 32, 34, 56, 72, 83]. The addition (in the simplest case) of a ‘fast moving’ perturbation to an equation can provoke well-posedness for otherwise non well-posed equations. A typical key step is to observe the Hölder (or even $\mathcal{C}^{1+\alpha}$ -) regularity of integrals of functions (with certain continuity or integrability properties) w.r.t. occupation measures, seen as a function of the starting point of the path. See for instance [32, Section 2.1 and in particular, Theorems 2.5 and 2.6].

The following consequence of Proposition 4.2 allows to define compositions of paths and BV -functions. It views (4.5) as a condition on X relative to φ .

Corollary 4.4. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ and suppose that X is $(s, 1)$ -variable with respect to φ for some $s \in (0, 1)$. Then the discontinuity set S_φ of φ is a zero set for the occupation measure $\mu_X^{[0, T]}$ of X ,*

$$\mu_X^{[0, T]}(S_\varphi) = 0. \quad (4.6)$$

In particular, we have

$$\mathcal{L}^1(\{t \in [0, T] : X_t \in S_\varphi\}) = 0 \quad (4.7)$$

and

$$\|\varphi\|_{L^\infty(X)} \leq \|\varphi\|_{L^\infty(\mathbb{R}^n)}. \quad (4.8)$$

Corollary 4.4 immediately implies Lemma 2.4, and therefore the correctness of Definition 2.5.

Proof of Lemma 2.4. If $\tilde{\varphi}^{(1)}$ and $\tilde{\varphi}^{(2)}$ both are (component-wise) Lebesgue representatives of $\varphi = (\varphi_1, \dots, \varphi_m) \in BV_{\text{loc}}(\mathbb{R}^n)^m$ and X is $(s, 1)$ -variable w.r.t. each φ_i , then by (2.2), (2.3) and (4.7) we have

$$\int_0^T |\tilde{\varphi}^{(1)}(X_t) - \tilde{\varphi}^{(2)}(X_t)| dt = \int_{\{t \in [0, T] : X_t \in \mathbb{R}^n \setminus S_\varphi\}} |\tilde{\varphi}^{(1)}(X_t) - \tilde{\varphi}^{(2)}(X_t)| dt = 0.$$

□

Recall that for any $\gamma \geq 0$ the upper γ -density $\Theta_\gamma^* \nu(x)$ of a Borel measure ν on \mathbb{R}^n at a point $x \in \mathbb{R}^n$ is defined by

$$\Theta_\gamma^* \nu(x) = \limsup_{r \rightarrow 0} \frac{\nu(B(x, r))}{r^\gamma},$$

see for instance [5, Definition 2.2.5] or [69, Definition 6.8]. We prove Corollary 4.4.

Proof of Corollary 4.4. Let $\mathcal{U} \subset \mathbb{R}^n$ be a relatively compact open set containing the support $X([0, T])$ of $\mu_X^{[0, T]}$ as in (4.5). Since the integral is finite we can find a Borel set $N \subset \mathbb{R}^n$ such that $\mu_X^{[0, T]}(N) = 0$ and, for all $x \in \mathbb{R}^n \setminus N$, we have $U^{1-s} \mu_{\varphi, \mathcal{U}}(x) < +\infty$. Therefore, by a standard conclusion, cf. [69, Chapter 8],

$$\begin{aligned} \Theta_{n-1+s}^* \mu_{\varphi, \mathcal{U}}(x) &= \limsup_{r \rightarrow 0} r^{-n+1-s} \mu_{\varphi, \mathcal{U}}(B(x, r)) \\ &\leq \lim_{r \rightarrow 0} \int_{B(x, r)} |x - y|^{-n+1-s} \mu_{\varphi, \mathcal{U}}(dy) = 0 \end{aligned} \quad (4.9)$$

for all such x . Since $\mu_{\varphi, \mathcal{U}} = D(\varphi|_{\mathcal{U}})$, where $\varphi|_{\mathcal{U}} \in BV(\mathcal{U})$ is the restriction of φ to \mathcal{U} , (4.9) trivially implies

$$\Theta_{n-1}^* \|D(\varphi|_{\mathcal{U}})\| (x) = \limsup_{r \rightarrow 0} r^{-n+1} \|D(\varphi|_{\mathcal{U}})\| (B(x, r)) = 0 \quad (4.10)$$

and, with suitable $r_0(x) > 0$,

$$\int_0^{r_0(x)} r^{-n} \|D(\varphi|_{\mathcal{U}})\| (B(x, r)) \leq \int_0^{r_0(x)} r^{s-1} dr < +\infty \quad (4.11)$$

for all $x \in \mathcal{U} \setminus N$. However, from (4.10) and (4.11) it then follows that $\varphi|_{\mathcal{U}}$ has an approximate limit at all $x \in \mathcal{U} \setminus N$, see [5, Remark 3.82]. In other words, $S_\varphi \cap \mathcal{U} \subset N$, and since $\mu_X^{[0, T]}(S_\varphi \setminus \mathcal{U}) \leq \mu_X^{[0, T]}(\mathbb{R}^n \setminus \mathcal{U}) = 0$, (4.6) follows. By monotone convergence (4.3) extends to all nonnegative Borel function $g : \mathbb{R}^n \rightarrow [0, +\infty]$, and for $g = \mathbf{1}_{S_\varphi}$ we obtain (4.7). Estimate (4.8) follows from (2.2): For any $x \in \mathbb{R}^n \setminus S_\varphi$ we have

$$|\varphi(x)| \leq \limsup_{r \rightarrow 0} \frac{1}{\mathcal{L}^n(B(x, r))} \int_{B(x, r)} |\varphi(y)| dy \leq \|\varphi\|_{L^\infty(\mathbb{R}^n)}.$$

□

Remark 4.5. Condition (4.5) may also be seen as a requirement on φ relative to the path X : The measure $\mu_X^{[0, T]}$ is concentrated on $\mathbb{R}^n \setminus N$ with N as in the above proof. Potential theory forbids $\mu_{\varphi, \mathcal{U}}$ to charge subsets of $\mathbb{R}^n \setminus N$ with finite \mathcal{H}^{n-1+s} -measure: Any such set A would have $(1-s)$ -Riesz capacity zero, [59, Chapter III, Theorem 3.14], and this would imply the existence of some point $\hat{x} \in A$ with $U^{1-s} \mu_X^{[0, T]}(\hat{x}) = +\infty$, [59, Chapter II, Theorem 2.4]. An alternative, geometric proof of this fact can be given in terms of upper densities: Suppose that A is a Borel subset of $\mathbb{R}^n \setminus N$ with $\mathcal{H}^{n-1+s}(A) < +\infty$. Then by (4.9) together with the density comparison theorem, [5, Theorem 2.56] (or [69, Chapter 6, Theorem 6.9]), we have

$$\mu_{\varphi, \mathcal{U}}(A) \leq \sup_{x \in A} \Theta_{n-1+s}^* \mu_{\varphi, K}(x) \mathcal{H}^{n-1+s}(A) = 0.$$

The following example gives a coarse sufficient condition for (4.5).

Example 4.6. If $s \in (0, 1)$,

$$I^{1-s} \mu_X^{[0, T]} < +\infty \quad (4.12)$$

and $I^{1-s} \mu_{\varphi, \mathcal{U}} < +\infty$, then (4.5) holds by the Cauchy-Schwarz inequality for the energy, see [59, Chapter I, Section 4].

4.2 Upper regularity and bounded potentials

The first maximum principle, [59, Chapter I, Theorem 1.10], states that, for any $s \in (0, 1)$ and any Borel measure ν , a bound

$$U^{1-s} \nu(x) \leq M, \quad (4.13)$$

valid with some $M > 0$ for ν -a.e. $x \in \mathbb{R}^n$, implies (4.13) for all $x \in \mathbb{R}^n$.

Proposition 4.7. *Let $s \in (0, 1)$. Suppose that there is a constant $M > 0$ such that*

$$U^{1-s} \mu_X^{[0,T]}(x) < M \quad (4.14)$$

for $\mu_X^{[0,T]}$ -a.e. $x \in X([0, T])$. Then (4.5) holds with any $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ and all relatively compact open \mathcal{U} , and $\mu_X^{[0,T]}$ cannot charge any set of A of finite \mathcal{H}^{n-1+s} -measure.

Proof. Since $\mu_X^{[0,T]}$ is supported in $X([0, T])$, the bound (4.14) holds for $\mu_X^{[0,T]}$ -a.e. $x \in \mathbb{R}^n$. By the maximum principle it then holds for all $x \in \mathbb{R}^n$ and as a consequence, (4.5) holds with any $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ and all relatively compact open \mathcal{U} . The second part of the statement follows as in Remark 4.5. \square

Note that (4.14) implies (4.12), but in contrast to Example 4.6 no assumption is made on the measures $\mu_{\varphi, \mathcal{U}}$.

We prepare a counterpart of Proposition 4.7 in terms of $\|D\varphi\|$. The gradient measure $D\varphi$ of a function $\varphi \in BV(\mathbb{R}^n)$ does not charge any \mathcal{H}^{n-1} -null set, [5, Lemma 3.76]. Let $D^a\varphi$ denote the part of $D\varphi$ which is absolutely continuous w.r.t. \mathcal{L}^n and $D^s\varphi$ the part that is singular w.r.t. \mathcal{L}^n . Clearly $D^a\varphi$ vanishes on \mathcal{L}^n -null sets. The Cantor part $D^c\varphi$ of $D\varphi$ is defined as $D^c\varphi = D^s\varphi|_{\mathbb{R}^n \setminus S_\varphi}$, [5, Definition 3.91], and in general it can, roughly speaking, charge subsets of any Hausdorff dimension between $n - 1$ and n . There is an $(n - 1)$ -dimensional Borel subset J_φ of the approximate discontinuity set S_φ , called the *set of approximate jump points*, [5, Definition 3.67], and the jump part of the gradient measure is defined to be $D^j\varphi = D^s\varphi|_{J_\varphi}$, [5, Definition 3.91]. The set $S_\varphi \setminus J_\varphi$ is of zero \mathcal{H}^{n-1} -measure, [5, Theorem 3.78]. The gradient measure of φ admits the unique decomposition

$$D\varphi = D^a\varphi + D^j\varphi + D^c\varphi. \quad (4.15)$$

A description of the different summands in terms of geometric scaling properties can be found in [5, Proposition 3.92].

Proposition 4.8. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ and $s \in (0, 1)$. Suppose that for any compact K there is a constant $M_K > 0$ such that*

$$U^{1-s} \mu_{\varphi, K}(x) < M_K \quad (4.16)$$

for $\mu_{\varphi, K}$ -a.e. $x \in K$. Then φ cannot have jumps and the Cantor part $D^c\varphi$ of $D\varphi$ cannot charge sets of finite \mathcal{H}^{n-1+s} -measure. The function φ has a unique continuous Borel version, Hölder of order s , and any Lebesgue representative coincides with this version on the Lebesgue set. Moreover, (4.4) holds for any path X , any relatively compact open $\mathcal{U} \supset X([0, T])$ and any $p \in [1, +\infty]$.

Condition (4.16) does not impose any restrictions on $D^a\varphi$.

Proof. Again by the maximum principle (4.16) holds for all $x \in \mathbb{R}^n$. Taking into account (4.15) we can conclude that if φ satisfies (4.16) for all compact K , then φ cannot have jumps: Since the set J_φ of approximate jump points has Hausdorff dimension $n - 1$ it

has zero \mathcal{H}^{n-1+s} -measure, and as before we can see such sets cannot be charged by $D\varphi$. The statement on Hölder continuity follows using large closed balls in place of K , a simple cut-off argument and Corollary C.3. \square

Remark 4.9. If $D\varphi$ is absolutely continuous, then the upper $n - 1 + s$ -regularity of φ for $s \in (0, 1)$ implies the local s -Hölder continuity of φ on \mathcal{C}^1 -smooth domains by a variant of Morrey's inequality in Sobolev-Morrey spaces by [57, Theorem 2.2].

Remark 4.10. Given a Borel measure ν on \mathbb{R}^n the *pointwise lower Hausdorff dimension of ν at $x \in \mathbb{R}^n$* is defined as

$$\underline{\dim}_H \nu(x) = \sup\{\gamma \geq 0 : \overline{D}^\gamma \nu(x) = 0\} = \inf\{\gamma \geq 0 : \overline{D}^\gamma \nu(x) = +\infty\}.$$

Its *lower Hausdorff dimension* is defined as $\underline{\dim}_H \nu = \text{ess inf}_{x \in \mathbb{R}^n} \underline{\dim}_H \nu(x)$, and it is well known that $\underline{\dim}_H \nu = \inf\{\dim_H A : A \subset \mathbb{R}^n \text{ Borel and } \nu(A) > 0\}$. See for instance [9, 70]. Conditions (4.14) and (4.16) imply that $\underline{\dim}_H \mu_X^{[0,T]} \geq n - 1 + s$ and, respectively, $\underline{\dim}_H \|D\varphi\| \geq n - 1 + s$.

The following is well known.

Proposition 4.11. *Let ν be a finite Borel measure on \mathbb{R}^n with compact support. If with some $M > 0$ we have $\int_{\mathbb{R}^n} |x - y|^{-s} \nu(dy) < M$ for all $x \in \text{supp } \nu$, then there is some $c > 0$ such that $\nu(B(x, r)) \leq cr^s$ for all $x \in \text{supp } \nu$ and $r > 0$. If there is some $r_0 > 0$ such that $\nu(B(x, r)) \leq cr^d$ for all $x \in \text{supp } \nu$ and $0 < r < r_0$, then for any $s < d$ there is some $M > 0$ such that $\int_{\mathbb{R}^n} |x - y|^{-s} \nu(dy) < M$ for all $x \in \text{supp } \nu$.*

We briefly sketch the folklore proof for the convenience of the reader.

Proof. The first statement is immediate from (C.1). To see the second claim note that since ν is finite, we can readjust c to obtain $\nu(B(x, r)) \leq cr^d$ for all $x \in \text{supp } \nu$ and $r > 0$. If $R > 0$ is such that $B(0, R/2)$ contains $\text{supp } \nu$, then a classical argument, [69, p. 109], shows that for any $x \in \text{supp } \nu$ we have

$$\int_{\mathbb{R}^n} |x - y|^{-s} \nu(dy) = s \int_0^{2R} r^{-s-1} \nu(B(x, r)) dr \leq cs \int_0^{2R} r^{d-s} dr.$$

\square

Corollary 4.12.

- (i) *If X is upper d -regular with $d > n - 1 + s$, then (4.14) holds.*
- (ii) *If φ is upper d -regular with $d > n - 1 + s$, then (4.16) holds for any compact K .*

For certain paths X the occupation measure $\mu_X^{[0,T]}$ of X is absolutely continuous. To the Radon-Nikodym derivative $\frac{d\mu_X^{[0,T]}}{d\mathcal{L}^n} = L_T^X$ one refers as *the local times* of X .

Proposition 4.13. *Suppose that $\mu_X^{[0,T]}$ of X is absolutely continuous with density $L_T^X \in L^p(\mathbb{R}^n)$ for some $p \in [1, \infty]$. Then X is upper $\frac{n}{q}$ -regular, where $\frac{1}{p} + \frac{1}{q} = 1$ with the agreement $\frac{1}{+\infty} = 0$. In particular, if the local times are bounded, $L_T^X \in L^\infty(\mathbb{R}^n)$, then X is upper n -regular.*

Proof. We have

$$\mu_X^{[0,T]}(B(x, r)) = \int_{B(x, r)} L_T^X(y) dy \leq \|L_T^X\|_{L^p(\mathbb{R}^n)} \mathcal{L}^n(B(x, r))^{\frac{1}{q}} \leq c \|L_T^X\|_{L^p(\mathbb{R}^n)} r^{\frac{n}{q}}.$$

□

Remark 4.14. Existence and regularity of local times are well-studied in the case of Gaussian processes, see e.g. [7, 10, 11, 39]. A key property is the so-called local non-determinism which guarantees that, roughly speaking, the increments of the process are not too degenerate.

In some situations it may be useful to localize the conditions on φ and X .

Corollary 4.15. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$, let X be a path and $s \in (0, 1)$. If $B \subset \mathbb{R}^n$ is a Borel set such that (4.14) holds for $\mu_X^{[0,T]}$ -a.e. $x \in B$ and (4.16) holds for $\mu_{\varphi, \mathcal{W}}$ -a.e. $x \in B^c$, then (4.5) holds.*

Of course this matters only if $B \cap X([0, T]) \neq \emptyset$ and $B^c \cap X([0, T]) \neq \emptyset$ and similarly for $\text{supp } D\varphi$. In the situation of Corollary 4.15 the Hausdorff dimension of $X([0, T]) \cap B$ must be greater or equal to $n - 1 + s$, but on B^c the path X may be arbitrary. On the other hand, φ may be arbitrary on B , in particular, it may have jumps there, but it cannot jump on B^c .

Proof. The claim follows from

$$\begin{aligned} I^{1-s}(\mu_{\varphi, \mathcal{W}}, \mu_X^{[0,T]}) &= I^{1-s}(\mu_{\varphi, \mathcal{W}}|_B, \mu_X^{[0,T]}|_B) + I^{1-s}(\mu_{\varphi, \mathcal{W}}|_B, \mu_X^{[0,T]}|_{B^c}) \\ &\quad + I^{1-s}(\mu_{\varphi, \mathcal{W}}|_{B^c}, \mu_X^{[0,T]}|_B) + I^{1-s}(\mu_{\varphi, \mathcal{W}}|_{B^c}, \mu_X^{[0,T]}|_{B^c}) \end{aligned}$$

and the maximum principle for the restrictions $\mu_{\varphi, \mathcal{W}}|_B$ etc. together with the preceding considerations. □

Example 4.16. Suppose that $\varphi_{\mathcal{E}} : \mathbb{R}^2 \rightarrow \mathbb{R}$ is as in Example 2.8 for $n = 2$. If X is a smooth unit speed curve outside $\mathbb{R} \setminus [0, 1] \times \mathbb{R}$ and a typical Brownian bridge path inside $[0, 1] \times \mathbb{R}$, then X is $(s, 1)$ -variable w.r.t. $\varphi_{\mathcal{E}}$. The same is true if $\varphi = \mathbf{1}_D$ and $D \subset [0, 1] \times \mathbb{R}$ is a set of finite perimeter.

We discuss details of Example 2.8 and 2.9.

Example 4.17. Let $\varphi_{\mathcal{E}} : \mathbb{R}^n \rightarrow \mathbb{R}$ be as in Example 2.8. If $s \in (0, d_{\mathcal{E}})$, then the fact that $\|D\varphi_{\mathcal{E}}\|(B(x, r)) \leq cr^{n-1+d_{\mathcal{E}}}$ together with Corollary 4.12 (ii) shows that any path X is

in $V(\varphi_\ell, s, \infty)$. From now on assume that $s \in (d_\ell, 1)$. For a constant path $X \equiv x$ with fixed $x \in \mathbb{R}^n$ we have

$$\mu_X^{[0,T]} = T\delta_x,$$

where δ_x is the point mass probability measure at x . If $x = (\frac{1}{2}, 0, \dots, 0)$, then x has distance $\frac{1}{6}$ to $\text{supp } \|D\varphi_\ell\|$, and hence $U^{1-s} \|D\varphi_\ell\| (x)$ is bounded and $X \in V(\varphi_\ell, s, \infty)$. On the other hand, if $x = (0, \dots, 0)$, then

$$I^{1-s}(\mu_{\varphi_\ell, \mathcal{U}}, \mu_X^{[0,T]}) = T \int_{\mathcal{U}} |y|^{-n+1-s} \|D\varphi_\ell\| (dy) = +\infty$$

for any open neighborhood \mathcal{U} of the origin, as follows from $\|D\varphi_\ell\| (B(y, r)) \geq cr^{n-1+d_\ell}$ and standard arguments ([69, p. 109]). Consequently X is not in $V(\varphi_\ell, s)$ in this case. For a non-constant smooth function X on \mathbb{R}

$$\mu_X^{[0,T]}((x-r, x+r)) \leq cr$$

implies that $X \in V(\varphi_\ell, s, \infty)$, in the case $n = 1$. For $n = 2$ we have $\|D\varphi_\ell\| (B(y, r)) \geq cr^{1+d_\ell}$, so that, again by standard arguments, $U^{1-s} \|D\varphi_\ell\| (y) = +\infty$ for any $y \in \{\frac{1}{3}\} \times [0, 1]$. For any $M > 0$ and any such y we can find an open neighborhood $U_y \subset \mathbb{R}^2$ of y on which we have $U^{1-s} \|D\varphi_\ell\| > M$, [59, Chapter I, Theorem 1.3], and consequently this must hold on all of $\{\frac{1}{3}\} \times [0, 1]$. If $T = 1$ and X is the unit speed motion along $X([0, 1]) = \{\frac{1}{3}\} \times [0, 1]$, then

$$\mu_X^{[0,1]} = \mathcal{H}^1|_{X([0,1])},$$

[29, Theorem 3.2.6], and $I^{1-s}(\|D\varphi_\ell\| |_{X([0,1])}, \mu_X^{[0,1]}) = +\infty$. Hence X is not in $V(\varphi_\ell, s)$.

Example 4.18. Let $D \subset \mathbb{R}^n$ be as in Example 2.9 and $s \in (0, 1)$. Suppose X is a unit speed smooth curve that hits ∂D finitely often. By the additivity of the occupation measures w.r.t. the time interval we may assume all hitting points are different. By the smoothness of ∂D and X it suffices to show that we have $I^{1-s}(\mu, \nu) < +\infty$, where $\mu = \mathcal{H}^1|_{\{(0, \dots, 0, x_n): |x_n| < 1\}}$ and $\nu = \mathcal{H}^{n-1}|_{\{(x_1, \dots, x_{n-1}, 0): |(x_1, \dots, x_{n-1})| < 1\}}$. For $n \geq 2$ and $\varepsilon > 0$ we have $\int_{-1}^1 (\varepsilon^2 + |t|)^{(-n+1-s)/2} dt \leq c\varepsilon^{-n+3-s}$. Therefore

$$\int_{B(0, 2^{-k}) \setminus B(0, 2^{-k-1})} \int_{\{(0, \dots, 0, y_n): |y_n| < 1\}} |x-y|^{-n+1-s} \mu(dy) \nu(dx) \leq c 2^{-k(2-s)},$$

which is summable over $k = 0, 1, 2, \dots$. For $n = 1$ the integral over $t \in [-1, 1]$ is even bounded. It follows that the mutual energy of μ and ν is finite, and hence for such D , X and s we have $X \in V(\mathbf{1}_D, s)$. If $n = 2$ and X spends \mathcal{L}^1 -positive time in ∂D , then we can find $0 \leq a < b \leq T$ such that $X([a, b]) \subset \partial D$ and $\mu_X^{[a,b]} = \mathcal{H}^1|_{[a,b]}$. Since $\|D\mathbf{1}_D\|$ equals a constant times $\mathcal{H}^{n-1}|_D$, it follows that

$$\begin{aligned} I^{1-s}(\|D\varphi\|_{X([0,T])}, \mu_X^{[0,T]}) \\ \geq c \int_{X([a,b])} \int_{X([a,b])} |x-y|^{-n+1-s} \mathcal{H}^1(dx) \mathcal{H}^{n-1}(dy) = +\infty, \end{aligned}$$

so that X is not in $V(\mathbf{1}_D, s)$.

4.3 Fourier transform and trading of regularity

Condition (4.12) requires that the Fourier transform

$$\hat{\mu}_X^{[0,T]}(\xi) = \int_{\mathbb{R}^n} e^{-i\langle \xi, x \rangle} \mu_X^{[0,T]}(dx), \quad \xi \in \mathbb{R}^n$$

of the occupation measure $\mu_X^{[0,T]}$ is square integrable w.r.t. $|\xi|^{s-1} d\xi$, note that

$$I^{1-s} \mu_X^{[0,T]} = c(s, n) \int_{\mathbb{R}^n} |\xi|^{s-1} |\hat{\mu}_X^{[0,T]}(\xi)|^2 d\xi \quad (4.17)$$

by [69, Lemma 12.12]. Increased activity of X allows larger $s \in (0, 1)$, see [50, Chapter 17, p. 250] for this classical discussion. Since $\hat{\mu}_X^{[0,T]}$ is bounded, integrability w.r.t. $|\xi|^{s-1} d\xi$ already implies square integrability, giving (4.12). (Alternatively, a Riemann-Lebesgue argument shows that integrability implies (4.14), which in turn implies (4.12).) The integral in (4.5) is a polarized version of (4.17),

$$I^{1-s}(\mu_{\varphi, \mathcal{U}}, \mu_X^{[0,T]}) = \operatorname{Re} \int_{\mathbb{R}^n} |\xi|^{s-1} \hat{\mu}_{\varphi, \mathcal{U}}(\xi) \overline{\hat{\mu}_X^{[0,T]}(\xi)} d\xi, \quad (4.18)$$

here $\hat{\mu}_{\varphi, \mathcal{U}}$ denotes the Fourier transform of the measure $\mu_{\varphi, \mathcal{U}}$.

Formulas (4.5) and (4.18) suggest a trading of regularity between the measures.

Proposition 4.19. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$, $X : [0, T] \rightarrow \mathbb{R}^n$ and $s \in (0, 1)$. For any $\gamma \in (0, 1 - s)$ we have*

$$\int_{\mathbb{R}^n} U^{1-s} \mu_{\varphi, \mathcal{U}}(x) \mu_X^{[0,T]}(dx) = \int_{\mathbb{R}^n} U^\gamma \mu_{\varphi, \mathcal{U}}(x) U^{1-s-\gamma} \mu_X^{[0,T]}(x) dx. \quad (4.19)$$

Proof. Proposition A.1 (ii), applied with $\nu_1 = \mu_{\varphi, \mathcal{U}}$, $\nu_2 = \mu_X^{[0,T]}$, $\gamma_1 = \gamma$ and $\gamma_2 = 1 - s - \gamma$, yields the result. \square

Since $1 - s$ marks the pivotal level of smoothness for the measures, formula (4.19) and upper regularity in the form considered above do not improve the statements in Corollary 4.15. However, Proposition 4.19 is used efficiently in Corollary 4.23 below.

4.4 Probabilistic examples

We connect to probabilistic examples. A condition of type (3.6) can ensure variability w.r.t. any BV -function.

Corollary 4.20. *Let $s \in (0, 1)$ and suppose that $Y = (Y_t)_{t \in [0, T]}$ is an \mathbb{R}^n -valued stochastic process with $Y_0 = 0$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. If*

$$\sup_{x \in \mathbb{R}^n} \mathbb{E} \int_0^T |Y_t - x|^{-n+1-s} dt < +\infty, \quad (4.20)$$

then we can find an event $\Omega_0 \in \mathcal{F}$ with $\mathbb{P}(\Omega_0) = 1$ such that for all $\omega \in \Omega_0$ the path $Y(\omega)$ is $(s, 1)$ -variable w.r.t. any $\varphi \in BV(\mathbb{R}^n)$.

Proof. Using Fubini and the above bound,

$$\mathbb{E} \int_{\mathbb{R}^n} \int_0^T |Y_t - x|^{-n+1-s} dt \|D\varphi\| (dx) \leq \|D\varphi\| (\mathbb{R}^n) \sup_{x \in \mathbb{R}^n} \mathbb{E} \int_0^T |Y_t - x|^{-n+1-s} dt < +\infty.$$

□

Example 4.21. Let $B^H = (B_t^H)_{t \geq 0}$ be an n -dimensional fractional Brownian motion with Hurst index $0 < H < 1$ over a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Let $s \in (0, 1)$. We claim that if

$$n - 1 + s < \frac{1}{H}, \quad (4.21)$$

then there is a constant $c(n, s, H) > 0$ such that for all $x \in \mathbb{R}^n$ we have

$$\mathbb{E} \int_0^T |B_t^H - x|^{-n+1-s} dt < c(n, s, H). \quad (4.22)$$

For Brownian motion - the special case $H = \frac{1}{2}$ - we have (4.21) in dimensions $n = 1$ and $n = 2$ for any $s \in (0, 1)$. For $n = 3$ a fractional Brownian motion with Hurst index $H < \frac{1}{2}$ satisfies (4.21) for $0 < s < \frac{1}{H} - 2$. A higher dimension n of the ambient space forces a higher irregularity of the path in order to have an occupation measure supported on a set of sufficiently small codimension. The arguments for (4.22) are rather standard, [28, 50]. The expectation equals

$$\begin{aligned} & \frac{1}{(2\pi)^{n/2}} \int_0^T t^{-nH} \int_{\mathbb{R}^n} \exp\left(-\frac{|y|^2}{2t^{2H}}\right) |x - y|^{-n+1-s} dy dt \\ &= \frac{1}{(2\pi)^{n/2}} \int_0^T t^{-nH} \int_{|\xi| > t^H} \exp\left(-\frac{|x - \xi|^2}{2t^{2H}}\right) |\xi|^{-n+1-s} d\xi dt \\ &+ \frac{1}{(2\pi)^{n/2}} \int_0^T t^{-nH} \int_{|\xi| \leq t^H} \exp\left(-\frac{|x - \xi|^2}{2t^{2H}}\right) |\xi|^{-n+1-s} d\xi dt. \end{aligned}$$

The first summand is bounded by

$$\begin{aligned} & \frac{1}{(2\pi)^{n/2}} \int_0^T t^{-2nH+(1-s)H} \int_{\mathbb{R}^n} \exp\left(-\frac{|x - \xi|^2}{2t^{2H}}\right) d\xi dt \\ &= \frac{1}{(2\pi)^{n/2}} \int_0^T t^{-2nH+(1-s)H} \int_{\mathbb{R}^n} \exp\left(-\frac{|z|^2}{2t^{2H}}\right) dz dt \\ &= \int_0^T t^{-(n-1+s)H} dt. \end{aligned}$$

The second summand does not exceed

$$\frac{1}{(2\pi)^{n/2}} \int_0^T t^{-nH} \int_{|\xi| \leq t^H} |\xi|^{-n+1-s} d\xi dt = \frac{1}{(2\pi)^{n/2}} \int_0^T t^{-(n-1+s)H} dt.$$

The above idea can be generalized to link $(s, 1)$ -variability into upper regularity of the underlying probability measures.

Corollary 4.22. *Let $Y = (Y_t)_{t \in [0, T]}$ be an \mathbb{R}^n -valued stochastic process with $Y_0 = 0$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. For any $t \in [0, T]$, let $\nu_t := P_{Y_t}$ denote the law of Y_t . Suppose that $s \in (0, 1)$ and that there exists some $g \in L^1(0, T)$ such that for \mathcal{L}^1 -a.e. $t \in (0, T)$ we have*

$$\nu_t(B(x, r)) \leq g(t)r^d, \quad x \in \mathbb{R}^n, \quad r > 0, \quad (4.23)$$

with some $d > n - 1 + s$. Then (4.20) holds.

Proof. By Fubini

$$\mathbb{E} \int_0^T |Y_t - x|^{-n+1-s} dt = \int_0^T \int_{\mathbb{R}^n} |y - x|^{-n+1-s} \nu_t(dy) dt,$$

and as in Proposition 4.11 it follows that for \mathcal{L}^1 -a.e. $t \in (0, T)$ the inner integral is bounded by

$$(n - 1 + s) \int_0^\infty r^{-n-s} \nu_t(B(x, r)) dr \leq 1 + (n - 1 + s) g(t) \int_0^1 r^{-n-s+d} dr < +\infty.$$

□

If a coefficient φ is fixed and $\|D\varphi\|$ satisfies a certain moment condition, the requirements on Y to be $(s, 1)$ -variable can be weakened substantially.

Corollary 4.23. *Let $\varphi \in BV(\mathbb{R}^n)$ and $s \in (0, 1)$. Suppose that $Y = (Y_t)_{t \in [0, T]}$ is an \mathbb{R}^n -valued stochastic process with $Y_0 = 0$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Suppose that there are $\varepsilon \in (0, 1 - s)$, $c > 0$, $x_1, \dots, x_N \in \mathbb{R}^n$ and $\delta_1, \dots, \delta_N \in (0, n - 1 + s - \varepsilon)$ such that we have*

$$\mathbb{E} \int_0^T |Y_t - x|^{-n+\varepsilon} dt < c \sum_{k=1}^N |x - x_k|^{-n+\delta_k}, \quad x \in \mathbb{R}^n,$$

and

$$\sum_{k=1}^N \int_{\mathbb{R}^n} |x - x_k|^{-n+1-s-\varepsilon+\delta_k} \|D\varphi\| (dx) < +\infty. \quad (4.24)$$

Then we can find an event $\Omega_0 \in \mathcal{F}$ with $\mathbb{P}(\Omega_0) = 1$ such that for all $\omega \in \Omega_0$ the path $Y(\omega)$ is $(s, 1)$ -variable w.r.t. φ .

Proof. Under (3.13) and (4.24) we have, by (4.19), Fubini and Proposition A.1 (i),

$$\begin{aligned}
\mathbb{E}I^{1-s}(\|D\varphi\|, \mu_Y^{[0,T]}) &= \int_{\mathbb{R}^n} \mathbb{E}U^\varepsilon \mu_Y^{[0,T]}(x) U^{1-s-\varepsilon} \|D\varphi\|(x) dx \\
&\leq c \sum_{k=1}^N \int_{\mathbb{R}^n} |x - x_k|^{-n+\delta_k} U^{1-s-\varepsilon} \|D\varphi\|(x) dx \\
&= c \sum_{k=1}^N \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |x - x_k|^{-n+\delta_k} |x - y|^{-n+1-s-\varepsilon} \|D\varphi\|(dy) \\
&= c \sum_{k=1}^N \int_{\mathbb{R}^n} |y - x_k|^{-n+1-s-\varepsilon+\delta_k} \|D\varphi\|(dy).
\end{aligned}$$

□

Example 4.24. For $n \geq 2$, the fractional Brownian motion with $H > \frac{1}{n}$ satisfies (3.13) with $s = \frac{1}{H}$, $N = 1$, $0 < \varepsilon < (1 - \frac{1}{H})$ and $\delta_1 = \frac{1}{H} + \varepsilon$: We have

$$\mathbb{E} \int_0^T |Y_t - x|^{-n+\varepsilon} dt = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}^n} |y - x|^{-n+\varepsilon} \int_0^T t^{-nH} \exp\left(-\frac{|y|^2}{2t^{2H}}\right) dt dy.$$

Substituting $u = \frac{1}{2}|y|^2 t^{-2H}$ we see that for any $y \in \mathbb{R}^n \setminus \{0\}$ the inner integral equals

$$\frac{2^{n/2}}{2^{1+\frac{1}{2H}H}} |y|^{\frac{1}{H}-n} \int_{\frac{1}{2}|y|^2 T^{-2H}}^{u^{\frac{n}{2}-\frac{1}{2H}-1}} \exp(-u) du \leq c(n, H) |y|^{\frac{1}{H}-n}.$$

Using the convolution identity for Riesz kernels,

$$\mathbb{E} \int_0^T |Y_t - x|^{-n+\varepsilon} dt \leq c(n, H) \int_{\mathbb{R}^n} |y - x|^{-n+\varepsilon} |y|^{\frac{1}{H}-n} dy = c(n, H) |x|^{\frac{1}{H}+\varepsilon-n}.$$

The following generalization of this example is immediate.

Corollary 4.25. *Let $Y = (Y_t)_{t \in [0, T]}$ be an \mathbb{R}^n -valued stochastic process with $Y_0 = 0$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. For any $t \in [0, T]$, let $\nu_t := P_{Y_t}$ denote the law of Y_t . Suppose that $s \in (0, 1)$, ε, c, x_k and δ_k are as in Corollary 4.23, and that there exists some $g \in L^1(0, T)$ such that for \mathcal{L}^1 -a.e. $t \in (0, T)$ we have*

$$\int_{\mathbb{R}^n} |y - x|^{-n+\varepsilon} \nu_t(dy) \leq g(t) \sum_{k=1}^N |x - x_k|^{-n+\delta_k}, \quad x \in \mathbb{R}^n.$$

Then (3.13) holds.

Remark 4.26. Corollary 4.23 can also be applied to Gaussian bridges: Suppose that $n \geq 2$, let B^H be a fractional Brownian motion with $H > \frac{1}{n}$, and let Y be the Gaussian

bridge obtained by conditioning to have $B_T^H = \mathbf{1}$, where $\mathbf{1} = (1, \dots, 1) \in \mathbb{R}^n$. Then using [78, Theorem 3.1] it can be shown that

$$\mathbb{E} \int_0^T |Y_t - x|^{-n+\varepsilon} dt < c \left[|x|^{\frac{1}{H}+\varepsilon-n} + |x-1|^{\frac{1}{H}+\varepsilon-n} \right], \quad x \in \mathbb{R}^n.$$

Similar arguments can be applied also to more general bridges conditioned, e.g., on several time marginals, cf. [78, Remark 3.6(ii)].

4.5 A brief discussion of variability versus irregularity

In [16] the authors studied ODEs of type $dx_t = b(t, x_t)dt + dw_t$, where b is an irregular vector field and w is a continuous fast moving perturbation. Although our goal and methods are different from theirs, our point of view upon irregular paths and occupation measures follow a similar spirit. In cases where $b = b(x)$ is a bounded continuous function and w is ‘active enough’ they can prove existence and uniqueness of a continuous solution x , [16, Theorem 1.9]. If b is a distribution only, $b(x)$ must be defined appropriately, and in [16] this was done for paths x that differ from the sufficiently fast moving perturbation w only by a Hölder signal, [16, Definition 1.10 and Theorem 1.11]. The needed activity of w is encoded in the boundedness and (temporal) Hölder continuity in a certain (spatial) Hölder norm of the image

$$T_t^w b(x) = \int_0^t b(x + w_u) du$$

of b under an operator T^w taking the average of b along w . The more active w , the stronger is the averaging effect of this operator, and under suitable quantitative assumptions the authors then obtain existence, uniqueness, and flow properties for solutions x even if b is only a distribution, [16, Theorems 1.12, 1.13, 1.14]. To quantify how ‘fast moving’ a path $w \in \mathcal{C}([0, T], \mathbb{R}^n)$ must be they look at the Fourier transform $\hat{\mu}_w^{[0,t]}(\xi)$ of the occupation measure $\mu_w^{[0,t]}$ of w up to time t and, given $\varrho > 0$ and $\gamma > 0$, say that w is (ϱ, γ) -irregular if

$$\sup_{a \in \mathbb{R}^n} \sup_{0 \leq u < t \leq T} (1 + |a|)^\varrho \frac{|\hat{\mu}_w^{[0,u]}(a) - \hat{\mu}_w^{[0,t]}(a)|}{|u - t|^\gamma} < +\infty, \quad (4.25)$$

[16, Definition 1.3]. If a path w satisfies (4.25), then the averaging operator T^w is bounded between certain Fourier-Lebesgue (or Hölder) spaces, [16, Corollary 1.5, Theorem 1.6, Theorem 1.7]. Increased activity of w implies higher regularity (diffusivity) of its occupation measure, and hence better decay of its Fourier transform (larger ϱ), which encodes a stronger regularization effect of T^w . Condition (4.25) is a condition for single paths, and it is later connected to the individual coefficient b via the mapping properties of T^w . In contrast, (2.1) is a condition on X relative to a given BV-function φ . In (4.17) and (4.18) the interval $[0, T]$ is fixed, and the decay of the Fourier transforms at infinity is quantified in terms of integrability properties.

4.6 Compositions of BV_{loc} -functions and Hölder paths

Corollary 4.4 allows a multidimensional version of [18, Proposition 4.6], which ensures that compositions $\varphi \circ X$ are elements of $W^{\beta,p}(0,T)$. We first provide a bound for the Gagliardo seminorm part in the norm $\|\varphi \circ X\|_{W^{\beta,p}(0,T)}$.

Proposition 4.27. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$. Let $X : [0, T] \rightarrow \mathbb{R}^n$ be a path which is Hölder continuous of order $\alpha \in (0, 1]$. Suppose that $s \in (0, 1)$, $p \in [1, +\infty)$ and $X \in V(\varphi, s, p)$. Then for any $\beta \in (0, \alpha s)$ there is a constant $c > 0$, depending only on α, β, n, p and s , such that*

$$[\varphi \circ X]_{\beta,p}^p \leq c [X]_{\alpha,\infty}^{sp} \int_0^T \left[\int_{X([0,T])} \frac{\|D\varphi\|(dy)}{|X_t - y|^{n-1+s}} \right]^p dt. \quad (4.26)$$

If φ, X, α and s are as before but $X \in V(\varphi, s, \infty)$, then for any $\beta \in (0, \alpha s)$ there is a constant $c > 0$, depending only on α, β, n and s , such that

$$\text{ess sup}_{t \in [0, T]} \int_0^t \frac{|\varphi(X_t) - \varphi(X_\tau)|}{(t - \tau)^{1+\theta}} d\tau \leq c [X]_{\alpha,\infty}^s \text{ess sup}_{t \in [0, T]} \int_{X([0, T])} \frac{\|D\varphi\|(dy)}{|X_t - y|^{n-1+s}}. \quad (4.27)$$

Proof. By (4.7) we have

$$\int_0^T g(X_\tau) d\tau = \int_0^T \mathbf{1}_{\mathbb{R}^n \setminus S_\varphi}(X_\tau) g(X_\tau) d\tau \quad (4.28)$$

for any nonnegative Borel function $g : \mathbb{R}^n \rightarrow [0, +\infty]$. For any fixed $t \in [0, T]$ we have

$$\int_0^t \frac{|\varphi(X_t) - \varphi(X_\tau)|}{(t - \tau)^{1+\beta p}} d\tau = \int_0^t \frac{|\varphi(X_t) - \varphi(X_\tau)|}{(t - \tau)^{1+\beta p}} \mathbf{1}_{\mathbb{R}^n \setminus S_\varphi}(X_\tau) d\tau$$

by (4.28), and viewing this as a nonnegative function of t , also

$$\begin{aligned} & \int_0^T \int_0^t \frac{|\varphi(X_t) - \varphi(X_\tau)|^p}{(t - \tau)^{1+\beta p}} d\tau dt \\ &= \int_0^T \int_0^t \frac{|\varphi(X_t) - \varphi(X_\tau)|^p}{(t - \tau)^{1+\beta p}} \mathbf{1}_{\mathbb{R}^n \setminus S_\varphi}(X_\tau) \mathbf{1}_{\mathbb{R}^n \setminus S_\varphi}(X_t) d\tau dt. \end{aligned} \quad (4.29)$$

Let $\mu := \|D\varphi\||_{X([0, T])}$ denote the restriction of $\|D\varphi\|$ to $X([0, T])$. By Proposition C.1 and the α -Hölder continuity of X the right hand side of (4.29) is seen to be bounded by

$$\begin{aligned} & c(n, s)^p \int_0^T \int_0^t \frac{|X_t - X_\tau|^{sp}}{(t - \tau)^{1+\beta p}} [\mathcal{M}_{1-s, 4|X_t - X_\tau|} \mu(X_t) + \mathcal{M}_{1-s, 4|X_t - X_\tau|} \mu(X_\tau)]^p d\tau dt \\ & \leq c(n, s)^p [X]_{\alpha,\infty}^{sp} \int_0^T \int_0^t (t - \tau)^{\alpha sp - \beta p - 1} \times \\ & \quad \times [\mathcal{M}_{1-s, 4|X_t - X_\tau|} \mu(X_t) + \mathcal{M}_{1-s, 4|X_t - X_\tau|} \mu(X_\tau)]^p d\tau dt. \end{aligned}$$

The trivial estimate (C.1) implies

$$\mathcal{M}_{1-s,4|X_t-X_s|}\mu(X_t) \leq \int_{\mathbb{R}^n} \frac{\mu(dy)}{|X_t - y|^{n-1+s}} \quad (4.30)$$

for any $t \in [0, T]$, and therefore

$$\begin{aligned} & \int_0^T \int_0^t (t - \tau)^{\alpha sp - \beta p - 1} [\mathcal{M}_{1-s,4|X_t-X_\tau|}\mu(X_t)]^p d\tau dt \\ & \leq \int_0^T \int_0^t (t - \tau)^{\alpha sp - \beta p - 1} d\tau \left[\int_{\mathbb{R}^n} \frac{\mu(dy)}{|X_t - y|^{n-1+s}} \right]^p dt \\ & \leq \frac{T^{(\alpha s - \beta)p}}{(\alpha s - \beta)p} \int_0^T \left[\int_{X([0,T])} \frac{\|Df\|(dy)}{|X_t - y|^{n-1+s}} \right]^p dt. \end{aligned}$$

Using Fubini we obtain the same upper bound for the summand with $\mu(X_\tau)$ in place of $\mu(X_t)$. Combining the estimates and using the symmetry of the integrand in the Gagliardo seminorm, we arrive at (4.26). The estimate (4.27) follows similarly. \square

Remark 4.28. In [17, Proposition 5.0.3 and Remark 5.0.3] it has been shown that for discontinuous $\varphi \in BV(\mathbb{R})$ one cannot expect $\varphi(X)$ to have finite p -variation for any $p \geq 1$, if X visits a point of discontinuity of φ infinitely many times. In particular, $\varphi(X)$ cannot be Hölder continuous of any order in this case. This motivates to use Sobolev norms and generalized Stieltjes type integrals rather than p -variation and Young integrals.

If $X \in V(\varphi, s, p)$, a pinning argument shows that φ must be in $L^p(X)$. This entails that, for appropriate β and p , the composition $\varphi \circ X$ is in $W^{\beta,p}(0, T)$ and, for large enough p , even Hölder continuous.

Lemma 4.29. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$. Let $X : [0, T] \rightarrow \mathbb{R}^n$ be a path which is Hölder continuous of order $\alpha \in (0, 1)$. Suppose that $s \in (0, 1)$, $p \in [1, +\infty]$ and $X \in V(\varphi, s, p)$. Then $\varphi \in L^p(X)$ and $\varphi \circ X \in W^{\beta,p}(0, T)$ for any $\beta \in (0, \alpha s)$. Moreover, if $\alpha s > \frac{1}{p}$, then $\varphi \circ X$ is Hölder continuous of any order smaller than $\alpha s - \frac{1}{p}$. (We again use the agreement that $\frac{1}{+\infty} := 0$.)*

Proof. Suppose first that $p \in [1, +\infty)$. Choose $t_0 \in [0, T]$ such that $X_{t_0} \in \mathbb{R}^n \setminus S_\varphi$. By Corollary 4.4 such t_0 clearly exists. Then we have

$$|\varphi(X_t)|^p \leq 2^{p-1} [|\varphi(X_t) - \varphi(X_{t_0})|^p + |\varphi(X_{t_0})|^p].$$

for a.e. $t \in [0, T]$. Using (4.30) we obtain

$$\begin{aligned} & |\varphi(X_t) - \varphi(X_{t_0})|^p \\ & \leq |X_t - X_{t_0}|^{sp} [\mathcal{M}_{1-s,4|X_t-X_{t_0}|}\mu(X_t) + \mathcal{M}_{1-s,4|X_t-X_{t_0}|}\mu(X_{t_0})]^p \\ & \leq 2^{p-1} [X]_{\alpha, \infty}^{sp} T^{\alpha sp} \left(\left[\int_{X([0,T])} \frac{\|D\varphi\|(dy)}{|X_t - y|^{n-1+s}} \right]^p + \left[\int_{X([0,T])} \frac{\|D\varphi\|(dy)}{|X_{t_0} - y|^{n-1+s}} \right]^p \right), \end{aligned}$$

and integration yields

$$\begin{aligned}
& \int_0^T (\varphi \circ X)(t)^p dt \\
& \leq 2^{p-1} T |(\varphi \circ X)(t_0)|^p + 4^{p-1} [X]_{\alpha, \infty}^{sp} T^{\alpha sp + 1} \left[\int_{X([0, T])} \frac{\|D\varphi\|(dy)}{|X_{t_0} - y|^{n-1+s}} \right]^p \\
& \quad + 4^{p-1} [X]_{\alpha, \infty}^{sp} T^{\alpha sp} \int_0^T \left[\int_{X([0, T])} \frac{\|D\varphi\|(dy)}{|X_t - y|^{n-1+s}} \right]^p dt. \quad (4.31)
\end{aligned}$$

In order to show that $\varphi \circ X \in L^p([0, T])$ it now suffices to prove that there is some $t_0 \in [0, T]$ such that $X_{t_0} \in \mathbb{R}^n \setminus S_\varphi$,

$$|\varphi(X_{t_0})|^p < +\infty, \quad (4.32)$$

and

$$\int_{X([0, T])} \frac{\|D\varphi\|(dy)}{|X_{t_0} - y|^{n-1+s}} < +\infty. \quad (4.33)$$

Let $N \subset [0, T]$ be the set of all t such that $X_t \in S_\varphi$. As seen before, N is a Lebesgue null set, and by Definition 2.5 and (2.2) we have $\tilde{\varphi}(X_t) = \lambda_\varphi(X_t) < \infty$ for all $t \in [0, T] \setminus N$ since $X_t \in \mathbb{R}^n \setminus S_\varphi$ for such t . By (s, p) -variability, (4.33) must hold for all $t \in [0, T] \setminus N'$, where N' is a Lebesgue null set. Thus both (4.32) and (4.33) must hold for any $t_0 \in [0, T] \setminus (N \cup N')$, i.e. for a.e. $t_0 \in [0, T]$. In the case that $p = +\infty$ the (s, ∞) -variability of X guarantees (4.33) for all $t_0 \in [0, T]$, so that the desired result follows by obvious modifications of the above arguments. The membership in $W^{\beta, p}(0, T)$ is seen using Proposition 4.27 and the statement on continuity using Sobolev embedding, [23, Theorem 8.2]. \square

Remark 4.30. By Remark 4.28 and Lemma 4.29 one cannot expect X to be (s, p) -variable with respect to φ for $p > \frac{1}{\alpha s}$ if X visits discontinuity points of φ infinitely often.

In the rest of this subsection we derive an estimate for the weighted L^p -term in the norm $\|\varphi \circ X\|_{W_0^{\beta, p}(0, T)}$. Only the special case $p = 1$ will be used later on.

Proposition 4.31. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$. Let $X : [0, T] \rightarrow \mathbb{R}^n$ be a path which is Hölder continuous of order $\alpha \in (0, 1]$ and $X \in V(\varphi, s, p)$ for some $s \in (0, 1)$ and $p \in [1, +\infty)$. Then for any $\beta \in (0, \alpha s \wedge \frac{1}{p})$ there is a constant $c > 0$, depending only on α, β, n, p and s , such that*

$$\begin{aligned}
& \int_0^T \frac{|(\varphi \circ X)(t)|^p}{t^{\beta p}} dt \\
& \leq c \left([X]_{\alpha, \infty}^{sp} \int_0^T \left[\int_{X([0, T])} \frac{\|D\varphi\|(dy)}{|X_t - y|^{n-1+s}} \right]^p dt + \int_0^T |(\varphi \circ X)(t)|^p dt \right). \quad (4.34)
\end{aligned}$$

To prove Proposition 4.31 we make use of the following fact.

Lemma 4.32. *Let $\beta > 0$ and $q > 0$ such that $\beta q < 1$. Then for all continuous functions u on $[0, T]$ we have*

$$\int_0^T \frac{|u(t)|^q}{t^{\beta q}} dt \leq c \left(\int_0^T \int_0^T \frac{|u(t) - u(\tau)|^q}{|t - \tau|^{1+\beta q}} dt d\tau + \int_0^T |u(t)|^q dt \right), \quad (4.35)$$

where $c > 0$ is a constant depending only on β and q .

Lemma 4.32 is a slight adaption of the following result in [19] and [25]: Suppose $\mathcal{D} \subset \mathbb{R}^d$ is a bounded Lipschitz domain and let $\delta_{\mathcal{D}}(x) = \inf\{|y - x| : y \in \mathcal{D}^c\}$ denote the distance of x to its complement \mathcal{D}^c . Then, by [25, Equation (17)], we have, for any $q > 0$ and $\alpha \in (0, 1)$, that

$$\int_{\mathcal{D}} \frac{|u(x)|^q}{[\delta_{\mathcal{D}}(x)]^\alpha} dx \leq c \left(\int_{\mathcal{D}} \int_{\mathcal{D}} \frac{|u(x) - u(y)|^q}{|x - y|^{d+\alpha}} dx dy + \int_{\mathcal{D}} |u(x)|^q dx \right) \quad (4.36)$$

for all $u \in \mathcal{C}_c(\mathcal{D})$.

Proof. An application of (4.36) to the case $d = 1$, $\alpha = \beta q$ and $\mathcal{D} = (0, T)$ yields

$$\int_0^T \frac{|u(t)|^q}{t^\beta} dt \leq c \left(\int_0^T \int_0^T \frac{|u(t) - u(\tau)|^q}{|t - \tau|^{1+\beta}} dt d\tau + \int_0^T |u(t)|^q dt \right)$$

for all $u \in \mathcal{C}_c(0, T)$. Note also that $t^{-\beta} \leq [\delta_{\mathcal{D}}(t)]^{-\beta}$. Now suppose that $u \in \mathcal{C}([0, T])$. For each n let u_n be the continuous function on $[0, T]$ such that $u_n = u$ on $[\frac{1}{n}, T - \frac{1}{n}]$, u_n is linear on $[\frac{1}{2n}, \frac{1}{n}] \cup [T - \frac{1}{n}, T - \frac{1}{2n}]$, and $u_n \equiv 0$ on $[0, \frac{1}{2n}] \cup [T - \frac{1}{2n}, T]$. Then obviously $u_n \in \mathcal{C}_c(0, T)$ and we have

$$\sup_{t \in [0, T]} |u_n(t)| \leq S, \quad (4.37)$$

where $S := \sup_{t \in [0, T]} |u(t)|$. Since bounded convergence implies

$$\lim_n \int_0^T \frac{|u_n(t)|^q}{t^{\beta q}} dt = \int_0^T \frac{|u(t)|^q}{t^{\beta q}} dt \quad \text{and} \quad \lim_n \int_0^T |u_n(t)|^q dt = \int_0^T |u(t)|^q dt,$$

it suffices to show that

$$\limsup_n \int_0^T \int_0^T \frac{|u_n(t) - u_n(\tau)|^q}{|t - \tau|^{1+\beta q}} dt d\tau \leq c \int_0^T \int_0^T \frac{|u(t) - u(\tau)|^q}{|t - \tau|^{1+\beta q}} dt d\tau.$$

On $[0, \frac{1}{n}]$ the function u_n obeys the Lipschitz bound $|u_n(t) - u_n(\tau)| \leq nS|t - \tau|$, which implies

$$\int_0^{n^{-1}} \int_0^{n^{-1}} \frac{|u_n(t) - u_n(\tau)|^q}{|t - \tau|^{1+\beta q}} dt d\tau \leq S^q n^q \int_0^{n^{-1}} \int_0^{n^{-1}} |t - \tau|^{q-1-\beta q} d\tau dt \leq c S^q n^{\beta q - 1}.$$

This goes to zero as $n \rightarrow \infty$. Similarly, using (4.37), and writing c for positive constants depending only on q and β and possibly changing from line to line,

$$\begin{aligned}
& \int_0^{n^{-1}} \int_{n^{-1}}^T \frac{|u_n(t) - u_n(\tau)|^q}{|t - \tau|^{1+\beta q}} dt d\tau \\
& \leq c \int_0^{n^{-1}} \int_{n^{-1}}^T \frac{|u_n(t) - u(t) + u(\tau) - u_n(\tau)|^q}{|t - \tau|^{1+\beta}} dt d\tau + c \int_0^{n^{-1}} \int_{n^{-1}}^T \frac{|u(t) - u(\tau)|^q}{|t - \tau|^{1+\beta q}} dt d\tau \\
& \leq cS^q \int_0^{n^{-1}} \int_{n^{-1}}^T |t - \tau|^{-\beta q - 1} dt d\tau + c \int_0^T \int_0^T \frac{|u(t) - u(\tau)|^q}{|t - \tau|^{1+\beta q}} dt d\tau \quad (4.38) \\
& \leq cS^q n^{\beta q - 1} + c \int_0^T \int_0^T \frac{|u(t) - u(\tau)|^q}{|t - \tau|^{1+\beta q}} dt d\tau.
\end{aligned}$$

Treating the regions involving intervals $[T - \frac{1}{n}, T]$ similarly concludes the proof. \square

We record another useful fact. See [4, 63] for more general chain rules.

Lemma 4.33. *If $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ and $\Phi : \mathbb{R} \rightarrow \mathbb{R}$ is a \mathcal{C}^1 -function with bounded derivative and $\Phi(0) = 0$, then $\Phi \circ \varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ and*

$$\|D(\Phi \circ \varphi)\| \leq \|\Phi'\|_{\text{sup}} \|D\varphi\|. \quad (4.39)$$

If $X \in V(\varphi, s, p)$, then also $X \in V(\Phi \circ \varphi, s, p)$.

Proof. By [5, Theorem 3.96 and its proof] we have $\Phi \circ \varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ and (4.39). This implies that for any compact $K \subset \mathbb{R}^n$ we have

$$U^{1-s}(\|D(\Phi \circ \varphi)\| |K)(x) \leq \|\Phi'\|_{\text{sup}} U^{1-s}(\|D\varphi\| |K)(x), \quad x \in \mathbb{R}^n.$$

Now the second statement follows from (4.4). \square

We prove Proposition 4.31.

Proof of 4.31. For any $N \geq 2$ let $\Phi_N \in \mathcal{C}^1(\mathbb{R})$ be an increasing function with $\|\Phi'\|_{\text{sup}} \leq 1$ and such that $\Phi_N(y) = -N$ for $y \leq -N$, $\Phi_N(y) = N$ for $y \geq N$ and $\Phi_N(y) = y$ for $-(N-1) < y < N-1$. Then, by Lemma 4.33 and the hypotheses of Proposition 4.31, we have $\Phi_N(\varphi) \in BV_{\text{loc}}(\mathbb{R}^n)$ and X is (s, p) -variable w.r.t. each $\Phi_N(\varphi)$. Suppose that (4.34) holds for all $\Phi_N(\varphi)$ in place of φ . Then, by (4.39),

$$\begin{aligned}
& \int_0^T \frac{|(\Phi_N(\varphi) \circ X)(t)|^p}{t^{\beta p}} dt \\
& \leq c[X]_{\alpha, \infty}^{sp} \left(\int_0^T \left[\int_{X([0, T])} \frac{\|D(\Phi_N(\varphi))\|(dy)}{|X_t - y|^{1-s}} \right]^p dt + \int_0^T |(\Phi_N(\varphi) \circ X)(t)|^p dt \right). \\
& \leq c[X]_{\alpha, \infty}^{sp} \left(\int_0^T \left[\int_{X([0, T])} \frac{\|D\varphi\|(dy)}{|X_t - y|^{1-s}} \right]^p dt + \int_0^T |(\varphi \circ X)(t)|^p dt \right),
\end{aligned}$$

and (4.34) for φ follows using Fatou's lemma. Consequently it suffices to prove (4.34) under the assumption that $\sup_{x \in \mathbb{R}^n} |\varphi(x)| \leq N$, and we do so in the sequel.

Let $(\eta_\varepsilon)_{\varepsilon>0}$ be a (radially symmetric) flat mollifier as in Appendix B. For each $\varepsilon > 0$ the composition $\varphi_\varepsilon \circ X$ of the mollified function $\varphi_\varepsilon := \varphi * \eta_\varepsilon$ with the path X is continuous on $[0, T]$, so that by (4.35),

$$\int_0^T \frac{|\varphi_\varepsilon(X_t)|^p}{t^{\beta p}} dt \leq c \left(\int_0^T \int_0^T \frac{|\varphi_\varepsilon(X_t) - \varphi_\varepsilon(X_\tau)|^p}{|t - \tau|^{1+\beta p}} dt d\tau + \int_0^T |\varphi_\varepsilon(X_t)|^p dt \right) \quad (4.40)$$

with a constant $c > 0$ depending only on β and p . By (B.2) we have

$$\lim_{\varepsilon \rightarrow 0} \varphi_\varepsilon(y) = \varphi(y), \quad y \in \mathbb{R}^n \setminus S_\varphi, \quad (4.41)$$

where S_φ is the approximate discontinuity set of φ . Since

$$\sup_{x \in \mathbb{R}^n} |\varphi_\varepsilon(x)| \leq \sup_{x \in \mathbb{R}^n} \int_{\mathbb{R}^n} |\varphi(y)| \eta_\varepsilon(x - y) dy \leq N$$

and $\mu_X^{[0, T]}$ is finite, we can use (4.41) together with Corollary 4.4 and bounded convergence to conclude that

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_0^T |\varphi_\varepsilon(X_t)|^p dt &= \lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^n} |\varphi_\varepsilon(x)|^p \mu_X^{[0, T]}(dx) \\ &= \int_{\mathbb{R}^n} |\varphi(x)|^p \mu_X^{[0, T]}(dx) = \int_0^T |\varphi(X_t)|^p dt. \end{aligned} \quad (4.42)$$

As before, let $\mu_{\varphi, \mathcal{U}}$ denote the restriction of $\|D\varphi\|$ to a relatively compact open set \mathcal{U} containing $X([0, T])$. Let $\mu_\varepsilon := \|D\varphi_\varepsilon\|_{|X([0, T])}$. For sufficiently small ε the open ε -parallel set $(X([0, T]))_\varepsilon$ of $X([0, T])$ is contained in \mathcal{U} , so that by Corollary B.3 we have

$$U^{1-s}(\mu_\varepsilon)(x) \leq c(n, s) U^{1-s} \mu_{\varphi, \mathcal{U}}(x), \quad x \in \mathbb{R}^n, \quad (4.43)$$

where $c(n, s) > 0$ is a constant depending only on n and s . Let next $0 < \tau < t \leq T$ be distinct and such that $X_t, X_\tau \in \mathbb{R}^n \setminus S_\varphi$. Then Proposition C.1 and bound (4.30) imply

$$|\varphi_\varepsilon(X_t) - \varphi_\varepsilon(X_\tau)| \leq c(n, s) |X_t - X_\tau|^s [U^{1-s} \mu_\varepsilon(X_t) + U^{1-s} \mu_\varepsilon(X_\tau)].$$

Combining with (4.43) gives

$$|\varphi_\varepsilon(X_t) - \varphi_\varepsilon(X_\tau)| \leq c(n, s) |X_t - X_\tau|^s [U^{1-s} \mu_{\varphi, \mathcal{U}}(X_t) + U^{1-s} \mu_{\varphi, \mathcal{U}}(X_\tau)]. \quad (4.44)$$

Therefore we have

$$\begin{aligned} \frac{|\varphi_\varepsilon(X_t) - \varphi_\varepsilon(X_\tau)|^p}{(t - \tau)^{1+\beta p}} \\ \leq c(n, s)^p [X]_{\alpha, \infty}^{sp} (t - \tau)^{\alpha sp - \beta p - 1} [U^{1-s} \mu_{\varphi, \mathcal{U}}(X_t) + U^{1-s} \mu_{\varphi, \mathcal{U}}(X_\tau)]^p \end{aligned}$$

for any such t and τ , and as at the end of the proof of Proposition 4.27 we see that, thanks to (s, p) -variability, the right hand side is integrable over $[0, T]^2$. Since by (4.41) we have

$$\lim_{\varepsilon \rightarrow 0} \frac{|\varphi_\varepsilon(X_t) - \varphi_\varepsilon(X_\tau)|^p}{(t - \tau)^{1+\beta p}} = \frac{|\varphi(X_t) - \varphi(X_\tau)|^p}{(t - \tau)^{1+\beta p}}$$

for any such t and τ , symmetry and dominated convergence imply

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \int_0^T \int_0^T \frac{|\varphi_\varepsilon(X_t) - \varphi_\varepsilon(X_\tau)|^p}{|t - \tau|^{1+\beta p}} d\tau dt \\ &= \lim_{\varepsilon \rightarrow 0} \int_0^T \int_0^T \frac{|\varphi_\varepsilon(X_t) - \varphi_\varepsilon(X_\tau)|^p}{|t - \tau|^{1+\beta p}} \mathbf{1}_{\mathbb{R}^n \setminus S_\varphi}(X_\tau) \mathbf{1}_{\mathbb{R}^n \setminus S_\varphi}(X_t) d\tau dt \\ &= \int_0^T \int_0^T \frac{|\varphi(X_t) - \varphi(X_\tau)|^p}{|t - \tau|^{1+\beta p}} d\tau dt. \end{aligned} \tag{4.45}$$

Applying Fatou's lemma to the left hand side of (4.40) and using (4.42) and (4.45), we obtain

$$\begin{aligned} \int_0^T \frac{|\varphi(X_t)|^p}{t^{\beta p}} dt &\leq \liminf_{\varepsilon \rightarrow 0} \int_0^T \frac{|\varphi_\varepsilon(X_t)|^p}{t^{\beta p}} dt \\ &\leq c \lim_{\varepsilon \rightarrow 0} \left(\int_0^T \int_0^T \frac{|\varphi_\varepsilon(X_t) - \varphi_\varepsilon(X_\tau)|^p}{|t - \tau|^{1+\beta p}} dt d\tau + \int_0^T |\varphi_\varepsilon(X_t)|^p dt \right) \\ &= c \left(\int_0^T \int_0^T \frac{|\varphi(X_t) - \varphi(X_\tau)|^p}{|t - \tau|^{1+\beta p}} dt d\tau + \int_0^T |\varphi(X_t)|^p dt \right). \end{aligned} \tag{4.46}$$

Using Proposition 4.27 on the first integral on the right hand side and readjusting constants, we arrive at (4.34). \square

4.7 Proof of existence and regularity of the integral

Now the proof of Theorem 2.13 follows easily.

Proof of Theorem 2.13. To show that $\varphi \circ X \in W_0^{\beta,1}(0, T)$ as claimed in (i) we have to show that

$$\|\varphi \circ X\|_{\beta,1} = \int_0^T \frac{|\varphi(X_t)|}{t^\beta} dt + [\varphi \circ X]_{\beta,1} < +\infty.$$

Here, the finiteness of the weighted L^1 -term follows from Proposition 4.31 and Lemma 4.29, and the finiteness of the Gagliardo seminorm follows from Proposition 4.27. Since $Y \in \mathcal{C}^\gamma([0, T], \mathbb{R})$ we have $Y \in W_T^{1-\beta, \infty}$ by (2.5), provided that $1 - \beta < \gamma$. Thus the existence of the integral (2.8) as claimed in (ii) follows from Proposition 2.12 by choosing $\beta \in (1 - \gamma, \alpha s)$. To conclude the Hölder regularity claimed in (iii) we can follow [73, Proposition 4.1 (II)] and [92, Proposition 6.2 (i)] and note that, with β and γ as stated

and $0 \leq \tau < t \leq T$, we have

$$\begin{aligned} & \left| \int_0^t \varphi(X_u) dY_u - \int_0^\tau \varphi(X_u) dY_u \right| \\ & \leq \|Y\|_{W_T^{1-\beta, \infty}(0, T)} \left(\int_\tau^t \frac{|\varphi(X_u)|}{(u-\tau)^\beta} du + \int_\tau^t \int_\tau^u \frac{|\varphi(X_u) - \varphi(X_r)|}{(u-r)^{1+\beta}} dr du \right). \end{aligned}$$

If $X \in V(\varphi, s, p)$, then $\varphi \circ X \in W^{\beta, p}(0, T)$ by Propositions 4.27 and 4.31 and Lemma 4.29. For $p = +\infty$ we have

$$\int_\tau^t \frac{|\varphi(X_u)|}{(u-\tau)^\beta} du \leq c \|\varphi \circ X\|_{W^{\beta, \infty}(0, T)} (t-\tau)^{1-\beta}$$

and

$$\int_\tau^t \int_\tau^u \frac{|\varphi(X_u) - \varphi(X_r)|}{(u-r)^{1+\beta}} dr du \leq c \|\varphi \circ X\|_{W^{\beta, \infty}(0, T)} (t-\tau)$$

as in [73]. For $1 \leq p < +\infty$ we can proceed similarly as in [92] and use Hölder's inequality to see that

$$\int_\tau^t \frac{|\varphi(X_u)|}{(u-\tau)^\beta} du \leq c \|\varphi \circ X\|_{W^{\beta, p}(0, T)} (t-\tau)^{(1-\beta)-1/p}.$$

Using $\beta < \beta' < \alpha s$ and $\frac{1}{p} + \frac{1}{q} = 1$ we also have

$$\begin{aligned} & \int_\tau^t \int_\tau^u \frac{|\varphi(X_u) - \varphi(X_r)|}{(u-r)^{1+\beta}} dr du \\ & \leq \int_\tau^t \int_\tau^u \frac{|\varphi(X_u) - \varphi(X_r)|}{(u-r)^{\beta'+1/p}} \frac{1}{(u-r)^{\beta-\beta'+1/q}} dr du \\ & \leq \left(\int_\tau^t \int_\tau^u \frac{|\varphi(X_u) - \varphi(X_r)|}{(u-r)^{\beta'p+1}} dr du \right)^{1/p} \left(\int_\tau^t \int_\tau^r (r-u)^{(\beta'-\beta)q-1} du dr \right)^{1/q} \\ & \leq c \|\varphi \circ X\|_{W^{\beta', p}(0, T)} (t-\tau)^{(1-\beta)-1/p+\beta'}. \end{aligned}$$

□

The following quantitative estimates are a byproduct of the above proof, Propositions 4.27 and 4.31 and (2.7).

Corollary 4.34. *If φ , X , and Y satisfy the hypotheses of Theorem 2.13 (ii), then for any $1 - \gamma < \beta < \alpha s$ and any $t \in [0, T]$ we have*

$$\begin{aligned} & \left| \int_0^t \varphi(X_u) dY_u \right| \\ & \leq c \|Y\|_{W_T^{1-\beta, \infty}(0, T)} \left([X]_{\alpha, \infty}^s \int_0^T \int_{X([0, T])} \frac{\|D\varphi\|(dy)}{|X_t - y|^{n-1+s}} dt + \int_0^T |\varphi(X_t)| dt \right). \end{aligned} \quad (4.47)$$

If φ , X , and Y satisfy the hypotheses of Theorem 2.13 (ii), then for any $1 - \gamma < \beta < \alpha s$ and any $t \in [0, T]$ we have

$$\begin{aligned} & \left\| \int_0^\cdot \varphi(X_u) dY_u \right\|_{\mathcal{C}^{1-\beta}([0, T])} \\ & \leq c \|Y\|_{W_T^{1-\beta, \infty}(0, T)} \left([X]_{\alpha, \infty}^{sp} \int_0^T \left[\int_{X([0, T])} \frac{\|D\varphi\|(dy)}{|X_t - y|^{n-1+s}} \right]^p dt + \int_0^T |\varphi(X_t)|^p dt \right) \end{aligned}$$

with straightforward modification for $p = +\infty$.

4.8 Interpretation as currents

Although it will not be used in the sequel, we briefly comment on an alternative interpretation of (4.47) which is close to the concept of stochastic currents investigated in [35] and [33]. Given a path $X : [0, T] \rightarrow \mathbb{R}^n$ and a number $s \in (0, 1)$, set

$$[\varphi]_{X, s} := \|U^{1-s} \|D\varphi\| \|_{L^1(X)}, \quad \varphi \in BV(\mathbb{R}^n).$$

Obviously this defines a seminorm on $BV(\mathbb{R}^n)$. Recall that a sequence $(\varphi_n)_n \subset BV(\mathbb{R}^n)$ is said to strictly converge to $\varphi \in BV(\mathbb{R}^n)$ if $\lim_n \varphi_n = \varphi$ in $L^1(\mathbb{R}^n)$ and $\lim_n \|D\varphi_n\|(\mathbb{R}^n) = \|D\varphi\|(\mathbb{R}^n)$, [5, Definition 3.14].

Proposition 4.35. *For any path X and any $s \in (0, 1)$ the seminorm $[\cdot]_{X, s}$ is lower semicontinuous on $BV(\mathbb{R}^n)$ w.r.t. strict convergence. Moreover,*

$$\{\varphi \in BV(\mathbb{R}^n) : [\varphi]_{X, s} < +\infty\} \tag{4.48}$$

is a subspace of $BV(\mathbb{R}^n)$, closed w.r.t. strict convergence.

Proof. If $(\varphi_n)_n \subset BV(\mathbb{R}^n)$ strictly converges to $\varphi \in BV(\mathbb{R}^n)$, then, by [5, Proposition 3.15], also $\lim_n \|D\varphi_n\| = \|D\varphi\|$ vaguely. Hence, by [59, Chapter I, Theorem 1.3], we have

$$U^{1-s} \|D\varphi\|(x) \leq \liminf_n U^{1-s} \|D\varphi_n\|(x), \quad x \in \mathbb{R}^n,$$

and by Fatou's lemma, used w.r.t. $\mu_X^{[0, T]}$, the lower semicontinuity follows. The closedness of the above subspace follows similarly. \square

Although it is too strict for most applications, let us mention that the norm $\|\varphi\|_{BV} := \|\varphi\|_{L^1(\mathbb{R}^n)} + \|D\varphi\|(\mathbb{R}^n)$ makes $BV(\mathbb{R}^n)$ a Banach space, and convergence in this norm implies strict convergence. Consequently (4.48) is also closed w.r.t. $\|\varphi\|_{BV}$, hence itself Banach with this norm.

We write $V_{X, s}(\mathbb{R}^n, \mathbb{R}^n)$ for the Banach space of all $\varphi = (\varphi_1, \dots, \varphi_n) \in (BV(\mathbb{R}^n))^n$ with $[\varphi_i]_{X, s} < +\infty$ for all i with norm $\|\varphi\|_{X, s} := \sum_{i=1}^n \left(\|\varphi_i\|_{BV(\mathbb{R}^n)} + [\varphi_i]_{X, s} \right)$.

The following interpretation of the integral as a bounded linear functional on $V_{X, s}(\mathbb{R}^n, \mathbb{R}^n)$ is a special case of (4.47) and seems close to [33, Remark 12].

Corollary 4.36. *Let $X \in \mathcal{C}^\gamma([0, T], \mathbb{R}^n)$ with $\gamma > \frac{1}{2}$ and let $s \in (\frac{1}{\gamma} - 1, 1)$. Then the integral $\int_0^T \varphi(X_u) dX_u$ exists for any $\varphi \in V_{X,s}(\mathbb{R}^n, \mathbb{R}^n)$ and satisfies*

$$\left| \int_0^T \varphi(X_u) dX_u \right| \leq c \|Y\|_{W_T^{1-\beta, \infty}(0, T, \mathbb{R}^n)} (1 + [X]_{\alpha, \infty}^s) \|\varphi\|_{X, s}$$

for any $\beta \in (1 - \gamma, \gamma s)$.

4.9 Proof of the change of variable formula

We provide a proof of Theorem 2.14 which follows by mollification of the coefficient and taking the limit. The following result yields convergence in $W_0^{\beta, 1}(0, T)$.

Lemma 4.37. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ and $X \in \mathcal{C}^\alpha([0, T], \mathbb{R}^n)$. Suppose that $s \in (0, 1)$, $p \in [1, +\infty)$, and $X \in V(\varphi, s, p)$. Set $\varphi_\varepsilon = \varphi * \eta_\varepsilon$, where $(\eta_\varepsilon)_{\varepsilon > 0}$ is a mollifier. Then for any $\beta \in (0, \alpha s)$ we have*

$$\lim_{\varepsilon \rightarrow 0} \|\varphi \circ X - \varphi_\varepsilon \circ X\|_{W_0^{\beta, p}} = 0. \quad (4.49)$$

Proof. Compare also with [31]. Consider first the seminorm part

$$[\varphi \circ X - \varphi_\varepsilon \circ X]_{\beta, p} = \int_0^T \int_0^T \frac{|\varphi(X_t) - \varphi(X_\tau) - \varphi_\varepsilon(X_t) + \varphi_\varepsilon(X_\tau)|^p}{|t - \tau|^{1+\beta p}} d\tau dt. \quad (4.50)$$

Since $X \in V(\varphi, s, p)$, (B.2) and Corollary 4.4 imply that

$$\lim_{\varepsilon \rightarrow 0} |\varphi(X_t) - \varphi_\varepsilon(X_t)| = 0 \quad (4.51)$$

for a.e. $t \in [0, T]$. Thus it suffices to find an integrable upper bound in order to apply dominated convergence theorem. For this we use

$$|\varphi(X_t) - \varphi(X_\tau) - \varphi_\varepsilon(X_t) + \varphi_\varepsilon(X_\tau)|^p \leq 2^{p-1} |\varphi(X_t) - \varphi(X_\tau)|^p + 2^{p-1} |\varphi_\varepsilon(X_t) - \varphi_\varepsilon(X_\tau)|^p.$$

For the first summand on the right hand side we can use Proposition 4.27. For the second, Proposition C.1, the bound (4.30), and Corollary B.3 imply that for ε sufficiently small and \mathcal{L}^1 -a.e. $t, \tau \in [0, T]$,

$$\begin{aligned} & \frac{|\varphi_\varepsilon(X_t) - \varphi_\varepsilon(X_\tau)|^p}{|t - \tau|^{1+\beta p}} \\ & \leq c \frac{|X_t - X_\tau|^{sp}}{|t - \tau|^{1+\beta p}} [U^{1-s}(\|D\varphi_\varepsilon\|_{X([0, T])})(X_t) + U^{1-s}(\|D\varphi_\varepsilon\|_{X([0, T])})(X_\tau)]^p \\ & \leq c \frac{|X_t - X_\tau|^{sp}}{|t - \tau|^{1+\beta p}} [U^{1-s}(\|D\varphi\|_{\mathcal{W}})(X_t) + U^{1-s}(\|D\varphi\|_{\mathcal{W}})(X_\tau)]^p \end{aligned} \quad (4.52)$$

with $\mathcal{U} \supset X([0, T])$ relatively compact open as in Definition 2.1. The quantity (4.50) goes to zero as $\varepsilon \rightarrow 0$ by the dominated convergence theorem. As in (4.40) and (4.46), we obtain

$$\int_0^T \frac{|\varphi(X_t) - \varphi_\varepsilon(X_t)|^p}{t^{\beta p}} dt \leq c \left(\int_0^T \int_0^T \frac{|\varphi(X_t) - \varphi_\varepsilon(X_t) + \varphi_\varepsilon(X_\tau) - \varphi(X_\tau)|^p}{|t - \tau|^{1+\beta p}} dt d\tau + \int_0^T |\varphi(X_t) - \varphi_\varepsilon(X_t)|^p dt \right). \quad (4.53)$$

Recycling the pinning argument, we choose $t_0 \in [0, T]$ such that $X_{t_0} \in \mathbb{R}^n \setminus S_\varphi$ to see that

$$|\varphi_\varepsilon(X_t)|^p \leq 2^{p-1} [|\varphi_\varepsilon(X_t) - \varphi_\varepsilon(X_{t_0})|^p + |\varphi_\varepsilon(X_{t_0})|^p],$$

and treat the difference $|\varphi_\varepsilon(X_t) - \varphi_\varepsilon(X_{t_0})|$ using the maximal inequalities and potential bounds as in the proof of Proposition 4.27, which gives an integrable upper bound. By (4.51) and dominated convergence we have

$$\lim_{\varepsilon \rightarrow 0} \int_0^T |\varphi(X_t) - \varphi_\varepsilon(X_t)|^p dt = 0,$$

and combining with the above, we see that also (4.53) goes to zero as $\varepsilon \rightarrow 0$. \square

We prove Theorem 2.14.

Proof of Theorem 2.14. By Theorem 2.13 all integrals $\int_0^t \partial_k F_\varepsilon(X_u) dX_u^k$ are well defined. Let $F_\varepsilon = F * \eta_\varepsilon$, where $(\eta_\varepsilon)_{\varepsilon>0}$ is a mollifier. Then $\partial_k F_\varepsilon = \partial_k F * \eta_\varepsilon$ and a standard Taylor approximation argument yields

$$F_\varepsilon(X_t) = F_\varepsilon(\hat{x}) + \sum_{k=1}^n \int_0^t \partial_k F_\varepsilon(X_u) dX_u^k. \quad (4.54)$$

By Corollary 4.4, $X_t \in \mathbb{R}^n \setminus \bigcup_{k=1}^n S_{\partial_k F}$ for \mathcal{L}^1 -a.e. $t \in [0, T]$. Thus, by Lemma B.4, $X_t \in \mathbb{R}^n \setminus S_F$ for \mathcal{L}^1 -a.e. $t \in [0, T]$ as well and we have

$$\lim_{\varepsilon \rightarrow 0} F_\varepsilon(X_t) = F(X_t) \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} F_\varepsilon(\hat{x}) = F(\hat{x}).$$

By Lemma 4.37 we have

$$\lim_{\varepsilon \rightarrow 0} \|\partial_k F_\varepsilon(X_\cdot) - \partial_k F(X_\cdot)\|_{W_0^{\beta,1}(0,T)} = 0$$

for each $k = 1, 2, \dots, n$ and for some $\beta \in (1 - \alpha, \alpha s)$. Using (2.7) we therefore arrive at (2.10). Finally, for continuous F we have $S_F = \emptyset$ concluding the proof. \square

5 Existence and uniqueness proofs

In this section we prove Theorems 3.8, 3.24, 3.25 and 3.28.

We make repeated use of the elementary facts that a function $f = (f_1, \dots, f_m)$ is in the space $W^{\beta,p}(0, T, \mathbb{R}^m)$ if and only if all its components f_i are in $W^{\beta,p}(0, T)$, and that, by the norm equivalence in \mathbb{R}^m , the norm $\|f\|_{W^{\beta,p}(0, T, \mathbb{R}^m)}$ is comparable to $\sum_{i=1}^m \|f_i\|_{W^{\beta,p}(0, T)}$ (similarly for other function spaces). We apply the fact that estimates like (2.7) or (2.9) remain valid for vector valued functions of compatible dimensions at the expense of having a different multiplicative constant in front. We also use the symbol $L^p(X, \mathbb{R}^m)$ for a vector valued version of $L^p(X)$.

5.1 Invertibility and BV-regularity

In this subsection we verify Lemma 3.14. Recall that by the Cayley-Hamilton theorem the inverse A^{-1} of an invertible $(n \times n)$ -matrix A satisfies

$$A^{-1} = \frac{(-1)^{n-1}}{\det(A)} (A^{n-1} + c_{n-1}A^{n-2} + \dots + c_1I_n), \quad (5.1)$$

where I_n is the $(n \times n)$ -identity matrix and, for $k = 0, \dots, n-1$, one has

$$c_{n-k} = B_k(s_1, -1!s_2, 2!s_3, \dots, (-1)^{k-1}(k-1)!s_k),$$

with B_k denoting the k -th complete exponential Bell polynomial and $s_l = \text{Tr}(A^l)$ being the trace of A^l . We prove Lemma 3.14.

Proof of Lemma 3.14. Suppose that σ satisfies Assumption 3.12. Recall that we always consider fixed Lebesgue representatives of the components σ_{ij} . Let $N \subset \mathbb{R}^n$ be a \mathcal{L}^n -null set such that $\det(\sigma(x)) > \varepsilon$ and $\sigma_{ij}(x) \leq \|\sigma\|_{L^\infty(\mathbb{R}^n, \mathbb{R}^{n \times n})}$ for all $x \in \mathbb{R}^n \setminus N$ and any i and j . For such x the matrix $\sigma(x)$ is invertible, and we set $\hat{\sigma}(x) := (\sigma(x))^{-1}$. For $x \in N$ we can set $\hat{\sigma}(x) := 0$. For $x \in \mathbb{R}^n \setminus N$ the matrices $\hat{\sigma}(x)$ of $\sigma(x)$ satisfy (5.1) in place of A^{-1} and A . In particular, since for all matrices A the elements of the matrix products A^k , traces $\text{tr}(A^k)$, and the determinant $\det(A)$ are polynomials of the elements of A , we observe that the coefficients $\hat{\sigma}_{ij}(x)$ of $\hat{\sigma}(x)$ are rational functions of the coefficients $\sigma_{ij}(x)$ of the form

$$\hat{\sigma}_{ij}(x) = \frac{P_{ij}(\sigma(x))}{\det(\sigma(x))}, \quad (5.2)$$

where for each i and j the function $P_{ij}(\sigma)$ is a polynomial of degree $n-1$ in the coefficients σ_{kl} , $k, l = 1, \dots, n$, of σ . By the boundedness of the σ_{kl} , clearly $P_{ij}(\sigma) \in L^\infty(\mathbb{R}^n)$ (in the sense of equivalence classes) and, by (5.2), also $\hat{\sigma}_{ij} \in L^\infty(\mathbb{R}^n)$. Since products of bounded $BV(\mathbb{R}^n)$ functions are again in $BV(\mathbb{R}^n)$, [5, Theorem 3.9 and the remarks following its proof], we have $P_{ij}(\sigma) - P_{ij}(0) \in BV(\mathbb{R}^n)$. By the same reason $\det(\sigma) \in BV(\mathbb{R}^n)$, and since its reciprocal is bounded by $\frac{1}{\varepsilon}$, the chain rule implies that $\frac{1}{\det(\sigma)} \in BV(\mathbb{R}^n)$ (recall

Lemma 4.33 or see [5, Theorem 3.96]). Therefore $\hat{\sigma}_{ij} \in BV(\mathbb{R}^n)$ by (5.2). For $x \in \mathbb{R}^n \setminus N$ we have $\det \sigma(x) \leq n! \|\sigma\|_{L^\infty(\mathbb{R}^n, \mathbb{R}^{n \times n})}$, so that

$$\det(\hat{\sigma}(x)) = \frac{1}{\det(\sigma(x))} \geq \frac{1}{n! \|\sigma\|_{L^\infty(\mathbb{R}^n, \mathbb{R}^{n \times n})}}.$$

□

5.2 Solutions to the deterministic equation

In this subsection we provide a proof for Proposition 3.18. Assumptions 3.12 and 3.15 allow to conclude that $\hat{\sigma}$ has a Lipschitz potential.

Proposition 5.1. *Suppose σ satisfies Assumptions 3.12 and 3.15. Then there exists a Lipschitz function $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that its Jacobian matrix ∇g , defined a priori in distributional sense, satisfies*

$$\nabla g = \hat{\sigma} \quad \mathcal{L}^n\text{-a.e.} \quad (5.3)$$

In particular, $g \in W^{1,\infty}(\mathcal{O})$ for any bounded domain $\mathcal{O} \subset \mathbb{R}^n$.

We record a short helpful argument to conclude local continuity and boundedness. By $\mathcal{S}(\mathbb{R}^n, \mathbb{R}^n)$ and $\mathcal{S}'(\mathbb{R}^n, \mathbb{R}^n)$ we denote the spaces of \mathbb{R}^n -valued Schwartz functions and tempered distributions, respectively.

Lemma 5.2. *If $n \geq 2$, $1 < p < +\infty$ and $G \in \mathcal{S}'(\mathbb{R}^n)$ is such that $\nabla G \in L^p(\mathbb{R}^n, \mathbb{R}^n)$, then $G \in L^q(\mathbb{R}^n)$, where $\frac{1}{q} = \frac{1}{p} - \frac{1}{n}$.*

Proof. Given $\varphi \in \mathcal{S}(\mathbb{R}^n)$ let $\psi := -\nabla(-\Delta)^{-1}\varphi$, where Δ denotes the Laplacian on \mathbb{R}^n and $(-\Delta)^{-1}$ the Newton potential. Then $\psi \in \mathcal{S}(\mathbb{R}^n, \mathbb{R}^n)$ and

$$\langle G, \varphi \rangle = -\langle G, \operatorname{div} \psi \rangle = \langle \nabla G, \psi \rangle = \int_{\mathbb{R}^n} (\nabla G) \cdot \psi \, dx,$$

so that

$$\begin{aligned} |\langle G, \varphi \rangle| &\leq \| \|\nabla G\| \|_{L^p(\mathbb{R}^n)} \| \|\nabla(-\Delta)^{-1}\varphi\| \|_{L^{p'}(\mathbb{R}^n)} \\ &\leq c \| \|\nabla G\| \|_{L^p(\mathbb{R}^n)} \| \|(-\Delta)^{-1/2}\varphi\| \|_{L^{p'}(\mathbb{R}^n)} \\ &\leq c \| \|\nabla G\| \|_{L^p(\mathbb{R}^n)} \| \|\varphi\| \|_{L^{q'}(\mathbb{R}^n)} \end{aligned}$$

by the $L^{p'}$ -boundedness of the Riesz transform, [79, Chapter II, Section 4, Theorem 3 and Chapter III, Section 1], and the fractional Sobolev inequality, [79, Chapter V, Section 1, Theorem 1]. Consequently $G \in L^q(\mathbb{R}^n)$ by duality. □

Remark 5.3. Recall that for $q > n$ and for any smooth bounded domain $\mathcal{O} \subset \mathbb{R}^n$ we have the Sobolev embedding $W^{1,q}(\mathcal{O}) \subset \mathcal{C}^\delta(\overline{\mathcal{O}})$, where $0 < \delta \leq 1 - \frac{n}{q}$. For instance, see [3, 4.27].

Using Lemma 5.2 and Remark 5.3 we can prove Proposition 5.1.

Proof. In the case that $n = 1$ we can simply integrate $\hat{\sigma}$ to find a locally bounded Lipschitz function g satisfying (5.3). Suppose $n \geq 2$. By Assumption 3.15 and a distributional version of Poincaré's Lemma, there exists $g = (g_1, \dots, g_n) \in \mathcal{S}'(\mathbb{R}^n, \mathbb{R}^n)$ such that $\nabla g = \hat{\sigma}$ holds in distributional sense. That is, the j th row $\nabla g_j = (D_1 g_j, \dots, D_n g_j)$ of ∇g equals the j th row $\hat{\sigma} = (\hat{\sigma}_{j1}, \dots, \hat{\sigma}_{jn})$ of $\hat{\sigma}$ in $\mathcal{S}'(\mathbb{R}^n, \mathbb{R}^n)$. See [48, Chapter 4, Section 3, Proposition 9] of [77, Chapter II, Section 6, Théorème VI] (the result is stated for the dual of smooth compactly supported functions, but the proof does not change for Schwartz functions and tempered distributions). For any j and k we have $D_k g_j = \hat{\sigma}_{jk} \in L^1(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ by Assumption 3.12 and Lemma 3.14. Now suppose that $\mathcal{O} \subset \mathbb{R}^n$ is a bounded domain. By Lemma 5.2 and the boundedness of \mathcal{O} we can conclude that $g \in W^{1,q}(\mathcal{O}, \mathbb{R}^n)$ for arbitrarily large $q < +\infty$, and Remark 5.3 implies that g is continuous on $\overline{\mathcal{O}}$, and hence also bounded on $\overline{\mathcal{O}}$. As a consequence, we have $g \in W^{1,\infty}(\mathcal{O}, \mathbb{R}^n)$, and a standard mollifier argument yields $|g(x) - g(y)| \leq \|\nabla g\|_{L^\infty(\mathbb{R}^n, \mathbb{R}^{n \times n})} |x - y|$ for all $x, y \in \mathcal{O}$, see [45, Theorem 4.1]. \square

Our proof of Proposition 3.18 is based on an inverse function theorem for Sobolev functions proved in [54, Theorem 1]. We quote a special case of this result sufficient for our purposes. See also [55].

Proposition 5.4. *Let $n \geq 2$. If $g \in W_{\text{loc}}^{1,n}(\mathbb{R}^n, \mathbb{R}^n)$ is non-constant and there exist $\delta > -1$ and $\kappa \in [1, +\infty)$ such that for \mathcal{L}^n -a.e. $x \in \mathbb{R}^n$ we have*

$$\langle \nabla g(x) \xi, \xi \rangle \geq \delta |\nabla g(x) \xi| |\xi|, \quad \xi \in \mathbb{R}^n, \quad (5.4)$$

and

$$|\nabla g(x)|^n \leq \kappa \det(\nabla g(x)), \quad (5.5)$$

then g is a homeomorphism of \mathbb{R}^n onto itself.

Remark 5.5. Condition (5.5) is usually rephrased by saying that g is κ -quasiregular or of bounded distortion, see for instance [47, Section 1.2] or [74]. A result similar to Proposition 5.4 is [21, Theorem 1]. There condition (5.4) is replaced by a \mathcal{L}^n -a.e. nonnegativity condition on the sums of principal minors of σ .

We prove Proposition 3.18.

Proof of Proposition 3.18. By Assumption 3.12, Lemma 3.14, Assumption 3.15, and Proposition 5.1 there exists a Lipschitz function $g \in W_{\text{loc}}^{1,n}(\mathbb{R}^n, \mathbb{R}^n)$ such that (5.3) holds. In the case $n = 1$, Assumption 3.12 and (5.3) imply that $\|\sigma\|_{L^\infty(\mathbb{R})}^{-1} \leq \hat{\sigma}(x) = g'(x)$ for \mathcal{L}^1 -a.e. $x \in \mathbb{R}$, and hence g is strictly monotone and bi-Lipschitz with inverse $f = g^{-1}$ having Lipschitz constant bounded by $\|\sigma\|_{L^\infty(\mathbb{R})}$. Assume $n \geq 2$. Again by Assumption 3.12 and Lemma 3.14 we have

$$\frac{1}{n! \|\sigma\|_{L^\infty(\mathbb{R}^n, \mathbb{R}^{n \times n})}^n} \leq \frac{1}{\det(\sigma(x))} = \det(\hat{\sigma}(x)) = \det(\nabla g(x))$$

for \mathcal{L}^n -a.e. $x \in \mathbb{R}^n$, and by (5.3) also $|\nabla g(x)|^n \leq \|\hat{\sigma}\|_{L^\infty(\mathbb{R}^n, \mathbb{R}^{n \times n})}^n$. Hence (5.5) holds with

$$\kappa = n! \|\hat{\sigma}\|_{L^\infty(\mathbb{R}^n, \mathbb{R}^{n \times n})}^n \|\sigma\|_{L^\infty(\mathbb{R}^n, \mathbb{R}^{n \times n})}^n.$$

Condition (5.4) is immediate from (3.10). Consequently g is a homeomorphism by Proposition 5.4, and we denote its inverse by $f = g^{-1}$. For any bounded domain \mathcal{O} the restriction $g|_{\mathcal{O}}$ is a homeomorphism with inverse $f|_{g(\mathcal{O})}$, and by Proposition 5.1 also $g|_{\mathcal{O}} \in W^{1, n-1}(\mathcal{O}, \mathbb{R}^n)$. This implies that $f \in W_{\text{loc}}^{1,1}(g(\mathcal{O}), \mathbb{R}^n)$, see [47, Theorem 5.2]. Consequently the chain rule can be applied to $g \circ f = \text{id}$, and taking into account (4.1) we obtain

$$I = \nabla(g \circ f)(y) = (\nabla g)(f(y))\nabla f(y) = \sigma^{-1}(f(y))\nabla f(y)$$

for \mathcal{L}^n -a.a. $y \in \mathbb{R}^n$. The invertibility of σ yields $\nabla f = \sigma(f)$ \mathcal{L}^n -a.e. and since $\sigma \in L^\infty(\mathbb{R}^n, \mathbb{R}^{n \times n})$, it follows that f is Lipschitz with Lipschitz constant bounded by $\|\sigma\|_{L^\infty(\mathbb{R}^n, \mathbb{R}^{n \times n})}$. \square

5.3 Existence of solutions in the invertible case

We observe the following stability property of conditions of upper regularity type.

Lemma 5.6. *Let $Y : [0, T] \rightarrow \mathbb{R}^n$ be a path, $B \subset \mathbb{R}^n$ Borel and $d > 0$. Suppose that $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a Borel function with Lipschitz inverse $g = f^{-1}$. Then there is a constant $M > 0$ such that for all $u \in [0, T]$ we have*

$$\int_0^T |f(Y_t) - f(Y_u)|^{-d} dt \leq M \int_0^T |Y_t - Y_u|^{-d} dt.$$

For $n = 1$ this implication remains true if f has an absolutely continuous inverse $g = f^{-1}$ only.

Proof. For any $t, u \in [0, T]$ we have

$$|Y_t - Y_u| = |g(f(Y_t)) - g(f(Y_u))| \leq \text{Lip}(g)|f(Y_t) - f(Y_u)|,$$

and consequently the claimed estimate holds with $M = \text{Lip}(g)^d$. In the case $n = 1$ the function g is absolutely continuous if and only if g' is locally integrable. Let $K \subset \mathbb{R}$ be a compact convex set containing $f(X([0, T]))$ and set $M := \int_K |g'(y)| dy$. Then we have

$$\begin{aligned} |g(f(Y_t)) - g(f(Y_u))| &\leq \int_0^1 |g'(\lambda f(Y_t) + (1-\lambda)f(Y_u))| d\lambda \cdot |f(Y_t) - f(Y_u)| \\ &\leq M |f(Y_t) - f(Y_u)|, \end{aligned}$$

which allows to obtain the same estimate as before. \square

In the next lemma we justify the application of the change of variable (2.10).

Lemma 5.7. *Let g and f be as in Lemma 5.6. Let $s \in (0, 1)$, $\gamma \in (\frac{1}{1+s}, 1)$, $Y \in \mathcal{C}^\gamma([0, T], \mathbb{R}^n)$ with $Y_0 = 0$, $B \subset \mathbb{R}^n$ Borel, and $\hat{x} \in \mathbb{R}^n$. Suppose that σ is upper d -regular on B with $d > n - 1 + s$ or that*

$$\sup_{x \in Y([0, T]) \cap (f^{-1}(B) - f^{-1}(\hat{x}))} \int_0^T |Y_t - x|^{-n+1-s} dt < +\infty, \quad (5.6)$$

and similarly for B^c . Then the path $X : [0, T] \rightarrow \mathbb{R}^n$, defined by $X_t := f(Y_t + g(\hat{x}))$, $t \in [0, T]$, is $(s, 1)$ -variable w.r.t. σ , and

$$X_t = \hat{x} + \int_0^t \sigma(X_u) dY_u, \quad t \in [0, T]. \quad (5.7)$$

Proof. If Y satisfies (5.6), then by Lemma 5.6 we have

$$\sup_{u \in [0, T], X_u \in B} \int_0^T |X_t - X_u|^{-n+1-s} dt < +\infty,$$

and a similar conclusion is true for B^c . Therefore the hypotheses on σ , together with Corollary 4.15 and Proposition 4.2, imply that X is $(s, 1)$ -variable w.r.t. σ . If f is bi-Lipschitz on \mathbb{R}^n it is proper, and applying [5, Theorem 3.16] component-wise, we may conclude that for each j and k the function $\partial_k f_j = \sigma_{jk}(f)$ is in $BV_{\text{loc}}(\mathbb{R}^n)$, see also [46]. By the same theorem and the bijectivity of f we also have

$$\|D(\partial_k f_j)\| = \|D(\sigma_{jk} \circ g^{-1})\| = \|D(g_{\#} \sigma_{jk})\| \leq (\text{Lip}(g))^{n-1} g_{\#} \|D\sigma_{jk}\|,$$

with the notation $g_{\#}$ from [5, Theorem 3.16], which is the pushforward operation on measures, however, defined differently when operating on functions. Using $|f(Y_t^{g(\hat{x})}) - z| \leq \text{Lip}(g) |Y_t^{g(\hat{x})} - g(z)|$, the fact that g is proper, and the fact that $X = f(Y^{g(\hat{x})})$ is $(s, 1)$ -variable w.r.t. σ ,

$$\begin{aligned} & \int_0^T U^{1-s} \|D(\partial_k f_j)\| |K(Y_t^{g(\hat{x})})| dt \\ & \leq (\text{Lip}(g))^{n-1} \int_0^T \int_K |Y_t^{g(\hat{x})} - a|^{-n+1-s} g_{\#} \|D\sigma_{jk}\| (da) dt \\ & = (\text{Lip}(g))^{n-1} \int_0^T \int_{g^{-1}(K)} |Y_t^{g(\hat{x})} - g(z)|^{-n+1-s} \|D\sigma_{jk}\| (dz) dt \\ & \leq (\text{Lip}(g))^{2n-2+s} \int_0^T \int_{f(K)} |f(Y_t^{g(\hat{x})}) - z|^{-n+1-s} \|D\sigma_{jk}\| (dz) dt \\ & < +\infty. \end{aligned}$$

That is, $Y^{g(\hat{x})}$ is $(s, 1)$ -variable w.r.t. $\partial_k f_j$. Consequently, by Theorem 2.14,

$$X_t = f(Y_t^{g(\hat{x})}) = f(Y_0^{g(\hat{x})}) + \int_0^t \nabla f(Y_u^{g(\hat{x})}) dY_u = \hat{x} + \int_0^t \sigma(X_u) dY_u$$

for all $t \in [0, T]$. The stated Hölder regularity is clear since f is Lipschitz. Finally, if $n = 1$ and g is as in Theorem 3.8, we can use the fact that $f'(Y_t) = \sigma(X_t)$ for \mathcal{L}^1 -a.e. $t \in [0, T]$ and arrive at the same conclusion. \square

Lemma 5.8. *Let g and f be as in Lemma 5.6. Let $s \in (0, 1)$, $\gamma \in (\frac{1}{1+s}, 1)$, let $Y = (Y_t)_{t \in [0, T]}$ be a process with \mathbb{P} -a.s. Hölder continuous of order γ as in Theorem 3.25, and let $\hat{x} \in \mathbb{R}^n$. Suppose that there are $\varepsilon \in (0, 1 - s)$, $c > 0$, and $\delta \in (0, n - 1 + s - \varepsilon)$ such that (3.13) and (3.14) hold. Then for \mathbb{P} -a.e. $\omega \in \Omega$ the path $X(\omega) : [0, T] \rightarrow \mathbb{R}^n$, defined by $X_t(\omega) := f(Y_t(\omega) + g(\hat{x}))$, $t \in [0, T]$, is $(s, 1)$ -variable w.r.t. σ , and satisfies*

$$X_t(\omega) = \hat{x} + \int_0^t \sigma(X_u(\omega)) dY_u(\omega), \quad t \in [0, T].$$

Proof. By Lemma 5.6 and the hypotheses we have

$$\begin{aligned} \mathbb{E} \int_0^T |X_t - x|^{-n+\varepsilon} dt &= \mathbb{E} \int_0^T |f(Y_t^{g(\hat{x})}) - x|^{-n+\varepsilon} dt \\ &\leq M \mathbb{E} \int_0^T |Y_t^{g(\hat{x})} - g(x)|^{-n+\varepsilon} dt \\ &= \mathbb{E} \int_0^T |Y_t - (g(x) - g(\hat{x}))|^{-n+\varepsilon} dt \\ &\leq cM |g(x) - g(\hat{x})|^{-n+\delta} \\ &\leq cM \text{Lip}(f)^{n-\delta} |x - \hat{x}|^{-n+\delta}. \end{aligned}$$

Consequently, Corollary 4.23 implies that for \mathbb{P} -a.e. $\omega \in \Omega$ the path $X(\omega)$ is $(s, 1)$ -variable w.r.t. σ . One can now follow the arguments in the proof of Lemma 5.7. \square

We can now verify our existence results.

Proofs of Theorems 3.8 and 3.24. Theorems 3.8 (i) is clear, and it ensures that f and g satisfy the hypotheses of Lemma 5.6. By Proposition 3.18 also the hypotheses on σ of Theorem 3.24 guarantee the existence of f and g are as required in Lemma 5.6. Consequently Lemma 5.7 applies and yields the desired statements. \square

Proof of Theorem 3.25. Proposition 3.18 yields functions f and g as in Lemma 5.6, and by Lemma 5.8 the theorem follows. \square

5.4 Uniqueness of solutions in the invertible case

In this subsection we prove Theorem 3.28. We follow the basic idea of [38, Lemma 3.5] and [81, Theorem 2.2], and employ a smoothing argument which permits to use Riemann sum approximations.

The following Proposition is a multidimensional version of the results proved in [90, Theorem 4.3.1] and straightforward to see from there.

Proposition 5.9. *Let $h = (h_1, \dots, h_n)$ be an element of $\mathcal{C}^2(\mathbb{R}^n, \mathbb{R}^n)$. Then for any $X \in \mathcal{C}^\alpha([0, T], \mathbb{R}^n)$ with $\alpha > \frac{1}{2}$ and any $t \in [0, T]$ we have*

$$h(X_t) = h(X_0) + \int_0^t \nabla h(X_u) dX_u.$$

Moreover, the integrals can be understood also as Riemann-Stieltjes integrals.

Suppose now that σ satisfies Assumptions 3.12 and 3.15, let $\hat{\sigma}$ be as in Lemma 3.14, and let g be the Lipschitz solution to (5.3). That is, $\nabla g = \hat{\sigma}$ \mathcal{L}^n -a.e. on \mathbb{R}^n . We denote by $g_\delta = (g_{1,\delta}, \dots, g_{n,\delta})$ the element-wise smooth approximation of $g = (g_1, \dots, g_n)$ defined by

$$g_{i,\delta} = g_i * \eta_\delta, \quad (5.8)$$

where $(\eta_\delta)_{\delta>0}$ is a mollifier as in Lemma B.4. From the identity $\nabla(g * \eta_\delta) = \nabla g * \eta_\delta$ it is immediate that

$$\nabla g_\delta = \hat{\sigma}_\delta \quad \mathcal{L}^n\text{-a.e.}, \quad (5.9)$$

where $\hat{\sigma}_\delta = ((\hat{\sigma}_{jk})_\delta)_{1 \leq j, k \leq n}$ is the element-wise mollified version of $\hat{\sigma}$ defined by

$$(\hat{\sigma}_{jk})_\delta = \hat{\sigma}_{jk} * \eta_\delta.$$

By the continuity of g , we have

$$\lim_{\delta \rightarrow 0} g_\delta(y) = g(y), \quad y \in \mathbb{R}^n, \quad (5.10)$$

and since $g_i \in W_{\text{loc}}^{1,1}(\mathbb{R}^n)$ for all i and $\hat{\sigma}_{jk} \in BV(\mathbb{R}^n)$ for all j and k by Lemma 3.14, we have

$$\lim_{\delta \rightarrow 0} \hat{\sigma}_\delta(y) = \hat{\sigma}(y), \quad y \in \mathbb{R}^n \setminus S_{\hat{\sigma}} \quad (5.11)$$

by Lemma B.4.

Proposition 5.10. *Suppose that σ satisfies Assumptions 3.12 and 3.15, $Y \in \mathcal{C}^\gamma([0, T], \mathbb{R}^n)$, and $\hat{x} \in \mathbb{R}^n$. If $\alpha \in (\frac{1}{2}, 1)$ and $X \in \mathcal{C}^\alpha([0, T], \mathbb{R}^n)$ is a variability solution for σ and Y started at \hat{x} , then for any $\delta > 0$ and $t \in [0, T]$ we have*

$$g_\delta(X_t) = g_\delta(\hat{x}) + \int_0^t \hat{\sigma}_\delta(X_u) \sigma(X_u) dY_u, \quad (5.12)$$

where g_δ and $\hat{\sigma}_\delta$ are as in (5.9).

To prove Proposition 5.10 we can follow the strategy of [90, Theorem 4.4.2] and proceed by Riemann sum approximation. We make use of the following elementary observation.

Lemma 5.11. *Suppose that $\beta \in (0, 1)$, $t \in [0, T]$ and that $B \in W^{\beta, 1}(0, t, \mathbb{R}^{n \times n}) \cap L^\infty(0, t, \mathbb{R}^{n \times n})$ is an $(n \times n)$ -matrix valued function. If $(A_m)_{\geq 1}$ is a sequence of $(n \times n)$ -matrix valued functions $A_m \in W^{\beta, 1}(0, t, \mathbb{R}^{n \times n}) \cap L^\infty(0, t, \mathbb{R}^{n \times n})$ satisfying*

$$\lim_{m \rightarrow \infty} \|A_m\|_{W^{\beta, 1}(0, t, \mathbb{R}^{n \times n})} = 0 \quad \text{and} \quad \lim_{m \rightarrow \infty} |A_m(u)| = 0 \quad \text{for } \mathcal{L}^1\text{-a.e. } u \in [0, t],$$

then

$$\lim_{m \rightarrow \infty} \|A_m B\|_{W^{\beta, 1}(0, t, \mathbb{R}^{n \times n})} = 0.$$

Proof. By dominated convergence we have

$$\limsup_{m \rightarrow \infty} \int_0^T \frac{|A_m(u)B(u)|}{u^\beta} du \leq \lim_{m \rightarrow \infty} \int_0^T \frac{|A_m(u)||B(u)|}{u^\beta} du = 0.$$

Using

$$|A_m(u)B(u) - A_m(r)B(r)| \leq |A_m(u)||B(u) - B(r)| + |B(r)||A_m(u) - A_m(r)|,$$

together with the facts that, by dominated convergence,

$$\lim_{m \rightarrow \infty} \int_0^t \int_0^t \frac{|A_m(u)||B(u) - B(r)|}{|u - r|^{\beta+1}} dudr = 0$$

and

$$\begin{aligned} \limsup_{m \rightarrow \infty} \int_0^t \int_0^t \frac{|B(r)||A_m(u) - A_m(r)|}{|u - r|^{\beta+1}} dudr \\ \leq \|B\|_{L^\infty(0, t, \mathbb{R}^{n \times n})} \lim_{m \rightarrow \infty} \|A_m\|_{W^{\beta, 1}(0, t, \mathbb{R}^{n \times n})} = 0 \end{aligned}$$

we obtain the result. \square

We prove Proposition 5.10.

Proof. To show that the integral is well-defined we first claim that for any $\beta < \alpha$ s we have

$$\|\hat{\sigma}_\delta(X)\sigma(X)\|_{W_0^{\beta, 1}(0, T, \mathbb{R}^{n \times n})} < \infty. \quad (5.13)$$

To see this, note that $\hat{\sigma}_k$ is bounded on a locally compact neighborhood \mathcal{O} containing $X([0, T])$, and since X is a variability solution, Proposition 4.31 implies that

$$\int_0^T \frac{|\hat{\sigma}_\delta(X_u)\sigma(X_u)|}{u^\beta} du \leq \sup_{x \in \mathcal{O}} |\hat{\sigma}_\delta(x)| \int_0^T \frac{|\sigma(X_u)|}{u^\beta} du < \infty.$$

Using (4.8) we obtain

$$\begin{aligned} \int_0^T \frac{|\hat{\sigma}_\delta(X_u)\sigma(X_u) - \hat{\sigma}_\delta(X_r)\sigma(X_r)|}{|u - r|^{\beta+1}} dudr \\ \leq \sup_{x \in \mathcal{O}} |\hat{\sigma}_\delta(x)| \int_0^T \frac{|\sigma(X_u) - \sigma(X_r)|}{|u - r|^{\beta+1}} dudr \\ + \|\sigma\|_{L^\infty(\mathbb{R}^n, \mathbb{R}^{n \times n})} \int_0^T \frac{|\hat{\sigma}_\delta(X_u) - \hat{\sigma}_\delta(X_r)|}{|u - r|^{\beta+1}} dudr. \end{aligned}$$

Proposition 4.27 implies the boundedness of the first integral on the right hand side, and the second integral is bounded by Hölder continuity of X and differentiability of $\hat{\sigma}_\delta$. Thus we have (5.13), and as a consequence, the integral in (5.12) is well-defined. To verify equation (5.12) we first note that, since g_δ and $\hat{\sigma}_\delta$ are smooth and X is Hölder continuous of order $\alpha > \frac{1}{2}$, Proposition 5.9 yields

$$g_\delta(X_t) = g_\delta(\hat{x}) + \int_0^t \hat{\sigma}_\delta(X_u) dX_u.$$

The integrals can be approximated by Riemann-Stieltjes sums

$$\int_0^t \hat{\sigma}_\delta(X_u) dX_u = \lim_{m \rightarrow \infty} \sum_{j=1}^{N_m} \hat{\sigma}_\delta(X_{t_{j-1}^{(m)}}) \cdot \left(X_{t_j^{(m)}} - X_{t_{j-1}^{(m)}} \right),$$

where $(\pi_m)_m$ is a refining sequence of finite partitions of $[0, t]$ with subinterval endpoints $t_j^{(m)}$ and mesh $|\pi_m| = \max_j |t_j^{(m)} - t_{j-1}^{(m)}|$. In particular, for

$$\Pi_X^m(u) = \sum_{j=1}^{N_m} \hat{\sigma}_\delta(X_{t_{j-1}^{(m)}}) \mathbf{1}_{(t_{j-1}^{(m)}, t_j^{(m)}]}(u), \quad u \in [0, t],$$

we have

$$\lim_{m \rightarrow \infty} \|\hat{\sigma}_\delta(X_\cdot) - \Pi_X^m(\cdot)\|_{W^{\beta,1}(0,t, \mathbb{R}^{n \times n})} = 0 \quad (5.14)$$

and

$$\lim_{m \rightarrow \infty} |\hat{\sigma}_\delta(X_u) - \Pi_X^m(u)| = 0, \quad u \in [0, t], \quad (5.15)$$

which can be seen similarly as in [90, Theorem 4.1.1]. Since X solves (3.1), we obtain

$$\begin{aligned} \int_0^t \hat{\sigma}_\delta(X_u) dX_u &= \lim_{m \rightarrow \infty} \sum_{j=1}^{N_m} \int_{t_{j-1}^{(m)}}^{t_j^{(m)}} \hat{\sigma}_\delta(X_{t_{j-1}^{(m)}}) \sigma(X_u) dY_u \\ &= \lim_{m \rightarrow \infty} \int_0^t \Pi_X^m(u) \sigma(X_u) dY_u. \end{aligned} \quad (5.16)$$

Since $\sigma(X) \in W_0^{\beta,1}(0, t, \mathbb{R}^{n \times n}) \cap L^\infty(0, t, \mathbb{R}^{n \times n})$ we may, by (5.14) and (5.15), apply Lemma 5.11 with

$$A_m(u) = \hat{\sigma}_\delta(X_u) - \Pi_X^m(u) \quad \text{and} \quad B(u) = \sigma(X_u), \quad u \in [0, t],$$

to obtain

$$\lim_{m \rightarrow \infty} \|\hat{\sigma}_\delta(X_\cdot) \sigma(X_\cdot) - \Pi_X^m(\cdot) \sigma(X_\cdot)\|_{W^{\beta,1}(0,t, \mathbb{R}^{n \times n})} = 0.$$

Then, by (2.7) and (5.16),

$$\int_0^t \hat{\sigma}_\delta(X_u) dX_u = \int_0^t \hat{\sigma}_\delta(X_u) \sigma(X_u) dY_u$$

concluding the proof. \square

Theorem 3.28 will follow from (5.12) as $\delta \rightarrow 0$ if taking limits can be justified. To provide this justification we show that variability w.r.t. σ implies variability w.r.t. $\hat{\sigma}$.

Proposition 5.12. *Suppose that $\sigma = (\sigma_{jk})_{1 \leq j, k \leq n}$ satisfies Assumption 3.12 and let $\hat{\sigma}$ be as in Lemma 3.14. If for $s \in (0, 1)$ and $p \in [1, +\infty]$ we have $X \in V(\sigma, s, p)$, then also $X \in V(\hat{\sigma}, s, p)$.*

Proof. Recall (5.2), i.e., that

$$\hat{\sigma}_{ij} = \frac{P_{ij}(\sigma)}{\det(\sigma)},$$

where $P_{ij}(\sigma)$ is a polynomial of degree $n - 1$ in the components σ_{kl} of σ . Writing $p := P_{ij} - P_{ij}(0)$ we recall from the proof of Lemma 3.14 that the functions $p(\sigma)$, $\det(\sigma)$, $\frac{1}{\det(\sigma)}$, and $\frac{p(\sigma)}{\det(\sigma)}$ are elements of $BV(\mathbb{R}^n)$. We provide quantitative bounds for their gradient measures. Consider $u = (p(\sigma), \det(\sigma)) \in BV(\mathbb{R}^n)^2$ and let $F \in \mathcal{C}_c^1(\mathbb{R}^2)$ be such that $F(a, b) = \frac{a}{b}$ for $(a, b) \in [-\|p(\sigma)\|_{L^\infty(\mathbb{R}^n)}, \|p(\sigma)\|_{L^\infty(\mathbb{R}^n)}] \times [\varepsilon, \|\det(\sigma)\|_{L^\infty(\mathbb{R}^n)}]$. Then by [5, Theorem 3.96 and its proof] we have

$$\|DF(u)\| \leq \|\nabla F\|_{\text{sup}} \|Du\| \leq \|\nabla F\|_{\text{sup}} (\|Dp(\sigma)\| + \|D \det(\sigma)\|).$$

With $G \in \mathcal{C}_c^1(\mathbb{R})$ such that $G(b) = \frac{1}{b}$ for $b \in [\varepsilon, \|\det(\sigma)\|_{L^\infty(\mathbb{R}^n)}]$ we similarly obtain

$$\|DG(\det(\sigma))\| \leq \|\nabla G\|_{\text{sup}} \|D \det(\sigma)\|.$$

Since $\hat{\sigma}_{ij} = F(u) + P_{ij}(0)G(\det(\sigma))$, this shows that

$$\|D\hat{\sigma}_{ij}\| \leq c_{ij}(\|Dp(\sigma)\| + \|D \det(\sigma)\|),$$

where $c_{ij} = \|\nabla F\|_{\text{sup}} + |P_{ij}(0)| \|\nabla G\|_{\text{sup}}$. For any $k \geq 1$ let now $F_k \in \mathcal{C}_c^1(\mathbb{R}^k)$ be such that $F_k(a_1, a_2, \dots, a_k) = a_1 a_2 \cdots a_k$ on $[-\|\sigma\|_{L^\infty(\mathbb{R}^n, \mathbb{R}^{n \times n})}, \|\sigma\|_{L^\infty(\mathbb{R}^n, \mathbb{R}^{n \times n})}]^k$. The polynomial $p(\sigma)$ can be written as a finite sum of constant multiples of products

$$\sigma_{i_1 j_1} \sigma_{i_2 j_2} \cdots \sigma_{i_k j_k} = F_k(\sigma_{i_1 j_1}, \sigma_{i_2 j_2}, \dots, \sigma_{i_k j_k}),$$

where $1 \leq k \leq n - 1$ and all factors are elements of the matrix σ . Again by [5, Theorem 3.96], we have

$$\begin{aligned} \|D(\sigma_{i_1 j_1} \sigma_{i_2 j_2} \cdots \sigma_{i_k j_k})\| &= \|\nabla F_k\|_{\text{sup}} \|D(\sigma_{i_1 j_1}, \sigma_{i_2 j_2}, \dots, \sigma_{i_k j_k})\| \\ &\leq \|\nabla F_k\|_{\text{sup}} \sum_{l=1}^k \|D\sigma_{i_l j_l}\|. \end{aligned}$$

Summing up, we obtain

$$\|Dp(\sigma)\| \leq c'_{ij} \sum_{k,l=1}^n \|D\sigma_{kl}\|.$$

Since $\det(\sigma)$ is a polynomial in the components σ_{kl} of σ of degree n , we similarly obtain

$$\|D \det(\sigma)\| \leq c \sum_{k,l=1}^n \|D\sigma_{kl}\|,$$

and combining,

$$\|D\hat{\sigma}_{ij}\| \leq C_{ij} \sum_{k,l=1}^n \|D\sigma_{kl}\|$$

with $C_{ij} = c_{ij}(c'_{ij} + c)$. Clearly this inequality remains true if we restrict the measures to a relatively compact open neighborhood \mathcal{U} of $X([0, T])$ such that X satisfies (2.1) for all σ_{kl} and with this very \mathcal{U} (we can take \mathcal{U} to be the intersection of the neighborhoods from (2.1) for the individual components σ_{kl}). Integrating the nonnegative functions $|X_t - \cdot|^{-n+1-s}$ with respect to these restricted measures, we obtain

$$\begin{aligned} U^{1-s}(\|D\hat{\sigma}_{ij}\| |_{\mathcal{U}})(X_t) &= \int_{\mathcal{U}} |X_t - y|^{-n+1-s} \|D\hat{\sigma}_{ij}\| (dy) \\ &\leq C_{ij} \sum_{k,l=1}^n \int_{\mathcal{U}} |X_t - y|^{-n+1-s} \|D\sigma_{kl}\| (dy) \\ &\leq C_{ij} \sum_{k,l=1}^n U^{1-s}(\|D\sigma_{kl}\| |_{\mathcal{U}})(X_t) \end{aligned}$$

for any $t \in [0, T]$. Taking L^p -norms and using the triangle inequality,

$$\|U^{1-s}(\|D\hat{\sigma}_{ij}\| |_{\mathcal{U}})(X.)\|_{L^p(0,T)} \leq C_{ij} \sum_{k,l=1}^n \|U^{1-s}(\|D\sigma_{kl}\| |_{\mathcal{U}})(X.)\|_{L^p(0,T)} < +\infty.$$

□

We prove Theorem 3.28.

Proof of Theorem 3.28. Since by Proposition 5.12 the path X is $(s, 1)$ -variable w.r.t. $\hat{\sigma}$, we have $X_u \in \mathbb{R}^n \setminus S_{\hat{\sigma}}$ for \mathcal{L}^1 -a.e. $u \in [0, t]$, and, together with (5.11),

$$\lim_{\delta \rightarrow 0} |\hat{\sigma}_{\delta}(X_u)\sigma(X_u) - I| = 0$$

for such u . Since also $\hat{\sigma}(X.) \in L^{\infty}(0, t, \mathbb{R}^{n \times n})$ by (4.8), dominated convergence, together with the boundedness of σ and $\hat{\sigma}$, implies

$$\lim_{\delta \rightarrow 0} \int_0^t \frac{|\hat{\sigma}_{\delta}(X_u)\sigma(X_u) - I|}{u^{\beta}} du = 0.$$

For the Gagliardo seminorm we first note that

$$\begin{aligned} |\hat{\sigma}_{\delta}(X_u)\sigma(X_u) - \hat{\sigma}_{\delta}(X_r)\sigma(X_r)| &\leq \|\hat{\sigma}(X.)\|_{L^{\infty}(0,t,\mathbb{R}^{n \times n})} |\sigma(X_u) - \sigma(X_r)| \\ &\quad + \|\sigma(X.)\|_{L^{\infty}(0,t,\mathbb{R}^{n \times n})} |\hat{\sigma}_{\delta}(X_u) - \hat{\sigma}_{\delta}(X_r)|. \end{aligned}$$

Since we have

$$[\sigma(X.)]_{\beta,1} \leq c[X]_{\alpha,\infty}^s \sum_{i,j=1}^n \|U^{1-s} \|D\hat{\sigma}_{ij}\| \|_{L^1(X)}$$

by (4.26) in Proposition 4.27 and, reasoning as in (4.52), also

$$[\sigma_\delta(X.)]_{\beta,1} \leq c[X]_{\alpha,\infty}^s \sum_{i,j=1}^n \|U^{1-s} \|D\hat{\sigma}_{ij}\| \|_{L^1(X)}$$

with a constant $c > 0$ independent of δ , we can again use dominated convergence to conclude that

$$\lim_{\delta \rightarrow 0} \int_0^t \int_0^t \frac{|\hat{\sigma}_\delta(X_u)\sigma(X_u) - \hat{\sigma}_\delta(X_r)\sigma(X_r)|}{|u-r|^{\beta+1}} dudr = 0.$$

This shows that

$$\lim_{\delta \rightarrow 0} \|\hat{\sigma}_\delta(X.)\sigma(X.) - I\|_{W_0^{\beta,1}(0,t,\mathbb{R}^{n \times n})} = 0,$$

and using (2.7) we obtain

$$\lim_{\delta \rightarrow 0} \int_0^t \hat{\sigma}_\delta(X_u)\sigma(X_u)dY_u = \int_0^t dY_u = Y_t - Y_0.$$

Taking into account (5.10) and (5.12), we arrive at

$$g(X_t) - g(\hat{x}) = Y_t - Y_0, \quad t \in [0, T]. \quad (5.17)$$

Now suppose that X and \tilde{X} are two solutions. By (5.17) we must have

$$g(X_t) = g(\tilde{X}_t), \quad t \in [0, T].$$

Since g is invertible with inverse f satisfying (3.7) (cf. Proof of Theorem 3.24), this implies that

$$X_t = \tilde{X}_t = f(Y_t), \quad t \in [0, T].$$

□

A Convolution of Riesz kernels

Statement (i) in the following proposition is well known, see [76, Lemma 25.2 and (25.38)], [59, Theorem 1.15 in Chapter I] or [79, Chapter V, Section 1.1], and it can also be obtained by a quick subordination argument. Statement (ii) is a direct consequence of (i) and Fubini.

Proposition A.1. *Let $\gamma_1, \gamma_2 > 0$ and $\gamma_1 + \gamma_2 < n$.*

(i) We have

$$\int_{\mathbb{R}^n} |x|^{-n+\gamma_1} |x-y|^{-n+\gamma_2} dx = |y|^{-n+\gamma_1+\gamma_2}, \quad y \in \mathbb{R}^n \setminus \{0\}.$$

(ii) If ν_1 and ν_2 are finite nonnegative Borel measures with compact support, then

$$\int_{\mathbb{R}^n} U^{\gamma_1} \nu_1(x) U^{\gamma_2} \nu_2(x) dx = \int_{\mathbb{R}^n} U^{\gamma_2} U^{\gamma_1} \nu_1(x) \nu_2(dx) = \int_{\mathbb{R}^n} U^{\gamma_1+\gamma_2} \nu_1(x) \nu_2(dx).$$

B Mollification results

We collect some useful known approximation results used in the main text.

We begin with an approximation lemma for Riesz potentials that is a slight variant of [59, Section I.3, Theorem 1.11 and its proof]. As usual we say that $(\eta_\varepsilon)_{\varepsilon>0}$ is a (radially symmetric) *mollifier* if $\eta \in \mathcal{C}_c^\infty(\mathbb{R}^n)$ is a nonnegative radial function, compactly supported inside the unit ball and such that $\int_{\mathbb{R}^n} \eta(x) dx = 1$, and $\eta_\varepsilon(x) := \varepsilon^{-n} \eta(\varepsilon^{-1}x)$, $x \in \mathbb{R}^n$, for any $\varepsilon > 0$, cf. [5, p. 41]. We say that a mollifier $(\eta_\varepsilon)_{\varepsilon>0}$ is a *flat mollifier* if for all $x \in \mathbb{R}^n$ with $|x| \leq \frac{1}{2}$ we have $\eta_\varepsilon(x) = c_\eta$ with a suitable constant $c_\eta > 0$. To have η constant in a small ball around the origin is useful to quickly see the following.

Lemma B.1. *Let $(\eta_\varepsilon)_{\varepsilon>0}$ be a flat mollifier, let ν be a non-negative Borel measure on \mathbb{R}^n , and $0 < \gamma < n$. Then:*

(i) *There is a constant $c(n, \gamma) > 0$ depending only on n and γ such that for any $\varepsilon > 0$ and all $x \in \mathbb{R}^n$ we have*

$$U^\gamma(\nu * \eta_\varepsilon)(x) \leq c(n, \gamma) U^\gamma \nu(x).$$

(ii) *For any $x \in \mathbb{R}^n$ we have*

$$\lim_{\varepsilon \rightarrow 0} U^\gamma(\nu * \eta_\varepsilon)(x) = U^\gamma \nu(x).$$

The proof is as in [59, Section I.3, Theorem 1.11], but since we use a slightly different mollifier, we repeat the short arguments for (i) for convenience.

Proof. We have

$$\begin{aligned} U^\gamma(\nu * \eta_\varepsilon)(x) &= \int_{\mathbb{R}^n} \eta_\varepsilon(z-x) \int_{\mathbb{R}^n} \frac{\nu(dy)}{|z-y|^{n-\gamma}} dz \\ &= \int_{\mathbb{R}^n} \frac{1}{|x-y|^{n-\gamma}} \left(\int_{\mathbb{R}^n} \eta_\varepsilon(z-x) \frac{|x-y|^{n-\gamma}}{|z-y|^{n-\gamma}} dz \right) \nu(dy). \end{aligned}$$

The inner integral is bounded by $\Phi(\frac{x}{\varepsilon}, \frac{y}{\varepsilon})$, where

$$\Phi(x, y) := |x-y|^{n-\gamma} \int_{|\zeta-x| \leq \frac{1}{2}} \frac{d\zeta}{|\zeta-y|^{n-\gamma}}.$$

The function Φ defines a continuous function of $|x - y|$ which is zero if $|x - y| = 0$ and tends to 1 for $|x - y| \rightarrow +\infty$, and denoting the maximum of this function by $c(n, \gamma)$, we obtain (i). For (ii) one can follow the proof in [59, p. 73]. \square

The next lemma contain versions of a statement from [5, Theorem 2.2]. Given a Borel set $E \subset \mathbb{R}^n$ and $\varepsilon > 0$, write $E_\varepsilon := \{x \in \mathbb{R}^n : \text{dist}(E, x) < \varepsilon\}$.

Lemma B.2. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ and let $(\eta_\varepsilon)_\varepsilon$ be a mollifier. Then for any Borel set $E \subset \mathbb{R}^n$ and any Borel function $\psi : \mathbb{R}^n \rightarrow [0, +\infty]$ we have*

$$\int_{\mathbb{R}^n} \psi(y) \|D(\varphi * \eta_\varepsilon)\| |_E(dy) \leq \int_{\mathbb{R}^n} \psi(y) (\|D\varphi\| |_{E_\varepsilon} * \eta_\varepsilon)(y) dy. \quad (\text{B.1})$$

In particular,

$$\int_E |\nabla(\varphi * \eta_\varepsilon)|(y) dy \leq \|D\varphi\| (E_\varepsilon).$$

The proof basically follows [5, Theorem 2.2].

Proof. Since $\|D(\varphi * \eta_\varepsilon)\| = |\nabla(\varphi * \eta_\varepsilon)| \cdot \mathcal{L}^n$ and $\nabla(\varphi * \eta_\varepsilon) = (D\varphi) * \eta_\varepsilon$, [5, Proposition 3.2], the left hand side of inequality (B.1) is seen to equal

$$\begin{aligned} \int_{\mathbb{R}^n} \psi(y) \mathbf{1}_E(y) \left| \int_{\mathbb{R}^n} \eta_\varepsilon(y - z) D\varphi(dz) \right| dy &\leq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \psi(y) \mathbf{1}_E(y) \eta_\varepsilon(y - z) dy \|D\varphi\| (dz) \\ &\leq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \psi(y) \eta_\varepsilon(y - z) dy \mathbf{1}_{E_\varepsilon}(z) \|D\varphi\| (dz), \end{aligned}$$

where we have used Fubini's theorem. Another application of the latter shows that the last integral equals the right hand side of (B.1). The case $\psi \equiv 1$ yields the second statement because the mollifier has integral equal to one. \square

We obtain the following consequence for potentials.

Corollary B.3. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$, $0 < \gamma < n$, let $K \subset \mathbb{R}^n$ be a compact set and $(\eta_\varepsilon)_{\varepsilon > 0}$ a symmetric flat mollifier. Then*

$$U^\gamma(\|D(\varphi * \eta_\varepsilon)\| |_K)(x) \leq c(n, \gamma) U^\gamma(\|D\varphi\| |_{K_\varepsilon})(x)$$

for any $x \in \mathbb{R}^n$ and $\varepsilon > 0$.

Proof. Given $x \in \mathbb{R}^n$ write $\psi_x(y) := |x - y|^{-n+\gamma}$. Then

$$\begin{aligned} U^\gamma(\|D(\varphi * \eta_\varepsilon)\| |_K)(x) &= \int_{\mathbb{R}^n} \psi_x(y) \|D(\varphi * \eta_\varepsilon)\| |_K(dy) \\ &\leq \int_{\mathbb{R}^n} \psi_x(y) (\|D\varphi\| |_{K_\varepsilon} * \eta_\varepsilon)(y) dy = U^\gamma(\|D\varphi\| |_{K_\varepsilon} * \eta_\varepsilon)(x) \end{aligned}$$

by Lemma B.2, and an application of Lemma B.1 yields the desired bound. \square

Let $(\eta_\varepsilon)_{\varepsilon>0}$ be a mollifier. Recall that if $\varphi \in L^1_{\text{loc}}(\mathbb{R}^n)$ and $\tilde{\varphi}$ is a Lebesgue representative of φ , then

$$\lim_{\varepsilon \rightarrow 0} \varphi * \eta_\varepsilon(x) = \tilde{\varphi}(x), \quad x \in \mathbb{R}^n \setminus S_\varphi, \quad (\text{B.2})$$

as shown in [5, Proposition 3.64 (b)].

Lemma B.4. *Let $F \in W^{1,1}_{\text{loc}}(\mathbb{R}^n)$, such that $\sigma_k := \partial_k F \in BV_{\text{loc}}(\mathbb{R}^n)$ for $k = 1, \dots, n$. Then $S_F \subset \bigcup_{k=1}^n S_{\sigma_k}$ for $k = 1, \dots, n$. If $(\eta_\varepsilon)_{\varepsilon>0}$ is a mollifier, then*

$$(i) \lim_{\varepsilon \rightarrow 0} F * \eta_\varepsilon = F \text{ pointwise on } \mathbb{R}^n \setminus S_F,$$

$$(ii) \lim_{\varepsilon \rightarrow 0} \sigma_k * \eta_\varepsilon = \lim_{\varepsilon \rightarrow 0} F * \partial_k \eta_\varepsilon = \sigma_k \text{ pointwise on } \mathbb{R}^n \setminus S_{\sigma_k}.$$

Proof. We show that $\bigcap_{k=1}^n (\mathbb{R}^n \setminus S_{\sigma_k}) \subset (\mathbb{R}^n \setminus S_F)$. Once this is shown, (i) and (ii) follow using (B.2) and $F * \partial_k \eta_\varepsilon = \partial_k(F * \eta_\varepsilon) = \sigma_k * \eta_\varepsilon$.

Let $x \in \bigcap_{k=1}^n (\mathbb{R}^n \setminus S_{\sigma_k})$. Then, writing $\sigma := (\sigma_1, \dots, \sigma_n)$, we see that $x \in \mathbb{R}^n \setminus S_{|\sigma|}$. Let $r > 0$. For any $\varepsilon > 0$ there exists $r > \delta > 0$ such that

$$\begin{aligned} & \int_0^r t^{-n} \int_{B(x,t)} |\sigma(y)| dy dt \\ &= \int_\delta^r t^{-n} \int_{B(x,t)} |\sigma(y)| dy dt + c(n) \int_0^\delta \frac{1}{\mathcal{L}^n(B(x,t))} \int_{B(x,t)} |\sigma(y)| dy dt \\ &\leq \frac{(r-\delta)}{\delta^n} \|\sigma\|_{L^1(B(x,r))} + \delta c(n) (\lambda_{|\sigma|}(x) + \varepsilon) \\ &< +\infty. \end{aligned}$$

Moreover, for sufficiently small $\varrho > 0$,

$$\frac{1}{\varrho^{n-1}} \int_{B(x,\varrho)} |\sigma(y)| dy = c(n) \frac{\varrho}{\mathcal{L}^n(B(x,\varrho))} \int_{B(x,\varrho)} |\sigma(y)| dy \leq c(n) \varrho (\lambda_{|\sigma|}(x) + \varepsilon)$$

that converges to zero as $\varrho \searrow 0$. Therefore, by [5, Remark 3.82 and Exercise 3.14], we have $x \in \mathbb{R}^n \setminus S_F$. \square

C Some properties of maximal functions

We record a version of [1, Corollary 4.3]. See also [44, 52]. For any given measure ν on $(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^d))$, any $\gamma \in [0, 1)$, and any $R \in (0, +\infty]$, let

$$\mathcal{M}_{\gamma,R}\nu(x) := \sup_{0 < r < R} r^{\gamma-n} \nu(B(x,r)), \quad x \in \mathbb{R}^n,$$

denote the (truncated) fractional Hardy-Littlewood maximal function of ν of order γ . In the case $R = +\infty$ we use the standard notation $\mathcal{M}_\gamma\nu = \mathcal{M}_{\gamma,+\infty}\nu$.

Proposition C.1. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ and $s \in (0, 1]$. Then there is a constant $c = c(n, s) > 0$ such that for all Lebesgue points $x, y \in \mathbb{R}^n$ of φ we have*

$$|\varphi(x) - \varphi(y)| \leq c|x-y|^s (\mathcal{M}_{1-s,4|x-y|} \|D\varphi\|(x) + \mathcal{M}_{1-s,4|x-y|} \|D\varphi\|(y)).$$

In order to prove the claim, we quote [1, Lemma 4.1 and its proof]. For any $0 < s < +\infty$, any $R > 0$, and any locally integrable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, the fractional sharp maximal function $f_{s,R}^\#$ of f is defined by

$$f_{s,R}^\#(x) := \sup_{0 < r < R} r^{-s-n} \int_{B(x,r)} |f - f_{B(x,r)}| dy.$$

Lemma C.2. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be locally integrable and $0 < s < +\infty$. Then there is a constant $c(n, s) > 0$ such that for all Lebesgue points $x, y \in \mathbb{R}^n$ of f we have*

$$|f(x) - f(y)| \leq c(n, s) |x - y|^s \left(f_{s,4|x-y|}^\#(x) + f_{s,4|x-y|}^\#(y) \right).$$

We prove Proposition C.1.

Proof. For any $x \in \mathbb{R}^n$ and $r > 0$ we have the 1-1-Poincaré inequality

$$\int_{B(x,r)} |f - f_{B(x,r)}| dy \leq c r \|Df\| (B(x, r)),$$

see [5, Remark 3.45]. Multiplying both sides by r^{-s-n} and taking suprema we obtain

$$\sup_{0 < r < R} r^{-s-n} \int_{B(x,r)} |f - f_{B(x,r)}| dy \leq c \mathcal{M}_{1-s,R} \|Df\| (x).$$

By [1, Lemma 4.1] there is a constant $c(n, s) > 0$ such that for all Lebesgue points $x, y \in \mathbb{R}^n$ of f we have

$$|f(x) - f(y)| \leq c(n, s) |x - y|^s \left(f_{s,4|x-y|}^\#(x) + f_{s,4|x-y|}^\#(y) \right).$$

□

It is trivial to see that for any Borel measure ν on \mathbb{R}^n , any $\gamma \in (0, 1)$, and any $R \in (0, +\infty]$ we have

$$\mathcal{M}_{\gamma,R} \nu(x) \leq U^\gamma \nu(x), \quad x \in \mathbb{R}^n. \quad (\text{C.1})$$

Together with Proposition C.1 we obtain the following immediate consequence.

Corollary C.3. *Let $\varphi \in BV_{\text{loc}}(\mathbb{R}^n)$ and $s \in (0, 1)$. If*

$$\sup_{x \in \mathbb{R}^n} U^{1-s} \|D\varphi\| (x) < +\infty,$$

then φ has a Borel version that is Hölder continuous of order s , and any Lebesgue representative of φ coincides with this version on the Lebesgue set of φ .

D Elements of fractional calculus

We briefly recall the definitions of fractional integrals and derivatives, [76], and the generalized Lebesgue-Stieltjes integral introduced by Zähle in [90] and used in [73, 91].

For fixed $T < \infty$, the fractional left and right Riemann–Liouville integrals of order $\theta > 0$ of a function $f \in L_1(0, T)$ are denoted by

$$I_{0+}^\theta f(t) = \frac{1}{\Gamma(\theta)} \int_0^t \frac{f(s)}{(t-s)^{1-\theta}} ds$$

and

$$I_{T-}^\theta f(t) = \frac{(-1)^{-\theta}}{\Gamma(\theta)} \int_t^T \frac{f(s)}{(t-s)^{1-\theta}} ds.$$

The integral operators $I_{0+}^\theta, I_{T-}^\theta : L_1(0, T) \rightarrow L_1(0, T)$ are linear and one-to-one, and the inverse operators, denoted by $I_{0+}^{-\theta} = (I_{0+}^\theta)^{-1}$ and $I_{T-}^{-\theta} = (I_{T-}^\theta)^{-1}$, are known as Riemann–Liouville fractional derivatives. Furthermore, for any $\theta \in (0, 1)$ and for any $f \in I_{0+}^\theta(L_1(0, T))$ and $g \in I_{T-}^\theta(L_1(0, T))$, the Weyl–Marchaud derivatives

$$D_{0+}^\theta f(t) = \frac{1}{\Gamma(1-\theta)} \left(\frac{f(t)}{t^\theta} + \theta \int_0^t \frac{f(t) - f(s)}{(t-s)^{\theta+1}} ds \right)$$

and

$$D_{T-}^\theta g(t) = \frac{(-1)^\theta}{\Gamma(1-\theta)} \left(\frac{g(t)}{(T-t)^\theta} + \theta \int_t^T \frac{g(t) - g(s)}{(s-t)^{\theta+1}} ds \right)$$

are well defined, and coincide with the Riemann–Liouville derivatives by relations $D_{0+}^\theta f(t) = I_{0+}^{-\theta} f(t)$ and $D_{T-}^\theta g(t) = I_{T-}^{-\theta} g(t)$ for almost every $t \in (0, T)$.

For functions f and g such that the limits $f(0+), g(0+), g(T-)$ exist in \mathbb{R} , set $f_{0+}(t) = f(t) - f(0+)$ and $g_{T-}(t) = g(t) - g(T-)$. If $f_{0+} \in I_{0+}^\theta(L_p(0, T))$ and $g_{T-} \in I_{T-}^{1-\theta}(L_q(0, T))$ for some $\theta \in [0, 1]$ and $p, q \in [1, \infty]$ such that $1/p + 1/q = 1$, the fractional version of the Stieltjes integral introduced by Zähle [90] is defined by

$$\begin{aligned} \int_0^T f_t dg_t := & (-1)^\theta \int_0^T D_{0+}^\theta(f - f(0+))(t) D_{T-}^{1-\theta}(g - g(T-))(t) dt \\ & + f(0+)(g(T-) - g(0+)), \end{aligned} \quad (\text{D.1})$$

It can be shown that here the right side does not depend on θ . Moreover, if $\theta p < 1$, then

$$\int_0^T f_t dg_t = (-1)^\theta \int_0^T D_{0+}^\theta f(t) D_{T-}^{1-\theta}(g - g(T-))(t) dt \quad (\text{D.2})$$

which coincides with Definition 2.11.

References

- [1] D. Aalto and J. Kinnunen. Maximal functions in Sobolev spaces. In *Sobolev spaces in mathematics. I*, volume 8 of *Int. Math. Ser. (N. Y.)*, pages 25–67. Springer, New York, 2009.
- [2] D. R. Adams and L. I. Hedberg. *Function spaces and potential theory*, volume 314 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1996.
- [3] R. A. Adams and J. J. F. Fournier. *Sobolev spaces*, volume 140 of *Pure and Applied Mathematics (Amsterdam)*. Elsevier/Academic Press, Amsterdam, second edition, 2003.
- [4] L. Ambrosio and G. Dal Maso. A general chain rule for distributional derivatives. *Proc. Amer. Math. Soc.*, 108(3):691–702, 1990.
- [5] L. Ambrosio, N. Fusco, and D. Pallara. *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2000.
- [6] W. Auzinger. Sectorial operators and normalized numerical range. *Appl. Numer. Math.*, 45(4):367–388, 2003.
- [7] A. Ayache, D. Wu, and Y. Xiao. Joint continuity of the local times of fractional Brownian sheets. *Ann. Inst. Henri Poincaré Probab. Stat.*, 44(4):727–748, 2008.
- [8] M. Barchiesi, D. Henao, and C. Mora-Corral. Local invertibility in Sobolev spaces with applications to nematic elastomers and magnetoelasticity. *Arch. Ration. Mech. Anal.*, 224(2):743–816, 2017.
- [9] L. Barreira, Y. Pesin, and J. Schmeling. Dimension and product structure of hyperbolic measures. *Ann. of Math. (2)*, 149(3):755–783, 1999.
- [10] S. M. Berman. Local times and sample function properties of stationary Gaussian processes. *Trans. Amer. Math. Soc.*, 137:277–299, 1969.
- [11] S. M. Berman. Gaussian processes with stationary increments: Local times and sample function properties. *Ann. Math. Statist.*, 41:1260–1272, 1970.
- [12] S. M. Berman. Local nondeterminism and local times of Gaussian processes. *Bull. Amer. Math. Soc.*, 79:475–477, 1973.
- [13] J. Bertoin. Sur la mesure d’occupation d’une classe de fonctions self-affines. *Japan J. Appl. Math.*, 5(3):431–439, 1988.
- [14] A. Bevilacqua and F. Flandoli. An occupation time formula for semimartingales in \mathbb{R}^N . *Stochastic Process. Appl.*, 124(10):3342–3361, 2014.

- [15] J. Bliedtner and W. Hansen. *Potential theory*. Universitext. Springer-Verlag, Berlin, 1986. An analytic and probabilistic approach to balayage.
- [16] R. Catellier and M. Gubinelli. Averaging along irregular curves and regularisation of ODEs. *Stochastic Process. Appl.*, 126(8):2323–2366, 2016.
- [17] Z. Chen. *On Pathwise Stochastic Integration of Processes with Unbounded Power Variation*. PhD thesis, Aalto University, 2016.
- [18] Z. Chen, L. Leskelä, and L. Viitasaari. Pathwise Stieltjes integrals of discontinuously evaluated stochastic processes. *Stochastic Process. Appl.*, 129(8):2723–2757, 2019.
- [19] Z.-Q. Chen and R. Song. Hardy inequality for censored stable processes. *Tohoku Math. J. (2)*, 55(3):439–450, 2003.
- [20] G. Crippa and C. De Lellis. Estimates and regularity results for the DiPerna-Lions flow. *J. Reine Angew. Math.*, 616:15–46, 2008.
- [21] M. Cristea. A note on the global injectivity of some light Sobolev mappings. *J. Math. Pures Appl. (9)*, 128:213–224, 2019.
- [22] A. M. Davie. Uniqueness of solutions of stochastic differential equations. *Int. Math. Res. Not. IMRN*, (24):Art. ID rnm124, 26, 2007.
- [23] E. Di Nezza, G. Palatucci, and E. Valdinoci. Hitchhiker’s guide to the fractional Sobolev spaces. *Bull. Sci. Math.*, 136(5):521–573, 2012.
- [24] H. Doss. Liens entre équations différentielles stochastiques et ordinaires. *Ann. Inst. H. Poincaré Sect. B (N.S.)*, 13(2):99–125, 1977.
- [25] B. Dyda. A fractional order Hardy inequality. *Illinois J. Math.*, 48(2):575–588, 2004.
- [26] H. J. Engelbert and W. Schmidt. On one-dimensional stochastic differential equations with generalized drift. In *Stochastic differential systems (Marseille-Luminy, 1984)*, volume 69 of *Lect. Notes Control Inf. Sci.*, pages 143–155. Springer, Berlin, 1985.
- [27] H. J. Engelbert and W. Schmidt. On solutions of one-dimensional stochastic differential equations without drift. *Z. Wahrsch. Verw. Gebiete*, 68(3):287–314, 1985.
- [28] K. Falconer. *Fractal geometry*. John Wiley & Sons, Ltd., Chichester, 1990. Mathematical foundations and applications.
- [29] H. Federer. *Geometric measure theory*. Die Grundlehren der mathematischen Wissenschaften, Band 153. Springer-Verlag New York Inc., New York, 1969.
- [30] A. Figalli. Existence and uniqueness of martingale solutions for SDEs with rough or degenerate coefficients. *J. Funct. Anal.*, 254(1):109–153, 2008.

- [31] A. Fiscella, R. Servadei, and E. Valdinoci. Density properties for fractional Sobolev spaces. *Ann. Acad. Sci. Fenn. Math.*, 40(1):235–253, 2015.
- [32] F. Flandoli. *Random perturbation of PDEs and fluid dynamic models*, volume 2015 of *Lecture Notes in Mathematics*. Springer, Heidelberg, 2011. Lectures from the 40th Probability Summer School held in Saint-Flour, 2010, École d’Été de Probabilités de Saint-Flour. [Saint-Flour Probability Summer School].
- [33] F. Flandoli, M. Gubinelli, M. Giaquinta, and V. M. Tortorelli. Stochastic currents. *Stochastic Process. Appl.*, 115(9):1583–1601, 2005.
- [34] F. Flandoli, M. Gubinelli, and E. Priola. Well-posedness of the transport equation by stochastic perturbation. *Invent. Math.*, 180(1):1–53, 2010.
- [35] F. Flandoli, M. Gubinelli, and F. Russo. On the regularity of stochastic currents, fractional Brownian motion and applications to a turbulence model. *Ann. Inst. Henri Poincaré Probab. Stat.*, 45(2):545–576, 2009.
- [36] I. Fonseca and W. Gangbo. *Degree theory in analysis and applications*, volume 2 of *Oxford Lecture Series in Mathematics and its Applications*. The Clarendon Press, Oxford University Press, New York, 1995. Oxford Science Publications.
- [37] I. Fonseca and W. Gangbo. Local invertibility of Sobolev functions. *SIAM J. Math. Anal.*, 26(2):280–304, 1995.
- [38] J. Garzón, J. A. León, and S. Torres. Fractional stochastic differential equation with discontinuous diffusion. *Stoch. Anal. Appl.*, 35(6):1113–1123, 2017.
- [39] D. Geman and J. Horowitz. Occupation densities. *Ann. Probab.*, 8(1):1–67, 1980.
- [40] C. C. Graham, K. E. Hare, and D. L. Ritter. The size of L^p -improving measures. *J. Funct. Anal.*, 84(2):472–495, 1989.
- [41] M. Gubinelli. Controlling rough paths. *J. Funct. Anal.*, 216(1):86–140, 2004.
- [42] D. P. Hardin and E. B. Saff. Minimal Riesz energy point configurations for rectifiable d -dimensional manifolds. *Adv. Math.*, 193(1):174–204, 2005.
- [43] K. E. Hare and M. Roginskaya. The energy of signed measures. *Proc. Amer. Math. Soc.*, 132(2):397–406, 2004.
- [44] T. Heikkinen, J. Lehrbäck, J. Nuutinen, and H. Tuominen. Fractional maximal functions in metric measure spaces. *Anal. Geom. Metr. Spaces*, 1:147–162, 2013.
- [45] J. Heinonen. *Lectures on Lipschitz analysis*, volume 100 of *Report. University of Jyväskylä Department of Mathematics and Statistics*. University of Jyväskylä, Jyväskylä, 2005.

- [46] S. Hencl. Bilipschitz mappings with derivatives of bounded variation. *Publ. Mat.*, 52(1):91–99, 2008.
- [47] S. Hencl and P. Koskela. *Lectures on mappings of finite distortion*, volume 2096 of *Lecture Notes in Mathematics*. Springer, Cham, 2014.
- [48] J. Horváth. *Topological vector spaces and distributions. Vol. I*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1966.
- [49] A. Jonsson and H. Wallin. Function spaces on subsets of \mathbf{R}^n . *Math. Rep.*, 2(1), 1984.
- [50] J.-P. Kahane. *Some random series of functions*, volume 5 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 1985.
- [51] T. Kamae. A characterization of self-affine functions. *Japan J. Appl. Math.*, 3(2):271–280, 1986.
- [52] J. Kinnunen and E. Saksman. Regularity of the fractional maximal function. *Bull. London Math. Soc.*, 35(4):529–535, 2003.
- [53] N. Kôno. On self-affine functions. *Japan J. Appl. Math.*, 3(2):259–269, 1986.
- [54] L. V. Kovalev and J. Onninen. On invertibility of Sobolev mappings. *J. Reine Angew. Math.*, 656:1–16, 2011.
- [55] L. V. Kovalev, J. Onninen, and K. Rajala. Invertibility of Sobolev mappings under minimal hypotheses. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 27(2):517–528, 2010.
- [56] N. V. Krylov and M. Röckner. Strong solutions of stochastic equations with singular time dependent drift. *Probab. Theory Related Fields*, 131(2):154–196, 2005.
- [57] P. D. Lamberti and V. Vespri. Remarks on Sobolev-Morrey-Campanato spaces defined on $C^{0,\gamma}$ domains. *Eurasian Math. J.*, 10(4):47–62, 2019.
- [58] J. Lamperti. A simple construction of certain diffusion porcesses. *J. Math. Kyoto Univ.*, 4:161–170, 1964.
- [59] N. S. Landkof. *Foundations of modern potential theory*. Springer-Verlag, New York-Heidelberg, 1972. Translated from the Russian by A. P. Doohovskoy, Die Grundlehren der mathematischen Wissenschaften, Band 180.
- [60] J.-F. Le Gall. Applications du temps local aux équations différentielles stochastiques unidimensionnelles. In *Seminar on probability, XVII*, volume 986 of *Lecture Notes in Math.*, pages 15–31. Springer, Berlin, 1983.

- [61] H. Lee and G. Trutnau. Well-posedness for a class of degenerate Itô-SDEs with fully discontinuous coefficients. *Preprint*, pages 1–59, 2019. <https://arXiv.org/abs/1909.09430>.
- [62] J. A. León, D. Nualart, and S. Tindel. Young differential equations with power type nonlinearities. *Stochastic Process. Appl.*, 127(9):3042–3067, 2017.
- [63] G. Leoni and M. Morini. Necessary and sufficient conditions for the chain rule in $W_{\text{loc}}^{1,1}(\mathbb{R}^N; \mathbb{R}^d)$ and $BV_{\text{loc}}(\mathbb{R}^N; \mathbb{R}^d)$. *J. Eur. Math. Soc. (JEMS)*, 9(2):219–252, 2007.
- [64] B. Lins, I. M. Spitkovsky, and S. Zhong. The normalized numerical range and the Davis-Wielandt shell. *Linear Algebra Appl.*, 546:187–209, 2018.
- [65] T. Lyons and Z. Qian. *System control and rough paths*. Oxford Mathematical Monographs. Oxford University Press, Oxford, 2002. Oxford Science Publications.
- [66] T. J. Lyons. Differential equations driven by rough signals. *Rev. Mat. Iberoamericana*, 14(2):215–310, 1998.
- [67] T. J. Lyons, M. Caruana, and T. Lévy. *Differential equations driven by rough paths*, volume 1908 of *Lecture Notes in Mathematics*. Springer, Berlin, 2007. Lectures from the 34th Summer School on Probability Theory held in Saint-Flour, July 6–24, 2004, With an introduction concerning the Summer School by Jean Picard.
- [68] B. Maslowski and D. Nualart. Evolution equations driven by a fractional Brownian motion. *J. Funct. Anal.*, 202(1):277–305, 2003.
- [69] P. Mattila. *Geometry of sets and measures in Euclidean spaces*, volume 44 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1995. Fractals and rectifiability.
- [70] P. Mattila, M. Morán, and J.-M. Rey. Dimension of a measure. *Studia Math.*, 142(3):219–233, 2000.
- [71] S. Nakao. On pathwise uniqueness and comparison of solutions of one-dimensional stochastic differential equations. *Osaka J. Math.*, 20(1):197–204, 1983.
- [72] D. Nualart and Y. Ouknine. Regularization of differential equations by fractional noise. *Stochastic Process. Appl.*, 102(1):103–116, 2002.
- [73] D. Nualart and A. Răşcanu. Differential equations driven by fractional Brownian motion. *Collect. Math.*, 53(1):55–81, 2002.
- [74] Y. G. Reshetnyak. *Space mappings with bounded distortion*, volume 73 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1989. Translated from the Russian by H. H. McFaden.

- [75] D. Revuz and M. Yor. *Continuous martingales and Brownian motion*, volume 293 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, third edition, 1999.
- [76] S. G. Samko, A. A. Kilbas, and O. I. Marichev. *Fractional integrals and derivatives*. Gordon and Breach Science Publishers, Yverdon, 1993.
- [77] L. Schwartz. *Théorie des distributions*. Publications de l'Institut de Mathématique de l'Université de Strasbourg, No. IX-X. Nouvelle édition, entièrement corrigée, refondue et augmentée. Hermann, Paris, 1966.
- [78] T. Sottinen and A. Yazigi. Generalized Gaussian bridges. *Stochastic Process. Appl.*, 124(9):3084–3105, 2014.
- [79] E. M. Stein. *Singular integrals and differentiability properties of functions*. Princeton Mathematical Series, No. 30. Princeton University Press, Princeton, N.J., 1970.
- [80] T. Tao and J. Wright. L^p improving bounds for averages along curves. *J. Amer. Math. Soc.*, 16(3):605–638, 2003.
- [81] S. Torres and L. Viitasaari. Stochastic differential equations with discontinuous coefficients. *Preprint*, pages 1–18, 2019. <https://arXiv.org/abs/1908.03183>.
- [82] H. Triebel. *Theory of function spaces*, volume 78 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1983.
- [83] A. J. Veretennikov. Strong solutions and explicit formulas for solutions of stochastic integral equations. *Mat. Sb. (N.S.)*, 111(153)(3):434–452, 480, 1980. transl. in *Math. USSR Sb.* 39(3):387–403, 1981.
- [84] Y. Xiao. Hölder conditions for the local times and the Hausdorff measure of the level sets of Gaussian random fields. *Probab. Theory Related Fields*, 109(1):129–157, 1997.
- [85] Y. Xiao. Random fractals and Markov processes. In *Fractal geometry and applications: a jubilee of Benoit Mandelbrot, Part 2*, volume 72 of *Proc. Sympos. Pure Math.*, pages 261–338. Amer. Math. Soc., Providence, RI, 2004.
- [86] Y. Xiao. Properties of local-nondeterminism of Gaussian and stable random fields and their applications. *Ann. Fac. Sci. Toulouse Math. (6)*, 15(1):157–193, 2006.
- [87] Y. Yamato. Stochastic differential equations and nilpotent Lie algebras. *Z. Wahrsch. Verw. Gebiete*, 47(2):213–229, 1979.
- [88] P. Yaskov. On pathwise Riemann-Stieltjes integrals. *Statist. Probab. Lett.*, 150:101–107, 2019.
- [89] L. C. Young. An inequality of the Hölder type, connected with Stieltjes integration. *Acta Math.*, 67(1):251–282, 1936.

- [90] M. Zähle. Integration with respect to fractal functions and stochastic calculus. I. *Probab. Theory Related Fields*, 111(3):333–374, 1998.
- [91] M. Zähle. Integration with respect to fractal functions and stochastic calculus. II. *Math. Nachr.*, 225:145–183, 2001.
- [92] M. Zähle and E. Schneider. Forward integrals and SDE with fractal noise. In *Horizons of fractal geometry and complex dimensions*, volume 731 of *Contemp. Math.*, pages 279–302. Amer. Math. Soc., Providence, RI, 2019.
- [93] X. Zhang. Stochastic flows of SDEs with irregular coefficients and stochastic transport equations. *Bull. Sci. Math.*, 134(4):340–378, 2010.
- [94] W. P. Ziemer. *Weakly differentiable functions*, volume 120 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1989. Sobolev spaces and functions of bounded variation.