

MEAN FIELD LIMITS OF PARTICLE-BASED STOCHASTIC REACTION-DIFFUSION MODELS

S. ISAACSON, J. MA AND K. SPILIOPOULOS

ABSTRACT. Particle-based stochastic reaction-diffusion (PBSRD) models are a popular approach for studying biological systems involving both noise in the reaction process and diffusive transport. In this work we derive coarse-grained deterministic partial integro-differential equation (PIDE) models that provide a mean field approximation to the volume reactivity PBSRD model, a model commonly used for studying cellular processes. We formulate a weak measure-valued stochastic process (MVSP) representation for the volume reactivity PBSRD model, demonstrating for a simplified but representative system that it is consistent with the commonly used Doi Fock Space representation of the corresponding forward equation. We then prove the convergence of the general volume reactivity model MVSP to the mean field PIDEs in the large-population (i.e. thermodynamic) limit.

1. INTRODUCTION

The dynamics of many biological processes rely on an interplay between spatial transport and chemical reaction. At the scale of a single cell, experiments have demonstrated that many such processes have stochastic dynamics. Particle-based stochastic reaction-diffusion (PBSRD) models are a widely used approach for studying such processes, explicitly modeling the diffusion of, and reactions between, individual molecules. PBSRD models are appropriate for studying chemical systems in cells containing up to hundreds of thousands to millions of molecules, over timescales of days. They are more macroscopic descriptions than millisecond-timescale quantum mechanical or molecular dynamics models of a few molecules [S09], but more microscopic descriptions than deterministic 3D reaction-diffusion PDEs for the average concentration of each species of molecule. One PBSRD model that has been widely used to study biological processes is the volume reactivity (VR) model of Doi [TS67, D76a, D76b]. In this model positions of *individual molecules* are typically represented as points undergoing Brownian motion. Bimolecular reactions between two reactant molecules occur with a probability per unit time based on their current positions [D76a, D76b]. Unimolecular reactions are typically assumed to represent internal processes, and as such are modeled as occurring with exponentially distributed times based on a specified reaction-rate constant.

Due to their mathematical complexity and high dimensionality, PBSRD models are almost entirely studied by Monte Carlo simulation approximating the underlying stochastic process of molecules diffusing and reacting. The computational expense of such methods can greatly limit the size of chemical systems (in each of number of molecules, number of reactions, or physical domain size) that can be studied. One approach to overcoming this challenge is to use more coarse-grained mathematical models that accurately capture the dynamics of the underlying PBSRD model in appropriate physical regimes. Deterministic and stochastic partial differential equation (PDE/SPDE) models are often postulated as coarse-grainings of PBSRD models in certain large-population or thermodynamic limits where the population size becomes unbounded but species concentrations are held fixed. However, for the PBSRD models commonly used in biological modeling, e.g. the VR model and the contact reactivity model, there is limited *rigorous* work identifying and proving the existence of such deterministic coarse-grained limits (i.e. law of large numbers).

To facilitate the development of rigorous coarse-grainings of PBSRD models, our work begins with formulating the dynamics of the diffusing and reacting molecules as measure-valued stochastic processes (MVSPs). These processes describe the evolution of the concentration fields of each chemical species as a sum of δ functions in each molecule's position. A weak formulation of the dynamics of these processes is then derived,

¹Boston University, Department of Mathematics and Statistics, 111 Cummington Mall, Boston, 02215. E-mails: isaacson@math.bu.edu, majw@bu.edu, kspiliop@math.bu.edu

Date: March 27, 2020.

K.S. was partially supported by the National Science Foundation DMS-1550918. J. M. and S. A. I. were partially supported by National Science Foundation DMS-1902854 and DMS-0920886.

giving the action of the processes on an arbitrary test function. The subsequent equations for the time-evolution of the pairing between a test function and the MVSP then involve both continuous noise processes that account for the diffusion of individual molecules, and state-dependent Poisson-random measures that encode the timing and occurrence of chemical reactions between molecules. We establish in a simplified, but representative, case that the MVSP is equivalent to the commonly used Doi Fock Space representation for the forward equation of the VR model.

We then investigate the large population limit of the MVSP dynamics in which the initial number of molecules of each chemical species becomes unbounded, but the concentrations of each species are held fixed. The latter can be achieved by considering molar concentrations, and treating Avogadro's number and/or the domain volume as a large "system size" parameter. As we work in freespace, here we consider the limit where Avogadro's number can be considered a large parameter. Such limits are typically considered one of the primary physical regimes in which PDE or SPDE models for biological systems arise as physical approximations to the underlying process of molecules diffusing and reacting [AT80, LLN19, HCDWS19].

To rigorously determine the limit of the VR model, we will generalize the martingale problem approach for studying solutions to stochastic differential equations developed by Stroock and Varadhan [EK86, SV06] to our weak MVSP representation. Adaptations of this method have been successfully used to study large-population limits in stochastic models for population dynamics, evolutionary dynamics, interacting particle systems, and financial models [GSS13, GSSS15, DH96, DH09, DIRT15, IT15, MS02, SCS88]. We first identify a macroscopic system of partial integro-differential equations (PIDEs) whose solution corresponds to the large population limit of the MVSP, and then rigorously prove the convergence (in a weak sense) of the MVSP to this solution.

Our approach is unique in using a bottom-up hierarchy to rigorously derive from spatial PBSRD models *new* macroscopic PIDEs that correspond to the true large-population limit, and which correctly account for chemical interactions between particles. This is in contrast to the standard macroscopic reaction-diffusion PDE models of chemical reaction systems used at the cellular scale [NTS08, MNKS09]. The latter are typically obtained by formally modifying reaction-rate equation ODE models for *non-spatial* chemical reaction systems by simply adding Laplacian terms to model molecular diffusion.

To illustrate our main result, consider the special case of the reversible reaction $A + B \rightleftharpoons C$ reaction. Let γ denote a system size parameter (i.e. Avogadro's number, or in bounded domains the product of Avogadro's number and the domain volume). We assume all molecules move by Brownian Motion in \mathbb{R}^d , with species-dependent diffusivities, D^A , D^B and D^C respectively. Let $K_1^\gamma(x, y) = K_1(x, y)/\gamma$ denote the probability per time an individual A molecule at x and B molecule at y can react, with $m_1(z|x, y)$ giving the probability density that when the A and B molecules react they produce a C molecule at z . We define $K_2^\gamma(z) = K_2(z)$ and $m_2(x, y|z)$ similarly for the reverse reaction. Finally, denote by $A(t)$ the stochastic process for the number of species A molecules at time t , and label the position of i th molecule of species A at time t by the stochastic process $\mathbf{Q}_i^{A(t)} \subset \mathbb{R}^d$. The random measure

$$A^\gamma(x, t) = \frac{1}{\gamma} \sum_{i=1}^{A(t)} \delta(x - \mathbf{Q}_i^{A(t)})$$

corresponds to the stochastic process for the the molar concentration of species A at x at time t . We can similarly define $B^\gamma(x, t)$ and $C^\gamma(x, t)$. In this work we study the large population (thermodynamic) limit where $\gamma \rightarrow \infty$ and $A^\gamma(x, 0)$ converges to a well defined limiting molar concentration field (with similar limits for the molar concentrations of species B and C). We prove, in a weak sense, that as $\gamma \rightarrow \infty$,

$$(A^\gamma(x, t), B^\gamma(x, t), C^\gamma(x, t)) \rightarrow (\bar{A}(x, t), \bar{B}(x, t), \bar{C}(x, t)),$$

where

$$\begin{aligned} \partial_t \bar{A}(x, t) &= D^A \Delta \bar{A}(x, t) - \int_{\mathbb{R}^d} K_1(x, y) \bar{A}(x, t) \bar{B}(y, t) dy + \int_{\mathbb{R}^{2d}} K_2(z) m_2(x, y|z) C(z, t) dy dz. \\ \partial_t \bar{B}(x, t) &= D^B \Delta \bar{B}(x, t) - \int_{\mathbb{R}^d} K_1(x, y) \bar{A}(y, t) \bar{B}(x, t) dy + \int_{\mathbb{R}^{2d}} K_2(z) m_2(x, y|z) C(z, t) dy dz. \\ \partial_t \bar{C}(x, t) &= D^C \Delta \bar{C}(x, t) - K_2(z) C(z, t) + \int_{\mathbb{R}^{2d}} K_1(x, y) m_1(z|x, y) \bar{A}(x, t) \bar{B}(y, t) dx dy. \end{aligned}$$

Our main result, Theorem 5.3, establishes this rigorous limit for the VR PBSRD model of general chemical reaction systems involving first and second order reactions. To simplify the (already detailed) exposition, we impose one constraint, assuming that the reaction network structure is such that the total concentration of molecules in the system has a strict upper bound. Theorem 5.3 therefore does not cover reaction systems containing reactions that can lead to unbounded population growth, ruling out creation reactions like $\emptyset \rightarrow A$ and $A \rightarrow 2A$. We note, however, that our basic approach should still be adaptable to such systems, but would require the introduction of a stopping time for when the total population of molecules reaches some threshold. This is similar to one approach used for proving the classical large-population limit of *non-spatial* stochastic chemical kinetic systems [DK15].

Upon completion of this work we became aware of the recent arXiv preprint [LLN19]. In [LLN19] the authors study the rigorous large-population limit for a subset of the reactions we allow in our VR PBSRD model, restricting to reactions of the form $A + B \rightarrow C + D$. This ensured that the total number of particles was always preserved in their system, effectively decoupling the equations for particle diffusion and particle-particle reactions. The benefit to this approach is that the authors were then able to work with a strong-form pathwise representation for the reaction and diffusion of particles, enabling them to use relative entropy methods to prove propagation of chaos. As far as we are aware, however, such a decoupling does not appear to be possible for more general reaction networks that are needed to model many biological processes, where particle numbers can change. As such, while [LLN19] could leverage relative entropy methods and large deviations estimates to establish their results, we will work with the general martingale problem formulation. The large-population limit was also studied in [O89] for PBSRD systems involving only birth reactions ($A \rightarrow 2A$), death reactions ($A \rightarrow \emptyset$), and first-order conversion reactions ($A \rightarrow B$). Our high level formulation for the system dynamics more closely follows [O89], which also studied a weak MVSP representation for the reaction and diffusion of particles. We stress, however, that a key difference in our work is in allowing for general second order reactions, i.e. reactions of the form $A + B \rightarrow \dots$, where the reaction dynamics critically depend on spatial interactions between two individual reactant particles. Such reactions are prevalent in most cellular signaling processes, and common in many chemical network models for biological systems.

The paper is organized as follows. In Section 2 we describe the problem in more mathematical terms, introducing basic notation for specifying chemical reaction systems, for describing system state as a MVSP for the (number) density of particles in the system, and for representing reactant (product) configuration spaces that encode possible positions of individual reactant (product) particles involved in a reaction. In Section 3 we define reaction kernels specifying the probability per time a reaction involving specific reactants can occur. For each reaction type we also specify placement measures. These give the probability density that product particles of a reaction between one or more reactants are placed at specific positions. We then introduce the stochastic equation describing the evolution of the empirical measure (MVSP) of the chemical species in path space. In Section 4 we summarize the basic assumptions we make about the form of the reaction rate functions and product placement measures. In Section 5 we present our main result, Theorem 5.3, describing for general reaction networks the evolution equation satisfied in the large-population limit by the empirical measures for the molar concentration of each species. We also present a number of illustrative examples showing the derived large-population limit for specific chemical systems. In Section 6 we prove that the MVSP formulation we study is equivalent to the more commonly used Fock-space (i.e. Kolmogorov forward equation) representation popularized by Doi [D76a, D76b], focusing on the simplified case of the reversible $A + B \rightleftharpoons C$ reaction. Finally, in Section 7 we give the proof of Theorem 5.3. The appendix includes proofs of a number of technical estimates as well as the existence, uniqueness and regularity statement for the forward Kolmogorov equation of the $A + B \rightleftharpoons C$ reaction system studied in Section 6.

2. NOTATION AND PRELIMINARY DEFINITIONS

We consider a collection of particles with J possible different types. Note, in the following we will interchangeably use particle or molecule and type or species. Let $S = \{S_1, \dots, S_J\}$ denote the set of different possible particle types, with $p_i \in S$ the value of the type of the i -th particle. In the remainder, we also assume an underlying probability triple, $(\Omega, \mathcal{F}, \mathbb{P})$, on which all random variables are defined.

The goal of this paper is to study the process that molecules diffuse in space \mathbb{R}^d freely and undergo at most L possible different type of reactions, denoted as $\mathcal{R}_1, \dots, \mathcal{R}_L$. We describe the \mathcal{R}_ℓ th reaction, $\ell \in \{1, \dots, L\}$,

by

$$\sum_{j=1}^J \alpha_{\ell j} S_j \rightarrow \sum_{j=1}^J \beta_{\ell j} S_j,$$

where we assume the stoichiometric coefficients $\{\alpha_{\ell j}\}_{j=1}^J$ and $\{\beta_{\ell j}\}_{j=1}^J$ are non-negative integers. Let $\boldsymbol{\alpha}^{(\ell)} = (\alpha_{\ell 1}, \alpha_{\ell 2}, \dots, \alpha_{\ell J})$ and $\boldsymbol{\beta}^{(\ell)} = (\beta_{\ell 1}, \beta_{\ell 2}, \dots, \beta_{\ell J})$ be multi-index vectors collecting the coefficients of the ℓ th reaction. We denote the reactant and product orders of the reaction by $|\boldsymbol{\alpha}^{(\ell)}| \doteq \sum_{i=1}^J \alpha_{\ell i} \leq 2$ and $|\boldsymbol{\beta}^{(\ell)}| \doteq \sum_{j=1}^J \beta_{\ell j} \leq 2$, assuming that at most two reactants and two products participate in any reaction. We therefore implicitly assume all reactions are at most second order. For subsequent notational purposes, we order the reactions such that the first \tilde{L} reactions correspond to those that have no products, i.e. annihilation reactions of the form

$$\sum_{j=1}^J \alpha_{\ell j} S_j \rightarrow \emptyset,$$

for $\ell \in \{1, \dots, \tilde{L}\}$. We assume the remaining $L - \tilde{L}$ reactions have one or more product particles.

Let D^i label the diffusion coefficient for the i th molecule, taking values in $\{D_1, \dots, D_J\}$, where D_j is the diffusion coefficient for species S_j , $j = 1, \dots, J$. We denote by $Q_t^i \in \mathbb{R}^d$ the position of the i th molecule, $i \in \mathbb{N}_+$, at time t . A particle's state can be represented as a vector in $\hat{P} = \mathbb{R}^d \times \mathcal{S}$, the combined space encoding particle position and type. This state vector is subsequently denoted by $\hat{Q}_t^i \stackrel{\text{def}}{=} (Q_t^i, p_i)$.

We now formulate our representation for the (number) concentration, equivalently number density, fields of each species. Let E be a complete metric space and $M(E)$ the collection of measures on E . For $f : E \mapsto \mathbb{R}$ and $\mu \in M(E)$, define

$$\langle f, \mu \rangle_E = \int_{x \in E} f(x) \mu(dx).$$

We will frequently have $E = \mathbb{R}^d$. In this case we omit the subscript E and simply write $\langle f, \mu \rangle$. For each $t \geq 0$, we define the concentration of particles in the system at time t by the distribution

$$\nu_t = \sum_{i=1}^{N(t)} \delta_{\hat{Q}_t^i} = \sum_{i=1}^{N(t)} \delta_{Q_t^i} \delta_{p_i},$$

where $N(t) = \langle 1, \nu_t \rangle_{\hat{P}}$ represents the stochastic process for the total number of particles at time t . To investigate the behavior of different type of particles, we denote the marginal distribution on the j th type, i.e. the concentration field for species j , by

$$\nu_t^j(\cdot) = \nu_t(\cdot \times \{S_j\}),$$

a distribution on \mathbb{R}^d . $N_j(t) = \langle 1, \nu_t^j \rangle$ will similarly label the total number of particles of type S_j at time t .

In addition to having notations for representing particle concentration fields, we will also often make use of state vectors for *all* particles in the system. Denote by

$$(2.1) \quad H(\nu_t) = \left((Q_t^{\sigma_1(1)}, S_1), \dots, (Q_t^{\sigma_1(N_1(t))}, S_1), \dots, (Q_t^{\sigma_J(1)}, S_J), \dots, (Q_t^{\sigma_J(N_J(t))}, S_J), 0, 0, \dots \right)$$

a state vector of the full particle system. Here, for each type $j = 1, \dots, J$, the particle index maps $\{\sigma_j(k)\}_{k=1}^{N_j(t)}$ encode an ordering for particles of species j , $Q^{\sigma_j(1)} \preceq \dots \preceq Q^{\sigma_j(N_j(t))}$, arising from an (assumed) underlying ordering on \mathbb{R}^d . $H^i(\nu_t) \in \hat{P}$ will label the i th entry of the vector $H(\nu_t)$. We denote by

$$(2.2) \quad H_Q(\nu_t^j) = \left(Q_t^{\sigma_j(1)}, \dots, Q_t^{\sigma_j(N_j(t))}, 0, 0, \dots \right)$$

an analogous *position-only* state vector for type j particles, with $H_Q^i(\nu_t^j) \in \mathbb{R}^d$ labeling the i th entry in $H_Q(\nu_t^j)$.

With the preceding definitions, we last introduce a system of notation to encode reactant and particle positions and configurations that are needed to later specify reaction processes.

Definition 2.1. For reaction \mathcal{R}_ℓ , define the reactant index space

$$\mathbb{I}^{(\ell)} = \{\mathbf{i} = (i_1^{(1)}, \dots, i_{\alpha_{\ell 1}}^{(1)}, \dots, i_1^{(J)}, \dots, i_{\alpha_{\ell J}}^{(J)}) \mid i_r^{(j)} \in \mathbb{N} \setminus \{0\}, \text{ for all } 1 \leq j \leq J, 1 \leq r \leq \alpha_{\ell j}\} = (\mathbb{N} \setminus \{0\})^{|\alpha^{(\ell)}|}.$$

In describing the dynamics of ν_t , we will sample vectors containing the indices of the specific reactant particles participating in a single ℓ -type reaction from $\mathbb{I}^{(\ell)}$. For $\mathbf{i} \in \mathbb{I}^{(\ell)}$ a particular sampled set of reactant indices, $i_r^{(j)} \in \mathbf{i}$ will label the r th sampled index of species type j . Note that when $\alpha_{\ell j} = 0$ species j is not a reactant for the ℓ th reaction, and hence there will be no indices for particles of species j within any $\mathbf{i} \in \mathbb{I}^{(\ell)}$.

Definition 2.2. For reaction \mathcal{R}_ℓ , similar to our definition of $\mathbb{I}^{(\ell)}$, we define the reactant position space

$$\mathbb{X}^{(\ell)} = \{\mathbf{x} = (x_1^{(1)}, \dots, x_{\alpha_{\ell 1}}^{(1)}, \dots, x_1^{(J)}, \dots, x_{\alpha_{\ell J}}^{(J)}) \mid x_r^{(j)} \in \mathbb{R}^d, \text{ for all } 1 \leq j \leq J, 1 \leq r \leq \alpha_{\ell j}\} = (\mathbb{R}^d)^{|\alpha^{(\ell)}|}.$$

For $\mathbf{x} \in \mathbb{X}^{(\ell)}$ a sampled reactant position configuration for one individual \mathcal{R}_ℓ reaction, $x_r^{(j)}$ then labels the sampled position for the r th reactant particle of species j involved in the reaction. Let $d\mathbf{x} = \left(\bigwedge_{j=1}^J \left(\bigwedge_{r=1}^{\alpha_{\ell j}} dx_r^{(j)} \right) \right)$ be the corresponding volume form on $\mathbb{X}^{(\ell)}$, which also naturally defines an associated Lebesgue measure.

Definition 2.3. For reaction \mathcal{R}_ℓ with $\tilde{L} + 1 \leq \ell \leq L$, i.e. having at least one product particle, define the product position space

$$\mathbb{Y}^{(\ell)} = \{\mathbf{y} = (y_1^{(1)}, \dots, y_{\beta_{\ell 1}}^{(1)}, \dots, y_1^{(J)}, \dots, y_{\beta_{\ell J}}^{(J)}) \mid y_r^{(j)} \in \mathbb{R}^d, \text{ for all } 1 \leq j \leq J, 1 \leq r \leq \beta_{\ell j}\} = (\mathbb{R}^d)^{|\beta^{(\ell)}|}.$$

Analogous to Definition 2.1, when $\beta_{\ell j} = 0$ species j is not a product for the ℓ th reaction, and hence there will be no indices for particles of species j within the product position space. For $\mathbf{y} \in \mathbb{Y}^{(\ell)}$ a sampled product position configuration for one individual \mathcal{R}_ℓ reaction, $y_r^{(j)}$ then labels the sampled position for the r th product particle of species j involved in the reaction. Let $d\mathbf{y} = \left(\bigwedge_{j=1}^J \left(\bigwedge_{r=1}^{\beta_{\ell j}} dy_r^{(j)} \right) \right)$ be the corresponding volume form on $\mathbb{Y}^{(\ell)}$, which also naturally defines an associated Lebesgue measure.

Definition 2.4. For reaction \mathcal{R}_ℓ , particle distribution $\nu = \sum_{j=1}^J \nu^j \delta_{S_j} \in M(\hat{P})$, corresponding marginal distribution $\nu^j \in M(\mathbb{R}^d)$ for particles of type j , and $\mathbf{i} \in \mathbb{I}^{(\ell)}$, define the ℓ th projection mapping $\mathcal{P}^{(\ell)} : M(\hat{P}) \times \mathbb{I}^{(\ell)} \rightarrow \mathbb{X}^{(\ell)}$ as

$$\mathcal{P}^{(\ell)}(\nu, \mathbf{i}) = (H_Q^{i_1^{(1)}}(\nu^1), \dots, H_Q^{i_{\alpha_{\ell 1}}^{(1)}}(\nu^1), \dots, H_Q^{i_1^{(J)}}(\nu^J), \dots, H_Q^{i_{\alpha_{\ell J}}^{(J)}}(\nu^J)).$$

When reactants with indices \mathbf{i} in particle distribution ν are chosen to undergo a reaction of type ℓ , $\mathcal{P}^{(\ell)}(\nu, \mathbf{i})$ then gives the vector of the corresponding reactant particles' positions. Note, for simplicity of notation, in the remainder we will sometimes evaluate $\mathcal{P}^{(\ell)}$ with inconsistent particle distributions and index vectors. In all such expressions other terms will be zero so that this inconsistency does not matter.

Definition 2.5. For reaction \mathcal{R}_ℓ , particle distribution $\nu = \sum_{j=1}^J \nu^j \delta_{S_j} \in M(\hat{P})$ and $\mathbf{i} \in \mathbb{I}^{(\ell)}$, define the effective reactant index sampling space $\Omega^{(\ell)}(\nu) \subset \mathbb{I}^{(\ell)}$ as

$$\Omega^{(\ell)}(\nu) = \{\mathbf{i} \in \mathbb{I}^{(\ell)} \mid 1 \leq j \leq J, i_1^{(j)} < \dots < i_{\alpha_{\ell j}}^{(j)} \leq \langle 1, \nu^j \rangle\}.$$

Definition 2.6. For reaction \mathcal{R}_ℓ and particle distribution $\nu = \sum_{j=1}^J \nu^j \delta_{S_j} \in M(\hat{P})$, define the ℓ th reactant measure mapping $\lambda^{(\ell)}[\cdot] : M(\hat{P}) \rightarrow M(\mathbb{X}^{(\ell)})$ evaluated at $\mathbf{x} \in \mathbb{X}^{(\ell)}$ via $\lambda^{(\ell)}[\nu](d\mathbf{x}) = \otimes_{j=1}^J (\otimes_{r=1}^{\alpha_{\ell j}} \nu^j(dx_r^{(j)}))$.

Definition 2.7. For reaction \mathcal{R}_ℓ , define a subspace $\tilde{\mathbb{X}}^{(\ell)} \subset \mathbb{X}^{(\ell)}$ by removing all particle reactant position vectors in $\mathbb{X}^{(\ell)}$ for which two particles of the same species have the same position. That is

$$\tilde{\mathbb{X}}^{(\ell)} = \mathbb{X}^{(\ell)} \setminus \{\mathbf{x} \in \mathbb{X}^{(\ell)} \mid x_r^{(j)} = x_k^{(j)} \text{ for some } 1 \leq j \leq J, 1 \leq k \neq r \leq \alpha_{\ell j}\}.$$

3. GENERATOR AND PROCESS LEVEL DESCRIPTION

Let us consider the time evolution of the process $\nu_t^\zeta = \sum_{i=1}^{N^\zeta(t)} \delta_{\hat{Q}_i}$ which gives the spatial distribution of all particles (i.e. number density or concentration). Here $N^\zeta(t) = \langle 1, \nu_t^\zeta \rangle_{\hat{P}}$ denotes the total number of particles at time t and $\zeta = (\frac{1}{\gamma}, \eta)$ is a two-vector consisting of a scaling parameter, γ , and a displacement range parameter, η . In the large population limit we consider γ plays the role of a system size, and is

considered to be large (e.g. Avogadro's number, or in bounded domains the product of Avogadro's number and the domain volume) [DK15]. On the other hand, η is a regularizing parameter allowing us to be able to consider and rigorously handle delta-function placement measures for reaction products (a common choice in many PBSRD simulation methods). We will further clarify these parameters later on, focusing on the (large-population) limit that $\gamma \rightarrow \infty$ and $\eta \rightarrow 0$ jointly, denoted as $\zeta \rightarrow 0$.

To formulate the process-level model, it is necessary to specify more concretely the reaction process between individual particles. For reaction \mathcal{R}_ℓ , denote by $K_\ell^\gamma(\mathbf{x})$ the rate (i.e. probability per time) that reactant particles with positions $\mathbf{x} \in \mathbb{X}^{(\ell)}$ react. As described in the next section, we assume this rate function has a specific scaling dependence on γ . Let $m_\ell^\eta(\mathbf{y} | \mathbf{x})$ be the placement measure when the reactants at positions $\mathbf{x} \in \mathbb{X}^{(\ell)}$ react and generate products at positions $\mathbf{y} \in \mathbb{Y}^{(\ell)}$. We assume this placement measure depends on the displacement range parameter η .

Stochastic particle dynamics involve both diffusive motion and chemical reactions. In describing particle motion we will make use of $\{W_t^n\}_{n \in \mathbb{N}_+}$, a countable collection of standard independent Brownian motions in \mathbb{R}^d . To describe a reaction \mathcal{R}_ℓ with no products, i.e. $1 \leq \ell \leq \tilde{L}$, we associate with it a Poisson point measure $dN_\ell(s, \mathbf{i}, \theta)$ on $\mathbb{R}_+ \times \mathbb{I}^{(\ell)} \times \mathbb{R}_+$. Here $\mathbf{i} \in \mathbb{I}^{(\ell)}$ gives the sampled reactant configuration, with $i_r^{(j)}$ labeling the r th sampled index of species j . The corresponding intensity measure of dN_ℓ is given by $d\tilde{N}_\ell(s, \mathbf{i}, \theta) = ds \left(\bigwedge_{j=1}^J \left(\bigwedge_{r=1}^{\alpha_{\ell j}} \left(\sum_{k \geq 0} \delta_k(d i_r^{(j)}) \right) \right) \right) d\theta$. Analogously, for each reaction \mathcal{R}_ℓ with products, i.e. $\tilde{L} + 1 \leq \ell \leq L$, we associate with it a Poisson point measure $dN_\ell(s, \mathbf{i}, \mathbf{y}, \theta_1, \theta_2)$ on $\mathbb{R}_+ \times \mathbb{I}^{(\ell)} \times \mathbb{Y}^{(\ell)} \times \mathbb{R}_+ \times \mathbb{R}_+$. Here $\mathbf{i} \in \mathbb{I}^{(\ell)}$ gives the sampled reactant configuration, with $i_r^{(j)}$ labeling the r th sampled index of species j . $\mathbf{y} \in \mathbb{Y}^{(\ell)}$ gives the sampled product configuration, with $y_r^{(j)}$ labeling the sampled position for the r th newly created particle of species j . The corresponding intensity measure is given by $d\tilde{N}_\ell(s, \mathbf{i}, \mathbf{y}, \theta_1, \theta_2) = ds \left(\bigwedge_{j=1}^J \left(\bigwedge_{r=1}^{\alpha_{\ell j}} \left(\sum_{k \geq 0} \delta_k(d i_r^{(j)}) \right) \right) \right) d\mathbf{y} d\theta_1 d\theta_2$.

The existence of the Poisson point measure follows as the intensity measure is σ -finite (see Chapter I - Theorem 8.1 in [NW14] or Corollary 9.7 in [K01]). Let $d\tilde{N}_\ell(s, \mathbf{i}, \mathbf{y}, \theta_1, \theta_2) = dN_\ell(s, \mathbf{i}, \mathbf{y}, \theta_1, \theta_2) - d\tilde{N}_\ell(s, \mathbf{i}, \mathbf{y}, \theta_1, \theta_2)$ be the compensated Poisson measure, for $\tilde{L} + 1 \leq \ell \leq L$. For any measurable set $A \in \mathbb{I}^{(\ell)} \times \mathbb{Y}^{(\ell)} \times \mathbb{R}_+ \times \mathbb{R}_+$, $N_\ell(\cdot, A)$ is a Poisson process and $\tilde{N}_\ell(\cdot, A)$ is a martingale (see Proposition 9.18 in [K01]). Similarly, we can define $d\tilde{N}_\ell(s, \mathbf{i}, \theta) = dN_\ell(s, \mathbf{i}, \theta) - d\tilde{N}_\ell(s, \mathbf{i}, \theta)$, for $1 \leq \ell \leq \tilde{L}$. In this case, given any measurable set $A \in \mathbb{I}^{(\ell)} \times \mathbb{R}_+$, we then have that $N_\ell(\cdot, A)$ is a Poisson process and $\tilde{N}_\ell(\cdot, A)$ is a martingale.

With the preceding definitions, we can formulate a weak representation for the dynamics of ν_t^ζ . We consider test functions denoted by $f \in C_b^2(\hat{P})$, which we define to mean $f(\cdot, S_j) \in C_b^2(\mathbb{R}^d)$ for each j . The time evolution for the process $\langle f, \nu_t^\zeta \rangle_{\hat{P}}$ can then be represented by

$$\begin{aligned}
\langle f, \nu_t^\zeta \rangle_{\hat{P}} &= \langle f, \nu_0^\zeta \rangle_{\hat{P}} + \sum_{i \geq 1} \int_0^t \mathbf{1}_{\{i \leq \langle 1, \nu_{s-}^\zeta \rangle_{\hat{P}}\}} \sqrt{2D^i} \frac{\partial f}{\partial Q} (H^i(\nu_{s-}^\zeta)) dW_s^i + \int_0^t \sum_{i=1}^{\langle 1, \nu_{s-}^\zeta \rangle_{\hat{P}}} D^i \frac{\partial^2 f}{\partial Q^2} (H^i(\nu_{s-}^\zeta)) ds \\
&+ \sum_{\ell=1}^{\tilde{L}} \int_0^t \int_{\mathbb{I}^{(\ell)}} \int_{\mathbb{R}_+} \left(\left\langle f, \nu_{s-}^\zeta - \sum_{j=1}^J \sum_{r=1}^{\alpha_{\ell j}} \delta_{(H_Q^{i_r^{(j)}}(\nu_{s-}^{\zeta, j}), S_j)} \right\rangle_{\hat{P}} - \langle f, \nu_{s-}^\zeta \rangle_{\hat{P}} \right) \\
&\quad \times \mathbf{1}_{\{i \in \Omega^{(\ell)}(\nu_{s-}^\zeta)\}} \times \mathbf{1}_{\{\theta \leq K_\ell^\gamma(\mathcal{P}^{(\ell)}(\nu_{s-}^\zeta, \mathbf{i}))\}} dN_\ell(s, \mathbf{i}, \theta) \\
&+ \sum_{\ell=\tilde{L}+1}^L \int_0^t \int_{\mathbb{I}^{(\ell)}} \int_{\mathbb{Y}^{(\ell)}} \int_{\mathbb{R}_+^2} \left(\left\langle f, \nu_{s-}^\zeta - \sum_{j=1}^J \sum_{r=1}^{\alpha_{\ell j}} \delta_{(H_Q^{i_r^{(j)}}(\nu_{s-}^{\zeta, j}), S_j)} + \sum_{j=1}^J \sum_{r=1}^{\beta_{\ell j}} \delta_{(y_r^{(j)}, S_j)} \right\rangle_{\hat{P}} \right. \\
(3.1) \quad &\left. - \langle f, \nu_{s-}^\zeta \rangle_{\hat{P}} \right) \times \mathbf{1}_{\{i \in \Omega^{(\ell)}(\nu_{s-}^\zeta)\}} \times \mathbf{1}_{\{\theta_1 \leq K_\ell^\gamma(\mathcal{P}^{(\ell)}(\nu_{s-}^\zeta, \mathbf{i}))\}} \times \mathbf{1}_{\{\theta_2 \leq m_\ell^\eta(\mathbf{y} | \mathcal{P}^{(\ell)}(\nu_{s-}^\zeta, \mathbf{i}))\}} dN_\ell(s, \mathbf{i}, \mathbf{y}, \theta_1, \theta_2)
\end{aligned}$$

Formula (3.1) captures the dynamics of our particle system. Each molecule $i \leq \langle 1, \nu_s^\zeta \rangle_{\hat{P}}$ (recall that $\langle 1, \nu_s^\zeta \rangle_{\hat{P}}$ denotes the total number of molecules at time s) diffuses with diffusion coefficient D^i . When the ℓ th reaction happens for $\ell = \tilde{L} + 1, \dots, L$ (and analogously for $\ell = 1, \dots, \tilde{L}$), as directed by the kernel K_ℓ^γ , the system loses reactants particles and gains product particles. The latter are placed according to the placement measure $m_\ell^\eta(\mathbf{y} | \mathcal{P}^{(\ell)}(\nu_{s-}^\zeta, \mathbf{i}))$. As such, the particle labeled by i in (3.1) will change dynamically as reactions occur.

For this reason, particle positions are accessed through the use of the state vectors, H^i and H_Q^i , as is also done in structured population models [BM15]. The Poisson measures N_ℓ are used to sample the times at which reactions occur, which reactant particles react, and where reaction products are placed.

We will subsequently assume that $\langle 1, \nu_s^\zeta \rangle$ is uniformly bounded in time in Assumption 5.1. The stochastic integral with respect to Brownian motion in (3.1) is then a martingale (for a fixed ζ). Taking the expectation, we obtain for the mean that

$$\begin{aligned}
\mathbb{E}[\langle f, \nu_t^\zeta \rangle_{\hat{P}}] &= \mathbb{E}[\langle f, \nu_0^\zeta \rangle_{\hat{P}}] + \mathbb{E}\left[\int_0^t \sum_{i=1}^{\langle 1, \nu_{s-}^\zeta \rangle_{\hat{P}}} D^i \frac{\partial^2 f}{\partial Q^2}(H^i(\nu_{s-}^\zeta)) ds\right] \\
&+ \sum_{\ell=1}^{\tilde{L}} \mathbb{E}\left[\int_0^t \sum_{\mathbf{i} \in \Omega^{(\ell)}(\nu_{s-}^\zeta)} \left(\left\langle f, \nu_{s-}^\zeta - \sum_{j=1}^J \sum_{r=1}^{\alpha_{\ell j}} \delta_{(H_Q^{i_r(j)}(\nu_{s-}^\zeta), S_j)} \right\rangle - \langle f, \nu_{s-}^\zeta \rangle_{\hat{P}} \right) \times K_\ell^\gamma(\mathcal{P}^{(\ell)}(\nu_{s-}^\zeta, \mathbf{i})) ds\right] \\
&+ \sum_{\ell=\tilde{L}+1}^L \mathbb{E}\left[\int_0^t \sum_{\mathbf{i} \in \Omega^{(\ell)}(\nu_{s-}^\zeta)} \int_{\mathbb{Y}^\ell} \left(\left\langle f, \nu_{s-}^\zeta - \sum_{j=1}^J \sum_{r=1}^{\alpha_{\ell j}} \delta_{(H_Q^{i_r(j)}(\nu_{s-}^\zeta), S_j)} + \sum_{j=1}^J \sum_{r=1}^{\beta_{\ell j}} \delta_{(y_r^{(j)}, S_j)} \right\rangle_{\hat{P}} \right. \right. \\
(3.2) \quad &\left. \left. - \langle f, \nu_{s-}^\zeta \rangle_{\hat{P}} \right) \times K_\ell^\gamma(\mathcal{P}^{(\ell)}(\nu_{s-}^\zeta, \mathbf{i})) \times m_\ell^\eta(\mathbf{y} | \mathcal{P}^{(\ell)}(\nu_{s-}^\zeta, \mathbf{i})) d\mathbf{y} ds\right].
\end{aligned}$$

4. ASSUMPTIONS ON REACTION FUNCTIONS AND PLACEMENT MEASURES

In studying the large population limit that $\gamma \rightarrow \infty$, we will constrain our choices of reaction kernels and placement measures through the following assumptions. Special cases of our choices include a variety of kernels and placement measures that are commonly used in modeling and simulation [D76b, EC09, LE11, IS13, IZ18, DYK18].

Assumption 4.1. *We assume that for all $1 \leq \ell \leq L$, the reaction rate kernel $K_\ell(\mathbf{x})$ is uniformly bounded for all $\mathbf{x} \in \mathbb{X}^{(\ell)}$. We denote generic constants that depend on this bound by $C(K)$.*

Assumption 4.2. *We assume that for any $\eta \geq 0$, $\tilde{L} + 1 \leq \ell \leq L$, $\mathbf{y} \in \mathbb{Y}^{(\ell)}$ and $\mathbf{x} \in \mathbb{X}^{(\ell)}$, the placement measure $m_\ell^\eta(\mathbf{y} | \mathbf{x})$ is a probability measure, i.e. $\int_{\mathbb{Y}^{(\ell)}} m_\ell^\eta(\mathbf{y} | \mathbf{x}) d\mathbf{y} = 1$.*

As previously mentioned, we want to allow for placement measures involving delta-functions. To do so in a mathematically rigorous way we introduced the smoothing parameter η , through which we can define a corresponding mollifier in a standard way, as given by Definition (4.1):

Definition 4.1. *For $x \in \mathbb{R}^d$, let $G(x)$ denote a standard positive mollifier and $G_\eta(x) = \eta^{-d}G(x/\eta)$. That is, $G(x)$ is a smooth function on \mathbb{R}^d satisfying the following four requirements*

- (1) $G(x) \geq 0$,
- (2) $G(x)$ is compactly supported in $B(0, 1)$, the unit ball in \mathbb{R}^d ,
- (3) $\int_{\mathbb{R}^d} G(x) dx = 1$,
- (4) $\lim_{\eta \rightarrow 0} G_\eta(x) = \lim_{\eta \rightarrow 0} \eta^{-d}G(x/\eta) = \delta_0(x)$, where $\delta_0(x)$ is the Dirac delta function and the limit is taken in the space of Schwartz distributions.

The allowable forms of the placement measure for each possible reaction are given by Assumptions 4.3-4.6:

Assumption 4.3. *If \mathcal{R}_ℓ is a first order reaction of the form $S_i \rightarrow S_j$, we assume that the placement measure $m_\ell^\eta(y | x)$ takes the mollified form of*

$$m_\ell^\eta(y | x) = G_\eta(y - x),$$

with the distributional limit as $\eta \rightarrow 0$ given by

$$m_\ell(y | x) = \delta_x(y).$$

This describes that the newly created S_j particle is placed at the position of the reactant S_i particle.

Assumption 4.4. *If \mathcal{R}_ℓ is a second order reaction of the form $S_i + S_k \rightarrow S_j$, we assume that the binding placement measure $m_\ell(z | x, y)$ takes the mollified form of*

$$m_\ell^\eta(z | x, y) = \sum_{i=1}^I p_i \times G_\eta(z - (\alpha_i x + (1 - \alpha_i)y)),$$

with the distributional limit as $\eta \rightarrow 0$ given by

$$m_\ell(z | x, y) = \sum_{i=1}^I p_i \times \delta(z - (\alpha_i x + (1 - \alpha_i)y)),$$

where I is a fixed finite integer and $\sum_i p_i = 1$. This describes that the creation of particle S_j is always on the segment connecting the reactant S_i and reactant S_k particles, but allows some random choice of position. A special case would be $I = 2$, $p_i = \frac{1}{2}$, $\alpha_1 = 0$ and $\alpha_2 = 1$, which corresponds to placing the particle randomly at the position of one of the two reactants. One common choice is taking $I = 1$, $p_1 = 1$ and choosing α_1 to be the diffusion weighted center of mass [IZ18].

Assumption 4.5. If \mathcal{R}_ℓ is a second order reaction of the form $S_i + S_k \rightarrow S_j + S_r$, we assume that the placement measure $m_\ell(z, w | x, y)$ takes the mollified form of

$$m_\ell^\eta(z, w | x, y) = p \times G_\eta(x - z) \otimes G_\eta(y - w) + (1 - p) \times G_\eta(x - w) \otimes G_\eta(y - z),$$

with the distributional limit as $\eta \rightarrow 0$ given by

$$m_\ell(z, w | x, y) = p \times \delta_{(x,y)}((z, w)) + (1 - p) \times \delta_{(x,y)}((w, z)).$$

This describes that newly created product S_j and S_r particles are always at the positions of the reactant S_i and S_k particles. p is typically either 0 or 1, depending on the underlying physics of the reaction.

Assumption 4.6. If \mathcal{R}_ℓ is a first order reaction of the form $S_i \rightarrow S_j + S_k$, we assume the unbinding displacement measure is in the mollified form of

$$m_\ell^\eta(x, y | z) = \rho(|x - y|) \sum_{i=1}^I p_i \times G_\eta(z - (\alpha_i x + (1 - \alpha_i)y)),$$

with the distributional limit as $\eta \rightarrow 0$ given by

$$m_\ell(x, y | z) = \rho(|x - y|) \sum_{i=1}^I p_i \times \delta(z - (\alpha_i x + (1 - \alpha_i)y)),$$

with $\sum_i p_i = 1$. Here we assume the relative separation of the product S_j and S_k particles, $|x - y|$, is sampled from the probability density $\rho(|x - y|)$. Their (weighted) center of mass is sampled from the density encoded by the sum of δ functions. Such forms are common for detailed balance preserving reversible bimolecular reactions [IZ18].

We further assume some regularity of the separation placement density, $\rho(r)$, introduced in Assumption 4.6:

Assumption 4.7. For Assumption 4.2 to be true, we'll need

$$\int_{\mathbb{R}^d} \rho(|w|) dw = 1.$$

Since ρ is a probability density and non-negative, this implies the tail estimate

$$\int_{r>R} r^{d-1} \rho(r) dr \leq \varepsilon,$$

which we will use in subsequent calculations.

Finally, to study the large-population limit of the population density measures, we must specify how the reaction kernels depend on the scaling parameter (i.e. system size parameter) γ . Motivated by the classical spatially homogeneous reaction network large-population limit [DK15], we choose

Assumption 4.8. The reaction kernel is assumed to have the explicit γ dependence that

$$K_\ell^\gamma(\mathbf{x}) = \gamma^{1-|\alpha^{(\ell)}|} K_\ell(\mathbf{x})$$

for any $\mathbf{x} \in \mathbb{X}^{(\ell)}$, $1 \leq \ell \leq L$.

When interpreting the scaling parameter γ as Avogadro's number, or in bounded domains as the product of Avogadro's number and the domain volume [DK15], such scalings can be derived by requiring the formal well-mixed (i.e. infinitely fast diffusion) limit of the volume reactivity PBSRD model to match the corresponding classical spatially homogeneous stochastic chemical kinetics model. See Appendix A for an illustrative example of how the chosen scalings arise in this case.

Recall that $|\alpha^{(\ell)}|$ represents the number of reactant particles needed for the ℓ -th reaction. As we assume $|\alpha|^\ell \leq 2$, we obtain three scalings for the three allowable reaction orders:

- $|\alpha^{(\ell)}| = 0$ corresponds to a pure birth reaction. By Assumption 4.8, the scaling is γ ; i.e. a larger system size implies more births. In a well-mixed model this would imply that as γ and the initial number of molecules are increased, we maintain a fixed rate *with units of molar concentration per time* for the birth reaction to occur.
- $|\alpha^{(\ell)}| = 1$ corresponds to a unimolecular reaction. By Assumption 4.8, there's no rescaling as it's linear. We assume the rates of first order reactions are internal processes to particles, and as such independent of the system size.
- $|\alpha^{(\ell)}| = 2$ corresponds to a bimolecular reaction. By Assumption 4.8, the scaling of reaction kernel is γ^{-1} . As the system size increases it is harder for two *individual* reactant particles to encounter each other.

5. MAIN RESULT AND EXAMPLES

We now formulate a weak representation for the time evolution of scaled empirical measures $\mu_t^{\zeta,j} = \frac{1}{\gamma} \nu_t^{\zeta,j}$ with $j = 1, \dots, J$ and $\mu_t^\zeta = \frac{1}{\gamma} \nu_t^\zeta = \sum_{j=1}^J \mu_t^{\zeta,j} \delta_{S_j}$. $\mu_t^{\zeta,j}$ physically corresponds to the molar concentration field for species j at time t .

For a test function $f \in C_b^2(\mathbb{R}^d)$ and for each species $j = 1, \dots, J$, let us define the generator

$$(\mathcal{L}_j f)(x) = D_j \Delta_x f(x).$$

We'll focus on proving the convergence as $\zeta \rightarrow 0$ of the marginal distribution vector

$$\left(\mu_t^{\zeta,1}, \mu_t^{\zeta,2}, \dots, \mu_t^{\zeta,J} \right).$$

We make two final assumptions before stating our main result. First, to simplify the analysis we assume the total molar concentration is bounded as $\zeta \rightarrow 0$:

Assumption 5.1. *We assume that the total (molar) population concentration satisfies $\sum_{j=1}^J \langle 1, \mu_t^{\zeta,j} \rangle \leq C(\mu)$, i.e. is uniformly in time bounded by some constant $C(\mu)$. In the remainder we abuse notation and also denote generic constants that depend on this bound by $C(\mu)$.*

Assumption 5.1 implies that our main result, Theorem 5.3, does not apply to reaction networks including zero order birth reactions (i.e. reactions of the form $\emptyset \rightarrow S_i$). Similarly, reactions of the form $S_i \rightarrow S_i + S_k$ would be excluded since they also allow the possibility of unbounded population growth. One way to overcome this limitation would be to introduce a stopping time for when the total molar population concentration first exceeds $C(\mu)$. Convergence up to this stopping time follows by essentially the same arguments.

Finally, we assume convergence of the initial molar concentrations of each type at $t = 0$ as $\gamma \rightarrow \infty$:

Assumption 5.2. *We assume that the initial distribution $\mu_0^{\zeta,j} \rightarrow \xi_0^j$ weakly as $\zeta \rightarrow 0$, where ξ_0^j is a compactly supported measure, for all $1 \leq j \leq J$.*

We are now ready to state our main result. Let $M_F(\mathbb{R}^d)$ be the space of finite measures endowed with the weak topology and $\mathbb{D}_{M_F(\mathbb{R}^d)}[0, T]$ be the space of cadlag paths with values in $M_F(\mathbb{R}^d)$ endowed with Skorokhod topology.

Theorem 5.3. *(Main result) Given Assumptions 4.1-4.7 for the reaction kernels and placement measures, and scaling Assumptions 4.8, 5.1 and 5.2, the sequence of measure-valued processes $\{(\mu_t^{\zeta,1}, \dots, \mu_t^{\zeta,J})\}_{t \in [0, T]} \in \mathbb{D}_{\otimes_{j=1}^J M_F(\mathbb{R}^d)}([0, T])$ is relatively compact in $\mathbb{D}_{\otimes_{j=1}^J M_F(\mathbb{R}^d)}([0, T])$ for each $j = 1, 2, \dots, J$. It converges in distribution to $\{(\xi_t^1, \dots, \xi_t^J)\}_{t \in [0, T]} \in C_{\otimes_{j=1}^J M_F(\mathbb{R}^d)}([0, T])$ as $\zeta \rightarrow 0$, respectively being with $j = 1, \dots, J$ the*

unique solution to the system

$$\begin{aligned}
\langle f, \xi_t^j \rangle &= \langle f, \xi_0^j \rangle + \int_0^t \langle (\mathcal{L}_j f)(x), \xi_s^j(dx) \rangle ds \\
&\quad - \sum_{\ell=1}^{\tilde{L}} \int_0^t \int_{\tilde{\mathbb{X}}(\ell)} \frac{1}{\alpha^{(\ell)!}} K_\ell(\mathbf{x}) \left(\sum_{r=1}^{\alpha_{\ell j}} f(x_r^{(j)}) \right) \lambda^{(\ell)}[\xi_s](d\mathbf{x}) ds \\
(5.1) \quad &\quad + \sum_{\ell=\tilde{L}+1}^L \int_0^t \int_{\tilde{\mathbb{X}}(\ell)} \frac{1}{\alpha^{(\ell)!}} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}(\ell)} \left(\sum_{r=1}^{\beta_{\ell j}} f(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} - \sum_{r=1}^{\alpha_{\ell j}} f(x_r^{(j)}) \right) \lambda^{(\ell)}[\xi_s](d\mathbf{x}) ds.
\end{aligned}$$

Remark 5.4. Given that weak convergence to a constant implies convergence in probability, we get that Theorem 5.3 actually implies convergence in probability. Namely, for any $\delta > 0$:

$$\lim_{\zeta \rightarrow 0} \mathbb{P} \left[d_{\otimes_{j=1}^J M_F(\mathbb{R}^d)}((\mu^{\zeta,1}, \dots, \mu^{\zeta,J}), (\xi^1, \dots, \xi^J)) \geq \delta \right] = 0,$$

where $d_{\otimes_{j=1}^J M_F(\mathbb{R}^d)}$ is the metric for $\mathbb{D}_{\otimes_{j=1}^J M_F(\mathbb{R}^d)}[0, T]$, see for example Section 3.2 of [CSY20] for an exposition in an analogous situation.

Proof of Theorem 5.3. Let π^ζ be the \mathbb{P} -law of $(\mu^{\zeta,1}, \dots, \mu^{\zeta,J})$, i.e.

$$\pi^\zeta(A) = \mathbb{P}[(\mu^{\zeta,1}, \dots, \mu^{\zeta,J}) \in A],$$

for all $A \in \mathcal{B}(\mathbb{D}_{\otimes_{j=1}^J M_F(\mathbb{R}^d)}([0, T]))$. This means that for all $\zeta \in (0, 1)^2$ we have that $\pi^\zeta \in M(\mathbb{D}_{\otimes_{j=1}^J M_F(\mathbb{R}^d)}([0, T]))$.

By the relative compactness of Theorem 7.7 we get that every subsequence π^{ζ_k} has a further sub-sequence $\pi^{\zeta_{k_m}}$ which converges weakly. Lemma 7.1 says that any limit point π of $\pi^{\zeta_{k_m}}$ is such that $\pi = \delta_{(\xi^1, \dots, \xi^J)}$ where ξ^j satisfies the evolution equation (5.1). Lemma 7.10 proves uniqueness of solutions to Eq (5.1). Therefore, by Prokhorov's theorem π^ζ converges weakly to π , where π is the distribution of (ξ^1, \dots, ξ^J) , the unique solution to (5.1). That is to say that $(\mu^{\zeta,1}, \dots, \mu^{\zeta,J})$ converges in distribution to (ξ^1, \dots, ξ^J) . Lemma 7.6 proves that for each $j = 1, \dots, J$, $\{\xi_t^j\}_{t \in [0, T]} \in C_{M_F(\mathbb{R}^d)}([0, T])$. This concludes the proof of the theorem. \square

Remark 5.5. If the limiting measures $(\xi_t^1(dx), \dots, \xi_t^J(dx))$ have marginal densities (i.e. molar concentrations) $(\rho_1(x, t), \dots, \rho_J(x, t))$, then the marginals must solve the following reaction-diffusion partial integro-differential equations (PIDEs) in a weak sense.

$$\begin{aligned}
\partial_t \rho_j(x, t) &= D_j \Delta_x \rho_j(x, t) - \sum_{\ell=1}^L \left(\frac{1}{\alpha^{(\ell)!}} \sum_{r=1}^{\alpha_{\ell j}} \int_{\tilde{\mathbf{x}} \in \tilde{\mathbb{X}}(\ell)} \delta_x(\tilde{x}_r^{(j)}) K_\ell(\tilde{\mathbf{x}}) \left(\prod_{k=1}^J \prod_{s=1}^{\alpha_{\ell k}} \rho_k(\tilde{x}_s^{(k)}, t) \right) d\tilde{\mathbf{x}} \right) \\
(5.2) \quad &\quad + \sum_{\ell=\tilde{L}+1}^L \left(\frac{1}{\alpha^{(\ell)!}} \sum_{r=1}^{\beta_{\ell j}} \int_{\tilde{\mathbf{x}} \in \tilde{\mathbb{X}}(\ell)} K_\ell(\tilde{\mathbf{x}}) \left(\int_{\mathbf{y} \in \mathbb{Y}(\ell)} \delta_x(y_r^{(j)}) m_\ell(\mathbf{y} | \tilde{\mathbf{x}}) d\mathbf{y} \right) \left(\prod_{k=1}^J \prod_{s=1}^{\alpha_{\ell k}} \rho_k(\tilde{x}_s^{(k)}, t) \right) d\tilde{\mathbf{x}} \right).
\end{aligned}$$

To illustrate our main result, we now present a few examples to illustrate the limiting PIDEs for basic reaction types:

Example 5.6. A system with birth and death reactions for one species, A . Let \mathcal{R}_1 be the death reaction $A \rightarrow \emptyset$ with probability per time $K_1^\gamma(x)$ to happen for a particle at x . Since \mathcal{R}_1 involves only one reactant and one species, $\alpha_{11} = 1$. As there are no products, $\beta_{11} = 0$. Let \mathcal{R}_2 be the birth reaction $\emptyset \rightarrow A$ with constant probability per time K_2^γ to happen. When one birth event occurs, the position of the new A particle is sampled from the placement measure $m_2(x)$. For \mathcal{R}_2 there are no reactants, so $\alpha_{21} = 0$. One product A particle is generated, so $\beta_{21} = 1$. There are two types of reactions in total, $L = 2$, but reaction \mathcal{R}_1 has no products so $\tilde{L} = 1$.

Let the spatial number distribution for particle A at time t be $\nu_t^{\zeta,1} \in M(\mathbb{R}^d)$, with $\nu_t^\zeta = \nu_t^{\zeta,1} \delta_{S_1} \in M(\hat{P})$. In this example, we would have $\lambda^{(1)}[\nu_t^\zeta](dx) = \nu_t^{\zeta,1}(dx)$. By (3.1) ν_t^ζ satisfies

$$\langle f, \nu_t^\zeta \rangle_{\hat{P}} = \langle f, \nu_0^\zeta \rangle_{\hat{P}} + \sum_{i \geq 1} \int_0^t \mathbf{1}_{\{i \leq \langle 1, \nu_{s-}^\zeta \rangle_{\hat{P}}\}} \sqrt{2D^i} \frac{\partial f}{\partial Q} (H^i(\nu_{s-}^\zeta)) dW_s^i + \int_0^t \sum_{i=1}^{\langle 1, \nu_{s-}^\zeta \rangle_{\hat{P}}} D^i \frac{\partial^2 f}{\partial Q^2} (H^i(\nu_{s-}^\zeta)) ds$$

$$\begin{aligned}
& - \int_0^t \int_{(\mathbb{N} \setminus \{0\})} \int_{\mathbb{R}_+} \left\langle f, \delta_{(H_Q^i(\nu_{s-}^{\zeta,1}), S_1)} \right\rangle_{\hat{P}} \times \mathbf{1}_{\{i \leq \langle 1, \nu_{s-}^{\zeta,1} \rangle\}} \times \mathbf{1}_{\{\theta \leq K_1^\gamma(H_Q^i(\nu_{s-}^{\zeta,1}))\}} dN_1(s, i, \theta) \\
(5.3) \quad & + \int_0^t \int_{\mathbb{R}^d} \int_{\mathbb{R}_+} \int_{\mathbb{R}_+} \left\langle f, \delta_{(x, S_1)} \right\rangle_{\hat{P}} \times \mathbf{1}_{\{\theta_1 \leq K_2^\gamma\}} \times \mathbf{1}_{\{\theta_2 \leq m_2^\eta(x)\}} dN_2(s, x, \theta_1, \theta_2).
\end{aligned}$$

If the limiting spatial distributed measure for species A has marginal density $\rho(x, t)$, by Remark 5.5 it must solve the following reaction-diffusion equation in a weak sense:

$$(5.4) \quad \partial_t \rho(x, t) = D_1 \Delta_x \rho(x, t) - K_1(x) \rho(x, t) + K_2 m_2(x).$$

Remark 5.7. Note that Theorem 5.3 does not apply to reaction networks including zero order birth reactions. However, the above convergence result holds by the same arguments up to a stopping time under the scaling assumptions in Assumption 4.8, the assumption that the placement measure for birth $m_2^\eta(x)$ is in $L^1(\mathbb{R}^d)$, and the assumption that m_2^η does not depend on η , i.e. $m_2^\eta(x) = m_2(x)$. The product particle is then most probably placed within a compact subset of \mathbb{R}^d .

Example 5.8. A system with three species, A , B and C that can undergo the reversible bimolecular reaction $A + B \rightleftharpoons C$. Let \mathcal{R}_1 be the reaction $A + B \rightarrow C$, with $K_1^\gamma(x, y)$ the probability per time one A particle at position x and one B particle at position y bind. Once reaction \mathcal{R}_1 fires, we generate a new particle C at position z following the placement measure $m_1(z|x, y)$. For \mathcal{R}_1 , the reactants are particle A and B , so $\alpha_{11} = \alpha_{12} = 1$ and $\alpha_{13} = 0$. The product is particle C , so that $\beta_{11} = \beta_{12} = 0$, while $\beta_{13} = 1$.

Let \mathcal{R}_2 be the reaction $C \rightarrow A + B$, with $K_2^\gamma(z)$ the probability per time one C particle at position z unbinds. Once reaction \mathcal{R}_2 fires, we generate a new A particle at position x and B particle at position y following the placement measure $m_2(x, y|z)$. For \mathcal{R}_2 , the reactant is a C particle, so $\alpha_{21} = \alpha_{22} = 0$ and $\alpha_{23} = 1$. The products are A and B particles, so that $\beta_{21} = \beta_{22} = 1$, while $\beta_{23} = 0$.

Let the spatial number distribution for A particles at time t be $\nu_t^{\zeta,1} \in M(\mathbb{R}^d)$, the spatial number distribution for B particles at time t be $\nu_t^{\zeta,2} \in M(\mathbb{R}^d)$ and the spatial number distribution for C particles at time t be $\nu_t^{\zeta,3} \in M(\mathbb{R}^d)$. Then $\nu_t^\zeta = \nu_t^{\zeta,1} \delta_{S_1} + \nu_t^{\zeta,2} \delta_{S_2} + \nu_t^{\zeta,3} \delta_{S_3} \in M(\hat{P})$. We have that $\lambda^{(1)}[\nu_t^\zeta](d\mathbf{x}) = \nu_t^{\zeta,1}(dx_1^{(1)}) \nu_t^{\zeta,2}(dx_1^{(2)})$ and $\lambda^{(2)}[\nu_t^\zeta](d\mathbf{x}) = \nu_t^{\zeta,3}(dx_1^{(3)})$. ν_t^ζ then satisfies

$$\begin{aligned}
\left\langle f, \nu_t^\zeta \right\rangle_{\hat{P}} &= \left\langle f, \nu_0^\zeta \right\rangle_{\hat{P}} + \sum_{i \geq 1} \int_0^t \mathbf{1}_{\{i \leq \langle 1, \nu_{s-}^\zeta \rangle_{\hat{P}}\}} \sqrt{2D^i} \frac{\partial f}{\partial Q} (H^i(\nu_{s-}^\zeta)) dW_s^i + \int_0^t \sum_{i=1}^{\langle 1, \nu_{s-}^\zeta \rangle_{\hat{P}}} D^i \frac{\partial^2 f}{\partial Q^2} (H^i(\nu_{s-}^\zeta)) ds \\
&+ \int_0^t \int_{(\mathbb{N} \setminus \{0\})^2} \int_{\mathbb{R}^d} \int_{\mathbb{R}_+} \int_{\mathbb{R}_+} \left\langle f, -\delta_{(H_Q^i(\nu_{s-}^{\zeta,1}), S_1)} - \delta_{(H_Q^j(\nu_{s-}^{\zeta,2}), S_2)} + \delta_{(z, S_3)} \right\rangle_{\hat{P}} \times \mathbf{1}_{\{i \leq \langle 1, \nu_{s-}^{\zeta,1} \rangle\}} \\
&\quad \times \mathbf{1}_{\{j \leq \langle 1, \nu_{s-}^{\zeta,2} \rangle\}} \times \mathbf{1}_{\{\theta_1 \leq K_1^\gamma(H_Q^i(\nu_{s-}^{\zeta,1}), H_Q^j(\nu_{s-}^{\zeta,2}))\}} \times \mathbf{1}_{\{\theta_2 \leq m_1^\eta(z|H_Q^i(\nu_{s-}^{\zeta,1}), H_Q^j(\nu_{s-}^{\zeta,2}))\}} dN_1(s, i, j, z, \theta_1, \theta_2) \\
&+ \int_0^t \int_{\mathbb{N} \setminus \{0\}} \int_{\mathbb{R}^d \times \mathbb{R}^d} \int_{\mathbb{R}_+} \int_{\mathbb{R}_+} \left\langle f, \delta_{(x, S_1)} + \delta_{(y, S_2)} - \delta_{(H_Q^k(\nu_{s-}^{\zeta,3}), S_3)} \right\rangle_{\hat{P}} \times \mathbf{1}_{\{k \leq \langle 1, \nu_{s-}^{\zeta,3} \rangle\}} \\
(5.5) \quad &\quad \times \mathbf{1}_{\{\theta_1 \leq K_2^\gamma(H_Q^k(\nu_{s-}^{\zeta,3}))\}} \times \mathbf{1}_{\{\theta_2 \leq m_2^\eta(x, y|H_Q^k(\nu_{s-}^{\zeta,3}))\}} dN_2(s, k, x, y, \theta_1, \theta_2).
\end{aligned}$$

If the limiting spatially distributed measures for species A , B and C have marginal densities $(\rho_1(x, t), \rho_2(x, t), \rho_3(x, t))$ respectively, by Remark 5.5 they must solve the following reaction-diffusion equations in a weak sense:

$$\begin{aligned}
\partial_t \rho_1(x, t) &= D_1 \Delta_x \rho_1(x, t) - \left(\int_{\mathbb{R}^d} K_1(x, y) \rho_2(y, t) dy \right) \rho_1(x, t) + \int_{\mathbb{R}^d} K_2(z) \left(\int_{\mathbb{R}^d} m_2(x, y|z) dy \right) \rho_3(z, t) dz \\
\partial_t \rho_2(y, t) &= D_2 \Delta_y \rho_2(y, t) - \left(\int_{\mathbb{R}^d} K_1(x, y) \rho_1(x, t) dx \right) \rho_2(y, t) + \int_{\mathbb{R}^d} K_2(z) \left(\int_{\mathbb{R}^d} m_2(x, y|z) dx \right) \rho_3(z, t) dz \\
\partial_t \rho_3(z, t) &= D_3 \Delta_z \rho_3(z, t) - K_2(z) \rho_3(z, t) + \int_{\mathbb{R}^{2d}} K_1(x, y) m_1(z|x, y) \rho_1(x, t) \rho_2(y, t) dx dy.
\end{aligned}$$

(5.6)

Example 5.9. A system with two species, A and B , that can undergo the reversible dimerization reaction $A + A \rightleftharpoons B$. Let \mathcal{R}_1 be the reaction $A + A \rightarrow B$ with $K_1^\gamma(x, y)$ the probability per time one A particle at position x and another A particle at position y bind. Once reaction \mathcal{R}_1 fires, we generate a new B particle

at position z by sampling from the placement measure $m_1(z|x, y)$. For \mathcal{R}_1 , the reactants are two A particles, so $\alpha_{11} = 2$ and $\alpha_{12} = 0$. The product is one B particle, so that $\beta_{11} = 0$ and $\beta_{12} = 1$.

Let \mathcal{R}_2 be the reaction $B \rightarrow A + A$, with $K_2^\gamma(z)$ the probability per time one B particle at position z unbinds. Once reaction \mathcal{R}_2 fires, we generate two new A particles at positions x and y by sampling from the placement measure $m_2(x, y|z)$. For \mathcal{R}_2 , the reactant is one B particle, so $\alpha_{21} = 0$ and $\alpha_{22} = 1$. The products are two A particles, so that $\beta_{21} = 2$ and $\beta_{22} = 0$.

Let the spatial number distribution for A particles at time t be $\nu_t^{\zeta,1} \in M(\mathbb{R}^d)$ and the spatial number distribution for B particles at time t be $\nu_t^{\zeta,2} \in M(\mathbb{R}^d)$. Then $\nu_t^\zeta = \nu_t^{\zeta,1} \delta_{S_1} + \nu_t^{\zeta,2} \delta_{S_2} \in M(\hat{P})$. We have that $\lambda^{(1)}[\nu_t^\zeta](d\mathbf{x}) = \nu_t^{\zeta,1}(dx_1^{(1)}) \nu_t^{\zeta,1}(dx_2^{(1)})$ and $\lambda^{(2)}[\nu_t^\zeta](d\mathbf{x}) = \nu_t^{\zeta,2}(dx_1^{(2)})$. ν_t^ζ then satisfies

$$\begin{aligned}
\langle f, \nu_t^\zeta \rangle_{\hat{P}} &= \langle f, \nu_0^\zeta \rangle_{\hat{P}} + \sum_{i \geq 1} \int_0^t \mathbf{1}_{\{i \leq \langle 1, \nu_{s-}^\zeta \rangle_{\hat{P}}\}} \sqrt{2D^i} \frac{\partial f}{\partial Q} (H^i(\nu_{s-}^\zeta)) dW_s^i + \int_0^t \sum_{i=1}^{\langle 1, \nu_{s-}^\zeta \rangle_{\hat{P}}} D^i \frac{\partial^2 f}{\partial Q^2} (H^i(\nu_{s-}^\zeta)) ds \\
&+ \int_0^t \int_{(\mathbb{N} \setminus \{0\})^2} \int_{\mathbb{R}^d} \int_{\mathbb{R}_+} \int_{\mathbb{R}_+} \langle f, -\delta_{(H_Q^i(\nu_{s-}^{\zeta,1}), S_1)} - \delta_{(H_Q^j(\nu_{s-}^{\zeta,1}), S_1)} + \delta_{(z, S_2)} \rangle_{\hat{P}} \times \mathbf{1}_{\{i < j \leq \langle 1, \nu_{s-}^{\zeta,1} \rangle\}} \\
&\quad \times \mathbf{1}_{\{\theta_1 \leq K_1^\gamma(H_Q^i(\nu_{s-}^{\zeta,1}), H_Q^j(\nu_{s-}^{\zeta,1}))\}} \times \mathbf{1}_{\{\theta_2 \leq m_1^\eta(z | H_Q^i(\nu_{s-}^{\zeta,1}), H_Q^j(\nu_{s-}^{\zeta,1}))\}} dN_1(s, i, j, z, \theta_1, \theta_2) \\
&+ \int_0^t \int_{\mathbb{N} \setminus \{0\}} \int_{\mathbb{R}^d \times \mathbb{R}^d} \int_{\mathbb{R}_+} \int_{\mathbb{R}_+} \langle f, \delta_{(x, S_1)} + \delta_{(y, S_1)} - \delta_{(H_Q^k(\nu_{s-}^{\zeta,2}), S_2)} \rangle_{\hat{P}} \times \mathbf{1}_{\{k \leq \langle 1, \nu_{s-}^{\zeta,2} \rangle\}} \\
(5.7) \quad &\quad \times \mathbf{1}_{\{\theta_1 \leq K_2^\gamma(H_Q^k(\nu_{s-}^{\zeta,2}))\}} \times \mathbf{1}_{\{\theta_2 \leq m_2^\eta(x, y | H_Q^k(\nu_{s-}^{\zeta,2}))\}} dN_2(s, k, x, y, \theta_1, \theta_2).
\end{aligned}$$

If the spatially distributed measures for species A and B have marginal densities $(\rho_1(x, t), \rho_2(z, t))$ respectively, then from Remark 5.5 they must solve the following reaction-diffusion equations in a weak sense:

$$\begin{aligned}
\partial_t \rho_1(x, t) &= D_1 \Delta_x \rho_1(x, t) - \left(\int_{\mathbb{R}^d} K_1(x, y) \rho_1(y, t) dy \right) \rho_1(x, t) + 2 \int_{\mathbb{R}^d} K_2(z) \left(\int_{\mathbb{R}^d} m_2(x, y|z) dy \right) \rho_2(z, t) dz \\
\partial_t \rho_2(z, t) &= D_2 \Delta_z \rho_2(z, t) - K_2(z) \rho_2(z, t) + \frac{1}{2} \int_{\mathbb{R}^{2d}} K_1(x, y) m_1(z|x, y) \rho_1(x, t) \rho_1(y, t) dx dy.
\end{aligned}
\tag{5.8}$$

6. EQUIVALENCE BETWEEN MEASURE VALUED FORMULATION AND FORWARD KOLMOGOROV EQUATION

In this section, we demonstrate equivalence of the measure-valued stochastic process formulation (3.1) to the forward Kolmogorov equation representation of the volume-reactivity model popularized by Doi [D76a, D76b]. For ease of notation, and brevity of presentation, we restrict attention to the special case of the reversible $A + B \rightleftharpoons C$ reaction, i.e. Example 5.8. Though we do not show here the general case, we note that this reversible example includes the key complicating components; two-body particle interactions and changing (total) numbers of particles. It is therefore illustrative of other reactions that may only involve particle creation (e.g. $\emptyset \rightarrow A$), or involve interactions but preserve particle numbers (e.g. $A + B \rightarrow C + D$ or $A \rightarrow B$).

Denote by $K_1^\gamma(x, y)$ the probability per time a particle of type A at x and a particle of type B at y react, and by $K_2^\gamma(z)$ the probability per time a particle of type C at z dissociates. We let $m_1^\eta(z|x, y)$ be the corresponding placement measure for the $A + B \rightarrow C$ reaction, producing a particle of type C at z , given a particle of type A at x and a particle of type B at y . Similarly, $m_2^\eta(x, y|z)$ is the placement measure for the $C \rightarrow A + B$ reaction, producing a particle of type A at x and a particle of type B at y given a particle of type C at z respectively.

6.1. Weak MVSP Formulation for the $A + B \rightleftharpoons C$ Reaction. The weak MVSP representation is given by (5.5). Taking expectation we obtain for the mean

$$\mathbb{E}[\langle f, \nu_t^\zeta \rangle_{\hat{P}}] = \mathbb{E}[\langle f, \nu_0^\zeta \rangle_{\hat{P}}] + \mathbb{E} \left[\int_0^t \sum_{i=1}^{\langle 1, \nu_{s-}^\zeta \rangle_{\hat{P}}} D^i \frac{\partial^2 f}{\partial Q^2} (H^i(\nu_{s-}^\zeta)) ds \right]$$

$$\begin{aligned}
& + \mathbb{E} \left[\int_0^t \sum_{i=1}^{\langle 1, \nu_{s-}^{\zeta,1} \rangle} \sum_{j=1}^{\langle 1, \nu_{s-}^{\zeta,2} \rangle} \int_{\mathbb{R}^d} \left\langle f, -\delta_{(H_Q^i(\nu_{s-}^{\zeta,1}), S_1)} - \delta_{(H_Q^j(\nu_{s-}^{\zeta,2}), S_2)} + \delta_{(z, S_3)} \right\rangle_{\hat{P}} \right. \\
& \quad \times K_1^\gamma \left(H_Q^i(\nu_{s-}^{\zeta,1}), H_Q^j(\nu_{s-}^{\zeta,2}) \right) m_1^\eta \left(z | H_Q^i(\nu_{s-}^{\zeta,1}), H_Q^j(\nu_{s-}^{\zeta,2}) \right) dz ds \Big] \\
& + \mathbb{E} \left[\int_0^t \sum_{k=1}^{\langle 1, \nu_{s-}^{\zeta,3} \rangle} \int_{\mathbb{R}^{2d}} \left\langle f, \delta_{(x, S_1)} + \delta_{(y, S_2)} - \delta_{(H_Q^k(\nu_{s-}^{\zeta,3}), S_3)} \right\rangle_{\hat{P}} \right. \\
& \quad \times K_2^\gamma \left(H_Q^k(\nu_{s-}^{\zeta,3}) \right) m_2^\eta \left(x, y | H_Q^k(\nu_{s-}^{\zeta,3}) \right) dx dy ds \Big] \\
& = \mathbb{E} \left[\left\langle f, \nu_0^\zeta \right\rangle_{\hat{P}} \right] + \mathbb{E} \left[\int_0^t \left\langle \mathcal{L}f, \nu_{s-}^\zeta \right\rangle_{\hat{P}} ds \right] \\
& + \mathbb{E} \left[\int_0^t \int_{\mathbb{R}^{3d}} (-f(x, S_1) - f(y, S_2) + f(z, S_3)) K_1^\gamma(x, y) m_1^\eta(z|x, y) \nu_{s-}^{\zeta,1}(dx) \nu_{s-}^{\zeta,2}(dy) dz ds \right] \\
(6.1) \quad & + \mathbb{E} \left[\int_0^t \int_{\mathbb{R}^{3d}} (f(x, S_1) + f(y, S_2) - f(z, S_3)) K_2^\gamma(z) m_2^\eta(x, y|z) dx dy \nu_{s-}^{\zeta,3}(dz) ds \right].
\end{aligned}$$

In Eq (6.1), we denote $\mathcal{L}f(Q, S_j) = D_j \Delta_Q f$, for all $j = 1, 2, 3$. As we shall demonstrate soon, this is consistent with what we expect from the forward equation (6.2).

6.2. Doi Forward Kolmogorov Equation for the $\mathbf{A} + \mathbf{B} \rightleftharpoons \mathbf{C}$ Reaction. We use a notation consistent with that introduced by Doi [D76a, D76b]. Suppose $A(t)$ is the stochastic process for the number of species A particles in the system at time t , with $B(t)$ and $C(t)$ defined similarly. Values of $A(t)$, $B(t)$ and $C(t)$ will be given by a , b and c (i.e. $A(t) = a$). When $A(t) = a$, we will let $\mathbf{Q}_l^a(t) \in \mathbb{R}^d$ label the stochastic process for the position of the l th molecule of species A. \mathbf{q}_l^a will denote a possible value of $\mathbf{Q}_l^a(t)$. The species A position vector when $A(t) = a$ is then given by

$$\mathbf{Q}^a(t) = (\mathbf{Q}_1^a(t), \dots, \mathbf{Q}_a^a(t)) \in \mathbb{R}^{da}.$$

Similarly, \mathbf{q}^a will denote a possible value of $\mathbf{Q}^a(t)$,

$$\mathbf{Q}^a(t) = \mathbf{q}^a = (\mathbf{q}_1^a, \dots, \mathbf{q}_a^a).$$

$\mathbf{Q}^b(t)$, $\mathbf{Q}_m^b(t)$, $\mathbf{Q}^c(t)$, $\mathbf{Q}_n^c(t)$, \mathbf{q}_m^b , \mathbf{q}_n^c , \mathbf{q}^b and \mathbf{q}^c will all be defined analogously. The state of the system is then a hybrid discrete–continuous state stochastic process given by $(A(t), B(t), C(t), \mathbf{Q}^{A(t)}, \mathbf{Q}^{B(t)}, \mathbf{Q}^{C(t)})$.

With this notation, denote by $p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t)$ the probability density that $A(t) = a$, $B(t) = b$ and $C(t) = c$ with $\mathbf{Q}^a(t) = \mathbf{q}^a$, $\mathbf{Q}^b(t) = \mathbf{q}^b$ and $\mathbf{Q}^c(t) = \mathbf{q}^c$. We assume that particles of the same species are indistinguishable, that is for $1 \leq l < l' \leq a$ fixed

$$\begin{aligned}
& p^{(a,b,c)}(\mathbf{q}_1^a, \dots, \mathbf{q}_{l-1}^a, \mathbf{q}_l^a, \mathbf{q}_{l+1}^a, \dots, \mathbf{q}_{l'-1}^a, \mathbf{q}_{l'}^a, \mathbf{q}_{l'+1}^a, \dots, \mathbf{q}_a^a, \mathbf{q}^b, \mathbf{q}^c, t) \\
& = p^{(a,b,c)}(\mathbf{q}_1^a, \dots, \mathbf{q}_{l-1}^a, \mathbf{q}_{l'}^a, \mathbf{q}_{l+1}^a, \dots, \mathbf{q}_{l'-1}^a, \mathbf{q}_l^a, \mathbf{q}_{l'+1}^a, \dots, \mathbf{q}_a^a, \mathbf{q}^b, \mathbf{q}^c, t),
\end{aligned}$$

with similar relations holding for permutations of the molecule orderings within \mathbf{q}^b and \mathbf{q}^c . With this assumption the $p^{(a,b,c)}$ are chosen to be normalized so that

$$\sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \left[\frac{1}{a! b! c!} \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right] = 1.$$

Here the bracketed term corresponds to the probability of having a given number of each species, i.e.

$$\Pr[A(t) = a, B(t) = b, C(t) = c] = \frac{1}{a! b! c!} \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a.$$

Let $\mathbf{P}(t) = \{p^{(a,b,t)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t)\}_{a,b,c=0}^{\infty}$ denote the vector of all the probabilities. The forward equation (see [IZ18]) is given by the coupled system of PIDEs that

$$(6.2) \quad \partial_t \mathbf{P}(t) = (\mathcal{L} + \mathcal{R}^+ + \mathcal{R}^-) \mathbf{P}(t).$$

Here the linear operators \mathcal{L} , \mathcal{R}^+ and \mathcal{R}^- correspond to diffusion, the forward association reaction and the reverse dissociation reaction respectively. The diffusion operator in the (a, b, c) equation is given by

$$(6.3) \quad (\mathcal{L}P(t))_{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) = \left(D_1 \sum_{l=1}^a \Delta_{\mathbf{q}_l^a} + D_2 \sum_{m=1}^b \Delta_{\mathbf{q}_m^b} + D_3 \sum_{n=1}^c \Delta_{\mathbf{q}_n^c} \right) p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t),$$

where $\Delta_{\mathbf{q}_l^a}$ denotes the d -dimensional Laplacian acting on the \mathbf{q}_l^a coordinate, and $\Delta_{\mathbf{q}_m^b}$ and $\Delta_{\mathbf{q}_n^c}$ are defined similarly. (Recall D_1, D_2, D_3 are the diffusivity of species A, B, and C respectively.) To define the reaction operators, \mathcal{R}^+ and \mathcal{R}^- , we introduce notations for adding or removing a particle from a given state, \mathbf{q}^a . Let

$$\mathbf{q}^a \cup \mathbf{x} = (\mathbf{q}_1^a, \dots, \mathbf{q}_a^a, \mathbf{x}), \quad \mathbf{q}^a \setminus \mathbf{q}_l^a = (\mathbf{q}_1^a, \dots, \mathbf{q}_{l-1}^a, \mathbf{q}_{l+1}^a, \dots, \mathbf{q}_a^a),$$

which correspond to adding a particle to species A at \mathbf{x} , and removing the l th particle of species A respectively. With these definitions, the reaction operator for the $A + B \rightarrow C$ association reaction in the (a, b, c) equation is

$$(6.4) \quad (\mathcal{R}^+P(t))_{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) = - \left(\sum_{l=1}^a \sum_{m=1}^b K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) \right) p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) \\ + \sum_{n=1}^c \left[\int_{\mathbb{R}^{2d}} m_1^\eta(\mathbf{q}_n^c | \mathbf{x}, \mathbf{y}) K_1^\gamma(\mathbf{x}, \mathbf{y}) p^{(a+1, b+1, c-1)}(\mathbf{q}^a \cup \mathbf{x}, \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t) d\mathbf{x} d\mathbf{y} \right],$$

while the reaction operator for the dissociation reaction $C \rightarrow A + B$ in the (a, b, c) equation is

$$(6.5) \quad (\mathcal{R}^-P(t))_{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) = - \left(\sum_{n=1}^c K_2^\gamma(\mathbf{q}_n^c) \right) p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) \\ + \sum_{l=1}^a \sum_{m=1}^b \left[\int_{\mathbb{R}^d} m_2^\eta(\mathbf{q}_l^a, \mathbf{q}_m^b | \mathbf{z}) K_2^\gamma(\mathbf{z}) p^{(a-1, b-1, c+1)}(\mathbf{q}^a \setminus \mathbf{q}_l^a, \mathbf{q}^b \setminus \mathbf{q}_m^b, \mathbf{q}^c \cup \mathbf{z}, t) d\mathbf{z} \right].$$

This representation is consistent with the classical second quantization representation of Doi [D76a, D76b].

Suppose the initial condition $P(0) = P_0 = \{p_0^{(a,b,c)}\}_{a,b,c=0}^\infty$ is fixed, and we have (a_0, b_0, c_0) particles of A, B and C respectively at time zero. We consider the evolution of $P(t)$ as a vector in a L^2 Fock Space $F = L^2(X)$ equipped with inner product defined by (C.1), where

$$X = \bigoplus_{\substack{a_0 \vee b_0 + c_0 \\ a, b, c \geq 0 \\ a+b+2c = a_0 + b_0 + 2c_0}} \mathbb{R}^{d(a+b+c)}.$$

Remark 6.1. For the $A + B \rightleftharpoons C$ reaction, the quantity $A(t) + B(t) + 2C(t) = a_0 + b_0 + 2c_0$ is always conserved. For our example, X is therefore a finite sum of Euclidean spaces over $a, b, c \in \mathbb{N}_+$ such that $a + b + 2c = a_0 + b_0 + 2c_0$.

To simplify the calculation of regularity results for (6.2) for comparison to the forward equation, in this section we make

Assumption 6.2. We assume the reaction kernel function $K(x, y)$ for the $A + B \rightarrow C$ reaction only depends on the separation of two reactant particles, $|x - y|$, denoted as $K(x, y) = \tilde{K}(|x - y|)$. Furthermore, we assume $\tilde{K}(|w|) \in L^2(\mathbb{R}^d)$, $w \in \mathbb{R}^d$.

Assumption 6.3. We assume the function $\rho(|w|)$, $w \in \mathbb{R}^d$, defined in Assumption 4.6 for the $C \rightarrow A + B$ reaction, is in $L^2(\mathbb{R}^d)$.

Under these two assumptions the following regularity theorem holds, for which the proof is given in Appendix D.

Theorem 6.4. Given Assumptions 6.2 and 6.3, there exists a unique global mild solution to (6.2), $P(\cdot) \in \mathcal{C}([0, \infty); H^2(X))$. That is, $P(t)$ satisfies

$$P(t) = e^{t\mathcal{L}}P_0 + \int_0^t e^{(t-s)\mathcal{L}}(\mathcal{R}^+P(s) + \mathcal{R}^-P(s)) ds,$$

with the initial condition $\mathbf{P}(0) = \mathbf{P}_0 \in H^1(X)$. Further, if $p_0^{(a,b,c)} \geq 0$ for each (a,b,c) , and satisfies the normalization condition

$$\sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \left[\frac{1}{a! b! c!} \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} p_0^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right] = 1,$$

then $\mathbf{P}(t)$ is always non-negative for all $t \geq 0$ and the same normalization condition holds,

$$\sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \left[\frac{1}{a! b! c!} \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right] = 1.$$

Note, as $a + b + 2c$ is conserved, see Remark 6.1, the above summation is only over a finite set of indices.

6.3. Equivalence of the two approaches. Now, we are in position to compare the two approaches as described in Sections 6.1 and 6.2. For this purpose we have Proposition 6.5 and Proposition 6.6 whose proofs are deferred to Appendix C.

Proposition 6.5. *For any $M \in \mathbb{N}$, any function $\varphi \in C^\infty(\mathbb{R}^M)$ and any set of functions $\{f_m\}_{m=1}^M \in C_b^2(\hat{P})$, the (derived) evolution equations satisfied by $\mathbb{E}[\varphi(\langle f_1, \nu_t^\xi \rangle_{\hat{P}}, \langle f_2, \nu_t^\xi \rangle_{\hat{P}}, \dots, \langle f_M, \nu_t^\xi \rangle_{\hat{P}})]$ are the same from the weak measure-valued process representation and the forward Kolmogorov equation. This implies that these two approaches produce the same statistics.*

By Proposition 6.5 we know that the measure-valued formulation and the forward Kolmogorov equation yield the same statistics, at least when the statistics involve smooth test functions. We next derive equations for the mean particle spatial field density of different species at time t in Proposition 6.6. Note, Proposition 6.6 can be viewed as a special case of the equations derived in Proposition 6.5 formally (which can be made rigorous after introducing appropriate mollifiers) by choosing $M = 1$, $\varphi = 1$ and $f_1 = \delta_{(x,S_1)}$, $f_1 = \delta_{(y,S_2)}$ or $f_1 = \delta_{(z,S_3)}$ respectively for representing the spatial field density for species A , B , and C . To be rigorous, the proof of Proposition (6.6) derives these equations directly from the forward Kolmogorov equation instead of by generalizing Proposition 6.5.

Proposition 6.6. *Let $A(\mathbf{x}, t)$, $B(\mathbf{y}, t)$ and $C(\mathbf{z}, t)$ denote the spatial field number density in the particle model at time t for species A at position x , B at position y and C at position z respectively. Under the assumptions of Theorem 6.4, the evolution equations for their expectations at time t satisfy*

$$\begin{aligned} \partial_t \mathbb{E}[A(\mathbf{x}, t)] &= D_1 \Delta_{\mathbf{x}} \mathbb{E}[A(\mathbf{x}, t)] - \int_{\mathbb{R}^d} K_1^\gamma(\mathbf{x}, \mathbf{y}) \mathbb{E}[A(\mathbf{x}, t) B(\mathbf{y}, t)] d\mathbf{y} \\ &\quad + \int_{\mathbb{R}^d} \left[\int_{\mathbb{R}^d} m_2(\mathbf{x}, \mathbf{y} | \mathbf{z}) K_2^\gamma(\mathbf{z}) d\mathbf{y} \right] \mathbb{E}[C(\mathbf{z}, t)] d\mathbf{z}, \\ \partial_t \mathbb{E}[B(\mathbf{y}, t)] &= D_2 \Delta_{\mathbf{y}} \mathbb{E}[B(\mathbf{y}, t)] - \int_{\mathbb{R}^d} K_1^\gamma(\mathbf{x}, \mathbf{y}) \mathbb{E}[A(\mathbf{x}, t) B(\mathbf{y}, t)] d\mathbf{x} \\ &\quad + \int_{\mathbb{R}^d} \left[\int_{\mathbb{R}^d} m_2(\mathbf{x}, \mathbf{y} | \mathbf{z}) K_2^\gamma(\mathbf{z}) d\mathbf{x} \right] \mathbb{E}[C(\mathbf{z}, t)] d\mathbf{z}, \\ \partial_t \mathbb{E}[C(\mathbf{z}, t)] &= D_3 \Delta_{\mathbf{z}} \mathbb{E}[C(\mathbf{z}, t)] - K_2^\gamma(\mathbf{z}) \mathbb{E}[C(\mathbf{z}, t)] + \int_{\mathbb{R}^{2d}} m_1(\mathbf{z} | \mathbf{x}, \mathbf{y}) K_1^\gamma(\mathbf{x}, \mathbf{y}) \mathbb{E}[A(\mathbf{x}, t) B(\mathbf{y}, t)] d\mathbf{x} d\mathbf{y}. \end{aligned} \tag{6.6}$$

7. DETAILS OF THE PROOF OF THEOREM 5.3

The purpose of this section is to prove the various lemmas and theorems cited in the proof of our main result, Theorem 5.3. Without loss of generality we assume that $\tilde{L} = 0$ in this section. The case when $\tilde{L} > 0$ follows by similar arguments as we now give in the $\tilde{L} = 0$ case.

To rigorously determine the large-population limit of the MVSP, we use the martingale problem approach for studying solutions to stochastic differential equations developed by Stroock and Varadhan [EK86, SV06]. The proof is organized as follows. In Subsection 7.1 we provide the path level description of $\mu_t^{\zeta, j}$, analogous to (3.1) for ν_t^ζ , and in Subsection 7.2 we derive equations for its expectation. Assuming that the large-population limit exists, its identification is presented in Subsection 7.3. Then, in Subsection 7.4 we prove

that the limit exists by proving that the sequence of measures is appropriately tight. We conclude in Subsection 7.5 by proving that the limit equation has a unique solution. Collectively, these results imply Theorem 5.3.

7.1. Path level description. Using that we can write the marginal distribution (molar concentration) of species j as

$$\mu_t^{\zeta,j}(dx) = \frac{1}{\gamma} \sum_{i=1}^{\gamma \langle 1, \mu_t^{\zeta,j} \rangle} \delta_{H_Q^i(\gamma \mu_t^{\zeta,j})}(dx), \quad j \in \{1, \dots, J\},$$

we have, analogously to (3.1), the coupled system

$$\begin{aligned} \langle f, \mu_t^{\zeta,j} \rangle &= \langle f, \mu_0^{\zeta,j} \rangle + \frac{1}{\gamma} \sum_{i \geq 1} \int_0^t \mathbf{1}_{\{i \leq \gamma \langle 1, \mu_{s-}^{\zeta,j} \rangle\}} \sqrt{2D_j} \frac{\partial f}{\partial Q} (H^i(\gamma \mu_{s-}^{\zeta,j})) dW_s^i + \frac{1}{\gamma} \int_0^t \sum_{i=1}^{\gamma \langle 1, \mu_{s-}^{\zeta,j} \rangle} D_j \frac{\partial^2 f}{\partial Q^2} (H^i(\gamma \mu_{s-}^{\zeta,j})) ds \\ &+ \sum_{\ell=1}^L \int_0^t \int_{\mathbb{I}^{(\ell)}} \int_{\mathbb{Y}^{(\ell)}} \int_{\mathbb{R}_+^2} \left(\left\langle f, \mu_{s-}^{\zeta,j} - \frac{1}{\gamma} \sum_{r=1}^{\alpha_{\ell j}} \delta_{H_Q^{i_r(j)}(\gamma \mu_{s-}^{\zeta,j})} + \frac{1}{\gamma} \sum_{r=1}^{\beta_{\ell j}} \delta_{y_r^{(j)}} \right\rangle - \langle f, \mu_{s-}^{\zeta,j} \rangle \right) \\ &\times \mathbf{1}_{\{i \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta,j})\}} \times \mathbf{1}_{\{\theta_1 \leq K_\ell^\gamma(\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i}))\}} \times \mathbf{1}_{\{\theta_2 \leq m_\ell^\eta(\mathbf{y} | \mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i}))\}} dN_\ell(s, \mathbf{i}, \mathbf{y}, \theta_1, \theta_2), \end{aligned} \quad (7.1)$$

for $j \in \{1, \dots, J\}$.

In this formulation, one important fact is that for fixed ζ , $\gamma \langle 1, \mu_{s-}^{\zeta,j} \rangle$ is finite by assumption, which provides exchangeability of the sum and Lebesgue integral.

7.2. Taking expectations. By taking expectation on (7.1) we obtain

$$\begin{aligned} \mathbb{E}[\langle f, \mu_t^{\zeta,j} \rangle] &= \mathbb{E}[\langle f, \mu_0^{\zeta,j} \rangle] + \mathbb{E}\left[\frac{1}{\gamma} \int_0^t \sum_{i=1}^{\gamma \langle 1, \mu_{s-}^{\zeta,j} \rangle} D_j \frac{\partial^2 f}{\partial Q^2} (H^i(\gamma \mu_{s-}^{\zeta,j})) ds\right] + \sum_{\ell=1}^L \mathbb{E}\left[\frac{1}{\gamma} \int_0^t \int_{\mathbb{Y}^{(\ell)}} \sum_{\mathbf{i} \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta,j})} \right. \\ &\left. \left(- \sum_{r=1}^{\alpha_{\ell j}} f(H_Q^{i_r(j)}(\gamma \mu_{s-}^{\zeta,j})) + \sum_{r=1}^{\beta_{\ell j}} f(y_r^{(j)}) \right) \times K_\ell^\gamma(\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i})) \times m_\ell(\mathbf{y} | \mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i})) d\mathbf{y} ds\right], \\ &= \mathbb{E}[\langle f, \mu_0^{\zeta,j} \rangle] + \mathbb{E}\left[\int_0^t \int_{\mathbb{R}^d} \frac{1}{\gamma} \sum_{i=1}^{\gamma \langle 1, \mu_{s-}^{\zeta,j} \rangle} D_j \frac{\partial^2 f}{\partial Q^2}(x) \delta_{H_Q^i(\gamma \mu_{s-}^{\zeta,j})}(dx) ds\right] \\ &- \sum_{\ell=1}^L \mathbb{E}\left[\int_0^t \frac{1}{\gamma} \sum_{\mathbf{i} \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta,j})} \left(\sum_{r=1}^{\alpha_{\ell j}} f(H_Q^{i_r(j)}(\gamma \mu_{s-}^{\zeta,j})) \right) K_\ell^\gamma(\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i})) \left(\int_{\mathbb{Y}^{(\ell)}} m_\ell^\eta(\mathbf{y} | \mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i})) d\mathbf{y} \right) ds\right], \\ &+ \sum_{\ell=1}^L \mathbb{E}\left[\int_0^t \frac{1}{\gamma} \sum_{\mathbf{i} \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta,j})} K_\ell^\gamma(\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i})) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f(y_r^{(j)}) \right) m_\ell^\eta(\mathbf{y} | \mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i})) d\mathbf{y} \right) ds\right], \\ &= \mathbb{E}[\langle f, \mu_0^{\zeta,j} \rangle] + \mathbb{E}\left[\int_0^t \langle (\mathcal{L}_j f)(x), \mu_{s-}^{\zeta,j}(dx) \rangle ds\right] \\ &- \sum_{\ell=1}^L \mathbb{E}\left[\int_0^t \frac{1}{\gamma} \int_{\mathbb{X}^{(\ell)}} \sum_{\mathbf{i} \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta,j})} \left(\sum_{r=1}^{\alpha_{\ell j}} f(x_r^{(j)}) \right) K_\ell^\gamma(\mathbf{x}) \delta_{\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i})}(\mathbf{d}\mathbf{x}) ds\right], \\ &+ \sum_{\ell=1}^L \mathbb{E}\left[\int_0^t \frac{1}{\gamma} \int_{\mathbb{X}^{(\ell)}} \sum_{\mathbf{i} \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta,j})} K_\ell^\gamma(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f(y_r^{(j)}) \right) m_\ell^\eta(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \delta_{\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i})}(\mathbf{d}\mathbf{x}) ds\right], \\ &= \mathbb{E}[\langle f, \mu_0^{\zeta,j} \rangle] + \mathbb{E}\left[\int_0^t \langle (\mathcal{L}_j f)(x), \mu_{s-}^{\zeta,j}(dx) \rangle ds\right] \end{aligned}$$

$$\begin{aligned}
& - \sum_{\ell=1}^L \mathbb{E} \left[\int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} \left(\sum_{r=1}^{\alpha_{\ell j}} f(x_r^{(j)}) \right) K_{\ell}(\mathbf{x}) \lambda^{(\ell)}[\mu_{s-}^{\zeta}](d\mathbf{x}) ds \right], \\
& + \sum_{\ell=1}^L \mathbb{E} \left[\int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_{\ell}(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f(y_r^{(j)}) \right) m_{\ell}^{\eta}(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s-}^{\zeta}](d\mathbf{x}) ds \right], \\
& = \mathbb{E}[\langle f, \mu_0^{\zeta, j} \rangle] + \mathbb{E} \left[\int_0^t \langle (\mathcal{L}_j f)(x), \mu_{s-}^{\zeta, j}(dx) \rangle ds \right] \\
& + \sum_{\ell=1}^L \mathbb{E} \left[\int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_{\ell}(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f(y_r^{(j)}) \right) m_{\ell}^{\eta}(\mathbf{y} | \mathbf{x}) d\mathbf{y} - \sum_{r=1}^{\alpha_{\ell j}} f(x_r^{(j)}) \right) \lambda^{(\ell)}[\mu_{s-}^{\zeta}](d\mathbf{x}) ds \right],
\end{aligned} \tag{7.2}$$

where in the third equality we use the assumption that $\int_{\mathbb{Y}^{(\ell)}} m_{\ell}^{\eta}(\mathbf{y} | \mathbf{x}) d\mathbf{y} = 1$. For the second to last equality we switch integrals of the form $\int_{\mathbb{X}^{(\ell)}} \sum_{\mathbf{i} \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta})} \cdots \delta_{\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta}, \mathbf{i})}(d\mathbf{x})$ to $\int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} \cdots \lambda^{(\ell)}[\mu_{s-}^{\zeta}](d\mathbf{x})$ using the definition of $\mu_s^{\zeta, j}(dx)$ and $\lambda^{(\ell)}[\cdot]$ (see Definition 2.6), and removing probability zero sets where two particles with the same type are simultaneously located at the same spatial location (see Definition 2.7). Note, by definition the effective reactant index sampling space $\Omega^{(\ell)}$ (see Definition 2.5) orders indices for particles of the same species. In converting from integrals involving the positions of individual particles (i.e. $\delta_{\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta}, \mathbf{i})}(d\mathbf{x})$) to integrals involving product measures ($\lambda^{(\ell)}[\mu_{s-}^{\zeta}](d\mathbf{x})$) we need to remove the "diagonal" indices by means of integrating on $\tilde{\mathbb{X}}^{(\ell)}$ (see Definition 2.7) and normalizing by the total number of index orderings, ($\boldsymbol{\alpha}^{(\ell)}!$).

7.3. Identification of the Limit. Inspired by Eq (7.2), we expect that if the weak limit, as ζ goes to zero, of the marginal distribution vector $\boldsymbol{\mu}_t^{\zeta} := (\mu_t^{\zeta, 1}, \mu_t^{\zeta, 2}, \dots, \mu_t^{\zeta, J})$ exists and is unique, then it will satisfy the analogous equation. For this purpose, denote $\boldsymbol{\xi}_t := (\xi_t^1, \xi_t^2, \dots, \xi_t^J)$ to be the corresponding limiting particle distribution on $\mathbb{R}^{J \times d}$ and $\xi_t = \sum_{j=1}^J \xi_t^j \delta_{S_j}$ to be the corresponding limiting particle distribution on \hat{P} . Then for each $1 \leq j \leq J$, the following must be satisfied,

$$\begin{aligned}
\langle f, \xi_t^j \rangle & = \langle f, \xi_0^j \rangle + \int_0^t \langle (\mathcal{L}_j f)(x), \xi_s^j(dx) \rangle ds \\
(7.3) \quad & + \sum_{\ell=1}^L \int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_{\ell}(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f(y_r^{(j)}) \right) m_{\ell}(\mathbf{y} | \mathbf{x}) d\mathbf{y} - \sum_{r=1}^{\alpha_{\ell j}} f(x_r^{(j)}) \right) \lambda^{(\ell)}[\xi_s](d\mathbf{x}) ds.
\end{aligned}$$

Existence of the limit is shown in the tightness Section 7.4, while uniqueness is shown in Section 7.5.

Let \mathcal{S} be the collection of elements Φ in $B(\otimes_{j=1}^J M_F(\mathbb{R}^d))$ of the form

$$(7.4) \quad \Phi(\boldsymbol{\mu}) = \varphi(\langle f_1, \boldsymbol{\mu} \rangle, \langle f_2, \boldsymbol{\mu} \rangle \dots \langle f_M, \boldsymbol{\mu} \rangle)$$

for some $M \in \mathbb{N}$, some $\varphi \in C^{\infty}(\mathbb{R}^{J \times M})$, $\langle f_m, \boldsymbol{\mu} \rangle = (\langle f_{1,m}, \mu^1 \rangle, \dots, \langle f_{J,m}, \mu^J \rangle)$ where each $\{f_{j,m}\} \in C_b^2(\mathbb{R}^d)$ for $j = 1, \dots, J$ and $m = 1, \dots, M$. Then \mathcal{S} separates points in $\otimes_{j=1}^J M_F(\mathbb{R}^d)$ (see Chapter 3.4 of [EK86] and Proposition 3.3 of [CSY20]). As long as the limiting process exists and is unique, to identify the limit, it thus suffices to show convergence of the martingale problem for functions of the form (7.4).

For $\Phi \in \mathcal{S}$ of the form (7.4), $\boldsymbol{\mu} := (\mu^1, \mu^2, \dots, \mu^J) \in \otimes_{j=1}^J M_F(\mathbb{R}^d)$ and $\mu = \sum_{j=1}^J \mu^j \delta_{S_j} \in M_F(\hat{P})$ with each $\mu^j \in M_F(\mathbb{R}^d)$, define

$$\begin{aligned}
(\mathcal{A}\Phi)(\boldsymbol{\mu}) & \stackrel{\text{def}}{=} \sum_{m=1}^M \sum_{j=1}^J \frac{\partial \varphi}{\partial x_{(m-1) * J + j}}(\langle f_1, \boldsymbol{\mu} \rangle, \langle f_2, \boldsymbol{\mu} \rangle \dots \langle f_M, \boldsymbol{\mu} \rangle) \left\{ \langle \mathcal{L}_{\ell} f_{j,m}, \mu^j \rangle \right. \\
& \quad \left. + \sum_{\ell=1}^L \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_{\ell}(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_{j,m}(y_r^{(j)}) \right) m_{\ell}(\mathbf{y} | \mathbf{x}) d\mathbf{y} - \sum_{r=1}^{\alpha_{\ell j}} f_{j,m}(x_r^{(j)}) \right) \lambda^{(\ell)}[\mu](d\mathbf{x}) \right\}
\end{aligned} \tag{7.5}$$

We claim that \mathcal{A} , which is the generator of the system described by (7.3) for $1 \leq j \leq J$, will be the generator of the limiting martingale problem.

Lemma 7.1 (Weak Convergence). *For any $\Phi \in \mathcal{S}$ and $0 \leq r_1 \leq r_2 \cdots \leq r_W = s < t < T$ and $\{\psi_w\}_{w=1}^W \subset B(\otimes_{j=1}^J M_F(\mathbb{R}^d))$, we have that*

$$(7.6) \quad \lim_{\zeta \rightarrow 0} \mathbb{E} \left[\left\{ \Phi(\mu_t^\zeta) - \Phi(\mu_s^\zeta) - \int_s^t (\mathcal{A}\Phi)(\mu_r^\zeta) dr \right\} \prod_{w=1}^W \psi_w(\mu_{r_w}^\zeta) \right] = 0.$$

Proof. For each $j = 1, \dots, J$, we can rewrite Eq (7.1) as

$$\langle f, \mu_t^{\zeta, j} \rangle = \langle f, \mu_0^{\zeta, j} \rangle + M_t^{f, j} + A_t^{f, j},$$

where

$$(7.7) \quad A_t^{f, j} = \int_0^t \left\langle (\mathcal{L}_j f)(x), \mu_{s-}^{\zeta, j}(dx) \right\rangle ds + \sum_{\ell=1}^L \int_0^t \int_{\tilde{\mathbb{X}}(\ell)} \frac{1}{\alpha^{(\ell)!}} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}(\ell)} \left(\sum_{r=1}^{\beta_{\ell j}} f(y_r^{(j)}) \right) m_\ell^\eta(\mathbf{y} | \mathbf{x}) d\mathbf{y} - \sum_{r=1}^{\alpha_{\ell j}} f(x_r^{(j)}) \right) \lambda^{(\ell)}[\mu_{s-}^{\zeta, j}](d\mathbf{x}) ds.$$

and

$$(7.8) \quad M_t^{f, j} = \frac{1}{\gamma} \sum_{i \geq 1} \int_0^t 1_{\{i \leq \gamma \langle 1, \mu_{s-}^{\zeta, j} \rangle\}} \sqrt{2D_j} \frac{\partial f}{\partial Q}(H^i(\gamma \mu_{s-}^{\zeta, j})) dW_s^i + \sum_{\ell=1}^L \int_0^t \int_{\mathbb{I}(\ell)} \int_{\mathbb{Y}(\ell)} \int_{\mathbb{R}_+^2} \left(\left\langle f, \mu_{s-}^{\zeta, j} - \frac{1}{\gamma} \sum_{r=1}^{\alpha_{\ell j}} \delta_{H_Q^{i_r^{(j)}}(\gamma \mu_{s-}^{\zeta, j})} + \frac{1}{\gamma} \sum_{r=1}^{\beta_{\ell j}} \delta_{y_r^{(j)}} \right\rangle - \langle f, \mu_{s-}^{\zeta, j} \rangle \right) \times 1_{\{i \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta, j})\}} \times 1_{\{\theta_1 \leq K_\ell^\gamma(\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta, j}), \mathbf{i})\}} \times 1_{\{\theta_2 \leq m_\ell^\eta(\mathbf{y} | \mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta, j}), \mathbf{i})\}} d\tilde{N}_\ell(s, \mathbf{i}, \mathbf{y}, \theta_1, \theta_2),$$

is a square integrable martingale (See Proposition 2.4 in [NW14]) with quadratic variation

$$(7.9) \quad \begin{aligned} \langle M^{f, j} \rangle_t &= \frac{1}{\gamma^2} \int_0^t \sum_{i=1}^{\gamma \langle 1, \mu_{s-}^{\zeta, j} \rangle} \left(\sqrt{2D_j} \frac{\partial f}{\partial Q}(H^i(\gamma \mu_{s-}^{\zeta, j})) \right)^2 ds \\ &+ \sum_{\ell=1}^L \int_0^t \int_{\mathbb{Y}(\ell)} \frac{1}{\gamma^2} \sum_{\{\mathbf{i} \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta, j})\}} \left(- \sum_{r=1}^{\alpha_{\ell j}} f(H_Q^{i_r^{(j)}}(\gamma \mu_{s-}^{\zeta, j})) + \sum_{r=1}^{\beta_{\ell j}} f(y_r^{(j)}) \right)^2 \\ &\quad \times K_\ell^\gamma(\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta, j}), \mathbf{i}) \times m_\ell^\eta(\mathbf{y} | \mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta, j}), \mathbf{i}) d\mathbf{y} ds, \\ &\leq \frac{1}{\gamma} \int_0^t \left\langle 2D_j \left(\frac{\partial f}{\partial x} \right)^2, \mu_{s-}^{\zeta, j} \right\rangle ds \\ &+ \sum_{\ell=1}^L \int_0^t \int_{\mathbb{Y}(\ell)} \frac{1}{\gamma^2} \sum_{\{\mathbf{i} \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta, j})\}} \|f\|_{C_b(\mathbb{R}^d)}^2 (\alpha_{\ell j} + \beta_{\ell j})^2 K_\ell^\gamma(\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta, j}), \mathbf{i}) m_\ell^\eta(\mathbf{y} | \mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta, j}), \mathbf{i}) d\mathbf{y} ds, \\ &\leq \frac{C(D_j)t \|f\|_{C_b^1(\mathbb{R}^d)}^2}{\gamma} C(\mu) + \sum_{\ell=1}^L \|f\|_{C_b(\mathbb{R}^d)}^2 (\alpha_{\ell j} + \beta_{\ell j})^2 \int_0^t \frac{1}{\gamma^2} \sum_{\{\mathbf{i} \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta, j})\}} K_\ell^\gamma(\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta, j}), \mathbf{i}) ds, \\ &\leq \frac{C(D_j)t \|f\|_{C_b^1(\mathbb{R}^d)}^2}{\gamma} C(\mu) + C(K)4^2 \|f\|_{C_b(\mathbb{R}^d)}^2 \sum_{\ell=1}^L \int_0^t \frac{1}{\gamma^2} \sum_{\{\mathbf{i} \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta, j})\}} \gamma^{1-|\alpha^{(\ell)}|} ds, \\ &\leq \frac{C(D_j)t \|f\|_{C_b^1(\mathbb{R}^d)}^2}{\gamma} C(\mu) + \frac{C(K)tL \|f\|_{C_b(\mathbb{R}^d)}^2}{\gamma} C(\mu). \end{aligned}$$

The quadratic variation is therefore uniformly bounded and goes to 0 as $\zeta \rightarrow 0$ ($\gamma \rightarrow \infty$) since f and its partial derivatives are uniformly bounded.

Now define $M_t^{f,j} = \mathcal{C}_t^{f,j} + \mathcal{D}_t^{f,j}$, where

$$\mathcal{C}_t^{f,j} = \frac{1}{\gamma} \sum_{i \geq 1} \int_0^t \mathbf{1}_{\{i \leq \gamma \langle 1, \mu_{s-}^{\zeta,j} \rangle\}} \sqrt{2D_j} \frac{\partial f}{\partial Q} (H^i(\gamma \mu_{s-}^{\zeta,j})) dW_s^i \quad (7.10)$$

is the continuous martingale part and

$$\begin{aligned} \mathcal{D}_t^{f,j} &= \sum_{\ell=1}^L \int_0^t \int_{\mathbb{I}^{(\ell)}} \int_{\mathbb{Y}^{(\ell)}} \int_{\mathbb{R}_+^2} \left(\left\langle f, \mu_{s-}^{\zeta,j} - \frac{1}{\gamma} \sum_{r=1}^{\alpha_{\ell j}} \delta_{H_Q^{i_r(j)}(\gamma \mu_{s-}^{\zeta,j})} + \frac{1}{\gamma} \sum_{r=1}^{\beta_{\ell j}} \delta_{y_r^{(j)}} \right\rangle - \langle f, \mu_{s-}^{\zeta,j} \rangle \right) \\ (7.11) \quad &\times \mathbf{1}_{\{i \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta,j})\}} \times \mathbf{1}_{\{\theta_1 \leq K_\ell^\gamma(\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i}))\}} \times \mathbf{1}_{\{\theta_2 \leq m_\ell^\eta(\mathbf{y} | \mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i}))\}} d\tilde{N}_\ell(s, \mathbf{i}, \mathbf{y}, \theta_1, \theta_2), \end{aligned}$$

is the martingale part coming from the stochastic integral with respect to the Poisson point processes. Here, for simplicity of notation, we let

$$\begin{aligned} g^{\ell, f, \mu^{\zeta,j}}(s, \mathbf{i}, \mathbf{y}, \theta_1, \theta_2) &= \left(\left\langle f, \mu_{s-}^{\zeta,j} - \frac{1}{\gamma} \sum_{r=1}^{\alpha_{\ell j}} \delta_{H_Q^{i_r(j)}(\gamma \mu_{s-}^{\zeta,j})} + \frac{1}{\gamma} \sum_{r=1}^{\beta_{\ell j}} \delta_{y_r^{(j)}} \right\rangle - \langle f, \mu_{s-}^{\zeta,j} \rangle \right) \\ &\times \mathbf{1}_{\{i \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta,j})\}} \times \mathbf{1}_{\{\theta_1 \leq K_\ell^\gamma(\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i}))\}} \times \mathbf{1}_{\{\theta_2 \leq m_\ell^\eta(\mathbf{y} | \mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i}))\}} \\ (7.12) \quad &= \frac{1}{\gamma} \left(- \sum_{r=1}^{\alpha_{\ell j}} f \left(H_Q^{i_r(j)}(\gamma \mu_{s-}^{\zeta,j}) \right) + \sum_{r=1}^{\beta_{\ell j}} f \left(y_r^{(j)} \right) \right) \\ &\times \mathbf{1}_{\{i \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\zeta,j})\}} \times \mathbf{1}_{\{\theta_1 \leq K_\ell^\gamma(\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i}))\}} \times \mathbf{1}_{\{\theta_2 \leq m_\ell^\eta(\mathbf{y} | \mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\zeta,j}, \mathbf{i}))\}}, \end{aligned}$$

which represents the jumps and is uniformly bounded by $\mathcal{O}(\frac{1}{\gamma})$. With some abuse of notation we shall write $\mathbf{g}^{\ell, f, \mu^\zeta}$ for the vector $(g^{\ell, f, \mu^{\zeta,1}}, \dots, g^{\ell, f, \mu^{\zeta,J}})$. Then (7.11) becomes

$$\mathcal{D}_t^{f,j} = \sum_{\ell=1}^L \int_0^t \int_{\mathbb{I}^{(\ell)}} \int_{\mathbb{Y}^{(\ell)}} \int_{\mathbb{R}_+^2} g^{\ell, f, \mu^{\zeta,j}}(s, \mathbf{i}, \mathbf{y}, \theta_1, \theta_2) d\tilde{N}_\ell(s, \mathbf{i}, \mathbf{y}, \theta_1, \theta_2).$$

Now we apply Itô's formula (See Theorem 5.1 in [NW14]) to $\Phi(\mu_t^\zeta)$. We obtain for $\theta = (\theta_1, \theta_2)$,

$$\begin{aligned} \Phi(\mu_t^\zeta) - \Phi(\mu_s^\zeta) - \int_s^t (\mathcal{A}\Phi)(\mu_r^\zeta) dr &= \int_s^t \sum_{m=1}^M \sum_{j=1}^J \frac{\partial \varphi}{\partial x_{(m-1)J+j}} (\langle f_1, \mu_r^\zeta \rangle, \langle f_2, \mu_r^\zeta \rangle, \dots, \langle f_M, \mu_r^\zeta \rangle) d\mathcal{C}_r^{f_j, m, j} \\ &+ \frac{1}{2} \int_s^t \sum_{m=1}^M \sum_{j=1}^J \frac{\partial^2 \varphi}{\partial x_{(m-1)J+j}^2} (\langle f_1, \mu_r^\zeta \rangle, \langle f_2, \mu_r^\zeta \rangle, \dots, \langle f_M, \mu_r^\zeta \rangle) d\langle \mathcal{C}^{f_j, m, j} \rangle_r \\ &+ \sum_{\ell=1}^L \int_s^t \int_{\mathbb{I}^{(\ell)}} \int_{\mathbb{Y}^{(\ell)}} \int_{\mathbb{R}_+^2} \left(\varphi(\langle f_1, \mu_r^\zeta \rangle + \mathbf{g}^{\ell, f_1, \mu^\zeta}(r, \mathbf{i}, \mathbf{y}, \theta), \dots, \langle f_M, \mu_r^\zeta \rangle + \mathbf{g}^{\ell, f_M, \mu^\zeta}(r, \mathbf{i}, \mathbf{y}, \theta)) \right. \\ &\quad \left. - \varphi(\langle f_1, \mu_r^\zeta \rangle, \langle f_2, \mu_r^\zeta \rangle, \dots, \langle f_M, \mu_r^\zeta \rangle) \right) d\tilde{N}_\ell(r, \mathbf{i}, \mathbf{y}, \theta) \\ &+ \sum_{\ell=1}^L \int_s^t \int_{\mathbb{I}^{(\ell)}} \int_{\mathbb{Y}^{(\ell)}} \int_{\mathbb{R}_+^2} \left(\varphi(\langle f_1, \mu_r^\zeta \rangle + \mathbf{g}^{\ell, f_1, \mu^\zeta}(s, \mathbf{i}, \mathbf{y}, \theta), \dots, \langle f_M, \mu_r^\zeta \rangle + \mathbf{g}^{\ell, f_M, \mu^\zeta}(s, \mathbf{i}, \mathbf{y}, \theta)) \right. \\ &\quad \left. - \varphi(\langle f_1, \mu_r^\zeta \rangle, \langle f_2, \mu_r^\zeta \rangle, \dots, \langle f_M, \mu_r^\zeta \rangle) - \sum_{m=1}^M \sum_{j=1}^J g^{\ell, f_j, m, \mu^{\zeta,j}}(s, \mathbf{i}, \mathbf{y}, \theta) \frac{\partial \varphi}{\partial x_{(m-1)J+j}} (\langle f_1, \mu_r^\zeta \rangle, \dots, \langle f_M, \mu_r^\zeta \rangle) \right) \\ &\quad d\tilde{N}_\ell(s, \mathbf{i}, \mathbf{y}, \theta) \\ &+ \sum_{m=1}^M \sum_{j=1}^J \sum_{\ell=1}^L \int_0^t \frac{\partial \varphi}{\partial x_{(m-1)J+j}} (\langle f_1, \mu_r^\zeta \rangle, \langle f_2, \mu_r^\zeta \rangle, \dots, \langle f_M, \mu_r^\zeta \rangle) \end{aligned}$$

$$\begin{aligned}
& \times \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\alpha^{(\ell)!}} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_{j,m}(y_r^{(j)}) \right) (m_\ell^\eta(\mathbf{y} | \mathbf{x}) - m_\ell(\mathbf{y} | \mathbf{x})) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s-}^\zeta](d\mathbf{x}) ds \\
& = \sum_{\kappa=1}^5 \Lambda_\kappa^\zeta(t).
\end{aligned}
\tag{7.13}$$

We now use the Skorokhod representation theorem (Theorem 1.8 in [EK86]) which, for the purposes of identifying the limit and proving (7.6), allows us to assume that the aforementioned claimed convergence of $\mu_t^\zeta = (\mu_t^{\zeta,1}, \mu_t^{\zeta,2}, \dots, \mu_t^{\zeta,J})$ holds with probability one in the topology of weak convergence of measures. The Skorokhod representation theorem involves the introduction of another probability space, but we ignore this distinction in the notation. To show (7.6), it is then sufficient to prove that the left hand side of (7.13) goes to zero in probability. With this goal in mind we proceed with proving convergence in probability to zero for $\Lambda_\kappa^\zeta(t)$ for $\kappa = 1, \dots, 5$.

By Lemma B.2, we immediately have that

$$\lim_{\zeta \rightarrow 0} \sup_{t \in [0, T]} \mathbb{E}|\Lambda_5^\zeta(t)| = 0.$$

In addition, notice that $\Lambda_1^\zeta(t)$ and $\Lambda_3^\zeta(t)$ are square integrable martingales. In fact, by (7.9), (7.12) and the fact that the jump size is uniformly bounded by $\mathcal{O}(\frac{1}{\gamma})$, we have that

$$\lim_{\zeta \rightarrow 0} \sup_{t \in [0, T]} \mathbb{E}|\Lambda_1^\zeta(t) + \Lambda_3^\zeta(t)|^2 = 0.$$

For similar reasons, we also have by (7.10) that

$$\lim_{\zeta \rightarrow 0} \sup_{t \in [0, T]} \mathbb{E}|\Lambda_2^\zeta(t)| = 0,$$

and by (7.12) that

$$\lim_{\zeta \rightarrow 0} \sup_{t \in [0, T]} \mathbb{E}|\Lambda_4^\zeta(t)| = 0.$$

We then have that the left hand side of (7.13) goes to zero in probability, concluding the proof of the lemma. \square

7.4. Tightness. Recall that $M_F(\mathbb{R}^d)$ denotes the space of finite measures endowed with the weak topology, and denote by $M'_F(\mathbb{R}^d)$ the space of finite measures endowed with the vague topology. In this section, we prove tightness of the measure-valued processes $\{\mu_t^{\zeta,j}\}_{t \in [0, T], j = 1, 2, \dots, J}$ on $\mathbb{D}_{M_F(\mathbb{R}^d)}[0, T]$, the space of cadlag paths with values in $M_F(\mathbb{R}^d)$ endowed with Skorokhod topology. Towards this aim, we first show that the processes $\{\mu_t^{\zeta,j}\}_{t \in [0, T], j = 1, \dots, J}$, are tight on $\mathbb{D}_{M'_F(\mathbb{R}^d)}[0, T]$.

7.4.1. Tightness in $\mathbb{D}_{M'_F(\mathbb{R}^d)}[0, T]$. It suffices to show that the real-valued processes $\{\langle f, \mu_t^{\zeta,j} \rangle\}$, $j = 1, \dots, J$, for any test function $f(x) \in C_0^2(\mathbb{R}^d)$, which is dense in $C_0(\mathbb{R}^d)$, are tight in $\mathbb{D}_{\mathbb{R}}[0, T]$, see [S89]. In establishing this we use the Rebolledo Criterion [AM86] (Lemma 7.4) and the Aldous Condition [D78] (Lemma 7.3).

Lemma 7.2. *For any $T > 0$ and $\delta > 0$, there exists constants C and C' such that for any pair of stopping times (σ, τ) with $0 \leq \sigma \leq \tau \leq \sigma + \delta \leq T$, we have*

$$\mathbb{E}[\langle M^{f,j} \rangle_\tau - \langle M^{f,j} \rangle_\sigma] \leq C\delta.$$

and

$$\mathbb{E}[|A_\tau^{f,j} - A_\sigma^{f,j}|^2] \leq C'\delta^2,$$

for $j = 1, \dots, J$ and $A^{f,j}$, $M^{f,j}$ follow the definitions (7.7), (7.8) respectively.

Proof. Following (7.9), we can obtain that

$$\begin{aligned}
\mathbb{E}[\langle M^{f,j} \rangle_\tau - \langle M^{f,j} \rangle_\sigma] &= \frac{1}{\gamma^2} \mathbb{E} \left[\int_\sigma^\tau \sum_{i=1}^{\gamma \langle 1, \mu_{s-}^{\xi,j} \rangle} \left(\sqrt{2D_j} \frac{\partial f}{\partial Q} (H^i(\gamma \mu_{s-}^{\xi,j})) \right)^2 ds \right] \\
&\quad + \sum_{\ell=1}^L \mathbb{E} \left[\int_\sigma^\tau \int_{\mathbb{Y}^{(\ell)}} \frac{1}{\gamma^2} \sum_{\{\mathbf{i} \in \Omega^{(\ell)}(\gamma \mu_{s-}^{\xi,j})\}} \left(- \sum_{r=1}^{\alpha_{\ell j}} f(H_Q^{i_r^j}(\gamma \mu_{s-}^{\xi,j})) + \sum_{r=1}^{\beta_{\ell j}} f(y_r^j) \right)^2 \right. \\
&\quad \left. \times K_\ell^\gamma \left(\mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\xi,j}, \mathbf{i}) \right) \times m_\ell^\eta \left(\mathbf{y} \mid \mathcal{P}^{(\ell)}(\gamma \mu_{s-}^{\xi,j}, \mathbf{i}) \right) d\mathbf{y} ds \right], \\
&\leq \frac{C(\mu) \mathbb{E}[\tau - \sigma] \|f\|_{C_0^1(\mathbb{R}^d)}^2}{\gamma} (C(D_j) + C(K)L), \\
&\leq C\delta.
\end{aligned}$$

From (7.7), we obtain

$$\begin{aligned}
\mathbb{E}[|A_\tau^{f,j} - A_\sigma^{f,j}|^2] &= \mathbb{E} \left[\left| \int_\sigma^\tau \langle (\mathcal{L}_j f)(x), \mu_{s-}^{\xi,j}(dx) \rangle ds + \sum_{\ell=1}^L \int_\sigma^\tau \int_{\mathbb{X}^{(\ell)}} \frac{1}{\alpha^{(\ell)!}} K_\ell(\mathbf{x}) \right. \right. \\
&\quad \left. \left. \times \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f(y_r^{(j)}) \right) m_\ell^\eta(\mathbf{y} \mid \mathbf{x}) d\mathbf{y} - \sum_{r=1}^{\alpha_{\ell j}} f(x_r^{(j)}) \right) \lambda^{(\ell)}[\mu_{s-}^{\xi,j}](d\mathbf{x}) ds \right|^2 \right]. \\
&\leq 2\mathbb{E} \left[\left| \int_\sigma^\tau \langle (\mathcal{L}_j f)(x), \mu_{s-}^{\xi,j}(dx) \rangle ds \right|^2 \right] + 2\mathbb{E} \left[\left| \sum_{\ell=1}^L \int_\sigma^\tau \int_{\mathbb{X}^{(\ell)}} \frac{1}{\alpha^{(\ell)!}} K_\ell(\mathbf{x}) \right. \right. \\
&\quad \left. \left. \times \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} |f(y_r^{(j)})| \right) m_\ell^\eta(\mathbf{y} \mid \mathbf{x}) d\mathbf{y} + \sum_{r=1}^{\alpha_{\ell j}} |f(x_r^{(j)})| \right) \lambda^{(\ell)}[\mu_{s-}^{\xi,j}](d\mathbf{x}) ds \right|^2 \right] \\
&\leq C(D_j) \|f\|_{C_0^1(\mathbb{R}^d)}^2 C(\mu)^2 \mathbb{E}[|\tau - \sigma|^2] + C(K) \|f\|_{C_0^2(\mathbb{R}^d)}^2 L^2 C(\mu)^4 \mathbb{E}[|\tau - \sigma|^2] \\
&\leq C'\delta^2.
\end{aligned}$$

□

Lemma 7.3 (Aldous condition). *For any $T > 0$, $\varepsilon_1 > 0$, $\varepsilon_2 > 0$, $j = 1, 2, \dots, J$, there exists $\delta > 0$ and n_0 such that for any sequence $(\sigma_n, \tau_n)_{n \in \mathbb{N}}$ of pairs of stopping times with $\sigma_n \leq \tau_n \leq T$,*

$$\sup_{n \geq n_0} \mathbb{P}\{|\langle f, \mu_{\sigma_n}^{n,j} \rangle - \langle f, \mu_{\tau_n}^{n,j} \rangle| \geq \varepsilon_2, \tau_n \leq \sigma_n + \delta\} \leq \varepsilon_1.$$

Proof. This follows as for $\tau_n \leq \sigma_n + \delta$.

$$\begin{aligned}
&\sup_{n \geq n_0} \mathbb{P}\{|M_{\sigma_n}^{f,j} - M_{\tau_n}^{f,j}| + |A_{\sigma_n}^{f,j} - A_{\tau_n}^{f,j}| \geq \varepsilon_2\} \\
&\leq \sup_{n \geq n_0} 2 \frac{1}{\varepsilon_2} \mathbb{E}[\langle M^{f,j} \rangle_{\tau_n} - \langle M^{f,j} \rangle_{\sigma_n} + |A_{\tau_n}^{f,j} - A_{\sigma_n}^{f,j}|^2] \quad (\text{by Markov inequality}) \\
&\leq \frac{2}{\varepsilon_2^2} (C\delta + C'\delta^2) \leq \varepsilon_1 \quad (\text{by Lemma 7.2 and when } \delta \text{ sufficiently small})
\end{aligned}$$

□

Lemma 7.4 (Rebolledo Criterion). *For each $j = 1, \dots, J$, the sequence of real-valued processes $\{\langle f, \mu_t^{\xi,j} \rangle\}_{\zeta \in (0,1)^2}$ is tight in $\mathbb{D}_{\mathbb{R}}[0, T]$.*

Proof. By assumption, we always have $\langle 1, \mu_t^{\xi,j} \rangle$, $j = 1, \dots, J$ are uniformly bounded. Since $f \in C_0^2(\mathbb{R}^d)$, we have

$$\sup_{\|\zeta\| \leq 1} \mathbb{E} \left[\sup_{t \in [0, T]} |\langle f, \mu_t^{\xi,j} \rangle| \right] < \infty$$

holds. Combined with the Aldous condition from Lemma 7.3, we obtain that, for each $j = 1, 2, \dots, J$, the sequence of real-valued processes $\{\langle f, \mu_t^{\zeta, j} \rangle\}_{\zeta \in (0,1)^2}$ is tight in $\mathbb{D}_{\mathbb{R}}[0, T]$ by the Rebolledo Criterion [AM86]. \square

7.4.2. *Tightness in $\mathbb{D}_{M_F(\mathbb{R}^d)}[0, T]$.* By the next Lemma 7.5, we're able to control the mass of measures outside of compact sets so that we can go from tightness in $\mathbb{D}_{M'_F(\mathbb{R}^d)}[0, T]$ to tightness in $\mathbb{D}_{M_F(\mathbb{R}^d)}[0, T]$.

Lemma 7.5. *There exists a sequence of $C_b^2(\mathbb{R}^d)$ functions $\{f_m(x)\}_{m \geq 0}$, in particular, $f_0 \equiv 1$, such that*

$$\begin{aligned} f_m(x) &= 0 \text{ when } \|x\| \leq m-1 \\ f_m(x) &= 1 \text{ when } \|x\| > m \\ 0 \leq f_m(x) &\leq 1 \text{ when } m-1 < \|x\| \leq m. \end{aligned}$$

and furthermore, $\sup_{m \geq 0} \|f_m(x)\|_{C_b^2(\mathbb{R}^d)} := \sup_{m \geq 0} \sup_{x \in \mathbb{R}^d, |\alpha| \leq 2} |D^\alpha f_m(x)| < \infty$. For such sequence of functions $\{f_m(x)\}_{m \geq 0}$,

$$(7.14) \quad \lim_{m \rightarrow \infty} \limsup_{\zeta \rightarrow 0} \mathbb{E} \left(\sup_{t \in [0, T]} \langle f_m, \mu_t^{\zeta, j} \rangle \right) = 0.$$

for all $j = 1, 2, \dots, J$.

Proof. Following [BMW12], consider the function $\psi(s) = 6s^5 - 15s^4 + 10s^3 \in C^2([0, 1])$. One can check that $\psi(0) = 1 - \psi(1) = \psi'(0) = \psi'(1) = \psi''(0) = \psi''(1) = 0$. Now we define our functions $\{f_m(x), x \in \mathbb{R}^d, m \geq 1\}$ as $f_m(x) = \psi(0 \vee (\|x\| - (m-1)) \wedge 1)$. The derivatives of f_m s are uniformly controlled by the derivatives of ψ , thus this choice satisfies our conditions. For any $\varepsilon > 0$, by Assumption 4.7, there exists a large enough integer-valued radius R such that $\int_{r>R} \rho(r)r^{d-1}dr < \varepsilon$. As a consequence, for m sufficiently large

$$\begin{aligned} \langle f_m, \mu_t^{\zeta, j} \rangle &= \langle f_m, \mu_0^{\zeta, j} \rangle + M_t^{f_m, j} + \int_0^t \langle (\mathcal{L}_j f_m)(x), \mu_{s-}^{\zeta, j}(dx) \rangle \\ &\quad + \sum_{\ell=1}^L \int_0^t \int_{\bar{\mathbb{X}}(\varepsilon)} \frac{1}{\alpha^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}(\varepsilon)} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell^\eta(\mathbf{y} | \mathbf{x}) d\mathbf{y} - \sum_{r=1}^{\alpha_{\ell j}} f_m(x_r^{(j)}) \right) \lambda^{(\ell)}[\mu_{s-}^{\zeta, j}](d\mathbf{x}) ds, \\ &\leq \langle f_m, \mu_0^{\zeta, j} \rangle + M_t^{f_m, j} + \|f_m\|_{C_b^2(\mathbb{R}^d)} \int_0^t \langle 1_{\{m-1 \leq \|x\| < m\}}, \mu_{s-}^{\zeta, j}(dx) \rangle \\ &\quad + \sum_{\ell=1}^L \int_0^t \int_{\bar{\mathbb{X}}(\varepsilon)} \frac{1}{\alpha^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}(\varepsilon)} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell^\eta(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s-}^{\zeta, j}](d\mathbf{x}) ds, \\ &\leq \langle f_m, \mu_0^{\zeta, j} \rangle + M_t^{f_m, j} + \|f_m\|_{C_b^2(\mathbb{R}^d)} \int_0^t \langle f_{m-1}(x), \mu_{s-}^{\zeta, j}(dx) \rangle \\ &\quad + \sum_{\ell=1}^L \int_0^t \int_{\bar{\mathbb{X}}(\varepsilon)} \frac{1}{\alpha^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}(\varepsilon)} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell^\eta(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s-}^{\zeta, j}](d\mathbf{x}) ds. \end{aligned} \tag{7.15}$$

Taking supremum over time and then expectation on Eq 7.15, we get

$$\begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} \langle f_m, \mu_t^{\zeta, j} \rangle \right] &\leq \mathbb{E} \left[\langle f_m, \mu_0^{\zeta, j} \rangle \right] + \mathbb{E} \left[\sup_{t \in [0, T]} |M_t^{f_m, j}| \right] + \|f_m\|_{C_b^2(\mathbb{R}^d)} \mathbb{E} \left[\sup_{t \in [0, T]} \int_0^t \langle f_{m-1}(x), \mu_{s-}^{\zeta, j}(dx) \rangle \right] \\ &\quad + \mathbb{E} \left[\sup_{t \in [0, T]} \sum_{\ell=1}^L \int_0^t \int_{\bar{\mathbb{X}}(\varepsilon)} \frac{1}{\alpha^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}(\varepsilon)} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell^\eta(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s-}^{\zeta, j}](d\mathbf{x}) ds \right]. \\ &\leq \mathbb{E} \left[\langle f_m, \mu_0^{\zeta, j} \rangle \right] + \mathbb{E} \left[\sup_{t \in [0, T]} |M_t^{f_m, j}| \right] + \|f_m\|_{C_b^2(\mathbb{R}^d)} \mathbb{E} \left[\sup_{t \in [0, T]} \int_0^t \langle f_{m-1}(x), \mu_{s-}^{\zeta, j}(dx) \rangle \right] \end{aligned}$$

$$\begin{aligned}
& + \sup_{1 \leq i \leq J} \mathbb{E} \left[\sup_{t \in [0, T]} C \int_0^t \left\langle f_{m-1-R}(x), \mu_{s-}^{\zeta, i}(dx) \right\rangle ds \right] + C_2 T (\varepsilon + C\eta), \\
& \quad (\text{by Lemma B.1 studying the different cases of allowed reactions}) \\
& \leq \mathbb{E} \left[\sup_{t \in [0, T]} \left\langle f_m, \mu_0^{\zeta, j} \right\rangle \right] + \mathbb{E} \left[\sup_{t \in [0, T]} \left| M_t^{f_m, j} \right| \right] \\
& \quad + C_1 \int_0^T \sup_{1 \leq i \leq J} \mathbb{E} \left[\sup_{s \in [0, t]} \left\langle f_{m-1-R}(x), \mu_{s-}^{\zeta, i}(dx) \right\rangle \right] dt + C_2 T (\varepsilon + C\eta),
\end{aligned}$$

(7.16)

where $C = 2LC(K)(C(\mu) \vee 1)$, $C_1 = 2(C \vee \|f_m\|_{C_b^2(\mathbb{R}^d)})$ and $C_2 = 2LC(K)C(\mu)\|f_m\|_{C_b^2(\mathbb{R}^d)}$.

Let $Y_T^{m, \zeta} := \sup_{1 \leq j \leq J} \mathbb{E} \left[\sup_{t \in [0, T]} \left(\left\langle f_m, \mu_t^{\zeta, j} \right\rangle \right) \right]$. By construction, we always have $Y_T^{m, \zeta} \leq Y_T^{m-1, \zeta} \leq Y_T^{0, \zeta}$. Due to the uniform boundedness of $\|f_m\|_{C_b^2(\mathbb{R}^d)}$, we have that $\mathbb{E} \langle M^{f_m, j} \rangle_T = \mathcal{O}(\frac{1}{\gamma})$, uniformly for each $1 \leq j \leq J$ and for all m based on Eq (7.9). Without loss of generality, let's consider the subsequence where m is divisible by $R+1$. Then Eq (7.16) gives

$$\begin{aligned}
Y_T^{m, \zeta} & \leq Y_0^{m, \zeta} + \sup_{1 \leq j \leq J} \mathbb{E} \left[\sup_{t \in [0, T]} |M_t^{f_m, j}| \right] + C_1 \int_0^T Y_t^{m-(R+1), \zeta} dt + C_2 T (\varepsilon + C\eta), \\
& \leq Y_0^{m, \zeta} + 2 \sup_{1 \leq j \leq J} \sqrt{\mathbb{E} \langle M^{f_m, j} \rangle_T} + C_1 \int_0^T Y_t^{m-(R+1), \zeta} dt + C_2 T (\varepsilon + C\eta) \\
& \quad (\text{by Jensen's inequality and Doob's inequality}) \\
& \leq Y_0^{m, \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T (\varepsilon + C\eta) + C_1 \int_0^T Y_t^{m-(R+1), \zeta} dt \\
& \leq Y_0^{m, \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T (\varepsilon + C\eta) \\
& \quad + C_1 \int_0^T \left(Y_0^{m-(R+1), \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T (\varepsilon + C\eta) + C_1 \int_0^t Y_{t_1}^{m-2(R+1), \zeta} dt_1 \right) dt \\
& = Y_0^{m, \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T (\varepsilon + C\eta) \\
& \quad + C_1 T \left(Y_0^{m-(R+1), \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T (\varepsilon + C\eta) \right) + (C_1)^2 \int_0^T \int_0^t Y_{t_1}^{m-2(R+1), \zeta} dt_1 dt \\
& \leq Y_0^{m, \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T (\varepsilon + C\eta) + C_1 T \left(Y_0^{m-(R+1), \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T (\varepsilon + C\eta) \right) \\
& \quad + (C_1)^2 \int_0^T \int_0^t \left(Y_0^{m-2(R+1), \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T (\varepsilon + C\eta) + C_1 \int_0^{t_1} Y_{t_2}^{m-3(R+1), \zeta} dt_2 \right) dt_1 dt \\
& = Y_0^{m, \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T (\varepsilon + C\eta) + C_1 T \left(Y_0^{m-(R+1), \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T (\varepsilon + C\eta) \right) \\
& \quad + \frac{(C_1 T)^2}{2!} \left(Y_0^{m-2(R+1), \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T (\varepsilon + C\eta) \right) + (C_1)^3 \int_0^T \int_0^t \int_0^{t_1} Y_{t_2}^{m-3(R+1), \zeta} dt_2 dt_1 dt \\
& \leq \sum_{l=0}^{m/(R+1)-1} \frac{(C_1 T)^l}{l!} \left(Y_0^{m-(R+1)l, \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T (\varepsilon + C\eta) \right) + \frac{(C_1 T)^{m/(R+1)}}{(m/(R+1))!} Y_T^{0, \zeta} \\
& \leq \sum_{l=0}^{\lfloor m/(2R+2) \rfloor} \frac{(C_1 T)^l}{l!} \left(Y_0^{m-(R+1)l, \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T (\varepsilon + C\eta) \right) \\
& \quad + \sum_{l=\lfloor m/(2R+2) \rfloor + 1}^{m/(R+1)-1} \frac{(C_1 T)^l}{l!} \left(Y_0^{m-(R+1)l, \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T (\varepsilon + C\eta) \right) + \frac{(C_1 T)^{m/(R+1)}}{(m/(R+1))!} Y_T^{0, \zeta}
\end{aligned}$$

$$\begin{aligned}
&\leq \left(Y_0^{m-\lfloor m/2 \rfloor - (R+1), \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T(\varepsilon + C\eta) \right) \left(\sum_{l=0}^{\lfloor m/(2R+2) \rfloor} \frac{(C_1 T)^l}{l!} \right) \\
&\quad + \left(Y_0^{0, \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T(\varepsilon + C\eta) \right) \left(\sum_{l=\lfloor m/(2R+2) \rfloor + 1}^{\infty} \frac{(C_1 T)^l}{l!} \right) + \frac{(C_1 T)^{m/(R+1)}}{(m/(R+1))!} Y_T^{0, \zeta} \\
&\leq \left(Y_0^{m-\lfloor m/2 \rfloor - (R+1), \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T(\varepsilon + C\eta) \right) e^{C_1 T} \\
&\quad + \left(Y_0^{0, \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T(\varepsilon + C\eta) \right) \left(\sum_{l=\lfloor m/(2R+2) \rfloor + 1}^{\infty} \frac{(C_1 T)^l}{l!} \right) + \frac{(C_1 T)^{m/(R+1)}}{(m/(R+1))!} Y_T^{0, \zeta}.
\end{aligned}$$

(7.17)

From Eq (7.17), we're able to obtain

$$\begin{aligned}
&\lim_{m \rightarrow \infty} \limsup_{\zeta \rightarrow 0} \sup_{1 \leq j \leq J} \mathbb{E} \left[\sup_{t \in [0, T]} \left(\langle f_m, \mu_t^{\zeta, j} \rangle \right) \right] \\
&\leq \lim_{m \rightarrow \infty} \limsup_{\zeta \rightarrow 0} \left[\left(Y_0^{m-\lfloor m/2 \rfloor - (R+1), \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T(\varepsilon + C\eta) \right) e^{C_1 T} \right. \\
&\quad \left. + \left(Y_0^{0, \zeta} + C_3 \frac{1}{\sqrt{\gamma}} + C_2 T(\varepsilon + C\eta) \right) \left(\sum_{l=\lfloor m/(2R+2) \rfloor + 1}^{\infty} \frac{(C_1 T)^l}{l!} \right) + \frac{(C_1 T)^{m/(R+1)}}{(m/(R+1))!} Y_T^{0, \zeta} \right] \\
&= C_2 T e^{C_1 T} \varepsilon + \lim_{m \rightarrow \infty} \left[\tilde{Y}_0^{m-\lfloor m/2 \rfloor - (R+1)} e^{CT} + \left(\tilde{Y}_0^0 + C_2 T \varepsilon \right) \left(\sum_{l=\lfloor m/(2R+2) \rfloor + 1}^{\infty} \frac{(CT)^l}{l!} \right) + \frac{(CT)^{m/(R+1)}}{(m/(R+1))!} \tilde{Y}_T^0 \right] \\
&\quad \left(\text{where we denote } \tilde{Y}_t^l = \limsup_{\zeta \rightarrow 0} Y_t^{l, \zeta} \right) \\
&= C_2 T e^{C_1 T} \varepsilon.
\end{aligned}$$

(7.18)

Here, in the second last row of Eq (7.18), inside the squared bracket, the first term vanishes as

$$\begin{aligned}
\lim_{m \rightarrow \infty} \tilde{Y}_0^{m-\lfloor m/2 \rfloor - (R+1)} &= \lim_{m \rightarrow \infty} \limsup_{\zeta \rightarrow 0} \sup_{1 \leq j \leq J} \mathbb{E} \left[\left(\langle f_{m-\lfloor m/2 \rfloor - (R+1)}, \mu_0^{\zeta, j} \rangle \right) \right] \\
&= \sup_{1 \leq j \leq J} \lim_{m \rightarrow \infty} \limsup_{\zeta \rightarrow 0} \mathbb{E} \left[\left(\langle f_{m-\lfloor m/2 \rfloor - (R+1)}, \mu_0^{\zeta, j} \rangle \right) \right] \\
(7.19) \quad &= \sup_{1 \leq j \leq J} \lim_{m \rightarrow \infty} \mathbb{E} \langle f_{m-\lfloor m/2 \rfloor - (R+1)}, \xi_0^j \rangle = 0.
\end{aligned}$$

The exchange of limits and supremum is allowed as the supremum is taken over the finite set $\{1, \dots, J\}$. We can get the third line of Eq (7.19) from the second line because of Assumption 5.2 that the initial distribution $\mu_0^{\zeta, j}$ converges weakly to ξ_0^j , for all $1 \leq j \leq J$. Finally, we obtain the limit as zero using Assumption 5.2 on the ξ_0^j 's, i.e. that they are compactly supported. For the second term, \tilde{Y}_0^0 , the initial concentration is bounded, whereas $\left(\sum_{l=\lfloor m/4 \rfloor + 1}^{\infty} \frac{(CT)^l}{l!} \right)$ is the remainder of exponential function expansion, which will go to 0 as $m \rightarrow \infty$. For the last term, as $\tilde{Y}_T^0 = \limsup_{\zeta \rightarrow 0} \sup_{1 \leq j \leq J} \mathbb{E} \left[\sup_{t \in [0, T]} \left(\langle 1, \mu_t^{\zeta, j} \rangle \right) \right]$ is bounded by $C(\mu)$, the whole third term will vanish as $m \rightarrow \infty$.

Note, C_1 and C_2 do not depend on η , R or ε . This implies

$$\lim_{m \rightarrow \infty} \limsup_{\zeta \rightarrow 0} \mathbb{E} \left[\sup_{t \in [0, T]} \left(\langle f_m, \mu_t^{\zeta, j} \rangle \right) \right]$$

is less than an arbitrary small number $C_2 T e^{C_1 T} \varepsilon$, i.e. the limit is zero, for all $1 \leq j \leq J$. \square

Let $(\xi_t^1, \xi_t^2, \dots, \xi_t^J)$ denote the weak limit of a subsequence of $(\mu_t^{\zeta,1}, \mu_t^{\zeta,2}, \dots, \mu_t^{\zeta,J})$ in $\mathbb{D}_{M'_F(\mathbb{R}^{3d})}([0, T])$ as $\zeta \rightarrow 0$, where we abuse notation and let $(\mu_t^{\zeta,1}, \mu_t^{\zeta,2}, \dots, \mu_t^{\zeta,J})$ also denote the corresponding subsequence. Then

Lemma 7.6. $\{\xi_t^j\}_{t \in [0, T]}$ is continuous process from $[0, T]$ to both $M'_F(\mathbb{R}^d)$ and $M_F(\mathbb{R}^d)$ for each $j = 1, 2, \dots, J$.

Proof. By construction, see for example the proof of Lemma 7.1 and in particular (7.11)-(7.12), we have that

$$\sup_{t \in [0, T]} \sup_{f \in C_0^2(\mathbb{R}^n), \|f\|_{L^\infty} \leq 1} |\langle f, \mu_t^{\zeta, j} \rangle - \langle f, \mu_{t-}^{\zeta, j} \rangle| \leq \frac{C}{\gamma}$$

holds for some constant C independent of γ . In addition, by Proposition 5.3 in Chapter 3 of [EK86], the mapping $\nu \mapsto \sup_{t \in [0, T]} |\langle f, \nu_t \rangle - \langle f, \nu_{t-} \rangle|$ is continuous on $\mathbb{D}_{M'_F(\mathbb{R}^d)}([0, T])$ for each $f \in C_0^2(\mathbb{R}^d)$. Then, by Theorem 10.2 in Chapter 3 of [EK86], we obtain as we take $\zeta \rightarrow 0$, that $\{\xi_t^j\}_{t \in [0, T]}$ is continuous process from $[0, T]$ to $M'_F(\mathbb{R}^n)$. Next, we'll show that $\{\xi_t^j\}_{t \in [0, T]} \in \mathbb{D}_{M_F(\mathbb{R}^d)}([0, T])$, is a continuous process from $[0, T]$ to $M_F(\mathbb{R}^d)$. To this end, we need to be able to control what happens to the total mass of the measures (see also [BMW12]).

Adapting the notations in Lemma 7.5, let's define compactly supported functions $f_{m,r} = f_m(1 - f_r)$. Notice that $f_{m,r}$ will converge monotonically to f_m as $r \rightarrow \infty$. Then

$$(7.20) \quad \mathbb{E} \left(\sup_{t \in [0, T]} \langle f_{m,r}, \xi_t^j \rangle \right) = \lim_{\zeta \rightarrow 0} \mathbb{E} \left(\sup_{t \in [0, T]} \langle f_{m,r}, \mu_t^{\zeta, j} \rangle \right) \leq \liminf_{\zeta \rightarrow 0} \mathbb{E} \left(\sup_{t \in [0, T]} \langle f_m, \mu_t^{\zeta, j} \rangle \right) < \infty$$

by the continuity of the mapping $\nu \mapsto \sup_{t \in [0, T]} \langle f, \nu_t \rangle$, giving the first equality, and monotonicity of $f_{m,r} \leq f_m$, providing the second inequality.

From Eq (7.20), taking the limit $r \rightarrow \infty$ first and using monotone convergence theorem, we'll get

$$\mathbb{E} \left(\sup_{t \in [0, T]} \langle f_m, \xi_t^j \rangle \right) \leq \liminf_{\zeta \rightarrow 0} \mathbb{E} \left(\sup_{t \in [0, T]} \langle f_m, \mu_t^{\zeta, j} \rangle \right) < \infty$$

which implies

$$(7.21) \quad \mathbb{E} \left(\sup_{t \in [0, T]} \langle 1, \xi_t^j \rangle \right) < \infty.$$

Using Lemma 7.5, it follows that

$$(7.22) \quad \lim_{m \rightarrow \infty} \mathbb{E} \left(\sup_{t \in [0, T]} \langle f_m, \xi_t^j \rangle \right) = 0.$$

This gives that there exists subsequence of $\sup_{t \in [0, T]} \langle f_m, \xi_t^j \rangle$, $m \geq 0$, such that $\sup_{t \in [0, T]} \langle f_m, \xi_t^j \rangle \rightarrow 0$ almost surely, $\{\xi_t^j\}_{t \leq T}$ is a tight sequence almost surely and $\{\xi_t^j\}_{t \in [0, T]}$ is in $\mathcal{C}_{M'_F(\mathbb{R}^d)}([0, T])$. Due to the latter fact and due to (7.22), $\{\xi_t^j\}_{t \in [0, T]}$ is in $\mathcal{C}_{M_F(\mathbb{R}^d)}([0, T])$ as well. \square

Theorem 7.7 (Tightness). *The measure-valued process $\{\mu_t^{\zeta, j}\}_{t \in [0, T]}$ is tight in $\mathbb{D}_{M_F(\mathbb{R}^d)}[0, T]$, for each $j = 1, 2, \dots, J$.*

Proof. Referring to Meleard and Roelly [MS93], to prove tightness in $\mathbb{D}_{M_F(\mathbb{R}^d)}[0, T]$, it suffices to prove $\langle 1, \mu_t^{\zeta, j} \rangle$ converges in law to $\langle 1, \xi_t^j \rangle$ in $\mathbb{D}_{\mathbb{R}}([0, T])$, where note that ξ_t^j is the limit point of $\mu_t^{\zeta, j}$ in $\mathbb{D}_{M'_F(\mathbb{R}^d)}[0, T]$, which also lies in $\mathcal{C}_{M_F(\mathbb{R}^d)}([0, T])$ by Lemma 7.6.

Let F be any globally Lipschitz continuous bounded function from \mathbb{R} to \mathbb{R} . Then

$$\begin{aligned} & \limsup_{\zeta \rightarrow 0} |\mathbb{E}[F(\langle 1, \mu_t^{\zeta, j} \rangle)] - F(\langle 1, \xi_t^j \rangle)]| \\ & \leq \lim_{m \rightarrow \infty} \limsup_{\zeta \rightarrow 0} |\mathbb{E}[F(\langle 1, \mu_t^{\zeta, j} \rangle)] - F(\langle 1 - f_m, \mu_t^{\zeta, j} \rangle)]| + \lim_{m \rightarrow \infty} \limsup_{\zeta \rightarrow 0} |\mathbb{E}[F(\langle 1 - f_m, \mu_t^{\zeta, j} \rangle)] - F(\langle 1 - f_m, \xi_t^j \rangle)]| \\ & \quad + \lim_{m \rightarrow \infty} |\mathbb{E}[F(\langle 1 - f_m, \xi_t^j \rangle)] - F(\langle 1, \xi_t^j \rangle)]|, \end{aligned}$$

$$= 0, \quad (7.23)$$

where on the righthand side of Eq (7.23), the first and third term become 0 as a result of Lemma 7.5, while the second term vanishes due to the continuity of the mapping $\nu \mapsto \sup_{t \in [0, T]} \langle 1 - f_m, \nu_t \rangle$ by noting that $1 - f_m$ is compactly supported. \square

7.5. Uniqueness of Limiting Solution. We've established tightness of the measure-valued processes $\{\mu_t^{\zeta, j}\}_{t \in [0, T]}$, for all $1 \leq j \leq J$ (See Theorem 7.7). We now show that the limiting measure is unique.

For a measurable complete metric space E , $\nu \in M_F(E)$, define the norm $\|\cdot\|_{M_F(E)}$ on $M_F(E)$ as

$$\|\nu\|_{M_F(E)} = \sup_{f \in L^\infty(E), \|f\|_{L^\infty} \leq 1} |\langle f, \nu \rangle_E|,$$

which is the variation norm of finite measures. Using density argument, one can show that this is equivalent to (See step 4 of Theorem 3.2. of [BMW12])

$$\|\nu\|_{M_F(E)} = \sup_{f \in C_b^2(E), \|f\|_{L^\infty} \leq 1} |\langle f, \nu \rangle_E|.$$

For our purpose, we'll use test function $f \in C_b^2(E)$. The following two results then imply uniqueness:

Lemma 7.8. *Let $E = (\mathbb{R}^d)^n$ be a product space of \mathbb{R}^d , $n \geq 1$. Let $\mu^1, \dots, \mu^n \in M_F(\mathbb{R}^d)$, $\nu^1, \dots, \nu^n \in M_F(\mathbb{R}^d)$ and $\otimes_{i=1}^n \mu^i$, $\otimes_{i=1}^n \nu^i$ be product measures on E . Then*

$$\|\otimes_{i=1}^n \mu^i - \otimes_{i=1}^n \nu^i\|_{M_F(E)} \leq \sum_{i=1}^n (\|\mu^i - \nu^i\|_{M_F(\mathbb{R}^d)} \times \prod_{j=1}^{i-1} \langle 1, \mu^j \rangle \times \prod_{j=i+1}^n \langle 1, \nu^j \rangle).$$

Proof. For any $f \in L^\infty(E)$, $\|f\|_{L^\infty} \leq 1$, we have

$$\begin{aligned} |\langle f, \otimes_{i=1}^n \mu^i - \otimes_{i=1}^n \nu^i \rangle_E| &= \left| \left\langle f, \sum_{i=1}^n ((\otimes_{j=1}^{i-1} \mu^j) \otimes (\mu^i - \nu^i) \otimes (\otimes_{j=i+1}^n \nu^j)) \right\rangle_E \right|, \\ &\leq \sum_{i=1}^n |\langle f, ((\otimes_{j=1}^{i-1} \mu^j) \otimes (\mu^i - \nu^i) \otimes (\otimes_{j=i+1}^n \nu^j)) \rangle_E|, \\ (7.24) \quad &\leq \sum_{i=1}^n (\|\mu^i - \nu^i\|_{M_F(\mathbb{R}^d)} \times \prod_{j=1}^{i-1} \langle 1, \mu^j \rangle \times \prod_{j=i+1}^n \langle 1, \nu^j \rangle). \end{aligned}$$

The last inequality is due to the assumption that $\|f\|_{L^\infty(E)} \leq 1$ and using the definition of signed measure norms.

Since Eq (7.24) is true for all $f \in L^\infty(E)$, $\|f\|_{L^\infty} \leq 1$, Lemma 7.8 is proved. \square

Corollary 7.9. *Let $E = (\mathbb{R}^d)^n$ be a product space of \mathbb{R}^d , $n \geq 1$. Let $\mu^1, \dots, \mu^n \in M_F(\mathbb{R}^d)$, $\nu^1, \dots, \nu^n \in M_F(\mathbb{R}^d)$ and $\otimes_{i=1}^n \mu^i$, $\otimes_{i=1}^n \nu^i$ be product measures on E . If there exists $M > 0$, such that $|\langle 1, \mu^i \rangle| \leq M$ and $|\langle 1, \nu^i \rangle| \leq M$ for all $1 \leq i \leq n$, then*

$$\|\otimes_{i=1}^n \mu^i - \otimes_{i=1}^n \nu^i\|_{M_F(E)} \leq M^{n-1} \sum_{i=1}^n \|\mu^i - \nu^i\|_{M_F(\mathbb{R}^d)}.$$

Proof. This is a consequence of Lemma 7.8 using the fact that $\langle 1, \mu^i \rangle$ or $\langle 1, \nu^i \rangle$ are uniform bounded by M . \square

Lemma 7.10 (Uniqueness). *The solution to (5.1) is unique in $\mathcal{C}_{M_F(\mathbb{R}^d)}([0, T])$.*

Proof. Suppose we have two different set of solutions to (5.1), $\{(\xi_t^1, \xi_t^2, \dots, \xi_t^J)\}_{t \in [0, T]}$ and $\{(\bar{\xi}_t^1, \bar{\xi}_t^2, \dots, \bar{\xi}_t^J)\}_{t \in [0, T]}$, with the same initial condition $(\xi_0^1, \xi_0^2, \dots, \xi_0^J) = (\bar{\xi}_0^1, \bar{\xi}_0^2, \dots, \bar{\xi}_0^J)$. In Eq (5.1), if we use a test function of the form of $\psi_t(x) \in C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^d)$, it becomes

$$\langle \psi_t, \xi_t^j \rangle = \langle \psi_0, \xi_0^j \rangle + \int_0^t \langle \partial_s \psi_s + (\mathcal{L}_j \psi_s)(x), \xi_s^j(dx) \rangle ds$$

$$(7.25) \quad + \sum_{\ell=1}^L \int_0^t \int_{\bar{\mathbb{X}}^{(\ell)}} \frac{1}{\alpha^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} \psi_s(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} - \sum_{r=1}^{\alpha_{\ell j}} \psi_s(x_r^{(j)}) \right) \lambda^{(\ell)}[\xi_s](d\mathbf{x}) ds.$$

Let $\mathcal{P}_{j,t}, t \geq 0$, be the semigroup generated by $\mathcal{L}_j, j = 1, 2, \dots, J$. Choose $\psi_s(x) = \mathcal{P}_{j,t-s}f(x)$, respectively for each $1 \leq j \leq J$, where $f \in C_b^2(\mathbb{R}^d), \|f\|_{L^\infty} \leq 1$, then Eq (7.25) becomes

$$(7.26) \quad \begin{aligned} \langle f, \xi_t^j \rangle &= \langle \mathcal{P}_{j,t}f, \xi_0^j \rangle + \sum_{\ell=1}^L \int_0^t \int_{\bar{\mathbb{X}}^{(\ell)}} \frac{1}{\alpha^{(\ell)}!} \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} \mathcal{P}_{\ell,t-s}f(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} - \sum_{r=1}^{\alpha_{\ell j}} \mathcal{P}_{\ell,t-s}f(x_r^{(j)}) \right) \\ &\times K_\ell(\mathbf{x}) \lambda^{(\ell)}[\xi_s](d\mathbf{x}) ds. \end{aligned}$$

From Eq (7.26), we obtain the following estimates for $\langle f, \xi_t^j - \bar{\xi}_t^j \rangle$. By (7.21) we have that

$$M = 1 \vee \sup_{\{t \in [0, T], j=1, 2, \dots, J\}} |\langle 1, \xi_t^j \rangle| \vee |\langle 1, \bar{\xi}_t^j \rangle| < \infty,$$

which then gives

$$(7.27) \quad \begin{aligned} |\langle f, \xi_t^j - \bar{\xi}_t^j \rangle| &\leq \sum_{\ell=1}^L \int_0^t \int_{\bar{\mathbb{X}}^{(\ell)}} \frac{1}{\alpha^{(\ell)}!} \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} \mathcal{P}_{\ell,t-s}f(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} - \sum_{r=1}^{\alpha_{\ell j}} \mathcal{P}_{\ell,t-s}f(x_r^{(j)}) \right) \\ &\times K_\ell(\mathbf{x}) \left(\lambda^{(\ell)}[\xi_s](d\mathbf{x}) - \lambda^{(\ell)}[\bar{\xi}_s](d\mathbf{x}) \right) | ds, \\ &\leq C(K) \sum_{\ell=1}^L \frac{\alpha_{\ell j} + \beta_{\ell j}}{\alpha^{(\ell)}!} \int_0^t \|\lambda^{(\ell)}[\xi_s] - \lambda^{(\ell)}[\bar{\xi}_s]\|_{M_F(\mathbb{X}^{(\ell)})} ds, \\ &\leq C(K) \sum_{\ell=1}^L \frac{\alpha_{\ell j} + \beta_{\ell j}}{\alpha^{(\ell)}!} \int_0^t \|\otimes_{i=1}^J (\otimes_{r=1}^{\alpha_{\ell i}} \xi_s^i) - \otimes_{i=1}^J (\otimes_{r=1}^{\alpha_{\ell i}} \bar{\xi}_s^i)\|_{M_F(\mathbb{X}^{(\ell)})} ds, \\ &\leq C(K) \sum_{\ell=1}^L \frac{\alpha_{\ell j} + \beta_{\ell j}}{\alpha^{(\ell)}!} \int_0^t M^{|\alpha^{(\ell)}|-1} \sum_{i=1, \dots, J} \alpha_{\ell i} \|\xi_s^i - \bar{\xi}_s^i\|_{M_F(\mathbb{R}^d)} ds, \end{aligned}$$

where the last inequality is due to Corollary 7.9 since $\langle 1, \xi_s^i \rangle$ or $\langle 1, \bar{\xi}_s^i \rangle$ are uniformly bounded by M for all $1 \leq i \leq J$. In the second to the last equality of Eq (7.27), we have used the following estimates.

$$(7.28) \quad \begin{aligned} & \left| \frac{1}{\alpha^{(\ell)}!} \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} \mathcal{P}_{\ell,t-s}f(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} - \sum_{r=1}^{\alpha_{\ell j}} \mathcal{P}_{\ell,t-s}f(x_r^{(j)}) \right) K_\ell(\mathbf{x}) \right| \\ & \leq \left| \frac{1}{\alpha^{(\ell)}!} \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} |\mathcal{P}_{\ell,t-s}f(y_r^{(j)})| \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} + \sum_{r=1}^{\alpha_{\ell j}} |\mathcal{P}_{\ell,t-s}f(x_r^{(j)})| \right) K_\ell(\mathbf{x}) \right| \\ & \leq \frac{\alpha_{\ell j} + \beta_{\ell j}}{\alpha^{(\ell)}!} C(K). \end{aligned}$$

(here we use the fact that $\|\mathcal{P}_t f\|_{L^\infty} \leq 1$ and $\int_{\mathbb{Y}^{(\ell)}} m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} = 1$)

Based on Eq (7.27), we obtain

$$\begin{aligned} \sum_{j=1}^J \|\xi_t^j - \bar{\xi}_t^j\|_{M_F(\mathbb{R}^d)} &\leq C(K) \sum_{\ell=1}^L \sum_{j=1}^J \frac{\alpha_{\ell j} + \beta_{\ell j}}{\alpha^{(\ell)}!} \int_0^t M^{|\alpha^{(\ell)}|-1} \sum_{i=1}^J \alpha_{\ell i} \|\xi_s^i - \bar{\xi}_s^i\|_{M_F(\mathbb{R}^d)} ds, \\ &\leq C(K) L J \max_{1 \leq \ell \leq L, 1 \leq j \leq J} \left\{ \frac{\alpha_{\ell j} + \beta_{\ell j}}{\alpha^{(\ell)}!} M^{|\alpha^{(\ell)}|-1} \alpha_{\ell j} \right\} \int_0^t \sum_{i=1}^J \|\xi_s^i - \bar{\xi}_s^i\|_{M_F(\mathbb{R}^d)} ds, \end{aligned}$$

Applying Gronwall's inequality, we get $\sum_{j=1}^J \|\xi_t^j - \bar{\xi}_t^j\|_{M_F(\mathbb{R}^d)} = 0$ for all $t \in [0, T]$, which proves the uniqueness of solution, concluding the proof of the lemma. \square

ACKNOWLEDGEMENTS

SAI thanks Romain Yvinec for first introducing him to the MVSP representation for particle systems and the work of [BM15].

A γ DEPENDENCY OF REACTION KERNELS

In this section we demonstrate one way in which the claimed γ scaling given in Assumption 4.8 arises for bimolecular reactions. We ignore the zeroth order case as our main result, Theorem 5.3, does not allow for such reactions. In the first order case the reaction rate kernel, $K^\gamma(\mathbf{x})$, is usually interpreted as an internal property of molecules, giving the probability per time an individual reactant particle can undergo the reaction. As such, it would not be expected to depend on γ . In contrast, the reaction rate kernel for a bimolecular reaction is often calibrated to agree with a known well-mixed reaction rate constant in the limit that the system is forced to be well-mixed (i.e. the limit that particle diffusivities are taken to be infinite), which ultimately gives rise to the γ dependence.

Consider an isolated system containing only two particles that can undergo a bimolecular annihilation reaction of the form $A + B \rightarrow \emptyset$. Assume we are considering the reaction within a bounded domain $\Omega \subset \mathbb{R}^d$ with hypervolume $|\Omega|$. In modeling chemical reaction systems, one is often given a spatially-homogeneous, well-mixed, macroscopic reaction rate, K_{wm} , with units of (molar concentration) $^{1-\alpha}$ (time) $^{-1}$ for a reaction of order α . The corresponding reaction rate used in a spatially-homogeneous well-mixed stochastic chemical kinetics model is then $\bar{K} = (\gamma |\Omega|)^{1-\alpha} K_{\text{wm}}$, with units of (time) $^{-1}$. For our second order reaction \bar{K} gives the probability per time for the pair of A and B molecules to react and annihilate in the well-mixed stochastic model.

Consider the PBSRD model's dynamics until the two reactants annihilate. Let $p(\mathbf{x}, \mathbf{y}, t)$ denote the probability density the particle positions are \mathbf{x} and \mathbf{y} respectively at time t , and no reaction has yet occurred. Then

$$\frac{\partial p}{\partial t}(\mathbf{x}, \mathbf{y}, t) = (D^A \Delta_{\mathbf{x}} + D^B \Delta_{\mathbf{y}})p(\mathbf{x}, \mathbf{y}, t) - K^\gamma(\mathbf{x}, \mathbf{y})p(\mathbf{x}, \mathbf{y}, t), \quad \mathbf{x} \in \Omega, \mathbf{y} \in \Omega, t > 0,$$

with reflecting no-flux boundary conditions for \mathbf{x} or \mathbf{y} in $\partial\Omega$. In the formal well-mixed limit that the particle diffusivities are taken to be infinite, we expect that $p(\mathbf{x}, \mathbf{y}, t) = p(t)$. Letting $P(t) = p(t) |\Omega|^2$ denote the probability the reaction has not yet occurred, we then have

$$\frac{dP}{dt} = - \left(\frac{1}{|\Omega|^2} \int_{\Omega^2} K^\gamma(\mathbf{x}, \mathbf{y}) d\mathbf{x} d\mathbf{y} \right) P(t).$$

To match the well-mixed stochastic model we would then require that

$$\frac{1}{|\Omega|^2} \int_{\Omega^2} K^\gamma(\mathbf{x}, \mathbf{y}) d\mathbf{x} d\mathbf{y} = \bar{K} = \frac{K_{\text{wm}}}{\gamma |\Omega|}.$$

If we assume that $K^\gamma(\mathbf{x}, \mathbf{y}) = \gamma^\beta K(\mathbf{x}, \mathbf{y})$, then we immediately obtain the scaling given in Assumption 4.8, i.e. $\beta = -1$.

More concretely, consider the widely used Doi interaction $K^\gamma(\mathbf{x}, \mathbf{y}) = \lambda 1_{[0, \varepsilon]}(|\mathbf{x} - \mathbf{y}|)$ [D76a, D76b]. We find that

$$\lambda = \frac{K_{\text{wm}} |\Omega|}{\gamma |\mathcal{R} \cap \Omega^2|},$$

where $\mathcal{R} = \{(\mathbf{x}, \mathbf{y}) \in \mathbb{R}^{2d} \mid |\mathbf{x} - \mathbf{y}| \leq \varepsilon\}$. As $\Omega \rightarrow \mathbb{R}^d$ we formally expect

$$\lambda \rightarrow \frac{K_{\text{wm}}}{\gamma |B_\varepsilon|},$$

where $|B_\varepsilon|$ denotes the hypervolume of the ball of radius ε . This demonstrates that the scaling of Assumption 4.8 persists in freespace.

B MASS CONTROL LEMMAS FOR DIFFERENT CASES OF REACTIONS

The goal of this section is to prove the following key estimate.

Lemma B.1. *For the ℓ -th reaction, $1 \leq \ell \leq L$, let η small enough, $\varepsilon > 0$ and $R \in \mathbb{N}$ as in Assumption 4.7. Then, the following estimates holds for m large enough,*

$$\begin{aligned} & \mathbb{E} \left[\sup_{t \in [0, T]} \int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\alpha^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell^\eta(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s-}^\zeta](d\mathbf{x}) ds \right] \\ & \leq 2C(K)(C(\mu) \vee 1) \sup_{1 \leq i \leq J} \mathbb{E} \left[\sup_{t \in [0, T]} \int_0^t \left\langle f_{m-1-R}(x), \mu_{s-}^{\zeta, i}(dx) \right\rangle ds \right] + 2C(K)C(\mu) \|f_m\|_{C_b^2(\mathbb{R}^d)} T (C\eta + \varepsilon). \end{aligned}$$

To do so we first prove some intermediate results and the proof of Lemma B.1 will then follow at the end of this section.

Lemma B.2. *For any $\eta \geq 0$ small enough, $\tilde{L} + 1 \leq \ell \leq L$, $\mathbf{y} \in \mathbb{Y}^{(\ell)}$, $\mathbf{x} \in \mathbb{X}^{(\ell)}$, and $f \in C_b^2(\mathbb{Y}^{(\ell)})$, there exists a constant $C\eta$ such that*

$$\left| \int_{\mathbb{Y}^{(\ell)}} f(\mathbf{y}) (m_\ell^\eta(\mathbf{y} | \mathbf{x}) - m_\ell(\mathbf{y} | \mathbf{x})) d\mathbf{y} \right| \leq \|f\|_{C_b^2(\mathbb{Y}^{(\ell)})} C\eta,$$

and this constant $C\eta \rightarrow 0$ as $\eta \rightarrow 0$ by Definition 4.1 on the choice of positive mollifier and placement measures based on Assumptions 4.3 - 4.7.

Proof. This is essentially a result from the definition of mollifiers. We'll discuss this estimate for each of the following different cases of reactions. The upper bound is always some constant of the order of η times $\|f\|_{C_b^1(\mathbb{Y}^{(\ell)})}$.

Case 1: Reaction of the form $S_i \rightarrow S_j$.

Plugging in the definitions of $m_\ell^\eta(\mathbf{y} | \mathbf{x})$ and $m_\ell(\mathbf{y} | \mathbf{x})$ from Assumption 4.3, we will get

$$\begin{aligned} \left| \int_{\mathbb{Y}^{(\ell)}} f(\mathbf{y}) (m_\ell^\eta(\mathbf{y} | \mathbf{x}) - m_\ell(\mathbf{y} | \mathbf{x})) d\mathbf{y} \right| &= \left| \int_{\mathbb{R}^d} f(y) G_\eta(y - x) dy - f(x) \right| \\ &= \left| \int_{\mathbb{R}^d} (f(y) - f(x)) G_\eta(y - x) dy \right| \\ &\leq \left| \int_{B(x, \eta)} |f(y) - f(x)| G_\eta(y - x) dy \right| \\ &\leq \left| \int_{B(x, \eta)} \|f\|_{C_b^1(\mathbb{R}^d)} \eta \times G_\eta(y - x) dy \right| \\ &\leq \|f\|_{C_b^1(\mathbb{R}^d)} \eta. \end{aligned}$$

Case 2: Reaction of the form $S_i \rightarrow S_j + S_k$.

Plugging in the definitions of $m_\ell^\eta(\mathbf{y} | \mathbf{x})$ and $m_\ell(\mathbf{y} | \mathbf{x})$ from Assumption 4.6, we will get

$$\begin{aligned} & \left| \int_{\mathbb{Y}^{(\ell)}} f(\mathbf{y}) (m_\ell^\eta(\mathbf{y} | \mathbf{x}) - m_\ell(\mathbf{y} | \mathbf{x})) d\mathbf{y} \right| \\ &= \left| \sum_{i=1}^I p_i \times \left[\int_{\mathbb{R}^{2d}} f(y_1, y_2) \rho(|y_1 - y_2|) G_\eta(x - (\alpha_i y_1 + (1 - \alpha_i) y_2)) dy_1 dy_2 \right. \right. \\ & \quad \left. \left. - \int_{\mathbb{R}^{2d}} f(y_1, y_2) \rho(|y_1 - y_2|) \delta(x - (\alpha_i y_1 + (1 - \alpha_i) y_2)) dy_1 dy_2 \right] \right| \quad (\text{Let } w = y_1 - y_2) \\ &= \left| \sum_{i=1}^I p_i \times \left[\int_{\mathbb{R}^{2d}} f(w + y_2, y_2) \rho(|w|) G_\eta(x - \alpha_i w - y_2) dw dy_2 \right. \right. \\ & \quad \left. \left. - \int_{\mathbb{R}^{2d}} f(w + y_2, y_2) \rho(|w|) \delta(x - \alpha_i w - y_2) dw dy_2 \right] \right| \end{aligned}$$

$$\begin{aligned}
&= \left| \sum_{i=1}^I p_i \times \left[\int_{\mathbb{R}^d} \rho(|w|) \left(\int_{\mathbb{R}^d} (f(w + y_2, y_2) - f(w + x - \alpha_i w, x - \alpha_i w)) G_\eta(x - \alpha_i w - y_2) dy_2 \right) dw \right] \right| \\
&\leq \left| \sum_{i=1}^I p_i \times \left[\int_{\mathbb{R}^d} \rho(|w|) \left(\int_{B(x - \alpha_i w, \eta)} |f(w + y_2, y_2) - f(w + x - \alpha_i w, x - \alpha_i w)| \times G_\eta(x - \alpha_i w - y_2) dy_2 \right) dw \right] \right| \\
&\leq \left| \sum_{i=1}^I p_i \times \left[\int_{\mathbb{R}^d} \rho(|w|) \left(\int_{B(x - \alpha_i w, \eta)} \|f\|_{C^1(\mathbb{R}^{2d})} \eta \times G_\eta(x - \alpha_i w - y_2) dy_2 \right) dw \right] \right| \\
&\leq \|f\|_{C^1(\mathbb{R}^{2d})} \eta \quad (\text{by noting that } \int_{\mathbb{R}^d} \rho(|w|) dw = 1 \text{ in Assumption 4.7.})
\end{aligned}$$

Case 3: Reaction of the form $S_i + S_k \rightarrow S_j$.

Plugging in the definitions of $m_\ell^\eta(\mathbf{y} | \mathbf{x})$ and $m_\ell(\mathbf{y} | \mathbf{x})$ from Assumption 4.4, we will get

$$\begin{aligned}
& \left| \int_{\mathbb{Y}^{(\ell)}} f(\mathbf{y}) (m_\ell^\eta(\mathbf{y} | \mathbf{x}) - m_\ell(\mathbf{y} | \mathbf{x})) d\mathbf{y} \right| \\
&= \left| \sum_{i=1}^I p_i \times \left[\int_{\mathbb{R}^d} f(y) G_\eta(y - (\alpha_i x_1 + (1 - \alpha_i) x_2)) dy - f(\alpha_i x_1 + (1 - \alpha_i) x_2) \right] \right| \\
&= \left| \sum_{i=1}^I p_i \times \int_{\mathbb{R}^d} (f(y) - f(\alpha_i x_1 + (1 - \alpha_i) x_2)) G_\eta(y - (\alpha_i x_1 + (1 - \alpha_i) x_2)) dy \right| \\
&\leq \left| \sum_{i=1}^I p_i \times \int_{B(\alpha_i x_1 + (1 - \alpha_i) x_2, \eta)} |f(y) - f(\alpha_i x_1 + (1 - \alpha_i) x_2)| G_\eta(y - (\alpha_i x_1 + (1 - \alpha_i) x_2)) dy \right| \\
&\leq \left| \sum_{i=1}^I p_i \times \int_{B(\alpha_i x_1 + (1 - \alpha_i) x_2, \eta)} \|f\|_{C_b^1(\mathbb{R}^d)} \eta \times G_\eta(y - (\alpha_i x_1 + (1 - \alpha_i) x_2)) dy \right| \\
&\leq \|f\|_{C_b^1(\mathbb{R}^d)} \eta
\end{aligned}$$

Case 4: Reaction of the form $S_i + S_k \rightarrow S_j + S_r$.

Plugging in the definitions of $m_\ell^\eta(\mathbf{y} | \mathbf{x})$ and $m_\ell(\mathbf{y} | \mathbf{x})$ from Assumption 4.5, we will get

$$\begin{aligned}
& \left| \int_{\mathbb{Y}^{(\ell)}} f(\mathbf{y}) (m_\ell^\eta(\mathbf{y} | \mathbf{x}) - m_\ell(\mathbf{y} | \mathbf{x})) d\mathbf{y} \right| \\
&= \left| p \times \left[\int_{\mathbb{R}^{2d}} f(y_1, y_2) G_\eta(y_1 - x_1) G_\eta(y_2 - x_2) dy_1 dy_2 - f(x_1, x_2) \right] \right. \\
&\quad \left. + (1 - p) \times \left[\int_{\mathbb{R}^{2d}} f(y_1, y_2) G_\eta(y_2 - x_1) G_\eta(y_1 - x_2) dy_1 dy_2 - f(x_2, x_1) \right] \right| \\
&= \left| p \times \int_{\mathbb{R}^{2d}} (f(y_1, y_2) - f(x_1, x_2)) G_\eta(y_1 - x_1) G_\eta(y_2 - x_2) dy_1 dy_2 \right. \\
&\quad \left. + (1 - p) \times \int_{\mathbb{R}^{2d}} (f(y_1, y_2) - f(x_2, x_1)) G_\eta(y_2 - x_1) G_\eta(y_1 - x_2) dy_1 dy_2 \right| \\
&\leq \left| p \times \int_{B((x_1, x_2), \sqrt{2}\eta)} |f(y_1, y_2) - f(x_1, x_2)| G_\eta(y_1 - x_1) G_\eta(y_2 - x_2) dy_1 dy_2 \right. \\
&\quad \left. + (1 - p) \times \int_{B((x_2, x_1), \sqrt{2}\eta)} |f(y_1, y_2) - f(x_2, x_1)| G_\eta(y_2 - x_1) G_\eta(y_1 - x_2) dy_1 dy_2 \right| \\
&\leq \left| p \times \int_{B((x_1, x_2), \sqrt{2}\eta)} \|f\|_{C_b^1(\mathbb{R}^{2d})} \times \sqrt{2}\eta \times G_\eta(y_1 - x_1) G_\eta(y_2 - x_2) dy_1 dy_2 \right. \\
&\quad \left. + (1 - p) \times \int_{B((x_2, x_1), \sqrt{2}\eta)} \|f\|_{C_b^1(\mathbb{R}^{2d})} \times \sqrt{2}\eta \times G_\eta(y_2 - x_1) G_\eta(y_1 - x_2) dy_1 dy_2 \right| \\
&\leq \sqrt{2} \|f\|_{C_b^1(\mathbb{R}^{2d})} \eta.
\end{aligned}$$

□

Lemma B.3. *If the ℓ -th reaction is a reaction of the form $S_i \rightarrow S_j$, then*

(B.1)

$$\int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s^-}^\zeta](d\mathbf{x}) ds \leq C(K) \int_0^t \langle f_m(x), \mu_{s^-}^{\zeta, i}(dx) \rangle ds$$

Proof. By plugging in the specific form of the reaction rate and placement measure as in Assumption 4.3, we have

$$\begin{aligned} & \int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s^-}^\zeta](d\mathbf{x}) ds \\ &= \int_0^t \int_{\mathbb{R}^d} K_\ell(x) \left(\int_{\mathbb{R}^d} f_m(y) \delta_x(y) dy \right) \mu_{s^-}^{\zeta, i}(dx) ds \\ &= \int_0^t \int_{\mathbb{R}^d} K_\ell(x) f_m(x) \mu_{s^-}^{\zeta, i}(dx) ds \\ &\leq C(K) \int_0^t \langle f_m(x), \mu_{s^-}^{\zeta, i}(dx) \rangle ds. \end{aligned}$$

□

Lemma B.4. *If the ℓ -th reaction is a reaction of the form $S_i \rightarrow S_j + S_k$, where i and k could be j , then for the choice of $\varepsilon \geq 0$ and $R \in \mathbb{N}$ in Assumption 4.7, we have for m large enough,*

$$\begin{aligned} & \int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s^-}^\zeta](d\mathbf{x}) ds \\ &\leq 2C(K) \int_0^t \langle f_{m-1-R}(x), \mu_{s^-}^{\zeta, i}(dx) \rangle ds + 2C(K) \|f_m\|_{C_b^2(\mathbb{R}^d)} C(\mu) t \varepsilon. \end{aligned}$$

Proof. Let $\varepsilon \geq 0$ and $R \in \mathbb{N}$ be such that Assumption 4.7 is satisfied. By plugging in the specific form of the reaction rate and placement measure as in Assumption 4.6, we have

$$\begin{aligned} & \int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s^-}^\zeta](d\mathbf{x}) ds \\ &\leq \int_0^t \int_{\mathbb{R}^d} K_\ell(x) \left(\int_{\mathbb{R}^{2d}} (f_m(y) + f_m(z)) m_\ell(y, z | x) dy dz \right) \mu_{s^-}^{\zeta, i}(dx) ds \\ &= \int_0^t \int_{\mathbb{R}^d} K_\ell(x) \left(\int_{\mathbb{R}^{2d}} (f_m(y) + f_m(z)) \rho(|y-z|) \sum_{i=1}^I p_i \delta(x - (\alpha_i y + (1-\alpha_i)z)) dy dz \right) \mu_{s^-}^{\zeta, i}(dx) ds \\ &\leq C(K) \int_0^t \left\langle \left(\int_{|y-z| \leq R} (f_m(y) + f_m(z)) \rho(|y-z|) \sum_{i=1}^I p_i \delta(x - (\alpha_i y + (1-\alpha_i)z)) dy dz \right), \mu_{s^-}^{\zeta, i}(dx) \right\rangle ds. \\ &\quad + C(K) \int_0^t \left\langle \left(\int_{|y-z| > R} (f_m(y) + f_m(z)) \rho(|y-z|) \sum_{i=1}^I p_i \delta(x - (\alpha_i y + (1-\alpha_i)z)) dy dz \right), \mu_{s^-}^{\zeta, i}(dx) \right\rangle ds. \\ &\leq C(K) \int_0^t \left\langle \left(\int_{|y-z| \leq R} 2f_{m-1-R}(x) \rho(|y-z|) \sum_{i=1}^I p_i \delta(x - (\alpha_i y + (1-\alpha_i)z)) dy dz \right), \mu_{s^-}^{\zeta, i}(dx) \right\rangle ds. \\ &\quad + 2C(K) \|f_m\|_{C_b^2(\mathbb{R}^d)} \int_0^t \left\langle \sum_{i=1}^I p_i \left(\int_{|\eta| > R} \int_{z \in \mathbb{R}^d} \rho(|\eta|) \delta(x - \alpha_i \eta - z) dz d\eta \right), \mu_{s^-}^{\zeta, i}(dx) \right\rangle ds. \\ &\leq 2C(K) \int_0^t \left\langle f_{m-1-R}(x) \left(\int_{|y-z| \leq R} m_\ell(y, z | x) dy dz \right), \mu_{s^-}^{\zeta, i}(dx) \right\rangle ds. \end{aligned}$$

$$\begin{aligned}
& + C(K) \|f_m\|_{C_b^2(\mathbb{R}^d)} \int_0^t \left\langle \sum_{i=1}^I p_i \left(\int_{|\eta|>R} \rho(|\eta|) d\eta \right), \mu_{s-}^{\zeta,i}(dx) \right\rangle ds. \\
& \leq 2C(K) \int_0^t \left\langle f_{m-1-R}(x), \mu_{s-}^{\zeta,i}(dx) \right\rangle ds + 2C(K) \|f_m\|_{C_b^2(\mathbb{R}^d)} C(\mu) t \varepsilon.
\end{aligned}$$

□

Lemma B.5. *If the ℓ -th reaction is a reaction of the form $S_i + S_k \rightarrow S_j$, then*

$$\begin{aligned}
& \int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s-}^{\zeta}](d\mathbf{x}) ds \\
& \leq C(K) C(\mu) \int_0^t \left(\left\langle f_{m-1}(x), \mu_{s-}^{\zeta,i}(dx) \right\rangle + \left\langle f_{m-1}(y), \mu_{s-}^{\zeta,k}(dy) \right\rangle \right) ds.
\end{aligned}$$

Proof. By plugging in the specific form of the reaction rate and placement measure as in Assumption 4.4, we have

$$\begin{aligned}
& \int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s-}^{\zeta}](d\mathbf{x}) ds \\
& \leq \int_0^t \left\langle \left\langle K_\ell(x, y) \left(\int_{\mathbb{R}^d} f_m(z) m_\ell(z | x, y) dz \right), \mu_{s-}^{\zeta,i}(dx) \right\rangle, \mu_{s-}^{\zeta,k}(dy) \right\rangle ds. \\
& \leq C(K) \int_0^t \left\langle \left\langle \left(\int_{\mathbb{R}^d} f_m(z) \sum_{i=1}^I p_i \delta_{\alpha_i x + (1-\alpha_i)y}(z) dz \right), \mu_{s-}^{\zeta,i}(dx) \right\rangle, \mu_{s-}^{\zeta,k}(dy) \right\rangle ds. \\
& \leq C(K) \int_0^t \left\langle \left\langle \sum_{i=1}^I p_i (f_{m-1}(x) + f_{m-1}(y)), \mu_{s-}^{\zeta,i}(dx) \right\rangle, \mu_{s-}^{\zeta,k}(dy) \right\rangle ds. \\
& \leq C(K) C(\mu) \int_0^t \left(\left\langle f_{m-1}(x), \mu_{s-}^{\zeta,i}(dx) \right\rangle + \left\langle f_{m-1}(y), \mu_{s-}^{\zeta,k}(dy) \right\rangle \right) ds.
\end{aligned}$$

□

Lemma B.6. *If the ℓ -th reaction is a reaction of the form $S_i + S_k \rightarrow S_j + S_r$, where i, k, r could be j , then*

$$\begin{aligned}
& \int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s-}^{\zeta}](d\mathbf{x}) ds \\
& \leq C(K) C(\mu) \int_0^t \left(\left\langle f_m(x), \mu_{s-}^{\zeta,i}(dx) \right\rangle + \left\langle f_m(y), \mu_{s-}^{\zeta,k}(dy) \right\rangle \right) ds.
\end{aligned}$$

Proof. By plugging in the specific form of the reaction rate and placement measure as in Assumption 4.5, we have

$$\begin{aligned}
& \int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s-}^{\zeta}](d\mathbf{x}) ds \\
& \leq \int_0^t \left\langle \left\langle K_\ell(x, y) \left(\int_{\mathbb{R}^{2d}} (f_m(z) + f_m(w)) m_\ell(z, w | x, y) dz dw \right), \mu_{s-}^{\zeta,i}(dx) \right\rangle, \mu_{s-}^{\zeta,k}(dy) \right\rangle ds \\
& \leq C(K) \int_0^t \left\langle \left\langle \left(\int_{\mathbb{R}^{2d}} (f_m(z) + f_m(w)) (p \times \delta_{(x,y)}((z, w)) + (1-p) \times \delta_{(x,y)}((w, z))) dz dw \right), \mu_{s-}^{\zeta,i}(dx) \right\rangle, \mu_{s-}^{\zeta,k}(dy) \right\rangle ds \\
& \leq C(K) \int_0^t \left\langle \left\langle (f_m(x) + f_m(y)), \mu_{s-}^{\zeta,i}(dx) \right\rangle, \mu_{s-}^{\zeta,k}(dy) \right\rangle ds. \\
& \leq C(K) C(\mu) \int_0^t \left(\left\langle f_m(x), \mu_{s-}^{\zeta,i}(dx) \right\rangle + \left\langle f_m(y), \mu_{s-}^{\zeta,k}(dy) \right\rangle \right) ds.
\end{aligned}$$

□

Now we are in position to give the proof of Lemma B.1.

Proof of Lemma B.1. We have that

$$\begin{aligned}
& \mathbb{E} \left[\sup_{t \in [0, T]} \int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell^\eta(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s-}^\zeta](d\mathbf{x}) ds \right] \\
&= \mathbb{E} \left[\sup_{t \in [0, T]} \int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s-}^\zeta](d\mathbf{x}) ds \right] \\
&\quad + \mathbb{E} \left[\sup_{t \in [0, T]} \int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) (m_\ell^\eta(\mathbf{y} | \mathbf{x}) - m_\ell(\mathbf{y} | \mathbf{x})) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s-}^\zeta](d\mathbf{x}) ds \right] \\
&\leq \mathbb{E} \left[\sup_{t \in [0, T]} \int_0^t \int_{\tilde{\mathbb{X}}^{(\ell)}} \frac{1}{\boldsymbol{\alpha}^{(\ell)}!} K_\ell(\mathbf{x}) \left(\int_{\mathbb{Y}^{(\ell)}} \left(\sum_{r=1}^{\beta_{\ell j}} f_m(y_r^{(j)}) \right) m_\ell(\mathbf{y} | \mathbf{x}) d\mathbf{y} \right) \lambda^{(\ell)}[\mu_{s-}^\zeta](d\mathbf{x}) ds \right] \\
&\quad + 2C(K)C(\mu) \|f_m\|_{C_b^2(\mathbb{R}^d)} C\eta T \quad (\text{by Lemma B.2}) \\
&\leq 2C(K)(C(\mu) \vee 1) \sup_{1 \leq i \leq J} \mathbb{E} \left[\sup_{t \in [0, T]} \int_0^t \langle f_{m-1-R}(x), \mu_{s-}^{\zeta, i}(dx) \rangle ds \right] \\
&\quad + 2C(K) \|f_m\|_{C_b^2(\mathbb{R}^d)} C(\mu) T \varepsilon + 2C(K)C(\mu) \|f_m\|_{C_b^2(\mathbb{R}^d)} C\eta T \quad (\text{by Lemma B.3 - B.6}) \\
&= 2C(K)(C(\mu) \vee 1) \sup_{1 \leq i \leq J} \mathbb{E} \left[\sup_{t \in [0, T]} \int_0^t \langle f_{m-1-R}(x), \mu_{s-}^{\zeta, i}(dx) \rangle ds \right] + 2C(K)C(\mu) \|f_m\|_{C_b^2(\mathbb{R}^d)} T (C\eta + \varepsilon),
\end{aligned}$$

concluding the proof of the lemma. \square

C PROOFS OF PROPOSITIONS 6.5 AND 6.6.

Let us recall the forward equation (6.2). For both proofs of Propositions 6.5 and 6.6 we need to define an appropriate L^2 space. In particular, define an appropriate L^2 (Fock) space, F , with inner product for two functions, $\mathbf{G}_1 = \{g_1^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c)\}_{a,b,c=0}^\infty$ and $\mathbf{G}_2 = \{g_2^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c)\}_{a,b,c=0}^\infty$, as

$$(C.1) \quad (\mathbf{G}_1, \mathbf{G}_2)_F = \sum_{a=0}^\infty \sum_{b=0}^\infty \sum_{c=0}^\infty \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} g_1^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c) g_2^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c) d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c,$$

we can interpret $\mathcal{T}^* = \mathcal{L} + \mathcal{R}^+ + \mathcal{R}^-$ as the adjoint of the generator, \mathcal{T} , for the process

$$(\mathbf{Q}^{A(t)}(t), \mathbf{Q}^{B(t)}(t), \mathbf{Q}^{C(t)}(t), A(t), B(t), C(t)).$$

Formally, we find

$$\begin{aligned}
(C.2) \quad (\mathcal{T}\mathbf{G})_{a,b,c}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c) &= (\mathcal{L}\mathbf{G})_{a,b,c}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c) \\
&\quad + \sum_{l=1}^a \sum_{m=1}^b \left(\int_{\mathbb{R}^d} m_1^\eta(\mathbf{z} | \mathbf{q}_l^a, \mathbf{q}_m^b) K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) g^{(a-1, b-1, c+1)}(\mathbf{q}^a \setminus \mathbf{q}_l^a, \mathbf{q}^b \setminus \mathbf{q}_m^b, \mathbf{q}^c \cup \mathbf{z}) d\mathbf{z} \right. \\
&\quad \left. - K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c) \right) \\
&\quad + \sum_{n=1}^c \left(\int_{\mathbb{R}^{2d}} m_2^\eta(\mathbf{x}, \mathbf{y} | \mathbf{q}_n^c) K_2^\gamma(\mathbf{q}_n^c) g^{(a+1, b+1, c-1)}(\mathbf{q}^a \cup \mathbf{x}, \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t) d\mathbf{x} d\mathbf{y} \right. \\
&\quad \left. - K_2^\gamma(\mathbf{q}_n^c) g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c) \right)
\end{aligned}$$

Note that here $\mathcal{L}^* = \mathcal{L}$.

Next we present the proofs of the two propositions.

Proof of Proposition 6.5. For simplicity of notation, without loss of generality, we will show the equivalence for the evolution of $\mathbb{E}[\varphi(\langle f, \nu_t^\zeta \rangle_{\hat{P}})]$. The same procedure follows for the more general multi-dimensional

case $\mathbb{E}[\varphi(\langle f_1, \nu_t^\xi \rangle_{\hat{P}}, \langle f_2, \nu_t^\xi \rangle_{\hat{P}}, \dots, \langle f_M, \nu_t^\xi \rangle_{\hat{P}})]$. By the definition of ν_t^ξ and adopting the notation of this section,

$$\begin{aligned}
\mathbb{E}[\varphi(\langle f, \nu_t^\xi \rangle_{\hat{P}})] &= \mathbb{E}[\varphi\left(\sum_{i=1}^{A(t)} f(\mathbf{Q}_i^{A(t)}(t), S_1) + \sum_{j=1}^{B(t)} f(\mathbf{Q}_j^{B(t)}(t), S_2) + \sum_{k=1}^{C(t)} f(\mathbf{Q}_k^{C(t)}(t), S_3)\right)] \\
&= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} \varphi\left(\sum_{i=1}^a f(\mathbf{q}_i^a, S_1) + \sum_{j=1}^b f(\mathbf{q}_j^b, S_2) + \sum_{k=1}^c f(\mathbf{q}_k^c, S_3)\right) \\
&\quad \times p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c, \\
\text{(C.3)} \quad &= (\mathbf{G}, \mathbf{P}(t))_F,
\end{aligned}$$

where we define $g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c) = \varphi\left(\sum_{i=1}^a f(\mathbf{q}_i^a, S_1) + \sum_{j=1}^b f(\mathbf{q}_j^b, S_2) + \sum_{k=1}^c f(\mathbf{q}_k^c, S_3)\right)$. For such a form of $g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c)$, plugging into Eq (C.2), we have

$$\begin{aligned}
(\mathcal{T}\mathbf{G})_{a,b,c}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c) &= \left(D_1 \sum_{l=1}^a \Delta_{\mathbf{q}_l^a} + D_2 \sum_{m=1}^b \Delta_{\mathbf{q}_m^b} + D_3 \sum_{n=1}^c \Delta_{\mathbf{q}_n^c}\right) \varphi\left(\sum_{i=1}^a f(\mathbf{q}_i^a, S_1) + \sum_{j=1}^b f(\mathbf{q}_j^b, S_2) + \sum_{k=1}^c f(\mathbf{q}_k^c, S_3)\right) \\
&+ \sum_{l=1}^a \sum_{m=1}^b \left(\int_{\mathbb{R}^d} \left[g^{(a-1,b-1,c+1)}(\mathbf{q}^a \setminus \mathbf{q}_l^a, \mathbf{q}^b \setminus \mathbf{q}_m^b, \mathbf{q}^c \cup \mathbf{z}) - g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c)\right] \right. \\
&\quad \left. \times m_1^\eta(\mathbf{z} | \mathbf{q}_l^a, \mathbf{q}_m^b) \times K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) d\mathbf{z}\right) \\
&+ \sum_{n=1}^c \left(\int_{\mathbb{R}^{2d}} \left[g^{(a+1,b+1,c-1)}(\mathbf{q}^a \cup \mathbf{x}, \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t) - g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c)\right] \right. \\
&\quad \left. \times m_2^\eta(\mathbf{x}, \mathbf{y} | \mathbf{q}_n^c) \times K_2^\gamma(\mathbf{q}_n^c) d\mathbf{x} d\mathbf{y}\right) \\
&= D_1 \sum_{i=1}^a \varphi'' \left(\sum_{i=1}^a f(\mathbf{q}_i^a, S_1) + \sum_{j=1}^b f(\mathbf{q}_j^b, S_2) + \sum_{k=1}^c f(\mathbf{q}_k^c, S_3)\right) |\nabla_{\mathbf{q}_i^a} f_1(\mathbf{q}_i^a)|^2 \\
&+ D_1 \sum_{i=1}^a \varphi' \left(\sum_{i=1}^a f(\mathbf{q}_i^a, S_1) + \sum_{j=1}^b f(\mathbf{q}_j^b, S_2) + \sum_{k=1}^c f(\mathbf{q}_k^c, S_3)\right) \Delta_{\mathbf{q}_i^a} f_1(\mathbf{q}_i^a) \\
&+ D_2 \sum_{j=1}^b \varphi'' \left(\sum_{i=1}^a f(\mathbf{q}_i^a, S_1) + \sum_{j=1}^b f(\mathbf{q}_j^b, S_2) + \sum_{k=1}^c f(\mathbf{q}_k^c, S_3)\right) |\nabla_{\mathbf{q}_j^b} f_2(\mathbf{q}_j^b)|^2 \\
&+ D_2 \sum_{j=1}^b \varphi' \left(\sum_{i=1}^a f(\mathbf{q}_i^a, S_1) + \sum_{j=1}^b f(\mathbf{q}_j^b, S_2) + \sum_{k=1}^c f(\mathbf{q}_k^c, S_3)\right) \Delta_{\mathbf{q}_j^b} f_2(\mathbf{q}_j^b) \\
&+ D_3 \sum_{k=1}^c \varphi'' \left(\sum_{i=1}^a f(\mathbf{q}_i^a, S_1) + \sum_{j=1}^b f(\mathbf{q}_j^b, S_2) + \sum_{k=1}^c f(\mathbf{q}_k^c, S_3)\right) |\nabla_{\mathbf{q}_k^c} f_3(\mathbf{q}_k^c)|^2 \\
&+ D_3 \sum_{k=1}^c \varphi' \left(\sum_{i=1}^a f(\mathbf{q}_i^a, S_1) + \sum_{j=1}^b f(\mathbf{q}_j^b, S_2) + \sum_{k=1}^c f(\mathbf{q}_k^c, S_3)\right) \Delta_{\mathbf{q}_k^c} f_3(\mathbf{q}_k^c) \\
&+ \sum_{l=1}^a \sum_{m=1}^b \left(\int_{\mathbb{R}^d} \left[\varphi\left(\sum_{i=1}^a f(\mathbf{q}_i^a, S_1) + \sum_{j=1}^b f(\mathbf{q}_j^b, S_2) + \sum_{k=1}^c f(\mathbf{q}_k^c, S_3) - f(\mathbf{q}_l^a, S_1) - f(\mathbf{q}_m^b, S_2) + f(\mathbf{z}, S_3)\right) \right. \right. \\
&\quad \left. \left. - \varphi\left(\sum_{i=1}^a f(\mathbf{q}_i^a, S_1) + \sum_{j=1}^b f(\mathbf{q}_j^b, S_2) + \sum_{k=1}^c f(\mathbf{q}_k^c, S_3)\right)\right] \times m_1^\eta(\mathbf{z} | \mathbf{q}_l^a, \mathbf{q}_m^b) \times K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) d\mathbf{z}\right)
\end{aligned}$$

$$\begin{aligned}
& + \sum_{n=1}^c \left(\int_{\mathbb{R}^{2d}} \left[\varphi \left(\sum_{i=1}^a f(\mathbf{q}_i^a, S_1) + \sum_{j=1}^b f(\mathbf{q}_j^b, S_2) + \sum_{k=1}^c f(\mathbf{q}_k^c, S_3) + f(\mathbf{x}, S_1) + f(\mathbf{y}, S_2) - f(\mathbf{q}_n^c, S_3) \right) \right. \right. \\
& \quad \left. \left. - \varphi \left(\sum_{i=1}^a f(\mathbf{q}_i^a, S_1) + \sum_{j=1}^b f(\mathbf{q}_j^b, S_2) + \sum_{k=1}^c f(\mathbf{q}_k^c, S_3) \right) \right] \times m_2^\eta(\mathbf{x}, \mathbf{y} | \mathbf{q}_n^c) \times K_2^\gamma(\mathbf{q}_n^c) d\mathbf{x} d\mathbf{y} \right) \\
\end{aligned} \tag{C.4}$$

Rewriting $\mathbb{E}[\varphi(\langle f, \nu_t^\xi \rangle_{\hat{P}})]$ in Eq (C.3) as an integral equation, we obtain

$$\begin{aligned}
\mathbb{E}[\varphi(\langle f, \nu_t^\xi \rangle_{\hat{P}})] &= (\mathbf{G}, \mathbf{P}(t))_F = (\mathbf{G}, \mathbf{P}(0))_F + \int_0^t \partial_s (\mathbf{G}, \mathbf{P}(s))_F ds \\
&= \mathbb{E}[\varphi(\langle f, \nu_0^\xi \rangle_{\hat{P}})] + \int_0^t (\mathbf{G}, \partial_s \mathbf{P}(s))_F ds \\
&= \mathbb{E}[\varphi(\langle f, \nu_0^\xi \rangle_{\hat{P}})] + \int_0^t (\mathbf{G}, \mathcal{T}^* \mathbf{P}(s))_F ds \quad (\text{by Eq (6.2)}) \\
&= \mathbb{E}[\varphi(\langle f, \nu_0^\xi \rangle_{\hat{P}})] + \int_0^t (\mathcal{T} \mathbf{G}, \mathbf{P}(s))_F ds \\
&= \mathbb{E}[\varphi(\langle f, \nu_0^\xi \rangle_{\hat{P}})] \\
&+ \mathbb{E} \left[\int_0^t \varphi'(\langle f, \nu_s^\xi \rangle_{\hat{P}}) \times \left[D_1 \sum_{i=1}^{A(s)} \Delta_{\mathbf{Q}_i} f(\mathbf{Q}_i^{A(s)}(s), S_1) + D_2 \sum_{j=1}^{B(s)} \Delta_{\mathbf{Q}_j} f(\mathbf{Q}_j^{B(s)}(s), S_2) \right. \right. \\
& \quad \left. \left. + D_3 \sum_{k=1}^{C(s)} \Delta_{\mathbf{Q}_k} f(\mathbf{Q}_k^{C(s)}(s), S_3) \right] + \varphi''(\langle f, \nu_s^\xi \rangle_{\hat{P}}) \times \left[D_1 \sum_{i=1}^{A(s)} |\nabla_{\mathbf{Q}_i} f(\mathbf{Q}_i^{A(s)}(s), S_1)|^2 \right. \right. \\
& \quad \left. \left. + D_2 \sum_{j=1}^{B(s)} |\nabla_{\mathbf{Q}_j} f(\mathbf{Q}_j^{B(s)}(s), S_2)|^2 + D_3 \sum_{k=1}^{C(s)} |\nabla_{\mathbf{Q}_k} f(\mathbf{Q}_k^{C(s)}(s), S_3)|^2 \right] ds \right] \\
&+ \mathbb{E} \left[\int_0^t \sum_{l=1}^{A(s)} \sum_{m=1}^{B(s)} \left(\int_{\mathbb{R}^d} K_1^\gamma(\mathbf{Q}_l^{A(s)}(s), \mathbf{Q}_m^{B(s)}(s)) m_1^\eta(\mathbf{z} | \mathbf{Q}_l^{A(s)}(s), \mathbf{Q}_m^{B(s)}(s)) \right. \right. \\
& \quad \left. \left. \times \left(\varphi(\langle f, \nu_s^\xi \rangle_{\hat{P}}) - f(\mathbf{Q}_l^{A(s)}(s), S_1) - f(\mathbf{Q}_m^{B(s)}(s), S_2) + f(\mathbf{z}, S_3) \right) - \varphi(\langle f, \nu_s^\xi \rangle_{\hat{P}}) \right) d\mathbf{z} \right] ds \\
&+ \mathbb{E} \left[\int_0^t \sum_{n=1}^{C(s)} \left(\int_{\mathbb{R}^{2d}} K_2^\gamma(\mathbf{Q}_n^{C(s)}(s)) m_2^\eta(\mathbf{x}, \mathbf{y} | \mathbf{Q}_n^{C(s)}(s)) \right. \right. \\
& \quad \left. \left. \times \left(\varphi(\langle f, \nu_s^\xi \rangle_{\hat{P}}) + f(\mathbf{x}, S_1) + f(\mathbf{y}, S_2) - f(\mathbf{Q}_n^{C(s)}(s), S_3) \right) - \varphi(\langle f, \nu_s^\xi \rangle_{\hat{P}}) \right) d\mathbf{x} d\mathbf{y} \right] ds. \\
\end{aligned} \tag{C.5}$$

From the measure-valued formulation in Eq (5.5), we obtain the following integral equation for $\mathbb{E}[\varphi(\langle f, \nu_t^\xi \rangle_{\hat{P}})]$ by applying Itô's formula on Eq (5.5),

$$\begin{aligned}
\mathbb{E}[\varphi(\langle f, \nu_t^\xi \rangle_{\hat{P}})] &= \mathbb{E}[\varphi(\langle f, \nu_0^\xi \rangle_{\hat{P}})] + \mathbb{E} \left[\int_0^t \varphi'(\langle f, \nu_s^\xi \rangle_{\hat{P}}) \sum_{i=1}^{\langle 1, \nu_{s-}^\xi \rangle_{\hat{P}}} D^i \frac{\partial^2 f}{\partial Q^2} (H^i(\nu_{s-}^\xi)) ds \right] \\
&+ \frac{1}{2} \mathbb{E} \left[\int_0^t \varphi''(\langle f, \nu_s^\xi \rangle_{\hat{P}}) \sum_{i=1}^{\langle 1, \nu_{s-}^\xi \rangle_{\hat{P}}} \left(\sqrt{2D^i} \frac{\partial f}{\partial Q} (H^i(\nu_{s-}^\xi)) \right)^2 ds \right] \\
&+ \mathbb{E} \left[\int_0^t \int_{(\mathbb{N} \setminus \{0\})^2} \int_{\mathbb{R}^d} \int_{\mathbb{R}_+^2} \varphi(\langle f, \nu_s^\xi \rangle_{\hat{P}}) + \left(-f(H_Q^i(\nu_{s-}^{\xi,1}), S_1) - f(H_Q^j(\nu_{s-}^{\xi,2}), S_2) + f(\mathbf{z}, S_3) \right) \right. \\
\end{aligned}$$

$$\begin{aligned}
& \times \mathbf{1}_{\{i \leq \langle 1, \nu_{s-}^{\zeta,1} \rangle\}} \times \mathbf{1}_{\{j \leq \langle 1, \nu_{s-}^{\zeta,2} \rangle\}} \times \mathbf{1}_{\{\theta_1 \leq K_1^\gamma(H_Q^i(\nu_{s-}^{\zeta,1}), H_Q^j(\nu_{s-}^{\zeta,2}))\}} \times \mathbf{1}_{\{\theta_2 \leq m_1^\eta(z|H_Q^i(\nu_{s-}^{\zeta,1}), H_Q^j(\nu_{s-}^{\zeta,2}))\}} \\
& - \varphi(\langle f, \nu_s^\zeta \rangle_{\hat{P}}) dN_1(s, i, j, z, \theta_1, \theta_2)] \\
& + \mathbb{E} \left[\int_0^t \int_{\mathbb{N} \setminus \{0\}} \int_{\mathbb{R}^{2d}} \int_{\mathbb{R}_+^2} \varphi(\langle f, \nu_s^\zeta \rangle_{\hat{P}} + (f(x, S_1) + f(y, S_2) - f(H_Q^k(\nu_{s-}^{\zeta,3}), S_3)) \right. \\
& \quad \left. \times \mathbf{1}_{\{k \leq \langle 1, \nu_{s-}^{\zeta,3} \rangle\}} \times \mathbf{1}_{\{\theta_1 \leq K_2^\gamma(H_Q^k(\nu_{s-}^{\zeta,3}))\}} \times \mathbf{1}_{\{\theta_2 \leq m_2^\eta(x, y|H_Q^k(\nu_{s-}^{\zeta,3}))\}} \right) \\
& - \varphi(\langle f, \nu_s^\zeta \rangle_{\hat{P}}) dN_2(s, k, x, y, \theta_1, \theta_2)] \\
& = \mathbb{E}[\varphi(\langle f, \nu_0^\zeta \rangle_{\hat{P}})] + \mathbb{E} \left[\int_0^t \varphi'(\langle f, \nu_s^\zeta \rangle_{\hat{P}}) \sum_{i=1}^{\langle 1, \nu_{s-}^\zeta \rangle_{\hat{P}}} D^i \frac{\partial^2 f}{\partial Q^2}(H^i(\nu_{s-}^\zeta)) ds \right] \\
& + \mathbb{E} \left[\int_0^t \varphi''(\langle f, \nu_s^\zeta \rangle_{\hat{P}}) \sum_{i=1}^{\langle 1, \nu_{s-}^\zeta \rangle_{\hat{P}}} D^i \left(\frac{\partial f}{\partial Q}(H^i(\nu_{s-}^\zeta)) \right)^2 ds \right] \\
& + \mathbb{E} \left[\int_0^t \sum_{i=1}^{\langle 1, \nu_{s-}^{\zeta,1} \rangle} \sum_{j=1}^{\langle 1, \nu_{s-}^{\zeta,2} \rangle} \int_{\mathbb{R}^d} K_1^\gamma(H_Q^i(\nu_{s-}^{\zeta,1}), H_Q^j(\nu_{s-}^{\zeta,2})) \times m_1^\eta(z|H_Q^i(\nu_{s-}^{\zeta,1}), H_Q^j(\nu_{s-}^{\zeta,2})) \right. \\
& \quad \left. \times \left[\varphi(\langle f, \nu_s^\zeta \rangle_{\hat{P}} + (-f(H_Q^i(\nu_{s-}^{\zeta,1}), S_1) - f(H_Q^j(\nu_{s-}^{\zeta,2}), S_2) + f(z, S_3))) - \varphi(\langle f, \nu_s^\zeta \rangle_{\hat{P}}) \right] dz ds \right] \\
& + \mathbb{E} \left[\int_0^t \sum_{k=1}^{\langle 1, \nu_{s-}^{\zeta,3} \rangle} \int_{\mathbb{R}^{2d}} K_2^\gamma(H_Q^k(\nu_{s-}^{\zeta,3})) \times m_2^\eta(x, y|H_Q^k(\nu_{s-}^{\zeta,3})) \right. \\
& \quad \left. \times \left[\varphi(\langle f, \nu_s^\zeta \rangle_{\hat{P}} + (f(x, S_1) + f(y, S_2) - f(H_Q^k(\nu_{s-}^{\zeta,3}), S_3))) - \varphi(\langle f, \nu_s^\zeta \rangle_{\hat{P}}) \right] dx dy ds \right]. \tag{C.6}
\end{aligned}$$

We observe that the integral equation for $\mathbb{E}[\varphi(\langle f, \nu_s^\zeta \rangle_{\hat{P}})]$ is the same when derived from the forward equation (C.5) and from the measure-valued formulation (C.6). \square

Proof of Proposition 6.6. We use the forward Kolmogorov equation to prove the proposition. We are interested in finding an equation for the average concentration field for A, B and C molecules, i.e. $\mathbb{E}[A(\mathbf{x}, t)]$, $\mathbb{E}[B(\mathbf{y}, t)]$ and $\mathbb{E}[C(\mathbf{z}, t)]$ from forward equation. This is defined by

$$\begin{aligned}
\mathbb{E}[A(\mathbf{x}, t)] &= \mathbb{E} \left[\sum_{i=1}^{A(t)} \delta(\mathbf{Q}_i^{A(t)}(t) - \mathbf{x}) \right] \\
&= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \sum_{i=1}^a \int_{\mathbb{R}^{(a+b+c)d}} \delta(\mathbf{q}_i^a - \mathbf{x}) p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c \\
&= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! b! c!} \int_{\mathbb{R}^{(a-1+b+c)d}} p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^{a-1} d\mathbf{q}^b d\mathbf{q}^c.
\end{aligned}$$

Similarly, for molecule B and C, we have

$$\begin{aligned}
\mathbb{E}[B(\mathbf{y}, t)] &= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! (b-1)! c!} \int_{\mathbb{R}^{(a+b-1+c)d}} p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^{b-1} \cup \mathbf{y}, \mathbf{q}^c, t) d\mathbf{q}^a d\mathbf{q}^{b-1} d\mathbf{q}^c. \\
\mathbb{E}[C(\mathbf{z}, t)] &= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! (c-1)!} \int_{\mathbb{R}^{(a+b+c-1)d}} p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^{c-1} \cup \mathbf{z}, t) d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^{c-1}.
\end{aligned}$$

In deriving this equation we will need to use the correlation in the A and B fields, given by

$$\mathbb{E}[A(\mathbf{x}, t)B(\mathbf{y}, t)] = \mathbb{E} \left[\sum_{i=1}^{A(t)} \sum_{j=1}^{B(t)} \delta(\mathbf{Q}_i^{A(t)}(t) - \mathbf{x}) \delta(\mathbf{Q}_j^{B(t)}(t) - \mathbf{y}) \right]$$

$$\begin{aligned}
&= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \sum_{i=1}^a \sum_{j=1}^b \int_{\mathbb{R}^{(a+b)d}} \delta(\mathbf{q}_i^a - \mathbf{x}) \delta(\mathbf{q}_j^b - \mathbf{y}) p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c \\
&= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! (b-1)! c!} \int_{\mathbb{R}^{(a+b+c-2)d}} p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^{b-1} \cup \mathbf{y}, \mathbf{q}^c, t) d\mathbf{q}^{a-1} d\mathbf{q}^{b-1} d\mathbf{q}^c.
\end{aligned}$$

Using these definitions, and assuming the probability densities vanish at infinity, we find that

$$\begin{aligned}
\partial_t \mathbb{E}[A(\mathbf{x}, t)] &= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! b! c!} \int_{\mathbb{R}^{(a-1+b+c)d}} ((\mathcal{L} + \mathcal{R}^+ + \mathcal{R}^-) \mathbf{P})_{a,b,c}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^{a-1} d\mathbf{q}^b d\mathbf{q}^c. \\
&= \text{(I)} + \text{(II)} + \text{(III)},
\end{aligned}$$

where

$$\begin{aligned}
\text{(I)} &= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! b! c!} \int_{\mathbb{R}^{(a-1+b+c)d}} (\mathcal{L} \mathbf{P})_{a,b,c}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^{a-1} d\mathbf{q}^b d\mathbf{q}^c \\
&= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! b! c!} \int_{\mathbb{R}^{(a-1+b+c)d}} \left[\left(D_1 \sum_{l=1}^{a-1} \Delta_{\mathbf{q}_l^a} + D_2 \sum_{m=1}^b \Delta_{\mathbf{q}_m^b} + D_3 \sum_{n=1}^c \Delta_{\mathbf{q}_n^c} \right) p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^c, t) \right. \\
&\quad \left. + D_1 \Delta_{\mathbf{x}} p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^c, t) \right] d\mathbf{q}^{a-1} d\mathbf{q}^b d\mathbf{q}^c \\
&= D_1 \Delta_{\mathbf{x}} \mathbb{E}[A(\mathbf{x}, t)]
\end{aligned}$$

where on the second to last line, the first term becomes zero due to integration by parts and the fact that probability density vanishes at infinity (recall that by Theorem 6.4 $p \in \mathcal{C}([0, \infty); H^2(X))$). Similarly,

$$\begin{aligned}
\text{(II)} &= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! b! c!} \int_{\mathbb{R}^{(a-1+b+c)d}} (\mathcal{R}^+ \mathbf{P})_{a,b,c}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^{a-1} d\mathbf{q}^b d\mathbf{q}^c. \\
&= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! b! c!} \int_{\mathbb{R}^{(a-1+b+c)d}} - \sum_{m=1}^b K_1^\gamma(\mathbf{x}, \mathbf{q}_m^b) p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^{a-1} d\mathbf{q}^b d\mathbf{q}^c. \\
&\quad + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! b! c!} \int_{\mathbb{R}^{(a-1+b+c)d}} \left\{ - \sum_{l=1}^{a-1} \sum_{m=1}^b K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^c, t) \right. \\
&\quad \left. + \sum_{n=1}^c \left[\int_{\mathbb{R}^{2d}} m_1(\mathbf{q}_n^c | \boldsymbol{\xi}, \boldsymbol{\eta}) K_1^\gamma(\boldsymbol{\xi}, \boldsymbol{\eta}) p^{(a+1,b+1,c-1)}(\mathbf{q}^{a-1} \cup \mathbf{x} \cup \boldsymbol{\xi}, \mathbf{q}^b \cup \boldsymbol{\eta}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t) d\boldsymbol{\xi} d\boldsymbol{\eta} \right] \right\} d\mathbf{q}^{a-1} d\mathbf{q}^b d\mathbf{q}^c. \\
&= - \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! b! c!} \int_{\mathbb{R}^{(a+b+c-2)d}} \left[b \times \int_{\mathbb{R}^d} K_1^\gamma(\mathbf{x}, \mathbf{y}) p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^{b-1} \cup \mathbf{y}, \mathbf{q}^c, t) d\mathbf{y} \right] d\mathbf{q}^{a-1} d\mathbf{q}^{b-1} d\mathbf{q}^c. \\
&\quad + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! b! c!} \left\{ - \int_{\mathbb{R}^{(a-1+b+c)d}} \sum_{l=1}^{a-1} \sum_{m=1}^b K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^{a-1} d\mathbf{q}^b d\mathbf{q}^c \right. \\
&\quad \left. + \int_{\mathbb{R}^{(a+b+c)d}} \int_{\mathbb{R}^d} \frac{c}{a(b+1)} \sum_{l=1}^a \sum_{m=1}^{b+1} m_1(\mathbf{z} | \mathbf{q}_l^a, \mathbf{q}_m^b) K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) \times \right. \\
&\quad \left. p^{(a+1,b+1,c-1)}(\mathbf{q}^a \cup \mathbf{x}, \mathbf{q}^{b+1}, \mathbf{q}^{c-1}, t) d\mathbf{z} d\mathbf{q}^a d\mathbf{q}^{b+1} d\mathbf{q}^{c-1} \right\} \\
&= - \int_{\mathbb{R}^d} K_1^\gamma(\mathbf{x}, \mathbf{y}) \mathbb{E}[A(\mathbf{x}, t) B(\mathbf{y}, t)] d\mathbf{y} \\
&\quad - \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! b! c!} \int_{\mathbb{R}^{(a-1+b+c)d}} \sum_{l=1}^{a-1} \sum_{m=1}^b K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^{a-1} d\mathbf{q}^b d\mathbf{q}^c \\
&\quad + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! (b+1)! (c-1)!} \int_{\mathbb{R}^{(a+b+c)d}} \sum_{l=1}^a \sum_{m=1}^{b+1} K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) p^{(a+1,b+1,c-1)}(\mathbf{q}^a \cup \mathbf{x}, \mathbf{q}^{b+1}, \mathbf{q}^{c-1}, t) d\mathbf{q}^a d\mathbf{q}^{b+1} d\mathbf{q}^{c-1}
\end{aligned}$$

$$= - \int_{\mathbb{R}^d} K_1^\gamma(\mathbf{x}, \mathbf{y}) \mathbb{E}[A(\mathbf{x}, t)B(\mathbf{y}, t)] d\mathbf{y}.$$

In the third equality of (II), we exchanged the orders of integrals and sums using that

$$\begin{aligned} & \int_{\mathbb{R}^{bd}} \sum_{m=1}^b K_1^\gamma(\mathbf{x}, \mathbf{q}_m^b) p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^b \\ &= \sum_{m=1}^b \int_{\mathbb{R}^{bd}} K_1^\gamma(\mathbf{x}, \mathbf{q}_m^b) p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, (\mathbf{q}^b \setminus \mathbf{q}_m^b) \cup \mathbf{q}_m^b, \mathbf{q}^c, t) d\mathbf{q}_m^b d(\mathbf{q}^b \setminus \mathbf{q}_m^b) \\ &= \sum_{m=1}^b \int_{\mathbb{R}^{bd}} K_1^\gamma(\mathbf{x}, \mathbf{y}) p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^{b-1} \cup \mathbf{y}, \mathbf{q}^c, t) d\mathbf{y} d\mathbf{q}^{b-1} \end{aligned}$$

(Here we replace \mathbf{q}_m^b by \mathbf{y} and $\mathbf{q}^b \setminus \mathbf{q}_m^b$ by \mathbf{q}^{b-1} due to that particles of the same type are indistinguishable)

$$= b \times \int_{\mathbb{R}^{bd}} K_1^\gamma(\mathbf{x}, \mathbf{y}) p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^{b-1} \cup \mathbf{y}, \mathbf{q}^c, t) d\mathbf{y} d\mathbf{q}^{b-1}.$$

Similar ideas also apply to the third term and to deriving (III). In the second to last equality of (II), we used that $\int_{\mathbb{R}^d} m_1(\mathbf{z}|\mathbf{x}, \mathbf{y}) d\mathbf{z} = 1$ and the second and third term cancel by shifting indexes. Similarly

$$\begin{aligned} \text{(III)} &= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! b! c!} \int_{\mathbb{R}^{(a-1+b+c)d}} (\mathcal{R}^- \mathbf{P})_{a,b,c}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^{a-1} d\mathbf{q}^b d\mathbf{q}^c \\ &= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! b! c!} \int_{\mathbb{R}^{(a-1+b+c)d}} \left\{ - \sum_{n=1}^c K_2^\gamma(\mathbf{q}_n^c) p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^c, t) \right. \\ &\quad \left. + \sum_{m=1}^b \left[\int_{\mathbb{R}^d} m_2(\mathbf{x}, \mathbf{q}_m^b | \mathbf{z}) K_2^\gamma(\mathbf{z}) p^{(a-1,b-1,c+1)}(\mathbf{q}^{a-1} \setminus \mathbf{q}_m^b \cup \mathbf{x}, \mathbf{q}^b \setminus \mathbf{q}_m^b, \mathbf{q}^c \cup \mathbf{z}, t) d\mathbf{z} \right] \right. \\ &\quad \left. + \sum_{m=1}^b \sum_{l=1}^{a-1} \left[\int_{\mathbb{R}^d} m_2(\mathbf{q}_l^{a-1}, \mathbf{q}_m^b | \mathbf{z}) K_2^\gamma(\mathbf{z}) p^{(a-1,b-1,c+1)}(\mathbf{q}^{a-1} \setminus \mathbf{q}_l^{a-1} \cup \mathbf{x}, \mathbf{q}^b \setminus \mathbf{q}_m^b, \mathbf{q}^c \cup \mathbf{z}, t) d\mathbf{z} \right] \right\} d\mathbf{q}^{a-1} d\mathbf{q}^b d\mathbf{q}^c \\ &= - \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! b! c!} \int_{\mathbb{R}^{(a+b+c-2)d}} c \int_{\mathbb{R}^d} K_2^\gamma(\mathbf{z}) p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^{c-1} \cup \mathbf{z}, t) d\mathbf{z} d\mathbf{q}^{a-1} d\mathbf{q}^b d\mathbf{q}^{c-1} \\ &\quad + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! b! c!} \left\{ \int_{\mathbb{R}^{(a+b+c-2)d}} b \left[\int_{\mathbb{R}^{2d}} m_2(\mathbf{x}, \mathbf{y} | \mathbf{z}) K_2^\gamma(\mathbf{z}) p^{(a-1,b-1,c+1)}(\mathbf{q}^{a-1}, \mathbf{q}^{b-1}, \mathbf{q}^c \cup \mathbf{z}, t) d\mathbf{y} d\mathbf{z} \right] \right. \\ &\quad \left. d\mathbf{q}^{a-1} d\mathbf{q}^{b-1} d\mathbf{q}^c + \int_{\mathbb{R}^{(a+b+c-3)d}} (a-1)b \right. \\ &\quad \left. \times \left[\int_{\mathbb{R}^{3d}} m_2(\boldsymbol{\xi}, \boldsymbol{\eta} | \mathbf{z}) K_2^\gamma(\mathbf{z}) p^{(a-1,b-1,c+1)}(\mathbf{q}^{a-2} \cup \mathbf{x}, \mathbf{q}^{b-1}, \mathbf{q}^c \cup \mathbf{z}, t) d\boldsymbol{\xi} d\boldsymbol{\eta} d\mathbf{z} \right] d\mathbf{q}^{a-2} d\mathbf{q}^{b-1} d\mathbf{q}^c \right\} \\ &= - \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)! (c-1)!} \int_{\mathbb{R}^{(a+b+c-2)d}} \int_{\mathbb{R}^d} K_2^\gamma(\mathbf{z}) p^{(a,b,c)}(\mathbf{q}^{a-1} \cup \mathbf{x}, \mathbf{q}^b, \mathbf{q}^{c-1} \cup \mathbf{z}, t) d\mathbf{z} d\mathbf{q}^{a-1} d\mathbf{q}^b d\mathbf{q}^{c-1} \\ &\quad + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} \left[\int_{\mathbb{R}^{2d}} m_2(\mathbf{x}, \mathbf{y} | \mathbf{z}) K_2^\gamma(\mathbf{z}) p^{(a,b,c+1)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c \cup \mathbf{z}, t) d\mathbf{y} d\mathbf{z} \right] d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c \\ &\quad + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-2)! (b-1)! c!} \int_{\mathbb{R}^{(a+b+c-3)d}} \left[\int_{\mathbb{R}^d} K_2^\gamma(\mathbf{z}) p^{(a-1,b-1,c+1)}(\mathbf{q}^{a-2} \cup \mathbf{x}, \mathbf{q}^{b-1}, \mathbf{q}^c \cup \mathbf{z}, t) d\mathbf{z} \right] \\ &\quad d\mathbf{q}^{a-2} d\mathbf{q}^{b-1} d\mathbf{q}^c \\ &= \int_{\mathbb{R}^d} \left[\int_{\mathbb{R}^d} m_2(\mathbf{x}, \mathbf{y} | \mathbf{z}) K_2^\gamma(\mathbf{z}) d\mathbf{y} \right] \mathbb{E}[C(\mathbf{z}, t)] d\mathbf{z}. \end{aligned}$$

In (III), we used that $\int_{\mathbb{R}^{2d}} m_2(\mathbf{x}, \mathbf{y} | \mathbf{z}) d\mathbf{x} d\mathbf{y} = 1$. In the second to last line, the first and third term cancel by shifting indexes.

In summary, the average concentration of species A satisfies

$$\begin{aligned} \partial_t \mathbb{E}[A(\mathbf{x}, t)] &= D_1 \Delta_{\mathbf{x}} \mathbb{E}[A(\mathbf{x}, t)] - \int_{\mathbb{R}^d} K_1^\zeta(\mathbf{x}, \mathbf{y}) \mathbb{E}[A(\mathbf{x}, t) B(\mathbf{y}, t)] d\mathbf{y} \\ &\quad + \int_{\mathbb{R}^d} \left[\int_{\mathbb{R}^d} m_2(\mathbf{x}, \mathbf{y} | \mathbf{z}) K_2^\zeta(\mathbf{z}) d\mathbf{y} \right] \mathbb{E}[C(\mathbf{z}, t)] d\mathbf{z}. \end{aligned}$$

Following similar arguments, one can derive equations for the average concentration of each species, given by (6.6) as claimed. This concludes the proof of the Proposition. \square

D PROOF OF THEOREM 6.4

Proof of Theorem 6.4. Due to the linearity of the equation, the proof of existence and uniqueness is standard here, so we only present a sketch of the argument for completeness.

Notice that the operator \mathcal{L} defined in (6.3) generates a contractive analytic semigroup on F , denoted by $\{e^{t\mathcal{L}}\}_{t \geq 0}$. Now, since by Lemma D.1, $\mathcal{R}^+, \mathcal{R}^-$ are Lipschitz continuous, existence of a unique local mild solution to (6.2), $\mathbf{P} \in \mathcal{C}([0, t_0]; F)$ follows by the standard Picard-Lindelöf theorem for equations with values in Banach spaces if the initial condition satisfies $\mathbf{P}_0 \in F$.

Next we establish global existence of a unique mild solution. The boundedness of the linear operators $\mathcal{R}^+, \mathcal{R}^-$ by Lemma D.1, together with the contraction property of the semigroup $t \mapsto e^{t\mathcal{L}}$, implies that

$$(D.1) \quad \|\mathbf{P}(t)\|_F \leq \|\mathbf{P}_0\|_F + C \int_0^t \|\mathbf{P}(s)\|_F ds$$

and a subsequent Gronwall lemma yields the bound $\|\mathbf{P}(t)\|_F \leq \|\mathbf{P}_0\|_F e^{Ct}$. This bound allows us, by choosing t_0 small enough, to extend the solution from the interval $[0, t_0)$ to the interval $[0, \infty)$. Hence, a unique global mild solution $\mathbf{P} \in \mathcal{C}([0, \infty); F)$ exists.

We actually have stronger regularity for the solution, $\mathbf{P}(t)$. This is a direct consequence of the contraction and regularization properties of the semigroup $\{e^{t\mathcal{L}}\}_{t \geq 0}$. Indeed, since $\mathbf{P}(t)$ is in $L^2(X)$, Lemma D.1 gives that $\mathcal{R}^+(\mathbf{P})$ and $\mathcal{R}^-(\mathbf{P})$ are both in $L^2(X)$. If, in addition, the initial condition $\mathbf{P}_0 \in H^1(X)$, then the mild form of the solution together with standard parabolic estimates (see estimates 3.1 in Chapter I.V, Section 3 of [LSU68]), gives that $\mathbf{P} \in \mathcal{C}([0, \infty); H^2(X))$.

Note, $\mathbf{P}(t)$ can be viewed as a probability density. Indeed, we have that if $\mathbf{P}_0 \geq 0$, $\mathbf{P}(t)$ is, by Lemma D.3, always non-negative for all $t \geq 0$. Second, if \mathbf{P}_0 satisfies the normalization condition,

$$\sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \left[\frac{1}{a! b! c!} \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} p_0^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right] = 1,$$

then the same normalization condition holds for $\mathbf{P}(t)$, $t \geq 0$,

$$\sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \left[\frac{1}{a! b! c!} \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right] = 1,$$

by Lemma D.2. This concludes the proof of the theorem. \square

Lemma D.1. *We have that the operators \mathcal{R}^+ and \mathcal{R}^- are bounded linear, Lipschitz continuous operators on F . Namely, for $\mathbf{G} = \{g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c)\}_{a,b,c=0}^{\infty} \in F$,*

$$(D.2) \quad \begin{aligned} \|\mathcal{R}^+(\mathbf{G})\|_F &\leq C \|\mathbf{G}\|_F, \\ \|\mathcal{R}^-(\mathbf{G})\|_F &\leq C \|\mathbf{G}\|_F, \\ \|\mathcal{R}^+(\mathbf{G}_1) - \mathcal{R}^+(\mathbf{G}_2)\|_F &\leq C \|\mathbf{G}_1 - \mathbf{G}_2\|_F, \\ \|\mathcal{R}^-(\mathbf{G}_1) - \mathcal{R}^-(\mathbf{G}_2)\|_F &\leq C \|\mathbf{G}_1 - \mathbf{G}_2\|_F. \end{aligned}$$

Proof. We'll only show the first two estimates hold. The Lipschitz conditions on \mathcal{R}^+ and \mathcal{R}^- follow directly from \mathcal{R}^+ and \mathcal{R}^- being bounded and linear. Assume that the initial number of particles are a_0, b_0 and c_0 for species A, B and C respectively. We then have that the following upper bounds hold for all times $0 \leq a \leq a_0 + c_0 := a_{max}$, $0 \leq b \leq b_0 + c_0 := b_{max}$ and $0 \leq c \leq a_0 \wedge b_0 + c_0 := c_{max}$.

By definition of the norm on Fock Space and the definition of \mathcal{R}^+ , we have that

$$\begin{aligned}
\|\mathcal{R}^+(\mathbf{G})\|_F^2 &= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} \left((\mathcal{R}^+ \mathbf{P})_{a,b,c}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c) \right)^2 d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c, \\
&= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} \left(- \left(\sum_{l=1}^a \sum_{m=1}^b K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) \right) g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) \right. \\
&\quad \left. + \sum_{n=1}^c \left[\int_{\mathbb{R}^{2d}} m_1(\mathbf{q}_n^c | \mathbf{x}, \mathbf{y}) K_1^\gamma(\mathbf{x}, \mathbf{y}) g^{(a+1,b+1,c-1)}(\mathbf{q}^a \cup \mathbf{x}, \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t) d\mathbf{x} d\mathbf{y} \right] \right)^2 d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c, \\
&\leq \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} \left(a_{\max} b_{\max} C(K) g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) \right)^2 d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c \\
&\quad + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} \left(\sum_{n=1}^c \left[\int_{\mathbb{R}^{2d}} m_1(\mathbf{q}_n^c | \mathbf{x}, \mathbf{y}) K_1^\gamma(\mathbf{x}, \mathbf{y}) \right. \right. \\
\text{(D.3)} \quad &\quad \left. \left. \times g^{(a+1,b+1,c-1)}(\mathbf{q}^a \cup \mathbf{x}, \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t) d\mathbf{x} d\mathbf{y} \right] \right)^2 d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c.
\end{aligned}$$

Without loss of generality, let us assume $0 < \alpha_i \leq 1$ in Assumption 4.4. Now denote $C_1 = (a_{\max} b_{\max} C(K))^2 < \infty$, and substitute the specific form of

$$m_1(\mathbf{q}_n^c | \mathbf{x}, \mathbf{y}) = \sum_{i=1}^I p_i \times \delta(\mathbf{q}_n^c - (\alpha_i \mathbf{x} + (1 - \alpha_i) \mathbf{y}))$$

into (D.3). We obtain

$$\begin{aligned}
\|\mathcal{R}^+(\mathbf{G})\|_F^2 &\leq C_1 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} \left(\sum_{n=1}^c \left[\int_{\mathbb{R}^{2d}} \sum_{i=1}^I p_i \times K_1^\gamma(\mathbf{x}, \mathbf{y}) \right. \right. \\
&\quad \left. \left. \times \delta(\mathbf{q}_n^c - (\alpha_i \mathbf{x} + (1 - \alpha_i) \mathbf{y})) \times g^{(a+1,b+1,c-1)}(\mathbf{q}^a \cup \mathbf{x}, \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t) d\mathbf{x} d\mathbf{y} \right] \right)^2 d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c. \\
&\leq C_1 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} \left(\sum_{n=1}^c \left[\int_{\mathbb{R}^d} \sum_{i=1}^I p_i \times K_1^\gamma \left(\frac{1}{\alpha_i} (\mathbf{q}_n^c - (1 - \alpha_i) \mathbf{y}), \mathbf{y} \right) \right. \right. \\
&\quad \left. \left. \times g^{(a+1,b+1,c-1)} \left(\mathbf{q}^a \cup \frac{1}{\alpha_i} (\mathbf{q}_n^c - (1 - \alpha_i) \mathbf{y}), \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t \right) d\mathbf{y} \right] \right)^2 d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c. \\
&\leq C_1 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c-1)d}} c \sum_{i=1}^I p_i^2 \sum_{n=1}^c \int_{\mathbb{R}^d} \left(\left[\int_{\mathbb{R}^d} K_1^\gamma \left(\frac{1}{\alpha_i} (\mathbf{q}_n^c - (1 - \alpha_i) \mathbf{y}), \mathbf{y} \right) \right. \right. \\
&\quad \left. \left. \times g^{(a+1,b+1,c-1)} \left(\mathbf{q}^a \cup \frac{1}{\alpha_i} (\mathbf{q}_n^c - (1 - \alpha_i) \mathbf{y}), \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t \right) d\mathbf{y} \right]^2 d\mathbf{q}_n^c \right) d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c \setminus \mathbf{q}_n^c. \\
&\leq C_1 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! (c-1)!} \int_{\mathbb{R}^{(a+b+c-1)d}} \sum_{i=1}^I p_i^2 \sum_{n=1}^c \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} \tilde{K}_1^\gamma \left(\left| \frac{1}{\alpha_i} (\mathbf{q}_n^c - (1 - \alpha_i) \mathbf{y}) - \mathbf{y} \right| \right) d\mathbf{y} \right)^2 \\
&\quad \times \left(\int_{\mathbb{R}^d} \left| g^{(a+1,b+1,c-1)} \left(\mathbf{q}^a \cup \frac{1}{\alpha_i} (\mathbf{q}_n^c - (1 - \alpha_i) \mathbf{y}), \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t \right) \right|^2 d\mathbf{y} \right) d\mathbf{q}_n^c d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c \setminus \mathbf{q}_n^c. \\
&\leq C_1 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! (c-1)!} \int_{\mathbb{R}^{(a+b+c-1)d}} \sum_{i=1}^I p_i^2 \left(\int_{\mathbb{R}^d} \alpha_i \tilde{K}_1^\gamma(|z|)^2 dz \right) \\
&\quad \times \sum_{n=1}^c \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} \left| g^{(a+1,b+1,c-1)} \left(\mathbf{q}^a \cup \frac{1}{\alpha_i} (\mathbf{q}_n^c - (1 - \alpha_i) \mathbf{y}), \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t \right) \right|^2 d\mathbf{y} \right) d\mathbf{q}_n^c d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c \setminus \mathbf{q}_n^c.
\end{aligned}$$

$$\begin{aligned}
&\leq C_1 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! (c-1)!} \int_{\mathbb{R}^{(a+b+c-1)d}} \sum_{i=1}^I p_i^2 \left(\int_{\mathbb{R}^d} \alpha_i \tilde{K}_1^{\gamma} (|z|)^2 dz \right) \\
&\quad \times \sum_{n=1}^c \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} \left| g^{(a+1, b+1, c-1)} \left(\mathbf{q}^a \cup \frac{1}{\alpha_i} (\mathbf{q}_n^c - (1 - \alpha_i) \mathbf{y}), \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t \right) \right|^2 d\mathbf{q}_n^c \right) d\mathbf{y} d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c \setminus \mathbf{q}_n^c. \\
&\leq C_1 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! (c-1)!} \int_{\mathbb{R}^{(a+b+c-1)d}} \sum_{i=1}^I p_i^2 \left(\int_{\mathbb{R}^d} \alpha_i \tilde{K}_1^{\gamma} (|z|)^2 dz \right) \\
&\quad \times \sum_{n=1}^c \alpha_i \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} \left| g^{(a+1, b+1, c-1)} \left(\mathbf{q}^a \cup \mathbf{x}, \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t \right) \right|^2 dx \right) d\mathbf{y} d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c \setminus \mathbf{q}_n^c. \\
&\leq C_1 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{Iabc}{(a+1)! (b+1)! (c-1)!} \left(\int_{\mathbb{R}^d} \tilde{K}_1^{\gamma} (|z|)^2 dz \right) \\
&\quad \times \int_{\mathbb{R}^{(a+b+c+1)d}} \left(\left| g^{(a+1, b+1, c-1)} \left(\mathbf{q}^{a+1}, \mathbf{q}^{b+1}, \mathbf{q}^{c-1}, t \right) \right|^2 \right) d\mathbf{q}^{a+1} d\mathbf{q}^{b+1} d\mathbf{q}^{c-1}. \\
&\leq C_1 \|\mathbf{G}\|_F^2 + C_2 \|\mathbf{G}\|_F^2 = (C_1 + C_2) \|\mathbf{G}\|_F^2, \\
&\text{(D.4)}
\end{aligned}$$

where $C_2 = I \times a_{max} b_{max} c_{max} \left(\int_{\mathbb{R}^d} \tilde{K}_1^{\gamma} (|z|)^2 dz \right) < \infty$ by Assumption 6.2.

Similarly, by definition of the norm on Fock Space and the definition of \mathcal{R}^- , we have that

$$\begin{aligned}
\|\mathcal{R}^-(\mathbf{G})\|_F^2 &= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} \left((\mathcal{R}^- \mathbf{G})_{a,b,c} (\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c) \right)^2 d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c, \\
&= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} \left(- \left(\sum_{n=1}^c K_2^{\gamma} (\mathbf{q}_n^c) \right) g^{(a,b,c)} (\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) \right. \\
&\quad \left. + \sum_{l=1}^a \sum_{m=1}^b \left[\int_{\mathbb{R}^d} m_2 (\mathbf{q}_l^a, \mathbf{q}_m^b | \mathbf{z}) K_2^{\gamma} (\mathbf{z}) g^{(a-1, b-1, c+1)} (\mathbf{q}^a \setminus \mathbf{q}_l^a, \mathbf{q}^b \setminus \mathbf{q}_m^b, \mathbf{q}^c \cup \mathbf{z}, t) dz \right] \right)^2 d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c, \\
&\leq \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} \left(c_{max} C(K) g^{(a,b,c)} (\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) \right)^2 d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c \\
&\quad + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} \left(\sum_{l=1}^a \sum_{m=1}^b \left[\int_{\mathbb{R}^d} m_2 (\mathbf{q}_l^a, \mathbf{q}_m^b | \mathbf{z}) K_2^{\gamma} (\mathbf{z}) \right. \right. \\
&\quad \left. \left. \times g^{(a-1, b-1, c+1)} (\mathbf{q}^a \setminus \mathbf{q}_l^a, \mathbf{q}^b \setminus \mathbf{q}_m^b, \mathbf{q}^c \cup \mathbf{z}, t) dz \right] \right)^2 d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c. \\
&\text{(D.5)}
\end{aligned}$$

Now denote $C_3 = (c_{max} C(K))^2 < \infty$, and substitute the specific form of

$$m_2 (\mathbf{q}_l^a, \mathbf{q}_m^b | \mathbf{z}) = \rho(|\mathbf{q}_l^a - \mathbf{q}_m^b|) \sum_{i=1}^I p_i \times \delta (\mathbf{z} - (\alpha_i \mathbf{q}_i^a + (1 - \alpha_i) \mathbf{q}_m^b)),$$

into (D.5). We obtain

$$\begin{aligned}
\|\mathcal{R}^-(\mathbf{G})\|_F^2 &\leq C_3 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} \left(\sum_{l=1}^a \sum_{m=1}^b \left[\int_{\mathbb{R}^d} m_2 (\mathbf{q}_l^a, \mathbf{q}_m^b | \mathbf{z}) K_2^{\gamma} (\mathbf{z}) \right. \right. \\
&\quad \left. \left. \times g^{(a-1, b-1, c+1)} (\mathbf{q}^a \setminus \mathbf{q}_l^a, \mathbf{q}^b \setminus \mathbf{q}_m^b, \mathbf{q}^c \cup \mathbf{z}, t) dz \right] \right)^2 d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c. \\
&\leq C_3 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \int_{\mathbb{R}^{(a+b+c)d}} \left(ab \sum_{l=1}^a \sum_{m=1}^b \left[\int_{\mathbb{R}^d} m_2 (\mathbf{q}_l^a, \mathbf{q}_m^b | \mathbf{z}) K_2^{\gamma} (\mathbf{z}) \right. \right. \\
&\quad \left. \left. \times g^{(a-1, b-1, c+1)} (\mathbf{q}^a \setminus \mathbf{q}_l^a, \mathbf{q}^b \setminus \mathbf{q}_m^b, \mathbf{q}^c \cup \mathbf{z}, t) dz \right] \right)^2 d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c.
\end{aligned}$$

$$\begin{aligned}
& \times g^{(a-1, b-1, c+1)} (\mathbf{q}^a \setminus \mathbf{q}_l^a, \mathbf{q}^b \setminus \mathbf{q}_m^b, \mathbf{q}^c \cup \mathbf{z}, t) dz \Big]^2) d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c. \\
\leq & C_3 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{(a-1)!(b-1)!c!} \int_{\mathbb{R}^{(a+b+c)d}} \left(\sum_{l=1}^a \sum_{m=1}^b C(K)^2 \left[\int_{\mathbb{R}^d} \sum_{i=1}^I p_i \times \rho(|\mathbf{q}_l^a - \mathbf{q}_m^b|) \right. \right. \\
& \left. \left. \times \delta(\mathbf{z} - (\alpha_i \mathbf{q}_i^a + (1 - \alpha_i) \mathbf{q}_m^b)) \times g^{(a-1, b-1, c+1)} (\mathbf{q}^a \setminus \mathbf{q}_l^a, \mathbf{q}^b \setminus \mathbf{q}_m^b, \mathbf{q}^c \cup \mathbf{z}, t) dz \right]^2 \right) d\mathbf{q}^a d\mathbf{q}^b d\mathbf{q}^c. \\
\leq & C_3 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{C(K)^2}{(a-1)!(b-1)!c!} \int_{\mathbb{R}^{(a+b+c)d}} \left(\sum_{l=1}^a \sum_{m=1}^b \sum_{i=1}^I p_i^2 \times \rho(|\mathbf{q}_l^a - \mathbf{q}_m^b|)^2 \right. \\
& \left. \times \left| g^{(a-1, b-1, c+1)} (\mathbf{q}^a \setminus \mathbf{q}_l^a, \mathbf{q}^b \setminus \mathbf{q}_m^b, \mathbf{q}^c \cup (\alpha_i \mathbf{q}_i^a + (1 - \alpha_i) \mathbf{q}_m^b), t) \right|^2 \right) d\mathbf{q}_l^a d\mathbf{q}_m^b d\mathbf{q}^a \setminus \mathbf{q}_l^a d\mathbf{q}^b \setminus \mathbf{q}_m^b d\mathbf{q}^c. \\
\leq & C_3 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{C(K)^2 ab}{(a-1)!(b-1)!c!} \sum_{i=1}^I p_i^2 \times \int_{\mathbb{R}^{(a+b+c)d}} (\rho(|\mathbf{x} - \mathbf{y}|)^2 \\
& \times \left| g^{(a-1, b-1, c+1)} (\mathbf{q}^{a-1}, \mathbf{q}^{b-1}, \mathbf{q}^c \cup (\alpha_i \mathbf{x} + (1 - \alpha_i) \mathbf{y}), t) \right|^2) d\mathbf{x} d\mathbf{y} d\mathbf{q}^{a-1} d\mathbf{q}^{b-1} d\mathbf{q}^c. \\
\leq & C_3 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{C(K)^2 ab}{(a-1)!(b-1)!c!} \sum_{i=1}^I p_i^2 \times \int_{\mathbb{R}^{(a+b+c)d}} (\rho(|\mathbf{w}|)^2 \\
& \times \left| g^{(a-1, b-1, c+1)} (\mathbf{q}^{a-1}, \mathbf{q}^{b-1}, \mathbf{q}^c \cup (\alpha_i \mathbf{w} + \mathbf{y}), t) \right|^2) d\mathbf{w} d\mathbf{y} d\mathbf{q}^{a-1} d\mathbf{q}^{b-1} d\mathbf{q}^c. \\
\leq & C_3 \|\mathbf{G}\|_F^2 + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{C(K)^2 ab(c+1)}{(a-1)!(b-1)!(c+1)!} \sum_{i=1}^I p_i^2 \times \left(\int_{\mathbb{R}^d} \rho(|\mathbf{w}|)^2 d\mathbf{w} \right) \\
& \times \int_{\mathbb{R}^{(a+b+c-1)d}} \left| g^{(a-1, b-1, c+1)} (\mathbf{q}^{a-1}, \mathbf{q}^{b-1}, \mathbf{q}^c \cup \mathbf{z}, t) \right|^2 dz d\mathbf{q}^{a-1} d\mathbf{q}^{b-1} d\mathbf{q}^c. \\
& \text{(where } \mathbf{z} = \alpha_i \mathbf{w} + \mathbf{y} \text{)} \\
\leq & C_3 \|\mathbf{G}\|_F^2 + C_4 \|\mathbf{G}\|_F^2 = (C_3 + C_4) \|\mathbf{G}\|_F^2, \\
\text{(D.6)} &
\end{aligned}$$

where $C_4 = I \times C(K)^2 a_{\max} b_{\max} (c_{\max} + 1) \left(\int_{\mathbb{R}^d} \rho(|\mathbf{w}|)^2 d\mathbf{w} \right) < \infty$ by Assumption 6.3. \square

Lemma D.2. Assume that the solution $\mathbf{P} \in \mathcal{C}([0, \infty); H^2(X))$ to (6.2) exists and is unique. If the normalization condition holds for the initial condition \mathbf{P}_0 , i.e. that

$$\sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \left[\frac{1}{a! b! c!} \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} p_0^{(a,b,c)} (\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right] = 1,$$

then we have that the normalization condition holds for $\mathbf{P}(t)$ for all $t \geq 0$, i.e. that

$$\text{(D.7)} \quad \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \left[\frac{1}{a! b! c!} \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} p^{(a,b,c)} (\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right] = 1.$$

Proof. By assumption, we have that the normalization condition holds for the initial condition. Furthermore, we have

$$\begin{aligned}
& \frac{\partial}{\partial t} \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \left[\frac{1}{a! b! c!} \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} p^{(a,b,c)} (\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right] \\
& = \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \left[\frac{1}{a! b! c!} \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} \frac{\partial}{\partial t} p^{(a,b,c)} (\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right]
\end{aligned}$$

$$\begin{aligned}
&= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \left[\frac{1}{a! b! c!} \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} (\mathcal{L}\mathbf{P})_{a,b,c}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right] \\
&+ \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \left[\frac{1}{a! b! c!} \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} (\mathcal{R}^+\mathbf{P})_{a,b,c}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right] \\
&+ \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \left[\frac{1}{a! b! c!} \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} (\mathcal{R}^-\mathbf{P})_{a,b,c} p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right].
\end{aligned}$$

Note, in the last equality the first term is zero using the divergence theorem and that $\mathbf{P} \in \mathcal{C}([0, \infty); H^2(X))$. The second term is

$$\begin{aligned}
&\sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \left[\frac{1}{a! b! c!} \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} (\mathcal{R}^+\mathbf{P})_{a,b,c}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right] \\
&= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \left[\int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} - \left(\sum_{l=1}^a \sum_{m=1}^b K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) \right) p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right. \\
&\quad \left. + \int_{\mathbb{R}^{da}} \int_{\mathbb{R}^{db}} \int_{\mathbb{R}^{dc}} \sum_{n=1}^c \left[\int_{\mathbb{R}^{2d}} m_1(\mathbf{q}_n^c | \mathbf{x}, \mathbf{y}) K_1^\gamma(\mathbf{x}, \mathbf{y}) p^{(a+1,b+1,c-1)}(\mathbf{q}^a \cup \mathbf{x}, \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t) d\mathbf{x} d\mathbf{y} \right] d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right] \\
&= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \left[\int_{\mathbb{R}^{d(a+b+c)}} - \left(\sum_{l=1}^a \sum_{m=1}^b K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) \right) p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right. \\
&\quad \left. + \sum_{n=1}^c \int_{\mathbb{R}^{d(a+b+c+1)}} \left[\int_{\mathbb{R}^d} m_1(\mathbf{q}_n^c | \mathbf{x}, \mathbf{y}) d\mathbf{q}_n^c \right] K_1^\gamma(\mathbf{x}, \mathbf{y}) p^{(a+1,b+1,c-1)}(\mathbf{q}^a \cup \mathbf{x}, \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t) d\mathbf{q}^c \setminus d\mathbf{q}_n^c d\mathbf{q}^b \cup d\mathbf{y}, d\mathbf{q}^a \cup d\mathbf{x} \right] \\
&= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \frac{1}{a! b! c!} \left[\int_{\mathbb{R}^{d(a+b+c)}} - \left(\sum_{l=1}^a \sum_{m=1}^b K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) \right) p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right. \\
&\quad \left. + \int_{\mathbb{R}^{d(a+b+c+1)}} \frac{c}{(a+1)(b+1)} \sum_{l=1}^{a+1} \sum_{m=1}^{b+1} K_1^\gamma(\mathbf{q}_l^{a+1}, \mathbf{q}_m^{b+1}) p^{(a+1,b+1,c-1)}(\mathbf{q}^{a+1}, \mathbf{q}^{b+1}, \mathbf{q}^{c-1}, t) d\mathbf{q}^{c-1} d\mathbf{q}^{b+1}, d\mathbf{q}^{a+1} \right] \\
&= \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} \left[\frac{1}{a! b! c!} \int_{\mathbb{R}^{d(a+b+c)}} - \left(\sum_{l=1}^a \sum_{m=1}^b K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) \right) p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) d\mathbf{q}^c d\mathbf{q}^b d\mathbf{q}^a \right. \\
&\quad \left. + \frac{1}{(a+1)!(b+1)!(c-1)!} \int_{\mathbb{R}^{d(a+b+c+1)}} \sum_{l=1}^{a+1} \sum_{m=1}^{b+1} K_1^\gamma(\mathbf{q}_l^{a+1}, \mathbf{q}_m^{b+1}) p^{(a+1,b+1,c-1)}(\mathbf{q}^{a+1}, \mathbf{q}^{b+1}, \mathbf{q}^{c-1}, t) d\mathbf{q}^{c-1} d\mathbf{q}^{b+1}, d\mathbf{q}^{a+1} \right] \\
&= 0,
\end{aligned}$$

by using the symmetry of $p^{(a,b,c)}$ with respect to permutations of the ordering of particle positions for particles of the *same* type. A similar calculation shows that the third term is zero. Thus, the time derivative of the left-hand-side (D.7) is always zero. Combining with the normalization condition for the initial condition, we have that (D.7) is true for all $t \geq 0$. \square

Lemma D.3. *Assuming there exists a unique solution $\mathbf{P} \in \mathcal{C}([0, \infty); H^2(X))$ and $\mathbf{P}_0 \geq 0$, the solution components $p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t)$ are always non-negative for all $t \geq 0$.*

Proof. First, we consider $g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t)$ satisfying the following decoupled linear PDEs with initial condition $g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, 0) = p_0^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c) \geq 0$,

$$(D.8) \quad \frac{\partial}{\partial t} g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) = (\mathcal{L} + \mathcal{R}_1^+ + \mathcal{R}_1^-) g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t),$$

where we define

$$\mathcal{R}_1^+ g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) = - \left(\sum_{l=1}^a \sum_{m=1}^b K_1^\gamma(\mathbf{q}_l^a, \mathbf{q}_m^b) \right) g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t)$$

and

$$\mathcal{R}_1^- g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) = - \left(\sum_{n=1}^c K_2^\gamma(\mathbf{q}_n^c) \right) g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t).$$

Since $0 \leq K_1^\gamma \leq C(K)$ and $0 \leq K_2^\gamma \leq C(K)$, we can then obtain via a comparison argument for semilinear equations that the solution to Eq (D.8), $g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) \geq 0$.

Let us further define

$$\mathcal{R}_2^+ g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) = \sum_{n=1}^c \left[\int_{\mathbb{R}^{2d}} m_1(\mathbf{q}_n^c | \mathbf{x}, \mathbf{y}) K_1^\gamma(\mathbf{x}, \mathbf{y}) g^{(a+1, b+1, c-1)}(\mathbf{q}^a \cup \mathbf{x}, \mathbf{q}^b \cup \mathbf{y}, \mathbf{q}^c \setminus \mathbf{q}_n^c, t) d\mathbf{x} d\mathbf{y} \right],$$

and

$$\mathcal{R}_2^- g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) = \sum_{l=1}^a \sum_{m=1}^b \left[\int_{\mathbb{R}^d} m_2(\mathbf{q}_l^a, \mathbf{q}_m^b | \mathbf{z}) K_2^\gamma(\mathbf{z}) g^{(a-1, b-1, c+1)}(\mathbf{q}^a \setminus \mathbf{q}_l^a, \mathbf{q}^b \setminus \mathbf{q}_m^b, \mathbf{q}^c \cup \mathbf{z}, t) d\mathbf{z} \right].$$

We then set $\mathcal{R}^+ = \mathcal{R}_1^+ + \mathcal{R}_2^+$ and $\mathcal{R}^- = \mathcal{R}_1^- + \mathcal{R}_2^-$. Due to the positive mapping property of the operators \mathcal{R}_2^+ and \mathcal{R}_2^- , we shall have for the function $g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t)$ that

$$\frac{\partial}{\partial t} g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) \leq (\mathcal{L} + \mathcal{R}^+ + \mathcal{R}^-) g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t).$$

Hence, again utilizing the comparison principle for semilinear PDEs, we obtain that

$$0 \leq g^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t) \leq p^{(a,b,c)}(\mathbf{q}^a, \mathbf{q}^b, \mathbf{q}^c, t),$$

i.e. the non-negativity of our solution, concluding the proof. \square

REFERENCES

- [D78] D. Aldous, "Stopping times and tightness." The Annals of Probability 6.2, (1978), pp. 335-340.
- [DK15] D. F. Anderson and T. G. Kurtz, Stochastic analysis of biochemical systems. Vol. 1. Berlin: Springer, 2015.
- [AT80] L. Arnold and M. Theodosopulu, *Deterministic limit of the stochastic model of chemical reactions with diffusion*, Advances in Applied Probability **12** (1980), no. 2, pp. 367-379.
- [BM15] V. Bansaye and S. Méléard, Stochastic models for structured populations. Springer, 2015.
- [CSY20] A. Capponi, X. Sun and D. Yao, *A Dynamic Network Model of Interbank Lending: Systemic Risk and Liquidity Provisioning*, Mathematics of Operations Research. Forthcoming, 2020.
- [CK49] F. C. Collins and G.E. Kimball, *Diffusion-controlled reaction rates*, J. Colloid. Sci. **4** (1949), no. 4, pp. 425-437.
- [DH96] P. Dai Pra and F. Hollander, *Mckean-vlasov limit for interacting random processes in random media*, Journal of Statistical Physics **84** (1996), no. 3-4, pp. 735-772.
- [DH09] P. Dai Pra and M. Tolotti, *Heterogeneous credit portfolios and the dynamics of the aggregate losses*, Stochastic Processes and their Applications **119** (2009), no. 9, pp. 2913-2944.
- [DIRT15] F. Delarue, J. Inglis, S. Rubenthaler, and E. Tanre, *Particle systems with a singular mean-field self-excitation. application to neuronal networks*, Stochastic Processes and their Applications **125** (2015), pp. 2451-2492.
- [D76a] M. Doi, *Second quantization representation for classical many-particle system*, J. Phys. A: Math. Gen. **9** (1976), no. 9, pp. 1465-1477.
- [D76b] M. Doi, *Stochastic theory of diffusion-controlled reaction*, J. Phys. A: Math. Gen. **9** (1976), no. 9, pp. 1479-1495.
- [DYK18] A. Donev, C.-Y. Yang and C. Kim, *Efficient reactive Brownian dynamics*, J. Chem. Phys. **148** (2018), 034103 (17pp).
- [EC09] R. Erban and S. J. Chapman, *Stochastic modelling of reaction-diffusion processes: algorithms for bimolecular reactions*, Phys. Biol. **6** (2009), no. 4, 046001 (18pp).
- [EK86] S. N. Ethier and T.G. Kurtz, *Markov Processes: Characterization and Convergence*. John Wiley & Sons Inc., New York, 1986.
- [HCDWS19] J. Helfmann, N.D. Conrad, A. Djurdjevic, S. Winkelmann, and C. Schütte, *From interacting agents to density-based modeling with stochastic pdes*, arXiv:1905.13525 (2019).
- [GSS13] K. Giesecke, K. Spiliopoulos, and R.B. Sowers, *Default clustering in large portfolios: Typical events*, The Annals of Applied Probability **23**, (2013), pp. 348-385.
- [GSSS15] K. Giesecke, K. Spiliopoulos, R.B. Sowers, and J. Sirignano, *Large portfolio asymptotics for loss from default*, Mathematical Finance **25**, (2015), pp. 77-114.
- [NW14] N. Ikeda and S. Watanabe, Stochastic differential equations and diffusion processes. Vol. 24. Elsevier, 2014.
- [IT15] J. Inglis and D. Talay, *Mean-field limit of a stochastic particle system smoothly interacting through threshold hitting-times and applications to neural networks with dendritic component*, SIAM Journal on Mathematical Analysis **47** (2015), pp. 3884-3916.
- [IS13] S. A. Isaacson, *A Convergent Reaction-Diffusion Master Equation*, J. Chem. Phys. **139** (2013), 054101 (12 pp).
- [IZ18] S. A. Isaacson and Y. Zhang, "An unstructured mesh convergent reactiondiffusion master equation for reversible reactions." Journal of Computational Physics 374, (2018), pp. 954-983.

- [K82] J. Keizer, *Nonequilibrium statistical thermodynamics and the effect of diffusion on chemical reaction rates*, J. Phys. Chem. **86**, (1982), pp. 5052–5067.
- [K01] T.G. Kurtz, "Lectures on stochastic analysis." Department of Mathematics and Statistics, University of Wisconsin, Madison, WI (2001): 53706-1388.
- [LSU68] O.A. Ladyženskaja, V.A. Solonnikov and N.N. Ural'ceva. Linear and Quasi-linear equations of parabolic type, *Transactions of Mathematical Monographs, American Mathematical Society*, Volume 23, (1968).
- [LLN19] T.S. Lim, Y. Lu, and J. Nolen, *Quantitative propagation of chaos in the bimolecular chemical reaction-diffusion model*, arXiv:1906.01051 (2019).
- [BMW12] B. Jourdain, S. Méléard, and W. A. Woyczynski, "Lvy flights in evolutionary ecology." Journal of mathematical biology 65.4 (2012), pp. 677-707.
- [AM86] A. Joffe and M. Métivier, "Weak convergence of sequences of semimartingales with applications to multitype branching processes." Advances in Applied Probability 18.1, (1986), pp. 20-65.
- [LE11] J. Lipkova, K.C. Zygalakis, S.J. Chapman and R. Erban, *Analysis of Brownian Dynamics Simulations of Reversible Bimolecular Reactions*, SIAM J. Appl Math **71**, (2011), no. 3, pp. 714-730.
- [MS93] S. Méléard and S. Roelly. "Sur les convergences troite ou vague de processus valeurs mesures." Comptes rendus de l'Acadmie des sciences. Srie 1, Mathmatique 317.8, (1993), pp. 785-788.
- [MNKS09] J. Muñoz-García, Z. Neufeld, B. N. Kholodenko, and H. M. Sauro, *Positional information generated by spatially distributed signaling cascades*, PLoS Comp. Biol. **5** (2009), no. 3, e1000330.
- [MS02] O. Moynot and M. Samuelides, *Large deviations and mean-field theory for asymmetric random recurrent neural networks*, Probability Theory and Related Fields **123**, (2002), pp. 41–75.
- [NTS08] S. Neves, P. Tsokas, A. Sarkar, E. Grace, P. Rangamani, S. Taubenfeld, C. Alberini, J. Schaff, R. Blitzer, I. Moraru, and R. Iyengar, *Cell shape and negative links in regulatory motifs together control spatial information flow in signaling networks*, Cell **133** (2008), no. 4, pp. 666–680.
- [O89] K. Oelschläger, *On the derivation of reaction-diffusion equations as limit dynamics of systems of moderately interacting stochastic processes*, Probab. Th. Rel. Fields **82** (1989), no. 4, pp. 565–586.
- [S89] S. Roelly-Coppoletta, "A criterion of convergence of measurevalued processes: application to measure branching processes." Stochastics: An International Journal of Probability and Stochastic Processes 17.1-2, (1986), pp. 43-65.
- [P05] P.E. Protter, "Stochastic differential equations." Stochastic integration and differential equations. Springer, Berlin, Heidelberg, 2005. 249-361.
- [S09] D.E. Shaw, R.O. Dror, J.K. Salmon, JP Grossman, K.M. Mackenzie, J.A. Bank, C. Young, M.M. Deneroff, B. Batson, and K.J. Bowers, *Millisecond-scale molecular dynamics simulations on Anton*, Proceedings of the Conference on High Performance Computing Networking, Storage and Analysis (2009), 39.
- [SV06] D.W. Stroock and S. R. S. Varadhan, "Multidimensional Diffusion Processes", Springer-Verlang Berlin Heidelberg, (2006).
- [SM17] M. V. Smoluchowski, *Mathematical theory of the kinetics of the coagulation of colloidal solutions*, Z. Phys. Chem. **92** (1917), pp. 129–168.
- [SCS88] H. Sompolinsky, A. Crisanti, and H. Sommers, *Chaos in random neural networks*, Physical Review Letters **61**, (1988), 259.
- [T67] E Teramoto and N Shigesada, *Theory of bimolecular reaction processes in liquids*, Prog. Theor. Phys. **37** (1967), no. 1, pp. 29–51.