

Harder–Narasimhan theory

These notes contain not one original idea. An abstract formulation of Harder–Narasimhan theory is stated without proof in [F], and I found it helpful to write it all out. For applications of this system, see [F] and [K], as well as their various references; we give a rather generic example in §8.

(Note: I wrote this document around 2010, and was recently encouraged to place it on the arXiv. I’ve taken the opportunity to correct some typos, sins of exposition and formatting, and actual errors. A contemporary literature search would reveal that this theory has since been generalized. — Jonathan Pottharst)

§1. Hypotheses and statements.

The inputs and axioms given here will be in force throughout this note.

(1.1) When working in exact categories, I like to say “strict mono/epimorphism” as opposed to “admissible mono/epimorphism”. A *strict subobject* is a subobject whose inclusion map is a strict monomorphism. I also call the distinguished (short) exact sequences “(short) strict exact sequences”.

Inputs (1.2). Our input consists of the data:

- a totally ordered abelian group V (written additively),
- an exact category \mathcal{C} (our target),
- an abelian category \mathcal{D} ,
- a functor $F: \mathcal{C} \rightarrow \mathcal{D}$,
- a rule $\mathrm{rk}: |\mathcal{D}| \rightarrow \mathbf{Z}_{\geq 0}$ (where $|\cdot|$ denotes isomorphism classes), and
- a rule $\mathrm{deg}: |\mathcal{C}| \rightarrow V$.

Axioms (1.3). We assume:

- F is exact and faithful, and for each object X of \mathcal{C} , F induces a bijection

$$\{\text{strict subobjects of } X\} \xrightarrow{\sim} \{\text{subobjects of } F(X)\};$$

- rk is additive over short exact sequences, and, for each object X of \mathcal{D} , $\mathrm{rk} X = 0$ if and only if $X = 0$; and
- deg is additive over short strict exact sequences, and, for each morphism $f: X \rightarrow Y$ of \mathcal{C} with $F(f)$ an isomorphism, one has $\mathrm{deg}(X) \leq \mathrm{deg}(Y)$, with equality if and only if f itself is an isomorphism.

(1.4) We call $V \otimes_{\mathbf{Z}} \mathbf{Q}$ the set of *slopes*. (Typically, $V \subseteq \mathbf{R}$.) We say that the strict subobject X' of X and the subobject $F(X')$ of $F(X)$ *correspond* to one another. We call rk the *rank*, \deg the *degree*, and, for nonzero objects X of \mathcal{C} , $\mu(X) := \deg(X)/\text{rk}(X) \in V \otimes_{\mathbf{Z}} \mathbf{Q}$ its *slope*. (Throughout, we abusively write $\text{rk}(X)$ for $\text{rk}(F(X))$ for objects X of \mathcal{C} .) Note that the second part of the axiom for \deg can be equivalently phrased with μ in place of \deg .

(1.5) One calls a nonzero object X of \mathcal{C} *semistable* (resp. *stable*) if for all nonzero proper subobjects X' of X one has $\mu(X') \leq \mu(X)$ (resp. $\mu(X') < \mu(X)$).

A *HN filtration* of X is an increasing, finite, separating and exhaustive filtration by strict subobjects

$$0 = X_0 \subsetneq X_1 \subsetneq \cdots \subsetneq X_N = X$$

with each $\text{Gr}_i = X_i/X_{i-1}$ semistable, and each $\mu(\text{Gr}_{i+1}) < \mu(\text{Gr}_i)$. The $\mu(\text{Gr}_i)$ are called the *HN slopes* of X , and the $\text{rk}(\text{Gr}_i)$ are called their respective ranks. (Note that, in applications, simple semistable objects might have rank greater than 1, so that the ranks of the graded pieces can be larger than the traditional “multiplicities” of the slopes.)

(1.6) These are the central results.

Theorem. *For any slope μ , the full subcategory $\mathcal{C}^{(\mu)}$ of \mathcal{C} consisting of semistable objects of slope μ (together with the zero object) is an abelian, exact subcategory of \mathcal{C} that is closed under extensions in \mathcal{C} .*

Theorem. *Each nonzero object of \mathcal{C} possesses a unique HN filtration. When the HN filtrations are reindexed “by slopes”, forming the HN filtration is functorial.*

The first theorem is (4.12) below. The second theorem is the combination of (6.3), (6.2.b), and (6.5.c).

§2. Properties of F .

It is impressive how much mileage one gets from only the knowledge of F .

Proposition (2.1). *For any object X of \mathcal{C} , the correspondence induced by F ,*

$$\{\text{strict subobjects of } X\} \xrightarrow{\sim} \{\text{subobjects of } F(X)\},$$

is an isomorphism of posets.

Proof. The above correspondence is a bijection by axiom. Let X', X'' be two strict subobjects of X . It is clear that if $X' \subseteq X''$ then $F(X') \subseteq F(X'')$; we show the converse. Consider the composite $X' \subseteq X \rightarrow X/X''$. Applying F , we get $F(X') \rightarrow F(X/X'') = F(X)/F(X'')$. If $F(X') \subseteq F(X'')$ then this is the zero map. Since F is faithful, $X' \rightarrow X/X''$ is the zero map, and therefore $X' \subseteq X''$. \square

(2.2) A morphism $f: X \rightarrow Y$ in \mathcal{C} is said to be an *F -isomorphism* (resp. *F -monomorphism*, *F -epimorphism*) if $F(f)$ is an isomorphism (resp. monomorphism, epimorphism).

Proposition (2.3). *If a morphism $f: X \rightarrow Y$ in \mathcal{C} is an F -monomorphism, then it is a monomorphism. In particular, if f is an F -isomorphism, then f is a monomorphism.*

Proof. Suppose that f is an F -monomorphism, and that $g: Z \rightarrow X$ is a map in \mathcal{C} with

$f \circ g = 0$. Then $F(f) \circ F(g) = 0$, and because f is an F -monomorphism we have $F(g) = 0$. But F is faithful, so $g = 0$. \square

(2.4) Let X be an object of \mathcal{C} , and X' a subobject. We let \widetilde{X}' denote the strict subobject of X corresponding to the subobject $F(X')$ of $F(X)$, i.e. the unique strict subobject of X satisfying $F(\widetilde{X}') = F(X')$. We call \widetilde{X}' the *saturation* of X' in X .

Proposition (2.5). *Let X be an object of \mathcal{C} , and X', X'' subobjects. Then:*

- (a) $X' \subseteq \widetilde{X}'$.
- (b) $\widetilde{X}' \subseteq \widetilde{X}''$ if and only if $F(X') \subseteq F(X'')$. In particular, $F(X') = F(X'')$ if and only if $\widetilde{X}' = \widetilde{X}''$, and $\widetilde{\widetilde{X}'} = \widetilde{X}'$.
- (c) Considering X' and X'' equivalent if $F(X') = F(X'')$, each equivalence class contains a unique strict object, equal to the saturation of any object in the class, and this object is the final member with respect to inclusion.
- (d) \widetilde{X}' is characterized among strict subobjects of X containing X' as being the initial one, or as the unique one such that F takes the inclusion $X' \subseteq \widetilde{X}'$ to an isomorphism.

Proof. (a): If it were true that $X' \not\subseteq \widetilde{X}'$, then the natural map $X' \rightarrow X/\widetilde{X}'$ would be nonzero. Since F is faithful, the induced map $F(X') \rightarrow F(X/\widetilde{X}') = F(X)/F(\widetilde{X}') = F(X)/F(X')$ would also be nonzero, but this is impossible.

(b,c,d): These are obvious, given (a). \square

(2.6) The process of modifying the subobject X' to \widetilde{X}' , i.e. one with better exactness properties, is similar to passing to a derived or homotopy category. From this optic, it would be tempting to localize \mathcal{C} with respect to F , i.e. formally invert all F -isomorphisms. The problem is that F -isomorphisms do not necessarily respect deg. In fact, by axiom, they only respect deg when they are actual isomorphisms!

Nonetheless, one still can perform most abelian category operations in \mathcal{C} , as we now show.

Proposition (2.7). *Every morphism $f: X \rightarrow Y$ has a kernel and image in \mathcal{C} . The kernel is a strict subobject of X , and $X/\ker f$ is canonically identified with the image.*

Proof. We write, temporarily, $\widetilde{\ker f}$ for the strict subobject of X corresponding to $\ker F(f)$ via (2.1). We claim it is the kernel of f . First, consider the composite $\widetilde{\ker f} \rightarrow X \rightarrow Y$. Applying F gives the composite $\ker F(f) \rightarrow F(X) \rightarrow F(Y)$, which is zero, but F is faithful, so our original composite must be zero. On the other hand, consider any subobject $X' \subseteq X$ such that the composite $X' \rightarrow X \rightarrow Y$ is zero. Applying F , we see that $F(\widetilde{X}') = F(X') \rightarrow F(X) \rightarrow F(Y)$ is zero, and therefore $F(\widetilde{X}') \subseteq \ker F(f)$ inside $F(X)$. By (2.5.c), we find that $\widetilde{X}' \subseteq \widetilde{\ker f}$, and therefore $X' \subseteq \widetilde{X}' \subseteq \widetilde{\ker f}$. This shows that the strict subobject $\widetilde{\ker f}$ is the kernel, so we henceforth denote it by $\ker f$.

Now f induces a map $\overline{f}: X/\ker f \rightarrow Y$, and applying F yields the monomorphism $F(X)/\ker F(f) \rightarrow F(Y)$ in \mathcal{D} . By (2.2), \overline{f} is a monomorphism. Therefore, $X/\ker f$ is canonically identified to a subobject of Y that we call $\text{img } f$. We claim it is an image for this morphism. We must show that any subobject $Y' \subseteq Y$ through which f factors must contain

$\text{img } f$. Because we have the factorization of f into $X \xrightarrow{f_0} Y' \subseteq Y$, with the second map a monomorphism, we have $\ker f = \ker f_0$. Therefore, composite $X/\ker f \xrightarrow{\bar{f}_0} Y' \subseteq Y$ is simply \bar{f} , i.e. \bar{f} and hence f factors through Y' . But \bar{f} is identified with the inclusion $\text{img } f \subseteq Y$, so we are done. \square

Proposition (2.8). *Let X be an object of \mathcal{C} , and let X', X'' be two subobjects. There exists a least upper bound $X' + X''$ and a greatest lower bound $X' \cap X''$ for X', X'' in the poset of all subobjects of X , and the canonical sequence*

$$0 \rightarrow X' \cap X'' \xrightarrow{(1, -1)} X' \oplus X'' \rightarrow X' + X'' \rightarrow 0$$

is short strict exact. If X', X'' are strict in X , then so is $X' \cap X''$.

Proof. Using (2.7), define $X' + X''$ to be the image of the map $X' \oplus X'' \rightarrow X$. It is immediate to verify that $X' + X''$ is a least upper bound for X', X'' in X .

Let K be the kernel of $X' \oplus X'' \rightarrow X$. We show that the compositions $K \rightarrow X' \oplus X'' \rightarrow X'$ and $K \rightarrow X' \oplus X'' \rightarrow X''$ are monomorphisms. Suppose given a map $Z \rightarrow K$ in \mathcal{C} , whose composite $Z \rightarrow K \rightarrow X' \oplus X'' \rightarrow X'$ is zero. Consider the two compositions

$$\begin{aligned} K &\rightarrow X' \oplus X'' \rightarrow X' \rightarrow X' + X'' \text{ and} \\ K &\rightarrow X' \oplus X'' \rightarrow X'' \rightarrow X' + X''. \end{aligned}$$

Their sum is the composition $K \rightarrow X' \oplus X'' \rightarrow X' + X''$, and therefore is zero. Because $X'' \rightarrow X' + X''$ is a monomorphism, the composite $Z \rightarrow K \rightarrow X' \oplus X'' \rightarrow X''$ is zero. Since $Z \rightarrow K \rightarrow X' \oplus X''$ becomes zero followed by each of the projections to X', X'' , it must be zero. But $K \rightarrow X' \oplus X''$ is a monomorphism, and so $Z \rightarrow K$ is zero. This shows that $K \rightarrow X' \oplus X'' \rightarrow X'$ is a monomorphism, and the same argument shows that $K \rightarrow X' \oplus X'' \rightarrow X''$ is a monomorphism as well.

We denote by $X' \cap X''$ the subobject of $X' + X''$ determined by the monomorphism $K \rightarrow X' \oplus X'' \rightarrow X' \rightarrow X' + X''$. By construction it is a subobject of X' , and the above argument shows that it is also a subobject of X'' . The same reasoning shows that the composition $X' \cap X'' \rightarrow X' \oplus X'' \rightarrow X''$ is the *negative* of the canonical inclusion $X' \cap X'' \subseteq X''$ as subobjects of $X' + X''$. Therefore, we have the short exact sequence as claimed.

We now show that the subobject $X' \cap X''$ is a greatest lower bound for X', X'' in X . Given any subobject Y of X whose inclusion into X is factored through both X', X'' , we must factor its inclusion into X through $X' \cap X''$. Combining the two given factorizations gives a map $(1, -1): Y \rightarrow X' \oplus X''$, whose projection to X' is the inclusion $Y \subseteq X'$ and whose sum to X is zero. Therefore, it uniquely factors through $X' \cap X'' \rightarrow X' \oplus X''$. But the composite $X' \cap X'' \rightarrow X' \oplus X'' \rightarrow X'$ is the canonical inclusion, so we have exhibited a factorization of $Y \subseteq X'$ through $X' \cap X'' \subseteq X'$.

Finally, suppose that X', X'' are strict. Denote by $X' \tilde{\cap} X''$ the strict subobject of X corresponding via (2.1) to $F(X') \cap F(X'')$. We show that the strict object $X' \tilde{\cap} X''$ is a greatest lower bound for X, X'' in X . By the isomorphism of posets, $X' \tilde{\cap} X''$ is contained in both X', X'' . Moreover, if Y is any subobject of X that is contained in X', X'' , then $F(Y) \subseteq F(X')$ and $F(Y) \subseteq F(X'')$, hence $F(Y) \subseteq F(X') \cap F(X'')$ in $F(X)$. By (2.5.b) this is equivalent to $\tilde{Y} \subseteq X' \tilde{\cap} X''$, and we conclude by (2.5.a). \square

(2.9) We warn the reader that, in the above lemma, even if X', X'' are strict subobjects of X , the subobject $X' + X''$ might not be strict.

Proposition (2.10). *Let X be an object of \mathcal{C} , and X', X'' two subobjects. Assume the inclusions $X' \cap X'' \subseteq X'$ and $X'' \subseteq X' + X''$ are strict. (This holds, in particular, if X', X'' are strict subobjects of X .) Then the canonical map*

$$f: \frac{X'}{X' \cap X''} \rightarrow \frac{X' + X''}{X''}$$

induced by $X' \subseteq X' + X''$ is an isomorphism. Moreover, these objects are canonically subobjects of X/X'' .

Proof. Applying F to f clearly yields an isomorphism, so by (2.2) f is a monomorphism. We must show that the subobject thus obtained is the whole thing. Considering that the compositions $X' \rightarrow X'/(X' \cap X'') \rightarrow (X' + X'')/X''$ and $X' \rightarrow X' + X'' \rightarrow (X' + X'')/X''$ are equal, we see that the image of f contains the image of X' under the latter of these two. The image of f contains 0, which is the image of the composite $X'' \rightarrow X' + X'' \rightarrow X' + X''/X''$. Therefore, the image of f contains the image of the composite $X' \oplus X'' \rightarrow X' + X'' \rightarrow (X' + X'')/X''$. But each of the latter two maps are strict surjections. The second claim follows from applying (2.3) to the canonical map $(X' + X'')/X'' \rightarrow X/X''$. \square

§3. Properties of rk.

The main use of rk is to bound chains of strict subobjects in \mathcal{C} .

Proposition (3.1). *Every object of \mathcal{D} has finite length. For any object \mathcal{D} , any nonempty collection of its subobjects closed under $+$ admits a final element.*

Proof. By dévissage and induction, one sees that length of an object X of \mathcal{D} is bounded above by its rank.

For the second claim, the ranks in the collection are bounded above by the rank of the parent object, and any member whose rank is maximal is the final element. \square

Proposition (3.2). *Let X be an object of \mathcal{C} , and $X' \subseteq X'' \subseteq X$ subobjects. Then $\text{rk}(X') = \text{rk}(\tilde{X}')$, and if $\text{rk}(X') = \text{rk}(X'')$ then $\tilde{X}' = \tilde{X}''$. In particular, any subobject X' of X with $\text{rk}(X') = \text{rk}(X)$ has $\tilde{X}' = X$.*

Proof. The first claim is clear, because $F(X') = F(\tilde{X}')$. For the second claim, we may therefore replace X' and X'' by \tilde{X}' and \tilde{X}'' , and assume that X' and X'' are strict. Now consider the short strict exact sequence $0 \rightarrow X' \rightarrow X'' \rightarrow X''/X' \rightarrow 0$. Applying F , we get the short exact sequence $0 \rightarrow F(X') \rightarrow F(X'') \rightarrow F(X''/X') \rightarrow 0$. We must have $\text{rk}(F(X''/X')) = 0$, which forces $0 = F(X''/X') = F(X'')/F(X')$. Therefore $F(X') = F(X'')$, and the axiom for F shows that $X' = X''$ because X' and X'' are strict. \square

§4. Properties of deg and μ .

With notions of deg and μ , an arithmetic of subobjects emerges.

Proposition (4.1). *Let X be an object of \mathcal{C} , and consider the equivalence relation on subobjects of X from (2.5.c). In any equivalence class, all members have the same rank, and*

the unique strict subobject is the unique member with maximal degree (resp. slope).

Proof. The claim for rk is trivial. In a given equivalence class, given that the strict object is final with respect to inclusion, the claim for deg is a restatement of our axiom for deg. The claim for μ follows. \square

Proposition (4.2). *Let X be an object of \mathcal{C} , and X' a subobject whose slope is maximal among the slopes of subobjects of X . Then X' is a strict subobject.*

Proof. By (2.5.d), one has $X' = \tilde{X}'$ if and only if X' is strict. Moreover, by (4.1) one has $\mu(X') \leq \mu(\tilde{X}')$, with equality if and only if $X' = \tilde{X}'$. But the slope of X' is maximal, so also $\mu(\tilde{X}') \leq \mu(X')$. \square

Proposition (4.3). *Let $0 = X_0 \subsetneq X_1 \subsetneq \cdots \subsetneq X_N = X$ be strict subobjects of X . Write $\text{Gr}_i = X_i/X_{i-1}$, $\text{rk}_i = \text{rk}(\text{Gr}_i)$, $\deg_i = \deg(\text{Gr}_i)$, and $\mu_i = \mu(\text{Gr}_i)$. Then:*

(a) *One has the identity*

$$\mu(X) = \sum_{i=1}^N \frac{\text{rk}_i}{\text{rk}(X)} \mu_i.$$

(b) *Either $\min \{\mu_i\}_{i=1\dots N} < \mu(X) < \max \{\mu_i\}_{i=1\dots N}$ holds, or all μ_i are equal to $\mu(X)$.*

Proof. (a): By induction, using the additivity of deg over short exact sequences, we find that $\deg(X) = \sum \deg_i$. Then we calculate:

$$\mu(X) = \frac{\deg(X)}{\mu(X)} = \frac{\sum \deg_i}{\text{rk}(X)} = \frac{\sum \text{rk}_i \mu_i}{\text{rk}(X)} = \sum \frac{\text{rk}_i}{\text{rk}(X)} \mu_i.$$

(b): By induction, using the additivity of rk over short exact sequences, we find that $\text{rk}(X) = \sum \text{rk}_i$. Therefore, each of the rational numbers $\text{rk}_i / \text{rk}(X)$ lies strictly between 0 and 1, and their sum is 1. Thus, (a) expresses $\mu(X)$ as a weighted average of the μ_i with weights strictly between 0 and 1 that sum to 1. The result follows. \square

(4.4) We will most often use (4.3.b) as follows: given a short strict exact sequence $0 \rightarrow X' \rightarrow X \rightarrow X'' \rightarrow 0$, exactly one of $\mu(X') < \mu(X) < \mu(X'')$, $\mu(X'') < \mu(X) < \mu(X')$, or $\mu(X') = \mu(X) = \mu(X'')$ holds. Therefore, to know one of the ordering relationships among these three slopes, it suffices to know either of the other two.

Proposition (4.5). *For a nonzero subobject X of \mathcal{C} , the following conditions are equivalent:*

- *For all nonzero proper subobjects X' of X , one has $\mu(X') \leq \mu(X)$ (resp. $\mu(X') < \mu(X)$).*
- *For all nonzero, proper, strict subobjects X' of X , one has $\mu(X') \leq \mu(X)$ (resp. $\mu(X') < \mu(X)$).*
- *For all nonzero, proper, strict subobjects X' of X , one has $\mu(X) \leq \mu(X/X')$ (resp. $\mu(X) < \mu(X/X')$).*

Proof. The first two conditions are equivalent by (4.1). The last two conditions are equivalent by (4.4). \square

(4.6) If the equivalent conditions of (4.5) hold for X , we say that X is *semistable*.

Proposition (4.7). *Let X be an object of \mathcal{C} , and X' a nonzero subobject whose slope is maximal among the slopes of nonzero subobjects of X . Then X' is semistable.*

In particular, if X has rank one, then X is semistable.

Proof. The first claim follows directly from the definition of semistability, since any subobject of X' is a subobject of X . For the second claim, assume X has rank one. By additivity and nonnegativity, any nonzero subobject of X has rank one, and by (3.2) it follows that the only nonzero strict subobject of X is X itself. By definition, then, X is semistable. \square

Proposition (4.8). *Let $0 \rightarrow X' \rightarrow X \rightarrow X'' \rightarrow 0$ be a short strict exact sequence in \mathcal{C} . If X' and X'' are both semistable of the same slope μ , then X is also semistable of slope μ . In particular, $X' \oplus X''$ is semistable of slope μ .*

Proof. By (4.4), $\mu(X) = \mu$. Let $Y \subseteq X$ be any strict subobject. Consider the short strict exact sequence

$$0 \rightarrow Y \cap X' \rightarrow Y \rightarrow Y/(Y \cap X') \rightarrow 0.$$

By (4.4), we must show that $\mu(Y \cap X') \leq \mu$ and $\mu(Y/(Y \cap X')) \leq \mu$. The first follows because X' is semistable of slope μ . The second follows because $Y/(Y \cap X')$ is a subobject of X'' by (2.10) and because X'' is semistable of slope μ . \square

Proposition (4.9). *Let X be an object of \mathcal{C} that is semistable of slope μ . Then any nonzero strict subobject or strict quotient of X that has slope μ is semistable.*

Proof. Let $X' \subseteq X$ be nonzero, strict, and of slope μ . Then any subobject of X' is also a subobject of X , and therefore has slope at most μ . Therefore X' is semistable. Dually, let $X \twoheadrightarrow X''$ be a nonzero, strict, and of slope μ . Then any strict quotient of X'' is also a strict quotient of X , and therefore has slope at least μ . Therefore X'' is semistable. \square

Proposition (4.10). *Let $f: X \rightarrow Y$ be a morphism in \mathcal{C} with X and Y semistable of the same slope μ . Then $\ker f$ and $\operatorname{img} f$ are semistable of slope μ , and $\operatorname{img} f$ is a strict subobject of Y .*

Proof. Since X (resp. Y) is semistable of slope μ , one has $\mu(\ker f) \leq \mu$ (resp. $\mu(\operatorname{img} f) \leq \mu$). But now (4.4) and the short strict exact sequence

$$0 \rightarrow \ker f \rightarrow X \rightarrow \operatorname{img} f \rightarrow 0$$

imply that $\mu(\ker f) = \mu(\operatorname{img} f) = \mu$. By (4.9), $\ker f$ and $\operatorname{img} f$ are semistable as well. Finally, by (4.2), $\operatorname{img} f \subseteq Y$ is strict if $\mu(\operatorname{img} f)$ is maximal among the slopes of subobjects of Y . Since Y is semistable of slope $\mu = \mu(\operatorname{img} f)$, this condition holds. \square

(4.11) Let μ be a slope. We denote by $\mathcal{C}^{(\mu)}$ the full subcategory of \mathcal{C} consisting of semistable objects of slope μ , together with the zero object.

Proposition (4.12). *For every slope μ , the category $\mathcal{C}^{(\mu)}$ is an abelian, exact subcategory of \mathcal{C} that is closed under extensions in \mathcal{C} .*

Proof. By (4.10), morphisms in $\mathcal{C}^{(\mu)}$ admit kernels and images. By (2.7) and (4.10), images are also coimages. The existence of cokernels follows. Hence it is abelian. By definition, a sequence is exact in $\mathcal{C}^{(\mu)}$ if and only if it is exact in \mathcal{C} ; this gives exactness. The final claim is (4.8). \square

§5. Subobjects of maximal slope.

This preparatory section contains the technical arguments that make HN filtrations work.

(5.1) Let X be an object of \mathcal{C} . We say a subobject X' of X is of *maximal slope* if it is nonzero and $\mu(X'') \leq \mu(X')$ for all nonzero subobjects X'' of X . For example, X is semistable if and only if X is itself a subobject of X of maximal slope. Recall that by (4.7) such a subobject X' is semistable, and by (4.2) it is a strict subobject of X .

Proposition (5.2). *Let X be an object of \mathcal{C} . Let X', X'' be subobjects of X of maximal slope. Then $X' + X''$ and $X' \cap X''$ are of maximal slope if nonzero.*

In particular, if X admits any subobjects of maximal slope, then X admits a unique final subobject of maximal slope.

Proof. By (4.2), X', X'' are strict. Since X' is semistable by (4.7), we have $\mu(X') \leq \mu(X'/(X' \cap X'')) = \mu((X' + X'')/X'')$ by (2.10). Since $\mu(X'')$ is maximal, we have $\mu(X' + X'') \leq \mu(X'')$, and by (4.4) we have $\mu((X' + X'')/X'') \leq \mu(X' + X'')$. Finally, since $\mu(X')$ is maximal, we have $\mu(X' + X'') \leq \mu(X')$. Summarizing, we have the inequalities

$$\mu(X') \leq \mu(X'/(X' \cap X'')) = \mu((X' + X'')/X'') \leq \mu(X' + X'') \leq \mu(X'),$$

which must all be equalities. The claim for $X' + X''$ immediately follows. On the other hand, when $X' \cap X''$ is nonzero, knowing that the first of these inequalities is an equality shows by (4.4) that we must also have $\mu(X' \cap X'') = \mu(X')$, whence the claim for $X' \cap X''$. The final claim follows from (2.1) and (3.1). \square

(5.3) Given an object X of \mathcal{C} , consider the condition on a nonzero subobject X' of X :

$$(5.3.1) \quad \text{for all subobjects } X'' \text{ of } X \text{ properly containing } X', \text{ one has } \mu(X'') < \mu(X').$$

By (4.1) such an X' is strict, and also it suffices to know (5.3.1) for strict X'' .

Proposition (5.4). *Let X be an object of \mathcal{C} , with nonzero strict subobjects X', X'' . Suppose X' satisfies (5.3.1), X'' is semistable, and $X'' \not\subseteq X'$. Then $\mu(X'') < \mu(X')$.*

Proof. Let $f: X'' \rightarrow X \rightarrow X/X'$ be the composition. By hypothesis, f is nonzero, and hence $F(f)$ is nonzero. Because X'' is semistable we have $\mu(X'') \leq \mu(\text{img } f)$. By (4.1), we have $\mu(\text{img } f) \leq \mu(\widetilde{\text{img } f})$. Let X^3 be the unique strict subobject of X (properly) containing X' with $X^3/X' = \widetilde{\text{img } f} \subseteq X/X'$. By (5.3.1), $\mu(X^3) < \mu(X')$. By (4.4), we must also have $\mu(\widetilde{\text{img } f}) < \mu(X^3)$. Putting everything together, we have $\mu(X'') \leq \mu(\text{img } f) \leq \mu(\widetilde{\text{img } f}) < \mu(X^3) < \mu(X')$. \square

Proposition (5.5). *Let X be an object of \mathcal{C} . The following conditions are equivalent for a nonzero subobject X' of X :*

- (a) X' satisfies (5.3.1) and is semistable.

(b) X' satisfies (5.3.1) and is of maximal slope.

(c) X' is the final subobject of X of maximal slope.

Proof. (a) \implies (b): Let X'' be a nonzero subobject of X . To show that $\mu(X'') \leq \mu(X')$, we may replace X'' by \tilde{X}'' and assume that X'' is a strict subobject. Since $X'' + X'$ contains X' , by (5.3.1) we have $\mu(X'' + X') \leq \mu(X')$. Since X' is semistable, by (2.10) we have $\mu(X') \leq \mu(X'/(X'' \cap X')) = \mu((X'' + X')/X'')$. Combining these, we have $\mu(X'' + X') \leq \mu((X'' + X')/X'')$, and now (4.4) implies that $\mu(X'') \leq \mu(X'' + X')$. To conclude, we combine the preceding inequality with another application of (5.3.1), $\mu(X'' + X') \leq \mu(X')$.

(b) \implies (a): This is (4.7).

(a,b) \implies (c): We must show that any nonzero subobject X'' of X having maximal slope is contained in X' . But this follows from the contrapositive (5.4), because (5.3.1) holds for X' , (4.7) shows X'' is semistable, and both X', X'' are of maximal slope so $\mu(X'') = \mu(X')$.

(c) \implies (a,b): We must check (5.3.1). Let X'' be a subobject of X properly containing X' . If it were possible that $\mu(X') \leq \mu(X'')$, then because $\mu(X')$ is maximal we would have $\mu(X'') = \mu(X')$. But X' is final among objects with slope $\mu(X')$, so this is impossible. \square

(5.6) Let X be an object of \mathcal{C} , and X' a subobject satisfying the equivalent conditions of (5.4). By (5.4.c), such an X' is uniquely determined, if it exists. By (4.2), X' is a strict subobject of X . It is easy to see that X is semistable if and only if $X' = X$. When $X' \subsetneq X$, by (5.4.c) X' is the maximal subobject that maximally contradicts the semistability of X . We call X' the *SCSS* for X , where “SCSS” stands for *strongly contradicting semistability*.

Proposition (5.7). *Every nonzero object X of \mathcal{C} admits an SCSS.*

Proof. We proceed by induction on the rank d of X . The base case is where X is semistable (which includes when $d = 1$ by (4.7)) is trivial, so we assume that $d > 1$ and X is not semistable. Because X is not semistable, there exist nonzero, proper submodules Y of X having $\mu(X) < \mu(Y)$. If we were to have $\text{rk}(Y) = \text{rk}(X)$, then by (3.2) we would have $\tilde{Y} = X$, and by (4.1) $\mu(Y) \leq \mu(X)$. Thus, any such Y must have rank strictly less than d .

Therefore, there is a maximal integer s among the ranks of nonzero proper submodules Y of X satisfying $\mu(X) < \mu(Y)$, and s satisfies $0 < s < d$. Fix any such submodule Y of rank s . Replacing Y by \tilde{Y} (which neither decreases $\mu(Y)$ nor changes $\text{rk}(Y)$), we may assume $Y \subseteq X$ is strict. Apply the inductive hypothesis to produce an SCSS Y' of Y . We verify that Y' is an SCSS for X , by showing that any nonzero strict subobject X' of X satisfying $\mu(Y') \leq \mu(X')$ is contained in Y' .

We have the short strict exact sequence by (2.8):

$$0 \rightarrow Y \cap X' \xrightarrow{(1,-1)} Y \oplus X' \rightarrow Y + X' \rightarrow 0.$$

Therefore, using $\mu(Y \cap X') \leq \mu(Y') \leq \mu(X')$, followed by $\mu(Y) \leq \mu(Y') \leq \mu(X')$, we have

$$\begin{aligned}
\mathrm{rk}(Y + X')\mu(Y + X') &= \deg(Y + X') = \deg(Y \oplus X') - \deg(Y \cap X') \\
&= \deg(Y) + \deg(X') - \deg(Y \cap X') \\
&= \mathrm{rk}(Y)\mu(Y) + \mathrm{rk}(X')\mu(X') - \mathrm{rk}(Y \cap X')\mu(Y \cap X') \\
&\geq \mathrm{rk}(Y)\mu(Y) + [\mathrm{rk}(X') - \mathrm{rk}(Y \cap X')]\mu(X') \\
&\geq [\mathrm{rk}(Y) + \mathrm{rk}(X') - \mathrm{rk}(Y \cap X')]\mu(Y) \\
&= \mathrm{rk}(Y + X')\mu(Y).
\end{aligned}$$

But $\mathrm{rk}(Y + X') \geq \mathrm{rk}(Y) = s > 0$, so $\mu(Y + X') \geq \mu(Y) > \mu(X)$. The maximality of s therefore implies that $\mathrm{rk}(Y + X') = \mathrm{rk}(Y)$. Now we have

$$Y \subseteq Y + X' \subseteq \widetilde{Y + X'} = \widetilde{Y} = Y,$$

with the penultimate identity coming from (3.2) and the final identity because Y is strict. But this forces $X' \subseteq Y$, and, Y' being an SCSS for Y , $\mu(Y') \leq \mu(X')$ forces $X' \subseteq Y'$. \square

Acknowledgment (5.8). We thank Ruochuan Liu for sharing the above argument with us.

§6. HN filtrations.

We now assemble the preceding facts to show the main results on HN filtrations: existence, uniqueness, and functoriality.

(6.1) Let X be a nonzero object of \mathcal{C} . By an *HN filtration* of X we mean an increasing, finite, separating and exhaustive filtration by strict subobjects

$$0 = X_0 \subsetneq X_1 \subsetneq \cdots \subsetneq X_N = X$$

with each $\mathrm{Gr}_i = X_i/X_{i-1}$ semistable, and each $\mu(\mathrm{Gr}_{i+1}) < \mu(\mathrm{Gr}_i)$. Showing the existence and uniqueness of the HN filtration is now straightforward, because the SCSS, studied in the preceding section, is its unique first step.

Proposition (6.2). *Let X be a nonzero object of \mathcal{C} . Then:*

- (a) *If X is not semisimple, and $0 = X_0 \subsetneq X_1 \subsetneq \cdots \subsetneq X_N = X$ is any HN filtration for X , then X_1 satisfies the equivalent conditions of (5.5) for an SCSS for X .*
- (b) *If a HN filtration for X exists, then it is unique.*

Proof. (a): We induct on the length N of the filtration. In the case $N = 1$, X is semistable, and we are done. Now assume $N > 1$. By hypothesis, X_1 is semistable. Given a strict subobject X'' of X properly containing X_1 , we must show that $\mu(X'') < \mu(X_1)$. Equivalently by (4.4), we must show that $\mu(X''/X_1) < \mu(X_1)$. Since $\mu(\mathrm{Gr}_2) < \mu(\mathrm{Gr}_1) = \mu(X_1)$, it suffices to show that any subobject of X/X_1 has slope at most $\mu(\mathrm{Gr}_2)$. But by inductive hypothesis, Gr_2 satisfies the equivalent conditions of (5.5) for X/X_1 , and in particular Gr_2 is a subobject of X/X_1 of maximal slope.

(b): By (a), for each $1 \leq i < N$, X_i/X_{i-1} is the SCSS of X/X_{i-1} , and then X_i is uniquely determined as the preimage of X_i/X_{i-1} in X . \square

Proposition (6.3). *Each nonzero object X of \mathcal{C} possesses a HN filtration.*

Proof. We perform induction on the rank of X . The base case is when X is semistable, which includes the case of rank one by (4.7), and we take $X_0 = 0$ and $X_1 = X$. Now suppose given X , assumed not semistable. Let X' be the SCSS of X , which exists by (5.7). By inductive hypothesis, X/X' has a HN filtration, $0 = Y_0 \subsetneq Y_1 \subsetneq \cdots \subsetneq Y_N = X/X'$. Take $X_0 = 0$, and for $1 \leq i \leq N+1$ let X_i be the preimage of Y_{i-1} in X . By construction, all the graded pieces are semistable. We only need to show that $\mu(\text{Gr}_2) < \mu(\text{Gr}_1)$. Considering the short strict exact sequence $0 \rightarrow \text{Gr}_1 \rightarrow X_2 \rightarrow \text{Gr}_2 \rightarrow 0$, by (4.4) it suffices to show that $\mu(X_2) < \mu(\text{Gr}_1)$, and this follows because $\text{Gr}_1 = X'$ satisfies (5.3.1). \square

(6.4) Each nonzero object X of \mathcal{C} has canonically associated to it a collection of slopes with ranks, called the *HN slopes* and their respective *ranks*: if $0 = X_0 \subsetneq X_1 \subsetneq \cdots \subsetneq X_N = X$ is the HN filtration, then we set $\text{Gr}_i = \text{Gr}_i(X) = X_i/X_{i-1}$, $\mu_i = \mu_i(X) = \mu(\text{Gr}_i)$, and $\text{rk}_i = \text{rk}_i(X) = \text{rk}(\text{Gr}_i)$. However, one only gets good functorial properties from using *slopes* to index the filtration. For any slope μ , let $X^{(\mu)}$ be the strict subobject of X defined by

$$X^{(\mu)} = \begin{cases} X = X_N & \text{if } \mu \leq \mu_N, \\ X_{N-1} & \text{if } \mu_N < \mu \leq \mu_{N-1}, \\ \vdots & \vdots \\ X_1 & \text{if } \mu_2 < \mu \leq \mu_1, \text{ and} \\ 0 & \text{if } \mu > \mu_1. \end{cases}$$

The $X^{(\mu)}$ define a decreasing, separating and exhaustive filtration indexed by the slopes. It has finitely many breaks, occurring precisely at the slopes of X . We let $\text{Gr}^{(\mu)} = \text{Gr}^{(\mu)}(X) = X^{(\mu)}/X^{(\mu+)}$, where $X^{(\mu+)}$ is the next smaller step in the HN filtration of X if μ is a break, and $X^{(\mu)}$ itself otherwise. Thus, the nonzero $\text{Gr}^{(\mu)}$ are precisely the Gr_i , indexed by their slopes: $\text{Gr}^{(\mu_i)} = \text{Gr}_i$.

Lemma (6.5). *Let X and Y be two nonzero objects of \mathcal{C} . Then:*

- (a) *If X and Y are semistable and $\text{Hom}_{\mathcal{C}}(X, Y) \neq 0$, then $\mu(X) \leq \mu(Y)$.*
- (b) *If all HN slopes of X are strictly greater than those of Y , then $\text{Hom}_{\mathcal{C}}(X, Y) = 0$.*
- (c) *Let $f \in \text{Hom}_{\mathcal{C}}(X, Y)$, and μ be a slope. The restriction of f to $X^{(\mu)}$ factors canonically through $Y^{(\mu)}$.*
- (d) *For each slope μ , the rule $X \mapsto \text{Gr}^{(\mu)}(X)$ gives a functor $\mathcal{C} \rightarrow \mathcal{C}^{(\mu)}$.*

Proof. (a): Suppose $f \in \text{Hom}_{\mathcal{C}}(X, Y)$ is nonzero. As X is semistable, one has $\mu(X) \leq \mu(\text{img } f)$, and since Y is semistable, one has $\mu(\text{img } f) \leq \mu(Y)$.

(b): Let X (resp. Y) have HN filtration $0 = X_0 \subsetneq X_1 \subsetneq \cdots \subsetneq X_N = X$ (resp. $0 = Y_0 \subsetneq Y_1 \subsetneq \cdots \subsetneq Y_M = Y$). We show by induction on N that any map $X \rightarrow Y$ must vanish. The base case, when X is semistable, is proved by induction on M . The base case on M , when Y is also semistable, is a restatement of (a). For the inductive step on M , given a map $X \rightarrow Y$ we apply the base case to see that the composite $X \rightarrow Y \rightarrow Y/Y_{M-1} = \text{Gr}_M(Y)$ vanishes,

hence the map $X \rightarrow Y$ factors through a map $X \rightarrow Y_{M-1}$, which also must vanish by inductive hypothesis. For the inductive step on N , we apply the base case to the composite $X_1 \rightarrow X \rightarrow Y$ to see that our given map $X \rightarrow Y$ factors through a map $X/X_1 \rightarrow Y$, which in turn vanishes by inductive hypothesis.

(c): Let f and μ be given. It suffices to replace X by $X^{(\mu)}$, f by its restriction to $X^{(\mu)}$, and Y by $Y/Y^{(\mu)}$, whence $X = X^{(\mu)}$ and $Y^{(\mu)} = 0$, and to show that $f = 0$. But now we are in the situation of (b), so indeed $f = 0$.

(d): This follows trivially from (c). \square

(6.6) Although (6.5) gives us, for any map $f: X \rightarrow Y$ in \mathcal{C} , a map $f^{(\mu)}: \text{Gr}^{(\mu)}(X) \rightarrow \text{Gr}^{(\mu)}(Y)$ in $\mathcal{C}^{(\mu)}$ for each slope μ , passing from f to the family of $f^{(\mu)}$ is very destructive in practice. For example, if X has rank one, and $\iota: X' \subseteq X$ is a nonzero proper subobject, then all the $\iota^{(\mu)}$ vanish.

§7. HN polygons.

We give some complements on HN polygons, which are an analogue of Newton polygons in this setting.

(7.1) Let $a, b \in \mathbf{Q}$. By $[a, b]_{\mathbf{Q}}$, we mean the interval $[a, b] \cap \mathbf{Q}$ of rational numbers. The set of slopes $V \otimes_{\mathbf{Z}} \mathbf{Q}$ is a \mathbf{Q} -vector space, so there is an obvious notion of a linear function on $[a, b]_{\mathbf{Q}}$ with values in $V \otimes_{\mathbf{Z}} \mathbf{Q}$. Similarly, we may speak of piecewise-linear functions on $[a, b]_{\mathbf{Q}}$ with values in $V \otimes_{\mathbf{Z}} \mathbf{Q}$, and we understand the breakpoints to be at rational numbers.

(7.2) Let X be a nonzero object of \mathcal{C} , and $0 = X_0 \subsetneq X_1 \subsetneq \cdots \subsetneq X_N = X$ its HN filtration. We define $\text{HN}(X)$ to be the unique concave-down, piecewise-linear function defined on the rational interval $[0, \text{rk}(X)]_{\mathbf{Q}}$, satisfying $\text{HN}(X)(0) = 0$, and whose i th slope is μ_i with horizontal width rk_i . The *HN polygon* of X is defined to be the graph of $\text{HN}(X)$. We call the point $(\text{rk}(X), \text{deg}(X))$ the *HN endpoint* of X ; it is easy to calculate that $\text{HN}(X)(\text{rk}(X)) = \text{deg}(X)$, thus explaining the terminology.

Note that $0 = X_0 \subsetneq X_1 \subsetneq \cdots \subsetneq X_i = X_i$ is the HN filtration of X_i . Therefore, $\text{HN}(X_i) = \text{HN}(X)|_{[0, \text{rk}(X_i)]_{\mathbf{Q}}}$. In particular, the breakpoints of the HN polygon of X are precisely the HN endpoints of the objects X_i .

Proposition (7.3). *Let X be a nonzero object of \mathcal{C} , with HN filtration $0 = X_0 \subsetneq X_1 \subsetneq \cdots \subsetneq X_N = X$. Then:*

- (a) *For any nonzero subobject X' of X , $\text{deg}(X') \leq \text{HN}(X)(\text{rk}(X'))$. In other words, the HN endpoint of any subobject of X lies on or below the HN polygon of X .*
- (b) *Moreover, if equality holds in (a), i.e. the HN endpoint of X' lies on the HN polygon of X , then $X_{i-1} \subseteq X' \subseteq X_i$ for i satisfying $\text{rk}(X_{i-1}) \leq \text{rk}(X') \leq \text{rk}(X_i)$.*
- (c) *The HN polygon of X is the upper convex hull of the points $(\text{rk}(X'), \text{deg}(X'))$, for X' ranging over its nonzero subobjects.*
- (d) *For any nonzero subobject X' of X , the HN polygon of X' lies on or below the HN polygon of X .*

Proof. (a,b): At any point in time, it suffices to replace X' by \tilde{X}' , so we may assume X' is a strict subobject of X . Let us induct on the length N of the HN filtration of X . In the case $N = 1$, X is semistable. But then $\mu(X') \leq \mu(X)$ and $\text{rk}(X') \leq \text{rk}(X)$ give (a); (b) is trivial.

Now assume $N > 1$. Note that if either $X' \subseteq X_1$ or $X_1 \subseteq X'$ holds, then we know (a,b) for X' . In the first case, one applies the base case above to X_1 , and, in the second case, the additivity of rk and \deg allow us to reduce to the case of X'/X_1 inside X/X_1 , to which the inductive hypothesis applies. By (6.2.a), we have $\mu(X') \leq \mu(X_1)$, and if equality holds then $X' \subseteq X_1$; thus we assume also that $\mu(X') < \mu(X_1)$. Also, since $\text{rk}(X' \cap X_1) = \text{rk}(X_1)$ implies $X_1 \subseteq X'$, we can assume that $\text{rk}(X' \cap X_1) < \text{rk}(X_1)$. And, since $X' + X_1$ contains X_1 , we know (a,b) for it.

Using, in order, (2.8), then the semistability of X_1 , then that $\mu(X') < \mu(X_1)$ but $\text{rk}(X_1) > \text{rk}(X' \cap X_1)$, we have

$$\begin{aligned} \text{rk}(X' + X_1)\mu(X' + X_1) &= \deg(X' + X_1) = \deg(X') + \deg(X_1) - \deg(X' \cap X_1) \\ &= \text{rk}(X')\mu(X') + \text{rk}(X_1)\mu(X_1) - \text{rk}(X' \cap X_1)\mu(X' \cap X_1) \\ &\geq \text{rk}(X')\mu(X') + [\text{rk}(X_1) - \text{rk}(X' \cap X_1)]\mu(X_1) \\ &> [\text{rk}(X') + \text{rk}(X_1) - \text{rk}(X' \cap X_1)]\mu(X') \\ &= \text{rk}(X' + X_1)\mu(X'). \end{aligned}$$

Since $\text{rk}(X' + X_1) > 0$, this implies $\mu(X') < \mu(X' + X_1)$. This inequality implies that the ray from the origin to the HN endpoint of X' lies strictly below the ray from the origin to the HN endpoint of $X' + X_1$, and the endpoint of X' clearly lies to the left of the endpoint of $X' + X_1$. By our knowledge of (a) for $X' + X_1$, we conclude that the HN endpoint of X' lies strictly beneath the HN polygon of X , whence (a) and (b).

(c): Given (a), it suffices to see that the breakpoints of the HN polygon are achieved. But the breakpoints are precisely the HN endpoints of the subobjects occurring in the HN filtration of X .

(d): Any subobject of X' is a subobject of X , so the claim follows from comparing the recipe of (c) as applied to X and X' . \square

(7.4) We give a common variant of the above. A *semistable filtration* of a nonzero object X of \mathcal{C} is a filtration F_\bullet : $0 = F_0 \subsetneq F_1 \subsetneq \cdots \subsetneq F_M = X$ by strict subobject such that each F_i/F_{i-1} is semistable. Its *polygon* $P(F_\bullet)$ is obtained just like the HN polygon is obtained from the HN filtration of X , except that one must first sort the slopes of the F_i/F_{i-1} into nonincreasing order.

Proposition (7.5). *Let X be a nonzero object of \mathcal{C} , and let $0 = F_0 \subsetneq F_1 \subsetneq \cdots \subsetneq F_M = X$ be a semistable filtration. Then $\text{HN}(X) \leq P(F_\bullet)$, and $\text{HN}(X)(\text{rk}(X)) = P(F_\bullet)(\text{rk}(X))$.*

Proof. The coincidence of endpoints of the two polygons follows immediately from the additivity of rk and \deg over short exact sequences. Let $0 = X_0 \subsetneq X_1 \subsetneq \cdots \subsetneq X_N = X$ denote the HN filtration of X . We first show that, for each index $i = 1, \dots, N$, one has $\text{HN}(X)(\text{rk}(X_i)) \leq P(F_\bullet)(\text{rk}(X_i))$.

Note that, if we count the slopes of F with multiplicities according to the ranks, then for an integer n with $0 \leq n \leq \text{rk}(X)$, $P(F_\bullet)(n)$ is characterized as the sum of the n largest slopes. Therefore, if we can find n slopes of F whose sum is greater or equal to C , then we know that $P(F_\bullet)(n) \geq C$. We apply this reasoning to $n = \text{rk}(X_i)$ and $C = \deg(X_i)$.

Fix i . For each $j = 1, \dots, M$, the subobject $F_j \cap X_i \subseteq X$ is strict by the last claim of (2.8). Using the strictness at $j-1$, we can form the quotient $(F_j \cap X_i)/(F_{j-1} \cap X_i)$. It is easy to check that the inclusion $F_j \cap X_i \subseteq F_j$ induces a morphism $(F_j \cap X_i)/(F_{j-1} \cap X_i) \rightarrow F_j/F_{j-1}$ that is an F -monomorphism, hence a monomorphism by (2.3). The semistability of F_j/F_{j-1} gives the first inequality in

$$\mu\left(\frac{F_j \cap X_i}{F_{j-1} \cap X_i}\right) \leq \mu(F_j/F_{j-1}) \quad \text{and} \quad d_j := \text{rk}\left(\frac{F_j \cap X_i}{F_{j-1} \cap X_i}\right) \leq \text{rk}(F_j/F_{j-1}).$$

We claim that choosing the j th slope of F with multiplicity d_j works. Indeed, the second inequality above shows that d_j does not exceed the multiplicity of the j th slope, and

$$\sum_{j=1}^M d_j = \sum_{j=1}^M [\text{rk}(F_j \cap X_i) - \text{rk}(F_{j-1} \cap X_i)] = \text{rk}(X_i).$$

Moreover,

$$\sum_{j=1}^M d_j \mu(F_j/F_{j-1}) \geq \sum_{j=1}^M d_j \mu\left(\frac{F_j \cap X_i}{F_{j-1} \cap X_i}\right) = \sum_{j=1}^M [\deg(F_j \cap X_i) - \deg(F_{j-1} \cap X_i)] = \deg(X_i).$$

Now, knowing that each $\text{HN}(X)(\text{rk}(X_i)) \leq \text{P}(F_\bullet)(\text{rk}(X_i))$, we conclude. For each i , consider the restrictions of $\text{HN}(X)$ and $\text{P}(F_\bullet)$ to interval $[\text{rk}(X_{i-1}), \text{rk}(X_i)]_{\mathbf{Q}}$. The restriction of $\text{HN}(X)$ is linear, and the restriction of $\text{P}(F_\bullet)$ is piecewise-linear, concave down, and with endpoints above those of $\text{HN}(X)$. Therefore, we must have $\text{HN}(X) \leq \text{P}(F_\bullet)$ over this interval. Since these intervals cover $[0, \text{rk}(X)]_{\mathbf{Q}}$, we are done. \square

Proposition (7.6). *Let $0 \rightarrow X' \rightarrow X \rightarrow X'' \rightarrow 0$ be a short strict exact sequence in \mathcal{C} , with X' and X'' nonzero. Then:*

(a) $\text{HN}(X) \leq \text{HN}(X' \oplus X'')$.

(b) *If all the slopes of X' are greater than or equal to all the slopes of X'' , then*

$$\text{HN}(X' \oplus X'')(x) = \begin{cases} \text{HN}(X')(x) & \text{if } 0 \leq x \leq \text{rk}(X'), \text{ and} \\ \text{HN}(X'')(x - \text{rk}(X')) + \deg(X') & \text{if } \text{rk}(X') \leq x \leq \text{rk}(X). \end{cases}$$

In other words, the HN polygon of $X' \oplus X''$ is obtained by joining the HN polygons of X' and X'' .

Proof. (a): By considering (7.3.c) applied to each of X and $X' \oplus X''$, it suffices to show that for any nonzero strict subobject Y of X , there exist subobjects Y' of X' and Y'' of X'' with $\text{rk}(Y) = \text{rk}(Y') + \text{rk}(Y'')$ and $\deg(Y) = \deg(Y') + \deg(Y'')$, because then the subobject $Y' \oplus Y''$ would bound the HN polygon of $X' \oplus X''$ above the HN polygon of X at $\text{rk}(Y)$. We may take $Y' = Y \cap X'$ and $Y'' = (Y + X')/X'$; the computation of ranks is immediate, and the computation of degrees follows from (2.10).

(b): The obvious filtration on $X' \oplus X''$ built from the HN filtrations on X' and X'' satisfies the characterizing properties of the HN filtration. (When X' and X'' share a common slope, one uses (4.8).) The claim follows from this. \square

(7.7) Given a sequence $0 \rightarrow X' \rightarrow X \rightarrow X'' \rightarrow 0$ as in (7.6), it is natural to ask if conditions on the slopes of X' and X'' generally guarantee that the extension X is split (resp. nonsplit). As far as I know, particular examples show that no such general conditions exist: for any slope combination, one can find both split and nonsplit extensions. In other words, although Harder–Narsimhan theory gives us information on Homs, such as in (6.5), it does not give us information on higher Exts.

§8. Examples: φ -modules.

We now give some variants of examples.

Common Inputs (8.1). All our examples will involve:

- a Bézout domain R with fraction field K , and
- an injective ring homomorphism $\varphi: R \rightarrow R$.

Note that φ extends uniquely to a ring homomorphism $K \rightarrow K$, via the rule $\varphi(r/s) = \varphi(r)/\varphi(s)$.

(8.2) Because R is a Bézout ring, any finitely generated, torsion-free R -module is free. In particular, a finitely generated submodule N of a finite free R -module M is free. Such N is moreover a strict subobject of M in the exact category of finite free R -modules if and only if it is a direct summand, if and only if it is *saturated*:

$$N = (K \otimes_R N) \cap M \text{ within } K \otimes_R M.$$

In other words, for finite free M , the natural maps

$$(8.2.1) \quad \left\{ \begin{array}{c} R\text{-direct summands} \\ \text{of } M \end{array} \right\} \subseteq \left\{ \begin{array}{c} \text{saturated} \\ R\text{-submodules} \\ \text{of } M \end{array} \right\} \xrightarrow{K \otimes_R -} \left\{ \begin{array}{c} K\text{-subspaces} \\ \text{of } K \otimes_R M \end{array} \right\}$$

are all bijections.

(8.3) If M and N are finite free R -modules and $f: M \rightarrow N$ is an R -module map, we say that f is an *isogeny* from M to N , written $f: M \xrightarrow{\sim} N$, if it induces an isomorphism $f: K \otimes_R M \xrightarrow{\sim} K \otimes_R N$. This is the case if and only if f is injective and there exists nonzero $r \in R$ such that $rN \subseteq f(M)$.

(8.4) For an R -module M , define $\varphi^*M = R_{\varphi} \otimes_R M$. The subscripts on the \otimes -symbol mean that $1 \otimes rm = \varphi(r) \otimes m$. We consider this object as an R -module using the left \otimes -factor. To give an R -linear map $\varphi^*M \rightarrow N$ is the same as to give a φ -linear map $f': M \rightarrow N$, that is, satisfying $f'(rm) = \varphi(r)f'(m)$. (Namely, $f'(m) = f(1 \otimes m)$ and $f(r \otimes m) = rf'(m)$.) If M is finite free, then so is φ^*M , of the same rank. Analogous definitions and claims apply to K -vector spaces instead of R -modules, compatibly with base change from R to K .

(8.4) We define a *weak φ -module over R* to be a finite free R -module M equipped with an isogeny $\varphi_M: \varphi^*M \xrightarrow{\sim} M$, and we let \mathcal{C}_w denote the exact category of these (with R -linear maps respecting the isogenies). Such (M, φ_M) is called a *φ -module over R* if φ_M is an isomorphism, and the strictly full, exact subcategory of these is written \mathcal{C} . Similarly, a *φ -module over K* is a finite-dimensional K -vector space M' equipped with an isomorphism $\varphi_{M'}: \varphi^*M' \xrightarrow{\sim} M'$, and the abelian category of these will be denoted \mathcal{D} .

(8.5) If M is an R -module and $k \geq 0$ an integer, we write $M^{\wedge k}$ for the k th exterior power of M . A map $f: M \rightarrow N$ of R -modules gives rise to a map $f^{\wedge k}: M^{\wedge k} \rightarrow N^{\wedge k}$ by the formula $f^{\wedge k}(m_1 \wedge \cdots \wedge m_k) = f(m_1) \wedge \cdots \wedge f(m_k)$. If M is a finite free R module of rank d , then $M^{\wedge k}$ is a finite free R -module of rank $\binom{d}{k}$. If f is an isogeny between free R -modules of rank d , then $f^{\wedge d}$ is an isogeny between free R -modules of rank 1.

In particular, if (M, φ_M) is a weak φ -module over R of rank d , then

$$\varphi_{M^{\wedge d}}: \varphi^*(M^{\wedge d}) \cong (\varphi^*M)^{\wedge d} \xrightarrow{\varphi_M^{\wedge d}} M^{\wedge d}$$

defines a weak φ -module $(M^{\wedge d}, \varphi_{M^{\wedge d}})$ over R of rank 1. Choosing a basis $v \in M^{\wedge d}$, we let $\det(\varphi_M) \in R$ be the nonzero element satisfying $\varphi_{M^{\wedge d}}(1 \otimes v) = \det(\varphi_M)v$. It is defined up to multiplication by $\varphi(r)/r$, for $r \in R^\times$. Note that (M, φ_M) is a φ -module if and only if $(M^{\wedge d}, \varphi_{M^{\wedge d}})$ is, if and only if $\det(\varphi_M) \in R^\times$.

Analogous definitions and claims apply to K -vector spaces instead of R -modules, compatibly with base change from R to K .

(8.6) We apply HN theory to:

$$\mathcal{C} = \mathcal{C}_w \text{ or } \mathcal{C}, \quad \mathcal{D} = \mathcal{D}, \quad F = K \otimes_R -, \quad \text{rk} = \dim_K,$$

and V and \deg to be specified below. Let us check the axioms that concern these data.

(8.7) We check the axiom for F . Faithfulness and exactness are clear, so we treat the remaining claim. Fix an object (M, φ_M) of \mathcal{C}_w or \mathcal{C} , and for brevity let $F(M, \varphi_M) = (M', \varphi_{M'})$. A subobject of (M, φ_M) in \mathcal{C}_w is uniquely determined by an R -submodule N of M such that $\varphi_M(\varphi^*N) \subseteq N$, and this subobject is strict if and only if N is saturated in M . Similarly, a subobject of $(M', \varphi_{M'})$ in \mathcal{D} is uniquely determined by a K -subspace N' of M' such that $\varphi_{M'}(\varphi^*N') \subseteq N'$ (whence comparing dimensions gives $\varphi_{M'}(\varphi^*N') = N'$). For a saturated R -submodule N of M that corresponds to the K -subspace N' of M' under (8.3.1), so that $N' = K \otimes_R N$ and $N = N' \cap M$, we must show that $\varphi_M(\varphi^*N) \subseteq N$ if and only if $\varphi_{M'}(\varphi^*N') \subseteq N'$. Indeed, if the latter holds then

$$\varphi_M(\varphi^*N) \subseteq \varphi_{M'}(\varphi^*N') \cap \varphi_M(\varphi^*M) \subseteq N' \cap M = N$$

as was desired, and the reverse implication is clear.

When (M, φ_M) belongs to \mathcal{C} , we must moreover show that when N' is a $\varphi_{M'}$ -stable K -subspace of M' , the saturated R -submodule $N = N' \cap M$ satisfies $\varphi_M(\varphi^*N) = N$. In fact, the above argument shows that N inherits a weak φ -module structure, and we also get a weak φ -module structure on M/N . Then one has $\det(\varphi_M) = \det(\varphi_N) \det(\varphi_{M/N})$, and $\det(\varphi_M)$ is a unit, so $\det(\varphi_N)$ is also a unit.

(8.8) The axiom for rk is trivial.

First Variant (8.9). We let V be a totally ordered abelian group (written additively), and $\text{ord}: K^\times \rightarrow V$ a homomorphism, satisfying:

- for all nonzero $r \in R$ one has $\text{ord}(\varphi(r)/r) \geq 0$, with equality if and only if $r \in R^\times$.

Then we take $\mathcal{C} = \mathcal{C}_w$, and $\deg(M, \varphi_M) = -\text{ord}(\det(\varphi_M))$.

(8.10) We verify the axiom for \mathcal{C}_w and this \deg . The additivity of \deg follows from multiplicativity of the determinant over short exact sequences, and that ord is a homomorphism. To verify the other claim, by passing to top exterior powers we may reduce to the case of an F -isomorphism $f: (M, \varphi_M) \rightarrow (N, \varphi_N)$ with M and N of rank 1; we identify M with its image in N under f . Choose a basis v of N and a nonzero $r \in R$ such that M is generated by rv . We have $\varphi_N(1 \otimes v) = \det(\varphi_N)v$ and

$$\det(\varphi_M) \cdot rv = \varphi_M(1 \otimes rv) = \varphi(r)\varphi_N(1 \otimes v) = \varphi(r)\det(\varphi_N)v = (\varphi(r)/r)\det(\varphi_N) \cdot rv,$$

so $\deg(M) = \deg(N) - \text{ord}(\varphi(r)/r)$. The claim is now seen to be equivalent to our axiom for ord .

Example (8.11). Suppose R is a valuation ring (and therefore a Bézout domain). The group $V = K^\times/R^\times$, called the value group of R , is totally ordered by $a \leq b$ if and only if $aR \supseteq bR$ within K . (We ignore that the operation of V is by default written multiplicatively.) The projection map $K^\times \rightarrow V$, called the valuation, will serve as our ord . For $r \in K^\times$ one has $\text{ord}(r) \geq 0$ if and only if $r \in R$, with equality if and only if $r \in R^\times$. For these data, our axiom becomes the following requirement on an injective ring homomorphism $\varphi: R \rightarrow R$: for all nonzero $r \in R$, one has $\varphi(r) \in rR$, and $\varphi(r)R = rR$ implies $r \in R^\times$.

Let us explicitly construct such an R and φ . Let k be a field with an automorphism $\sigma: k \xrightarrow{\sim} k$, X an indeterminate, and $q \geq 2$ an integer. Let $R = k[[X]]$, and $\varphi: \sum_{n \geq 0} a_n X^n \mapsto \sum_{n \geq 0} \sigma(a_n) X^{qn}$. Then R is a valuation ring for the X -adic valuation and the identity $\text{ord} \circ \varphi = q \text{ord}$ implies the required axiom.

Second Variant (8.12). Let V be a totally ordered abelian group (written additively), and $\text{ord}: R^\times \rightarrow V$ a homomorphism, satisfying:

- for all nonzero $r \in R$ such that $\varphi(r)/r \in R^\times$, one has $\text{ord}(\varphi(r)/r) \geq 0$, with equality if and only if $r \in R^\times$.

Then we take $\mathcal{C} = \mathcal{C}$ and $\deg(M, \varphi_M) = -\text{ord}(\det(\varphi_M))$.

(8.13) The axiom for \mathcal{C} and this \deg are verified just as in (8.10). (In the notation there, the fact that both $\det(\varphi_M)$ and $\det(\varphi_N)$ are units implies $\varphi(r)/r$ is also a unit, so $\deg(\varphi(r)/r)$ is defined.)

Example (8.14). Fix a prime p , and let $R \subset \mathbf{Q}_p[[X, X^{-1}]]$ be the Robba ring. This consists of Laurent series $f = \sum_{n \in \mathbf{Z}} a_n X^n$ convergent in some p -adic annulus of the form $\epsilon \leq |X| < 1$, where ϵ is allowed to depend on f . There is a unique continuous ring endomorphism φ that is the identity on coefficients and sends $X \mapsto (1 + X)^p - 1$. (The ϵ of convergence for $\varphi(f)$ may be closer to 1 than the ϵ of convergence for f .)

Some nontrivial facts include that R is a Bézout ring, and the following description of its units. Consider the subring $E \subset R$ of series f that, after perhaps adjusting ϵ , are not only convergent but bounded on the annulus $\epsilon \leq |X| < 1$. Then the inclusion $E^\times \subseteq R^\times$ is a bijection. Moreover, E is a (Henselian) discretely valued field with uniformizer p , and its valuation $\text{ord}: R^\times = E^\times \rightarrow \mathbf{Z}$ satisfies the hypothesis of (8.12). This example is the subject of [K].

Non-Example (8.15). Let k be a perfect field of characteristic $p > 0$. We take $V = \mathbf{Z}$, $R = W(k)$ the Witt vectors of k , φ to be the automorphism induced from the p th power Frobenius map on k , and ord to be the p -adic valuation. In this setting, \mathcal{C}_w is the category of so-called *F-crystals* over k . Then $\text{ord} \circ \varphi = \text{ord}$, so neither variant of this section applies. In fact, with our choice of \mathcal{D} and F the axiom for \deg always fails: for any nonzero object (M, φ_M) , the inclusion $pM \subseteq M$ is an F -isomorphism but not an isomorphism, whereas the two isomorphic objects $pM \cong M$ would have the same degree.

Nonetheless, there is an adaptation of HN theory to \mathcal{C}_w resembling the variant (8.9), using the p -adic valuation for ord_p to define $\deg(M, \varphi_M) = -\text{ord}_p(\det(\varphi_M))$, but replacing the use of $F: \mathcal{C}_w \rightarrow \mathcal{D}$ with other means. This, and more, is the subject of the Dieudonné–Manin Theorem.

References.

- [F] L. Fargues, La filtration de Harder–Narasimhan des schémas en groupes finis et plats. *J. Reine Angew. Math.* **645** (2010), 1–39.
- [K] K. S. Kedlaya, Slope filtrations revisited. *Doc. Math.* **10** (2005), 447–525.