

# LINEAR PULLBACK COMPONENTS OF THE SPACE OF CODIMENSION ONE FOLIATIONS

V. FERRER AND I. VAINSENER

**ABSTRACT.** The space of holomorphic foliations of codimension one and degree  $d \geq 2$  in  $\mathbb{P}^n$  ( $n \geq 3$ ) has an irreducible component whose general element can be written as a pullback  $F^*\mathcal{F}$ , where  $\mathcal{F}$  is a general foliation of degree  $d$  in  $\mathbb{P}^2$  and  $F : \mathbb{P}^n \dashrightarrow \mathbb{P}^2$  is a general rational linear map. We give a polynomial formula for the degrees of such components.

## INTRODUCTION

Codimension one holomorphic foliations in  $\mathbb{P}^n$  are defined by nonzero integrable twisted 1-forms, *i.e.*,  $\omega \in H^0(\mathbb{P}^n, \Omega_{\mathbb{P}^n}(d+2))$  satisfying

$$(1) \quad \omega \wedge d\omega = 0$$

Since  $\omega, \lambda\omega$  yield the same foliation for any  $\lambda \in \mathbb{C}^*$ , the space of such foliations is in fact a closed subscheme  $\mathbb{F}(d, n)$  of the projective space  $\mathbb{P}(H^0(\mathbb{P}^n, \Omega_{\mathbb{P}^n}(d+2)))$  defined by the equations derived from (1).

Explicitly, these equations are as follows. Any element  $\omega$  of  $H^0(\mathbb{P}^n, \Omega_{\mathbb{P}^n}(d+2))$  can be written  $\omega = A_0 dZ_0 + \cdots + A_n dZ_n$ , where  $A_i$  are homogeneous polynomials of degree  $d+1$  in the variables  $Z_0, \dots, Z_n$ , such that  $A_0 Z_0 + \cdots + A_n Z_n = 0$ . The integrability condition (1) imposes relations arising from

$$A_i \left( \frac{\partial A_k}{\partial Z_j} - \frac{\partial A_j}{\partial Z_k} \right) + A_j \left( \frac{\partial A_i}{\partial Z_k} - \frac{\partial A_k}{\partial Z_i} \right) + A_k \left( \frac{\partial A_j}{\partial Z_i} - \frac{\partial A_i}{\partial Z_j} \right) = 0.$$

with  $0 \leq i < j < k \leq n$ . These equations are quadratic in the coefficients of the polynomials  $A_i$ . For  $n = 2$  the space of foliations is a projective space: the integrability condition is automatically satisfied.

The geometry of the space of codimension one foliations in  $\mathbb{P}^n$  for  $n \geq 3$  is a rich field of research. In particular the problem of describing the irreducible components of these spaces has received many contributions, cf. [2], [4], [5], [12] and references therein just to quote a few.

In [11] Jouanolou describes the irreducible components for the space of foliations of degrees  $d = 0, 1$ . We have that  $\mathbb{F}(0, n)$  is naturally isomorphic to the grassmannian of subspaces of codimension 2 in  $\mathbb{P}^n$ . The space of foliations of degree  $d = 1$  has two irreducible components.

For foliations of degree 2 in  $\mathbb{P}^n$ ,  $n \geq 3$  Cerveau and Lins Neto [2] have shown that there are just 6 components. We recall some known components of  $\mathbb{F}(d, n)$  pinpointing those for which the degree has been found.

---

1991 *Mathematics Subject Classification.* 14N10, 14H40, 14K05.

*Key words and phrases.* holomorphic foliation, irreducible components, enumerative geometry. The authors were partially supported by CNPQ..

(1) **Logarithmic components.** Let  $F_1, \dots, F_m$  denote homogeneous polynomials in  $\mathbb{C}^{n+1}$ ,  $m \geq 3$ . Set  $d_i = \deg(F_i)$ , and let  $\lambda_1, \dots, \lambda_m \in \mathbb{C}^*$  be such that  $\sum_i \lambda_i d_i = 0$ . Then the form  $\omega = F_1 \dots F_m \sum \lambda_i \frac{dF_i}{F_i}$  is an integrable 1-form in  $\mathbb{P}^n$ . The foliation induced by  $\omega$  is called a logarithmic foliation. Suppose that  $F_1, \dots, F_m$  are irreducible and relatively prime. The space of this type of foliations is denoted by  $\mathcal{L}(d_1, \dots, d_m)$ . Its closure  $\overline{\mathcal{L}(d_1, \dots, d_m)}$  is an irreducible component of  $\mathbb{F}(d, n)$ , where  $d = d_1 + \dots + d_m - 2$ , cf. [1]. In the case  $d = 2$  we have two possibilities:  $\overline{\mathcal{L}(1, 1, 1, 1)}$  and  $\overline{\mathcal{L}(1, 1, 2)}$ . Their degrees have not been found to the best of our knowledge.

(2) **Rational components.** Let  $F, G$  be homogeneous polynomials such that

- $\deg(F) = l, \deg(G) = m, m + l - 2 = d$ ;
- $dF \wedge dG \neq 0$  along  $\{F = G = 0\} \setminus \{0\}$  and
- the hypersurface  $\{F = 0\} \subset \mathbb{P}^n$  is smooth.

Then  $\omega = mGdF - lFdG$  defines a foliation in  $\mathbb{P}^n$  of degree  $d$ . Denote by  $R(l, m)$  the set of foliations of this type. The closure  $\overline{R(l, m)}$  is an irreducible component of  $\mathbb{F}(d, n)$ , cf. Gómez-Mont and Lins Neto [8]. Observe that for  $d = 2$  we have two possibilities:  $l = m = 2$  or  $l = 1, m = 3$ . In this way we obtain two more components of  $\mathbb{F}(2, n)$ :  $\overline{R(2, 2)}$  and  $\overline{R(1, 3)}$ . The degrees of these components were computed in Cukierman-Pereira-Vainsencher [5].

(3) The exceptional component  $\mathcal{E}$  of  $\mathbb{F}(2, 3)$ . Here the leaves of a typical foliation are the orbits of a linear action of the affine group in one variable in  $\mathbb{P}^3$ . It is shown in [2] that for  $n > 3$ , in addition to the components described above there exists a component  $\mathcal{E}_n$  obtained by linear pullback  $\mathbb{P}^n \dashrightarrow \mathbb{P}^3$  of foliations in  $\mathcal{E}$ . The degree of  $\mathcal{E}$  has been found in [14].

(4) **Pullback components.** Let  $\mathcal{F}$  be a foliation of degree  $k$  in  $\mathbb{P}^2$ , defined by a 1-form  $\omega$ , and  $F : \mathbb{P}^n \dashrightarrow \mathbb{P}^2$  a dominant rational map of degree  $m$ . Then the 1-form  $F^*\omega$  defines a foliation  $\mathcal{F}^*$  of degree  $d = (k+2)m - 2$  in  $\mathbb{P}^n$ . In [3] Cerveau, Lins Neto and Edixhoven prove that for  $n \geq 3$  the closure of the set of foliations obtained by such  $F^*\omega$  form an irreducible component  $PB(m, k, n)$  of  $\mathbb{F}((k+2)m - 2, n)$ .

Our goal in this work is to compute the degrees and dimensions of the linear pullback components,  $LPB(d, n) := PB(1, d, n)$ . Note that for  $d = (k+2)m - 2 \leq 6$  the only pullback components in  $\mathbb{F}(d, n)$  are linear or quadratic. For  $d = 7$  there exists a component of cubic pullbacks. For  $d = 8, 9$  the only pullback components in  $\mathbb{F}(d, n)$  are linear or quadratic. If  $d+2$  is prime, then the only pullback component of  $\mathbb{F}(d, n)$  is linear pullback. Our main result is a polynomial formula for  $\deg LPB(d, n)$ , see 3.5.

## 1. PULL-BACK OF PROJECTIVE 1-FORMS.

**1.1. Projective 1-forms.** The main reference for this material is [11]. A projective 1-form of degree  $d$  in  $\mathbb{P}^n$  is a global section of  $\Omega_{\mathbb{P}^n}^1(d+2)$ , for some  $d \geq 0$ .

We denote by  $S_d$  the space  $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(d)) = \text{Sym}_d(\mathbb{C}^{n+1})^\vee$  of homogeneous polynomials of degree  $d$  in the variables  $Z_0, \dots, Z_n$ . We write  $\partial_i = \partial/\partial Z_i$  thought of as a vector field basis for  $\mathbb{C}^{n+1}$ . The dual basis will also be written as  $dZ_0, \dots, dZ_n$ .

whenever we think of differential forms. Twisting the Euler exact sequence (cf. [10, Thm 8.13, p. 176]): we get

$$0 \rightarrow \Omega_{\mathbb{P}^n}^1(d+2) \rightarrow \mathcal{O}_{\mathbb{P}^n}(d+1) \otimes S_1 \rightarrow \mathcal{O}_{\mathbb{P}^n}(d+2) \rightarrow 0.$$

Taking global sections we find the exact sequence

$$(2) \quad 0 \rightarrow V_d^n := H^0(\mathbb{P}^n, \Omega_{\mathbb{P}^n}^1(d+2)) \longrightarrow S_{d+1} \otimes S_1 \xrightarrow{\iota_{R_n}} S_{d+2} \rightarrow 0$$

where  $\iota_{R_n}(\sum A_i dZ_i) = \sum A_i Z_i$  is the contraction by the radial vector field. Thus a 1-form  $\omega \in V_d^n$  can be written in homogeneous coordinates as

$$\omega = A_0 dZ_0 + \cdots + A_n dZ_n$$

where the  $A_i$ 's are homogeneous polynomials of degree  $d+1$  satisfying

$$A_0 Z_0 + \cdots + A_n Z_n = 0.$$

**1.2. Linear Pullback.** Let  $F : \mathbb{P}^n \dashrightarrow \mathbb{P}^2$  be a linear projection, i.e.,  $F = [F_0 : F_1 : F_2]$ , with  $F_i$  linearly independent homogeneous polynomials of degree 1. Pick  $\omega \in H^0(\mathbb{P}^2, \Omega_{\mathbb{P}^2}^1(d+2))$ . Write  $\omega = B_0 dX_0 + B_1 dX_1 + B_2 dX_2$  where  $B_i \in S_{d+1}(X_0, X_1, X_2)$ . The pullback  $F^*(\omega)$  is the 1-form

$$(3) \quad F^*(\omega) = F^*B_0 dF_0 + F^*B_1 dF_1 + F^*B_2 dF_2.$$

A simple application of Euler relation shows that  $\iota_{R_n}(F^*(\omega)) = F^*(\iota_{R_2}(w)) = 0$ . On the other hand, as any projective 1-form in  $\mathbb{P}^2$  is integrable we have  $0 = F^*(\omega \wedge d\omega) = F^*(\omega) \wedge dF^*(\omega)$  i.e.,  $F^*(\omega)$  is a projective integrable 1-form of degree  $d$ ,

$$F^*(\omega) \in H^0(\mathbb{P}^n, \Omega_{\mathbb{P}^n}^1((d+2))).$$

**1.3. Remark.** For a fixed map  $F$  as above we obtain injective linear maps

$$H^0(\mathbb{P}^2, \Omega_{\mathbb{P}^2}^1(d+2)) \longrightarrow H^0(\mathbb{P}^n, \Omega_{\mathbb{P}^n}^1(d+2)),$$

$$\mathbb{F}(d, 2) \longrightarrow \mathbb{F}(d, n).$$

**1.4. Parameter space for rational maps.** Let  $F = [F_0 : F_1 : F_2]$  be a rational map as above, with  $F_i \in S_1$ . Note that if we change the basis of the linear system  $\langle F_0, F_1, F_2 \rangle$  then we obtain projectively equivalent pullbacks. So the natural parameter space for rational linear maps is the Grassmannian of dimension 3 subspaces of the space  $S_1$  of forms of degree 1,

$$(4) \quad \mathbb{G} := \mathbb{G}(3, S_1).$$

## 2. LINEAR PULLBACK COMPONENT

We show next that the locus in  $\mathbb{P}(V_d^n)$  (cf. 2) corresponding to codimension one foliations obtained by linear pullbacks of foliations in  $\mathbb{P}^2$  is the birational image of a natural projective bundle over the Grassmannian  $\mathbb{G}(3, S_1)$ .

**2.1. Proposition.** *Notation as in (2) and (4), let*

$$\mathcal{V}_d := \{(F, \mu) \in \mathbb{G} \times V_d^n \mid \mu = F^*\omega \text{ for some } \omega \in V_d^2\}.$$

*Then*

(i)  $\mathcal{V}_d$  is a vector subbundle of  $\mathbb{G} \times V_d^n$  of rank  $(d+1)(d+3)$ .

Let  $q_2 : \mathbb{P}(\mathcal{V}_d) \subset \mathbb{G} \times \mathbb{P}(V_d^n) \rightarrow \mathbb{P}(V_d^n)$  be the projection and set

$$\mathbb{Y} := q_2(\mathbb{P}(\mathcal{V}_d)), \quad g := \dim \mathbb{G} = 3(n-2).$$

*Then*

- (ii) the dimension of  $\mathbb{Y}$  is  $g + (d+1)(d+3) - 1$  and  
 (iii) the degree of  $\mathbb{Y}$  is given by the Segre class  $s_g(\mathcal{V}_d)$ .

*Proof.* Consider the tautological exact sequence of vector bundles over  $\mathbb{G}$ , (cf. 4)

$$(5) \quad 0 \rightarrow \mathcal{T} \rightarrow \mathbb{G} \times S_1 \rightarrow \mathcal{Q} \rightarrow 0$$

where  $\mathcal{T}$  is of rank 3 with fiber  $\mathcal{T}_F = \langle F_0, F_1, F_2 \rangle, F_i \in S_1$ . We obtain a natural rational map,

$$\begin{array}{ccc} \mathbb{G} \times \mathbb{P}^n - \overline{F} & \xrightarrow{\quad} & \mathbb{P}(\mathcal{T}^\vee) \\ & \searrow p_1 & \downarrow \psi \\ & & \mathbb{G} \end{array}$$

We interpret the map  $\overline{F}$  as the family of  $F$ 's as in 1.2 and  $\mathbb{P}(\mathcal{T}^\vee)$  as the family of  $\mathbb{P}^2$ 's obtained as the fibers of  $\psi$ . The relative cotangent bundle of  $\psi$  fits into the relative Euler exact sequence (cf. [6, B.5.8, p.435]):

$$0 \rightarrow \Omega_\psi(1) \rightarrow \psi^* \mathcal{T} \rightarrow \mathcal{O}_{\mathcal{T}^\vee}(1) \rightarrow 0.$$

Twisting by  $\mathcal{O}_{\mathcal{T}^\vee}(d+1)$  we find

$$0 \rightarrow \Omega_\psi(d+2) \rightarrow \psi^* \mathcal{T}(d+1) \rightarrow \mathcal{O}_{\mathcal{T}^\vee}(d+2) \rightarrow 0.$$

Taking direct image yields the following exact sequence over  $\mathbb{G}$

$$(6) \quad 0 \rightarrow \psi_* \Omega_\psi(d+2) \rightarrow \text{Sym}_{d+1} \mathcal{T} \otimes \mathcal{T} \rightarrow \text{Sym}_{d+2}(\mathcal{T}) \rightarrow 0.$$

Define

$$(7) \quad \mathcal{V}_d := \psi_* \Omega_\psi(d+2).$$

The fiber of  $\mathcal{V}_d$  over each  $F \in \mathbb{G}$ , is

$$H^0(\mathbb{P}(\mathcal{T}_F^\vee), \Omega_{\mathbb{P}(\mathcal{T}_F^\vee)}(d+2))$$

i.e., the space of 1-forms defining foliations of degree  $d$  in the varying  $\mathbb{P}^2 \simeq \mathbb{P}(\mathcal{T}_F^\vee)$ .

On the other hand, we obtain from (5) injective maps of vector bundles over  $\mathbb{G}$ ,

$$\iota_1 : \text{Sym}_{d+1} \mathcal{T} \otimes \mathcal{T} \longrightarrow S_{d+1} \otimes S_1$$

and

$$\iota_2 : \text{Sym}_{d+2} \mathcal{T} \longrightarrow S_{d+2}.$$

These two maps fit into the following diagram of exact sequences

$$\begin{array}{ccccc} \ker(a) = \mathcal{V}_d & \xrightarrow{j} & \ker(b) = V_d^n & & \\ \downarrow & & \downarrow & & \\ 0 \longrightarrow & \text{Sym}_{d+1} \mathcal{T} \otimes \mathcal{T} & \xrightarrow{\iota_1} & S_{d+1} \otimes S_1 & \\ \downarrow a & & & \downarrow b & \\ 0 \longrightarrow & \text{Sym}_{d+2} \mathcal{T} & \xrightarrow{\iota_2} & S_{d+2} & \end{array}$$

In this way we obtain an injective map of vector bundles over  $\mathbb{G}$ ,

$$(8) \quad j : \mathcal{V}_d \longrightarrow \mathbb{G} \times V_d^n.$$

The vector subbundle  $\mathcal{V}_d \subset \mathbb{G} \times V_d^n$  is as stated in (i).

Let  $q_1 : \mathbb{P}(\mathcal{V}_d) \rightarrow \mathbb{G}$  and  $q_2 : \mathbb{P}(\mathcal{V}_d) \rightarrow \mathbb{P}(V_d^n)$  be the maps induced by projection:

$$\begin{array}{ccc} & \mathbb{P}(\mathcal{V}_d) & \\ q_1 \swarrow & & \searrow q_2 \\ \mathbb{G} & & \mathbb{Y} \subset \mathbb{P}(V_d^n) \end{array}.$$

We prove in Lemma 2.3 below that  $q_2 : \mathbb{P}(\mathcal{V}_d) \rightarrow \mathbb{P}(V_d^n)$  is generically injective. Set  $u := \dim \mathbb{Y} = \dim \mathbb{P}(\mathcal{V}_d)$ . Write  $H$  for the hyperplane class of  $\mathbb{P}(V_d^n)$ . We have  $q_2^*H = c_1 \mathcal{O}_{\mathcal{V}_d}(1) =: h$ . Using [6, §3.1, p. 47, Prop. 4.4, p. 83 and Ex. 8.3.14, p. 143], we may compute

$$\deg \mathbb{Y} = \int_{\mathbb{P}(V_d^n)} H^u \cap \mathbb{Y} = \int_{\mathbb{P}(\mathcal{V}_d)} h^u = \int_{\mathbb{G}} q_{1*}(h^u) = \int_{\mathbb{G}} s_g(\mathcal{V}_d).$$

□

**2.2. Remark** From sequence (6) we obtain  $s(\mathcal{V}_d) = s(\text{Sym}_{d+1} \mathcal{T} \otimes \mathcal{T})c(\text{Sym}_{d+2}(\mathcal{T}))$ .

**2.3. Lemma.** For general  $(F, \omega) \in \mathbb{G} \times H^0(\mathbb{P}^2, \Omega_{\mathbb{P}^2}^1(d+2))$ ,  $F^*(\omega)$  determines uniquely  $\omega$  and  $F$ .

*Proof.* With notation as in 1.2, let  $I(F) = Z(F_0, F_1, F_2) \subset \mathbb{P}^n$  denote the indeterminacy locus of  $F$ . For generic  $(F, \omega)$ , the singular set of  $F^*(\omega)$  consists of linear components of codimension two of the form  $F^{-1}(q)$  where  $q \in \text{Sing}(\omega)$ . These linear components intersect in  $I(F)$ . Indeed, we can suppose  $q_1 = [0 : 0 : 1]$  and  $q_2 = [0 : 1 : 0]$ , so  $F^{-1}(q_1) = Z(F_0, F_1)$  and  $F^{-1}(q_2) = Z(F_0, F_2)$ . From  $I(F)$  we retrieve the linear system  $\langle F_0, F_1, F_2 \rangle$  i.e., the point  $F \in \mathbb{G}$ .

On the other hand, consider the blow-up  $\pi : \mathbb{B} \rightarrow \mathbb{P}^n$  of  $\mathbb{P}^n$  in  $I(F)$ . We proceed to show that it is possible to recover the 1-form  $\omega$  from the strict transform  $\tilde{\omega}$  of  $F^*(\omega)$ . Indeed, recall  $\mathbb{B} = \{(p, [x_0 : x_1 : x_2]) \in \mathbb{P}^n \times \mathbb{P}^2 \mid x_i F_j(p) - x_j F_i(p) = 0\}$ . Therefore over  $U := \{x_0 \neq 0\}$  we have  $\mathbb{B}|_U = \{(p, [1 : t : s]) \mid F_1(p) = tF_0(p); F_2(p) = sF_0(p)\}$  and the equation of the exceptional divisor is  $F_0 = 0$ . In this chart we have  $dF_1 = F_0 dt + t dF_0$ ,  $dF_2 = F_0 ds + s dF_0$ , therefore

$$\pi^* F^*(\omega) = B_0(F_0, tF_0, sF_0) dF_0 + B_1(F_0, tF_0, sF_0) dF_1 + B_2(F_0, tF_0, sF_0) dF_2 =$$

$$F_0^{d+1} [B_0(1, t, s) dF_0 + B_1(1, t, s)(F_0 dt + t dF_0) + B_2(1, t, s)(F_0 ds + s dF_0)] =$$

$$F_0^{d+1} [(B_0(1, t, s) + tB_1(1, t, s) + sB_2(1, t, s)) dF_0 + B_1(1, t, s) F_0 dt + B_2(1, t, s) F_0 ds]$$

Recalling  $\omega$  is a projective 1-form, so  $B_0(1, t, s) + tB_1(1, t, s) + sB_2(1, t, s) = 0$ , we obtain

$$\pi^* F^*(\omega) = F_0^{d+2} [B_1(1, t, s) dt + B_2(1, t, s) ds].$$

Therefore the strict transform of  $F^*\omega$  is  $\tilde{\omega} = B_1(1, t, s) dt + B_2(1, t, s) ds$ . Ditto for the other local charts of the blowup. This shows that we may recover  $\omega$  from  $\tilde{\omega}$ . □

**2.4. Corollary.** The component  $LPB(d, n) \subset \mathbb{F}(d, n)$  is rational.

### 3. COMPUTATION OF THE DEGREE OF $LPB(d, n)$

**3.1. Proposition.** *Notation as in 2.1,  $\deg(LP B(d, n))$  is a polynomial in  $d$  of degree  $3g = 9(n - 2)$ .*

We need some preliminary results.

**3.2. Lemma.** *Let  $E$  be a vector bundle of rank  $r$  on a variety  $X$ . The  $k$ -Segre class*

$$s_k(\mathrm{Sym}_d(E)) = \sum_{|\lambda|=k} p_\lambda(d) c_\lambda(E)$$

where  $p_\lambda(d)$  is a polynomial in  $d$  of degree  $\leq rk$ , and there exists  $p_\lambda$  of degree  $rk$ . Here the sum runs over the partitions of  $k$ , and if  $\lambda = (\lambda_1, \dots, \lambda_l)$  is a partition of  $k$ ,  $c_\lambda(E) := c_{\lambda_1}(E) \cdots c_{\lambda_l}(E)$ .

*Proof.* First we prove the following assertion relating the Chern characters to the Segre classes of  $\mathrm{Sym}_d(E)$ .

**3.3. Claim A(k):** *Assume the Chern character*

$$ch_j(\mathrm{Sym}_d(E^\vee)) = \sum_{|\mu|=j} q_\mu(d) c_\mu(E)$$

where  $q_\mu(d)$  are polynomials in  $d$  of degree  $\leq r + j - 1$  for all  $j \geq 0$ . Then  $s_k(\mathrm{Sym}_d(E)) = \sum_{|\lambda|=k} p_\lambda(d) c_\lambda(E)$  where  $p_\lambda(d)$  are polynomials in  $d$  of degree  $\leq rk$ . Moreover, if  $ch_1(\mathrm{Sym}_d(E)) = q_1(d) c_1(E)$  and  $\deg(q_1) = r$ , then  $s_k(\mathrm{Sym}_d(E))$  has a coefficient of degree  $rk$ .

For a vector bundle  $F$  of rank  $r$ , let  $x_1, \dots, x_r$  be the Chern roots of  $F^\vee$ . Then for all  $k \geq 0$ , the Segre class

$$s_k(F) = \sum_{1 \leq i_1 \leq \dots \leq i_k \leq r} x_{i_1} \cdots x_{i_k}$$

is the  $k$ -complete symmetric function (cf. [7, p. 28]).

On the other hand the complete symmetric functions can be expressed in term of the power sum symmetric functions,  $p_k = \sum_i x_i^k$ . We borrow from ([13, p. 25]) the explicit relations:

$$(9) \quad s_k(F) = \sum_{|\lambda|=k} w_\lambda ch_\lambda(F^\vee)$$

where  $\lambda = (\lambda_1, \dots, \lambda_l)$  is a partition of  $k$ . Following the notation in [13] we write  $\lambda = (1^{m_1}, 2^{m_2}, \dots)$  where  $m_i := \#\{j \mid \lambda_j = i\}$  and  $w_\lambda = \prod_{i \geq 1} \frac{\lambda_i!}{i^{m_i} m_i!}$ .

Write for short  $ch_j = ch_j(\mathrm{Sym}_d(E^\vee))$ . Whenever the coefficients of  $ch_j$  are polynomials in  $d$  of degree  $m$  we will write  $\deg(ch_j) = m$ , by abuse of notation. We observe that  $\deg(ch_\lambda) = \deg(ch_{\lambda_1} \cdots ch_{\lambda_l}) \leq l(r - 1) + k \leq rk$  and the equality holds if and only if  $l = k$ , i.e. if  $\lambda = (1, \dots, 1)$  in which case the coefficient of  $ch_1^k$  is  $\frac{1}{k!}$ . In other words

$$s_k(\mathrm{Sym}_d(E)) = \frac{1}{k!} ch_1^k + \text{l.o.t}$$

Hence  $s_k(\mathrm{Sym}_d(E))$  is a linear combination of monomials in the Chern classes of  $E$  whose coefficients are polynomials in  $d$  of degree  $\leq rk$ .

Next we prove the following claim by induction on  $r = \mathrm{rank}(E)$  and on  $k$ :

**3.4. Claim  $\mathbf{P}(\mathbf{k})$ :** For  $k \geq 0$ ,  $ch_k(\text{Sym}_d(E)) = \sum_{|\mu|=j} q_\mu(d) c_\mu(E)$ , where  $q_\mu(d)$  are polynomials in  $d$  of degree  $\leq r + k - 1$ . Moreover,  $ch_1(\text{Sym}_d(E)) = q_1(d) c_1(E)$  where  $q_1(d)$  is a polynomial of degree  $r$ .

For  $r = 1$  we have  $ch_k(\text{Sym}_d(E)) = \frac{1}{k!} d^k c_1(E)^k$ .

Suppose that  $\mathbf{P}(\mathbf{k})$  is true for vector bundles of rank  $r - 1$ . Let  $\pi : E \rightarrow X$  be a vector bundle of rank  $(E) = r$  and  $p : \mathbb{P}(E) \rightarrow X$  the induced projective bundle. For  $k = 0$  we have  $ch_0(\text{Sym}_d(E)) = \text{rank}(\text{Sym}_d(E)) = \binom{d+r-1}{r-1}$ , a polynomial in  $d$  of degree  $r - 1$ . Suppose that  $k \geq 1$ , and that  $\mathbf{P}(\mathbf{s})$  holds for  $\mathbf{s} < \mathbf{k}$ .

Over  $\mathbb{P}(E)$  we have the tautological exact sequence:

$$(10) \quad 0 \rightarrow \mathcal{O}_E(-1) \rightarrow p^*E \rightarrow Q \rightarrow 0.$$

It induces the following exact sequence for  $d \geq 1$

$$0 \rightarrow \mathcal{O}_E(-1) \otimes \text{Sym}_{d-1}(p^*E) \rightarrow \text{Sym}_d(p^*E) \rightarrow \text{Sym}_d(Q) \rightarrow 0.$$

Hence we may write the relation for the Chern characters

$$(11) \quad ch(\text{Sym}_d(p^*E)) = ch(\mathcal{O}_E(-1) \otimes \text{Sym}_{d-1}(p^*E)) + ch(\text{Sym}_d(Q)).$$

On the other hand,  $ch(\mathcal{O}_E(-1) \otimes \text{Sym}_{d-1}(p^*E)) = ch(\mathcal{O}_E(-1)) ch(\text{Sym}_{d-1}(p^*E))$ . So each graded part satisfies

$$(12) \quad ch_k(\mathcal{O}_E(-1) \otimes \text{Sym}_{d-1}(p^*E)) = \sum_{i=0}^k \frac{\alpha^i}{i!} ch_{k-i}(\text{Sym}_{d-1}(p^*E))$$

where  $\alpha := c_1(\mathcal{O}_E(-1))$ . It follows from (11) and (12) that

$$ch_k(\text{Sym}_d(p^*E)) = \sum_{i=0}^k \frac{\alpha^i}{i!} ch_{k-i}(\text{Sym}_{d-1}(p^*E)) + ch_k(\text{Sym}_d(Q)).$$

Hence

$$(13) \quad ch_k(\text{Sym}_d(p^*E)) - ch_k(\text{Sym}_{d-1}(p^*E)) = \sum_{i=1}^k \frac{\alpha^i}{i!} ch_{k-i}(\text{Sym}_{d-1}(p^*E)) + ch_k(\text{Sym}_d(Q)).$$

Observe that the right hand side of (13) involves:

- $ch_k(\text{Sym}_d(Q))$ , which by induction, since  $\text{rank}(Q) = r - 1$ , is of the form  $\sum_{|\mu|=k} q_\mu(d) c_\mu(Q)$  where  $q_\mu(d)$  are polynomials in  $d$  of degree  $\leq r - 1 + k - 1 = r + k - 2$ . Moreover, by (10)  $c_s(Q) = c_s(p^*E) - \alpha c_{s-1}(p^*E)$ . Thus  $ch_k(\text{Sym}_d(Q))$  is a linear combination of monomials in the Chern classes of  $p^*E$  and in  $\alpha$  whose coefficients are polynomials in  $d$  of degree  $\leq r + k - 2$ .
- $ch_s(\text{Sym}_{d-1}(p^*E))$  with  $s < k$  which, by induction on  $k$  is of the form  $\sum_{|\mu|=s} q_\mu(d) c_\mu(p^*(E))$  where  $q_\mu(d)$  are polynomials in  $d$  of degree  $\leq r + s - 1$ ; the maximal degree appearing is  $r + k - 2$  (coming from the coefficients in  $\alpha ch_{k-1}(\text{Sym}_{d-1}(p^*E))$ ).

Recall that the pullback  $p^* : A_*(X) \rightarrow A_*(\mathbb{P}(E))$  is a monomorphism with left inverse  $\beta \mapsto p_*(c_1(\mathcal{O}_E(1))^{r-1} \cap (\beta))$  ([6, p. 49]). Applying this inverse to (13) we conclude that

$$ch_k(\text{Sym}_d(E)) - ch_k(\text{Sym}_{d-1}(E))$$

is a linear combination of monomials in the Chern classes of  $E$  whose coefficients are polynomials in  $d$  of degree  $\leq r + k - 2$ , and this implies that the coefficients in  $ch_k(\text{Sym}_d(E))$  are polynomials in  $d$  of degree  $\leq r + k - 1$ .

Observe that for  $k = 1$  we obtain that  $ch_1(\text{Sym}_d(E)) - ch_1(\text{Sym}_{d-1}(E)) = \binom{d+r-2}{r-1}c_1(E)$ , a polynomial of degree  $r - 1$ . So  $ch_1(\text{Sym}_d(E)) = q_1(d)c_1(E)$  where  $q_1(d)$  is polynomial of degree  $r$ .

Using **A(k)** 3.3, we deduce that  $s_k(\text{Sym}_d(E))$  is a linear combination of monomials in the Chern classes of  $E$  whose coefficients are polynomials in  $d$  of degree  $\leq \text{rank}(E)k$  and there exists a coefficient of degree  $\text{rank}(E)k$ .  $\square$

Next we prove Proposition 3.1.

*Proof.* From the exact sequence

$$0 \rightarrow \mathcal{V}_d \rightarrow \text{Sym}_{d+1} \mathcal{T} \otimes \mathcal{T} \rightarrow \text{Sym}_{d+2}(\mathcal{T}) \rightarrow 0$$

we obtain the relation for Chern characters

$$ch(\mathcal{V}_d^\vee) = ch(\text{Sym}_{d+1} \mathcal{T}^\vee \otimes \mathcal{T}^\vee) - ch(\text{Sym}_{d+2}(\mathcal{T}^\vee)).$$

Thus for each  $k \geq 1$  we have

$$ch_k(\mathcal{V}_d^\vee) = \sum_{i=0}^k ch_{k-i}(\text{Sym}_{d+1} \mathcal{T}^\vee) ch_i(\mathcal{T}^\vee) - ch_k(\text{Sym}_{d+2}(\mathcal{T}^\vee)).$$

By assertion **P(k)** (3.4)  $ch_k(\text{Sym}_d(\mathcal{T}^\vee))$  is a polynomial in  $d$  of degree  $\leq k + 2$ . Therefore  $ch_k(\mathcal{V}_d^\vee)$  is a polynomial in  $d$  of degree  $\leq k + 2$ . Moreover,

$$ch_1(\text{Sym}_d(\mathcal{T}^\vee)) = q_1(d)c_1(\mathcal{T}^\vee)$$

where  $q_1(d)$  is a polynomial of degree 3. Therefore

$$\begin{aligned} ch_1(\mathcal{V}_d^\vee) &= ch_1(\text{Sym}_{d+1} \mathcal{T}^\vee) ch_0(\mathcal{T}^\vee) + ch_0(\text{Sym}_{d+1} \mathcal{T}^\vee) ch_1(\mathcal{T}^\vee) - ch_1(\text{Sym}_{d+2}(\mathcal{T}^\vee)) = \\ &= (3q_1(d+1) + \binom{d+3}{2} - q_1(d+2))c_1(\mathcal{T}^\vee) \end{aligned}$$

is a polynomial in  $d$  of degree 3.

As in the proof of assertion **A(k)** (3.3), we conclude that  $s_k(\mathcal{V}_d)$  is polynomial in  $d$  of degree  $3k$ .  $\square$

**3.5. Some formulas.** To get explicit formulas for  $s_g(\mathcal{V}_d)$  for any fixed  $d, n$  we use sequence (6), and Macaulay2, [9]. We find (cf. script §5 below)

$$\deg(LPB(d, 3)) = \frac{20}{27} \binom{d+4}{5} (d^2 + 6d + 11)(d^2 + 2d + 3).$$

$$\begin{aligned} \deg(LPB(d, 4)) &= \frac{1}{839808} \frac{(d+4)!}{(d-1)!} (8d^{12} + 192d^{11} + 2176d^{10} + 15360d^9 + 75090d^8 + \\ &+ 267552d^7 + 711859d^6 + 1423716d^5 + 2119892d^4 + 2279136d^3 + 1662291d^2 + 730188d + \\ &+ 125388)(2 + d). \end{aligned}$$

#### 4. HIGHER DEGREE PULLBACK COMPONENTS

As stated in the Introduction, the set of foliations obtained by pullback of foliations in  $\mathbb{P}^2$  by rational maps of degree  $m$  also form an irreducible component  $PB(m, k, n)$  of  $\mathbb{F}(d, n)$ ,  $d := (k + 2)m - 2$ .

As in the linear case, a natural parameter space of rational maps  $F : \mathbb{P}^n \dashrightarrow \mathbb{P}^2$  of degree  $m$  is the grassmannian  $\mathbb{G}(3, S_m)$ . Mimicking 2.1, we can construct a fiber bundle

$$\mathcal{V}_{m,k} := \{(F, \mu) \in \mathbb{G}(3, S_m) \times V_k^n \mid \mu = F^* \omega \text{ for some } \omega \in V_k^2\}.$$



However, for  $m \geq 2$ , the map  $j : \mathcal{V}_{m,k} \rightarrow V_d^n$  (analogous to the map in (8)) is injective only over the open subset consisting of dominant maps. So its image is not a subbundle of  $V_d^n$ . In fact, for a non-dominant map  $F \in \mathbb{G}(3, S_m)$ ,  $j_F(\omega) = F^*(\omega) = 0$  for all  $\omega \in H^0(\mathbb{P}^2, \Omega_{\mathbb{P}^2}^1(k+2))$  defining a foliation that leaves invariant the closure of  $\text{Im}(F)$ .

We could in principle find the locus  $Z \subset \mathbb{G}(3, S_m)$  where the rank of  $\text{Im}(j)$  drops and then blow-up  $\mathbb{G}(3, S_m)$  along  $Z$ . Doing this we expect to build a subbundle  $\tilde{\mathcal{V}}_k$  of  $V_d^n$  over the blow-up that coincides with  $\text{Im}(j)$  over the open set of dominant maps. The projection of  $\mathbb{P}(\tilde{\mathcal{V}}_k)$  to  $\mathbb{P}(V_d^n)$  is the space of foliations obtained as pullback by dominant maps. To make it work, we'd need to know how to get our hands on the Segre classes of  $Z$ . We hope to report on this elsewhere.

## 5. SCRIPTS

**5.1. Scripts for Macaulay2.** In order to compute  $\deg(LPB(d, n))$  for any given value of  $n \geq 3$ , just set  $N = n$  at the beginning of the script below. It can be fed into <http://habanero.math.cornell.edu:3690/>.

```
loadPackage "Schubert2"
N=3 --plug-in 3,4...
pt= base d
-- set d to be a free parameter in the "intersection ring"
--of the base variety
G=flagBundle({3,N-2}, pt)
-- Grassmannian of 3-planes in N+1-space,
(S,Q)=G.Bundles
-- names the sub and quotient bundles on G
A=symmetricPower(d+2, S)
B=symmetricPower(d+1, S)*S
integral(chern(A-B))
```

**Acknowledgments.** We wish to thank Luca Scala for the suggestions that led to a significant improvement of the proof of Proposition 3.1.

## REFERENCES

- [1] O. Calvo-Andrade. *Irreducible components of the space of holomorphic foliations*. Math. Ann. 299 (1994), no. 4, 751-767.
- [2] D. Cerveau & A. Lins Neto. *Irreducible components of the space of holomorphic foliations of degree two in  $\mathbb{P}^n$ ,  $n \geq 3$* . Ann. of Math., 143, (1996), 577-612 .
- [3] D. Cerveau, A. Lins Neto & , S. J. Edixhoven. *Pull-back components of the space of holomorphic foliations on  $CP(n)$ ,  $n \geq 3$* . J. Algebraic Geom. 10 (2001), no. 4, 695-711.
- [4] F. Cukierman & , J.V. Pereira. *Stability of holomorphic foliations with split tangent sheaf*. Amer. J. Math. 130 (2008), no. 2, 413-439.
- [5] F. Cukierman , J.V. Pereira & I. Vainsencher. *Stability of foliations induced by rational maps*. Ann. Fac. Sci. Toulouse Math. (6) 18 (2009), no. 4, 685-715.
- [6] W. Fulton. *Intersection Theory*. Springer-Verlag. New York. 1985.
- [7] W. Fulton, P. Pragacz. *Schubert Varieties and Degeneracy Loci*. Lecture Notes in Mathematics. Vol 1689. Springer-Verlag Berlin Heidelberg. (2006)
- [8] X. Gómez-Mont & A. Lins Neto, *Structural stability of singular holomorphic foliations having a meromorphic first integral*. Topology 30 (1991), no. 3, 315-334.

- [9] D. Grayson and M. E. Stillman. Macaulay2 version 1.9.2, a software system for research in algebraic geometry. available at <https://faculty.math.illinois.edu/Macaulay2/>.
- [10] R. Hartshorne. *Algebraic Geometry*. Graduate Texts in Mathematics. **52**. New York - Heidelberg - Berlin: Springer-Verlag. (1977).
- [11] J. P. Jouanolou. *Équations de Pfaff algébriques*. Lecture Notes in Mathematics, 708. Springer, Berlin. 1979.
- [12] F. Loray, J.V. Pereira & F. Touzet. *Foliations with trivial canonical bundle on Fano 3-folds*. Math. Nachr. 286 (2013), no. 8-9, 921-940.
- [13] I. G. Macdonald, *Symmetric functions and Hall polynomials*. Bull. Amer. Math. Soc. (N.S.) 4 (1981), no. 2, 254–265.
- [14] A. Rossini, & I. Vainsencher. *Degree of the exceptional component of foliations in  $\mathbb{P}^3$* . RACSAM (2019). <https://doi.org/10.1007/s13398-019-00627-2>. arXiv:1806.04814, (2018).

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UFF – RUA PROF. MARCOS WALDEMAR DE FREITAS REIS, S/N, BLOCO H, 5 ANDAR. 24210-201, NITERÓI, RJ - BRAZIL.

ICEX-DEPTO. MATEMÁTICA, UFMG – AV. ANTONIO CARLOS 6627. 31270-901 BELO HORIZONTE BRAZIL.