

Interference as an information-theoretic game

Sebastian Horvat*

Faculty of Physics, University of Vienna, Vienna, Austria

Borivoje Dakić†

Vienna Center for Quantum Science and Technology (VCQ),

Faculty of Physics, University of Vienna, Vienna, Austria and

Institute for Quantum Optics and Quantum Information (IQOQI), Austrian Academy of Sciences, Vienna, Austria

(Dated: April 6, 2020)

The double slit experiment provides a clear demarcation between classical and quantum theory, while multi-slit experiments demarcate quantum and higher-order interference theories. In this work we show that these experiments pertain to a broader class of processes, which can be formulated as information-processing tasks, providing a clear cut between classical, quantum and higher-order theories. The tasks involve two parties and communication between them with the goal of winning certain parity games. We show that the order of interference is in one-to-one correspondence with the parity order of these games. Furthermore, we prove the order of interference to be additive under composition of systems both in classical and quantum theory. The latter result can be used as a (semi)device-independent witness of the number of particles in the quantum setting. Finally, we extend our game formulation within the generalized probabilistic framework and prove that tomographic locality implies the additivity of the order of interference under composition. These results shed light on the operational meaning of the order of interference and can provide clues for the identification of the information-theoretic principles behind second-order interference in quantum theory.

I. INTRODUCTION

As Richard Feynman famously put it, "the double slit experiment is absolutely impossible to explain in any classical way and has in it the heart of quantum mechanics. In reality, it contains the only mystery" [1]. Indeed, the most common way of introducing quantum theory and its basic tenets is via the double-slit experiment, in which a single particle is shown to produce an interference pattern when sent through two slits, in contrast to classical theory in which this effect is missing. Much later, Sorkin [2] analyzed multi-slit experiments (i.e. generalizations of the double slit experiment to three or more slits) and noticed that quantum mechanics exhibits second-order interference only, meaning that any measurement pattern produced by a quantum system is reducible to the combination of double-slit interferences. The latter served as a motivation for introducing higher-order interference theories which are defined with respect to the order of interference produced in generalized multi-slit experiments [3]. Such theories are usually formulated within the framework of *generalized probabilistic theories* (GPT-s) [4, 5], with quantum theory being only a particular example. This further motivated the search for a set of intuitive physical principles which could explain why Nature should exhibit at most second-order interference [6–9].

In this work we study higher-order interference from the information-theoretic perspective. Firstly, we reconsider the definitions of multi-slit experiments and Sorkin's classification and notice that: (a) the framework is defined only for single particles (or single systems), (b) the interference order is defined with respect to simple operations of blocking/opening the slits, and (c) the (final) measurement refers

exclusively to intensity measurements (average number of particles at a particular location on the screen). We proceed by generalizing this setting to an arbitrary number of particles (systems), arbitrary set of (local) operations and arbitrary final measurements. The framework for higher-order interference is formulated as an information-theoretic game where we analyse generic scenarios involving classical and quantum systems. Our generalization involves many particles (systems) and their ability to win certain parity games. The order of these games and their winning probabilities are directly related to the order of interference (in Sorkin's classification) and to the number of systems involved. This generalization offers a shift in the perspective: higher order interference theories are not defined by which phenomena are allowed (e.g. by the structure of interference patterns), but by which tasks can or cannot be accomplished within the theory (i.e. their information-processing capacity). Furthermore, our study provides a direct relation between the order of interference and number of systems used in the protocol, and can thus provide a (semi) device-independent way of witnessing the number of particles (systems) present in the process.

In the final section we provide the information-theoretic formulation within the GPT formalism and we study how the order of interference behaves under the composition of systems. We construct a lower bound on the interference order which is additive under composition. In classical and quantum theory, this lower bound coincides with the upper bound, which in turn shows an interesting connection between the order of interference and the composition law. Finally, we prove that in generic local-tomographic theories [10], with the restriction to single-system operations, the order of interference is additive.

Together with an information-theoretic perspective, our findings can be seen in light of paving an alternative way towards understanding the physical principles behind the

* sebastian.horvat@univie.ac.at

† borivoje.dakic@univie.ac.at




Figure 1. Alice and Bob are located at opposite sides of a pierced plate with two slits, each of which can be either open or blocked depending on Alice's input bits. Alice sends her particle through the slits towards Bob, who, upon receiving the particle, generates an output bit b .

order of interference of quantum theory: we suspect that an important clue might be provided by the composition of systems and by the tensor-product structure. The latter would thus contribute to the operational and physical understanding of quantum theory [10–13] and provide indications for potential developments of post-quantum theories. Moreover, our work deepens the connection between interference and information processing which has already been alluded to in various contexts involving two-way communication [14–16], information speed [17], quantum acausal processes [18], superposition of orders [19] and directions [20], quantum combs [21], quantum switch [22] and quantum causal models [23].

II. INFORMATION-THEORETIC FORMULATION: PARITY GAMES

In the standard double-slit experiment a particle is sent on a plate pierced by two parallel slits. After passing through the plate, the particle can be detected on a screen. Each slit can be either open (which we denote with 0), or blocked (which we denote with 1). The figure of merit is the interference term

$$I_2 = P_{00} - P_{10} - P_{01} + P_{11}, \quad (1)$$

where $P_{x_1x_2}$ denotes the probability of detecting the particle at a point on the screen given that the slits are in states x_1 and x_2 . Notice that we explicitly included the term which corresponds to the situation in which both slits are closed, despite it being necessarily zero (i.e. $P_{11} = 0$). Classical mechanics predicts $I_2 = 0$, while quantum theory allows the particle to be in spatial superposition, thereby enabling the possibility of generating a non-vanishing I_2 .

In order to reformulate this experiment as an information-theoretic task, let us consider the scenario involving two parties, Alice and Bob, located at opposite sides of the plate, as shown in Figure 1. Alice possesses a single particle that she can send towards Bob and has control of the slits, i.e. she can decide whether to block them or not. On the other side, Bob

receives (or not) the particle, performs an arbitrary measurement and outputs a bit $b \in \{0, 1\}$. We introduce the redefined second order interference term as follows

$$\begin{aligned} \tilde{I}_2 &= \frac{1}{4} [P(0|11) - P(0|10) - P(0|01) + P(0|00)] = \\ &= \frac{1}{4} \sum_{x_1, x_2=0}^1 P(b = x_1 \oplus x_2 | x_1 x_2) - \frac{1}{2}, \end{aligned} \quad (2)$$

where $P(b|x_1x_2)$ is the probability that Bob outputs b given that the two slits are in states x_1 and x_2 . Notice that the term corresponding to both slits being closed can now be non-zero, since Bob's measurement is arbitrary, i.e. does not necessarily refer to measuring the probability of the particle inflicting on a point of the screen (intensity measurement). The probability $P(b|x_1x_2)$ is thus a generalization of $P_{x_1x_2}$ from (1) and reduces to the latter only in the case in which Bob performs the intensity measurement. The redefined interference term \tilde{I}_2 measures the probability of successfully accomplishing the following task (*parity game*):

- (a) in each run, Alice prepares the slits in state $\vec{x} \equiv \{x_1, x_2\}$ and sends her particle through the slits towards Bob;
- (b) in order to complete the task, Bob must output the parity $s_{\vec{x}} \equiv x_1 \oplus x_2$.

Bob does not have any prior information about Alice's inputs, which is why a uniform average is taken.

Notice that expression (2) is formulated in a device-independent way [24], relating only inputs and outputs, without any mention of their underlying physical realization. It is therefore natural to generalize the scenario by replacing slit operations with generic black boxes, which implement arbitrary local operations depending on their inputs. Alice can thus perform any operation (e.g. in quantum theory, general completely-positive maps): the only restriction stems from the operations being local. Moreover, instead of constraining Alice to sending only single particles (as it is also the case for standard interference experiments), we can generalize her resources to an arbitrary number of systems and analyse how the winning probability depends on the number of systems used in the process (as we will see in the next section, the game can be won even in classical theory, by using two particles). Additionally, the particles/systems sent by Alice can have any internal structure (e.g. spin) that can be potentially accessed by the black boxes.

Analogously to the generalization of the standard double-slit experiment to multi-slit experiments, we can further generalize the scenario to an arbitrary number of boxes m . In this case, Alice sends her resources, which consist of k systems, towards Bob, whose task is to output the overall parity of the boxes' inputs, as depicted in Figure 2. The figure of merit is thus

$$\tilde{I}_m(k) = \frac{1}{2^m} \sum_{x_1, \dots, x_m=0}^1 P(b = s_{\vec{x}} | x_1 \dots x_m) - \frac{1}{2}, \quad (3)$$

where $\{x_1, \dots, x_m\}$ are input bits encoded in the m boxes and $s_{\vec{x}} \equiv \bigoplus_{i=1}^m x_i$ is the overall parity. In the case in which Alice is constrained to sending only single particles, the boxes are implemented as slits and Bob's final operation consists of




Figure 2. Alice possesses a finite amount of resources that she can send towards Bob through m black boxes. The latter implement arbitrary local operations depending on Alice's inputs $\{x_1, \dots, x_m\}$. Upon receiving the resources, Bob performs an arbitrary operation and generates an output bit b .

intensity measurements, the generalized higher-order interference term $\tilde{I}_m(k)$ reduces to the standard higher-order term defined in [2] (up to normalization):

$$I_m = \sum_{x_1, \dots, x_m=0}^1 (-1)^{\sum_j x_j} P_{\vec{x}}, \quad (4)$$

where $x_j = 0(1)$ corresponds to j -th slit being open(closed) and $P_{\vec{x}}$ is the probability of detecting the particle on the screen given that the slits are in state \vec{x} .

Juxtaposed to the standard definition of m -th order interference theories involving the structure of interference patterns produced by single particles in m -slit experiments, our information-theoretic extension provides a definition in terms of the probability of winning the parity game involving m boxes by using a finite amount of resources. In addition, the latter formulation enables an analysis of the relation of the order of interference and the number of systems involved in the process (e.g., as we will see in the next section, the m -th order parity game can be won using m classical particles). We therefore introduce the characterization of higher-order theories in terms of functions $n(k)$, where n refers to the maximum order of interference that can be exhibited using k systems, i.e. $\tilde{I}_{m=n(k)}(k) \neq 0$ and $\tilde{I}_{m>n(k)}(k) = 0$.

To recapitulate, the modified multi-slit experiment consists of the following generalizations: (a) instead of one particle, Alice can possess arbitrarily many particles/systems of arbitrary internal structure, (b) the slits are replaced by black boxes which implement generic local operations depending on Alice's input bits, (c) the screen is replaced by Bob who is supposed to generate an output corresponding to the overall parity of the inputs encoded by Alice.

Finally, we can generalize Alice's input bits $\{x_1, \dots, x_m\}$ to arbitrary *dits*, i.e. elements of a set with cardinality d . In this case we introduce a class of games in which Bob is supposed to decode one of the *weighted sums modulo d* of the inputs, i.e. $s_{\vec{x}, \vec{\nu}}^{(d)} \equiv (\sum_{i=1}^m \nu_i x_i) \bmod d$, where $\vec{\nu}$ is a dit-string with components $\nu_i \in \{1, \dots, d-1\}$. Given that Alice is using k

systems, the generalized interference terms are then

$$\tilde{I}_{m, \vec{\nu}, f}^{(d)}(k) = \frac{1}{d^m} \sum_{x_1, \dots, x_m=0}^{d-1} P(b = f(s_{\vec{\nu}, \vec{x}}^{(d)} | x_1 \dots x_m)) - \frac{1}{d}, \quad (5)$$

which are defined for all *reversible* functions f that map dits into dits. Operationally, these functions take into account the potential relabelling of Bob's outputs. Notice that for $d = 2$ there was no need for specifying this, since relabelling Bob's output in Eq. (3) introduces only a minus sign. The games defined in (5) offer a definition of higher order interference theories for arbitrary d . Namely, we define an $n(k)$ -theory to satisfy the following: (i) all processes produce $\tilde{I}_{m>n(k), \vec{\nu}, f}^{(d)}(k) = 0$ for all f and all $\vec{\nu}$, and (ii) for each dit-string $\vec{\nu}$, there exists at least one process that produces $\tilde{I}_{m=n(k), \vec{\nu}, f}^{(d)}(k) \neq 0$. Notice that the (non)existence of the required process for a specific outputs' labelling fixed by f implies the (non)existence of the analogous process for any other labelling f' (since Bob can always relabel his outcomes independently of the process). Therefore, throughout this manuscript, we will only construct proofs of existence of processes which exhibit $\tilde{I}_{m=n(k), \vec{\nu}, f}^{(d)}(k) \neq 0$ for f fixed to be the identity map (i.e. no final relabelling) and we will thereby omit the index f . Moreover, whenever we omit the index $\vec{\nu}$, we refer to the unit dit-string, i.e. $\nu_i = 1, \forall i$. Therefore, the term $\tilde{I}_m^{(d)}(k)$ will correspond to the specific game in which Bob's goal is to output $s_{\vec{x}} = (\sum_i x_i) \bmod d$. As we will show later, all conclusions regarding the order of interference of classical and quantum theory hold for arbitrary $d \geq 2$. Winning the parity games can be seen as a truly "global" task, in the sense that Bob needs to produce an output that depends on all the local inputs in *each run*. For suppose that Alice manages to communicate to Bob all but one input, say the j -th one: in this case, the overall modulo $s_{\vec{x}} = (\sum_i x_i) \bmod d$ is completely unknown to Bob. The same conclusion holds if Alice omits to send more inputs or even if she does not send any information to Bob. Therefore, the interference term $\tilde{I}_{m, \vec{\nu}, f}^{(d)}(k)$ is zero irrespectively of whether Bob lacks knowledge about one or about all inputs.

In the subsequent sections, we are going to focus on classical and quantum theory and investigate their power to win the parity games. We will show that the order of interference $n(k)$ satisfies the relation

$$n(k) = kn(1), \quad (6)$$

where $n(1)$ is the order of single-system interference (for classical theory, $n(1) = 1$, while for quantum theory, $n(1) = 2$) and $n(k)$ is the interference order achievable using k systems. Moreover, we will dwell on possible generalizations of the latter relation to generic higher-order-interference theories within the GPT framework.

Before proceeding further, we will first show an important mathematical property that relates the order of interference to the algebraic order of the probability distributions $P(b|\vec{x})$.

III. ALGEBRAIC ORDER OF THE PROBABILITY DISTRIBUTIONS

In this section we are going to show that any distribution which does not exhibit higher than n -th order interference can be written as a linear combination of functions of at most n different inputs, or, in other words, that the algebraic order of $P(b|\vec{x})$ is at most n . Here we will consider the $d = 2$ case, while the general case is addressed in Appendix A.

Let us consider the parity game involving m boxes and assume that Alice uses one system of order $n < m$. By definition, this means that the interference terms (3) vanish for all parity games involving more than n boxes, which can be written in a concise form as follows:

$$\begin{aligned} \tilde{I}_{m,\vec{\sigma}}^{(2)} &= \frac{1}{2^m} \sum_{\vec{x}} (-1)^{\sum_j \sigma_j x_j} P(0|\vec{x}) = 0, \\ \forall \sigma_j \in \{0, 1\}, \quad \text{s.t.} \quad \sum_j \sigma_j &> n. \end{aligned} \quad (7)$$

For compactness, we introduced the m -component bit string $\vec{\sigma}$, that specifies which interference term the latter equation refers to: if $\sigma_j = 1$ for $j = i_1, \dots, i_l$, the equation states that the order of interference involving boxes i_1, \dots, i_l is equal to zero (l can be any integer between $n + 1$ and m).

Let us regard $P(0|\vec{x})$ as one component of a vector \vec{P} in a 2^m -dimensional vector space formed by the tensor product of m two-dimensional spaces, i.e. $P(0|\vec{x}) = \vec{e}_{x_1} \otimes \dots \otimes \vec{e}_{x_m} \cdot \vec{P}$, where $\vec{e}_{x_j=0,1}$ span the j -th two-dimensional space. Equations (7) then imply:

$$\begin{aligned} \tilde{I}_{m,\vec{\sigma}}^{(2)}(k=1) &= \frac{1}{2^m} \sum_{\vec{x}} (-1)^{\sum_j \sigma_j x_j} \vec{e}_{x_1} \otimes \dots \otimes \vec{e}_{x_m} \cdot \vec{P} = \\ &= \frac{1}{2^{m/2}} \vec{f}_{\sigma_1} \otimes \dots \otimes \vec{f}_{\sigma_m} \cdot \vec{P} = \frac{1}{2^{m/2}} \lambda_{\vec{\sigma}} = 0, \end{aligned} \quad (8)$$

where we introduced the rotated vectors

$$\vec{f}_{\sigma_j} \equiv \frac{1}{\sqrt{2}} (\vec{e}_0 + (-1)^{\sigma_j} \vec{e}_1). \quad (9)$$

The terms $\lambda_{\vec{\sigma}}$ are components of vector \vec{P} in the new basis spanned by the rotated vectors \vec{f}_{σ_j} . Equation (8) then states that the components $\lambda_{\vec{\sigma}}$ are zero if $\sum_j \sigma_j > n$. The probabilities $P(0|\vec{x})$ can thus be expressed as

$$\begin{aligned} P(0|\vec{x}) &= \left[\bigotimes_{i=1}^m \vec{e}_{x_i} \right] \cdot \left[\sum_{\vec{\sigma}} \lambda_{\vec{\sigma}} \bigotimes_{j=1}^m \vec{f}_{\sigma_j} \right] = \\ &= \frac{1}{2^{m/2}} \sum_{\substack{\sigma_1, \dots, \sigma_m \\ \sum_j \sigma_j \leq n}} (-1)^{\sum_j \sigma_j x_j} \lambda_{\vec{\sigma}}, \end{aligned} \quad (10)$$

where we used $\vec{e}_{x_i} \cdot \vec{f}_{\sigma_i} = 2^{-1/2} (-1)^{\sigma_i x_i}$. Therefore, $P(0|\vec{x})$ is a linear combination of functions of at most n different inputs:

$$P(0|\vec{x}) = \sum_{l=0}^n \sum_{i_1, \dots, i_l} c_{i_1, \dots, i_l}^{(l)} g^{(l)}(x_{i_1}, \dots, x_{i_l}), \quad (11)$$

where we introduced the functions

$$g^{(l)}(x_{i_1}, \dots, x_{i_l}) = (-1)^{\sum_{a=1}^l x_{i_a}}, \quad (12)$$

and the coefficients

$$c_{i_1, \dots, i_l}^{(l)} = \frac{1}{2^{m/2}} \vec{\lambda}_{\vec{\sigma}_{(i_1, \dots, i_l)}^{(l)}}, \quad \left(\vec{\sigma}_{(i_1, \dots, i_l)}^{(l)} \right)_s = \sum_{r=1}^l \delta_{s, i_r}. \quad (13)$$

Next, in order to tackle the problem for arbitrary d , for each set of generalized interference terms $\left\{ \tilde{I}_{m,\vec{\nu},f}^{(d)}, \forall \vec{\nu}, f \right\}$ we introduce its *dual* terms $\left\{ \tilde{J}_{m,\vec{\nu},\alpha,b}^{(d)}, \forall \vec{\nu}; \forall b \in \{0, \dots, d-1\} \right\}$ as

$$\tilde{J}_{m,\vec{\nu},b}^{(d)} = \frac{1}{d^m} \sum_{\vec{x}} (\omega_d)^{\sum_a \nu_a x_a} P(b|\vec{x}), \quad (14)$$

where $\omega_d = e^{i2\pi/d}$ is the d -th root of unity. For the $d = 2$ case, the dual terms coincide with the ones defined in terms of the game formulation (3); however, this is not the case for $d > 2$. Nevertheless, in Appendix A we show that the generalized interference term $\tilde{I}_{m,\vec{\nu},f}^{(d)}$ from (5) vanishes for all relabellings f and all dit-strings $\vec{\nu}$ if and only if $\tilde{J}_{m,\vec{\nu},b}^{(d)} = 0$, for all $b = 0, \dots, d-1$ and for all $\vec{\nu}$. This allows us to characterize the order of interference by analysing the behaviour of its dual, which can often be more tractable. Indeed, by this method we show in Appendix B that the conclusion derived in (12) holds for arbitrary d , i.e. that any distribution that exhibits at most m -th order interference can be written as a linear sum of functions which depend on at most m dits.

IV. CLASSICAL RESOURCES

Now we are going to analyse the generalized interference terms achievable within classical theory. *Single systems.* For a start, let us focus on the simplest scenario, i.e. the generalization of the double slit experiment, in which Alice sends a single particle through two boxes towards Bob (for now we stick to binary inputs, i.e. $d = 2$). Recall that the two boxes can implement arbitrary local operations (not only blocking/opening the slits) labelled by x_1 and x_2 . If the particle is classical, i.e. has a definite trajectory, Bob's output can depend on the state of at most one box and can be decomposed as follows:

$$P(b|x_1 x_2) = q_1 P_1(b|x_1) + q_2 P_2(b|x_2), \quad (15)$$

where $q_{1,2}$ are probabilities and $P_{1,2}(b|x_{1,2})$ each depend on the state of at most one box. The modified interference term therefore vanishes:

$$\begin{aligned} \tilde{I}_2^{(2)}(k=1) &= \frac{1}{4} \sum_{x_1, x_2=0}^1 P(b = x_1 \oplus x_2 | x_1 x_2) - \frac{1}{2} \\ &= \frac{1}{4} \sum_{x_1, x_2=0}^1 (-1)^{x_1 + x_2} P(0|x_1 x_2) = 0. \end{aligned} \quad (16)$$

Intuitively, the knowledge about the state of one box does not increase the probability of correctly guessing the inputs' parity with respect to a random guess. The same reasoning holds for arbitrary d , implying that $\tilde{I}_2^{(d)}(k=1) = 0$ for one classical particle.

Multiple systems. Now, what if Alice possesses two classical particles? Let us for the moment assume that the boxes are implemented by slits (as in the original double-slit experiment) and that Alice in each run sends deterministically one particle towards the first slit and the other towards the second slit. If the parity of the slits' states is $x_1 \oplus x_2 = 0$, then Bob either receives both particles on his side or he receives no particles at all; alternatively, if the parity is 1, Bob receives exactly one particle. Thus, by simply counting the number of received particles, Bob can in each run determine the inputs' parity and can perfectly accomplish the required task, thereby generating $\tilde{I}_2^{(2)}(k=2) = 1/2$. On the other hand, the standard interference definition (1) remains null also for two particles, since the average number of particles received by Bob (or inflicted on the screen) is equal to 1 regardless of the inputs' parity. This shows the difference between the standard formulation and our game formulation, as the latter allows Bob to measure coincidences, and not only the average particle number (i.e. intensity).

We proceed by analysing the fully general scenario with m boxes that implement arbitrary local transformations and assume that Alice's resources consist of k classical objects, be it particles, conglomerates of particles or localized wave packets. For this reason, it is clear that Alice's resources cannot interact with more than k boxes, which means that Bob's output can depend on at most k inputs: if $k < m$, Bob's output is equivalent to a random guess and $\tilde{I}_m^{(d)}(k) = 0$, while if $k \geq m$, he can in principle deterministically accomplish the required task and thus generate $\tilde{I}_m^{(d)}(k) = 1 - 1/d$ (if Alice sends one system per box, Bob can in principle retrieve all inputs). The same reasoning holds for all the other interference terms $\tilde{I}_{m,\vec{\nu},f}^{(d)}$.

Therefore, classical theory satisfies $n(k) = k$, meaning that k classical systems can exhibit at most k -th order interference.

V. QUANTUM RESOURCES

Contrasted to classical mechanics, quantum theory allows spatial superpositions of physical objects, which can generate a non-zero interference term I_2 , even with a single particle. On the other hand, higher order interference terms $I_{j>2}$ defined in multi-slit experiments remain null even in quantum theory (see for instance [2]). In this section we show that analogous statements hold for the generalized interference terms $\tilde{I}_{m,\vec{\nu},f}^{(d)}(k=1)$ and provide an extension to more particles (systems). Let us first look into the simplest case, i.e. one particle and two boxes.

VI. Two boxes

We start by considering the case $d = 2$. Let us suppose that Alice possesses a single quantum particle and sends it in spatial superposition towards the two boxes, which implement binary inputs. The quantum state is given by

$$|\psi\rangle_0 = \frac{1}{\sqrt{2}} (|1\rangle + |2\rangle), \quad (17)$$

where $|1\rangle$ and $|2\rangle$ are states corresponding to the two trajectories. Next, suppose that each box interacts with the particle in a simple way, e.g. by providing a local phase-shift $\phi_i = x_i\pi$. After passing through the boxes, the state is

$$|\psi\rangle_{x_1,x_2} = \frac{1}{\sqrt{2}} (e^{ix_1\pi} |1\rangle + e^{ix_2\pi} |2\rangle). \quad (18)$$

Therefore, Bob receives the particle in the following states

$$|\psi\rangle = \begin{cases} \frac{1}{\sqrt{2}} (|1\rangle + |2\rangle), & \text{if } x_1 \oplus x_2 = 0, \\ \frac{1}{\sqrt{2}} (|1\rangle - |2\rangle), & \text{if } x_1 \oplus x_2 = 1, \end{cases} \quad (19)$$

which are orthogonal and thus perfectly distinguishable, thereby enabling Bob to deterministically decode the parity $x_1 \oplus x_2$ and to produce $\tilde{I}_2^{(2)}(k=1) = 1/2$.

The previous result holds for binary inputs; now suppose that the parties are playing the modulo game (5) specified by dit string $\vec{\nu} = (\nu_1, \nu_2)$, i.e. Bob's goal is to output $(\nu_1 x_1 + \nu_2 x_2) \bmod d$ (we assume there is no final relabelling, i.e. that f is the identity; as we already argued, all other cases follow automatically). The players can employ the following strategy. For inputs $(x_1, x_2) \in \{(0,0), (0,1), (1,0)\}$, Alice uses the same encoding as in the binary case (i.e. by applying local phases $x_i\pi$); for any other combination of inputs, Alice does not send the particle. On the other side, Bob performs the same measurement as in the binary case. If he receives the particle, he can infer the parity $x_1 \oplus x_2$ with unit probability. If the parity is zero, Bob outputs 0, since $\nu_1 x_1 + \nu_2 x_2 = 0$. If the parity is one, then the desired output is either ν_1 or ν_2 and Bob can thus produce the correct outcome with 50% probability. On the other hand, if he does not receive the particle, he outputs a random *dit*. The interference term is then

$$\begin{aligned} \tilde{I}_{2,\vec{\nu}}^{(d)}(k=1) &= \frac{1}{d^2} \sum_{x_1,x_2=0}^{d-1} P(b = s_{x,\vec{\nu}}^{(d)} | x_1 x_2) - \frac{1}{d} \\ &= \frac{1}{d^2} \left[2 + (d^2 - 3) \frac{1}{d} \right] - \frac{1}{d} > 0. \end{aligned} \quad (20)$$

We have therefore shown that a single quantum particle can be used to exhibit second order interference for any d .

VII. General case

Here we analyse the fully general scenario involving arbitrary local operations and arbitrary measurements. We start by constraining Alice's resources to a single quantum system and afterwards we extend our considerations to the case of

multiple systems.

Single systems. For a start, let us fix the resources to one quantum system, without restricting its internal degrees of freedom. The resource can thus be an electron, an atom or any localized quantum system that can be prepared in coherent spatial superposition using a beam splitter or some more sophisticated mechanism. The local operations implemented by the m boxes are completely arbitrary, examples being local unitary operators and CP maps acting on the internal degrees of freedom of the system.

The most general state Alice can prepare is thus given by

$$\rho_0 = \sum_{i,j,k,l} c_{ijkl} |i\rangle \langle j| \otimes |\phi_k\rangle \langle \phi_l|, \quad (21)$$

where $\{|i\rangle, i = 1, \dots, m\}$ denote states representing defined paths directed towards the m boxes, while $\{|\phi_k\rangle\}$ span the Hilbert space of the internal degrees of freedom of the system. The matrix elements c_{ijkl} need to satisfy certain conditions in order for ρ_0 to be a legitimate quantum state, but we will not specify them since this constraint is not relevant for what follows. After the system passes through the boxes, the state is transformed to

$$\rho_{x_1, \dots, x_m} = \sum_{i,j,k,l} c_{ijkl} |i\rangle \langle j| \otimes M_{x_i, x_j} \{|\phi_k\rangle \langle \phi_l|\}, \quad (22)$$

where $M_{x_i, x_j} \{\dots\}$ are arbitrary local maps, i.e. they depend only on their respective inputs x_i and x_j . For instance, in case of local unitary operations, they are given by $M_{x_i, x_j} \{|\phi_k\rangle \langle \phi_l|\} = U_i(x_i) |\phi_k\rangle \langle \phi_l| U_j^\dagger(x_j)$.

To simplify the expression, let us introduce the set of tensors $\{C_{ij}(x_i, x_j) \equiv \sum_{kl} c_{ijkl} M_{x_i, x_j} \{|\phi_k\rangle \langle \phi_l|\}, \forall i, j\}$ reducing expression (22) to

$$\rho_{x_1, \dots, x_m} = \sum_{i,j} |i\rangle \langle j| \otimes C_{ij}(x_i, x_j). \quad (23)$$

Next, let us suppose that Alice and Bob are playing the modulo game (5) specified by a dit-string \vec{v} and with no final relabelling (all other cases follow automatically). Notice that for $m > 2$ and for any $\alpha \in \{1, \dots, d-1\}$, the following property holds

$$\begin{aligned} \sum_{\vec{x}} (\omega_d)^\alpha \sum_k \nu_k x_k \rho_{\vec{x}} &= \\ &= \sum_{i,j} |i\rangle \langle j| \otimes \sum_{\vec{x}} (\omega_d)^\alpha \sum_k \nu_k x_k C_{ij}(x_i, x_j) = 0, \end{aligned} \quad (24)$$

where $\omega_d \equiv e^{i2\pi/d}$. This is so because the tensors C_{ij} depend on at most two inputs and because $\sum_{x_k=0}^{d-1} (\omega_d)^{\alpha \nu_k x_k} = 0$. Let us define the average taken over states with equal modulo as follows

$$\rho^{(S)} \equiv \frac{1}{d^{m-1}} \sum_{(\sum_k \nu_k x_k) \bmod d = S} \rho_{\vec{x}}. \quad (25)$$

As we show in Appendix C, equations (24) then imply

$$\rho^{(S)} = \rho^{(S')} \equiv \bar{\rho}, \quad \forall S, S' = 0, \dots, d-1. \quad (26)$$

Since all average states $\rho^{(S)}$ are equal, Bob cannot distinguish them by any means. Formally, Bob performs a measurement consisting of d outcomes and represented by a generic POVM $\{\Pi_1, \dots, \Pi_d\}$. If Alice sends $k = 1$ quantum systems, the generalized interference term specified by the dit-string \vec{v} necessarily vanishes for $m > 2$:

$$\begin{aligned} \tilde{I}_{m>2, \vec{v}}^{(d)}(k=1) &= \frac{1}{d} \text{Tr} \left[\sum_S \Pi_S \rho^{(S)} \right] - 1/d \\ &= \frac{1}{d} \text{Tr}[\mathbb{1}\bar{\rho}] - 1/d = 0. \end{aligned} \quad (27)$$

A resource consisting of one quantum system is enough to produce non-vanishing second order interference (for binary inputs, it can even raise the interference term to its maximum possible value); on the other hand, for $m > 2$, there is no difference between sending one quantum system and sending no resource at all. This drastic difference can be traced down to (24), where the tensors C_{ij} couple at most two inputs x_i and x_j . This is because quantum states are described by (1, 1) tensors, i.e. density matrices. The latter constraint arises through the Born rule, which essentially sets quantum interference to the second order [3].

Multiple systems. Next, let us suppose that Alice's resources consist of more than one quantum system, say, in general, k of them. We assume that the systems are distinguishable and thus described by the tensor product of the single-system Hilbert spaces¹. The most general state prepared by Alice is then

$$\rho_0 = \sum_{\mathbf{i}, \mathbf{j}, \mathbf{r}, \mathbf{l}} c_{\mathbf{i}\mathbf{j}\mathbf{r}\mathbf{l}} \bigotimes_{p=1}^k \left[|n_{i_p}^{(p)}\rangle \langle n_{j_p}^{(p)}| \otimes |\phi_{r_p}^{(p)}\rangle \langle \phi_{l_p}^{(p)}| \right], \quad (28)$$

where \mathbf{i} is short for (i_1, \dots, i_k) and analogously for the other indices. The vectors $\{|n_{i_p}^{(p)}\rangle, \forall i_p = 1, \dots, m\}$ span the spatial Hilbert space of the p -th system, while $\{|\phi_{r_p}^{(p)}\rangle, \forall r_p\}$ span its internal degrees of freedom (the dimensionality of which is arbitrary).

After passing through the boxes, the state is transformed to

$$\rho_{\vec{x}} = \sum_{\mathbf{i}, \mathbf{j}} \bigotimes_{p=1}^k \left[|n_{i_p}^{(p)}\rangle \langle n_{j_p}^{(p)}| \right] \otimes C_{\mathbf{i}\mathbf{j}}(x_{i_1}, x_{j_1}, \dots, x_{i_k}, x_{j_k}), \quad (29)$$

where $\{C_{\mathbf{i}\mathbf{j}}\}$ are tensors depending on at most $2k$ inputs (defined in the same fashion as in the single-system case, see Eq. (23)). The crucial difference with respect to the single system case is that the expressions (24) are now valid only if $m > 2k$. Analogously to the single-system case, one comes to the following conclusion

$$\tilde{I}_{m>2k, \vec{v}}^{(d)}(k) = 0. \quad (30)$$

¹ In the case of identical particles, the allowed space of states is spanned by the (anti)symmetric subspace of the tensor product of the single-system Hilbert spaces. Therefore, besides the relations derived for distinguishable systems, further constraints might hold for processes involving identical particles.

We therefore showed that k quantum systems *cannot* produce more than $2k$ -th order interference. Now we are going to show that k systems *can* produce $2k$ -th order interference. For binary inputs (i.e. $d = 2$), Alice and Bob can partition the $2k$ boxes into pairs $\{(x_1, x_2), \dots, (x_{2k-1}, x_{2k})\}$ and for each pair use the protocol described in Section V VI; Bob can thus perfectly decode the parity of each pair, which enables him to win the game with unit probability. For $d > 2$, the proof is not so straightforward, since the protocol involving one particle and two boxes does not raise the interference term to its maximum value as it does for binary inputs. Nevertheless, in Appendix D we derive a general result which shows that k systems of single-system orders $\{n_1, \dots, n_k\}$ can produce $(\sum_{i=1}^k n_i)$ -th order interference, for any $d \geq 2$ (this statement is independent of the underlying physical theory). Therefore, k quantum systems (i.e. systems of order two) can produce $2k$ -th order interference.

According to our classification, the latter results implies that quantum theory satisfies $n(k) = 2k$, meaning that k (distinguishable) quantum systems can produce at most $2k$ -th order interference.

VI. HIGHER-ORDER INTERFERENCE THEORIES

In the preceding sections we saw that if Alice's resources consist of k classical systems, the interference terms $\tilde{I}_{m,\vec{v},f}^{(d)}(k)$ vanish for $m > k$ (for any d). On the other hand, by using k quantum systems, one can produce at most $2k$ -th order interference, essentially due to the locality of the encoding operations and Born's rule. It is tempting to conjecture that the analogous statement holds for generic higher order theories, i.e. $n(k) = kn(1)$, where $n(1)$ is the single-system order of interference.

VI.I. Single systems

Our information-theoretic formulation given in Section II can be studied within the GPT framework, as definition (5) does not contain any information about the structure of the underlying theory. However, the subtle point here is to define the notion of local operations specified by the action of the boxes. In what follows, we will provide this definition for *single systems*. The latter reduces to the standard GPT definition given in [4], if the set of transformations and measurements are restricted to projectors (i.e. opening/closing the "slits" and detecting the particle).

We adopt the standard formalism of GPTs involving states, transformations and effects [4]. The state space of Alice's system will be denoted with $\text{St}(A)$, the set of effects with $\text{Eff}(A)$ and the set of transformations with $\text{Transf}(A)$. The transformations are linear operators on the state space, mapping states into states, while the effects are linear function-

als on the state space that map states into probabilities². We first define the set of "box" effects $\{\vec{e}_1, \dots, \vec{e}_m\}$, where \vec{e}_j produces the probability of "finding the system at box j " (as in the definition provided in [4]). These effects naturally induce the following decomposition of the state space:

$$S_1, \dots, S_m \subset \text{St}(A) : \forall \vec{s} \in S_i, \vec{e}_j \cdot \vec{s} = \delta_{i,j}. \quad (31)$$

With this partition, we are ready to define the set of transformations generated by the boxes. We denote the set of transformations representing the actions of the i -th box with $\{T_{x_i}^{(i)} \in \text{Transf}(A), \forall x_i = 0, \dots, d-1\}$. In order to formalize the notion of locality of the operations, it is reasonable to assume that the action of box i leaves invariant all states localized at boxes $j \neq i$, i.e. $\forall \vec{s} \in S_{j \neq i} : T_{x_j}^{(j)} \vec{s} = \vec{s}$ (this condition is known as "branch locality" [25]). Furthermore, we assume that the ordering of the boxes' actions is irrelevant, i.e. $[T_{x_i}^{(i)}, T_{x_j}^{(j)}] = 0$, if $i \neq j$. In quantum (field) theory, if "locality" of the boxes' actions refers to real space, the latter condition is ensured via the axiom of microcausality (see also [26]).

The total set of transformations generated by the boxes is thus

$$\tau = \left\{ T_{\vec{x}} \equiv T_{x_1}^{(1)} T_{x_2}^{(2)} \dots T_{x_m}^{(m)}, \quad \forall x_1, \dots, x_m = 0, 1, \dots, d-1 \right\}. \quad (32)$$

The interference term is defined with respect to the following process: Alice prepares an arbitrary state $\vec{s} \in \text{St}(A)$ and sends the system through the boxes which implement the transformation $T_{\vec{x}} \in \tau$; Bob receives the system and performs an arbitrary measurement with d outcomes represented by the following effects $\mathbf{F} = \{\vec{f}_1, \dots, \vec{f}_d \in \text{Eff}(A), \sum_j \vec{f}_j = \vec{u}\}$, where \vec{u} is the unit effect. Higher order interference theories for single systems are then defined as follows.

Definition. We say that the theory is of n -th order if and only if it satisfies the following:

(a) $\forall d \geq 2, \forall \vec{v}, \forall f$, the following holds:

$$\tilde{I}_{m,\vec{v},f}^{(d)}(k=1) = \frac{1}{d^m} \sum_{x_1, \dots, x_m} \vec{f}_{\vec{s}^{(d)}} \cdot T_{\vec{x}} \vec{s} - \frac{1}{d} = 0, \quad \forall m > n, \quad (33)$$

for any state $\vec{s} \in \text{St}(A)$, set of transformations τ and measurement \mathbf{F} ;

(b) $\forall d \geq 2, \forall \vec{v}, \forall f$ there exist a state, a set of transformations and a measurement such that

$$\tilde{I}_{m,\vec{v},f}^{(d)}(k=1) \neq 0, \quad m = n. \quad (34)$$

VI.II. Multiple systems

In Appendix D we constructed a proof that k generic systems of orders $\{n_1, \dots, n_k\}$ can exhibit $\sum_i n_i$ -th order interference³, for any $d \geq 2$. The latter provides a lower bound

² For example, $\vec{e} \cdot (T\vec{s})$ denotes the probability of obtaining the measurement result corresponding to the effect \vec{e} , given that the measurement is performed on the transformed state $T\vec{s}$.

³ Under the condition that the composition of systems is well defined.

which is additive in the interference order. On the other hand, we saw that in classical and quantum theory, this lower bound coincides with the upper bound, i.e. these theories satisfy $n(k) = kn(1)$. In what follows, we will discuss the extendability of this relation to a broader class of theories.

Let us start by assuming the principle of Local Tomography [10], which states that a physical state is fully characterized by the marginals of its subsystems and correlations among thereof. Under this assumption, the state space of a composite system is isomorphic to the tensor product of the subsystems' state spaces (as it holds for example in quantum and classical theory [10]). We label the k subsystems with A_1, \dots, A_k ; the composite state space is then $\text{St}(A) = \text{St}(A_1) \otimes \dots \otimes \text{St}(A_k)$. The space of effects is a subset of the single-systems' spaces of effects, i.e. $\text{Eff}(A) \subseteq \text{Eff}(A_1) \otimes \dots \otimes \text{Eff}(A_k)$ ⁴. On the other hand, the space of transformations is generally larger than the tensor product of the subsystems' sets of transformations. However, here we will focus exclusively on boxes which act independently on the single systems (thereby excluding e.g. entangling gates in quantum theory). We want to show that under these assumptions, k systems of orders $\{n_1, \dots, n_k\}$ cannot exhibit more than $\sum_i n_i$ -th order interference. Let us first focus on binary inputs, i.e. $d = 2$.

An arbitrary state $\vec{s} \in \text{St}(A)$ prepared by Alice can be written as a linear combination of the tensor product of single system states: $\vec{s} = \sum_{\mathbf{i}} a_{\mathbf{i}} \vec{s}_{i_1}^{(1)} \otimes \dots \otimes \vec{s}_{i_k}^{(k)}$, where \mathbf{i} is short for i_1, \dots, i_k . The same holds for an arbitrary effect $\vec{f} \in \text{Eff}(A)$ representing Bob's final measurement, i.e. $\vec{f} = \sum_{\mathbf{i}} b_{\mathbf{i}} \vec{f}_{i_1}^{(1)} \otimes \dots \otimes \vec{f}_{i_k}^{(k)}$. As we already said, we assume that the set of transformations is generated by the single system transformations defined in Section VI VI.I, i.e. $\tau = \left\{ T_{\vec{x}}^{(1)} \otimes \dots \otimes T_{\vec{x}}^{(k)}, \forall \vec{x} \right\}$, where the locality and no-signaling conditions hold for transformations acting on each subsystem. For $d = 2$, the m -th order interference term for an arbitrary process is then:

$$\begin{aligned} \tilde{I}_m^{(2)}(k) &= \frac{1}{2^m} \sum_{\vec{x}} (-1)^{\sum_a x_a} P(0|\vec{x}) \\ &= \frac{1}{2^m} \sum_{\mathbf{i}, \mathbf{j}} a_{\mathbf{i}} b_{\mathbf{j}} \sum_{\vec{x}} (-1)^{\sum_a x_a} \left[\bigotimes_{p=1}^k \vec{f}_{i_p}^{(p)} \right] \cdot \left[\bigotimes_{p'=1}^k T_{\vec{x}}^{(p')} \vec{s}_{j_{p'}}^{(p')} \right] \\ &= \frac{1}{2^m} \sum_{\mathbf{i}, \mathbf{j}} a_{\mathbf{i}} b_{\mathbf{j}} \sum_{\vec{x}} (-1)^{\sum_a x_a} \prod_{p=1}^k P_{\vec{x}}^{(p)}, \end{aligned} \quad (35)$$

where $P_{\vec{x}}^{(p)} \equiv \vec{f}_{i_p}^{(p)} \cdot T_{\vec{x}}^{(p)} \vec{s}_{j_p}^{(p)}$ are probabilities arising from the single system processes involving p -th system (for simplicity we omitted indices \mathbf{i}, \mathbf{j}). Therefore, we see that due to Local Tomography and the restriction to single-system operations, the interference term decouples into a linear combination of products of single-system processes. As it was shown in Section III, the probability distribution pertaining to any process

involving a system of order n_p can be written as a linear combination of functions of at most n_p inputs:

$$P_{\vec{x}}^{(p)} = \sum_{i_1, \dots, i_{n_p}} c_{i_1, \dots, i_{n_p}}^{(p)} f^{(p)}(x_{i_1}, \dots, x_{i_{n_p}}). \quad (36)$$

Therefore, if $N \equiv \sum_p n_p < m$, the interference term necessarily vanishes:

$$\tilde{I}_m^{(2)}(k) = \sum_{i_1, \dots, i_N} a_{i_1, \dots, i_N} \sum_{\vec{x}} (-1)^{\sum_a x_a} g(x_{i_1}, \dots, x_{i_N}) = 0, \quad (37)$$

where we introduced for simplicity the coefficients a_{i_1, \dots, i_N} and functions $g(x_{i_1}, \dots, x_{i_N})$, the exact forms of which are of no relevance.

In order to show that the latter results holds for arbitrary d , we just need to take a look at the dual interference terms defined in Section III. Given our assumptions, one arrives to the generalization of (35):

$$\tilde{J}_{m, \vec{v}, b}^{(d)}(k) = \frac{1}{d^m} \sum_{\mathbf{i}, \mathbf{j}} a_{\mathbf{i}} b_{\mathbf{j}} \sum_{\vec{x}} (\omega_d)^{\sum_a \nu_a x_a} \prod_{p=1}^k P_{b, \vec{x}}^{(p)}. \quad (38)$$

By assumption, the probabilities $P_{b, \vec{x}}^{(p)} = \vec{f}_{b, i_p}^{(p)} \cdot T_{\vec{x}}^{(p)} \vec{s}_{i_p}^{(p)}$ pertaining to p -th system do not exhibit more than n_p -th order interference and can thus be expressed as a linear combination of functions depending on at most n_p inputs, as we showed in Appendix B. Therefore, if $N \equiv \sum_p n_p < m$, then $\tilde{J}_{m, \vec{v}, b}^{(d)}(k)$ vanishes for all \vec{v}, b . Since the dual formulation is equivalent to the game formulation (see Appendix A), this implies that the k systems cannot be used to achieve more than $\sum_j n_j$ -th order interference for any d .

To summarize, under the assumptions of (i) Local Tomography, and (ii) independence of single-system transformations, we proved that a system composed of k systems of orders (n_1, \dots, n_k) is a $(\sum_j n_j)$ -th order system. This holds trivially in classical theory, while in quantum theory we proved a stronger statement: additivity follows without assumption (ii). It might be the case that this assumption is redundant for any GPT, since the multipartite operations are still restricted to being local; however, in order to prove this, one ought to introduce a generalization of Definition VI VI.I, which we leave for future considerations.

VII. CONCLUSION AND OUTLOOK

In this work we introduced a class of information-theoretic games, which generalize standard multi-slit interference experiments. The order of interference of a theory is then seen as the (im)possibility of accomplishing certain information-processing tasks using finite resources. These games essentially characterize how much information can be decoded from a physical system given that the information was encoded in a global property (parity or modulo of the inputs) of local pieces of information (local inputs). We showed that within quantum theory, the order of interference of k systems is $2k$; it would be interesting to inspect potential connections

⁴ Under the *no-restriction hypothesis* [27], the space of effects is isomorphic to the tensor product structure; however, this assumption is not necessary in our proof.

of this result to superdense coding, which states that k qubits can be used to send at most $2k$ bits. Moreover, the game formulation can provide a (semi) device-independent witness of the particle number.

So far there have been several attempts at explaining the order of interference of quantum theory; however, a physically intuitive explanation has not yet been obtained. The reason for this might be that the sole question is leading us in the wrong direction. Instead of asking ourselves why does quantum theory behave in a particular way in multi-slit experiments (or GPT generalizations thereof), we could ask why does quantum theory restrict the amount of globally-encoded information (parity/modulo of locally encoded bits/dits) that can be retrieved from a system. Moreover, what is the relation between the (im)possibility of such a decoding and the number of systems used for encoding the information? Is there any physical argument for why interference should be additive

under composition (e.g. violation of the no-signaling principle)? Ultimately, how does one even define the notion of *number of systems* in a device-independent scenario? We believe that our work provides a framework fit for potentially finding answers to these and similar questions, which can consequently lead us to a fuller understanding of quantum and post-quantum theories.

ACKNOWLEDGMENTS

B.D. acknowledges support from an ESQ Discovery Grant of the Austrian Academy of Sciences (OAW) and the Austrian Science Fund (FWF) through BeyondC (F71). S.H. acknowledges support from an ESQ Discovery Grant of the Austrian Academy of Sciences (OAW). Both authors thank Markus P. Müller, Joshua Morris and Nicolás M. Sánchez for helpful discussions.

-
- [1] Feynman, R.P., Leighton, R.B. and Sands, M., 2005. The Feynman Lectures on Physics: Definitive Edition. Pearson Addison-Wesley.
- [2] Sorkin, R.D., 1994. Quantum mechanics as quantum measure theory. *Modern Physics Letters A*, 9(33), pp.3119-3127.
- [3] Dakic, B., Paterek, T. and Brukner, C., 2014. Density cubes and higher-order interference theories. *New Journal of Physics*, 16(2), p.023028.
- [4] Barnum, H., Mueller, M.P. and Ududec, C., 2014. Higher-order interference and single-system postulates characterizing quantum theory. *New Journal of Physics*, 16(12), p.123029.
- [5] Lee, C.M. and Selby, J.H., 2017. Higher-order interference in extensions of quantum theory. *Foundations of Physics*, 47(1), pp.89-112.
- [6] Barnum, H., Lee, C.M., Scandolo, C.M. and Selby, J.H., 2017. Ruling out higher-order interference from purity principles. *Entropy*, 19(6), p.253.
- [7] Ududec, C., Barnum, H. and Emerson, J., 2011. Three slit experiments and the structure of quantum theory. *Foundations of Physics*, 41(3), pp.396-405.
- [8] Niestegge, G., 2013. Three-slit experiments and quantum non-locality. *Foundations of Physics*, 43(6), pp.805-812.
- [9] Lee, C.M. and Selby, J.H., 2016. Generalised phase kick-back: the structure of computational algorithms from physical principles. *New Journal of Physics*, 18(3), p.033023.
- [10] Hardy, L., 2001. Quantum theory from five reasonable axioms. arXiv preprint quant-ph/0101012.
- [11] Dakic, B. and Brukner, C., 2009. Quantum theory and beyond: Is entanglement special?. arXiv preprint arXiv:0911.0695.
- [12] Masanes, L. and Mueller, M.P., 2011. A derivation of quantum theory from physical requirements. *New Journal of Physics*, 13(6), p.063001.
- [13] Chiribella, G., D Ariano, G.M. and Perinotti, P., 2011. Informational derivation of quantum theory. *Physical Review A*, 84(1), p.012311.
- [14] Del Santo, F. and Dakić, B., 2018. Two-way communication with a single quantum particle. *Physical review letters*, 120(6), p.060503.
- [15] Massa, F., Moqanaki, A., Baumeler, Ä., Del Santo, F., Kettlewell, J.A., Dakić, B. and Walther, P., 2019. Experimental two-way communication with one photon. *Advanced Quantum Technologies*, 2(11), p.1900050.
- [16] Hsu, L.Y., Lai, C.Y., Chang, Y.C., Wu, C.M. and Lee, R.K., 2020. Infinite information can be carried using a single quantum particle. arXiv preprint arXiv:2002.10374.
- [17] Horvat, S. and Dakić, B., 2019. Quantum enhancement to information speed. arXiv preprint arXiv:1911.11803.
- [18] Oreshkov, O., Costa, F. and Brukner, C., 2012. Quantum correlations with no causal order. *Nature communications*, 3, p.1092.
- [19] Feix, A., Araujo, M. and Brukner, C., 2016. Causally non-separable processes admitting a causal model. *New Journal of Physics*, 18(8), p.083040.
- [20] Guerin, P.A., Feix, A., Araujo, M. and Brukner, C., 2016. Exponential communication complexity advantage from quantum superposition of the direction of communication. *Physical review letters*, 117(10), p.100502.
- [21] Chiribella, G., D Ariano, G.M. and Perinotti, P., 2008. Quantum circuit architecture. *Physical review letters*, 101(6), p.060401.
- [22] Chiribella, G., D Ariano, G.M., Perinotti, P. and Valiron, B., 2013. Quantum computations without definite causal structure. *Physical Review A*, 88(2), p.022318.
- [23] Allen, J.M.A., Barrett, J., Horsman, D.C., Lee, C.M. and Spekkens, R.W., 2017. Quantum common causes and quantum causal models. *Physical Review X*, 7(3), p.031021.
- [24] Brunner, N., Cavalcanti, D., Pironio, S., Scarani, V. and Wehner, S., 2014. Bell nonlocality. *Reviews of Modern Physics*, 86(2), p.419.
- [25] Dahlsten, O.C., Garner, A.J. and Vedral, V., 2014. The uncertainty principle enables non-classical dynamics in an interferometer. *Nature communications*, 5(1), pp.1-7.
- [26] Garner, A.J., Mueller, M.P. and Dahlsten, O.C., 2017. The complex and quaternionic quantum bit from relativity of simultaneity on an interferometer. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 473(2208), p.20170596.
- [27] Chiribella, G., D Ariano, G.M. and Perinotti, P., 2010. Probabilistic theories with purification. *Physical Review A*, 81(6), p.062348.
- [28] Lomonosov, V. and Rosenthal, P., 2004. The simplest proof of Burnside's theorem on matrix algebras. *Linear algebra and its*

Appendix A: Equivalence between the game formulation and its dual

In this section we will show that the formulation of higher order interference theories in terms of winning probabilities of the "modulo" games

$$\tilde{I}_{m,\vec{v},f}^{(d)}(k) = \frac{1}{d^m} \sum_{x_1, \dots, x_m=0}^{d-1} P\left(b = f(s_{\vec{v},\vec{x}}^{(d)}) | x_1 \dots x_m\right) - \frac{1}{d} = 0, \quad \forall \vec{v}, \forall f \quad (\text{A1})$$

is equivalent to its dual formulation in terms of the dual interference terms

$$\tilde{J}_{m,\vec{v},b}^{(d)} = \frac{1}{d^m} \sum_{\vec{x}} (\omega_d)^{\sum_a \nu_a x_a} P(b | \vec{x}) = 0, \quad \forall \vec{v}, \quad \forall b \in \{0, \dots, d-1\}, \quad (\text{A2})$$

where $\omega_d = e^{i2\pi/d}$ is the d -th root of unity.

For convenience, let us first introduce the dual terms with an added index $\alpha \in \{1, \dots, d-1\}$:

$$\tilde{J}_{m,\vec{v},b,\alpha}^{(d)} \equiv \frac{1}{d^m} \sum_{\vec{x}} (\omega_d)^\alpha \sum_a \nu_a x_a P(b | \vec{x}). \quad (\text{A3})$$

We will show that for any dit-string \vec{v} , the following equivalence holds:

$$\left\{ \tilde{I}_{m,\vec{v},f}^{(d)} = 0, \quad \forall f \right\} \longleftrightarrow \left\{ \tilde{J}_{m,\vec{v},b,\alpha}^{(d)} = 0, \forall b \in \{0, \dots, d-1\}, \forall \alpha \in \{1, \dots, d-1\} \right\} \quad (\text{A4})$$

Since the order of interference is defined as the vanishing of the interference terms (A1) for all \vec{v} , (A4) would imply that the dual formulation of interference (A2) is equivalent to the game formulation (A1) (this is so because if the RHS of equivalence (A4) holds for all \vec{v} , then the dual conditions (A2) also hold, since the index α becomes redundant). In what follows, we will prove that (A4) does hold indeed.

In order to simplify the notation, let us introduce the $d \times d$ matrix P , whose elements are defined as

$$P_{bs} = \frac{1}{2^{m-1}} \sum_{\substack{\vec{x} \\ s_{\vec{v},\vec{x}}^{(d)}=s}} P(b | \vec{x}). \quad (\text{A5})$$

The normalization of probabilities implies that P is a stochastic matrix, i.e.

$$\sum_b P_{bs} = 1, \quad \forall s. \quad (\text{A6})$$

The game formulation equations on the left side of the equivalence in (A4) can be written as

$$\frac{1}{d^m} \sum_{x_1, \dots, x_m=0}^{d-1} P\left(b = f(s_{\vec{v},\vec{x}}^{(d)}) | x_1 \dots x_m\right) = \frac{1}{d}, \quad \forall f \quad \rightarrow \quad \sum_s P_{f(s)s} = 1, \quad \forall f, \quad (\text{A7})$$

which we rewrite succinctly as

$$\text{Tr}[\Pi_f P] = 1, \quad \forall f, \quad (\text{A8})$$

where $\Pi_f \equiv \sum_l |f(s)\rangle \langle s|$ ranges over all d -dimensional permutation matrices (the most general finite dimensional reversible transformations are indeed permutations).

On the other hand, the dual conditions on the right side of equivalence (A4) assume the following form

$$(PF)_{lk} = \delta_{k,0} \frac{1}{\sqrt{d}} \sum_j P_{lj}, \quad (\text{A9})$$

where F is the d -dimensional Fourier matrix with elements

$$(F)_{lk} = \frac{1}{\sqrt{d}}(\omega_d)^{lk}. \quad (\text{A10})$$

We will first show that the dual conditions (A9) imply the information-theoretic conditions (A8). Assuming that the dual conditions (A9) hold, we evaluate the trace

$$\text{Tr}[\Pi_f P] = \text{Tr}[F^\dagger \Pi_f P F] = \sum_{kl} (F^\dagger \Pi_f)_{kl} (P F)_{lk} = \sum_l (F^\dagger \Pi_f)_{0l} (P F)_{l0} = 1, \forall f. \quad (\text{A11})$$

The first step follows from the unitarity of F and the cyclicity of the trace, the third step is due to the dual conditions (A9), and the last step follows from $(F^\dagger \Pi_f)_{0l} = 1/\sqrt{d}$. Therefore, conditions (A9) imply (A8).

In order to prove the converse statement, let us introduce the following new basis $\beta = \{|E\rangle, |e_1\rangle, \dots, |e_{d-1}\rangle\}$, where $|E\rangle$ has the following form

$$|E\rangle = 1/\sqrt{d}(1, 1, \dots, 1)^T, \quad (\text{A12})$$

and the remaining vectors span the orthogonal subspace (they can for instance be chosen as the rows/columns of the Fourier matrix). The normalization conditions (A6) can then be written as

$$\langle E|P = \langle E|, \quad (\text{A13})$$

which implies that the matrix P in the new basis has the following form

$$P = |E\rangle \langle E| + \sum_j q_j |e_j\rangle \langle E| + \bar{E}, \quad (\text{A14})$$

where q_j are arbitrary coefficients and \bar{E} is a matrix that has support only on the subspace orthogonal to $|E\rangle$. In the new basis β , all permutation matrices have the following form

$$\Pi_f = |E\rangle \langle E| + \Delta_f, \quad (\text{A15})$$

where Δ_f is again a matrix with no support on $|E\rangle$. The latter follows from the fact that the representation of the permutation group that we are using is reducible to the direct sum of a one-dimensional representation (spanned by the vector $|E\rangle$ which is invariant under all permutations) and a $(d-1)$ -dimensional irreducible representation (here given by Δ_f). Assuming that the game-conditions (A8) hold, we obtain the following

$$1 = \text{Tr}[\Pi_f P] = 1 + \text{Tr}[\Delta_f \bar{E}] \rightarrow \text{Tr}[\Delta_f \bar{E}] = 0, \quad \forall f. \quad (\text{A16})$$

Since the matrices Δ_f provide an irreducible representation of the permutation group, the Burnside Theorem [28] implies that they span the set of all $(d-1)$ -dimensional matrices; therefore, Equation (A16) implies that \bar{E} is necessarily zero. The matrix P is thus

$$P = |E\rangle \langle E| + \sum_j q_j |e_j\rangle \langle E| = |q\rangle \langle E|, \quad (\text{A17})$$

where we introduced the vector $|q\rangle \equiv |E\rangle + \sum_j q_j |e_j\rangle$. Therefore, the sought matrix $P F$ is

$$(P F)_{kl} = \langle l|q\rangle \langle E|F|k\rangle = \delta_{k,0} \frac{1}{\sqrt{d}} \sum_j P_{lj}, \quad (\text{A18})$$

where we used $\sum_{k=0}^{d-1} (\omega_d)^k = 0$ and $\{|k\rangle; k = 0, \dots, d-1\}$ are the original basis vectors. Hence, equivalence (A4) holds, and thus the game formulation (A1) is equivalent to the dual formulation (A2). Except for being an interesting mathematical and conceptual result, the exact equivalence between these two formulations will be useful for showing that any distribution that exhibits at most m -th order interference can be written as a linear combination of functions with at most m inputs. This will consequently be the necessary ingredient for proving that Local Tomography implies the additivity of the order of interference under the composition of systems.

Appendix B: Algebraic order for arbitrary d

In this Appendix we are going to extend the mathematical property of Section III to arbitrary d .

Let us consider the modulo games involving m boxes and assume that Alice uses a system of order $n < m$. In Appendix A we showed that this is equivalent to the following constraints

$$\begin{aligned} \tilde{J}_{m,\vec{\nu},b}^{(d)} &= \frac{1}{d^m} \sum_{\vec{x}} (\omega_d)^{\sum_j \nu_j x_j} P(b|\vec{x}) = 0, \\ \forall b \in \{0, 1, \dots, d-1\}, \nu_j \in \{0, 1, \dots, d-1\}, \quad \text{s.t.} \quad \sum_j \delta_{\nu_j,0} < (m-n). \end{aligned} \quad (\text{B1})$$

For compactness, we allow the dit string $\vec{\nu}$ to range over all dit-strings with less than $(m-n)$ null components. This notation specifies which interference terms the equation refers to: if $\nu_j \neq 0$ for $j = i_1, \dots, i_l$, the equation states that the order of interference involving boxes i_1, \dots, i_l is equal to zero (l can be any integer between $n+1$ and m).

Let us regard $P(b|\vec{x})$ as one component of a vector \vec{P}_b in a d^m -dimensional vector space formed by the tensor product of m d -dimensional spaces, i.e. $P(b|\vec{x}) = \bigotimes_{i=1}^m \vec{e}_{x_i} \cdot \vec{P}_b$, where $\{\vec{e}_{x_j}, x_j = 0, \dots, d-1\}$ span the j -th d -dimensional space. Equations (B1) then imply:

$$\tilde{J}_{m,\vec{\nu},b}^{(d)} = \frac{1}{d^m} \sum_{\vec{x}} (\omega_d)^{\sum_j \nu_j x_j} \bigotimes_{i=1}^m \vec{e}_{x_i} \cdot \vec{P}_b = \frac{1}{d^{m/2}} \bigotimes_{i=1}^m \vec{f}_{\nu_i} \cdot \vec{P}_b = \frac{1}{d^{m/2}} \lambda_{\vec{\nu},b} = 0, \quad (\text{B2})$$

where we introduced the rotated vectors (rows/columns of the d -dimensional Fourier matrix)

$$\vec{f}_{\nu_j} \equiv \frac{1}{\sqrt{d}} \sum_{x_j=0}^{d-1} (\omega_d)^{\nu_j x_j}, \quad (\text{B3})$$

and components $\lambda_{\vec{\nu},b}$ in the corresponding basis. Equation (B2) states that the components $\lambda_{\vec{\nu},b}$ vanish for all $\vec{\nu}$ with more than n non-zero components. The probabilities $P(b|\vec{x})$ can thus be expressed as

$$P(b|\vec{x}) = \left[\bigotimes_{i=1}^m \vec{e}_{x_i} \right] \cdot \left[\sum_{\vec{\nu}} \lambda_{\vec{\nu},b} \bigotimes_{j=1}^m \vec{f}_{\nu_j} \right] = \frac{1}{d^{m/2}} \sum_{\substack{\nu_1, \dots, \nu_m \\ \sum_j \delta_{\nu_j,0} \geq (m-n)}} (\omega_d)^{\sum_j \nu_j x_j} \lambda_{\vec{\nu},b}, \quad (\text{B4})$$

where we used $\vec{e}_{x_i} \cdot \vec{f}_{\nu_i} = \frac{1}{\sqrt{d}} (\omega_d)^{\nu_i x_i}$. Since the sum runs only over strings ν with at most n non-zero components, $P(b|\vec{x})$ can be written as a function of at most n inputs, for every b .

Therefore we have proved that the n -th order interference dual conditions (B1), or equivalently, the information-theoretic formulation (5), provide a set of necessary and sufficient conditions for a distribution to be decomposable into a linear combination of at most n inputs. Hence, the algebraic order of the distribution $P(b|\vec{x})$ is at most n .

Appendix C: Proof of Equation (26)

In this Appendix we will show that equations

$$\sum_{\vec{x}} (\omega_d)^\alpha \sum_k \nu_k x_k \rho_{\vec{x}} = 0, \quad \forall \alpha \in \{1, \dots, d-1\} \quad (\text{C1})$$

imply the following relations

$$\rho^{(S)} = \rho^{(S')}, \quad \forall S, S' = 0, \dots, d-1, \quad (\text{C2})$$

where $\rho^{(S)}$ is defined as

$$\rho^{(S)} \equiv \frac{1}{d^{m-1}} \sum_{\substack{x_1, \dots, x_m \\ (\sum_k \nu_k x_k) \bmod d = S}} \rho_{\vec{x}}. \quad (\text{C3})$$

For a start, Equations (C1) can be written as

$$\sum_{S=0}^{d-1} (\omega_d)^{\alpha S} \rho^{(S)} = 0, \quad \forall \alpha \in \{1, \dots, d-1\}. \quad (\text{C4})$$

Next, if we introduce the vector $\vec{\rho} = (\rho^{(0)}, \rho^{(1)}, \dots, \rho^{(d-1)})^T$, the latter equations can be cast in the following form

$$F\vec{\rho} = \sqrt{d}\bar{\rho}(1, 0, \dots, 0)^T, \quad (\text{C5})$$

where $F_{kl} = 1/\sqrt{d}(\omega_d)^{kl}$ is the d -dimensional Fourier matrix and $\bar{\rho} = 1/d \sum_S \rho^{(S)}$. Taking the inverse of Equation C5 (i.e. applying F^\dagger from the left) we obtain

$$\vec{\rho} = \bar{\rho}(1, 1, \dots, 1)^T, \quad (\text{C6})$$

which is equivalent to

$$\rho^{(S)} = \rho^{(S')} = \bar{\rho}, \quad \forall S, S' = 0, \dots, d-1. \quad (\text{C7})$$

Appendix D: A lower bound on interference of generic composite systems

In this Appendix we are going to show that two systems, labelled by (A, B) and respectively of single-system orders (n_A, n_B) , can be used to achieve $(n_A + n_B)$ -th order interference.

By definition, there exist two processes involving single-systems A and B that can win the modulo games with n_A and n_B boxes with some probabilities $q_A > 1/d$ and $q_B > 1/d$. For concreteness, suppose that Alice and Bob play the modulo game for the unit dit-string $\nu_i = 1, \forall i$, with $(n_A + n_B)$ boxes, i.e. Bob is supposed to output $s_{\vec{x}}^{(d)} = (\sum_j x_j) \bmod d$. The proof for any other $\vec{\nu}$ is completely analogous. The protocol proceeds as follows. Alice sends system A to the first n_A boxes (i.e. containing inputs $\{x_1, \dots, x_{n_A}\}$) and system B through the other n_B boxes (which encode inputs $\{x_{n_A+1}, \dots, x_{n_A+n_B}\}$). Upon receiving the systems, Bob performs two separate measurements with outcomes $b^{(A)}$ and $b^{(B)}$ and produces a final output $b = (b^{(A)} + b^{(B)}) \bmod d$. Here the two systems are treated completely independently: the only ‘‘mixing’’ between them happens in the final step, since Bob’s output depends on both systems’ processes. The $(n_A + n_B)$ -th order interference term is then

$$\tilde{I}_{n_A+n_B}^{(d)} = \frac{1}{d^{n_A+n_B}} \sum_{x_1, \dots, x_{n_A+n_B}} P(b = s_{\vec{x}}^{(d)} | \vec{x}) - \frac{1}{d} = \frac{1}{d^{n_A+n_B}} \sum_{x_1, \dots, x_{n_A+n_B}} \sum_l P(b_l^{(A)}, b_l^{(B)} | \vec{x}) - \frac{1}{d}, \quad (\text{D1})$$

where the sum runs over all two-outcome measurements which produce the correct modulo, i.e. $(b_l^{(A)} + b_l^{(B)}) \bmod d = (\sum_{j=1} x_j) \bmod d$. Intuitively, Bob can produce the correct overall modulo, even if he measures wrong single-system moduli: e.g. for $d = 2$, if Bob’s single-system outcomes are $s_A \oplus 1$ and $s_B \oplus 1$, where s_A and s_B are the correct parities, he will still produce the correct overall parity, since $(s_A \oplus 1) \oplus (s_B \oplus 1) = s_A \oplus s_B$. Indeed, for arbitrary d , there are in total $(d-1)$ wrong single-system outcomes which provide the correct overall output. The interference term is thus

$$\tilde{I}_{n_A+n_B}^{(d)} = q_A q_B + \sum_{i=1}^{d-1} p_i^{(A)} p_{d-1}^{(B)} - \frac{1}{d}, \quad (\text{D2})$$

where q_A and q_B are average winning probabilities for the two systems, while the distribution averages over wrong outcomes are defined as

$$p_i^{(A)} \equiv \frac{1}{d^{n_A}} \sum_{x_1, \dots, x_{n_A}} P^{(A)}(b^{(A)} = (s_{\vec{x}}^{(d)} + i) \bmod d | \vec{x}), \quad (\text{D3})$$

and analogously for system B. Next, we define weights α_j and β_j in the following way

$$\begin{aligned} p_i^{(A)} &= (1 - q_A)\alpha_i, \\ p_{d-i}^{(B)} &= (1 - q_B)\beta_i, \end{aligned} \quad (\text{D4})$$

which implies $\sum_j \alpha_j = \sum_j \beta_j = 1$. Plugging these expressions into the interference term we obtain

$$\tilde{I}_{n_A+n_B}^{(d)} = q_A q_B + (1 - q_A)(1 - q_B)\lambda - \frac{1}{d}, \quad \lambda \equiv \sum_j \alpha_j \beta_j. \quad (\text{D5})$$

By a straightforward application of the method of Lagrange multipliers with constraints $\sum_j \alpha_j = 1$ and $\sum_j \beta_j = 1$, one obtains that the minimum value of λ is $\frac{1}{d-1}$ (which is achieved for uniform α_j, β_j). Therefore the following inequality holds

$$\tilde{I}_{n_A+n_B}^{(d)} \geq q_A q_B + \frac{1}{d-1}(1-q_A)(1-q_B) - \frac{1}{d}. \quad (\text{D6})$$

The RHS of the latter inequality is equal to zero for $q_A = q_B = 1/d$ and is monotonically increasing for $q_A > 1/d, q_B > 1/d$. Thus we constructed a process such that, if the two systems exhibit n_A -th and n_B -th order interference, the composite system exhibits $(n_A + n_B)$ -th order interference.

Using the same reasoning, the discussion can be generalized to an arbitrary number of systems, since systems can be composed two-by-two. Therefore, by the same construction, k systems of orders $\{n_1, \dots, n_k\}$ can produce $(\sum_{i=1}^k n_i)$ -th order interference.
