

# HYPERSURFACES WITH LIGHT-LIKE POINTS IN A LORENTZIAN MANIFOLD II

M. UMEHARA AND K. YAMADA

**ABSTRACT.** In the authors' previous work, it was shown that if a zero mean curvature  $C^4$ -differentiable hypersurface in an arbitrarily given Lorentzian manifold admits a degenerate light-like point, then the hypersurface contains a light-like geodesic segment passing through the point. The purpose of this paper is to point out that the same conclusion holds with just  $C^3$ -differentiability of the hypersurfaces.

## INTRODUCTION

This paper is a remark on authors' previous work [3]. We denote by  $M$  a  $C^\infty$ -differentiable Lorentzian  $(n+1)$ -manifold ( $n \geq 2$ ). Let  $U$  be a domain of  $(\mathbf{R}^n; u_1, \dots, u_n)$  and  $o \in U$  an arbitrary fixed point. We let  $F : U \rightarrow M$  be a  $C^r$ -immersion ( $r \geq 3$ ). A point  $q \in U$  which is neither time-like nor space-like is called a *light-like point*, which is a zero of the function  $B_F$  defined in [3, (1.2)]. A light-like point  $q$  is called *degenerate* if the exterior derivative of  $B_F$  vanishes at  $q$ . In [3], a class  $\mathcal{X}_0^r(M, \delta)$  of germs of  $C^r$ -immersion at  $\delta := F(o)$  is defined. Zero mean curvature immersions in  $M$  locally belong to this class. It was shown that:

**Fact A** ([3, Theorem A]). *Suppose that  $F \in \mathcal{X}_0^r(M, \delta)$  is a germ of a  $C^r$ -immersion ( $r \geq 4$ ) at a degenerate light-like point  $o$ . Then the image of  $F$  contains a light-like geodesic segment in  $M$  passing through  $\delta (= F(o))$  consisting of only degenerate light-like points.*

When  $M = \mathbf{R}_1^3$  and  $F$  is a zero mean curvature immersion, Fact A was shown by Klyachin [2] under the assumption that  $F$  is  $C^3$ -differentiable. So it is expected that the fact also holds for any  $C^3$ -differentiable  $F$ . The above fact was applied to prove a Bernstein-type theorem in [1] for entire zero mean curvature graphs without time-like points. So by the following theorem, the main theorem of [1] holds under  $C^3$ -differentiability of the graphs:

**Theorem B.** *In Fact A, the same conclusion holds even if  $F$  is a germ of a  $C^3$ -immersion. In particular, if  $F$  is a  $C^3$ -differentiable zero mean curvature hypersurface in  $M$  containing a degenerate light-like point  $o$ , then  $F$  contains a light-like geodesic segment passing through  $F(o)$  consisting of only degenerate light-like points.*

In [3], we proved Fact A by showing the local Lipschitz property of a certain system of ordinary differential equations (cf. [3, (4.8)]). However, by a careful tracing of the proof of [3, Theorem A], we obtain the same conclusion under merely  $C^3$ -differentiability of  $F$ , as in the discussions in the next section. It should be

---

*Date:* March 27, 2020.

*2010 Mathematics Subject Classification.* Primary 53A10; Secondary 53B30, 35M10.

*Key words and phrases.* maximal hypersurface, light-like geodesic, zero mean curvature, Lorentzian manifolds.

The second author was partially supported by the Grant-in-Aid for Scientific Research (B) No. 17H02839 from Japan Society for the Promotion of Science.

remarked that the authors do not know whether the theorem holds under  $C^2$ -differentiability of  $F$  or not, which should remain an open problem.

### 1. PROOF OF THE THEOREM

Since  $o \in U$  is a light-like point, there exists a non-zero tangent vector  $\mathbf{v}$  of  $U$  at  $o$  such that  $dF(\mathbf{v})$  is a light-like vector. We let  $\sigma$  be a light-like geodesic passing through  $F(o)$  such that  $dF(\mathbf{v})$  points in the tangential direction of  $\sigma$  at  $F(o)$ . We then take a Fermi-coordinate system  $(x_0, \dots, x_n)$  of signature  $(- + \dots +)$  centered at  $F(o) \in M$  along  $\sigma$  (cf. [3, Appendix A]). When  $M$  is the Lorentz-Minkowski space  $\mathbf{R}_1^{n+1}$ , the Fermi coordinate system  $(x_0, \dots, x_n)$  can be taken as the canonical coordinate system.

Using this coordinate system,  $\sigma$  can be expressed as

$$\sigma(t) = (t, 0, \dots, 0, t).$$

Moreover, all of the Christoffel symbols of the Lorentzian metric  $g$  of  $M$  vanish along  $\sigma$ . We can express  $F$  as (cf. [3, (4.1)])

$$F(u_1, \dots, u_n) = \left( f(x_1(u_1, \dots, u_n), \dots, x_n(u_1, \dots, u_n)), \right. \\ \left. x_1(u_1, \dots, u_n), \dots, x_n(u_1, \dots, u_n) \right),$$

where  $f(x_1, \dots, x_n)$  is a  $C^3$ -function of variables  $x_1, \dots, x_n$  defined on a neighborhood of the origin  $o \in \mathbf{R}^n$  ( $o$  corresponds to the degenerate light-like point of  $F$ ). In this expression of  $F$ ,  $(x_1, \dots, x_n)$  can be considered as a local coordinate system of  $U$  at  $o$  without loss of generality. We set  $y := x_n$  and use the notation  $' = d/dy$ , and set

$$(1.1) \quad a(y) := f(0, \dots, 0, y), \quad \text{and} \quad b_i(y) := f_{x_i}(0, \dots, 0, y) \quad (i = 1, \dots, n-1),$$

where  $f_{x_i} := \partial f / \partial x_i$ . Then  $a$  (resp.  $b_i$  for each  $i = 1, \dots, n-1$ ) is a  $C^3$  (resp.  $C^2$ )-differentiable function of single variable  $y$ . Using these, we define a function  $c$  by (cf. [3, (4.2)]) by

$$(1.2) \quad f(x_1, \dots, x_{n-1}, y) = a(y) + \left( \sum_{i=1}^{n-1} b_i(y) x_i \right) + c(x_1, \dots, x_{n-1}, y).$$

Then  $c$  is a function of class  $C^2$ , and the third derivatives

$$c_{yx_i x_j} = c_{x_i y x_j} = f_{yx_i x_j}, \quad c_{yy x_i} = f_{yy x_i} - b'_i$$

exist for  $i, j = 1, \dots, n-1$ , which are continuous functions. By (1.1) and (1.2), one can easily show that

$$(1.3) \quad c(0, \dots, 0, y) = c_{x_i}(0, \dots, 0, y) = c_y(0, \dots, 0, y) \\ = c_{yx_i}(0, \dots, 0, y) = c_{yy x_i}(0, \dots, 0, y) = 0$$

and that

$$(1.4) \quad f_{x_i x_j x_k}(0, \dots, 0, y) = c_{x_i x_j x_k}(0, \dots, 0, y), \\ f_{yx_i x_j}(0, \dots, 0, y) = c_{yx_i x_j}(0, \dots, 0, y) = c_{x_i y x_j}(0, \dots, 0, y), \\ f_{yy x_i}(0, \dots, 0, y) = b''_i(y)$$

for  $i, j, k = 1, \dots, n-1$ .

*Remark 1.* In [3] and [1], we used the function  $h_{ij}$  satisfying

$$(1.5) \quad f(x_1, \dots, x_{n-1}, y) = a(y) + \sum_{i=1}^{n-1} b_i(y) x_i + \sum_{i,j=1}^{n-1} \frac{h_{ij}(x_1, \dots, x_{n-1}, y)}{2} x_i x_j.$$

Moreover, we wrote that  $h_{ij}$  are  $C^1$ -functions. However it may not be true, in general. We should avoid this expression. In fact, for example, even if  $h_{ii}$  is a  $C^1$ -function, the following computation is not allowed

$$\left(h_{ii}x_i^2\right)_{x_ix_i} = (h_{ii})_{x_ix_i}x_i^2 + 2(h_{ii})_{x_i}x_i + 2h_{ii}.$$

So it should be replaced (1.5) by (1.2) in [3] and the final remark of [1].

The condition that  $o$  is a degenerate light-like point is written as (cf. [3, (4.3) and (4.5)])

$$(1.6) \quad a(0) = 0, \quad a'(0) = 1, \quad b_i(0) = b'_i(0) = 0 \quad (i = 1, \dots, n-1),$$

where  $' = d/dy$ .

Since  $F \in \mathcal{X}_0^3(M, \delta)$ , there exists a  $C^1$ -function  $\varphi$  such that (cf. [3, (3.1) and (3.3)])

$$(1.7) \quad \tilde{A} := A - \varphi B$$

vanishes identically on  $U$ , where

$$B := B_F, \quad A := A_F$$

are functions associated with  $F$  defined in [3, (2.4) and (2.6)]. We remark that  $B, \nabla B$  and  $A$  can be expressed without use of derivatives of the Christoffel symbols, where  $\nabla B := (B_{x_1}, \dots, B_{x_n})$ . Since  $(x_0, \dots, x_n)$  is a Fermi coordinate system, the expressions of  $B, \nabla B$  and  $A$  along  $\sigma$  coincide with those of  $\mathbf{R}_1^{n+1}$  given in [3, Appendix B]. For the Lorentzian metric  $g$  of  $M$ , set

$$g_{\alpha\beta} := g(\partial_{x_\alpha}, \partial_{x_\beta}), \quad \hat{g}_{\alpha\beta} := g_{\alpha\beta} \circ F \quad (\alpha, \beta = 0, \dots, n),$$

where  $\partial_{x_\alpha} := \partial/\partial x_\alpha$ . Since we can write

$$(1.8) \quad F_{x_i} = f_{x_i}\partial_0 + \partial_{x_i} \quad (i = 1, \dots, n),$$

the coefficients  $s_{i,j}$  of the induced metric  $ds^2$  with respect to the local coordinate system  $(U; x_1, \dots, x_n)$  given in [3, (2.3)] can be computed as

$$s_{i,j} = f_{x_i}f_{x_j}\hat{g}_{00} + f_{x_i}\hat{g}_{0j} + f_{x_j}\hat{g}_{i0} + \hat{g}_{ij} \quad (i, j = 1, \dots, n).$$

Since  $\{\hat{g}_{\alpha\beta}\}_{\alpha,\beta=0}^n$  are functions of  $f$ , each  $s_{i,j}$  is also a function of  $f$  and  $f_I$ , where

$$f_I := \{f_{x_i}\}_{i=1}^n.$$

In particular,  $B$  is a function of  $f$  and  $f_I$ . Also each component  $\tilde{s}^{i,j}$  ( $i, j = 1, \dots, n$ ) of the cofactor matrix of the  $n \times n$ -matrix  $(s_{i,j})$  is also a function of  $f$  and  $f_I$ .

By (1.8),  $\mu_i^\beta$  ( $i = 1, \dots, n, \beta = 0, \dots, n$ ) in [3, Page 3410] are given by

$$\mu_i^\beta = \begin{cases} 1 & \text{if } \beta = i, \\ f_{x_i} & \text{if } \beta = 0, \\ 0 & \text{if } \beta \neq 0, i, \end{cases}$$

and

$$\mu_{i,\beta} = \sum_{\alpha=0}^n \mu_i^\alpha \hat{g}_{\alpha\beta} \quad (i = 1, \dots, n, \beta = 0, \dots, n)$$

are functions of  $f$  and  $f_I$ . In particular, each component  $\tilde{\nu}_i$  of the normal vector field  $\tilde{\nu}$  (cf. [3, (2.5)]) is also a function of  $f$  and  $f_I$ . We denote by  $\{\Gamma_{\alpha\beta}^\gamma\}_{\alpha,\beta,\gamma=0}^n$  the Christoffel symbols of  $g$  with respect to the coordinate system  $(x_0, \dots, x_n)$  of  $M$ , and set

$$\hat{\Gamma}_{\alpha\beta}^\gamma := \Gamma_{\alpha\beta}^\gamma \circ F \quad (\alpha, \beta, \gamma = 0, \dots, n),$$

each of which can be considered as a function of  $f$ . Since  $F_{x_j} = f_{x_j} \partial_{x_0} + \partial_{x_j}$ , we can write

$$D_{\partial_{x_i}} F_{x_j} = f_{x_i x_j} \partial_{x_0} + \sum_{\alpha=0}^n \left( f_{x_i} f_{x_j} \hat{\Gamma}_{00}^\alpha + f_{x_i} \hat{\Gamma}_{0j}^\alpha + f_{x_j} \hat{\Gamma}_{i0}^\alpha + \hat{\Gamma}_{ij}^\alpha \right) \partial_{x_\alpha},$$

where  $D$  is the Levi-Civita connection induced by  $g$ . In particular, by [3, (2.6)],  $A$  is also a function of  $f$ ,  $f_I$ ,  $f_{IJ}$  and  $f_{In}$ , where

$$f_{IJ} := \{f_{x_i x_j}\}_{i,j=1}^{n-1}, \quad f_{In} := \{f_{x_i x_n}\}_{i=1}^{n-1}.$$

As a consequence,  $\tilde{A}$  (cf. (1.7)) has the following expression:

$$(1.9) \quad \tilde{A} = \mathcal{R}(f, f_I) f_{x_n x_n} + \mathcal{S}(f_I, f_{IJ}, f_{In}, \varphi),$$

where  $\mathcal{R}$  is a function of  $f$ ,  $f_I$ , and  $\mathcal{S}$  is a function of  $f$ ,  $f_I$ ,  $f_{IJ}$ ,  $f_{In}$  and  $\varphi$ . For example, if  $M = \mathbf{R}_1^{n+1}$  and  $\varphi = 0$ , then  $\tilde{A} = A$  and

$$(1.10) \quad \mathcal{R} = 1 - \sum_{j=1}^{n-1} f_{x_j}^2,$$

$$(1.11) \quad \mathcal{S} = \sum_{k=1}^{n-1} \left( 1 - f_{x_n}^2 - \sum_{j=1, j \neq k}^{n-1} f_{x_j}^2 \right) f_{x_k x_k} \\ + 2 \sum_{1 \leq j < k \leq n-1} f_{x_j} f_{x_k} f_{x_j x_k} + 2 \sum_{j=1}^{n-1} f_{x_j} f_{x_n} f_{x_j x_n}.$$

Although  $f_I$  (resp.  $f_{IJ}$  and  $f_{In}$ ) has (resp. have) only  $C^2$ -differentiability (resp.  $C^1$ -differentiability),  $\mathcal{R}$  and  $\mathcal{S}$  depend smoothly on variables  $f, f_I, f_{IJ}, f_{In}$  and  $\varphi$ , because  $M$  itself is a  $C^\infty$ -manifold. Differentiating (1.9), we have the following expression of the derivative  $\tilde{A}_{x_i}$  with respect to  $\partial_{x_i}$ :

$$\tilde{A}_{x_i} = \mathcal{R}(f, f_I) f_{x_n x_n x_i} + \mathcal{T}(f_I, f_{IJ}, f_{In}, f_{IJK}, f_{IJn}, \varphi, \varphi_I),$$

where  $\mathcal{T}$  is a function of  $f_I, f_{IJ}, f_{In}, f_{IJK}, f_{IJn}, \varphi, \varphi_I$  and

$$f_{IJK} := \{f_{x_i x_j x_k}\}_{i,j=1}^{n-1}, \quad f_{IJn} := \{f_{x_n x_i x_j}\}_{i,j=1}^{n-1}$$

For example, if  $n = 2$ ,  $\mathcal{R}$  is a function of  $f, f_{x_1}$  and  $\mathcal{S}$  is a function of  $\varphi, f, f_{x_1}, f_{x_1 x_1}, f_{x_1 x_2}$ . So we have the following:

$$A_{x_1} = \mathcal{R}_f(f, f_{x_1}) f_{x_1} f_{x_2 x_2} + \mathcal{R}_{f_{x_1}}(f, f_{x_1}) f_{x_1 x_1} f_{x_2 x_2} + \mathcal{S}_\varphi(f, f_{x_1}, f_{x_1 x_1}, f_{x_1 x_2}, \varphi) \varphi_{x_1} \\ + \mathcal{S}_f(f, f_{x_1}, f_{x_1 x_1}, f_{x_1 x_2}, \varphi) f_{x_1} + \mathcal{S}_{f_{x_1}}(f, f_{x_1}, f_{x_1 x_1}, f_{x_1 x_2}, \varphi) f_{x_1 x_1} \\ + \mathcal{S}_{f_{x_1 x_1}}(f, f_{x_1}, f_{x_1 x_1}, f_{x_1 x_2}, \varphi) f_{x_1 x_1 x_1} + \mathcal{S}_{f_{x_1 x_2}}(f, f_{x_1}, f_{x_1 x_1}, f_{x_1 x_2}, \varphi) f_{x_1 x_2 x_1},$$

where, for example,

$$\mathcal{R}_f(f, f_{x_1}) := \frac{\partial \mathcal{R}(f, f_{x_1})}{\partial f}, \quad \mathcal{R}_{f_{x_1}}(f, f_{x_1}) := \frac{\partial \mathcal{R}(f, f_{x_1})}{\partial f_{x_1}}.$$

Since  $\tilde{A}$  vanishes identically, we have  $\tilde{A} = 0$  and  $\tilde{A}_{x_i} = 0$  for  $i = 1, \dots, n-1$ . We set  $\mathbf{x} := (x_1, \dots, x_{n-1})$  and substitute  $\mathbf{x} = (0, \dots, 0)$  to them. Then they are functions of one variable  $y$ , and we have

$$(1.12) \quad (\mathbf{a} :=) \tilde{A}|_{\mathbf{x}=(0, \dots, 0)} = 0,$$

$$(1.13) \quad (\mathbf{a}_i :=) \tilde{A}_{x_i}|_{\mathbf{x}=(0, \dots, 0)} = 0 \quad (i = 1, \dots, n-1).$$

Although  $\mathbf{a}$  and  $\mathbf{a}_i$  ( $i = 1, \dots, n-1$ ) are all functions of  $y$ , we now would like to think that they are functions of  $a, a', a'', b_I, b'_I, b''_I, c_{IJ}, c_{IJK}$  and  $\varphi$ , where

$$\begin{aligned} c_{ij}(y) &:= c_{x_i x_j}(0, \dots, 0, y), & c_{ijk}(y) &:= c_{x_i x_j x_k}(0, \dots, 0, y), \\ \hat{\varphi}(y) &:= \varphi(0, \dots, 0, y), & \hat{\varphi}_i(y) &:= \varphi_{x_i}(0, \dots, 0, y) \end{aligned}$$

and

$$\begin{aligned} b_I &:= \{b_i\}_{i=1}^{n-1}, & b'_I &:= \{b'_i\}_{i=1}^{n-1}, & b''_I &:= \{b''_i\}_{i=1}^{n-1}, \\ \hat{\varphi}_I &:= \{\hat{\varphi}_i\}_{i=1}^{n-1}, & c_{IJ} &:= \{c_{ij}\}_{i,j=1}^{n-1}, & c_{IJK} &:= \{c_{ijk}\}_{i,j,k=1}^{n-1}. \end{aligned}$$

Then we have the following expressions

$$(1.14) \quad \mathbf{a} = \Lambda^1(a, a', b_I, b'_I, c_{IJ}, c_{IJK}, \hat{\varphi})a'' + \Lambda^2(a, a', b_I, b'_I, c_{IJ}, c_{IJK}, \hat{\varphi}),$$

$$(1.15) \quad \mathbf{a}_i = \Lambda_i^3(a, a', b_I, b'_I, c_{IJ}, c_{IJK}, \hat{\varphi}, \hat{\varphi}_I)a'' + \Lambda_i^4(a, a', b_I, b'_I, c_{IJ}, c_{IJK}, \hat{\varphi}, \hat{\varphi}_I)b''_i + \Lambda_i^5(a, a', b_I, b'_I, c_{IJ}, c_{IJK}, \hat{\varphi}, \hat{\varphi}_I),$$

where  $\Lambda^1, \Lambda^2$  and  $\Lambda_i^l$  ( $l = 3, \dots, 5, i = 1, \dots, n-1$ ) are functions which are  $C^\infty$ -differentiable with respect to the parameters  $a, a', b_I, b'_I, c_{IJ}, c_{IJK}, \hat{\varphi}, \hat{\varphi}_I$ . For example, the explicit expressions of  $\mathbf{a}$  and  $\mathbf{a}_i$  for  $M = \mathbf{R}_1^3$  are given in [4, (4.4) and (4.5)].

Since  $\mathcal{R} = 1$  at  $o$ ,  $\Lambda^1$  and  $\Lambda_i^4$  ( $i = 1, \dots, n-1$ ) do not vanish at  $o$  under the initial condition (1.6). Regarding  $c_{IJ}, c_{IJK}, \hat{\varphi}$  and  $\hat{\varphi}_I$  are functions of  $y$ , we can rewrite (1.12) and (1.13) in the following forms

$$(1.16) \quad a'' = P(y, a, a', b_I, b'_I),$$

$$(1.17) \quad b''_i = Q_i(y, a, a', b_I, b'_I) \quad (i = 1, 2, \dots, n-1),$$

where  $P, Q_i$  are functions of  $y, a, a', b_I, b'_I$ .

For example, we consider the case  $M = \mathbf{R}_1^{n+1}$ . If we set

$$\begin{aligned} c_{ij}(y) &:= c_{x_i x_j}(0, \dots, 0, y), & c_{ijk}(y) &:= c_{x_i x_j x_k}(0, \dots, 0, y), \\ c'_{ij}(y) &:= c_{x_i x_j y}(0, \dots, 0, y), \end{aligned}$$

then

$$(1.18) \quad \alpha = Ca'' + \sum_{j=1}^{n-1} c_{jj}(D + b_j^2) + 2 \sum_{i < j} b_i b_j c_{ij} + 2 \sum_{i=1}^{n-1} a' b_i b'_i,$$

$$(1.19) \quad \begin{aligned} \alpha_l &= Cb''_l - 2 \left( \sum_{j=1}^{n-1} b_j c_{jl} \right) a'' + \sum_{j=1}^{n-1} (D + b_j^2) c_{jjl} \\ &\quad - 2 \sum_{j=1}^{n-1} \left( a' b'_l + \sum_{i=1, i \neq j}^{n-1} b_i c_{il} \right) c_{jj} + 2 \sum_{i < j} (b_j c_{il} c_{ij} + b_i c_{jl} c_{ij} + b_i b_j c_{ijl}) \\ &\quad + 2 \sum_{i=1}^{n-1} (c_{il} a' b'_i + b_i b'_l b'_i + b_i a' c'_{il}) \end{aligned}$$

hold for  $l = 1, \dots, n-1$ , where  $C := 1 - \sum_{i=1}^{n-1} b_i^2$  and  $D := 1 - (a')^2 - \sum_{i=1}^{n-1} b_i^2$ .

In particular, (1.16) and (1.17) give a normal form for a system of ordinary differential equations thinking  $a, b_i$  ( $i = 1, \dots, n-1$ ) are unknown functions, and can be considered as precise expressions of [3, (4.12)]. Since  $P, Q_i$  ( $i = 1, \dots, n-1$ ) are  $C^\infty$ -differentiable with respect to the variables  $a, a', b_I, b'_I$ , this system of ordinary differential equations satisfies the local Lipschitz condition. So the uniqueness of the solution follows. On the other hand, by applying [3, Proposition 4.2],

$$(1.20) \quad a(y) = y, \quad b_i(y) = 0 \quad (i = 1, \dots, n-1)$$

gives a solution of this system satisfying the initial condition (1.6). However, we should remark that in the proof of [3, Proposition 4.2], we set (cf. (4.14))

$$f_0(x_1, \dots, x_n) := x_n + \sum_{1 \leq j \leq k \leq n-1} c_{j,k}(x_1, \dots, x_n) x_j x_k.$$

However, by the same reason as Remark 1, we should replace it by

$$f_0(x_1, \dots, x_n) := x_n + c(x_1, \dots, x_n),$$

where  $c$  is a  $C^3$ -function satisfying (1.3) and (1.4). Then

$$x_n \mapsto (f_0(0, \dots, 0, x_n), 0, \dots, 0, x_n)$$

gives a light-like geodesic of  $M$  consisting of degenerate light-like points.

So, we can conclude that (1.6) implies (1.20), that is, we have proven the theorem.

*Remark.* We point out that there is a minor typographical error in the proof of Theorem D in Section 6 in [3]. In fact, in the beginning of the proof, we had set  $f(x, y) = a(y) + b(y)x + h(x, y)x^3$ , but it should be  $f(x, y) = a(y) + b(y)x + h(x, y)x^2$ . The remaining arguments can be read without any need for corrections.

*Acknowledgements.* The authors thank Professor Udo Hertrich-Jeromin for valuable comments.

#### REFERENCES

- [1] S. Akamine, A. Honda, M. Umehara and K. Yamada, *Bernstein-type theorem for zero mean curvature hypersurfaces without time-like points in Lorentz-Minkowski space*, Bull. Braz. Math. Soc. (N.S.), (2020), DOI 10.1007/s00574-020-00196-8.
- [2] V. A. Klyachin, *Zero mean curvature surfaces of mixed type in Minkowski space*, Izv. Math., **67** (2003), 209–224.
- [3] M. Umehara and K. Yamada, *Hypersurfaces with light-like points in a Lorentzian manifold*, J. Geom. Anal., **29** (2019), 3405–3437.
- [4] M. Umehara and K. Yamada, *Surfaces with light-like points in Lorentz-Minkowski space with applications*, in “Lorentzian Geometry and Related Topics”, Springer Proceedings of Mathematics & Statics **211**, 2017, pp. 253–273, Springer-Verlag.

(Umehara) DEPARTMENT OF MATHEMATICAL AND COMPUTING SCIENCES, TOKYO INSTITUTE OF TECHNOLOGY, TOKYO 152-8552, JAPAN

*E-mail address:* umehara@is.titech.ac.jp

(Yamada) DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, TOKYO 152-8551, JAPAN

*E-mail address:* kotaro@math.titech.ac.jp