

SOME INEQUALITIES ON h -CONVEX FUNCTIONS

M. ABBASI¹, A. MORASSAEI¹ AND F. MIRZAPOUR¹

ABSTRACT. In this paper, we state some characterizations of h -convex function is defined on a convex set in a linear space. By doing so, we extend the Jensen-Mercer inequality for h -convex function. We will also define h -convex function for operators on a Hilbert space and present the operator version of the Jensen-Mercer inequality. Lastly, we propound the complementary inequality of Jensen's inequality for h -convex functions.

1. INTRODUCTION

Assume that I is an interval in \mathbb{R} . Let us recall definitions of some special classes of functions.

We say that [6] $f : I \rightarrow \mathbb{R}$ is a Godunova-Levin function, or that f belongs to the class $Q(I)$ if f is non-negative and for all $x, y \in I$ and $t \in (0, 1)$ we have

$$f(tx + (1-t)y) \leq \frac{f(x)}{t} + \frac{f(y)}{1-t}.$$

For $s \in (0, 1]$, a function $f : [0, \infty) \rightarrow [0, \infty)$ is said to be s -convex function, or that f belongs to the class K_s^2 , if

$$f(tx + (1-t)y) \leq t^s f(x) + (1-t)^s f(y)$$

for every $x, y \in [0, \infty)$ and $t \in [0, 1]$ (see [1]). Also, we say that $f : I \rightarrow [0, \infty)$ is a P -function [4], or that f belongs to the class $P(I)$, if for all $x, y \in I$ and $t \in [0, 1]$ we have

$$f(tx + (1-t)y) \leq f(x) + f(y).$$

Throughout this paper, suppose that I and J are intervals in \mathbb{R} , $(0, 1) \subseteq J$ and functions h and f are real non-negative functions defined on J and I , respectively.

In [10], Varošanec defined the h -convex function as follows:
Let $h : J \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a non-negative function, $h \not\equiv 0$. We say that $f : I \rightarrow \mathbb{R}$

2010 *Mathematics Subject Classification.* Primary 47A63; Secondary 26D15.

Key words and phrases. h -convex function; Jensen-Mercer inequality; operator inequality; Hilbert space.

is a h -convex function, or that f belongs to the class $SX(h, I)$, if f is non-negative and for all $x, y \in I$, $t \in (0, 1)$ we have

$$f(tx + (1 - t)y) \leq h(t)f(x) + h(1 - t)f(y). \quad (1.1)$$

If inequality (1.1) is reversed, then f is said to be h -concave, that is $f \in SV(h, I)$.

If $h(t) = t$, then all non-negative convex functions belong to $SX(h, I)$ and all non-negative concave functions belong to $SV(h, I)$. If $h(t) = \frac{1}{t}$, then $SX(h, I) = Q(I)$; if $h(t) = 1$, then $SX(h, I) \supseteq P(I)$; and if $h(t) = t^s$, where $s \in (0, 1)$, then $SX(h, I) \supseteq K_s^2$.

A function $h : J \rightarrow \mathbb{R}$ is said to be a *super-additive function* if

$$h(x + y) \geq h(x) + h(y), \quad (1.2)$$

for all $x, y \in J$. If inequality (1.2) is reversed, then h is said to be a *sub-additive function*. If the equality holds in (1.2), then h is said to be an *additive function*.

Function h is called a *super-multiplicative function* if

$$h(xy) \geq h(x)h(y), \quad (1.3)$$

for all $x, y \in J$ [10]. If inequality (1.3) is reversed, then h is called a *sub-multiplicative function*. If the equality holds in (1.3), then h is called a *multiplicative function*.

Example 1.1. [10] Consider the function $h : [0, +\infty) \rightarrow \mathbb{R}$ by $h(x) = (c + x)^{p-1}$. If $c = 0$, then the function h is multiplicative. If $c \geq 1$, then for $p \in (0, 1)$ the function h is super-multiplicative and for $p > 1$ the function h is sub-multiplicative.

2. PRELIMINARIES

In what follows we assume that \mathcal{H} and \mathcal{K} are Hilbert spaces, $\mathbb{B}(\mathcal{H})$ and $\mathbb{B}(\mathcal{K})$ are C^* -algebras of all bounded linear operators on the appropriate Hilbert space with identities $I_{\mathcal{H}}$ and $I_{\mathcal{K}}$, $\mathbb{B}_h(\mathcal{H})$ denotes the algebra of all self-adjoint operators in $\mathbb{B}(\mathcal{H})$. An operator $A \in \mathbb{B}_h(\mathcal{H})$ is called *positive*, if $\langle Ax, x \rangle \geq 0$ holds for every $x \in \mathcal{H}$ and then we write $A \geq 0$. For $A, B \in \mathbb{B}_h(\mathcal{H})$, we say $A \leq B$ if $B - A \geq 0$. We write $A > 0$ and say A is *strictly positive operator*, if A is a positive invertible operator. Let f be a continuous real valued function defined on an interval I . The function f is called *operator monotone* if $A \leq B$

implies $f(A) \leq f(B)$ for all A, B with spectra in I . A function f is said to be *operator convex* on I if

$$f(tA + (1-t)B) \leq tf(A) + (1-t)f(B),$$

for all $A, B \in \mathbb{B}_h(\mathcal{H})$ with spectra in I and all $t \in [0, 1]$. A map $\Phi : \mathbb{B}(\mathcal{H}) \rightarrow \mathbb{B}(\mathcal{K})$ is called *positive* if $\Phi(A) \geq 0$, whenever $A \geq 0$ and is said to be *normalized* if $\Phi(I_{\mathcal{H}}) = I_{\mathcal{K}}$. We denote by $\mathbf{P}[\mathbb{B}(\mathcal{H}), \mathbb{B}(\mathcal{K})]$ the set of all positive linear maps $\Phi : \mathbb{B}(\mathcal{H}) \rightarrow \mathbb{B}(\mathcal{K})$ and by $\mathbf{P}_N[\mathbb{B}(\mathcal{H}), \mathbb{B}(\mathcal{K})]$ the set of all normalized positive linear maps $\Phi \in \mathbf{P}[\mathbb{B}(\mathcal{H}), \mathbb{B}(\mathcal{K})]$. If $\Phi \in \mathbf{P}_N[\mathbb{B}(\mathcal{H}), \mathbb{B}(\mathcal{K})]$ and f is an operator convex function on an interval I , then

$$f(\Phi(A)) \leq \Phi(f(A)) \quad (\text{Davis-Choï-Jensen's inequality}) \quad (2.1)$$

for every self-adjoint operator A on \mathcal{H} , whose spectrum is contained in I , see [5].

3. CHARACTERIZATIONS

Assume that C is a convex subset of a linear space X and f is an arbitrary real-valued function on C . The non-negative function $f : C \rightarrow \mathbb{R}$ is called *h -convex function* on C , if $f(tx + (1-t)y) \leq h(t)f(x) + h(1-t)f(y)$ for every $x, y \in C$ and $t \in [0, 1]$.

Let x and y be two fixed elements in C . Define the map $f_{x,y}$ as follows:

$$f_{x,y} : [0, 1] \rightarrow \mathbb{R}, \quad f_{x,y}(t) = f(tx + (1-t)y).$$

The following theorem is a characterization of h -convex functions.

Theorem 3.1 (First characterization). *With the above assumptions, the following statements are equivalent:*

- (i) f is a h -convex function on C .
- (ii) The mapping $f_{x,y}$ is a h -convex function on $[0, 1]$, for any $x, y \in C$.

Proof. First, assume that (i) holds. Let $\alpha, \beta \in [0, 1]$ such that $\alpha + \beta = 1$ and $t_1, t_2 \in [0, 1]$. Hence

$$\begin{aligned} f_{x,y}(\alpha t_1 + \beta t_2) &= f((\alpha t_1 + \beta t_2)x + (1 - \alpha t_1 - \beta t_2)y) \\ &= f\left(\alpha(t_1x + (1-t_1)y) + \beta(t_2x + (1-t_2)y)\right) \\ &\leq h(\alpha)f(t_1x + (1-t_1)y) + h(\beta)f(t_2x + (1-t_2)y) \\ &= h(\alpha)f_{x,y}(t_1) + h(\beta)f_{x,y}(t_2), \end{aligned}$$

this means that $f_{x,y}$ is a h -convex function on $[0, 1]$.

Conversely, suppose that (ii) holds. For $t \in [0, 1]$ and $x, y \in C$, we have

$$\begin{aligned} f(tx + (1-t)y) &= f_{x,y}(t) = f_{x,y}((1-t)0 + t1) \\ &\leq h(1-t)f_{x,y}(0) + h(t)f_{x,y}(1) \\ &= h(1-t)f(y) + h(t)f(x), \end{aligned}$$

that is, f is a h -convex function on C . \square

Now, for fixed $t \in [0, 1]$, we define the function $f_t : C^2 \rightarrow \mathbb{R}$ by $f_t(x, y) = f(tx + (1-t)y)$.

In the next theorem, we state a new characterization of h -convex functions.

Theorem 3.2 (Second characterization). *The following statements of h -convex functions hold:*

- (i) *If f is a h -convex function on C , then f_t is a h -convex function on C^2 for every $t \in [0, 1]$.*
- (ii) *If C is a cone in X and f_t is a h -convex function on C^2 for every $t \in (0, 1)$, then f is a h -convex function on C .*

Proof. (i) For fixed $t \in [0, 1]$ and $(x, y), (u, v) \in C^2$. Then for every $\alpha \in [0, 1]$

$$\begin{aligned} f_t(\alpha(x, y) + (1-\alpha)(u, v)) &= f_t(\alpha x + (1-\alpha)u, \alpha y + (1-\alpha)v) \\ &= f(t(\alpha x + (1-\alpha)u) + (1-t)(\alpha y + (1-\alpha)v)) \\ &= f(\alpha(tx + (1-t)y) + (1-\alpha)(tu + (1-t)v)) \\ &\leq h(\alpha)f(tx + (1-t)y) + h(1-\alpha)f(tu + (1-t)v) \\ &= h(\alpha)f_t(x, y) + h(1-\alpha)f_t(u, v), \end{aligned}$$

that is, f_t is a h -convex function on C^2 .

(ii) Let $x, y \in C$ and $t \in (0, 1)$. Since C is cone in X , $C+C \subseteq C$ and $\alpha C \subseteq C$ for every $\alpha \geq 0$, then $t^{-1}x, (1-t)^{-1}y \in C$ and $(t^{-1}x, 0), (0, (1-t)^{-1}y) \in C^2$. On the other hand, by h -convexity of f_t on C^2 , we have

$$\begin{aligned} f(tx + (1-t)y) &= f_t(x, y) \\ &= f_t(t(t^{-1}x, 0) + (1-t)(0, (1-t)^{-1}y)) \\ &\leq h(t)f_t(t^{-1}x, 0) + h(1-t)f_t(0, (1-t)^{-1}y) \\ &= h(t)f(x) + h(1-t)f(y), \end{aligned}$$

therefore, f is a h -convex function on C .

□

Theorem 3.3 (Third characterization). *Let h be a strictly positive multiplicative function, then the following statements are equivalent:*

- (i) f is a h -convex function.
- (ii) If $(1+s)x - sy \in C$, for every $x, y \in C$ and $s \geq 0$, then

$$f((1+s)x - sy) \geq h(1+s)f(x) - h(s)f(y). \quad (3.1)$$

Proof. First, note that multiplicity of h implies that $h\left(\frac{1}{t}\right) = \frac{1}{h(t)}$, for every $t > 0$ and $h(1) = 1$.

Assume that (i) holds. By using $x = \frac{1}{s+1}[(1+s)x - sy] + \frac{s}{s+1}y$, we have

$$\begin{aligned} f(x) &= f\left(\frac{1}{s+1}[(1+s)x - sy] + \frac{s}{s+1}y\right) \\ &\leq h\left(\frac{1}{s+1}\right) f[(1+s)x - sy] + h\left(\frac{s}{s+1}\right) f(y) \\ &= \frac{h(1)}{h(s+1)} f[(1+s)x - sy] + \frac{h(s)}{h(s+1)} f(y) \quad (\text{by multiplicity of } h) \end{aligned}$$

and therefore,

$$f[(1+s)x - sy] \geq h(1+s)f(x) - h(s)f(y).$$

Now, suppose that (ii) holds. If $\alpha, \beta \in [0, 1]$ and $\alpha + \beta = 1$, then there exists $s \geq 0$ such that $\alpha = \frac{1}{s+1}$ and $\beta = \frac{s}{s+1}$. Put $z = \alpha x + \beta y$, hence $x = (1+s)z - sy$ and so

$$f(x) = f((1+s)z - sy) \geq h(1+s)f(z) - h(s)f(y).$$

Consequently,

$$\begin{aligned} f(\alpha x + \beta y) &= f(z) \leq \frac{1}{h(1+s)} f(x) + \frac{h(s)}{h(1+s)} f(y) \\ &= h\left(\frac{1}{1+s}\right) f(x) + h\left(\frac{s}{1+s}\right) f(y) \\ &= h(\alpha)f(x) + h(\beta)f(y), \end{aligned}$$

this show that f is h -convex function.

□

Theorem 3.4. (i) Assume that X is a real vector space and $f : X \rightarrow \mathbb{R}$ is an even h -convex function. Then

$$\begin{aligned} \frac{f((1-2t)x) + f((2t-1)y)}{h(t) + h(1-t)} &\leq f((1-t)x + ty) + f(tx + (1-t)y) \\ &\leq [h(t) + h(1-t)][f(x) + f(y)]. \end{aligned} \quad (3.2)$$

(ii) Let X be a topological vector space, h be an integrable strictly positive function and f be a continuous even h -convex function, then

$$\frac{1}{2} \int_0^1 [f(tx) + f(ty)] dt \leq \int_0^1 [h(t) + h(1-t)] f(tx + (1-t)y) dt. \quad (3.3)$$

In addition, if h is super-additive, then

$$\frac{1}{2h(1) \left(\int_0^1 h(t) dt \right)} \int_0^1 [f(tx) + f(ty)] dt \leq f(x) + f(y). \quad (3.4)$$

Proof. (i) Since f is a h -convex function, hence the right inequality of (3.2) is clear. Set $a = (1-t)x + ty$ and $b = -tx - (1-t)y$, we have $(1-t)a + tb = (1-2t)x$ and $ta + (1-t)b = (2t-1)y$. Therefore

$$\begin{aligned} &f((1-2t)x) + f((2t-1)y) \\ &= f((1-t)a + tb) + f(ta + (1-t)b) \\ &\leq [h(t) + h(1-t)][f(a) + f(b)] \quad (\text{by the right inequality of (3.2)}) \\ &= [h(t) + h(1-t)][f((1-t)x + ty) + f(-tx - (1-t)y)] \\ &= [h(t) + h(1-t)][f((1-t)x + ty) + f(tx + (1-t)y)]. \quad (f \text{ is a even function}) \end{aligned}$$

We therefore deduce the desired inequality in (3.2).

(ii) By using (3.2) inequality, we get

$$f((1-2t)x) + f((2t-1)y) \leq [h(t) + h(1-t)] [f((1-t)x + ty) + f(tx + (1-t)y)]. \quad (3.5)$$

Integrating each side of (3.5), we have

$$\begin{aligned} &\int_0^1 f((1-2t)x) dt + \int_0^1 f((2t-1)y) dt \\ &\leq \int_0^1 [h(t) + h(1-t)] f((1-t)x + ty) dt + \int_0^1 [h(t) + h(1-t)] f(tx + (1-t)y) dt \\ &= 2 \int_0^1 [h(t) + h(1-t)] f(tx + (1-t)y) dt. \end{aligned} \quad (3.6)$$

Since f is even and by changing of variables $u = 1 - 2t$, yield

$$\int_0^1 f((1-2t)x) dt = \int_0^1 f(tx) dt,$$

and similarly

$$\int_0^1 f((2t-1)y) \, dt = \int_0^1 f(ty) \, dt.$$

Consequently, by (3.6), the proof of (3.3) completes.

Now, assume that h is super-additive. Hence by (3.3), we have

$$\begin{aligned} & \frac{1}{2} \int_0^1 [f(tx) + f(ty)] \, dt \\ & \leq \int_0^1 [h(t) + h(1-t)] f(tx + (1-t)y) \, dt \\ & \leq h(1) \int_0^1 [h(t)f(x) + h(1-t)f(y)] \, dt \quad (h \text{ is super-additive and } f \text{ is } h\text{-convex}) \\ & = h(1) \left[f(x) \left(\int_0^1 h(t) \, dt \right) + f(y) \left(\int_0^1 h(1-t) \, dt \right) \right] \\ & = h(1) \left(\int_0^1 h(t) \, dt \right) [f(x) + f(y)], \end{aligned}$$

this show that (3.4) holds. \square

Corollary 3.5. (i) *Assume that X is a real vector space and $f : X \rightarrow \mathbb{R}$ is an even convex function. Then*

$$\begin{aligned} f((1-2t)x) + f((2t-1)y) & \leq f((1-t)x + ty) + f(tx + (1-t)y) \\ & \leq f(x) + f(y). \end{aligned} \quad (3.7)$$

(ii) *Let X be a topological vector space and f be a continuous even convex function, then*

$$\frac{1}{2} \int_0^1 [f(tx) + f(ty)] \, dt \leq \int_0^1 f(tx + (1-t)y) \, dt \leq f(x) + f(y). \quad (3.8)$$

Proof. Enough put in Theorem 3.4, $h(t) = t$. \square

Corollary 3.6. [9, Lemma 3.2]

(i) *Assume that X is a real vector space and $f : X \rightarrow \mathbb{R}$ is an even function in $P(I)$. Then*

$$\begin{aligned} \frac{f((1-2t)x) + f((2t-1)y)}{2} & \leq f((1-t)x + ty) + f(tx + (1-t)y) \\ & \leq 2(f(x) + f(y)). \end{aligned} \quad (3.9)$$

(ii) Let X be a topological vector space and f be a continuous even function in $P(I)$, then

$$\frac{1}{4} \int_0^1 [f(tx) + f(ty)] dt \leq \int_0^1 f(tx + (1-t)y) dt \leq f(x) + f(y). \quad (3.10)$$

Proof. In Theorem 3.4, put $h(t) = 1$. \square

Example 3.7. [9, Theorem 3.3] Let $(X, \|\cdot\|)$ be a normed space, $x, y \in X$ and $0 < p < 1$. Since $f(x) = \|x\|^p$ is an even continuous P -convex function, we have the following Hermit-Hadamard inequality

$$\frac{\|x\|^p + \|y\|^p}{4(p+1)} \leq \int_0^1 \|(1-t)x + ty\|^p dt \leq \|x\|^p + \|y\|^p.$$

4. JENSEN-MERCER TYPE INEQUALITY

In [8], Mercer proved that

$$f\left(x_1 + x_n - \sum_{j=1}^n t_j x_j\right) \leq f(x_1) + f(x_n) - \sum_{j=1}^n t_j f(x_j). \quad (4.1)$$

where x_j 's also satisfy in the condition $0 < x_1 \leq x_2 \leq \cdots \leq x_n$, $t_j \geq 0$ and $\sum_{j=1}^n t_j = 1$.

In this section, we present the Jensen-Mercer inequality for h -convex functions.

Theorem 4.1. [10, Theorem 19] Let t_1, \dots, t_n be positive real numbers ($n \geq 2$). If h is a non-negative super-multiplicative function, f is a h -convex function on I and $x_1, \dots, x_n \in I$, then

$$f\left(\frac{1}{T_n} \sum_{j=1}^n t_j x_j\right) \leq \sum_{j=1}^n h\left(\frac{t_j}{T_n}\right) f(x_j), \quad (4.2)$$

where $T_n = \sum_{j=1}^n t_j$.

Lemma 4.2. Let $0 < x \leq y$ and f be a h -convex function, then for every $z \in [x, y]$, there exists $\lambda \in [0, 1]$ such that

$$f(x + y - z) \leq [h(\lambda) + h(1 - \lambda)][f(x) + f(y)] - f(z). \quad (4.3)$$

Moreover, if h is super-additive, then

$$f(x + y - z) \leq h(1)[f(x) + f(y)] - f(z).$$

Proof. Since $z \in [x, y]$, there exists $\lambda \in [0, 1]$ such that

$$z = \lambda x + (1 - \lambda)y.$$

By using h -convexity of f , we have

$$\begin{aligned} f(x + y - z) &= f((1 - \lambda)x + \lambda y) \\ &\leq h(1 - \lambda)f(x) + h(\lambda)f(y) \\ &= [h(\lambda) + h(1 - \lambda)][f(x) + f(y)] - [h(\lambda)f(x) + h(1 - \lambda)f(y)] \\ &\leq [h(\lambda) + h(1 - \lambda)][f(x) + f(y)] - f(\lambda x + (1 - \lambda)y) \\ &= [h(\lambda) + h(1 - \lambda)][f(x) + f(y)] - f(z). \end{aligned}$$

If h is super-additive, then $h(\lambda) + h(1 - \lambda) \leq h(1)$. So the end part of theorem holds. \square

Theorem 4.3. *Let f be a h -convex function on an interval containing the x_j ($j = 1, \dots, n$) such that $0 < x_1 \leq \dots \leq x_n$, then*

$$f\left(x_1 + x_n - \sum_{j=1}^n t_j x_j\right) \leq \left(\sum_{j=1}^n h(t_j)[h(\lambda_j) + h(1 - \lambda_j)]\right) (f(x_1) + f(x_n)) - \sum_{j=1}^n h(t_j)f(x_j),$$

where for every $j = 1, \dots, n$, there exists $\lambda_j \in [0, 1]$ such that $x_j = \lambda_j x_1 + (1 - \lambda_j)x_n$.

Proof. With the above assumption, we have

$$\begin{aligned} &f\left(x_1 + x_n - \sum_{j=1}^n t_j x_j\right) \\ &= f\left(\sum_{j=1}^n t_j (x_1 + x_n - x_j)\right) \\ &\leq \sum_{j=1}^n h(t_j)f(x_1 + x_n - x_j) \\ &\leq \sum_{j=1}^n h(t_j)\left([h(\lambda_j) + h(1 - \lambda_j)](f(x_1) + f(x_n)) - f(x_j)\right) \text{ (by Lemma 4.2)} \\ &= \sum_{j=1}^n h(t_j)[h(\lambda_j) + h(1 - \lambda_j)](f(x_1) + f(x_n)) - \sum_{j=1}^n h(t_j)f(x_j), \end{aligned}$$

this completes the proof. \square

Corollary 4.4. *With the assumptions of previous theorem, if h is a super-additive function such that for every probability vector (t_1, \dots, t_n) , $\sum_{j=1}^n h(t_j) \leq 1$, then*

$$f\left(x_1 + x_n - \sum_{j=1}^n t_j x_j\right) \leq h(1)(f(x_1) + f(x_n)) - \sum_{j=1}^n h(t_j)f(x_j).$$

Moreover, if h is multiplicative, then

$$f\left(x_1 + x_n - \sum_{j=1}^n t_j x_j\right) \leq f(x_1) + f(x_n) - \sum_{j=1}^n h(t_j)f(x_j).$$

5. OPERATOR h -CONVEX FUNCTIONS

In this section, we present the definition of operator h -convex function for operators acting on a Hilbert space.

Definition 5.1. Let $h : J \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a non-negative function, $h \not\equiv 0$. We say that $f : I \rightarrow \mathbb{R}$ is an *operator h -convex function*, if f is non-negative and for all $A, B \in \mathbb{B}(\mathcal{H})$ with $\sigma(A), \sigma(B) \subseteq I$ and $t \in (0, 1)$,

$$f(tA + (1-t)B) \leq h(t)f(A) + h(1-t)f(B), \quad (5.1)$$

where $\sigma(A)$ and $\sigma(B)$ are spectrum of A and B , respectively.

If inequality (5.1) is reversed, then f is said to be *operator h -concave*.

If $t = \frac{1}{2}$ in (5.1), then f is called *h -mid-convex function*.

Example 5.2. Assume that h is a function on $[0, \infty)$ such that $h(t) \geq t$ and $f(t) = t^2$ on an interval $I \subseteq \mathbb{R}$. Then f is operator h -mid-convex function. Because,

$$\begin{aligned} & h\left(\frac{1}{2}\right) (A^2 + B^2) - \left(\frac{A+B}{2}\right)^2 \\ &= h\left(\frac{1}{2}\right) (A^2 + B^2) - \frac{A^2 + AB + BA + B^2}{4} \\ &= \frac{(4h(1/2) - 1)A^2 - AB - BA + (4h(1/2) - 1)B^2}{4} \\ &\geq \frac{1}{4} (A^2 - AB - BA - B^2) \\ &= \frac{1}{4} (A - B)^2 \geq 0. \end{aligned}$$

Now, we can prove the following theorem as Theorem 1.9 in [5]. So, we omit the proof it.

Theorem 5.3 (Jensen's type operator inequality). *Let \mathcal{H} and \mathcal{K} be Hilbert space. Assume that h is non-negative super-multiplicative function and f is a real valued function on an interval $I \subseteq \mathbb{R}$. Suppose that $A, A_j \in \mathbb{B}_h(\mathcal{H})$ such that $\sigma(A), \sigma(A_j) \subseteq I$ ($j = 1, \dots, n$). Then the following conditions are equivalent:*

- (i) f is operator h -mid-convex on I ;
- (ii) $f(C^*AC) \leq 2h(\frac{1}{2})C^*f(A)C$ for every self-adjoint operator $A : \mathcal{H} \rightarrow \mathcal{H}$ and isometry $C : \mathcal{K} \rightarrow \mathcal{H}$, i.e.; $C^*C = 1_{\mathcal{K}}$;
- (iii) $f(C^*AC) \leq 2h(\frac{1}{2})C^*f(A)C$ for every self-adjoint operator $A : \mathcal{H} \rightarrow \mathcal{H}$ and isometry $C : \mathcal{H} \rightarrow \mathcal{H}$;
- (iv) $f\left(\sum_{j=1}^n C_j^* A_j C_j\right) \leq 2h(\frac{1}{2}) \sum_{j=1}^n C_j^* f(A_j) C_j$ for every self-adjoint operator $A_j : \mathcal{H} \rightarrow \mathcal{H}$ and bounded linear operators $C_j : \mathcal{K} \rightarrow \mathcal{H}$; with $\sum_{j=1}^n C_j^* C_j = 1_{\mathcal{K}}$ ($j = 1, \dots, n$);
- (v) $f\left(\sum_{j=1}^n C_j^* A_j C_j\right) \leq 2h(\frac{1}{2}) \sum_{j=1}^n C_j^* f(A_j) C_j$ for every self-adjoint operator $A_j : \mathcal{H} \rightarrow \mathcal{H}$ and bounded linear operators $C_j : \mathcal{H} \rightarrow \mathcal{H}$; with $\sum_{j=1}^n C_j^* C_j = 1_{\mathcal{H}}$ ($j = 1, \dots, n$);
- (vi) $f\left(\sum_{j=1}^n P_j A_j P_j\right) \leq 2h(\frac{1}{2}) \sum_{j=1}^n P_j f(A_j) P_j$ for every self-adjoint operator $A_j : \mathcal{H} \rightarrow \mathcal{H}$ and projection $P_j : \mathcal{H} \rightarrow \mathcal{H}$; with $\sum_{j=1}^n P_j = 1_{\mathcal{H}}$ ($j = 1, \dots, n$).

Using an idea of [5] we prove the following result.

Theorem 5.4 (Davis-Choi-Jensen's inequality). *Let Φ be a normalized positive linear map and f be an operator h -convex function on an interval I , then*

$$f(\Phi(A)) \leq 2h\left(\frac{1}{2}\right) \Phi(f(A)), \quad (5.2)$$

for every self-adjoint operator A with $\sigma(A) \subseteq I$.

Proof. We know that a self-adjoint operator A can be approximated uniformly by a simple function $A' = \sum_j t_j E_j$ where $\{E_j\}$ is a decomposition of the unit $I_{\mathcal{H}}$. By using normality of Φ , we get $\sum_j \Phi(E_j) = I_{\mathcal{K}}$. By applying (iv) of

Theorem 5.3 to $C_j = \sqrt{\Phi(E_j)}$, we have

$$\begin{aligned}
f(\Phi(A')) &= f\left(\sum_j t_j \Phi(E_j)\right) = f\left(\sum_j C_j t_j C_j\right) \\
&\leq 2h\left(\frac{1}{2}\right) \sum_j C_j f(t_j) C_j = 2h\left(\frac{1}{2}\right) \sum_j f(t_j) \Phi(E_j) \\
&= 2h\left(\frac{1}{2}\right) \Phi\left(\sum_j f(t_j) E_j\right) = 2h\left(\frac{1}{2}\right) \Phi(f(A')).
\end{aligned}$$

Since Φ is continuous, the proof is complete. \square

Theorem 5.5. *Let t_1, t_2, \dots, t_n be positive real numbers ($n \geq 2$) such that $\sum_{j=1}^n t_j = 1$. If h is non-negative super-multiplicative function and if f is h -convex function on an interval $I \subseteq \mathbb{R}$, A_1, \dots, A_n are self-adjoint operators in $\mathbb{B}(\mathcal{H})$ such that $\sigma(A_j) \subseteq I$, then*

$$f\left(\sum_{j=1}^n t_j A_j\right) \leq \sum_{j=1}^n h(t_j) f(A_j). \quad (5.3)$$

If h is sub-multiplicative and f is operator h -concave on I , then inequality (5.3) is reversed.

Proof. We prove this theorem by induction on n . If $n = 2$, then inequality (5.3) is equivalent to inequality (5.1) with $t = t_1$ and $1 - t = t_2$. Assume that inequality (5.3) holds for $n - 1$. Then for n , we have

$$\begin{aligned}
f\left(\sum_{j=1}^n t_j A_j\right) &= f\left(t_n A_n + \sum_{j=1}^{n-1} t_j A_j\right) \\
&= f\left(t_n A_n + (t_1 + \dots + t_{n-1}) \sum_{j=1}^{n-1} \frac{t_j}{t_1 + \dots + t_{n-1}} A_j\right) \\
&\leq h(t_n) f(A_n) + h(t_1 + \dots + t_{n-1}) f\left(\sum_{j=1}^{n-1} \frac{t_j}{t_1 + \dots + t_{n-1}} A_j\right) \\
&\leq h(t_n) f(A_n) + h(t_1 + \dots + t_{n-1}) \sum_{j=1}^{n-1} h\left(\frac{t_j}{t_1 + \dots + t_{n-1}}\right) f(A_j) \\
&\leq h(t_n) f(A_n) + \sum_{j=1}^{n-1} h(t_j) f(A_j) \\
&= \sum_{j=1}^n h(t_j) f(A_j).
\end{aligned}$$

This completes the proof. \square

Corollary 5.6. *By assumptions of Theorem 5.5, if $\Phi \in \mathbf{P}_N[\mathbb{B}(\mathcal{H}), \mathbb{B}(\mathcal{K})]$, then*

$$f\left(\sum_{j=1}^n t_j \Phi(A_j)\right) \leq \sum_{j=1}^n 2h\left(\frac{t_j}{2}\right) \Phi(f(A_j)). \quad (5.4)$$

If h is sub-multiplicative and f is operator h -concave on I , then inequality (5.4) is reversed.

Proof. If $\Phi \in \mathbf{P}_N[\mathbb{B}(\mathcal{H}), \mathbb{B}(\mathcal{K})]$, then

$$\begin{aligned} f\left(\sum_{j=1}^n t_j \Phi(A_j)\right) &\leq \sum_{j=1}^n h(t_j) f(\Phi(A_j)) \\ &\leq \sum_{j=1}^n 2h(t_j) h\left(\frac{1}{2}\right) \Phi(f(A_j)) \quad (\text{by (5.2)}) \\ &\leq \sum_{j=1}^n 2h\left(\frac{t_j}{2}\right) \Phi(f(A_j)) \quad (\text{by super-multiplicity of } h). \end{aligned}$$

\square

For convenience, let $\varphi(t)$ be a real valued continuous function on the interval $[m, M]$. Define

$$\mu_\varphi = \frac{\varphi(M) - \varphi(m)}{M - m}, \quad \nu_\varphi = \frac{M\varphi(m) - m\varphi(M)}{M - m}.$$

We remark that a straight line $\ell(t) = \mu_\varphi t + \nu_\varphi$ is a line through two points $(m, \varphi(m))$ and $(M, \varphi(M))$.

Notice that, if $\varphi(t) = t$, then $\mu_\varphi = 1$ and $\nu_\varphi = 0$, if $\varphi(t) = 1$, then $\mu_\varphi = 0$ and $\nu_\varphi = 1$, and if $\varphi(t) = \frac{1}{t}$, then $\mu_\varphi = -\frac{1}{mM}$ and $\nu_\varphi = \frac{m+M}{mM}$.

Theorem 5.7. *Let $A_1, A_2, \dots, A_n \in \mathbb{B}(\mathcal{H})$ be self-adjoint operators with spectra in $[m, M]$ for some scalars m, M and $\Phi_1, \Phi_2, \dots, \Phi_n \in \mathbf{P}_N[\mathbb{B}(\mathcal{H}), \mathbb{B}(\mathcal{K})]$ and t_1, \dots, t_n non-negative real numbers with $\sum_{j=1}^n t_j = 1$. If f on $[m, M]$ is operator h -convex function and h on the interval J is super-multiplicative*

operator convex function, then

$$\begin{aligned}
& f \left(mI_{\mathcal{K}} + MI_{\mathcal{K}} - \sum_{j=1}^n t_j \Phi_j(A_j) \right) \\
& \leq h \left(\frac{\sum_{j=1}^n t_j \Phi_j(A_j) - mI_{\mathcal{K}}}{M - m} \right) f(m) + h \left(\frac{MI_{\mathcal{K}} - \sum_{j=1}^n t_j \Phi_j(A_j)}{M - m} \right) f(M) \\
& \leq (\mu_h + 2\nu_h)(f(m) + f(M))I_{\mathcal{K}} - \sum_{j=1}^n h(t_j)\Phi_j(f(A_j)).
\end{aligned} \tag{5.5}$$

Proof. Define the function $g : [m, M] \rightarrow \mathbb{R}$ by $g(t) = f(m + M - t)$. Since f is continuous and h -convex on $[m, M]$, so the same is true for g . Consequently, for every $t \in [m, M]$

$$f(t) \leq h \left(\frac{t - m}{M - m} \right) f(M) + h \left(\frac{M - t}{M - m} \right) f(m), \tag{5.6}$$

and

$$g(t) \leq h \left(\frac{t - m}{M - m} \right) g(M) + h \left(\frac{M - t}{M - m} \right) g(m). \tag{5.7}$$

Since $\sum_{j=1}^n t_j = 1$, $\Phi_j(I_{\mathcal{H}}) = I_{\mathcal{K}}$ and $mI_{\mathcal{H}} \leq A_j \leq MI_{\mathcal{H}}$ ($j = 1, \dots, n$), we conclude that $mI_{\mathcal{K}} \leq \sum_{j=1}^n t_j \Phi_j(A_j) \leq MI_{\mathcal{K}}$. Now, by using functional calculus and (5.6) and super-multiplicity of h , we get

$$\begin{aligned}
h(t_j)f(A_j) & \leq h(t_j)h \left(\frac{A_j - mI_{\mathcal{H}}}{M - m} \right) f(M) + h(t_j)h \left(\frac{MI_{\mathcal{H}} - A_j}{M - m} \right) f(m) \\
& \leq h \left(\frac{t_j A_j - mt_j I_{\mathcal{H}}}{M - m} \right) f(M) + h \left(\frac{Mt_j I_{\mathcal{H}} - t_j A_j}{M - m} \right) f(m) \\
& \leq \left[\mu_h \left(\frac{t_j A_j - mt_j I_{\mathcal{H}}}{M - m} \right) + \nu_h I_{\mathcal{H}} \right] f(M) \\
& \quad + \left[\mu_h \left(\frac{Mt_j I_{\mathcal{H}} - t_j A_j}{M - m} \right) + \nu_h I_{\mathcal{H}} \right] f(m).
\end{aligned} \tag{5.8}$$

Hence, by linearity of Φ_j for every $j = 1, \dots, n$ and the inequality (5.8), we have

$$\begin{aligned}
h(t_j)\Phi_j(f(A_j)) & \leq \left[\mu_h \left(\frac{t_j \Phi_j(A_j) - mt_j I_{\mathcal{K}}}{M - m} \right) + \nu_h I_{\mathcal{K}} \right] f(M) \\
& \quad + \left[\mu_h \left(\frac{Mt_j I_{\mathcal{K}} - t_j \Phi_j(A_j)}{M - m} \right) + \nu_h I_{\mathcal{K}} \right] f(m).
\end{aligned}$$

By summing of all $j = 1, \dots, n$

$$\begin{aligned}
 \sum_{j=1}^n h(t_j) \Phi_j(f(A_j)) &\leq \left[\mu_h \left(\frac{\sum_{j=1}^n t_j \Phi_j(A_j) - m I_{\mathcal{K}}}{M - m} \right) + \nu_h I_{\mathcal{K}} \right] f(M) \quad (5.9) \\
 &\quad + \left[\mu_h \left(\frac{M I_{\mathcal{K}} - \sum_{j=1}^n t_j \Phi_j(A_j)}{M - m} \right) + \nu_h I_{\mathcal{K}} \right] f(m) \\
 &= \mu_h \left[\mu_f \sum_{j=1}^n t_j \Phi_j(A_j) + \nu_f I_{\mathcal{K}} \right] + \nu_h (f(m) + f(M)) I_{\mathcal{K}}
 \end{aligned}$$

Also, using similar way and the (5.7) inequality, we have

$$\begin{aligned}
 &g \left(\sum_{j=1}^n t_j \Phi_j(A_j) \right) \\
 &\leq h \left(\frac{\sum_{j=1}^n t_j \Phi_j(A_j) - m I_{\mathcal{K}}}{M - m} \right) f(m) + h \left(\frac{M I_{\mathcal{K}} - \sum_{j=1}^n t_j \Phi_j(A_j)}{M - m} \right) f(M)
 \end{aligned}$$

or equivalently,

$$\begin{aligned}
 &f \left(m I_{\mathcal{K}} + M I_{\mathcal{K}} - \sum_{j=1}^n t_j \Phi_j(A_j) \right) = g \left(\sum_{j=1}^n t_j \Phi_j(A_j) \right) \\
 &\leq h \left(\frac{\sum_{j=1}^n t_j \Phi_j(A_j) - m I_{\mathcal{K}}}{M - m} \right) f(m) + h \left(\frac{M I_{\mathcal{K}} - \sum_{j=1}^n t_j \Phi_j(A_j)}{M - m} \right) f(M) \\
 &\leq \left[\mu_h \left(\frac{\sum_{j=1}^n t_j \Phi_j(A_j) - m I_{\mathcal{K}}}{M - m} \right) + \nu_h I_{\mathcal{K}} \right] f(m) \\
 &\quad + \left[\mu_h \left(\frac{M I_{\mathcal{K}} - \sum_{j=1}^n t_j \Phi_j(A_j)}{M - m} \right) + \nu_h I_{\mathcal{K}} \right] f(M) \\
 &= \mu_h \left[f(m) I_{\mathcal{K}} + f(M) I_{\mathcal{K}} - \left(\frac{M I_{\mathcal{K}} - \sum_{j=1}^n t_j \Phi_j(A_j)}{M - m} f(m) \right. \right. \\
 &\quad \left. \left. + \frac{\sum_{j=1}^n t_j \Phi_j(A_j) - m I_{\mathcal{K}}}{M - m} f(M) \right) \right] + \nu_h (f(m) + f(M)) I_{\mathcal{K}} \\
 &= \mu_h \left[(f(m) + f(M)) I_{\mathcal{K}} - \left(\mu_f \sum_{j=1}^n t_j \Phi_j(A_j) + \nu_f I_{\mathcal{K}} \right) \right] + \nu_h (f(m) + f(M)) I_{\mathcal{K}} \\
 &\leq (\mu_h + 2\nu_h) (f(m) + f(M)) I_{\mathcal{K}} - \sum_{j=1}^n h(t_j) \Phi_j(f(A_j)).
 \end{aligned}$$

In the final inequality, we use the inequality (5.9), and we obtain desired inequalities (5.5). \square

Corollary 5.8. (i) *Let $h(t) = t$ in Theorem 5.7, then we have*

$$\begin{aligned} & f \left(mI_{\mathcal{K}} + MI_{\mathcal{K}} - \sum_{j=1}^n t_j \Phi_j(A_j) \right) \\ & \leq \left(\frac{\sum_{j=1}^n t_j \Phi_j(A_j) - mI_{\mathcal{K}}}{M - m} \right) f(m) + \left(\frac{MI_{\mathcal{K}} - \sum_{j=1}^n t_j \Phi_j(A_j)}{M - m} \right) f(M) \\ & \leq f(m)I_{\mathcal{K}} + f(M)I_{\mathcal{K}} - \sum_{j=1}^n t_j \Phi_j(f(A_j)). \end{aligned}$$

(ii) *Let $h(t) = 1$ in Theorem 5.7, then we have*

$$f \left(mI_{\mathcal{K}} + MI_{\mathcal{K}} - \sum_{j=1}^n t_j \Phi_j(A_j) \right) \leq 2(f(m) + f(M))I_{\mathcal{K}} - \sum_{j=1}^n t_j \Phi_j(f(A_j)).$$

With similar proof of Theorem 5.7, we have the following proposition.

Proposition 5.9. *Let $A_1, A_2, \dots, A_n \in \mathbb{B}(\mathcal{H})$ be self-adjoint operators with spectra in $[m, M]$ for some scalars and $\Phi_1, \Phi_2, \dots, \Phi_n \in \mathbf{P}[\mathbb{B}(\mathcal{H}), \mathbb{B}(\mathcal{K})]$ positive linear maps with $\sum_{j=1}^n \Phi_j(I_{\mathcal{H}}) = I_{\mathcal{K}}$. If f on $[m, M]$ is operator h -convex function and h on the interval J is operator convex function, then*

$$\begin{aligned} & f \left(mI_{\mathcal{K}} + MI_{\mathcal{K}} - \sum_{j=1}^n \Phi_j(A_j) \right) \\ & \leq h \left(\frac{\sum_{j=1}^n \Phi_j(A_j) - mI_{\mathcal{K}}}{M - m} \right) f(m) + h \left(\frac{MI_{\mathcal{K}} - \sum_{j=1}^n \Phi_j(A_j)}{M - m} \right) f(M) \\ & \leq (\mu_h + 2\nu_h)(f(m) + f(M))I_{\mathcal{K}} - \sum_{j=1}^n \Phi_j(f(A_j)). \end{aligned}$$

Corollary 5.10. [7, Theorem 1] *Let $h(t) = t$ in Proposition 5.9, then we have*

$$\begin{aligned} & f \left(mI_{\mathcal{K}} + MI_{\mathcal{K}} - \sum_{j=1}^n \Phi_j(A_j) \right) \\ & \leq \left(\frac{\sum_{j=1}^n \Phi_j(A_j) - mI_{\mathcal{K}}}{M - m} \right) f(m) + \left(\frac{MI_{\mathcal{K}} - \sum_{j=1}^n \Phi_j(A_j)}{M - m} \right) f(M) \\ & \leq f(m)I_{\mathcal{K}} + f(M)I_{\mathcal{K}} - \sum_{j=1}^n \Phi_j(f(A_j)). \end{aligned}$$

Theorem 5.11. Suppose that $A_j \in \mathbb{B}_h(\mathcal{H})$ with $\sigma(A_j) \in [m, M]$ ($m < M$), $\Phi_j \in \mathbf{P}_N[\mathbb{B}(\mathcal{H}), \mathbb{B}(\mathcal{K})]$ ($j = 1, \dots, n$) and $t_1, \dots, t_n \geq 0$ such that $\sum_{j=1}^n t_j = 1$. If $f, g \in \mathcal{C}([m, M])$ and $F(u, v)$ is a real valued continuous function defined on $U \times V$, where $f[m, M] \subset U, g[m, M] \subset V$ and F is an operator monotone function relative to the first component u and f is a non-negative operator monotone h -convex and h is super-multiplicative operator convex function on J , then

$$\begin{aligned} & F \left[\sum_{j=1}^n h(t_j) \Phi_j(f(A_j)), g \left(\sum_{j=1}^n t_j \Phi_j(A_j) \right) \right] \\ & \leq \max_{m \leq t \leq M} F [\mu_h \mu_f t + \mu_h \nu_f + \nu_h (f(m) + f(M)), g(t)] I_{\mathcal{K}}. \end{aligned} \quad (5.10)$$

Proof. With the above assumptions and similar proof of previous theorem, we have

$$\sum_{j=1}^n h(t_j) \Phi_j(f(A_j)) \leq \mu_h \left[\mu_f \sum_{j=1}^n t_j \Phi_j(A_j) + \nu_f I_{\mathcal{K}} \right] + \nu_h (f(m) + f(M)) I_{\mathcal{K}}.$$

Consequently,

$$\begin{aligned} & F \left[\sum_{j=1}^n h(t_j) \Phi_j(f(A_j)), g \left(\sum_{j=1}^n t_j \Phi_j(A_j) \right) \right] \\ & \leq F \left[\mu_h \left[\mu_f \sum_{j=1}^n t_j \Phi_j(A_j) + \nu_f I_{\mathcal{K}} \right] + \nu_h (f(m) + f(M)) I_{\mathcal{K}}, g \left(\sum_{j=1}^n t_j \Phi_j(A_j) \right) \right] \\ & \leq \max_{m \leq t \leq M} F [\mu_h \mu_f t + \mu_h \nu_f + \nu_h (f(m) + f(M)), g(t)] I_{\mathcal{K}}, \end{aligned}$$

and we have the desired inequality (5.10). \square

Theorem 5.12. With assumptions of previous theorem, if f is operator h -convex on $[m, M]$ and h is operator convex function on J , then for every $\alpha \in \mathbb{R}$

$$\sum_{j=1}^n h(t_j) \Phi_j(f(A_j)) \leq \alpha g \left(\sum_{j=1}^n t_j \Phi_j(A_j) \right) + \beta I_{\mathcal{K}}, \quad (5.11)$$

where

$$\beta = \max_{m \leq t \leq M} \{ \mu_h \mu_f t + \mu_h \nu_f + \nu_h (f(m) + f(M)) - \alpha g(t) \}.$$

In addition,

(i) If αg is concave, then

$$\beta \geq \max_{s \in \{m, M\}} \{ \mu_h f(s) + \nu_h (f(m) + f(M)) - \alpha g(s) \}.$$

(ii) If αg is strictly convex differentiable, then

$$\beta \leq \mu_h f(s) - \alpha g(s) + |\mu_h \mu_f - \alpha g'(s)|(M - m) + \nu_h(f(m) + f(M)),$$

where $s \in \{m, M\}$.

Proof. Let $\alpha \in \mathbb{R}$. Define $F(u, v) = u - \alpha v$. Since F is operator monotone on a first variable u , hence by Theorem 5.11, we have

$$\begin{aligned} & \sum_{j=1}^n h(t_j) \Phi_j(f(A_j)) - \alpha g \left(\sum_{j=1}^n t_j \Phi_j(A_j) \right) \\ & \leq \max_{m \leq t \leq M} \{ \mu_h \mu_f t + \mu_h \nu_f + \nu_h(f(m) + f(M)) - \alpha g(t) \} I_K, \end{aligned}$$

and we have the desired inequality (5.11).

Put $\Psi(t) = \mu_h(\mu_f t + \nu_f) + \nu_h(f(m) + f(M)) - \alpha g(t)$. In the case (i), if $\alpha g(t)$ is concave on $[m, M]$, then Ψ is convex on $[m, M]$ and therefore

$$\begin{aligned} \beta &= \max_{m \leq t \leq M} \Psi(t) = \max\{\Psi(m), \Psi(M)\} \\ &= \max_{s \in \{m, M\}} \{ \mu_h f(s) + \nu_h(f(m) + f(M)) - \alpha g(s) \}. \end{aligned}$$

In the case (ii), if αg is strictly convex differentiable, then there exists $t_0 \in [m, M]$ such that $\beta = \max_{m \leq t \leq M} \Psi(t) = \Psi(t_0)$. So, if $s \in \{m, M\}$, then

$$\begin{aligned} \beta &= \mu_h(\mu_f t_0 + \nu_f) + \nu_h(f(m) + f(M)) - \alpha g(t_0) \\ &= \mu_h f(s) + \mu_h \mu_f(t_0 - s) - \alpha g(t_0) + \nu_h(f(m) + f(M)) \\ &= \mu_h f(s) - \alpha g(s) + \mu_h \mu_f(t_0 - s) - \alpha g(t_0) + \alpha g(s) + \nu_h(f(m) + f(M)) \\ &\leq \mu_h f(s) - \alpha g(s) + \mu_h \mu_f(t_0 - s) - \alpha g'(s)(t_0 - s) + \nu_h(f(m) + f(M)) \\ &\quad (\alpha g \text{ is strictly convex differentiable}) \\ &\leq \mu_h f(s) - \alpha g(s) + |\mu_h \mu_f - \alpha g'(s)|(M - m) + \nu_h(f(m) + f(M)). \end{aligned}$$

□

Corollary 5.13 (Complementary inequality of Jensen's inequality). *Let A_j , Φ_j and t_j ($j = 1, \dots, n$) be as in Theorem 5.12. If $f \in \mathcal{C}([m, M])$ is a function which is nonnegative, strictly h -convex and twice differentiable, then for every $\alpha \in \mathbb{R}^+$*

$$\sum_{j=1}^n h(t_j) \Phi_j(f(A_j)) \leq \alpha f \left(\sum_{j=1}^n t_j \Phi_j(A_j) \right) + \beta I_K, \quad (5.12)$$

where $\beta = \mu_h \mu_f t_0 + \mu_h \nu_f + \nu_h (f(m) + f(M)) - \alpha f(t_0)$ and

$$t_0 = \begin{cases} f'^{-1} \left(\frac{1}{\alpha} \mu_h \mu_f \right) & ; \text{ if } \alpha f'(m) < \mu_h \mu_f < \alpha f'(M), \\ m & ; \text{ if } \alpha f'(m) \geq \mu_h \mu_f, \\ M & ; \text{ if } \alpha f'(M) \leq \mu_h \mu_f. \end{cases}$$

REFERENCES

- [1] W.W. Breckner, Stetigkeitsaussagen für eine Klasse verallgemeinerter konvexer funktionen in topologischen linearen Räumen, Publ. Inst. Math. **23** (1978) 13–20.
- [2] V. Darvish, S.S. Dragomir, H.M. Nazari and A. Taghavi, Some inequalities associated with the Hermite-Hadamard inequalities for operators h -convex functions, Acta et Commentationes Universitatis Tratuensis de Mathematica, **21** (2017), 287–297.
- [3] T.H. Dinh and K.T.B. Vo, Some inequalities for operator (p, h) -convex functions, Linear Multilinear A. **66** (2018), 580–592.
- [4] S.S. Dragomir, J. Pečarić and L.E. Persson, Some inequalities of Hadamard type, Soochow J. Math. **21** (1995) 335–341.
- [5] T. Furuta, J. Mićić Hot, J.E. Pečarić and Y. Seo, Mond-Pečarić method in operator inequalities, Element, Zagreb, 2005.
- [6] E.K. Godunova and V.I. Levin, Neravenstva dlja funkci širokogo klassa, soderžaščego vypuklye, monotonnnye i nekotorye drugie vidy funkci, in: Vyčislitel. Mat. i. Mat. Fiz. Mežvuzov. Sb. Nauč. Trudov, MGPI, Moskva, (1985) 138–142.
- [7] A. Matković, J. Pečarić and I. Perić, A variant of Jensen's inequality of Mercer's type for operators with applications, Linear Algebra Appl. **418** (2006), 551–564.
- [8] A. McD. Mercer, A variant of Jensen's inequality, J. Ineq. Pure and Appl. Math., **4** (2003), Article 73. [ONLINE: <http://www.emis.de/journals/JIPAM/article314.html>].
- [9] J. Rooin, S. Habibzadeh and M.S. Moslehian, Jensen inequalities for P -class functions, Period Math. Hung. **77** (2018), 261–273.
- [10] S. Varošanec, On h -convexity, J. Math. Anal. Appl., **326** (2007), 303–311.

¹ DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, UNIVERSITY OF ZANJAN, UNIVERSITY BLVD., ZANJAN 45371-38791, IRAN.

E-mail address: mostafa.abasi5754@gmail.com

E-mail address: morassaei@znu.ac.ir

E-mail address: f.mirza@znu.ac.ir