

# Trajectory Poisson multi-Bernoulli filters

Ángel F. García-Fernández, Lennart Svensson, Jason L. Williams, Yuxuan Xia, Karl Granström

**Abstract**—This paper presents two trajectory Poisson multi-Bernoulli (TPMB) filters for multi-target tracking: one to estimate the set of alive trajectories at each time step and another to estimate the set of all trajectories, which includes alive and dead trajectories, at each time step. The filters are based on propagating a Poisson multi-Bernoulli (PMB) density on the corresponding set of trajectories through the filtering recursion. After the update step, the posterior is a PMB mixture (PMBM) so, in order to obtain a PMB density, a Kullback-Leibler divergence minimisation on an augmented space is performed. The developed filters are computationally lighter alternatives to the trajectory PMBM filters, which provide the closed-form recursion for sets of trajectories with Poisson birth model, and are shown to outperform previous multi-target tracking algorithms.

**Index Terms**—Multitarget tracking, sets of trajectories, Poisson multi-Bernoulli filter.

## I. INTRODUCTION

Multitarget tracking (MTT) consists of inferring the trajectories of an unknown number of targets that appear and disappear from a scene of interest based on noisy sensor data [1], [2]. Multitarget tracking is a fundamental process of numerous applications including advanced driver assistance systems, self-driving vehicles [3], air traffic monitoring [4] and maritime surveillance [5]. There are many approaches to perform multitarget tracking such as multiple hypothesis tracking [6], [7], joint probabilistic data association [8] and the random finite set (RFS) framework [9].

The traditional RFS approach to MTT is mainly concerned with multi-target filtering, in which one aims to estimate the current set of targets, without attempting to estimate target trajectories. With Poisson birth model, the solution to the multi-target filtering problem is given by the Poisson multi-Bernoulli mixture (PMBM) filter [10], [11]. If the birth model is multi-Bernoulli instead of Poisson, the filtering density is given by the multi-Bernoulli mixture (MBM) filter, which corresponds to the PMBM filtering recursion by setting the intensity of the Poisson process to zero and adding Bernoulli components for newborn targets in the prediction [11], [12].

An MBM can also be written as a mixture in which Bernoulli components have deterministic existence instead of probabilistic, giving rise to the MBM<sub>01</sub> filter [11, Sec. IV]. Deterministic existence leads to an exponential growth in the number of mixture components, which is undesirable from a computational point of view. In general, a PMBM is

preferred over MBM/MBM<sub>01</sub> forms due to a more efficient representation of the information on undetected targets, via the intensity of a Poisson RFS, and not limiting a priori the maximum number of new born targets at each time step [12].

Even though the PMBM filter provides a closed-form solution to the multi-target filtering problem, it is also relevant to consider Poisson multi-Bernoulli (PMB) filters [10], [13], which are computationally lighter and also have very good performance [14]. Relations between the PMB filter and the joint integrated data association filter [15] were given in [10].

Track building procedures for PMBM/MBM/PMB filters can be obtained based on filter meta-data, i.e., information contained in the hypothesis trees. However, the posterior itself only provides information about the current set of targets, and not their trajectories. One approach to building trajectories from posterior densities is to add unique labels to the target states and form trajectories by linking target state estimates with the same label [16]–[18]. With labelled multi-Bernoulli birth, the  $\delta$ -generalised labelled multi-Bernoulli ( $\delta$ -GLMB) filter [17] provides the corresponding filtering density, via a recursion that is similar to the MBM<sub>01</sub> filter recursion [11, Sec. IV]. A computationally lighter alternative to the  $\delta$ -GLMB filter is the labelled multi-Bernoulli (LMB) filter [19], which propagates an LMB density. Sequential track building approaches based on labelling can work well in many cases but it is not always adequate [20], [21].

The above track building problems can be solved by computing (multi-object) densities on sets of trajectories [20], rather than sets of labelled targets. This approach has led to the following filters: trajectory PMBM (TPMBM) filter [21], [22], trajectory MBM (TMBM) filter [23], trajectory MBM<sub>01</sub> (TMBM<sub>01</sub>) filter [20], and trajectory PHD (TPHD) and CPHD (TCPHD) filters [24]. These filters are analogous to their set of targets counterparts, but have the ability to estimate trajectories from first principles, and the possibility of improving the estimation of past states in the trajectories.

This paper proposes two trajectory PMB (TPMB) filters that approximate the trajectory PMBM filters [21] using track-oriented MBM merging [10, Sec. IV.A]. One TPMB filter aims to estimate the set of the alive trajectories at each time step, while the other aims to estimate the set of all trajectories (alive and already dead) at each time step. In the TPMB filters, the Poisson component represents information regarding trajectories that have not been detected and the multi-Bernoulli component represents information on trajectories that have been detected at some point in the past. As the true posterior is a TPMBM, the TPMB filter is derived by making use of a KLD minimisation, on a trajectory space with auxiliary variables, after each update step to obtain the TPMB approximation, see Figure 1. The resulting TPMB density also matches the

A. F. García-Fernández is with the Department of Electrical Engineering and Electronics, University of Liverpool, Liverpool L69 3GJ, United Kingdom (angel.garcia-fernandez@liverpool.ac.uk). L. Svensson, Y. Xia, and K. Granström are with the Department of Electrical Engineering, Chalmers University of Technology, SE-412 96 Gothenburg, Sweden (first-name.lastname@chalmers.se). J. L. Williams is with the Commonwealth Scientific and Industrial Research Organization (jason.williams@data61.csiro.au).




Figure 1: Diagram of the two TPMB filters. The first TPMB filter propagates a PMB density on the set of alive trajectories at the current time. After each update step, the posterior is a PMBM so the TPMB filter performs a KLD minimisation, in an augmented trajectory space with an auxiliary variable. The second TPMB filter works equivalently but propagating a PMB density on the set of all trajectories.

PHD of the updated TPMBM. As the TPMB filtering posterior is defined over the set of trajectories, one can estimate set of trajectories directly from this density. In this paper, we also propose a Gaussian implementation of the TPMB filters for linear/Gaussian models. Simulation results show that the TPMB filters have a performance close to the TPMBM filters, with a decrease in computational complexity, and outperform other filters in the literature.

The rest of the paper is organised as follows. We formulate the considered multitarget tracking problems in Section II. The considered TPMB approximation to a TPMBM density using KLD minimisation is obtained in Section III. The resulting TPMB filters are proposed in Section IV and their Gaussian implementations in Section V. Simulation results are shown in Section VI and conclusions are drawn in Section VII.

## II. PROBLEM FORMULATION

We tackle two multi-target tracking problems [21]:

- 1) The estimation of the set of alive trajectories at the current time step.
- 2) The estimation of the set of all trajectories that have ever existed up to the current time step. We refer to this set as the set of all trajectories.

These problems can be solved by calculating the (multi-trajectory) density over the considered set of trajectories. In this paper, we consider a computationally appealing approximation based on Poisson multi-Bernoulli densities.

### A. Set of trajectories

A single target state  $x \in \mathbb{R}^{n_x}$  contains information of interest about the target, e.g., its position and velocity. A set of single target states  $\mathbf{x}$  belongs to  $\mathcal{F}(\mathbb{R}^{n_x})$  where  $\mathcal{F}(\mathbb{R}^{n_x})$  denotes the set of all finite subsets of  $\mathbb{R}^{n_x}$ . We are interested in estimating target trajectories, where a trajectory consists of a sequence of target states that can start at any time step and end any time later on. A trajectory is therefore represented as a variable  $X = (t, x^{1:\nu})$  where  $t$  is the initial time step of the trajectory,  $\nu$  is its length and  $x^{1:\nu} = (x^1, \dots, x^\nu)$  denotes a sequence of length  $\nu$  that contains the target states at consecutive time steps.

We consider trajectories up to the current time step  $k$ . As a trajectory  $(t, x^{1:\nu})$  exists from time step  $t$  to  $t + \nu - 1$ , the variable  $(t, \nu)$  belongs to the set  $I_{(k)} = \{(t, \nu) : 0 \leq t \leq k \text{ and } 1 \leq \nu \leq k - t + 1\}$ . A single trajectory  $X$  up to time step  $k$  therefore belongs to the space  $T_{(k)} = \uplus_{(t, \nu) \in I_{(k)}} \{t\} \times \mathbb{R}^{\nu n_x}$ , where  $\uplus$  stands for union of sets that are mutually disjoint. Similarly to the set  $\mathbf{x}$  of targets, we denote a set of trajectories up to time step  $k$  as  $\mathbf{X} \in \mathcal{F}(T_{(k)})$ .

1) *Integrals and densities:* Given a real-valued function  $\pi(\cdot)$  on the single trajectory space  $T_{(k)}$ , its integral is [20]

$$\int \pi(X) dX = \sum_{(t, \nu) \in I_{(k)}} \int \pi(t, x^{1:\nu}) dx^{1:\nu}. \quad (1)$$

This integral goes through all possible start times, lengths and target states of the trajectory. Given a real-valued function  $\pi(\cdot)$  on the space  $\mathcal{F}(T_{(k)})$  of sets of trajectories, its set integral is

$$\int \pi(\mathbf{X}) \delta \mathbf{X} = \sum_{n=0}^{\infty} \frac{1}{n!} \int \pi(\{X_1, \dots, X_n\}) dX_{1:n} \quad (2)$$

where  $X_{1:n} = (X_1, \dots, X_n)$ . A function  $\pi(\cdot)$  is a multitarget density of a random finite set of trajectories if  $\pi(\cdot) \geq 0$  and its set integral is one.

### B. Multi-target Bayesian models

The multi-target state evolves according to a Markov system with the following modelling:

- P1 Given the set  $\mathbf{x}_k$  of targets at time step  $k$ , each target  $x \in \mathbf{x}_k$  survives with probability  $p^S(x)$  and moves to a new state with a transition density  $g(\cdot|x)$ , or dies with probability  $1 - p^S(x)$ .
- P2 The multitarget state  $\mathbf{x}_{k+1}$  is the union of the surviving targets and new targets, where new targets are born independently following a PPP with intensity  $\lambda^B(\cdot)$ .

Set  $\mathbf{x}_k$  is observed through the set  $\mathbf{z}_k$  of measurements, which is modelled as

- U1 Each target state  $x \in \mathbf{x}_k$  is either detected with probability  $p^D(x)$  and generates one measurement with density  $l(\cdot|x)$ , or missed with probability  $1 - p^D(x)$ .
- U2 The set  $\mathbf{z}_k$  is the union of the target-generated measurements and Poisson clutter with intensity  $\lambda^C(\cdot)$ .

U1 and U2 imply that a measurement is generated by at most one target. The objective is to compute the posterior density of the set  $\mathbf{X}_k$  of trajectories at time step  $k$  given the sequence  $(\mathbf{z}_1, \dots, \mathbf{z}_k)$  of measurements up to time step  $k$ . Depending on the problem formulation, see the beginning of Section II, this set can correspond to the set of all trajectories or the set of alive trajectories, and the meaning will be clear from context so we use the same notation. If  $\mathbf{X}_k$  denotes the set of alive trajectories at time step  $k$ , then  $t + \nu - 1 = k$  for each  $(t, x^{1:\nu}) \in \mathbf{X}_k$ . If  $\mathbf{X}_k$  denotes the set of all trajectories at time step  $k$ , then  $t + \nu - 1 \leq k$  for each  $(t, x^{1:\nu}) \in \mathbf{X}_k$ .

### C. Poisson multi-Bernoulli mixture

Given the sequence of measurements  $(\mathbf{z}_1, \dots, \mathbf{z}_k)$  and the models in Section II-B, the density  $f_{k'|k}(\cdot)$  of the set of trajectories at time step  $k' \in \{k, k+1\}$  is a Poisson multi-Bernoulli mixture [21]. This holds for the two problem formulations we consider, though the specific recursions to compute  $f_{k'|k}(\cdot)$  vary. In both cases,  $f_{k'|k}(\cdot)$  is a PMBM with

$$\begin{aligned} f_{k'|k}(\mathbf{X}_{k'}) &= \sum_{\mathbf{Y} \uplus \mathbf{W} = \mathbf{X}_{k'}} f_{k'|k}^p(\mathbf{Y}) f_{k'|k}^{mbm}(\mathbf{W}) \\ f_{k'|k}^p(\mathbf{X}_{k'}) &= e^{-\int \lambda_{k'|k}(X) dX} [\lambda_{k'|k}(\cdot)]^{\mathbf{X}_{k'}} \end{aligned} \quad (3)$$

$$f_{k'|k}^{mbm}(\mathbf{X}_{k'}) = \sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a \sum_{\uplus_{l=1}^{n_{k'|k}} \mathbf{X}^l = \mathbf{X}_{k'}} \prod_{i=1}^{n_{k'|k}} f_{k'|k}^{i,a^i}(\mathbf{X}^i)$$

where the sum in (3) goes through all disjoint and possibly empty sets  $\mathbf{Y}$  and  $\mathbf{W}$  such that  $\mathbf{Y} \cup \mathbf{W} = \mathbf{X}_{k'}$ , the multi-object exponential is defined as  $h^{\mathbf{X}} = \prod_{X \in \mathbf{X}} h(X)$  where  $h$  is a real-valued function and  $h^\emptyset = 1$  by convention, and

$$f_{k'|k}^{i,a^i}(\mathbf{X}) = \begin{cases} 1 - r_{k'|k}^{i,a^i} & \mathbf{X} = \emptyset \\ r_{k'|k}^{i,a^i} p_{k'|k}^{i,a^i}(X) & \mathbf{X} = \{X\} \\ 0 & \text{otherwise.} \end{cases} \quad (4)$$

We proceed to explain the aspects of (3) that are relevant to our contribution, for details, we refer the reader to [10], [21].

From (3), the PMBM is the union of two independent RFS, a Poisson RFS with (multi-trajectory) density  $f_{k'|k}^p(\cdot)$  that represents undetected trajectories, and a mixture of multi-Bernoulli RFS  $f_{k'|k}^{mbm}(\cdot)$  that represents trajectories that have been detected at some point up to time step  $k$ . The intensity of the Poisson RFS is denoted as  $\lambda_{k'|k}(\cdot)$ . The number of Bernoulli components is  $n_{k'|k}$ ,  $i$  is an index over the Bernoulli components and a global hypothesis is  $a = (a^1, \dots, a^{n_{k'|k}})$ , where  $a^i$  is the index to the single-trajectory hypothesis for the  $i$ -th Bernoulli component. The weight of global hypothesis  $a$  is  $w_{k'|k}^a$  and the set  $\mathcal{A}_{k'|k}$  contains all possible global hypotheses [10]. The function  $f_{k'|k}^{i,a^i}(\cdot)$  is the density of the  $i$ -th Bernoulli component with single trajectory hypothesis  $a^i$ . Its probability of existence and single-trajectory density are given by  $r_{k'|k}^{i,a^i}$  and  $p_{k'|k}^{i,a^i}(\cdot)$ . Equation (3) can also be written as

$$f_{k'|k}(\mathbf{X}_{k'}) = \sum_{\uplus_{l=1}^{n_{k'|k}} \mathbf{X}^l \uplus \mathbf{Y} = \mathbf{X}_{k'}} f_{k'|k}^p(\mathbf{Y}) \sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a \prod_{i=1}^{n_{k'|k}} \left[ f_{k'|k}^{i,a^i}(\mathbf{X}^i) \right]. \quad (5)$$

### III. TRAJECTORY PMB APPROXIMATION

In this section we first introduce auxiliary variables in the PMBM (5) in Section III-A. Then, we provide the best PMB approximation to the PMBM with auxiliary variables, in the sense of minimising the resulting KLD in Section III-B. Finally, in Section III-C, we show the relation between the KLDs with and without auxiliary variables.

#### A. PMBM with auxiliary variables

In this section, we introduce an auxiliary variable in the PMBM representation in (5) that will be useful to obtain the PMB approximation. Auxiliary variables are commonly used in Bayesian inference to deal with mixtures of densities, especially in sampling-based methods [16], [25]–[31] and latent variable models, e.g., expectation-maximisation [32]. Given (5), we extend the single trajectory space with an auxiliary variable  $u \in \mathbb{U}_{k'|k} = \{0, 1, \dots, n_{k'|k}\}$ , such that  $(u, X) \in \mathbb{U}_{k'|k} \times T_{(k')}$ . As will become clearer at the end of this section, variable  $u = 0$  implies that the single trajectory has not yet been detected, so it corresponds to the PPP, and

$u = i$  indicates that the single trajectory corresponds to the  $i$ -th Bernoulli component. We denote a set of trajectories with auxiliary variables as  $\tilde{\mathbf{X}}_{k'} \in \mathcal{F}(\mathbb{U}_{k'|k} \times T_{(k')})$ .

**Definition 1.** Given  $f_{k'|k}(\cdot)$  of the form (5), we define the density  $\tilde{f}_{k'|k}(\cdot)$  on the space  $\mathcal{F}(\mathbb{U}_{k'|k} \times T_{(k')})$  of sets of trajectories with auxiliary variable as

$$\begin{aligned} \tilde{f}_{k'|k}(\tilde{\mathbf{X}}_{k'}) &= \sum_{\uplus_{l=1}^{n_{k'|k}} \tilde{\mathbf{X}}^l \uplus \tilde{\mathbf{Y}} = \tilde{\mathbf{X}}_{k'}} \tilde{f}_{k'|k}^p(\tilde{\mathbf{Y}}) \sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a \prod_{i=1}^{n_{k'|k}} \left[ \tilde{f}_{k'|k}^{i,a^i}(\tilde{\mathbf{X}}^i) \right] \end{aligned} \quad (6)$$

$$= \tilde{f}_{k'|k}^p(\tilde{\mathbf{Y}}_{k'}) \sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a \prod_{i=1}^{n_{k'|k}} \left[ \tilde{f}_{k'|k}^{i,a^i}(\tilde{\mathbf{X}}_{k'}^i) \right] \quad (7)$$

where, for a given  $\tilde{\mathbf{X}}_{k'}$ ,  $\tilde{\mathbf{Y}}_{k'} = \{(u, X) \in \tilde{\mathbf{X}}_{k'} : u = 0\}$  and  $\tilde{\mathbf{X}}_{k'}^i = \{(u, X) \in \tilde{\mathbf{X}}_{k'} : u = i\}$ , and

$$\begin{aligned} \tilde{f}_{k'|k}^p(\tilde{\mathbf{X}}) &= e^{-\int \lambda_{k'|k}(x) dx} \left[ \tilde{\lambda}_{k'|k}(\cdot) \right]^{\tilde{\mathbf{X}}} \\ \tilde{\lambda}_{k'|k}(u, X) &= \delta_0[u] \lambda_{k'|k}(X) \\ \tilde{f}_{k'|k}^{i,a^i}(\tilde{\mathbf{X}}) &= \begin{cases} 1 - r_{k'|k}^{i,a^i} & \tilde{\mathbf{X}} = \emptyset \\ r_{k'|k}^{i,a^i} p_{k'|k}^{i,a^i}(X) \delta_i[u] & \tilde{\mathbf{X}} = \{(u, X)\} \\ 0 & \text{otherwise} \end{cases} \end{aligned} \quad (8)$$

where the Kronecker delta  $\delta_i[u] = 1$  if  $u = i$  and  $\delta_i[u] = 0$ , otherwise. Note that the sum over sets disappears in (7) as there is only one possible partition of  $\tilde{\mathbf{X}}_{k'}$  into  $\tilde{\mathbf{Y}}$ ,  $\tilde{\mathbf{X}}^1, \dots, \tilde{\mathbf{X}}^{n_{k'|k}}$  that provides a non-zero density. In addition, if two or more trajectories in  $\tilde{\mathbf{X}}_{k'}$  have non-zero auxiliary variables that are equal, then  $\tilde{f}_{k'|k}(\tilde{\mathbf{X}}_{k'}) = 0$ , as the corresponding Bernoulli component (8) evaluated for more than one trajectory is zero. On the contrary, there can be multiple trajectories in  $\tilde{\mathbf{X}}_{k'}$  whose auxiliary variable is zero, without implying that  $\tilde{f}_{k'|k}(\tilde{\mathbf{X}}_{k'}) = 0$ . Also, the density  $\tilde{f}_{k'|k}(\cdot)$  has domain  $\mathcal{F}(\mathbb{U}_{k'|k} \times T_{(k')})$  which implies that we only consider auxiliary variables  $u \in \mathbb{U}_{k'|k} = \{0, 1, \dots, n_{k'|k}\}$ .

The definition of  $\tilde{f}_{k'|k}(\cdot)$  is mathematically sound as it defines a density in  $\mathcal{F}(\mathbb{U}_{k'|k} \times T_{(k')})$  and, as we prove in Appendix A, integrating out the auxiliary variables, we recover the original density. This procedure of introducing auxiliary variables for PMBM for sets of trajectories is also directly applicable for PMBM for sets of targets. In the target case, the proposed use of auxiliary variables bears resemblance to the approaches in [33] [5, Sec. IX], in which targets that have never been detected are indistinguishable, in our case represented by the PPP with  $u = 0$ , and targets that have been detected are distinguishable, in our case represented by a unique  $u > 0$  and a Bernoulli density.

#### B. KLD minimisation with auxiliary variables

In this section, we derive the best PMB fit of a PMBM using KLD minimisation on the space with auxiliary variables.

1) *Kullback-Leibler divergence*: Given a real-valued function  $\pi(\cdot)$  on the single trajectory space  $\mathbb{U}_{k'|k} \times T_{(k')}$  with auxiliary variable, its single-trajectory integral is

$$\int \pi(\tilde{X}) d\tilde{X} = \sum_{u \in \mathbb{U}_{k'|k}} \sum_{(t, \nu) \in I_{(k')}} \int \pi(u, t, x^{1:\nu}) dx^{1:\nu}. \quad (9)$$

Given two densities  $\tilde{f}(\cdot)$  and  $\tilde{q}(\cdot)$  on the space  $\mathcal{F}(\mathbb{U}_{k'|k} \times T_{(k')})$  of sets of trajectories with auxiliary variable, the KLD  $D(\tilde{f} \parallel \tilde{q})$  is

$$D(\tilde{f} \parallel \tilde{q}) = \int \tilde{f}(\tilde{\mathbf{X}}) \log \frac{\tilde{f}(\tilde{\mathbf{X}})}{\tilde{q}(\tilde{\mathbf{X}})} \delta \tilde{\mathbf{X}}. \quad (10)$$

2) *PMB approximation*: We aim to obtain a PMB approximation  $\tilde{q}(\cdot)$  on the space  $\mathcal{F}(\mathbb{U}_{k'|k} \times T_{(k')})$  such that

$$\tilde{q}(\tilde{\mathbf{X}}_{k'}) = \tilde{q}^p(\tilde{\mathbf{Y}}_{k'}) \prod_{i=1}^{n_{k'|k}} [\tilde{q}^{i,1}(\tilde{\mathbf{X}}_{k'}^i)] \quad (11)$$

where

$$\tilde{q}^p(\tilde{\mathbf{X}}_{k'}) = e^{-\int \lambda^q(X) dX} [\tilde{\lambda}^q(\cdot)]^{\tilde{\mathbf{X}}_{k'}} \\ \tilde{\lambda}^q(u, X) = \delta_0[u] \lambda^q(X)$$

and

$$\tilde{q}^{i,1}(\tilde{\mathbf{X}}) = \begin{cases} 1 - r^i & \tilde{\mathbf{X}} = \emptyset \\ r^i p^i(X) \delta_i[u] & \tilde{\mathbf{X}} = \{(u, X)\} \\ 0 & \text{otherwise.} \end{cases} \quad (12)$$

In (11),  $\lambda^q(\cdot)$  represents the intensity of the PPP,  $r^i$  and  $p^i(\cdot)$  denote the existence probability and single-trajectory density of the  $i$ -th Bernoulli component.

**Proposition 2.** *Given a PMBM density  $\tilde{f}_{k'|k}(\cdot)$  of the form (7), the parameters of the PMB density  $\tilde{q}(\cdot)$ , see (11), that minimises the KLD  $D(\tilde{f}_{k'|k} \parallel \tilde{q})$  are given by*

$$\lambda^q(X) = \lambda_{k'|k}(X) \\ r^i = \sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a r_{k'|k}^{i,a^i} \\ p^i(X) = \frac{\sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a r_{k'|k}^{i,a^i} p_{k'|k}^{i,a^i}(X)}{\sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a r_{k'|k}^{i,a^i}}.$$

Proposition 2 is proved in Appendix B. In addition, in Appendix B, we also show that the PHD of  $\tilde{q}(\cdot)$  matches the PHD of  $\tilde{f}_{k'|k}(\cdot)$ .

### C. Relation between KLDs

In this section, we establish that the KLD on the space of sets of trajectories with auxiliary variables is an upper bound on the KLD for sets of trajectories without auxiliary variables.

**Lemma 3.** *Let  $f_{k'|k}(\cdot)$  and  $\tilde{f}_{k'|k}(\cdot)$  be the PMBMs in (5) and (7). Let  $q(\cdot)$  denote a multi-trajectory density and  $\tilde{q}(\cdot)$  an extension of  $q(\cdot)$  with auxiliary variables. Then,*

$$D(f_{k'|k} \parallel q) \leq D(\tilde{f}_{k'|k} \parallel \tilde{q}).$$

The proof of Lemma 3 is given in Appendix B. The KLD  $D(f_{k'|k} \parallel q)$  is the one we are primarily interested in, as it does not include auxiliary variables. Nevertheless, introducing auxiliary variables enables us to minimise the resulting KLD in closed-form, which is an upper bound on  $D(f_{k'|k} \parallel q)$ .

## IV. TRAJECTORY PMB FILTERS

In Section IV-A, we explain the dynamic and measurement models written for sets of alive trajectories and sets of all trajectories. The filtering recursions of the TPMB filters are provided in Section IV-B.

### A. Bayesian models for sets of trajectories

We proceed to write the Bayesian dynamic/measurement models for the two types of problem formulations [21]. These models are required for the filtering recursions in Section IV-B. In particular, we specify the intensity  $\lambda_{k+1}^B(X)$  of new born trajectories, the single trajectory transition density  $g_{k+1}(\cdot | X)$  and the probability  $p^S(X)$  of survival as a function of a trajectory  $X$ . In this paper,  $p^S(X)$  refers to the probability that a trajectory remains in the considered set of trajectories (either all trajectories or alive trajectories) and is different from  $p^S(x)$  as a function on a target state  $x$ .

1) *Dynamic model for set of alive trajectories*: The set of alive trajectories evolves according to this Markov system

- P1T Given the set  $\mathbf{X}_k$  of alive trajectories at time step  $k$ , each trajectory  $X = (t, x^{1:\nu}) \in \mathbf{X}_k$ , where  $t + \nu - 1 = k$ , either survives with probability  $p^S(X) = p^S(x^\nu)$  with a transition density

$$g_{k+1}(t_y, y^{1:\nu_y} | X) = \delta_t[t_y] \delta_{\nu+1}[\nu_y] \delta_{x^{1:\nu}}(y^{1:\nu_y-1}) \\ \times g(y^{\nu_y} | x^\nu)$$

or dies with probability  $1 - p^S(X)$ .

- P2T The set  $\mathbf{X}_{k+1}$  is the union of the surviving trajectories and new trajectories, which are born independently following a PPP with intensity

$$\lambda_{k+1}^B(t, x^{1:\nu}) = \delta_{k+1}[t] \delta_1[\nu] \lambda^B(x^\nu).$$

2) *Dynamic model for set of all trajectories*: The set of all trajectories evolves according to this Markov system

- P3T Given the set  $\mathbf{X}_k$  of all trajectories at time step  $k$ , each trajectory  $X = (t, x^{1:\nu}) \in \mathbf{X}_k$ , where  $t + \nu - 1 \leq k$ , survives with probability 1,  $p^S(X) = 1$ , with a transition density

$$g_{k+1}(t_y, y^{1:\nu_y} | X) = \delta_t[t_y] \\ \times \begin{cases} \delta_\nu[\nu_y] \delta_{x^{1:\nu}}(y^{1:\nu_y}) & \omega_y < k \\ \delta_\nu[\nu_y] \delta_{x^{1:\nu}}(y^{1:\nu_y}) (1 - p^S(x^\nu)) & \omega_y = k \\ \delta_{\nu+1}[\nu_y] \delta_{x^{1:\nu}}(y^{1:\nu_y-1}) p^S(x^\nu) \\ \quad \times g(y^{\nu_y} | x^\nu) & \omega_y = k + 1 \end{cases}$$

where  $\omega_y = t_y + \nu_y - 1$ .

- Same birth model as in P2T.

It should be noted that  $g_{k+1}(\cdot)$  does not depend on  $k$  in P1T (alive trajectories) but it depends on  $k$  in P3T (all trajectories). Nevertheless, we write the dependence on  $k$  in both settings to have a unified notation for both transition densities.

3) *Measurement model for sets of trajectories*: The measurement model U1 and U2 can be written for sets of all trajectories and sets of alive trajectories in the same manner:

- U1T Each trajectory  $(t, x^{1:\nu}) \in \mathbf{X}_k$ , where  $\mathbf{X}_k$  is the set of (all or alive) trajectories, is detected with probability

$$p_k^D(t, x^{1:\nu}) = \begin{cases} p^D(x^\nu) & t + \nu - 1 = k \\ 0 & \text{otherwise} \end{cases}$$

and generates one measurement with density  $l(\cdot|t, x^{1:\nu}) = l(\cdot|x^\nu)$  or misdetected with probability  $1 - p_k^D(t, x^{1:\nu})$ .

- Same clutter model as in U2.

### B. Filtering recursions

In this section, we present the prediction and update steps of the TPMB filters in Sections IV-B1 and IV-B2. We make use of the following notation. Given two real-valued functions  $a(\cdot)$  and  $b(\cdot)$  on the space of sets of trajectories, we denote

$$\langle a, b \rangle = \int a(X) b(X) dX.$$

1) *Prediction*: Let us consider the filtering density at time step  $k - 1$  over the set of trajectories is a Poisson multi-Bernoulli density of the form

$$f_{k-1|k-1}(\mathbf{X}_{k-1}) = \sum_{\uplus_{l=1}^{n_{k-1|k-1}} \mathbf{X}^l \uplus \mathbf{Y} = \mathbf{X}_{k-1}} f_{k-1|k-1}^p(\mathbf{Y}) \times \prod_{i=1}^{n_{k-1|k-1}} [f_{k-1|k-1}^i(\mathbf{X}^i)] \quad (13)$$

where the intensity of the Poisson component is  $\lambda_{k-1|k-1}(\cdot)$ ,  $n_{k-1|k-1}$  is the number of Bernoulli components and the probability of existence and single target density of the  $i$ -th Bernoulli component are  $r_{k-1|k-1}^i$  and  $p_{k-1|k-1}^i(\cdot)$ .

**Lemma 4** (TPMB prediction). *Given the PMB filtering density on the set of trajectories at time step  $k-1$  in (13), the predicted density at time step  $k$  is a PMB*

$$f_{k|k-1}(\mathbf{X}_{k-1}) = \sum_{\uplus_{l=1}^{n_{k|k-1}} \mathbf{X}^l \uplus \mathbf{Y} = \mathbf{X}_{k-1}} f_{k|k-1}^p(\mathbf{Y}) \times \prod_{i=1}^{n_{k|k-1}} [f_{k|k-1}^i(\mathbf{X}^i)] \quad (14)$$

where  $n_{k|k-1} = n_{k-1|k-1}$  and

$$\begin{aligned} \lambda_{k|k-1}(X) &= \lambda_k^B(X) \\ &+ \int g_k(X|X') p^S(X') \lambda_{k-1|k-1}(X') dX' \\ r_{k|k-1}^i &= r_{k-1|k-1}^i \langle p_{k-1|k-1}^i, p^S \rangle \\ p_{k|k-1}^i(X) &= \frac{\int g_k(X|X') p^S(X') p_{k-1|k-1}^i(X') dX'}{\langle p_{k-1|k-1}^i, p^S \rangle} \end{aligned}$$

where  $g_k(\cdot|\cdot)$ ,  $p^S(\cdot)$  and  $\lambda_{B,k}(\cdot)$  are chosen depending on the problem formulation: for alive trajectories, see Section IV-A1, and for all trajectories, see Section IV-A2.

Lemma 4 is a particular case of the TPMBM predictor [21], as a PMB is a PMBM with only one mixture component.

2) *Update*: The update of the TPMB filter is obtained by first doing a Bayesian update, which yields a PMBM distribution, followed by a KLD minimisation, on the space with auxiliary variables, as was illustrated in Figure 1.

**Lemma 5** (TPMB update). *Given the PMB predicted density on the set of trajectories at time step  $k$  in (14) and a measurement set  $\mathbf{z}_k = \{z_k^1, \dots, z_k^{m_k}\}$ , the updated distribution is a PMBM with parameters*

$$f_{k|k}(\mathbf{X}_k) \propto \sum_{\uplus_{l=1}^{n_{k'|k}} \mathbf{X}^l \uplus \mathbf{Y} = \mathbf{X}_k} f_{k|k}^p(\mathbf{Y}) \times \sum_{a \in \mathcal{A}_k} \prod_{i=1}^{n_{k|k}} [w_{k|k}^{i,a} f_{k|k}^{i,a}(\mathbf{X}^i)] \quad (15)$$

where  $n_{k|k} = n_{k|k-1} + m_k$  and

$$\lambda_{k|k}(X) = (1 - p_k^D(X)) \lambda_{k|k-1}(X).$$

For each Bernoulli component in  $f_{k|k-1}(\cdot)$ ,  $i \in \{1, \dots, n_{k|k-1}\}$ , there are  $h_i = m_k + 1$  new single-trajectory hypotheses, corresponding to a misdetection or and update with one of the measurements. The misdetection hypothesis for Bernoulli component  $i \in \{1, \dots, n_{k|k-1}\}$  is given by  $\mathcal{M}(i, 1) = \emptyset$ ,

$$\begin{aligned} w_{k|k}^{i,1} &= 1 - r_{k|k-1}^i \langle p_{k|k-1}^i, p_k^D \rangle \\ r_{k|k}^{i,1} &= \frac{r_{k|k-1}^i \langle p_{k|k-1}^i, 1 - p_k^D \rangle}{1 - r_{k|k-1}^i \langle p_{k|k-1}^i, p_k^D \rangle} \\ p_{k|k}^{i,1}(X) &= \frac{(1 - p_k^D(X)) p_{k|k-1}^i(X)}{\langle p_{k|k-1}^i, 1 - p_k^D \rangle}. \end{aligned}$$

The detection hypothesis for Bernoulli component  $i \in \{1, \dots, n_{k|k-1}\}$  and measurement  $z_k^j$  is given by  $\mathcal{M}(i, j) = \{j\}$ ,  $r_{k|k}^{i,1+j} = 1$ ,

$$\begin{aligned} w_{k|k}^{i,1+j} &= r_{k|k-1}^i \langle p_{k|k-1}^i, l(z_k^j|\cdot) p_k^D \rangle \\ p_{k|k}^{i,1+j}(X) &= \frac{l(z_k^j|X) p_k^D(X) p_{k|k-1}^i(X)}{\langle p_{k|k-1}^i, l(z_k^j|\cdot) p_k^D \rangle}. \end{aligned}$$

For a new Bernoulli component  $i \in \{n_{k|k-1} + j\}$ ,  $j \in \{1, \dots, m_k\}$ , which is initiated by measurement  $z_k^j$ , there are  $h_i = 2$  single-trajectory hypotheses

$$\mathcal{M}(i, 1) = \emptyset, w_{k|k}^{i,1} = 1, r_{k|k}^{i,1} = 0 \quad (16)$$

$$\mathcal{M}(i, 2) = \{j\}$$

$$w_{k|k}^{i,2} = \lambda^C(z_k^j) + \langle \lambda_{k|k-1}, l(z_k^j|\cdot) p_k^D \rangle \quad (17)$$

$$r_{k|k}^{i,2} = \frac{\langle \lambda_{k|k-1}, l(z_k^j|\cdot) p_k^D \rangle}{\lambda^C(z_k^j) + \langle \lambda_{k|k-1}, l(z_k^j|\cdot) p_k^D \rangle}$$

$$p_{k|k}^{i,2}(X) = \frac{l(z_k^j|X) p_k^D(X) \lambda_{k|k-1}(X)}{\langle \lambda_{k|k-1}, l(z_k^j|\cdot) p_k^D \rangle}.$$

The set  $\mathcal{M}(i, j)$  indicates the measurement index for Bernoulli component  $i$  and single-trajectory hypothesis  $j$ . The set of global data association hypotheses is then

$$\mathcal{A}_k = \left\{ (a_1, \dots, a_{n_{k|k}}) : a_i \in \mathbb{N}_{h_i}, \bigcup_{i=1}^{n_{k|k}} \mathcal{M}(i, a_i) = \mathbb{N}_{m_k}, \right. \\ \left. \mathcal{M}(i, a_i) \cap \mathcal{M}(j, a_j) = \emptyset \forall i \neq j \right\}. \quad (18)$$

where  $\mathbb{N}_{m_k} = \{1, \dots, m_k\}$ .

Lemma 5 corresponds to the TPMBM update [21] when the predicted density is a PMB. It should be noted that Lemma 5 is alike for both problem formulations. Finally, the projection of this PMBM density to a PMB density is obtained by Proposition 2.

## V. GAUSSIAN TPMB FILTERS

In this section, we explain the Gaussian implementation of the TPMB filter for alive trajectories and all trajectories in Section V-A and V-B, respectively. Practical considerations are provided in Section V-C. Trajectory estimation is explained in Section V-D. Finally, a discussion is provided in Section V-E.

We use the notation

$$\mathcal{N}(t, x^{1:\nu}; \tau, \bar{x}, P) = \begin{cases} \mathcal{N}(x^{1:\nu}; \bar{x}, P) & t = \tau, \nu = \iota \\ 0 & \text{otherwise} \end{cases} \quad (19)$$

where  $\iota = \dim(\bar{x})/n_x$ . Equation (19) represents a single trajectory Gaussian density with start time  $\tau$ , duration  $\iota$ , mean  $\bar{x} \in \mathbb{R}^{\iota n_x}$  and covariance matrix  $P \in \mathbb{R}^{\iota n_x \times \iota n_x}$  evaluated at  $(t, x^{1:\nu})$ . We use  $\otimes$  to indicate the Kronecker product and  $0_{m,n}$  is the  $m \times n$  zero matrix.

We make the additional assumptions

- A1 The survival and detection probabilities are constants:  $p^S(x) = p^S$  and  $p^D(x) = p^D$ , see P1 and U1.
- A2  $g(\cdot|x) = \mathcal{N}(\cdot; Fx, Q)$  and  $l(\cdot|x) = \mathcal{N}(\cdot; Hx, R)$ .
- A3 The PHD of the birth density at time step  $k$  is

$$\lambda_k^B(X) = \sum_{j=1}^{n_k^b} w_k^{b,j} \mathcal{N}\left(X; k, \bar{x}_k^{b,j}, P_k^{b,j}\right) \quad (20)$$

where  $n_k^b \in \mathbb{N}$  is the number of components,  $w_k^{b,j}$  is the weight of the  $j$ th component,  $\bar{x}_k^{b,j} \in \mathbb{R}^{n_x}$  its mean and  $P_k^{b,j} \in \mathbb{R}^{n_x \times n_x}$  its covariance matrix.

### A. Gaussian implementation for alive trajectories

In the Gaussian implementation for alive trajectories, the single-trajectory density of the  $i$ -th Bernoulli component, see (13), is of the form

$$p_{k'|k}^i(X) = \mathcal{N}\left(X; t^i, \bar{x}_{k'|k}^i, P_{k'|k}^i\right) \quad (21)$$

where  $t^i$  is the start time,  $\bar{x}_{k'|k}^i$  is the mean,  $P_{k'|k}^i$  the covariance matrix and  $t^i + \dim(\bar{x}_{k'|k}^i)/n_x - 1 = k'$ , which implies that the trajectory is alive at time step  $k'$ ,

The PPP has a Gaussian mixture intensity

$$\lambda_{k'|k}(X) = \sum_{q=1}^{n_{k'|k}^p} w_{k'|k}^{p,q} \mathcal{N}\left(X; t_{k'|k}^{p,q}, \bar{x}_{k'|k}^{p,q}, P_{k'|k}^{p,q}\right). \quad (22)$$

where  $n_{k'|k}^p$  is the number of components,  $w_{k'|k}^{p,q}$  is the weight of the  $q$ th component,  $t_{k'|k}^{p,q}$  is starting time,  $\bar{x}_{k'|k}^{p,q}$  its mean and  $P_{k'|k}^{p,q}$  its covariance matrix. It holds that  $t_{k'|k}^{p,q} + \dim(\bar{x}_{k'|k}^{p,q})/n_x - 1 = k'$ , which implies that we consider intensities over alive targets [24].

The prediction step is given by the following lemma.

**Lemma 6** (GTPMB prediction, alive trajectories). *Assume the filtering density for the alive trajectories is a PMB (13) with  $p_{k-1|k-1}^i(\cdot)$  and  $\lambda_{k-1|k-1}(\cdot)$  given by (21) and (22). Then, the predicted density is a PMB of the form (14) with parameters*

$$r_{k|k-1}^i = p^S r_{k-1|k-1}^i \\ \bar{x}_{k|k-1}^i = \left[ \left( \bar{x}_{k-1|k-1}^i \right)^T, \left( \bar{F}_i \bar{x}_{k-1|k-1}^i \right)^T \right]^T \quad (23)$$

$$P_{k|k-1}^i = \begin{bmatrix} P_{k-1|k-1}^i & P_{k-1|k-1}^i \bar{F}_i^T \\ \bar{F}_i P_{k-1|k-1}^i & \bar{F}_i P_{k-1|k-1}^i \bar{F}_i^T + Q \end{bmatrix} \quad (24)$$

$$\bar{F}_i = [0_{1, \iota^i - 1}, 1] \otimes F,$$

$$\lambda_{k|k-1}(X) = \sum_{q=1}^{n_k^b} w_k^{b,q} \mathcal{N}\left(X; k, \bar{x}_k^{b,q}, P_k^{b,q}\right) + p^S \sum_{q=1}^{n_{k-1|k-1}^p} w_{k-1|k-1}^{p,q} \mathcal{N}\left(X; t_{k-1|k-1}^{p,q}, \bar{x}_{k-1|k-1}^{p,q}, P_{k-1|k-1}^{p,q}\right) \quad (25)$$

where  $\iota^i = \dim(\bar{x}_{k-1|k-1}^i)/n_x$  and  $\bar{x}_{k|k-1}^{p,q}$  and  $P_{k|k-1}^{p,q}$  are obtained using (23) and (24) using  $\bar{x}_{k-1|k-1}^{p,q}$  and  $P_{k-1|k-1}^{p,q}$  instead of  $\bar{x}_{k-1|k-1}^i$  and  $P_{k-1|k-1}^i$ .

The update step is given by the following lemma.

**Lemma 7** (GTPMB update, alive trajectories). *Assume the predicted density is a PMB (14) with  $p_{k|k-1}^i(\cdot)$  and  $\lambda_{k|k-1}(\cdot)$  given by (21) and (22). Then, the updated density is a PMBM of the form (15). In the Gaussian case for alive trajectories, the expressions of Lemma 5 can be written as follows. The PHD of the Poisson component is*

$$\lambda_{k|k}(X) = (1 - p^D) \lambda_{k|k-1}(X). \quad (26)$$

The misdetection hypothesis for Bernoulli component  $i \in \{1, \dots, n_{k|k-1}\}$  has parameters

$$w_{k|k}^{i,1} = 1 - r_{k|k-1}^i p^D \\ r_{k|k}^{i,1} = \frac{r_{k|k-1}^i (1 - p^D)}{1 - r_{k|k-1}^i p^D} \\ p_{k|k}^{i,1}(X) = \mathcal{N}\left(X; t^i, \bar{u}_{k|k}^{i,1}, W_{k|k}^{i,1}\right) \\ = \mathcal{N}\left(X; t^i, \bar{x}_{k|k-1}^i, P_{k|k-1}^i\right).$$

The detection hypothesis for Bernoulli component  $i \in \{1, \dots, n_{k|k-1}\}$  and measurement  $z_k^j$  has  $r_{k|k}^{i,1+j} = 1$ ,

$$w_{k|k}^{i,1+j} = r_{k|k-1}^i p^D \mathcal{N}\left(z_k^j; \bar{z}_i, S_i\right)$$

$$p_{k|k}^{i,1+j}(X) = \mathcal{N}\left(X; t^i, \bar{u}_{k|k}^{i,j}, W_{k|k}^{i,j}\right)$$

where

$$\bar{z}_i = \bar{H}_i \bar{x}_{k|k-1}^i, \quad S_i = \bar{H}_i P_{k|k-1}^i \bar{H}_i^T + R \quad (27)$$

$$\bar{H}_i = [0_{1, \ell^i-1}, 1] \otimes H \quad (28)$$

$$\bar{u}_{k|k}^{i,j} = \bar{x}_{k|k-1}^i + P_{k|k-1}^i \bar{H}_i^T S_i^{-1} (z_k^j - \bar{z}_i) \quad (29)$$

$$W_{k|k}^{i,j} = P_{k|k-1}^i - P_{k|k-1}^i \bar{H}_i^T S_i^{-1} \bar{H}_i P_{k|k-1}^i. \quad (30)$$

and  $\ell^i = \dim(\bar{x}_{k|k-1}^i) / n_x$ .

For the new Bernoulli component  $i \in \{n_{k|k-1} + j\}$  initiated by measurement  $z_k^j$ , the first single-trajectory hypothesis has the parameters in (16). For the second hypothesis, we first calculate  $\langle \lambda_{k|k-1}, l(z_k^j \cdot) \rangle p_k^D$  in (17) for each PHD component (22)  $q \in \{1, \dots, n_{k|k-1}^p\}$ , which is denoted as  $v^q$ ,

$$v^q = p^D \mathcal{N}\left(z_k^j; \bar{H}_q \bar{x}_{k|k-1}^{p,q}, S_q\right) \quad (31)$$

$$\bar{H}_q = [0_{1, \ell^q-1}, 1] \otimes H, \quad S_q = \bar{H}_q P_{k|k-1}^{p,q} \bar{H}_q^T + R$$

Then, we obtain  $q^* = \max_q (v^q)$  and set

$$w_{k|k}^{i,2} = \lambda^C(z_k^j) + \sum_{q=1}^{n_{k|k-1}^p} v^q \quad (32)$$

$$r_{k|k}^{i,2} = \frac{\sum_{q=1}^{n_{k|k-1}^p} v^q}{w_{k|k}^{i,2}} \quad (33)$$

$$p_{k|k}^{i,2}(X) = \mathcal{N}\left(X; t_{k|k-1}^{p,q^*}, \bar{u}_{k|k}^{i,2}, W_{k|k}^{i,2}\right) \quad (34)$$

where

$$\bar{u}_{k|k}^{i,2} = \bar{x}_{k|k-1}^{p,q^*} + P_{k|k-1}^{p,q^*} \bar{H}_{q^*}^T S_{q^*}^{-1} (z_k^j - \bar{H}_{q^*} \bar{x}_{k|k-1}^{p,q^*}) \quad (35)$$

$$W_{k|k}^{i,2} = P_{k|k-1}^{p,q^*} - P_{k|k-1}^{p,q^*} \bar{H}_{q^*}^T S_{q^*}^{-1} \bar{H}_{q^*} P_{k|k-1}^{p,q^*}. \quad (36)$$

In Lemma 7, all equations are closed-form expressions obtained from Lemma 5, except the single trajectory density for the new Bernoulli components (34). The closed-form formula of the single trajectory densities of new Bernoulli components is actually a Gaussian mixture, with potentially different starting times and lengths. The filter becomes computationally more efficient by making a Gaussian approximation. To do so, we take the Gaussian component with highest weight (whose index is  $q^*$ ) to obtain (34). This procedure was referred to as absorption in [24].

After the Bayesian update, the updated density is a PMBM so a PMB density is obtained by applying Proposition 2, see Figure 1. The resulting PMB has the same PPP intensity as the updated PMBM. The resulting single-trajectory densities  $p_{k|k}^i(\cdot)$  are Gaussian mixtures so we perform another Kullback-Leibler minimisation to fit a Gaussian distribution, which is achieved by moment matching [32]. Therefore, the updated Bernoulli components of the resulting PMB density are characterised by

$$w_{k|k}^a = \frac{\prod_{i=1}^{n_{k|k}} w_{k|k}^{i,a^i}}{\sum_{a \in \mathcal{A}_k} \prod_{i=1}^{n_{k|k}} w_{k|k}^{i,a^i}} \quad (37)$$

$$r_{k|k}^i = \sum_{a \in \mathcal{A}_k} w_{k|k}^a r_{k|k}^{i,a^i} \quad (38)$$

$$\bar{x}_{k|k}^i = \sum_{a \in \mathcal{A}_k: r_{k|k}^{i,a^i} > 0} \left[ \frac{w_{k|k}^a r_{k|k}^{i,a^i}}{r_{k|k}^i} \bar{u}_{k|k}^{i,a^i} \right] \quad (39)$$

$$P_{k|k}^i = \sum_{a \in \mathcal{A}_k: r_{k|k}^{i,a^i} > 0} \left[ \frac{w_{k|k}^a r_{k|k}^{i,a^i}}{r_{k|k}^i} \left( W_{k|k}^{i,a^i} + \bar{u}_{k|k}^{i,a^i} (\bar{u}_{k|k}^{i,a^i})^T \right) \right] - \bar{x}_{k|k}^i (\bar{x}_{k|k}^i)^T \quad (40)$$

where we recall that  $\mathcal{A}_k$  is given by (18). Single trajectory hypotheses, whose  $r_{k|k}^{i,a^i} = 0$ , have no associated  $\bar{u}_{k|k}^{i,a^i}$  and  $W_{k|k}^{i,a^i}$  so they are not considered in the above sums.

### B. Gaussian implementation for all trajectories

In the Gaussian implementation for the set of all trajectories, we consider information over all trajectories that have ever been detected and information regarding alive trajectories that have not been detected. That is, as in most practical cases, trajectories that have never been detected and no longer exist are not of importance, the PPP only considers alive trajectories. The intensity of the PPP has the form (22) and the  $i$ -th Bernoulli component has a single-trajectory density

$$p_{k|k}^i(X) = \sum_{l=t^i}^{k'} \beta_{k'|k}^i(l) \mathcal{N}\left(X; t^i, \bar{x}_{k'|k}^i(l), P_{k'|k}^i(l)\right) \quad (41)$$

where  $t^i$  is the start time,  $\beta_{k'|k}^i(l)$  is the probability that the corresponding trajectory terminates at time step  $l$ , and  $\bar{x}_{k'|k}^i(l) \in \mathbb{R}^{\ell^i l n_x}$  and  $P_{k'|k}^i(l) \in \mathbb{R}^{\ell^i l n_x \times \ell^i l n_x}$ , with  $\ell^i l = k' - t^i + 1$ , are the mean and the covariance matrix of the trajectory given that it has length  $l$ .

The prediction step is given by the following lemma.

**Lemma 8** (GTPMB prediction, all trajectories). *Assume the filtering density for the alive trajectories is a PMB of the form (13) with  $p_{k-1|k-1}^i(\cdot)$  and  $\lambda_{k-1|k-1}(\cdot)$  given by (41) and (22). Then, the predicted density is a PMB of the form (14) with  $\lambda_{k|k-1}(\cdot)$  given by (25) and  $r_{k|k-1}^i = r_{k-1|k-1}^i$ . The means and covariance matrices of (41), for  $l \in \{t^i, \dots, k-1\}$ , are  $\bar{x}_{k|k-1}^i(l) = \bar{x}_{k-1|k-1}^i(l)$  and  $P_{k|k-1}^i(l) = P_{k-1|k-1}^i(l)$ , and, for  $l = k$ ,  $\bar{x}_{k|k-1}^i(k)$  and  $P_{k|k-1}^i(k)$  are obtained substituting  $\bar{x}_{k|k-1}^i(k-1)$  and  $P_{k|k-1}^i(k-1)$  into (23) and (24). Finally,*

$$\beta_{k|k-1}^i(l) = \begin{cases} \beta_{k-1|k-1}^i(l) & l \in \{t^i, \dots, k-2\} \\ (1-p^S) \beta_{k-1|k-1}^i(l) & l = k-1 \\ p^S \beta_{k-1|k-1}^i(k-1) & l = k. \end{cases}$$

It is important to notice that the prediction step does not modify the single trajectory density of hypotheses that consider dead trajectories  $l \leq k-2$ . For the hypothesis that considers that the trajectory dies,  $l = k-1$ , the single trajectory density remains unchanged but there is a change in the probability  $\beta_{k'|k}^i(l)$ , as one has to take into account the probability of death  $1 - p^S$ . For the alive hypothesis,  $l = k$ , the mean and covariance matrix are propagated as in the case

of alive trajectories, see Lemma 6, and its corresponding probability  $\beta_{k|k-1}^i(l)$  is obtained using the probability of survival and the probability  $\beta_{k-1|k-1}^i(k-1)$  that the trajectory was alive at the previous time step.

The update step is given by the following lemma.

**Lemma 9** (GTPMB update, all trajectories). *Assume the predicted density is a PMB of the form (14) with  $p_{k|k-1}^i(\cdot)$  and  $\lambda_{k|k-1}(\cdot)$  given by (41) and (22). Then, the updated density is a PMBM of the form (15). In the Gaussian case for all trajectories, the expressions of Lemma 5 can be written as follows. The PPP intensity  $\lambda_{k|k}(\cdot)$  is given by (26). The misdetection hypothesis for Bernoulli component  $i \in \{1, \dots, n_{k|k-1}\}$  has the following parameters. The mean and covariance matrices for  $l \in \{t^i, \dots, k\}$  are  $\bar{u}_{k|k}^{i,1}(l) = \bar{x}_{k|k-1}^i(l)$  and  $W_{k|k}^{i,1}(l) = P_{k|k-1}^i(l)$ , and*

$$\begin{aligned} w_{k|k}^{i,1} &= 1 - r_{k|k-1}^i \beta_{k|k-1}^i(k) p^D \\ r_{k|k}^{i,1} &= \frac{r_{k|k-1}^i (1 - \beta_{k|k-1}^i(k) p^D)}{1 - r_{k|k-1}^i \beta_{k|k-1}^i(k) p^D} \\ \beta_{k|k}^{i,1}(l) &\propto \begin{cases} \beta_{k|k-1}^i(l) & l \in \{t^i, \dots, k-1\} \\ (1 - p^D) \beta_{k|k-1}^i(l) & l = k. \end{cases} \end{aligned}$$

The detection hypothesis for Bernoulli component  $i \in \{1, \dots, n_{k|k-1}\}$  and measurement  $z_k^j$  has  $r_{k|k}^{i,1+j} = 1$

$$\begin{aligned} w_{k|k}^{i,1+j} &= r_{k|k-1}^i \beta_{k|k-1}^i(k) p^D \mathcal{N}(z_k^j; \bar{z}_i, S_i) \\ \beta_{k|k}^{i,1+j}(l) &= \begin{cases} 0 & l \in \{t^i, \dots, k-1\} \\ 1 & l = k \end{cases} \\ p_{k|k}^{i,1+j}(X) &= \mathcal{N}(X; t^i, \bar{u}_{k|k}^{i,j}, W_{k|k}^{i,j}) \end{aligned} \quad (42)$$

where  $\bar{z}_i$ ,  $S_i$ ,  $\bar{u}_{k|k}^{i,j}(k)$  and  $W_{k|k}^{i,j}(k)$  are given by substituting  $\bar{x}_{k|k-1}^i(k)$  and  $P_{k|k-1}^i(k)$  into (27)-(30).

As we only consider alive trajectories in the PPP, for the new Bernoulli component  $i \in \{n_{k|k-1} + j\}$  initiated by measurement  $z_k^j$ , the update is done as in Lemma 7, which uses (31)-(36), and setting  $\beta_{k|k}^{i,2}(k) = 1$  and  $\beta_{k|k}^{i,2}(l) = 0$   $l \neq k$ .

As happened when we considered sets of alive trajectories, after the update we get a PMBM so we fit a PMB by applying Proposition 2, which keeps the PPP unaltered. The existence probability of the  $i$ -th Bernoulli component is also given by (37)-(38). The parameter  $\beta_{k|k}^i(\cdot)$  in (41) is given by

$$\beta_{k|k}^i(l) = \sum_{a \in \mathcal{A}_k: r_{k|k}^{i,a^i} > 0} \left[ \frac{w_{k|k}^a r_{k|k}^{i,a^i} \beta_{k|k}^{i,a^i}(l)}{r_{k|k}^{i,a^i}} \right] \quad (43)$$

for  $l \in \{t^i, \dots, k\}$  and  $\beta_{k|k}^i(l) = 0$  otherwise. For each Bernoulli component, the hypotheses that represent dead trajectories  $l \in \{t^i, \dots, k-1\}$  are the same for all  $a \in \mathcal{A}_k$ , with mean  $\bar{u}_{k|k}^{i,1}(l)$  and covariance matrix  $W_{k|k}^i(l)$ , so the output of Proposition 2 is already in Gaussian form for  $l \in \{t^i, \dots, k-1\}$ . For alive trajectories,  $l = k$ , we perform

moment matching to obtain  $\bar{x}_{k|k}^i(k)$  and  $P_{k|k}^i(k)$  in (41), which yields

$$\begin{aligned} \bar{x}_{k|k}^i(k) &= \sum_{a \in \mathcal{A}_k: r_{k|k}^{i,a^i} > 0} \left[ \frac{w_{k|k}^a r_{k|k}^{i,a^i} \beta_{k|k}^{i,a^i}(k)}{\tilde{r}_{k|k}^i} \bar{u}_{k|k}^{i,a^i}(k) \right] \\ P_{k|k}^i(k) &= \sum_{a \in \mathcal{A}_k: r_{k|k}^{i,a^i} > 0} \left[ \frac{w_{k|k}^a r_{k|k}^{i,a^i} \beta_{k|k}^{i,a^i}(k)}{\tilde{r}_{k|k}^i} \left( W_{k|k}^{i,a^i}(k) \right. \right. \\ &\quad \left. \left. + \bar{u}_{k|k}^{i,a^i}(k) \left( \bar{u}_{k|k}^{i,a^i}(k) \right)^T \right) \right] - \bar{x}_{k|k}^i(k) \left( \bar{x}_{k|k}^i(k) \right)^T \end{aligned} \quad (44)$$

where the normalising constant  $\tilde{r}_{k|k}^i$  is

$$\tilde{r}_{k|k}^i = \sum_{a \in \mathcal{A}_k: r_{k|k}^{i,a^i} > 0} w_{k|k}^a r_{k|k}^{i,a^i} \beta_{k|k}^{i,a^i}(k).$$

We should note that, for each Bernoulli, means and covariance matrices for dead hypotheses ( $\bar{x}_{k|k}^i(l)$  and  $P_{k|k}^i(l)$  for  $l < k$ ) do not need any further updating, and multiple hypotheses for the current time are blended into a single Gaussian.

### C. Practical considerations

In this section, we consider the practical considerations to make the TPMB filters computationally efficient. As time goes on, the lengths of the trajectories increase, the sizes of the covariance matrices scale quadratically and the filtering recursion becomes computationally demanding. A solution to deal with increasingly long trajectories is to use an  $L$ -scan implementation [24] in which, in the prediction step, we approximate the covariance matrices of the PPP components and the Bernoulli components with the form

$$P_{k|k} \approx \text{diag} \left( \tilde{P}_{k|k}^{t^k}, \tilde{P}_{k|k}^{t^k+1}, \dots, \tilde{P}_{k|k}^{k-L}, \tilde{P}_{k|k}^{k-L+1:k} \right) \quad (46)$$

where matrix  $\tilde{P}_{k|k}^{k-L+1:k} \in \mathbb{R}^{L \cdot n_x \times L \cdot n_x}$  represents the joint covariance of the  $L$  last time instants, and  $\tilde{P}_{k|k}^{t^k} \in \mathbb{R}^{n_x \times n_x}$  represents the covariance matrix of the target state at time  $k$ . Thus, states outside the  $L$ -scan window are considered independent and remain unchanged with new measurements.

The update step requires obtaining the weights for all global hypotheses  $a \in \mathcal{A}_k$ . In practice, many of these weights are close to zero and can be pruned before evaluating them by using ellipsoidal gating and solving the corresponding ranked assignment problem. In the implementations, we choose the global hypotheses with  $N_h$  highest weights via Murty's algorithm [34], in combination with the Hungarian algorithm, as in [11]. Other data association solvers are discussed in [35]. A different approach is to directly approximate the marginal data association probabilities of  $w_{k|k}^a$ , which are sufficient to apply Proposition 2, for example as in [36] [37].

We also discard Bernoulli components whose existence probability  $r_{k|k}^i$  is lower than a threshold, and it is also possible to recycle them [38]. When we consider sets of all trajectories, Bernoulli components that represent hypotheses of trajectories that have been detected in the past always have existence probability equal to one. However, the probability



that these trajectories are alive currently, which is given by  $\beta_{k|k}^i(k)$ , can be very low, which implies that the weights (42) of their detection hypotheses is very low. Therefore, in order to avoid computing the weights, means and covariances of these hypotheses, which have negligible weight when  $\beta_{k|k}^i(k)$  is low enough, we set  $\beta_{k|k}^i(k) = 0$  if  $\beta_{k|k}^i(k)$  is less than a threshold  $\Gamma_a$ . In other words, if the probability that a Bernoulli component that was once detected is alive at the current time step is lower than  $\Gamma_a$ , it is considered dead,  $\beta_{k|k}^i(k) = 0$ , which implies that it is no longer updated or predicted, but it is still a component of the multi-Bernoulli of the posterior, see (13). In addition, we discard PPP intensity components, see (22), whose weight is less than a threshold  $\Gamma_p$ .

#### D. Estimation

Given the PMB posterior (13) and a threshold  $\Gamma_d$ , we use the following computationally efficient estimators. For the set of alive trajectories, the estimated set of trajectories at time step  $k$  is  $\left\{ \left( t^i, \bar{x}_{k|k}^i \right) : r_{k|k}^i > \Gamma_d \right\}$ , which reports the starting times and means of the Bernoulli components whose probability of existence is greater than  $\Gamma_d$ . For the set of all trajectories, the estimated set of trajectories at time step  $k$  is  $\left\{ \left( t^i, \bar{x}_{k|k}^i(l^*) \right) : r_{k|k}^i > \Gamma_d, l^* = \arg \max_l \beta_{k|k}^i(l) \right\}$ , which reports the starting times and the means with most likely duration of the Bernoulli components whose probability of existence is greater than  $\Gamma_d$ . Finally, the pseudocodes of the TPMB filters is given in Algorithm 1.

---

#### Algorithm 1 Gaussian TPMB filters pseudocode

---

```

- Set  $\lambda_{0|0}(\cdot) = 0, n_{0|0} = 0$ .
for  $k = 1$  to final time step do
  - Prediction:
    o For alive trajectories: use Lemma 6.
    o For all trajectories, use Lemma 8.
    o Apply  $L$ -scan approximation (46) to all covariance matrices.
  - Update:
    o For alive trajectories, use Lemma 7.
    o For all trajectories, use Lemma 9.
  - Use Proposition 7 to obtain a PMB:
    o For alive trajectories, use (37)-(40), see Section V-A.
    o For all trajectories, use (37)-(38) and (43)-(45), see Sec. V-B
  - Estimate the current set of trajectories, see Section V-D.
  - Perform pruning, see Section V-C.
end for

```

---

#### E. Discussion

We proceed to discuss several aspects of the proposed algorithms. The TPMB filter for alive trajectories has a similar recursion to the track-oriented PMB filter for sets of targets in [10], with the difference that past trajectory states are not integrated out. This allows the TPMB filter to estimate sets of trajectories from first principles. In particular, for the case  $L = 1$ , the track-oriented PMB filter and the TPMB filter perform the same filtering computations, though the TPMB stores the past means, and possibly the covariances, for each Bernoulli component. It should also be noted that, in this paper, we have derived the PMB approximation from direct KLD minimisation on the densities with auxiliary variables.

Instead, the derivation in [10] makes use of KLD minimisation on the data association variables, which are not explicit in the posterior. The TPMB filter for all trajectories requires the propagation of more variables, as single trajectory densities have more parameters (41), and the recursions differ.

We have presented the TPMB filters for Poisson birth, as it is generally more suitable than multi-Bernoulli birth [12], as non-detected targets are represented efficiently via the intensity of a Poisson RFS. Nevertheless, the above derivation is also valid for multi-Bernoulli birth. In this case, we just need to set  $\lambda_{k'|k}(\cdot) = 0$  and add the Bernoulli components of new born targets in the prediction step [11], [12]. The resulting algorithm would correspond to the trajectory multi-Bernoulli filter, which can include target labels in the state without modifying the recursion, as the trajectory MBM filter [20], [23].

We have presented the Gaussian implementation of the TPMB filter due to its simplicity and performance. Nevertheless, it is also possible to use Gaussian mixtures and particles to represent the single-trajectory densities in the Bernoulli components.

Another relevant algorithm is the labelled multi-Bernoulli (LMB) filter [19]. As the  $\delta$ -GLMB filter, the LMB filter does not work well, unless practical modifications are used, if there is more than one birth Bernoulli component with the same mean and covariance matrix, which enables more than one target to be born in a certain region [20, Sec. II.B]. In addition, in each update, the LMB filter (also the version in [39]) makes use of the  $\delta$ -GLMB update, which requires an exponential growth in the number of global hypotheses due to the MBM<sub>01</sub> parameterisation [11, Sec. IV]. The TPMB avoids these drawbacks by creating Bernoulli components directly from the measurements, and performing the update directly in PMBM form, without using MBM<sub>01</sub> parameterisation. In addition, the posterior of the TPMB filter has full trajectory information of each Bernoulli component.

## VI. SIMULATIONS

We analyse the performance of the two TPMB filters in comparison with the trajectory filters: the TPMBM filter [21], the trajectory global nearest neighbour PMB (TGNPMB) filter, the TPHD filter and the TCPHD filter [24]. The TGNPMB filter corresponds to the TPMBM filter but only propagating the global hypothesis with highest weight, as the global nearest neighbor approach [1]. The TPMB filters have been implemented with the following parameters: maximum number of hypotheses  $N_h = 200$ , threshold for pruning the PPP weights  $\Gamma_p = 10^{-5}$ , threshold for pruning Bernoulli components  $\Gamma_b = 10^{-5}$  and  $L = 5$ , and  $\Gamma_d = 0.5$ . For the set of alive trajectories, we use the same parameters as before and also parameter  $\Gamma_a = 10^{-4}$ . The TPMBM filters have been implemented with the same parameters and also with a threshold  $10^{-4}$  for pruning global hypothesis weights and Estimator 1 in [11] with threshold 0.4. The TPHD and TCPHD filters use a pruning threshold  $10^{-8}$ , absorption threshold 4, and limit the number of PHD components to 30 [24].

The previous trajectory filters are structured to exploit smoothing, though for  $L = 1$  no trajectory smoothing is performed. We have also considered three MTT algorithms that




Figure 2: True target trajectories of the four considered targets. The blue one and the red one are born at time step 1, whereas the green one and the black one are born at time step 21. The only target that dies during the simulation is the blue target, which dies at time step 40, when all targets are in close proximity. Targets positions every 10 time steps are marked with a circle, and their initial positions with a filled circle.

do not exploit smoothing and estimate the set of trajectories sequentially by linking the previous estimated set of trajectories with the newly estimated set of targets. The first one is the PMBM filter [10], [11] where the trajectory estimates are formed by linking target state estimates that originate from the same first detection (same Bernoulli component). The PMBM filter implementation parameters are the same as in the TPMBM filter. The  $\delta$ -GLMB filter [40] considers joint prediction and update using Murty’s algorithm, as in [41], with 1000 global hypotheses and the standard estimator that first computes the maximum a posterior of the cardinality [17]. The LMB filter [19] also considers 1000 global hypotheses in the update, maximum number of Gaussians per Bernoulli is 10, pruning threshold  $10^{-5}$  for each Gaussian and merging threshold for each Gaussian 4. The code for  $\delta$ -GLMB and LMB codes was obtained from <http://ba-tuong.vo-au.com>.

We consider a target state  $x = [p_x, \dot{p}_x, p_y, \dot{p}_y]^T$ , which contains position and velocity in a two-dimensional plane. All the units in this section are given in the international system. We use the nearly-constant velocity model with

$$F = I_2 \otimes \begin{pmatrix} 1 & \tau \\ 0 & 1 \end{pmatrix}, \quad Q = qI_2 \otimes \begin{pmatrix} \tau^3/3 & \tau^2/2 \\ \tau^2/2 & \tau \end{pmatrix}$$

where  $\tau = 1$  and  $q = 0.01$ . We also consider  $p_S = 0.99$ . The sensor measures target positions with parameters

$$H = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad R = \sigma^2 I_2,$$

where  $\sigma^2 = 1$ , and  $p_D = 0.9$ . The clutter intensity is  $\lambda^C(z) = \bar{\lambda}^C \cdot u_A(z)$  where  $u_A(z)$  is a uniform density in region  $A = [0, 300] \times [0, 300]$  and  $\bar{\lambda}^C = 10$  is the average number of clutter measurements per scan. The intensity of the birth process is Gaussian with  $\bar{x}_k^{b,1} = [100, 0, 100, 0]^T$  and  $P_k^{b,1} = \text{diag}([150^2, 1, 150^2, 1])$ , with weight  $w_1^{b,1} = 3$  and  $w_k^{b,1} = 0.005$  for  $k > 1$ .

The  $\delta$ -GLMB filter requires a (labelled) multi-Bernoulli birth model, not Poisson. For  $k > 1$ , we use one Bernoulli component with probability of existence 0.005, mean  $\bar{x}_k^{b,1}$  and




Figure 3: Trajectory metric error against time for the alive trajectories. Error increases at time step 40, when a target dies. On the whole, the TPMBM filter is the best performing filter followed by the TPMB filter.

covariance  $P_k^{b,1}$ . For  $k = 1$ , the expected number of targets is 3 according to the Poisson process, so we consider five Bernoulli components with the same probability of existence and spatial locations such that the support of the multi-Bernoulli birth covers the expected number of targets.

We consider the ground truth trajectories in Figure 2 and  $N_s = 81$  time steps, as in [12]. We assess filter performance using Monte Carlo simulation with 100 runs. At each time step  $k$ , we measure the error between the true set  $\mathbf{X}^k$  of trajectories and its estimate  $\hat{\mathbf{X}}^k$ , which differ depending on the problem formulation, see Section II. The error is determined by the linear programming metric  $d(\cdot, \cdot)$  for sets of trajectories in [42] with parameters  $p = 2$ ,  $c = 10$  and  $\gamma = 1$ . In our results, we only use the position elements to compute  $d(\cdot, \cdot)$  and normalise the error by the considered time window such that the squared error at time  $k$  becomes  $d^2(\mathbf{X}^k, \hat{\mathbf{X}}^k)/k$ . The root mean square (RMS) error at a given time step is

$$d(k) = \sqrt{\frac{1}{N_{mc}k} \sum_{i=1}^{N_{mc}} d^2(\mathbf{X}^k, \hat{\mathbf{X}}_i^k)}, \quad (47)$$

where  $\hat{\mathbf{X}}_i^k$  is the estimate of the set of trajectories at time  $k$  in the  $i$ th Monte Carlo run.

We first proceed to analyse the error in estimating the set of alive trajectories. The RMS trajectory error against time is shown in Figure 3. For all filters, estimation error increases after time step 40, in which a target dies, and the rest of the targets are in close proximity. The TPMBM filter is the most accurate filter to estimate the alive trajectories. This is to be expected, as without approximations, the TPMBM filtering recursion provides the true posterior over the set of trajectories. The second best performing filter in general is the proposed TPMB filter. Its performance is slightly worse than TGNPMB after time step 40 but it is considerably better before time step 40. TPHPD and TPHPD filters perform considerably worse, which is to be expected as they are less accurate approximations. The LMB filters performs worse than




Figure 4: Decomposition of the trajectory metric error against time for the alive trajectories.

$\delta$ -GLMB.  $\delta$ -GLMB is less accurate than PMBM, which is outperformed by the TPMBM and TPMB filter.

The squared trajectory metric  $d^2(\cdot, \cdot)$  can be decomposed into the square costs for missed targets, false targets, localisation error of properly detected targets, and track switches [42]. The resulting RMS errors for the decomposed costs are shown in Figure 4. Before time step 40, all filters except LMB, TPHD and TCPHD have a negligible cost for false targets. Also,  $\delta$ -GLMB and PMBM show a higher error for missed targets, followed by TPHD, TCPHD and TGNPMBM. After time step 40, these filters increase their error mainly due to false target errors, as one target is removed from the scene. Track switching cost are small and quite similar for all filters based on sets of trajectories.  $\delta$ -GLMB and LMB are the only filters with track switches before targets get in close proximity and provides the highest switching costs. PMBM filter has the third highest switching costs after time step 40. We would also like to remark that the errors obtained by using the square sum of the generalised optimal sub-pattern assignment (GOSPA) metric ( $\alpha = 2$ ) [43] at each time step instead of  $d^2(\cdot, \cdot)$  in (47) are quite similar to the trajectory metric errors, as, due to the choice of  $\gamma$ , the switching costs are small compared to the other costs, especially for the filters based on sets of trajectories.

The average execution times in seconds of a single run of our Matlab implementations with a 3.5 GHz Intel Xeon E5 processor are: 2.4 (TPMB), 7.0 (TPMBM), 0.7 (TGNPMB), 1.1 (TPHD), 1.1 (TCPHD), 5.8 (PMBM), 13.1 ( $\delta$ -GLMB) and 10.5 (LMB). The fastest algorithm is TGNPMB, though its performance is worse than TPMB. The TPMBM is slower than the other trajectory filters and PMBM but TPMBM is also the one with highest performance, as expected. There is a trade-off between computational complexity and accuracy in the selection of TPMBM, TPMB and TGNPMB.

We proceed to analyse the performances of the filters for different values of  $L$  and different  $p_D$  and  $\bar{\lambda}^C$ . In Table I, we show the resulting RMS error considering all time steps,

which is calculated by

$$d_T = \sqrt{1/N_s \cdot \sum_{k=1}^{N_s} d^2(k)}. \quad (48)$$

We can see that increasing  $L$  for the trajectory filters lowers the error, mainly due to improved localisation of past states. By looking at these errors, in general, we can see that the best performing filter is TPMBM, followed by TPMB and TGNPMB. As expected, if the clutter intensity increases, performance decreases for all filters. The higher the probability of detection, performance increases for all filters.

Finally, we consider the estimation of the set of all trajectories at each time step, using the same scenario. TPHD and TCPHD filters are not included as they are not really suitable for this problem [24]. The RMS errors (48) are shown in Table II. As before, error decreases by increasing  $L$ , and TPMBM is generally the best algorithm followed by TPMB and TGNPMB. PMBM performs worse than these filters, but better than  $\delta$ -GLMB and LMB.

The average execution time in seconds for  $p_D = 0.9$  and  $\bar{\lambda}^C = 10$  and the estimation of the set of all trajectories are: 3.3 (TPMB), 7.9 (TPMBM), 0.9 (TGNPMB), 5.8 (PMBM), 13.1 ( $\delta$ -GLMB) and 10.5 (LMB). Compared to tracking alive trajectories, there is an increase in the execution time in the trajectory filters, as these versions of the filters perform more computations. The sequential track estimators, PMBM,  $\delta$ -GLMB and LMB have the same computational burden to solve both problems, as they only differ in what estimated set of trajectories is reported.

## VII. CONCLUSIONS

In this paper, we have proposed two Poisson multi-Bernoulli filters for sets of trajectories to perform multiple target tracking. One TPMB filter contains information on the set of alive trajectories and the other contains information on the set of all trajectories, which include alive and dead trajectories. We have also proposed Gaussian implementations of the filters for linear/Gaussian dynamic measurement models and constant probability of detection and survival.

The resulting filters offer a trade-off between computational complexity and accuracy. They are faster than TPMBM filters but with a decrease in performance. On the other hand, the TPMB filters are considerably more accurate than TPHD and TCPHD filters, but with higher computational complexity.

## REFERENCES

- [1] S. Blackman and R. Popoli, *Design and Analysis of Modern Tracking Systems*. Artech House, 1999.
- [2] S. Challa, M. R. Morelande, D. Musicki, and R. J. Evans, *Fundamentals of Object Tracking*. Cambridge University Press, 2011.
- [3] T. Chen, R. Wang, B. Dai, D. Liu, and J. Song, "Likelihood-field-model-based dynamic vehicle detection and tracking for self-driving," *IEEE Transactions on Intelligent Transportation Systems*, vol. 17, no. 11, pp. 3142–3158, Nov. 2016.
- [4] C. Hurter, N. H. Riche, S. M. Drucker, M. Cordeil, R. Alligier, and R. Vuillemot, "FiberClay: Sculpting three dimensional trajectories to reveal structural insights," *IEEE Transactions on Visualization and Computer Graphics*, vol. 25, no. 1, pp. 704–714, Jan. 2019.

Table I: Trajectory metric errors (alive trajectories)

$L$	TPMB			TPMBM			TGNPMB			TPHD			TCPHD			PMBM	GLMB	LMB
	1	5	10	1	5	10	1	5	10	1	5	10	1	5	10			
No change	5.10	4.91	4.90	4.80	4.60	4.59	5.23	5.04	5.03	7.87	7.69	7.69	7.50	7.30	7.30	5.74	7.62	8.64
$p_D = 0.99$	4.91	4.74	4.73	4.53	4.34	4.34	4.66	4.47	4.46	7.17	7.06	7.06	7.07	6.96	6.96	5.45	7.38	7.48
$p_D = 0.80$	5.36	5.15	5.13	5.22	5.02	5.01	5.82	5.64	5.63	8.74	8.52	8.52	8.04	7.80	7.79	6.27	7.52	9.48
$\bar{\lambda}^C = 20$	5.18	4.99	4.98	4.90	4.70	4.70	6.35	6.21	6.20	8.04	7.87	7.86	7.55	7.36	7.36	5.91	8.19	9.12
$\bar{\lambda}^C = 30$	5.20	5.01	5.00	4.89	4.70	4.69	7.50	7.39	7.39	8.11	7.95	7.95	7.62	7.44	7.44	5.97	8.51	9.44

Table II: Trajectory metric errors (all trajectories)

$L$	TPMB			TPMBM			TGNPMB			PMBM	GLMB	LMB
	1	5	10	1	5	10	1	5	10			
No change	2.96	2.54	2.51	2.94	2.54	2.52	4.22	3.95	3.94	4.58	7.05	8.79
$p_D = 0.99$	2.51	2.08	2.06	2.49	2.08	2.06	3.11	2.78	2.77	4.17	6.68	6.74
$p_D = 0.80$	3.46	3.05	3.02	3.42	3.03	3.02	5.06	4.83	4.82	5.14	7.52	9.63
$\bar{\lambda}^C = 20$	3.09	2.68	2.65	3.03	2.65	2.63	6.18	6.02	6.01	4.81	7.75	9.07
$\bar{\lambda}^C = 30$	3.16	2.77	2.75	3.12	2.76	2.74	7.63	7.53	7.52	5.02	8.20	9.52

- [5] F. Meyer, T. Kropfreiter, J. L. Williams, R. Lau, F. Hlawatsch, P. Braca, and M. Z. Win, "Message passing algorithms for scalable multitarget tracking," *Proceedings of the IEEE*, vol. 106, no. 2, pp. 221–259, Feb. 2018.
- [6] D. Reid, "An algorithm for tracking multiple targets," *IEEE Transactions on Automatic Control*, vol. 24, no. 6, pp. 843–854, Dec. 1979.
- [7] E. Brekke and M. Chitre, "Relationship between finite set statistics and the multiple hypothesis tracker," *IEEE Transactions on Aerospace and Electronic Systems*, vol. 54, no. 4, pp. 1902–1917, Aug. 2018.
- [8] Y. Bar-Shalom and E. Tse, "Tracking in a cluttered environment with probabilistic data association," *Automatica*, vol. 11, no. 5, pp. 451–460, 1975.
- [9] R. P. S. Mahler, *Advances in Statistical Multisource-Multitarget Information Fusion*. Artech House, 2014.
- [10] J. L. Williams, "Marginal multi-Bernoulli filters: RFS derivation of MHT, JIPDA and association-based MeMBer," *IEEE Transactions on Aerospace and Electronic Systems*, vol. 51, no. 3, pp. 1664–1687, July 2015.
- [11] A. F. García-Fernández, J. L. Williams, K. Granström, and L. Svensson, "Poisson multi-Bernoulli mixture filter: direct derivation and implementation," *IEEE Transactions on Aerospace and Electronic Systems*, vol. 54, no. 4, pp. 1883–1901, Aug. 2018.
- [12] A. F. García-Fernández, Y. Xia, K. Granström, L. Svensson, and J. L. Williams, "Gaussian implementation of the multi-Bernoulli mixture filter," in *Proceedings on the 22nd International Conference on Information Fusion*, 2019.
- [13] J. L. Williams, "An efficient, variational approximation of the best fitting multi-Bernoulli filter," *IEEE Transactions on Signal Processing*, vol. 63, no. 1, pp. 258–273, Jan. 2015.
- [14] Y. Xia, K. Granström, L. Svensson, and A. F. García-Fernández, "Performance evaluation of multi-Bernoulli conjugate priors for multi-target filtering," in *20th International Conference on Information Fusion*, July 2017, pp. 1–8.
- [15] D. Musicki and R. Evans, "Joint integrated probabilistic data association: JIPDA," *IEEE Transactions on Aerospace and Electronic Systems*, vol. 40, no. 3, pp. 1093–1099, July 2004.
- [16] A. F. García-Fernández, J. Grajal, and M. R. Morelande, "Two-layer particle filter for multiple target detection and tracking," *IEEE Transactions on Aerospace and Electronic Systems*, vol. 49, no. 3, pp. 1569–1588, July 2013.
- [17] B. T. Vo and B. N. Vo, "Labeled random finite sets and multi-object conjugate priors," *IEEE Transactions on Signal Processing*, vol. 61, no. 13, pp. 3460–3475, July 2013.
- [18] E. H. Aoki, P. K. Mandal, L. Svensson, Y. Boers, and A. Bagchi, "Labeling uncertainty in multitarget tracking," *IEEE Transactions on Aerospace and Electronic Systems*, vol. 52, no. 3, pp. 1006–1020, June 2016.
- [19] S. Reuter, B.-T. Vo, B.-N. Vo, and K. Dietmayer, "The labeled multi-Bernoulli filter," *IEEE Transactions on Signal Processing*, vol. 62, no. 12, pp. 3246–3260, June 2014.
- [20] A. F. García-Fernández, L. Svensson, and M. R. Morelande, "Multiple target tracking based on sets of trajectories," *accepted for publication in IEEE Transactions on Aerospace and Electronic Systems*, 2019. [Online]. Available: <https://arxiv.org/abs/1605.08163>
- [21] K. Granström, L. Svensson, Y. Xia, J. L. Williams, and A. F. García-Fernández, "Poisson multi-Bernoulli mixture trackers: continuity through random finite sets of trajectories," in *21st International Conference on Information Fusion*, 2018, pp. 973–981.
- [22] K. Granström, L. Svensson, Y. Xia, J. Williams, and A. F. García-Fernández, "Poisson multi-Bernoulli mixtures for sets of trajectories," 2019. [Online]. Available: <https://arxiv.org/abs/1912.08718>
- [23] Y. Xia, K. Granström, L. Svensson, A. F. García-Fernández, and J. L. Williams, "Multi-scan implementation of the trajectory Poisson multi-Bernoulli mixture filter," *Journal of Advances in Information Fusion. Special Issue on Multiple Hypothesis Tracking*, vol. 14, no. 2, pp. 213–235, Dec. 2019.
- [24] A. F. García-Fernández and L. Svensson, "Trajectory PHD and CPHD filters," *IEEE Transactions on Signal Processing*, vol. 67, no. 22, pp. 5702–5714, Nov 2019.
- [25] M. K. Pitt and N. Shephard, "Filtering via simulation: Auxiliary particle filters," *Journal of the American Statistical Association*, vol. 94, no. 446, pp. 590–599, Jun. 1999.
- [26] V. Elvira, L. Martino, D. Luengo, and M. F. Bugallo, "Generalized multiple importance sampling," *Statistical Science*, vol. 34, no. 1, pp. 129–155, 2019.
- [27] A. F. García-Fernández, "A track-before-detect labeled multi-Bernoulli particle filter with label switching," *IEEE Transactions on Aerospace and Electronic Systems*, vol. 52, no. 5, pp. 2123–2138, Oct. 2016.
- [28] L. Úbeda-Medina, A. F. García-Fernández, and J. Grajal, "Adaptive auxiliary particle filter for track-before-detect with multiple targets," *IEEE Transactions on Aerospace and Electronic Systems*, vol. 53, no. 5, pp. 2317–2330, Oct. 2017.
- [29] M. R. Morelande, C. M. Kreucher, and K. Kastella, "A Bayesian approach to multiple target detection and tracking," *IEEE Transactions on Signal Processing*, vol. 55, no. 5, pp. 1589–1604, May. 2007.
- [30] W. Yi, M. R. Morelande, L. Kong, and J. Yang, "A computationally efficient particle filter for multitarget tracking using an independence approximation," *IEEE Transactions on Signal Processing*, vol. 61, no. 4, pp. 843–856, Feb. 2013.
- [31] M. Úney, B. Mulgrew, and D. E. Clark, "Latent parameter estimation in fusion networks using separable likelihoods," *IEEE Transactions on Signal and Information Processing over Networks*, vol. 4, no. 4, pp. 752–768, Dec 2018.
- [32] C. M. Bishop, *Pattern Recognition and Machine Learning*. Springer, 2006.
- [33] J. Houssineau and D. E. Clark, "Multitarget filtering with linearized complexity," *IEEE Transactions on Signal Processing*, vol. 66, no. 18, pp. 4957–4970, Sept. 2018.
- [34] K. G. Murty, "An algorithm for ranking all the assignments in order of increasing cost," *Operations Research*, vol. 16, no. 3, pp. 682–687, 1968.
- [35] D. F. Crouse, "On implementing 2D rectangular assignment algorithms," *IEEE Transactions on Aerospace and Electronic Systems*, vol. 52, no. 4, pp. 1679–1696, August 2016.
- [36] J. Williams and R. Lau, "Approximate evaluation of marginal association probabilities with belief propagation," *IEEE Transactions on Aerospace and Electronic Systems*, vol. 50, no. 4, pp. 2942–2959, Oct. 2014.

- [37] P. Horridge and S. Maskell, "Real-time tracking of hundreds of targets with efficient exact JPDAF implementation," in *9th International Conference on Information Fusion*, 2006.
- [38] J. L. Williams, "Hybrid Poisson and multi-Bernoulli filters," in *15th International Conference on Information Fusion*, 2012, pp. 1103–1110.
- [39] J. Olofsson, C. Veibäck, and G. Hendeby, "Sea ice tracking with a spatially indexed labeled multi-Bernoulli filter," in *20th International Conference on Information Fusion*, July 2017.
- [40] B. N. Vo, B. T. Vo, and H. G. Hoang, "An efficient implementation of the generalized labeled multi-Bernoulli filter," *IEEE Transactions on Signal Processing*, vol. 65, no. 8, pp. 1975–1987, April 2017.
- [41] J. Correa, M. Adams, and C. Perez, "A Dirac delta mixture-based random finite set filter," in *International Conference on Control, Automation and Information Sciences*, Oct. 2015, pp. 231–238.
- [42] A. S. Rahmathullah, A. F. García-Fernández, and L. Svensson, "A metric on the space of finite sets of trajectories for evaluation of multi-target tracking algorithms," 2016. [Online]. Available: <http://arxiv.org/abs/1605.01177>
- [43] —, "Generalized optimal sub-pattern assignment metric," in *20th International Conference on Information Fusion*, 2017, pp. 1–8.
- [44] T. M. Cover and J. A. Thomas, *Elements of Information Theory*. John Wiley & Sons, 2006.

# Trajectory Poisson multi-Bernoulli filters: Supplementary material

## APPENDIX A

In this appendix, we prove that if we integrate the auxiliary variables in  $\tilde{f}_{k'|k}(\cdot)$ , see (6), we recover  $f_{k'|k}(\cdot)$ . That is, we prove that

$$\sum_{u_{1:n} \in \mathbb{U}_{k'}^n} \tilde{f}_{k'|k}(\{(u_1, X_1), \dots, (u_n, X_n)\}) = f_{k'|k}(\{X_1, \dots, X_n\}). \quad (49)$$

We first obtain two preliminary results with only a Poisson component and a multi-Bernoulli mixture. Then, we proceed to prove the PMBM case.

### A. PPP

Integrating out the auxiliary variables in  $\tilde{f}_{k'|k}^p(\cdot)$ , we obtain

$$\begin{aligned} & \sum_{u_{1:n} \in \mathbb{U}_{k'|k}^n} \tilde{f}_{k'|k}^p(\{(u_1, X_1), \dots, (u_n, X_n)\}) \\ &= e^{-\int \lambda_{k'|k}(X) dX} \left[ \prod_{i=1}^n \lambda_{k'|k}(X_i) \right] \left[ \sum_{u_{1:n} \in \mathbb{U}_{k'|k}^n} \prod_{i=1}^n \delta_0[u_i] \right] \\ &= f_{k'|k}^p(\{X_1, \dots, X_n\}). \end{aligned} \quad (50)$$

### B. Multi-Bernoulli mixture

We first note that if  $\lambda_{k'|k} = 0$ , then, the PMBM (5) is an MBM. Using [9, Eq. (4.127)], for existence probabilities smaller than one, the MBM with auxiliary variables can be written as

$$\begin{aligned} & \tilde{f}_{k'|k}^{mbm}(\{(u_1, X_1), \dots, (u_n, X_n)\}) \\ &= \sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a \left[ \prod_{i=1}^{n_{k'|k}} (1 - r_{k'|k}^{i, a^i}) \right] \\ & \quad \sum_{1 \leq i_1 \neq \dots \neq i_n \leq n_{k'|k}} \prod_{p=1}^n \frac{r_{k'|k}^{i_p, a^i}}{1 - r_{k'|k}^{i_p, a^i}} p_{k'|k}^{i_p, a^i}(X_p) \delta_{i_p}[u_p] \end{aligned} \quad (51)$$

If we integrate out the auxiliary variables in (51), we obtain the MBM without auxiliary variables

$$\begin{aligned} & \sum_{u_{1:n} \in \mathbb{U}_{k'|k}^n} \tilde{f}_{k'|k}^{mbm}(\{(u_1, X_1), \dots, (u_n, X_n)\}) \\ &= \sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a \left[ \prod_{i=1}^{n_{k'|k}} (1 - r_{k'|k}^{i, a^i}) \right] \sum_{1 \leq i_1 \neq \dots \neq i_n \leq n_{k'|k}} \\ & \quad \prod_{p=1}^n \frac{r_{k'|k}^{i_p, a^i}}{1 - r_{k'|k}^{i_p, a^i}} p_{k'|k}^{i_p, a^i}(X_p) \left[ \sum_{u_{1:n} \in \mathbb{U}_{k'|k}^n} \delta_{i_p}[u_p] \right] \\ &= f_{k'|k}^{mbm}(\{X_1, \dots, X_n\}). \end{aligned}$$

If some existence probabilities are equal to one, the derivation is analogous, but removing the corresponding  $(1 - r_{k'|k}^{i, a^i})$  in the numerator and denominator in (51).

### C. PMBM

We can write the PMBM (5) as

$$\begin{aligned} f_{k'|k}(\{X_1, \dots, X_n\}) &= \sum_{\mathbf{Y} \subseteq \{X_1, \dots, X_n\}} f_{k'|k}^p(\mathbf{Y}) f_{k'|k}^{mbm}(\mathbf{X}_{k'} \setminus \mathbf{Y}) \\ &= \sum_{j=0}^n \sum_{\sigma \in \Gamma_{n,j}} f_{k'|k}^p(\{X_{\sigma_1}, \dots, X_{\sigma_j}\}) \\ & \quad \times f_{k'|k}^{mbm}(\{X_{\sigma_{j+1}}, \dots, X_{\sigma_n}\}) \end{aligned}$$

where  $\Gamma_{n,j}$  is the set that contains all possible sets  $\sigma = \{\sigma_1, \dots, \sigma_j\}$  of  $j$  elements from  $\{1, \dots, n\}$  and  $\{\sigma_{j+1}, \dots, \sigma_n\} = \{1, \dots, n\} \setminus \{\sigma_1, \dots, \sigma_j\}$ . The cardinality of set  $\Gamma_{n,j}$  is

$$|\Gamma_{n,j}| = \binom{n}{j}.$$

Using these formulas on the PMBM density with auxiliary variables, we have

$$\begin{aligned} & \tilde{f}_{k'|k}(\{(u_1, X_1), \dots, (u_n, X_n)\}) \\ &= \sum_{j=0}^n \sum_{\sigma \in \Gamma_{n,j}} \tilde{f}_{k'|k}^p(\{(u_{\sigma_1}, X_{\sigma_1}), \dots, (u_{\sigma_j}, X_{\sigma_j})\}) \\ & \quad \times \tilde{f}_{k'|k}^{mbm}(\{(u_{\sigma_{j+1}}, X_{\sigma_{j+1}}), \dots, (u_{\sigma_n}, X_{\sigma_n})\}) \end{aligned}$$

Then, integrating out the auxiliary variables

$$\begin{aligned} & \sum_{u_{1:n} \in \mathbb{U}_{k'|k}^n} \tilde{f}_{k'|k}(\{(u_1, X_1), \dots, (u_n, X_n)\}) \\ &= \sum_{u_{1:n} \in \mathbb{U}_{k'|k}^n} \sum_{j=0}^n \sum_{\sigma \in \Gamma_{n,j}} \tilde{f}_{k'|k}^p(\{(u_{\sigma_1}, X_{\sigma_1}), \dots, (u_{\sigma_j}, X_{\sigma_j})\}) \\ & \quad \times \tilde{f}_{k'|k}^{mbm}(\{(u_{\sigma_{j+1}}, X_{\sigma_{j+1}}), \dots, (u_{\sigma_n}, X_{\sigma_n})\}) \\ &= \sum_{j=0}^n \sum_{\sigma \in \Gamma_{n,j}} \left[ \sum_{u_{\sigma_{1:j}} \in \mathbb{N}_0^j} \tilde{f}_{k'|k}^p(\{(u_{\sigma_1}, X_{\sigma_1}), \dots, (u_{\sigma_j}, X_{\sigma_j})\}) \right] \\ & \quad \times \left[ \sum_{u_{\sigma_{j+1:n}} \in \mathbb{N}_0^{n-j}} \tilde{f}_{k'|k}^{mbm}(\{(u_{\sigma_{j+1}}, X_{\sigma_{j+1}}), \dots, (u_{\sigma_n}, X_{\sigma_n})\}) \right] \end{aligned}$$

Applying the results in the previous two subsections, we finish the proof of (49).

## APPENDIX B

In Section B-A, we prove Proposition 2. We also prove in Section B-B that the resulting density from the KLD minimisation also matches the PHD. Finally, we prove Lemma 3 in Section B-C.

### A. KLD minimisation

The augmented single trajectory space  $\mathbb{U}_{k'|k} \times T_{(k')}$  can be written as the union of disjoint spaces  $\mathbb{U}_{k'|k} \times T_{(k')} = \uplus_{u=0}^{n_{k'|k}} \{u\} \times T_{(k')}$ . Therefore, given a finite set  $\tilde{\mathbf{X}}_{k'} \subset \mathbb{U}_{k'|k} \times T_{(k')}$ , we can write  $\tilde{\mathbf{X}}_{k'} = \tilde{\mathbf{Y}}_{k'} \uplus \tilde{\mathbf{X}}_{k'}^1 \uplus \dots \uplus \tilde{\mathbf{X}}_{k'}^{n_{k'|k}}$ , where

$\tilde{\mathbf{Y}}_{k'} \subset \{0\} \times T_{(k')}$  and  $\tilde{\mathbf{X}}_{k'}^i \subset \{i\} \times T_{(k')}$ , to obtain [9, Eq. (3.53)]

$$\begin{aligned}
& D(\tilde{f} \parallel \tilde{q}) \\
&= \int \tilde{f}(\tilde{\mathbf{X}}_{k'}) \log \frac{\tilde{f}(\tilde{\mathbf{X}}_{k'})}{\tilde{q}(\tilde{\mathbf{X}}_{k'})} \delta \tilde{\mathbf{X}}_{k'} \\
&= \int \int \dots \int \tilde{f}(\tilde{\mathbf{Y}}_{k'} \uplus \tilde{\mathbf{X}}_{k'}^1 \uplus \dots \uplus \tilde{\mathbf{X}}_{k'}^{n_{k'}|k}) \\
&\quad \times \log \frac{\tilde{f}(\tilde{\mathbf{Y}}_{k'} \uplus \tilde{\mathbf{X}}_{k'}^1 \uplus \dots \uplus \tilde{\mathbf{X}}_{k'}^{n_{k'}|k})}{\tilde{q}(\tilde{\mathbf{Y}}_{k'} \uplus \tilde{\mathbf{X}}_{k'}^1 \uplus \dots \uplus \tilde{\mathbf{X}}_{k'}^{n_{k'}|k})} \delta \tilde{\mathbf{Y}}_{k'} \delta \tilde{\mathbf{X}}_{k'}^1 \dots \delta \tilde{\mathbf{X}}_{k'}^{n_{k'}|k} \\
&= c + \int \tilde{f}_{k'|k}^p(\tilde{\mathbf{Y}}_{k'}) \log \frac{\tilde{f}_{k'|k}^p(\tilde{\mathbf{Y}}_{k'})}{\tilde{q}^p(\tilde{\mathbf{Y}}_{k'})} \delta \tilde{\mathbf{Y}}_{k'} \\
&\quad - \sum_{i=1}^{n_{k'}|k} \int \sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a \tilde{f}_{k'|k}^{i,a^i}(\tilde{\mathbf{X}}_{k'}^i) \log \tilde{q}^{i,1}(\tilde{\mathbf{X}}_{k'}^i) \delta \tilde{\mathbf{X}}_{k'}^i
\end{aligned}$$

where  $c$  is a constant that does not depend on  $\tilde{q}(\cdot)$ . Maximising with respect to  $\tilde{q}^p(\cdot)$ ,  $\tilde{q}^{1,1}(\cdot)$ ,  $\dots$ ,  $\tilde{q}^{n_{k'}|k,1}(\cdot)$ , we get

$$\begin{aligned}
\tilde{q}^p(\tilde{\mathbf{Y}}_{k'}) &= \tilde{f}_{k'|k}^p(\tilde{\mathbf{Y}}_{k'}) \\
\tilde{q}^{i,1}(\tilde{\mathbf{X}}_{k'}^i) &= \sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a \tilde{f}_{k'|k}^{i,a^i}(\tilde{\mathbf{X}}_{k'}^i)
\end{aligned}$$

Using (8) and identifying terms w.r.t. (12), we have that the existence probability  $r^i$  and single-trajectory density  $p^i(\cdot)$  of  $\tilde{q}^{i,1}(\cdot)$  are those given in Proposition 2, which finishes the proof.

### B. Matching the PHD

In this section, we show that, for KLD minimisation in the previous section, it holds that the PHD of  $\tilde{f}(\cdot)$  is the same as the PHD of  $\tilde{q}(\cdot)$ . In Sections B-B1 and B-B2, we calculate the PHD of  $\tilde{f}(\cdot)$  and  $\tilde{q}(\cdot)$ , respectively.

1) *PHD of  $\tilde{f}(\cdot)$* : The PHD  $D_{\tilde{f}}(\cdot)$  of density  $\tilde{f}(\cdot)$  for sets of trajectories, which is given by (7), is [9], [24]

$$\begin{aligned}
D_{\tilde{f}}(\tilde{X}) &= \int \tilde{f}(\{\tilde{X}\} \cup \tilde{\mathbf{X}}_{k'}) \delta \tilde{\mathbf{X}}_{k'} \\
&= \int \tilde{f}(\{\tilde{X}\} \cup \tilde{\mathbf{Y}}_{k'} \uplus \tilde{\mathbf{X}}_{k'}^1 \uplus \dots \uplus \tilde{\mathbf{X}}_{k'}^{n_{k'}|k}) \\
&\quad \delta \tilde{\mathbf{Y}}_{k'} \delta \tilde{\mathbf{X}}_{k'}^1 \dots \delta \tilde{\mathbf{X}}_{k'}^{n_{k'}|k}
\end{aligned} \tag{52}$$

where we have applied the decomposition of the set integral into disjoint spaces, see Section B-A.

If  $\tilde{X} = (0, X) \in \{0\} \times T_{(k')}$ , then the PHD is

$$\begin{aligned}
D_{\tilde{f}}(\tilde{X}) &= \int \tilde{f}_{k'|k}^p(\{\tilde{X}\} \cup \tilde{\mathbf{Y}}_{k'}) \sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a \\
&\quad \times \prod_{i=1}^{n_{k'}|k} [\tilde{f}_{k'|k}^{i,a^i}(\tilde{\mathbf{X}}_{k'}^i)] \delta \tilde{\mathbf{Y}}_{k'} \delta \tilde{\mathbf{X}}_{k'}^1 \dots \delta \tilde{\mathbf{X}}_{k'}^{n_{k'}|k} \\
&= \int \tilde{f}_{k'|k}^p(\{\tilde{X}\} \cup \tilde{\mathbf{Y}}_{k'}) \delta \tilde{\mathbf{Y}}_{k'} \\
&= \lambda_{k'|k}(X)
\end{aligned} \tag{53}$$

where the last line follows directly as it corresponds to the PHD of the PPP  $\tilde{f}_{k'|k}^p(\cdot)$ .

If  $\tilde{X} = (u, X) \in \{u\} \times T_{(k')}$ ,  $u \in \{1, \dots, n_{k'}|k\}$ , then the PHD is

$$\begin{aligned}
D_{\tilde{f}}(\tilde{X}) &= \int \tilde{f}_{k'|k}^p(\tilde{\mathbf{Y}}_{k'}) \sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a \tilde{f}_{k'|k}^{u,a^u}(\{\tilde{X}\} \cup \tilde{\mathbf{X}}_{k'}^u) \\
&\quad \times \prod_{i=1, i \neq u}^{n_{k'}|k} [\tilde{f}_{k'|k}^{i,a^i}(\tilde{\mathbf{X}}_{k'}^i)] \delta \tilde{\mathbf{Y}}_{k'} \delta \tilde{\mathbf{X}}_{k'}^1 \dots \delta \tilde{\mathbf{X}}_{k'}^{n_{k'}|k} \\
&= \sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a \int \tilde{f}_{k'|k}^{u,a^u}(\{\tilde{X}\} \cup \tilde{\mathbf{X}}_{k'}^u) \delta \tilde{\mathbf{X}}_{k'}^u \\
&= \sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a r_{k'|k}^{u,a^u} p_{k'|k}^{u,a^u}(X)
\end{aligned} \tag{54}$$

2) *PHD of  $\tilde{q}(\cdot)$* : We calculate the PHD of  $\tilde{q}(\cdot)$ , which is given by (11) and Proposition 2, using (52). If  $\tilde{X} = (0, X) \in \{0\} \times T_{(k')}$ , then the PHD is

$$\begin{aligned}
D_{\tilde{q}}(\tilde{X}) &= \int \int \dots \int \tilde{q}^p(\{\tilde{X}\} \cup \tilde{\mathbf{Y}}_{k'}) \\
&\quad \times \prod_{i=1}^{n_{k'}|k} [\tilde{q}^{i,1}(\tilde{\mathbf{X}}_{k'}^i)] \delta \tilde{\mathbf{Y}}_{k'} \delta \tilde{\mathbf{X}}_{k'}^1 \dots \delta \tilde{\mathbf{X}}_{k'}^{n_{k'}|k} \\
&= \int \tilde{q}^p(\{\tilde{X}\} \cup \tilde{\mathbf{Y}}_{k'}) \delta \tilde{\mathbf{Y}}_{k'} \\
&= \lambda_{k'|k}(X).
\end{aligned} \tag{55}$$

If  $\tilde{X} = (u, X) \in \{u\} \times T_{(k')}$ ,  $u \in \{1, \dots, n_{k'}|k\}$ , then the PHD is

$$\begin{aligned}
D_{\tilde{q}}(\tilde{X}) &= \int \int \dots \int \tilde{q}^p(\tilde{\mathbf{Y}}_{k'}) \tilde{q}^{u,1}(\{\tilde{X}\} \cup \tilde{\mathbf{X}}_{k'}^u) \\
&\quad \times \prod_{i=1, i \neq u}^{n_{k'}|k} [\tilde{q}^{i,1}(\tilde{\mathbf{X}}_{k'}^i)] \delta \tilde{\mathbf{Y}}_{k'} \delta \tilde{\mathbf{X}}_{k'}^1 \dots \delta \tilde{\mathbf{X}}_{k'}^{n_{k'}|k} \\
&= \int \tilde{q}^{u,1}(\{\tilde{X}\} \cup \tilde{\mathbf{X}}_{k'}^u) \delta \tilde{\mathbf{X}}_{k'}^u \\
&= \sum_{a \in \mathcal{A}_{k'|k}} w_{k'|k}^a r_{k'|k}^{u,a^i} p_{k'|k}^{u,a^i}(X).
\end{aligned} \tag{56}$$

We can see that the PHD of  $\tilde{q}(\cdot)$ , which is given by (55) and (56), coincides with the PHD of  $\tilde{f}(\cdot)$ , which is given by (53) and (54).

### C. KLD bound

In this section, we prove Lemma 3. We have

$$\begin{aligned}
D(f_{k'|k} \parallel q) &= \sum_{n=0}^{\infty} \frac{1}{n!} \int f_{k'|k}(\{X_1, \dots, X_n\}) \\
&\quad \times \log \frac{f_{k'|k}(\{X_1, \dots, X_n\})}{q(\{X_1, \dots, X_n\})} dX_{1:n} \\
&= \sum_{n=0}^{\infty} \frac{1}{n!} \int \sum_{u_{1:n}} \tilde{f}_{k'|k}(\{(u_1, X_1), \dots, (u_n, X_n)\}) \\
&\quad \times \log \frac{\sum_{u_{1:n}} \tilde{f}_{k'|k}(\{(u_1, X_1), \dots, (u_n, X_n)\})}{\sum_{u_{1:n}} \tilde{q}(\{(u_1, X_1), \dots, (u_n, X_n)\})} dX_{1:n}
\end{aligned}$$

where we have used (49). Applying the log sum inequality [44] inside the integral, we obtain

$$\begin{aligned}
D(f_{k'|k} \| q) &\leq \sum_{n=0}^{\infty} \frac{1}{n!} \int \sum_{u_{1:n}} \tilde{f}_{k'|k}(\{(u_1, X_1), \dots, (u_n, X_n)\}) \\
&\quad \times \log \frac{\tilde{f}_{k'|k}(\{(u_1, X_1), \dots, (u_n, X_n)\})}{\tilde{q}(\{(u_1, X_1), \dots, (u_n, X_n)\})} dX_{1:n} \\
&= D(\tilde{f}_{k'|k} \| \tilde{q}).
\end{aligned}$$

This completes the proof of Lemma 3.