

MULTIPLICATIVE VECTOR FIELDS ON BUNDLE GERBES

DEREK KREPSKI AND JENNIFER VAUGHAN

ABSTRACT. This paper makes use of multiplicative vector fields on Lie groupoids to model infinitesimal symmetries of S^1 -bundle gerbes. It is shown that a connective structure on a bundle gerbe gives rise to a natural horizontal lift of multiplicative vector fields to the bundle gerbe, and that the 3-curvature presents the obstruction to the horizontal lift being a morphism of Lie 2-algebras. Connection-preserving multiplicative vector fields on a bundle gerbe with connective structure are shown to inherit a natural Lie 2-algebra structure; moreover, this Lie 2-algebra is quasi-isomorphic to the Poisson-Lie 2-algebra of the 2-plectic base manifold (M, χ) , where χ is the 3-curvature of the connective structure. As an application of this result, we give an analogue of a formula of Kostant in the 2-plectic and quasi-Hamiltonian context.

1. INTRODUCTION

Let M be a smooth manifold. A well-known Theorem of Weil [38] shows that the set of isomorphism classes of principal S^1 -bundles over M are in one-to-one correspondence with $H^2(M; \mathbb{Z})$. The analogous objects corresponding to cohomology classes in $H^3(M; \mathbb{Z})$ are known as S^1 -gerbes, introduced in differential geometry by Brylinski [6] using the formalism of stacks (presheaves of groupoids), following Giraud [15]. Some later models for S^1 -gerbes include S^1 -central extensions of Lie groupoids (e.g., see Behrend-Xu [4]), S^1 -bundle gerbes (equivalent to S^1 -central extensions of submersion groupoids), due to Murray [27] (see also Hitchin [17] and Chatterjee [8]), Dixmier-Douady bundles [12], differential cocycles [23], and principal Lie 2-group bundles [3, 31, 39]. The present paper primarily adopts the bundle gerbe perspective (see Section 3.1 for a review of bundle gerbes) to establish some results for S^1 -gerbes that are analogous to well-known results for their degree-2 counterparts, principal S^1 -bundles.

For example, consider a principal S^1 -bundle $P \rightarrow M$ with connection $\gamma \in \Omega^1(P)$ and let $\text{curv}(\gamma) \in \Omega^2(M)$ denote the curvature of the connection. It is well known that the connection determines a horizontal lift of vector fields on M to vector fields on P ,

$$\text{Lift}_\gamma : \mathfrak{X}(M) \rightarrow \mathfrak{X}(P),$$

Date: March 31, 2020.

The authors acknowledge the support of the Natural Sciences and Engineering Research Council of Canada (RGPIN-2015-05833), and the Pacific Institute of Mathematical Sciences.

which is a Lie algebra homomorphism if and only if $\text{curv}(\gamma) = 0$. What is the analogue for S^1 -gerbes?

To arrive at the analogue, we employ *multiplicative vector fields* on Lie groupoids. Recall from [24] that a multiplicative vector field on a Lie groupoid $\mathbf{G} = \{G_1 \rightrightarrows G_0\}$ is a Lie groupoid morphism (i.e., a functor) $\xi : \mathbf{G} \rightarrow T\mathbf{G} = \{TG_1 \rightrightarrows TG_0\}$ such that $\pi_{\mathbf{G}} \circ \xi = \text{id}_{\mathbf{G}}$. Such a functor can be defined by a multiplicative vector field is a pair of vector fields ξ_0 and ξ_1 on G_0 and G_1 , respectively, compatible with the Lie groupoid structure (so that the pair defines such a functor). The multiplicative vector fields on \mathbf{G} are the objects of a category $\mathbb{X}(\mathbf{G})$, whose morphisms are natural transformations $\eta : \xi \Rightarrow \xi'$ satisfying $\pi_{\mathbf{G}}(\eta(x)) = \epsilon(x)$ for every object $x \in G_0$, where $\epsilon : G_0 \rightarrow G_1$ denotes the unit map. By a Theorem of Hepworth [16, Theorem 4.15], multiplicative vector fields on a Lie groupoid \mathbf{G} correspond to vector fields on the associated differentiable stack $B\mathbf{G}$.

An S^1 -gerbe on a smooth manifold M is a differentiable stack, which may be presented as a bundle gerbe over M , or equivalently an S^1 -central extension $P \rightrightarrows X$ of the submersion Lie groupoid $X \times_M X \rightrightarrows X$ associated to a surjective submersion $X \rightarrow M$ (see Section 3.2). Therefore, vector fields on the S^1 -gerbe may be modelled by multiplicative vector fields on the Lie groupoid $P \rightrightarrows X$. In fact, as we show in Theorem 3.22, the category $\mathbb{X}(P \rightrightarrows X)$ of multiplicative vector fields on $P \rightrightarrows X$ is equivalent to the category described by Collier [10] of *infinitesimal symmetries* of the corresponding S^1 -gerbe, viewed as a stack.

Additionally, by a recent result of Berwick-Evans and Lerman [5], multiplicative vector fields on a Lie groupoid naturally have a structure of a Lie 2-algebra. In this work, a Lie 2-algebra is a 2-term L_∞ -algebra (e.g., see Baez-Crans [2])—that is, a 2-term chain complex $L_1 \rightarrow L_0$, equipped with a graded bracket and a chain homotopy coherently measuring the failure of the bracket to satisfy the Jacobi identity. For morphisms, we work with a localization of the bicategory of 2-term L_∞ -algebras with respect to quasi-isomorphisms—namely, Noohi’s bicategory with *butterflies* as 1-morphisms (see Section 4.1 for a review of the bicategory of Lie 2-algebras).

We thus find a natural setting in which to formulate the bundle gerbe analogue of the aforementioned result on horizontal lifts. Indeed, suppose an S^1 -gerbe over M is presented by a bundle gerbe $P \rightrightarrows X$, an S^1 -central extension of $X \times_M X \rightrightarrows X$, where $X \rightarrow M$ is a surjective submersion. The Lie 2-algebra of multiplicative vector fields $\mathbb{X}(X \times_M X \rightrightarrows X)$ is naturally quasi-isomorphic to $\mathfrak{X}(M)$, and we show in Theorem 4.6 that a connective structure on $P \rightrightarrows X$ (i.e., a multiplicative connection $\gamma \in \Omega^1(P)$ and curving $B \in \Omega^2(X)$) gives rise to a map

$$\text{Lift}_\gamma : \mathbb{X}(X \times_M X \rightrightarrows X) \rightarrow \mathbb{X}(P \rightrightarrows X),$$

which is a morphism of Lie 2-algebras if and only if the associated 3-curvature $3\text{-curv}(B, \gamma) \in \Omega^3(M)$ of the connective structure vanishes.

The Lie 2-algebra of *connection-preserving* multiplicative vector fields on a bundle gerbe plays a natural role in establishing some other results that may be viewed as 2-plectic and quasi-Hamiltonian analogues of classical results in symplectic geometry. A multiplicative vector field on a bundle gerbe $P \rightrightarrows X$, given by a pair of vector fields $\tilde{\xi}$ and ξ on X and P , respectively, is said to preserve a connective structure (B, γ) if

$$L_{\tilde{\xi}}B = d\alpha, \quad \text{and} \quad L_{\xi} = \delta\alpha, \quad \text{for some } \alpha \in \Omega^1(X),$$

where $\delta = \Sigma(-1)^j \delta_j^*$ is the simplicial differential for $P \rightrightarrows X$. We show that the collection $\mathbb{X}(P; B, \gamma)$ of connection-preserving multiplicative vector fields is a subcategory $\mathbb{X}(P \rightrightarrows X)$ (see Corollary 3.18 for the precise statement), and it naturally inherits a Lie 2-algebra structure (see Proposition 4.8).

Recall that for a 2-plectic manifold (M, χ) , the analogue of the Poisson algebra of smooth functions on a symplectic manifold is the Poisson-Lie 2-algebra $\mathbb{L}(M, \chi)$ (see Definition 4.5, due to Rogers [34]). Consider a bundle gerbe $P \rightrightarrows X$ over M , equipped with a connective structure (B, γ) as above, with $3\text{-curv}(B, \gamma) = \chi$ (i.e., a *prequantization* of (M, χ)). In Theorem 5.1, we show that $\mathbb{L}(M, \chi)$ is quasi-isomorphic to the Lie 2-algebra $\mathbb{X}(P \rightrightarrows X; B, \gamma)$ of *connection-preserving* multiplicative vector fields on $P \rightrightarrows X$. In other words, the Poisson Lie 2-algebra $\mathbb{L}(M, \chi)$ of a 2-plectic manifold is quasi-isomorphic to the Lie 2-algebra of infinitesimal symmetries of its prequantization, sometimes called *infinitesimal quantomorphisms* in the literature—e.g., see [13] wherein the authors prove a similar result (*cf.* Remark 5.2.)

As an application of Theorem 5.1, we propose 2-plectic and quasi-Hamiltonian analogues of a formula of Kostant. Given a Hamiltonian Lie algebra action $\mathfrak{g} \rightarrow \mathfrak{X}(M, \omega)$ on a symplectic manifold, Kostant gives a lift to a \mathfrak{g} -action on a prequantization of (M, ω) (a principal S^1 -bundle $P \rightarrow M$ with connection whose curvature equals ω) by infinitesimal quantomorphisms. In Section 5, we show that a Hamiltonian \mathfrak{g} -action on a 2-plectic manifold (M, χ) (i.e., one admitting a *homotopy moment map* as in [7]) naturally lifts to an infinitesimal \mathfrak{g} -action on a prequantization of (M, χ) . For quasi-Hamiltonian spaces with Lie group-valued moment maps, a similar lift is obtained.

This paper is organized as follows. In Section 2, we review some preliminaries to establish notation used throughout the paper. In Section 3, we recall some foundational aspects of bundle gerbes and begin our study of multiplicative vector fields on bundle gerbes. We discuss their basic properties, and provide a sketch of the equivalence between the category of multiplicative vector fields on a bundle gerbe and the category of infinitesimal symmetries of the underlying gerbe, and similarly for multiplicative vector fields and infinitesimal symmetries that preserve a connective structure. In Section 4, we review the 2-category of Lie 2-algebras used in this work, and establish Theorem 4.6 characterizing the 3-curvature as the obstruction to the

horizontal lift defining a morphism of Lie 2-algebras. We also introduce (Proposition 4.8) the Lie 2-algebra of connection-preserving multiplicative vector fields on a bundle gerbe. In Section 5, we establish some results in 2-plectic geometry and quasi-Hamiltonian geometry that are analogous to some classical results in symplectic geometry. In particular, we show (Theorem 5.1) that the Lie 2-algebra of multiplicative vector fields preserving the connective structure on a bundle gerbe is quasi-isomorphic to the Poisson-Lie-2-algebra of the 2-plectic manifold (M, χ) , where χ denotes the 3-curvature of the connective structure. We end the paper with a quasi-Hamiltonian analogue (Theorem 5.7) of a formula in symplectic geometry due to Kostant.

2. PRELIMINARIES AND NOTATION

Throughout this paper, all manifolds and maps between manifolds are assumed to be smooth. We assume the reader has some familiarity with Lie groupoids, though we recall some aspects next to establish notation. Recall that a Lie groupoid $G_1 \rightrightarrows G_0$ gives rise to a simplicial manifold G_\bullet : for $k \geq 2$, write

$$G_k = \underbrace{G_1 \times_{G_0} G_1 \times_{G_0} \cdots \times_{G_0} G_1}_{k \text{ factors}}$$

whose elements are k -tuples (g_1, \dots, g_k) of composable arrows (with $s(g_i) = t(g_{i+1})$). For $0 \leq i \leq k$, let $\partial_i : G_k \rightarrow G_{k-1}$ be the *face maps* given by

$$\partial_i(g_1, \dots, g_k) = \begin{cases} (g_2, \dots, g_k) & \text{if } i = 0 \\ (g_1, \dots, g_i g_{i+1}, \dots, g_k) & \text{if } 0 < i < k \\ (g_1, \dots, g_{k-1}) & \text{if } i = k. \end{cases}$$

For convenience, we set $\partial_0 = s$ and $\partial_1 = t$ on G_1 . The face maps satisfy the simplicial identities

$$\partial_i \partial_j = \partial_{j-1} \partial_i \quad \text{for } i < j, \quad (2.1)$$

and hence $\partial = \sum (-1)^i \partial_i^*$ defines a differential for the complexes $\Omega^k(G_\bullet)$, $k \geq 0$.

In particular, for a surjective submersion, $\pi : X \rightarrow M$, we may consider the *submersion groupoid* $X \times_M X \rightrightarrows X$, and obtain the simplicial manifold $X^{[\bullet+1]}$, where

$$X^{[n]} = \{(x_1, \dots, x_n) \mid \pi(x_1) = \cdots = \pi(x_n)\}.$$

As shown in [27], the following complexes are exact:

$$0 \longrightarrow \Omega^k(M) \xrightarrow{\pi^*} \Omega^k(X) \xrightarrow{\partial} \Omega^k(X^{[2]}) \xrightarrow{\partial} \cdots \quad (2.2)$$

Similarly, for proper Lie groupoids $G_1 \rightrightarrows G_0$, the sequence

$$\Omega^0(G_0) \xrightarrow{\partial} \Omega^0(G_1) \xrightarrow{\partial} \Omega^0(G_2) \xrightarrow{\partial} \cdots$$

is exact [11].

For a compact Lie group K acting on manifolds X and M , and a K -equivariant surjective submersion $\pi : X \rightarrow M$, there are natural K -equivariant analogues of the above. Recall briefly that for a K -manifold M , the Cartan model of K -equivariant differential forms is given by $\Omega_K(M) = (S(\mathfrak{g}^*) \otimes \Omega(M))^K$, the algebra of K -equivariant polynomials $\alpha : \mathfrak{g} \rightarrow \Omega(M)$, with differential $(d_K\alpha)(\xi) = d\alpha(\xi) - \iota_{\xi_M}\alpha(\xi)$. Here, $\xi_M \in \mathfrak{X}(M)$ denotes the generating vector field for $\xi \in \mathfrak{g}$: $\xi_M(m) = \frac{d}{dt}|_{t=0} \exp(-t\xi) \cdot m$. The map ∂ extends to K -equivariant forms, and the K -equivariant version of (2.2) is also exact—see [36].

Finally, for a principal S^1 -bundle $P \rightarrow M$, we identify $\text{Lie}(S^1) \cong \mathbb{R}$ and write $\frac{\partial}{\partial \theta}$ for the generating vector field of the principal S^1 -action.

3. S^1 -GERBES OVER MANIFOLDS

This section employs multiplicative vector fields on Lie groupoids as infinitesimal symmetries of S^1 -bundle gerbes, and introduces their connection-preserving counterparts. After a brief review of S^1 -bundle gerbes, we consider some basic properties of the category of multiplicative vector fields on bundle gerbes, showing that the connection-preserving multiplicative vector fields form a full subcategory (Corollary 3.18). In Section 3.5, we sketch an equivalence of categories between (connection-preserving) multiplicative vector fields on bundle gerbes (with connective structure) and the category of (connection-preserving) infinitesimal symmetries of S^1 -banded gerbes [9].

3.1. Bundle gerbes with connective structure. We review some elementary aspects of S^1 -bundle gerbes over a manifold, following [30]. For a detailed introduction to bundle gerbes, we refer the reader to Murray's original paper [27] as well as [29] and [35].

Definition 3.1. Let M be a manifold. An S^1 -bundle gerbe (or simply, *bundle gerbe*) $\mathcal{G} = (X, P, \mu)$ over M consists of a surjective submersion $\pi : X \rightarrow M$, a principal S^1 -bundle $P \rightarrow X^{[2]}$, and a bundle isomorphism $\mu : \partial_2^*P \otimes \partial_0^*P \rightarrow \partial_1^*P$ over $X^{[3]}$, called the gerbe product, satisfying the associativity condition $\partial_1^*\mu \circ (\partial_3^*\mu \otimes \text{id}) = \partial_2^*\mu \circ (\text{id} \otimes \partial_0^*\mu)$ over $X^{[4]}$.

Definition 3.2. Let $\mathcal{G} = (X, P, \mu)$ be a bundle gerbe over a manifold M . A *connection* on \mathcal{G} is a connection 1-form $\gamma \in \Omega^1(P)$ such that the bundle isomorphism μ is connection-preserving: $\mu^*(\hat{\partial}_1^*\gamma) = \hat{\partial}_2^*\gamma \otimes \hat{\partial}_0^*\gamma$, where $\hat{\partial}_j : \partial_j^*P \rightarrow P$ denotes the natural map covering ∂_j . A *curving* for the connection γ is a 2-form $B \in \Omega^2(X)$ satisfying $\partial B = \text{curv}(\gamma)$. We will call the pair (B, γ) a *connective structure* for \mathcal{G} .

Since $\partial dB = d\partial B = d\text{curv}(\gamma) = 0$, there exists a unique closed form $\chi \in \Omega^3(M)$, called the *3-curvature* of the connective structure, satisfying $\pi^*\chi = dB$.

Example 3.3. The trivial bundle gerbe \mathcal{I} over M is the bundle gerbe with $\pi = \text{id}_M$, $P = M \times S^1$, and canonical gerbe product μ . For $\omega \in \Omega^2(M)$, \mathcal{I}_ω is the bundle gerbe \mathcal{I} with trivial connection on $M \times S^1 \rightarrow M$ and curving ω . Note that the 3-curvature of this trivial connection with curving ω is $d\omega$. \diamond

Example 3.4. Let $\{U_i\}$ denote a good open cover of a smooth manifold M , and let $X = \bigsqcup U_i \rightarrow M$ denote the map whose restrictions $U_i \hookrightarrow M$ are the inclusions. Then we may identify $X^{[2]} = \bigsqcup U_{ij}$, where U_{ij} denotes the intersection $U_i \cap U_j$ (and similarly for $X^{[n]}$ and n -tuples of indices, e.g., $X^{[3]} = \bigsqcup U_{ijk}$). Suppose $\{g_{ijk} : U_{ijk} \rightarrow S^1\}$ denotes a collection of smooth functions. Let $P = \bigsqcup U_{ij} \times S^1 \rightarrow \bigsqcup U_{ij}$ and set

$$\mu(x, z, w) = (x, zwg_{ijk}(x)),$$

where we are identifying $x \in M$ with its copies in the U_{ij} 's. It's straightforward to see that μ defines a bundle gerbe product as in Definition 3.1, provided g_{ijk} defines a Čech 2-cocycle. Denote the resulting bundle gerbe by $\mathcal{G}_{g_{ijk}}$.

A connective structure on such a gerbe is thus specified by a collection $\{\gamma_{ij} = \text{pr}^*A_{ij} + z^{-1}dz\}$, or more simply by the forms $A_{ij} \in \Omega^1(U_{ij})$ that must satisfy $A_{jk} - A_{ik} + A_{ij} = g_{ijk}^{-1}dg_{ijk}$ on each U_{ijk} , together with a collection $B_i \in \Omega^2(U_i)$ satisfying $B_j - B_i = dA_{ij}$ on each U_{ij} . In other words, a bundle gerbe with connective structure with respect to the covering $\{U_i\}$ is specified by a Čech-Deligne 2-cocycle (g, A, B) [6]. \diamond

Recall that a *common refinement* of two surjective submersions $\pi : X \rightarrow M$ and $\pi' : X' \rightarrow M$ is a surjective submersion $\varpi : Y \rightarrow M$ together with surjective submersions $\lambda : Y \rightarrow X$ and $\rho : Y \rightarrow X'$ such that $\pi \circ \lambda = \varpi = \pi' \circ \rho$. For example, the fibre product $Y = X \times_M X'$ is such a common refinement.

Definition 3.5. [30] Let $\mathcal{G} = (X, P, \mu)$ and $\mathcal{G}' = (X', P', \mu')$ be bundle gerbes over a manifold M . An *isomorphism of bundle gerbes* $\mathcal{G} \dashrightarrow \mathcal{G}'$ consists of a common refinement $Y \rightarrow M$ of π and π' , together with a principal S^1 -bundle $Q \rightarrow Y$ and an isomorphism of S^1 -bundles $\varphi : \rho^*P' \otimes \partial_0^*Q \rightarrow \partial_1^*Q \otimes \lambda^*P$ over $Y^{[2]}$ satisfying the coherence condition

$$(\text{id} \otimes \lambda^*\mu) \circ (\partial_2^*\varphi \otimes \text{id}) \circ (\text{id} \otimes \partial_0^*\varphi) = \partial_1^*\varphi \circ (\rho^*\mu' \otimes \text{id})$$

over $Y^{[3]}$. We will denote such an isomorphism by (Q, φ) , or sometimes simply Q .

An *isomorphism of bundle gerbes with connective structure* $(\mathcal{G}, B, \gamma) \dashrightarrow (\mathcal{G}', B', \gamma')$ is an isomorphism (Q, φ) as above, together with a connection 1-form $\gamma_Q \in \Omega^1(Q)$ such that $\text{curv}(\gamma_Q) = \rho^*B' - \lambda^*B$ and φ is connection-preserving. We will denote such an isomorphism by (Q, φ, γ_Q) , or sometimes simply (Q, γ_Q) .

Example 3.6. In the literature, a trivialization of a bundle gerbe (\mathcal{G}, B, γ) with connective structure is sometimes presented as follows (e.g., see [29] or [25]). Let

(Q, γ_Q) be a principal S^1 -bundle with connection over X together with a connection-preserving isomorphism $\partial_1^* Q^* \otimes \partial_0^* Q \rightarrow P$ satisfying

$$B - \text{curv}(\gamma_Q) = \pi^* \omega,$$

for some $\omega \in \Omega^2(M)$, sometimes called the *error 2-form*. It is straightforward to repackage this data as an isomorphism $(Q, \gamma_Q) : \mathcal{I}_\omega \dashrightarrow (\mathcal{G}, B, \gamma)$. Note that such a trivialization requires that the 3-curvature of (\mathcal{G}, B, γ) equals $d\omega$. \diamond

Bundle gerbes on a manifold M are classified by their *Dixmier-Douady* class (or DD-class) in $H^3(M; \mathbb{Z})$, analogous to the Chern-Weil classification of principal S^1 -bundles by their Chern class in $H^2(M; \mathbb{Z})$. Moreover, the DD-class maps to the cohomology class of the 3-curvature via the coefficient homomorphism $H^3(M; \mathbb{Z}) \rightarrow H^3(M; \mathbb{R}) \cong H_{dR}^3(M)$. For bundle gerbes on M with connective structure, the classification takes values in the group of differential characters of degree 3 (e.g., see [23]).

Let G be a compact Lie group. For manifolds equipped with a smooth G -action, there are various notions of equivariant bundle gerbes in the literature—see, for instance, [25], [36], [28], [30]. Moreover, there are G -equivariant analogues of the aforementioned classifications: G -equivariant bundle gerbes are classified by their G -equivariant DD-class, while G -equivariant bundle gerbes with connective structure are classified by G -equivariant differential characters.

Example 3.7. An important family of examples of bundle gerbes naturally occur on Lie groups. Let G be a compact simple Lie group, with an invariant inner product $\langle -, - \rangle$ on its Lie algebra \mathfrak{g} . Let θ^L and θ^R denote the left and right invariant Maurer-Cartan forms. Let $\eta = \frac{1}{12} \langle \theta^L, [\theta^L, \theta^L] \rangle$ denote the Cartan 3-form, and recall that η has an equivariant extension $\eta_G(\xi) = \eta + \frac{1}{2} \langle \theta^L + \theta^R, \xi \rangle \in \Omega_G^3(G)$, with respect to the conjugation action. For certain choices of inner product (parameterized by the so-called *level*), there are a number of explicit constructions in the literature of a bundle gerbe with connective structure (\mathcal{G}, B, γ) over G whose 3-curvature is η , as well as G -equivariant versions—e.g., see [25], [14] and [21]. \diamond

3.2. Bundle gerbes as S^1 -central extensions. Bundle gerbes may be recast as S^1 -central extensions of Lie groupoids. This interpretation is briefly reviewed next. Given a bundle gerbe $\mathcal{G} = (X, P, \mu)$, we consider the Lie groupoid $P \rightrightarrows X$, with source and target maps, $s = \pi_0 := \partial_0 \circ \pi_P$ and $t =: \pi_1 = \partial_1 \circ \pi_P$, respectively, where $\pi_P : P \rightarrow X^{[2]}$ denotes the bundle projection. The groupoid multiplication $m : P \times_X P \rightarrow P$ (also denoted by \cdot) is induced by the gerbe product,

$$\mu(x, y, z; p \otimes q) = (x, y, z; m(p, q)),$$

as are the unit $\epsilon : X \rightarrow P$ and inverse $i : P \rightarrow P$, by setting

$$\mu^{-1}(x, x, y; p) = (x, x, y; \epsilon(x) \otimes p), \quad \text{and} \quad \mu^{-1}(x, y, x; \epsilon(x)) = (x, y, x; p \otimes i(p))$$

for all $p \in P$ lying over $(x, y) \in X^{[2]}$. In this viewpoint, we see that $P \rightrightarrows X$ is an S^1 -central extension of the submersion groupoid $X^{[2]} \rightrightarrows X$. That is, the bundle projection $P \rightarrow X^{[2]}$ is a morphism of Lie groupoids,

$$\begin{array}{ccc} P & \longrightarrow & X^{[2]} \\ \downarrow & & \downarrow \\ X & \xlongequal{\quad} & X \end{array}$$

and the S^1 -action on P is compatible with the groupoid multiplication:

$$(zp) \cdot (wq) = (zw)(p \cdot q)$$

for all composable $p, q \in P$ and $z, w \in S^1$.

Suppose (B, γ) defines a connective structure on \mathcal{G} . Then $\pi_P^* \partial B = d\gamma$, where recall ∂ denotes the simplicial differential on $X^{[\bullet+1]}$. Letting δ denote the simplicial differential on P_\bullet , the simplicial manifold associated to $P \rightrightarrows X$, we have $\delta B = d\gamma$. Also, since the bundle isomorphism μ is connection-preserving, we see that $\delta\gamma = 0$. In other words, γ is *multiplicative*: $m^* \gamma = \text{pr}_1^* \gamma + \text{pr}_2^* \gamma$.

Remark 3.8. (Notation) Throughout the paper, we shall use ∂ and δ to distinguish between the simplicial maps of $X^{[\bullet+1]}$ and P_\bullet , respectively, as in the preceding paragraph.

3.3. Multiplicative vector fields on bundle gerbes. Recall from [24] that a *multiplicative vector field* on a Lie groupoid $\mathbf{G} = \{G_1 \rightrightarrows G_0\}$ is a functor $\xi : \mathbf{G} \rightarrow T\mathbf{G}$ such that $\pi_{\mathbf{G}} \circ \xi = \text{id}_{\mathbf{G}}$. Such a functor ξ is therefore given by a pair of vector fields $(\xi_0, \xi_1) \in \mathfrak{X}(G_0) \times \mathfrak{X}(G_1)$ that must be compatible with units and the groupoid multiplications on \mathbf{G} and $T\mathbf{G}$. The collection of multiplicative vector fields on \mathbf{G} are the objects of a category $\mathbb{X}(\mathbf{G})$, whose morphisms are natural transformations $\eta : \xi \Rightarrow \xi'$ satisfying $\pi_{\mathbf{G}}(\eta(x)) = \epsilon(x)$ for every object $x \in G_0$, where $\epsilon : G_0 \rightarrow G_1$ denotes the unit map [16].

Let us now specialize to the Lie groupoid $\mathbf{P} = \{P \rightrightarrows X\}$ associated to a bundle gerbe (X, P, μ) . Consider a multiplicative vector field $(\tilde{\xi}, \check{\xi})$ on \mathbf{P} . Since $(\tilde{\xi}, \check{\xi})$ defines a functor, we have that $\check{\xi} \sim_{\pi_j} \tilde{\xi}$ for $j = 0, 1$, and hence $d\pi(\tilde{\xi}(x)) = d\pi(\check{\xi}(y))$ for all $(x, y) \in X^{[2]}$. Hence $\tilde{\xi}$ descends to a vector field ξ on M .

Moreover, $(\tilde{\xi}, \check{\xi})$ defines a vector field on $X^{[2]}$, and similarly on $X^{[n]}$ for any $n > 1$; denote such vector fields by $\tilde{\xi}^{[n]}$. In particular, we have that $\check{\xi} \sim_{\pi_P} \tilde{\xi}^{[2]}$. Similarly, there is a naturally defined vector field $(\check{\xi}, \tilde{\xi})$ on $P \times_X P$ (denoted by $\check{\xi} * \tilde{\xi}$ in [24]) such that $(\check{\xi}, \tilde{\xi}) \sim_m \check{\xi}$ and hence $(\check{\xi}, \tilde{\xi}) \sim_{\delta_j} \check{\xi}$ for $j = 0, 1, 2$.

Proposition 3.9. *Let $\mathcal{G} = (X, P, \mu)$ be a bundle gerbe, and let $(\tilde{\xi}, \check{\xi})$ be a multiplicative vector field on the corresponding Lie groupoid $P \rightrightarrows X$.*

- (1) *The vector field $\check{\xi}$ is S^1 -invariant.*

- (2) $\check{\xi} \sim_{\pi_j} \check{\xi}$ for $j = 0, 1$.
 (3) $\check{\xi}_2 \otimes \check{\xi}_0 \sim_{\mu} \check{\xi}_1$, where $\check{\xi}_j$ denotes the vector field on $\partial_j^* P$ induced by $(\check{\xi}^{[3]}, \check{\xi})$.

Moreover, any pair of vector fields $(\check{\xi}, \check{\xi}) \in \mathfrak{X}(X) \times \mathfrak{X}(P)$ satisfying the above conditions is a multiplicative vector field on $P \rightrightarrows X$.

Proof. To show (1), we verify that $[\frac{\partial}{\partial\theta}, \check{\xi}] = 0$. Choose a connection γ as in Definition 3.2. Since $[\frac{\partial}{\partial\theta}, \check{\xi}]$ is vertical, it suffices to show $\iota_{[\frac{\partial}{\partial\theta}, \check{\xi}]} \gamma = 0$. Note that since $\delta\gamma = 0$, it follows that $\delta\iota_{\check{\xi}}\gamma = 0$ and since \mathbf{P} is proper there exists $g : X \rightarrow \mathbb{R}$ such that $\delta g = \iota_{\check{\xi}}\gamma$, which is S^1 -invariant. Hence $\iota_{[\frac{\partial}{\partial\theta}, \check{\xi}]} \gamma = L_{\frac{\partial}{\partial\theta}} \iota_{\check{\xi}}\gamma - \iota_{\check{\xi}} L_{\frac{\partial}{\partial\theta}} \gamma = L_{\frac{\partial}{\partial\theta}} \delta g = 0$, as required.

Statement (2) has already been established, and (3) is a straightforward consequence of (1) and the discussion preceding the proposition. To verify the last statement, it is straightforward to see that conditions (2) and (3) show that $(\check{\xi}, \check{\xi})$ defines a functor, provided it respects the unit maps. It thus suffices to check that $d\epsilon(\check{\xi})(x) = \check{\xi}(\epsilon(x))$, which follows immediately from (2) and the observation that ϵ is a section of π_j (for $j = 0$ or 1). \square

Example 3.10. Consider a multiplicative vector field $(\check{\xi}, \check{\xi})$ on a bundle gerbe P given in terms of Čech data, as in Example 3.4. In this case, we have that $\check{\xi}$ is specified by a collection of local vector fields ξ_i on U_i and by the paragraph preceding Proposition 3.9, $\xi_i|_{U_{ij}} = \xi_j|_{U_{ij}}$ for all i, j and thus the local vector fields are restrictions of a vector field ξ on M . Denote the restriction of ξ to $U_{i_1, i_2, \dots}$ by $\xi_{i_1, i_2, \dots}$.

The vector field $\check{\xi}$ must take the form $(\xi_{ij}(x), f_{ij}(x, z))$, where $f_{ij}(x, z) \in \mathbb{R}$ is the component in the S^1 direction (where we implicitly trivialize the tangent bundle of S^1). By Proposition 3.9 (1), the functions f_{ij} are independent of $z \in S^1$; hence we abuse notation and write $f_{ij} : U_{ij} \rightarrow \mathbb{R}$. The composition

$$\begin{array}{ccccc} & & (\xi_{ij}, f_{ij}(x)) & & (\xi_{jk}, f_{jk}(x)) & & \\ & & \curvearrowright & & \curvearrowleft & & \\ \xi_i(x) & & & \xi_j(x) & & & \xi_k(x) \end{array}$$

in the tangent groupoid $TP \rightrightarrows TX$ is $(\xi_{jk}, f_{ik}(x))$, as $\check{\xi}$ is multiplicative. By definition, this composition is obtained by applying the tangent map of the gerbe multiplication. Therefore,

$$f_{ik} = g_{ijk}^{-1} dg_{ijk}(\xi) + f_{ij} + f_{jk}, \quad (3.1)$$

on U_{ijk} .

In [9], Collier studies the infinitesimal symmetries of an S^1 -gerbe considered as a stack over M . In this context, the author defines a category $\mathcal{L}_{g_{ijk}}$ of *infinitesimal symmetries* of the gerbe with Čech cocycle g_{ijk} . The objects of this category are

given by pairs $(\xi, \{-f_{ij}\})$ consisting of a vector field ξ on M and a collection of functions f_{ij} satisfying (3.1). \diamond

In [5], the authors show that the category $\mathbb{X}(\mathbf{G})$ of multiplicative vector fields on a Lie groupoid \mathbf{G} is a category internal to the category of vector spaces (i.e., a 2-vector space in the language of [2]). In fact, they show it is a Lie 2-algebra—we shall return to this in Section 4. Though we are mainly interested in the case of Lie groupoids $P \rightrightarrows X$ arising from bundle gerbes, there is no added difficulty in describing the morphisms in the general case, as we recall next, following [5].

For a general Lie groupoid \mathbf{G} , morphisms of multiplicative vector fields are given by sections of the Lie algebroid $A \rightarrow G_0$ of \mathbf{G} . (We shall implicitly make use of the equivalence of categories between 2-term chain complexes and 2-vector spaces—see [5, Remark 2.9] and [2].) To review some of the details, recall that the Lie algebroid $A = \ker ds|_{G_0}$, with anchor $\rho = dt : A \rightarrow TG_0$.

A section $a \in \Gamma(A)$ gives rise to a morphism of multiplicative vector fields as follows. Let $\tilde{a} = dt(a)$ and $\check{a} = \overrightarrow{a} + \overleftarrow{a}$, where $\overrightarrow{a}(g) = dR_g(a(t(g)))$ and $\overleftarrow{a}(g) = d(L_g \circ i)(a(s(g)))$. (Here, for $g \in G_1$, L_g and R_g denote left and right multiplication by g , and $i : G_1 \rightarrow G_1$ denotes inversion.) It follows that (\tilde{a}, \check{a}) is a multiplicative vector field [24, Example 3.4], and we may view a as a morphism $a : (\tilde{\xi}, \check{\xi}) \rightarrow (\tilde{\zeta}, \check{\zeta})$ whenever $(\tilde{\zeta}, \check{\zeta}) - (\tilde{\xi}, \check{\xi}) = (\tilde{a}, \check{a})$, with addition as composition.

Below we record some additional facts about multiplicative vector bundles for the case of bundle gerbes that we will use in Section 4. In particular, we see from Proposition 3.12 (4) that the multiplicative vector fields isomorphic to the zero vector are lifts of the zero vector on M .

Lemma 3.11. *Let (X, P, μ) be a bundle gerbe over M with connection (B, γ) . Let $a \in \Gamma(A_P)$, where $A_P \rightarrow X$ denotes the Lie algebroid of the corresponding Lie groupoid $P \rightrightarrows X$. Set $\mathbf{v}_a = \epsilon^* \iota_{\tilde{a}} \gamma$. Then $\iota_{\check{a}} \gamma = -\delta \mathbf{v}_a$.*

Proof. The proof is a straightforward computation that uses the multiplicativity of the connection form γ . Indeed, for $p \in P$, we have

$$\begin{aligned} (\iota_{\tilde{a}} \gamma)(p) &= \gamma_p dR_p(a(t(p))) \\ &= \gamma_{\epsilon(t(p))_p} dR_p(a(t(p))) \\ &= \gamma_{\epsilon(t(p))}(a(t(p))) \\ &= \mathbf{v}_a((t(p))). \end{aligned}$$

Similarly, $(\iota_{\check{a}} \gamma)(p) = -\mathbf{v}_a(s(p))$, as required. \square

Proposition 3.12. *Let (X, P, μ) be a bundle gerbe over M with connection (B, γ) , and let A_P denote the Lie algebroid of \mathbf{P} .*

- (1) *For any $g : X \rightarrow \mathbb{R}$, $a = g \frac{\partial}{\partial \theta} \Big|_X$ defines a section of A_P with $\check{a} = (\delta g) \frac{\partial}{\partial \theta}$.*

- (2) Let $(\tilde{\xi}, \check{\xi})$ be a multiplicative vector field. If $\tilde{\xi} \sim_{\pi} 0$, then there exists a section $a \in \Gamma(A_P)$ with $(\tilde{\xi}, \check{\xi}) = (\tilde{a}, \check{a})$.

Proof. The first item is straightforward. To verify (2), it suffices to consider the case that $\check{\xi}$ is horizontal, which is done next. Let $a = \text{Lift}_{\gamma}(\tilde{\xi}, 0)|_X$ denote the restriction of the horizontal lift of $(\tilde{\xi}, 0)$ on $X^{[2]}$ to P . Since $\tilde{a} = \tilde{\xi}$, it remains to check $\check{a} = \check{\xi}$. By Lemma 3.11, \check{a} is horizontal, and it is straightforward to check that $\check{a} \sim_{\pi_P} (\tilde{\xi}, \check{\xi})$. Therefore, $\check{a} = \check{\xi}$, as required. \square

Example 3.13. Consider, as in Examples 3.4 and 3.10, a bundle gerbe P given by Čech data. The Lie algebroid A_P in this case consists of vectors in $\bigsqcup TU_i \times TS^1$ of the form $(0, u)$. Therefore, we may identify a section $a \in \Gamma(A_P)$ with a collection of functions $a = \{u_i : U_i \rightarrow \mathbb{R}\}$ that record the vertical components. Hence $\tilde{a} = 0$, and a straightforward computation shows that (after identifying vertical vectors with functions) $\check{a} = \{u_j - u_i\}$. These are the morphisms of the category $\mathcal{L}_{g_{ijk}}$ in [9]. \diamond

3.4. Multiplicative vector fields on bundle gerbes with connective structure. Let $\mathcal{G} = (X, P, \mu)$ be a bundle gerbe over M with connective structure (B, γ) . In this Section, we introduce the category of multiplicative vector fields on the corresponding groupoid $P \rightrightarrows X$ that preserve the connective structure.

We begin with the following Proposition, which gives an interpretation of what the connective structure (B, γ) on \mathcal{G} measures in terms of multiplicative vector fields, analogous to the interpretation given in [9, Section 9].

Proposition 3.14. *Let $\mathcal{G} = (X, P, \mu)$ be a bundle gerbe over M with connective structure (B, γ) . Let $\tilde{\xi}$ and $\tilde{\zeta}$ be vector fields on X that are lifts of vector fields ξ and ζ on M .*

- (1) *Let $\text{Lift}_{\gamma}\tilde{\xi}^{[2]}$ be the horizontal lift of $\tilde{\xi}^{[2]}$ to P with respect to γ . Then $(\tilde{\xi}, \text{Lift}_{\gamma}\tilde{\xi}^{[2]})$ is a multiplicative vector field.*
- (2) *Let $g = \iota_{\tilde{\xi}}\iota_{\tilde{\zeta}}B$ and $a = g\frac{\partial}{\partial\theta}|_X \in \Gamma(A_P)$. Then a defines an isomorphism*

$$([\tilde{\xi}, \tilde{\zeta}], [\text{Lift}_{\gamma}\tilde{\xi}^{[2]}, \text{Lift}_{\gamma}\tilde{\zeta}^{[2]}]) \longrightarrow ([\tilde{\xi}, \tilde{\zeta}], \text{Lift}_{\gamma}[\tilde{\xi}, \tilde{\zeta}]^{[2]})$$

of multiplicative vector fields.

Proof. To verify (1), observe that the first two items of Proposition 3.9 are easily checked for $(\tilde{\xi}, \text{Lift}_{\gamma}\tilde{\xi}^{[2]})$. It remains to check compatibility with the groupoid multiplication. This follows from the observation that the product of horizontal vectors in TP is horizontal, since the connection γ is multiplicative.

To verify (2), we note

$$\begin{aligned}
\delta g &= \delta \iota_{\tilde{\zeta}} \iota_{\tilde{\zeta}} B \\
&= \iota_{\text{Lift}_\gamma \tilde{\xi}^{[2]}} \iota_{\text{Lift}_\gamma \tilde{\zeta}^{[2]}} \delta B \\
&= \iota_{\text{Lift}_\gamma \tilde{\xi}^{[2]}} \iota_{\text{Lift}_\gamma \tilde{\zeta}^{[2]}} d\gamma \\
&= \iota_{[\text{Lift}_\gamma \tilde{\xi}^{[2]}, \text{Lift}_\gamma \tilde{\zeta}^{[2]}]} \gamma
\end{aligned}$$

Therefore, by Proposition 3.12 (3), $a \in \Gamma(A_P)$ satisfies

$$(\tilde{a}, \check{a}) = (0, [\text{Lift}_\gamma \tilde{\xi}^{[2]}, \text{Lift}_\gamma \tilde{\zeta}^{[2]}] - \text{Lift}_\gamma[\tilde{\xi}, \tilde{\zeta}]^{[2]}),$$

as required. \square

Let $(\tilde{\xi}, \check{\xi})$ be a multiplicative vector field on a bundle gerbe $\mathcal{G} = (X, P, \mu)$ over M with connective structure (B, γ) . In Definition 3.15 below, we propose what it means for $(\tilde{\xi}, \check{\xi})$ to preserve the connective structure. To give a heuristic motivation for the definition, suppose we are given a diffeomorphism $F : M \rightarrow M$. A bundle gerbe morphism covering F is a *Morita morphism*—that is, an isomorphism $(Q, \gamma_Q) : (\mathcal{G}, B, \gamma) \dashrightarrow (\mathcal{G}, \tilde{F}^*B, \check{F}^*\gamma)$ of bundle gerbes with connection. (Note that $F^*\mathcal{G}$ is naturally isomorphic to \mathcal{G} since F is a diffeomorphism.) Let us take such an isomorphism to be of the form $Q = X \times S^1$, $\gamma_Q = \pi_Q^*\alpha + d\theta$, where $\alpha \in \Omega^1(X)$. Hence,

$$\tilde{F}^*B - B = d\alpha \quad \text{and} \quad \check{F}^*\gamma - \gamma = \delta\alpha.$$

Thinking of \tilde{F} and \check{F} as the flows of $\tilde{\xi}$ and $\check{\xi}$, respectively, we arrive at the Definition below.

Definition 3.15. Let $(\tilde{\xi}, \check{\xi})$ be a multiplicative vector field on a bundle gerbe $\mathcal{G} = (X, P, \mu)$ over M with connective structure (B, γ) . We say $(\tilde{\xi}, \check{\xi})$ *preserves* (B, γ) if there exists a 1-form $\alpha \in \Omega^1(X)$ such that

$$L_{\tilde{\zeta}}B = d\alpha \quad \text{and} \quad L_{\check{\zeta}}\gamma = \delta\alpha. \quad (3.2)$$

Remark 3.16. One could also consider multiplicative vector fields $(\tilde{\xi}, \check{\xi})$ on a bundle gerbe \mathcal{G} that preserve a connection γ (without necessarily preserving a curving). In this case, α is required to satisfy only the condition $L_{\check{\zeta}}\gamma = \delta\alpha$ from (3.2).

Proposition 3.17. *Let $\mathcal{G} = (X, P, \mu)$ be a bundle gerbe over M with connective structure (B, γ) , and let $a \in \Gamma(A_P)$. The multiplicative vector field (\tilde{a}, \check{a}) preserves (B, γ) .*

Proof. Let $\alpha = \iota_{\tilde{a}}B - d\nu_a$, and use Lemma 3.11 and the equality $\delta B = d\gamma$ to check that $L_{\tilde{a}}B = d\alpha$, and $L_{\check{a}}\gamma = \delta\alpha$. \square

Corollary 3.18. *Let (X, P, μ) be a bundle gerbe over M with connective structure (B, γ) . There is a category $\mathbb{X}(\mathbf{P}; B, \gamma)$ whose objects are pairs $(\tilde{\xi}, \check{\xi}; \alpha) \in \mathbb{X}(\mathbf{P})_0 \times$*

$\Omega^1(X)$ satisfying equation (3.2), and whose morphisms are sections $a \in \Gamma(A_P)$:

$$a : (\tilde{\xi}, \tilde{\xi}; \alpha) \rightarrow (\tilde{\zeta}, \tilde{\zeta}; \beta), \quad \text{iff} \quad (\tilde{\zeta}, \tilde{\zeta}; \beta) - (\tilde{\xi}, \tilde{\xi}; \alpha) = (\tilde{a}, \tilde{a}, \iota_{\tilde{a}}B - d\nu_a).$$

Moreover, the forgetful functor $\mathbb{X}(\mathbf{P}; B, \gamma) \rightarrow \mathbb{X}(\mathbf{P})$ is full and faithful.

Example 3.19. Consider the trivial bundle gerbe \mathcal{T}_ω with trivial connection and curving ω . A multiplicative vector field is simply a vector field on M . Indeed, for a multiplicative vector field, the component on $P = M \times S^1$ must be the trivial lift of a vector field ξ on M , since its vertical component would (by virtue of being multiplicative) define a continuous homomorphism $S^1 \rightarrow \mathbb{R}$, which must be trivial. Therefore, a multiplicative vector field $(\xi, \xi + 0)$ on $P \rightrightarrows M$ preserves the connective structure if and only if $L_\xi \omega$ is exact. \diamond

Example 3.20. Continuing from Examples 3.10 and 3.13, let (ξ_{ij}, f_{ij}) be a multiplicative vector field on the bundle gerbe P given in terms of Čech data. Let $(\{B_i\}, \{A_{ij}\})$ be a collection of 2-forms on the U_i and 1-forms on the U_{ij} that specify a connective structure on P , and suppose that (ξ_{ij}, f_{ij}) preserves the connective structure. Then there exists a collection of 1-forms $a_i \in \Omega^1(U_i)$ satisfying

$$\begin{aligned} \text{pr}^*(a_j - a_i) &= L_{\xi_{ij} + \text{pr}^* f_{ij} \frac{\partial}{\partial \theta}}(\text{pr}^* A_{ij} + d\theta) \\ &= \text{pr}^* L_{\xi_{ij}} A_{ij} + L_{\text{pr}^* f_{ij} \frac{\partial}{\partial \theta}} d\theta \\ &= \text{pr}^*(L_{\xi_{ij}} A_{ij} + df) \end{aligned}$$

on U_{ij} . That is,

$$a_j - a_i = L_{\xi_{ij}} A_{ij} + df. \quad (3.3)$$

In [9], Collier defines the category $\mathcal{L}_{(g_{ijk}, A_{ij})}$ whose objects $(\xi, \{-f_{ij}\}, \{-a_i\})$ are triples consisting of a vector field ξ on M , functions f_{ij} satisfying (3.1), and 1-forms a_i satisfying (3.3), as well as a (full) subcategory $\mathcal{L}_{(g_{ijk}, B_i, A_{ij})}$ whose objects satisfy the additional constraint $L_\xi B_i = da_i$ (i.e., infinitesimal symmetries that preserve the connection and the curving). As in Example 3.13, the morphisms are given by collections of functions $\{u_i : U_i \rightarrow \mathbb{R}\}$, which, as noted in that Example, correspond to sections a of the underlying Lie algebroid. Therefore, $\{u_i\}$ is a morphism $(\xi, \{-f_{ij}\}, \{-a_i\}) \rightarrow (\xi, \{-g_{ij}\}, \{-b_i\})$ if and only if $f_{ij} - g_{ij} = u_j - u_i$ and $b_i - a_i = du_i$, and we recover the category of connection-preserving infinitesimal symmetries from *op. cit.* \diamond

3.5. Multiplicative vector fields on bundle gerbes as infinitesimal symmetries of the underlying gerbe. By the discussion in Examples 3.10 and 3.13, the category of multiplicative vector fields on a bundle gerbe $\mathcal{G}_{g_{ijk}}$ over a manifold M , given in terms of a Čech cocycle $\{g_{ijk}\}$ associated to an open cover $\{U_i\}$ of M , coincides with the category $\mathcal{L}_{g_{ijk}}$ of *infinitesimal symmetries* of the gerbe as defined by Collier in [9]. A similar discussion in Example 3.20 considers infinitesimal symmetries that preserve the connective structure of the gerbe. In this section, we verify more generally that the category of multiplicative vector fields $\mathbb{X}(\mathbf{P})$ on a bundle

gerbe $\mathcal{G} = (X, P, \mu)$ is equivalent to the category $\mathcal{L}_{\mathcal{G}}$ in *op. cit.* of infinitesimal symmetries of the underlying gerbe (i.e., viewed as a *stack* over M), in the perspective of *op. cit.* A similar equivalence of categories incorporating a connective structure on \mathcal{G} is also given.

The purpose of this section is to connect our work to Collier's notion of infinitesimal symmetries of gerbes. In the interest of brevity, and since the content of this section is not used elsewhere in the paper, this section assumes a familiarity with the *presheaf of groupoids* (i.e., stack) viewpoint of S^1 -gerbes (e.g., as in Brylinski's book [6]). We shall follow Collier's notation closely, and review some elements in [9] to state our results precisely in that context.

To begin, recall how a bundle gerbe $\mathcal{G} = (X, P, \mu)$ over M gives rise to a gerbe, which by abuse of notation will also be denoted by \mathcal{G} . For each open set $U \subset M$, $\mathcal{G}(U)$ is the category of local trivializations of the bundle gerbe restricted to U . More precisely, objects of $\mathcal{G}(U)$ are pairs (Q, ϕ) consisting of an S^1 -bundle $\pi_Q : Q \rightarrow X_U = \pi^{-1}(U)$ together with a bundle isomorphism $\phi : \partial Q = \partial_1^* Q^* \otimes \partial_0^* Q \rightarrow P_U = P|_{X_U^{[2]}}$, satisfying a compatibility condition with bundle gerbe product μ (*cf.* Definition 3.5). A morphism $(Q, \phi) \rightarrow (Q', \phi')$ is given by a bundle isomorphism $u : Q \rightarrow Q'$ that is compatible with ϕ and ϕ' , $\phi \circ \partial u = \phi'$. The resulting presheaf of groupoids \mathcal{G} is seen to be a *stack*; moreover, it naturally has *band* equal to the sheaf of S^1 -valued functions on M . That is, \mathcal{G} is an S^1 -(banded) gerbe (see [35] for details).

Infinitesimal symmetries of S^1 -gerbes. Let $\underline{\mathbb{R}}_M$ and $\underline{S^1}_M$ denote the sheaves of smooth real and circle valued functions on M , respectively. For a vector field ξ on M and a function $g : M \rightarrow S^1$, let $\iota_{\xi} d \log(g) = \iota_{\xi} g^{-1} dg$, which defines a morphism of sheaves $\iota_{\xi} d \log : \underline{S^1}_M \rightarrow \underline{\mathbb{R}}_M$.

Given a vector field ξ on M , Collier defines a category $\mathcal{L}_{\mathcal{G}}(\xi)$ whose objects are *infinitesimal symmetries* of an S^1 -gerbe \mathcal{G} that *lift* ξ . Specifically, an object in $\mathcal{L}_{\mathcal{G}}(\xi)$ is a morphism of gerbes (i.e., a morphism of presheaves or a pseudo-natural transformation)

$$F : \mathcal{G} \rightarrow B\underline{\mathbb{R}}_M,$$

intertwining the morphisms of sheaves, $-\iota_{\xi} d \log$. Here, $B\underline{\mathbb{R}}_M$ denotes the trivial gerbe with band $\underline{\mathbb{R}}_M = C^{\infty}(M, \mathbb{R})$: given an open set $U \subset M$, $B\underline{\mathbb{R}}_M(U)$ is the category of $\underline{\mathbb{R}}_U$ -torsors. In particular, we obtain a family of functors $F_U : \mathcal{G}(U) \rightarrow B\underline{\mathbb{R}}_M(U)$ (or simply F when the underlying open set U is understood) such that for all objects Q in $\mathcal{G}(U)$, $F(Q)$ is an $\underline{\mathbb{R}}_U$ -torsor; moreover, for all morphisms $u : Q \rightarrow Q$, and all $g : U \rightarrow S^1$, $F(u) : F(Q) \rightarrow F(Q)$ satisfies $F(u \cdot g) = F(u) - \iota_{\xi} g^{-1} dg$.

A morphism $F \Rightarrow F'$ of infinitesimal symmetries lifting ξ is a modification of the underlying pseudo-natural transformations. In particular, such a morphism is specified by a map of $\underline{\mathbb{R}}_U$ -torsors $T : F(Q) \rightarrow F'(Q)$ for every object Q in $\mathcal{G}(U)$. Let $\mathcal{L}_{\mathcal{G}}$ denote the category whose objects consist of the collection infinitesimal lifts F of vector fields ξ on M as above, and whose morphisms are exactly those in $\mathcal{L}_{\mathcal{G}}(\xi)$

described above (i.e., there are no morphisms between infinitesimal lifts of different vector fields on M). Theorem 3.22 below shows $\mathcal{L}_{\mathcal{G}}$ is equivalent to the category of multiplicative vector fields $\mathbb{X}(\mathbf{P})$.

Lemma 3.21. *Let $\pi : X \rightarrow M$ be a surjective submersion, let $\pi_Q : Q \rightarrow X$ be a principal S^1 -bundle, and let $\partial Q = \partial_1^* Q^* \otimes \partial_0^* Q \rightarrow X^{[2]}$ and $(X, \partial Q, \mu)$ be the resulting canonically trivialized bundle gerbe. Let $a \in \Gamma(A_{\partial Q})$, and recall $\tilde{a} = dt(a)$, and $\check{a} = \overrightarrow{\tilde{a}} + \overleftarrow{\tilde{a}}$.*

- (1) *There is a unique vector field $\hat{a} \in \mathfrak{X}(Q)^{S^1}$ such that the section a is of the form $a(x) = [(\tilde{a}(x), 0, \hat{a}(q), 0)]$, where $\pi_Q(q) = x$.*
- (2) *$\check{a}(x_1, x_2, q_1 \otimes q_2) = [(\tilde{a}(x_1), \tilde{a}(x_2), \hat{a}(q_1), \hat{a}(q_2))]$ for all $(x_1, x_2, q_1 \otimes q_2) \in \partial Q$.*
- (3) *Let γ_Q denote a connection form on $Q \rightarrow X$, inducing a bundle gerbe connection $\partial\gamma_Q$ on ∂Q . Then $\pi_Q^* \mathbf{v}_a = -\iota_{\hat{a}} \gamma_Q$.*

Proof. We begin by reviewing elementary aspects of the Lie algebroid $A_{\partial Q}$. For a point $\mathbf{q} = (x_1, x_2, q_1 \otimes q_2) \in \partial Q$, where $(x_1, x_2) \in X^{[2]}$ and $\pi_Q(q_i) = x_i$, vectors in $T_{\mathbf{q}} \partial Q$ may be represented by equivalence classes $\mathbf{v} = [(u_1, u_2, v_1, v_2)]$ where $u_i \in T_{x_i} X$, $v_i \in T_{q_i} Q$ satisfy $d\pi(u_1) = d\pi(u_2)$ and $d\pi_Q(v_i) = u_i$ for $i = 1, 2$.

For $\mathbf{v} \in \ker ds$, where $s(\mathbf{q}) = x_2$ is the source map of $\partial Q \rightrightarrows X$, then $u_2 = 0$. Thus there exists a unique representative for \mathbf{v} of the form $\mathbf{v} = [(u_1, 0, w, 0)]$. Therefore, letting $\tilde{s} : \partial_1^* Q^* \times_{X^{[2]}} \partial_0^* Q \rightarrow X$ and $\tilde{\partial}_1 : \partial_1^* Q^* \times_{X^{[2]}} \partial_0^* Q \rightarrow Q$ denote the maps $\tilde{s}(x_1, x_2, q_1, q_2) = x_2$ and $\tilde{\partial}_1(x_1, x_2, q_1, q_2) = q_2$, respectively, we see that the map

$$\begin{aligned} (\ker d\tilde{s} \cap \ker d\tilde{\partial}_1)|_Q &\longrightarrow Q \times_X A_{\partial Q} \\ (x, x, q, q; u, 0, w, 0) &\longmapsto (q; x, x, q \otimes q; [u, 0, w, 0]) \end{aligned}$$

is a diffeomorphism.

To prove (1), let $a \in \Gamma(A_{\partial Q})$. We obtain the vector field $\hat{a} \in \mathfrak{X}(Q)$ as the composition

$$Q \xrightarrow{(\text{id}, a \circ \pi_Q)} Q \times_X A_{\partial Q} \longrightarrow (\ker d\tilde{s} \cap \ker d\tilde{\partial}_1)|_Q \xrightarrow{\text{pr}} TQ.$$

The vector field \hat{a} is S^1 -invariant:

$$[(\tilde{a}(x), 0, \hat{a}(q \cdot z), 0)] = [(\tilde{a}(x), 0, \hat{a}(q), 0)] = [(\tilde{a}(x), 0, dR_z \hat{a}(q), 0)].$$

Item (2) is a straightforward calculation. Similarly, since $\overrightarrow{\tilde{a}}(x_1, x_2, q_1 \otimes q_2) = [(\tilde{a}(x_1), 0, \hat{a}(q_1), 0)]$, and the induced connection on the bundle gerbe ∂Q is $\partial\gamma_Q = \partial_1^* \gamma_Q^* \otimes \partial_0^* \gamma_Q$, we see that for $q \in Q$,

$$\begin{aligned} \mathbf{v}_a(\pi_Q(q)) &= (\partial\gamma_Q)_{(\pi_Q(q), \pi_Q(q); q \otimes q)} [(\tilde{a}(\pi_Q(q)), 0, \hat{a}(q), 0)] \\ &= -(\gamma_Q)_q(\hat{a}(q)), \end{aligned}$$

as required. \square

Theorem 3.22. *Let $\mathcal{G} = (X, P, \mu)$ be a bundle gerbe over M . There is an equivalence of categories $\mathbb{X}(\mathbf{P}) \rightarrow \mathcal{L}_{\mathcal{G}}$.*

Proof. We construct the functor $\mathbb{X}(\mathbf{P}) \rightarrow \mathcal{L}_{\mathcal{G}}$ directly and provide a sketch that it is an equivalence of categories. Let $\bar{\xi} = (\tilde{\xi}, \check{\xi})$ be a multiplicative vector field that descends to a vector field ξ on M . Let $F_{\bar{\xi}} : \mathcal{G} \rightarrow B\mathbb{R}_M$ be given as follows. To each object (Q, ϕ) in $\mathcal{G}(U)$, let

$$F_{\bar{\xi}}(Q) = \{\hat{\xi} \in \mathfrak{X}(Q)^{S^1} \mid \hat{\xi} \sim_{\pi_Q} \tilde{\xi}|_{X_U}, \text{ and } d\phi(\delta\hat{\xi}) = \check{\xi}|_{P_U}\}$$

be the corresponding \mathbb{R}_U -torsor. Here, using the S^1 -invariance of $\tilde{\xi}$, $\delta\hat{\xi}$ denotes the vector field on $\partial_0^*Q \otimes \partial_1^*Q$ induced from $(\tilde{\xi}, \check{\xi})$ on each factor ∂_i^*Q , $i = 0, 1$.

Let $a \in \Gamma(A_P)$, and let $a_U \in \Gamma(A_{P_U})$ denotes its restriction to X_U . Observe that $\phi : \partial Q \rightarrow P_U$ induces a Lie groupoid isomorphism

$$\begin{array}{ccc} \partial_0^*Q \otimes \partial_1^*Q & \xrightarrow{\phi} & P_U \\ \Downarrow & & \Downarrow \\ X_U & \xlongequal{\quad} & X_U \end{array}$$

and hence we may view a_U as a section of the Lie algebroid of $\partial Q \rightrightarrows X_U$. The section a_U gives rise to an S^1 -invariant vector field \hat{a} on Q , as in Lemma 3.21, which satisfies $\hat{a} \sim_{\pi_Q} \check{a}|_{X_U}$, and $d\phi(\delta\hat{a}) = \check{a}|_{P_U}$.

Viewing $a : (\tilde{\xi}, \check{\xi}) \rightarrow (\tilde{\zeta}, \check{\zeta})$ as a morphism in $\mathbb{X}(\mathbf{P})$, the vector fields \hat{a} from the previous paragraph give rise to a morphism $T_a : F_{\bar{\xi}} \Rightarrow F_{\bar{\zeta}}$. Specifically, for each $(Q, \phi) \in \mathcal{G}(U)$, we obtain a map of \mathbb{R}_U -torsors $T_a : F_{\bar{\xi}}(Q) \rightarrow F_{\bar{\zeta}}(Q)$, by setting $T_a(\hat{\xi}) = \hat{\xi} + \hat{a}$. Hence, we obtain a functor $\mathbb{X}(\mathbf{P}) \rightarrow \mathcal{L}_{\mathcal{G}}$.

To show that the functor constructed above is an equivalence of categories, note first that, by [9, Corollary 6.1.8 (2)], it is immediate that the functor is essentially surjective. Faithfulness follows from the uniqueness of the vector field \hat{a} , as in Lemma 3.21. To see that the functor is full, suppose we are given an equivalence of lifts $T : F_{\bar{\zeta}}(Q) \rightarrow F_{\bar{\xi}}(Q)$. There exists $a' \in \Gamma(A_P)$ with $(\tilde{a}', \check{a}') = (\tilde{\zeta} - \tilde{\xi}, \check{\zeta} - \check{\xi})$, and the collection of such sections a' is a torsor for $C^\infty(M)$ (i.e., sections of the isotropy Lie algebra)—this follows from Prop 3.12 (3) and (4) and the exact sequence 2.2. For $\hat{\xi} \in F_{\bar{\xi}}(Q)$, $T(\hat{\xi}) - T_{a'}(\hat{\xi})$ is an S^1 -invariant lift of 0 (restricted to X_U), which may thus identify with a (basic) function $f : U \rightarrow \mathbb{R}$ that records its vertical component. Since $d\phi(\delta(T(\hat{\xi}) - T_{a'}(\hat{\xi}))) = 0$, $\delta f = 0$, and we have that such functions patch together to give a global function $g : M \rightarrow \mathbb{R}$. Letting $a = a' + (\pi^*g)\frac{\partial}{\partial\theta}$, it is straightforward to check that $T = T_a$, as required. \square

Infinitesimal symmetries of S^1 -gerbes preserving a connective structure. Recall that for an S^1 -gerbe \mathcal{G} , viewed as a presheaf of groupoids, a connection on \mathcal{G} is a morphism

of gerbes $\mathcal{A} : \mathcal{G} \rightarrow B\underline{\Omega}_M^1$ intertwining the morphism of sheaves $-d \log : \underline{S}_M^1 \rightarrow \Omega_M^1$. Here, $B\underline{\Omega}_M^1$ denotes the trivial gerbe with band $\underline{\Omega}_M^1$: specifically, given an open set $U \subset M$, $\underline{\Omega}_M^1(U)$ is the category of $\underline{\Omega}_U^1$ -torsors. In particular, we obtain a family of functors $\mathcal{A}_U : \mathcal{G}(U) \rightarrow B\underline{\Omega}_M^1(U)$ (or simply \mathcal{A} when the underlying open set U is understood) such that for all objects $Q \in \mathcal{G}(U)$, $\mathcal{A}(Q)$ is a $\underline{\Omega}_U^1$ -torsor; moreover, for all morphisms $u : Q \rightarrow Q$, and all $g : U \rightarrow S^1$, $\mathcal{A}(u) : \mathcal{A}(Q) \rightarrow \mathcal{A}(Q)$ satisfies $\mathcal{A}(u \cdot g) = \mathcal{A}(u) - d \log g$. A *curving* K for the connection \mathcal{A} assigns to each $Q \in \mathcal{G}(U)$ and each $\gamma \in \mathcal{A}(Q)$, a 2-form $K(\gamma) \in \Omega^2(U)$. This assignment is compatible with restrictions of open subsets, and morphisms $u : Q \rightarrow Q'$ —that is, $K((\mathcal{A}(u)(\gamma))) = K(\gamma)$. Moreover, K is $\Omega^1(U)$ -equivariant: for all $\alpha \in \Omega^1(U)$, $K(\gamma \cdot \alpha) = K(\gamma) + d\alpha$. We will call the pair (\mathcal{A}, K) a *connective structure* for \mathcal{G} .

Observe that for an S^1 -gerbe \mathcal{G} associated to a bundle gerbe (X, P, μ) , a connective structure (B, γ) for the bundle gerbe gives rise to a connective structure in the sense described above. Namely, for $U \subset M$ and $(Q, \phi) \in \mathcal{G}(U)$, we set

$$\mathcal{A}(Q) = \{\text{connection 1-forms } \gamma_Q \in \Omega^1(Q) : \phi^*(\gamma|_{P_U}) = \partial\gamma_Q\},$$

where $\partial\gamma_Q = \partial_1^* \gamma_Q^* \otimes \partial_0^* \gamma_Q$ is the induced connection on ∂Q . The curving K is then obtained as follows: observe that $\partial \text{curv}(\gamma_Q) = \text{curv}(\gamma|_{P_U}) = \partial B|_U$ and hence there exists a unique 2-form $K(\gamma_Q)$ satisfying $\pi^* K(\gamma_Q) = \text{curv}(\gamma_Q) - B|_U$. Note that it follows that $dK(\gamma_Q) = -\chi|_U$.

In [9, 10], Collier defines categories $\mathcal{L}_{(\mathcal{G}, \mathcal{A})}$ and $\mathcal{L}_{(\mathcal{G}, \mathcal{A}, K)}$ of infinitesimal symmetries of \mathcal{G} that preserve the connection and the connective structure, respectively. Recall that an infinitesimal symmetry lifting $\xi \in \mathfrak{X}(M)$ that preserves the connection \mathcal{A} , also called a *connective lift* of ξ , is a pair (F, Θ) consisting of an infinitesimal lift $F : \mathcal{G} \rightarrow B\underline{\mathbb{R}}_M$ in $\mathcal{L}_{\mathcal{G}}(\xi)$ and a (2-)morphism Θ for the diagram:

$$\begin{array}{ccc} \mathcal{G} & \xrightarrow{\mathcal{A}} & B\underline{\Omega}_M^1 \\ \downarrow F & \not\rightarrow_{\Theta} & \downarrow L_{-\xi}[\] \\ B\underline{\mathbb{R}}_M & \xrightarrow{-d[\]} & B\underline{\Omega}_M^1 \end{array}$$

That is, Θ is a modification for the two composite pseudo-natural transformations represented in the diagram. Recall that in this context, $-d[\] : B\underline{\mathbb{R}}_M \rightarrow B\underline{\Omega}_M^1$ may be described as follows: for each open set $U \subset M$ we obtain a family of functors $B\underline{\mathbb{R}}_M(U) \rightarrow B\underline{\Omega}_M^1(U)$ associating to each $\underline{\mathbb{R}}_U$ -torsor, the $\underline{\Omega}_U^1$ -torsor associated to it via $-d : C^\infty(U) \rightarrow \Omega^1(U)$. That is, for any open $V \subset U$ and any $C^\infty(V)$ -torsor A over V , $-d[A] = (A \times \Omega^1(V)) / \sim$, where by definition $(a \cdot f, \mu) \sim (a, \mu - df)$ for any $f : V \rightarrow \mathbb{R}$. The map $L_{-\xi}[\]$ may be described similarly.

A connection-preserving multiplicative vector field $(\tilde{\xi}, \check{\xi}; \alpha) \in \mathbb{X}(\mathbf{P}; B, \gamma)$ naturally gives rise to a connective lift $(F_{\tilde{\xi}}, \Theta_{(\tilde{\xi}, \alpha)})$ as follows. The infinitesimal symmetry $F_{\tilde{\xi}}$ is as above. To describe $\Theta_{(\tilde{\xi}, \alpha)}$, we define for each $U \subset M$ and $(Q, \phi) \in \mathcal{G}(U)$, a map of $\Omega^1(U)$ -torsors $\Theta_{(\tilde{\xi}, \alpha)}(Q) : L_{-\xi}[\mathcal{A}(Q)] \rightarrow -d[F_{\tilde{\xi}}(Q)]$. To that end, since $\delta \iota_{\tilde{\xi}} \gamma = 0$,

there exists a function $g \in C^\infty(X)$ satisfying $\iota_\xi \gamma = \delta g$, and we may set

$$\widehat{\xi}_g = \text{Lift}_{\gamma_Q}(\widetilde{\xi}) + (\pi_Q^* g|_{X_U}) \frac{\partial}{\partial \theta} \in F_{\widetilde{\xi}}(Q).$$

Since $\delta(\alpha - \iota_{\widetilde{\xi}} B - dg) = 0$, there exists a unique 1-form $\varepsilon_g \in \Omega^1(M)$ such that $\pi^* \varepsilon_g = \alpha - \iota_{\widetilde{\xi}} B - dg$. Now for $[(\gamma_Q, \tau)] \in L_{-\xi}[\mathcal{A}(Q)]$, let

$$\Theta_{(\widetilde{\xi}, \alpha)}(Q)[(\gamma_Q, \tau)] = [(\widehat{\xi}_g, \tau - \iota_\xi K(\gamma_Q) - \varepsilon_g)].$$

Observe that the above map is independent of the choice of g : any other such choice g' must be of the form $g' = g + \pi^* h$ for some $h \in C^\infty(M)$, which yields $\varepsilon_{g'} = \varepsilon_g - dh$, and $\widehat{\xi}_{g'} = \widehat{\xi}_g + (\pi_Q^* \pi^* h|_{X_U}) \frac{\partial}{\partial \theta}$. Hence, $(\widehat{\xi}_{g'}, \tau - \iota_\xi K(\gamma_Q) - \varepsilon_{g'}) \sim (\widehat{\xi}_g, \tau - \iota_\xi K(\gamma_Q) - \varepsilon_g)$.

To check $\Theta_{(\widetilde{\xi}, \alpha)}(Q)$ is well-defined,

$$\begin{aligned} & \Theta_{(\widetilde{\xi}, \alpha)}(Q)[(\gamma_Q + \pi_Q^* \pi^* \nu, \tau + L_\xi \nu)] \\ &= [(\widehat{\xi}_g - \pi_Q^* \pi^* \iota_\xi \nu \frac{\partial}{\partial \theta}, \tau + L_\xi \nu - \iota_\xi K(\gamma_Q + \pi_Q^* \pi^* \nu) - \varepsilon_g)] \\ &= [(\widehat{\xi}_g, \tau + L_\xi \nu - \iota_\xi (K(\gamma_Q) + d\nu) - \varepsilon_g - d\iota_\xi \nu)] \\ &= \Theta_{(\widetilde{\xi}, \alpha)}(Q)[(\gamma_Q, \tau)]. \end{aligned}$$

A connective lift (F, Θ) preserves the curving K if for each $(Q, \phi) \in \mathcal{G}(U)$ and each $\gamma \in \mathcal{A}(Q)$,

$$d\Theta(Q)(\gamma) = -L_\xi K(\gamma).$$

In this context, recall that if we write $\Theta(Q)[(\gamma, 0)] = [(\widehat{\xi}, \mu)]$, then $d\mu \in \Omega^2(U)$ is well-defined, and hence we set $d\Theta(Q)(\gamma) = d\mu$. Therefore, continuing from the above discussion, the connective lift $(F_{\widetilde{\xi}}, \Theta_{(\widetilde{\xi}, \alpha)})$ associated to $(\widetilde{\xi}, \check{\xi}; \alpha) \in \mathbb{X}(\mathbf{P}; B, \gamma)$ preserves the curving K if $d(-\iota_\xi K(\gamma_Q) - \varepsilon_g) = -L_\xi K(\gamma_Q)$, or equivalently $d\varepsilon_g = \iota_\xi dK(\gamma_Q)$. Since $dK(\gamma_Q) = -\chi|_U$, we see that $(F_{\widetilde{\xi}}, \Theta_{(\widetilde{\xi}, \alpha)})$ preserves the curving if ξ is Hamiltonian, with Hamiltonian 1-form ε_g .

A morphism $(F, \Theta) \Rightarrow (F', \Theta')$ of connective lifts is a so-called *connective equivalence*; i.e., a morphism of $F \Rightarrow F'$ of the underlying infinitesimal symmetries that is compatible with Θ and Θ' . A morphism of connective lifts that preserve the curving is simply a morphism of the underlying connective lifts. (*cf.* [10, Definition 4.1.16]).

For a morphism $a : (\widetilde{\xi}, \check{\xi}; \alpha) \rightarrow (\widetilde{\zeta}, \check{\zeta}; \beta)$ of multiplicative vector fields preserving the connective structure, we may associate a morphism $T_a : F_{\widetilde{\xi}} \Rightarrow F_{\widetilde{\zeta}}$, as in the proof of Theorem 3.22. It is straightforward to see that T_a is compatible with $\Theta_{(\widetilde{\xi}, \alpha)}$ and $\Theta_{(\widetilde{\zeta}, \beta)}$. In this context, compatibility amounts to the commutativity of the following

diagram,

$$\begin{array}{ccc}
 L_{-\xi}[\mathcal{A}(Q)] & \xrightarrow{\Theta_{(\bar{\xi}, \alpha)}} & -d[F_{\bar{\xi}}(Q)] \\
 & \searrow \Theta_{(\bar{\zeta}, \beta)} & \downarrow -d[T_a] \\
 & & -d[F_{\bar{\zeta}}(Q)]
 \end{array}$$

Before this is verified, we first choose $g \in C^\infty(X)$ satisfying $\iota_\xi \gamma = \delta g$, and set $h = g - \mathbf{v}_a$. Then, it is easy to see that $\varepsilon_g = \varepsilon_h \in \Omega^1(M)$ (as defined earlier). Moreover, by Lemma 3.21 (3), we have $\widehat{\zeta}_h = \widehat{\xi}_g + \widehat{a}$. Therefore,

$$\begin{aligned}
 -d[T_a] \circ \Theta_{(\bar{\xi}, \alpha)}[(\gamma_Q, \tau)] &= -d[T_a][(\widehat{\xi}_g, \tau - \iota_\xi K(\gamma_Q) - \varepsilon_g)] \\
 &= [(T_a(\widehat{\xi}_g), \tau - \iota_\xi K(\gamma_Q) - \varepsilon_g)] \\
 &= [(\widehat{\zeta}_h, \tau - \iota_\xi K(\gamma_Q) - \varepsilon_h)] \\
 &= \Theta_{(\bar{\zeta}, \beta)}[(\gamma_Q, \tau)],
 \end{aligned}$$

as required.

Similar to Theorem 3.22, we see that the above discussion describes a functor $\mathbb{X}(\mathbf{P}; \gamma) \rightarrow \mathcal{L}_{(\mathcal{G}, \mathcal{A})}$, which is an equivalence of categories.

Theorem 3.23. *Let $\mathcal{G} = (X, P, \mu)$ be a bundle gerbe over M . There is an equivalence of categories $\mathbb{X}(\mathbf{P}; \gamma) \rightarrow \mathcal{L}_{(\mathcal{G}, \mathcal{A})}$, which restricts to an equivalence of categories $\mathbb{X}(\mathbf{P}; B, \gamma) \rightarrow \mathcal{L}_{(\mathcal{G}, \mathcal{A}, K)}$.*

Proof. That the functor $\mathbb{X}(\mathbf{P}; \gamma) \rightarrow \mathcal{L}_{(\mathcal{G}, \mathcal{A})}$ is full and faithful is straightforward. It remains to show it is essentially surjective.

Let (F, Θ) be a connective lift of some vector field ξ on M . By Theorem 3.22, there exists a multiplicative vector field $\bar{\xi} = (\widetilde{\xi}, \check{\xi})$ and an isomorphism $S : F \Rightarrow F_{\bar{\xi}}$. Moreover, setting $\Theta' = (S * \text{id}_{-d[\]}) \circ \Theta$, we extend $F_{\bar{\xi}}$ to a connective lift $(F_{\bar{\xi}}, \Theta')$, which is isomorphic to (F, Θ) .

Let $g \in C^\infty(X)$ satisfy $\delta g = \iota_\xi \gamma$, and consider $\alpha_g = \iota_{\bar{\zeta}} B + dg \in \Omega^1(X)$. It is easy to see that $L_{\bar{\zeta}} \gamma = \delta \alpha_g$, and that the collection of α for which $(\widetilde{\xi}, \check{\xi}; \alpha) \in \mathbb{X}(\mathbf{P}; \gamma)$ is a $\Omega^1(M)$ -torsor. Since $(F_{\bar{\zeta}}, \Theta_{(\bar{\zeta}, \alpha_g)})$ is a connective lift extending $F_{\bar{\zeta}}$, and the set of such connective lifts is also a $\Omega^1(M)$ -torsor (by [9, Corollary 10.46]), $(F_{\bar{\zeta}}, \Theta')$ is isomorphic to some $(F_{\bar{\zeta}}, \Theta_{(\bar{\zeta}, \alpha)})$, as required.

The statement about restricting the functor follows from the observation that if there exists $(F, \Theta) \Rightarrow (F', \Theta')$, then (F, Θ) preserves the curving iff (F', Θ') does (cf. [10, Def. 4.1.16]). \square

4. LIE 2-ALGEBRAS AND 2-PECTIC GEOMETRY

Following a review of the bicategory of Lie 2-algebras, we introduce the main examples of interest in this paper—namely, the Poisson Lie 2-algebra of a 2-plectic manifold and the connection-preserving multiplicative vector fields on (i.e., infinitesimal symmetries of) a bundle gerbe (Proposition 4.8).

4.1. Review of Lie 2-algebras. In this paper, we work with Lie 2-algebras using Noohi’s bicategory $2\text{Term}L_\infty^b$ of 2-term L_∞ -algebras, with butterflies as 1-arrows, and morphisms of butterflies as 2-arrows [32]. As shown in *op. cit.*, $2\text{Term}L_\infty^b$ is the localization of the bicategory $2\text{Term}L_\infty$ of 2-term L_∞ -algebras defined by Baez and Crans [2] with respect to quasi-isomorphisms. We begin with a brief review of $2\text{Term}L_\infty$, using definitions from [33]; see also [2, Lemma 4.3.3].

Definition 4.1. A *Lie 2-algebra* is a 2-term chain complex $\mathbb{V} = [V_1 \xrightarrow{d} V_0]$ equipped with a skew-symmetric chain map $[-, -] : \mathbb{V} \otimes \mathbb{V} \rightarrow \mathbb{V}$ and a skew symmetric chain homotopy $J : \mathbb{V}^{\otimes 3} \rightarrow \mathbb{V}$, called the *Jacobiator*, from the chain map

$$\mathbb{V}^{\otimes 3} \rightarrow \mathbb{V}, \quad x \otimes y \otimes z \mapsto [x, [y, z]],$$

to the chain map

$$\mathbb{V}^{\otimes 3} \rightarrow \mathbb{V}, \quad x \otimes y \otimes z \mapsto [[x, y], z] + [y, [x, z]],$$

satisfying

$$\begin{aligned} & [x, J(y, z, w)] + J(x, [y, z], w) + J(x, z, [y, w]) + [J(x, y, z), w] + [z, J(x, y, w)] \\ & = J(x, y, [z, w]) + J([x, y], z, w) + [y, J(x, z, w)] + J(y, [x, z], w) + J(y, z, [x, w]). \end{aligned}$$

A Lie 2-algebra with vanishing Jacobiator is called a strict Lie-2-algebra.

Definition 4.2. A *morphism of Lie 2-algebras* $\mathbf{F} : \mathbb{V} \rightarrow \mathbb{W}$ consists of a chain map $\mathbf{F}_\bullet : V_\bullet \rightarrow W_\bullet$ together with a chain homotopy $F : \mathbb{V} \otimes \mathbb{V} \rightarrow \mathbb{W}$, from the chain map

$$\mathbb{V} \otimes \mathbb{V} \rightarrow \mathbb{W} \quad x \otimes y \mapsto \mathbf{F}([x, y])$$

to the chain map

$$\mathbb{V} \otimes \mathbb{V} \rightarrow \mathbb{W} \quad x \otimes y \mapsto [\mathbf{F}(x), \mathbf{F}(y)],$$

satisfying

$$\begin{aligned} \mathbf{F}_1(J(x, y, z)) - J(\mathbf{F}_0(x), \mathbf{F}_0(y), \mathbf{F}_0(z)) &= F(x, [y, z]) - F([x, y], z) - F(y, [x, z]) \\ &\quad - [F(x, y), \mathbf{F}_0(z)] + [\mathbf{F}_0(x), F(y, z)] - [\mathbf{F}_0(y), F(x, z)]. \end{aligned} \tag{4.1}$$

For completeness, we note that given a pair of morphisms of Lie 2-algebras, $\mathbf{F}, \mathbf{G} : \mathbb{V} \rightarrow \mathbb{W}$, a *2-morphism* between \mathbf{F} and \mathbf{G} is a chain homotopy between them that is compatible with the underlying chain homotopies F and G . The 2-category of Lie 2-algebras with the above morphisms and 2-morphisms is denoted by $2\text{Term}L_\infty$ [2].

We will work with the bicategory 2TermL_∞^b defined by Noohi [32], which we recall next. The objects remain the same; however, the arrows are replaced with butterflies.

Definition 4.3. A *butterfly* $\mathbf{E} : \mathbb{V} \dashrightarrow \mathbb{W}$ of Lie 2-algebras is a vector space E equipped with a skew-symmetric bracket $[-, -]$, together with a commutative diagram

$$\begin{array}{ccccc} V_1 & & & & W_1 \\ & \searrow \kappa & & \swarrow \lambda & \\ & & E & & \\ & \swarrow \sigma & & \searrow \rho & \\ V_0 & & & & W_0 \end{array}$$

such that:

- $\rho \circ \kappa = 0$ and the NE–SW sequence $0 \rightarrow W_1 \rightarrow E \rightarrow V_0 \rightarrow 0$ is short exact;
- ρ and σ preserve brackets;
- for every $a \in E$, $u \in W_1$, $x \in V_1$,

$$[a, \lambda(u)] = \lambda([\rho(a), u]), \quad \text{and} \quad [a, \kappa(x)] = \kappa([\sigma(a), x]);$$

- for every $a, b, c \in E$,

$$\lambda J(\rho(a), \rho(b), \rho(c)) + \kappa J(\sigma(a), \sigma(b), \sigma(c)) = [a, [b, c]] + [b, [c, a]] + [c, [a, b]].$$

Definition 4.4. Given a pair of butterflies $\mathbf{E}, \mathbf{F} : \mathbb{V} \dashrightarrow \mathbb{W}$, a *morphism of butterflies* $\varphi : \mathbf{E} \Rightarrow \mathbf{F}$ is a linear map $E \rightarrow F$ that commutes with the brackets and all the structure maps of the butterflies. (As noted in [32], the linear map in a morphism of butterflies is necessarily an isomorphism.)

The composition of butterflies $\mathbb{U} \xrightarrow{\mathbf{E}} \mathbb{V} \xrightarrow{\mathbf{E}'} \mathbb{W}$ is given by

$$\begin{array}{ccccc} U_1 & & & & W_1 \\ & \searrow (\kappa, 0) & & \swarrow (0, \lambda') & \\ & & E \oplus_{V_0} E' & & \\ & \swarrow \sigma \circ \text{pr} & & \searrow \rho' \circ \text{pr} & \\ U_0 & & & & W_0 \end{array}$$

where $E \oplus_{V_0} E'$ denotes the quotient of $E \oplus_{V_0} E'$ by the image of V_1 under (λ, κ') , with bracket defined on components.

As shown in [32], Lie 2-algebras with butterflies as 1-arrows and morphism of butterflies as 2-arrows form a bicategory 2TermL_∞^b . Moreover, Noohi explicitly defines a functor $2\text{TermL}_\infty \rightarrow 2\text{TermL}_\infty^b$, and shows that this descends to an equivalence of categories on the localization of 2TermL_∞ with respect to quasi-isomorphisms (i.e., morphisms inducing an isomorphism on homology). In particular, quasi-isomorphisms correspond to invertible butterflies—butterflies whose NW–SE sequence is also short

exact. (Recall that the inverse of an invertible butterfly is obtained by simply ‘flipping’ the butterfly diagram along the vertical axis.)

We will implicitly make use of this functor in what follows—for instance, when writing down 1-arrows of Lie 2-algebras, we may write a morphism of Lie 2-algebras, rather than the corresponding butterfly. For clarity, however, we will distinguish between the two kinds of 1-arrows by employing either a solid or dashed arrow for morphisms or butterflies, respectively.

4.2. Lie 2-algebras associated to pre-2-plectic manifolds and bundle gerbes.

Lie 2-algebras appear naturally in 2-plectic geometry [34], and as infinitesimal symmetries of bundle gerbes [9]. In this section, we recall the 2-plectic analogue of the Poisson-Lie algebra of smooth functions on a symplectic manifold. Continuing (or ‘quantizing’) this analogy, we recall the Lie 2-algebra of multiplicative vector fields on the Lie groupoid $P \rightrightarrows X$ associated to a bundle gerbe $\mathcal{G} = (X, P, \mu)$, and give the Lie 2-algebra structure for connection-preserving multiplicative vector fields in Proposition 4.8. We see that a connective structure on a bundle gerbe \mathcal{G} gives rise to a natural horizontal lift of multiplicative vector fields on $X^{[2]} \rightrightarrows X$ to multiplicative vector fields on $P \rightrightarrows X$, and that the resulting 3-curvature measures the failure of the horizontal lift to be a morphism of Lie 2-algebras (Theorem 4.6).

4.2.1. *The Poisson-Lie 2-algebra.* Recall that a *2-plectic manifold* is a manifold equipped with a closed non-degenerate 3-form (in analogy with symplectic manifolds and closed non-degenerate 2-forms). If we omit the non-degeneracy condition, we say the manifold is *pre-2-plectic* [34].

Definition 4.5. Let (M, χ) be a pre-2-plectic manifold. The *Poisson-Lie 2-algebra* $\mathbb{L}(M, \chi)$ has underlying 2-term complex given by

$$C^\infty(M) \xrightarrow{(0,d)} \{(\xi, \beta) \in \mathfrak{X}(M) \times \Omega^1(M) \mid \iota_\xi \chi = -d\beta\} .$$

with bracket given by

$$[(\xi_1, \beta_1), (\xi_2, \beta_2)] = ([\xi_1, \xi_2], \iota_{\xi_2} \iota_{\xi_1} \chi)$$

on $L_0 \otimes L_0$ and zero otherwise. The Jacobiator $J : L_0^{\otimes 3} \rightarrow L_1$ is given by

$$J(\xi_1, \beta_1; \xi_2, \beta_2; \xi_3, \beta_3) = -\iota_{\xi_3} \iota_{\xi_2} \iota_{\xi_1} \chi .$$

4.2.2. *Lie 2-algebras of infinitesimal symmetries of S^1 -gerbes.* As mentioned in Section 3.3, the category $\mathbb{X}(\mathbf{G})$ of multiplicative vector fields on a Lie groupoid $\mathbf{G} = G_1 \rightrightarrows G_0$ is a (strict) Lie 2-algebra [5]. Indeed, viewing $\mathbb{X}(\mathbf{G})$ as a 2-term chain complex

$$\Gamma(A) \longrightarrow \mathbb{X}(\mathbf{G})_0,$$

the bracket on $\mathbb{X}(\mathbf{G})_0 \otimes \mathbb{X}(\mathbf{G})_0$ is given on components: $[(\tilde{\xi}, \check{\xi}), (\tilde{\zeta}, \check{\zeta})] = ([\tilde{\xi}, \tilde{\zeta}], [\check{\xi}, \check{\zeta}])$, while on $\mathbb{X}(\mathbf{G})_0 \otimes \Gamma(A)$, it is given by $[(\tilde{\xi}, \check{\xi}), a] = [\check{\xi}, \vec{a}]|_{G_0}$. (The bracket on

$\Gamma(A) \otimes \mathbb{X}(\mathbf{G})_0$ is determined by skew-symmetry.) Here, recall that for $a \in \Gamma(A)$, and $(\tilde{\xi}, \check{\xi})$ multiplicative, $[\check{\xi}, \vec{a}] \in \ker ds$ and is right-invariant [24], and hence its restriction to G_0 defines a section in $\Gamma(A)$.

Therefore, associated to a bundle gerbe $\mathcal{G} = (X, P, \mu)$ over a manifold M , we naturally obtain (strict) Lie 2-algebras $\mathbb{X}(X^{[2]} \rightrightarrows X)$ and $\mathbb{X}(P \rightrightarrows X)$ as above. Recall from Proposition 3.14 (1) that a connective structure (B, γ) on \mathcal{G} give rise to a map of multiplicative vector fields $\text{Lift}_\gamma : \mathbb{X}(X^{[2]} \rightrightarrows X)_0 \rightarrow \mathbb{X}(\mathbf{P})_0$, which (cf. the proof of Proposition 3.12 (4)) may be completed to a chain map

$$\text{Lift}_{\gamma, \bullet} : \mathbb{X}(X^{[2]} \rightrightarrows X)_\bullet \longrightarrow \mathbb{X}(\mathbf{P})_\bullet. \quad (4.2)$$

By Proposition 3.14 (2), the curving B gives rise to a map

$$F_B : \mathbb{X}(X^{[2]} \rightrightarrows X)_0 \times \mathbb{X}(X^{[2]} \rightrightarrows X)_0 \longrightarrow \Gamma(A_P) \quad (4.3)$$

by setting $F_B(\tilde{\xi}, \tilde{\xi}^{[2]}; \tilde{\zeta}, \tilde{\zeta}^{[2]}) = \iota_{\tilde{\zeta}} \iota_{\tilde{\zeta}} B \frac{\partial}{\partial \theta} \Big|_X$, which we may view as defining a chain homotopy that controls the failure of Lift_γ to preserve the bracket (as in Definition 4.2). However, Lift_γ fails to be a morphism of Lie 2-algebras (as in Definition 4.2). Indeed, analogous to the horizontal lift for principal S^1 -bundles, the failure of the horizontal lift to be a morphism of Lie 2-algebras (i.e., Condition (4.1) in Definition 4.2) is measured by the 3-curvature of the bundle gerbe, as the next Theorem shows.

Theorem 4.6. *Let (\mathcal{G}, B, γ) be a bundle gerbe over M with connective structure and 3-curvature χ . The horizontal lift Lift_γ and chain homotopy F_B from (4.2) and (4.3) define a Lie 2-algebra morphism if and only if $\chi = 0$.*

Proof. Following the paragraph preceding the Proposition, it remains to consider Condition (4.1) from Definition 4.2. To that end, consider vector fields ξ_1, ξ_2 , and ξ_3 on M with corresponding lifts $\tilde{\xi}_1, \tilde{\xi}_2$, and $\tilde{\xi}_3$ on X . Write $\bar{\xi} = (\tilde{\xi}, \tilde{\xi}^{[2]})$.

Since the Lie 2-algebras under consideration are strict (i.e., with vanishing Jacobiator), Lift_γ and F_B will define a Lie 2-algebra morphism if and only if

$$\begin{aligned} 0 = & F_B(\bar{\xi}_1, [\bar{\xi}_2, \bar{\xi}_3]) - F_B([\bar{\xi}_1, \bar{\xi}_2], \bar{\xi}_3) - F_B(\bar{\xi}_2, [\bar{\xi}_1, \bar{\xi}_3]) \\ & - [F_B(\bar{\xi}_1, \bar{\xi}_2), \text{Lift}_\gamma(\bar{\xi}_3)] + [\text{Lift}_\gamma(\bar{\xi}_1), F_B(\bar{\xi}_2, \bar{\xi}_3)] - [\text{Lift}_\gamma(\bar{\xi}_2), F_B(\bar{\xi}_1, \bar{\xi}_3)]. \end{aligned} \quad (4.4)$$

Since the image of F_B consists of vertical sections of A_P , each of the last three terms above is also a vertical section. To determine the vertical components of these terms, let $a_{jk} = F_B(\bar{\xi}_j, \bar{\xi}_k)$, and $\check{\xi}_i = \text{Lift}_\gamma(\bar{\xi}_i)$, and observe that $\iota_{[\check{\xi}_i, \vec{a}_{jk}]} \gamma = L_{\check{\xi}_i} \iota_{\vec{a}_{jk}} \gamma$, since $\check{\xi}_i$ is horizontal and \vec{a}_{jk} is vertical. Restricting to X , we see that

$$\iota_{[\check{\xi}_i, \vec{a}_{jk}]} \gamma \Big|_X = L_{\tilde{\xi}_i} \iota_{\tilde{\xi}_j} \iota_{\tilde{\xi}_k} B$$

Therefore, taking vertical components in (4.4), we obtain

$$\begin{aligned} 0 = & \iota_{\tilde{\xi}_1} \iota_{[\tilde{\xi}_2, \tilde{\xi}_3]} B - \iota_{[\tilde{\xi}_1, \tilde{\xi}_2]} \iota_{\tilde{\xi}_3} B - \iota_{\tilde{\xi}_2} \iota_{[\tilde{\xi}_1, \tilde{\xi}_3]} B + L_{\tilde{\xi}_3} \iota_{\tilde{\xi}_1} \iota_{\tilde{\xi}_2} B + L_{\tilde{\xi}_1} \iota_{\tilde{\xi}_2} \iota_{\tilde{\xi}_3} B - L_{\tilde{\xi}_2} \iota_{\tilde{\xi}_1} \iota_{\tilde{\xi}_3} B \\ = & \iota_{\tilde{\xi}_1} \iota_{\tilde{\xi}_2} \iota_{\tilde{\xi}_3} dB \end{aligned}$$

Hence Condition (4.1) holds if and only if $\pi^*\chi = dB = 0$, which completes the proof, since $\pi : X \rightarrow M$ is a submersion. \square

The category $\mathbb{X}(\mathbf{P}; B, \gamma)$ from Corollary 3.18 of connection-preserving multiplicative vector fields is easily seen to be a 2-vector space. Proposition 4.8 below shows it is a Lie 2-algebra, whose brackets are inherited from $\mathbb{X}(\mathbf{P})$.

Lemma 4.7. *Let $\mathcal{G} = (X, P, \mu)$ be a bundle gerbe over M with connective structure (B, γ) , and $P \rightrightarrows X$ the associated Lie groupoid. Let $(\tilde{\xi}, \check{\xi}; \alpha) \in \mathbb{X}(\mathbf{P}; B, \gamma)_0$ and $a \in \Gamma(A_P)$. Then $\mathbf{v}_{[\tilde{\xi}, \vec{a}]}|_X = L_{\tilde{\xi}}\mathbf{v}_a + \iota_{\check{a}}\alpha$.*

Proof. First, recall that $\vec{a} \sim_{\delta_1} \tilde{a}$ and $\vec{a} \sim_{\delta_0} 0$. Therefore,

$$\mathbf{v}_{[\tilde{\xi}, \vec{a}]}|_X = \epsilon^* \iota_{[\tilde{\xi}, \vec{a}]} \gamma = \epsilon^* (L_{\tilde{\xi}} \iota_{\vec{a}} \gamma - \iota_{\vec{a}} L_{\tilde{\xi}} \gamma) = L_{\tilde{\xi}} \mathbf{v}_a - \epsilon^* \iota_{\vec{a}} \delta \alpha = L_{\tilde{\xi}} \mathbf{v}_a + \iota_{\check{a}} \alpha.$$

\square

Proposition 4.8. *Let $\mathcal{G} = (X, P, \mu)$ be a bundle gerbe over M with connective structure (B, γ) . Let $\mathbb{X}(\mathcal{G}; B, \gamma) = \{V_1 \rightarrow V_0\}$ denote the 2-term complex*

$$\Gamma(A_P) \xrightarrow{d} \mathbb{X}(\mathbf{P}; B, \gamma)_0,$$

with differential given by $da = (\tilde{a}, \check{a}; \iota_{\tilde{a}} B - d\mathbf{v}_a)$. Define a bracket on $V_0 \otimes V_0$ by

$$[(\tilde{\xi}, \check{\xi}; \alpha), (\tilde{\zeta}, \check{\zeta}; \beta)] = ([\tilde{\xi}, \tilde{\zeta}], [\check{\xi}, \check{\zeta}]; L_{\tilde{\xi}} \beta - L_{\check{\zeta}} \alpha)$$

and on $V_0 \otimes V_1$ by

$$[(\tilde{\xi}, \check{\xi}; \alpha), a] = [\check{\xi}, \vec{a}]|_X.$$

Then $\mathbb{X}(\mathcal{G}; B, \gamma)$ is a strict Lie 2-algebra (i.e., with Jacobiator $J \equiv 0$).

Proof. Let $\mathbb{V} = \{V_1 \rightarrow V_0\}$ as in the statement of the Proposition. It is straightforward to see that the bracket and differential are well-defined. To show the resulting bracket is compatible with the differential (i.e., that it gives a chain map $\mathbb{V} \otimes \mathbb{V} \rightarrow \mathbb{V}$), we must verify that for $(\tilde{\xi}, \check{\xi}; \alpha) \in V_0$ and $a, b \in V_1$:

$$d[(\tilde{\xi}, \check{\xi}; \alpha), a] = [(\tilde{\xi}, \check{\xi}; \alpha), da], \quad \text{and} \quad (4.5)$$

$$[da, b] = -[db, a]. \quad (4.6)$$

For (4.5), first recall that $([\check{\xi}, \vec{a}]|_X)^\sim = [\tilde{\xi}, \tilde{a}]$ and $([\check{\xi}, \vec{a}]|_X)^\vee = [\check{\xi}, \check{a}]$ (e.g., see the proof of [5, Theorem 3.4]). Therefore, using Lemma 4.7,

$$\begin{aligned} d[(\tilde{\xi}, \check{\xi}; \alpha), a] &= ([\tilde{\xi}, \tilde{a}], [\check{\xi}, \check{a}], \iota_{[\tilde{\xi}, \tilde{a}]} B - d\mathbf{v}_{[\check{\xi}, \vec{a}]}|_X) \\ &= ([\tilde{\xi}, \tilde{a}], [\check{\xi}, \check{a}], L_{\tilde{\xi}} \iota_{\tilde{a}} B - \iota_{\check{a}} L_{\tilde{\xi}} B - d(L_{\tilde{\xi}} \mathbf{v}_a + \iota_{\check{a}} \alpha)) \\ &= ([\tilde{\xi}, \tilde{a}], [\check{\xi}, \check{a}], L_{\tilde{\xi}} (\iota_{\tilde{a}} B - d\mathbf{v}_a) - L_{\check{a}} \alpha) \\ &= [(\tilde{\xi}, \check{\xi}; \alpha), da]. \end{aligned}$$

For (4.6), we first recall that $[\check{a}, \vec{b}] = [\vec{a} + \check{a}, \vec{b}] = [\vec{a}, \vec{b}]$, since left and right invariant vector fields commute. Therefore,

$$[da, b] = [\check{a}, \vec{b}]|_X = -[\check{b}, \vec{a}]|_X = -[db, a],$$

as required. The vanishing of the Jacobiator is straightforward. \square

Example 4.9. Consider the connection-preserving multiplicative vector fields for the trivial bundle gerbe \mathcal{I}_ω from Example 3.19. As a Lie 2-algebra $\mathbb{X}(\mathcal{I}_\omega) = \{W_1 \rightarrow W_0\}$ has $W_1 \cong C^\infty(M)$, since vectors in the Lie algebroid must necessarily be vertical vectors, and W_0 consists of pairs (ξ, A) such that $L_\xi\omega = dA$. The differential $d : W_1 \rightarrow W_0$ is $f \mapsto (0, -df)$. The bracket on $W_0 \otimes W_0$ is given by

$$[(\xi_1, A_1), (\xi_2, A_2)] = ([\xi_1, \xi_2], L_{\xi_1}A_2 - L_{\xi_2}A_1),$$

and on $W_0 \otimes W_1$ it is given by

$$[(\xi, A), f] = \xi(f).$$

Note that there is a morphism of Lie 2-algebras $F : \mathbb{X}(\mathcal{I}_\omega) \rightarrow \mathbb{L}(M, d\omega)$, which in degree 0 sends $(\xi, A) \mapsto (\xi, \iota_\xi\omega + A)$. It is straightforward to see that F is a quasi-isomorphism. Theorem 5.1 generalizes this observation. \diamond

Remark 4.10. In Example 3.13, we saw that the category of multiplicative vector fields $\mathbb{X}(\mathbf{P})$ on a bundle gerbe P given by a Čech 2-cocycle g_{ijk} recovers the category $\mathcal{L}_{g_{ijk}}$ of infinitesimal symmetries from [9]. In *op. cit.*, Collier also shows $\mathcal{L}_{g_{ijk}}$ is a Lie 2-algebra—the Lie 2-algebra structure coincides with $\mathbb{X}(\mathbf{P})$ from [5]. The analogous statement holds as well for the category $\mathcal{L}_{(g_{ijk}, B_i, A_{ij})}$ of infinitesimal symmetries preserving the curving and $\mathbb{X}(\mathcal{G}_{g_{ijk}, \{B_i\}, \{A_{ij}\}})$. \diamond

5. APPLICATIONS TO 2-PLECTIC AND QUASI-HAMILTONIAN GEOMETRY

In this section, we establish some 2-plectic and quasi-Hamiltonian analogues of classical results in symplectic geometry. In Theorem 5.1, we show the Poisson Lie 2-algebra of a 2-plectic manifold (M, χ) is quasi-isomorphic to the connection-preserving multiplicative vector fields on a bundle gerbe with connective structure and 3-curvature χ .

Let G be a compact Lie group with Lie algebra \mathfrak{g} . We apply the aforementioned quasi-isomorphism to the framework of quasi-Hamiltonian G -actions with G -valued moment maps and find an analogue of Kostant's theorem (Theorem 5.7) lifting a quasi-Hamiltonian \mathfrak{g} -action to a given 'prequantization' in that setting.

5.1. Infinitesimal symmetries of S^1 -gerbes and pre-2-plectic manifolds.

Let (M, ω) be a symplectic manifold, and suppose we are given a prequantization: a principal S^1 -bundle $\pi : P \rightarrow M$ with connection $\gamma \in \Omega^1(P)$ whose curvature $\text{curv}(\gamma) = \omega$. Recall that the formula

$$f \mapsto \text{Lift}_\gamma(X_f) + \pi^* f \frac{\partial}{\partial \theta}$$

defines a Lie algebra isomorphism $C^\infty(M) \xrightarrow{\cong} \mathcal{Q}(P, \gamma)$, where $\text{Lift}_\gamma(X_f)$ denotes the horizontal lift of X_f , the Hamiltonian vector field of f , and

$$\mathcal{Q}(P, \gamma) = \{X \in \mathfrak{X}(P) : L_X \gamma = 0\}$$

denotes the so-called *infinitesimal quantomorphisms* of P [18, 37]. In Theorem 5.1 below, we give the 2-plectic analogue:

Theorem 5.1. *Let (M, χ) be a pre-2-plectic manifold, and let $\mathcal{G} = (X, P, \mu)$ be a bundle gerbe with connective structure (B, γ) and 3-curvature χ . There is an invertible butterfly $\mathbf{E} : \mathbb{L}(M, \chi) \dashrightarrow \mathbb{X}(\mathcal{G}; B, \gamma)$.*

Proof. For a multiplicative vector field $(\tilde{\xi}, \check{\xi})$, observe that $\delta(\iota_{\tilde{\xi}} \gamma) = \iota_{(\tilde{\xi}, \check{\xi})} \delta \gamma = 0$; therefore, since $P \rightrightarrows X$ is proper, there exists $g : X \rightarrow \mathbb{R}$ such that $\delta g = \iota_{\tilde{\xi}} \gamma$. We will show that $E = \{(\tilde{\xi}, \check{\xi}; \alpha, g) \in V_0 \times C^\infty(X) \mid \delta g = \iota_{\tilde{\xi}} \gamma\}$ is the required invertible butterfly:

$$\begin{array}{ccccc} L_1 & & & & V_1 \\ & \searrow \kappa & & \swarrow \lambda & \\ & & E & & \\ & \swarrow \sigma & & \searrow \rho & \\ L_0 & & & & V_0 \end{array}$$

The indicated structure maps are defined as follows. Let $\rho = \text{pr}_1$ be the obvious projection, $\lambda(a) = (da, -\mathbf{v}_a)$, and $\kappa(f) = (0, 0; 0, \pi^* f)$. To define σ , first observe that $\delta(\alpha - \iota_{\tilde{\xi}} B - dg) = 0$, and hence there exists a unique 1-form $\varepsilon \in \Omega^1(M)$ such that $\pi^* \varepsilon = \alpha - \iota_{\tilde{\xi}} B - dg$. Hence $\pi^* \iota_{\tilde{\xi}} \chi = \iota_{\tilde{\xi}} dB = \pi^* d\varepsilon$. Set $\sigma(\tilde{\xi}, \check{\xi}; \alpha, g) = (\xi, -\varepsilon)$.

Define the bracket on E by

$$[(\tilde{\xi}, \check{\xi}, \alpha, g), (\tilde{\zeta}, \check{\zeta}, \beta, h)] = ([(\tilde{\xi}, \check{\xi}, \alpha), (\tilde{\zeta}, \check{\zeta}, \beta)], \iota_{\tilde{\xi}} \beta - \iota_{\tilde{\zeta}} \alpha + \iota_{\tilde{\zeta}} \iota_{\tilde{\xi}} B).$$

Next, we verify the required compatibility of the bracket with the structure maps. Clearly σ preserves the brackets. Also,

$$[(\tilde{\xi}, \check{\xi}, \alpha, g), \kappa(h)] = [(\tilde{\xi}, \check{\xi}, \alpha, g), (\bar{0}, 0, \pi^* h)] = 0 = \kappa[(\xi, \varepsilon), h],$$

since brackets of elements of mixed degree vanish in $\mathbb{L}(M, \chi)$. By Lemma 4.7 and the properties of the bracket for $\mathbb{X}(\mathcal{G}, B, \gamma)$, we have $[(\tilde{\xi}, \check{\xi}, \alpha, g), \lambda(a)] = \lambda[(\tilde{\xi}, \check{\xi}, \alpha), a]$. Finally, to see that σ preserves the brackets, it suffices to verify that

$$-\pi^* \iota_{\zeta} \iota_{\xi} \chi = L_{\tilde{\zeta}} \beta - L_{\tilde{\zeta}} \alpha - \iota_{[\tilde{\xi}, \tilde{\zeta}]} B - d(\iota_{\tilde{\xi}} \beta - \iota_{\tilde{\zeta}} \alpha + \iota_{\tilde{\zeta}} \iota_{\tilde{\xi}} B),$$

which follows from standard applications of Cartan calculus on differential forms.

Since the Jacobiator for $\mathbb{X}(\mathcal{G}, B, \gamma)$ is trivial, the Jacobiator condition on \mathbf{E} simplifies to

$$\kappa J_{\mathbb{L}}(\xi_1, \varepsilon_1; \xi_2, \varepsilon_2; \xi_3, \varepsilon_3) = [(\tilde{\xi}_1, \check{\xi}_1; \alpha_1), [(\tilde{\xi}_2, \check{\xi}_2; \alpha_2), (\tilde{\xi}_3, \check{\xi}_3; \alpha_3)]] + \text{cyc. perm.}$$

The V_0 -component in the above equality follows from the vanishing of the Jacobiator for $\mathbb{X}(\mathcal{G}, B, \gamma)$. To verify the component in $C^\infty(X)$, it should be verified that

$$-\pi^* \iota_{\xi_3} \iota_{\xi_2} \iota_{\xi_1} \chi = \iota_{\tilde{\xi}_1} (L_{\tilde{\xi}_2} \alpha_3 - L_{\tilde{\xi}_3} \alpha_2) - \iota_{[\tilde{\xi}_2, \tilde{\xi}_3]} \alpha_1 + \iota_{[\tilde{\xi}_2, \tilde{\xi}_3]} \iota_{\tilde{\xi}_1} B + \text{cyc. perm.},$$

which is a straightforward, albeit tedious, application of Cartan calculus on differential forms—see Proposition A.1

To show \mathbf{E} is an invertible butterfly, we must check that the diagonal sequences are short exact. That the NW–SE sequence is short exact follows easily from properness of the Lie groupoid $P \rightrightarrows X$. Exactness of the NE–SW sequence is verified next.

To check surjectivity of σ : given $(\xi, \varepsilon) \in L_0$, choose a multiplicative lift of ξ and take $\alpha' = -\pi^* \varepsilon + \iota_{\tilde{\xi}} B$. Next, find a function f with $\delta f = \iota_{\xi} \gamma$ and set $\alpha = \alpha' + df$. Then $\sigma(\tilde{\xi}, \check{\xi}; \alpha, f) = (\xi, \varepsilon)$

To check injectivity of λ : suppose $a \in \Gamma(A_P)$ is a section of the Lie algebroid and $da = 0$ and $\mathbf{v}_a = 0$. From $da = 0$, we get that $\tilde{a} = 0$, which gives that a lies in the isotropy Lie algebra (in this case trivial, i.e., $X \times \mathbb{R}$), whence we view a as a function on X , and we get that $\check{a} = 0$, which gives that $\delta a = 0$ and hence a descends to M . From $\mathbf{v}_a = 0$, we see that the vertical part of a must vanish, but since $\tilde{a} = 0$ all of a is vertical, so a must vanish.

Exactness at E : It's easy to see that $\sigma \circ \lambda = 0$ since π is a submersion. Suppose $\sigma(\tilde{\xi}, \check{\xi}, \alpha, g) = 0$. Then $\xi = 0$ and $\alpha - \iota_{\tilde{\xi}} B - dg = 0$. The first of those gives $(\tilde{\xi}, \check{\xi}) = (\tilde{a}, \check{a})$ for some $a \in \Gamma(A_P)$ (See Proposition 3.12.) Since $\delta \mathbf{v}_a = \delta g$, there exists a (unique) $f : M \rightarrow \mathbb{R}$ such that $\pi^* f = g - \mathbf{v}_a$. We may view f as defining a section of the isotropy Lie algebra, as noted in the previous paragraph, by setting $b = \pi^* f$. That is, $b : X \rightarrow A_P$ is a vertical vector in $TP|_X$ given by $b(x) = f(\pi(x)) \frac{\partial}{\partial \theta} \Big|_{\epsilon(x)}$. Note that $\mathbf{v}_b = \pi^* f$, and $\tilde{b} = 0$. Moreover, $\check{b} = 0$ by Proposition 3.12. Therefore, $\lambda(a + b) = (\tilde{\xi}, \check{\xi}, \alpha, g)$. \square

Remark 5.2. In [13, Thm 4.6], the authors prove, in the more general context of n -plectic geometry and L_∞ -algebras, a result similar to Theorem 5.1. Specializing to $n = 2$, the result in *op. cit.* shows that $\mathbb{L}(M, \chi)$ is quasi-isomorphic to a Lie 2-algebra the authors call the *infinitesimal quantomorphisms* of a principal $U(1)$ -2-bundle with connection. In that context, a principal $U(1)$ -2-bundle is represented by a Čech-Deligne cocycle, which is equivalent to the data of an S^1 -gerbe with connective structure given in terms of Čech data, while the corresponding Lie 2-algebra of

infinitesimal quantomorphisms is essentially the Lie 2-algebra of infinitesimal symmetries preserving the connective structure from [9]. (*Cf.* Example 3.20 and Remark 4.10 .)

Consider a Lie group G acting on a symplectic manifold (M, ω) , and suppose $\pi : P \rightarrow M$ is a prequantization. Let \mathfrak{g} denote the Lie algebra of G . If the G -action on (M, ω) is Hamiltonian, with moment map $\mu : M \rightarrow \mathfrak{g}^*$, we may compose the Kostant Lie algebra isomorphism $C^\infty(M) \cong \mathcal{Q}(P, \gamma)$ with the (co)moment map $\langle \mu, - \rangle : \mathfrak{g} \rightarrow C^\infty(M)$, to obtain a lift of the infinitesimal action $\mathfrak{g} \rightarrow \mathfrak{X}_{\text{Ham}}(M, \omega)$, $\xi \mapsto \xi_M$, to a connection-preserving \mathfrak{g} -action on P , according to Kostant's formula [18],

$$\mathfrak{g} \longrightarrow \mathcal{Q}(P, \gamma), \quad \xi \mapsto \text{Lift}_\gamma(\xi_M) + \pi^* \langle \mu, \xi \rangle \frac{\partial}{\partial \theta}. \quad (5.1)$$

Analogously, consider a Lie group G acting on a pre-2-plectic manifold (M, χ) . If the G -action preserves χ and the corresponding \mathfrak{g} -action $\xi \mapsto \xi_M$ is via Hamiltonian vector fields (i.e., there exist 1-forms $\varepsilon \in \Omega^1(M)$ such that $\iota_{\xi_M} \chi = -d\varepsilon$), the authors in [7] define a *homotopy moment map* to be a morphism of Lie 2-algebras that lifts the natural projection $\mathbb{L}(M, \chi) \rightarrow \mathfrak{X}(M, \chi)$, as in the following diagram.

$$\begin{array}{ccc} & & \mathbb{L}(M, \chi) \\ & \nearrow & \downarrow \\ \mathfrak{g} & \longrightarrow & \mathfrak{X}(M, \chi). \end{array}$$

Suppose (\mathcal{G}, B, γ) is a bundle gerbe over M with 3-curvature χ . If the \mathfrak{g} -action on (M, χ) admits a homotopy moment map, we may compose it with the invertible butterfly of Theorem 5.1 and obtain a Lie 2-algebra butterfly

$$\mathfrak{g} \dashrightarrow \mathbb{X}(\mathcal{G}; B, \gamma),$$

which is analogous to Kostant's formula (5.1) above in 2-plectic geometry.

The following Theorem shows that the invertible butterfly from Theorem 5.1 is compatible with trivializations, which will be useful in Section 5.2 where we discuss the quasi-Hamiltonian context.

Theorem 5.3. *A trivialization $(Q, \gamma_Q) : \mathcal{I}_\omega \dashrightarrow (\mathcal{G}, B, \gamma)$ of bundle gerbes with connection induces an invertible butterfly $Q : \mathbb{X}(\mathcal{I}_\omega) \dashrightarrow \mathbb{X}(\mathcal{G}; B, \gamma)$. Additionally, there is a canonical 2-isomorphism $\varphi : Q \circ E' \Longrightarrow E$,*

$$\begin{array}{ccc} & & \mathbb{L}(M, d\omega) \\ & \swarrow & \searrow \\ & E' & \varphi \Uparrow \\ \mathbb{X}(\mathcal{I}_\omega) & \dashrightarrow & \mathbb{X}(\mathcal{G}; B, \gamma) \\ & Q & \end{array}$$

where \mathbf{E} and \mathbf{E}' are the invertible butterflies from Theorem 5.1 corresponding to (\mathcal{G}, B, γ) and \mathcal{I}_ω , respectively.

Proof. Let $(Q, \gamma_Q) : \mathcal{I}_\omega \dashrightarrow (\mathcal{G}, B, \gamma)$ denote a trivialization (see Example 3.6). Denote the underlying 2-term chain complexes by $\mathbb{X}(\mathcal{I}_\omega) = \{W_1 \rightarrow W_0\}$ (see Example 4.9) and $\mathbb{X}(\mathcal{G}; B, \gamma) = \{V_1 \rightarrow V_0\}$ (see Proposition 4.8). As a vector space, let

$$F = \{(\tilde{\xi}, \check{\xi}, \alpha, g) \in V_0 \times C^\infty(X) \mid \delta g = \iota_{\tilde{\xi}} \gamma\}.$$

We will show that F gives rise to an invertible butterfly \mathbf{Q} :

$$\begin{array}{ccc} W_1 & & V_1 \\ & \searrow \kappa & \swarrow \lambda \\ & F & \\ & \swarrow \sigma & \searrow \rho \\ W_0 & & V_0 \end{array}$$

The structure maps are defined as follows. Let $\kappa(f) = (0, 0; 0, \pi^* f)$, $\lambda(a) = (da, -\mathbf{v}_a)$, $\rho = \text{pr}$, and $\sigma(\tilde{\xi}, \check{\xi}, \alpha, g) = (\xi, A)$, where ξ is the vector field on M onto which $\tilde{\xi}$ descends, and $A \in \Omega^1(M)$ is determined by $\pi^* A = \alpha - \iota_{\tilde{\xi}} \text{curv}(\gamma_Q) - dg$. Finally, the bracket on F is given by

$$[(\tilde{\xi}, \check{\xi}, \alpha, g), (\tilde{\zeta}, \check{\zeta}, \beta, h)] = ([(\tilde{\xi}, \check{\xi}, \alpha), (\tilde{\zeta}, \check{\zeta}, \beta)], L_{\tilde{\xi}} h - L_{\tilde{\zeta}} g + \iota_{\tilde{\xi}} \iota_{\tilde{\zeta}} \text{curv}(\gamma_Q)).$$

Next, we verify the required compatibility of the bracket with the structure maps. Clearly, ρ preserves brackets. Let $(\tilde{\xi}, \check{\xi}, \alpha, g)$ and $(\tilde{\zeta}, \check{\zeta}, \beta, h)$ in F . To check that σ preserves brackets, let $\sigma(\tilde{\xi}, \check{\xi}, \alpha, g) = (\xi, A_\alpha)$ and $\sigma(\tilde{\zeta}, \check{\zeta}, \beta, h) = (\zeta, A_\beta)$. Then it suffices to check that

$$\begin{aligned} \pi^*(L_\xi A_\beta - L_\zeta A_\alpha) &= L_{\tilde{\xi}}(\beta - \iota_{\tilde{\zeta}} \text{curv}(\gamma_Q) - dh) - L_{\tilde{\zeta}}(\alpha - \iota_{\tilde{\xi}} \text{curv}(\gamma_Q) - dg) \\ &= L_{\tilde{\xi}} \beta - L_{\tilde{\zeta}} \alpha - \iota_{[\tilde{\xi}, \tilde{\zeta}]} \text{curv}(\gamma_Q) - d(L_{\tilde{\xi}} h - L_{\tilde{\zeta}} g + \iota_{\tilde{\xi}} \iota_{\tilde{\zeta}} \text{curv}(\gamma_Q)). \end{aligned}$$

For $f \in W_1$,

$$[(\tilde{\xi}, \check{\xi}, \alpha, g), \kappa(f)] = (0, 0; 0, L_{\tilde{\xi}}(\pi^* f)) = \kappa(\xi(f)) = \kappa[\sigma(\tilde{\xi}, \check{\xi}, \alpha, g), f],$$

as required. For $a \in V_1$,

$$[(\tilde{\xi}, \check{\xi}, \alpha, g), \lambda(a)] = ([(\tilde{\xi}, \check{\xi}, \alpha), da], -L_{\tilde{\xi}}(\mathbf{v}_a) - L_{\tilde{\xi}} a + \iota_{\tilde{\xi}} \iota_{\tilde{\xi}} \text{curv}(\gamma_Q)),$$

while

$$\lambda[\rho(\tilde{\xi}, \check{\xi}, \alpha, g), a] = \lambda[(\tilde{\xi}, \check{\xi}, \alpha), a] = (d[(\tilde{\xi}, \check{\xi}, \alpha), a], -\mathbf{v}_{[\tilde{\xi}, \tilde{\xi}]}|_X).$$

Since

$$\begin{aligned} -L_{\tilde{\xi}}(\mathbf{v}_a) - L_{\tilde{\xi}} a + \iota_{\tilde{\xi}} \iota_{\tilde{\xi}} \text{curv}(\gamma_Q) &= -L_{\tilde{\xi}}(\mathbf{v}_a) - \iota_{\tilde{\xi}}(dg + \iota_{\tilde{\xi}} \text{curv}(\gamma_Q)) \\ &= -L_{\tilde{\xi}}(\mathbf{v}_a) - \iota_{\tilde{\xi}}(\alpha - \pi^* A_\alpha) \\ &= -\mathbf{v}_{[\tilde{\xi}, \tilde{\xi}]}|_X, \end{aligned}$$

by Lemma 4.7, the desired compatibility is shown.

Finally, since both Lie 2-algebras are strict, we must show that the bracket on F satisfies the Jacobi identity. Since the Jacobiator vanishes on $V_0^{\otimes 3}$, we need only check the component in $C^\infty(X)$ vanishes. Hence it remains to show that for $(\tilde{\xi}_i, \check{\xi}_i, \alpha_i; g_i)$ ($i = 1, 2, 3$):

$$\begin{aligned} L_{\tilde{\xi}_1}(L_{\tilde{\xi}_2}g_3 - L_{\tilde{\xi}_3}g_2 + \iota_{\tilde{\xi}_2}\iota_{\tilde{\xi}_1}\text{curv}(\gamma_Q)) - L_{[\tilde{\xi}_2, \tilde{\xi}_3]}g_1 \\ + \iota_{\tilde{\xi}_1}\iota_{[\tilde{\xi}_2, \tilde{\xi}_3]}\text{curv}(\gamma_Q) + \text{cyc. perm.} = 0 \end{aligned}$$

This follows from the fact that $d\text{curv}(\gamma_Q) = 0$.

The exactness of the diagonal sequences of the butterfly are verified in the same way as in the proof of Theorem 5.1.

Next we verify the statement about the compatibility of \mathbf{Q} with the invertible butterflies exhibited in (the proof of) Theorem 5.1. Let $\mathbf{E} : \mathbb{X}(\mathcal{G}, B, \gamma) \dashrightarrow \mathbb{L}(M, d\omega)$ and $\mathbf{E}' : \mathbb{X}(\mathcal{I}_\omega) \dashrightarrow \mathbb{L}(M, d\omega)$ be the resulting invertible butterflies. Given the identification of $\mathbb{X}(\mathcal{I}_\omega) = \{W_1 \rightrightarrows W_0\}$ used above (i.e., as in Example 4.9), observe that \mathbf{E}' may be written as

$$\begin{array}{ccccc} W_1 & & & & L_1 \\ & \searrow \kappa' & & \swarrow \lambda' & \\ & & E' & & \\ & \swarrow \sigma' & & \searrow \rho' & \\ W_0 & & & & L_0 \end{array}$$

where $E' = W_0 \times C^\infty(M)$, with structure maps $\kappa'(f) = (0, -df, -f)$, $\sigma' = \text{pr}$, $\lambda'(f) = (0, 0, f)$ and $\rho'(\xi, A, f) = -A + \iota_\xi\omega + df$.

The composition $\mathbb{L}(M, d\omega) \xrightarrow{E'} \mathbb{X}(\mathcal{I}_\omega) \xrightarrow{\mathbf{Q}} \mathbb{X}(\mathcal{G}, B, \gamma)$ is given by $E' \overset{W_1}{\oplus}_{W_0} F$.

Writing elements of the fibre product $E' \oplus_{W_0} F$ as $(\xi, A, f; \tilde{\xi}, \check{\xi}, \alpha; g)$, we see that the map

$$E' \overset{W_1}{\oplus}_{W_0} F \rightarrow E, \quad [(\xi, A, f; \tilde{\xi}, \check{\xi}, \alpha; g)] \mapsto (\tilde{\xi}, \check{\xi}, \alpha; g + \pi^* f)$$

is the desired 2-isomorphism. \square

5.2. Kostant's formula for Lie group-valued moment maps. In this section, we apply the results of Section 5.1 to the setting of quasi-Hamiltonian group actions with Lie group-valued moment maps to obtain an analogue of Kostant's formula for lifting (infinitesimally) a Hamiltonian Lie group action on a symplectic manifold to a given prequantization. For the remainder of this section, G shall denote a connected, compact Lie group with Lie algebra \mathfrak{g} , equipped with an invariant inner product $\langle -, - \rangle$.

Lie group-valued moment maps. Let θ^L and θ^R denote the left and right invariant Maurer-Cartan forms on a compact Lie group G , and let $\eta = \frac{1}{12}\langle \theta^L, [\theta^L, \theta^L] \rangle \in$

$\Omega^3(G)$ denote the Cartan 3-form. We view G as a G -manifold equipped with the conjugation action.

Definition 5.4. [1] A *quasi-Hamiltonian G -space* is a triple (M, ω, Φ) consisting of a G -manifold M , a G -invariant 2-form $\omega \in \Omega^2(M)^G$ and a G -equivariant map $\Phi : M \rightarrow G$, satisfying

- (i) $d\omega + \Phi^*\eta = 0$,
- (ii) $\iota_{\xi_M}\omega + \frac{1}{2}\Phi^*\langle \theta^L + \theta^R, \xi \rangle = 0$ for all $\xi \in \mathfrak{g}$, and
- (iii) at every point $x \in M$, $\ker \omega_x \cap \ker d\Phi_x = \{0\}$.

Analogous to coadjoint orbits in the dual of the Lie algebra \mathfrak{g}^* as fundamental examples of symplectic manifolds with Hamiltonian G -actions, conjugacy classes $C \hookrightarrow G$ are important examples of quasi-Hamiltonian G -spaces. Another important example is the double $G \times G$, with G acting by conjugation on each factor and G -valued moment map given by the commutator $(g, h) \mapsto ghg^{-1}h^{-1}$. Using the *fusion product* operation, these examples form the building blocks for a (finite dimensional) quasi-Hamiltonian description of the moduli space of flat G -bundles over a compact surface Σ , with prescribed holonomies along the boundary curves. See [1] for details.

In [26], Meinrenken defines prequantization of quasi-Hamiltonian G -spaces using Dixmier-Douady bundles (or simply DD-bundles), which are equivalent to S^1 -bundle gerbes. More precisely, the bicategory of DD-bundles over a fixed manifold is equivalent to the bicategory of bundle gerbes (and similarly for their G -equivariant counterparts) [23]. We recall the definition of prequantization next, providing a formulation using bundle gerbes.

To begin, observe that for a quasi-Hamiltonian G -space (M, ω, Φ) , the first condition in Definition 5.4 (together with the fact that $d\eta = 0$) shows that the pair (ω, η) defines a closed *relative* differential form in $\Omega^3(\Phi)$. In fact, together with the second condition in Definition 5.4, we have that (ω, η_G) defines a closed relative G -equivariant differential form in $\Omega_G^3(\Phi)$. A *prequantization* of a quasi-Hamiltonian G -space (M, ω, Φ) is a *relative G -equivariant bundle gerbe* (Q, \mathcal{G}) for Φ —i.e., a G -equivariant bundle gerbe $\mathcal{G} \rightarrow G$, together with a G -equivariant trivialization $Q : \mathcal{I} \dashrightarrow \Phi^*\mathcal{G}$, whose relative DD-class maps to $[(\omega, \eta_G)] \in H_G^3(\Phi; \mathbb{R})$ via the map induced in cohomology from the coefficient homomorphism $\mathbb{Z} \rightarrow \mathbb{R}$.

Let $(\mathcal{G}, B_G, \gamma)$ be a G -equivariant bundle gerbe over G with connective structure and G -equivariant 3-curvature η_G . Then a G -equivariant trivialization $(Q, \gamma_Q) : \mathcal{I}_{-\omega} \dashrightarrow (\mathcal{G}, B_G, \gamma)$ gives a prequantization for (M, ω, Φ) .

Remark 5.5. Note that a prequantization for (M, ω, Φ) exists if and only if the relative cohomology class $[(\omega, \eta_G)] \in H_G^3(\Phi; \mathbb{R})$ is integral (i.e., in the image of the coefficient homomorphism $H_G^3(\Phi; \mathbb{R}) \rightarrow H^3(G; \mathbb{Z})$). For related work on the integrality conditions required for prequantization of the moduli space of flat G -bundles, see [19, 20, 22, 25].

A quasi-Hamiltonian analogue of Kostant's formula. For Hamiltonian G -spaces (M, ω) with prequantization $P \rightarrow M$, Kostant's formula (5.1) provides a lift of the \mathfrak{g} -action on M to a connection-preserving \mathfrak{g} -action on P . In this Section, we propose an analogue of such a lift for quasi-Hamiltonian G -spaces.

Let (M, ω, Φ) be a quasi-Hamiltonian G -space. Suppose we are given a (not necessarily G -equivariant) bundle gerbe (\mathcal{G}, B, γ) over G with connective structure whose 3-curvature is η , together with a trivialization $(Q, \gamma_Q) : \mathcal{I}_{-\omega} \dashrightarrow (\mathcal{G}, B, \gamma)$. By Theorem 5.3, the trivialization (Q, γ_Q) gives rise to an invertible butterfly of Lie 2-algebras $\mathbf{Q} : \mathbb{X}(\mathcal{I}_{-\omega}) \dashrightarrow \mathbb{X}(\Phi^*(\mathcal{G}, B, \gamma))$, and such a butterfly is compatible with the butterflies resulting from Theorem 5.1. This proves the following Proposition.

Proposition 5.6. *Let (M, ω, Φ) be a quasi-Hamiltonian G -space. Suppose (\mathcal{G}, B, γ) is a bundle gerbe over G with connective structure whose 3-curvature is η , and let $(Q, \gamma_Q) : \mathcal{I}_{-\omega} \dashrightarrow \Phi^*(\mathcal{G}, B, \gamma)$ be a trivialization. Then the diagram*

$$\begin{array}{ccc} \mathbb{L}(M, -d\omega) & \dashrightarrow & \mathbb{X}(\mathcal{I}_{-\omega}) \\ \parallel & \not\cong & \downarrow \mathbf{Q} \\ \mathbb{L}(M, \Phi^*\eta) & \dashrightarrow & \mathbb{X}(\Phi^*(\mathcal{G}; B, \gamma)) \end{array}$$

2-commutes, where the horizontal morphisms are the invertible butterflies from Theorem 5.1, and \mathbf{Q} is as in Theorem 5.3.

In [7], the authors show that a G -equivariant extension of a closed G -invariant 3-form χ on a G -manifold M gives rise to a homotopy moment map $\mathfrak{g} \rightarrow \mathbb{L}(M, \chi)$. For a quasi-Hamiltonian G -space (M, ω, Φ) , we may view $(M, -d\omega)$ as a pre-2-plectic manifold. From condition (i) of Definition 5.4, it follows that $\mathbb{L}(M, -d\omega) = \mathbb{L}(M, \Phi^*\eta)$, and condition (ii) shows that the two G -equivariant extensions of $-d\omega$,

$$-(d_G\omega)(\xi) = -d\omega + \iota_{\xi_M}\omega, \quad \text{and} \quad (\Phi^*\eta_G)(\xi) = \Phi^*\eta + \frac{1}{2}\Phi^*\langle\theta^L + \theta^R, \xi\rangle$$

agree. Therefore, the resulting homotopy moment maps agree. Combining this observation with Proposition 5.6, we obtain the diagram in the following Theorem, which we consider a quasi-Hamiltonian analogue of the Kostant formula (5.1). In other words, we may lift the \mathfrak{g} -actions on M and G to infinitesimal symmetries on $\mathcal{I}_{-\omega}$ and (\mathcal{G}, B, γ) that are compatible with the given trivialization (Q, γ_Q) .

Theorem 5.7. *Let (M, ω, Φ) be a quasi-Hamiltonian G -space. Suppose (\mathcal{G}, B, γ) is a bundle gerbe over G with connective structure whose 3-curvature is η , and let $(Q, \gamma_Q) : \mathcal{I}_{-\omega} \dashrightarrow \Phi^*(\mathcal{G}, B, \gamma)$ be a trivialization. The following diagram 2-commutes:*

$$\begin{array}{ccc} \mathfrak{g} & \begin{array}{l} \nearrow \\ \searrow \end{array} & \begin{array}{ccc} \mathbb{L}(M, -d\omega) & \dashrightarrow & \mathbb{X}(\mathcal{I}_{-\omega}) \\ \parallel & \not\cong & \downarrow \mathbf{Q} \\ \mathbb{L}(M, \Phi^*\eta) & \dashrightarrow & \mathbb{X}(\Phi^*(\mathcal{G}; B, \gamma)) \end{array} \end{array}$$

APPENDIX A. A FORMULA USED IN THE PROOF OF THEOREM 5.1

We include the verification of a formula used in the proof of Theorem 5.1.

Proposition A.1. *Let N be a manifold, and let $B \in \Omega^2(N)$. Let X, Y , and Z denote vector fields on N , and suppose that there exist differential 1-forms a, b , and c such that $L_X B = da$, $L_Y B = db$, and $L_Z B = dc$. Then,*

$$-\iota_Z \iota_Y \iota_X dB = \iota_X (L_Y c - L_Z b) - \iota_{[Y, Z]} a + \iota_{[Y, Z]} \iota_X B + \text{cyc. perm.},$$

Proof. This is a straightforward application of the given relations and the Cartan calculus. Indeed,

$$\begin{aligned} \iota_X \iota_Y \iota_Z dB &= \iota_X \iota_Y L_Z B - \iota_X \iota_Y d \iota_Z B \\ &= \iota_X \iota_Y dc - \iota_X L_Y \iota_Z B + \iota_X d \iota_Y \iota_Z B \\ &= \iota_X \iota_Y dc - \iota_X \iota_{[Y, Z]} B - \iota_X \iota_Z L_Y B + \iota_X d \iota_Y \iota_Z B \\ &= \iota_X \iota_Y dc - \iota_X \iota_{[Y, Z]} B - L_X \iota_Y \iota_Z B \\ &= \iota_X \iota_Y dc - \iota_X \iota_Z db - \iota_X \iota_{[Y, Z]} B + \iota_{[X, Y]} \iota_Z B + \iota_Y L_X \iota_Z B \\ &= \iota_X \iota_Y dc - \iota_X \iota_Z db - \iota_X \iota_{[Y, Z]} B + \iota_{[X, Y]} \iota_Z B + \iota_Y \iota_{[X, Z]} B + \iota_Y \iota_Z da \\ &= \iota_X L_Y c - L_X \iota_Y c - \iota_X L_Z b + L_X \iota_Z b + \iota_Y L_Z a - L_Y \iota_Z a - \iota_X \iota_{[Y, Z]} B \\ &\quad - \iota_Z \iota_{[X, Y]} B - \iota_Y \iota_{[Z, X]} B \\ &= \iota_X L_Y c - \iota_{[X, Y]} c - \iota_Y L_X c - \iota_X L_Z b + \iota_{[X, Z]} b + \iota_Z L_X b + \iota_Y L_Z a - \iota_{[Y, Z]} a \\ &\quad - \iota_Z L_Y a - \iota_X \iota_{[Y, Z]} B - \iota_Z \iota_{[X, Y]} B - \iota_Y \iota_{[Z, X]} B \end{aligned}$$

□

REFERENCES

- [1] Anton Alekseev, Anton Malkin, and Eckhard Meinrenken. Lie group valued moment maps. *Journal of Differential Geometry*, 48(3):445–495, 1998.
- [2] John Baez and Alissa Crans. Higher-dimensional algebra VI: Lie 2-algebras. *Theory Appl. Categ.*, 12(15):492–528, 2004.
- [3] John C. Baez and Urs Schreiber. Higher gauge theory. *Contemporary Mathematics*, 431:7–30, 2007.
- [4] Kai Behrend and Ping Xu. Differentiable stacks and gerbes. *Journal of Symplectic Geometry*, 9(3):285–341, 2011.
- [5] Daniel Berwick-Evans and Eugene Lerman. Lie 2-algebras of vector fields. *arXiv:1609.03944*, 2016.
- [6] Jean-Luc Brylinski. *Loop spaces, characteristic classes and geometric quantization*, volume 107 of *Progress in mathematics*. Birkhäuser Boston, 1993.
- [7] Martin Callies, Yael Fregier, Christopher L Rogers, and Marco Zambon. Homotopy moment maps. *Advances in Mathematics*, 303:954–1043, 2016.
- [8] David Saumitra Chatterjee. *On the construction of abelian gerbs*. PhD thesis, University of Cambridge, 1998.
- [9] Braxton L Collier. Infinitesimal symmetries of Dixmier-Douady gerbes. *arXiv:1108.1525*, 2011.
- [10] Braxton L Collier. *Infinitesimal symmetries of Dixmier-Douady gerbes*. PhD thesis, University of Texas at Austin, 2012.

- [11] Marius Crainic. Differentiable and algebroid cohomology, van Est isomorphisms, and characteristic classes. *Commentarii Mathematici Helvetici*, 78(4):681–721, 2003.
- [12] Jacques Dixmier and Adrien Douady. Champs continus d’espaces Hilbertiens et de C^* -algèbres. *Bulletin de la Société mathématique de France*, 91:227–284, 1963.
- [13] Domenico Fiorenza, Christopher L Rogers, and Urs Schreiber. L_∞ -algebras of local observables from higher prequantum bundles, 2013; e-print. *Homology, Homotopy and Applications*, 16(2):107–142, 2014.
- [14] Krzysztof Gawędzki and Nuno Reis. Basic gerbe over non-simply connected compact groups. *Journal of Geometry and Physics*, 50(1):28–55, 2004.
- [15] Jean Giraud. *Cohomologie non abélienne*. Springer-Verlag, Berlin-New York, 1971. Die Grundlehren der mathematischen Wissenschaften, Band 179.
- [16] Richard Hepworth. Vector fields and flows on differentiable stacks. *Theory and Applications of Categories*, 22(21):542–587, 2009.
- [17] Nigel Hitchin. Lectures on special lagrangian submanifolds. *AMS IP Studies in Advanced Mathematics*, 23:151–182, 2001.
- [18] Bertram Kostant. Quantization and unitary representations. In *Lectures in Modern Analysis and Applications III*, pages 87–208. Springer, 1970.
- [19] Derek Krepski. Pre-quantization of the moduli space of flat G -bundles over a surface. *Journal of Geometry and Physics*, 58(11):1624–1637, 2008.
- [20] Derek Krepski. Prequantization of the moduli space of flat $PU(p)$ -bundles with prescribed boundary holonomies. *SIGMA. Symmetry, Integrability and Geometry: Methods and Applications*, 10:109, 2014.
- [21] Derek Krepski. Basic equivariant gerbes on non-simply connected compact simple Lie groups. *J. Geom. Phys.*, 133:30–41, 2018.
- [22] Derek Krepski and Eckhard Meinrenken. On the Verlinde formulas for $SO(3)$ -bundles. *Quarterly Journal of Mathematics*, 64(1):235–252, 2013.
- [23] Derek Krepski and Jordan Watts. Differential cocycles and Dixmier–Douady bundles. *Journal of Geometry and Physics*, 130:168–183, 2018.
- [24] Kirill CH Mackenzie and Ping Xu. Classical lifting processes and multiplicative vector fields. *Quarterly Journal of Mathematics*, 49(193):59–85, 1998.
- [25] Eckhard Meinrenken. The basic gerbe over a compact simple Lie group. *L’Enseignement Mathématique*, 49:307–333, 2003.
- [26] Eckhard Meinrenken. Twisted K-homology and group-valued moment maps. *International Mathematics Research Notices*, 2012(20):4563–4618, 2012.
- [27] Michael K. Murray. Bundle gerbes. *Journal of the London Mathematical Society*, 54(2):403–416, 1996.
- [28] Michael K Murray, David Michael Roberts, Danny Stevenson, and Raymond F Vozzo. Equivariant bundle gerbes. *Advances in Theoretical and Mathematical Physics*, 21(4):921–975, 2017.
- [29] Michael K Murray and Daniel Stevenson. Bundle gerbes: stable isomorphism and local theory. *Journal of the London Mathematical Society*, 62(03):925–937, 2000.
- [30] Thomas Nikolaus and Christoph Schweigert. Equivariance in higher geometry. *Advances in Mathematics*, 226(4):3367–3408, 2011.
- [31] Thomas Nikolaus and Konrad Waldorf. Four equivalent versions of nonabelian gerbes. *Pacific Journal of Mathematics*, 264(2):355–420, 2013.
- [32] Behrang Noohi. Integrating morphisms of Lie 2-algebras. *Compositio Mathematica*, 149(2):264–294, 2013.
- [33] Christopher L Rogers. *Higher symplectic geometry*. PhD thesis, University of California, Riverside, 2011.

- [34] Christopher L Rogers et al. 2-plectic geometry, Courant algebroids, and categorified prequantization. *Journal of Symplectic Geometry*, 11(1):53–91, 2013.
- [35] Daniel Stevenson. *The geometry of bundle gerbes*. PhD thesis, University of Adelaide, 2000.
- [36] Mathieu Stiénon. Equivariant Dixmier-Douady classes. *Mathematical Research Letters*, 17(1):127–145, 2010.
- [37] Jennifer Vaughan. Metaplectic-c quantomorphisms. *Symmetry, integrability and geometry: methods and applications*, 11:25–15, 2015.
- [38] André Weil. Sur les théorèmes de de Rham. *Comment. Math. Helv.*, 26:119–145, 1952.
- [39] Christoph Wockel. Principal 2-bundles and their gauge 2-groups. *Forum mathematicum*, 23(3):565–610, 2011.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANITOBA, WINNIPEG, MB, CANADA

E-mail address: `Derek.Krepski@umanitoba.ca`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANITOBA, WINNIPEG, MB, CANADA

E-mail address: `Jennifer.Vaughan@umanitoba.ca`