

BEILINSON-DRINFELD SCHUBERT VARIETIES AND GLOBAL DEMAZURE MODULES

ILYA DUMANSKI, EVGENY FEIGIN, AND MICHAEL FINKELBERG

ABSTRACT. We compute the spaces of sections of powers of the determinant line bundle on the spherical Schubert subvarieties of the Beilinson-Drinfeld affine Grassmannians. The answer is given in terms of global Demazure modules over the current Lie algebra.

INTRODUCTION

Let \mathfrak{g} be a simple complex Lie algebra. To simplify the notation, in the introduction we assume that \mathfrak{g} is simply-laced. We drop this restriction in the main body of the paper.

The central objects of the algebraic representation theory of \mathfrak{g} are finite-dimensional irreducible representations V_λ of \mathfrak{g} labeled by the dominant integral weights $\lambda \in P_+$. The geometric objects responsible for these representations are the flag varieties. In particular, flag varieties are naturally embedded into projectivizations of irreducible \mathfrak{g} -modules and the celebrated Borel-Weil theorem states that finite-dimensional \mathfrak{g} modules are realized as (dual) spaces of sections of line bundles on flag varieties (see e.g. [Fu, Kum2]). These properties are still valid after passing to the Demazure submodules inside V_λ and to the Schubert subvarieties in flag varieties.

In this paper we are interested in the representation theory (algebraic and geometric) of two natural infinite-dimensional analogues of the Lie algebra \mathfrak{g} – the current algebra $\mathfrak{g}[t] = \mathfrak{g} \otimes \mathbb{C}[t]$ and the (untwisted) affine Kac-Moody Lie algebra $\widehat{\mathfrak{g}}$ with the natural embedding $\mathfrak{g}[t] \subset \widehat{\mathfrak{g}}$. The $\widehat{\mathfrak{g}}$ -analogues of the \mathfrak{g} -modules V_λ are (infinite-dimensional) integrable highest weight representations $L(\Lambda)$ (see [Kac]). The central element of $\widehat{\mathfrak{g}}$ acts on $L(\Lambda)$ by a constant called the level of representation. In particular, there are finitely many level one integrable modules $L(\Lambda_i)$, $i = 0, \dots, m$, where $L(\Lambda_0)$ is the basic representation. In this paper we will only consider modules $L(\ell\Lambda_i)$ for $\ell \in \mathbb{Z}_{>0}$. The projectivization $\mathbb{P}(L(\Lambda_i))$ contains a partial affine flag variety $\text{Gr}(\Lambda_i)$ as the closure of the orbit of the highest weight line with respect to the action of the affine Kac-Moody group ([Kum2]). The disjoint union $\sqcup_{i=0}^m \text{Gr}(\Lambda_i)$ is isomorphic to the affine Grassmannian Gr for the adjoint Lie group of the Lie algebra \mathfrak{g} .

The Demazure submodules in integrable representations $L(\Lambda)$ are labeled by the elements of the extended affine Weyl group. We will only consider the $\mathfrak{g}[t]$ -invariant Demazure modules inside $L(\ell\Lambda_i)$ (note that in general a

Demazure module is only acted upon by the Iwahori subalgebra that is strictly contained in the current algebra). In particular, the $\mathfrak{g}[t]$ -invariant Demazure modules $D_{1,\lambda}$ inside the level one integrable representations are labeled by the dominant integral weights λ . We denote by $D_{\ell,\lambda}$, $\ell \geq 1$ the level ℓ affine Demazure modules contained in the ℓ -th tensor power of level one module $D_{1,\lambda}$. The projectivized Demazure module $\mathbb{P}(D_{1,\lambda})$ contains the spherical Schubert variety $\overline{\text{Gr}}^\lambda$ as the orbit closure of the current group action. Thanks to the embedding $\overline{\text{Gr}}^\lambda \subset \mathbb{P}(D_{1,\lambda})$ the Schubert varieties are equipped with ample line bundle \mathcal{L} , such that the dual space of sections of $\mathcal{L}^{\otimes \ell}$ is isomorphic to $D_{\ell,\lambda}$ for any ℓ . The line bundle \mathcal{L} on a Schubert variety can be also obtained as the restriction of the determinant line bundle on the affine Grassmannian (see [Kum2, Z2]).

The current algebra $\mathfrak{g}[t]$ possesses a remarkable family of cyclic finite-dimensional modules W_λ called the local Weyl modules (see [CL, CP, FL2, KN, Naoi]). In particular, as \mathfrak{g} -module, W_λ is isomorphic to the tensor product of fundamental local Weyl modules, where the number of factors of the form W_ω is exactly the coefficient of ω in the decomposition of λ . We note that in the simply-laced case one has an isomorphism $W_\lambda \simeq D_{1,\lambda}$. The global Weyl modules \mathbb{W}_λ are infinite-dimensional cyclic representations of $\mathfrak{g}[t]$ (see [BF1, CFK, CI, FeMa1, Kato1]). One of the most important properties of the global Weyl modules is the existence of free action of the commutative highest weight algebra \mathcal{A}_λ , commuting with the $\mathfrak{g}[t]$ -action. In particular, one obtains a family of (finite-dimensional) $\mathfrak{g}[t]$ -modules, labeled by the closed points in $\text{Spec}(\mathcal{A}_\lambda)$, obtained as fibers of \mathbb{W}_λ with respect to \mathcal{A}_λ ; the local Weyl module is the fiber at the origin.

A generalization of this picture was suggested in [DF]. The authors introduced a family of cyclic (infinite-dimensional) global Demazure modules $\mathbb{D}(\ell, \underline{\lambda})$ (that were denoted $R(D_{\ell,\lambda_1}, \dots, D_{\ell,\lambda_k})$ in [DF]) corresponding to a collection of dominant integral nonzero weights $\underline{\lambda} \in P_+^k$ and an integer $\ell > 0$; in particular, if all λ_i are fundamental and $\ell = 1$, then one gets back the global Weyl module (this is no longer true in the non simply-laced case). The global Demazure modules arise naturally in connection with the study of the projective arc spaces (see [Mu1, Mu2, Nash]). The modules $\mathbb{D}(\ell, \underline{\lambda})$ are acted upon by a commutative (highest weight) algebra $\mathcal{A}(\underline{\lambda}) = \mathcal{A}(\lambda_1, \dots, \lambda_k)$ whose action commutes with the $\mathfrak{g}[t]$ action (see [BCES, EGL, KMSV, SV] for the examples of similar algebras). The spectrum $\text{Spec}(\mathcal{A}(\underline{\lambda}))$ is the closure of a stratum of the diagonal stratification of a colored configuration space of the affine line (see section 2.1 for precise definitions). In particular, a closed point $\mathbf{c} \in \mathbb{A}^k$ defines the same named closed point in $\text{Spec}(\mathcal{A}(\underline{\lambda}))$. For a point $\mathbf{c} \in \text{Spec}(\mathcal{A}(\underline{\lambda}))$ we denote by $\mathbb{D}(\ell, \underline{\lambda})_{\mathbf{c}}$ the fiber of the global Demazure module at \mathbf{c} . Our first theorem is as follows:

Theorem A. (a) *Assume that $\lambda_i \neq 0$ for all i . One has an isomorphism of $\mathfrak{g}[t]$ -modules*

$$\mathbb{D}(\ell, \underline{\lambda})_0 \simeq D_{\ell, \lambda_1 + \dots + \lambda_k}.$$

(b) Let $\underline{\lambda} \in P_+^k$, $\underline{\mu} \in P_+^l$. If $\mathbf{c} \in \mathbb{A}^k$ and $\mathbf{d} \in \mathbb{A}^l$ have no common entries, then the following factorization property (an isomorphism of $\mathfrak{g}[t]$ -modules) holds:

$$\mathbb{D}(\ell, \underline{\lambda} \sqcup \underline{\mu})_{(\mathbf{c}, \mathbf{d})} \simeq \mathbb{D}(\ell, \underline{\lambda})_{\mathbf{c}} \otimes \mathbb{D}(\ell, \underline{\mu})_{\mathbf{d}}.$$

(c) The global Demazure module $\mathbb{D}(\ell, \underline{\lambda})$ is free over $\mathcal{A}(\underline{\lambda})$.

(d) The direct sum of $\mathcal{A}(\underline{\lambda})$ -dual modules $\bigoplus_{\ell \geq 0} \mathbb{D}(\ell, \underline{\lambda})^\vee$ carries a natural structure of $\mathcal{A}(\underline{\lambda})$ -algebra.

The properties of the global Demazure modules collected in Theorem A are parallel to the properties of the Beilinson-Drinfeld spherical Schubert varieties over the affine line. The main goal of this paper is to describe this relation explicitly. More precisely, we:

- identify the projective spectrum of the algebra $\bigoplus_{\ell \geq 0} \mathbb{D}(\ell, \underline{\lambda})^\vee$ with the partially symmetrized BD spherical Schubert varieties;
- embed symmetrized BD spherical Schubert varieties into the projectivization of the vector bundle $\mathcal{D}(\ell, \underline{\lambda})$ obtained as the localization of the (free) $\mathcal{A}(\underline{\lambda})$ -module $\mathbb{D}(\ell, \underline{\lambda})$;
- identify the dual sections of the determinant line bundle on symmetrized BD spherical Schubert varieties with global Demazure modules.

Let us state our results in more detail. Recall that the Beilinson-Drinfeld Grassmannians (BD Grassmannians for short) are global versions of the affine Grassmannians [BD1, BD2, FBZ, Z2] defined over the powers of an algebraic curve X ; in this paper we only consider the case $X = \mathbb{A}^1$ and denote the corresponding BD Grassmannians by $\text{Gr}_{\mathbb{A}^k}$ (see e.g. [BKK, CK, CW, Kam, MVy] for various applications in geometric representation theory). The Grassmannians $\text{Gr}_{\mathbb{A}^k}$ are ind-varieties over the configuration space \mathbb{A}^k , and the ind-structure is provided by the BD spherical Schubert varieties $\overline{\text{Gr}}^\lambda$, labeled by k -tuples of dominant coweights $\underline{\lambda} = (\lambda_1, \dots, \lambda_k) \in P_+^k$. A group scheme $\mathbb{G}(k)$ over \mathbb{A}^k (the global analogue of the current group $G(\mathbb{C}[[t]])$) acts on $\text{Gr}_{\mathbb{A}^k}$ fiberwise, and the BD spherical Schubert varieties are the closures of orbits of $\mathbb{G}(k)$ in the generic fiber of $\text{Gr}_{\mathbb{A}^k}$ (we note that the same group scheme acts on $\mathbb{P}(\mathcal{D}(\ell, \underline{\lambda}))$). The fibers of the projection $\overline{\text{Gr}}^\lambda \rightarrow \mathbb{A}^k$ are products of the spherical Schubert subvarieties of the affine Grassmannian (this is a manifestation of the crucial factorization property of the BD Grassmannians).

BD Grassmannians carry the ample determinant line bundle \mathcal{L} ; we keep the same notation for the restriction of this line bundle to the BD spherical Schubert varieties. The space of sections $H^0(\overline{\text{Gr}}^\lambda, \mathcal{L}^{\otimes \ell})$ is naturally a $\mathfrak{g}[t] - \mathbb{C}[\mathbb{A}^k]$ -bimodule (we note that the higher cohomology $H^{>0}(\overline{\text{Gr}}^\lambda, \mathcal{L}^{\otimes \ell})$ vanish). However, as a module over the current algebra it is not cyclic and hence hard to describe. In order to resolve this problem we consider a partially symmetrized version $\overline{\text{Gr}}^{(\underline{\lambda})}$ of the BD spherical Schubert varieties (see Section 2 for precise definition). The variety $\overline{\text{Gr}}^{(\underline{\lambda})}$ is equipped with

a natural projection onto $\mathrm{Spec}(\mathcal{A}(\underline{\lambda}))$ and the determinant line bundle descends to the symmetrized BD spherical Schubert varieties. We prove the following theorem.

Theorem B. *For all $\ell \geq 1$ and $\underline{\lambda} = (\lambda_1, \dots, \lambda_k) \in P_+^k$, such that all λ_i are nonzero, one has:*

(a) *a $\mathbb{G}(k)$ -equivariant embedding:*

$$\overline{\mathrm{Gr}}^{(\underline{\lambda})} \subset \mathbb{P}(\mathcal{D}(\ell, \underline{\lambda}));$$

(b) *an isomorphism of $\mathrm{Spec}(\mathcal{A}(\underline{\lambda}))$ -schemes:*

$$\overline{\mathrm{Gr}}^{(\underline{\lambda})} \simeq \mathrm{Proj}\left(\bigoplus_{\ell \geq 0} \mathbb{D}(\ell, \underline{\lambda})^\vee\right);$$

(c) *an isomorphism of $\mathfrak{g}[t] - \mathcal{A}(\underline{\lambda})$ -bimodules:*

$$H^0(\mathrm{Gr}^{(\underline{\lambda})}, \mathcal{L}^{\otimes \ell}) \simeq \mathbb{D}(\ell, \underline{\lambda})^\vee;$$

(d) *an isomorphism of $\mathfrak{g}[t] - \mathbb{C}[\mathbb{A}^k]$ -bimodules:*

$$H^0(\mathrm{Gr}^{\underline{\lambda}}, \mathcal{L}^{\otimes \ell}) \simeq \mathbb{D}(\ell, \underline{\lambda})^\vee \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}[\mathbb{A}^k],$$

where M^\vee stands for the $\mathcal{A}(\underline{\lambda})$ -dual to an $\mathcal{A}(\underline{\lambda})$ -module M .

Let us close with the following remark. In the main body of the paper, we denote the weights and roots of \mathfrak{g} by the checked letters (like λ^\vee and α^\vee) and reserve the non-checked notation for the dual data (coroots and coweights). The reason is that the central role in our paper is played by the spherical Schubert varieties in the affine (Beilinson-Drinfeld) Grassmannians. These varieties are naturally labeled by the coweights (rather than weights), which explains our choice of notation. Note that in the ADE case all the checks can be removed without any harm. Also, in the simply laced case, if $\lambda_1, \dots, \lambda_k$ are all fundamental, and $\lambda = \lambda_1 + \dots + \lambda_k$, the global Demazure modules $\mathbb{D}(1, \underline{\lambda})$ are nothing but the global Weyl modules \mathbb{W}_λ . However, if \mathfrak{g} is not simply laced, there is no such coincidence anymore. That is why we choose to call $\mathbb{D}(\ell, \underline{\lambda})$ global Demazure modules as opposed to “higher-level global Weyl modules”.

Our paper is organized as follows. In Section 1 we collect notation and recall main definitions. In Section 2 we introduce the symmetrized version of the Beilinson-Drinfeld Grassmannians and Schubert varieties over the spectrum of the highest weight algebras. In Section 3 we study the properties of the global Demazure modules. In particular, we prove that they are free over the highest weight algebras. In Section 4 we compute the spaces of sections of the powers of the determinant line bundle on BD Schubert varieties. In Appendix, we discuss a connection between global modules and the associativity of the fusion product. We also collect the key objects of the paper.

ACKNOWLEDGMENTS

The authors would like to thank Syu Kato for his suggestion to study the algebra of dual global Demazure modules. We are indebted to Roman Travkin for his careful reading, corrections, and improvements of the first draft. We are also grateful to Alexander Braverman and Boris Feigin for useful discussions. Finally, our thanks go to the anonymous referee for a very careful reading of the manuscript and many useful suggestions. The work was partially supported by the grant RSF 19-11-00056. I.D. is supported in part by the Simons Foundation.

1. GENERALITIES

We start with describing the notation for the key objects of the paper.

1.1. Classical objects. Let \mathfrak{g} be a simple Lie algebra over \mathbb{C} . The corresponding simply connected (resp. adjoint) complex Lie group will be denoted G^{sc} (resp. G^{ad}). Let $\mathfrak{g} = \mathfrak{n}_+ \oplus \mathfrak{h} \oplus \mathfrak{n}_-$ be the Cartan decomposition and let $r = \dim \mathfrak{h}$ be the rank of \mathfrak{g} . We denote by $\omega_1^\vee, \dots, \omega_r^\vee$ the fundamental weights of \mathfrak{g} and by $\alpha_1^\vee, \dots, \alpha_r^\vee$ its simple roots. Let $P^\vee = \bigoplus_{i=1}^r \mathbb{Z} \omega_i^\vee$ be the weight lattice of G^{sc} containing the root lattice $Q^\vee = \bigoplus_{i=1}^r \mathbb{Z} \alpha_i^\vee$ (that coincides with the weight lattice of G^{ad}). Let $P_+^\vee = \bigoplus_{i=1}^r \mathbb{Z}_{\geq 0} \omega_i^\vee \subset P^\vee$ be the set of dominant integral weights. Given $\lambda^\vee = \sum_{i=1}^r m_i \omega_i^\vee \in P_+^\vee$ we set $|\lambda^\vee| = \sum_{i=1}^r m_i$.

For a weight $\lambda^\vee \in P_+^\vee$, let V_{λ^\vee} be the highest weight λ^\vee irreducible \mathfrak{g} -module; in particular, the highest weight vector of V_{λ^\vee} is of the \mathfrak{h} weight λ^\vee and is killed by \mathfrak{n}_+ . Let W be the (finite) Weyl group of \mathfrak{g} with the longest element w_0 . In particular, the lowest weight vector in V_{λ^\vee} is of weight $w_0 \lambda^\vee$.

We denote by $P = P_{\text{ad}}$ (resp. $Q = P_{\text{sc}}$) the coweight lattice of G^{ad} (resp. of G^{sc}). Thus we have perfect pairings $P_{\text{ad}} \times Q^\vee \rightarrow \mathbb{Z}$, $P_{\text{sc}} \times P^\vee \rightarrow \mathbb{Z}$. The minimal invariant integral bilinear form on P_{sc} (such that the square length of a short coroot is 2) gives rise to a linear map $\iota: P_{\text{sc}} \rightarrow Q^\vee$. It extends by linearity to the same named map $P_{\text{sc}} \subset P_{\text{ad}} \xrightarrow{\iota} Q^\vee \otimes_{\mathbb{Z}} \mathbb{Q}$, and $\iota(P_{\text{ad}}) \subset P^\vee \subset Q^\vee \otimes_{\mathbb{Z}} \mathbb{Q}$. The resulting map $P = P_{\text{ad}} \rightarrow P^\vee$ will be also denoted by ι . In the simply laced case $\iota: P \rightarrow P^\vee$ is an isomorphism.

The fundamental coweights in P are denoted $\omega_1, \dots, \omega_r$; the simple coroots in P are denoted by $\alpha_1, \dots, \alpha_r$. We set $P_+ = \bigoplus_{i=1}^r \mathbb{Z}_{\geq 0} \omega_i \subset P$.

1.2. Current algebra modules. Let $\mathfrak{g}[t] = \mathfrak{g} \otimes \mathbb{C}[t]$ be the current algebra of \mathfrak{g} . In what follows we consider graded $\mathfrak{g}[t]$ -modules M , i.e. $M = \bigoplus_{i \geq 0} M_i$, $\mathfrak{g} \otimes t^k: M_i \rightarrow M_{i+k}$. If all M_i are finite-dimensional, then the graded character $\text{ch}_q(M)$ is a generating function $\sum_{i \geq 0} q^i \text{ch} M_i$ of the characters of the $\mathfrak{g} \otimes 1$ -modules.

A module M is called cyclic if it is generated by a single vector. The cyclic product of two cyclic $\mathfrak{g}[t]$ -modules M_1 and M_2 with fixed cyclic vectors

$w_1 \in M_1$ and $w_2 \in M_2$ is defined as

$$M_1 \odot M_2 = U(\mathfrak{g}[t]).w_1 \otimes w_2 \subset M_1 \otimes M_2.$$

For a collection of pairwise distinct complex numbers c_1, \dots, c_n and cyclic graded $\mathfrak{g}[t]$ -modules M_1, \dots, M_n the module $M_1(c_1) \otimes \dots \otimes M_n(c_n)$ is known to be cyclic with the cyclic vector being the tensor product of cyclic vectors of M_i . Here a $\mathfrak{g}[t]$ -module $M_i(c_i)$ is defined to be isomorphic to M_i as a vector space and the action of the current algebra on it is twisted by the automorphism $x \otimes t^i \mapsto x \otimes (t - c)^i$. We note that if one starts with graded cyclic modules M_i , the tensor product $\bigotimes_{i=1}^n M_i(c_i)$ is not graded in general. The fusion product (graded tensor product) $M_1 * \dots * M_k$ is defined as the associated graded of $\bigotimes_{i=1}^n M_i(c_i)$ with respect to the filtration induced by the action of the (t -degree graded) universal enveloping algebra $U(\mathfrak{g}[t])$ on the tensor product of cyclic vectors of M_i ([FeLo]).

Let $\mathbb{W}_{\lambda^\vee}$ and W_{λ^\vee} be the global and local Weyl modules of highest weight λ^\vee over the Lie algebra $\mathfrak{g}[t]$ (see [CP, CL, FL2, Kato1, Naoi]). Let D_λ be the level one affine Demazure module with highest weight $\iota(\lambda)$; in particular, for simply laced algebras ι is an isomorphism and $W_{\iota(\lambda)} \simeq D_\lambda$ for any coweight λ . For $\lambda \in P_+$ we denote $\mathbb{W}_{\iota(\lambda)}$ (resp. $W_{\iota(\lambda)}$) simply by \mathbb{W}_λ (resp. W_λ).

For $\ell \in \mathbb{Z}_{>0}$ we denote by $D_{\ell, \lambda}$ the level ℓ affine Demazure module with highest weight $\ell\iota(\lambda)$ (see subsection 1.3 for details).

Let M_1, \dots, M_k be graded cyclic $\mathfrak{g}[t]$ -modules with cyclic vectors w_i of dominant nonzero weights such that $t\mathfrak{h}[t]$ annihilates their cyclic vectors. Then we define the global module (see [DF])

$$R(M_1, \dots, M_k) = M_1[t] \odot \dots \odot M_k[t],$$

where $M_i[t]$ is defined as a module isomorphic to $M_i \otimes \mathbb{C}[t]$ as a vector space with the action of $\mathfrak{g}[t]$ given by

$$(1.1) \quad xt^l.v \otimes t^k = \sum_{j=0}^l (-1)^{l-j} \binom{l}{j} (xt^j.v) \otimes t^{l+k-j}$$

for $l, k \in \mathbb{Z}_{\geq 0}$, $x \in \mathfrak{g}$, $v \in M_i$.

Remark 1.1. The analogous formula used in [DF, FeMa2] has no sign $(-1)^{l-j}$. We introduce it here in order to match the formulas in the Beilinson-Drinfeld setup, where the minus sign pops up via the change of coordinates $t \mapsto t - x$. This sign change obviously produces no harm (simply changing $t \mapsto -t$ in the current algebra parametrization).

Remark 1.2. The modules $R(M_1, \dots, M_k)$ do depend on the choice of the cyclic vectors w_i of M_i .

The global module $R(M_1, \dots, M_k)$ admits the right action of $U(\mathfrak{h}[t])$, which commutes with the $\mathfrak{g}[t]$ -action. The highest weight algebra is defined as a quotient of $U(\mathfrak{h}[t])$ by the annihilator of the cyclic vector $\bigotimes_{i=1}^k w_i$ of $R(M_1, \dots, M_k)$. It turns out that the highest weight algebra depends only on

the weights of cyclic vectors w_i of M_i (not on a particular choice of modules). If the weight of w_i is $\iota(\lambda_i)$, then we denote the highest weight algebra of $R(M_1, \dots, M_k)$ by $\mathcal{A}(\lambda_1, \dots, \lambda_k)$. We will use a short-hand notation $\mathcal{A}(\underline{\lambda}) = \mathcal{A}(\lambda_1, \dots, \lambda_k)$, where $\underline{\lambda} = (\lambda_1, \dots, \lambda_k)$.

Since the weight $\iota(\lambda_i)$ subspace of a module $M_i[t]$ is isomorphic to a polynomial algebra in one variable, the algebra $\mathcal{A}(\underline{\lambda})$ is naturally embedded into $\bigotimes_{i=1}^k \mathcal{A}(\lambda_i) \simeq \mathbb{C}[z_1, \dots, z_k]$. More precisely, $\mathcal{A}(\underline{\lambda})$ is isomorphic to the subalgebra of the polynomial algebra $\mathbb{C}[z_1, \dots, z_k]$ generated by the polynomials

$$\langle \iota(\lambda_1), h \rangle z_1^l + \langle \iota(\lambda_2), h \rangle z_2^l + \dots + \langle \iota(\lambda_k), h \rangle z_k^l, \quad l \geq 1, h \in \mathfrak{h}.$$

Indeed, for $h \in \mathfrak{h}$ and $l > 0$ formula (1.1) gives

$$ht^l \cdot \bigotimes_{i=1}^k w_i = \sum_{i=1}^k \bigotimes_{j=1}^{i-1} w_i \otimes \langle \iota(\lambda_j), h \rangle w_j t^l \bigotimes_{j=i+1}^k w_j.$$

In particular, if all λ_i are fundamental coweights, $m_i = \#\{j : \lambda_j = \omega_i\}$, and $\lambda = \sum_{j=1}^k \lambda_j = \sum_{i=1}^r m_i \omega_i$, then

$$\mathcal{A}(\underline{\lambda}) \cong \mathbb{C}[z_1, \dots, z_k]^{S_{m_1} \times \dots \times S_{m_r}} =: \mathcal{A}_\lambda.$$

We note that

- $\mathcal{A}(\lambda_1, \dots, \lambda_k) \simeq \mathcal{A}(\ell\lambda_1, \dots, \ell\lambda_k)$ for any $\ell \in \mathbb{N}$.
- If \mathfrak{g} is simply laced, all weights $\lambda_1, \dots, \lambda_k$ are fundamental, and $\lambda = \sum_{i=1}^k \lambda_i$, then

$$R(D_{\lambda_1}, \dots, D_{\lambda_k}) \simeq \mathbb{W}_\lambda.$$

If all the coweights λ_i are fundamental, and they sum up to λ , then we denote by $\mathbb{D}_{\ell, \lambda}$ (global Demazure module) the module $R(D_{\ell, \lambda_1}, \dots, D_{\ell, \lambda_k})$. In particular, for simply-laced \mathfrak{g} one has $\mathbb{D}_{1, \lambda} \simeq \mathbb{W}_\lambda$.

If all the coweights λ_i are fundamental, then the algebra $\mathcal{A}(\lambda_1, \dots, \lambda_k)$ acts freely on $R(D_{\lambda_1}, \dots, D_{\lambda_k})$ and the fiber at the origin of the global Demazure module is isomorphic to the fusion product $D_{\lambda_1} * \dots * D_{\lambda_k} \simeq D_{\lambda_1 + \dots + \lambda_k}$. The higher level analogue still holds with fundamental λ_i replaced by $\ell\lambda_i$, see [DF].

As we will prove in Proposition 3.9, for arbitrary dominant coweights $\lambda_1, \dots, \lambda_k$, the module $R(D_{\ell, \lambda_1}, \dots, D_{\ell, \lambda_k})$ is free over $\mathcal{A}(\lambda_1, \dots, \lambda_k)$. We use the notation

$$\mathbb{D}(\ell, \underline{\lambda}) = R(D_{\ell, \lambda_1}, \dots, D_{\ell, \lambda_k}).$$

Remark 1.3. We note that $\mathbb{D}_{\ell, \lambda} = \mathbb{D}(\ell, \underbrace{\omega_1, \dots, \omega_1}_{m_1}, \dots, \underbrace{\omega_r, \dots, \omega_r}_{m_r})$ for a coweight $\lambda = \sum_{j=1}^r m_j \omega_j$.

In what follows we will need the following $\mathcal{A}(\underline{\lambda})$ -analog of the cyclic power. Namely, let

$$\underbrace{R(M_1, \dots, M_k) \odot_{\mathcal{A}(\underline{\lambda})} \dots \odot_{\mathcal{A}(\underline{\lambda})} R(M_1, \dots, M_k)}_{\ell}$$

be the $U(\mathfrak{g}[t])$ -span of the ℓ -th tensor power of the cyclic (highest weight) vector of $R(M_1, \dots, M_k)$ inside the ℓ -th tensor power over $\mathcal{A}(\underline{\lambda})$ of the module $R(M_1, \dots, M_k)$. We denote this cyclic tensor power by $R(M_1, \dots, M_k)_{\mathcal{A}(\underline{\lambda})}^{\odot \ell}$. This object will be important in Proposition 3.3.

1.3. Affine Lie algebras and Demazure modules. The details on the material below can be found in [Kac, Kum2].

Let $\widehat{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}K \oplus \mathbb{C}d$ be the untwisted affine Kac-Moody Lie algebra attached to \mathfrak{g} . Here K is central element and d is (negated) degree operator (i.e. $[d, x \otimes t^i] = -ix \otimes t^i$). The algebra $\widehat{\mathfrak{g}}$ enjoys the Cartan decomposition $\widehat{\mathfrak{g}} = \mathfrak{n}_+^a \oplus \mathfrak{h}^a \oplus \mathfrak{n}_-^a$, where $\mathfrak{h}^a = \mathfrak{h} \otimes 1 \oplus \mathbb{C}K \oplus \mathbb{C}d$ and $\mathfrak{n}_+^a = \mathfrak{g} \otimes t\mathbb{C}[t] \oplus \mathfrak{n}_+ \otimes 1$. We denote by $\mathfrak{b}^a = \mathfrak{h}^a \oplus \mathfrak{n}_+^a$ the Iwahori subalgebra.

Let Λ_0^\vee be the level one basic integrable weight of $\widehat{\mathfrak{g}}$ (in particular, $\Lambda_0^\vee(\mathfrak{h} \otimes 1) = 0$). We also denote by Λ_i^\vee , $i = 0, \dots, m$, the set of all integrable level one weights of $\widehat{\mathfrak{g}}$ and by $L(\Lambda_i^\vee)$ the corresponding highest weight $\widehat{\mathfrak{g}}$ -modules. In particular, the number m of the level one modules is equal to the cardinality of $P/Q \simeq \pi_1(G^{\text{ad}})$.

Let $\text{Gr}(\Lambda_i^\vee) \subset \mathbb{P}(L(\Lambda_i^\vee))$, $i = 0, \dots, m$, be the partial affine flag varieties corresponding to maximal parabolic subgroups of the affine Kac-Moody group \widehat{G}^{sc} (i.e. $\text{Gr}(\Lambda_i^\vee) \simeq \widehat{G}^{\text{sc}}/P_i$, where P_i is the stabilizer of the highest weight line in $\mathbb{P}(L(\Lambda_i^\vee))$). By the very definition, each $\text{Gr}(\Lambda_i^\vee)$ is equipped with the very ample line bundle \mathcal{L} (the pullback of $\mathcal{O}(1)$ from $\mathbb{P}(L(\Lambda_i^\vee))$) and one has the affine analog of the Borel-Weil theorem

$$H^0(\text{Gr}(\Lambda_i^\vee), \mathcal{L}^{\otimes \ell})^* \simeq L(\ell \Lambda_i^\vee), \quad \ell \geq 1,$$

where $L(\ell \Lambda_i^\vee)$ is the weight $\ell \Lambda_i^\vee$ integrable (level ℓ) $\widehat{\mathfrak{g}}$ -module and the superscript star denotes the restricted dual space.

Remark 1.4. The union $\sqcup_{i=0}^m \text{Gr}(\Lambda_i^\vee)$ is isomorphic to the affine Grassmannian of G^{ad} , see the details below.

Let $W^a = W \ltimes P$ be the extended affine Weyl group (recall that P is the coweight lattice of G^{ad}). Then for any $\lambda \in P_+$ there exists an element $w_\lambda \in W^a$ such that the \mathfrak{h} -weight of $w_\lambda \Lambda_0^\vee$ is equal to $w_0 \iota(\lambda)$. Let Λ_i^\vee be the unique integrable level one weight such that $w_\lambda \Lambda_0^\vee - \Lambda_i^\vee$ belongs to the root lattice of \mathfrak{g} . Let $u_{w_0 \iota(\lambda)} \in L(\Lambda_i^\vee)$ be a weight $w_0 \iota(\lambda)$ vector. We define the Demazure module $D_{1,\lambda} \subset L(\Lambda_i^\vee)$ as the $U(\mathfrak{b}^a)$ span of the vector $u_{w_0 \iota(\lambda)}$. An important property of the Demazure modules $D_{1,\lambda}$ is that they are invariant with respect to the whole current algebra $\mathfrak{g}[t] \supset \mathfrak{b}^a$. In particular, $D_{1,\lambda}$ contains the irreducible \mathfrak{g} -module $V_{\iota(\lambda)}$ as the $U(\mathfrak{n}_+)$ span of $u_{w_0 \iota(\lambda)}$.

The level ℓ Demazure module $D_{\ell,\lambda}$ is defined as the $U(\mathfrak{b}^a)$ span of the vector $u_{w_0 \iota(\lambda)}^{\otimes \ell}$. By definition, $D_{\ell,\lambda}$ is a subspace of $L(\Lambda_i^\vee)^{\otimes \ell}$. However, it is easy to see that

$$D_{\ell,\lambda} \subset L(\ell \Lambda_i^\vee) \subset L(\Lambda_i^\vee)^{\otimes \ell}.$$

By definition, one gets a natural structure of algebra on the space $D_{\bullet,\lambda}^* = \bigoplus_{\ell \geq 0} D_{\ell,\lambda}^*$ generated by the degree one component $D_{1,\lambda}^*$ (we set $D_{0,\lambda} = \mathbb{C}$).

We define a spherical Schubert variety $\overline{\text{Gr}}^\lambda$ as the closure of the $G^{\text{sc}}(\mathcal{O})$ -orbit of the line containing the lowest weight vector $u_{w_0\iota(\lambda)}$ (here $\mathcal{O} = \mathbb{C}[[t]]$). Then $\overline{\text{Gr}}^\lambda$ is embedded as a closed subscheme into the projectivization $\mathbb{P}(D_{1,\lambda})$ of the Demazure module $D_{1,\lambda}$ (see e.g. [Ma2, Théorème 2.Σ of Chapter X] or [Kum1]). Moreover, $\overline{\text{Gr}}^\lambda$ is also embedded as a closed subscheme into the projectivization of an arbitrary level Demazure module $D_{\ell,\lambda}$ as the closure of the lowest weight line.

Remark 1.5. Let $t^\lambda \in \overline{\text{Gr}}^\lambda \subset \mathbb{P}(D_{1,\lambda})$ be the point corresponding to the weight $\iota(\lambda)$ line. Then $\overline{\text{Gr}}^\lambda$ is the closure of the $G^{\text{sc}}(\mathcal{O})$ -orbit of t^λ .

The embedding $\overline{\text{Gr}}^\lambda \subset \mathbb{P}(D_{1,\lambda})$ endows $\overline{\text{Gr}}^\lambda$ with a very ample line bundle \mathcal{L} , the pullback of $\mathcal{O}(1)$. The line bundle \mathcal{L} is a generator of the Picard group of $\overline{\text{Gr}}^\lambda$, and one has the isomorphism of $\mathfrak{g}[t]$ -modules:

$$H^0(\overline{\text{Gr}}^\lambda, \mathcal{L}^{\otimes \ell})^* \simeq D_{\ell,\lambda} \text{ for all } \ell \geq 1.$$

We obtain a presentation of $\overline{\text{Gr}}^\lambda$ as the projective spectrum of the algebra of dual Demazure modules, i.e. $\overline{\text{Gr}}^\lambda \simeq \text{Proj}(\bigoplus_{\ell \geq 0} D_{\ell,\lambda}^*)$.

We have

$$\text{Gr}(\Lambda_i^\vee) = \bigcup_{\lambda: \Lambda_i^\vee - \iota(\lambda) \in Q^\vee + \Lambda_0^\vee} \overline{\text{Gr}}^\lambda.$$

Also, $\overline{\text{Gr}}^\lambda \subset \overline{\text{Gr}}^\mu$ if and only if $\mu - \lambda \in \bigoplus_{j=1}^r \mathbb{Z}_{\geq 0} \alpha_j$.

1.4. Affine Grassmannians. The affine Grassmannian of G^{ad} is $\text{Gr} := \text{Gr}_{G^{\text{ad}}} = G^{\text{ad}}(\mathcal{K})/G^{\text{ad}}(\mathcal{O})$, where $\mathcal{K} = \mathbb{C}((t))$ is the ring of Laurent series and $\mathcal{O} = \mathbb{C}[[t]]$ is the ring of Taylor series. The following properties of Gr can be found in [Z1, Z2, Kum2].

- The connected components of Gr are in bijection with P/Q , i.e. $\pi_0(\text{Gr}) \simeq \pi_1(G^{\text{ad}})$.
- $\text{Gr} = \sqcup_{i=0}^m \text{Gr}(\Lambda_i^\vee)$.
- For any $i = 0, \dots, m$, $\text{Pic}(\text{Gr}(\Lambda_i^\vee))$ is generated by the class of the ample determinant line bundle \mathcal{L} .

Recall (see, for example, [BL]) that the affine Grassmannian Gr is the moduli space of pairs (\mathcal{P}, β) , where

\mathcal{P} is a G^{ad} -torsor on \mathbb{A}^1 , $\beta: \mathcal{P}_{\mathbb{A}^1 \setminus 0} \rightarrow G \times (\mathbb{A}^1 \setminus 0)$ is a trivialization on $\mathbb{A}^1 \setminus 0$.

Replacing the point 0 with an arbitrary $c \in \mathbb{A}^1$, one gets a version Gr_c of the affine Grassmannian. Clearly, the isomorphism $\mathcal{O} \simeq \mathcal{O}_c = \mathbb{C}[[t - c]]$ induces the isomorphism $\text{Gr}_c \simeq \text{Gr}$ for any c . The schemes Gr_c glue together to the (trivial) bundle $\text{Gr}_{\mathbb{A}^1}$ over the affine line.

Remark 1.6. $\text{Gr}_{\mathbb{A}^1}$ is the simplest example of a Beilinson-Drinfeld Grassmannian, the general case is discussed in the next section.

2. BEILINSON-DRINFELD SCHUBERT VARIETIES

We will need several versions of the Beilinson-Drinfeld Schubert varieties (see [Z1, Z2]). Let us stress from the very beginning that the Beilinson-Drinfeld Grassmannians are defined over a (power of) a smooth curve X , but in this paper, we only consider the case $X = \mathbb{A}^1$. The standard Beilinson-Drinfeld definition produces schemes over affine spaces. We will also need the symmetrized versions with the natural projections to the spectrum of the highest weight algebras. So we first discuss the properties of the highest weight algebras and then introduce the symmetrized Beilinson-Drinfeld Schubert varieties.

2.1. The highest weight algebras. Let $\underline{\lambda} = (\lambda_1, \dots, \lambda_k)$ be a multiset of dominant coweights. Let $\lambda = \sum_{i=1}^k \lambda_i = \sum_{j=1}^r m_j \omega_j$ and $N = \sum_{j=1}^r m_j = |\lambda|$. We set

$$S_\lambda = S_{m_1} \times \dots \times S_{m_r}.$$

Recall the algebras $\mathcal{A}(\lambda_1, \dots, \lambda_k)$ and

$$\mathcal{A}_\lambda \simeq \mathcal{A}(\underbrace{\omega_1, \dots, \omega_1}_{m_1}, \dots, \underbrace{\omega_r, \dots, \omega_r}_{m_r}) \simeq \mathbb{C}[z_1, \dots, z_N]^{S_\lambda}$$

of Section 1.2.

Lemma 2.1. *There exists a natural surjection of algebras $\mathcal{A}_\lambda \twoheadrightarrow \mathcal{A}(\underline{\lambda})$.*

Proof. Note that $\mathcal{A}_\lambda = \mathcal{A}(\omega_{a_1}, \dots, \omega_{a_N})$, where $\sum_{i=1}^N \omega_{a_i} = \lambda$. Now it suffices to note that there exists a natural surjection

$$\mathcal{A}(\mu, \lambda_1, \dots, \lambda_k) \rightarrow \mathcal{A}(\mu + \lambda_1, \dots, \lambda_k)$$

induced by the surjection of the larger polynomial algebras

$$\mathbb{C}[z_1, \dots, z_{k+1}] \rightarrow \mathbb{C}[z_1, \dots, z_k], \quad z_1 \mapsto z_1, \quad z_2 \mapsto z_1, \quad z_i \mapsto z_{i-1}, \quad i > 2.$$

□

Let $\mathbb{A}^\lambda = \mathbb{A}^N / S_\lambda = \text{Spec } \mathcal{A}_\lambda$ be the space of configurations of colored points on the line \mathbb{A}^1 (m_i points of color ω_i). We have the main diagonal $\mathbb{A}^\lambda \supset \mathbb{A}^{(\lambda)} \simeq \mathbb{A}^1$ formed by all the configurations where all the points coincide. We have a finite morphism of addition of configurations

$$\text{add}: \mathbb{A}^\nu \times \mathbb{A}^\mu \rightarrow \mathbb{A}^{\nu+\mu}.$$

Iterating it we obtain

$$\text{add}: \mathbb{A}^{\lambda_1} \times \dots \times \mathbb{A}^{\lambda_k} \rightarrow \mathbb{A}^\lambda.$$

We define a closed subscheme $\mathbb{A}^{(\underline{\lambda})} \subset \mathbb{A}^\lambda$ as the add-image of the closed subscheme $\mathbb{A}^{(\lambda_1)} \times \dots \times \mathbb{A}^{(\lambda_k)} \subset \mathbb{A}^{\lambda_1} \times \dots \times \mathbb{A}^{\lambda_k}$:

$$\mathbb{A}^{(\underline{\lambda})} = \text{add}(\mathbb{A}^{(\lambda_1)} \times \dots \times \mathbb{A}^{(\lambda_k)}) \subset \mathbb{A}^\lambda.$$

Lemma 2.2. *One has*

$$\mathbb{C}[\mathbb{A}^{(\underline{\lambda})}] = \mathcal{A}(\underline{\lambda}).$$

Proof. We denote the coordinates on \mathbb{A}^N by $x_{i,j}$, where $i = 1, \dots, r$ and $j = 1, \dots, m_i$. The group S_λ acts by permuting the second indices. Let $\lambda_a = \sum_{b=1}^r m_{a,b} \omega_b$ for $a = 1, \dots, k$. In particular, $\sum_{a=1}^k m_{a,b} = m_b$ for all $b = 1, \dots, r$. Thus the coordinates with a fixed first index are divided into k groups. For each $a = 1, \dots, k$, we combine the corresponding groups for all the possible first indices into one big group Γ_a . Now all the coordinates in \mathbb{A}^N are divided into groups Γ_a , $1 \leq a \leq k$. We consider the linear subspace V in \mathbb{A}^N given by equations $x_{i,j} = x_{i',j'}$ whenever (i,j) and (i',j') lie in the same group Γ_a . We consider the saturation $S_\lambda V$ (a union of a few vector subspaces in \mathbb{A}^N). Finally, $\mathbb{A}^{(\Delta)} = (S_\lambda V)/S_\lambda$. Now the same argument as in the proof of [BCES, Proposition 2.2] finishes our proof. \square

Remark 2.3. By construction, $\mathbb{A}^k = \mathbb{A}^{(\lambda_1)} \times \dots \times \mathbb{A}^{(\lambda_k)}$ is finite over $\mathbb{A}^{(\Delta)}$, cf. [DF, BCES]. For a closed point $\mathbf{c} = (c_1, \dots, c_k) \in \mathbb{A}^k$ we sometimes keep the same notation for its image in $\mathbb{A}^{(\Delta)}$. For instance, by $\mathbb{C}_{\mathbf{c}}$ we usually mean the one-dimensional $\mathbb{C}[\mathbb{A}^{(\Delta)}]$ -module corresponding to the point \mathbf{c} .

2.2. BD Grassmannians and spherical Schubert varieties. The Beilinson-Drinfeld Grassmannian $\mathrm{Gr}_{\mathbb{A}^k}$ (BD Grassmannian for short) is the moduli space of collections consisting of the points $(c_1, \dots, c_k) \in \mathbb{A}^k$, a G^{ad} -torsor \mathcal{P} over \mathbb{A}^1 , and a trivialization of \mathcal{P} outside the points c_i .

Example 2.4. If $k = 1$ then $\mathrm{Gr}_{\mathbb{A}^1}$ is fibered over the affine line with a fiber isomorphic to the affine Grassmannian Gr .

In general, the fiber of the natural projection $\pi: \mathrm{Gr}_{\mathbb{A}^k} \rightarrow \mathbb{A}^k$ over a point (c_1, \dots, c_k) is isomorphic to the product of a copies of Gr , where a is the number of distinct entries c_i .

Example 2.5. Let $\mathring{\mathbb{A}}^k$, $k \geq 2$ be the open subvariety of \mathbb{A}^k consisting of points with pairwise distinct coordinates. Then

$$(2.1) \quad \pi^{-1}(\mathring{\mathbb{A}}^k) \simeq \mathring{\mathbb{A}}^k \times \mathrm{Gr}^k.$$

The BD Grassmannian $\mathrm{Gr}_{\mathbb{A}^k}$ enjoys the key *factorization property*. We have the addition of configurations morphism $\mathrm{add}: \mathbb{A}^k \times \mathbb{A}^l \rightarrow \mathbb{A}^{k+l}$ and an open subset $(\mathbb{A}^k \times \mathbb{A}^l)_{\mathrm{disj}} \subset \mathbb{A}^k \times \mathbb{A}^l$ formed by all the pairs of disjoint effective divisors. Then there is a canonical isomorphism

$$(\mathrm{Gr}_{\mathbb{A}^k} \times \mathrm{Gr}_{\mathbb{A}^l})|_{(\mathbb{A}^k \times \mathbb{A}^l)_{\mathrm{disj}}} \cong \mathrm{Gr}_{\mathbb{A}^{k+l}} \times_{\mathbb{A}^{k+l}} (\mathbb{A}^k \times \mathbb{A}^l)_{\mathrm{disj}}.$$

The BD Grassmannian $\mathrm{Gr}_{\mathbb{A}^k}$ is an ind-scheme, i.e. it is an inductive limit of the finite-dimensional BD Schubert varieties $\overline{\mathrm{Gr}}^\lambda$ for k -tuples of dominant coweights $\lambda = (\lambda_1, \dots, \lambda_k)$. More precisely, we consider a group scheme $\mathbb{G}(k)$ over \mathbb{A}^k , whose fiber over a point $\mathbf{c} = (c_1, \dots, c_k) \in \mathbb{A}^k$ is equal to the inverse limit ($n \rightarrow \infty$)

$$(2.2) \quad \mathbb{G}(k)_{\mathbf{c}} = \varprojlim_n G^{\mathrm{sc}}(\mathbb{C}[t]/P(t)^n), \quad P(t) = \prod_{i=1}^k (t - c_i).$$

Clearly, a fiber $\mathbb{G}(k)_{\mathbf{c}}$ is isomorphic to the a -th power of the group $G^{\text{sc}}(\mathcal{O})$, where a is the number of distinct elements among c_i . The group $\mathbb{G}(k)$ naturally acts on $\text{Gr}_{\mathbb{A}^k}$ fiberwise.

The spherical Schubert varieties in the BD Grassmannian are the closures of the $\mathbb{G}(k)$ -orbits in the fiber over the generic point of \mathbb{A}^k . The orbits are parametrized by the k -tuples $\underline{\lambda} \in P_+^k$. Given such a collection, let $t^{\underline{\lambda}}: \mathbb{A}^k \rightarrow \text{Gr}_{\mathbb{A}^k}$ be a section of π such that for $\mathbf{c} \in \mathring{\mathbb{A}}^k$ one has

$$t^{\underline{\lambda}}(\mathbf{c}) = ((t - c_1)^{\lambda_1}, \dots, (t - c_k)^{\lambda_k}) \in \prod_{i=1}^k \text{Gr}_{c_i}$$

(so the total section is the closure of $t^{\underline{\lambda}}(\mathring{\mathbb{A}}^k)$). Now the BD Schubert varieties are defined as the closures of the $\mathbb{G}(k)$ -orbits:

$$\overline{\text{Gr}}^{\underline{\lambda}} = \overline{\mathbb{G}(k) \cdot t^{\underline{\lambda}}} \subset \text{Gr}_{\mathbb{A}^k}.$$

The restriction of $\pi: \text{Gr}_{\mathbb{A}^k} \rightarrow \mathbb{A}^k$ to $\overline{\text{Gr}}^{\underline{\lambda}}$ is denoted by $\pi_{\underline{\lambda}}: \overline{\text{Gr}}^{\underline{\lambda}} \rightarrow \mathbb{A}^k$. This is a flat morphism, and all the fibers are reduced [Z1, Proposition 1.2.4] (it is proved for $k = 2$ in *loc. cit.*, but the proof works for arbitrary k). The fiber $\overline{\text{Gr}}_{\mathbf{c}}^{\underline{\lambda}} = \pi_{\underline{\lambda}}^{-1}(\mathbf{c})$ over a point $\mathbf{c} \in \mathbb{A}^k$ with

$$(2.3) \quad c_1 = \dots = c_{i_1} \neq c_{i_1+1} = \dots = c_{i_1+i_2} \neq \dots \neq c_{k-i_s+1} = \dots = c_k$$

is isomorphic to the product

$$\overline{\text{Gr}}^{\lambda_1+\dots+\lambda_{i_1}} \times \overline{\text{Gr}}^{\lambda_{i_1+1}+\dots+\lambda_{i_1+i_2}} \times \dots \times \overline{\text{Gr}}^{\lambda_{k-i_s+1}+\dots+\lambda_k}$$

of spherical Schubert varieties in the affine Grassmannian Gr . In particular, the fiber of $\pi_{\underline{\lambda}}$ over the origin (or any other point of the total diagonal) is isomorphic to the spherical Schubert variety $\overline{\text{Gr}}^{\lambda_1+\dots+\lambda_k}$.

The BD Grassmannians and the BD Schubert varieties carry the relatively very ample determinant line bundle \mathcal{L} . In particular, for any $\ell \geq 1$

$$H^0(\overline{\text{Gr}}_{\mathbf{c}}^{\underline{\lambda}}, \mathcal{L}_{\mathbf{c}}^{\otimes \ell})^* \simeq D_{\ell, \lambda_1+\dots+\lambda_{i_1}} \otimes \dots \otimes D_{\ell, \lambda_{k-i_s+1}+\dots+\lambda_k},$$

where $\mathcal{L}_{\mathbf{c}}$ is the restriction of the line bundle \mathcal{L} to the fiber $\overline{\text{Gr}}_{\mathbf{c}}^{\underline{\lambda}}$.

We introduce also the partially symmetrized (colored) version $\overline{\text{Gr}}^{(\underline{\lambda})}$ of the BD Schubert varieties. To define it we first consider the case of fundamental coweights λ_i . So assume that all λ_i are fundamental, i.e.

$$\lambda_1 = \dots = \lambda_{m_1} = \omega_1, \dots, \lambda_{k-m_r+1} = \dots = \lambda_k = \omega_r.$$

Let $\lambda = \sum_{i=1}^k \lambda_i = \sum_{j=1}^r m_j \omega_j$, and $N = |\lambda| = k$. The action of $S_{\lambda} = S_{m_1} \times \dots \times S_{m_r}$ on \mathbb{A}^N lifts to an action of S_{λ} on $\text{Gr}_{\mathbb{A}^N}$ such that $\pi_{\underline{\lambda}}$ is S_{λ} -equivariant. We define

$$\text{Gr}_{\mathbb{A}^{\lambda}} := \text{Gr}_{\mathbb{A}^N} / S_{\lambda}.$$

It is the moduli space of G^{ad} -torsors on \mathbb{A}^1 trivialized away from an N -tuple of points (x_1, \dots, x_N) , but we disregard the order within the groups

$(x_1, \dots, x_{m_1}), \dots, (x_{N-m_r+1}, \dots, x_N)$. We introduce the closed subvariety $\overline{\text{Gr}}^{(\underline{\lambda})} \subset \text{Gr}_{\mathbb{A}^\lambda}$ as the categorical quotient

$$\overline{\text{Gr}}^{(\underline{\lambda})} = \overline{\text{Gr}}^\lambda / S_\lambda \subset \text{Gr}_{\mathbb{A}^N} / S_\lambda = \text{Gr}_{\mathbb{A}^\lambda}.$$

Since the collection $\underline{\lambda}$ of fundamental weights is uniquely determined by their sum λ , we also use the notation $\overline{\text{Gr}}_\lambda$ for $\overline{\text{Gr}}^{(\underline{\lambda})}$.

Now we consider an arbitrary k -tuple $\underline{\lambda} = (\lambda_1, \dots, \lambda_k)$ (so that λ_i are not necessarily fundamental coweights). We set again $\lambda = \lambda_1 + \dots + \lambda_k$. Recall the closed subscheme $\mathbb{A}^{(\underline{\lambda})} \subset \mathbb{A}^\lambda$ introduced in Section 2.1. We set

$$\overline{\text{Gr}}^{(\underline{\lambda})} := \overline{\text{Gr}}_\lambda \times_{\mathbb{A}^\lambda} \mathbb{A}^{(\underline{\lambda})}.$$

The natural projection $\overline{\text{Gr}}^{(\underline{\lambda})} \rightarrow \mathbb{A}^{(\underline{\lambda})}$ is denoted $\pi_{(\underline{\lambda})}$.

Note that in case when $(\underline{\lambda}) = (\lambda, \lambda, \dots, \lambda)$, $\text{Gr}^{(\underline{\lambda})}$ is a Schubert variety in the symmetrized version of Beilinson-Drinfeld Grassmannian (see [Z2]).

The determinant line bundle \mathcal{L} descends from $\text{Gr}_{\mathbb{A}^N}$ to $\text{Gr}_{\mathbb{A}^\lambda}$. We will keep the same notation \mathcal{L} for its restriction to $\overline{\text{Gr}}_\lambda$ and to $\overline{\text{Gr}}^{(\underline{\lambda})}$.

Proposition 2.6. (a) *Let $\lambda_1, \dots, \lambda_k$ be fundamental coweights, $\lambda = \sum_{i=1}^k \lambda_i$. Then one has the base change isomorphism:*

$$H^0(\overline{\text{Gr}}^\lambda, \mathcal{L}^{\otimes \ell}) \cong H^0(\overline{\text{Gr}}_\lambda, \mathcal{L}^{\otimes \ell}) \otimes_{\mathcal{A}^\lambda} \mathbb{C}[\mathbb{A}^k].$$

(b) *Let $\lambda_1, \dots, \lambda_k$ be arbitrary dominant coweights, $\lambda = \sum_{i=1}^k \lambda_i$. Then one has the base change isomorphism:*

$$H^0(\overline{\text{Gr}}^\lambda, \mathcal{L}^{\otimes \ell}) \cong H^0(\overline{\text{Gr}}^{(\underline{\lambda})}, \mathcal{L}^{\otimes \ell}) \otimes_{\mathcal{A}^{(\underline{\lambda})}} \mathbb{C}[\mathbb{A}^k].$$

(c) *The $\mathbb{C}[\mathbb{A}^k]$ -module $H^0(\overline{\text{Gr}}^\lambda, \mathcal{L}^{\otimes \ell})$ is free.*

(d) *The $\mathcal{A}^{(\underline{\lambda})}$ -module $H^0(\overline{\text{Gr}}^{(\underline{\lambda})}, \mathcal{L}^{\otimes \ell})$ is free.*

Proof. We have a cartesian square

$$\begin{array}{ccc} \overline{\text{Gr}}^\lambda & \longrightarrow & \overline{\text{Gr}}^{(\underline{\lambda})} \\ \downarrow & & \downarrow \\ \mathbb{A}^{(\lambda_1)} \times \dots \times \mathbb{A}^{(\lambda_k)} & \longrightarrow & \mathbb{A}^{(\underline{\lambda})}, \end{array}$$

and the determinant line bundle \mathcal{L} on $\overline{\text{Gr}}^\lambda$ is the pullback of the determinant line bundle \mathcal{L} on $\overline{\text{Gr}}^{(\underline{\lambda})}$. The pushforward of the relatively very ample line bundle $\mathcal{L}^{\otimes \ell}$ from $\overline{\text{Gr}}^\lambda$ to \mathbb{A}^k is a locally free sheaf \mathcal{V} . Indeed, we already know that $\pi_\lambda: \overline{\text{Gr}}^\lambda \rightarrow \mathbb{A}^k$ is flat and all the fibers are reduced. But the dimension of the space of sections of $\mathcal{L}^{\otimes \ell}$ restricted to any fiber is independent of the choice of fiber by [FL1, Theorem 1] or [Z1, Theorem 1.2.2].

Furthermore, the pushforward of $\mathcal{L}^{\otimes \ell}$ from $\overline{\text{Gr}}_\lambda$ to \mathbb{A}^λ is a direct summand of S_λ -invariants in the pushforward of \mathcal{V} from \mathbb{A}^k to \mathbb{A}^λ . Hence the pushforward of $\mathcal{L}^{\otimes \ell}$ from $\overline{\text{Gr}}_\lambda$ to \mathbb{A}^λ is a locally free sheaf \mathcal{W} as well. Finally, $\pi_{(\underline{\lambda})}^* \mathcal{L}^{\otimes \ell}$ is the restriction of \mathcal{W} to $\mathbb{A}^{(\underline{\lambda})} \subset \mathbb{A}^\lambda$, and hence $\pi_{(\underline{\lambda})}^* \mathcal{L}^{\otimes \ell}$ is a locally free sheaf on $\mathbb{A}^{(\underline{\lambda})}$ as well. In particular, it is flat, and it remains to apply the base change for the above cartesian square. This proves (a) and (b).

To prove (c) and (d) note that $H^0(\overline{\text{Gr}}(\lambda), \mathcal{L}^{\otimes \ell})$ is projective over $\mathcal{A}(\lambda)$ because its fibers have the same dimension at every closed point. Since both $\mathcal{A}(\lambda)$ and $H^0(\overline{\text{Gr}}(\lambda), \mathcal{L}^{\otimes \ell})$ are non-negatively graded, and the degree 0 part of $\mathcal{A}(\lambda)$ is \mathbb{C} , we conclude by the graded Nakayama lemma that $H^0(\overline{\text{Gr}}(\lambda), \mathcal{L}^{\otimes \ell})$ is free over $\mathcal{A}(\lambda)$.¹ \square

3. GLOBAL MODULES

In this section, we prove several statements on the global modules defined in [DF]. Although we will be mainly interested in the global Demazure modules, we start in a more general setup.

So let $\{M_i\}_{i=1}^k$ be cyclic graded $\mathfrak{g}[t]$ -modules with cyclic vectors of dominant nonzero weights $\{\iota(\lambda_i)\}_{i=1}^k$, such that $t\mathfrak{h}[t]$ annihilates these cyclic vectors. Recall that it was proved in [DF] that for $\mathbf{c} = (c_1, \dots, c_k)$ lying in some Zariski-open subset of \mathbb{C}^k one has

$$(3.1) \quad R(M_1, \dots, M_k) \otimes_{\mathcal{A}(\lambda)} \mathbb{C}_{\mathbf{c}} \simeq \bigotimes_{i=1}^k M_i(c_i)$$

(see Remark 2.3).

It was also shown that the fiber of $R(M_1, \dots, M_k)$ at 0 surjects to the fusion product:

$$(3.2) \quad R(M_1, \dots, M_k) \otimes_{\mathcal{A}(\lambda)} \mathbb{C}_0 \twoheadrightarrow M_1 * \dots * M_k.$$

In particular, this surjection is an isomorphism if and only if the equality of dimensions $\dim(R(M_1, \dots, M_k) \otimes_{\mathcal{A}(\lambda)} \mathbb{C}_0) = \prod_{i=1}^k \dim M_i$ holds.

The next proposition shows that if the above surjection is an isomorphism, then the Zariski-open subset for which (3.1) holds can be described explicitly.

Proposition 3.1. *Suppose an isomorphism $R(M_1, \dots, M_k) \otimes_{\mathcal{A}(\lambda)} \mathbb{C}_0 \simeq M_1 * \dots * M_k$ holds. Then an isomorphism $R(M_1, \dots, M_k) \otimes_{\mathcal{A}(\lambda)} \mathbb{C}_{\mathbf{c}} \simeq \bigotimes_{i=1}^k M_i(c_i)$ holds for any \mathbf{c} with pairwise distinct coordinates $c_i \neq c_j$ (not just for \mathbf{c} in some open subset).*

Proof. As explained in [DF], an isomorphism

$$R(M_1, \dots, M_k) \otimes_{\mathcal{A}(\lambda)} \mathbb{C}_0 \simeq M_1 * \dots * M_k$$

implies by the semi-continuity theorem the equality

$$\dim(R(M_1, \dots, M_k) \otimes_{\mathcal{A}(\lambda)} \mathbb{C}_{\mathbf{c}}) = \prod_{i=1}^k \dim M_i$$

for any \mathbf{c} . Hence, it suffices to construct a surjection

$$R(M_1, \dots, M_k) \otimes_{\mathcal{A}(\lambda)} \mathbb{C}_{\mathbf{c}} \twoheadrightarrow \bigotimes_{i=1}^k M_i(c_i)$$

¹This last observation is due to Roman Travkin.

for \mathbf{c} with $c_i \neq c_j$. For any i clearly one has $M_i[t] \twoheadrightarrow M_i[t] \otimes_{\mathcal{A}(\lambda_i)} \mathbb{C}_{c_i} \simeq M_i(c_i)$ and hence

$$R(M_1, \dots, M_k) = \bigodot_{i=1}^k M_i[t] \twoheadrightarrow \bigodot_{i=1}^k M_i(c_i) \simeq \bigotimes_{i=1}^k M_i(c_i)$$

(the last isomorphism is proved in [FeLo, Proposition 1.4]). To show that this surjection factors through $R(M_1, \dots, M_k) \otimes \mathbb{C}_{\mathbf{c}}$ one needs to show that the relations $ht^n - (\langle \iota(\lambda_1), h \rangle c_1^n + \dots + \langle \iota(\lambda_k), h \rangle c_k^n)$ hold in the right-hand side module for any $h \in \mathfrak{h}$, which is clearly true. \square

For a global module $R(M_1, \dots, M_k)$ we denote by $R(M_1, \dots, M_k)^\vee$ its $\mathcal{A}(\underline{\lambda})$ -dual, i.e.

$$R(M_1, \dots, M_k)^\vee = \text{Hom}_{\mathcal{A}(\underline{\lambda})}(R(M_1, \dots, M_k), \mathcal{A}(\underline{\lambda})).$$

Remark 3.2. We note that $R(M_1, \dots, M_k)^\vee$ carries a natural structure of $\mathfrak{g}[t]$ -module. However, while $R(M_1, \dots, M_k)$ is cyclic, $R(M_1, \dots, M_k)^\vee$ does not have to be cyclic or cocyclic. The simplest example pops up for $\mathfrak{g} = \mathfrak{sl}_2$, $k = 2$ and $\lambda_1 = \lambda_2 = \omega$. We note that if $R(M_1, \dots, M_k)$ is free over $\mathcal{A}(\underline{\lambda})$, then the q -character of $R(M_1, \dots, M_k)^\vee$ is computed as $\text{ch}_q(R(M_1, \dots, M_k) \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}_0)|_{q \rightarrow q^{-1}} \cdot \text{ch}_q \mathcal{A}(\underline{\lambda})$. Hence in our special case one has

$$\text{ch}_q \mathbb{W}_{2\omega}^\vee = q^{-1} + (z^2 + 2 + z^{-2}) + q \dots,$$

showing that $\mathbb{W}_{2\omega}^\vee$ is neither cyclic nor cocyclic.

Proposition 3.3. *Assume that the weights of the cyclic vectors of $\mathfrak{g}[t]$ -modules M_i are nonzero. Then there is an isomorphism of $\mathfrak{g}[t] - \mathcal{A}(\underline{\lambda})$ -bimodules:*

$$R(M_1, \dots, M_k)_{\mathcal{A}(\underline{\lambda})}^{\odot \ell} \simeq R(M_1^{\odot \ell}, \dots, M_k^{\odot \ell})$$

(notation of Section 1.2).

Proof. Recall that the action of the highest weight algebra comes from the $U(\mathfrak{h}[t])$ -action. In this proof, we consider global modules with different highest weight algebras. So, we use the notation $\otimes_{U(\mathfrak{h}[t])}$ instead of $\otimes_{\mathcal{A}(\underline{\lambda})}$, although formally there is no difference.

We first consider an isomorphism

$$\begin{aligned} M_i[t] \otimes_{U(\mathfrak{h}[t])} M_i[t] &\xrightarrow{\sim} (M_i \otimes M_i)[t], \\ v_1 t^{k_1} \otimes_{U(\mathfrak{h}[t])} v_2 t^{k_2} &\mapsto (v_1 \otimes v_2) t^{k_1+k_2} \end{aligned}$$

(one can easily check that it is bijective and $\mathfrak{g}[t]$ -equivariant). Then we extend it to an isomorphism

$$\underbrace{\left(\bigotimes_{i=1}^k M_i[t] \right) \otimes_{U(\mathfrak{h}[t])} \dots \otimes_{U(\mathfrak{h}[t])}}_{\ell} \left(\bigotimes_{i=1}^k M_i[t] \right) \xrightarrow{\sim} \bigotimes_{i=1}^k M_i^{\otimes \ell}[t].$$

Considering the $\mathfrak{g}[t]$ -envelopes of the tensor products of cyclic vectors in both sides, we obtain the desired isomorphism

$$R(M_1, \dots, M_k)_{\mathcal{A}(\underline{\lambda})}^{\odot \ell} \simeq R(M_1^{\odot \ell}, \dots, M_k^{\odot \ell}).$$

□

Corollary 3.4. *There is a natural structure of a graded $\mathcal{A}(\underline{\lambda})$ -algebra on the space*

$$\bigoplus_{\ell \geq 0} R(M_1^{\odot \ell}, \dots, M_k^{\odot \ell})^\vee$$

(we set $R(M_1^{\odot 0}, \dots, M_k^{\odot 0})^\vee = \mathcal{A}(\underline{\lambda})$).

Proof. The multiplication structure is given by the dual of the map

$$R(M_1^{\odot \ell_1 + \ell_2}, \dots, M_k^{\odot \ell_1 + \ell_2}) \hookrightarrow R(M_1^{\odot \ell_1}, \dots, M_k^{\odot \ell_1}) \otimes_{\mathcal{A}(\underline{\lambda})} R(M_1^{\odot \ell_2}, \dots, M_k^{\odot \ell_2}).$$

□

To make a link to the Beilinson-Drinfeld setup we consider the global modules with all M 's being Demazure modules of the same level ℓ and nonzero highest weights. Let us recall the notation

$$\begin{aligned} \mathbb{D}(\ell, \underline{\lambda}) &= R(D_{\ell, \lambda_1}, \dots, D_{\ell, \lambda_k}), \\ \mathbb{D}_{\ell, \lambda} &= \mathbb{D}(\ell, \underbrace{\omega_1, \dots, \omega_1}_{m_1}, \dots, \underbrace{\omega_r, \dots, \omega_r}_{m_r}), \end{aligned}$$

where $\lambda = \sum_{i=1}^k \lambda_i = \sum_{j=1}^r m_j \omega_j$. Recall also that $D_{1, \lambda}^{\odot \ell} \simeq D_{\ell, \lambda}$. We get the following version of Corollary 3.4:

Corollary 3.5. *There is a natural structure of a graded $\mathcal{A}(\underline{\lambda})$ -algebra on the space*

$$\bigoplus_{\ell \geq 0} \mathbb{D}(\ell, \underline{\lambda})^\vee.$$

In particular, if all λ_i are fundamental, we have an \mathcal{A}_λ -algebra

$$\mathbb{D}_\lambda^\vee = \bigoplus_{\ell \geq 0} \mathbb{D}_{\ell, \lambda}^\vee.$$

Remark 3.6. Note that by construction this algebra is generated by its first homogeneous component, or, in other words, one has a surjection from the symmetric algebra:

$$\mathrm{Sym}_{\mathcal{A}(\underline{\lambda})}^\bullet \mathbb{D}(1, \underline{\lambda})^\vee \twoheadrightarrow \bigoplus_{\ell \geq 0} \mathbb{D}(\ell, \underline{\lambda})^\vee.$$

This means that the $\mathbb{A}^{(\underline{\lambda})}$ -scheme

$$\mathrm{Proj}(\bigoplus_{\ell \geq 0} \mathbb{D}(\ell, \underline{\lambda})^\vee)$$

is a closed subscheme of the projective space

$$\mathbb{P}_{\mathcal{A}(\underline{\lambda})}(\mathbb{D}(1, \underline{\lambda})) = \mathrm{Proj} \left(\mathrm{Sym}_{\mathcal{A}(\underline{\lambda})}^\bullet \mathbb{D}(1, \underline{\lambda})^\vee \right).$$

Remark 3.7. Assume that \mathfrak{g} is simply laced, all λ_i are fundamental, $\lambda = \sum_{i=1}^k \lambda_i$. Then $\mathbb{D}(1, \underline{\lambda}) \simeq \mathbb{W}_\lambda$ is the global Weyl module. It is proved in [Kato2] that the projective spectrum of the algebra $\bigoplus_{\lambda \in P_+} \mathbb{W}_\lambda^*$ is isomorphic to the (formal version of) semi-infinite flag variety (see also [Kato1, KNS, BF1, BF2, BF3, FiMi, FeMa1]). There are two important differences between the algebras $\bigoplus_{\lambda \in P_+} \mathbb{W}_\lambda^*$ and $\bigoplus_{\ell \geq 0} \mathbb{D}(\ell, \underline{\lambda})^\vee$. First, the sum in the first algebra runs over the dominant integral weights, while in the second case the summation is performed over the nonnegative integers. Second, the dual in the first algebra is taken with respect to the ground field, while in the second algebra one considers the duals with respect to the highest weight algebra.

Now we will prove that an arbitrary global Demazure module $\mathbb{D}(\ell, \underline{\lambda})$ is free over $\mathcal{A}(\underline{\lambda})$. We start with the simply-laced case.

Lemma 3.8. *Let \mathfrak{g} be of simply-laced type, $\underline{\lambda} = (\lambda_1, \dots, \lambda_k)$, and $\lambda = \lambda_1 + \dots + \lambda_k$. Then*

$$\mathbb{D}(\ell, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}_0 \simeq D_{\ell, \lambda}.$$

Proof. Note that this Lemma was already proved in [DF, Proposition 3.2] in the case of all λ_i being fundamental coweights. Thus, fundamental Demazure modules satisfy the condition of Proposition A.1. Using the associativity (Proposition A.1 (c)), we obtain the Lemma for arbitrary coweights. \square

We proceed to an arbitrary type.

Proposition 3.9. (a) *One has an isomorphism of $\mathfrak{g}[t]$ -modules*

$$\mathbb{D}(\ell, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}_0 \simeq D_{\ell, \lambda_1 + \dots + \lambda_k}.$$

(b) *The global Demazure module $\mathbb{D}(\ell, \underline{\lambda})$ is free over $\mathcal{A}(\underline{\lambda})$.*

Our proof uses ideas of [FL2, Theorem 8].

Proof. We reduce the general case to the case of \mathfrak{sl}_2 , which is simply laced and hence follows from Lemma 3.8.

As we know (recall (3.2)), $\dim(\mathbb{D}(\ell, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}_0) \geq \dim(D_{\ell, \lambda})$, so it suffices to construct a surjection

$$(3.3) \quad \mathbb{D}(\ell, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}_0 \leftarrow D_{\ell, \lambda}.$$

As it was shown in [FL2, J], the defining relations of $D_{\ell, \lambda}$ are

$$\mathfrak{n}^+[t].v = 0, \quad h.v = \ell \langle \iota(\lambda), h \rangle v, \quad t\mathfrak{h}[t].v = 0, \quad (f_\beta t^s)^{k_\beta+1}.v = 0.$$

Here $h \in \mathfrak{h}$, $s \in \mathbb{Z}_{\geq 0}$, and f_β is the Chevalley generator corresponding to a positive root β^\vee . Finally, $k_\beta = \ell \max\{0, \langle \Lambda_0^\vee + \iota(\lambda), -\beta + s \frac{(\beta, \beta)}{2} K \rangle\}$.

The first three relations obviously hold in the left-hand side of (3.3), so it remains to show that the last one also does. Consider the \mathfrak{sl}_2 -triple \mathfrak{sl}_2^β corresponding to β^\vee .

It was shown in [FL2, Lemma 7] that there is an $\mathfrak{sl}_2^\beta[t]$ -submodule of D_{1,λ_i} , that is isomorphic to $\widehat{\mathfrak{sl}_2^\beta}$ Demazure module $D_{\epsilon, m\omega^\vee}$, where $\epsilon = (\beta, \beta)/2$, $m = \langle \iota(\lambda), \beta \rangle / \epsilon$, and ω^\vee is the fundamental weight of \mathfrak{sl}_2^β . It follows that there is an $\mathfrak{sl}_2^\beta[t]$ -submodule of D_{ℓ, λ_i} , that is isomorphic to $\widehat{\mathfrak{sl}_2^\beta}$ Demazure module $D_{\ell\epsilon, m\omega^\vee}$. Denote it by $M(\ell, \iota(\lambda_i))$.

This induces an embedding $M(\ell, \iota(\lambda_i))[t] \hookrightarrow D_{\ell, \lambda_i}[t]$ and hence, denoting the highest vector of D_{ℓ, λ_i} by v_i we have

$$\begin{aligned} R(M(\ell, \iota(\lambda_1)), \dots, M(\ell, \iota(\lambda_k))) &\simeq U(\mathfrak{sl}_2^\beta[t]) \cdot \bigotimes_{i=1}^k v_i \hookrightarrow \\ &U(\mathfrak{g}[t]) \cdot \bigotimes_{i=1}^k v_i \simeq R(D_{\ell, \lambda_1}, \dots, D_{\ell, \lambda_k}) \simeq \mathbb{D}(\ell, \underline{\lambda}). \end{aligned}$$

Thereby, one has a map

$$(3.4) \quad R(M(\ell, \iota(\lambda_1)), \dots, M(\ell, \iota(\lambda_k))) \otimes_{\mathcal{A}} \mathbb{C}_0 \rightarrow \mathbb{D}(\ell, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}_0,$$

where \mathcal{A} is the highest weight algebra of the global $\mathfrak{sl}_2^\beta[t]$ Demazure module $R(M(\ell, \iota(\lambda_1)), \dots, M(\ell, \iota(\lambda_k)))$ and the map (3.4) is induced by the natural inclusion $\mathcal{A} \subset \mathcal{A}(\underline{\lambda})$. Now the left hand side of (3.4) is isomorphic to $M(\ell, \iota(\lambda))$, since we are in the simply laced case $\mathfrak{g} = \mathfrak{sl}_2^\beta$. The required relations $(f_\beta t^s)^{k_\beta+1} \cdot v = 0$, $s \geq 0$, hold in this module, and hence they hold in $\mathbb{D}(\ell, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}_0$ as well.

The end of the proof repeats the one of Proposition 2.6: $\mathbb{D}(\ell, \underline{\lambda})$ is projective over $\mathcal{A}(\underline{\lambda})$ because its fibers have the same dimension at every closed point. Now since both $\mathbb{D}(\ell, \underline{\lambda})$ and $\mathcal{A}(\underline{\lambda})$ are non-negatively graded, and the degree 0 part of $\mathcal{A}(\underline{\lambda})$ is \mathbb{C} , we conclude by the graded Nakayama lemma that $\mathbb{D}(\ell, \underline{\lambda})$ is free over $\mathcal{A}(\underline{\lambda})$. \square

Now we describe one more relation between global modules that will be used later in the paper:

Proposition 3.10. *One has an isomorphism of $\mathfrak{g}[t]$ -modules:*

$$\mathbb{D}(\ell, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \bigotimes_{i=1}^k \mathcal{A}(\ell\lambda_i) \simeq U(\mathfrak{g}[t]) \cdot \left(\bigotimes_{i=1}^k D_{\ell, \lambda_i}[t]_{\ell\iota(\lambda_i)} \right) \subset \bigotimes_{i=1}^k D_{\ell, \lambda_i}[t],$$

where $D_{\ell, \lambda_i}[t]_{\ell\iota(\lambda_i)}$ denotes the highest weight part of $D_{\ell, \lambda_i}[t]$.

Note that each algebra $\mathcal{A}(\ell\lambda_i)$ is isomorphic to the algebra of polynomials in one variable. We write $\mathcal{A}(\ell\lambda_i)$ (as opposed to just $\mathbb{C}[z_i]$) to point out that these algebras come as the highest weight spaces of the modules $D_{\ell, \lambda_i}[t]$. Further in the paper in the Beilinson-Drinfeld context we use the notation $\mathbb{C}[\mathbb{A}^{(\underline{\lambda})}] \subset \mathbb{C}[\mathbb{A}^k]$ instead of $\mathcal{A}(\underline{\lambda}) \subset \bigotimes_{i=1}^k \mathcal{A}(\ell\lambda_i)$, although it is the same, as explained in Subsection 2.1.

Proof. Let v be the cyclic vector of $\mathbb{D}(\ell, \underline{\lambda})$ and let v_i be the cyclic vector of $D_{\ell, \lambda_i}[t]$. We define the desired morphism by setting:

$$\begin{aligned} \phi: \mathbb{D}(\ell, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \bigotimes_{i=1}^k \mathcal{A}(\ell \lambda_i) &\rightarrow \bigotimes_{i=1}^k D_{\ell, \lambda_i}[t], \\ u.v \otimes_{\mathcal{A}(\underline{\lambda})} (h_1 t^{s_1} \otimes \dots \otimes h_n t^{s_k}) &\mapsto u(h_1 t^{s_1} v_1 \otimes \dots \otimes h_n t^{s_k} v_k), \end{aligned}$$

for $u \in U(\mathfrak{g}[t])$.

It is well defined because $\mathcal{A}(\ell \lambda_i)$ acts on $D_{\ell, \lambda_i}[t]$, and hence $\bigotimes_{i=1}^k \mathcal{A}(\ell \lambda_i)$ acts on $\bigotimes_{i=1}^k D_{\ell, \lambda_i}[t]$ commuting with the $\mathfrak{g}[t]$ -action.

It remains to prove injectivity. Consider both sides as $\bigotimes_{i=1}^k \mathcal{A}(\ell \lambda_i)$ -modules. Fibers of both sides at any point $\mathbf{c} = (c_1, \dots, c_k)$ with $c_i \neq c_j$ are $\bigotimes_{i=1}^k D_{\ell, \lambda_i}(c_i)$. Thereby, ϕ is injective on fibers in an open subset, and hence injective. \square

Example 3.11. Let $\mathfrak{g} = \mathfrak{sl}_2$, $k = 2$, $\lambda_1 = \lambda_2 = \omega$. Then one has two embeddings:

$$\begin{aligned} \mathbb{W}_{2\omega} &\hookrightarrow \mathbb{W}_\omega \otimes \mathbb{W}_\omega, \\ \mathbb{W}_{2\omega} \otimes_{\mathbb{C}[z_1, z_2]^{S_2}} \mathbb{C}[z_1, z_2] &\hookrightarrow \mathbb{W}_\omega \otimes \mathbb{W}_\omega. \end{aligned}$$

The image of the first embedding (the special case of Kato's theorem [Kato1, Corollary 3.5]) is the $U(\mathfrak{g}[t])$ -envelope of the tensor product of the highest vectors, while the image of the second embedding is the $U(\mathfrak{g}[t])$ -envelope of the tensor product of the highest weight components.

Remark 3.12. Due to Proposition 3.9, $\mathbb{D}(\ell, \underline{\lambda})$ is free over $\mathcal{A}(\underline{\lambda})$. Hence, $\mathbb{D}(\ell, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \bigotimes_{i=1}^k \mathcal{A}(\ell \lambda_i)$ is free over $\bigotimes_{i=1}^k \mathcal{A}(\ell \lambda_i)$. In particular, this implies:

$$\text{ch}_q(\mathbb{D}(\ell, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \bigotimes_{i=1}^k \mathcal{A}(\ell \lambda_i)) = \text{ch}_q D_{\ell, \lambda_1 + \dots + \lambda_k} \times (1 - q)^{-k}.$$

Remark 3.13. In fact, the proof of Proposition 3.10 hold for arbitrary global modules $R(M_1, \dots, M_n)$ without any changes. The isomorphism in the general case is of the form

$$R(M_1, \dots, M_k) \otimes_{\mathcal{A}(\underline{\lambda})} \bigotimes_{i=1}^k \mathcal{A}(\lambda_i) \simeq U(\mathfrak{g}[t]) \cdot \left(\bigotimes_{i=1}^k U(\mathfrak{h}[t]) v_i \right) \subset \bigotimes_{i=1}^k R(M_i).$$

4. GLOBAL DEMAZURE MODULES AND BD SCHUBERT VARIETIES

4.1. Sections of the determinant line bundle. The goal of this section is to identify the global Demazure modules $\mathbb{D}(\ell, \underline{\lambda})$ with the $\mathcal{A}(\underline{\lambda})$ -dual of the space of sections $H^0(\overline{\text{Gr}}(\underline{\lambda}), \mathcal{L}^{\otimes \ell})$ (we note that the higher cohomology $H^{>0}(\overline{\text{Gr}}(\underline{\lambda}), \mathcal{L}^{\otimes \ell})$ vanish: as in the proof of Proposition 2.6, it follows from

the flatness of $\pi_{(\underline{\lambda})}$ and the fact that the restriction of $\mathcal{L}^{\otimes \ell}$ to any fiber of $\pi_{(\underline{\lambda})}$ is very ample). To this end, we first establish an isomorphism

$$H^0(\overline{\text{Gr}}^{\underline{\lambda}}, \mathcal{L}^{\otimes \ell})^\vee \simeq \mathbb{D}(\ell, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}[\mathbb{A}^k],$$

where $\mathbb{A}^k = \mathbb{A}^{(\lambda_1)} \times \dots \times \mathbb{A}^{(\lambda_k)}$ and the notation M^\vee stands for the $\mathbb{C}[\mathbb{A}^k]$ -dual module to a $\mathbb{C}[\mathbb{A}^k]$ -module M . In order to compare these two spaces we make the following observation.

Lemma 4.1. *There is a homomorphism of Lie groups*

$$G^{\text{sc}}[t] \rightarrow \Gamma(\mathbb{A}^k, \mathbb{G}(k)),$$

where $\Gamma(\mathbb{A}^k, \mathbb{G}(k))$ is the group of sections of the group scheme $\mathbb{G}(k)$ over \mathbb{A}^k .

Proof. Recall that $\mathbb{G}(k)$ is defined as a scheme over \mathbb{A}^k whose fiber over a point \mathbf{c} is equal to the inverse limit ($n \rightarrow \infty$) of the groups $G^{\text{sc}}(\mathbb{C}[t]/P(t)^n)$, where $P(t) = \prod_{i=1}^k (t - c_i)$. Now the desired homomorphism is induced by sending the coordinate t in $\mathbb{C}[t]$ to $t \pmod{P(t)^n}$. \square

Corollary 4.2. *The space of sections $H^0(\overline{\text{Gr}}^{\underline{\lambda}}, \mathcal{L}^{\otimes \ell})$ is a $\mathfrak{g}[t]$ -module. The $\mathfrak{g}[t]$ -action commutes with the natural action of $\mathbb{C}[\mathbb{A}^k]$.*

Proof. The first claim is a direct consequence of Lemma 4.1. The second claim is clear since the group scheme $\mathbb{G}(k)$ acts fiberwise. \square

Now we prove our claim for the case of one weight.

Lemma 4.3. *Let $k = 1$, i.e. $\underline{\lambda} = (\lambda)$, $\lambda \in P_+$. Then for any $\ell \geq 1$ we have an isomorphism of $\mathfrak{g}[t]$ -modules*

$$H^0(\overline{\text{Gr}}^{(\lambda)}, \mathcal{L}^{\otimes \ell})^\vee \simeq D_{\ell, \lambda}[t].$$

Proof. Recall formula (1.1) for the action of the Lie algebra $\mathfrak{g}[x]$ on $D_{\ell, \lambda}[t] \simeq D_{\ell, \lambda} \otimes \mathbb{C}[t]$:

$$(4.1) \quad (g \otimes x^s)(v \otimes t^a) = \sum_{i=0}^s (-1)^{s-i} \binom{s}{i} (g \otimes x^i.v) \otimes t^{a+s-i}, \quad g \in \mathfrak{g}.$$

Here we deliberately replaced the variable t in $\mathfrak{g}[t]$ with an auxiliary variable x in order to make the picture similar to the BD context.

Now let us identify x with the global coordinate on \mathbb{A}^1 . Then one gets an isomorphism of vector spaces

$$H^0(\overline{\text{Gr}}^{(\lambda)}, \mathcal{L}^{\otimes \ell})^\vee \simeq D_{\ell, \lambda} \otimes \mathbb{C}[x],$$

where $D_{\ell, \lambda}$ is considered as a $\mathfrak{g}[t]$ -module. The action of $\mathfrak{g}[x]$ is induced by the map $x \mapsto x - t$, meaning that the result of the action of $g \otimes x^s$ on $v \otimes t^a$ is given by the right hand side of (4.1). \square

Theorem 4.4. *Let $\underline{\lambda} = (\lambda_1, \dots, \lambda_k) \in P_+^k$, all λ_i are nonzero. Then one has an isomorphism of $\mathfrak{g}[t] - \mathbb{C}[\mathbb{A}^k]$ -bimodules:*

$$H^0(\overline{\text{Gr}}^{\underline{\lambda}}, \mathcal{L}^{\otimes \ell})^\vee \simeq \mathbb{D}(\ell, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}[\mathbb{A}^k],$$

where M^\vee stands for the $\mathbb{C}[\mathbb{A}^k]$ -dual module to a $\mathbb{C}[\mathbb{A}^k]$ -module M .

Proof. According to Proposition 2.6(c), $H^0(\overline{\text{Gr}}^{(\lambda)}, \mathcal{L}^{\otimes \ell})^\vee$ is free as a $\mathbb{C}[\mathbb{A}^k]$ -module. In particular, its \mathfrak{g} -highest weight part $H^0(\overline{\text{Gr}}^{\underline{\lambda}}, \mathcal{L}^{\otimes \ell})_{\ell(\lambda)}^\vee$ is isomorphic to the free rank one module over $\mathbb{C}[\mathbb{A}^k]$. We also conclude that

$$(4.2) \quad \text{ch}_q H^0(\overline{\text{Gr}}^{\underline{\lambda}}, \mathcal{L}^{\otimes \ell})^\vee = \text{ch}_q D_{\ell, \lambda} \cdot (1 - q)^{-k}.$$

Due to Proposition 3.10 and Lemma 4.3, in order to prove the Theorem, it is enough to show:

$$(4.3) \quad H^0(\overline{\text{Gr}}^{\underline{\lambda}}, \mathcal{L}^{\otimes \ell})^\vee \simeq U(\mathfrak{g}[t]) \cdot \bigotimes_{i=1}^k H^0(\overline{\text{Gr}}^{(\lambda_i)}, \mathcal{L}^{\otimes \ell})_{\ell(\lambda_i)}^\vee,$$

where the lower index denotes the corresponding \mathfrak{g} -weight subspace.

By Lemma 4.3, $H^0(\overline{\text{Gr}}^{(\lambda_i)}, \mathcal{L}^{\otimes \ell})_{\ell(\lambda_i)}^\vee$ is isomorphic to the polynomial ring in one variable as a vector space. We consider the embedding

$$(4.4) \quad H^0(\overline{\text{Gr}}^{\underline{\lambda}}, \mathcal{L}^{\otimes \ell})^\vee \hookrightarrow H^0(\overline{\text{Gr}}^{\underline{\lambda}}(\mathring{\mathbb{A}}^k), \mathcal{L}^{\otimes \ell})^\vee,$$

where $\overline{\text{Gr}}^{\underline{\lambda}}(\mathring{\mathbb{A}}^k) \subset \overline{\text{Gr}}^{\underline{\lambda}}$ is $\pi_{\underline{\lambda}}^{-1}(\mathring{\mathbb{A}}^k)$, and the embedding (4.4) is induced by the open embedding $\overline{\text{Gr}}^{\underline{\lambda}}(\mathring{\mathbb{A}}^k) \hookrightarrow \overline{\text{Gr}}^{\underline{\lambda}}$. By the factorization property,

$$(4.5) \quad H^0(\overline{\text{Gr}}^{\underline{\lambda}}(\mathring{\mathbb{A}}^k), \mathcal{L}^{\otimes \ell})^\vee \cong \mathbb{C}[\mathring{\mathbb{A}}^k] \otimes \bigotimes_{i=1}^k D(\ell, \lambda_i).$$

In particular, the highest weight part $H^0(\overline{\text{Gr}}^{\underline{\lambda}}(\mathring{\mathbb{A}}^k), \mathcal{L}^{\otimes \ell})_{\ell(\lambda)}^\vee$ (with $\lambda = \sum_{i=1}^k \lambda_i$) is a free rank one module over the localization $\mathbb{C}[\mathring{\mathbb{A}}^k]$ of the polynomial algebra $\mathbb{C}[\mathbb{A}^k]$.

Thus we have the embeddings

$$\begin{aligned} H^0(\overline{\text{Gr}}^{\underline{\lambda}}, \mathcal{L}^{\otimes \ell})_{\ell(\lambda)}^\vee &\hookrightarrow H^0(\overline{\text{Gr}}^{\underline{\lambda}}(\mathring{\mathbb{A}}^k), \mathcal{L}^{\otimes \ell})_{\ell(\lambda)}^\vee \cong \mathbb{C}[\mathring{\mathbb{A}}^k] \otimes \bigotimes_{i=1}^k D(\ell, \lambda_i)_{\ell(\lambda_i)} \\ &\hookleftarrow \mathbb{C}[\mathbb{A}^k] \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{D}(\ell, \underline{\lambda})_{\ell(\lambda)} \end{aligned}$$

arising from the factorization property.

We claim that the images of these embeddings coincide. First we consider the case $k = 2$. Then the image I_{left} of the left embedding and the image I_{right} of the right embedding both are the free rank one modules over $\mathbb{C}[\mathbb{A}^2]$ inside the free rank one module over $\mathbb{C}[\mathring{\mathbb{A}}^2]$. If we denote the coordinates in \mathbb{A}^2 by z_1, z_2 , then necessarily $I_{\text{left}} = (z_1 - z_2)^a I_{\text{right}}$ for some $a \in \mathbb{Z}$, and we have to prove $a = 0$. Otherwise either $I_{\text{left}} \subsetneq I_{\text{right}}$ (if $a > 0$) or $I_{\text{right}} \subsetneq I_{\text{left}}$ (if $a < 0$). In the first case the graded character of I_{left} is *strictly less*

than the graded character of I_{right} (termwise) that contradicts the equality $\text{ch}_q H^0(\overline{\text{Gr}}^\lambda, \mathcal{L}^{\otimes \ell})^\vee = \text{ch}_q D_{\ell, \lambda} \cdot (1 - q)^{-k} = \text{ch}_q(\mathbb{C}[\mathbb{A}^k] \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{D}(\ell, \underline{\lambda}))$ by (4.2) and Remark 3.12 (in particular, the graded characters of the $\ell\iota(\lambda)$ -weight components must coincide as well). The second case similarly leads to a contradiction.

The coincidence of images for general k now follows after localization at generic points of diagonals in \mathbb{A}^k by factorization. Since we know the coincidence generically and in codimension one, it follows everywhere by algebraic Hartogs' lemma: given two locally free sheaves on \mathbb{A}^k , an isomorphism between them defined off a codimension 2 closed subset of \mathbb{A}^k necessarily extends to the whole of \mathbb{A}^k .

We conclude the equality

$$H^0(\overline{\text{Gr}}^\lambda, \mathcal{L}^{\otimes \ell})_{\ell\iota(\lambda)}^\vee = \mathbb{C}[\mathbb{A}^k] \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{D}(\ell, \underline{\lambda})_{\ell\iota(\lambda)}$$

inside $\mathbb{C}[\mathbb{A}^k] \otimes \bigotimes_{i=1}^k D(\ell, \lambda_i)$. But

$$\begin{aligned} H^0(\overline{\text{Gr}}^\lambda, \mathcal{L}^{\otimes \ell})^\vee &\supset U(\mathfrak{g}[t]) H^0(\overline{\text{Gr}}^\lambda, \mathcal{L}^{\otimes \ell})_{\ell\iota(\lambda)}^\vee \\ &= U(\mathfrak{g}[t]) (\mathbb{C}[\mathbb{A}^k] \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{D}(\ell, \underline{\lambda})_{\ell\iota(\lambda)}) = \mathbb{C}[\mathbb{A}^k] \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{D}(\ell, \underline{\lambda}). \end{aligned}$$

The equality of characters $\text{ch}_q H^0(\overline{\text{Gr}}^\lambda, \mathcal{L}^{\otimes \ell})^\vee = \text{ch}_q(\mathbb{C}[\mathbb{A}^k] \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{D}(\ell, \underline{\lambda}))$ once again guarantees that the above inclusion is actually an equality.

The theorem is proved. \square

Now to establish a relation between the global Demazure modules and the spaces of sections of determinant line bundles on the symmetrized Schubert varieties, we prove the following theorem.

Theorem 4.5. *Let $\lambda_1, \dots, \lambda_k \in P_+$, all λ_i are nonzero. Then one has an isomorphism of $\mathfrak{g}[t]$ -modules:*

$$H^0(\overline{\text{Gr}}^{(\underline{\lambda})}, \mathcal{L}^{\otimes \ell})^\vee \simeq \mathbb{D}(\ell, \underline{\lambda}),$$

where M^\vee stands for the $\mathcal{A}(\underline{\lambda})$ -dual module to an $\mathcal{A}(\underline{\lambda})$ -module M .

Proof. Using Proposition 2.6 and Theorem 4.4 we get an isomorphism:

$$(4.6) \quad \mathbb{D}(\ell, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}[\mathbb{A}^k] \simeq H^0(\overline{\text{Gr}}^{(\underline{\lambda})}, \mathcal{L}^{\otimes \ell})^\vee \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}[\mathbb{A}^k].$$

Let v be the cyclic vector of $\mathbb{D}(\ell, \underline{\lambda})$. The vector $v \otimes_{\mathcal{A}(\underline{\lambda})} 1$ in the left hand side of (4.6) is mapped to some vector of the form $w \otimes_{\mathcal{A}(\underline{\lambda})} 1$ in the right hand side. Using the $\mathfrak{g}[t]$ -equivariance we obtain that

$$\mathbb{D}(\ell, \underline{\lambda}) \simeq (U(\mathfrak{g}[t]) \cdot w) \otimes_{\mathcal{A}(\underline{\lambda})} 1 \subset H^0(\overline{\text{Gr}}^{(\underline{\lambda})}, \mathcal{L}^{\otimes \ell})^\vee \otimes_{\mathcal{A}(\underline{\lambda})} 1.$$

Hence there is an embedding $\mathbb{D}(\ell, \underline{\lambda}) \hookrightarrow H^0(\overline{\text{Gr}}^{(\underline{\lambda})}, \mathcal{L}^{\otimes \ell})^\vee$. Using Proposition 3.9, we see that the fibers at 0 of both sides are isomorphic to $D_{\ell, \lambda_1 + \dots + \lambda_k}$. The graded version of Nakayama lemma implies that the above injective map is a surjection, and thus an isomorphism. \square

Remark 4.6. *If the highest weight of a cyclic $\mathfrak{g}[t]$ module M is zero, then the module $M[z]$ is not cyclic. That is why we impose the condition $\lambda_i \neq 0$ in Theorems 4.4 and 4.5.*

It is an easy consequence of these Theorems that for the case $\underline{\lambda} = (\underline{\mu}, \underbrace{0, \dots, 0}_n)$

with $\mu_i \neq 0$ one has

$$\begin{aligned} H^0(\overline{\text{Gr}}^{\underline{\lambda}}, \mathcal{L}^{\otimes \ell})^\vee &\simeq (\mathbb{D}(\ell, \underline{\mu}) \otimes_{\mathcal{A}(\underline{\mu})} \mathbb{C}[\mathbb{A}^k]) \otimes \mathbb{C}[\mathbb{A}^n]; \\ H^0(\overline{\text{Gr}}^{(\underline{\lambda})}, \mathcal{L}^{\otimes \ell})^\vee &\simeq \mathbb{D}(\ell, \underline{\mu}) \otimes \mathbb{C}[\mathbb{A}^{(n)}]. \end{aligned}$$

Corollary 4.7. *One has an isomorphism of $\mathbb{A}^{(\underline{\lambda})}$ -schemes:*

$$\overline{\text{Gr}}^{(\underline{\lambda})} \simeq \text{Proj} \left(\bigoplus_{\ell \geq 0} \mathbb{D}(\ell, \underline{\lambda})^\vee \right).$$

Let us consider the special case when all λ_i are fundamental coweights. In particular, $\mathcal{A}(\underline{\lambda}) \simeq \mathcal{A}_\lambda$. We obtain the following corollary.

Corollary 4.8. *Assume that all λ_i are fundamental coweights and let $\lambda = \sum_{i=1}^k \lambda_i$. Then*

- (i) $H^0(\overline{\text{Gr}}^{(\underline{\lambda})}, \mathcal{L}^{\otimes \ell})^\vee \simeq \mathbb{D}(\ell, \lambda);$
- (ii) $\overline{\text{Gr}}^{(\underline{\lambda})} \simeq \text{Proj} \left(\bigoplus_{\ell \geq 0} \mathbb{D}(\ell, \lambda)^\vee \right).$

4.2. Embeddings of the BD Schubert varieties. The goal of this section is to show that the global Demazure modules provide projective embeddings of Beilinson-Drinfeld Schubert varieties (generalizing a relation between the affine Demazure modules and Schubert varieties).

Thanks to Proposition 3.9 the global Demazure module $\mathbb{D}(\ell, \underline{\lambda})$ is free over $\mathcal{A}(\underline{\lambda})$. Hence one gets a vector bundle $\mathcal{D}(\ell, \underline{\lambda})$ on $\mathbb{A}^{(\underline{\lambda})} = \text{Spec}(\mathcal{A}(\underline{\lambda}))$, whose fiber is given by the fiber of $\mathbb{D}(\ell, \underline{\lambda})$ at a point of the base. We will need the following lemma in order to embed the BD Schubert varieties into the fiberwise projectivized vector bundle $\mathcal{D}(1, \underline{\lambda})$.

Lemma 4.9. *The group scheme $\mathbb{G}(k)$ acts on $\mathcal{D}(\ell, \underline{\lambda})$ fiberwise.*

Proof. Recall (see (2.2)) that the fiber of $\mathbb{G}(k)$ over a point $\mathbf{c} = (c_1, \dots, c_k) \in \mathbb{A}^k$ is equal to the inverse limit

$$\mathbb{G}(k)_{\mathbf{c}} = \varprojlim_m G^{\text{sc}}(\mathbb{C}[t]/P(t)^m), \quad P(t) = \prod_{i=1}^k (t - c_i).$$

We also know that for $\mathbf{c} = (\underbrace{c_1, \dots, c_1}_{i_1}, \dots, \underbrace{c_n, \dots, c_n}_{i_n}) \in \mathbb{C}^k$ such that $c_p \neq c_q$

for $p \neq q$ one has

$$\mathbb{D}(\ell, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}_{\mathbf{c}} \simeq \bigotimes_{p=1}^n D_{\ell, \lambda_{i_1} + \dots + i_{p-1} + \dots + \lambda_{i_1} + \dots + i_p}(c_{i_p}).$$

We conclude that $\mathbb{G}(k)$ acts on $\mathcal{D}(\ell, \underline{\lambda})$ fiberwise. □

One has a section $s^\lambda: \mathbb{A}^{(\lambda)} \rightarrow \mathbb{P}(\mathcal{D}(1, \underline{\lambda}))$ of the natural projection map $\mathbb{P}(\mathcal{D}(1, \underline{\lambda})) \rightarrow \mathbb{A}^{(\lambda)}$ sending a point \mathbf{c} to the highest weight line of $\mathbb{D}(1, \underline{\lambda}) \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}_{\mathbf{c}}$. By Lemma 4.9 the group of sections of the group scheme $\mathbb{G}(k)$ naturally acts on $\mathbb{P}(\mathcal{D}(1, \underline{\lambda}))$. We obtain the following corollary.

Corollary 4.10. $\overline{\text{Gr}}^{(\lambda)}$ is equal to the closure of the $\mathbb{G}(k)$ -orbit of the section s^λ .

Proof. Follows from definition of $\overline{\text{Gr}}^\lambda$ in Section 2 and Theorem 4.5. \square

APPENDIX A. ON THE ASSOCIATIVITY OF FUSION PRODUCT

It was conjectured in [DF] that

$$(A.1) \quad R(M_1, \dots, M_k) \otimes_{\mathcal{A}(\underline{\lambda})} \mathbb{C}_0 \simeq M_1 * \dots * M_k.$$

The existence of the isomorphism (A.1) implies that fusion product does not depend on the choice of constants. Now we prove that (A.1) also implies the associativity of the fusion product.

Proposition A.1. *Let $N_1, \dots, N_k, M_1, \dots, M_m$ be finite-dimensional graded cyclic $\mathfrak{g}[t]$ -modules with cyclic vectors of weights $\lambda'_1, \dots, \lambda'_k, \mu'_1, \dots, \mu'_m$ such that*

$$R(N_1, \dots, N_k, M_1, \dots, M_m) \otimes_{\mathcal{A}(\underline{\lambda}^\vee, \underline{\mu}^\vee)} \mathbb{C}_0 \simeq N_1 * \dots * N_k * M_1 * \dots * M_m.$$

Then

(a)

$$R(N_1, \dots, N_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathbb{C}_0 \simeq N_1 * \dots * N_k,$$

(b)

$$N_1 * \dots * N_k * M_1 * \dots * M_m \simeq (N_1 * \dots * N_k) * M_1 * \dots * M_m,$$

(c)

$$\begin{aligned} R(R(N_1, \dots, N_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathbb{C}_0, M_1, \dots, M_m) \otimes_{\mathcal{A}(\lambda'_1 + \dots + \lambda'_k, \underline{\mu}^\vee)} \mathbb{C}_0 \\ \simeq R(N_1, \dots, N_k, M_1, \dots, M_m) \otimes_{\mathcal{A}(\underline{\lambda}^\vee, \underline{\mu}^\vee)} \mathbb{C}_0. \end{aligned}$$

Proof. We consider pairwise distinct $c_0, c_1, \dots, c_m \in \mathbb{C}$. Let

$$\mathbf{c} = (\underbrace{c_0, \dots, c_0}_k, c_1, \dots, c_m) \in \mathbb{C}^{k+m}.$$

Then clearly

$$\begin{aligned} (A.2) \quad R(N_1, \dots, N_k, M_1, \dots, M_m) \otimes_{\mathcal{A}(\underline{\lambda}^\vee, \underline{\mu}^\vee)} \mathbb{C}_{\mathbf{c}} \twoheadrightarrow \\ \left(R(N_1, \dots, N_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathbb{C}_{(c_0, \dots, c_0)} \right) \odot \left(R(M_1, \dots, M_m) \otimes_{\mathcal{A}(\underline{\mu}^\vee)} \mathbb{C}_{(c_1, \dots, c_m)} \right) \\ \simeq \left(R(N_1, \dots, N_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathbb{C}_0 \right) (c_0) \odot \left(M_1(c_1) \otimes \dots \otimes M_m(c_m) \right) \\ \simeq \left(R(N_1, \dots, N_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathbb{C}_0 \right) (c_0) \otimes (M_1(c_1) \otimes \dots \otimes M_m(c_m)). \end{aligned}$$

The last isomorphism holds because of [FeLo, Proposition 1.4].

Because of our assumption on the fiber at 0 of $R(N_1, \dots, N_k, M_1, \dots, M_m)$ we conclude that the fibers at all the points have the same dimension, and surjection (A.2) implies

$$\begin{aligned} \prod_{i=1}^k \dim N_i \times \prod_{i=1}^m \dim M_i &\geq \dim (R(N_1, \dots, N_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathbb{C}_0) \times \prod_{i=1}^m \dim M_i; \\ \prod_{i=1}^k \dim N_i &\geq \dim (R(N_1, \dots, N_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathbb{C}_0), \end{aligned}$$

Comparing with (3.2), we obtain part (a) of Proposition. We also conclude that (A.2) is an isomorphism.

Now, as proved in [DF, Proposition 2.11], there is a surjection

$$\begin{aligned} N_1 * \dots * N_k * M_1 * \dots * M_m &\simeq R(N_1, \dots, N_k, M_1, \dots, M_m) \otimes_{\mathcal{A}(\underline{\lambda}^\vee, \underline{\mu}^\vee)} \mathbb{C}_0 \twoheadrightarrow \\ &\quad \text{gr}(R(N_1, \dots, N_k, M_1, \dots, M_m) \otimes_{\mathcal{A}(\underline{\lambda}^\vee, \underline{\mu}^\vee)} \mathbb{C}_{\mathbf{c}}) \\ &\simeq \text{gr}\left((R(N_1, \dots, N_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathbb{C}_0)(c_0) \otimes (M_1(c_1) \otimes \dots \otimes M_m(c_m))\right) \\ &\simeq (N_1 * \dots * N_k) * M_1 * \dots * M_m. \end{aligned}$$

Comparing dimensions of both sides, we obtain part (b) of the Proposition.

To prove the remaining part, we first prove that

$$(A.3) \quad (R(N_1, \dots, N_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathbb{C}_0)[z] \simeq R(N_1, \dots, N_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathcal{A}',$$

where the algebra \mathcal{A}' is the algebra of polynomials in one variable, obtained by gluing all the variables in the algebra $\mathcal{A}(\underline{\lambda}^\vee)$:

$$\begin{array}{ccc} \mathbb{C}[z_1, \dots, z_k] & \xrightarrow{z_i \mapsto z} & \mathbb{C}[z] \\ \uparrow & & \parallel \\ \mathcal{A}(\underline{\lambda}^\vee) & \longrightarrow & \mathcal{A}'. \end{array}$$

Indeed, it follows from part (a) of the Proposition that $R(N_1, \dots, N_k, \cdot)$ is a free $\mathcal{A}(\underline{\lambda}^\vee)$ -module, and hence $R(N_1, \dots, N_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathcal{A}'$ is a free \mathcal{A}' -module. Therefore, we obtain (A.3) as an isomorphism of vector spaces. We note that the fibers of the left and right hand sides of (A.3) at a point $c \in \mathbb{C}$ are isomorphic as $\mathfrak{g}[t]$ -modules to $(N_1 * \dots * N_k)(c)$. It follows that both sides of (A.3) are isomorphic as $\mathfrak{g}[t]$ -modules. In particular, there is a surjection

$$(R(N_1, \dots, N_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathbb{C}_0)[z] \leftarrow R(N_1, \dots, N_k).$$

Using it, we obtain:

$$\begin{aligned}
& R(R(N_1, \dots, N_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathbb{C}_0, M_1, \dots, M_m) \otimes_{\mathcal{A}(\lambda_1^\vee + \dots + \lambda_k^\vee, \underline{\mu}^\vee)} \mathbb{C}_0 \\
& \simeq \left((R(N_1, \dots, N_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathbb{C}_0)[z] \odot R(M_1, \dots, M_m) \right) \otimes_{\mathcal{A}(\lambda_1^\vee + \dots + \lambda_k^\vee, \underline{\mu}^\vee)} \mathbb{C}_0 \\
& \quad \leftarrow \left(R(N_1, \dots, N_k) \odot R(M_1, \dots, M_m) \right) \otimes_{\mathcal{A}(\underline{\lambda}^\vee, \underline{\mu}^\vee)} \mathbb{C}_0 \\
& \quad \simeq R(N_1, \dots, N_k, M_1, \dots, M_m) \otimes_{\mathcal{A}(\underline{\lambda}^\vee, \underline{\mu}^\vee)} \mathbb{C}_0.
\end{aligned}$$

Comparing the dimensions of the leftmost and the rightmost terms, we obtain part c) of the Proposition. \square

Corollary A.2 (From the proof of Proposition A.1). *Suppose M_1, \dots, M_k are cyclic graded $\mathfrak{g}[t]$ modules such that*

$$R(M_1, \dots, M_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathbb{C}_0 \simeq M_1 * \dots * M_k.$$

Let $\mathbf{c} = (\underbrace{c_1, \dots, c_1}_{i_1}, \dots, \underbrace{c_n, \dots, c_n}_{i_n}) \in \mathbb{C}^k$ be such that $c_i \neq c_j$ for $i \neq j$. Then

$$\begin{aligned}
& R(M_1, \dots, M_k) \otimes_{\mathcal{A}(\underline{\lambda}^\vee)} \mathbb{C}_{\mathbf{c}} \\
& \simeq (M_1 * \dots * M_{i_1})(c_1) * \dots * (M_{i_1 + \dots + i_{n-1} + 1} * \dots * M_k)(c_n).
\end{aligned}$$

APPENDIX B. KEY OBJECTS OF THE PAPER

Simple Lie algebras:

\mathfrak{g} – simple Lie algebra of rank r with Cartan decomposition $\mathfrak{g} = \mathfrak{n}_+ \oplus \mathfrak{h} \oplus \mathfrak{n}_-$;
 G^{sc} (resp. G^{ad}) – simply connected (resp. adjoint) complex Lie group of \mathfrak{g} ;
 $\alpha_1^\vee, \dots, \alpha_r^\vee$ – simple roots, $\omega_1^\vee, \dots, \omega_r^\vee$ – fundamental weights;
 $\alpha_1, \dots, \alpha_r$ – simple coroots, $\omega_1, \dots, \omega_r$ – fundamental coweights;
 $P = \bigoplus_{i=1}^r \mathbb{Z}\omega_i \supset \bigoplus_{i=1}^r \mathbb{Z}_{\geq 0}\omega_i = P_+$ – coweight lattice and its dominant cone;
 $P^\vee = \bigoplus_{i=1}^r \mathbb{Z}\omega_i^\vee \supset \bigoplus_{i=1}^r \mathbb{Z}_{\geq 0}\omega_i^\vee = P_+^\vee$ – weight lattice and its dominant cone;
 $\iota: P \rightarrow P^\vee$ – the linear map from the coweight lattice to the weight lattice corresponding to the minimal invariant even bilinear form on the coroot lattice (“level 1”);
for $\lambda = \sum_{i=1}^r m_i \omega_i \in P_+$ we let $|\lambda| = \sum_{i=1}^r m_i$;
 V_{λ^\vee} – irreducible \mathfrak{g} -module with highest weight $\lambda^\vee \in P_+^\vee$.

Current and affine algebras:

$\mathfrak{g}[t] = \mathfrak{g} \otimes \mathbb{C}[t]$ – current algebra;
 $W_{\lambda^\vee}, \mathbb{W}_{\lambda^\vee}$ ($\lambda^\vee \in P_+^\vee$) – local and global Weyl modules for $\mathfrak{g}[t]$;
 $S_\lambda = \times_{i=1}^r S_{m_i}$ – symmetric group attached to $\lambda = \sum_{i=1}^r m_i \omega_i \in P_+$;
 \odot – cyclic product;
 $\widehat{\mathfrak{g}}$ – affine Kac-Moody Lie algebra;
 W and W^a – finite Weyl group and extended affine Weyl group;
 $D_{\ell, \lambda}$ ($\ell \in \mathbb{Z}_{\geq 1}$, $\lambda \in P_+$) – level ℓ weight $\ell\iota(\lambda)$ affine Demazure module;
 $\underline{\lambda} = (\lambda_1, \dots, \lambda_k)$ – collection of integral dominant coweights;
 $\mathbb{D}(\ell, \underline{\lambda}) \simeq R(D(\ell, \lambda_1), \dots, D(\ell, \lambda_k))$ – global Demazure module;

$\mathbb{D}_{\ell, \lambda} \simeq \mathbb{D}(\ell, \underbrace{\omega_1, \dots, \omega_1}_{m_1}, \dots, \underbrace{\omega_r, \dots, \omega_r}_{m_r})$, where $\lambda = \sum_{i=1}^r m_i \omega_i$;
 $\mathcal{A}(\underline{\lambda})$ – highest weight algebra of $\mathbb{D}(\ell, \underline{\lambda})$;
 $\mathcal{A}_\lambda \simeq \mathcal{A}(\underbrace{\omega_1, \dots, \omega_1}_{m_1}, \dots, \underbrace{\omega_r, \dots, \omega_r}_{m_r})$ – highest weight algebra of $\mathbb{D}_{\ell, \lambda}$;
 $M^\vee = \text{Hom}_{\mathcal{A}(\underline{\lambda})}(M, \mathcal{A}(\underline{\lambda}))$ – $\mathcal{A}(\underline{\lambda})$ -dual of an $\mathcal{A}(\underline{\lambda})$ -module M .

Geometry:

$\overline{\text{Gr}}^\lambda \subset \mathbb{P}(D(1, \lambda))$ – spherical affine Schubert variety;
 $\mathbb{A}^\lambda = \text{Spec}(\mathcal{A}_\lambda)$ – colored configuration space on the affine line;
 $\mathbb{A}^{(\underline{\lambda})} = \text{Spec}(\mathcal{A}(\underline{\lambda}))$ – closure of a diagonal stratification stratum in a colored configuration space on the affine line;
 Λ_0^\vee – basic level one integrable affine weight;
 $\Lambda_0^\vee, \Lambda_1^\vee, \dots, \Lambda_m^\vee$ – all level one integrable affine weights;
 $\text{Gr} := \text{Gr}_{G^{\text{ad}}} = G^{\text{ad}}(\mathbb{C}((t)))/G^{\text{ad}}(\mathbb{C}[[t]])$ – affine Grassmannian of G^{ad} ;
 $\text{Gr} \simeq \sqcup_{i=0}^m \text{Gr}(\Lambda_i^\vee)$ – decomposition into irreducible components;
 $\text{Gr}_{\mathbb{A}^k}$ – Beilinson-Drinfeld Grassmannian over \mathbb{A}^k ;
 $\text{Gr}^{\underline{\lambda}} \subset \text{Gr}_{\mathbb{A}^k}$ – Beilinson-Drinfeld spherical Schubert variety;
 $\text{Gr}^{(\underline{\lambda})}$ – partially symmetrized Beilinson-Drinfeld spherical Schubert variety over $\mathbb{A}^{(\underline{\lambda})}$;
 $\mathbb{G}(k)$ – group scheme acting on the Beilinson-Drinfeld Grassmannian;
 \mathcal{L} – very ample determinant line bundle;
 $\mathcal{D}(\ell, \underline{\lambda})$ – locally free sheaf on $\mathbb{A}^{(\underline{\lambda})}$, corresponding to the free $\mathcal{A}(\underline{\lambda})$ -module $\mathbb{D}(\ell, \underline{\lambda})$.

REFERENCES

- [BF1] A.Braverman, M.Finkelberg, *Weyl modules and q -Whittaker functions*, Math. Ann., vol. 359 (1), 2014, pp 45–59.
- [BF2] A.Braverman, M.Finkelberg, *Twisted zastava and q -Whittaker functions*, J. Lond. Math. Soc. (2) 96 (2017), no. 2, 309–325.
- [BF3] A.Braverman, M.Finkelberg, *Semi-infinite Schubert varieties and quantum K -theory of flag manifolds*, J. Amer. Math. Soc. 27 (2014), no. 4, 1147–1168.
- [BCES] A. Brookner, D. Corwin, P. Etingof, S.V. Sam, *On Cohen-Macaulayness of S_n -invariant subspace arrangements*, Int. Math. Res. Not. IMRN 2016, no. 7, 2104–2126.
- [BD1] A. Beilinson, V. Drinfeld, *Chiral algebras*, American Mathematical Society Colloquium Publications, 51, AMS, 2004.
- [BD2] A. Beilinson, V. Drinfeld, *Quantization of Hitchin’s integrable system and Hecke eigensheaves*, Preprint, available at www.math.uchicago.edu/~mitya/langlands
- [BKK] P. Baumann, J. Kamnitzer, A. Knutson, *The Mirkovic-Vilonen basis and Duistermaat-Heckman measures*, <https://arxiv.org/abs/1905.08460>.
- [BL] A. Beauville, Y. Laszlo, *Un lemme de descente*, C. R. Acad. Sci. Paris Sér. I Math. 320 (1995), no. 3, 335–340.
- [BLS] A. Beauville, Y. Laszlo, C. Sorger, *The Picard group of the moduli of G -bundles on a curve*, Compositio Math. 112, 183–216, 1998.
- [CFK] V. Chari, G. Fourier, and T. Khandai, *A categorical approach to Weyl modules*, Transform. Groups, 15(3):517–549, 2010.
- [CI] V.Chari, B.Ion, *BGG reciprocity for current algebras*, Compos. Math. 151 (2015), pp. 1265–1287.

- [CK] S. Cautis, J. Kamnitzer, *Categorical geometric symmetric Howe duality*, Selecta Math. (N.S.) 24 (2018), no. 2, 1593–1631.
- [CL] V. Chari, S. Loktev, *Weyl, Demazure and fusion modules for the current algebra of \mathfrak{sl}_{r+1}* , Adv. Math. 207 (2006), no. 2, 928–960.
- [CP] V. Chari, A. Pressley, *Weyl Modules for Classical and Quantum Affine Algebras*, Represent. Theory 5 (2001), 191–223.
- [CW] S. Cautis, H. Williams, *Cluster theory of the coherent Satake category*, J. Amer. Math. Soc. 32 (2019), no. 3, 709–778.
- [DF] I. Dumanski, E. Feigin, *Reduced arc schemes for Veronese embeddings and global Demazure modules*, arXiv:1912.07988.
- [FBZ] E. Frenkel, D. Ben-Zvi, *Vertex algebras and algebraic curves*, 2nd Edition, Mathematical Surveys and Monograph 88, AMS, 2004.
- [EGL] P. Etingof, E. Gorsky, I. Losev, *Representations of rational Cherednik algebras with minimal support and torus knots*, Advances in Mathematics, Volume 277, 2015.
- [Fu] W. Fulton, *Young Tableaux, with Applications to Representation Theory and Geometry*. Cambridge University Press, 1997.
- [FKL] B. Feigin, A.M. Kirillov, S. Loktev, *Combinatorics and Geometry of Higher Level Weyl Modules*, Advances in the Mathematical Sciences, Ser. 2, 221 (2007) math.QA/0503315
- [FeLo] B. Feigin, S. Loktev, *On generalized Kostka polynomials and the quantum Verlinde rule*, Amer. Math. Soc. Transl. Ser. 2, 194, Adv. Math. Sci., 44, Amer. Math. Soc., Providence, RI, 1999.
- [FeMa1] E. Feigin and I. Makedonskyi, *Semi-infinite Plücker relations and Weyl modules*, International Mathematics Research Notices, rny121, 2018.
- [FeMa2] E. Feigin and I. Makedonskyi, *Vertex algebras and coordinate rings of semi-infinite flags*, Commun. Math. Phys. (2019) 369: 221.
- [FL1] G. Fourier, P. Littelmann, *Tensor product structure of affine Demazure modules and limit constructions*, Nagoya Math. J. 182 (2006), 171–198.
- [FL2] G. Fourier, P. Littelmann, *Weyl modules, Demazure modules, KR-modules, crystals, fusion products and limit constructions*, Adv. Math. 211 (2007), no. 2, 566–593.
- [FiMi] M. Finkelberg, I. Mirković, *Semi-infinite flags I. Case of global curve \mathbb{P}^1* . In Differential topology, infinite-dimensional Lie algebras, and applications, volume 194 of Amer. Math. Soc. Transl. Ser. 2, pages 81–112. Amer. Math. Soc., Providence, RI, 1999.
- [J] A. Joseph, *On the Demazure character formula*, Ann. Sci. École Norm. Sup. (4) 18 (1985), no. 3, 389–419.
- [KMSV] M. Kasatani, T. Miwa, A. N. Sergeev, A. P. Veselov, *Coincident root loci and Jack and Macdonald polynomials for special values of the parameters*, Contemp. Math. 417, Amer. Math. Soc., Providence, RI, 2006, 207–225.
- [Kac] V. Kac, *Infinite-dimensional Lie algebras*, Cambridge: Cambridge University Press (1993).
- [Kam] J. Kamnitzer, *The Beilinson-Drinfeld Grassmannian and symplectic knot homology*, Grassmannians, moduli spaces and vector bundles, 81–94, Clay Math. Proc., 14, Amer. Math. Soc., Providence, RI, 2011.
- [Katol] S. Kato, *Demazure character formula for semi-infinite flag manifolds*, Math. Ann., 2018, 1–33.
- [Kato2] S. Kato, *Frobenius splitting of Schubert varieties of semi-infinite flag manifolds*, arXiv:1810.07106.
- [KN] R. Kodera, K. Naoi, *Loewy series of Weyl modules and the Poincaré polynomials of quiver varieties*, Publ. Res. Inst. Math. Sci. 48 (2012), no. 3, 477–500.
- [KNS] S. Kato, S. Naito, D. Sagaki, *Pieri-Chevalley type formula for equivariant K-theory of semi-infinite flag manifolds*, arXiv:1702.02408

- [Kum1] S. Kumar, *Demazure character formula in arbitrary Kac-Moody setting*, Inv. Math., 89 (1987), 395–423.
- [Kum2] S. Kumar, *Kac-Moody groups, their flag varieties and representation theory*, Progr. Math., 204. Birkhäuser Boston, Inc., Boston, MA, 2002.
- [LS] Y. Laszlo, C. Sorger, *The line bundles on the moduli of parabolic G -bundles over curves and their sections*, Ann. Sci. Ecole Norm. Sup. (4), 30(4), 499–525, 1997.
- [Ma1] O. Mathieu, *Construction du groupe de Kac-Moody et applications*, C. R. Acad. Sci. Paris Sér. I Math. 306 (1988), no. 5, 227–230.
- [Ma2] O. Mathieu, *Formules de caractères pour les algèbres de Kac-Moody générales*, Astérisque 159-160, Société mathématique de France (1988).
- [MVy] I. Mirković, M. Vybornov, *Quiver varieties and Beilinson-Drinfeld Grassmannians of type A* , <https://arxiv.org/abs/0712.4160>
- [Mu1] M. Mustata, *Moduli Spaces and Arcs in Algebraic Geometry*, Cologne, 2006, <http://www.math.lsa.umich.edu/~mmustata/lectures-arcs.pdf>.
- [Mu2] M. Mustata, *Jet schemes of locally complete intersection canonical singularities*, with an appendix by David Eisenbud and Edward Frenkel, Invent. Math. 145 (2001) 397–424.
- [Nash] J.F.Nash, *Arc structure of singularities*, Duke Math. J., 81 (1995), pp. 31–38.
- [Naoi] K.Naoi, *Weyl modules, Demazure modules and finite crystals for non-simply laced type*, Adv. Math. 229 (2012), no. 2, 875–934.
- [Polo] P. Polo, *Variétés de Schubert et excellentes filtrations. Orbites unipotentes et représentations, III*, Astérisque No. 173-174 (1989), 10–11, 281–311.
- [SV] A. Sergeev, A. Veselov, *Deformed quantum Calogero–Moser systems and Lie superalgebras*, Comm. Math. Phys. 245(2004), 249–278.
- [Z1] X. Zhu, *Affine Demazure modules and T -fixed point subschemes in the affine Grassmannian*, Adv. Math. 221 (2009), no. 2, 570–600.
- [Z2] X. Zhu, *An introduction to affine Grassmannians and the geometric Satake equivalence*, In: Geometry of Moduli Spaces and Representation Theory. IAS/Park City Mathematics Series. No.24, 2017. American Mathematical Society , Providence, RI, pp. 59–154.

ILYA DUMANSKI:

DEPARTMENT OF MATHEMATICS, NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF
ECONOMICS, RUSSIAN FEDERATION, USACHEVA STR. 6, 119048, MOSCOW,

and

INDEPENDENT UNIVERSITY OF MOSCOW, 119002, BOLSHOY VLASYEVSKIY PEREULOK
11, RUSSIA, MOSCOW.

Email address: ilyadumnsk@gmail.com

EVGENY FEIGIN:

DEPARTMENT OF MATHEMATICS, NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF
ECONOMICS, RUSSIAN FEDERATION, USACHEVA STR. 6, 119048, MOSCOW,

and

SKOLKOVO INSTITUTE OF SCIENCE AND TECHNOLOGY, SKOLKOVO INNOVATION CENTER,
BUILDING 3, MOSCOW 143026, RUSSIA.

Email address: evgfeig@gmail.com

MICHAEL FINKELBERG:

DEPARTMENT OF MATHEMATICS, NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF
ECONOMICS, RUSSIAN FEDERATION, USACHEVA STR. 6, 119048, MOSCOW,

SKOLKOVO INSTITUTE OF SCIENCE AND TECHNOLOGY, SKOLKOVO INNOVATION CENTER,
BUILDING 3, MOSCOW 143026, RUSSIA, *and*

INSTITUTE FOR INFORMATION TRANSMISSION PROBLEMS OF RAS, MOSCOW, RUSSIA.

Email address: fnklberg@gmail.com