

The CARMENES search for exoplanets around M dwarfs

A super-Earth planet orbiting HD 79211 (GJ 338 B) [★]

E. González-Álvarez¹, M. R. Zapatero Osorio¹, J. A. Caballero¹, J. Sanz-Forcada¹, V. J. S. Béjar^{2,3},
L. González-Cuesta^{2,3}, S. Dreizler⁴, F. F. Bauer⁵, E. Rodríguez⁵, L. Tal-Or^{6,4}, M. Zechmeister⁴, D. Montes⁷,
M. J. López-González⁵, I. Ribas^{8,9}, A. Reiners⁴, A. Quirrenbach¹⁰, P. J. Amado⁵, G. Anglada-Escudé^{5,11}, M. Azzaro¹²,
M. Cortés-Contreras¹, A. P. Hatzes¹³, T. Henning¹⁴, S. V. Jeffers⁴, A. Kaminski¹⁰, M. Kürster¹⁴, M. Lafarga^{8,9},
J. C. Morales^{8,9}, E. Pallé^{2,3}, M. Perger^{8,9}, and J. H. M. M. Schmitt¹⁵

¹ Centro de Astrobiología (CSIC-INTA), Carretera de Ajalvir km 4, 28850 Torrejón de Ardoz, Madrid, Spain

² Instituto de Astrofísica de Canarias, Av. Vía Láctea s/n, 38205 La Laguna, Tenerife, Spain

³ Departamento de Astrofísica, Universidad de La Laguna, 38206 La Laguna, Tenerife, Spain

⁴ Institut für Astrophysik, Georg-August-Universität, Friedrich-Hund-Platz 1, 37077 Göttingen, Germany

⁵ Instituto de Astrofísica de Andalucía (IAA-CSIC), Glorieta de la Astronomía s/n, 18008 Granada, Spain

⁶ Department of Physics, Ariel University, Ariel, 40700, Israel

⁷ Departamento de Física de la Tierra y Astrofísica & UPARCOS-UCM (Ins. de Física de Partículas y del Cosmos de la UCM), Facultad de Ciencias Físicas, Universidad Complutense de Madrid, 28040 Madrid, Spain

⁸ Institut de Ciències de l'Espai (ICE, CSIC), Campus UAB, c/ de Can Magrans s/n, 08193 Bellaterra, Barcelona, Spain

⁹ Institut d'Estudis Espacials de Catalunya (IEEC), 08034 Barcelona, Spain

¹⁰ Landessternwarte, Zentrum für Astronomie der Universität Heidelberg, Königstuhl 12, 69117 Heidelberg, Germany

¹¹ School of Physics and Astronomy, Queen Mary University of London, 327 Mile End Road, London E1 4NS, UK

¹² Centro Astronómico Hispano-Alemán (CSIC-MPG), Observatorio Astronómico de Calar Alto, Sierra de los Filabres, 04550 Górgal, Almería, Spain

¹³ Thüringer Landessternwarte Tautenburg, Sternwarte 5, 07778 Tautenburg, Germany

¹⁴ Max-Planck-Institut für Astronomie, Königstuhl 17, 69117 Heidelberg, Germany

¹⁵ Hamburger Sternwarte, Gojenbergsweg 112, 21029 Hamburg, Germany

Received 05 November 2019 / Accepted 06 March 2020

ABSTRACT

Aims. We report on radial velocity time series for two M0.0 V stars, GJ 338 B and GJ 338 A, using the CARMENES spectrograph, complemented by ground-telescope photometry from Las Cumbres and Sierra Nevada observatories. We aim to explore the presence of small planets in tight orbits using the spectroscopic radial velocity technique.

Methods. We obtained 159 and 70 radial velocity measurements of GJ 338 B and A, respectively, with the CARMENES visible channel between 2016 January and 2018 October. We also compiled additional relative radial velocity measurements from the literature and a collection of astrometric data that cover 200 a of observations to solve for the binary orbit.

Results. We found dynamical masses of $0.64 \pm 0.07 M_{\odot}$ for GJ 338 B and $0.69 \pm 0.07 M_{\odot}$ for GJ 338 A. The CARMENES radial velocity periodograms show significant peaks at 16.61 ± 0.04 d (GJ 338 B) and $16.3^{+3.5}_{-1.3}$ d (GJ 338 A), which have counterparts at the same frequencies in CARMENES activity indicators and photometric light curves. We attribute these to stellar rotation. GJ 338 B shows two additional, significant signals at 8.27 ± 0.01 and 24.45 ± 0.02 d, with no obvious counterparts in the stellar activity indices. The former is likely the first harmonic of the star's rotation, while we ascribe the latter to the existence of a super-Earth planet with a minimum mass of $10.27^{+1.47}_{-1.38} M_{\oplus}$ orbiting GJ 338 B. We have not detected signals of likely planetary origin around GJ 338 A.

Conclusions. GJ 338 Bb lies inside the inner boundary of the habitable zone around its parent star. It is one of the least massive planets ever found around any member of stellar binaries. The masses, spectral types, brightnesses, and even the rotational periods are very similar for both stars, which are likely coeval and formed from the same molecular cloud, yet they differ in the architecture of their planetary systems.

Key words. techniques: radial velocities – stars: binaries: visual: STF 1321 – stars: late-type – stars: planetary systems

1. Introduction

Despite the significant number of discoveries of planetary systems with M dwarf primaries (e.g., Anglada-Escudé et al. 2012; Bonfils et al. 2013; Astudillo-Defru et al. 2015; Suárez Mascareño et al. 2017; Affer et al. 2019), the properties and the statistics of planets hosted by low-mass stars remain poorly constrained. We are far from understanding such fundamental ques-

Send offprint requests to: Esther González-Álvarez

e-mail: egonzalez@cab.inta-csic.es

[★] Tables B.1–B.6 are only available in electronic form at the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/>

tions as how planetary systems form and how their architecture changes with the mass of the central star.

Various existing radial velocity (RV) surveys focus on M dwarfs for many reasons, including: (i) they are the most abundant stellar population in the solar vicinity (Henry et al. 2006), (ii) the occurrence of small planets (typically with sizes 1–4 times that of the Earth) is increasingly higher toward late spectral types at all orbital periods explored by the *Kepler* mission (Dressing & Charbonneau 2013; Morton & Swift 2014; Dressing & Charbonneau 2015; Mulders et al. 2015; Gaidos et al. 2016), (iii) planets of low-mass stars can be detected easily with the radial velocity technique, and (iv) the appealing possibility of spatially resolving terrestrial planets in the habitable zone of the nearest stars using next-generation instrumentation (e.g., Howard et al. 2012; Dressing & Charbonneau 2013; Sozzetti et al. 2013).

However, M dwarfs also have their own issues, for example, they have large convective regions and are on average more active than solar-like stars (Leto et al. 1997; Osten et al. 2005). It is crucial to study the stellar activity of M dwarfs together with the analysis of the presence of a planetary signal in their RV time series data. Many works (e.g., Hartman et al. 2009, 2010; Suárez Mascareño et al. 2015; Newton et al. 2016; Díez Alonso et al. 2019) have discussed how the rotational periods of M dwarfs often coincide with the orbital periods of planets in the expected habitable zone of these stars. The closeness between these periodicities represent an observational challenge since the signal of the planet has to be disentangled from the magnetic activity contribution of the star (Damasso et al. 2018).

Instruments such as CARMENES (Quirrenbach et al. 2016) and GIARPS (GIANO-B+HARPS-N, Claudi et al. 2017) minimize the stellar activity problem by obtaining simultaneous optical (VIS) and near-infrared (NIR) data. The RV signal of planetary origin is independent of the observed wavelength, whereas the amplitude of the signal due to stellar activity may strongly depend on wavelength (typically larger at shorter wavelengths; e.g., Martín et al. 2006; Huélamo et al. 2008; Carleo et al. 2018). The main goal of CARMENES is to discover and characterize Earth-like planets around an initial sample of about 300 M dwarfs (Reiners et al. 2018b). To date, the program has already confirmed eight planet candidates from large-scale photometric and spectroscopic surveys (e.g., Trifonov et al. 2018; Sarkis et al. 2018) and detected more than ten new planets (e.g., Reiners et al. 2018a; Kaminski et al. 2018; Luque et al. 2018, 2019; Ribas et al. 2018; Nagel et al. 2019; Perger et al. 2019; Zechmeister et al. 2019; Morales et al. 2019).

Here, we present the detailed CARMENES RV analysis of the M0.0 V stars GJ 338 A (HD 79210) and GJ 338 B (HD 79211), a wide binary system with similar mass stellar components. In Section 2, we introduce the binary and the known properties of each stellar member. Section 3 presents all new and literature RV data employed in this paper for the study of the binary orbit and the exploration of the presence of small planets around each star. We also present the recently obtained optical photometry that helped us confirm the rotation period of the stars independently of the spectroscopic measurements. In Section 4, we provide the analysis of the stellar binary orbit by combining all available astrometry and literature RVs with the main goal of determining the dynamical masses of the two stars, and the detailed study of the CARMENES RV data and the new photometry aimed at identifying low-mass planets. The properties of the newly discovered planet orbiting GJ 338 B are given in Section 5. A brief discussion on the implications of this finding and the conclusions of this paper appear in Sections 6 and 7.

2. Target stars

GJ 338 B is a nearby, bright M0.0 V star at a distance of 6.334 pc (*Gaia* Data Release 2, DR2; Gaia Collaboration et al. 2018). As described in detail by Abt & Levy (1973) and Morbey & Griffin (1987), this star, together with the twin M0.0 V star HD 79210 (GJ 338 A), forms the high common proper motion pair Σ Struve STF 1321 (WDS J09144+5214, ADS 7251). Occasionally, they have been classified as K7 V stars, but here we follow the latest classification by Alonso-Floriano et al. (2015). These authors determined spectral type M0.0 V for both stars (actually, GJ 338 A is the M0.0 V spectral standard star used in that work). The most updated stellar parameters of GJ 338 B and A are compiled in Table 1.

The projected separation between GJ 338 A and B is 108.54 au (17''.2). Cortés-Contreras et al. (2017) did not find any close companions to either GJ 338 A or B during their high-resolution imaging survey. The Washington Double Star catalog (Mason et al. 2001) tabulates two additional components C and D in the system STF 1321, located to date at slightly over 2 arcmin to the northeast and southeast, respectively, of the central, much brighter pair AB. With the latest *Gaia* DR2 data we confirm that C and D are background stars located further away and with very different proper motions as compared to our system.¹

According to van den Bos (1955), the first orbital parameter determinations of the pair STF 1321 were done by Hopmann (1954) and Güntzel-Lingner (1954). The former author estimated a long orbital period, $P_{\text{orb}} = 1555$ a, and a large semimajor axis, $a = 22''36$, whereas the latter author published a short period, $P_{\text{orb}} = 687$ a and a small semimajor axis $a = 16''56$. Since these first determinations, the orbital parameters of the visual binary have been revisited on a few occasions (Kiselev & Kiyeva 1980; Romanenko & Chentsov 1994; Kiselev & Romanenko 1996; Malkov et al. 2012) but none has surpassed the quality of the fit found by Chang (1972), who measured an intermediate period, $P_{\text{orb}} = 975$ a and semimajor axis $a = 16''72$ (ORB6 orbit grade = 4, following Hartkopf et al. 2001). Chang (1972) also derived stellar masses of $0.41 \pm 0.03 M_{\odot}$ and $0.73 \pm 0.05 M_{\odot}$ for GJ 338 A and B, respectively. These mass derivations, where the secondary is significantly more massive than the primary, is reversed with respect to what is expected from the brightness of stars: GJ 338 B, supposedly the most massive component, is ~ 0.1 mag fainter than GJ 338 A in all optical and near-infrared filters. These results are also in apparent disagreement with the latest, more similar mass determinations by Schweitzer et al. 2019 (see below). In particular, these authors determined masses of $0.58\text{--}0.61 M_{\odot}$ for each stellar component using different methods (see also Boyajian et al. 2012, Gaidos et al. 2014, and Reiners et al. 2018b).

The chemical composition analysis available in the literature reveals that GJ 338 A and B have a slightly sub-solar metallicity with reported measurements ranging from $[\text{Fe}/\text{H}] = 0.0$ to -0.35 dex (Jenkins et al. 2009; Gaidos & Mann 2014; Neves et al. 2013; Sharma et al. 2016). The recent study by Passegger et al. (2018) reported a metallicity of 0.07 ± 0.16 and -0.03 ± 0.16 for GJ 338 A and B, which lies close to the solar metallicity but also agrees with previous metallicity derivations within the quoted error bars. Based on these results, no significant difference in the chemical composition of GJ 338 A and B is expected.

Both stars were cataloged as probable kinematically young stars in the Local Association by Tetzlaff et al. (2011). This is consistent with their membership in the Galactic young disk

¹ "C": 2MASS J09143546+5242095; "D": TYC 3806–1033–1.

Table 1: Stellar parameters of GJ 338 binary system.

Parameters	GJ 338 A	GJ 338 B	Ref. ^a
Other name	HD 79210	HD 79211	
Karmn	J09143+526	J09144+526	AF15a
α (J2000)	09:14:22.78	09:14:24.68	<i>Gaia</i> DR2
δ (J2000)	+52:41:11.8	+52:41:10.9	<i>Gaia</i> DR2
G (mag)	6.9689 \pm 0.0005	7.0477 \pm 0.0004	<i>Gaia</i> DR2
J (mag)	4.89 \pm 0.04	4.779 \pm 0.174	2MASS
Spectral type	M0.0 V	M0.0 V	AF15a
d (pc)	6.334 \pm 0.002	6.334 \pm 0.002	<i>Gaia</i> DR2
$\mu_\alpha \cos \delta$ (mas a ⁻¹)	-1546.10 \pm 0.06	-1573.12 \pm 0.06	<i>Gaia</i> DR2
μ_δ (mas a ⁻¹)	-569.13 \pm 0.06	-660.12 \pm 0.06	<i>Gaia</i> DR2
U (km s ⁻¹)	-42.20 \pm 0.36	-44.01 \pm 0.36	CC16
V (km s ⁻¹)	-14.99 \pm 0.10	-17.44 \pm 0.10	CC16
W (km s ⁻¹)	-23.73 \pm 0.34	-23.10 \pm 0.34	CC16
T_{eff} (K)	4024 \pm 51	4005 \pm 51	Schw19
$\log g$ (cgs)	4.68 \pm 0.07	4.68 \pm 0.07	Schw19
[Fe/H] (dex)	-0.05 \pm 0.16	-0.03 \pm 0.16	Schw19
L (L_\odot)	0.0789 \pm 0.0038	0.0792 \pm 0.0031	Schw19
M (M_\odot)	0.69 \pm 0.07	0.64 \pm 0.07	This work
	0.591 \pm 0.047	0.596 \pm 0.042	Schw19
R (R_\odot)	0.58 \pm 0.02	0.58 \pm 0.03	Schw19
$v \sin i$ (km s ⁻¹)	2.9 \pm 1.2	2.3 \pm 1.5	GG05, Rein18
P_{rot} (d)	16.3 ^{+3.5} _{-1.3}	16.61 \pm 0.04	This work
L_X (10 ²⁷ erg s ⁻¹)	5.2 \pm 0.3	5.0 \pm 0.1	This work
$\log R'_{\text{HK}}$...	-4.4	MA10
Age (Ga)	1–7	1–7	This work

Notes. ^a *Gaia* DR2: Gaia Collaboration et al. (2018); 2MASS: Skrutskie et al. (2006); AF15a: Alonso-Floriano et al. (2015); CC16: Cortés-Contreras (2016); GG05: Głębcki & Gnaniński (2005); Rein18: Reiners et al. (2018b); Schw19: Schweitzer et al. (2019); MA10: Martínez-Arnáiz et al. (2010).

(age ≤ 1 Ga) reported by Cortés-Contreras (2016), who pointed out that, despite their Galactic kinematics indicating youth, other photometric and spectroscopic observables of GJ 338 A and B do not support a young age. Eiroa et al. (2013) reported the non-detection of mid-infrared flux excesses in GJ 338 B using *Herschel* data, thus discarding the presence of a dusty disk, which is compatible with the most likely stellar age of a few Ga.

GJ 338 A and GJ 338 B are active and show intense chromospheric fluxes (Moutou et al. 2017). This is consistent with the slightly fast rotation of components A and B, which have a projected rotational velocity $v \sin i = 2.9 \pm 1.2$ and 2.3 ± 1.5 km s⁻¹ (Głębcki & Gnaniński 2005; Reiners et al. 2018b). GJ 338 A and B have been observed in X-rays by *Chandra* on 29/12/2012, and we derived their luminosities at $L_X = 5.2 \pm 0.3 \times 10^{27}$ and $5.0 \pm 0.1 \times 10^{27}$ erg s⁻¹, respectively. From their X-ray luminosities, both components have a similar activity level with values typically found among active stars of their spectral type.

This X-ray luminosity is compatible with an age of 1 to 7 Ga following the relation by Sanz-Forcada et al. (2011). Based on the activity index $\log R'_{\text{HK}} \sim -4.4$ for GJ 338 B, Martínez-Arnáiz et al. (2010) and Isaacson & Fischer (2010) estimated that this star is active with an expected radial velocity jitter of about 4–10 m s⁻¹.

3. Observations

3.1. Radial velocities from the literature

Absolute radial velocities of each star are available in a number of publications and archives: Abt & Levy (1973), Beavers & Eit-

ter (1986), Bopp & Meredith (1986), and the *Gaia* DR2 catalog (Soubiran et al. 2018). With the only exception of the *Gaia* DR2 data, all other absolute velocities are affected by rather large error bars (typically ≥ 1 km s⁻¹) and dispersion, and we did not consider them in our analysis. More recent radial velocities with smaller associated uncertainties (typically from several to tens of m s⁻¹) come from the ELODIE (Baranne et al. 1996) and SOPHIE (Perruchot et al. 2008) high-resolution spectrographs. These data were taken as part of the ground-based efforts intended for the calibration of the velocities of the *Gaia* DR2 catalog (Soubiran et al. 2018). The ELODIE and SOPHIE radial velocities were obtained using different masks at the pipeline level, which had an impact on the determination of the velocity zero points. Unfortunately, the SOPHIE data of GJ 338 A and B were analyzed using different masks per star, thus preventing us from a direct comparison between the two stellar components of the binary. As pointed out by Soubiran et al. (2018), the error associated with the determination of the velocity zero point offsets due to the different masks employed during the velocity computations was large for the M dwarfs ($\approx \pm 0.4$ km s⁻¹), and it was not precisely quantified. The ELODIE radial velocities, however, were reduced with the same mask for the two stars, and we used these measurements in our study of the orbital parameters of the stellar pair. Eleven ELODIE radial velocities covered the time interval between January 1995 and February 2000. They have a dispersion of 10–30 m s⁻¹, which is about one order of magnitude worse than the CARMENES data (Section 3.2). Therefore, we do not use these data to explore the presence of small-mass planets around GJ 338 A and B. However, these measurements have sufficient quality for obtaining the orbital solution fit of the stellar binary.

In addition to the ELODIE data, and for the characterization of the stellar binary orbit, we employed the more than 30 relative radial velocities obtained for each member of the stellar pair with the High Resolution (HIRES) spectrograph (Vogt et al. 1994) of the Keck telescope published by Butler et al. (2017). These velocities span a period of 15.7 a and were later corrected for the presence of systematic effects affecting the nightly zero points (a discontinuous jump caused by interventions on the instrument in 2004 and a long-term drift; see Tal-Or et al. 2019). The HIRES RVs have associated a mean error bar value of 1.6 m s⁻¹. The relative HIRES RVs of GJ 338 A and B given by Tal-Or et al. (2019) are displayed in Fig. 1, where the trend due to the orbital motion of the pair around the center of mass of the system is clearly seen for each star. To bring the HIRES relative RVs given by Tal-Or et al. (2019) to an absolute RV calibration, we used the corresponding values reported by Nidever et al. (2002), which fully overlap with the first years of HIRES observations. The ELODIE and the HIRES data altogether cover over 20 a of observations. ELODIE absolute velocities were taken to the HIRES reference by considering the measurements of the two datasets at common observing epochs.

The radial velocity survey of Gagné et al. (2016) using the CSHELL spectrograph (Tokunaga et al. 1990) at the NASA Infrared Telescope Facility (IRTF) included observations of GJ 338 A and B at near-infrared wavelengths (*K* band) between 2014 March and 2014 December. They do not overlap with any of the ELODIE, HIRES, or CARMENES data. Gagné et al. (2016) concluded that hot, warm and cool Jupiter planets more massive than 2.3, 4.2, and 20 M_{Jup} and with orbital periodicities of 1–10, 10–100, and 100–1000 d, respectively, can be excluded around the B component (our main CARMENES target) with a confidence of 95%. The CSHELL/IRTF relative velocities are not considered in our planetary search of Section 4 because they

have associated uncertainties of the order of $6\text{--}18\text{ m s}^{-1}$, that is, more than three times larger than the typical CARMENES error bar. In addition, they are not included in our analysis of the stellar binary because no absolute velocities were reported by the authors and no calibration to absolute data is possible due to the lack of overlapping dates with any other data in hand.

3.2. CARMENES radial velocity time series

We obtained new RVs for each member of the stellar binary using the CARMENES fibre-fed, échelle spectrograph. CARMENES is installed at the 3.5 m telescope of the Calar Alto Observatory in Almería (Spain). It was specifically designed to deliver high-resolution spectra at optical (resolving power $R = 94,600$) and near-infrared ($R = 80,500$) wavelengths covering from 520 to 1710 nm without any significant gap, using two different spectrograph arms: the VIS and NIR channels, respectively (Quirrenbach et al. 2016).

The CARMENES M-dwarf survey started in January 2016. The original sample of stars contained 324 M dwarfs (Caballero et al. 2016a; Reiners et al. 2018a,b), including GJ 338 B. Raw data are automatically reduced with the CARACAL pipeline (Caballero et al. 2016b). Relative radial velocities are extracted separately for the VIS and NIR channels using the SERVAL software (Zechmeister et al. 2018), which computes the average of all individually derived RVs order by order of the échelle observations. The results of the CARMENES planetary survey around M dwarfs reveal that the instrument is able to deliver time series of relative RVs with root mean square (*rms*) values of 3.5 and 9.0 m s^{-1} for the CARMENES VIS and NIR channels (Reiners et al. 2018b).

The systematic RV errors in spectral orders highly contaminated by telluric contribution, which is the case of CARMENES NIR data, can be increased by unmasked detector defects (Reiners et al. 2018b). Therefore, for the NIR channel, we carefully selected the orders that were included in the computation of the NIR RVs. These orders, selected to provide the smallest *rms* of the NIR RVs, were chosen after analyzing the entire CARMENES M-dwarf sample.

3.2.1. GJ 338 B

GJ 338 B is the main target of this paper. We used 159 CARMENES VIS and 120 NIR relative RVs obtained between 2016 January and 2018 October. There are fewer NIR RVs because we selected only the NIR RVs taken after a technical intervention in the NIR channel in October 2016. GJ 338 B was observed with a typical cadence of one RV measurement every few to several days to sample short periodicities. All measurements employed in this work are provided in Tables B.1 (VIS) and B.4 (NIR). All data are corrected from barycentric motion, secular acceleration, instrumental drift, and nightly zero points (Trifonov et al. 2018). CARMENES VIS relative RVs of GJ 338 B as given in the CARMENES catalog of observations are depicted as a function of observing epoch in the middle panel of Fig. 1. The decreasing trend due to the orbital motion of the binary agrees with the trend delineated by the HIRES relative RVs. The right panel of Fig. 1 shows CARMENES NIR relative RVs of GJ 338 B. The slope of the NIR data is not as clear as that of the VIS RVs likely due to the larger uncertainties (and therefore, dispersion) of the NIR RVs.

Table 2: Photometric seasons available for GJ 338 A and B.

Obs. ^a	Filter	Season	ΔT	N_{obs}	σ
			(d)		(mmag)
LCO	V	2018 Mar 5–27	22	1830	14.4
LCO	i	2018 Mar 5–27	22	2100	13.4
LCO	V	2018 Dec 2 – 2019 Feb 27	87	350	10.7
LCO	B	2018 Dec 2 – 2019 Feb 27	87	330	8.0
SNO	V	2019 Feb 22 – 2019 Apr 13	50	819	6.1
SNO	B	2019 Feb 22 – 2019 Apr 13	50	839	5.9

Notes. ^a Observatories – LCO: Las Cumbres; SNO: Sierra Nevada.

3.2.2. GJ 338 A

The primary component of the stellar binary was also observed as part of the CARMENES M-dwarf program with a total of 70 RV measurements in the VIS channel taken between 2016 March and 2019 January. These VIS RVs, also relative and corrected from barycentric motion, secular acceleration, instrumental drift, and nightly zero points, are provided in Table B.2. GJ 338 A was not observed regularly at the beginning of the CARMENES M-dwarf project: the first three RVs were obtained more than one year before the remaining RV time series data. After 2017 December 18, GJ 338 A was monitored with a similar observing cadence as GJ 338 B. CARMENES VIS relative RVs are depicted as a function of observing epoch in the left panel of Fig. 1; they are shown together with the HIRES RV data of Tal-Or et al. (2019). Because the time coverage of the CARMENES data of GJ 338 A with a good time sampling is significantly shorter than that of GJ 338 B, we could not explore the same orbits as those of our study of GJ 338 B. These RVs were also used to study the orbital motion of the stellar pair (see Section 4.1) and to act as a reference (for its similarity in spectral type) in our search for low-mass planets around GJ 338 B.

3.3. Photometric time series

We obtained optical photometric time series of both GJ 338 A and B contemporaneous to the CARMENES data, aiming to derived the stellar rotational periods via the differential photometry technique. Images were acquired with telescopes of Las Cumbres Observatory Global Telescope (LCO, Brown et al. 2013) and Sierra Nevada Observatory (SNO). The 40-cm LCO telescopes used in our program are equipped with a $3k \times 2k$ SBIG CCD camera with a pixel scale of $0''.571$ providing a field of view of $29''.2 \times 19''.5$. The 90-cm Ritchey-Chrétien SNO telescope has a VersArray $2k \times 2k$ CCD camera with a square field of view of $13''.2 \times 13''.2$ (pixel scale of $0''.39$ on sky, Rodríguez et al. 2010). We applied a 2×2 binning to the detector. Raw frames were reduced following standard procedures at optical wavelengths, that is, bias subtraction and flat-field correction. Bad pixels were conveniently masked using well-tested masks provided by the observatories. In the case of LCO images, we used the *banzai* reduction products (McCully et al. 2018).

There were two photometric campaigns using LCO telescopes between 2018 March and 2019 February, and a third campaign using the SNO telescope from 2019 February to April. All three are summarized in Table 2, where we provide observing filters, the number of days covered per observing season, the number of photometric measurements, and the dispersion of the differential photometry. In the first photometric season, from 2018 March 5 to 27 (22 days of continuous monitoring), we acquired




Fig. 1: *Left panel:* GJ 338 A HIRES (blue) and CARMENES (red) relative RV measurements at visible wavelengths. *Middle panel:* GJ 338 B HIRES (blue) and CARMENES (red) relative RV measurements at visible wavelengths. The time series data of each instrument has been normalized to a mean null velocity. Open circles indicate measurements that deviate more than 2σ from the observed velocity trend. *Right panel:* CARMENES relative RVs of GJ 338 B in the near-infrared.

a total of 1800 and 2300 usable images of GJ 338 A and B using the V and i filters and a 2×2 binning of the detector. The on-source integration time was set at 1 s, and 100 consecutive images were typically acquired per filter and per night. On a few nights, the 100-frame sequence was acquired twice or three times. Weather conditions were variable from night to night as was also the seeing, with values varying between $1''.0$ and $5''.0$. The root-mean-square (rms) of the data are 14.4 mmag for the V -band light curve and 13.4 mmag for the i -band time series. The two light curves artificially normalized to zero magnitude are depicted in the top panels of Fig. 2.

In the second LCO campaign, from 2018 December through 2019 February (almost three months of continuous monitoring), we followed a slightly different observing strategy: between three and five images were typically obtained in the B and V filters throughout the observing nights resulting in more than 350 and 330 usable B - and V -band images whose differential photometry is shown in the middle panel of Fig. 2 (no normalization to zero magnitude was applied here). In this second season, the rms of the V -band light curve (10.7 mmag) is smaller than that of the first season. The third photometric season (2019 February 22 through April 13, that is, 50 days of nearly continuous monitoring) was acquired using the B and V filters immediately after the last LCO campaign. Typically, a total of 50 images were acquired with the SNO telescope per night. Exposure time changed between 6 s and 20 s, depending on the seeing conditions. The SNO light curves are illustrated in the bottom panels of Fig. 2, and have an rms of 6.1 mmag (V) and 5.9 mmag (B). Also the LCO B -band light curve shows a smaller dispersion than the V -band data.

We used a circular apertures with radius of seven and 20 pixels to determine the LCO and SNO photometry of GJ 338 A and B, respectively. We checked that these apertures minimized the rms of the resulting light curves. The field of view registered by the LCO and SNO detectors is unfortunately devoid of bright stars that can act as reference sources for the differential photometry. Therefore, we derived the differential magnitude between the two components of the stellar binary (component B – component A). Because both stars share very similar spectral type, we did not apply any color correction to the differential photometry. With such a procedure, it is not possible to discern which star is the source of any photometric variability, or if the two stars are variable simultaneously.

Given the large number of contemporaneous B , V , and i individual measurements available for both stellar components, we

also derived differential $B - V$ and $V - i$ color light curves and found that GJ 338 A and B indeed show different colors, and that these colors appear rather stable throughout all monitoring campaigns. The relative color differences, although small, are significant in terms of the achieved photometric precision. GJ 338 B has $B - V$ and $V - i$ indices that are, on average, 12.5 ± 0.6 and 34.2 ± 0.7 mmag redder than those of GJ 338 A, where the error bars were obtained as the quadratically-summed photometric dispersions of both bands divided by the squared root of the number of measurements. These differences can be interpreted as both stars having slightly different temperatures (GJ 338 B would be cooler than GJ 338 A). Another possibility is a differing chemical composition of the stellar atmospheres (higher metallicity induces redder colors), but we give less credit to this scenario because the two stars are likely coeval and were probably formed from the same molecular cloud. GJ 338 B is also slightly fainter than the A component ($\Delta B = 110.2 \pm 0.4$, $\Delta V = 97.7 \pm 0.6$, and $\Delta i = 65.6 \pm 0.2$ mmag), which combined with its redder nature may indicate that both stars have similar but not identical masses. The photometric properties of GJ 338 B (fainter and redder) suggest that this star is slightly less massive than GJ 338 A; this is compatible with the results presented in Section 4.1 but strongly disagrees with the mass derivations of Chang (1972).

GJ 338 A and B were also photometrically observed by the All-Sky Automated Survey for Supernovae survey (ASAS-SN, Kochanek et al. 2017). The ASAS-SN light curves show strong peak to peak variations. This behavior is far from what is seen in the LCO and SNO data. The ASAS-SN photometry was automatically derived from a 2-pixel radius aperture (i.e., about two FWHM in diameter) and the background was estimated from a 7–10 pixel radius annulus surrounding each of the two stars. The pixel scale of ASAS-SN detector is $8''.0$. Given the projected spatial separation of the stellar binary ($\rho \sim 17''.2$), it becomes obvious that the ASAS-SN parameters for extracting the photometry of the system were not adequate and yield contaminated and useless light curves for GJ 338 A and B.

4. Analysis

4.1. Stellar binary orbital solution

In an attempt to constrain the orbital parameters of the stellar pair GJ 338 A and B, we considered all astrometric and spectroscopic data available to us in the pre-CARMENES era. As for the astrometry, we used the angular separations and position angles




Fig. 2: Differential photometry (GJ 338 B – GJ 338 A) light curves obtained with the LCO and SNO telescopes. First and second panels correspond to the first photometric observing season covering a time interval of 22 days; the third panel illustrates the light curve for the second observing season covering 87 days of continuous monitoring; and the last observing season spanning 50 days is shown in the fourth and fifth panels. With the exception of the third panel, all photometry is artificially set to a mean value of zero magnitude.

compiled by the Washington Double Star Catalog² (WDS, Mason et al. 2001). This dataset provides astrometric measurements from mid 1821 through the first quarter of 2019. We completed the astrometric data using the SNO images from Section 3.3. By employing the 2MASS (Skrutskie et al. 2006) coordinates of all sources detected in the SNO frames, with the exception of the high proper motion targets GJ 338 A and B, we obtained the astrometric solution for seven different epochs in 2019, thus extending the time coverage of the Washington Double Star Catalog by two additional years. We found a plate scale of 0.388094 ± 0.000030 arcsec/pix (the pixels have the same length in the x - and y -axis within the quoted uncertainty) and that the SNO frames are 1.5 deg off the standard north–east orientation. This was corrected from the final measurements. We derived that the angular position of GJ 338 B with respect to the A component has changed by about 58 deg in 198 a of available astrometric observations. All astrometric data, including our latest derivations,




Fig. 3: *Top panel*: observed astrometric positions of GJ 338 B (blue dots) around the star GJ 338 A (black star located at 0,0 coordinates) expressed in $\Delta\delta$ vs. $\Delta\alpha \cos \delta$. The black ellipse (dashed line) corresponds to the adopted Keplerian solution, which lies close to the region of the largest concentration of possible MCMC solutions. The oldest data and most recent astrometric data correspond to the year 1821 and 2019, respectively. The inner panel is a zoom-in around the astrometric data. Only the adopted orbital solution is shown with the black dashed line. *Bottom panel*: residuals (± 0.11 in $\Delta\alpha \cos \delta$ and ± 0.14 in $\Delta\delta$) after removing the adopted orbital solution plotted as a function of observing epoch ($\Delta\alpha \cos \delta$: blue dots; $\Delta\delta$: red dots).

in the form of angular separation (ρ) and position angle (θ) are given in Table B.5 and displayed in Fig. 3. As for the RVs, we used the ELODIE and HIRES absolute RVs of GJ 338 A and B as explained in Section 3.1.

To find the best Keplerian solution to the orbit of the stellar pair, and based on our knowledge of the system, any orbital solution has to reproduce the observed astrometric and radial velocity data and to be compliant with the following: a) The total mass of the system should fall in the interval 1.03 – $1.38 M_{\odot}$, which is obtained from the 1σ smallest and largest individual values assigned to each component in Table B.1 of Schweitzer et al. (2019). According to these authors, the most likely total mass of the system is $1.187 \pm 0.063 M_{\odot}$. b) The two stellar components have very similar spectral type, effective temperature, magnitude, and luminosity (Passegger et al. 2018, 2019). Therefore, we expect the mass ratio, q , to be close to unity. In fact, Schweitzer et al. (2019) determined averaged masses of $0.591 \pm 0.047 M_{\odot}$ for GJ 338 A and $0.596 \pm 0.042 M_{\odot}$ for GJ 338 B (errors are also averaged from those given by the authors). These determinations were based on the analysis of CARMENES spectra and do not rely on astrometric or radial velocity observations.

² <https://www.usno.navy.mil/USNO/astrometry/optical-IR-prod/wds/WDS>




Fig. 4: *First panel*: radial velocity data and the joint astrometric and spectroscopic solution (black line) of the stellar binary system formed by GJ 338 A and GJ 338 B. The uncertainties are depicted with a gray color. ELODIE and HIRES absolute RVs are also shown. *Second and third panels*: zoom-in around the RVs of GJ 338 A and GJ 338 B, respectively. The straight line fit to the data is plotted as a green line. Given the high level of agreement between the adopted Keplerian solution and the straight line, the latter remains nearly hidden by the former in the two panels. *Fourth panel*: residuals of the RV observations minus the adopted Keplerian solution are shown with different symbols for the two stars. *Bottom panel*: ELODIE, HIRES, SOPHIE and CARMENES differential radial velocity curve ($v_A - v_B$) as a function of observing epoch. The values of the adopted Keplerian solution, which were fit using the ELODIE and HIRES data only, are shown with a black dashed line, errors in gray, and the straight line fit to the data is depicted with a green line. All four panels, from the second to the bottom one, have the same scale in the x-axis.

The orbital parameters of the stellar pair GJ 338 A and GJ 338 B were obtained by simultaneously fitting the astrometric and RV data. Our code is based on PyMC3, which is a flexible and high-performance model building language and

inference engine that scales well to problems with a large number of parameters, and on the exoplanet toolkit (e.g., Theano Development Team 2016), which extends PyMC3’s language to support many of the custom functions and distributions required when fitting our datasets (Salvatier et al. 2016). Exoplanet is also designed to provide the building blocks for fitting many datasets using the Markov Chain Monte Carlo (MCMC) simulation approach. We set to 100,000 the number of MCMC simulations to derive the parameter distribution for the orbit of the binary; we allowed MCMC to explore values of orbital parameters in all possible ranges (e.g., eccentricity from 0.0 to almost 1.0), except for the orbital period and the amplitude of the RV curves, where we explored values in the intervals 800–5000 a and $0.4\text{--}5.0\text{ km s}^{-1}$. The posterior distributions of the fitted parameters are graphically shown in the corner plot of Fig. A.1, where it becomes apparent that the distribution of the solutions is narrow with clear peaks for most of the orbital parameters. The eccentricity distribution shows a peak at 0.0 with a long tail towards slightly larger values, yet our solution strongly indicates that the orbit is nearly circular with an eccentricity likely smaller than $e = 0.1$. More degenerated is the determination of the orbital inclination angle. This is not a surprising result because the available astrometric data cover less than 20% of the actual orbit of the stellar binary, which obviously is insufficient to find a tight solution.

Here, we adopted the orbital parameters mean values of the MCMC posterior distributions as the “best” Keplerian solution describing the observed astrometry and absolute RVs altogether. All adopted orbital parameters are given in Table 3. The mean values are very close to the peaks of the distributions shown in the corner plot of Fig. A.1. We found that the binary has a likely orbital period of 1295 ± 184 a, which is significantly larger than that of Chang (1972) but closer to the determination by Hopmann (1954), and a semi-major axis of 130.9 ± 5.1 au (or $20''.66 \pm 0''.80$). The error bars were determined by differentiating the various mean solutions obtained when removing data points from the long astrometric dataset. Figure 3 shows the observed astrometric positions of GJ 338 B around GJ 338 A between 1821 and 2019, the adopted Keplerian ellipse, and the residuals after removing the adopted orbital solution from the data. The residuals have a dispersion of $\pm 0''.11$ and $\pm 0''.14$ in right ascension (α) and declination (δ), respectively, which are close to the uncertainties of the best quality astrometry in our dataset. Furthermore, the residuals are flat over time, which indicates that the solution is acceptable for the covered time interval. With this solution, a total mass of $1.33 \pm 0.17 M_\odot$ is obtained for the stellar pair, which is consistent at the 1σ level with the most likely value obtained by Schweitzer et al. (2019). The individual mass of each binary component was then determined from the total mass and amplitude of the RV curves (mass ratio, $q = K_A/K_B = M_B/M_A = 0.93 \pm 0.13$). We found dynamical masses of $0.69 \pm 0.07 M_\odot$ for GJ 338 A and $0.64 \pm 0.07 M_\odot$ for GJ 338 B.

Figure 4 shows the ELODIE and HIRES absolute RVs and the best Keplerian fit. The top panel illustrates 1.5 full orbital periods; the absolute RV data considered in this work cover a tiny fraction ($<10\%$) of the actual orbital period. The second and third panels of Fig. 4 show an enlargement of the previous panel centered on the observed velocities of GJ 338 A and B. Both the adopted Keplerian solution and a straight line fit to the RV data are included in the two panels. Given the small time coverage of the data, there is no perceptible difference between the Keplerian fit and the straight line (at the level of less than 1 m s^{-1} , that is, about one order of magnitude smaller than the RV uncertainties).

The fourth panel of Fig. 4 illustrates the velocity residuals (observations minus the fit), which have a dispersion of 11.13 m s^{-1} (GJ 338 A) and 11.30 m s^{-1} (GJ 338 B).

As a consistency check of the quality of our adopted Keplerian fit, we produced the bottom panel of Fig. 4 that displays the differential velocity ($v_A - v_B$) of the binary members as a function of time using ELODIE, HIRES, SOPHIE, and CARMENES data, thus expanding the time coverage of the RV dataset by an additional 5.7 a for a total of 23.7 a. We assumed that the RV measurements of GJ 338 A and B are coeval if they were acquired on the same night or within a week time. With this condition, we identified 29 HIRES, 5 ELODIE, 39 SOPHIE, and 23 CARMENES RV pairs for GJ 338 A and B. This plot is in principle independent (or at least it minimizes the systematics) of the absolute RVs provided that all RVs are obtained using the same mask. For this purpose, we re-reduced all publicly available ELODIE, SOPHIE and CARMENES spectra using the SERVAT software (Zechmeister et al. 2018) and the same template of GJ 338 B for the two stars, thus removing the velocity zero point problem mentioned in Section 3.1. The adopted Keplerian solution nicely reproduces the differential velocity curve of the ELODIE and HIRES RVs and also extrapolates equally well towards the more recent SOPHIE and CARMENES data.

Another consistency check of the validity of the adopted Keplerian solution is given by the determination of the binary mass ratio through the measurement of the slopes of the combined ELODIE and HIRES RV curves of GJ 338 A and B. The mass ratio obtained in this way is $q = 0.91 \pm 0.11$, which is compatible at the 1σ level with the value derived from the ratio of the RV amplitudes (K_A, K_B) of the adopted Keplerian fit.

Another method to derive the mass ratio of the binary is based on the equation $q = (v_A - \gamma)/(\gamma - v_B)$ (Wilson 1941), where γ stands for the systemic radial velocity. Because the mass ratio has to be a constant, we forced a flat (zero slope) q versus time as shown in Figure 5. We employed the ELODIE and HIRES absolute RVs previously used in the Keplerian solution determination and enlarged the data coverage adding the CARMENES relative RVs calibrated to absolute value using the adopted Keplerian solution. The derived values are as follows: $q = 0.930 \pm 0.021$, $\gamma = 11.797 \pm 0.001 \text{ km s}^{-1}$. The small uncertainties associated with q and γ reflect the precision of the method, which leads to the conclusion that GJ 338 A is likely the most massive component of the stellar binary, as expected from its higher luminosity and bluer optical colors. This result is opposed to that of Chang (1972). By using the astrometric total mass of the stellar system, we obtained individual masses of $0.69 \pm 0.03 M_\odot$ (GJ 338 A) and $0.64 \pm 0.03 M_\odot$ (GJ 338 B), which are basically identical to the results from the first approach (combined astrometric and spectroscopic fit) and are consistent with the mass determinations by Schweitzer et al. (2019) at the 1σ level, but slightly larger by 8.3–9.4 % than those reported by these authors.

From now on, we use the stellar mass determinations from the combined astrometric and spectroscopic fit. Table 3 summarizes the derived orbital parameters, including the individual mass determinations and the mass ratio determinations from the spectroscopic data only.

4.2. RV detrending

Before investigating the presence of planets around each member of the GJ 338 stellar system, the Keplerian orbital solution (Table 3) was subtracted from the observed HIRES and CARMENES optical RVs. As a consistency check, we determined the negative (and positive) slope independently for the




Fig. 5: Determination of the stellar mass ratio, $q = M_B/M_A$, using ELODIE, HIRES, and CARMENES VIS differential radial velocities (see Section 4.1). The horizontal, solid line corresponds to $q = 0.929 \pm 0.019$, where the uncertainty gray area accounts for the observed dispersion of the q values.

HIRES and CARMENES datasets, finding very similar results at the 1σ level. We concluded that given the short time coverage of the RV data and the long orbital period of the binary, the subtraction of a straight line instead of the Keplerian curve would not change the results reported next.

Our second step was to remove RV outliers from the GJ 338 B datasets by applying a 2σ -clipping algorithm, resulting in 1 HIRES RV and 6 CARMENES VIS RVs rejected from the lists of 32 and 159 independent measurements, respectively. Outliers typically have large error bars (due to, e.g., short exposures) and are marked in Fig. 1 and all other RV figures of this paper. The “flattened” and 2σ -clipping corrected CARMENES VIS RV dataset of GJ 338 B has a standard deviation of 4.2 m s^{-1} , which is slightly smaller than the dispersion of the HIRES velocities (5.4 m s^{-1}). The mean error bar value of the individual CARMENES VIS measurements provided by SERVAT is 2.4 m s^{-1} , that is, almost a factor of two lower than the standard deviation. The origin of this RV excess variations is explored in the following section.

We also removed the orbital trend from the CARMENES NIR data of GJ 338 B and applied the 2σ -clipping for eliminating RV outliers after which we find a standard deviation of 9.2 m s^{-1} . This value coincides with the typical dispersion of CARMENES NIR RVs of other M-type stars. Because of the larger *rms* of the NIR data, this channel does not deliver more convincing results than the VIS channel. Consequently, the NIR RVs were not used except when necessary.

The same procedure was applied to the CARMENES dataset of GJ 338 A resulting in the 2σ -clipping removal of 4 VIS RVs out of a total of 70 measurements. The detrended RV curve of GJ 338 A has a standard deviation of 5.5 m s^{-1} .

Besides RV precision, the cadence of the observations is also a critical ingredient for performing a proper analysis of temporal variability. In this regard, the CARMENES data were taken every few to several days while the target was visible from the Calar Alto Observatory. This observing frequency contrasts with the cadence of the HIRES observations (3–4 RV measurements per year). As a result, the CARMENES scheduled observations

Table 3: Orbital parameters for the GJ 338 stellar binary system.

Combined astrometric and spectroscopic solution		
Parameter	Description (Unit)	Adopted solution
α	Semi-major axis (arcsec)	20.66 ± 0.80
a	Semi-major axis (au)	130.9 ± 5.1
T_0	Time of periastron passage (a)	1513 ± 61
e	Eccentricity	$0.01^{+0.15}_{-0.01}$
i	Inclination (deg)	37 ± 12
Ω	Argument of ascending node (deg)	32.4 ± 1.8
ω	Argument of periastron (deg)	109.1 ± 6.0
P	Period (a)	1295 ± 180
K_A	Amplitude (km/s)	0.93 ± 0.12
K_B	Amplitude (km/s)	0.99 ± 0.12
M	Total mass (M_\odot)	1.33 ± 0.17
q	Mass ratio, K_A/K_B	0.93 ± 0.10
M_A	Mass of the primary (M_\odot)	0.69 ± 0.07
M_B	Mass of the secondary (M_\odot)	0.64 ± 0.07
γ	Systemic velocity (km s^{-1})	11.798 ± 0.001
Mass ratio ^(a) from equation $q = (v_A - \gamma)/(\gamma - v_B)$		
q	Mass ratio, M_B/M_A	0.930 ± 0.021
γ	Systemic velocity (km s^{-1})	11.797 ± 0.001

Notes. ^(a) Based on the HIRES absolute RV calibration.

are ideal for exploring planets in short- to intermediate-period orbits (García-Piquer et al. 2017), while the HIRES data are useful for identifying planets at longer orbital periods. Mixing the two datasets might not produce better results given the different data quality and temporal cadence. In what follows, we analyze the CARMENES VIS data independently and employ the HIRES and CARMENES NIR RVs to check the results.

4.3. Stellar activity

The profile of the stellar lines can be changed by photospheric and chromospheric activity, which has an impact on accurate RV measurements. M-type dwarfs typically have intense magnetic fields (Reiners et al. 2012, 2014; Newton et al. 2016, 2017; Shulyak et al. 2019), and it is expected that the related magnetic activity leaves a detectable trace in the high resolution spectra. Various groups have focused on the identification of spectral indicators of stellar activity (Montes et al. 2000; Gomes da Silva et al. 2011, 2012; Tal-Or et al. 2018) with the main goal of disentangling activity-induced RV variations and planetary signals. Two commonly-used chromospheric activity indicators are based on measurements of the $H\alpha$ $\lambda 6562.82 \text{ \AA}$ and Ca II H and $\text{K } \lambda \lambda 3933.7, 3968.5 \text{ \AA}$ lines. The blue Ca II H and K lines are not covered by the CARMENES data; instead, the Ca II infrared triplet $\lambda \lambda 8498, 8542, 8662 \text{ \AA}$ (IRT) is fully covered by the VIS channel. This triplet is also well known for its sensitivity to stellar activity as reported by Martínez-Arnáiz et al. (2011).

The CARMENES SERVAL pipeline provides measurements for a number of spectral features that can reveal stellar activity, including $H\alpha$, Ca II IRT, the differential line width (dLW), and the chromatic index (CRX), as defined by Zechmeister et al. (2018). The latter measures the $\text{RV} - \log \lambda$ correlation, and it is used as an indicator of the presence of stellar active regions. In addition, the pipeline computes the cross-correlation function

(CCF) for each CARMENES spectrum using a weighted binary mask that was built from coadded observations of the star itself. As explained by Reiners et al. (2018a), the CCFs are fitted with a Gaussian function to determine the contrast, the full width at half maximum (FWHM), and the bisector span (BIS), all of which are useful for studying stellar activity-related properties.

GJ 338 B. All CARMENES spectroscopic activity indicators of GJ 338 B are given in Table B.1. Their associated Generalized Lomb-Scargle periodograms (GLS; Zechmeister & Kürster 2009) are depicted in various panels of Fig. 6, where we also include the false alarm probability (FAP) levels of 10, 1, and 0.1 % from 10,000 bootstrap randomizations of the input data. Peaks of the GLS periodograms with FAP level $< 0.1\%$ are considered significant. The GLS method of analysis is particularly indicated for unevenly time series. With the exception of the CCF contrast, FWHM, and CRX indices, there is one single significant peak in all other activity indicators centered at $\sim 0.060 \text{ d}^{-1}$ ($\sim 16.6 \text{ d}$). The Ca II IRT and $H\alpha$ indices folded in phase with this period are shown in Fig. 7: all of them have a sinusoid-like variation (dLW and BIS indices also show similar patterns).

We used the GLS peak at 16.6 d as the initial period for finding the best sinusoidal curve that minimizes the residuals of the activity indices presented in Fig. 7. We found that the best period is $16.61 \pm 0.04 \text{ d}$, after averaging all individual determinations. We also checked that the variations of the activity indices do not correlate with the RVs. The shape of these variations and the fact that CRX and FWHM indices do not show the 16.6-d peak may be related to the origin and nature of the stellar variability that is inducing this signal in the CARMENES VIS RVs (see below and Schöfer et al. 2019). $H\alpha$ and Ca II IRT are mainly chromospheric indicators while CRX is related to the photosphere. We ascribed the 16.6-d period to the chromospheric rotation cycle




Fig. 6: GLS periodograms for GJ 338 B activity and RV data. In all panels, the horizontal dashed lines indicate FAP levels of 10% (red), 1% (orange), and 0.1% (green). Panel (a): CARMENES RVs from the VIS channel. The three highest peaks at 8.3, 16.6, and 24.4 d are labeled and their positions are marked in all other panels with vertical dashed lines. Panel (b): V-band photometry. Panels (c–e): chromospheric line indices of the infrared Ca II triplet, H α , and dLW. Panels (f–h): FWHM, contrast, and bisector velocity span from the CCF analysis. Panel (i): chromatic RV index (CRX).

of GJ 338 B. This period determination differs from the values tabulated by Wright et al. (2011), who measured a photometric period of 10.17 d. However, later Fuhrmeister et al. (2019) reported a tentative period of 16.6 ± 0.5 d and 17.4 ± 1.0 d using CARMENES H α and Ca II IRT activity indices, respectively. These values coincide with the 16.61 ± 0.04 d determined here and we agree with these authors that CARMENES Ca II IRT and H α activity indices are well suited for stellar period searches.

As a side note, Abt & Levy (1973) measured the Ca II emission-line velocities of GJ 338 B and A and found them variable. They discussed that these variations were caused by the intrinsic multiple nature of each star, and determined a “binary” period of ~ 16.47 d for GJ 338 B (the one with the highest significance and amplitude of ~ 5 km s $^{-1}$ according to the authors). Later on, Morbey & Griffin (1987) obtained more data that did not confirm the proposed periods and did not suggest any variation in velocity at all, thus rejecting the notion that GJ 338 B is a binary star itself. The CARMENES RVs support this latter result. However, it is possible that Abt & Levy (1973) detected the rotation cycle of GJ 338 B because they were using emission lines very sensitive to stellar activity.

GJ 338 A. We analyzed the CARMENES VIS RVs of GJ 338 A in a similar manner as for the B component and found that the GLS periodograms of the H α , dLW, and Ca II IRT activity indices (see Fig. 8) show a significant peak at $16.3^{+3.5}_{-1.3}$ d (above FAP level $< 0.1\%$). All RV and activity indices measurements are given in Table B.2. We verified that there is no confusion in the identification of the two stellar sources in the CARMENES catalog. The first evidence is that GJ 338 A RV data show a trend that is opposite to that of GJ 338 B, thus indicating that we are dealing with the two stellar components of the binary. The second piece of evidence is that the absolute RVs differ between the two stars. We attributed the periodicity of 16.3 d to the rotational cycle of GJ 338 A. Indeed, this rotation period is surprisingly close to that of GJ 338 B. It might hint that both M0-type stars, which were likely born from the same molecular cloud and are consequently coeval, have followed a very similar angular momentum evolution.

Photometry. For the present analysis, we studied the light curves separately per filter and per observing campaign and all seasons altogether (V-band data only, which is the only filter in common to the three campaigns). The differential light curve was obtained by dividing the fluxes of GJ 338 B and GJ 338 A because there is no other bright star in the field that can act as a reference source (see Section 3.3). The GLS periodogram of the entire V-band light curve is shown in the second panel (from the top) of Fig. 6, where a peak at around 18 d is detectable. The highest peaks in the GLS periodograms of the B- and i-band light curves lie between 16 and 20 d, but they are not significant. Also apparent in these periodograms is the peak at 8–10 d, which is likely due to the first harmonic of the primary signal. With the exception of the V-band data, we concluded that no obvious, significant peak can be extracted from the periodograms of the LCO and SNO light curves. From the analysis of the separated seasons of the V-filter data, we found a characteristic frequency in the interval 15.8–26.9 d, which covers the likely rotation periods of the two stars. The lack of an intense peak can be explained by photometric stability (or stellar variability that is below our detectability limit), by anti-correlated stellar variability of similar amplitude (thus producing a net amplitude close to zero in the differential light curve), or by the combination of two stellar




Fig. 7: GJ 338 B. Phase-folded time series of activity indicators Ca II $\lambda 8498$ Å (left) and H α $\lambda 6563$ Å (right) and residuals. The solid curve depicts the best sinusoidal fit that modulates the rotation period at 16.6 d.




Fig. 8: GLS periodograms for GJ 338 A. In all panels, the horizontal dashed lines indicate FAP levels of 10% (red), 1% (orange), and 0.1% (green). *Top panel:* CARMENES VIS RVs. *Second through fourth panels:* CARMENES Ca II $\lambda 8498$ Å, H α , and dLW spectroscopic activity indicators. All activity indicators and the RV data show a significant, broad peak at around $16.3^{+3.5}_{-1.3}$ d. This is likely associated with the rotation period of GJ 338 A.

variabilities of similar low amplitude and close, but not equal, periodicities, which could attenuate the true individual periodicities. The first hypothesis is the least likely one because there is stellar activity from the CARMENES spectroscopic data. With the current photometry, we could not discern the true scenario

and therefore, we were not able to derive individual photometric periodicities for GJ 338 B and GJ 338 A.

4.4. Frequency content of the radial velocities of GJ 338 B

The GLS periodogram of the CARMENES VIS RVs of GJ 338 B is illustrated in the top panels of Figs. 6 and 9. The highest peak is located at ~ 24.4 d followed by two other significant peaks at ~ 8.3 and ~ 16.6 d. The three signals lie above FAP level 0.1 % and are marked in all panels of Figs. 6 and 9. The GLS periodogram of the CARMENES NIR RVs of GJ 338 B is shown in the second panel of Fig. 9. Only the peak at 16.6 d exceeds FAP ~ 10 % (significance is not as high as in the optical probably because there are less NIR RVs and they have larger associated uncertainties). This signal at 16.6 d is common to all RV and many activity index GLS periodograms of GJ 338 B, and as discussed in the previous Section, it is very likely related to the stellar rotation. Even though it is believed that the activity-induced signal is the strongest at blue wavelengths and is attenuated at long wavelengths in the RV time series (Martín et al. 2006; Huélamo et al. 2008), GJ 338 B clearly shows this peak in the near-infrared despite the larger error bars of the CARMENES NIR channel. This implies that the mechanism responsible for the activity in GJ 338 B is capable of imprinting its signal at both optical and near-infrared wavelengths. Figure 10 illustrates the CARMENES NIR RVs folded in phase with the rotation period of ~ 16.6 d; the curve shows a amplitude of 4.3 ± 1.7 m s $^{-1}$.

For completeness, we also provide the GLS periodogram of the HIRES data of GJ 338 B in the third panel of Fig. 9. No significant signal is seen at any of the three peaks provided by the CARMENES VIS RV dataset. This is quite likely due to the insufficient cadence and long time coverage of the HIRES data. Interestingly, the combination of the HIRES and CARMENES VIS RVs recovers the 16.6 d signal well above FAP level 0.1 % (fourth panel of Fig. 9). The second most prominent signal is located at ~ 24.4 d, which agrees with the results of the CARMENES VIS data.

In what follows, we demonstrate that the three signals (~ 8.3 , ~ 16.6 , and ~ 24.4 d) of the CARMENES VIS RVs of GJ 338 B are not related to each other by an aliasing effect, which is typically caused by the gaps in the time coverage of the observations (e.g., Dawson & Fabrycky 2010). To identify the presence of possible aliasing phenomena, the spectral window




Fig. 9: GLS periodograms of the RV data of GJ 338 B. *Panel (a)*: periodogram of the CARMENES VIS RVs is plotted as a blue, solid line and the window function of the RV data is shown with an orange line. The three strongest signals are marked in this panel and in all other panels. The horizontal dashed lines indicate FAP levels of 10 % (red), 1 % (orange), and 0.1 % (green). *Panel (b)*: periodogram of the CARMENES NIR RVs. *Panel (c)*: periodogram of the HIRES RV measurements. *Panel (d)*: periodogram of the combined HIRES + CARMENES VIS RV data. *Panel (e)*: periodogram of the CARMENES VIS RV residuals after removing the best-fit that modulates the stellar activity at $P_{\text{rot}} = 16.6$ d. *Panel (f)*: periodogram of the CARMENES VIS RV residuals after removing the stellar rotation and its first harmonic at 8.3 d. *Panel (g)*: periodogram of the CARMENES VIS RV residuals after removing all three significant signals.

has to be considered. If peaks are seen in the window function, their corresponding aliases will be present in the RV periodograms as $f_{\text{alias}} = f_{\text{true}} \pm m f_{\text{window}}$, where m is an integer, f_{true} is the frequency identified in the RV periodogram and f_{window} the frequency from the window function (Deeming 1975). Typ-




Fig. 10: Phase-folded CARMENES NIR RVs (blue dots) to 16.54 ± 0.06 d. *Top panel*: the black line depicts the best-fit that modulate the activity. Open circles stand for those points that deviate more than 2σ . The amplitude is 4.3 ± 1.7 m/s. *Bottom panel*: residuals' rms is 7.5 m/s.




Fig. 11: Enlargement of the GLS periodogram of the CARMENES VIS RV data of GJ 338 B (blue line) around the three strongest signals (marked with color-coded dots). The window function is plotted as the orange line. Horizontal dashed lines indicate the different FAPs: 0.1 % (green), 1 % (orange) and 10 % (red). The 1-sidereal-year aliases around each of the strongest signals are indicated with vertical solid lines.

ical aliases are those associated with the sidereal year, synodic month, sidereal day, and solar day. The window function of the CARMENES VIS data is depicted in the top panel of Fig. 9 together with the GLS periodogram of the RVs. There are two significant peaks in the window function at the sidereal year and day. Using this information, we plotted the predicted aliases corresponding to each of the three significant peaks of the CARMENES VIS RVs in Fig. 11. From this Figure, none of the three signals are related to each other by an alias of the observing window.

To summarize, based on the CARMENES VIS and NIR RV data we identified one strong signal at 16.61 ± 0.04 d, which is




Fig. 12: Stacked BGLS periodogram for the CARMENES RVs data zoomed around the likely planetary orbital period of 24.4 days. The cumulative number of CARMENES observations is plotted as a function of periodicity; the color scale indicates the logarithm of the probability, where the red color stands for the most likely period.

also significant in some activity indicators, and thus we relate it to the rotation period of GJ 338 B. The signals at ~ 8.3 and ~ 24.4 d do not have a counterpart in the activity indicators and are not aliases of the rotation signal. The ~ 8.3 -d peak is exactly at half the rotation period, and we attribute it to the first harmonic of the stellar rotation. The ~ 24.4 -d signal is not an exact multiple of the star's rotation, and we attribute it to a planetary origin. We investigate its nature in Sect. 5.

To explore the stability of the ~ 24.4 -d period, we produced the stacked periodogram of the CARMENES data shown in Fig. 12. For this purpose, we computed the Bayesian Generalized Lomb-Scargle periodogram (BGLS, Mortier et al. 2015). This formalism determines the probability between peaks of the periodogram. The signal at 24.4 d is very well detected with a probability above 10^{30} (with the minimum probability set to 1). This signal becomes more significant by adding observations as expected for a real signal.

4.5. Frequency content of the radial velocities of GJ 338 A

The GLS periodogram of GJ 338 A CARMENES VIS RVs is illustrated in the top panel of Fig. 8. The highest peak is located at $16.3^{+3.5}_{-1.3}$ d and shows a broad structure. No other signal lies above FAP 0.1%. As discussed in Section 4.3, this periodicity is also present in the GLS periodograms of the CARMENES activity indicators (Ca II IRT, H α , and dLW) of GJ 338 A and it is very likely related to the stellar rotation.

This activity signal can be modeled by a periodic curve with an amplitude of 5.6 ± 0.7 m s $^{-1}$ (VIS RV data). After its removal from the CARMENES VIS data, we found an *rms* of the residuals of 4.6 m s $^{-1}$ and no additional significant peak in the corresponding periodogram above a FAP of 1 %. We calculated the minimum mass of any putative planet around GJ 338 A in an orbital period range between 7 and 50 d using the third Kepler's law, the mass of the parent star given in Table 1, and different values of eccentricity. The minimum period is given by four times the typical Nyquist frequency of the RV time series, and the maximum period is arbitrarily chosen. Figure 14 depicts the planetary minimum masses where we adopted, at the 1σ level, the *rms* of the CARMENES VIS RVs multiplied by a factor of 0.92 as the minimum detectable RV amplitude. Bartlett et al. (2009) argued that a signal must be about 92 % of the magnitude of the




Fig. 13: CARMENES VIS RV data of GJ 338 B folded in phase with different periods. Each panel has been corrected for the others signals. *Top panel*: RV data folded in phase with the stellar rotation period of 16.6 d. Overplotted it is the best sinusoid function reproducing the data with an amplitude of 2.7 ± 0.5 m s $^{-1}$. *Second panel*: RV folded in phase with the 8.3 d signal. Another sinusoid function is fit to these data with a RV amplitude of 1.8 ± 0.6 m s $^{-1}$. *Third panel*: RV data folded in phase with the planetary orbital period of 24.4 d. The best Keplerian solution for the planet, with an RV amplitude of 2.8 ± 0.4 m s $^{-1}$, is shown with the black line. *Bottom panel*: final RV residuals after removing the three signals (16.6, 8.3, and 24.4 d) have an *rms* = 3.0 m s $^{-1}$. Open circles represent the RVs that deviate more than 2σ from the trend of the original data and that were not included in the analysis but are shown in this figure.

average residuals to be measurable through Lomb-Scargle-based periodograms. We are able to discard the presence of planets




Fig. 14: Minimum detectable mass of any putative planet surrounding GJ 338 A as a function of the orbital period. We adopted the *rms* of the CARMENES VIS RV residuals as the amplitude of the Keplerian signals. The color code stands for the different values of the eccentricity.

more massive than $10\text{--}19 M_{\oplus}$ at $7\text{--}50$ d orbits with eccentricity ~ 0.1 , respectively. For more eccentric orbits, the planetary minimum masses decreases down to $8\text{--}14 M_{\oplus}$ at $7\text{--}50$ d orbits.

Following the same approach as for GJ 338 B (see Section 5), we also analyzed the GLS periodograms of the HIRES data and the combined HIRES+CARMENES VIS RVs of GJ 338 A, finding no characteristic frequency in the former and recovering the 16.3-d peak in the latter. The broad structure of the 16.3-d peak could be explained by the existence of differential rotation and/or blending of a real (very likely activity) signal at a similar period with an aliasing effect (due to one and two year aliases, time gaps between data, etc.). When removing the signal at 16.3 d from the HIRES+CARMENES dataset, another signal at 17 d became apparent above FAP 0.1% in the GLS periodogram of the residuals. Both 16.3 d and 17 d periodicities fall within the characteristic frequency interval of $15.8\text{--}26.9$ d obtained from the analysis of the photometric V-band data. No more significant signals are present when removing the 16.3 d and 17 d periodic variations. Using current data, we cannot discard the presence of super-Earth or less massive planets “hidden” in the broad 16.3 d peak of the RV periodogram.

5. Planet orbiting GJ 338 B

5.1. Pre-whitening method

We obtained the best-fit orbital parameters of the planet at ~ 24.4 d using the RVLIN code³ (Wright & Howard 2009). This code is based on the Levenberg-Marquardt (Levenberg 1944; Marquardt 1963) algorithm and contains a set of routines that fits an arbitrary number of Keplerian curves to RV data. Our approach was as follows: first, we removed the “stellar contribution”, that is, activity, with a rotational period of 16.6 d from the observations by fitting a sinusoid function to the CARMENES VIS data. The fit, shown in the upper panel of Fig. 13, yielded an RV amplitude of 2.7 m s^{-1} (smaller than that obtained from the analysis of the NIR RVs, see Section 4.4). The residuals had

Table 4: Comparison of different solutions for GJ 338 B.

Model	Prior	GP kernel	$\ln \mathcal{L}$	BIC ⁽¹⁾
BM ⁽²⁾	$\mathcal{U}_{\text{offset}}(-100, 100)$ $\mathcal{U}_{RV\text{ jitter}}(-10, 2)$ $\mathcal{U}_{\text{trend}}(-1, 1)$...	-569	1153
BM+GP	$\mathcal{U}_{\omega_0}(-10, 5)$ $\mathcal{U}_Q(-10, 5)$ $\mathcal{U}_{S_0}(-10, 5)$	SHO	-547	1124
BM+1pl	$\mathcal{U}_{\text{pl}}(24.39, 24.5)$...	-546	1132
BM+GP+1pl	$\mathcal{U}_{\text{pl}}(24.39, 24.5)$	SHO	-517	1089
BM+2GP+1pl	$\mathcal{U}_{\text{pl}}(24.39, 24.5)$	SHO	-513	1096
BM+GP+2pl	$\mathcal{U}_{\text{pl}_1}(24.39, 24.5)$ $\mathcal{U}_{\text{pl}_2}(8.3, 8.4)$	SHO	-497	1075

Notes. ⁽¹⁾ BIC corresponds to the Bayesian Information Criterion. ⁽²⁾ BM stands for the base model containing offsets, RV jitter, and a linear trend.

an *rms* of 3.9 m s^{-1} , which was slightly smaller than the initial standard deviation of the CARMENES VIS data, but still larger than the typical uncertainty associated with the individual RV measurements.

A new GLS periodogram of the residuals was obtained (it is depicted in panel (e) of Fig. 9). The planetary signal at ~ 24.4 d, which was also obvious in the original GLS periodogram, is now enhanced. The ~ 8.3 d peak still remains, although its significance is at the FAP level 10 %. Because it is likely related to stellar activity, we removed its contribution from the VIS RV data by fitting a second sinusoid with a period of 8.26 d and an RV amplitude of 1.8 m s^{-1} (middle upper panel of Fig. 13). The new residuals have an *rms* of 3.6 m s^{-1} . With these subsequent sinusoidal subtractions from the CARMENES VIS RV data, the only significant signal of the residuals has a characteristic period of 24.4 d (panel (f) of Fig. 9).

With the RVLIN code, we found the best Keplerian solution, which is illustrated in the third panel of Fig. 13, where the RV data are folded in phase with the planetary orbital period. The CARMENES VIS RV measurements that were not included in the analysis (open symbols in the RV plots) follow the trend and lie at the expected phases. The corresponding best-fit orbital parameters and their error bars are reported in Table 5. The best-fit orbital period is $P = 24.40 \pm 0.04$ d, and the planet has a minimum mass of $9.2 \pm 1.1 M_{\oplus}$ (computed by adopting the mass of $0.64 \pm 0.07 M_{\odot}$ for the parent star, as determined here). This planetary minimum mass would be $8.9 \pm 1.2 M_{\oplus}$ if the stellar mass determined by Schweitzer et al. (2019) were adopted instead. The planet candidate GJ 338 Bb is located at an orbital separation of 0.142 ± 0.014 au from its parent M0-type star, and despite its proximity, the planetary orbit appears to be slightly eccentric ($e = 0.25 \pm 0.13$), although given the error bar of the eccentricity, the orbit is also compatible with an almost circular orbit. The amplitude of the planet RV curve is $2.8 \pm 0.4 \text{ m s}^{-1}$, about 38 % smaller than the dispersion of the residuals data and similar to the mean error bar of individual CARMENES VIS RV data. The RV amplitudes of the stellar activity and the planet-induced signal are quite similar.

The RV residuals obtained after removing the stellar and planetary contribution from the CARMENES VIS RV data are shown in the bottom panel of Fig. 13. They have an *rms* of 3.0 m s^{-1} , still larger than the mean error bar of individual RV measurements, but similar to the typical *rms* average of CARMENES data. The GLS periodogram of these residuals is

³ The RVLIN code is available at <http://exoplanets.org/code/>

Table 5: Keplerian orbital parameters of GJ 338 Bb from pre-whitening and Gaussian process regression methods .

Parameter	Pre-whitening	GP regression	
	GJ 338B b	Prior	GJ 338B b
P (d)	24.40 ± 0.04	$\mathcal{U}(24.39, 24.50)$	24.45 ± 0.02
T_0 (BJD-2,400,000) ⁽¹⁾	57831.2 ± 2.0	$\mathcal{U}(0, 30)$	$57517.06^{+6.15}_{-4.72}$
e	0.25 ± 0.13	$\mathcal{U}(0, 0.8)$	$0.11^{+0.11}_{-0.08}$
ω (deg)	144.9 ± 31.8	$\mathcal{U}(-2\pi, 2\pi)$	$204.3^{+94.4}_{-70.0}$
K (m/s)	2.8 ± 0.4	$\mathcal{U}(0, 10)$	3.07 ± 0.37
v_0 (m/s) ⁽²⁾	0.09 ± 0.23	$\mathcal{U}(-100, 100)$	3.16 ± 0.32
$m_p \sin i$ (M_\oplus) ⁽³⁾	9.15 ± 1.11		$10.27^{+1.47}_{-1.38}$
$m_p \sin i$ (M_\oplus) ⁽⁴⁾	8.90 ± 1.20		$9.97^{+1.47}_{-1.38}$
a (au)	0.142 ± 0.014		0.141 ± 0.005
T_{eq} (K) ⁽⁵⁾	$300\text{--}390 \pm 30$		$301\text{--}391 \pm 20$

Notes. ⁽¹⁾ T_0 corresponds to the periastron passage. The reference time for the Gaussian process (GP) regression is 2457500.0. ⁽²⁾ Arbitrary zero point applied to CARMENES VIS RVs. ⁽³⁾ Derived by adopting the mass of $0.64 \pm 0.07 M_\odot$ for the parent star. ⁽⁴⁾ Derived with stellar mass determined by Schweitzer et al. (2019). ⁽⁵⁾ For Bond albedo in the interval 0.65–0.0.

displayed in panel (g) of Fig. 9; it shows no other significant peak. The results are not modified by changing the order of the pre-whitening method, thus giving robustness to the discovery of the small-mass planet around GJ 338 B. Table 5 summarises the orbital parameters.

For completeness, we combined the CARMENES VIS and HIRES RV datasets of GJ 338 B. The fourth panel of Fig. 9 illustrates the corresponding GLS periodogram. We performed the same pre-whitening analysis as previously described. The amplitude of the stellar activity of the combined RV curve was determined at 2.7 m s^{-1} , which fully agrees with that from the CARMENES-data only. The first residuals have an *rms* of 4.1 m s^{-1} , which is indeed slightly higher than the first residuals of the CARMENES VIS RV data only. However, from this point of the analysis to the end of the process, the noise added by the HIRES data and the poor temporal coverage of this dataset prevented us from finding a consistent orbital solution ($P = 24.40$ d) despite the fact that the periodograms of the combined HIRES+CARMENES VIS RV residuals (not shown here) do not differ significantly from Fig. 9. The eccentricity value that we found for the planet orbital solution with the combined CARMENES VIS and HIRES RV data sets is $e = 0.1$. The lessons learnt from this exercise is that the HIRES RVs do not contradict the findings of the CARMENES data and that the eccentricity of the orbit of the planet around GJ 338 B is not well constrained with the current data.

5.2. Gaussian process regression method

We also followed a different approach to study in detail the CARMENES VIS RV curve of GJ 338 B. This approach has the advantage of simultaneously fitting the stellar activity and the planetary signal(s). The match between the dominant period in dLW and H α indices (Section 4.3) with one of the strongest signals in the RVs (Section 4.4) motivated a simultaneous fit of these activity indicators together with the RVs in order to account for the correlated RV noise. We used the *celerite* package (Foreman-Mackey et al. 2017) and its damped simple harmonic oscillator (SHO) kernel of the form

$$S(\omega) = \sqrt{2} \frac{S_0 \omega_0^4}{\pi (\omega^2 - \omega_0^2)^2 + \omega_0^2 \omega^2 / Q^2} \quad (1)$$

The parameters of the model are the oscillator eigen-frequency (ω_0), the quality factor (Q , corresponding to the inverse of the damping time scale), and the amplitude of the damped oscillator (S_0). The former two are constrained by the activity indicator and an SHO Gaussian process (GP) kernel with these parameters, but with its own amplitude, is added to the planetary model. In addition to this, we used a model with individual offsets and RV jitter, as well as a linear trend, as the base line model (BM). The RV data and the activity indicators were then solved at the same time.

We used the Bayesian Information Criterion (BIC, Liddle 2007) to identify the optimal solution. When fitting models, it is possible to increase the log-likelihood ($\ln \mathcal{L}$) by adding parameters, but the model may result in overfitting. The BIC resolves this problem by introducing a penalty term of the number of parameters in the model. It can be mathematically defined as follows:

$$\text{BIC} = k \ln(n) - 2 \ln(\mathcal{L}), \quad (2)$$

where n is the number of data points, k the number of free parameters to be estimated, and \mathcal{L} the maximized value of the likelihood function of the model. The lower the BIC value, the better the model is.

In a first approach, we modeled the base model together with one SHO GP kernel (which picks the 16.6-d activity simultaneously in the RVs and in H α or dLW), and the base model plus one planet with an orbital period of 24.4 d. With respect to the base model, these two models represented an improvement of the BIC value. Next, we explored how to lower the BIC parameter even further.

In the second approach, we modeled the base model, one Keplerian signal at 24.4 d, and one SHO GP kernel to account for the stellar activity with a cycle of 16.6 d. The fitted stochastic oscillator has a very high quality factor, that is, the life time is of the order of the duration of the observation. With this high quality factor, the GP kernel only acts in a very limited period/frequency range around its base frequency. The signal at 8.3 d ($=1/2 P_{\text{rot}}$) is not removed. The planetary parameters of amplitude and orbital periodicity remained unaffected by the GP model.

In a third approach, we assumed that the peaks at 8.3 d and 16.3 d are due to stellar rotation and the peak at 24.4 d is originated by a planet. We run the base model and added the planet




Fig. 15: *Left panels*: top panel illustrates the CARMENES VIS RVs (blue dots) together with a Keplerian model combined with a simple-harmonic oscillator (SHO) model shown in orange. The Keplerian-only curve is shown with a light blue color. The second panel depicts the RV residuals after subtracting the combined Keplerian + simple-harmonic oscillator model from the original data. The third and bottom panels show the GLS periodograms of the original data (light blue line) and of the RV residuals (orange) as a function of frequency and period. The model accounts for the two strongest peaks at 16.6 d and 24.4 d. *Right panels*: CARMENES H α data compared to a simple harmonic oscillator (SHO) model (see Section 5.2) and the residuals (top two panels), as well as the periodogram as a function of frequency and period (lower two panels). In the top panel, the best fit model is shown in orange, the 1σ uncertainty in blue. The periodograms show the data in light blue and the residuals in orange.

and two SHO kernels that correspond to the stellar rotation period and its first harmonic ($P_{\text{rot}}/2$). We forced the damping time of the two SHO kernels to be identical. The derived planetary parameters are compatible with the findings of previous models within the quoted error bars.

Our last approach consisted in treating the 8.3 d ($P_{\text{rot}}/2$) signal as an additional Keplerian signal. This is, the base model, two planets at 8.3 and 24.4 d, and the stellar activity modeled with a SHO GP kernel at 16.6 d. Also this time, the planetary parameters of the 24.4-d planet remain unaffected. All models, including their priors, log-likelihood and BIC values, are summarized in Table 4.

According to the BIC criterion, the best solution for the CARMENES VIS RV curve of GJ 338 B is given by the base model, a SHO GP kernel modeling the stellar activity at 16.6 d, and two planets with orbital periods of 8.3 d and 24.4 d (BM+GP+2pl in Table 4). The orbital parameters of the outer-

most planet are listed in Table 5. The 8.3-d planet would have the following derived parameters: minimum mass of $4.3 \pm 1.5 M_{\oplus}$ and semi-major axis of 0.067 ± 0.015 au. However, as mentioned in Section 4.4, the 8.3-d period is exactly half of the stellar rotation cycle; with the current data we cannot confirm the presence of this small planet around GJ 338 B. Therefore, we adopted the model given by the second approach, this is the base model, one SHO kernel for the stellar activity at the star's rotation, and the planet at 24.4 d (BM+GP+1pl in Table 4). This is the solution for the GJ 338 B planetary system that we follow in this paper.

Summarizing, the eigen-frequency and the quality factor are converted to period and damping time, the uncertainties are derived from an MCMC procedure using *emcee* (Foreman-Mackey et al. 2013) with 200 walkers and 10,000 iterations after the burn-in phase. The autocorrelation length averaged over the walkers is 550, and the acceptance rate is 22 % and during burn-in it is 35 %




Fig. 16: *Top panel*: planetary signal of GJ 338 B CARMENES VIS RVs (blue dots) folded in phase with the orbital period derived from the GP regression analysis (the base model and the stellar activity were removed). The best Keplerian solution (black line) has an RV amplitude of $3.07 \pm 0.37 \text{ m s}^{-1}$. The red dots correspond to the binned data. *Bottom panel*: RV residuals after the subtraction of the model. The weighted *rms* of the residuals is 3.2 m s^{-1} .

(as described by Foreman-Mackey et al. 2013). We also checked the chains of each parameter as a function of the step and after the burn-in phase, and the walkers were wandering around the best parameter. The MCMC posterior distributions for the fitted parameters are presented in Fig. A.2. The derived orbital parameters using the GP regression are reported in Table 5 and they are compatible within the quoted error bars with the values obtained in the previous section using the pre-whitening approach and the RVLIN code.

In the top-left panel of Fig. 15, the solution combining stellar activity (as measured from the $H\alpha$ index, see right panels of Fig. 15) and the planetary signal is depicted as a function of time (with an arbitrary zero date). The stellar activity was modeled with a damped oscillator with a period of 16.6 d and a long damping time scale of about 250 d. The weighted *rms* of the CARMENES VIS RVs after model subtraction is 3.2 m s^{-1} , and 0.0084 for the $H\alpha$ index. The CARMENES VIS RV residuals are illustrated in the second panel of Fig. 15, while the third and four panels stand for the GLS periodograms of the RV original and residual data as a function of frequency and period. The RV data and the residuals folded in phase at the best Keplerian solution obtained with the GP regression are illustrated in Fig. 16. The RV amplitude of Keplerian signal is $3.07 \pm 0.37 \text{ m s}^{-1}$.

6. Discussion

The two stars are very similar in mass, spectral type, brightness, and even rotation period, but they appear to differ in the architecture of their planetary systems. Given the rotation period of GJ 338 B ($16.61 \pm 0.04 \text{ d}$), its projected rotational velocity

($v \sin i = 2.3 \pm 1.5 \text{ km s}^{-1}$), and the star's radius ($0.58 \pm 0.03 R_{\odot}$) as taken from the literature, it is not possible to determine a precise inclination angle of the stellar rotation angle. It is likely that the projected rotational velocity is smaller than the values published in the literature. The rotation period agrees with the observed relation between X-ray luminosity and stellar activity or rotation from, for example, Wright et al. (2011).

Here, we have used two mass determinations for the M0V-type star GJ 338 B. One was obtained from the most likely astrometric solution of the stellar binary combined with the mass ratio determination from absolute radial velocity measurements. This value (Table 3) is fully independent of any evolutionary model and mass-luminosity calibrations. Despite the degeneracy of the astrometric solution (Section 4.1), the individual masses of the stellar components are well constrained. The second measurement provided by Schweitzer et al. (2019) is based on various well known calibrations that are valid for low-mass stars. The two determinations differ by less than 10 %, thus having little impact in the mass determination of the planet orbiting this star.

The top panel of Fig. 17 shows the location of GJ 338 Bb in the planetary minimum mass versus planetary orbital period diagram together with other Neptune-size and super-Earth planets known around M-type dwarfs. Our planet does not stand out in this diagram, neither it does when we select only those planets that are orbiting one of the stars of the binary systems (S-type configuration; Rabl & Dvorak 1988). The bottom panel of Fig. 17 displays such comparison, where we included the ~ 70 discovered planets in S-type configuration with stellar projected physical separations $s \leq 1000 \text{ au}$ and known planet orbital periods and minimum masses. The majority of these planets have been identified around stars in binary systems with large physical separation, typically mean values $s > 400 \text{ au}$. GJ 338 Bb is one of the least massive planets identified in one component of a stellar binary with separation well below of 400 au. The super-Earth planet around GJ 338 B adds to the increasing list of planets orbiting one of the stars of stellar binary systems. We remark the similarity of the two stellar components of the GJ 338 binary and also provide various existing examples with very similar components such as: WASP-2 AB, XO-2N, XO-2S, HD 177830 AB, Kepler 296 AB, WASP-94 AB and Gliese 15 AB (Collier Cameron et al. 2007; Torres et al. 2008; Meschiari et al. 2011; Desidera et al. 2014; Rowe et al. 2014; Neveu-VanMalle et al. 2014; Pinamonti et al. 2018).

Following commonly used expressions (e.g., Borucki & Summers 1984; Sackett 1999; Stevens & Gaudi 2013), we derived the probability that GJ 338 Bb transits in front of the disk of its parent star to be $5^{+2}_{-1} \%$. Using the unbiased forecasting model presented by Chen & Kipping (2017), we predict the planetary radius to be $3.13^{+1.37}_{-0.90} R_{\oplus}$; with this, the depth of any putative transit signal would be $0.24^{+0.05}_{-0.01} \%$ as computed for a planetary mass of $10.27^{+1.47}_{-1.38} M_{\oplus}$. *TESS* observed the GJ 338 binary system in sector 21 between 2020 January 21 and February 18. Due to the pixel scale of *TESS* ($21''/\text{pixel}$), the two stars GJ 338 A and B are not resolved.

We adopted the traditionally defined habitable zone as the circumstellar region in which a terrestrial-mass planet with a $\text{CO}_2\text{--H}_2\text{O--NO}_2$ atmosphere can sustain liquid water on its surface (Selsis et al. 2007). We estimated the theoretical equilibrium temperature of GJ 338 Bb by using the Stefan–Boltzmann equation, the stellar parameters of Table 1, and two extreme values of the albedo ($A = 0.0$ and 0.65 ; the highest albedo corresponds to the case of Venus in our Solar System). The results are $T_{\text{eq}} = 391 \pm 20 \text{ K}$ for a non-reflecting planet and $301 \pm 20 \text{ K}$ for the




Fig. 17: *Top panel*: planetary minimum mass versus orbital period for known Neptune-sized and super-Earth planets (blue dots) around M dwarfs. The planet GJ 338 Bb is plotted as the red star symbol. *Bottom panel*: planetary minimum mass versus semi-major axis for all known planets (color dots) in S-type configuration around binary systems, where the stellar binaries have projected physical separation $s \leq 1000$ au (color bar). The stellar binary separation between the A and B members of our binary system has a semi-major axis of 130.9 au and is plotted as star symbol.

high-reflectance planet. The error bars come from the uncertainties in the stellar luminosity and semi-major axis of the planet. Kopparapu et al. (2013) calculated conservative and optimistic estimates of the habitable zone (water-loss and maximum greenhouse limits); by using their determinations of the boundaries of the habitable zone, we found them to be in the interval 0.29–0.57 au around GJ 338 B. This implies that the planet GJ 338 Bb lies inside the inner boundary of the habitable zone which is consistent with the hot equilibrium temperature estimated above. Previous works (e.g., Kasting et al. 1993; Rodríguez et al. 2012; Barnes 2017) established that low-mass stellar hosts can induce strong tidal effects on potentially habitable planets since they tend to be on close-in orbits. True planet habitability may be

compromised for planets orbiting within the tidal lock radius of the star. Figure 16 reported by Kasting et al. (1993) shows the tidal lock radius for various stellar spectral classes, including M0 V. From this Figure, it becomes apparent that GJ 338 Bb is very likely tidally locked given its short orbit.

7. Conclusions

With the CARMENES spectrograph, we monitored the very bright M0.0 V stars GJ 338 B (HD 79211) and GJ 338 A (HD 79210), which form a wide binary in the nearby solar vicinity. We also provided a refined astrometric and spectroscopic solution of the stellar binary orbit by using all data from the literature available to us. The adopted Keplerian solution for the two stars yielded an orbital period of 1295 a and a semi-major axis of 130.9 au. The orbital period value reported here is significantly larger than those previously given in the literature (e.g., Chang 1972). Despite the great similarity between the two stars, we determined slightly different mass values for the A and B stellar components (0.69 ± 0.07 and $0.64 \pm 0.07 M_{\odot}$, respectively), suggesting that the A stellar member is about 7% more massive than the B component. Furthermore, they are also consistent with the masses derived by Schweitzer et al. (2019).

All activity indicators provided by CARMENES VIS data (Ca II infrared triplet, H α line, chromatic spectral index, the differential line width and cross correlation function analysis), together with the new photometry acquired by our group at LCO and SNO, indicate that GJ 338 B and A are active stars and rotate with periodicities of $P = 16.61 \pm 0.04$ d and $P = 16.3^{+3.5}_{-1.3}$ d, respectively, which agrees with the expected X-ray luminosity–activity relations.

The detailed analysis of CARMENES RVs led to the discovery of a super-Earth planet orbiting GJ 338 B with an orbital period of 24.45 ± 0.02 d, and a minimum mass of $10.27^{+1.47}_{-1.38} M_{\oplus}$, and a semi-major axis of 0.141 ± 0.005 au. Another period was found in the CARMENES RV data at ~ 8.3 d, but without sufficient number of arguments to prove its Keplerian nature and because it lies at half of the stellar rotation period, we attributed it to the first harmonic of the stellar activity. A related analysis concluded that GJ 338 A does not have planets of similar mass or more massive than GJ 338 Bb. The majority of the stellar binary systems with planet hosts have larger separations than the pair GJ 338 A and B and their planets are also larger in mass than GJ 338 Bb. Therefore, GJ 338 Bb has become one of the least massive planets ever discovered in one star of a binary system of relatively small separation.

Acknowledgements. We wish to thank the anonymous referee for helpful comments and suggestions, which helped to improve the manuscript. We are grateful to Prof. B. D. Mason for providing Washington Double Star astrometric data for the stellar binary. Based on observations collected at the German-Spanish Astronomical Center, Calar Alto, jointly operated by the Junta de Andalucía and the Instituto de Astrofísica de Andalucía (CSIC), and observations from Las Cumbres Observatory Global Telescope (LCOGT) network. LCOGT observations were partially acquired via program number TAU2019A-002 of the Wise Observatory, Tel-Aviv University, Israel. CARMENES is an instrument for the Centro Astronómico Hispano-Alemán de Calar Alto (CAHA, Almería, Spain). CARMENES is funded by the German Max-Planck-Gesellschaft (MPG), the Spanish Consejo Superior de Investigaciones Científicas (CSIC), the European Union through FEDER/ERF funds, and the members of the CARMENES Consortium (Max-Planck-Institut für Astronomie, Instituto de Astrofísica de Andalucía, Landessternwarte Königstuhl, Institut de Ciències de l'Espai, Institut für Astrophysik Göttingen, Universidad Complutense de Madrid, Thüringer Landessternwarte Tautenburg, Instituto de Astrofísica de Canarias, Hamburger Sternwarte, Centro de Astrobiología and Centro Astronómico Hispano-Alemán), with additional contributions by the Spanish Ministry of Economy, the state of Baden-Württemberg, the German Science Foundation (DFG), the Klaus Tschira Foundation (KTS), and by the Junta de Andalucía. This work is supported by

the Spanish Ministry for Science, Innovation, and Universities through projects AYA-2016-79425-C3-1/2/3-P, AYA2015-69350-C3-2-P, ESP2017-87676-C5-2-R, ESP2017-87143-R. The Instituto de Astrofísica de Andalucía is a Centre of Excellence “Severo Ochoa” (SEV-2017-0709). The Centro de Astrobiología (CAB, CSIC-INTA) is a Center of Excellence “Maria de Maeztu”.

References

- Abt, H. A. & Levy, S. G. 1973, *AJ*, 78, 1093
- Affer, L., Damasso, M., Micela, G., et al. 2019, *A&A*, 622, A193
- Alonso-Floriano, F. J., Morales, J. C., Caballero, J. A., et al. 2015, *A&A*, 577, A128
- Anglada-Escudé, G., Arriagada, P., Vogt, S. S., et al. 2012, *ApJ*, 751, L16
- Astudillo-Defru, N., Bonfils, X., Delfosse, X., et al. 2015, *A&A*, 575, A119
- Baranne, A., Queloz, D., Mayor, M., et al. 1996, *A&AS*, 119, 373
- Barnes, R. 2017, *Celestial Mechanics and Dynamical Astronomy*, 129, 509
- Bartlett, J. L., Ianna, P. A., & Begam, M. C. 2009, *PASP*, 121, 365
- Beavers, W. I. & Eitter, J. J. 1986, *ApJS*, 62, 147
- Bonfils, X., Lo Curto, G., Correia, A. C. M., et al. 2013, *A&A*, 556, A110
- Bopp, B. W. & Meredith, R. 1986, *PASP*, 98, 772
- Borucki, W. J. & Summers, A. L. 1984, *Icarus*, 58, 121
- Boyajian, T. S., von Braun, K., van Belle, G., et al. 2012, *ApJ*, 757, 112
- Brown, T. M., Baliber, N., Bianco, F. B., et al. 2013, *PASP*, 125, 1031
- Butler, R. P., Vogt, S. S., Laughlin, G., et al. 2017, *AJ*, 153, 208
- Caballero, J. A., Cortés-Contreras, M., Alonso-Floriano, F. J., et al. 2016a, in 19th Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun (CS19), 148
- Caballero, J. A., Guàrdia, J., López del Fresno, M., et al. 2016b, in *Proc. SPIE*, Vol. 9910, *Observatory Operations: Strategies, Processes, and Systems VI*, 99100E
- Carleo, I., Benatti, S., Lanza, A. F., et al. 2018, *A&A*, 613, A50
- Chang, K. 1972, *AJ*, 77, 759
- Chen, J. & Kipping, D. 2017, *ApJ*, 834, 17
- Claudi, R., Benatti, S., Carleo, I., et al. 2017, *European Physical Journal Plus*, 132, 364
- Collier Cameron, A., Bouchy, F., Hébrard, G., et al. 2007, *MNRAS*, 375, 951
- Cortés-Contreras, M. 2016, PhD thesis, Universidad Complutense de Madrid, Spain
- Cortés-Contreras, M., Béjar, V. J. S., Caballero, J. A., et al. 2017, *A&A*, 597, A47
- Damasso, M., Bonomo, A. S., Astudillo-Defru, N., et al. 2018, *A&A*, 615, A69
- Dawson, R. I. & Fabrycky, D. C. 2010, *ApJ*, 722, 937
- Deeming, T. J. 1975, *Ap&SS*, 36, 137
- Desidera, S., Bonomo, A. S., Claudi, R. U., et al. 2014, *A&A*, 567, L6
- Díez Alonso, E., Caballero, J. A., Montes, D., et al. 2019, *A&A*, 621, A126
- Dressing, C. D. & Charbonneau, D. 2013, *ApJ*, 767, 95
- Dressing, C. D. & Charbonneau, D. 2015, *ApJ*, 807, 45
- Eiroa, C., Marshall, J. P., Mora, A., et al. 2013, *A&A*, 555, A11
- Foreman-Mackey, D., Agol, E., Ambikasaran, S., & Angus, R. 2017, *AJ*, 154, 220
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, 125, 306
- Fuhrmeister, B., Czesla, S., Schmitt, J. H. M. M., et al. 2019, *A&A*, 623, A24
- Gagné, J., Plavchan, P., Gao, P., et al. 2016, *ApJ*, 822, 40
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, *A&A*, 616, A1
- Gaidos, E. & Mann, A. W. 2014, *ApJ*, 791, 54
- Gaidos, E., Mann, A. W., Kraus, A. L., & Ireland, M. 2016, *MNRAS*, 457, 2877
- Gaidos, E., Mann, A. W., Lépine, S., et al. 2014, *MNRAS*, 443, 2561
- García-Piquer, A., Morales, J. C., Ribas, I., et al. 2017, *A&A*, 604, A87
- Głębicki, R. & Gnaniński, P. 2005, in *ESA Special Publication*, Vol. 560, 13th Cambridge Workshop on Cool Stars, Stellar Systems and the Sun, ed. F. Favata, G. A. J. Hussain, & B. Battrick, 571
- Gomes da Silva, J., Santos, N. C., Bonfils, X., et al. 2011, *A&A*, 534, A30
- Gomes da Silva, J., Santos, N. C., Bonfils, X., et al. 2012, *A&A*, 541, A9
- Güntzel-Lingner, 1954, *IAU Commission Binary and Multiple Star Systems Information Bulletin* 4, ed. P. Muller
- Hartkopf, W. I., Mason, B. D., & Worley, C. E. 2001, *AJ*, 122, 3472
- Hartman, J. D., Bakos, G. Á., Kovács, G., & Noyes, R. W. 2010, *MNRAS*, 408, 475
- Hartman, J. D., Gaudi, B. S., Pinsonneault, M. H., et al. 2009, *ApJ*, 691, 342
- Henry, T. J., Jao, W.-C., Subasavage, J. P., et al. 2006, *AJ*, 132, 2360
- Hopmann, 1954, *IAU Commission Binary and Multiple Star Systems Information Bulletin* 3, ed. P. Muller
- Howard, A. W., Marcy, G. W., Bryson, S. T., et al. 2012, *ApJS*, 201, 15
- Huélamo, N., Figueira, P., Bonfils, X., et al. 2008, *A&A*, 489, L9
- Isaacson, H. & Fischer, D. 2010, *ApJ*, 725, 875
- Jenkins, J. S., Ramsey, L. W., Jones, H. R. A., et al. 2009, *ApJ*, 704, 975
- Kaminski, A., Trifonov, T., Caballero, J. A., et al. 2018, *A&A*, 618, A115
- Kasting, J. F., Whitmire, D. P., & Reynolds, R. T. 1993, *Icarus*, 101, 108
- Kiselev, A. A. & Kiyayeva, O. V. 1980, *AZh*, 57, 1227
- Kiselev, A. A. & Romanenko, L. G. 1996, *Astronomy Reports*, 40, 795
- Kochanek, C. S., Shappee, B. J., Stanek, K. Z., et al. 2017, *PASP*, 129, 104502
- Kopparapu, R. K., Ramirez, R., Kasting, J. F., et al. 2013, *ApJ*, 765, 131
- Leto, G., Pagano, I., Buemi, C. S., & Rodono, M. 1997, *A&A*, 327, 1114
- Levenberg, K. 1944, *Quarterly of Applied Mathematics*, 2, 164
- Liddle, A. R. 2007, *MNRAS*, 377, L74
- Luque, R., Nowak, G., Pallé, E., et al. 2018, *A&A*, 620, A171
- Luque, R., Pallé, E., Kossakowski, D., et al. 2019, *A&A*, 628, A39
- Malkov, O. Y., Tamazian, V. S., Docobo, J. A., & Chulkov, D. A. 2012, *A&A*, 546, A69
- Marquardt, D. 1963, *SIAM Journal on Applied Mathematics*, 11, 431
- Martín, E. L., Guenther, E., Zapatero Osorio, M. R., Bouy, H., & Wainscoat, R. 2006, *ApJ*, 644, L75
- Martínez-Arnáiz, R., López-Santiago, J., Crespo-Chacón, I., & Montes, D. 2011, *MNRAS*, 414, 2629
- Martínez-Arnáiz, R., Maldonado, J., Montes, D., Eiroa, C., & Montesinos, B. 2010, *A&A*, 520, A79
- Mason, B. D., Wycoff, G. L., Hartkopf, W. I., Douglass, G. G., & Worley, C. E. 2001, *AJ*, 122, 3466
- McCully, C., Volgenau, N. H., Harbeck, D.-R., et al. 2018, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 10707, *Software and Cyberinfrastructure for Astronomy V*, 107070K
- Meschiari, S., Laughlin, G., Vogt, S. S., et al. 2011, *ApJ*, 727, 117
- Montes, D., Fernández-Figueroa, M. J., De Castro, E., et al. 2000, *A&AS*, 146, 103
- Morales, J. C., Mustill, A. J., Ribas, I., et al. 2019, *Science*, 365, 1441
- Morbey, C. L. & Griffin, R. F. 1987, *ApJ*, 317, 343
- Mortier, A., Faria, J. P., Correia, C. M., Santerne, A., & Santos, N. C. 2015, *A&A*, 573, A101
- Morton, T. D. & Swift, J. 2014, *ApJ*, 791, 10
- Moutou, C., Hébrard, E. M., Morin, J., et al. 2017, *MNRAS*, 472, 4563
- Mulders, G. D., Pascucci, I., & Apai, D. 2015, *ApJ*, 798, 112
- Nagel, E., Czesla, S., Schmitt, J. H. M. M., et al. 2019, *A&A*, 622, A153
- Neves, V., Bonfils, X., Santos, N. C., et al. 2013, *A&A*, 551, A36
- Neveu-VanMalle, M., Queloz, D., Anderson, D. R., et al. 2014, *A&A*, 572, A49
- Newton, E. R., Irwin, J., Charbonneau, D., et al. 2017, *ApJ*, 834, 85
- Newton, E. R., Irwin, J., Charbonneau, D., Berta-Thompson, Z. K., & Dittmann, J. A. 2016, *ApJ*, 821, L19
- Nidever, D. L., Marcy, G. W., Butler, R. P., Fischer, D. A., & Vogt, S. S. 2002, *ApJS*, 141, 503
- Osten, R. A., Hawley, S. L., Allred, J. C., Johns-Krull, C. M., & Roark, C. 2005, *ApJ*, 621, 398
- Passegger, V. M., Reiniers, A., Jeffers, S. V., et al. 2018, *A&A*, 615, A6
- Passegger, V. M., Schweitzer, A., Shulyak, D., et al. 2019, *A&A*, 627, A161
- Perger, M., Scandariato, G., Ribas, I., et al. 2019, *A&A*, 624, A123
- Perruchot, S., Kohler, D., Bouchy, F., et al. 2008, in *Ground-based and Airborne Instrumentation for Astronomy II*, Vol. 7014, 70140J
- Pinamonti, M., Damasso, M., Marzari, F., et al. 2018, *A&A*, 617, A104
- Quirrenbach, A., Amado, P. J., Caballero, J. A., et al. 2016, in *Proc. SPIE*, Vol. 9908, *Ground-based and Airborne Instrumentation for Astronomy VI*, 990812
- Rabl, G. & Dvorak, R. 1988, *A&A*, 191, 385
- Reiners, A., Joshi, N., & Goldman, B. 2012, *AJ*, 143, 93
- Reiners, A., Ribas, I., Zechmeister, M., et al. 2018a, *A&A*, 609, L5
- Reiners, A., Schüssler, M., & Passegger, V. M. 2014, *ApJ*, 794, 144
- Reiners, A., Zechmeister, M., Caballero, J. A., et al. 2018b, *A&A*, 612, A49
- Ribas, I., Tuomi, M., Reiniers, A., et al. 2018, *Nature*, 563, 365
- Rodríguez, A., Callegari, N., Michtchenko, T. A., & Hussmann, H. 2012, *MNRAS*, 427, 2239
- Rodríguez, E., García, J. M., Costa, V., et al. 2010, *MNRAS*, 408, 2149
- Romanenko, L. G. & Chentsov, E. L. 1994, *AZh*, 71, 278
- Rowe, J. F., Bryson, S. T., Marcy, G. W., et al. 2014, *ApJ*, 784, 45
- Sackett, P. D. 1999, in *NATO Advanced Science Institutes (ASI) Series C*, Vol. 532, *NATO Advanced Science Institutes (ASI) Series C*, ed. J.-M. Mariotti & D. Alloin, 189
- Salvatier, J., Wiecki, T. V., & Fonnesbeck, C. 2016, *PeerJ Computer Science*, 2, e55
- Sanz-Fornada, J., Micela, G., Ribas, I., et al. 2011, *A&A*, 532, A6
- Sarkis, P., Henning, T., Kürster, M., et al. 2018, *AJ*, 155, 257
- Schöfer, P., Jeffers, S. V., Reiniers, A., et al. 2019, *A&A*, 623, A44
- Schweitzer, A., Passegger, V. M., Cifuentes, C., et al. 2019, *A&A*, 625, A68
- Selsis, F., Kasting, J. F., Lévrard, B., et al. 2007, *A&A*, 476, 1373
- Sharma, K., Prugniel, P., & Singh, H. P. 2016, *A&A*, 585, A64
- Shulyak, D., Reiniers, A., Nagel, E., et al. 2019, *A&A*, 626, A86
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, 131, 1163
- Soubiran, C., Jasiewicz, G., Chemin, L., et al. 2018, *A&A*, 616, A7

- Sozzetti, A., Bernagozzi, A., Bertolini, E., et al. 2013, in *European Physical Journal Web of Conferences*, Vol. 47, *European Physical Journal Web of Conferences*, 03006
- Stevens, D. J. & Gaudi, B. S. 2013, *PASP*, 125, 933
- Suárez Mascareño, A., González Hernández, J. I., Rebolo, R., et al. 2017, *A&A*, 605, A92
- Suárez Mascareño, A., Rebolo, R., González Hernández, J. I., & Esposito, M. 2015, *MNRAS*, 452, 2745
- Tal-Or, L., Trifonov, T., Zucker, S., Mazeh, T., & Zechmeister, M. 2019, *MNRAS*, 484, L8
- Tal-Or, L., Zechmeister, M., Reiners, A., et al. 2018, *A&A*, 614, A122
- Tetzlaff, N., Neuhauser, R., & Hohle, M. M. 2011, *MNRAS*, 410, 190
- Theano Development Team. 2016, *arXiv e-prints*, abs/1605.02688
- Tokunaga, A. T., Toomey, D. W., Carr, J., Hall, D. N. B., & Epps, H. W. 1990, in *Proc. SPIE*, Vol. 1235, *Instrumentation in Astronomy VII*, ed. D. L. Crawford, 131
- Torres, G., Winn, J. N., & Holman, M. J. 2008, *ApJ*, 677, 1324
- Trifonov, T., Kürster, M., Zechmeister, M., et al. 2018, *A&A*, 609, A117
- van den Bos, W. H. 1955, *MNRAS*, 115, 187
- Vogt, S. S., Allen, S. L., Bigelow, B. C., et al. 1994, in *Proc. SPIE*, Vol. 2198, *Instrumentation in Astronomy VIII*, ed. D. L. Crawford & E. R. Craine, 362
- Wilson, O. C. 1941, *ApJ*, 93, 29
- Wright, J. T. & Howard, A. W. 2009, *ApJS*, 182, 205
- Wright, N. J., Drake, J. J., Mamajek, E. E., & Henry, G. W. 2011, *ApJ*, 743, 48
- Zechmeister, M., Dreizler, S., Ribas, I., et al. 2019, *A&A*, 627, A49
- Zechmeister, M. & Kürster, M. 2009, *A&A*, 496, 577
- Zechmeister, M., Reiners, A., Amado, P. J., et al. 2018, *A&A*, 609, A12

Appendix A: Figures




Fig. A.1: MCMC distribution of the combined astrometric and spectroscopic analysis of the GJ 338 AB binary system. The vertical dashed lines indicate the 16%, 50%, and 84% quantiles of the fitted parameters; this corresponds to 1σ uncertainty. The reference time of T_0 distribution is 2,200,000.




Fig. A.2: MCMC distribution of the planet orbiting GJ 338 B. The vertical dashed lines indicate the 16%, 50%, and 84% quantiles of the fitted parameters; this corresponds to 1σ uncertainty. The best fit is colored.

Appendix B: Tables

Tables B.1–B.6 are only available at the CDS (see title footnote on page 1).

Table B.4: Relative radial velocity measurements of GJ 338 B with CARMENES in the NIR.

Epoch [BJD] (d)	RV _{NIR} (m s ⁻¹)
2457695.74104	-3.06 ± 20.96
2457698.53211	-52.24 ± 31.33
2457699.58824	-7.92 ± 20.26
2457701.65194	-0.48 ± 8.62
2457703.73873	-6.41 ± 7.35
2457704.53247	8.08 ± 6.78
2457705.58085	-144.95 ± 90.90
2457712.55064	1.31 ± 7.00
2457752.72317	1.65 ± 10.47
2457754.3795	18.47 ± 8.39
2457756.6115	5.85 ± 7.08
2457757.54099	10.02 ± 12.10
2457759.6843	4.13 ± 8.54
2457760.4353	-5.99 ± 9.74
2457761.41194	8.45 ± 12.05
2457762.54788	-13.7 ± 11.37
2457763.58288	2.72 ± 11.26
2457766.5783	15.78 ± 7.83
2457768.59089	3.84 ± 6.63
2457771.64125	3.27 ± 9.55
2457779.43054	-8.54 ± 12.41
2457787.5346	10.99 ± 8.01
2457788.44339	18.98 ± 8.67
2457790.50189	11.54 ± 8.36
2457791.32384	10.69 ± 10.33
2457792.60432	-4.73 ± 14.37
2457793.34107	3.40 ± 9.11
2457794.64241	-3.31 ± 8.84
2457798.43155	-4.73 ± 7.57
2457799.65549	-6.61 ± 13.76
2457800.35454	9.48 ± 8.64
2457802.65848	-8.5 ± 7.44
2457806.46081	9.94 ± 7.39
2457808.60857	16.68 ± 31.66
2457817.6022	20.28 ± 10.71
2457818.56409	2.07 ± 7.49
2457819.32766	1.09 ± 7.28
2457821.35015	3.42 ± 7.15
2457822.61006	-0.31 ± 15.39
2457824.38691	4.44 ± 5.95
2457828.43287	-10.05 ± 12.89
2457829.36917	-18.36 ± 11.07
2457833.45451	-2.84 ± 5.64
2457849.37139	-0.12 ± 8.55
2457855.33463	-3.02 ± 20.03
2457855.3403	-2.82 ± 7.48
2457857.37306	10.56 ± 6.97
2457859.34242	7.25 ± 5.97
2457860.34365	4.33 ± 6.60
2457862.36904	-3.92 ± 6.88
2457863.35123	14.76 ± 9.28
2457864.42355	-5.33 ± 9.67
2457875.44635	23.60 ± 40.47
2457879.36289	-5.86 ± 17.79
2457883.4567	-1.78 ± 8.00
2457886.3354	-3.04 ± 10.32
2457889.45065	9.30 ± 17.61
2457896.40923	0.99 ± 7.76
2458002.69546	11.39 ± 8.82
2458018.65934	14.09 ± 7.62
2458022.72064	-1.73 ± 8.05
2458023.70637	-14.57 ± 6.02

Doppler measurements for GJ 338 B in the NIR – continued.

Epoch [BJD] (d)	RV _{NIR} (m s ⁻¹)
2458027.64945	0.79 ± 7.42
2458028.70023	-7.12 ± 7.45
2458060.60622	-20.71 ± 11.15
2458064.58946	-28.85 ± 7.74
2458084.51992	3.03 ± 6.03
2458085.61449	-0.01 ± 14.99
2458088.63985	-7.87 ± 12.55
2458089.56545	-17.31 ± 11.87
2458093.57994	-1.44 ± 7.13
2458094.64509	-6.38 ± 5.91
2458095.55142	4.50 ± 5.62
2458097.63023	2.38 ± 11.33
2458109.48135	-2.08 ± 6.77
2458110.42561	-0.52 ± 8.08
2458114.39908	-18.81 ± 11.97
2458117.45108	-0.15 ± 8.71
2458118.4391	-4.04 ± 7.04
2458119.59615	12.19 ± 12.92
2458120.44254	5.82 ± 14.75
2458127.51091	18.34 ± 29.77
2458136.36142	6.12 ± 5.13
2458141.67104	5.60 ± 11.28
2458149.44293	0.70 ± 6.50
2458160.3289	-1.62 ± 28.6
2458161.36664	-10.58 ± 7.50
2458166.44431	3.51 ± 8.69
2458166.4492	-14.39 ± 15.30
2458166.46016	5.69 ± 7.07
2458167.31522	-1.05 ± 10.82
2458169.61017	0.13 ± 14.68
2458171.4728	3.29 ± 8.03
2458172.39829	-1.09 ± 7.14
2458175.42501	-4.63 ± 6.50
2458187.50091	9.67 ± 10.06
2458199.3126	9.66 ± 8.57
2458206.39496	-4.28 ± 9.71
2458207.3833	-1.01 ± 14.65
2458209.46715	-9.56 ± 8.38
2458211.4099	-20.60 ± 7.87
2458212.40058	-19.62 ± 8.55
2458215.29871	-3.56 ± 6.89
2458217.36205	6.60 ± 10.01
2458218.36262	9.32 ± 16.48
2458225.34727	-1.91 ± 6.84
2458229.45336	-4.05 ± 10.66
2458237.44359	-10.96 ± 10.63
2458238.43597	-8.05 ± 10.11
2458245.41564	-10.63 ± 7.19
2458246.46316	-11.49 ± 11.04
2458249.47983	0.23 ± 9.36
2458258.37302	0.0 ± 9.13
2458260.3685	-9.11 ± 6.50
2458261.37097	-9.07 ± 9.07
2458262.34507	5.46 ± 12.18
2458263.40113	3.60 ± 10.16
2458264.3803	-7.61 ± 7.45
2458266.42172	8.41 ± 9.59
2458273.39345	4.43 ± 7.92
2458284.34815	-2.21 ± 12.39
2458291.35922	-19.62 ± 8.76

Table B.5: Relative astrometric position of the binary companion GJ 338 B to GJ 338 A.

Epoch (a)	θ (deg)	ρ (")	Ref. ^(a)
1821.50	43.8	21.12	StF1837
1824.46	44.2	20.80	S__1906A
1831.13	51.5	20.	HJ_1833c
1833.54	48.5	20.03	StF1837
1840.32	50.6	20.15	Stt1878
1841.44	50.5	16.68	Mad1842a
1841.54	50.6	20.32	Da_1867
1842.44	50.4	19.74	Kai1865
1846.29	51.4	20.10	Mad1906A
1848.29	51.8	19.87	Da_1867
1848.57	52.4	19.94	Stt1878
1851.09	53.0	19.47	Mad1856
1852.41	53.0	19.24	Mad1856
1855.17	53.7	19.59	D__1883
1856.30	54.7	20.06	Wnc1869
1858.38	54.9	19.91	Mad1859
1859.98	53.6	19.65	Mad1906A
1863.12	55.7	19.73	D__1884
1863.25	53.45	19.33	Mai1863
1863.25	53.4	19.33	Mad1906A
1866.79	56.2	19.93	Tal1868
1871.24	57.3	19.75	Du_1876
1871.66	57.0	20.08	Tal1873
1874.22	57.0	20.0	Gld1875
1874.23	57.0	20.30	Tal1878
1874.81	58.3	19.62	D__1884
1876.12	58.6	19.64	D__1883
1876.46	59.6	19.90	Plm1878
1877.38	58.9	19.65	Fla1906A
1877.975	59.2	19.64	HI_1877a
1878.01	59.1	19.62	Stt1893a
1878.379	59.7	19.650	Je_1880i
1878.92	58.2	19.52	Gny1879
1880.206	59.65	19.50	Nst1904
1880.375	60.7	19.527	Je_1880i
1881.12	60.2	19.51	Big1883
1883.01	61.2	19.35	Eng1885a
1884.38	61.5	19.72	Sbk1892
1885.23	60.8	19.45	Nst1904
1891.09	62.5	19.50	Bu_1894
1891.12	62.4	19.38	Bu_1894
1893.25	63.33	19.615	Enh1895
1895.05	63.5	19.28	Rnz1935
1896.98	64.1	19.25	Vil1961
1899.28	64.5	19.30	Bry1899
1900.08	65.05	18.96	Es_1902b
1901.11	65.55	18.77	Es_1902b
1901.30	65.3	19.05	Bry1901
1902.19	65.13	18.60	Es_1902b
1903.08	65.7	19.301	WFC1998
1903.12	65.8	19.203	WFC1998
1903.18	65.6	19.031	WFC1998
1903.19	65.7	19.12	Dob1927
1904.31	65.9	19.24	L__1904
1905.17	65.4	19.05	Bu_1906
1905.25	65.9	20.19	Frm1907
1905.833	66.6	19.15	VBs1906
1907.08	66.9	19.12	Sma1931
1907.17	66.3	19.49	Lau1907a
1907.20	66.5	18.82	Es_1908a

Relative astrometric position – continued.

Epoch (a)	θ (deg)	ρ (")	Ref. ^(a)
1908.32	66.8	19.26	Lau1908c
1908.32	66.9	19.07	Hgz1909
1908.34	66.6	19.10	Prz1908
1909.30	67.7	18.95	Dob1927
1909.39	66.9	19.17	Lau1909b
1911.23	67.4	19.10	Lau1911b
1912.22	67.8	18.88	Vou1922
1912.26	67.8	19.04	Neu1923
1913.131	67.8	19.02	VBs1920
1913.16	68.1	18.93	Vou1922
1913.19	68.2	18.87	Jam1921
1913.32	68.6	18.74	Sla1913
1913.37	68.9	19.16	Bry1921
1914.15	70.4	19.44	Frk1914a
1914.38	68.3	19.07	Vys1936
1914.68	68.4	18.91	Dob1927
1915.10	69.1	18.936	WFD1931
1915.28	69.1	19.31	Ptt1917
1915.32	68.5	18.96	Com1929
1915.34	70.9	18.52	Rab1923
1916.752	69.01	18.916	Hgz1920
1917.7	69.2	18.171	WFD1926a
1918.10	69.5	18.89	Gui1931
1918.19	69.1	18.596	WFD1925
1918.34	69.1	19.09	Es_1919
1918.35	69.4	19.07	Bea1930
1919.18	69.2	18.88	VvS1919
1919.46	67.8	19.279	WFD1933b
1920.222	70.0	18.83	B_1923a
1920.28	69.5	18.69	Cha1925b
1920.57	69.85	18.844	Chg1972
1921.23	70.7	19.36	Abt1922
1923.208	70.6	18.84	VBs1927a
1923.30	70.17	18.817	Lbz1929
1923.36	70.10	18.892	Prz1926
1924.08	71.3	18.28	Blo1931
1924.95	71.2	18.76	Sha1927
1925.28	71.21	18.738	Prz1926
1925.35	71.7	18.67	Dob1927
1925.60	71.21	18.625	Chg1972
1925.95	72.3	18.249	WFD1935
1926.217	74.0	18.92	Baz1927b
1926.24	71.43	18.674	Lbz1929
1926.31	71.55	18.799	Prz1926
1926.33	71.59	18.793	Prz1926
1927.34	71.72	18.697	Lbz1929
1928.17	72.05	18.69	Frw1936
1928.35	72.6	18.67	Kom1935a
1928.35	72.24	18.713	Prz1926
1928.37	72.22	18.618	Lbz1929
1930.03	71.8	19.097	WFC1958a
1930.09	72.6	18.50	Bon1938
1930.27	74.2	18.52	Baz1932b
1931.30	73.0	18.65	Kom1933
1932.24	73.5	18.62	Kom1933
1934.23	73.3	18.74	Pok1936
1934.36	74.2	18.57	Baz1936b
1934.43	74.1	18.65	Dob1934
1937.26	74.5	18.45	Vat1939
1937.99	74.76	18.503	Jef1951

Relative astrometric position – continued.

Epoch (a)	θ (deg)	ρ (")	Ref. ^(a)
1939.27	74.9	18.40	Scd1940a
1939.62	75.03	18.477	Chg1972
1940.04	75.34	18.444	Str1946
1941.13	75.63	18.428	Str1946
1942.27	76.7	18.43	Dur1944b
1943.00	76.3	18.14	Baz1945a
1944.16	76.30	18.400	Chg1972
1946.82	77.6	18.43	Mun1948
1946.924	77.10	18.325	DeO1954
1947.01	76.8	18.190	WFC1959a
1947.110	77.28	18.336	DeO1954
1947.20	77.5	18.57	Mlr1947
1948.38	77.1	18.63	Jan1955
1949.256	77.91	18.258	DeO1954
1949.258	77.90	18.275	DeO1954
1949.278	77.90	18.269	DeO1954
1949.283	77.79	18.246	DeO1954
1949.955	78.01	18.267	DeO1954
1950.	93.3	17.68	Skf2006c
1950.018	77.93	18.235	DeO1954
1950.029	78.06	18.264	DeO1954
1950.19	78.02	18.266	Chg1972
1950.395	78.22	18.259	Jef1978
1950.88	78.35	18.219	Jan1953
1951.050	78.20	18.249	DeO1954
1951.145	78.34	18.214	DeO1954
1951.145	78.39	18.205	DeO1954
1951.172	78.36	18.256	DeO1954
1951.306	78.38	18.205	DeO1954
1951.97	78.80	18.263	Jan1953
1952.079	78.63	18.220	DeO1954
1952.13	78.0	18.17	Wie1956c
1952.156	78.70	18.219	DeO1954
1952.205	78.68	18.206	DeO1954
1953.05	78.96	18.258	Jan1953
1953.150	87.760	17.620	PkO2016
1953.250	78.99	18.184	Jef1978
1954.04	79.19	18.170	Chg1972
1954.181	79.19	18.183	Gzl1962
1954.181	79.45	18.192	Gzl1962
1954.225	79.47	18.161	Gzl1962
1954.33	77.9	18.31	Jan1955
1954.952	79.50	18.151	Gzl1962
1955.064	79.48	18.151	Gzl1962
1955.146	79.72	18.152	Gzl1962
1955.162	79.74	18.121	Gzl1962
1955.26	81.0	18.17	Wor1956b
1955.28	79.9	18.02	Rab1961b
1956.232	80.02	18.174	Gzl1962
1956.28	80.0	18.04	Rab1961b
1956.36	79.88	18.158	Bot1962
1957.097	80.09	18.128	Gzl1962
1957.14	80.12	18.115	Bot1962
1957.19	77.9	18.55	Fle1958
1957.248	80.14	18.120	Gzl1962
1957.258	80.20	18.128	Gzl1962
1957.272	80.18	18.126	Gzl1962
1958.11	81.6	18.159	WFC1975
1958.21	80.35	18.107	Bot1962
1958.261	80.44	18.103	Ge_1964

Relative astrometric position – continued.

Epoch (a)	θ (deg)	ρ (")	Ref. ^(a)
1958.283	80.37	18.063	Ge_1964
1958.299	80.39	18.087	Ge_1964
1959.86	80.95	18.041	Kpr1992
1960.67	81.18	18.050	Chg1972
1961.44	81.23	18.01	Hau1965
1962.173	81.820	18.022	PkO2003
1962.173	81.82	18.022	PkO1988
1962.20	81.5	18.12	B__1962d
1963.111	81.833	17.996	PkO2014b
1963.114	81.786	18.006	PkO2016
1963.115	82.130	18.007	PkO2003
1963.115	82.13	18.007	PkO1988
1963.271	81.77	17.967	USN1969
1963.60	81.94	17.999	Chg1972
1963.890	82.01	17.939	USN1969
1963.895	82.02	17.950	USN1969
1963.914	81.97	17.968	USN1969
1963.933	81.92	17.947	USN1969
1964.220	82.177	17.982	PkO2016
1964.221	82.410	17.965	PkO2003
1964.221	82.41	17.965	PkO1988
1964.221	82.183	17.979	PkO2014b
1965.247	82.450	17.957	PkO2016
1965.261	82.751	17.961	PkO2003
1965.261	82.75	17.961	PkO1988
1965.266	82.428	17.951	PkO2014b
1965.35	82.6	17.80	Baz1967
1965.58	82.49	17.965	Chg1972
1966.130	82.723	17.939	PkO2016
1966.134	82.977	17.943	PkO2003
1966.134	82.98	17.943	PkO1988
1966.179	82.744	17.926	PkO2014b
1966.896	82.88	17.912	USN1969
1967.117	83.007	17.924	PkO2016
1967.120	83.259	17.906	PkO2003
1967.120	83.26	17.906	PkO1988
1967.121	83.013	17.925	PkO2014b
1967.21	82.98	17.928	Chg1972
1968.047	83.27	17.902	USN1974
1968.047	83.32	17.892	USN1974
1968.151	83.31	17.901	USN1974
1968.151	83.32	17.893	USN1974
1968.171	83.260	17.898	PkO2014b
1968.187	83.574	17.846	PkO2003
1968.190	83.290	17.895	PkO2016
1968.193	83.57	17.894	PkO1988
1968.214	83.31	17.910	USN1974
1968.214	83.31	17.912	USN1974
1968.940	83.51	17.882	USN1974
1968.940	83.51	17.885	USN1974
1969.074	83.60	17.864	USN1974
1969.074	83.61	17.874	USN1974
1969.087	83.570	17.882	PkO2016
1969.184	83.591	17.859	PkO2014b
1969.185	83.858	17.890	PkO2003
1969.185	83.86	17.890	PkO1988
1969.200	83.51	17.878	USN1974
1969.200	83.54	17.875	USN1974
1969.61	83.70	17.879	Chg1972
1970.032	83.71	17.858	USN1974

Relative astrometric position – continued.

Epoch (a)	θ (deg)	ρ (")	Ref. ^(a)
1970.032	83.76	17.861	USN1974
1970.032	83.76	17.864	USN1974
1970.032	83.78	17.869	USN1974
1970.035	83.81	17.875	USN1974
1970.035	83.87	17.865	USN1974
1970.082	83.83	17.852	USN1974
1970.082	83.88	17.863	USN1974
1970.120	83.85	17.863	USN1974
1970.142	83.84	17.861	USN1974
1970.155	83.84	17.852	USN1974
1970.158	83.88	17.859	USN1974
1970.158	83.96	17.855	USN1974
1970.158	83.98	17.857	USN1974
1970.161	83.86	17.856	USN1974
1970.161	83.92	17.858	USN1974
1970.180	83.81	17.888	USN1974
1970.180	83.81	17.898	USN1974
1970.180	83.85	17.852	USN1974
1970.180	83.91	17.859	USN1974
1970.180	83.93	17.863	USN1974
1970.180	83.97	17.863	USN1974
1970.199	83.85	17.869	USN1974
1970.199	83.85	17.870	USN1974
1970.199	83.86	17.863	USN1974
1970.230	84.17	17.869	PkO1988
1970.233	83.914	17.860	PkO2016
1970.234	83.93	17.863	USN1974
1970.234	83.944	17.847	PkO2014b
1970.270	83.90	17.860	USN1974
1970.278	83.94	17.855	USN1974
1970.559	84.202	17.866	PkO2003
1971.049	84.33	17.858	PkO1988
1971.051	84.14	17.837	USN1974
1971.051	84.17	17.831	USN1974
1971.062	84.163	17.843	PkO2016
1971.083	84.13	17.835	USN1974
1971.083	84.14	17.833	USN1974
1971.083	84.15	17.835	USN1974
1971.083	84.16	17.833	USN1974
1971.084	84.16	17.825	USN1974
1971.084	84.17	17.832	USN1974
1971.084	84.17	17.834	USN1974
1971.084	84.19	17.816	USN1974
1971.125	84.186	17.834	PkO2014b
1971.129	84.367	17.855	PkO2003
1971.174	84.17	17.846	USN1974
1971.174	84.18	17.846	USN1974
1971.174	84.19	17.844	USN1974
1971.184	84.18	17.845	USN1974
1971.184	84.19	17.830	USN1974
1971.184	84.19	17.832	USN1974
1971.184	84.20	17.834	USN1974
1971.184	84.22	17.832	USN1974
1971.184	84.23	17.834	USN1974
1971.225	84.19	17.838	USN1974
1971.225	84.21	17.839	USN1974
1971.234	84.22	17.835	USN1974
1971.234	84.23	17.839	USN1974
1971.234	84.24	17.835	USN1974
1971.242	84.17	17.837	USN1974

Relative astrometric position – continued.

Epoch (a)	θ (deg)	ρ (")	Ref. ^(a)
1971.242	84.18	17.833	USN1974
1971.242	84.19	17.838	USN1974
1971.264	84.18	17.839	USN1974
1971.264	84.22	17.837	USN1974
1971.264	84.22	17.845	USN1974
1971.264	84.24	17.839	USN1974
1971.277	84.16	17.840	USN1974
1971.277	84.25	17.835	USN1974
1971.283	84.20	17.853	USN1974
1971.283	84.23	17.829	USN1974
1971.283	84.24	17.836	USN1974
1971.286	84.21	17.831	USN1974
1971.286	84.25	17.844	USN1974
1971.288	84.22	17.828	USN1974
1971.288	84.25	17.840	USN1974
1971.288	84.25	17.842	USN1974
1971.288	84.26	17.840	USN1974
1971.854	84.39	17.832	USN1974
1971.925	84.31	17.813	USN1974
1971.925	84.31	17.833	USN1974
1971.925	84.38	17.820	USN1974
1971.925	84.39	17.830	USN1974
1971.925	84.39	17.832	USN1974
1971.925	84.41	17.823	USN1974
1972.050	84.412	17.803	PkO2014b
1972.108	84.67	17.841	PkO1988
1972.108	84.672	17.841	PkO2003
1972.111	84.495	17.819	PkO2016
1972.162	84.50	17.820	USN1974
1972.255	84.50	17.804	USN1974
1972.255	84.56	17.812	USN1974
1972.999	79.0	19.00	CIJ2003
1973.047	84.74	17.806	USN1978
1973.047	84.75	17.803	USN1978
1973.047	84.76	17.808	USN1978
1973.047	84.79	17.822	USN1978
1973.095	84.785	17.807	PkO2016
1973.109	84.953	17.813	PkO2003
1973.110	84.95	17.813	PkO1988
1973.115	84.770	17.804	PkO2014b
1973.118	84.74	17.790	USN1978
1973.118	84.77	17.800	USN1978
1973.132	84.68	17.794	USN1978
1973.19	84.7	17.68	Ary1977
1973.214	84.81	17.796	USN1978
1973.214	84.86	17.806	USN1978
1973.227	84.74	17.805	USN1978
1973.227	84.79	17.795	USN1978
1974.156	85.07	17.792	Kpr1979
1974.180	85.05	17.789	USN1978
1974.180	85.11	17.769	USN1978
1974.203	85.094	17.786	PkO2016
1974.206	85.33	17.809	PkO1988
1974.207	85.328	17.809	PkO2003
1974.213	85.086	17.785	PkO2014b
1974.232	85.07	17.782	USN1978
1974.232	85.09	17.793	USN1978
1975.092	85.26	17.767	USN1978
1975.092	85.37	17.769	USN1978
1975.103	85.37	17.755	USN1978

Relative astrometric position – continued.

Epoch (a)	θ (deg)	ρ (")	Ref. ^(a)
1975.103	85.37	17.758	USN1978
1975.149	85.599	17.780	PkO2003
1975.150	85.60	17.777	PkO1988
1975.151	85.375	17.765	PkO2016
1975.179	85.381	17.767	PkO2014b
1975.182	85.38	17.768	USN1978
1975.272	85.41	17.745	USN1978
1975.272	85.41	17.746	USN1978
1975.302	85.40	17.764	USN1978
1975.302	85.41	17.760	USN1978
1975.876	85.58	17.744	USN1978
1975.876	85.61	17.737	USN1978
1975.908	85.57	17.750	USN1978
1975.909	85.65	17.748	USN1978
1975.925	85.61	17.745	USN1978
1975.925	85.62	17.743	USN1978
1975.928	85.58	17.757	USN1978
1975.928	85.59	17.747	USN1978
1976.154	85.65	17.754	USN1978
1976.154	85.67	17.749	USN1978
1976.165	85.948	17.784	PkO2003
1976.165	85.95	17.784	PkO1988
1976.166	85.647	17.748	PkO2014b
1976.167	85.659	17.749	PkO2016
1976.220	85.66	17.743	USN1978
1976.228	85.64	17.746	USN1978
1976.228	85.75	17.747	USN1978
1976.250	85.66	17.746	USN1978
1976.250	85.67	17.753	USN1978
1976.274	85.71	17.753	USN1978
1976.280	85.68	17.739	USN1978
1976.280	85.72	17.735	USN1978
1976.283	85.68	17.743	USN1978
1976.283	85.76	17.754	USN1978
1976.930	85.88	17.736	USN1978
1976.930	85.89	17.721	USN1978
1976.930	85.91	17.732	USN1978
1977.113	85.92	17.731	USN1978
1977.113	85.93	17.720	USN1978
1977.113	85.96	17.727	USN1978
1977.113	85.96	17.737	USN1978
1977.177	85.971	17.725	PkO2016
1977.183	86.21	17.753	PkO1988
1977.184	86.147	17.753	PkO2003
1977.190	85.985	17.723	PkO2014b
1977.203	85.95	17.729	USN1978
1977.203	86.02	17.729	USN1978
1977.230	85.96	17.730	USN1978
1977.230	85.98	17.725	USN1978
1977.230	85.99	17.730	USN1978
1977.230	86.00	17.716	USN1978
1977.230	86.02	17.707	USN1978
1977.233	85.95	17.721	USN1978
1977.233	86.01	17.729	USN1978
1977.233	86.02	17.731	USN1978
1977.285	85.99	17.718	USN1978
1977.285	85.99	17.732	USN1978
1977.956	86.13	17.700	USN1984
1977.956	86.17	17.716	USN1984
1977.956	86.19	17.710	USN1984

Relative astrometric position – continued.

Epoch (a)	θ (deg)	ρ (")	Ref. ^(a)
1977.956	86.19	17.711	USN1984
1977.956	86.20	17.702	USN1984
1978.080	86.244	17.707	PkO2016
1978.083	86.548	17.735	PkO2003
1978.087	86.51	17.728	PkO1988
1978.092	84.0	18.10	Scn2003
1978.096	86.234	17.707	PkO2014b
1978.109	86.24	17.711	USN1984
1978.109	86.26	17.709	USN1984
1978.112	86.25	17.703	USN1984
1978.112	86.25	17.710	USN1984
1978.175	86.28	17.703	USN1984
1979.191	86.586	17.686	PkO2016
1979.195	86.860	17.707	PkO2003
1979.195	86.86	17.707	PkO1988
1979.196	86.542	17.679	PkO2014b
1979.999	87.0	17.00	ClI2003
1980.135	87.127	17.692	PkO2003
1980.135	87.13	17.692	PkO1988
1980.165	86.877	17.673	PkO2016
1980.181	86.875	17.674	PkO2014b
1981.154	87.37	17.664	PkO1988
1981.154	87.375	17.664	PkO2003
1981.157	87.170	17.652	PkO2016
1981.166	87.223	17.653	PkO2014b
1981.252	87.21	17.659	USN1984
1981.252	87.22	17.657	USN1984
1982.054	87.626	17.673	PkO2003
1982.054	87.63	17.673	PkO1988
1982.056	87.428	17.631	PkO2016
1982.072	87.398	17.634	PkO2014b
1982.10	88.1	17.38	Stu1984
1982.29	88.	19.2	Wat1984
1983.149	87.744	17.618	PkO2014b
1983.156	87.94	17.643	PkO1988
1983.157	87.957	17.643	PkO2003
1983.20	87.75	17.605	Tor1985
1984.157	88.089	17.598	PkO2014b
1984.164	88.084	17.597	PkO2016
1984.177	94.5	17.29	Tob2005g
1984.269	87.2	17.71	Csa1985
1984.276	86.0	17.60	Doc1985b
1984.276	87.2	17.80	Lin1985b
1984.342	88.26	17.595	PkO2010
1985.072	88.369	17.583	PkO2014b
1985.096	88.336	17.585	PkO2016
1985.161	88.49	17.582	PkO2010
1986.02	89.3	17.033	WFD1985
1986.226	88.702	17.568	PkO2016
1986.230	88.702	17.564	PkO2014b
1986.232	88.78	17.564	PkO2010
1987.177	88.993	17.547	PkO2014b
1987.182	88.984	17.545	PkO2016
1987.184	89.07	17.546	PkO2010
1988.177	89.32	17.528	PkO2010
1988.179	89.318	17.534	PkO2016
1988.202	89.254	17.528	PkO2014b
1989.242	89.66	17.510	PkO2010
1989.242	89.601	17.505	PkO2014b
1989.244	89.632	17.509	PkO2016

Relative astrometric position – continued.

Epoch (a)	θ (deg)	ρ (")	Ref. ^(a)
1990.045	89.5	17.10	Tob2003
1990.260	89.939	17.499	PkO2016
1990.262	89.96	17.495	PkO2010
1990.262	89.919	17.490	PkO2014b
1990.33	89.0	17.69	Kzn1994
1991.25	90.1	17.47	HIP1997a
1991.81	90.5	17.476	TYC2000b
1992.170	90.617	17.440	PkO2014b
1992.178	90.66	17.449	PkO2010
1992.178	90.531	17.458	PkO2016
1992.45	91.2	17.12	Ary1994
1993.130	90.681	17.433	PkO2014b
1993.133	90.87	17.431	PkO2010
1993.138	90.804	17.442	PkO2016
1994.199	95.8	17.31	Tob2003
1994.214	91.121	17.434	PkO2014b
1994.233	91.16	17.431	PkO2010
1994.259	91.141	17.422	PkO2016
1995.022	91.37	17.33	ADP1998
1995.100	91.80	17.39	ADP1998
1995.229	91.8	17.58	Ctt2003b
1995.23	91.8	17.58	Ctt1995
1995.254	91.465	17.401	PkO2016
1995.254	91.479	17.398	PkO2014b
1995.256	91.51	17.398	PkO2010
1996.220	91.761	17.392	PkO2016
1996.223	91.737	17.396	PkO2014b
1996.240	91.77	17.390	PkO2010
1997.102	92.007	17.357	PkO2016
1997.107	92.01	17.340	PkO2010
1997.370	92.64	17.32	ADP1998
1997.370	92.31	17.36	ADP1998
1997.98	92.1	17.32	Ary2000
1998.88	92.4	17.27	Alz2003b
1998.93	92.3	17.05	TMA2003
1999.222	92.77	17.352	PkO2010
1999.235	92.696	17.333	PkO2016
2002.288	93.4	17.39	Dal2003a
2002.32	93.7	17.67	Bdn2003
2003.209	93.96	17.31	Los2005
2004.161	94.28	17.258	Izm2010
2004.306	94.31	17.261	PkO2010
2005.040	94.3	17.28	Arn2006d
2005.160	94.61	17.252	Izm2010
2005.315	94.5	17.12	Dal2006a
2006.146	94.93	17.227	Izm2010
2006.256	94.90	17.229	PkO2010
2007.128	95.1	17.28	Arn2007c
2007.19	95.4	17.18	Los2009b
2007.245	95.2	17.11	WSI2008
2008.0590	95.486	17.1913	Izm2015
2009.0710	95.760	17.1831	Izm2015
2009.282	95.4	17.20	Ctt2010
2009.926	95.8	16.9	Ant2010b
2010.0470	96.056	17.1679	Izm2015
2010.213	96.3	17.21	WSI2011b
2011.1310	96.402	17.1513	Izm2015
2012.0410	96.701	17.1349	Izm2015
2012.271	96.7	17.27	Dal2014
2012.3184	97.10	17.11	Cve2015

Relative astrometric position – continued.

Epoch (a)	θ (deg)	ρ (")	Ref. ^(a)
2012.954	96.4	17.2	CrC2014
2013.392	97.8	16.99	Ary2014
2014.2070	97.373	17.1103	Izm2015
2014.237	97.5	17.08	WSI2015
2015.278	97.2	17.22	Ctt2016
2015.283	97.73	17.184	CDa2016
2015.2937	97.80	17.09	Cve2017
2016.213	98.1	17.102	WSI2017b
2016.233	98.19	17.007	Loc2017a
2016.369	97.4	17.2	Wbt2017
2017.130	99.0	17.078	WSI2018a
2017.266	97.9	17.28	Ctt2018
2017.268	97.1	17.1	Wbt2018
2017.28	98.17	17.17	CC_2018a
2019.142	101.95	16.99	This work
2019.169	102.14	17.01	This work
2019.185	102.14	16.98	This work
2019.145	102.24	17.03	This work
2019.251	101.27	17.02	This work
2019.256	101.31	17.02	This work
2019.281	101.40	17.02	This work

Notes. ^(a) References listed in Table B.6.

Table B.1: GJ 338 B data of the CARMENES observations.

BJD (d)	RV _{vis} (m s ⁻¹)	dLW (m ² s ⁻²)	H α	FWHM (km s ⁻¹)	Contrast	BIS (km s ⁻¹)	IRT1	IRT2	IRT3	CRX (m s ⁻¹ dex ⁻¹)
2457395.64114	4.62 ± 1.27	-5.67 ± 1.73	0.7326 ± 0.0006	5.58 ± 0.01	24.49 ± 0.04	0.043 ± 0.003	0.581 ± 0.001	0.467 ± 0.001	0.45 ± 0.001	-13.17 ± 9.96
2457398.53654	-5.57 ± 2.07	-2.36 ± 1.8	0.7267 ± 0.0008	5.59 ± 0.01	24.54 ± 0.04	0.043 ± 0.003	0.579 ± 0.001	0.468 ± 0.001	0.448 ± 0.001	-18.66 ± 13.59
2457400.60073	0.76 ± 1.4	4.63 ± 1.48	0.7434 ± 0.0006	5.58 ± 0.01	24.38 ± 0.04	0.035 ± 0.003	0.589 ± 0.001	0.477 ± 0.001	0.456 ± 0.001	-2.77 ± 10.42
2457408.64378	-1.63 ± 3.46	2.49 ± 2.04	0.7474 ± 0.001	5.59 ± 0.01	24.34 ± 0.04	0.052 ± 0.005	0.599 ± 0.001	0.485 ± 0.001	0.466 ± 0.001	-20.92 ± 14.25
2457408.65378	0.9 ± 3.25	13.02 ± 1.7	0.7481 ± 0.0005	5.6 ± 0.01	24.35 ± 0.04	0.043 ± 0.003	0.597 ± 0.001	0.485 ± 0.001	0.465 ± 0.001	8.31 ± 7.33
2457410.52863	-3.46 ± 1.57	6.78 ± 1.36	0.7501 ± 0.0004	5.61 ± 0.01	24.38 ± 0.04	0.046 ± 0.002	0.593 ± 0.001	0.481 ± 0.001	0.461 ± 0.001	9.48 ± 8.76
2457410.53606	-1.92 ± 1.5	7.35 ± 1.5	0.7492 ± 0.0004	5.6 ± 0.01	24.34 ± 0.04	0.046 ± 0.002	0.593 ± 0.001	0.481 ± 0.001	0.461 ± 0.001	4.16 ± 8.07
2457413.51735	1.4 ± 2.18	-3.37 ± 1.91	0.7522 ± 0.0006	5.57 ± 0.01	24.45 ± 0.04	0.035 ± 0.003	0.592 ± 0.001	0.483 ± 0.001	0.462 ± 0.001	5.59 ± 8.99
2457414.4621	7.2 ± 1.82	-11.24 ± 3.08	0.7396 ± 0.0013	5.58 ± 0.01	24.58 ± 0.04	0.03 ± 0.003	0.586 ± 0.001	0.471 ± 0.002	0.457 ± 0.002	-27.27 ± 16.03
2457415.41656	11.8 ± 2.67	-13.69 ± 3.22	0.7422 ± 0.002	5.59 ± 0.01	24.65 ± 0.05	0.04 ± 0.011	0.588 ± 0.003	0.481 ± 0.003	0.452 ± 0.003	-30.27 ± 26.11
2457418.51102	4.86 ± 1.24	12.28 ± 2.09	0.7525 ± 0.0007	5.61 ± 0.01	24.45 ± 0.04	0.044 ± 0.004	0.596 ± 0.001	0.483 ± 0.001	0.466 ± 0.001	9.11 ± 10.24
2457422.49104	12.68 ± 1.38	25.44 ± 1.72	0.762 ± 0.0006	5.62 ± 0.01	24.41 ± 0.04	0.037 ± 0.003	0.601 ± 0.001	0.495 ± 0.001	0.473 ± 0.001	5.88 ± 8.38
2457423.36589	7.97 ± 1.43	23.68 ± 2.17	0.7686 ± 0.0008	5.57 ± 0.01	24.23 ± 0.04	0.037 ± 0.004	0.607 ± 0.001	0.498 ± 0.001	0.476 ± 0.001	5.93 ± 11.01
2457424.42513	8.34 ± 2.85	27.96 ± 2.24	0.7757 ± 0.0006	5.59 ± 0.01	24.13 ± 0.04	0.04 ± 0.003	0.615 ± 0.001	0.509 ± 0.001	0.482 ± 0.001	5.35 ± 11.56
2457425.35299	-0.88 ± 1.65	27.49 ± 2.11	0.7663 ± 0.0005	5.59 ± 0.01	24.14 ± 0.04	0.05 ± 0.003	0.606 ± 0.001	0.502 ± 0.001	0.477 ± 0.001	12.21 ± 9.69
2457426.41015	-2.32 ± 1.57	13.32 ± 2.13	0.7513 ± 0.0009	5.57 ± 0.01	24.13 ± 0.05	0.05 ± 0.005	0.598 ± 0.001	0.489 ± 0.001	0.467 ± 0.001	-19.23 ± 13.57
2457427.40809	-2.41 ± 1.23	15.16 ± 1.71	0.7446 ± 0.0006	5.61 ± 0.01	24.5 ± 0.04	0.045 ± 0.003	0.595 ± 0.001	0.486 ± 0.001	0.464 ± 0.001	-3.58 ± 10.27
2457430.47261	1.79 ± 1.8	-12.23 ± 2.16	0.7244 ± 0.001	5.58 ± 0.01	24.6 ± 0.04	0.043 ± 0.006	0.584 ± 0.001	0.469 ± 0.001	0.451 ± 0.001	12.8 ± 13.96
2457440.30135	12.52 ± 1.89	20.23 ± 2.35	0.7498 ± 0.0009	5.65 ± 0.01	24.81 ± 0.04	0.038 ± 0.005	0.598 ± 0.001	0.491 ± 0.002	0.468 ± 0.002	-33.19 ± 16.36
2457440.31597	11.29 ± 1.66	21.89 ± 1.89	0.747 ± 0.0006	5.55 ± 0.01	24.05 ± 0.04	0.037 ± 0.003	0.598 ± 0.001	0.489 ± 0.001	0.468 ± 0.001	-32.03 ± 12.56
2457444.47613	2.95 ± 1.41	15.66 ± 1.72	0.7483 ± 0.0007	5.6 ± 0.01	24.27 ± 0.04	0.044 ± 0.004	0.596 ± 0.001	0.481 ± 0.001	0.468 ± 0.001	-6.97 ± 9.11
2457441.47018	7.73 ± 2.07	44.76 ± 3.1	0.7369 ± 0.0034	5.5 ± 0.01	23.94 ± 0.05	0.025 ± 0.019	0.582 ± 0.003	0.469 ± 0.004	0.454 ± 0.003	-30.74 ± 10.84
2457442.41755	6.31 ± 1.12	7.32 ± 2.1	0.7608 ± 0.0007	5.68 ± 0.01	25.03 ± 0.04	0.038 ± 0.004	0.598 ± 0.001	0.489 ± 0.001	0.471 ± 0.001	-17.36 ± 42.28
2457488.39857	4.07 ± 1.47	13.22 ± 2.8	0.7664 ± 0.0012	5.61 ± 0.01	24.41 ± 0.04	0.045 ± 0.006	0.606 ± 0.001	0.501 ± 0.001	0.476 ± 0.001	12.61 ± 10.85
2457491.37557	6.78 ± 2.29	19.99 ± 1.61	0.7507 ± 0.0004	5.61 ± 0.01	24.46 ± 0.04	0.046 ± 0.002	0.601 ± 0.001	0.49 ± 0.001	0.471 ± 0.001	-3.56 ± 15.71
2457494.39418	-1.45 ± 1.31	11.03 ± 1.9	0.7329 ± 0.0007	5.6 ± 0.01	24.5 ± 0.04	0.044 ± 0.003	0.588 ± 0.001	0.476 ± 0.001	0.459 ± 0.001	25.22 ± 7.63
2457499.3582	2.12 ± 2.58	20.2 ± 1.81	0.7489 ± 0.0007	5.6 ± 0.01	24.5 ± 0.04	0.044 ± 0.003	0.587 ± 0.001	0.488 ± 0.001	0.469 ± 0.001	-4.4 ± 7.91
2457509.31697	-3.28 ± 1.48	3.18 ± 3.72	0.7505 ± 0.0014	5.58 ± 0.01	24.44 ± 0.04	0.036 ± 0.007	0.603 ± 0.002	0.492 ± 0.002	0.466 ± 0.002	44.23 ± 10.52
2457555.35907	6.34 ± 2.13	-21 ± 4.8	0.7441 ± 0.0012	5.58 ± 0.01	24.75 ± 0.04	0.039 ± 0.006	0.588 ± 0.002	0.474 ± 0.002	0.451 ± 0.002	35.17 ± 18.09
2457650.68397	3.06 ± 2.12	4.66 ± 19.98	0.7497 ± 0.0097	5.47 ± 0.03	23.02 ± 0.1	-0.137 ± 0.063	0.596 ± 0.011	0.508 ± 0.014	0.462 ± 0.013	27.78 ± 19.22
2457653.67217	3.05 ± 10.23	-37.01 ± 19.73	0.7809 ± 0.0114	5.55 ± 0.03	23.89 ± 0.09	0.02 ± 0.077	0.598 ± 0.013	0.512 ± 0.016	0.458 ± 0.016	-118.77 ± 105.4
2457657.69273	11.79 ± 13.14	12.31 ± 2.23	0.7678 ± 0.0001	5.59 ± 0.01	24.38 ± 0.04	0.033 ± 0.005	0.604 ± 0.001	0.49 ± 0.001	0.471 ± 0.001	249.25 ± 131.74
2457658.70763	1.21 ± 1.99	13.45 ± 3.71	0.7743 ± 0.0002	5.57 ± 0.01	23.96 ± 0.04	0.069 ± 0.01	0.61 ± 0.002	0.497 ± 0.002	0.478 ± 0.002	2.76 ± 16.97
2457673.6	-0.42 ± 2.04	14.93 ± 3.48	0.7714 ± 0.0019	5.59 ± 0.01	24.33 ± 0.04	0.015 ± 0.019	0.606 ± 0.002	0.499 ± 0.002	0.473 ± 0.002	9.92 ± 19.13
2457688.70235	10.17 ± 2.85	-2.58 ± 6.04	0.7656 ± 0.0035	5.54 ± 0.01	24.31 ± 0.05	0.015 ± 0.017	0.606 ± 0.002	0.485 ± 0.004	0.47 ± 0.004	40.75 ± 25.92
2457695.74187	1.37 ± 3.48	-18.27 ± 7.36	0.7578 ± 0.0034	5.61 ± 0.01	24.7 ± 0.05	0.043 ± 0.017	0.591 ± 0.004	0.487 ± 0.005	0.468 ± 0.005	62.29 ± 32.51
2457698.53295	-2.99 ± 4.91	18.27 ± 7.36	0.7578 ± 0.0034	5.61 ± 0.01	24.61 ± 0.04	0.035 ± 0.006	0.605 ± 0.001	0.495 ± 0.002	0.472 ± 0.001	11.44 ± 50.09
2457699.58911	5.7 ± 3.12	-5.13 ± 4.4	0.7489 ± 0.0023	5.58 ± 0.01	24.43 ± 0.04	0.022 ± 0.012	0.592 ± 0.003	0.48 ± 0.003	0.46 ± 0.003	2.75 ± 17.63
2457701.65253	1.97 ± 1.88	-2.11 ± 2.82	0.7511 ± 0.0011	5.61 ± 0.01	24.61 ± 0.04	0.041 ± 0.006	0.596 ± 0.001	0.483 ± 0.001	0.464 ± 0.001	13.22 ± 14.52
2457703.73957	5.08 ± 1.97	2.28 ± 2.87	0.7706 ± 0.0013	5.61 ± 0.01	24.61 ± 0.04	0.035 ± 0.006	0.605 ± 0.001	0.495 ± 0.002	0.472 ± 0.001	3.53 ± 12.95
2457704.53336	1.97 ± 1.39	15.48 ± 2	0.7734 ± 0.0012	5.55 ± 0.01	24.06 ± 0.04	0.023 ± 0.006	0.603 ± 0.001	0.493 ± 0.001	0.469 ± 0.001	34.96 ± 108.4
2457705.58152	6.18 ± 10.45	42.74 ± 13.14	0.7561 ± 0.0076	5.47 ± 0.02	23.43 ± 0.07	0.015 ± 0.046	0.595 ± 0.009	0.476 ± 0.011	0.458 ± 0.01	-15.65 ± 19.49
2457712.55149	6.39 ± 2.23	-6.46 ± 5.58	0.757 ± 0.0014	5.57 ± 0.01	24.5 ± 0.04	0.035 ± 0.007	0.595 ± 0.001	0.484 ± 0.002	0.463 ± 0.001	2.17 ± 13.77
2457752.72374	2.19 ± 1.75	13.91 ± 2.71	0.7647 ± 0.0011	5.63 ± 0.01	24.62 ± 0.04	0.054 ± 0.006	0.598 ± 0.001	0.494 ± 0.001	0.471 ± 0.001	-6.67 ± 12.59
2457754.38006	2.02 ± 1.4	19.46 ± 2.47	0.755 ± 0.0011	5.63 ± 0.01	24.7 ± 0.04	0.048 ± 0.006	0.596 ± 0.002	0.486 ± 0.002	0.47 ± 0.002	12.08 ± 15.46
2457756.61231	5.01 ± 1.74	18.93 ± 2.98	0.766 ± 0.0012	5.62 ± 0.01	24.54 ± 0.04	0.056 ± 0.006	0.602 ± 0.001	0.495 ± 0.001	0.473 ± 0.001	12.84 ± 14.19
2457757.53817	22.38 ± 12.63	-91.79 ± 19.03	0.7546 ± 0.0112	5.37 ± 0.03	24.12 ± 0.12	0.133 ± 0.08	0.613 ± 0.011	0.473 ± 0.014	0.463 ± 0.014	-218.99 ± 120.25
2457757.54142	2.35 ± 1.88	17.1 ± 2.35	0.763 ± 0.0012	5.62 ± 0.01	24.44 ± 0.04	0.045 ± 0.006	0.6 ± 0.001	0.491 ± 0.001	0.472 ± 0.001	-12.84 ± 14.19
2457759.68529	6.42 ± 2.07	14.17 ± 2.83	0.758 ± 0.0012	5.64 ± 0.01	24.67 ± 0.04	0.044 ± 0.006	0.593 ± 0.001	0.489 ± 0.001	0.464 ± 0.001	7.14 ± 15.37
2457760.43644	-1.57 ± 1.79	8.36 ± 2.48	0.7563 ± 0.0013	5.62 ± 0.01	24.61 ± 0.04	0.038 ± 0.007	0.591 ± 0.002	0.482 ± 0.002	0.463 ± 0.002	12.73 ± 16.2
2457761.41273	-1.51 ± 1.62	6.65 ± 2.37	0.7439 ± 0.0011	5.58 ± 0.01	24.5 ± 0.05	0.052 ± 0.006	0.591 ± 0.001	0.478 ± 0.001	0.457 ± 0.001	-0.06 ± 14.25
2457762.54871	-0.77 ± 2.06	-12.5 ± 3.15	0.7452 ± 0.0012	5.58 ± 0.01	24.7 ± 0.04	0.042 ± 0.006	0.588 ± 0.001	0.476 ± 0.001	0.457 ± 0.001	-15.39 ± 17.38
2457763.58385	1.36 ± 1.51	-18.22 ± 3.75	0.7475 ± 0.0013	5.58 ± 0.01	24.67 ± 0.04	0.04 ± 0.007	0.592 ± 0.001	0.477 ± 0.002	0.455 ± 0.001	3.39 ± 10.43
2457766.57912	7.87 ± 2	-3.64 ± 3.03	0.7517 ± 0.0011	5.62 ± 0.01	24.73 ± 0.04	0.031 ± 0.006	0.595 ± 0.001	0.48 ± 0.001	0.462 ± 0.001	-14.53 ± 16.45
2457768.59153	2.87 ± 2.16	8.66 ± 3.09	0.752 ± 0.0017	5.58 ± 0.01	24.37 ± 0.04	0.022 ± 0.009	0.597 ± 0.002	0.485 ± 0.002	0.467 ± 0.002	22.28 ± 18.91
2457771.64173	0.14 ± 2.58	13.68 ± 3.96	0.7635 ± 0.0021	5.62 ± 0.01	24.33 ± 0.04	0.039 ± 0.011	0.598 ± 0.002	0.489 ± 0.002	0.473 ± 0.002	16.11 ± 22.32
2457779.43116	0.85 ± 2.02	-5.05 ± 3.07	0.7402 ± 0.0016	5.58 ± 0.01	24.63 ± 0.04	0.032 ± 0.008	0.585 ± 0.002	0.476 ± 0.002	0.459 ± 0.002	-24.85 ± 18.02

Table B.1: continued.

BJD (d)	RV _{vis} (m s ⁻¹)	dLW (m ² s ⁻²)	H α	FWHM (km s ⁻¹)	Contrast	BIS (km s ⁻¹)	IRT1	IRT2	IRT3	CRX (m s ⁻¹ dex ⁻¹)
2457787.53524	8.41 ± 2.22	22.38 ± 2.57	0.7768 ± 0.0013	5.63 ± 0.01	24.48 ± 0.04	0.037 ± 0.007	0.613 ± 0.001	0.508 ± 0.002	0.484 ± 0.002	-23.95 ± 18.09
2457788.44445	7.53 ± 1.89	12.78 ± 3.54	0.7611 ± 0.0013	5.62 ± 0.01	24.57 ± 0.05	0.024 ± 0.007	0.6 ± 0.001	0.495 ± 0.002	0.474 ± 0.002	-4.42 ± 13.44
2457790.50252	-1.03 ± 1.74	14.56 ± 3.1	0.7631 ± 0.0016	5.6 ± 0.01	24.73 ± 0.05	0.041 ± 0.008	0.602 ± 0.002	0.492 ± 0.002	0.472 ± 0.002	-9.06 ± 13.85
2457791.32467	1.04 ± 1.75	12.89 ± 3.14	0.7898 ± 0.0013	0.613 ± 0.002	0.511 ± 0.002	0.49 ± 0.002	-1.59 ± 15.67
2457792.60552	-0.33 ± 2.34	12.35 ± 3.1	0.7558 ± 0.0014	0.601 ± 0.002	0.492 ± 0.002	0.47 ± 0.002	-10.11 ± 18.15
2457793.34201	-2.07 ± 2.42	7.7 ± 4.6	0.7626 ± 0.0021	5.61 ± 0.01	24.74 ± 0.05	0.065 ± 0.011	0.601 ± 0.002	0.49 ± 0.002	0.471 ± 0.002	-22.92 ± 21.18
2457794.64321	-4.66 ± 2.02	-0.62 ± 2.52	0.7488 ± 0.0012	5.57 ± 0.01	24.58 ± 0.04	0.051 ± 0.007	0.592 ± 0.001	0.483 ± 0.001	0.46 ± 0.001	-15.04 ± 15.21
2457798.43226	2.93 ± 2.25	-10.85 ± 3.17	0.7491 ± 0.0019	5.6 ± 0.01	24.61 ± 0.04	0.018 ± 0.01	0.592 ± 0.002	0.477 ± 0.002	0.461 ± 0.002	-5.23 ± 20.79
2457799.65634	-5.78 ± 2.02	-6.85 ± 2.95	0.7499 ± 0.0015	5.57 ± 0.01	24.58 ± 0.04	0.047 ± 0.008	0.588 ± 0.002	0.474 ± 0.002	0.456 ± 0.002	2.11 ± 17.85
2457800.35509	-6.99 ± 2.18	-1.74 ± 3.48	0.7491 ± 0.0015	5.6 ± 0.01	24.7 ± 0.05	0.053 ± 0.008	0.59 ± 0.001	0.476 ± 0.002	0.461 ± 0.002	-17.83 ± 20.71
2457802.65925	2.6 ± 1.94	8.46 ± 2.03	0.7449 ± 0.0011	5.57 ± 0.01	24.31 ± 0.04	0.034 ± 0.005	0.59 ± 0.001	0.482 ± 0.001	0.462 ± 0.001	-30.81 ± 14.04
2457806.46145	4.77 ± 2.07	3.45 ± 3.28	0.7578 ± 0.0014	5.6 ± 0.01	24.49 ± 0.04	0.031 ± 0.007	0.6 ± 0.002	0.491 ± 0.002	0.469 ± 0.002	-11.73 ± 18.49
2457808.60944	9.54 ± 5.59	-2.93 ± 7.49	0.7576 ± 0.0036	5.56 ± 0.01	24.28 ± 0.05	0.044 ± 0.02	0.605 ± 0.004	0.499 ± 0.005	0.468 ± 0.004	-21.45 ± 48.75
2457817.60324	-0.9 ± 2.18	-2.07 ± 2.39	0.7355 ± 0.0013	5.58 ± 0.01	24.5 ± 0.04	0.041 ± 0.007	0.588 ± 0.002	0.479 ± 0.002	0.459 ± 0.002	17.75 ± 17.96
2457818.56492	-0.44 ± 1.98	5.25 ± 1.79	0.7466 ± 0.0009	5.6 ± 0.01	24.57 ± 0.04	0.038 ± 0.005	0.597 ± 0.001	0.481 ± 0.001	0.462 ± 0.001	15.21 ± 12.31
2457819.32832	-0.77 ± 1.84	3.91 ± 2.09	0.7392 ± 0.0011	5.58 ± 0.01	24.44 ± 0.04	0.028 ± 0.005	0.591 ± 0.001	0.479 ± 0.001	0.458 ± 0.001	15.81 ± 13.66
2457821.35089	3.15 ± 1.92	6.51 ± 2.22	0.7736 ± 0.0012	0.605 ± 0.001	0.498 ± 0.001	0.474 ± 0.001	36.64 ± 14.61
2457822.6108	-0.49 ± 1.24	18.99 ± 2.09	0.7595 ± 0.0007	5.6 ± 0.01	24.35 ± 0.04	0.043 ± 0.004	0.602 ± 0.001	0.49 ± 0.001	0.469 ± 0.001	8.56 ± 9.11
2457824.38807	0.54 ± 1.47	16.26 ± 2.4	0.7522 ± 0.0008	5.63 ± 0.01	24.49 ± 0.04	0.044 ± 0.004	0.596 ± 0.001	0.486 ± 0.001	0.468 ± 0.001	9.68 ± 12.49
2457828.4336	-1.83 ± 3	-3.56 ± 4.52	0.7466 ± 0.0023	5.61 ± 0.01	24.5 ± 0.04	0.027 ± 0.012	0.587 ± 0.003	0.478 ± 0.003	0.462 ± 0.003	-14.48 ± 28.51
2457829.37008	0.64 ± 2.94	-3.69 ± 4.4	0.7544 ± 0.0022	5.59 ± 0.01	24.62 ± 0.04	0.011 ± 0.011	0.595 ± 0.002	0.482 ± 0.003	0.465 ± 0.003	-19.99 ± 27.73
2457833.45546	2.11 ± 1.55	6.57 ± 1.79	0.7338 ± 0.0007	5.6 ± 0.01	24.6 ± 0.04	0.038 ± 0.004	0.589 ± 0.001	0.473 ± 0.001	0.456 ± 0.001	5.53 ± 11.73
2457849.37225	0.54 ± 2.49	11.02 ± 2.3	0.7516 ± 0.0011	5.6 ± 0.01	24.46 ± 0.04	0.039 ± 0.006	0.596 ± 0.001	0.486 ± 0.001	0.466 ± 0.001	15.51 ± 14.25
2457855.33555	1.25 ± 4.79	11.95 ± 7.23	0.7584 ± 0.0036	5.54 ± 0.02	24.11 ± 0.06	0.056 ± 0.02	0.603 ± 0.004	0.487 ± 0.004	0.468 ± 0.004	-3.72 ± 45.72
2457855.3413	2.36 ± 1.66	15.35 ± 2.21	0.7524 ± 0.0011	5.61 ± 0.01	24.45 ± 0.04	0.051 ± 0.006	0.599 ± 0.001	0.489 ± 0.001	0.467 ± 0.001	12.09 ± 14.01
2457857.37404	4.5 ± 1.4	12.14 ± 2.46	0.7488 ± 0.0008	5.62 ± 0.01	24.62 ± 0.04	0.04 ± 0.004	0.596 ± 0.001	0.486 ± 0.001	0.465 ± 0.001	-6.53 ± 11.56
2457859.34327	0.41 ± 1.33	9.38 ± 1.88	0.7518 ± 0.0007	5.62 ± 0.01	24.55 ± 0.04	0.043 ± 0.004	0.597 ± 0.001	0.486 ± 0.001	0.467 ± 0.001	-0.45 ± 9.56
2457860.34447	1.74 ± 1.33	11.28 ± 1.84	0.7522 ± 0.0008	5.61 ± 0.01	24.53 ± 0.04	0.043 ± 0.004	0.597 ± 0.001	0.484 ± 0.001	0.466 ± 0.001	1.68 ± 9.52
2457862.36996	1.9 ± 1.79	-1.68 ± 2.94	0.7536 ± 0.0014	5.56 ± 0.01	24.49 ± 0.04	0.031 ± 0.007	0.6 ± 0.002	0.485 ± 0.002	0.465 ± 0.002	-9.24 ± 15.48
2457863.35221	1.27 ± 2.55	-2.53 ± 3.22	0.7541 ± 0.0013	5.59 ± 0.01	24.63 ± 0.04	0.032 ± 0.007	0.593 ± 0.002	0.479 ± 0.002	0.462 ± 0.002	-8.73 ± 14.81
2457864.42416	1.57 ± 1.74	3.22 ± 1.7	0.7518 ± 0.0008	5.59 ± 0.01	24.46 ± 0.04	0.045 ± 0.004	0.595 ± 0.001	0.483 ± 0.001	0.463 ± 0.001	18.77 ± 12.86
2457875.44711	2.47 ± 6.77	-0.33 ± 9.79	0.7545 ± 0.0043	5.55 ± 0.02	24.17 ± 0.06	0.087 ± 0.025	0.594 ± 0.005	0.496 ± 0.006	0.467 ± 0.006	11.03 ± 68.23
2457879.36369	3.56 ± 5.06	-22.58 ± 6.85	0.7602 ± 0.0035	5.53 ± 0.02	24.5 ± 0.05	0.039 ± 0.019	0.589 ± 0.004	0.484 ± 0.004	0.459 ± 0.004	-7.61 ± 47.99
2457883.45768	1.12 ± 2.07	0.65 ± 2.65	0.7491 ± 0.0015	5.56 ± 0.01	24.31 ± 0.04	0.027 ± 0.008	0.593 ± 0.002	0.482 ± 0.002	0.463 ± 0.002	-39.31 ± 16.65
2457886.33653	2.41 ± 2.48	-3.58 ± 3.31	0.7488 ± 0.0015	5.57 ± 0.01	24.54 ± 0.04	0.041 ± 0.008	0.592 ± 0.002	0.476 ± 0.002	0.461 ± 0.002	9.58 ± 17.59
2457889.45141	-2.74 ± 2.64	0.44 ± 3.27	0.7576 ± 0.0018	5.5 ± 0.01	24.05 ± 0.05	0.036 ± 0.009	0.595 ± 0.002	0.484 ± 0.002	0.463 ± 0.002	23.8 ± 23.96
2457896.40968	1.36 ± 1.92	-12.02 ± 2.35	0.7385 ± 0.0007	0.588 ± 0.001	0.473 ± 0.001	0.455 ± 0.001	8.72 ± 11.84
2458018.66037	0.3 ± 2.02	-12.4 ± 3.15	0.7484 ± 0.0008	5.58 ± 0.01	24.7 ± 0.03	0.027 ± 0.004	0.592 ± 0.001	0.473 ± 0.001	0.455 ± 0.001	16.2 ± 16.51
2458022.72147	-6.5 ± 1.69	-10.82 ± 4.49	0.7438 ± 0.0011	5.58 ± 0.01	24.64 ± 0.04	0.047 ± 0.006	0.589 ± 0.001	0.476 ± 0.001	0.457 ± 0.001	-5.1 ± 12.25
2458023.70754	-1.28 ± 2.44	-40.07 ± 7.13	0.7403 ± 0.0017	5.57 ± 0.01	24.81 ± 0.05	0.054 ± 0.009	0.587 ± 0.002	0.473 ± 0.002	0.455 ± 0.002	-16.07 ± 22.29
2458027.65052	-1.06 ± 1.94	-32.44 ± 3.57	0.7454 ± 0.0009	5.56 ± 0.01	24.76 ± 0.04	0.046 ± 0.004	0.576 ± 0.001	0.462 ± 0.001	0.445 ± 0.001	1.02 ± 18.58
2458028.70078	-3.43 ± 2.27	-38.3 ± 4.07	0.7673 ± 0.0001	5.59 ± 0.01	24.98 ± 0.03	0.045 ± 0.005	0.593 ± 0.001	0.482 ± 0.001	0.459 ± 0.001	-10.92 ± 20.59
2458060.60718	-3.57 ± 2.47	-15.05 ± 3.42	0.7526 ± 0.0016	5.56 ± 0.01	24.45 ± 0.03	0.023 ± 0.006	0.585 ± 0.002	0.474 ± 0.002	0.452 ± 0.002	-29.07 ± 22.97
2458064.59034	-12.82 ± 1.63	-20.81 ± 2.3	0.7399 ± 0.0011	5.55 ± 0.01	24.65 ± 0.04	0.037 ± 0.006	0.578 ± 0.001	0.46 ± 0.001	0.444 ± 0.001	3.82 ± 13.79
2458084.52062	3.64 ± 1.89	-5.65 ± 1.97	0.7407 ± 0.0009	5.56 ± 0.01	24.47 ± 0.04	0.036 ± 0.005	0.582 ± 0.001	0.465 ± 0.001	0.447 ± 0.001	-6.77 ± 11.22
2458085.61559	3.2 ± 3.74	-37.38 ± 11.24	0.7579 ± 0.0025	5.58 ± 0.01	24.74 ± 0.04	0.051 ± 0.013	0.591 ± 0.003	0.49 ± 0.003	0.468 ± 0.003	-5.67 ± 33.59
2458088.64082	-6.13 ± 3.85	-52.52 ± 16.05	0.7521 ± 0.0035	5.55 ± 0.01	24.89 ± 0.04	0.012 ± 0.019	0.602 ± 0.004	0.482 ± 0.004	0.466 ± 0.004	-27.77 ± 32.97
2458089.56637	-4.47 ± 4.5	-59.93 ± 13.88	0.7522 ± 0.0031	5.49 ± 0.01	24.85 ± 0.04	-0.003 ± 0.016	0.6 ± 0.003	0.482 ± 0.003	0.459 ± 0.003	5.17 ± 42.55
2458093.58096	-6.76 ± 1.42	-11.72 ± 2.1	0.7419 ± 0.0008	5.59 ± 0.01	24.67 ± 0.04	0.042 ± 0.004	0.583 ± 0.001	0.466 ± 0.001	0.445 ± 0.001	4.24 ± 11.3
2458094.64595	-3.52 ± 1.25	-21.09 ± 2.86	0.7422 ± 0.0008	5.6 ± 0.01	24.81 ± 0.04	0.034 ± 0.004	0.579 ± 0.001	0.466 ± 0.001	0.445 ± 0.001	-21.57 ± 6.95
2458095.55215	-5.89 ± 1.32	-27.81 ± 3.12	0.7415 ± 0.0009	5.57 ± 0.01	24.71 ± 0.04	0.041 ± 0.004	0.578 ± 0.001	0.465 ± 0.001	0.448 ± 0.001	-14.65 ± 9.74
2458097.63142	-7.68 ± 1.85	-26.25 ± 2.81	0.7434 ± 0.0008	5.59 ± 0.01	24.86 ± 0.04	0.04 ± 0.006	0.58 ± 0.001	0.464 ± 0.001	0.446 ± 0.001	-14.08 ± 15.37
2458109.48233	-8.08 ± 1.85	-31.69 ± 4.09	0.7331 ± 0.0011	5.59 ± 0.01	24.86 ± 0.04	0.04 ± 0.006	0.577 ± 0.001	0.463 ± 0.001	0.446 ± 0.001	-1.7 ± 11.85
2458110.42661	-6.56 ± 1.84	-30.95 ± 4.04	0.7413 ± 0.0001	5.56 ± 0.01	24.78 ± 0.03	0.039 ± 0.005	0.582 ± 0.001	0.464 ± 0.001	0.445 ± 0.001	-3.63 ± 16.17
2458114.40019	-15.13 ± 3.3	-70.87 ± 8.41	0.7286 ± 0.0023	5.53 ± 0.01	24.77 ± 0.04	0.062 ± 0.012	0.571 ± 0.003	0.458 ± 0.003	0.446 ± 0.003	8.72 ± 29.99
2458117.45171	-0.3 ± 1.86	-41.41 ± 4.43	0.7326 ± 0.0012	5.61 ± 0.01	25.19 ± 0.04	0.038 ± 0.006	0.572 ± 0.002	0.459 ± 0.002	0.444 ± 0.002	-15.05 ± 15.44
2458118.43947	-0.97 ± 1.58	-18.33 ± 2.82	0.743 ± 0.0008	5.6 ± 0.01	24.98 ± 0.03	0.025 ± 0.004	0.581 ± 0.001	0.465 ± 0.001	0.448 ± 0.001	-14.48 ± 11.02
2458119.59697	3.24 ± 2.59	-52.85 ± 8.05	0.7318 ± 0.0022	5.58 ± 0.01	24.93 ± 0.05	0.056 ± 0.011	0.581 ± 0.002	0.465 ± 0.002	0.445 ± 0.002	29.08 ± 21.43
2458120.44339	-0.19 ± 2.21	-42.94 ± 7	0.74 ± 0.0019	5.6 ± 0.01	25.14 ± 0.04	0.047 ± 0.01	0.577 ± 0.002	0.469 ± 0.002	0.455 ± 0.002	-30.99 ± 19.62
2458127.51188	10.56 ± 7.15	-94.62 ± 20.2	0.7282 ± 0.0061	5.53 ± 0.02	24.56 ± 0.06	0.09 ± 0.037	0.594 ± 0.006	0.474 ± 0.007	0.457 ± 0.007	-30.84 ± 69.25

Table B.1: continued.

BJD (d)	RV _{vis} (m s ⁻¹)	dLW (m ² s ⁻²)	H α	FWHM (km s ⁻¹)	Contrast	BIS (km s ⁻¹)	IRT1	IRT2	IRT3	CRX (m s ⁻¹ dex ⁻¹)
2458136.36257	-2.48 ± 1.61	-10.23 ± 2.44	0.7543 ± 0.0008	5.57 ± 0.01	24.67 ± 0.04	0.048 ± 0.004	0.589 ± 0.001	0.476 ± 0.001	0.456 ± 0.001	-49.3 ± 8.62
2458141.67195	-5.35 ± 1.29	-16.15 ± 2.26	0.7379 ± 0.0008	5.6 ± 0.01	24.8 ± 0.04	0.046 ± 0.004	0.58 ± 0.001	0.464 ± 0.001	0.447 ± 0.001	-6.17 ± 8.74
2458149.44415	3.91 ± 1.73	-31.16 ± 3.81	0.7303 ± 0.001	5.59 ± 0.01	24.97 ± 0.04	0.039 ± 0.005	0.577 ± 0.001	0.46 ± 0.001	0.443 ± 0.001	-14.25 ± 14.75
2458160.3298	-0.45 ± 5.1	-91.03 ± 13.54	0.7344 ± 0.0036	5.52 ± 0.02	25.14 ± 0.06	0.043 ± 0.019	0.58 ± 0.004	0.474 ± 0.005	0.447 ± 0.005	-24.93 ± 48.99
2458161.3676	-9.88 ± 1.58	-27.53 ± 2.75	0.7269 ± 0.0009	5.59 ± 0.01	24.86 ± 0.04	0.041 ± 0.005	0.575 ± 0.001	0.456 ± 0.001	0.441 ± 0.001	-15.21 ± 11.25
2458166.46121	-1.65 ± 1.62	-24.94 ± 3.47	0.7265 ± 0.0009	5.58 ± 0.01	24.82 ± 0.04	0.035 ± 0.005	0.577 ± 0.001	0.458 ± 0.001	0.443 ± 0.001	3.14 ± 11.95
2458167.31605	-0.6 ± 2.76	-30.95 ± 5.33	0.7424 ± 0.0013	5.6 ± 0.01	25.06 ± 0.04	0.044 ± 0.007	0.582 ± 0.002	0.47 ± 0.002	0.452 ± 0.002	-23.11 ± 17.57
2458169.61102	-2.31 ± 1.97	-13.12 ± 3.5	0.7387 ± 0.0009	5.58 ± 0.01	24.63 ± 0.04	0.043 ± 0.005	0.586 ± 0.001	0.473 ± 0.001	0.456 ± 0.001	-15.75 ± 14.99
2458171.47374	-0.66 ± 1.83	-37.5 ± 5.06	0.7408 ± 0.0016	5.57 ± 0.01	24.85 ± 0.05	0.025 ± 0.008	0.579 ± 0.002	0.469 ± 0.002	0.452 ± 0.002	-31.29 ± 11.97
2458172.3992	-4.15 ± 2.04	-26.27 ± 5.19	0.7357 ± 0.0013	5.59 ± 0.01	24.79 ± 0.04	0.04 ± 0.007	0.582 ± 0.001	0.465 ± 0.001	0.452 ± 0.001	5.75 ± 13.72
2458175.426	-2.86 ± 1.79	-8.45 ± 1.89	0.7438 ± 0.001	5.59 ± 0.01	24.67 ± 0.04	0.038 ± 0.005	0.582 ± 0.001	0.47 ± 0.001	0.45 ± 0.001	-43.54 ± 11.95
2458187.50148	-6.08 ± 3.68	-14.33 ± 5.01	0.7344 ± 0.0026	5.57 ± 0.01	24.58 ± 0.04	0.034 ± 0.014	0.583 ± 0.003	0.475 ± 0.003	0.457 ± 0.003	-36.2 ± 32.95
2458199.31335	4.75 ± 2.65	-30.97 ± 3.89	0.7273 ± 0.0018	5.55 ± 0.01	24.69 ± 0.04	0.027 ± 0.009	0.573 ± 0.002	0.454 ± 0.002	0.437 ± 0.002	-20.16 ± 24.52
2458206.39619	-3.98 ± 1.88	-5.14 ± 2.67	0.7299 ± 0.0014	5.59 ± 0.01	24.6 ± 0.04	0.037 ± 0.007	0.583 ± 0.002	0.47 ± 0.002	0.455 ± 0.002	-24.09 ± 14.92
2458207.38432	-6.87 ± 2.4	-12.43 ± 3.42	0.7346 ± 0.0016	5.58 ± 0.01	24.6 ± 0.04	0.042 ± 0.008	0.583 ± 0.002	0.468 ± 0.002	0.449 ± 0.002	-29.14 ± 16.69
2458209.46798	-7.27 ± 1.66	-7.46 ± 2.18	0.7284 ± 0.0011	5.59 ± 0.01	24.64 ± 0.04	0.039 ± 0.006	0.58 ± 0.001	0.468 ± 0.001	0.449 ± 0.001	-8.97 ± 13.64
2458211.41096	-12.24 ± 1.99	-18.3 ± 2.87	0.7261 ± 0.0014	5.56 ± 0.01	24.7 ± 0.04	0.046 ± 0.007	0.58 ± 0.002	0.462 ± 0.002	0.449 ± 0.002	-20.94 ± 16.25
2458212.40137	-9.26 ± 1.59	-42.83 ± 5.14	0.7273 ± 0.0013	5.56 ± 0.01	24.92 ± 0.04	0.04 ± 0.007	0.576 ± 0.001	0.459 ± 0.002	0.443 ± 0.002	9.45 ± 13.08
2458215.29963	-8.13 ± 1.68	-37.71 ± 3.28	0.7168 ± 0.0008	5.56 ± 0.01	24.89 ± 0.04	0.04 ± 0.004	0.566 ± 0.001	0.452 ± 0.001	0.436 ± 0.001	21.97 ± 13.06
2458217.36295	-3.54 ± 2.65	-53.02 ± 7.19	0.7251 ± 0.0016	5.55 ± 0.01	24.99 ± 0.04	0.022 ± 0.008	0.575 ± 0.002	0.458 ± 0.002	0.444 ± 0.002	-5.49 ± 22.34
2458218.36353	1.26 ± 4.03	-69.75 ± 11.86	0.7286 ± 0.0029	5.5 ± 0.01	24.88 ± 0.05	0.004 ± 0.015	0.586 ± 0.003	0.467 ± 0.003	0.452 ± 0.003	-26.75 ± 32.61
2458225.34812	1.56 ± 1.57	-8.43 ± 1.75	0.7466 ± 0.0007	5.61 ± 0.01	24.78 ± 0.04	0.044 ± 0.004	0.589 ± 0.001	0.474 ± 0.001	0.458 ± 0.001	-7.3 ± 10.2
2458229.45421	-10.41 ± 2.87	-28.95 ± 3.34	0.7268 ± 0.0013	5.53 ± 0.01	24.62 ± 0.04	0.037 ± 0.007	0.577 ± 0.001	0.462 ± 0.002	0.443 ± 0.001	3.59 ± 15.19
2458237.44478	-8.14 ± 2.61	-26.81 ± 4.05	0.7475 ± 0.0019	5.56 ± 0.01	24.65 ± 0.05	0.031 ± 0.01	0.579 ± 0.002	0.477 ± 0.002	0.455 ± 0.002	-26.6 ± 23.15
2458238.43669	-6.44 ± 2.09	-10.3 ± 2.67	0.7392 ± 0.0016	5.56 ± 0.01	24.46 ± 0.05	0.042 ± 0.009	0.584 ± 0.002	0.471 ± 0.002	0.451 ± 0.002	43.93 ± 16.87
2458245.41633	-5.46 ± 1.84	-7.3 ± 2.15	0.7226 ± 0.0008	5.56 ± 0.01	24.43 ± 0.04	0.038 ± 0.004	0.581 ± 0.001	0.462 ± 0.001	0.446 ± 0.001	-8.05 ± 12.31
2458246.46411	-7.16 ± 2.41	-13.33 ± 2.04	0.7167 ± 0.0007	5.52 ± 0.01	24.36 ± 0.04	0.044 ± 0.004	0.573 ± 0.001	0.455 ± 0.001	0.44 ± 0.001	-1.24 ± 11.71
2458249.48059	-1.26 ± 2	-36.78 ± 2.5	0.7136 ± 0.0008	5.5 ± 0.01	24.48 ± 0.04	0.035 ± 0.004	0.57 ± 0.001	0.449 ± 0.001	0.433 ± 0.001	-14.23 ± 13.67
2458258.37405	-7.73 ± 2.63	-6.91 ± 2.57	0.7394 ± 0.0011	5.56 ± 0.01	24.45 ± 0.05	0.034 ± 0.006	0.588 ± 0.001	0.471 ± 0.001	0.454 ± 0.001	-0.99 ± 12.76
2458260.36916	-7.71 ± 1.86	0.1 ± 1.96	0.7332 ± 0.0007	5.57 ± 0.01	24.5 ± 0.04	0.049 ± 0.004	0.585 ± 0.001	0.474 ± 0.001	0.455 ± 0.001	-5.06 ± 10.19
2458261.37184	-9.56 ± 1.64	-5.92 ± 2.29	0.7278 ± 0.0008	5.57 ± 0.01	24.51 ± 0.04	0.039 ± 0.004	0.582 ± 0.001	0.466 ± 0.001	0.448 ± 0.001	10.01 ± 11.13
2458262.34593	-13.09 ± 2.44	-17.42 ± 2.34	0.759 ± 0.001	5.55 ± 0.01	24.59 ± 0.05	0.043 ± 0.005	0.597 ± 0.001	0.485 ± 0.001	0.464 ± 0.001	22.28 ± 16.78
2458263.40211	-9.17 ± 2.26	-31.91 ± 2.57	0.7096 ± 0.0009	5.51 ± 0.01	24.56 ± 0.05	0.041 ± 0.005	0.574 ± 0.001	0.455 ± 0.001	0.438 ± 0.001	-4.38 ± 17.28
2458264.38102	-8.32 ± 1.74	-35.8 ± 2.46	0.7152 ± 0.0009	5.52 ± 0.01	24.52 ± 0.04	0.036 ± 0.005	0.564 ± 0.001	0.444 ± 0.001	0.432 ± 0.001	9.86 ± 12.99
2458266.42253	2.82 ± 2.97	-62.33 ± 6.64	0.7119 ± 0.0016	5.51 ± 0.01	24.7 ± 0.05	0.025 ± 0.008	0.567 ± 0.002	0.449 ± 0.002	0.434 ± 0.002	40.9 ± 20.56
2458273.39447	2.58 ± 1.85	-22.92 ± 3.27	0.739 ± 0.0009	5.54 ± 0.01	24.49 ± 0.04	0.04 ± 0.005	0.59 ± 0.001	0.474 ± 0.001	0.456 ± 0.001	22.7 ± 13.99
2458284.34913	-0.57 ± 2.16	-32.06 ± 3.24	0.7216 ± 0.0008	5.54 ± 0.01	24.53 ± 0.04	0.038 ± 0.004	0.579 ± 0.001	0.46 ± 0.001	0.446 ± 0.001	18.9 ± 14.66
2458291.36014	4.4 ± 2.29	-35.81 ± 5.3	0.7451 ± 0.0012	5.53 ± 0.01	24.36 ± 0.04	0.039 ± 0.006	0.587 ± 0.001	0.469 ± 0.001	0.451 ± 0.001	-6.8 ± 20.31
2458396.66991	-4.92 ± 1.92	5.52 ± 0.01	24.61 ± 0.04	0.038 ± 0.005
2458398.61908	-3.57 ± 2.49	5.5 ± 0.01	24.37 ± 0.04	0.038 ± 0.008

Table B.2: GJ 338 A data of the CARMENES observations.

BJD (d)	RV _{VIS} (m s ⁻¹)	dLW (m ² s ⁻²)	H α	IRT1	IRT2	IRT3
2457467.41671	-14.18 \pm 2.44	-13.71 \pm 4.53	0.7289 \pm 0.0018	0.594 \pm 0.002	0.476 \pm 0.002	0.461 \pm 0.002
2457499.37135	-17.72 \pm 2.12	-18.41 \pm 3.33	0.7309 \pm 0.0007	0.587 \pm 0.001	0.477 \pm 0.001	0.46 \pm 0.001
2457555.37464	-13.82 \pm 2.46	-9.81 \pm 4.14	0.7373 \pm 0.0011	0.598 \pm 0.001	0.493 \pm 0.001	0.467 \pm 0.001
2458105.57328	-2.32 \pm 1.64	0.45 \pm 2.19	0.7587 \pm 0.001	0.618 \pm 0.001	0.517 \pm 0.001	0.491 \pm 0.001
2458122.45286	-2.15 \pm 1.87	11.09 \pm 2.31	0.7512 \pm 0.0009	0.621 \pm 0.001	0.519 \pm 0.001	0.494 \pm 0.001
2458130.66117	-3.15 \pm 2.09	14.76 \pm 3	0.755 \pm 0.0013	0.621 \pm 0.001	0.519 \pm 0.002	0.496 \pm 0.001
2458131.49145	6.73 \pm 4.94	-38.47 \pm 12.38	0.7491 \pm 0.0038	0.62 \pm 0.004	0.52 \pm 0.005	0.493 \pm 0.005
2458132.39372	-7.3 \pm 2.73	-8.62 \pm 6.7	0.748 \pm 0.002	0.617 \pm 0.002	0.518 \pm 0.002	0.491 \pm 0.002
2458134.41383	-3.18 \pm 1.83	5.44 \pm 2.63	0.749 \pm 0.0011	0.613 \pm 0.001	0.511 \pm 0.001	0.487 \pm 0.001
2458160.33274	3.48 \pm 2.95	-9.93 \pm 5.51	0.7489 \pm 0.002	0.614 \pm 0.002	0.507 \pm 0.002	0.491 \pm 0.002
2458161.37036	1.42 \pm 2.04	14.51 \pm 2.21	0.7579 \pm 0.0009	0.618 \pm 0.001	0.519 \pm 0.001	0.493 \pm 0.001
2458167.31995	-6.74 \pm 2.63	18.79 \pm 3.98	0.8101 \pm 0.0017	0.641 \pm 0.002	0.546 \pm 0.002	0.521 \pm 0.002
2458187.50542	-2.61 \pm 4.38	-25.79 \pm 10.62	0.7302 \pm 0.0032	0.616 \pm 0.004	0.519 \pm 0.004	0.493 \pm 0.004
2458199.3235	9.83 \pm 5.27	-34.02 \pm 15.82	0.7429 \pm 0.0044	0.627 \pm 0.005	0.522 \pm 0.005	0.492 \pm 0.005
2458206.40375	-0.85 \pm 2.31	-6.45 \pm 5.94	0.7423 \pm 0.0017	0.613 \pm 0.002	0.516 \pm 0.002	0.488 \pm 0.002
2458211.41433	2.37 \pm 2.29	2.2 \pm 3.52	0.7486 \pm 0.0012	0.623 \pm 0.001	0.522 \pm 0.001	0.492 \pm 0.001
2458217.36617	-3.05 \pm 2.79	-24.47 \pm 7.23	0.747 \pm 0.002	0.613 \pm 0.002	0.511 \pm 0.002	0.492 \pm 0.002
2458229.47009	-6.19 \pm 3.73	-10.69 \pm 4.77	0.7468 \pm 0.0016	0.621 \pm 0.002	0.523 \pm 0.002	0.493 \pm 0.002
2458237.44785	0.63 \pm 2.56	-28.21 \pm 7.65	0.7352 \pm 0.0022	0.605 \pm 0.003	0.503 \pm 0.003	0.481 \pm 0.003
2458258.37791	9.49 \pm 3.73	-7.65 \pm 5.47	0.7562 \pm 0.0017	0.627 \pm 0.002	0.526 \pm 0.002	0.499 \pm 0.002
2458263.43879	-7.98 \pm 2.52	9.22 \pm 4.9	0.7435 \pm 0.0015	0.622 \pm 0.002	0.519 \pm 0.002	0.495 \pm 0.002
2458265.44641	-6.09 \pm 2.13	-9.72 \pm 2.62	0.72 \pm 0.0009	0.608 \pm 0.001	0.5 \pm 0.001	0.479 \pm 0.001
2458270.3468	-3.35 \pm 2.59	-9.29 \pm 2.91	0.709 \pm 0.0009	0.604 \pm 0.001	0.493 \pm 0.001	0.472 \pm 0.001
2458271.43819	-0.73 \pm 2.23	-23.29 \pm 4.3	0.7152 \pm 0.0015	0.604 \pm 0.002	0.498 \pm 0.002	0.478 \pm 0.002
2458290.36545	3.55 \pm 2.18	-11.48 \pm 2.51	0.7364 \pm 0.0011	0.615 \pm 0.001	0.51 \pm 0.001	0.488 \pm 0.001
2458299.35326	-4.37 \pm 2.71	5.27 \pm 2.28	0.7506 \pm 0.0009	0.619 \pm 0.001	0.517 \pm 0.001	0.488 \pm 0.001
2458383.70118	-0.12 \pm 1.77	0.14 \pm 2.74	0.7529 \pm 0.001	0.614 \pm 0.001	0.508 \pm 0.001	0.488 \pm 0.001
2458385.68178	1.71 \pm 2.13	-8.38 \pm 2.97	0.7908 \pm 0.0011	0.634 \pm 0.001	0.533 \pm 0.001	0.506 \pm 0.001
2458386.6702	0.54 \pm 2.07	11.16 \pm 4.57	0.7471 \pm 0.0011	0.607 \pm 0.001	0.504 \pm 0.001	0.481 \pm 0.001
2458391.66718	3.1 \pm 1.64	-2.81 \pm 2.85	0.7559 \pm 0.0012	0.613 \pm 0.001	0.509 \pm 0.001	0.487 \pm 0.001
2458392.69892	-4.12 \pm 5.96	-34.35 \pm 16.29	0.7871 \pm 0.0042	0.647 \pm 0.004	0.548 \pm 0.005	0.519 \pm 0.005
2458393.7162	8.73 \pm 1.82	5.06 \pm 2.33	0.7559 \pm 0.0012	0.616 \pm 0.001	0.516 \pm 0.001	0.491 \pm 0.001
2458399.66111	-3.02 \pm 1.95	7.11 \pm 3.2	0.7719 \pm 0.0014	0.622 \pm 0.002	0.524 \pm 0.002	0.499 \pm 0.002
2458410.64141	17.46 \pm 9.25	-69.46 \pm 29.68	0.7664 \pm 0.0079	0.641 \pm 0.008	0.548 \pm 0.009	0.53 \pm 0.009
2458413.69185	1.43 \pm 1.66	6.82 \pm 2.2	0.7522 \pm 0.001	0.612 \pm 0.001	0.51 \pm 0.001	0.489 \pm 0.001
2458417.56694	11.44 \pm 2.85	-10.25 \pm 6.97	0.7406 \pm 0.002	0.615 \pm 0.003	0.51 \pm 0.003	0.492 \pm 0.003
2458424.69799	-0.4 \pm 2.95	-2.72 \pm 3.12	0.7531 \pm 0.0015	0.618 \pm 0.002	0.518 \pm 0.002	0.494 \pm 0.002
2458430.70433	-2.3 \pm 4.73	-11.28 \pm 3.21	0.7404 \pm 0.0014	0.606 \pm 0.002	0.503 \pm 0.002	0.484 \pm 0.002
2458433.57916	-2.24 \pm 2.27	-5.86 \pm 3.13	0.7477 \pm 0.0012	0.611 \pm 0.001	0.508 \pm 0.002	0.484 \pm 0.001
2458451.51207	5.59 \pm 1.72	-0.9 \pm 2.71	0.7318 \pm 0.0011	0.605 \pm 0.001	0.502 \pm 0.001	0.483 \pm 0.001
2458454.63129	5.01 \pm 1.99	7.65 \pm 2.63	0.7576 \pm 0.0013	0.625 \pm 0.002	0.52 \pm 0.002	0.493 \pm 0.002
2458455.75059	0.55 \pm 2.22	21.23 \pm 2.69	0.7412 \pm 0.0011	0.613 \pm 0.001	0.513 \pm 0.001	0.486 \pm 0.001
2458458.68105	-1.17 \pm 2.14	9.35 \pm 2.59	0.7594 \pm 0.0009	0.619 \pm 0.001	0.518 \pm 0.001	0.493 \pm 0.001
2458462.69461	2.06 \pm 2.9	-11.24 \pm 2.45	0.7363 \pm 0.0011	0.599 \pm 0.001	0.493 \pm 0.001	0.474 \pm 0.001
2458466.75157	1.13 \pm 4.09	-63.3 \pm 9.5	0.7354 \pm 0.0035	0.607 \pm 0.004	0.495 \pm 0.004	0.478 \pm 0.004
2458469.61903	16.69 \pm 2.18	13.52 \pm 2.64	0.7539 \pm 0.0012	0.62 \pm 0.001	0.516 \pm 0.001	0.494 \pm 0.001
2458470.62218	15.66 \pm 2.02	15.08 \pm 2.93	0.7526 \pm 0.0012	0.618 \pm 0.001	0.52 \pm 0.001	0.496 \pm 0.001
2458471.74714	10.16 \pm 2.32	10.11 \pm 4.02	0.7812 \pm 0.0016	0.631 \pm 0.002	0.54 \pm 0.002	0.51 \pm 0.002
2458472.47359	3.76 \pm 4.5	36.17 \pm 4.51	0.7771 \pm 0.0016	0.631 \pm 0.002	0.532 \pm 0.002	0.506 \pm 0.002
2458473.41844	3.67 \pm 1.9	30.68 \pm 2.76	0.7743 \pm 0.001	0.634 \pm 0.001	0.536 \pm 0.001	0.508 \pm 0.001
2458475.44393	1.87 \pm 2.5	1.41 \pm 4.25	0.7742 \pm 0.0016	0.628 \pm 0.002	0.531 \pm 0.002	0.505 \pm 0.002
2458476.40495	-6.35 \pm 2.03	9.55 \pm 2.64	0.7721 \pm 0.0011	0.625 \pm 0.002	0.527 \pm 0.002	0.5 \pm 0.002
2458477.43512	-3.86 \pm 2.38	-16.8 \pm 3.94	0.7517 \pm 0.0017	0.616 \pm 0.002	0.511 \pm 0.002	0.489 \pm 0.002
2458478.46223	0.94 \pm 10.14	-74.72 \pm 25.34	0.7589 \pm 0.0085	0.631 \pm 0.009	0.519 \pm 0.011	0.519 \pm 0.011
2458479.71038	-1.11 \pm 1.69	-11.57 \pm 2.89	0.751 \pm 0.0011	0.605 \pm 0.001	0.502 \pm 0.001	0.48 \pm 0.001
2458480.73582	-7.78 \pm 1.76	-22.81 \pm 2.5	0.7349 \pm 0.0011	0.599 \pm 0.001	0.493 \pm 0.001	0.472 \pm 0.001
2458485.76598	10.35 \pm 1.63	18.75 \pm 1.81	0.7459 \pm 0.0009	0.613 \pm 0.001	0.511 \pm 0.001	0.486 \pm 0.001
2458486.40406	15.25 \pm 1.99	14.27 \pm 3.2	0.7624 \pm 0.0013	0.621 \pm 0.001	0.523 \pm 0.001	0.498 \pm 0.001

Table B.2: continued.

BJD (d)	RV_{VIS} (m s^{-1})	dLW ($\text{m}^2 \text{s}^{-2}$)	$H\alpha$	IRT1	IRT2	IRT3
2458487.42853	12.78 ± 1.76	21.95 ± 2.64	0.7701 ± 0.0013	0.628 ± 0.002	0.533 ± 0.002	0.504 ± 0.002
2458488.62244	6.74 ± 1.65	38.45 ± 2.68	0.7734 ± 0.0011	0.635 ± 0.001	0.539 ± 0.001	0.511 ± 0.001
2458490.62585	-1.09 ± 1.8	33.25 ± 3.12	0.7631 ± 0.0011	0.628 ± 0.001	0.529 ± 0.001	0.503 ± 0.001
2458491.77262	-1.94 ± 1.72	18.38 ± 3.44	0.7794 ± 0.0013	0.631 ± 0.001	0.533 ± 0.001	0.508 ± 0.001
2458492.51898	-4.43 ± 1.59	18.25 ± 2.55	0.763 ± 0.0012	0.626 ± 0.001	0.525 ± 0.001	0.498 ± 0.001
2458494.36674	-3.03 ± 3.51	-28.83 ± 7.76	0.7489 ± 0.0029	0.615 ± 0.003	0.511 ± 0.004	0.486 ± 0.003
2458495.76284	-8.11 ± 1.81	-1.02 ± 1.92	0.7399 ± 0.0009	0.61 ± 0.001	0.503 ± 0.001	0.476 ± 0.001
2458496.46758	-9.01 ± 1.59	-8.45 ± 2.25	0.7503 ± 0.001	0.609 ± 0.001	0.506 ± 0.001	0.483 ± 0.001
2458497.54743	-7.16 ± 1.76	-16.11 ± 2.34	0.7259 ± 0.0011	0.598 ± 0.001	0.492 ± 0.001	0.472 ± 0.001
2458498.63931	-6.82 ± 1.69	-17.99 ± 2.18	0.7299 ± 0.001	0.6 ± 0.001	0.491 ± 0.001	0.466 ± 0.001
2458499.36021	-2.89 ± 2.19	-24.21 ± 3.3	0.7295 ± 0.0013	0.598 ± 0.001	0.491 ± 0.002	0.471 ± 0.002
2458500.35752	7.77 ± 2.31	-41.64 ± 6.63	0.7374 ± 0.0022	0.605 ± 0.003	0.497 ± 0.003	0.481 ± 0.003

Table B.3: Relative and differential RVs for binary GJ338AB computed with common template (GJ338B).

BJD (d)	RV _A (m s ⁻¹)	eRV _A (m s ⁻¹)	RV _B (m s ⁻¹)	eRV _B (m s ⁻¹)	RV _A – RV _B (m s ⁻¹)	e(RV _A – RV _B) (m s ⁻¹)
CARMENES						
2457467.40826	-1232.61	2.55	6.42	4.62	-1239.02	5.27
2457499.35162	-1236.58	1.97	-5.43	2.44	-1231.14	3.13
2457555.35128	-1232.69	2.51	-0.60	2.27	-1232.08	3.38
2458118.80384	-1221.67	1.71	-5.11	2.43	-1216.55	2.97
2458125.93723	-1222.39	2.08	2.074	7.08	-1224.46	7.38
2458135.3882	-1222.66	1.72	-9.93	1.55	-1212.72	2.31
2458160.32833	-1215.61	2.93	-6.86	5.27	-1208.74	6.03
2458161.36622	-1217.42	2.08	-17.08	1.57	-1200.33	2.60
2458167.55802	-1226.01	2.67	-7.81	2.09	-1218.19	3.39
2458187.49951	-1220.96	3.95	-12.57	3.74	-1208.38	5.44
2458199.30827	-1208.06	5.21	-2.60	2.64	-1205.45	5.84
2458206.647	-1219.65	2.41	-11.77	2.11	-1207.87	3.20
2458210.93298	-1215.81	2.36	-16.31	1.78	-1199.49	2.95
2458216.82996	-1220.65	2.83	-9.33	2.82	-1211.31	3.99
2458229.44627	-1224.36	3.62	-16.74	2.82	-1207.61	4.59
2458237.69429	-1218.07	2.41	-13.99	2.37	-1204.07	3.38
2458258.87475	-1209.4	3.68	-14.03	2.11	-1195.36	4.24
2458262.59343	-1225.97	2.62	-16.88	2.04	-1209.08	3.32
2458265.93446	-1224.45	2.13	-4.49	2.96	-1219.95	3.65
2458272.41633	-1218.37	2.22	-4.13	1.88	-1214.23	2.91
2458290.86279	-1214.41	2.22	-1.76	2.30	-1212.64	3.19
2458395.19305	-1209.79	1.81	-11.71	1.82	-1198.07	2.57
2458398.09806	-1222.02	1.95	-10.84	2.39	-1211.17	3.08
ELODIE						
2449730.591135	-1227.38	4.87	134.89	5.12	-1362.27	7.07
2449751.337934	-1213.74	4.85	154.00	2.85	-1367.75	5.63
2450065.654345	-1264.87	7.85	93.93	4.04	-1358.81	8.83
2450070.679460	-1251.67	5.60	96.36	3.50	-1348.03	6.61
2450476.5263	-1235.34	4.17	176.57	4.48	-1411.92	6.12
SOPHIE						
2455478.65251	-1603.90	2.76	-336.81	2.94	-1267.09	4.03
2457021.51007	-1232.64	1.37	9.48	1.29	-1242.12	1.89
2457033.51057	-1240.13	2.43	3.32	3.12	-1243.46	3.96
2457035.51045	-1560.82	1.21	-309.59	1.03	-1251.22	1.59
2457037.51058	-1476.14	1.10	-222.25	0.87	-1253.88	1.40
2457046.51065	-1497.41	1.33	-242.06	1.18	-1255.35	1.78
2457047.51170	-1292.69	2.98	-29.39	1.57	-1263.29	3.38
2457052.51264	-1224.14	1.34	37.43	1.05	-1261.58	1.70
2457054.51039	-1202.83	1.39	59.57	1.09	-1262.41	1.77
2457062.51009	-1254.04	1.46	-0.14	1.13	-1253.90	1.85
2457064.51043	-1262.99	1.35	-1.92	0.88	-1261.06	1.61
2457065.51102	-1202.60	1.21	60.14	1.45	-1262.75	1.89
2457070.51020	-1333.29	1.26	-57.53	1.15	-1275.76	1.71
2457071.51021	-1268.89	1.11	-7.31	1.09	-1261.58	1.55
2457078.51005	-1416.19	1.53	-201.24	1.36	-1214.95	2.05
2457080.50964	-1316.16	1.31	-60.82	1.28	-1255.34	1.84
2457081.50966	-1346.63	1.12	-92.05	0.97	-1254.58	1.48
2457083.51003	-1335.56	1.34	-67.24	1.08	-1268.32	1.72
2457086.50899	-1388.19	1.36	-138.77	1.70	-1249.42	2.19
2457087.50743	-1323.69	1.24	-79.89	1.91	-1243.80	2.28
2457088.51516	-1309.00	0.97	-30.22	0.83	-1278.78	1.28
2457090.50935	-1473.75	1.31	-220.37	1.08	-1253.37	1.70
2457092.009	-1273.91	1.25	-329.23	1.27	-944.684	1.79
2457093.50906	-1304.81	1.14	-58.11	1.00	-1246.69	1.52
2457094.50896	-1336.30	1.18	-83.20	1.06	-1253.10	1.59
2457107.50820	-1521.29	1.60	-301.29	1.39	-1220.00	2.12

Table B.3: continued.

BJD (d)	RV_A (m s^{-1})	eRV_A (m s^{-1})	RV_B (m s^{-1})	eRV_B (m s^{-1})	$RV_A - RV_B$ (m s^{-1})	$e(RV_A - RV_B)$ (m s^{-1})
2457111.50672	-1635.85	1.58	-394.89	3.19	-1240.96	3.56
2457112.508	-1484.62	1.44	-232.56	1.13	-1252.05	1.83
2457114.50941	-1650.43	1.83	-382.52	1.69	-1267.90	2.50
2457120.50713	-1470.88	1.66	-219.69	1.14	-1251.18	2.02
2457121.50694	-1653.90	1.27	-395.16	1.24	-1258.73	1.78
2457124.50697	-1653.50	1.27	-399.61	0.97	-1253.89	1.61
2457407.01064	-1508.29	1.77	-48.40	1.23	-1459.88	2.15
2458073.50663	-1354.98	1.23	-173.43	1.24	-1181.54	1.75
2458140.01064	-1374.14	1.65	-93.65	1.26	-1280.48	2.08
2454435.70113	-1606.36	1.87	-316.09	1.62	-1290.26	2.47
2454463.60953	-1595.02	1.99	-304.10	1.91	-1290.92	2.76
2454952.35061	-1619.89	2.76	-341.21	3.04	-1278.68	4.11
2455327.34803	-1605.61	2.24	-342.29	2.11	-1263.31	3.08

Table B.6: References in Table B.5.

Ref.	
AbH1973	Abt, H.A. & Levy, S.G. AJ 78, 1093, 1973
Abt1922	Abetti, G. Pub. R. Obs. Astrof. Arcetri 39, 3, 1922
ADP1998	Abad, C., Docobo, J., & Della Prugna, F. A& AS 133, 71, 1998
Alz2003b	Alzner, A. Webb Soc., Double Star Circ. 11, 2003
Ant2010b	Anton, R. JDSO 6, 180, 2010
Arn2006d	Arnold, D. JDSO 2, 61, 2006
Arn2007c	Arnold, D. JDSO 3, 119, 2007
Ary1977	Argyle, R.W. Webb Society Handbook 1, 1977
Ary1994	Argyle, R.W. Webb Soc., Double Star Circ. 6, 1994
Ary2000	Argyle, R.W. Webb Soc., Double Star Circ. 8, 2000
Ary2014	Argyle, R.W. Webb Soc., Double Star Circ. 22, 2014
B__1923a	van den Bos, W.H. BAN 1, 193, 1923
B__1962d	van den Bos, W.H. AJ 67, 555, 1962
Baz1927b	Baize, P. J. Obs. 10, 164, 1927
Baz1932b	Baize, P. & Igounet, Mme. J. Obs. 15, 145, 1932
Baz1936b	Baize, P. J. Obs. 19, 85-92, 1936
Baz1945a	Baize, P. J. Obs. 28, 1-16, 1945
Baz1967	Baize, P. J. Obs. 50, 7-32, 1967
Bdn2003	Bodin, S. Double Star Observer 9, # 2, 16, 2003
Bea1930	Beal, W.O. Pub. Univ. Minnesota Obs. 1, 93, 1930 (includes all data from Beal, AJ 31, 136, 1918 and Beal, AJ 35, 53, 1923)
Big1883	Bigourdan, G. Ann. Obs. Paris - Observations H1, 1883
Bll1913A	Ball, R.S. cited by Burnham (Carnegie Inst. Wash. # 168, 1913)
Blo1931	Bloch, M. Bull. Obs. Lyon 13, 151, 1931
Bon1938	Bonnet, R. Paris Obs. 1938
Bot1962	Boettger, G. Potsdam Obs. 30, Pt.4, 1962
Bry1899	Bryant, W.W. Greenwich Observations, 1899 (data also in MNRAS 60, 494, 1900)
Bry1901	Bryant, W.W. Greenwich Observations, 1901 (data also in MNRAS 62, 383, 1902)
Bry1921	Bryant, W.W. Greenwich Cat. of Double Stars, 1921
Bu_1894	Burnham, S.W. Pub. Lick Obs. 2, 3, 1894 (includes most data from Bu_1893a)

Table B.6: Continued.

Ref.	
Bu_1906	Burnham, S.W. Burnham Double Star Cat., 1906
Bu_1913	Burnham, S.W. Carnegie Inst. Wash. # 168, 1913
CC_2018a	Bush, M., Del Medico, E., Boulais, H., Lang, C., Segura, J.E., D'Ambrosia, S., Rowe, D., Freed, R. & Genet, R. JDSO 14, 38, 2018
CDa2016	Chiron, D. El Observador de Estrellas Dobles, Issue # 17, 28, 2016
Cha1925b	Chandon, E. J. Obs. 8, 159, 1925
Chg1972	Chang, K. AJ 77, 759, 1972
Chm2004	Chanamé, J. & Gould, A. ApJ 601, 289, 2004
CIA2012f	Boyajian, T.S., von Braun, K., van Belle, G., McAlister, H.A., ten Brummelaar, T.A., Kane, S.R., Muirhead, P.S., Jones, J., White, R., Schaefer, G., Ciardi, D., Henry, T., Lopez-Morales, M., Ridgway, S., Gies, D., Jao, W.-C., Rojas-Ayala, B., Parks, J.R., Sturmann, L., Sturmann, J., Turner, N.H., Farrington, C., Goldfinger, P.J. & Berger, D.H. ApJ 757, 112, 2012
CII2003	Comellas, J.L. OAG Gen. Cat. 10,000 vis. double star meas. 1970-2003 (J2000.0), 2003
Com1929	Comstock, G.C. Pub. Washburn Obs. 14, Pt.1, 1929
CrC2014	Cortés-Contreras, M., Caballero, J.A. & Montes, D. Observatory 134, 348, 2014
Csa1985	Costa, J.M. (Docobo et al.) Bol. Acad. Galega de Ciencias 2, 131, 1985
Ctt1995	Courtot, J.-F. Obs. et Travaux 44, 22, 1995
Ctt2003b	Courtot, J.-F. OAG Gen. Cat. 10,000 vis. double star meas. 1970-2003 (J2000.0), 2003
Ctt2010	Courtot, J.-F. Webb Soc., Double Star Circ. 18, 5, 2010
Ctt2016	Courtot, J.-F. Webb Soc., Double Star Circ. 24, 6, 2016
Ctt2018	Courtot, J.-F. Webb Soc., Double Star Circ. 26, 9, 2018
Cve2015	Cvetkovic, Z., Pavlovic, R. & Boeva, S. AJ 149, 150, 2015
Cve2017	Cvetkovic, Z., Pavlovic, R. & Boeva, S. AJ 153, 195, 2017
D__1883	Dembowski, H. Misure micrometriche di Stelle Doppie e multiple... (Rome, Coi tipi del Salviucci), vol. I, 1883
D__1884	Dembowski, H. Misure micrometriche di Stelle Doppie e multiple... (Rome, Coi tipi del Salviucci), vol. II, 1884
Da_1867	Dawes, W.R. MemRAS 35, 137, 1867
Dal2003a	Daley, J.A. Pub. Ludwig Schupmann Obs, # 2, 2003
Dal2006a	Daley, J.A. JDSO 2, 44, 2006
Dal2014	Daley, J. JDSO 10, 136, 2014
Dam2014	Damm, F. private communication, 2014
DeO1954	Dearborn Obs.

Table B.6: Continued.

Ref.	
	Ann. Dearborn Obs. 7, 1, 1954
Dob1927	Doberck, W.A.
	Erganz. AN 7, 1927
Dob1934	Doberck, W.A.
	AN 254, 41, 1934
Doc1985b	Docobo, J.A. (Docobo et al.)
	Bol. Acad. Galega de Ciencias 2, 131, 1985
Du_1876	Duner, N.C.
	Mesures Microm. Lund, 1876
Dur1944b	Duruy, M.V.
	J. Obs. 27, 50, 1944
Eng1885a	Engelmann, R.
	AN 111, 337, 1885
Enh1895	Engelhardt, B. von
	Dresden, Engelhardt Sternwarte, part 3, 1895
Es_1902b	Espin, T.E.
	MNRAS 62, 528, 1902
Es_1908a	Espin, T.E.
	MNRAS 68, 202, 1908
Es_1919	Espin, T.E.
	MNRAS 79, 211, 1919
Fla1906A	Flammarion, C.
	cited by Lewis (MemRAS 56, 1, 1906)
Fle1958	Fleckenstein, J.
	Milan Merate New Ser, # 115, 1958
Frk1914a	Franks, W.S.
	MNRAS 74, 517, 1914
Frm1907	Farman, M.
	Obs. de Chevreuse, 1907
Frw1936	Ferwerda, J.G.
	BAN 7, 308, 1936
Ge_1964	Gasteyer, C. & Eichhorn, H.
	AJ 69, 762, 1964
Gld1875	Gledhill, J.
	MemRAS 42, 101, 1875
Gny1879	Goldney, G.A.
	AN 95, 1, 1879
Gui1931	Guillaume, J.
	Bull. Obs. Lyon 13, 151, 1931
Gzl1962	Guntzel-Lingner, U.
	Potsdam Obs. 30, Part 3, 101, 1962
Hau1965	Haupt, H.
	Mitt. Sternw. Wien 12, # 16, 1965
HIP1997a	Hipparcos
	Hipparcos Catalog, ESA SP-1200, 1997 (incl. Hp photometry, vol. 10)
HJ_1833c	Herschel, J.F.W.
	MemRAS 6, 1, 1833
HI_1877a	Hall, A.
	Washington Observations, Appendix 6, 1877
Hrt2017a	Hartkopf, W.I. & Mason, B.D.
	Catalog of Rectilinear Elements (v 2017.09.20)
	(http://www.usno.navy.mil/USNO/astrometry/optical-IR-prod/wds/lin1)
Hzg1909	Hertzprung, E.
	AN 180, 39, 1909
Hzg1920	Hertzprung, E.
	Pub. Potsdam Obs. 24, Pt.2, 1920
Izm2010	Izmailov, I.S., Khovrichева, M.L., Khovrichев, M. Yu., Kiyaeva, O.V., Khrutskaya, E.V., Romanenko, L.G., Grosheva, E.A., Maslennikov, K.L. & Kalinichenko, O.A.
	SvAL 36, 349, 2010 (Pisma Astron. Zhur. 36, 365, 2010)
Izm2015	Izmailov, I.S. & Roshchina E. A.

Table B.6: Continued.

Ref.	
	Ast, Bull. 71, 225, 2016
Jam1921	Jameson, H. MNRAS 81, 567, 1921
Jan1953	Luplau-Janssen, C. & Engelstoft, M.O. J. Obs. 36, 107, 1953
Jan1955	Luplau-Janssen, C. J. Obs. 38, 106, 1955
Je_1880i	Jedrzejewicz, J. AN 98, 289, # 2347, 1880
Jef1951	Jeffers, H.M. Lick Obs. Bull. 20, 51, 1951
Jef1978	Jeffers, H.M. & Vasilevskis, S. AJ 83, 411, 1978
Kai1865	Kaiser, F. AN 64, 97, 1865
Kom1933	Komendantov, N.V. Pulkova Obs. Circ. # 9, 26, 1933
Kom1935a	Komendantov, N.V. Pulkova Obs. Pub. Ser. 2, # 47, 29, 1935
Kpr1979	Kamper, K.W. PASP 91, 678, 1979
Kpr1992	Kamper, K.W. AJ 104, 231, 1992
Kzn1994	Kaznica, J.J. Webb Soc., Double Star Circ. 6, 1994
L__1904	Lewis, T. Greenwich Observations, 1904 (data also in MNRAS 66, 17, 1905)
Lau1907a	Lau, H.E. AN 175, 65, 1907
Lau1908c	Lau, H.E. AN 179, 101, 1908
Lau1909b	Lau, H.E. AN 183, 145, 1909
Lau1911b	Lau, H.E. AN 189, 197, 1911
Lbz1929	Labitzke, P. & Przbyllok, E. Koenigsberg Obs. 45, 65, 1929
Lin1985b	Ling, J.F. (Docobo et al.) Bol. Acad. Galega de Ciencias 2, 131, 1985
Loc2017a	Locatelli, G. Il Bollettino Delle Stelle Doppie 20, 19, 2017
Los2005	Losse, F. Obs. et Travaux 59, 27, 2005
Los2009b	Losse, F. El Observador de Estrellas Dobles, Issue # 3, 45, 2009
Mad1842a	Maedler, J.H. Dorpat Observations 9, 37, 1842
Mad1843	Maedler, J.H. Dorpat Observations 10, 29, 1843
Mad1856	Maedler, J.H. Dorpat Observations 13, 1, 1856
Mad1859	Maedler, J.H. Dorpat Observations 15, 1, 1859
Mad1906A	Maedler, J.H. cited by Lewis (MemRAS 56, 1, 1906)
Mai1863	Main, R. Radcliffe Obs. 23, 186, 1863
Mlr1947	Muller, P. J. Obs. 30, 104, 1947
Mun1948	Mundler, M.

Table B.6: Continued.

Ref.	
Neu1923	Pub. Koenigsstuhl Obs. 15, 87, 1948 Neumin, G.E.
Nst1904	Bull. Obs. Pulkova 9, 1923 Niessen, L.
PkO1988	Ann. R. Obs. Belgium 8, 1, 1904 Pulkova Obs.
PkO2003	Pulkova Visual Double Star Catalogue, 1988 Kiselev A.A., Kalinichenko O.A., Kiyaeva O.V., Shakht N.A., Romanenko L.G., Izmailov I.S., Bykov O.P., Maslennikov K.L., Bukhmastova Yu.L.
PkO2010	Pulkova Visual Double Star Catalogue, 2003 Vizieer Catalog I/292 Shakht, N.A., Gorshanov, D.L., Grosheva, E.A., Kiselev, A.A. & Polyakov, E.V.
PkO2014b	Astrophysics 53, 227, 2010 Kiselev, A.A. Kiyaeva, O.V., Izmailov, I.S., Romanenko, L.G., Kalinichenko, O.A., Vasilkova, O.O., Vasileva, T.A., Shakht, N.A., Gorshanov, D.L. & Roschina, E.A.
PkO2016	Astron. Reports 58, 78, 2014 (SvA 91, 130, 2014) Izmailov, I.S., Roshchinca, E.A., Kiselev, A.A., Kiseleva, T.P., Kalinichenko, O., Bykov, O.P., Kiyaeva, O.V., Romanenko, L.G., Shakht, N.A., Maslennikov, K.L. & Vasileva, T.A.
Plm1878	SvAL 42, 41, 2016 (Pisma Astron. Zhur. 42, 46, 2016) Plummer, W.E.
Pok1936	Oxford Astron. Obs. 1, 17, 1878 Pokrovsky, K.D.
Prz1908	Pulkova Obs. Circ. # 20, 1936 Przbyllok, E.
Prz1926	Veroff. Sternw. Heidelberg 5, 1, 1908 Przbyllok, E., Labitzke, P., & Meier, P.
Ptt1917	Beob. Sternw. Koenigsberg # 44, 1926 Pettit, E.
Rab1923	AJ 30, 167, 1917 Rabe, W.
Rab1961b	AN 217, 413, 1923 Rabe, W.
Rak1982	Pub. Munich Obs. 6, # 2, 1961 Rakos, K.D., Albrecht, R., Jenkner, H., Kreidl, T., Michalke, R., Oberlerchner, D., Santos, E., Schermann, A., et al.
Rnz1935	A&AS 47, 221, 1982 Renz, F.
S__1906A	Pulkova Obs. Circ. # 15, 1935 South, J.
Sbk1892	cited by Lewis (MemRAS 56, 1, 1906) Seabroke, G.M.
Scd1940a	MemRAS 50, 1, 1892 Schmeidler, F.
Scn2003	AN 270, 234, 1940 Schnabel, C.
Sha1927	OAG Gen. Cat. 10,000 vis. double star meas. 1970-2003 (J2000.0), 2003 Shajn, G.
Skf2006c	AN 230, 369, 1927 Skiff, B.A.
Sla1913	private communication, 2006 (may not be properly precessed) (data mining of Joy & Abt, ApJS 28, 1, 1974) Slater, R.C.
Sma1931	J. British Astron. Assoc. 24, 218, 1913 Smart, W.M., Green, H.E., & Fletcher, A.
StF1837	MNRAS 92, 37, 1931 Struve, F.G.W.
Str1946	Stellarum duplicium et multiplicium mensurae micrometricae... (Dorpat, Academiae scientiarum caesareae petropolitaneifque), 1837 Strand, K.A.

Table B.6: Continued.

Ref.	
Str1969b	AJ 52, 1, 1946 Strand, K.A. Pub. USNO 18, Pt. 5, 1969
Stt1878	Struve, O. Pulkova Observations 9, 1, 1878
Stt1893a	Struve, O. Pulkova Obs. Observations 10, 1, 1893
Stu1984	Sturdy, K. Webb Soc., Double Star Circ. 3, 1984, part 2
Tal1868	Talmage, C.G. Barclay Obs. 2, 1868
Tal1873	Talmage, C.G. Barclay Obs. 3, 1873
Tal1878	Talmage, C.G. Barclay Obs. 4, 1878
TMA2003	2MASS Catalog 2MASS Point Src Cat., 2003 all-sky release (http://pegasus.phast.umass.edu/)
Tob2003	Tobal, T. OAG Gen. Cat. 10,000 vis. double star meas. 1970-2003 (J2000.0), 2003
Tob2005g	Tobal, T. Webb Soc. Double Star Circ. 13, 2005 (also OAG General Cat. Suppl. 7, 2004)
Tor1985	Pannunzio, R. & Morbidelli, R. A& AS 61, 37, 1985
TYC2000b	Hog, E., Fabricius, C., Makarov, V.V., Urban, S., Corbin, T., Wycoff, G., Bastian, U., Schwkendiek, P., & Wicenc, A. Tycho-2 Cat., A& A 357, 367, 2000
TYC2002	Fabricius, C., Hog, E., Makarov, V.V., Mason, B.D., Wycoff, G.L., & Urban, S.E. A& A 384, 180-189, 2002
UC_2013a	Zacharias, N., Finch, C.T., Girard, T.M., Henden, A., Bartlett, J.L., Monet, D.G., & Zacharias, M.I. (UCAC4 Catalog) AJ 145, 44, 2013
USN1969	Kallarakal, V.V., Lindenblad, I.W., Josties, F.J., Riddle, R.K., Miranian, M., Mintz, B.F., & Klugh, A.P. Pub. USNO 18, Pt.7, 1969
USN1974	Josties, F.J., Dahn, C.C., Kallarakal, V.V., Miranian, M., & Douglass, G.G. Pub. USNO 22, Pt.6, 1974
USN1978	Josties, F.J., Kallarakal, V.V., Douglass, G.G., & Christy, J.W. Pub. USNO 24, Pt.5, 1978
USN1984	Josties, F.J. & Harrington, R.S. ApJS 54, 103, 1984
Vat1939	Tibor, M. et al. Vatican Obs. Astron. Res. 1, 4, 1939
VBs1906	Van Biesbroeck, G. AN 172, 33, 1906
VBs1920	Van Biesbroeck, G. Ann. R. Obs. Belgium 14, 1920
VBs1927a	Van Biesbroeck, G. Pub. Yerkes Obs. 5, pt. 1, 1927
Vil1961	Villiger, W. Pub. Munich Obs. 6, 124, 1961
Vou1922	Voute, J.G. Ann. Leiden Obs. 10, Pt.2, 1922
VvS1919	de Vos van Steenwijk, J.E. MNRAS 80, 223, 1919
Vys1936	Vyssotsky, A.N. AN 258, 401, 1936
Wat1984	Waterworth, R. Webb Soc., Double Star Circ. 3, 1984, part 2
Wbt2017	Webster, N. Webb Soc., Double Star Circ. 25, 51, 2017

Table B.6: Continued.

Ref.	
Wbt2018	Webster, N. Webb Soc., Double Star Circ. 26, 32, 2018
Wen1913	Wendell, O.C. Harvard Ann. 69, 180, 1913
WFC1958a	Schorr, R. & Kohlschuetter, A. AGK2 Cat. North of -2.5 deg, Vol. 1-15, 1958 (Bergedorf)
WFC1959a	Barney, I., Hoffleit, D., & Jones, R.B. Trans. Astron. Obs. Yale Univ., 26, 1, 1959 (+50 to +55 deg zones)
WFC1975	Dieckvoss, W. AGK3 Cat. North of -2.5 deg dec, Vol. 1-8, 1975
WFC1998	Urban, S.E., Corbin, T.E., Wycoff, G.L., Martin, J.C., Jackson, E.S., Zacharias, M.I., & Hall, D.M. AJ 115, 1212, 1998 (Astrographic Cat. 2000)
WFD1925	Tucker, R.H. Pub. Lick Obs. 15, 1925
WFD1926a	Przybyllok, E. Astr. Obs. Konigsberg Univ. Obs. 44, 1926
WFD1931	Boss, B. Albany Cat. of 20,811 stars for epoch 1910, 1931
WFD1933b	Morgan, H.R. Pub. USNO, 2nd ser., v 13, 1933
WFD1935	Dyson, F.W. Second Greenwich Cat. of stars for 1925.0, 1935
WFD1985	CUO, RGO, ROA Carlsberg Meridian Cat. - La Palma, # 1-8, 1985-1994
Wie1956c	Wieth-Knudsen, N. J. Obs. 39, 201, 1956
Wnc1869	Winnecke, A. AN 73, 145, 1869
Wor1956b	Worley, C.E. AJ 61, 162, 1956
WSI2008	Mason, B.D., Hartkopf, W.I., & Wycoff, G.L. AJ 136, 2223, 2008
WSI2011b	Mason, B.D., Hartkopf, W.I., & Wycoff, G.L. AJ 142, 46, 2011
WSI2015	Hartkopf, W.I. & Mason, B.D. AJ 150, 136, 2015
WSI2017b	Mason, B.D. & Hartkopf, W.I. AJ 143, 183, 2017
WSI2018a	Mason, B.D., Hartkopf, W.I. & Urban, S.E. 2017 USNO 26" speckle measures, in preparation