

Singular limit for reactive diffusive transport through an array of thin channels in case of critical diffusivity

Markus Gahn*, Maria Neuss-Radu†

Abstract

We consider a nonlinear reaction–diffusion equation in a domain consisting of two bulk regions connected via small channels periodically distributed within a thin layer. The height and the thickness of the channels are of order ϵ , and the equation inside the layer depends on the parameter ϵ . We consider the critical scaling of the diffusion coefficients in the channels and nonlinear Neumann-boundary condition on the channels' lateral boundaries. We derive effective models in the limit $\epsilon \rightarrow 0$, when the channel-domain is replaced by an interface Σ between the two bulk-domains. Due to the critical size of the diffusion coefficients, we obtain jumps for the solution and its normal fluxes across Σ , involving the solutions of local cell problems on the reference channel in every point of the interface Σ .

Keywords: Array of channels; homogenization; two-scale convergence; reaction-diffusion equation; effective transmission conditions; nonlinear boundary conditions

AMS Subject Classification: 35K57; 35B27, 35Q92

1 Introduction

In this paper we consider reaction-diffusion processes in a microscopic domain Ω_ϵ consisting of two bulk-domains Ω_ϵ^+ and Ω_ϵ^- , which are connected via small periodically distributed channels $\Omega_{*,\epsilon}^M$ obtained by scaled and shifted reference elements Z^* . The height and the thickness of the channels is of order ϵ , where the parameter ϵ is small compared to the size of the bulk-domains. Within the

*Corresponding Author; Interdisciplinary Center for Scientific Computing, University of Heidelberg, Im Neuenheimer Feld 205, Heidelberg, 69120, Germany, markus.gahn@iwr.uni-heidelberg.de

†Department Mathematics, Friedrich-Alexander-Universität Erlangen-Nürnberg Cauerstr. 11, Erlangen, 91058, Germany, maria.neuss-radu@math.fau.de

microscopic domain Ω_ϵ we consider a reaction-diffusion equation with nonlinear reaction-kinetics and nonlinear Neumann-boundary condition on the lateral boundary of the channels. This boundary condition describes for example reactions taking place at the boundary of the channels or exchange with the surrounding medium. Within the channels we assume low diffusivity of order ϵ . The aim is the derivation of a macroscopic model with effective interface conditions in the singular limit $\epsilon \rightarrow 0$.

Reaction-diffusion transport plays an important role in many applications. We want to mention gap junctions connecting biological cells and also ion channels in the membranes of cells and organells. Other important examples are filters arising in engineering sciences. In all these applications nonlinear effects on the boundary of the channels play an important role. However, even for relatively simple models, just taking into account reaction and diffusion processes, numerical simulations are very expensive. Therefore, macroscopic approximations of the solutions, obtained in the limit $\epsilon \rightarrow 0$, are highly demanded.

In the singular limit $\epsilon \rightarrow 0$ we obtain two bulk-domains Ω^+ and Ω^- , which are separated by the interface Σ . The crucial point is the derivation of the interface conditions across Σ in the macroscopic model. These interface conditions carry information about the microscopic processes in the channels $\Omega_{*,\epsilon}^M$. We obtain a jump for the solution and its normal fluxes across Σ , involving the solution of local cell problems of reaction-diffusion type on the reference element Z^* in every point of the interface Σ . This coupling between the microscopic variable from Z^* and the macroscopic variable from Σ arises due to critical scaling for the diffusion in the microscopic model, and leads to additional difficulties in the limit process. Here, the most challenging step is to pass to the limit in the thin channels, where we have to cope simultaneously with the singular limit and the periodicity of the channels. Therefor we use the method of two-scale convergence for thin channels and their oscillating surface, which was defined in [3] and is closely related to the two-scale convergence in thin heterogeneous layers introduced in [18]. Due to the specific scaling in the microscopic equation in the channels, the macroscopic two-scale limit of the microscopic solution is depending on both, the macroscopic variable $\bar{x} \in \Sigma$ and the microscopic variable $y \in Z^*$.

As a first step in the derivation of the macroscopic model we derive ϵ -dependent a priori estimates for the microscopic solutions, which imply (weak) two-scale convergence for the solutions in the thin layer. These compactness results are enough to pass to the limit in the linear terms of the microscopic equation, but not for the nonlinear terms, especially on the boundary of the channels. For those terms we need strong two-scale compactness results. Using the unfolding method for thin channels, which gives us an equivalent characterization of the two-scale convergence, we prove a general strong two-scale compactness result based on error estimates for the discrete shifts of the microscopic solution. An additional difficulty in our problem arises due to the fact, that the time-derivative of the microscopic solution is only a functional on a function space defined on the whole domain Ω_ϵ and it is not straightforward to obtain the existence of the time-derivative of the unfolded sequence together with suitable

a priori estimates. To overcome this problem we use a duality argument to show that the time-derivative of the unfolded sequence in the channels exists and can be controlled by the time-derivative of the microscopic solution on the whole domain. To exhibit the form of the macroscopic model in the limit $\epsilon \rightarrow 0$, we construct test-functions adapted to the structure of the transmission problem and which are admissible for the definition of the two-scale convergence in the channel domain. These test-functions lie in a function space which we show to be dense in the space of macroscopic solutions.

First homogenization results for problems with a geometrical framework related to our setting were given in [20]. In [7, 8], the Stokes-equation was considered in two-bulk domains separated by a sieve of thickness zero. Contributions to the homogenization of the Laplace equation in domains connected by thin channels have been given in [9, 19, 21, 22], where the asymptotic behavior of the solution is investigated for different ratio of the thickness of the layer and the radius of the cylindrical channels. The homogenization of an elliptic Steklov type spectral problem in domains connected by thin channels was considered in [1, 10]. Reaction-diffusion problems a thin layer instead of channels were considered in [14, 15, 18]. In this paper we extend those results to channels, nonlinear boundary conditions, and low regularity for the time-derivative, for the case of low diffusion in the channels, see also [11]. The case of moderate and high diffusivity, however for linear problems with Neumann boundary conditions, can be found in [3]. The more challenging problem concerning the ion transport through channels of biological membranes was announced in [17].

The structure of the paper is as follows. In Section 2 we introduce the microscopic model with its underlying geometrical structure, and give the definition of a weak solution and the necessary function spaces. Further, we prove existence and uniqueness of a unique solution and establish a priori estimates for these solutions and their shifts. In Section 3 we give the definition of the two-scale convergence and the unfolding operator for thin channels. Further, we derive weak and strong two-scale compactness results associated with the specific scaling in our microscopic model. Finally in Section 4 we derive the macroscopic model.

2 The microscopic model

Let $\epsilon > 0$ be a sequence of positive numbers tending to zero such that $\epsilon^{-1} \in \mathbb{N}$ and let $H > 0$ be a fixed real number. Let $n \in \mathbb{N}, n \geq 2$. For $x \in \mathbb{R}^n$, we write $x = (\bar{x}, x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$. Let Ω be a subset of \mathbb{R}^n defined as

$$\Omega := \Sigma \times (-H, H),$$

where $\Sigma \subset \mathbb{R}^{n-1}$ is a connected and open domain with Lipschitz-boundary.

We consider the domain $\Omega_\epsilon \subset \Omega$ consisting of three subdomains: the bulk regions Ω_ϵ^+ and Ω_ϵ^- which are connected by channels periodically distributed within a thin layer constituting the domain $\Omega_{*,\epsilon}^M$, see Figure 1. The bulk regions




Figure 1: Microscopic domain Ω_ϵ for the case $\epsilon = \frac{1}{3}$ and $n = 3$.




Figure 2: Standard channel domain Z^* in the standard cell Z .

are given by:

$$\Omega_\epsilon^+ = \Sigma \times (\epsilon, H), \quad \Omega_\epsilon^- = \Sigma \times (-H, -\epsilon).$$

Furthermore, we denote by

$$S_\epsilon^+ = \Sigma \times \{\epsilon\}, \quad S_\epsilon^- = \Sigma \times \{-\epsilon\},$$

the bottom and the top of Ω_ϵ^+ and Ω_ϵ^- , respectively. The thin layer separating the two bulk domains is given by

$$\Omega_\epsilon^M := \Sigma \times (-\epsilon, \epsilon).$$

To define the channels, which are periodically distributed within Ω_ϵ^M , we first define the standard cell

$$Z := Y \times (-1, 1) := (0, 1)^{n-1} \times (-1, 1)$$

with the upper and lower boundaries

$$S^\pm := Y \times \{\pm 1\}.$$

Let $Z^* \subset Z$ be a connected and open Lipschitz domain representing the standard channel domain, see Figure 2, such that

$$S_*^\pm := \{y \in \partial Z^* : y_n = \pm 1\}$$

is a Lipschitz-domain in \mathbb{R}^{n-1} with positive measure. Let the lateral boundary of the standard channel be denoted by

$$N := \partial Z^* \setminus (S_*^+ \cup S_*^-).$$

We assume that this lateral boundary has a finite distance to the lateral boundary $\partial Z \setminus (S^+ \cup S^-)$ of the standard cell Z . The domain consisting of the channels is then given by

$$\Omega_{*,\epsilon}^M := \bigcup_{\bar{k} \in I_\epsilon} \epsilon(Z^* + (\bar{k}, 0))$$

where $I_\epsilon = \{\bar{k} \in \mathbb{Z}^{n-1} : \epsilon(Z + (\bar{k}, 0)) \subset \Omega_\epsilon^M\}$. The interfaces between the channel domain and the bulk domains are defined by

$$S_{*,\epsilon}^\pm := \bigcup_{\bar{k} \in I_\epsilon} \epsilon(S_*^\pm + (\bar{k}, 0)).$$

The microscopic domain Ω_ϵ is thus defined by

$$\Omega_\epsilon = \Omega_\epsilon^+ \cup \Omega_\epsilon^- \cup \Omega_{*,\epsilon}^M \cup S_{*,\epsilon}^+ \cup S_{*,\epsilon}^-.$$

We assume Ω_ϵ to be Lipschitz. The boundary of Ω_ϵ is given by $\partial\Omega_\epsilon = N_\epsilon \cup \partial_N\Omega_\epsilon$, where

$$N_\epsilon := \bigcup_{k \in I_\epsilon} \epsilon(N + (k, 0)), \quad \partial_N\Omega_\epsilon := \partial\Omega_\epsilon \setminus N_\epsilon.$$

For a function defined on Ω_ϵ , we usually add superscripts $+$, $-$, M to denote its restriction to the sub-domains Ω_ϵ^+ , Ω_ϵ^- and $\Omega_{*,\epsilon}^M$ respectively. Finally, we define the domains

$$\Omega^+ := \Sigma \times (0, H), \quad \Omega^- := \Sigma \times (-H, 0).$$

which are separated by the interface Σ .

Let $p \in [1, \infty]$ and p' denotes the dual exponent of p , i. e., $\frac{1}{p} + \frac{1}{p'} = 1$. For a suitable subset $G \subset \mathbb{R}^m$ with $m \in \mathbb{N}$ we use the following notation for $u \in L^p(G)$ and $v \in L^{p'}(G)$:

$$(u, v)_G := \int_G u(z)v(z)dz,$$

and for $p = p' = 2$ we just obtain the inner product on $L^2(G)$. With the subscript per we indicate function spaces of functions defined on subsets of \mathbb{R}^n which are periodic with respect to the first $(n-1)$ components. For example we write $C_{\text{per}}(\bar{Z}^*)$ for the space of functions $C(\bar{Z}^*)$ periodically extended in \bar{y} -direction.

2.1 The microscopic model

We study the following reaction-diffusion problem for the unknown function $u_\epsilon = (u_\epsilon^+, u_\epsilon^M, u_\epsilon^-) : (0, T) \times \Omega_\epsilon \rightarrow \mathbb{R}$:

$$\begin{aligned} \partial_t u_\epsilon^\pm - D^\pm \Delta u_\epsilon^\pm &= f^\pm(u_\epsilon^\pm) && \text{in } (0, T) \times \Omega_\epsilon^\pm, \\ \frac{1}{\epsilon} \partial_t u_\epsilon^M - \nabla \cdot \left(\epsilon D^M \left(\frac{x}{\epsilon} \right) \nabla u_\epsilon^M \right) &= \frac{1}{\epsilon} g_\epsilon(u_\epsilon^M) && \text{in } (0, T) \times \Omega_\epsilon^M, \end{aligned} \quad (1a)$$

with the boundary conditions

$$\begin{aligned} \nabla u_\epsilon^\pm \cdot \nu &= 0 && \text{on } (0, T) \times \partial_N \Omega_\epsilon, \\ -\epsilon D^M \left(\frac{x}{\epsilon} \right) \nabla u_\epsilon^M \cdot \nu &= h_\epsilon(u_\epsilon^M) && \text{on } (0, T) \times N_\epsilon, \end{aligned} \quad (1b)$$

where ν denotes the outer unit normal with respect to Ω_ϵ , and the initial conditions

$$u_\epsilon(0) = u_{\epsilon,i} \quad \text{on } \Omega_\epsilon. \quad (1c)$$

At the interfaces $S_{*,\epsilon}^\pm$, we impose the natural transmission conditions, i.e., the continuity of the solution and of the normal flux, namely

$$\begin{aligned} u_\epsilon^\pm &= u_\epsilon^M && \text{on } (0, T) \times S_{*,\epsilon}^\pm, \\ D^\pm \nabla u_\epsilon^\pm \cdot \nu &= \epsilon D^M \left(\frac{x}{\epsilon} \right) \nabla u_\epsilon^M \cdot \nu && \text{on } (0, T) \times S_{*,\epsilon}^\pm, \end{aligned} \quad (1d)$$

where here ν denotes an arbitrary unit normal on $S_{*,\epsilon}^\pm$. We emphasize that we consider constant diffusion coefficients D^\pm in the bulk-domains Ω_ϵ^\pm just for an easier notation. The results can be easily extended to more general problems, for example oscillating diffusion coefficients.

2.1.1 Assumptions on the Data:

(A1) We assume $D^\pm > 0$ and $D^M : Z^* \rightarrow \mathbb{R}^{n \times n}$ with $D^M \in L^\infty(Z^*)^{n \times n}$ Y -periodic and coercive, that means for almost every $y \in Z^*$ and all $\xi \in \mathbb{R}^n$ it holds for $c_0 > 0$ that

$$D^M(y)\xi \cdot \xi \geq c_0 \|\xi\|^2.$$

(A2) For the reaction rates in the bulk domains we assume $f \in C^0([0, T] \times \bar{\Omega} \times \mathbb{R})$ is Lipschitz-continuous with respect to the last variable.

(A3) For the reaction rate in the channels domain we assume $g_\epsilon(t, x, z) = g\left(t, \frac{x}{\epsilon}, z\right)$ for $(t, x, z) \in (0, T) \times \Omega_{*,\epsilon}^M \times \mathbb{R}$, with $g \in C^0([0, T] \times \bar{Z}^* \times \mathbb{R})$ Y -periodically extended in the second variable. Further, we assume that g is Lipschitz-continuous with respect to the last variable.

(A4) For the reaction rate on the lateral surface of the channels we assume $h_\epsilon(t, x, z) := h\left(t, \frac{x}{\epsilon}, z\right)$ for $(t, x, z) \in (0, T) \times N_\epsilon \times \mathbb{R}$, with $h \in C^0([0, T] \times \overline{N} \times \mathbb{R})$ extended Y -periodically with respect to the second variable, and Lipschitz-continuous with respect to the last variable.

(A5) For the initial conditions, we assume that

$$u_{\epsilon,i}(x) = \begin{cases} u_i^+(x) & \text{for } x \in \Omega_\epsilon^+, \\ u_i^M\left(\bar{x}, \frac{x_n}{\epsilon}\right) & \text{for } x \in \Omega_{*,\epsilon}^M, \\ u_i^-(x) & \text{for } x \in \Omega_\epsilon^-, \end{cases}$$

where $(u_i^+, u_i^M, u_i^-) \in L^2(\Omega^+) \times L^2(\Sigma, C^0(\overline{Z^*})) \times L^2(\Omega^-)$. Especially, it holds that

$$\frac{1}{\sqrt{\epsilon}} \left\| u_i^M\left(\cdot, \frac{x_n}{\epsilon}\right) \right\|_{L^2(\Omega_{*,\epsilon}^M)} \leq \|u_i^M\|_{L^2(\Sigma, C^0(\overline{Z^*}))} \leq C.$$

2.2 Weak solution of the microscopic problem

In this section we define a weak solution of the microscopic model (1). Therefore, we introduce Hilbert spaces with inner products adapted to the scaling in the equation for u_ϵ^M in the channels. First of all, we define

$$\mathcal{L}_\epsilon := L^2(\Omega_\epsilon) = L^2(\Omega_\epsilon^+) \times L^2(\Omega_{*,\epsilon}^M) \times L^2(\Omega_\epsilon^-),$$

together with the inner product

$$(w_\epsilon, v_\epsilon)_{\mathcal{L}_\epsilon} := (w_\epsilon, v_\epsilon)_{\Omega_\epsilon^+} + (w_\epsilon, v_\epsilon)_{\Omega_\epsilon^-} + \frac{1}{\epsilon} (w_\epsilon, v_\epsilon)_{\Omega_{*,\epsilon}^M}.$$

The second equality in the definition of \mathcal{L}_ϵ means that we always identify $L^2(\Omega_\epsilon)$ with the product space on the different compartments. Further, we define

$$\mathcal{H}_\epsilon := H^1(\Omega_\epsilon) = \left\{ (u_\epsilon^+, u_\epsilon^M, u_\epsilon^-) \in H^1(\Omega_\epsilon^+) \times H^1(\Omega_{*,\epsilon}^M) \times H^1(\Omega_\epsilon^-) : u_\epsilon^\pm = u_\epsilon^M \text{ on } S_{*,\epsilon}^\pm \right\},$$

with the inner product

$$(w_\epsilon, v_\epsilon)_{\mathcal{H}_\epsilon} := (w_\epsilon, v_\epsilon)_{\mathcal{L}_\epsilon} + (\nabla w_\epsilon, \nabla v_\epsilon)_{\Omega_\epsilon^+} + (\nabla w_\epsilon, \nabla v_\epsilon)_{\Omega_\epsilon^-} + \epsilon (\nabla w_\epsilon, \nabla v_\epsilon)_{\Omega_{*,\epsilon}^M}.$$

The associated norms on \mathcal{L}_ϵ and \mathcal{H}_ϵ are denoted by $\|\cdot\|_{\mathcal{L}_\epsilon}$ and $\|\cdot\|_{\mathcal{H}_\epsilon}$. We immediately obtain the Gelfant-triple

$$\mathcal{H}_\epsilon \hookrightarrow \mathcal{L}_\epsilon \hookrightarrow \mathcal{H}'_\epsilon. \quad (2)$$

Definition 2.1. *We say that $u_\epsilon = (u_\epsilon^+, u_\epsilon^M, u_\epsilon^-)$ is a weak solution of the microscopic model (1), if*

$$u_\epsilon \in L^2((0, T), \mathcal{H}_\epsilon) \cap H^1((0, T), \mathcal{H}'_\epsilon),$$

for all $\phi_\epsilon \in \mathcal{H}_\epsilon$ it holds the variational equation

$$\begin{aligned} \langle \partial_t u_\epsilon, \phi_\epsilon \rangle_{\mathcal{H}'_\epsilon, \mathcal{H}_\epsilon} + \sum_{\pm} (D^\pm \nabla u_\epsilon^\pm, \nabla \phi_\epsilon)_{\Omega_\epsilon^\pm} + \epsilon \left(D^M \left(\frac{\cdot}{\epsilon} \right) \nabla u_\epsilon^M, \nabla \phi_\epsilon \right)_{\Omega_{*,\epsilon}^M} \\ = \sum_{\pm} (f^\pm(u_\epsilon^\pm), \phi_\epsilon)_{\Omega_\epsilon^\pm} + \frac{1}{\epsilon} (g_\epsilon(u_\epsilon^M), \phi_\epsilon)_{\Omega_{*,\epsilon}^M} - (h_\epsilon(u_\epsilon^M), \phi_\epsilon)_{N_\epsilon}, \end{aligned} \quad (3)$$

and u_ϵ fulfills the initial condition (1c).

The duality pairing in (3) is connected to the scaling in the microscopic equation in (1a) in the following way. For $\partial_t u_\epsilon^\pm \in L^2((0, T), H^1(\Omega_\epsilon^\pm)')$ and $\partial_t u_\epsilon^M \in L^2((0, T), H^1(\Omega_{*,\epsilon}^M)')$ we obtain from the Gelfand-triplet (2) and the definition of the inner product on \mathcal{L}_ϵ

$$\langle \partial_t u_\epsilon, \phi_\epsilon \rangle_{\mathcal{H}'_\epsilon, \mathcal{H}_\epsilon} = \sum_{\pm} \langle \partial_t u_\epsilon^\pm, \phi_\epsilon \rangle_{H^1(\Omega_\epsilon^\pm)', H^1(\Omega_\epsilon^\pm)} + \frac{1}{\epsilon} \langle \partial_t u_\epsilon^M, \phi_\epsilon \rangle_{H^1(\Omega_{*,\epsilon}^M)', H^1(\Omega_{*,\epsilon}^M)}.$$

However, in our case the time-derivative $\partial_t u_\epsilon$ is a functional defined on the whole space \mathcal{H}_ϵ , i. e., a space of functions defined on the whole domain Ω_ϵ , and it is not straightforward to restrict such functionals to $H^1(\Omega_\epsilon^\pm)'$ and $H^1(\Omega_{*,\epsilon}^M)'$. For the derivation of the strong compactness results in the channels we have to control the time-derivative in the domain $\Omega_{*,\epsilon}^M$. Therefore, we introduce the space of functions with zero traces on the interface between the channels and the bulk-domains

$$\mathcal{H}_{\epsilon,0}^M := \left\{ v_\epsilon^M \in H^1(\Omega_{*,\epsilon}^M) : v_\epsilon^M|_{S_{*,\epsilon}^\pm} = 0 \right\},$$

together with the inner product

$$(w_\epsilon^M, v_\epsilon^M)_{\mathcal{H}_{\epsilon,0}^M} := \frac{1}{\epsilon} (w_\epsilon^M, v_\epsilon^M)_{\Omega_{*,\epsilon}^M} + \epsilon (\nabla w_\epsilon^M, \nabla v_\epsilon^M)_{\Omega_{*,\epsilon}^M},$$

and the associated norm denoted by $\|\cdot\|_{\mathcal{H}_{\epsilon,0}^M}$. This leads to the Gelfand-triplet $\mathcal{H}_{\epsilon,0}^M \hookrightarrow L^2(\Omega_{*,\epsilon}^M) \hookrightarrow (\mathcal{H}_{\epsilon,0}^M)'$. By extending functions $\phi_\epsilon^M \in \mathcal{H}_{\epsilon,0}^M$ by zero to Ω_ϵ , we obtain for $u_\epsilon \in L^2((0, T), \mathcal{H}_\epsilon) \cap H^1((0, T), \mathcal{H}'_\epsilon)$

$$\langle \partial_t u_\epsilon^M, \phi_\epsilon^M \rangle_{(\mathcal{H}_{\epsilon,0}^M)', \mathcal{H}_{\epsilon,0}^M} = \epsilon \langle \partial_t u_\epsilon, \phi_\epsilon^M \rangle_{\mathcal{H}'_\epsilon, \mathcal{H}_\epsilon},$$

and

$$\|\partial_t u_\epsilon^M\|_{(\mathcal{H}_{\epsilon,0}^M)'} \leq \epsilon \|\partial_t u_\epsilon\|_{\mathcal{H}'_\epsilon}. \quad (4)$$

2.3 Existence of a weak solution and a priori estimates

In this section we give the existence result for the microscopic model (1) and show a priori estimates depending explicitly on the parameter ϵ . These estimates form the basis for the derivation of the macroscopic model.

Proposition 2.2. *There exists a unique weak solution of the microscopic model (1).*

Proof. The claim follows from the Galerkin-method and a Leray-Schauder fixed-point argument, together with a priori estimates similar as in Lemma 2.3. Since this is quite standard we skip the details. \square

In the next lemma, we state the a priori estimates for the microscopic solution u_ϵ . Therefore, we make use of the following Poincaré-inequality: For all $\theta > 0$ there exists $C(\theta) > 0$, such that for all $\phi_\epsilon^M \in H^1(\Omega_{*,\epsilon}^M)$ it holds that

$$\|\phi_\epsilon^M\|_{L^2(N_\epsilon)} \leq \frac{C(\theta)}{\sqrt{\epsilon}} \|\phi_\epsilon^M\|_{L^2(\Omega_{*,\epsilon}^M)} + \theta\sqrt{\epsilon} \|\nabla\phi_\epsilon^M\|_{L^2(\Omega_{*,\epsilon}^M)}. \quad (5)$$

This result is easily obtained by decomposing $\Omega_{*,\epsilon}^M$ into microscopic cells $\epsilon(Z^* + (\bar{k}, 0))$ for $\bar{k} \in I_\epsilon$, using a scaling argument, and the usual trace estimate on ∂Z^* .

Lemma 2.3. *The solution u_ϵ of the microscopic model (1) fulfills the following a priori estimate*

$$\|\partial_t u_\epsilon\|_{L^2((0,T),\mathcal{H}_\epsilon)} + \|u_\epsilon\|_{L^2((0,T),\mathcal{H}_\epsilon)} \leq C.$$

Especially, we obtain

$$\|\partial_t u_\epsilon\|_{L^2((0,T),(\mathcal{H}_{\epsilon,0}^M)')} \leq C\epsilon.$$

Proof. We test the variational equation (3) with u_ϵ to obtain with the coercivity of D^M

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|u_\epsilon\|_{\mathcal{L}_\epsilon}^2 + \sum_{\pm} D^\pm \|\nabla u_\epsilon^\pm\|_{L^2(\Omega_\epsilon^\pm)}^2 + c_0\epsilon \|\nabla u_\epsilon^M\|_{L^2(\Omega_{*,\epsilon}^M)}^2 \\ \leq \sum_{\pm} (f^\pm(u_\epsilon^\pm), u_\epsilon^\pm)_{\Omega_\epsilon^\pm} + \frac{1}{\epsilon} (g_\epsilon(u_\epsilon^M), u_\epsilon^M)_{\Omega_{*,\epsilon}^M} - (h_\epsilon(u_\epsilon^M), u_\epsilon^M)_{N_\epsilon}. \end{aligned} \quad (6)$$

We only consider the boundary term on N_ϵ in more detail, since the other terms can be treated in a similar way. The Lipschitz-continuity of h_ϵ and the trace-inequality (5) imply for $\theta > 0$ (remember $|N_\epsilon| \leq C$)

$$\begin{aligned} |(h_\epsilon(u_\epsilon^M), u_\epsilon^M)_{N_\epsilon}| &\leq C \|1 + u_\epsilon^M\|_{L^2(N_\epsilon)} \|u_\epsilon^M\|_{L^2(N_\epsilon)} \leq C(1 + \|u_\epsilon^M\|_{L^2(N_\epsilon)}^2) \\ &\leq C(\theta) \left(1 + \frac{1}{\epsilon} \|u_\epsilon^M\|_{L^2(\Omega_{*,\epsilon}^M)}^2\right) + \theta\epsilon \|\nabla u_\epsilon^M\|_{L^2(\Omega_{*,\epsilon}^M)}^2. \end{aligned}$$

For $\theta > 0$ small enough the last term can be absorbed from the left-hand side in the inequality (6). Integration with respect to time and the Gronwall inequality implies the estimate for u_ϵ in the norm of $L^2((0, T), \mathcal{H}_\epsilon)$. For the estimate of the time-derivative we just test equation (3) with $\phi_\epsilon \in \mathcal{H}_\epsilon$ and $\|\phi_\epsilon\|_{\mathcal{H}_\epsilon} \leq 1$. We skip the details. The last inequality in the Lemma follows from (4). \square

To pass to the limit in the nonlinear terms of (3), we need strong two-scale compactness results. Due to the critical scaling of the equation in the channels, we will see that the two-scale limit u_0^M of u_ϵ^M is depending on a macroscopic variable $\bar{x} \in \Sigma$ and a microscopic variable $y \in Z^*$, see Theorem 4.2. Hence, usual compactness arguments, for example a direct application of the Aubin-Lions lemma, fail. To overcome this problem we use the unfolding operator for thin layers and apply a Kolmogorov-type compactness result to the unfolded sequence. This argument is based on error estimates for the difference of discrete shifts of the microscopic solution. We introduce the following notation:

For an arbitrary set $U \subset \mathbb{R}^n$ and a function $v_\epsilon : U \rightarrow \mathbb{R}$, we define for $l \in \mathbb{Z}^n$ the shifted function

$$v_\epsilon^l(x) := v_\epsilon(x + \epsilon l).$$

Here, if not stated otherwise, we extend v_ϵ by zero to \mathbb{R}^n . Then, we define the difference between the shifted function and the function itself by

$$\delta_l v_\epsilon(x) := \delta v_\epsilon(x) := v_\epsilon^l(x) - v_\epsilon(x) = v_\epsilon(x + \epsilon l) - v_\epsilon(x).$$

If it is clear from the context we neglect the index l and just write δv_ϵ . Now, for $0 < h \ll 1$ we define

$$\Sigma_h := \{x \in \Sigma : \text{dist}(\partial\Sigma, x) > h\},$$

and

$$\Omega_{\epsilon,h} := \Omega_\epsilon \cap (\Sigma_h \times (-H, H)), \quad \Omega_{\epsilon,h}^\pm := \Omega_{\epsilon,h} \cap \Omega_\epsilon^\pm.$$

We emphasize that in the definition of $\Omega_{\epsilon,h}$ channels can be intersected by the boundary of $\Sigma_h \times (-H, H)$ and we would lose the Lipschitz-regularity of $\Omega_{\epsilon,h}$. Therefore we use the domain $\widehat{\Omega}_{\epsilon,h}$ defined in the following. Let

$$I_{\epsilon,h} := \{\bar{k} \in \mathbb{Z}^{n-1} : \epsilon(Y + \bar{k}) \subset \Sigma_h\}, \quad \widehat{\Sigma}_h := \text{int} \left(\bigcup_{\bar{k} \in I_{\epsilon,h}} \epsilon(\bar{Y} + \bar{k}) \right),$$

and

$$\widehat{\Omega}_{*,\epsilon,h}^M := \Omega_{*,\epsilon}^M \cap (\widehat{\Sigma}_h \times (-\epsilon, \epsilon)), \quad \widehat{\Omega}_{\epsilon,h}^+ := \widehat{\Sigma}_h \times (\epsilon, H), \quad \widehat{\Omega}_{\epsilon,h}^- := \widehat{\Sigma}_h \times (-H, -\epsilon).$$

The top and the bottom of the channels in $\widehat{\Omega}_{*,\epsilon,h}$ are denoted by $\widehat{S}_{*,\epsilon,h}^\pm$, and the lateral boundary of the channels by $\widehat{N}_{\epsilon,h}$. Then we define

$$\widehat{\Omega}_{\epsilon,h} := \widehat{\Omega}_{\epsilon,h}^+ \cup \widehat{\Omega}_{\epsilon,h}^- \cup \widehat{\Omega}_{*,\epsilon,h}^M \cup \widehat{S}_{*,\epsilon,h}^+ \cup \widehat{S}_{*,\epsilon,h}^-.$$

On this domain we define the function space

$$\mathcal{H}_{\epsilon,h,0} := \left\{ v_\epsilon \in H^1(\widehat{\Omega}_{\epsilon,h}) : v_\epsilon|_{\partial\widehat{\Sigma}_h \times (-H, -\epsilon)} = v_\epsilon|_{\partial\widehat{\Sigma}_h \times (\epsilon, H)} = 0 \right\} \subset \mathcal{H}_\epsilon,$$

where we consider $\mathcal{H}_{\epsilon,h,0}$ as a subset of \mathcal{H}_ϵ by extending functions from $\mathcal{H}_{\epsilon,h,0}$ by zero to the whole domain Ω_ϵ . In a similar way we define

$$\mathcal{L}_{\epsilon,h} := L^2(\widehat{\Omega}_{\epsilon,h}) = L^2(\widehat{\Omega}_{\epsilon,h}^+) \times L^2(\widehat{\Omega}_{*,\epsilon,h}^M) \times L^2(\widehat{\Omega}_{\epsilon,h}^-) \subset \mathcal{L}_\epsilon,$$

what leads to the Gelfand-triple

$$\mathcal{H}_{\epsilon,h,0} \hookrightarrow \mathcal{L}_{\epsilon,h} \hookrightarrow \mathcal{H}'_{\epsilon,h,0}.$$

We emphasize that on $\mathcal{L}_{\epsilon,h}$ we consider the same inner product as on \mathcal{L}_ϵ , just by extending function in $\mathcal{L}_{\epsilon,h}$ by zero to the whole domain Ω_ϵ .

Lemma 2.4. *Let u_ϵ be the solution of the microscopic model (1). Then, for every $0 < h \ll 1$, there exists a $C = C(h) > 0$, such that for all $l \in \mathbb{Z}^{n-1} \times \{0\}$ with $|\epsilon l| \ll h$ it holds that*

$$\begin{aligned} & \frac{1}{\sqrt{\epsilon}} \|\delta u_\epsilon\|_{L^\infty((0,T), L^2(\widehat{\Omega}_{*,\epsilon,2h}^M))} + \sqrt{\epsilon} \|\nabla \delta u_\epsilon\|_{L^2((0,T) \times \widehat{\Omega}_{*,\epsilon,2h}^M)} \\ & \leq C \left(\epsilon + \|\delta u_\epsilon(0)\|_{\mathcal{L}_{\epsilon,h}} + \sum_{\pm} \|\delta u_\epsilon^\pm\|_{L^2((0,T) \times \Omega_{\epsilon,h}^\pm)} \right). \end{aligned} \quad (7)$$

Proof. We have $u_\epsilon|_{\widehat{\Omega}_{\epsilon,h}}, u_\epsilon^l|_{\widehat{\Omega}_{\epsilon,h}} \in H^1((0,T), \mathcal{H}'_{\epsilon,h,0})$ and for all $\phi_{\epsilon,h} \in \mathcal{H}_{\epsilon,h,0}$ it holds almost everywhere in $(0,T)$ that

$$\begin{aligned} & \langle \partial_t \delta u_\epsilon, \phi_{\epsilon,h} \rangle_{\mathcal{H}'_{\epsilon,h,0}, \mathcal{H}_{\epsilon,h,0}} + \sum_{\pm} (D^\pm \nabla \delta u_\epsilon^\pm, \nabla \phi_{\epsilon,h})_{\widehat{\Omega}_{\epsilon,h}^\pm} + \epsilon \left(D^M \left(\frac{x}{\epsilon} \right) \nabla \delta u_\epsilon^M, \nabla \phi_{\epsilon,h} \right)_{\widehat{\Omega}_{*,\epsilon,h}^M} \\ & = \sum_{\pm} (f^\pm((u_\epsilon^\pm)^l) - f^\pm(u_\epsilon^\pm), \phi_{\epsilon,h})_{\widehat{\Omega}_{\epsilon,h}^\pm} + \frac{1}{\epsilon} (g_\epsilon((u_\epsilon^M)^l) - g_\epsilon(u_\epsilon^M), \phi_{\epsilon,h})_{\widehat{\Omega}_{*,\epsilon,h}^M} \\ & \quad - (h_\epsilon((u_\epsilon^M)^l) - h_\epsilon(u_\epsilon^M), \phi_{\epsilon,h})_{\widehat{N}_{\epsilon,h}}. \end{aligned}$$

Now, let $\eta \in C_0^\infty(\widehat{\Sigma}_h)$ be a cut-off function with $0 \leq \eta \leq 1$ and $\eta = 1$ in $\widehat{\Sigma}_{2h}$, and $\|\nabla_{\bar{x}} \eta\|_{L^\infty(\widehat{\Sigma}_h)} \leq C = C(h)$. As a test-function in the equation above we choose $\phi_{\epsilon,h} = \eta^2 \delta u_\epsilon$ and obtain with the coercivity of D^M

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|\eta \delta u_\epsilon\|_{\mathcal{L}_{\epsilon,h}}^2 + \sum_{\pm} D^\pm \|\eta \nabla \delta u_\epsilon^\pm\|_{L^2(\widehat{\Omega}_{\epsilon,h}^\pm)}^2 + c_0 \epsilon \|\eta \nabla \delta u_\epsilon^M\|_{L^2(\widehat{\Omega}_{*,\epsilon,h}^M)}^2 \\ & \leq \sum_{\pm} \left\{ -2 (D^\pm \nabla \delta u_\epsilon^\pm, \eta \delta u_\epsilon^\pm \nabla \eta)_{\widehat{\Omega}_{\epsilon,h}^\pm} + (\delta f^\pm(u_\epsilon^\pm), \eta^2 \delta u_\epsilon^\pm)_{\widehat{\Omega}_{\epsilon,h}^\pm} \right\} \\ & \quad - 2\epsilon \left(D^M \left(\frac{x}{\epsilon} \right) \nabla \delta u_\epsilon^M, \eta \delta u_\epsilon^M \nabla \eta \right)_{\widehat{\Omega}_{*,\epsilon,h}^M} + \frac{1}{\epsilon} (\delta g_\epsilon(u_\epsilon^M), \eta^2 \delta u_\epsilon^M)_{\widehat{\Omega}_{*,\epsilon,h}^M} \\ & \quad - (\delta h_\epsilon(u_\epsilon^M), \eta^2 \delta u_\epsilon^M)_{\widehat{N}_{\epsilon,h}} \\ & = : \sum_{\pm} [I_{\epsilon,1}^\pm + I_{\epsilon,2}^\pm] + \sum_{j=3}^5 I_{\epsilon,j}^M. \end{aligned}$$

Integration with respect to time gives us for almost every $t \in (0, T)$

$$\begin{aligned} & \|\eta \delta u_\epsilon(t)\|_{\mathcal{L}_{\epsilon,h}}^2 + \sum_{\pm} \|\eta \nabla \delta u_\epsilon^\pm\|_{L^2((0,t) \times \widehat{\Omega}_{\epsilon,h}^\pm)}^2 + \epsilon \|\eta \nabla \delta u_\epsilon^M\|_{L^2((0,t) \times \widehat{\Omega}_{*,\epsilon,h}^M)}^2 \\ & \leq C \int_0^t \left\{ \sum_{\pm} [I_{\epsilon,1}^\pm + I_{\epsilon,2}^\pm] + \sum_{j=3}^5 I_{\epsilon,j}^M \right\} dt + \sum_{\pm} I_{\epsilon,6}^\pm + I_{\epsilon,7}^M, \end{aligned}$$

with

$$I_{\epsilon,6}^\pm := \|\eta \delta u_\epsilon^\pm(0)\|_{L^2(\widehat{\Omega}_{\epsilon,h}^\pm)}^2, \quad I_{\epsilon,7}^M := \frac{1}{\epsilon} \|\eta \delta u_\epsilon^M(0)\|_{L^2(\widehat{\Omega}_{*,\epsilon,h}^M)}^2.$$

We start to estimate the term $I_{\epsilon,1}^\pm$. For $\theta > 0$ we obtain

$$\begin{aligned} \int_0^t I_{\epsilon,1}^\pm & \leq C \|\eta \nabla \delta u_\epsilon^\pm\|_{L^2((0,t) \times \widehat{\Omega}_{\epsilon,h}^\pm)} \|\delta u_\epsilon^\pm\|_{L^2((0,t) \times \widehat{\Omega}_{\epsilon,h}^\pm)} \\ & \leq C(\theta) \|\delta u_\epsilon^\pm\|_{L^2((0,t) \times \widehat{\Omega}_{\epsilon,h}^\pm)}^2 + \theta \|\eta \nabla \delta u_\epsilon^\pm\|_{L^2((0,t) \times \widehat{\Omega}_{\epsilon,h}^\pm)}^2, \end{aligned}$$

for a constant $C(\theta) > 0$ depending on θ . In a similar way and using the a priori estimate from Lemma 2.3, we obtain for $I_{\epsilon,3}^M$

$$\begin{aligned} \int_0^t I_{\epsilon,3}^M dt & \leq C \epsilon \|\nabla \delta u_\epsilon^M\|_{L^2((0,t) \times \widehat{\Omega}_{*,\epsilon,h}^M)} \|\eta \nabla \delta u_\epsilon^M\|_{L^2((0,t) \times \widehat{\Omega}_{*,\epsilon,h}^M)} \\ & \leq C \left(\epsilon^3 \|\nabla \delta u_\epsilon^M\|_{L^2((0,t) \times \widehat{\Omega}_{*,\epsilon,h}^M)}^2 + \frac{1}{\epsilon} \|\eta \delta u_\epsilon^M\|_{L^2((0,t) \times \widehat{\Omega}_{*,\epsilon,h}^M)}^2 \right) \\ & \leq C \left(\epsilon^2 + \frac{1}{\epsilon} \|\eta \delta u_\epsilon^M\|_{L^2((0,t) \times \widehat{\Omega}_{*,\epsilon,h}^M)}^2 \right). \end{aligned}$$

For $I_{\epsilon,5}^M$ we use the Lipschitz-continuity of h_ϵ , the trace-inequality (5), and again the a priori estimate from Lemma 2.3 to obtain for $\theta > 0$

$$\begin{aligned} \int_0^t I_{\epsilon,5}^M dt & \leq C \|\eta \delta u_\epsilon^M\|_{L^2((0,t) \times \widehat{N}_{\epsilon,h})}^2 \\ & \leq \frac{C(\theta)}{\epsilon} \|\eta \delta u_\epsilon^M\|_{L^2((0,t) \times \widehat{\Omega}_{*,\epsilon,h}^M)}^2 + C \epsilon \|\delta u_\epsilon^M\|_{L^2((0,t) \times \widehat{\Omega}_{*,\epsilon,h}^M)}^2 + \theta \epsilon \|\eta \nabla \delta u_\epsilon^M\|_{L^2((0,t) \times \widehat{\Omega}_{*,\epsilon,h}^M)}^2 \\ & \leq \frac{C(\theta)}{\epsilon} \|\eta \delta u_\epsilon^M\|_{L^2((0,t) \times \widehat{\Omega}_{*,\epsilon,h}^M)}^2 + C \epsilon^2 + \theta \epsilon \|\eta \nabla \delta u_\epsilon^M\|_{L^2((0,t) \times \widehat{\Omega}_{*,\epsilon,h}^M)}^2. \end{aligned}$$

For $I_{\epsilon,2}^\pm$ and $I_{\epsilon,4}^M$ we obtain directly from the Lipschitz-continuity of f^\pm and g_ϵ

$$\int_0^t \sum_{\pm} I_{\epsilon,2}^\pm + I_{\epsilon,4}^M dt \leq C \|\delta u_\epsilon\|_{L^2((0,t), \mathcal{L}_{\epsilon,h})}^2.$$

Choosing $\theta > 0$ small enough, the desired result follows from the Gronwall-inequality. \square

Remark 2.5. *The error term $I_{\epsilon,1}^{\pm}$ arises due to the cut-off function η in the proof of Lemma 2.4. Hence, with the method used above we are not able to get rid of the norms of $\delta u_{\epsilon}^{\pm}$ on the right-hand side of inequality (7). For specific boundary conditions, like zero Dirichlet-boundary conditions, or in case of a rectangle Σ , zero Neumann-boundary conditions or periodic boundary conditions on the lateral boundary, it is easily possible to extend the solution u_{ϵ} in \bar{x} -direction and obtain an estimate of the form*

$$\|\delta u_{\epsilon}\|_{L^2((0,T),\mathcal{H}_{\epsilon})} \leq C(\epsilon + \|\delta u_{\epsilon}(0)\|_{\mathcal{L}_{\epsilon}}).$$

However, the interior estimate in Lemma 2.4 is enough to obtain strong two-scale compactness results in the channel domain and the method presented in the proof has the advantage that it is applicable for more general boundary conditions.

3 Two-scale convergence and the unfolding operator for thin channels

In this section we define the two-scale convergence for thin channels and give some weak two-scale compactness results based on a priori estimates in $L^2((0,T),\mathcal{H}_{\epsilon})$. Further, we derive strong two-scale convergence results based on error estimates for the discrete shifts as in Lemma 2.4. Therefore, we make use of the unfolding operator and a Kolmogorov-type compactness result. We start with the definition of the two-scale convergence for channels, see also [3].

Definition 3.1. *Let $p \in [1, \infty)$ and p' the dual exponent of p .*

(i) *We say the sequence $v_{\epsilon} \in L^p((0,T) \times \Omega_{*,\epsilon}^M)$ converges (weakly) in the two-scale sense to a limit function $v_0 \in L^p((0,T) \times \Sigma \times Z^*)$, if*

$$\lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \int_0^T \int_{\Omega_{*,\epsilon}^M} v_{\epsilon}(t,x) \psi\left(t, \bar{x}, \frac{x}{\epsilon}\right) dx dt = \int_0^T \int_{\Sigma} \int_{Z^*} v_0(t, \bar{x}, y) \psi(t, \bar{x}, y) dy d\bar{x} dt,$$

for all $\psi \in L^{p'}((0,T) \times \Sigma, C_{\text{per}}(\overline{Z^}))$. The sequence converges strongly in the two-scale sense (in L^p) if it holds that*

$$\lim_{\epsilon \rightarrow 0} \epsilon^{-\frac{1}{p}} \|v_{\epsilon}\|_{L^p((0,T) \times \Omega_{*,\epsilon}^M)} = \|v_0\|_{L^p((0,T) \times \Sigma \times Z^*)}.$$

(ii) *We say the sequence $v_{\epsilon} \in L^p((0,T) \times N_{\epsilon})$ converges (weakly) in the two-scale sense to a limit function $v_0(t, \bar{x}, y) \in L^p((0,T) \times \Sigma \times N)$ on N_{ϵ} , if*

$$\lim_{\epsilon \rightarrow 0} \int_0^T \int_{N_{\epsilon}} v_{\epsilon}(t,x) \psi\left(t, \bar{x}, \frac{x}{\epsilon}\right) dx dt = \int_0^T \int_{\Sigma} \int_N v_0(t, \bar{x}, y) \psi(t, \bar{x}, y) dy d\bar{x} dt,$$

for all $\psi \in L^{p'}((0,T), C(\overline{\Sigma}, C_{\text{per}}(\overline{N})))$. The sequence converges strongly in the two-scale sense (in L^p) on N_{ϵ} if it holds that

$$\lim_{\epsilon \rightarrow 0} \|v_{\epsilon}\|_{L^p((0,T) \times N_{\epsilon})} = \|v_0\|_{L^p((0,T) \times \Sigma \times N)}.$$

We just say a sequence converges in the two-scale sense, if it converges in the two-scale sense in L^p .

In the following Lemma we give some weak two-scale compactness results in the microscopic channels, based on a priori estimates of the microscopic solution.

Lemma 3.2. *Let $p \in (1, \infty)$ and p' the dual exponent of p .*

(i) *Let v_ϵ be a sequence of functions in $L^p((0, T) \times \Omega_{*,\epsilon}^M)$ such that*

$$\epsilon^{-\frac{1}{p}} \|v_\epsilon\|_{L^p((0, T) \times \Omega_{*,\epsilon}^M)} \leq C.$$

Then, there exists $v_0 \in L^p((0, T) \times \Sigma \times Z^)$ such that, up to a subsequence, v_ϵ two-scale converges to v_0 .*

(ii) *Let v_ϵ be a sequence of functions in $L^p((0, T) \times W^{1,p}(\Omega_{*,\epsilon}^M))$ such that*

$$\epsilon^{-\frac{1}{p}} \|v_\epsilon\|_{L^p((0, T) \times \Omega_{*,\epsilon}^M)} + \epsilon^{\frac{1}{p'}} \|\nabla v_\epsilon\|_{L^p((0, T) \times \Omega_{*,\epsilon}^M)} \leq C.$$

Then, there exists $v_0 \in L^p((0, T) \times \Sigma, W^{1,p}(Z^))$, such that up to a subsequence $v_\epsilon \rightarrow v_0$ and $\epsilon \nabla v_\epsilon \rightarrow \nabla_y v_0$ in the two-scale sense.*

(iii) *Let v_ϵ be a sequence of functions in $L^p((0, T) \times N_\epsilon)$ such that*

$$\|v_\epsilon\|_{L^p((0, T) \times N_\epsilon)} \leq C.$$

Then, there exists $v_0 \in L^p((0, T) \times \Sigma \times N)$, such that $v_\epsilon \rightarrow v_0$ in the two-scale sense on N_ϵ .

Proof. Statements (i) and (iii) were shown in [3, Theorem 4.4] for $p = 2$. The general case works the same lines. To prove (ii) we notice that

$$\epsilon^{-\frac{1}{p}} \left(\|v_\epsilon\|_{L^p((0, T) \times \Omega_{*,\epsilon}^M)} + \|\epsilon \nabla v_\epsilon\|_{L^p((0, T) \times \Omega_{*,\epsilon}^M)} \right) \leq C.$$

Hence, there exist $v_0 \in L^p((0, T) \times \Sigma \times Z^*)$ and $\xi_0 \in L^p((0, T) \times \Sigma \times Z^*)^n$, such that up to a subsequence

$$\begin{aligned} v_\epsilon &\rightarrow v_0 && \text{in the two-scale sense,} \\ \epsilon \nabla v_\epsilon &\rightarrow \xi_0 && \text{in the two-scale sense.} \end{aligned}$$

By integration by parts we obtain for all $\Phi \in C_0^\infty((0, T) \times \Sigma \times Z^*)^n$

$$\begin{aligned} \int_0^T \int_\Sigma \int_{Z^*} v_0 \nabla_y \cdot \Phi(t, \bar{x}, y) dy d\bar{x} dt &= \lim_{\epsilon \rightarrow 0} v_\epsilon \left[\epsilon \nabla_{\bar{x}} \cdot \Phi \left(t, \bar{x}, \frac{x}{\epsilon} \right) + \nabla_y \cdot \Phi \left(t, \bar{x}, \frac{x}{\epsilon} \right) \right] dx dt \\ &= - \lim_{\epsilon \rightarrow 0} \epsilon \int_0^T \int_{\Omega_{*,\epsilon}^M} \nabla v_\epsilon \cdot \Phi \left(t, \bar{x}, \frac{x}{\epsilon} \right) dx dt \\ &= - \int_0^T \int_\Sigma \int_{Z^*} \xi_0 \cdot \Phi(t, \bar{x}, y) dy d\bar{x} dt, \end{aligned}$$

which yields the desired result. \square

Lemma 3.2 and the a priori estimates for the microscopic solution from Lemma 2.3 are enough to pass to the limit in the linear terms in the channels $\Omega_{*,\epsilon}^M$ in the variational equation (3). To pass to the limit in the nonlinear terms, we need strong two-scale convergence. To establish these convergence results, we use the unfolding operator for channels defined below.

Definition 3.3. *Let $(G_\epsilon, G) \in \{(\Omega_{*,\epsilon}^M, Z^*), (N_\epsilon, N)\}$. Then for $p \in [1, \infty)$ we define the unfolding operator*

$$\begin{aligned} \mathcal{T}_\epsilon &: L^p((0, T) \times G_\epsilon) \rightarrow L^p((0, T) \times \Sigma \times G), \\ \mathcal{T}_\epsilon v_\epsilon(t, \bar{x}, y) &= v_\epsilon\left(t, \epsilon \left(\left[\frac{\bar{x}}{\epsilon} \right], 0 \right) + \epsilon y\right), \end{aligned}$$

where $[\cdot]$ denotes the Gauß-bracket.

Here it makes sense to use the same notation for the unfolding operator on $\Omega_{*,\epsilon}^M$ and N_ϵ , since the unfolding operator commutes with the trace operator in the following sense: For $v_\epsilon \in L^p((0, T), W^{1,p}(\Omega_{*,\epsilon}^M))$ it holds that

$$\mathcal{T}_\epsilon(v_\epsilon)|_N = \mathcal{T}_\epsilon(v_\epsilon|_{N_\epsilon}).$$

We summarize some properties of \mathcal{T}_ϵ :

Lemma 3.4. *Let $p \in [1, \infty)$. Then it holds that:*

(i) *For $v_\epsilon \in L^p((0, T) \times \Omega_{*,\epsilon}^M)$ and $w_\epsilon \in L^{p'}((0, T) \times \Omega_{*,\epsilon}^M)$, where p' denotes the dual exponent of p , we have*

$$\begin{aligned} (\mathcal{T}_\epsilon v_\epsilon, \mathcal{T}_\epsilon w_\epsilon)_{(0,T) \times \Sigma \times Z^*} &= \frac{1}{\epsilon} (v_\epsilon, w_\epsilon)_{(0,T) \times \Omega_{*,\epsilon}^M}, \\ \|\mathcal{T}_\epsilon v_\epsilon\|_{L^p((0,T) \times \Sigma \times Z^*)} &= \epsilon^{-\frac{1}{p}} \|v_\epsilon\|_{L^p((0,T) \times \Omega_{*,\epsilon}^M)}, \end{aligned}$$

and for $v_\epsilon \in L^p((0, T), W^{1,p}(\Omega_{*,\epsilon}^M))$ it holds that

$$\nabla_y \mathcal{T}_\epsilon v_\epsilon = \epsilon \mathcal{T}_\epsilon \nabla v_\epsilon.$$

(ii) *For $v_\epsilon \in L^p((0, T) \times N_\epsilon)$ and $w_\epsilon \in L^{p'}((0, T) \times N_\epsilon)$, where p' denotes the dual exponent of p , we have*

$$\begin{aligned} (\mathcal{T}_\epsilon v_\epsilon, \mathcal{T}_\epsilon w_\epsilon)_{(0,T) \times \Sigma \times N} &= (v_\epsilon, w_\epsilon)_{(0,T) \times N_\epsilon}, \\ \|\mathcal{T}_\epsilon v_\epsilon\|_{L^p((0,T) \times \Sigma \times N)} &= \|v_\epsilon\|_{L^p((0,T) \times N_\epsilon)}. \end{aligned}$$

Proof. These results are obtained by a simple calculation. For the main ideas of the proof see [6]. \square

We have the following relation between the two-scale convergence and the unfolding operator.

Lemma 3.5.

- (i) The sequence $v_\epsilon \in L^p((0, T) \times \Omega_{*,\epsilon}^M)$ for $p \in (1, \infty)$ converges weakly/strongly in the two-scale sense in L^p , if and only if $\mathcal{T}_\epsilon v_\epsilon$ converges weakly/strongly in $L^p((0, T) \times \Sigma \times Z^*)$ to the same limit.
- (ii) The sequence $v_\epsilon \in L^p((0, T) \times N_\epsilon)$ for $p \in (1, \infty)$ converges weakly/strongly in the two-scale sense in L^p , if and only if $\mathcal{T}_\epsilon v_\epsilon$ converges weakly/strongly in $L^p((0, T) \times \Sigma \times N)$ to the same limit.

Proof. This result was obtained for $p = 2$ and weak convergences in [4] for bulk domains. However, the proof can easily be extended to our setting. The strong convergence results follow from the properties of the unfolding operator in Lemma 3.4. \square

The following Lemma shows, that the strong two-scale convergence is sufficient to pass to the limit in the nonlinear terms.

Lemma 3.6. *Let g_ϵ and h_ϵ satisfy the Assumptions (A3) and (A4).*

- (i) Let v_ϵ be a sequence in $L^2((0, T) \times \Omega_{*,\epsilon}^M)$ such that

$$\frac{1}{\sqrt{\epsilon}} \|v_\epsilon\|_{L^2((0, T) \times \Omega_{*,\epsilon}^M)} \leq C,$$

and v_ϵ converges to $v_0 \in L^2((0, T) \times \Sigma \times Z^*)$ strongly in the two-scale sense in L^p for $p \in [1, 2]$. Then it holds that

$$g_\epsilon(v_\epsilon) \rightarrow g(v_0) \quad \text{in the two-scale sense.}$$

- (ii) Let v_ϵ be a sequence in $L^2((0, T) \times N_\epsilon)$ such that

$$\|v_\epsilon\|_{L^2((0, T) \times N_\epsilon)} \leq C,$$

and v_ϵ converges to $v_0 \in L^2((0, T) \times \Sigma \times N)$ strongly in the two-scale sense in L^p on N_ϵ for $p \in [1, 2]$. Then it holds that

$$h_\epsilon(v_\epsilon) \rightarrow h(v_0) \quad \text{in the two-scale sense.}$$

Proof. We only prove the second result, since the first one follows in a similar way. Let $\phi \in L^{p'}((0, T), C(\bar{\Sigma}, C_{\text{per}}(\bar{N})))$, $v^n \in C_0^\infty((0, T) \times \Sigma, C_{\text{per}}(\bar{N}))$ for $n \in \mathbb{N}$ such that $v^n \rightarrow v_0$ in $L^2((0, T) \times \Sigma \times N)$, and we define $v_\epsilon^n(t, x) := v^n(t, \bar{x}, \frac{x}{\epsilon})$.

$$\begin{aligned} \int_0^T \int_{N_\epsilon} h_\epsilon(v_\epsilon) \phi \left(t, \bar{x}, \frac{x}{\epsilon} \right) d\sigma dt &= \int_0^T \int_{N_\epsilon} [h_\epsilon(v_\epsilon) - h_\epsilon(v_\epsilon^n)] \phi \left(t, \bar{x}, \frac{x}{\epsilon} \right) d\sigma dt \\ &\quad + \int_0^T \int_{N_\epsilon} h_\epsilon(v_\epsilon^n) \phi \left(t, \bar{x}, \frac{x}{\epsilon} \right) d\sigma dt =: I_{\epsilon,1} + I_{\epsilon,2}. \end{aligned}$$

For the first term $I_{\epsilon,1}$ we use the Lipschitz continuity of h_ϵ and the properties of the unfolding operator to obtain (with $\mathcal{T}_\epsilon v_\epsilon^n(t, \bar{x}, y) = v^n(t, \epsilon \left[\frac{\bar{x}}{\epsilon} \right] + \epsilon \bar{y}, y)$)

$$\begin{aligned} |I_{\epsilon,1}| &\leq C \|v_\epsilon - v_\epsilon^n\|_{L^p((0,T) \times N_\epsilon)} = C \|\mathcal{T}_\epsilon v_\epsilon - \mathcal{T}_\epsilon v_\epsilon^n\|_{L^p((0,T) \times \Sigma \times N)} \\ &\leq C (\|\mathcal{T}_\epsilon v_\epsilon - v_0\|_{L^p((0,T) \times \Sigma \times N)} + \|v_0 - v^n\|_{L^p((0,T) \times \Sigma \times N)} + \|v^n - \mathcal{T}_\epsilon v_\epsilon^n\|_{L^p((0,T) \times \Sigma \times N)}). \end{aligned}$$

The first term convergence to zero for $\epsilon \rightarrow 0$, due to the strong two-scale convergence of v_ϵ and Lemma 3.5. The second term goes to zero for $n \rightarrow \infty$, and the last term vanishes for $\epsilon \rightarrow 0$, due to the dominated convergence theorem of Lebesgue, since $\mathcal{T}_\epsilon v_\epsilon^n \rightarrow v^n$ almost everywhere in $(0, T) \times \Sigma \times N$.

Let us estimate $I_{\epsilon,2}$:

$$\begin{aligned} I_{\epsilon,2} &= \left[\int_0^T \int_{N_\epsilon} h_\epsilon(v_\epsilon^n) \phi \left(t, \bar{x}, \frac{x}{\epsilon} \right) d\sigma dt - \int_0^T \int_\Sigma \int_N h(v^n) \phi(t, \bar{x}, y) d\sigma_y d\bar{x} dt \right] \\ &\quad + \int_0^T \int_\Sigma \int_N [h(v^n) - h(v_0)] \phi d\sigma_y d\bar{x} dt + \int_0^T \int_\Sigma \int_N h(v_0) \phi d\sigma_y d\bar{x} dt. \end{aligned}$$

The term in the brackets converges to zero for $\epsilon \rightarrow 0$, due to the oscillation lemma, see [3, Lemma 4.3]. The second term vanishes for $n \rightarrow \infty$, due to the Lipschitz continuity of h . This gives the desired result. \square

To establish the strong two-scale convergence of u_ϵ^M we will show the strong convergence in L^p of $\mathcal{T}_\epsilon u_\epsilon^M$. This requires to control the dependence on the time-variable. Since the time-derivative of u_ϵ respectively u_ϵ^M only exists in a weak sense, in fact we have $\partial_t u_\epsilon^M \in L^2((0, T), (\mathcal{H}_{\epsilon,0}^M)')$, it is not obvious in which space $\partial_t \mathcal{T}_\epsilon u_\epsilon^M$ lies and how its norm can be estimated with respect to ϵ . To overcome this problem we use a functional analytical argument. We consider the L^2 -adjoint of \mathcal{T}_ϵ , the so called averaging operator \mathcal{U}_ϵ , to obtain a representation of $\partial_t \mathcal{T}_\epsilon$ via the averaging operator. Therefore, we have to restrict the domain of definition for \mathcal{T}_ϵ and \mathcal{U}_ϵ . This idea was already used in [15] and here we put in a more general framework. First of all, let us give a general functional analytic result:

Lemma 3.7. *Let V, W reflexive, separable Banach-spaces, and Y, X Hilbert-spaces, such that we have the Gelfand-triples*

$$V \hookrightarrow Y \hookrightarrow V', \quad W \hookrightarrow X \hookrightarrow W',$$

with continuous and dense embeddings. Here, we identify Y and X with their dual spaces Y' and X' via the Riesz-representation theorem. Let $A \in \mathcal{L}(Y, X)$ and we denote by $A^ \in \mathcal{L}(X, Y)$ the adjoint operator of A . If $A^*(W) \subset V$ with $\|A^*w\|_V \leq C \|w\|_W$ for all $w \in W$, and $u \in L^2((0, T), Y) \cap H^1((0, T), V')$, then it holds $\partial_t Au \in L^2((0, T), W')$ with*

$$\langle \partial_t Au, w \rangle_{W', W} = \langle \partial_t u, A^*w \rangle_{V', V} \quad \text{for all } w \in W.$$

Here we apply the operator A pointwise to u with respect to $t \in (0, T)$.

Proof. This is just a consequence of the definition of the adjoint operator and the generalized time-derivative. In fact, we have for all $w \in W$ and $\psi \in \mathcal{D}(0, T)$

$$\int_0^T (Au, w)_X \psi'(t) dt = \int_0^T (u, A^*w)_Y \psi'(t) dt = - \int_0^T \langle \partial_t u, A^*w \rangle_{V', V} \psi(t) dt.$$

□

Let us define $\epsilon^{-1}\mathcal{U}_\epsilon$ as the L^2 -adjoint of \mathcal{T}_ϵ , i. e., let

$$\mathcal{U}_\epsilon : L^2((0, T) \times \Sigma \times Z^*) \rightarrow L^2((0, T) \times \Omega_{*, \epsilon}^M),$$

such that

$$(\mathcal{T}_\epsilon v_\epsilon, \phi)_{(0, T) \times \Sigma \times Z^*} = \frac{1}{\epsilon} (v_\epsilon, \mathcal{U}_\epsilon \phi)_{(0, T) \times \Omega_{*, \epsilon}^M},$$

for all $v_\epsilon \in L^2((0, T) \times \Omega_{*, \epsilon}^M)$ and $\phi \in L^2((0, T) \times \Sigma \times Z^*)$. It is easy to check that

$$\mathcal{U}_\epsilon(\phi)(t, x) = \int_Y \phi \left(t, \epsilon \left(\bar{z} + \left[\frac{\bar{x}}{\epsilon} \right] \right), \left(\left\{ \frac{\bar{x}}{\epsilon} \right\}, \frac{x_n}{\epsilon} \right) \right) d\bar{z} \quad \text{for } (t, x) \in (0, T) \times \Omega_{*, \epsilon}^M,$$

and $x = [x] + \{x\}$, but we will not use this explicit formula for $\mathcal{U}_\epsilon(\phi)$.

Corollary 3.8. *For all $\phi \in L^2((0, T) \times \Sigma \times Z^*)$ it holds that*

$$\|\mathcal{U}_\epsilon \phi\|_{L^2((0, T) \times \Omega_{*, \epsilon}^M)} \leq \sqrt{\epsilon} \|\phi\|_{L^2((0, T) \times \Sigma \times Z^*)}.$$

Proof. This follows by a simple duality argument, see also [12, Corollary 2.15] for more details. □

Concerning the regularity of $\mathcal{U}_\epsilon(\phi)$ with respect to the spatial variable, we have that

$$\mathcal{U}_\epsilon : L^2((0, T) \times \Sigma, H^1(Z^*)) \rightarrow L^2(0, T), H^1(\Omega_{*, \epsilon}^M)$$

with

$$\epsilon \nabla \mathcal{U}_\epsilon(\phi) = \mathcal{U}_\epsilon(\nabla_y \phi) \quad \text{for all } \phi \in L^2((0, T) \times \Sigma, H^1(Z^*)). \quad (8)$$

This result uses the fact that Z^* is not touching the lateral boundary of Z and can be shown by similar arguments like in the proof of [15, Proposition 6]. We emphasize that the situation gets more delicate if the channel Z^* touches the lateral boundary of Z and in that case one has to restrict to function spaces with vanishing traces on ∂Z , see also [15].

Next, we apply Lemma 3.7 to obtain a representation of $\partial_t \mathcal{T}_\epsilon u_\epsilon^M$ by means of the $\partial_t u_\epsilon^M$ and \mathcal{U}_ϵ . However, since we have just $\partial_t u_\epsilon^M \in L^2((0, T), (\mathcal{H}_{\epsilon, 0}^M)')$, we have to restrict the operator \mathcal{U}_ϵ . We define

$$\mathcal{H}_0 := \{v \in H^1(Z^*) : v|_{S_\pm^*} = 0\} \subset H^1(Z^*).$$

and consider

$$\mathcal{U}_\epsilon : L^2((0, T) \times \Sigma, \mathcal{H}_0) \rightarrow L^2((0, T), \mathcal{H}_{\epsilon, 0}^M).$$

Proposition 3.9. *Let $v_\epsilon \in L^2((0, T), L^2(\Omega_{*,\epsilon}^M)) \cap H^1((0, T), (\mathcal{H}_{\epsilon,0}^M)')$. Then we have $\mathcal{T}_\epsilon v_\epsilon \in H^1((0, T), L^2(\Sigma, \mathcal{H}_0)')$ with*

$$\langle \partial_t \mathcal{T}_\epsilon v_\epsilon(t), \phi \rangle_{L^2(\Sigma, \mathcal{H}_0)', L^2(\Sigma, \mathcal{H}_0)} = \frac{1}{\epsilon} \langle \partial_t v_\epsilon(t), \mathcal{U}_\epsilon \phi \rangle_{(\mathcal{H}_{\epsilon,0}^M)', \mathcal{H}_{\epsilon,0}^M}$$

for all $\phi \in L^2(\Sigma, \mathcal{H}_0)$ and almost every $t \in (0, T)$. Additionally, we have

$$\|\partial_t \mathcal{T}_\epsilon v_\epsilon\|_{L^2((0, T), L^2(\Sigma, \mathcal{H}_0)')} \leq \frac{1}{\epsilon} \|\partial_t v_\epsilon\|_{(\mathcal{H}_{\epsilon,0}^M)'}. \quad (9)$$

Proof. $\partial_t \mathcal{T}_\epsilon v_\epsilon \in L^2((0, T), L^2(\Sigma, \mathcal{H}_0)')$ follows directly from Lemma 3.7, if we choose $V = \mathcal{H}_{\epsilon,0}^M$, $Y = L^2(\Omega_{*,\epsilon}^M)$, $W = L^2(\Sigma, \mathcal{H}_0)$, $X = L^2(\Sigma \times Z^*)$, and $A = \mathcal{T}_\epsilon$. Then we have $A^* = \epsilon^{-1} \mathcal{U}_\epsilon$, and Lemma 3.7 implies the desired result for the existence of the time-derivative $\partial_t \mathcal{T}_\epsilon v_\epsilon$. For the estimate (9) we choose $\phi \in L^2(\Sigma, \mathcal{H}_0)$ with $\|\phi\|_{L^2(\Sigma, \mathcal{H}_0)} \leq 1$ and obtain

$$\begin{aligned} \langle \partial_t \mathcal{T}_\epsilon v_\epsilon, \phi \rangle_{L^2(\Sigma, \mathcal{H}_0)', L^2(\Sigma, \mathcal{H}_0)} &= \frac{1}{\epsilon} \langle \partial_t v_\epsilon, \mathcal{U}_\epsilon \phi \rangle_{(\mathcal{H}_{\epsilon,0}^M)', \mathcal{H}_{\epsilon,0}^M} \\ &\leq \frac{1}{\epsilon} \|\partial_t v_\epsilon\|_{(\mathcal{H}_{\epsilon,0}^M)'} \|\mathcal{U}_\epsilon \phi\|_{\mathcal{H}_{\epsilon,0}^M} \leq \frac{1}{\epsilon} \|\partial_t v_\epsilon\|_{(\mathcal{H}_{\epsilon,0}^M)'}, \end{aligned}$$

where the last inequality follows from (8) and Corollary 3.8. \square

Lemma 3.10. *For all $\phi_\epsilon \in L^2((0, T) \times \Omega_{*,\epsilon}^M)$, $0 < h \ll 1$, and $\bar{\xi} \in \mathbb{R}^{n-1}$ with $|\bar{\xi}| \ll h$, it holds for ϵ small enough that*

$$\|\mathcal{T}_\epsilon \phi_\epsilon(\cdot, \cdot + \bar{\xi}, \cdot) - \mathcal{T}_\epsilon \phi_\epsilon\|_{L^2((0, T) \times \Sigma_{2h} \times Z)}^2 \leq \frac{1}{\epsilon} \sum_{\bar{j} \in \{0, 1\}^{n-1}} \|\delta_l \phi_\epsilon\|_{L^2((0, T) \times \widehat{\Omega}_{*,\epsilon,h}^M)}^2$$

with $l = l(\epsilon, \bar{\xi}, \bar{j}) = \bar{j} + \left\lceil \frac{\bar{\xi}}{\epsilon} \right\rceil$.

Proof. The idea of the proof can be found in [18, page 709] for a thin layer and can be extended in an obvious way to our setting. \square

In the next theorem we formulate a general strong two-scale compactness result for sequences $v_\epsilon \in L^2((0, T), H^1(\Omega_{*,\epsilon}^M))$ and their traces $v_\epsilon|_{N_\epsilon}$. This result allows us to pass to the limit in the nonlinear terms in (3). Similar ideas have been used in [15, Theorem 7.5], where however, they were carried-out for the sequence of solutions of a microscopic problem in a thin layer with oscillating diffusion coefficients.

Theorem 3.11. *Let $v_\epsilon \in L^2((0, T), H^1(\Omega_{*,\epsilon}^M)) \cap H^1((0, T), (\mathcal{H}_{\epsilon,0}^M)')$ such that*

(i) *we have the estimate*

$$\frac{1}{\epsilon} \|\partial_t v_\epsilon\|_{L^2((0, T), (\mathcal{H}_{\epsilon,0}^M)')} + \frac{1}{\sqrt{\epsilon}} \|v_\epsilon\|_{L^2((0, T) \times \Omega_{*,\epsilon}^M)} + \sqrt{\epsilon} \|\nabla v_\epsilon\|_{L^2((0, T) \times \Omega_{*,\epsilon}^M)} \leq C,$$

(ii) for all $0 < h \ll 1$ and $l \in \mathbb{Z}^{n-1} \times \{0\}$ and $|\epsilon l| \ll h$ it holds that

$$\frac{1}{\sqrt{\epsilon}} \|\delta v_\epsilon\|_{L^2((0,T) \times \widehat{\Omega}_{*,\epsilon,h}^M)} + \sqrt{\epsilon} \|\nabla \delta v_\epsilon\|_{L^2((0,T) \times \widehat{\Omega}_{*,\epsilon,h}^M)} \xrightarrow{\epsilon l \rightarrow 0} 0.$$

Then there exists $v_0 \in L^2((0,T) \times \Sigma, H^1(Z^*))$ such that up to a subsequence it holds for $p \in [1, 2)$ and $\beta \in (\frac{1}{2}, 1)$

$$\begin{aligned} v_\epsilon &\rightarrow v_0 && \text{in the two-scale sense,} \\ \epsilon \nabla v_\epsilon &\rightarrow \nabla_y v_0 && \text{in the two-scale sense,} \\ \mathcal{T}_\epsilon v_\epsilon &\rightarrow v_0 && \text{strongly in } L^p(\Sigma, L^2((0,T), H^\beta(Z^*))). \end{aligned}$$

Especially, we have $v_\epsilon \rightarrow v_0$ strongly in the two-scale sense in L^p and $v_\epsilon|_{N_\epsilon} \rightarrow v_0|_N$ strongly in the two-scale sense on N_ϵ in L^p .

Proof. The weak two-scale convergences of v_ϵ and $\epsilon \nabla v_\epsilon$ follow directly from Lemma 3.2 and the estimates in (i). The strong two-scale convergence of v_ϵ and $v_\epsilon|_{N_\epsilon}$ in L^p follow from the strong convergence of $\mathcal{T}_\epsilon v_\epsilon$ in $L^p(\Sigma, L^2((0,T), H^\beta(Z^*)))$ by the imbedding $H^\beta(Z^*) \hookrightarrow L^2(N)$, and Lemma 3.5. So it remains to prove the strong convergence of $\mathcal{T}_\epsilon v_\epsilon$. Therefore, we use the Kolmogorov-type compactness result [13, Corollary 2.5] for the sequence

$$\mathcal{T}_\epsilon v_\epsilon \in L^2(\Sigma, L^2((0,T), H^1(Z^*))) \hookrightarrow L^p(\Sigma, L^2((0,T), H^\beta(Z^*))).$$

We have to check the following three conditions:

(K1) For every $A \subset \Sigma$ measurable, the sequence

$$v_\epsilon^A(t, y) := \int_A \mathcal{T}_\epsilon v_\epsilon d\bar{x}$$

is relatively compact in $L^2((0,T), H^\beta(Z^*))$.

(K2) For $0 < h \ll 1$ and $\bar{\xi} \in \mathbb{R}^{n-1}$ with $|\bar{\xi}| < h$ it holds that

$$\sup_\epsilon \|\mathcal{T}_\epsilon v_\epsilon(\cdot, \cdot + \bar{\xi}, \cdot) - \mathcal{T}_\epsilon v_\epsilon\|_{L^p(\Sigma_h, L^2((0,T), H^\beta(Z^*)))} \xrightarrow{\bar{\xi} \rightarrow 0} 0.$$

(K3) For $0 < h \ll 1$ it holds that

$$\sup_\epsilon \|\mathcal{T}_\epsilon v_\epsilon\|_{L^p(\Sigma \setminus \Sigma_h, L^2((0,T), H^\beta(Z^*)))} \xrightarrow{h \rightarrow 0} 0.$$

Let us start with (K3). We obtain from the Hölder-inequality since $p < 2$ that

$$\|\mathcal{T}_\epsilon v_\epsilon\|_{L^p(\Sigma \setminus \Sigma_h, L^2((0,T), H^\beta(Z^*)))} \leq C|h|^{\frac{2-p}{2p}} \|\mathcal{T}_\epsilon v_\epsilon\|_{L^2(\Sigma \setminus \Sigma_h, L^2((0,T), H^1(Z^*)))} \leq C|h|^{\frac{2-p}{2p}},$$

where the last inequality follows from (i) and the properties of the unfolding operator from Lemma 3.4. This gives (K3). To prove (K1) we first notice that we have

$$v_\epsilon^A \in L^2((0, T), H^1(Z^*)) \cap H^1((0, T), \mathcal{H}'_0)$$

with

$$\langle \partial_t v_\epsilon^A, \phi \rangle_{\mathcal{H}'_0, \mathcal{H}_0} = \langle \partial_t \mathcal{T}_\epsilon v_\epsilon, \chi_A(\cdot, \bar{x}) \phi(\cdot, y) \rangle_{L^2(\Sigma, \mathcal{H}_0)', L^2(\Sigma, \mathcal{H}_0)}$$

for all $\phi \in \mathcal{H}_0$. Obviously, due to (i) and Lemma 3.4, the sequence v_ϵ^A is bounded in $L^2((0, T), H^1(Z^*))$. Proposition 3.9 and the estimate of $\partial_t v_\epsilon$ in (i) imply the boundedness of $\partial_t \mathcal{T}_\epsilon v_\epsilon^A$ in $L^2((0, T), L^2(\Sigma, \mathcal{H}_0)')$. Since $H^1(Z^*) \hookrightarrow H^\beta(Z^*)$ is compact for $\frac{1}{2} < \beta < 1$ and $H^\beta(Z^*) \hookrightarrow \mathcal{H}'_0$ is continuous, the Aubin-Lions lemma, see [16], implies (K1). Now, we choose $0 < h \ll 1$ and obtain for $|\bar{\xi}| < h$ with Lemma 3.10

$$\begin{aligned} & \|\mathcal{T}_\epsilon v_\epsilon(\cdot, \cdot + \bar{\xi}, \cdot) - \mathcal{T}_\epsilon v_\epsilon\|_{L^2(\Sigma_{2h}, L^2((0, T), H^1(Z^*)))} \\ & \leq C \sum_{\bar{j} \in \{0, 1\}^{n-1}} \left(\frac{1}{\sqrt{\epsilon}} \|\delta v_\epsilon\|_{L^2((0, T) \times \widehat{\Omega}_{*, \epsilon, h}^M)} + \sqrt{\epsilon} \|\nabla \delta v_\epsilon\|_{L^2((0, T) \times \widehat{\Omega}_{*, \epsilon, h}^M)} \right), \end{aligned}$$

for $l = \bar{j} + \left\lceil \frac{\bar{\xi}}{\epsilon} \right\rceil$. Due to assumption (ii), the right-hand side converges to zero for $\epsilon, \bar{\xi} \rightarrow 0$. The uniform convergence with respect to ϵ in (K2) is obtained by the convergence of the shifts to 0 for every fixed ϵ . The result follows from [13, Corollary 2.5]. \square

4 Derivation of the macroscopic model

The aim of this section is the derivation of the macroscopic model for $\epsilon \rightarrow 0$ with the methods developed in Section 3, which are based on the a priori estimates for the microscopic solutions established in Section 2.3. First of all we give a convergence result for the sequences in the bulk-domains:

Proposition 4.1. *Let u_ϵ be the sequence of solutions of the microscopic problem (1). Then there exists $u_0^\pm \in L^2((0, T), H^1(\Omega^\pm))$ such that up to a subsequence*

$$\begin{aligned} \chi_{\Omega_\epsilon^\pm} u_\epsilon^\pm &\rightarrow u_0^\pm && \text{strongly in } L^2((0, T) \times \Omega^\pm), \\ u_\epsilon^\pm(\cdot, \bar{x}, \pm\epsilon) &\rightarrow u_0^\pm|_\Sigma && \text{strongly in } L^2((0, T) \times \Sigma), \\ \chi_{\Omega_\epsilon^\pm} \nabla u_\epsilon^\pm &\rightharpoonup \nabla u_0^\pm && \text{weakly in } L^2((0, T) \times \Omega^\pm). \end{aligned}$$

Proof. This result was shown in [18, Proposition 2.1 and 2.2] for time-derivatives $\partial_t u_\epsilon^\pm$ in $L^2((0, T) \times \Omega_\epsilon^\pm)$. In our case, we have that $\partial_t u_\epsilon^\pm$ are functionals on the space

$$\{\phi_\epsilon^\pm \in H^1(\Omega_\epsilon^\pm) : \phi_\epsilon^\pm = 0 \text{ on } \Sigma \times \{\pm\epsilon\}\}.$$

However, the methods from [18] can easily be extended to our setting and we skip the details. \square

In the next theorem we give the convergence results for the sequences in the channels.

Theorem 4.2. *Let u_ϵ be the sequence of solutions of the microscopic problem (1). Then there exists $u_0^M \in L^2((0, T) \times \Sigma, H^1(Z^*))$ such that up to a subsequence it holds for $p \in [1, 2)$*

$$\begin{aligned} u_\epsilon^M &\rightarrow u_0^M && \text{strongly in the two-scale sense in } L^p, \\ \epsilon \nabla u_\epsilon^M &\rightarrow \nabla_y u_0^M && \text{in the two-scale sense,} \\ u_\epsilon^M|_{N_\epsilon} &\rightarrow u_0^M|_N && \text{strongly in the two-scale sense in } L^p. \end{aligned}$$

Additionally, u_ϵ^M and $u_\epsilon^M|_{N_\epsilon}$ also converges weakly in the two-scale sense in L^2 . Further, u_0^M fulfills the following boundary condition on the top and the bottom of Z^* :

$$u_0^M(t, \bar{x}, y) = u_0^\pm(t, \bar{x}, 0) \quad \text{for almost every } (t, \bar{x}, y) \in (0, T) \times \Sigma \times S_*^\pm, \quad (10)$$

i. e., u_0^M is constant on S_*^\pm .

Proof. For the convergence results we only have to check the conditions in Theorem 3.11. Condition (i) is just Lemma 2.3. For (ii) we use inequality (7) from Lemma 2.4. The initial terms in (7) tend to 0 for $\epsilon l \rightarrow 0$, due to the standard Kolmogorov-compactness result, see for example [5] and the Assumption (A5). The last term on the right-hand side in (7) goes to 0, because of the strong convergence of u_ϵ^\pm from Proposition 4.1 and again the Kolmogorov-compactness result.

It remains to show the equation on S_*^\pm . Therefore we choose functions $\phi \in C^\infty((0, T) \times \Sigma \times \overline{Z^*})^n$ such that $\phi(t, x, \cdot)$ has compact support in $Z^* \cup S_*^+ \cup S_*^-$, and extend this function by zero to Z and then Y -periodically in \bar{y} -direction. From the two-scale results for u_ϵ^M and the strong convergence of $u_\epsilon^\pm(\cdot, \bar{x}, \pm\epsilon)$ we obtain by integration by parts

$$\begin{aligned} \int_0^T \int_\Sigma \int_{Z^*} \nabla_y u_0^M \phi dy d\bar{x} dt &= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \int_0^T \int_{\Omega_{*,\epsilon}^M} \epsilon \nabla u_\epsilon^M \phi \left(t, \bar{x}, \frac{x}{\epsilon} \right) dx dt \\ &= \lim_{\epsilon \rightarrow 0} \left\{ -\frac{1}{\epsilon} \int_0^T \int_{\Omega_{*,\epsilon}^M} u_\epsilon^M \nabla_y \cdot \phi \left(t, \bar{x}, \frac{x}{\epsilon} \right) dx dt + \sum_{\pm} \int_0^T \int_{S_{*,\epsilon}^\pm} u_\epsilon^\pm \phi \left(t, \bar{x}, \frac{x}{\epsilon} \right) \cdot \nu d\sigma dt \right\} \\ &= - \int_0^T \int_\Sigma \int_{Z^*} u_0^M \nabla_y \cdot \phi dy d\bar{x} dt + \sum_{\pm} \int_0^T \int_\Sigma \int_{S_*^\pm} u_0^\pm \phi \cdot \nu d\sigma_y d\bar{x} dt. \end{aligned}$$

Using again the integration by parts formula gives us the desired result. \square

Next, we will state the macroscopic model, which is solved by the limit function $u_0 := (u_0^+, u_0^M, u_0^-)$ from Proposition 4.1 and Theorem 4.2. The appropriate solution space is $L^2((0, T), \mathcal{H})$, where \mathcal{H} is given by

$$\begin{aligned} \mathcal{H} := \{ u = (u^+, u^M, u^-) \in H^1(\Omega^+) \times L^2(\Sigma, H^1(Z^*)) \times H^1(\Omega^-) : \\ u^\pm|_\Sigma = u^M|_{S_*^\pm} \text{ on } \Sigma \times S_*^\pm \}, \end{aligned}$$

with the inner product

$$(u, \phi)_{\mathcal{H}} = \sum_{\pm} (u^{\pm}, \phi^{\pm})_{H^1(\Omega^{\pm})} + (u^M, \phi^M)_{\Sigma \times Z^*} + (\nabla_y u^M, \nabla_y \phi^M)_{\Sigma \times Z^*}. \quad (11)$$

For the derivation of the macroscopic problem in the limit $\epsilon \rightarrow 0$, we have to exploit the convergence properties of the microscopic solutions u_{ϵ} . Especially this requires to choose in the variational equation (3) test-functions adapted to the structure of the transmission problem and whose restrictions to the channel domain is admissible for the definition of two-scale convergence. In fact, in general, for $u \in \mathcal{H}$ the function $u^M(\bar{x}, \frac{x}{\epsilon})$ is not well-defined in $\Omega_{*,\epsilon}^M$ and N_{ϵ} . Therefore, we consider the subspace \mathcal{H}^{∞} of \mathcal{H} of smooth functions

$$\mathcal{H}^{\infty} := C^{\infty}(\overline{\Omega^+}) \times C_0^{\infty}(\Sigma, C^{\infty}(\overline{Z^*})) \times C^{\infty}(\overline{\Omega^-}) \cap \mathcal{H}.$$

The following density result holds.

Proposition 4.3. *The space \mathcal{H}^{∞} is dense in \mathcal{H} with respect to the norm induced by (11).*

Proof. We have $\mathcal{H} = \overline{\mathcal{H}^{\infty}} \oplus \overline{\mathcal{H}^{\infty}}^{\perp}$. Hence, for $u = (u^+, u^M, u^-) \in \overline{\mathcal{H}^{\infty}}^{\perp}$ we have

$$(u^M, \phi^M)_{\Sigma \times Z^*} + (\nabla_y u^M, \nabla_y \phi^M)_{\Sigma \times Z^*} = 0$$

for all $\phi \in C_0^{\infty}(\Sigma, C^{\infty}(\overline{Z^*}))$ with $\phi^M = 0$ on S_*^{\pm} . This implies $\Delta_y u^M = u^M$, $\nabla_y u^M \in L^2(\Sigma, H(\operatorname{div}, Z^*))$, and $\nabla_y u^M \cdot \nu = 0$ in the distributional sense, i. e., for all $\phi^M \in C_0^{\infty}(\Sigma, C^{\infty}(\overline{Z^*}))$ with $\phi^M = 0$ on S_*^{\pm} it holds that

$$\langle \nabla_y u^M \cdot \nu, \phi^M \rangle_{L^2(\Sigma, H^{-\frac{1}{2}}(\partial Z^*)), L^2(\Sigma, H^{\frac{1}{2}}(\partial Z^*))} = 0. \quad (12)$$

By density, see [2, Theorem 3.1], equation (12) is also true for all $\phi^M \in L^2(\Sigma, H^1(Z^*))$ with $\phi^M = 0$ on S_*^{\pm} . Since $u \in \overline{\mathcal{H}^{\infty}}^{\perp}$, we obtain for arbitrary $\phi = (\phi^+, \phi^M, \phi^-) \in \mathcal{H}^{\infty}$ by the divergence theorem

$$\langle \nabla_y u^M \cdot \nu, \phi^M \rangle_{L^2(\Sigma, H^{-\frac{1}{2}}(\partial Z^*)), L^2(\Sigma, H^{\frac{1}{2}}(\partial Z^*))} = - \sum_{\pm} (u^{\pm}, \phi^{\pm})_{H^1(\Omega^{\pm})}. \quad (13)$$

Due to (12) we have

$$\begin{aligned} & - \sum_{\pm} (u^{\pm}, \phi^{\pm})_{H^1(\Omega^{\pm})} \\ &= \left\langle \nabla_y u^M \cdot \nu, \frac{y_n + 1}{2} \phi^+(\bar{x}, 0) - \frac{y_n - 1}{2} \phi^-(\bar{x}, 0) \right\rangle_{L^2(\Sigma, H^{-\frac{1}{2}}(\partial Z^*)), L^2(\Sigma, H^{\frac{1}{2}}(\partial Z^*))} \end{aligned}$$

Hence, by density of $C^{\infty}(\overline{\Omega^{\pm}})$ in $H^1(\Omega^{\pm})$, the equation above holds for all $\phi^{\pm} \in H^1(\Omega^{\pm})$, and using again (12), we get (13) for all $\phi \in \mathcal{H}$ and therefore $u = 0$. \square

Theorem 4.4. *The limit function $u_0 = (u_0^+, u_0^M, u_0^-)$ from Proposition 4.1 and Theorem 4.2 is the unique weak solution of the following problem:*

$$u_0 \in L^2((0, T), \mathcal{H}) \cap H^1((0, T), \mathcal{H}')$$

and

$$\begin{aligned} \partial_t u_0^\pm - D^\pm \Delta u_0^\pm &= f^\pm(u_0^\pm) && \text{in } (0, T) \times \Omega^\pm, \\ -D^\pm \nabla u_0^\pm \cdot \nu^\pm &= 0 && \text{on } (0, T) \times \partial\Omega^\pm \setminus \Sigma, \\ u_0^\pm(0) &= u_i^\pm && \text{in } \Omega^\pm, \end{aligned}$$

with the interface conditions

$$\begin{aligned} u_0^\pm|_\Sigma &= u_0^M|_{S_*^\pm} && \text{on } (0, T) \times \Sigma \times S_*^\pm, \\ [D^\pm \nabla u_0^\pm \cdot \nu^\pm]_\Sigma &= \sum_{\pm} \int_{S_*^\pm} D^M \nabla_y u_0^M \cdot \nu^M d\sigma && \text{on } (0, T) \times \Sigma, \end{aligned}$$

with the jump of the normal fluxes across the interface $[D^\pm \nabla u_0^\pm \cdot \nu^\pm]_\Sigma = (D^+ \nabla u_0^+ - D^- \nabla u_0^-) \cdot \nu^+$, ν^M is the outer-unit normal on ∂Z^* , and u_0^M solves the local cell problem

$$\begin{aligned} \partial_t u_0^M - \nabla_y (D^M \nabla_y u_0^M) &= g(u_0^M) && \text{in } (0, T) \times \Sigma \times Z^*, \\ -D^M \nabla_y u_0^M \cdot \nu &= h(u_0^M) && \text{on } (0, T) \times \Sigma \times N, \\ u_0^M &= u_i^M && \text{in } \Sigma \times Z^*. \end{aligned}$$

Proof. From Proposition 4.1 and Theorem 4.2 immediately follows that $u_0 \in L^2((0, T), \mathcal{H})$. Let $\phi \in C_0^\infty([0, T], \mathcal{H}^\infty)$ and choose as a test-function in (3)

$$\phi_\epsilon(t, x) := \begin{cases} \phi^+(t, \bar{x}, x_n - \epsilon) & \text{for } x \in \Omega_\epsilon^+, \\ \phi^M(t, \bar{x}, \frac{x}{\epsilon}) & \text{for } x \in \Omega_{*,\epsilon}^M, \\ \phi^-(t, \bar{x}, x_n + \epsilon) & \text{for } x \in \Omega_\epsilon^-. \end{cases}$$

Integration with respect to time, integration by parts in time, and Proposition 4.1 and Theorem 4.2 together with Lemma 3.6 imply for $\epsilon \rightarrow 0$

$$\begin{aligned} &\sum_{\pm} \{ -(u_0^\pm, \partial_t \phi^\pm)_{(0,T) \times \Omega^\pm} + (D^\pm \nabla u_0^\pm, \nabla \phi^\pm)_{(0,T) \times \Omega^\pm} \} \\ &\quad - (u_0^M, \phi^M)_{(0,T) \times \Sigma \times Z^*} + (D^M \nabla_y u_0^M, \nabla_y \phi^M)_{(0,T) \times \Sigma \times Z^*} \\ &= \sum_{\pm} \{ (f^\pm(u_0^\pm), \phi^\pm)_{(0,T) \times \Omega^\pm} + (u_i^\pm, \phi^\pm(0))_{\Omega^\pm} \} \\ &\quad + (g(u_0^M), \phi^M)_{(0,T) \times \Sigma \times Z^*} - (h(u_0^M), \phi^M)_{(0,T) \times \Sigma \times N} + (u_i^M, \phi^M(0))_{\Sigma \times Z^*}. \end{aligned}$$

By density, see Proposition 4.3, this equation also holds for all $\phi \in C_0^\infty([0, T], \mathcal{H})$. This, together with the relation (10) is just the weak formulation of the macroscopic problem in the Theorem. Especially, this equation implies $\partial_t u_0 \in L^2((0, T), \mathcal{H}')$ and $u_0(0) = (u_i^+, u_i^M, u_i^-)$. Uniqueness follows by standard arguments. \square

Acknowledgements

The first author was supported by the SCIDATOS project, funded by the Klaus Tschira Foundation (Grant number 00.0277.2015).

References

- [1] Y. Amirat, O. Bodart, G. A. Chechkin, and A. L. Piatnitski. Asymptotics of a spectral-sieve problem. *Journal of Mathematical Analysis and Applications*, 435(2):1652–1671, 2016.
- [2] J.-M. E. Bernard. Density results in Sobolev spaces whose elements vanish on a part of the boundary. *Chin. Ann. Math.*, 32B(6):823–846, 2011.
- [3] A. Bhattacharya, M. Gahn, and M. Neuss-Radu. Effective transmission conditions for reaction-diffusion processes in domains separated by thin channels. *Preprint arXiv:2003.11946*, 2020.
- [4] A. Bourgeat, S. Luckhaus, and A. Mikelić. Convergence of the homogenization process for a double-porosity model of immiscible two-phase flow. *SIAM J. Math. Anal.*, 27:1520–1543, 1996.
- [5] H. Brezis. *Functional Analysis, Sobolev Spaces and Partial Differential Equations*. Springer-Verlag, New York, 2010.
- [6] D. Cioranescu, A. Dalhamian, and G. Griso. Periodic unfolding and homogenization. *C. R. Acad. Sci. Paris Sér. 1*, 335:99–104, 2002.
- [7] C. Conca. Étude d’un fluid traversant une paroi perforée I. Comportement limite près de la paroi. *J. Math. pures et appl.*, 66:1–43, 1987.
- [8] C. Conca. Étude d’un fluid traversant une paroi perforée II. Comportement limite loin de la paroi. *J. Math. pures et appl.*, 66:45–69, 1987.
- [9] T. Del Vecchio. The thick neumann’s sieve. *Annali di Matematica Pura ed Applicata*, 147(1):363–402, 1987.
- [10] R. R. Gadyly’shin, A. L. Piatnitskii, and G. A. Chechkin. On the asymptotic behaviour of eigenvalues of a boundary-value problem in a planar domain of steklov sieve type. *Izvestiya: Mathematics*, 82(6):1108, 2018.
- [11] M. Gahn. Homogenisierung von Reaktions-Diffusionsgleichungen in Gebieten, die durch ein Netzwerk von Kanälen getrennt sind. *Diploma Thesis (Supervisor: W. Jäger, M. Neuss-Radu)*, University of Heidelberg, 2012.
- [12] M. Gahn. *Derivation of effective models for reaction-diffusion processes in multi-component media*. PhD thesis (Friedrich-Alexander-Universität Erlangen-Nürnberg); Shaker Verlag, 2017.

- [13] M. Gahn and M. Neuss-Radu. A characterization of relatively compact sets in $L^p(\Omega, B)$. *Stud. Univ. Babeş-Bolyai Math.*, 61(3):279–290, 2016.
- [14] M. Gahn, M. Neuss-Radu, and P. Knabner. Derivation of effective transmission conditions for domains separated by a membrane for different scaling of membrane diffusivity. *Discrete & Continuous Dynamical Systems-Series S*, 10(4):773–797, 2017.
- [15] M. Gahn, M. Neuss-Radu, and P. Knabner. Effective interface conditions for processes through thin heterogeneous layers with nonlinear transmission at the microscopic bulk-layer interface. *Networks and Heterogeneous Media*, 13(4):609–640, 2018.
- [16] J. L. Lions. *Quelques méthodes de résolution des problèmes aux limites non linéaires*. Dunod, Paris, 1969.
- [17] M. Neuss-Radu and W. Jäger. Homogenization of thin porous layers and applications to ion transport through channels of biological membranes. *Oberwolfach Reports*, 49:2809–2812, 2005.
- [18] M. Neuss-Radu and W. Jäger. Effective transmission conditions for reaction-diffusion processes in domains separated by an interface. *SIAM J. Math. Anal.*, 39:687–720, 2007.
- [19] D. Onofrei. The unfolding operator near a hyperplane and its applications to the Neumann sieve model. *Advances in Mathematical Sciences and Applications*, 16(1):239–258, 2006.
- [20] E. Sanchez-Palencia. Boundary value problems in domains containing perforated walls, séminaire college de france. *Research Notes in Mathematics*, 70:309–325, 1982.
- [21] T. A. Shaposhnikova. Homogenization of the Neumann problem in a domain a part of which is a set of channels. *Differential Equations*, 37(9):1315–1323, 2001.
- [22] V. V. Yablokov. The problem on homogenization of solutions of second order elliptic equations in the domain whose part is a union of thin cylinders. *Vestnik Moskovskogo Universiteta. Seriya 1. Matematika. Mekhanika*, 2:15–21, 2004.