

# Spectral Graph Clustering via the Expectation-Solution Algorithm

Zachary M. Pisano\*

Joshua S. Agterberg<sup>†</sup>

Carey E. Priebe<sup>‡</sup>

and

Daniel Q. Naiman<sup>§</sup>

Department of Applied Mathematics and Statistics  
Johns Hopkins University

March 31, 2020

## Abstract

The stochastic blockmodel (SBM) models the connectivity within and between disjoint subsets of nodes in networks. Prior work demonstrated that the rows of an SBM's adjacency spectral embedding (ASE) and Laplacian spectral embedding (LSE) both converge in law to Gaussian mixtures where the components are curved exponential families. Maximum likelihood estimation via the Expectation-Maximization (EM) algorithm for a full Gaussian mixture model (GMM) can then perform the task of clustering graph nodes, albeit without appealing to the components' curvature. Noting that EM is a special case of the Expectation-Solution (ES) algorithm, we propose two ES algorithms that allow us to take full advantage of these curved structures. After presenting the ES algorithm for the general curved-Gaussian mixture, we develop those corresponding to the ASE and LSE limiting distributions. Simulating from artificial SBMs and a brain connectome SBM reveals that clustering graph nodes via our ES algorithms improves upon that of EM for a full GMM.

*Keywords:* EM algorithm, random graph, curved exponential family, estimating equations, mixture model

---

\*Zachary M. Pisano is Candidate for a PhD in Statistics, Whitehead Hall, 3400 North Charles Street, Baltimore, MD 21218 (e-mail: zpisano1@jhu.edu).

<sup>†</sup>Joshua S. Agterberg is Candidate for a PhD in Statistics, Whitehead Hall, 3400 North Charles Street, Baltimore, MD 21218.

<sup>‡</sup>Carey E. Priebe is Professor of Statistics, Clark Hall, 3400 North Charles Street, Baltimore, MD 21218

<sup>§</sup>Daniel Q. Naiman is Professor of Statistics, Whitehead Hall, 3400 North Charles Street, Baltimore, MD 21218. This work is partially supported by the D3M program of the Defense Advanced Research Projects Agency (DARPA).

# 1 Introduction

Statistical inference on graphs is a burgeoning field of study in statistics with applications in neuroscience (Lyzinski et al., 2017 and Priebe et al., 2019) and social networks (Karrer and Newman, 2011), among other areas of research. Given a random graph  $G$  on  $n$  vertices generated by some model  $F$ , a typical procedure is to embed its adjacency matrix  $\mathbf{A} \in \{0, 1\}^{n \times n}$  into a lower dimensional space  $\mathbb{R}^d$  as a collection of  $n$  points. Commonly chosen embeddings include the *adjacency spectral embedding* (ASE) and *Laplacian spectral embedding* (LSE), obtained via the truncated eigendecomposition of  $\mathbf{A}$  and its normalized Laplacian  $\mathcal{L}(\mathbf{A})$ . From here one may seek to perform the task of clustering these points and — by extension — their corresponding vertices.

The spectral graph clustering problem has been extensively studied for settings in which the graph (or graphs) on hand are posited to have been generated by a stochastic block-model (SBM) (Holland et al., 1983), with many results regarding consistent recovery of the block assignments being known (Fishkind et al., 2013). Athreya et al. (2016) and Tang and Priebe (2018) showed that the distribution of the points of the ASE and LSE both converge to curved normal mixture distributions. Moreover, they demonstrated that clustering via the Expectation-Maximization (EM) algorithm (Dempster et al., 1977) for a Gaussian mixture model (GMM) performs better than doing so via the  $K$ -Means algorithm. However, their implementation of the EM algorithm failed to take into account the curved structure of the mixture’s component distributions, and may have therefore resulted in an increased number of clustering errors.

This paper seeks to improve upon their results by introducing an Expectation-Solution (ES) algorithm (Elashoff and Ryan, 2004) that makes full use of the ASE and LSE limiting distributions’ curved-normal structure. We accomplish this by noting that the complete-data likelihood equations from the EM algorithm may be rethought of as complete-data estimating equations used by ES. Since each component of a curved-Gaussian mixture model (C-GMM) encodes its variance as a function of the component means and potentially the mixing proportions, the estimating equations corresponding to the component variances are rendered superfluous. This leads us to an iterative scheme where the com-

ponent means and mixing proportions are updated as in EM for a full GMM, but the component variances are updated by simply plugging in the new means and proportions into the variance functions.

To the best of our knowledge this paper represents the first time the ES algorithm has been utilized for a C-GMM, and owing to its relative simplicity we cannot help but recommend it as a novel tool for the practitioner’s toolbox, one that is not only limited in its usefulness to the spectral clustering problem. We structure the rest of the paper as follows: In Section 2 we review the setting and background for random dot product graphs and SBMs. In Section 3 we present the ES algorithm for the general incomplete-data setting, a generic C-GMM, and for both the ASE and LSE limiting distributions. In Sections 4 and 5 we present and discuss the results of our algorithm for simulated data from artificial SBMs and a brain connectome SBM.

## 1.1 Notation

Except where otherwise specified we use emboldened capital letters such as  $\mathbf{A}$ ,  $\mathbf{B}$ ,  $\mathbf{X}$ , etc. to denote matrices, un-emboldened capital letters with one indexed subscript to denote a row of the corresponding matrix as a column vector, as in  $A_i$ ,  $B_j$ ,  $X_k$ . Emboldened lowercase Greek letters  $\boldsymbol{\pi}$ ,  $\boldsymbol{\tau}$  are vectors. We take  $\boldsymbol{\Delta}^K := \{(\pi_1, \pi_2, \dots, \pi_K) \in \mathbb{R}_{\geq 0}^K \mid \sum_{k=1}^K \pi_k = 1\}$  to be the unit simplex in  $\mathbb{R}^K$ , and  $\delta(\cdot)$  to be the probability distribution assigning point mass to its argument. The vector of all ones and all zeros in  $\mathbb{R}^n$  are given by  $\mathbf{1}_n$  and  $\mathbf{0}_n$ , respectively. We omit subscripts where the dimension is obvious.

Given a real symmetric matrix  $\mathbf{M}$  we write its spectral decomposition as

$$\mathbf{M} = \mathbf{U}_M \mathbf{D}_M \mathbf{U}_M^\top,$$

where  $\mathbf{U}_M$  is a unitary matrix and  $\mathbf{D}_M = \text{diag}(\lambda_1^{(M)}, \lambda_2^{(M)}, \dots, \lambda_n^{(M)})$  such that  $\lambda_1^{(M)} \geq \lambda_2^{(M)} \geq \dots \geq \lambda_n^{(M)}$  are the ordered eigenvalues of  $\mathbf{M}$ . We define the normalized Laplacian of  $\mathbf{M}$  as  $\mathcal{L}(\mathbf{M}) = (\text{diag}(\mathbf{M}\mathbf{1}_n)^{-\frac{1}{2}})\mathbf{M}(\text{diag}(\mathbf{M}\mathbf{1}_n)^{-\frac{1}{2}})$ , with spectral decomposition  $\tilde{\mathbf{U}}_M \tilde{\mathbf{D}}_M \tilde{\mathbf{U}}_M^\top$ .

We use the shorthand

$$\{K\text{-means}, EM, ES\} \circ \{ASE, LSE\} \quad (1)$$

to refer to each clustering method; e.g.,  $EM \circ ASE$  is to be read as “EM for the ASE.” Finally, in sections where we are explicitly concerned with GMMs, any reference to the EM algorithm is understood to refer to the EM algorithm for a full GMM as outlined in Fraley and Raftery (2002).

## 2 Background and Setting

The  $K$ -block SBM encodes the probabilistic connectivity within and between disjoint subsets of graph nodes. For  $K \geq 2$ ,  $\boldsymbol{\pi} \in \boldsymbol{\Delta}^K$ , and  $\mathbf{B} \in (0, 1)^{K \times K}$  a rank- $d$  symmetric matrix with distinct rows, we write  $(\mathbf{A}, \boldsymbol{\tau}) \sim \text{SBM}(n, \mathbf{B}, \boldsymbol{\pi})$  with sparsity factor  $\rho_n \in (0, 1]$  provided the following:

$$\begin{aligned} \boldsymbol{\tau} &:= (\tau_1, \dots, \tau_n) \\ \tau_i &\stackrel{i.i.d.}{\sim} \text{Cate}(\boldsymbol{\pi}) \\ A_{ij} | \boldsymbol{\tau} &\stackrel{ind.}{\sim} \text{Bern}(\rho_n B_{\tau_i \tau_j}), i < j \\ A_{ij} &= A_{ji} \\ \text{diag}(\mathbf{A}) &= \mathbf{0}_n. \end{aligned}$$

If only  $\mathbf{A}$  is observed, we write  $\mathbf{A} \sim \text{SBM}(n, \mathbf{B}, \boldsymbol{\pi})$ .

The sparsity factor  $\rho_n$  indexes a sequence of models where the edge probabilities change with  $n$ . We mention the sparsity factor purely for the sake of completeness, but common assumptions include taking  $\rho_n \equiv 1$  for all  $n$  or  $\rho_n \rightarrow 0$  such that  $\rho_n > \log^4 n/n$ . The former assumption is equivalent to the assumption that there exists  $c > 0$  such that  $\rho_n \rightarrow c$ ; Tang and Priebe (2018) used the latter assumption to establish concentration in spectral norm of  $\mathbf{A}$  and  $\mathcal{L}(\mathbf{A})$  around  $\rho_n \mathbf{P}$  and  $\mathcal{L}(\mathbf{P})$  (where  $\mathbf{P}$  is discussed below). As we are primarily concerned with SBM settings in which the sparsity factor is identically 1 we suppress it throughout the remainder of the paper.

The SBM is a special case of the random dot product graph (RDPG) (Sussman et al., 2012), which encodes the probability that two nodes in a random graph share an edge as defined by the  $n \times n$  symmetric edge probability matrix  $\mathbf{P}$  such that  $A_{ij} \stackrel{\text{ind.}}{\sim} \text{Bern}(P_{ij})$  for  $i \leq j$  (Young and Scheinerman, 2007). It is mathematically convenient to allow for self-loops, as that allows us to rewrite  $\mathbf{P} = \mathbf{X}\mathbf{X}^\top$ , where  $\mathbf{X} \in \mathbb{R}^{n \times \text{rank}(\mathbf{P})}$  such that the row magnitudes are at most 1 and the dot product between any two rows falls in the unit interval. We refer to the rows of  $\mathbf{X}$  as the *latent positions* of the random graph model. Note that the latent positions are inherently unidentifiable, since for any  $d \times d$  orthogonal matrix  $\mathbf{W}$  we have that  $\mathbf{X}\mathbf{X}^\top = (\mathbf{X}\mathbf{W})(\mathbf{X}\mathbf{W})^\top$ .

For a given realization of the block assignments we can expand the rank- $d$  block-probability matrix  $\mathbf{B}$  up to the full  $\mathbf{P}$  to perform the same deconstruction. In doing so we find that there exist  $\boldsymbol{\nu}_1, \boldsymbol{\nu}_2, \dots, \boldsymbol{\nu}_K \in \mathbb{R}^d$  such that

$$\mathbf{X} = \begin{bmatrix} \mathbf{1}_{n_1} \otimes \boldsymbol{\nu}_1^\top \\ \mathbf{1}_{n_2} \otimes \boldsymbol{\nu}_2^\top \\ \vdots \\ \mathbf{1}_{n_K} \otimes \boldsymbol{\nu}_K^\top \end{bmatrix}, \quad (2)$$

where  $n_k := |\{i \leq n \mid \tau_i = k\}|$ . The  $\boldsymbol{\nu}_k$  are the latent positions of the SBM, and we can reduce this to the simple statement  $\mathbf{B} = \mathbf{x}\mathbf{x}^\top$  where

$$\mathbf{x} = \begin{bmatrix} \boldsymbol{\nu}_1^\top \\ \boldsymbol{\nu}_2^\top \\ \vdots \\ \boldsymbol{\nu}_K^\top \end{bmatrix}.$$

For  $d \leq n$ , the  $d$ -dimensional *adjacency spectral embedding* (ASE) of  $\mathbf{A}$  is given by the  $n \times d$  matrix  $\hat{\mathbf{X}} := \mathbf{U}_\mathbf{A}^{(d)}(\mathbf{D}_\mathbf{A}^{(d)})^{\frac{1}{2}}$ , where the columns of  $\mathbf{U}_\mathbf{A}^{(d)}$  are the first  $d$  columns of  $\mathbf{U}_\mathbf{A}$  and  $\mathbf{D}_\mathbf{A}^{(d)}$  is the  $d \times d$  principal minor of  $\mathbf{D}_\mathbf{A}$  consisting of the  $d$  largest ordered eigenvalues on the diagonal (Lyzinski et al., 2014). The  $n$  rows of  $\hat{\mathbf{X}}$  can thus be thought of as a collection of points in  $\mathbb{R}^d$  that estimate the true latent positions up to orthogonal transformation. We can, therefore, touch upon the notion of how a random choice of an unobservable latent position  $X_i$  (say, via some distribution  $F$  on  $\mathbb{R}^d$ ), informs the

distribution of the corresponding latent position of node  $i$ . For a  $K$ -block SBM we have that the distribution  $F$  on the latent positions is a mixture of point masses, with mixture weights  $\pi_k$ . Athreya et al. (2016) obtained the following result as a corollary to a more general theorem regarding RDPGs and SBMs with  $\rho_n \rightarrow 0$  such that  $\rho_n > \log^4 n/n$ .

**Theorem 2.1.** *Let  $X_i \stackrel{i.i.d.}{\sim} F = \sum_{k=1}^K \pi_k \delta(\boldsymbol{\nu}_k)$ , where  $\boldsymbol{\pi} \in \boldsymbol{\Delta}^K$ . Define  $\Delta = \mathbb{E}[X_1 X_1^\top]$ . Also let  $\boldsymbol{\Sigma}(x) = \Delta^{-1} \mathbb{E}[X_1 X_1^\top (x^\top X_1 - x^\top X_1 X_1^\top x)] \Delta^{-1}$ . If  $\rho_n \equiv 1$ , there exists a sequence of orthogonal matrices  $\mathbf{W}_n$  such that for any fixed index  $i$*

$$\sqrt{n}(\mathbf{W}_n \hat{X}_i - X_i) | X_i = \boldsymbol{\nu}_k \xrightarrow{d} \mathcal{N}(\mathbf{0}, \boldsymbol{\Sigma}(\boldsymbol{\nu}_k)).$$

That is to say that the optimally rotated rows of the ASE for a graph generated by an SBM are approximately distributed as a mixture of curved multivariate normal distributions each centered at the latent positions of  $\mathbf{B}$ . Owing to the method by which the ASE is computed, we note that the rows are identically distributed but not independent.

The *Laplacian spectral embedding* (LSE) is to the normalized Laplacian  $\mathcal{L}(\mathbf{A})$  as the ASE is to  $\mathbf{A}$ . I.e., for  $d \leq n$  the  $d$ -dimensional LSE is given by  $\check{\mathbf{X}} = \tilde{\mathbf{U}}_{\mathbf{A}}^{(d)} (\tilde{\mathbf{D}}_{\mathbf{A}}^{(d)})^{\frac{1}{2}}$ , where  $\tilde{\mathbf{U}}_{\mathbf{A}}^{(d)}$  and  $\tilde{\mathbf{D}}_{\mathbf{A}}^{(d)}$  are analogous to  $\mathbf{U}_{\mathbf{A}}^{(d)}$  and  $\mathbf{D}_{\mathbf{A}}^{(d)}$ . Like  $\hat{\mathbf{X}}$ , the  $n$  rows of  $\check{\mathbf{X}}$  can be thought of as a collection of points in  $\mathbb{R}^d$ . The LSE is preferred in settings where the adjacency matrix is sparse or the edge probability matrix is believed to be of the form  $\rho_n \mathbf{P}_n$  for some positive sequence  $\rho_n \rightarrow 0$  as  $n \rightarrow \infty$  since the normalized Laplacian is the same for  $\mathbf{P}$  as it is for  $\rho \mathbf{P}$ ; i.e.,

$$\begin{aligned} \mathcal{L}(\rho \mathbf{P}) &= (\text{diag}((\rho \mathbf{P}) \mathbf{1}_n)^{-\frac{1}{2}}) (\rho \mathbf{P}) (\text{diag}((\rho \mathbf{P}) \mathbf{1}_n)^{-\frac{1}{2}}) \\ &= \rho^{-1} \rho (\text{diag}(\mathbf{P} \mathbf{1}_n)^{-\frac{1}{2}}) \mathbf{P} (\text{diag}(\mathbf{P} \mathbf{1}_n)^{-\frac{1}{2}}) \\ &= \mathcal{L}(\mathbf{P}). \end{aligned}$$

Tang and Priebe (2018) obtained the following result for the LSE of an SBM, analogous to the previous theorem.

**Theorem 2.2.** *Let the setting be as in Theorem 2.1, and also let  $\boldsymbol{\mu} = \mathbb{E}[X_1]$  and*

$$\tilde{\Delta} = \mathbb{E} \left[ \begin{array}{c} X_1 X_1^\top \\ X_1^\top \boldsymbol{\mu} \end{array} \right].$$

Define

$$\tilde{\Sigma}(x) = \mathbb{E} \left[ \left( \frac{\tilde{\Delta}^{-1} X_1}{X_1^\top \boldsymbol{\mu}} - \frac{x}{2x^\top \boldsymbol{\mu}} \right) \left( \frac{X_1^\top \tilde{\Delta}_{-1}}{X_1^\top \boldsymbol{\mu}} - \frac{x^\top}{x^\top \boldsymbol{\mu}} \right) \left( \frac{x^\top X_1 - x^\top X_1 X_1^\top x}{x^\top \boldsymbol{\mu}} \right) \right].$$

If  $\rho_n \equiv 1$  and  $n_k = |\{i \leq n | X_i = \boldsymbol{\nu}_k\}|$ , then there exists a sequence of orthogonal matrices  $\mathbf{W}_n$  such that for any fixed index  $i$

$$n \left( \mathbf{W}_n \check{X}_i - \frac{\boldsymbol{\nu}_k}{\sqrt{\sum_l n_l \boldsymbol{\nu}_k^\top \boldsymbol{\nu}_l}} \right) | X_i = \boldsymbol{\nu}_k \xrightarrow{d} \mathcal{N}(0, \tilde{\Sigma}(\boldsymbol{\nu}_k)).$$

The limiting distributions justify the use of full GMMs to cluster the rows of either spectral embedding. Doing so improves upon the clustering performance of  $K$ -means, since the inherent nonidentifiability of the RDPG model can lead to different partitions of the point cloud depending on the orthogonal transformation of either spectral embedding; moreover, in utilizing  $K$ -means we impose the assumption of equal, spherical covariances on the clusters, yet the clusters can be non-spherical and of varying spread (Rubin-Delanchy et al., 2020 and Priebe et al., 2019). By contrast, clustering via the EM algorithm accounts for the possibility of unequal covariances and is otherwise invariant to orthogonal transformation of the data; however, the curvature of the components remains unaddressed.

In practice, both  $\text{rank}(\mathbf{B})$  and  $K$  are unknown. A principled method of estimating the former is to inspect the scree-plot of the singular values of  $\mathbf{A}$  and look for “elbows” defining the cut-off between singular values corresponding to signal dimensions and those corresponding to noise dimensions (Priebe et al., 2017). We can estimate the latter by means of maximizing a fitness criterion penalized by model complexity, a la the Akaike Information Criterion (AIC) (Akaike, 1974) and the Bayesian Information Criterion (BIC) (Schwarz, 1978).

For a comprehensive look at the proofs of Theorems 2.1 and 2.2, as well as fuller details of the setting we describe, we refer the reader to Athreya et al. (2018).

### 3 The Expectation-Solution Algorithms

First introduced by Elashoff and Ryan (2004), the Expectation-Solution (ES) algorithm arises as a generalization of the EM algorithm (Dempster et al., 1977), where instead of

updating the parameter estimates according to a collection of complete-data likelihood equations, we update the parameters as solutions to complete-data estimating equations. We will demonstrate that implementation of ES in a curved-Gaussian mixture setting allows us to make full use of the mixture components' curved structure without sacrificing the relative simplicity of EM.

### 3.1 The Expectation-Solution Algorithm for the General Incomplete Data Setting

We model our treatment of the ES algorithm on that found in McLachlan and Krishnan (2008).

Suppose  $\mathbf{X} \sim f_{\Psi}$  is observed,  $\Psi \in \mathbb{R}^p$ , and there exists a natural unobserved extension of the observed data  $\mathbf{Y} = (\mathbf{X}, \mathbf{Z})$ . Let

$$U_c(\mathbf{Y}, \Psi) = \mathbf{0}$$

be the  $p$ -dimensional complete-data estimating equation that could be solved if  $\mathbf{X}$  were totally observable. To apply ES, we first re-express the complete-data estimating equation in terms of a linear function of a  $q$ -dimensional vector function  $S(\mathbf{Y})$ , a  $(p \times q)$ -dimensional matrix function  $\mathbf{A}_{\Psi}$ , and a  $p$ -dimensional vector function  $b_{\Psi}(\mathbf{X})$ :

$$\begin{aligned} U_c(\mathbf{Y}, \Psi) &= U^{(1)}(\mathbf{X}, S(\mathbf{Y}), \Psi) \\ &= \sum_{j=1}^q a_j(\Psi) S_j(\mathbf{Y}) + b_{\Psi}(\mathbf{X}) \\ &= \mathbf{A}_{\Psi} S(\mathbf{Y}) + b_{\Psi}(\mathbf{X}) \\ &= \mathbf{0}, \end{aligned}$$

where  $a_j(\Psi)$  is the  $j$ th column of  $\mathbf{A}_{\Psi}$ . Here  $S(\mathbf{Y})$  is the complete data summary statistic.

The algorithm is:



<b>Algorithm 1: The ES Algorithm for the General Incomplete Data Setting</b>
--

1. Initialize  $\Psi^* = \Psi_0$ .
2. **E-Step:** Compute  $h(\Psi|\Psi^*) := \mathbb{E}_{\Psi^*}[S(\mathbf{Y})|\mathbf{X} = \mathbf{y}]$ .
3. **S-Step:** Find  $\hat{\Psi}$  that solves  $\mathbf{A}_{\hat{\Psi}}h(\hat{\Psi}|\Psi^*) + b_{\hat{\Psi}}(\mathbf{X}) = \mathbf{0}$ .
4. Take  $\Psi^* = \hat{\Psi}$ .
5. Repeat steps 2–4 until some convergence criterion is satisfied.

**Remark.** *When the complete-data estimating equations for a given model coincide exactly with the complete-data likelihood equations for the same model, then the ES and EM algorithms are equivalent. Thus, any comparisons drawn between EM and ES for a particular setting are actually between two different ES algorithms, a fact which allows us to use comparable convergence criteria. See the first paragraph of section 4 for more details.*

### 3.2 The ES Algorithm for Mixtures of Curved Gaussians

Consider the usual  $K$ -component  $d$ -variate Gaussian mixture model (GMM):

$$X_i \stackrel{i.i.d.}{\sim} \sum_{k=1}^K \pi_k \mathcal{N}(\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k), i = 1, \dots, n;$$

such that  $\boldsymbol{\pi} := (\pi_1, \dots, \pi_K) \in \Delta^K$  and  $(\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \in \mathbb{R}^d \times \mathbb{M}_d$  for all  $k$ , where  $\mathbb{M}_d$  is the space of  $d \times d$  positive definite matrices. Maximum likelihood estimation of the parameters upon observing  $x_1, \dots, x_n$  can be handily accomplished by the EM algorithm. Here, the component distributions are by themselves smooth exponential family distributions, a fact that allows us to utilize various results related to the computation of maximum likelihood estimates in the M-step.

Suppose instead that  $X_1, X_2, \dots, X_n$  are identically distributed, and our model is

$$X_i \sim \sum_{k=1}^K \pi_k \mathcal{N}(\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k(\boldsymbol{\mu}_k))$$

$$\boldsymbol{\mu}_k \in \mathcal{M} \subset \mathbb{R}^d$$

$$\boldsymbol{\Sigma}_k : \mathcal{M} \rightarrow \mathbb{M}_d, \forall k$$

and each  $\boldsymbol{\Sigma}_k$  is continuous and differentiable. Here each component distribution is a curved exponential family distribution, where the component variances are functions of the means

(Bickel and Doksum, 2015); hence we refer to such a setting as a curved-Gaussian mixture model (C-GMM). If the  $X_i$  are also independent and we attempt to derive complete-data likelihood equations to derive an EM algorithm, we may become swiftly enmired in non-linear equations which we must solve to update the means at each iteration, depending on the structure of each  $\Sigma_k$ .

Here the ES algorithm circumvents the potential difficulty afforded by the variance functions. We begin by considering the natural unobserved data extension of  $\mathbf{X} = (X_1, \dots, X_n)$ , which should come as no surprise to those familiar with the EM algorithm:

$$Z_i = (Z_{i1}, \dots, Z_{iK}) \stackrel{i.i.d.}{\sim} \text{Multinomial}(1, K, \boldsymbol{\pi})$$

$$X_i | Z_{ik} = 1 \sim \mathcal{N}(\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k(\boldsymbol{\mu}_k)).$$

In the complete-data setting  $(\mathbf{X}, \mathbf{Z})$  we immediately note the following:

$$\mathbb{E}_{\Psi}[Z_{ik}] = \pi_k$$

$$\mathbb{E}_{\Psi}[Z_{ik}X_i] = \pi_k \boldsymbol{\mu}_k,$$

which hold regardless of the dependencies among the  $X_i$ . These give rise to the natural estimating equations

$$\frac{\sum_{i=1}^n Z_{ik}}{n} - \pi_k = 0;$$

$$\frac{\sum_{i=1}^n Z_{ik}X_i}{n} - \pi_k \boldsymbol{\mu}_k = 0,$$

which are solved by

$$\hat{\pi}_k = \frac{\sum_{i=1}^n Z_{ik}}{n};$$

$$\hat{\boldsymbol{\mu}}_k = \frac{\sum_{i=1}^n Z_{ik}X_i}{\sum_{i=1}^n Z_{ik}}.$$

If we take  $\Psi$  to consist of  $\pi_1, \dots, \pi_{K-1}$  and the entries of  $\boldsymbol{\mu}_1, \dots, \boldsymbol{\mu}_K$  then the complete-data

estimating equation  $U_c((\mathbf{X}, \mathbf{Z}), \Psi) = 0$  is characterized by

$$\begin{aligned}
 \mathbf{A}_\Psi &= \mathbf{I}_{(d+1)K-1} \\
 S(\mathbf{X}, \mathbf{Z}) &= \frac{1}{n} \begin{bmatrix} \sum_{i=1}^n Z_{i1} \\ \vdots \\ \sum_{i=1}^n Z_{i(K-1)} \\ \sum_{i=1}^n Z_{i1} X_i \\ \vdots \\ \sum_{i=1}^n Z_{iK} X_i \end{bmatrix} \\
 b_\Psi(\mathbf{X}) &= - \begin{bmatrix} \pi_1 \\ \vdots \\ \pi_{K-1} \\ \pi_1 \boldsymbol{\mu}_1 \\ \vdots \\ (1 - \sum_{k=1}^{K-1} \pi_k) \boldsymbol{\mu}_K \end{bmatrix}.
 \end{aligned}$$

In computing  $h(\Psi|\Psi^*) = \mathbb{E}_{\Psi^*}[S(\mathbf{X}, \mathbf{Z})|\mathbf{X} = \mathbf{x}]$ , we find that we must only obtain

$$\begin{aligned}
 Z_{ik}^* &:= \mathbb{E}_{\Psi^*}[Z_{ik}|X_i = x_i] \\
 &= \frac{\pi_k^* \phi(x_i | \boldsymbol{\mu}_k^*, \boldsymbol{\Sigma}_k(\boldsymbol{\mu}_k^*))}{\sum_{j=1}^K \pi_j^* \phi(x_i | \boldsymbol{\mu}_j^*, \boldsymbol{\Sigma}_j(\boldsymbol{\mu}_j^*))}.
 \end{aligned}$$

Thus  $h(\Psi|\Psi^*) = S(\mathbf{x}, \mathbf{Z}^*)$  by linearity of expectation. The next iterate  $\hat{\Psi}$  is then obtained by solving  $\mathbf{A}_\Psi S(\mathbf{x}, \mathbf{Z}^*) + b_\Psi(\mathbf{X}) = 0$ , which yields

$$\begin{aligned}
 \hat{\pi}_k &= \frac{\sum_{i=1}^n Z_{ik}^*}{n}, \\
 \hat{\boldsymbol{\mu}}_k &= \frac{\sum_{i=1}^n Z_{ik}^* x_i}{\sum_{i=1}^n Z_{ik}^*}.
 \end{aligned}$$

Here is the algorithm written concisely:

**Algorithm 2: The ES Algorithm for the Curved-Gaussian Mixture Setting**

1. Initialize  $\Psi^* = \Psi_0$ .
2. **E-Step:** For each  $i$  and  $k$  compute

$$Z_{ik}^* = \frac{\pi_k^* \phi(x_i | \boldsymbol{\mu}_k^*, \boldsymbol{\Sigma}_k(\boldsymbol{\mu}_k^*))}{\sum_{j=1}^K \pi_j^* \phi(x_i | \boldsymbol{\mu}_j^*, \boldsymbol{\Sigma}_j(\boldsymbol{\mu}_j^*))}.$$

3. **S-Step:** Compute the entries of  $\hat{\Psi}$  as

$$\hat{\pi}_k = \frac{\sum_{i=1}^n Z_{ik}^*}{n}$$

$$\hat{\boldsymbol{\mu}}_k = \frac{\sum_{i=1}^n Z_{ik}^* x_i}{\sum_{i=1}^n Z_{ik}^*}.$$

4. Take  $\Psi^* = \hat{\Psi}$ .
5. Repeat steps 2–4 until some convergence criterion is satisfied.

Note that the only difference between this ES algorithm and the usual EM algorithm for GMMs is that we “update” the component variances by plugging the usual updates for the component means into the respective  $\boldsymbol{\Sigma}_k(\cdot)$  instead of computing

$$\hat{\boldsymbol{\Sigma}}_k = \frac{\sum_{i=1}^n Z_{ik}^* (x_i - \hat{\boldsymbol{\mu}}_k)(x_i - \hat{\boldsymbol{\mu}}_k)^\top}{\sum_{i=1}^n Z_{ik}^*}$$

for each  $k$ . As a result we estimate only  $(d+1)K - 1$  parameters, whereas in the full GMM setting we estimate another  $K(d + \binom{d}{2})$  parameters comprising the component covariances. Since the model complexity penalties used to compute both AIC and BIC increase in magnitude with the number of parameters, the severe reduction in the number of parameters needed to be estimated in a C-GMM can vastly decrease these penalties. I.e., if we are deciding between a GMM and C-GMM with approximately equal likelihoods for a given collection of data, then it is clear that the C-GMM will achieve the higher (and therefore more desirable) AIC or BIC.

If we are interested in clustering the observations we perform the usual GMM clustering procedure, which is to assign each observation to the cluster with the highest posterior probability.

**Remark.** We note here the distinction between two classes of C-GMMs. We call a C-

*GMM separable* if each component variance is solely a function of its respective mean and possibly its mixing proportion. Likewise, if one or more of the variances take other components' means or mixing proportions as arguments, then that C-GMM is **tied** — as in the cases of both spectral embeddings' limiting distributions.

### 3.3 ES Algorithm for the ASE

Let us assume the setting of Theorem 2.1 with  $\mathbf{B} = \mathbf{xx}^\top$ , where the  $k^{\text{th}}$  row of  $\mathbf{x}$  is  $\boldsymbol{\nu}_k$ , and that we have observed  $\mathbf{A}$  and computed its  $d$ -dimensional ASE  $\hat{\mathbf{X}}$ . Since  $F = \sum_{k=1}^K \pi_k \delta(\boldsymbol{\nu}_k)$ , we have that  $\Delta = \sum_{k=1}^K \pi_k \boldsymbol{\nu}_k \boldsymbol{\nu}_k^\top$  and

$$\boldsymbol{\Sigma}(\boldsymbol{\nu}_k) = \Delta^{-1} \left( \sum_{j=1}^K \pi_j \boldsymbol{\nu}_j \boldsymbol{\nu}_j^\top (\boldsymbol{\nu}_k^\top \boldsymbol{\nu}_j - \boldsymbol{\nu}_k^\top \boldsymbol{\nu}_j \boldsymbol{\nu}_j^\top \boldsymbol{\nu}_k) \right) \Delta^{-1}.$$

The variance function thus takes every latent position and mixture weight as arguments, and our iterative scheme must reflect this. Therefore, let  $\boldsymbol{\Sigma}(\cdot | \mathbf{x}, \boldsymbol{\pi})$  denote the covariance function instead. The normal mixture model we thus consider is

$$\hat{X}_i \sim \sum_{k=1}^K \pi_k \mathcal{N} \left( \boldsymbol{\nu}_k, \frac{\boldsymbol{\Sigma}(\boldsymbol{\nu}_k | \mathbf{x}, \boldsymbol{\pi})}{n} \right),$$

where each  $\hat{X}_i$  is identically distributed.

The algorithm is thus:

#### Algorithm 3: The ES Algorithm for the ASE

1. Initialize  $\boldsymbol{\Psi}^* = \boldsymbol{\Psi}_0$ .
2. **E-Step:** Compute

$$Z_{ik}^* = \frac{\pi_k^* \phi(\hat{X}_i | \boldsymbol{\nu}_k^*, \boldsymbol{\Sigma}(\boldsymbol{\nu}_k^* | \mathbf{x}^*, \boldsymbol{\pi}^*)/n)}{\sum_{j=1}^K \pi_j^* \phi(\hat{X}_i | \boldsymbol{\nu}_j^*, \boldsymbol{\Sigma}(\boldsymbol{\nu}_j^* | \mathbf{x}^*, \boldsymbol{\pi}^*)/n)}$$

3. **S-Step:** Compute

$$\hat{\pi}_k = \frac{\sum_{i=1}^n Z_{ik}^*}{n}$$

$$\hat{\boldsymbol{\nu}}_k = \frac{\sum_{i=1}^n Z_{ik}^* \hat{X}_i}{\sum_{i=1}^n Z_{ik}^*}.$$

4. Take  $\boldsymbol{\Psi}^* = \hat{\boldsymbol{\Psi}}$ .
5. Repeat steps 2-4 until some convergence criterion is satisfied.

**Remark.** We can construct a separable analogue of Algorithm 3 by updating each variance with the newest iterate of the corresponding mean and proportion while holding all other arguments as their previous iterates, only updating them every  $\iota$  iterations, or holding them constant. Such schemes (particularly the last) greatly alter the model at hand, hence we would not recommend them for use; rather, we only mention them to fill in the middle ground between algorithms 2 and 3.

### 3.4 ES Algorithm for the LSE

Let us assume the setting of Theorem 2.2. We have

$$\begin{aligned}\boldsymbol{\mu} &= \sum_{k=1}^K \pi_k \boldsymbol{\nu}_k \\ \tilde{\Delta} &= \sum_{k=1}^K \pi_k \frac{\boldsymbol{\nu}_k \boldsymbol{\nu}_k^\top}{\boldsymbol{\nu}_k^\top \boldsymbol{\mu}} \\ \tilde{\Sigma}(\boldsymbol{\nu}_k | \mathbf{x}, \boldsymbol{\pi}) &= \sum_{j=1}^K \pi_j \left( \frac{\tilde{\Delta}^{-1} \boldsymbol{\nu}_j}{\boldsymbol{\nu}_j^\top \boldsymbol{\mu}} - \frac{\boldsymbol{\nu}_k}{2 \boldsymbol{\nu}_k^\top \boldsymbol{\mu}} \right) \left( \frac{\boldsymbol{\nu}_j^\top \tilde{\Delta}^{-1}}{\boldsymbol{\nu}_j^\top \boldsymbol{\mu}} - \frac{\boldsymbol{\nu}_k^\top}{\boldsymbol{\nu}_k^\top \boldsymbol{\mu}} \right) \left( \frac{\boldsymbol{\nu}_k^\top \boldsymbol{\nu}_j - \boldsymbol{\nu}_k^\top \boldsymbol{\nu}_j \boldsymbol{\nu}_j^\top \boldsymbol{\nu}_k}{\boldsymbol{\nu}_k^\top \boldsymbol{\mu}} \right).\end{aligned}$$

The normal mixture model we consider here is

$$\check{X}_i \sim \sum_{k=1}^K \pi_k \mathcal{N} \left( \frac{\boldsymbol{\nu}_k}{\sqrt{\sum_l n_l \boldsymbol{\nu}_l^\top \boldsymbol{\nu}_k}}, \frac{\tilde{\Sigma}(\boldsymbol{\nu}_k | \mathbf{x}, \boldsymbol{\pi})}{n^2} \right),$$

where each  $\check{X}_i$  is identically distributed. Owing to the presence of the unobserved  $n_k$  in the component means, the fact that the ASE and LSE are defined by the 1-1 transformation

$$\check{\mathbf{X}} = (\text{diag}(\mathbf{A}\mathbf{1}_n)^{-1/2}) \hat{\mathbf{X}}, \quad (3)$$

and the fact that the LSE covariance function takes the ASE component means as arguments, we cannot simply invoke an analogue of the above algorithm. We propose expanding the parameter vector  $\boldsymbol{\Psi}$  by treating the  $n_k$  as parameters that can be obtained by the  $K$  estimating equations

$$\mathbb{E}_{\boldsymbol{\Psi}} \left[ \sum_{i=1}^n Z_{ik} \right] - n_k = 0,$$

which would be solved by  $\hat{n}_k := \sum_{i=1}^n Z_{ik} = n \hat{\pi}_k$  if we observed the cluster labels of each row of  $\check{\mathbf{X}}$ .

Upon observing  $\mathbf{A}$  and computing both  $\hat{\mathbf{X}}$  and  $\check{\mathbf{X}}$ , we propose the following algorithm:

**Algorithm 4: The ES Algorithm for the LSE**

1. Initialize  $\Psi^* = \Psi_0$  and take  $\Sigma_k(\cdot) = \tilde{\Sigma}(\cdot|\mathbf{x}, \boldsymbol{\pi})/n^2$ .
2. **E-Step:** Compute

$$\boldsymbol{\mu}_k^* = \frac{\boldsymbol{\nu}_k^*}{\sqrt{\sum_l n_l^* \boldsymbol{\nu}_l^{*T} \boldsymbol{\nu}_k^*}}$$

$$Z_{ik}^* = \frac{\pi_k^* \phi(\check{X}_i | \boldsymbol{\mu}_k^*, \tilde{\Sigma}(\boldsymbol{\nu}_k^* | \mathbf{x}^*, \boldsymbol{\pi}^*)/n^2)}{\sum_{j=1}^K \pi_j^* \phi(\check{X}_i | \boldsymbol{\mu}_j^*, \tilde{\Sigma}(\boldsymbol{\nu}_j^* | \mathbf{x}^*, \boldsymbol{\pi}^*)/n^2)}.$$

3. **S-Step:** Compute

$$\hat{\pi}_k = \frac{\sum_{i=1}^n Z_{ik}^*}{n}$$

$$\hat{\boldsymbol{\nu}}_k = \frac{\sum_{i=1}^n Z_{ik}^* \hat{X}_i}{\sum_{i=1}^n Z_{ik}^*}$$

$$\hat{n}_k = n \hat{\pi}_k.$$

4. Take  $\Psi^* = \hat{\Psi}$ .
5. Repeat steps 2-4 until some convergence criterion is satisfied.

Note that this algorithm makes full use of both the LSE and ASE; the  $\check{X}_i$  are used in the E-Step, but the  $\hat{X}_i$  are used in the S-Step. Even so, clustering is to be done based on the rows of  $\check{\mathbf{X}}$ , as they correspond exactly to the posterior probabilities we compute while the algorithm runs its course.

## 4 Simulations

For each simulation setting we generated 100 samples of size  $n = 200, 300, \dots, 900$  and compared the performance of our ES algorithms with their respective EM analogues. The specific EM algorithm we used was the function *em* from the R package *Mclust* (Scrucca et al., 2016). All four procedures were initialized at the true parameter values. Since EM is a maximum likelihood procedure, the default convergence criterion is to terminate when updates to the loglikelihood are less than  $1 \times 10^{-5}$ . By contrast there is no convenient objective function associated with the ES algorithm, since the iterates arise as solutions

to complete-data estimating equations; moreover, it was observed anecdotally in a few of our settings that the loglikelihood may decrease after an ES iteration. With that in mind, we noted that the EM algorithm actually is an ES algorithm where the complete-data estimating equations are the complete-data likelihood equations; and we altered the code of *em* to reflect this. Therefore convergence of EM and ES for the ASE was assumed when the Euclidean distance between successive estimates of the parameter vector  $\Psi$  consisting of the mixing proportions and the entries of  $\mathbf{x}$  was less than  $1 \times 10^{-5}$ , and convergence of EM and ES for the LSE was assumed when the Euclidean distance between successive estimates of the parameter vector  $\tilde{\Psi}$  consisting of the mixing proportions and the scaled latent positions was less than  $1 \times 10^{-6}$  for the LSE, or the number of iterations exceeded 10,000. The lower threshold for  $ES \circ LSE$  was based on the fact that in computing the LSE we effectively shrink the rows of the ASE towards the origin.

To evaluate clustering performance we computed the adjusted Rand index (ARI) (Hubert and Arabie, 1985) for the cluster assignments from each method with the true cluster labels. For each clustering method  $m$  found in (1), we let  $ARI_m$  denote the ARI produced by that method. For  $\ell = A, L$  we performed Wilcoxon rank-sum tests (Mann and Whitney, 1947) with hypotheses

$$\begin{aligned} H_o &: \text{med}(ARI_{EM \circ LSE}) = \text{med}(ARI_{ES \circ LSE}) \\ H_a &: \text{med}(ARI_{EM \circ LSE}) \neq \text{med}(ARI_{ES \circ LSE}). \end{aligned}$$

We tabulate all results in the form of 95% confidence intervals for the median of each  $\Delta_\ell := ARI_{EM \circ LSE} - ARI_{ES \circ LSE}$ .

To evaluate accuracy of the parameter estimates we computed the squared error of each method's terminating estimate from the true parameter vector  $\Psi$ . As will be seen in subsection 4.2 we circumvented the issue of the latent positions' non-identifiability by rotating and centering the our simulated data over the "canonical" latent positions in the first orthant by way of a Procrustes transformation. We take  $\Psi_{ASE}$  to consist of the entries of  $\pi$  and  $\mathbf{x}$ , as well as those of the covariance matrices  $\frac{\Sigma(\nu_k|\mathbf{x},\pi)}{n}$ . As  $EM \circ LSE$  does not outright produce estimates for the  $\nu_k$ , we take  $\Psi_{LSE}$  to consist of the entries of  $\pi$ ,  $\mu_k := \frac{\nu_k}{\sqrt{\sum_l n_l \nu_l^\top \nu_k}}$ , and the covariance matrices  $\frac{\tilde{\Sigma}(\nu_k|\mathbf{x},\pi)}{n^2}$ . We then performed Wilcoxon



rank sum tests for the paired collection of squared errors for  $\{EM, ES\} \circ ASE$ , as well as  $\{EM, ES\} \circ LSE$ . We let  $p_A$  denote the  $p$ -value corresponding to the test

$$\begin{aligned} H_o &: \text{med}(\|\hat{\Psi}_{EM \circ ASE} - \Psi_{ASE}\|) \leq \text{med}(\|\hat{\Psi}_{ES \circ ASE} - \Psi_{ASE}\|) \\ H_a &: \text{med}(\|\hat{\Psi}_{EM \circ ASE} - \Psi_{ASE}\|) > \text{med}(\|\hat{\Psi}_{ES \circ ASE} - \Psi_{ASE}\|), \end{aligned}$$

and  $p_L$  be that corresponding to the test

$$\begin{aligned} H_o &: \text{med}(\|\hat{\Psi}_{EM \circ LSE} - \Psi_{LSE}\|) \leq \text{med}(\|\hat{\Psi}_{ES \circ LSE} - \Psi_{LSE}\|) \\ H_a &: \text{med}(\|\hat{\Psi}_{EM \circ LSE} - \Psi_{LSE}\|) > \text{med}(\|\hat{\Psi}_{ES \circ LSE} - \Psi_{LSE}\|). \end{aligned}$$

In all tables we embolden entries which indicate strict improvement of our algorithms over EM.

## 4.1 Non-Graph Setting

We first tested  $ES \circ \{ASE, LSE\}$  on random data generated directly from the mixture distributions given in Theorems 2.1 and 2.2. We considered the following models:

$$\begin{aligned} \mathbf{x}_1 &= \begin{bmatrix} 0.6210 & 0.3382 \\ 0.3382 & 0.6210 \end{bmatrix} & \mathbf{x}_2 &= \begin{bmatrix} 0.4076 & 0.1840 \\ 0.1840 & 0.4076 \end{bmatrix} \\ \mathbf{x}_3 &= \begin{bmatrix} 0.6024 & 0.3703 \\ 0.3703 & 0.5319 \end{bmatrix} & \mathbf{x}_4 &= \begin{bmatrix} 0.3962 & 0.2074 \\ 0.2074 & 0.3721 \end{bmatrix} \end{aligned}$$

with  $\boldsymbol{\pi} = (\frac{1}{2}, \frac{1}{2})$  in all cases. It will be seen that these particular choices of  $\mathbf{x}$  correspond to the four SBMs from which we simulate in the following two subsections. Each randomly generated sample was taken to comprise  $\hat{\mathbf{X}}$  and the simulated LSE was computed via the 1–1 relationship (3), taking  $\mathbf{A} = \hat{\mathbf{X}} \hat{\mathbf{X}}^\top$ .

For models 1 and 2 the EM algorithms approximately match the ES algorithms' performance for large  $n$ . As  $n$  increases, the component variances of both mixtures shrink around the component means, pulling the points into tight, distinct clusters. EM tended to cluster more accurately than ES in models 3 and 4, and we credit this to the fact that in these settings the point clouds overlap significantly for small to moderate  $n$ , and it is not until around  $n = 700$  that distinct clusters start to form.

Table 1: Model 1 Mixture. Negative  $\text{med}(\Delta)$  indicates superiority of ES over EM.

$n$	95% CI for $\text{med}(\Delta_A)$	95% CI for $\text{med}(\Delta_L)$	$p_A$	$p_L$
200	<b>(-0.0598, -0.0252)</b>	<b>(-0.0435, -0.0165)</b>	<b>3e-5</b>	<b>0.0002</b>
300	<b>(-0.0123, -4e-5)</b>	<b>(-0.0122, -0.0001)</b>	<b>5e-7</b>	<b>1e-5</b>
400	(-0.0091, 7e-6)	(-0.0095, 6e-7)	<b>0.0072</b>	<b>0.0002</b>
500	<b>(-0.0079, -2e-5)</b>	<b>(-0.0079, -6e-5)</b>	<b>0.0014</b>	<b>0.0003</b>
600	(-0.0065, 1e-6)	<b>(-0.0065, -1e-5)</b>	<b>0.0032</b>	<b>0.0008</b>
700	(-0.0029, 8e-5)	(-0.0028, 0.0028)	<b>0.0059</b>	<b>0.0011</b>
800	(-0.0050, 7e-5)	(-0.0025, 5e-5)	<b>0.0123</b>	<b>0.0015</b>
900	(-0.0044, 0.0025)	(-0.0044, 0.0022)	0.0796	<b>0.0370</b>

Table 2: Model 2 Mixture. Negative  $\text{med}(\Delta)$  indicates superiority of ES over EM.

$n$	95% CI for $\text{med}(\Delta_A)$	95% CI for $\text{med}(\Delta_L)$	$p_A$	$p_L$
200	(0.0062, .1097)	(0.0085, 0.1029)	1.0000	0.9998
300	(-0.0207, 0.0202)	(-0.0091, 0.0244)	0.7730	0.9940
400	<b>(-0.0234, -0.0048)</b>	<b>(-0.0126, -0.0008)</b>	<b>0.0002</b>	0.4324
500	<b>(-0.0138, -0.0034)</b>	<b>(-0.0102, -7e-5)</b>	<b>0.0012</b>	0.0989
600	(-0.0060, 0.0001)	<b>(-0.0059, -3e-5)</b>	<b>0.0005</b>	<b>0.0014</b>
700	(-0.0052, 4e-7)	(-0.0052, 0.0001)	<b>0.0005</b>	<b>0.0019</b>
800	<b>(-0.0048, -0.0023)</b>	<b>(-0.0047, -0.0001)</b>	<b>1e-6</b>	<b>2e-6</b>
900	(-0.0042, 3e-5)	(-0.0042, 4e-5)	<b>0.0001</b>	<b>0.0005</b>

In model 1, the ES algorithms tend to more accurately estimate the component means, variances, and weights than do the EM algorithms. As  $n$  increases the accuracy of the EM algorithms begins to match that of the ES algorithms; again, we suspect that this is due to the emergence of distinct clusters. In model 2, the EM algorithms perform more accurate estimation for small  $n$ , but are overtaken by the ES algorithms as  $n$  increases. Owing to the fact that  $\mathbf{x}_2$  arises as the latent positions for a sparse SBM (relative to that corresponding to  $\mathbf{x}_1$ ), we suspect that as  $n$  increases beyond 900 parameter estimation by either algorithm will be comparable as this model's clusters further separate.

Table 3: Model 3 Mixture. Negative  $\text{med}(\Delta)$  indicates superiority of ES over EM.

$n$	95% CI for $\text{med}(\Delta_A)$	95% CI for $\text{med}(\Delta_L)$	$p_A$	$p_L$
200	(0.0309, 0.0849)	(0.0433, 0.0941)	1.0000	0.9999
300	(0.0252, 0.0738)	(0.0477, 0.0923)	0.9998	0.9999
400	(0.0233, 0.0651)	(0.0215, 0.0631)	0.9999	0.9993
500	(0.0031, 0.0372)	(0.0063, 0.0371)	0.9108	0.9797
600	(-0.0029, 0.0103)	(-0.0001, 0.0087)	0.7889	0.6208
700	(-0.0046, 0.0036)	(-0.0049, 0.0024)	0.1181	<b>0.0174</b>
800	<b>(-0.0065, -2e-5)</b>	(-0.0045, 3e-5)	<b>3e-5</b>	<b>2e-5</b>
900	(-0.0020, 0.0020)	(-4e-5, 0.0022)	0.1889	0.0731

Table 4: Model 4 Mixture. Negative  $\text{med}(\Delta)$  indicates superiority of ES over EM.

$n$	95% CI for $\text{med}(\Delta_A)$	95% CI for $\text{med}(\Delta_L)$	$p_A$	$p_L$
200	(0.0410, 0.0863)	(0.0256, 0.0733)	1.0000	1.0000
300	(0.0821, 0.1459)	(0.0561, 0.1030)	1.0000	1.0000
400	(0.1039, 0.1828)	(0.0770, 0.1303)	1.0000	1.0000
500	(0.0574, 0.1422)	(0.0461, 0.0965)	1.0000	0.9998
600	(0.0255, 0.0797)	(0.0241, 0.0658)	0.9999	0.9983
700	(0.0046, 0.0200)	(0.0027, 0.0177)	0.9944	0.8045
800	(0.0020, 0.0140)	(0.0002, 0.0121)	0.9661	0.3431
900	(-0.0018, 0.0055)	(-0.0014, 0.0055)	0.8110	<b>0.0423</b>

In models 3 and 4 the ES algorithms largely fail to estimate the parameters more accurately than the EM algorithms, save for around  $n = 800$  in model 3. Due to the decreasing trend in the confidence bounds in table 4 as  $n$  increases, as well as the fact that  $\mathbf{x}_4$  corresponds to a sparser SBM than does  $\mathbf{x}_3$ , we suspected that this phenomenon is replicated at a range of  $n > 900$  where the ES algorithms outperform the EM algorithms in model 4. Hence we repeated the experiment for  $n = 1000, 1100, \dots, 1700$  and output the results to table 5, which does indeed indicate a range for which  $ES \circ LSE$  beats  $EM \circ LSE$ , both in terms of clustering accuracy and parameter estimation.

Table 5: Model 4 Mixture,  $n > 900$ 

$n$	95% CI for $\text{med}(\Delta_A)$	95% CI for $\text{med}(\Delta_L)$	$p_A$	$p_L$
1000	(-0.0034, 0.0018)	<b>(-0.0068, -0.0001)</b>	0.4541	<b>0.0044</b>
1100	(-0.0031, 0.0016)	(-0.0032, 0.0016)	0.3857	<b>0.0065</b>
1200	(-0.0031, 0.0014)	(-0.0030, 0.0014)	0.4555	<b>0.0211</b>
1300	(-0.0026, 0.0014)	(-0.0027, 0.0001)	0.0965	<b>0.0320</b>
1400	(-0.0013, 0.0013)	<b>(-0.0050, -0.0011)</b>	<b>0.0338</b>	<b>0.0001</b>
1500	(-0.0024, 5e-6)	<b>(-0.0036, -0.0001)</b>	<b>0.0019</b>	<b>1e-6</b>
1600	(-0.0023, 4e-5)	(-0.0023, 4e-5)	0.1926	<b>0.0137</b>
1700	<b>(-0.0022, -1e-5)</b>	<b>(-0.0032, -0.0010)</b>	0.0885	<b>0.0098</b>

## 4.2 Balanced Affinity Network Structure

A  $K$ -block SBM is said to possess *homogeneous balanced affinity structure* if  $B_{ii} = a$  for all  $i$ ,  $B_{ij} = b$  for all  $i \neq j$ ,  $0 < b < a < 1$ , and  $\pi_k = \frac{1}{K}$  for all  $k$  (Cape et al., 2019). For the purposes of our simulations we considered the 2-block homogeneous balanced affinity SBMs characterized by  $(a_1, b_1) = (.5, .4)$  and  $(a_2, b_2) = (.2, .15)$ .

Since the latent position matrix  $\mathbf{x}$  is only identifiable up to orthogonal transformation, we took the canonical latent position matrix to be that centered in the first quadrant of  $\mathbb{R}^2$  via the following transformation:

$$\mathbf{x} = \mathbf{U}_B \mathbf{D}_B^{\frac{1}{2}} \mathbf{U}_B^\top.$$

For each model we used the R package *igraph* to sample from the SBM with the class assignments fixed, computed the ASE  $\hat{\mathbf{X}}$  from its definition, then centered the rows over the canonical latent positions with the  $d \times d$  orthogonal transformation  $\hat{\mathbf{W}}$  that solved the Procrustes problem

$$\begin{aligned} & \min_{\mathbf{W}} \|\hat{\mathbf{X}}\mathbf{W} - \mathbf{X}\|_F \\ & \text{subject to } \mathbf{W}^\top \mathbf{W} = \mathbf{I} \end{aligned}$$

where  $\mathbf{X}$  is as in (2). Following this the LSE was computed via the 1–1 relationship (3). We then performed the procedure as described in the previous subsection and output the

results in tables 6–7. Overall we observe the same results as in tables 1 and 2; however, it’s clear that the degree to which ES out-clusters EM has increased, particularly in model 2.

Table 6: Model 1 SBM. Negative  $\text{med}(\Delta)$  indicates superiority of ES over EM.

$n$	95% CI for $\text{med}(\Delta_A)$	95% CI for $\text{med}(\Delta_L)$	$p_A$	$p_L$
200	<b>(-0.0634, -0.0139)</b>	<b>(-0.0504, -0.0201)</b>	<b>0.0080</b>	<b>8e-10</b>
300	<b>(-0.0237, -0.0104)</b>	<b>(-0.0225, -0.0108)</b>	<b>8e-5</b>	<b>2e-12</b>
400	<b>(-0.0140, -0.0044)</b>	<b>(-0.0094, -0.0040)</b>	<b>5e-9</b>	<b>1e-12</b>
500	<b>(-0.0076, -2e-5)</b>	(-0.0039, 0.0001)	<b>7e-6</b>	<b>1e-7</b>
600	(-0.0034, 4e-6)	(-0.0032, 0.0001)	<b>1e-6</b>	<b>3e-9</b>
700	(-0.0028, 5e-5)	(-0.0028, 8e-6)	<b>0.0001</b>	<b>2e-5</b>
800	(-0.0049, 7e-6)	<b>(-0.0049, -1e-5)</b>	<b>1e-5</b>	<b>0.0001</b>
900	(-0.0043, 3e-5)	(-0.0043, 2e-5)	<b>9e-7</b>	<b>0.0005</b>

Table 7: Model 2 SBM. Negative  $\text{med}(\Delta)$  indicates superiority of ES over EM.

$n$	95% CI for $\text{med}(\Delta_A)$	95% CI for $\text{med}(\Delta_L)$	$p_A$	$p_L$
200	(0.0115, 0.0338)	(-0.0028, 0.0060)	1.0000	0.0967
300	(0.0077, 0.0603)	(-0.0095, 0.0019)	0.9999	<b>0.0003</b>
400	<b>(-0.0308, -0.0006)</b>	<b>(-0.0180, -0.0060)</b>	0.0919	<b>7e-11</b>
500	<b>(-0.0227, -0.0082)</b>	<b>(-0.0118, -0.0029)</b>	<b>2e-5</b>	<b>7e-11</b>
600	<b>(-0.0155, -0.0053)</b>	<b>(-0.0078, -0.0001)</b>	<b>4e-5</b>	<b>2e-10</b>
700	<b>(-0.0115, -0.0026)</b>	<b>(-0.0091, -0.0024)</b>	<b>4e-11</b>	<b>3e-11</b>
800	<b>(-0.0088, -0.0022)</b>	<b>(-0.0064, -0.0001)</b>	<b>2e-9</b>	<b>8e-9</b>
900	<b>(-0.0061, -0.0020)</b>	<b>(-0.0059, -0.0020)</b>	<b>1e-9</b>	<b>3e-13</b>

### 4.3 Core-Periphery Network Structure

A 2-block SBM is said to possess *core-periphery structure* if  $\mathbf{B} = \begin{bmatrix} a & b \\ b & b \end{bmatrix}$  and  $\boldsymbol{\pi} = (\pi_1, 1 - \pi_1)$  (Cape et al., 2019). We considered the 2-block balanced core-periphery SBMs characterized by  $(a_3, b_3) = (.2, .15)$  and  $(a_4, b_4) = (.5, .42)$ . We then performed the same procedure as

described in the previous subsections and output the results to tables 8–9. As in the mixture setting, ES largely fails to outperform EM, except for small to moderate  $n$  in model 3. However, the CIs in tables 3–4 indicate that EM tended to strictly outperform ES; but in tables 7–8,  $ES_L$  is approximately on par with  $EM_L$ . As before, we repeated the model 4 simulations for  $n = 1000, 1100, \dots, 1700$  and output the results to table 10.

Both here and in the mixture setting, it was observed that ES tended to vastly overestimate the entries of the covariance matrices, hence the algorithms’ seeming inability to more accurately estimate  $\Psi$ , except for  $ES \circ LSE$  in model 3 when  $700 \leq n \leq 900$ .

Table 8: Model 3 SBM. Negative med( $\Delta$ ) indicates superiority of ES over EM.

$n$	95% CI for med( $\Delta_A$ )	95% CI for med( $\Delta_L$ )	$p_A$	$p_L$
200	(-0.0388, 0.0090)	(0.0009, 0.0348)	1.0000	1.0000
300	<b>(-0.0735, -0.0279)</b>	(-0.0149, 0.0136)	1.0000	1.0000
400	<b>(-0.0790, -0.0217)</b>	<b>(-0.0290, -0.0060)</b>	1.0000	0.9999
500	<b>(-0.0350, -0.0026)</b>	(-0.0116, 0.0053)	1.0000	0.9661
600	(-0.0036, 0.0161)	(-0.0052, 0.0090)	1.0000	0.1257
700	(0.0062, 0.0225)	(-0.0042, 0.0086)	1.0000	<b>0.0356</b>
800	(0.0123, 0.0246)	(0.0020, 0.0124)	1.0000	<b>0.0231</b>
900	(0.0034, 0.0128)	(-0.0018, 0.0041)	1.0000	<b>3e-7</b>

#### 4.4 Synthetic Analysis of MRI Connectome SBM

Priebe et al. (2019) investigated clustering via the EM algorithm for the ASE and LSE of a 4-block SBM used to model neural connectivity in the human brain. They noted that the 2-dimensional LSE captures left-hemisphere/right-hemisphere connectivity and the 2-dimensional ASE captures gray-matter/white-matter connectivity. That is, clustering according to either embedding resulted in one of “two-truths.”

To compare clustering performance for this setting we considered the model  $\mathbf{A} \sim$

Table 9: Model 4 SBM. Negative med( $\Delta$ ) indicates superiority of ES over EM.

$n$	95% CI for med( $\Delta_A$ )	95% CI for med( $\Delta_L$ )	$p_A$	$p_L$
200	(0.0433, 0.0709)	(-0.0008, 0.0324)	1.0000	1.0000
300	(0.0724, 0.1082)	(-0.0131, 0.0150)	1.0000	1.0000
400	(0.0260, 0.0745)	(-0.0099, 0.0138)	1.0000	1.0000
500	(-0.0037, 0.0481)	(-0.0112, 0.0064)	1.0000	1.0000
600	(0.0095, 0.0527)	(-0.0128, 0.0006)	1.0000	0.9999
700	(0.0248, 0.0490)	(-0.0039, 0.0008)	1.0000	1.0000
800	(-0.0001, 0.0219)	(-0.0042, 0.0072)	1.0000	0.9999
900	(-0.0001, 0.0167)	(0.0001, 0.0102)	1.0000	0.9999

Table 10: Model 4 SBM,  $n > 900$

$n$	95% CI for med( $\Delta_A$ )	95% CI for med( $\Delta_L$ )	$p_A$	$p_L$
1000	(0.0075, 0.0207)	(0.0066, 0.0172)	1.0000	0.9959
1100	(0.0102, 0.0216)	(0.0070, 0.0164)	1.0000	0.8689
1200	(0.0102, 0.0215)	(0.0083, 0.0178)	1.0000	0.9849
1300	(0.0088, 0.0176)	(0.0089, 0.0162)	1.0000	0.4216
1400	(0.0035, 0.0103)	(0.0023, 0.0075)	1.0000	0.4856
1500	(0.0001, 0.0069)	(0.0001, 0.0058)	0.9999	0.6867
1600	(-0.00014, 0.0044)	(0.0010, 0.0044)	0.8484	<b>0.0129</b>
1700	(-0.0021, 0.0011)	(-0.0001, 0.0031)	<b>0.0211</b>	0.0745

SBM( $\mathbf{B}, \boldsymbol{\pi}$ ) where  $\boldsymbol{\pi} = (.28, 0.22, 0.28, 0.22)$ ,  $\mathbf{B}$  is the rank-4 matrix

$$\mathbf{B} = \begin{bmatrix} .020 & .044 & .002 & .009 \\ .044 & .115 & .010 & .042 \\ .002 & .010 & .020 & .045 \\ .009 & .042 & .045 & .117 \end{bmatrix},$$

and blocks 1–4 correspond to left/gray, left/white, right/gray, and right/white neurons, respectively. The eigendecomposition  $\mathbf{B} = \mathbf{U}_B \mathbf{D}_B \mathbf{U}_B^\top$  gives the latent position matrix

(after rotating properly to the first orthant in  $\mathbb{R}^4$ ) as

$$\mathbf{x} = \mathbf{U}_B \mathbf{D}_B^{\frac{1}{2}} \mathbf{U}_B^\top = \begin{bmatrix} \nu_1^\top \\ \nu_2^\top \\ \nu_3^\top \\ \nu_4^\top \end{bmatrix} = \begin{bmatrix} .0915 & 0.1076 & 0.0057 & 0.0034 \\ .1076 & 0.3149 & 0.0056 & 0.0649 \\ .0057 & 0.0056 & 0.0886 & 0.1099 \\ .0034 & 0.0649 & 0.1099 & 0.3173 \end{bmatrix}.$$

For each  $n = 500, 600, \dots, 1200$  we generated 100 graphs, computed their ASEs and LSEs, carried out the procedures as otherwise described, and output the results to table 9. We opted not to compare accuracy of parameter estimation, since we observed that our algorithms tended to vastly overestimate the covariances as described in the previous subsection. Here ES tended to more successfully cluster than EM for all values of  $n$ .

Table 11: Brain Connectome SBM

$n$	95% CI for med( $\Delta_A$ )	95% CI for med( $\Delta_L$ )
500	<b>(-0.0384, -0.0285)</b>	<b>(-0.0365, -0.02669)</b>
600	<b>(-0.0302, -0.0221)</b>	<b>(-0.0229, -0.0160)</b>
700	<b>(-0.0166, -0.0110)</b>	<b>(-0.0151, -0.0097)</b>
800	<b>(-0.0114, -0.0080)</b>	<b>(-0.0078, -0.0034)</b>
900	<b>(-0.0099, -0.0070)</b>	<b>(-0.0058, -0.0029)</b>
1000	<b>(-0.0068, -0.0042)</b>	<b>(-0.0050, -0.0023)</b>
1100	<b>(-0.0037, -0.0023)</b>	<b>(-0.0025, -0.0011)</b>
1200	<b>(-0.0033, -0.0011)</b>	<b>(-0.0021, -6e-6)</b>

## 5 Discussion

We have described an algorithm that estimates the parameters of a curved-normal mixture model and accounts for the components' curvature. The algorithm itself is an adaptation of the usual EM algorithm for smooth-normal mixture models, where instead of updating the component variance estimates in the usual way we simply plug the usual updates to the component means into the variance function(s). Even though we developed this algorithm



purely for the purpose of spectral clustering for SBMs, we suspect that we can easily adapt it to mixtures of curved exponential families in which the component distributions are not normal. We hope to explore this, as well as sufficient conditions for consistent and asymptotically normal estimates as outlined in the appendix of Elashoff and Ryan (2004), in a future paper.

The simulation results demonstrate that taking into account the curvature of the spectral embeddings’ limiting distributions allows us to improve node clustering for SBMs, and — in some settings — by a vast margin. The dominance our algorithms display in outperforming EM for the brain connectome model lead us to highly recommend our method for that application. Moreover, there existed at least one sample size for which at least one of our proposed algorithms clustered significantly more accurately than vanilla EM in all but one of the models from which we simulated. Since we initialized all simulations at the true parameter values, we have omitted any discussion of sufficient conditions for local or global convergence of  $ES \circ \{ASE, LSE\}$ . However, it was observed anecdotally in the brain connectome setting that initializing ES far from the canonical latent positions still resulted in accurate clusters.

Though the simulations are presented through the lens of evaluating ES against EM, we also compared both methods to  $K$ -means (Appendix). When all three algorithms were compared to each other, we determined that their clustering performance was approximately equal for the homogeneous balanced affinity models, and that EM and ES performed vastly better than  $K$ -means in the other settings. This corroborates the prior work done by Athreya et al. (2016) and Tang and Priebe (2018); since the canonical latent positions of a 2-dimensional full-rank homogeneous balanced affinity model lie equally spaced in the first quadrant with equal covariances, the  $K$ -means assumption of spherical covariances does not particularly hinder the clustering problem. As the other settings are far more varied in the spacing of their latent positions and the shape and orientation of their covariances, the flexibility afforded by EM and ES renders them far more effective than  $K$ -means. Moreover, the fact that ES eclipsed EM for the connectome SBM but not the core-periphery models suggests that ES is more useful for a wider range of models,

while EM’s superiority may be limited to that particular submodel.

The major drawback of our algorithms in the SBM setting is their seeming inability to accurately estimate the component variances. We noted that ES occasionally yielded values of  $\Delta^{-1}$  and  $\tilde{\Delta}^{-1}$  with diagonal entries far exceeding 1. A possible alteration to the algorithm that may reduce or eliminate this issue entirely would be to replace  $\Delta$ ,  $\boldsymbol{\mu}$ , and  $\tilde{\Delta}$  with

$$\begin{aligned}\hat{\Delta} &= \frac{\sum_{i=1}^n \hat{X}_i \hat{X}_i^\top}{n} \\ \hat{\boldsymbol{\mu}} &= \frac{\sum_{i=1}^n \hat{X}_i}{n} \\ \hat{\tilde{\Delta}} &= \frac{\sum_{i=1}^n \frac{\hat{X}_i \hat{X}_i^\top}{\hat{X}_i^\top \hat{\boldsymbol{\mu}}}}{n},\end{aligned}$$

respectively, since doing so holds those terms constant and would prevent slight perturbations to the component means from causing excessive inflation to the entries of the component variances.

Nonetheless, the strong evidence that spectral clustering based on ES can dominate that of EM opens up avenues for future research. During the writing of this article, Xie and Xu (2019) developed one-step estimators for the latent position matrices of both the ASE and LSE, the rows of which also asymptotically converge to curved normal mixtures. However, their mixture possesses component variances that are locally more efficient than those of the mixtures we have considered thus far; and they find that EM clustering based on this estimator can improve upon  $EM \circ \{ASE, LSE\}$ . In a future article we hope to compare performance of  $ES \circ \{ASE, LSE\}$  with EM for their one-step estimators, as well as implement an ES algorithm based on the latter.

## References

- Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic Control* 16(6), 716–723.
- Athreya, A., D. E. Fishkind, M. Tang, Y. Park, J. T. Vogelstein, K. Levin, V. Lyzinski,

- Y. Qin, and D. L. Sussman (2018). Statistical inference on random dot product graphs: A survey. *Journal of Machine Learning Research* 18, 1–92.
- Athreya, A., C. E. Priebe, M. Tang, D. J. Marchette, and D. L. Sussman (2016, February). A limit theorem for scaled eigenvectors of random dot product graphs. *Sankhya A* 78(1), 1–18.
- Bickel, P. J. and K. A. Doksum (2015). *Mathematical Statistics: Basic Ideas and Selected Topics* (2nd ed.), Volume 1. CRC Press.
- Cape, J., M. Tang, and C. E. Priebe (2019, September). On spectral embedding performance and elucidating network structure in stochastic block model graphs. *Network Science* 7(3), 269–291.
- Dempster, A. P., N. M. Laird, and D. B. Rubin (1977). Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society. Series B (Methodological)* 39(1), 1–38.
- Elashoff, M. and L. Ryan (2004, January). An em algorithm for estimating equations. *Journal of Computational and Graphical Statistics* 13(1), 48–65.
- Fishkind, D. E., D. L. Sussman, J. T. Vogelstein, and C. E. Priebe (2013). Consistent adjacency-spectral partitioning for the stochastic block model when the model parameters are unknown. *SIAM Journal on Matrix Analysis and Applications* 34, 23–39.
- Fraley, C. and A. E. Raftery (2002, June). Model-based clustering, discriminant analysis, and density estimation. *Journal of the American Statistical Association* 97(458), 611–631.
- Holland, P. W., K. B. Askey, and S. Leinhardt (1983). Stochastic blockmodels: First steps. *Social Networks* 5(2), 109–137.
- Hubert, L. and P. Arabie (1985). Comparing partitions. *Journal of Classification* 2(1), 193–218.

- Karrer, B. and M. E. J. Newman (2011). Stochastic blockmodels and community structure in networks. *Physical Review E* 83.
- Lyzinski, V., D. L. Sussman, M. Tang, A. Athreya, and C. E. Priebe (2014). Perfect clustering for stochastic blockmodel graphs via adjacency spectral embedding. *Electronic Journal of Statistics* 8, 2905–2922.
- Lyzinski, V., M. Tang, A. Athreya, Y. Park, and C. E. Priebe (2017). Community detection and classification in hierarchical stochastic blockmodels. *IEEE Transactions on Network Science and Engineering* 4(1), 13–26.
- Mann, H. B. and D. R. Whitney (1947). On a test of whether one of two random variables is stochastically larger than the other. *Annals of Mathematical Statistics* 18(1), 50–60.
- McLachlan, G. J. and T. Krishnan (2008). *The EM Algorithm and Extensions* (2nd ed.). Wiley-Interscience.
- Priebe, C. E., Y. Park, M. Tang, A. Athreya, V. Lyzinski, and J. T. Vogelstein (2017, May). Semiparametric spectral modeling of the *drosophila* connectome. *arXiv*.
- Priebe, C. E., Y. Park, J. T. Vogelstein, J. M. Conroy, V. Lyzinski, M. Tang, A. Athreya, J. Cape, and E. Bridgeford (2019). On a two-truths phenomenon in spectral graph clustering. In *Proceedings of the National Academy of Sciences of the United States of America*, Number 13 in 116, pp. 5995–6000.
- Rubin-Delanchy, P., J. Cape, M. Tang, and C. E. Priebe (2020, January). A statistical interpretation of spectral embedding: the generalised random dot product graph. *arXiv*.
- Schwarz, G. E. (1978). Estimating the dimension of a model. *The Annals of Statistics* 6(2), 464–464.
- Scrucca, L., M. Fop, T. B. Murphy, and A. E. Raftery (2016, August). mclust 5: Clustering, classification, and density estimation using Gaussian finite mixture models. *The R Journal* 8(1), 289–317.

Sussman, D. L., M. Tang, D. E. Fishkind, and C. E. Priebe (2012). A consistent adjacency spectral embedding for stochastic blockmodel graphs. *Journal of the American Statistical Association* 107(499), 1119–1128.

Tang, M. and C. E. Priebe (2018). Limit theorems for eigenvectors of the normalized laplacian for random graphs. *The Annals of Statistics* 46(5), 2360–2415.

Xie, F. and Y. Xu (2019). Efficient estimation for random dot product graphs via a one-step procedure. *arXiv*.

Young, S. J. and E. Scheinerman (2007). Random dot product graph models for social networks. In *Proceedings of the 5th international conference on algorithms and models for the web-graph*, pp. 138–149.

## Appendix: Comparison of EM and ES to $K$ -Means

In our simulations we also compared the clustering performance of EM and ES against that of  $K$ -means. As before we present the results in terms of 95% confidence intervals for the median difference in ARI, but due to the presence of multiple ties in the homogeneous balanced affinity settings we elected to use the SIGN test with a two-sided alternative. For  $\ell = A, L$  as in section 4 and  $Z = M, S$  we define

$$\Delta_{\ell}^{KZ} := ARI_{KM\circ\ell SE} - ARI_{EZ\circ\ell SE}.$$

As in Section 4, we embolden all entries indicating strict, significant improvement of our ES algorithms over  $K$ -means.

Table 12: Model 1 Mixture

$n$	95% CI for med( $\Delta_A^{KM}$ )	95% CI for med( $\Delta_L^{KM}$ )	95% CI for med( $\Delta_A^{KS}$ )	95% CI for med( $\Delta_L^{KS}$ )
200	(0.0167, 0.0493)	(0, 0.0220)	(0, 0.0048)	(0, 0.015)
300	(0, 0.0122)	(0, 0.0119)	(0, 0)	(0, 0)
400	(0, 0)	(0, 0)	(0, 0)	(0, 0)
500	(0, 0)	(0, 0)	(0, 0)	(0, 0)
600	(0, 0)	(0, 0)	(0, 0)	(0, 0)
700	(0, 0)	(0, 0)	(0, 0)	(0, 0)
800	(0, 0)	(0, 0)	(0, 0)	(0, 0)
900	(0, 0)	(0, 0)	(0, 0)	(0, 0)

Table 13: Model 2 Mixture

$n$	95% CI for med( $\Delta_A^{KM}$ )	95% CI for med( $\Delta_L^{KM}$ )	95% CI for med( $\Delta_A^{KS}$ )	95% CI for med( $\Delta_L^{KS}$ )
200	(0.0407, 0.0915)	(0.0139, 0.0439)	(0.0127, 0.0761)	(0.0140, 0.0774)
300	(0.0098, 0.0313)	(0.0061, 0.0197)	(1e-7, -0.0190)	(0, 0.0105)
400	(0.0131, 0.0168)	(-2e-7, 0.0106)	(-1e-7, 0.0023)	(0, 0.0023)
500	(-0.0067, 0.0134)	(-0.0066, 0.0068)	(0, 2e-7)	(0, 0.0066)
600	(-2e-8, 0.0059)	(0, 0.0059)	(0, 5e-9)	(0, 0.0018)
700	(0, 0.0053)	(0, 0.0051)	(0, 0)	(0, 0)
800	(0, 0.0046)	(0, 0.0046)	(0, 4e-8)	(0, 0)
900	(-2e-9, 0.0041)	(-7e-10, 0.0013)	(0, 0)	(0, 0)

Table 14: Model 3 Mixture

$n$	95% CI for med( $\Delta_A^{KM}$ )	95% CI for med( $\Delta_L^{KM}$ )	95% CI for med( $\Delta_A^{KS}$ )	95% CI for med( $\Delta_L^{KS}$ )
200	(-0.1070, -0.0537)	(-0.1450, -0.0964)	<b>(-0.0256, -8e-5)</b>	<b>(-0.026, -0.0080)</b>
300	(-0.1756, -0.1317)	(-0.2068, -0.1662)	<b>(-0.0998, -0.0318)</b>	<b>(-0.1273, -0.0888)</b>
400	(-0.2019, -0.1712)	(-0.2215, -0.1966)	<b>(-0.1682, -0.1204)</b>	<b>(-0.1762, -0.1242)</b>
500	(-0.2306, -0.2002)	(-0.2476, -0.2263)	<b>(-0.2288, -0.1986)</b>	<b>(-0.2370, -0.2103)</b>
600	(-0.2514, -0.2305)	(-0.2671, -0.2437)	<b>(-0.2547, -0.2296)</b>	<b>(-0.2656, -0.2434)</b>
700	(-0.2573, -0.2380)	(-0.2772, -0.2555)	<b>(-0.2657, -0.2386)</b>	<b>(-0.2755, -0.2577)</b>
800	(-0.2534, -0.2388)	(-0.2716, -0.2562)	<b>(-0.2619, -0.2370)</b>	<b>(-0.2760, -0.2574)</b>
900	(-0.2495, -0.2370)	(-0.2686, -0.2586)	<b>(-0.2528, -0.2426)</b>	<b>(-0.2722, 0.2577)</b>

Table 15: Model 4 Mixture

$n$	95% CI for med( $\Delta_A^{KM}$ )	95% CI for med( $\Delta_L^{KM}$ )	95% CI for med( $\Delta_A^{KS}$ )	95% CI for med( $\Delta_L^{KS}$ )
200	(-0.0543, 0.0101)	(-0.0777, -0.0244)	<b>(-0.0066, 0.0428)</b>	<b>(-0.0099, -0.0077)</b>
300	(-0.1207, -0.0883)	(-0.1306, -0.0908)	<b>(-0.0316, 0.1010)</b>	<b>(-0.0606, -0.0346)</b>
400	(-0.1704, -0.1367)	(-0.1772, -0.1520)	<b>(-0.0887, -0.0466)</b>	<b>(-0.1018, -0.0688)</b>
500	(-0.2116, -0.1890)	(-0.2191, -0.1912)	<b>(-0.1310, -0.943)</b>	<b>(-0.1502, -0.0961)</b>
600	(-0.2270, -0.1970)	(-0.2412, -0.2150)	<b>(-0.1992, -0.1556)</b>	<b>(-0.2252, -0.1858)</b>
700	(-0.2436, -0.2149)	(-0.2607, -0.2342)	<b>(-0.2345, -0.2051)</b>	<b>(-0.2505, -0.2301)</b>
800	(-0.2476, -0.2252)	(-0.2678, -0.2462)	<b>(-0.2420, -0.2214)</b>	<b>(-0.2625, -0.2399)</b>
900	(-0.2512, -0.2372)	(-0.2777, -0.2572)	<b>(-0.2540, -0.2363)</b>	<b>(-0.2785, -0.2611)</b>

Table 16: Model 1 SBM

$n$	95% CI for med( $\Delta_A^{KM}$ )	95% CI for med( $\Delta_L^{KM}$ )	95% CI for med( $\Delta_A^{KS}$ )	95% CI for med( $\Delta_L^{KS}$ )
200	(0.0148, 0.0530)	(7e-6, 0.0257)	(1e-5, 0.0148)	(0, 0.0026)
300	(0, 0.0227)	(0, 0.0113)	(0, 0)	(0, 0)
400	(0, 0.0088)	(0, 0.0086)	(0, 0)	(0, 0)
500	(0, 0)	(0, 3e-9)	(0, 0)	(0, 0)
600	(0, 0)	(0, 0.0020)	(0, 0)	(0, 0)
700	(0, 0)	(0, 0)	(0, 0)	(0, 0)
800	(0, 0)	(0, 0)	(0, 0)	(0, 0)
900	(0, 0)	(0, 0)	(0, 0)	(0, 0)

Table 17: Model 2 SBM

$n$	95% CI for med( $\Delta_A^{KM}$ )	95% CI for med( $\Delta_L^{KM}$ )	95% CI for med( $\Delta_A^{KS}$ )	95% CI for med( $\Delta_L^{KS}$ )
200	(0.0033, 0.0135)	(0.0010, 0.0109)	(0.0094, 0.0265)	(0, 0.0015)
300	(0.0235, 0.0510)	(0.0073, 0.0216)	(0.0131, 0.0830)	(0, 0.0008)
400	(0.0137, 0.0430)	(5e-6, 0.0186)	(-8e-7, 0.0065)	(0, 5e-6)
500	(0.0122, 0.0325)	(0.0038, 0.0115)	(0, 0.0054)	(-0.0014, 9e-7)
600	(-6e-6, 0.0098)	(1e-7, 0.0055)	(-1e-6, 8e-8)	(-0.0051, 7e-8)
700	(0, 0.0063)	(0, 0.0047)	(-4e-8, 2e-7)	(-0.0014, 2e-7)
800	(0, 0.0087)	(-1e-7, 0.0044)	(0, 0)	(-2e-9, 0)
900	(0, 0.0079)	(0, 0.0040)	(0, 0)	(0, 5e-9)

Table 18: Model 3 SBM

$n$	95% CI for med( $\Delta_A^{KM}$ )	95% CI for med( $\Delta_L^{KM}$ )	95% CI for med( $\Delta_A^{KS}$ )	95% CI for med( $\Delta_L^{KS}$ )
200	(-0.1061, -0.0559)	(-0.1834, -0.1487)	<b>(-0.1552, -0.0954)</b>	<b>(-0.1868, -0.1220)</b>
300	(-0.2065, -0.1188)	(-0.2767, -0.2429)	<b>(-0.2323, -0.1825)</b>	<b>(-0.2796, -0.2473)</b>
400	(-0.3298, -0.2676)	(-0.3666, -0.3380)	<b>(-0.3401, -0.2991)</b>	<b>(-0.3709, -0.3437)</b>
500	(-0.3872, -0.3664)	(-0.4409, -0.4039)	<b>(-0.3981, -0.3631)</b>	<b>(-0.4373, -0.4038)</b>
600	(-0.4284, -0.4022)	(-0.4673, -0.4512)	<b>(-0.4409, -0.4151)</b>	<b>(-0.4816, -0.4551)</b>
700	(-0.4900, -0.4617)	(-0.5192, -0.4958)	<b>(-0.4835, -0.4501)</b>	<b>(-0.5214, -0.4889)</b>
800	(-0.5236, -0.4878)	(-0.5515, -0.5234)	<b>(-0.5133, -0.4663)</b>	<b>(-0.5452, -0.5111)</b>
900	(-0.5145, -0.4678)	(-0.5365, -0.5034)	<b>(-0.4927, -0.4568)</b>	<b>(-0.5307, -0.4972)</b>

Table 19: Model 4 SBM

$n$	95% CI for med( $\Delta_A^{KM}$ )	95% CI for med( $\Delta_L^{KM}$ )	95% CI for med( $\Delta_A^{KS}$ )	95% CI for med( $\Delta_L^{KS}$ )
200	(-0.0781, -0.0312)	(-0.1298, -0.0814)	<b>(-0.0050, -5e-5)</b>	<b>(-0.1049, -0.0718)</b>
300	(-0.1356, -0.0636)	(-0.2158, -0.1760)	<b>(-0.0233, -0.0090)</b>	<b>(-0.2061, -0.1693)</b>
400	(-0.1975, -0.1219)	(-0.2891, -0.2402)	<b>(-0.0916, -0.0621)</b>	<b>(-0.2826, -0.2527)</b>
500	(-0.2809, -0.2366)	(-0.3505, -0.3271)	<b>(-0.2265, -0.1744)</b>	<b>(-0.3466, -0.3235)</b>
600	(-0.3477, -0.3134)	(-0.4117, -0.3841)	<b>(-0.3084, -0.2622)</b>	<b>(-0.4078, -0.3756)</b>
700	(-0.4164, -0.3799)	(-0.4611, -0.4339)	<b>(-0.3875, -0.3591)</b>	<b>(-0.4630, -0.4344)</b>
800	(-0.4417, -0.4144)	(-0.4962, -0.4755)	<b>(-0.4413, -0.4067)</b>	<b>(-0.5070, -0.4767)</b>
900	(-0.4970, -0.4754)	(-0.5504, -0.5230)	<b>(-0.4965, -0.4661)</b>	<b>(-0.5469, -0.5286)</b>

Table 20: Brain Connectome SBM

$n$	95% CI for med( $\Delta_A^{KM}$ )	95% CI for med( $\Delta_L^{KM}$ )	95% CI for med( $\Delta_A^{KS}$ )	95% CI for med( $\Delta_L^{KS}$ )
500	(-0.5238, -0.5044)	(-0.5679, -0.5493)	<b>(-0.5135, -0.4820)</b>	<b>(-0.5643, -0.5336)</b>
600	(-0.5514, -0.5200)	(-0.5909, -0.5672)	<b>(-0.5322, -0.5055)</b>	<b>(-0.5811, -0.5472)</b>
700	(-0.5676, -0.5432)	(-0.6148, -0.5929)	<b>(-0.5453, -0.5199)</b>	<b>(-0.5953, -0.5665)</b>
800	(-0.5430, -0.5032)	(-0.5907, -0.5580)	<b>(-0.5229, -0.4929)</b>	<b>(-0.5722, -0.5447)</b>
900	(-0.5442, -0.4975)	(-0.5824, -0.5468)	<b>(-0.5303, -0.4956)</b>	<b>(-0.5744, -0.5439)</b>
1000	(-0.5119, -0.4769)	(-0.5498, -0.5247)	<b>(-0.5029, -0.4747)</b>	<b>(-0.5461, -0.5212)</b>
1100	(-0.4952, -0.4709)	(-0.5363, -0.5170)	<b>(-0.4938, -0.4682)</b>	<b>(-0.5362, -0.5093)</b>
1200	(-0.4701, -0.4462)	(-0.5155, -0.4904)	<b>(-0.4719, -0.4489)</b>	<b>(-0.5123, -0.4914)</b>